

ITERATES OF QUANTUM OPERATIONS

JÓZSEF ZSOLT BERNÁD¹

ABSTRACT. Iterates of quantum operations and their convergence are investigated in the context of mean ergodic theory. We discuss in detail the convergence of the iterates and show that the uniform ergodic theorem plays an essential role. Our results will follow from some general theorems concerning completely positive maps, mean ergodic operators, and operator algebras on Hilbert spaces. A few examples of both finite and infinite dimensional Hilbert spaces are presented as well.

1. INTRODUCTION AND PRELIMINARIES

Quantum system are never perfectly closed and therefore interactions always take place with certain parts of the environment. This is the open quantum system concept, where we wish to neglect up to some extent the dynamics of the environment but follow the state changes of the central system. These processes of state changes are called quantum operations. Quantum operations in the Hilbert space formulation of quantum mechanics are obtained in the following way [1]: we start with an uncorrelated joint state of the central system and the environment; it is followed by a joint unitary evolution; and then an observer measures a property of the environment, described by a projective operations acting only on the environment. It has been shown by Kraus [2] that quantum operations are completely positive. They play an important role in the theory and applications of uniformly continuous completely positive dynamical semigroups [3, 4, 5]. Physical applications are present in various subfields of quantum information processing, like quantum computing [6] or quantum control theory [7], but also in questions related to the foundations of quantum mechanics [8].

In this paper we have the purpose to give a consistent approach to iterations of quantum operations. Fixed points of quantum operations has already been studied [9, 10] and in the case of finite dimensional Hilbert spaces the properties of the set of fixed points have also been discussed [11]. Here, we intend to characterize the convergence of the iterated operations and finally to investigate the dynamics on the asymptotic space. Our work takes advantage of the results in mean ergodic theory [12].

In the whole manuscript we consider a separable Hilbert space \mathcal{H} [13] with inner product $\langle \cdot, \cdot \rangle$ which is conjugate linear in the first and linear in the second variable. The norm $\langle x, x \rangle^{1/2}$ of any element x in \mathcal{H} will be denoted by $\|x\|$. The adjoint of a linear operator A on \mathcal{H} is the unique operator A^\dagger satisfying $\langle A^\dagger x, y \rangle = \langle x, Ay \rangle$ for all x, y in \mathcal{H} . The set of all bounded linear operators on \mathcal{H} is denoted by $\mathcal{B}(\mathcal{H})$. $\mathcal{B}(\mathcal{H})$ is a Banach space with respect to the operator norm

$$\|A\| = \sup\{\|Ax\| : x \in \mathcal{H}, \|x\| \leq 1\}.$$

Introducing the Banach space of trace class operators

$$\mathcal{B}_1(\mathcal{H}) = \{X \in \mathcal{B}(\mathcal{H}) : \|X\|_1 = \text{Tr}\{\sqrt{X^\dagger X}\} < \infty\}$$

the set of states or density operators is given by

$$\mathcal{D}(\mathcal{H}) = \{\rho \in \mathcal{B}_1(\mathcal{H}) : \rho \geq 0, \quad \text{Tr}\{\rho\} = 1\}.$$

Let us consider the n -dimensional Hilbert space \mathbb{C}^n . Operators of $\mathcal{B}(\mathcal{H} \otimes \mathbb{C}^n)$ are $n \times n$ matrices with entries A_{ij} acting on the Hilbert space \mathcal{H} for each $1 \leq i, j \leq n$. We define the linear operator $\phi^{(n)} : \mathcal{B}(\mathcal{H} \otimes \mathbb{C}^n) \rightarrow \mathcal{B}(\mathcal{H} \otimes \mathbb{C}^n)$ by

$$\phi^{(n)} \begin{pmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \ddots & \vdots \\ A_{n1} & \cdots & A_{nn} \end{pmatrix} \mapsto \begin{pmatrix} \phi(A_{11}) & \cdots & \phi(A_{1n}) \\ \vdots & \ddots & \vdots \\ \phi(A_{n1}) & \cdots & \phi(A_{nn}) \end{pmatrix}.$$

If $\phi^{(n)}$ is positive for all $n \geq 1$ then ϕ is a completely positive map. A quantum operation is a completely positive map due to its derivation, which includes an arbitrary ancilla Hilbert space [1]. Let ϕ be a trace non-increasing quantum operation, then there exist a family of operators $V_i \in \mathcal{B}(\mathcal{H})$ such that [2]

$$\sum_i V_i^\dagger V_i \leq 1$$

and

$$\phi(\rho) = \sum_i V_i \rho V_i^\dagger, \quad \forall \rho \in \mathcal{D}(\mathcal{H}). \quad (1.1)$$

The summation $\sum_i V_i^\dagger V_i$ is convergent with respect to the ultraweak topology on $\mathcal{B}(\mathcal{H})$ generated by the seminorms

$$\mathcal{B}(\mathcal{H}) \rightarrow \mathbb{R}_+, \quad X \rightarrow |\text{Tr}\{XA\}| \quad \text{with } A \in \mathcal{B}_1(\mathcal{H}).$$

The proof can be found in Lemma 2.1 of Ref. [2].

Any $X \in \mathcal{B}_1(\mathcal{H})$ can be written as

$$X/\|X\|_1 = \rho_1 - \rho_2 + i(\rho_3 - \rho_4)$$

with four density operators $\rho_i \in \mathcal{D}(\mathcal{H})$ ($i \in 1, 2, 3, 4$). Therefore, ϕ can be extended to a linear mapping of $\mathcal{B}_1(\mathcal{H})$ into itself and the extension, denoted by ϕ again, is defined by

$$\phi(X) = \sum_i V_i X V_i^\dagger, \quad \forall X \in \mathcal{B}_1(\mathcal{H}).$$

Every bounded linear functional on $\mathcal{B}_1(\mathcal{H})$ is of the form $X \rightarrow \text{Tr}\{XA\}$ with $A \in \mathcal{B}(\mathcal{H})$. This establishes a one-to-one correspondence preserving linearity and the norm between $\mathcal{B}(\mathcal{H})$ and the dual of $\mathcal{B}_1(\mathcal{H})$ [14]. Thus, $\mathcal{B}(\mathcal{H})$ is isometrically isomorphic to the dual of $\mathcal{B}_1(\mathcal{H})$. Hence, the adjoint map ϕ^* to ϕ of $\mathcal{B}_1(\mathcal{H})$ is defined for arbitrary $A \in \mathcal{B}(\mathcal{H})$ by

$$\text{Tr}\{\phi(X)A\} = \text{Tr}\{X\phi^*(A)\},$$

where

$$\phi^*(A) = \sum_i V_i^\dagger A V_i. \quad (1.2)$$

ϕ^* is also a completely positive map [2].

The purpose of this paper is to investigate the behavior of the iterates ϕ^n and $(\phi^*)^n$ with the help of mean ergodic theorems. These theorems describe the averages of the iterates and throw some light on the fix points of ϕ and ϕ^* . This is the subject of Sec. 2. It will be shown in Sec. 3 that this approach plays an important role in the asymptotic properties of the iterates for large n . As our problem is formulated on Banach spaces of linear operators on general separable Hilbert spaces we need to use several deep results from the theory of operators.

2. MEAN ERGODIC THEOREMS

Let us fix some notations. $T : \mathcal{B} \rightarrow \mathcal{B}$ is called power bounded operator on a Banach space \mathcal{B} if the norms of the powers T^n ($n \geq 0$) are uniformly bounded.

Proposition 2.1. *The two completely positive maps $\phi : \mathcal{B}_1(\mathcal{H}) \rightarrow \mathcal{B}_1(\mathcal{H})$ and $\phi^* : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ are power bounded.*

Proof. Since the matrix

$$\begin{pmatrix} I & X \\ X^\dagger & X^\dagger X \end{pmatrix} = \begin{pmatrix} I & 0 \\ X^\dagger & 0 \end{pmatrix} \begin{pmatrix} I & X \\ 0 & 0 \end{pmatrix} \geq 0, \quad \forall X \in \mathcal{B}(\mathcal{H})$$

is positive with I being the identity operator on \mathcal{H} and $\phi^{*(2)}$ is a positive map, i.e., complete positivity of ϕ^* implies that ϕ^* is also a 2-positive map,

$$\phi^*(X^\dagger)\phi^*(X) \leq \phi^*(I)\phi^*(X^\dagger X) \leq \|\phi^*(I)\|\phi^*(X^\dagger X), \quad (2.1)$$

where we have used $\|\phi^*(I)\|I \geq \phi^*(I)$, because $\phi^*(I)$ is a positive operator. The statement in Eq. (2.1) is well known, for more details see Refs. [15, 16].

From Corollary 1 of Ref. [17] we know that the operator norm $\|\cdot\|_{\mathcal{B}(\mathcal{H})}$ of ϕ^* can be obtained in the following way

$$\|\phi^*\|_{\mathcal{B}(\mathcal{H})} = \sup_{U \in \mathcal{B}(\mathcal{H})} \|\phi^*(U)\|$$

where supremum is over the unitary operators. The property $\|XX^\dagger\| = \|X\|^2$ of the norm of $\mathcal{B}(\mathcal{H})$ yields

$$\|\phi^*(U)\|^2 = \|[\phi^*(U)]^\dagger \phi^*(U)\| = \|\phi^*(U^\dagger)\phi^*(U)\|,$$

and by the inequality (2.1)

$$\|\phi^*(U)\|^2 \leq \|\phi^*(I)\|\|\phi^*(U^\dagger U)\| = \|\phi^*(I)\|^2 = \left\| \sum_i V_i^\dagger V_i \right\|^2 \leq 1.$$

Hence, $\|\phi^*\|_{\mathcal{B}(\mathcal{H})} \leq 1$, i.e., ϕ^* is a contraction. Contractions are obviously power-bounded.

It remains to be shown that ϕ is also a power bounded map. Since

$$\|X\|_1 = \sup_{\|A\|=1} |\text{Tr}\{AX\}|, \quad X \in \mathcal{B}_1(\mathcal{H}) \quad \text{and} \quad A \in \mathcal{B}(\mathcal{H}),$$

we get

$$\begin{aligned} \|\phi(X)\|_1 &= \sup_{\|A\|=1} |\text{Tr}\{A\phi(X)\}| = \sup_{\|A\|=1} |\text{Tr}\{\phi^*(A)X\}| \leq \sup_{\|A\|=1} \{\|\phi^*(A)\|\|X\|_1\} \\ &\leq \|X\|_1, \end{aligned}$$

which yields $\|\phi\|_{\mathcal{B}_1(\mathcal{H})} \leq 1$ and by thus ϕ is also a contraction. \square

The supporting references of the above proof are considering general C^* -algebras and in our case we deal only with special C^* -algebras. In the subsequent part we discuss very general mean ergodic theorems for a power bounded operator $T : \mathcal{B} \rightarrow \mathcal{B}$ on a Banach space \mathcal{B} . We are interested in the convergence of the averages $A_n(T) = n^{-1} \sum_{i=1}^n T^i$, Cesàro mean of the first n iterates of T . We introduce the following two closed linear subspaces [12]

$$\mathcal{B}_{\text{me}}(T) = \{x \in \mathcal{B} : \lim A_n(T)x \text{ exists}\},$$

and fixed space of T

$$F(T) = \{x \in \mathcal{B} : Tx = x\} = \text{Ker}(I - T).$$

Let \mathcal{B}' be the space of all continuous functionals on \mathcal{B} . \mathcal{B}' is called the dual space of \mathcal{B} and it is a Banach space. $T^* : \mathcal{B}' \rightarrow \mathcal{B}'$ is the adjoint operator of T defined by $f(Tx) = (T^*f)(x)$ for $f \in \mathcal{B}'$ and $x \in \mathcal{B}$. Then we have the following splitting theorem [18]

Theorem 2.2. *Let T be a power bounded linear operator on a Banach space \mathcal{B} . Then*

$$\mathcal{B}_{\text{me}}(T) = F(T) \oplus \overline{\text{Rng}(I - T)}.$$

The linear operator $\mathcal{P}x = \lim A_n(T)x$ assigned to $x \in \mathcal{B}_{\text{me}}(T)$ is the projection of $\mathcal{B}_{\text{me}}(T)$ onto $F(T)$. We have $\mathcal{P} = \mathcal{P}^2 = T\mathcal{P} = \mathcal{P}T$ and for any $y \in \mathcal{B}$ the assertions

- a) $\lim A_n(T)y = 0$
- b) $y \in \overline{\text{Rng}(I - T)}$
- c) $f(y) = 0$ for all $f \in \{g \in \mathcal{B}' : T^*g = g\}$

are equivalent.

Proof. The theorem is mostly due to K. Yosida [18] and proved in [12], see Theorem 1.3. \square

T is called mean ergodic if $\mathcal{B}_{\text{me}}(T) = \mathcal{B}$. Power bounded linear operators on a reflexive Banach space \mathcal{B} , the canonical embedding map from \mathcal{B} into the double dual \mathcal{B}'' is surjective, are mean ergodic [19]. The approach of K. Yosida is based on weakly compact linear operators, which map the unit ball of the Banach space on a weakly compact set [18]. The reflexivity of the Banach space comes again in picture, because every bounded linear operator defined on a reflexive Banach space is weakly compact. However, the mean ergodic property is not necessarily related only to the reflexivity of Banach spaces [20]. As both $\mathcal{B}_1(\mathcal{H})$ and $\mathcal{B}(\mathcal{H})$ are not reflexive in general, therefore the maps ϕ and ϕ^* are not always mean ergodic.

The set of Hilbert-Schmidt operators

$$\mathcal{B}_2(\mathcal{H}) = \{X \in \mathcal{B}(\mathcal{H}) : \|X\|_2 = \text{Tr}\{X^\dagger X\} < \infty\}$$

is a Banach space and with the scalar product $\langle X, Y \rangle_{\text{HS}} = \text{Tr}\{X^\dagger Y\}$ becomes a Hilbert space, which means that it is a reflexive space. In fact, $\mathcal{B}_2(\mathcal{H})$ is a two

sided *-closed ideal in $\mathcal{B}(\mathcal{H})$ and

$$\mathcal{B}_1(\mathcal{H}) \subseteq \mathcal{B}_2(\mathcal{H}) \subseteq \mathcal{B}(\mathcal{H}). \quad (2.2)$$

Corollary 2.3. *If the summation in (1.2) is finite then ϕ_2^* , the restriction of ϕ^* to $\mathcal{B}_2(\mathcal{H})$, is mean ergodic.*

Proof. ϕ^* is a power bounded map and ϕ_2^* inherits this property. By $\mathcal{B}_2(\mathcal{H})$ being a two sided ideal in $\mathcal{B}(\mathcal{H})$

$$AX, XA \in \mathcal{B}_2(\mathcal{H}), \quad \forall A \in \mathcal{B}(\mathcal{H}) \quad \text{and} \quad \forall X \in \mathcal{B}_2(\mathcal{H}).$$

Hence for any $X \in \mathcal{B}_2(\mathcal{H})$

$$\phi_2^*(X) = \sum_i V_i^\dagger X V_i \in \mathcal{B}_2(\mathcal{H})$$

where we have used that the summation is finite. Therefore, $\text{Rng}(\phi_2^*) \subseteq \mathcal{B}_2(\mathcal{H})$. In case of infinite summation the set $\text{Rng}(\phi_2^*)$ can be larger than $\mathcal{B}_2(\mathcal{H})$, because $\sum_i V_i^\dagger V_i$ converges only in the ultraweak topology.

$\mathcal{B}_2(\mathcal{H})$ is a Hilbert space, hence is reflexive. Thus, the properties of ϕ_2^* together with the theorem of Lorch [19] implies mean ergodicity. \square

Remark. It is interesting to note that if the summation in (1.2) is finite then the adjoint of ϕ_2^* , let us simply call ϕ_2 , is

$$\phi_2(X) = \sum_i V_i X V_i^\dagger,$$

where we have used the properties of the inner product $\langle X, Y \rangle_{\text{HS}}$. Thus, the map in (1.1) involving only finite summation can also be extended to the space of Hilbert-Schmidt operators.

Corollary 2.4. *If the summation in (1.2) is finite then $F(\phi_2^*) = F(\phi_2)$.*

Proof. Due to the finite summation both ϕ_2^* and ϕ_2 are contractions on $\mathcal{B}_2(\mathcal{H})$. Take now $X \in F(\phi_2^*)$ which implies

$$\langle X, \phi_2(X) \rangle_{\text{HS}} = \langle \phi_2^*(X), X \rangle_{\text{HS}} = \|X\|_2^2.$$

Since ϕ_2 is a contraction, it follows that

$$\begin{aligned} \|\phi_2(X) - X\|_2^2 &= \|\phi_2(X)\|_2^2 - 2\text{Re}\langle X, \phi_2(X) \rangle_{\text{HS}} + \|X\|_2^2 \\ &= \|\phi_2(X)\|_2^2 - \|X\|_2^2 \leq 0. \end{aligned}$$

Thus, $\phi_2(X) = X$ or $X \in F(\phi_2)$. Consequently, $F(\phi_2^*) \subseteq F(\phi_2)$. The opposite inclusion is proven in the same fashion, but starting with $X \in F(\phi_2)$. \square

Now let us give examples of fixed spaces.

Example 2.5. Let $\mathcal{H} = \mathbb{C}^2$ with its standard basis $e_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $e_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. We consider the following two Pauli matrices

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix},$$

and the quantum operation

$$\begin{aligned}\phi(X) &= V_1 X V_1^\dagger + V_2 X V_2^\dagger, \\ V_1 &= \sqrt{p}\sigma_x, \quad V_2 = \sqrt{1-p}\sigma_y,\end{aligned}$$

where $p \in [0, 1]$ and $X \in \mathcal{M}_2(\mathbb{C})$, i.e., the set of all $n \times n$ matrices over \mathbb{C} . Note that $\phi^* = \phi$.

If $p = 0$ then

$$F(\phi) = \left\{ \begin{pmatrix} a & -b \\ b & a \end{pmatrix}, \quad \forall a, b \in \mathbb{C} \right\}.$$

If $p \in (0, 1)$ then

$$F(\phi) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}, \quad \forall a \in \mathbb{C} \right\}.$$

If $p = 1$ then

$$F(\phi) = \left\{ \begin{pmatrix} a & b \\ b & a \end{pmatrix}, \quad \forall a, b \in \mathbb{C} \right\}.$$

Example 2.6. Let $\mathcal{H} = \ell^2$, the space of square-summable sequences. The left and right shift operators are defined by

$$\begin{aligned}S_L &: (a_1, a_2, a_3, a_4 \dots) \mapsto (a_2, a_3, a_4, a_5 \dots), \\ S_R &: (a_1, a_2, a_3, a_4 \dots) \mapsto (0, a_1, a_2, a_3, \dots),\end{aligned}$$

and using these contractions we construct the following quantum operation

$$\begin{aligned}\phi(X) &= V_1 X V_1^\dagger + V_2 X V_2^\dagger, \\ V_1 &= \sqrt{p}S_L, \quad V_2 = \sqrt{1-p}S_R, \quad p \in (0, 1),\end{aligned}$$

where $X \in \mathcal{B}_1(\ell^2)$. It is immediate that ϕ is a contraction and

$$\sum_{i=1,2} V_i^\dagger V_i = pS_L^\dagger S_L + (1-p)S_R^\dagger S_R < I$$

with I being the identity operator on ℓ^2 . We consider first X to be

$$X : (a_1, a_2, a_3, a_4 \dots) \mapsto (x_1 a_1, x_2 a_2, x_3 a_3, \dots) \quad \text{with} \quad \sum_{i=1}^{\infty} |x_i| < \infty$$

and p has a fixed value. Then $\phi(X) = X$ has the following solution

$$x_i = x_1 \frac{f^{(i)}(p)}{p^{i-1}}, \quad i > 1,$$

where

$$\begin{aligned}f^{(i)}(p) &= \sum_{j=0}^{i-1} (-1)^j p^j a_j^{(i)}, \\ a_j^{(i)} &= \begin{cases} 1, & j = 0, \\ a_j^{(i-1)} + a_{j-1}^{(i-1)}, & 0 < j < i-1, \\ 1, & j = i-1 \quad \text{and} \quad i \text{ is odd}, \\ 0, & j = i-1 \quad \text{and} \quad i \text{ is even.} \end{cases}\end{aligned}$$

We find that $f^{(i)}(p)/p^{i-1} > 1$ and one obtains by induction the following relation

$$\frac{f^{(i+1)}(p)}{p^i} - \frac{f^{(i)}(p)}{p^{i-1}} = \frac{(1-p)^i}{p^i} > 0, \quad i > 1.$$

It is immediate that the solution to $\phi(X) = X$ is not a trace class operator, and even more so is not bounded. Extending the presented approach to more general X operators, we find

$$F(\phi) = \{0\},$$

here 0 denotes the null operator.

Remark. Elements of $\mathcal{D}(\mathcal{H})$ are subject to both averages $A_n(\phi)$ and $A_n(\phi^*)$ due to $\mathcal{D}(\mathcal{H}) \subset \mathcal{B}_1(\mathcal{H}) \subseteq \mathcal{B}(\mathcal{H})$. However, there might be cases when

$$\begin{aligned} \mathcal{D}(\mathcal{H}) \cap F(\phi) &= \emptyset, \\ \mathcal{D}(\mathcal{H}) \cap F(\phi^*) &= \emptyset. \end{aligned}$$

If $\sum_i V_i^\dagger V_i = 1$ then

$$\{X \in \mathcal{B}_1(\mathcal{H}) : A_i X = X A_i \text{ for all } i\} \subseteq F(\phi)$$

and

$$\{X \in \mathcal{B}(\mathcal{H}) : A_i X = X A_i \text{ for all } i\} \subseteq F(\phi^*).$$

For more details, see Ref. [10].

We shall now discuss a particular property of completely positive maps. As the following argumentation is the same for both ϕ and ϕ^* , we consider the case of ϕ^* . The following result is known and we formulate it in the context discussed here in this manuscript.

Lemma 2.7. *Let p_i be strictly positive numbers with $\sum_i p_i = 1$ and let ϕ_i^* be commuting completely positive maps. If $\phi^* = \sum_i p_i \phi_i^*$ then*

$$F(\phi^*) = \bigcap_i F(\phi_i^*).$$

Proof. The proof is mostly due to A. Brunel [21] and M. Falkowitz [22]. An elegant way of proving it is to use that the identity is an extreme point in the convex set consisting of all contractions, an application of Krein-Milman Theorem. For further details, see Lemma 1.14 in [12]. \square

Mean ergodic theorems enables us to understand the convergence of the averages $A_n(T)$ and the fixed space of T . In the subsequent section we show how these averages are used in evaluations of iterates ϕ^n and $(\phi^*)^n$.

3. MAIN RESULT

In this chapter we are concerned with the iterates ϕ^n and $(\phi^*)^n$. In order to characterize them we need the following slightly modified theorem of K. Yosida and S. Kakutani in [23].

Theorem 3.1. *Let T be a power bounded linear operator on a Banach space \mathcal{B} and λ any complex number with $|\lambda| = 1$. Then $T_\lambda = T/\lambda$ is a power bounded operator such that*

- a) $A_n(T_\lambda)$ converges strongly to \mathcal{P}_λ , the projection of $\mathcal{B}_{\text{me}}(T_\lambda)$ onto $F(T_\lambda)$,
- b) $\mathcal{P}_\lambda = \mathcal{P}_\lambda^2 = T_\lambda \mathcal{P}_\lambda = \mathcal{P}_\lambda T_\lambda$,
- c) $\lambda \neq \mu$ implies $F(T_\lambda) \cap F(T_\mu) = \{0\}$,
- d) $\mathcal{P}_\lambda \neq 0$ iff λ is an eigenvalue of T .

Proof. It is immediate from the relation

$$\sup_k \|T^k\|_{\text{op}} = \sup_k \|T_\lambda^k\|_{\text{op}}$$

that T_λ is power bounded, where $\|\cdot\|_{\text{op}}$ is the operator norm. The strong convergence

$$\lim_{n \rightarrow \infty} \|A_n(T_\lambda)x - \mathcal{P}_\lambda x\| = 0, \quad \forall x \in \mathcal{B}_{\text{me}}(T_\lambda)$$

with the norm $\|\cdot\|$ of \mathcal{B} and the relations

$$\mathcal{P}_\lambda = \mathcal{P}_\lambda^2 = T_\lambda \mathcal{P}_\lambda = \mathcal{P}_\lambda T_\lambda$$

are clear from Theorem 2.2. Furthermore, we also have

$$\mathcal{P}_\lambda T = T \mathcal{P}_\lambda = \lambda \mathcal{P}_\lambda. \quad (3.1)$$

Then, we have by a)

$$\lim_{n \rightarrow \infty} \|A_n(T_\lambda) \mathcal{P}_\mu x - \mathcal{P}_\lambda \mathcal{P}_\mu x\| = 0, \quad \forall x \in \mathcal{B}_{\text{me}}(T_\lambda) \cap \mathcal{B}_{\text{me}}(T_\mu),$$

and

$$A_n(T_\lambda) \mathcal{P}_\mu = \frac{1}{n} \left(\frac{T}{\lambda} + \frac{T^2}{\lambda^2} + \cdots + \frac{T^n}{\lambda^n} \right) \mathcal{P}_\mu. \quad (3.2)$$

Putting 3.1 and 3.2 together, we obtain that

$$\frac{1}{n} \left(\frac{\mu}{\lambda} + \frac{\mu^2}{\lambda^2} + \cdots + \frac{\mu^n}{\lambda^n} \right) \mathcal{P}_\mu$$

converges to $\mathcal{P}_\lambda \mathcal{P}_\mu$. When $\lambda \neq \mu$

$$\frac{1}{n} \left(\frac{\mu}{\lambda} + \frac{\mu^2}{\lambda^2} + \cdots + \frac{\mu^n}{\lambda^n} \right) \mathcal{P}_\mu = \frac{1}{n} \frac{1 - \left(\frac{\mu}{\lambda}\right)^{n+1}}{1 - \frac{\mu}{\lambda}} \mathcal{P}_\mu$$

converges uniformly to the null operator. Thus,

$$\mathcal{P}_\lambda x = 0, \quad \forall x \in F(T_\mu) \cap \mathcal{B}_{\text{me}}(T_\lambda).$$

Hence, the statement in c) is proven. Since \mathcal{P}_λ is projecting onto

$$F(T_\lambda) = \text{Ker}(I - T/\lambda) = \text{Ker}(\lambda I - T)$$

and $\text{Ker}(\lambda I - T) \neq \{0\}$ when λ is an eigenvalue, we arrive at the statement in d). \square

We introduce the following two sets

$$\sigma(T) = \{\lambda \in \mathbb{C} : \text{Ker}(\lambda I - T) \neq \{0\}, |\lambda| = 1\} \quad (3.3)$$

and

$$\mathcal{B}(T) = \bigcap_{\lambda \in \sigma(T)} \mathcal{B}_{\text{me}}(T_\lambda). \quad (3.4)$$

After all these preparations we return to the iterates of ϕ and ϕ^* . The next result is essentially a consequence of Proposition 2.1 and Theorem 3.1.

Corollary 3.2. *Let $\{P_\lambda\}_{\lambda \in \sigma(\phi)}$ be the set of projectors associated according to Theorem 3.1 with $\phi : \mathcal{B}_1(\mathcal{H}) \rightarrow \mathcal{B}_1(\mathcal{H})$ and similarly $\{Q_\lambda\}_{\lambda \in \sigma(\phi^*)}$ with $\phi^* : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$. Set*

$$S_\phi(X) = \phi(X) - \sum_{\lambda \in \sigma(\phi)} \lambda P_\lambda(X) \quad \forall X \in \mathcal{B}(\phi) \subseteq \mathcal{B}_1(\mathcal{H}),$$

$$S_{\phi^*}(X) = \phi^*(X) - \sum_{\lambda \in \sigma(\phi^*)} \lambda Q_\lambda(X) \quad \forall X \in \mathcal{B}(\phi^*) \subseteq \mathcal{B}(\mathcal{H}),$$

then the iterates of ϕ and ϕ^* are given by the formulas

$$\phi^n(X) = \sum_{\lambda \in \sigma(\phi)} \lambda^n P_\lambda(X) + S_\phi^n(X) \quad \forall X \in \mathcal{B}(\phi),$$

$$(\phi^*)^n(X) = \sum_{\lambda \in \sigma(\phi^*)} \lambda^n Q_\lambda(X) + S_{\phi^*}^n(X) \quad \forall X \in \mathcal{B}(\phi^*)$$

for $n \geq 1$. λ is an eigenvalue of S_ϕ (S_{ϕ^*}) with eigenvector in $\mathcal{B}(\phi)$ ($\mathcal{B}(\phi^*)$) iff is an eigenvalue of ϕ (ϕ^*) and $\lambda \notin \sigma(\phi)$ ($\lambda \notin \sigma(\phi^*)$).

Proof. We consider only the case of ϕ , because the proof for ϕ^* is exactly along the same lines. By Theorem 3.1 we have for all $X \in \mathcal{B}(\phi)$

$$P_\lambda S_\phi(X) = P_\lambda \left(\phi(X) - \sum_{\lambda \in \sigma(\phi)} \lambda P_\lambda(X) \right) = \lambda P_\lambda(X) - \lambda P_\lambda(X) = 0$$

and similarly $S_\phi P_\lambda(X) = 0$ for all $\lambda \in \sigma(\phi)$. It is immediate also that

$$\phi S_\phi(X) = S_\phi \phi(X) = S_\phi^2(X).$$

Then, for all $X \in \mathcal{B}(\phi)$

$$\begin{aligned} \phi^2(X) &= \left(\sum_{\lambda \in \sigma(\phi)} \lambda P_\lambda + S_\phi \right)^2 (X) \\ &= \left(\sum_{\lambda \in \sigma(\phi)} \lambda P_\lambda \right)^2 (X) + S_\phi^2(X), \end{aligned}$$

but according to Theorem 3.1 $P_\lambda P_\mu(X) = 0$ whenever $\lambda \neq \mu$, which yields

$$\phi^2(X) = \sum_{\lambda \in \sigma(\phi)} \lambda^2 P_\lambda(X) + S_\phi^2(X).$$

By induction we obtain the required result for all powers of n . To prove the last statement, let $\lambda \neq 0$ be such an eigenvalue of ϕ with eigenvector $X_\lambda \neq 0$ that

$$\phi(X_\lambda) = \lambda X_\lambda \quad X_\lambda \in \mathcal{B}(\phi)$$

and $\lambda \notin \sigma(\phi)$. Then, for all $\mu \in \sigma(\phi)$

$$\mu P_\mu(X_\lambda) = P_\mu \phi(X_\lambda) = \lambda P_\mu(X_\lambda)$$

and since $\mu \neq \lambda$, we obtain $P_\mu(X_\lambda) = 0$. Therefore,

$$\lambda X_\lambda = \mu P_\mu(X_\lambda) = \sum_{\mu \in \sigma(\phi)} \lambda P_\mu(X_\lambda) + S_\phi(X_\lambda) = S_\phi(X_\lambda).$$

Hence λ is an eigenvalue of S_ϕ . Conversely, let $\lambda \neq 0$ be such an eigenvalue of S_ϕ with eigenvector $X_\lambda \neq 0$ that

$$S_\phi(X_\lambda) = \lambda X_\lambda \quad X_\lambda \in \mathcal{B}(\phi).$$

We have already shown that $S_\phi \phi(X) = S_\phi^2(X)$ and using this relation we obtain

$$\phi(X_\lambda) = \phi \left(\frac{S_\phi}{\lambda} \right) (X_\lambda) = \frac{1}{\lambda} S_\phi^2(X_\lambda) = \lambda X_\lambda,$$

λ is also an eigenvalue of ϕ . In order to show that $\lambda \notin \sigma(\phi)$, we assume that there exists a $\mu \in \sigma(\phi)$ such that $\lambda = \mu$. Since

$$A_n(\phi/\mu)(X_\lambda) = \frac{1}{n} \left(\frac{\phi}{\mu} + \frac{\phi^2}{\mu^2} + \cdots + \frac{\phi^n}{\mu^n} \right) (X_\lambda) = X_\lambda \quad n \geq 1$$

and the limit $n \rightarrow \infty$ with the help of Theorem 2.2 results $P_\mu(X_\lambda) = X_\lambda$. It is a contradiction, because

$$P_\mu(X_\lambda) = P_\mu \left(\frac{S_\phi}{\lambda} \right) (X_\lambda) = 0$$

where we have used the already proved relation $P_\mu S_\phi = 0$. Thus, $\lambda \notin \sigma(\phi)$. \square

Remark. It is important to note that the sets $\mathcal{B}(\phi)$ and $\mathcal{B}(\phi^*)$ play an essential role in the above result. Therefore, it is natural to ask under which conditions $\mathcal{B}(\phi) = \mathcal{B}_1(\mathcal{H})$ and $\mathcal{B}(\phi^*) = \mathcal{B}(\mathcal{H})$. This is the case, when ϕ and ϕ^* are mean ergodic. Let us see a few examples: $\mathcal{B}_1(\mathcal{H})$ and $\mathcal{B}(\mathcal{H})$ are reflexive Banach spaces [19]; ϕ and ϕ^* are weakly compact operators [18]; $\mathcal{B}_1(\mathcal{H})$ and $\mathcal{B}(\mathcal{H})$ are separable and quasi-reflexive Banach spaces of order one [20]. In general, the mean ergodicity of both ϕ and ϕ^* may be not simultaneously fulfilled.

Corollary 3.3. *If the summation in (1.2) is finite then the iterates of $\phi_2^* : \mathcal{B}_2(\mathcal{H}) \rightarrow \mathcal{B}_2(\mathcal{H})$ and its adjoint map ϕ_2 are given by the formulas*

$$\begin{aligned} (\phi_2^*)^n(X) &= \sum_{\lambda \in \sigma(\phi_2^*)} \lambda^n Q_{\lambda,2}(X) + S_{\phi_2^*}^n(X) \quad \forall X \in \mathcal{B}_2(\mathcal{H}) \\ \phi_2^n(X) &= \sum_{\lambda \in \sigma(\phi_2)} \lambda^n P_{\lambda,2}(X) + S_{\phi_2}^n(X) \quad \forall X \in \mathcal{B}_2(\mathcal{H}). \end{aligned}$$

On the sets

$$\bigcup_{\lambda \in \sigma(\phi_2^*)} F(\phi_2^*/\lambda) \quad \text{and} \quad \bigcup_{\lambda \in \sigma(\phi_2)} F(\phi_2/\lambda)$$

ϕ_2^* and respectively ϕ_2 restrict to unitary operators.

Proof. This is immediate from Theorem 3.1 and Corollary 3.2. One has to use the following facts: $\mathcal{B}_2(\mathcal{H})$ is a Hilbert space; $F(T_\lambda) = \text{Ker}(\lambda I - T)$; ϕ_2^* and ϕ_2 are mean ergodic. \square

If

$$\bigcup_{\lambda \in \sigma(\phi_2^*)} F(\phi_2^*/\lambda) = \bigcup_{\lambda \in \sigma(\phi_2)} F(\phi_2/\lambda)$$

then $\phi_2^*\phi_2 = \phi_2\phi_2^* = I$. This result is similar to the findings of Szókefalvi-Nagy and Foias, a contraction on a Hilbert space defines a decomposition of the Hilbert space into two parts, where on one of them the contraction acts as a unitary operator [24]. A special example: \mathcal{H} is finite dimensional, which also means $\mathcal{B}_1(\mathcal{H}) = \mathcal{B}_2(\mathcal{H}) = \mathcal{B}(\mathcal{H})$.

In order to investigate the limit $n \rightarrow \infty$ of the formulas obtained in Corollary 3.2 one has to determine the spectrum of ϕ or ϕ^* . It is of interest that $\|S_\phi\|_{\mathcal{B}_1(\mathcal{H})} < 1$ and $\|S_{\phi^*}\|_{\mathcal{B}(\mathcal{H})} < 1$, because then the asymptotic space can be identified through the projectors $\{P_\lambda\}_{\lambda \in \sigma(\phi)}$ and $\{Q_\lambda\}_{\lambda \in \sigma(\phi^*)}$. An obvious choice is that ϕ^* and ϕ are compact operators, the image of the unit ball under the map is relatively compact, because the spectrum of a compact operator contains the cluster point $\{0\}$ and only eigenvalues, which form the point spectrum. According to Theorem 3.1 an eigenvalue λ with $|\lambda| = 1$ is not an eigenvalue of either S_ϕ or S_{ϕ^*} . As both ϕ^* and ϕ are contractions, their spectrum is contained in the closed unit disc, we have

$$\lim_{n \rightarrow \infty} S_\phi^n = \lim_{n \rightarrow \infty} S_{\phi^*}^n = 0.$$

It turns out that one can make an even more general statement. But first we have to introduce the concept of quasi-compact operator. An operator T on a Banach space \mathcal{B} is quasi-compact if there exists an integer n and a compact operator V with $\|T^n - V\|_{\mathcal{B}} < 1$. This leads to the uniform ergodic theory of K. Yosida and S. Kakutani, which is the consequence of applying Theorem 3.1 to quasi-compact operators [23]. Thus, we are able to state the following result.

Corollary 3.4. *Let ϕ and ϕ^* be quasi-compact operators. Then, there exists constants $\epsilon_\phi, \epsilon_{\phi^*} > 0$ and $M_\phi, M_{\phi^*} > 0$ such that*

$$\|S_\phi^n\|_{\mathcal{B}_1(\mathcal{H})} \leq \frac{M_\phi}{(1 + \epsilon_\phi)^n} \quad \text{and} \quad \|S_{\phi^*}^n\|_{\mathcal{B}(\mathcal{H})} \leq \frac{M_{\phi^*}}{(1 + \epsilon_{\phi^*})^n}.$$

Proof. As the proof can be found in [23] we show only the cornerstones of the argumentation. Quasi-compactness implies weakly compact property of ϕ and ϕ^* , which means that they are mean ergodic. Furthermore, the ranges of the projectors $\{P_\lambda\}_{\lambda \in \sigma(\phi)}$ and $\{Q_\lambda\}_{\lambda \in \sigma(\phi^*)}$ are finite. Therefore, both S_ϕ and S_{ϕ^*} are quasi-compact and the unit circle belongs to their resolvent set, which is equivalent to the existence of $\epsilon_\phi, \epsilon_{\phi^*} > 0$ and $M_\phi, M_{\phi^*} > 0$, which fulfill the relations of the statement. \square

Remark. It may happen that both ϕ and ϕ^* can not be simultaneously quasi-compact operators. The question, under which conditions is a completely positive map compact or quasi-compact, is left open for now. Although the uniform ergodic theory is an old result, there are recent developments with respect to the iterates of quasi-compact operators, see for example [25].

Example 3.5. Let us reconsider the quantum operation in Example 2.6. $\mathcal{M}_2(\mathbb{C})$ with the scalar product $\langle X, Y \rangle_{\text{HS}} = \text{Tr}\{X^\dagger Y\}$ is a Hilbert space. The orthonormal

basis is chosen to be

$$\begin{aligned} X_1 &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} / \sqrt{2}, & X_2 &= \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} / \sqrt{2}, \\ X_3 &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} / \sqrt{2}, & X_4 &= \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} / \sqrt{2}. \end{aligned}$$

Then

$$\phi(X) = \lambda_1 P_{\lambda_1}(X) + \lambda_2 P_{\lambda_2}(X) + S_\phi(X),$$

where

$$\begin{aligned} \lambda_1 &= 1 & \text{with } P_{\lambda_1}(X) &= \text{Tr}\{X_1^\dagger X\} X_1, \\ \lambda_2 &= -1 & \text{with } P_{\lambda_2}(X) &= \text{Tr}\{X_2^\dagger X\} X_2, \\ S_\phi(X) &= (2p-1)\text{Tr}\{X_3^\dagger X\} X_3 + (1-2p)\text{Tr}\{X_4^\dagger X\} X_4. \end{aligned}$$

It is immediate

$$\phi^n(X) = P_{\lambda_1}(X) + (-1)^n P_{\lambda_2}(X) + (2p-1)^n \text{Tr}\{X_3^\dagger X\} X_3 + (1-2p)^n \text{Tr}\{X_4^\dagger X\} X_4$$

and

$$\|S_\phi^n\|_{\mathcal{M}_2(\mathbb{C})} \leq |1-2p|^n,$$

where $\|\cdot\|_{\mathcal{M}_2(\mathbb{C})}$ is the operator norm on $\mathcal{M}_2(\mathbb{C})$. When $p \in (0, 1)$ we have the following relations $|1-2p| < 1$ and $S_\phi^n \rightarrow 0$ as $n \rightarrow \infty$.

Example 3.6. Let \mathcal{H} be a separable Hilbert space with orthonormal basis e_n ($n = 0, 1, 2, \dots$). We define the operators a and a^\dagger by

$$ae_n = \sqrt{n}e_{n-1}, \quad a^\dagger e_n = \sqrt{n+1}e_{n+1}.$$

For the sake of simplicity we set $|n\rangle = e_n$ and then a typical vector x has the unique expansion $x = \sum_{i=0}^{\infty} x_n |n\rangle$. The orthonormal basis e'_n ($n = 0, 1, 2, \dots$) of the dual of \mathcal{H} has the property $e'_n(e_m) = 0$ when $n \neq m$ and $e'_n(e_n) = 1$. We define $\langle n| = e'_n$. The operator a^\dagger is the adjoint of a , because

$$\langle a^\dagger x, y \rangle = \sum_{n=0}^{\infty} \sqrt{n+1} x_n^* y_{n+1} = \langle x, ay \rangle.$$

a^\dagger and a are the creation and annihilation operators of bosons. Furthermore, they are unbounded operators and have the same domain of definition, which is dense in \mathcal{H} . We consider the following quantum operation

$$\begin{aligned} \phi(X) &= V_1 X V_1^\dagger + V_2 X V_2^\dagger, \\ V_1 &= \sqrt{p}I, \quad V_2 = \sqrt{1-pe^{-i\pi a^\dagger a}}, \quad p \in (0, 1), \end{aligned}$$

where $X \in \mathcal{B}_1(\mathcal{H})$ and I is the identity map. Although a^\dagger and a are unbounded, V_2 is a bounded operator with spectral radius one.

As there are only two terms $V_1 \cdot V_1^\dagger$ and $V_2 \cdot V_2^\dagger$ in ϕ , we can extend the quantum operation to $\mathcal{B}_2(\mathcal{H})$. Now, with the help of the Hilbert-Schmidt scalar product we define the following orthogonal projections

$$P_{n,m}(X) = \text{Tr}\{|m\rangle\langle n|X\}|n\rangle\langle m| \quad X \in \mathcal{B}_2(\mathcal{H})$$

with $n, m = 0, 1, 2, \dots$. Then

$$\begin{aligned} \phi^n(X) &= \sum_{n=0}^{\infty} P_{n,n}(X) + \sum_{k=1}^{\infty} \sum_{n=0}^{\infty} \left[P_{n+2k,n}(X) + P_{n,n+2k}(X) \right] \\ &+ (2p-1) \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \left[P_{n+2k+1,n}(X) + P_{n,n+2k+1}(X) \right]. \end{aligned}$$

Thus,

$$S_\phi(X) = (2p-1) \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \left[P_{n+2k+1,n}(X) + P_{n,n+2k+1}(X) \right]$$

and

$$\|S_\phi^n\|_{\mathcal{B}_2(\mathcal{H})} \leq |1-2p|^n,$$

where $\|\cdot\|_{\mathcal{B}_2(\mathcal{H})}$ is the operator norm on $\mathcal{B}_2(\mathcal{H})$.

In conclusion, we have investigated iterations of quantum operations in the context of mean ergodic theorems. In the case of finite dimensional Hilbert spaces the iterations and their convergence are simply an application of the uniform ergodic theorem of K. Yosida and S. Kakutani. The operator in the limit $n \rightarrow \infty$ is unitary on the asymptotic space. As long as the uniform ergodic theorem can not be applied one has to investigate the spectrum of quantum operation together with the mean ergodic property.

ACKNOWLEDGEMENT

This work is supported by the European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 732894 (FET Proactive HOT).

REFERENCES

1. K.-E. Hellwig and K. Kraus, *Commun. Math. Phys.* **11**, 214 (1969).
2. K. Kraus, *Ann. Phys.* **64**, 311 (1971).
3. V. Gorini, A. Kossakowski and E. C. G. Sudarshan, *J. Math. Phys.* **17**, 821 (1976).
4. G. Lindblad, *Comm. Math. Phys.* **48**, 119 (1976).
5. H. Spohn, *Rev. Mod. Phys.* **52**, 569 (1980).
6. M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2000).
7. H. M. Wiseman and G. J. Milburn, *Quantum Measurement and Control* (Cambridge University Press, Cambridge, UK, 2010).
8. E. Joos, H. D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer-Verlag, Berlin, 1996).
9. O. Bratteli, P. Jørgensen, A. Kishimoto and R. Werner, *J. Operator Theory* **43**, 97 (2000).
10. A. Arias, A. Gheondea, and S. Gudder, *J. Math. Phys.* **43**, 5872 (2002).
11. J. Novotný, G. Alber, and I. Jex, *J. Phys. A: Math. Theor.* **45**, 485301 (2012).
12. U. Krengel, *Ergodic Theorems* (de Gruyter, Berlin, 1985).
13. It is not necessary that the Hilbert space is separable, because properties of quantum operations are not related to the separability criteria, see pg. 318 in Ref. [2]. Furthermore, relations between different operator spaces are true for all Hilbert spaces [14]. However, there are two reasons to not consider nonseparable Hilbert spaces. First, quantum mechanics and its applications are built around separable Hilbert spaces which is due to the history of the theory. Second, Gleason's Theorem in *J. Math. and Mech.* **6**, 885 (1957), which

characterizes all states on a separable Hilbert space, cannot be generalized to nonseparable Hilbert spaces without assuming the continuum hypothesis, see Eilers–Horst Theorem in *Int. J. Theor. Phys.* **13**, 419 (1975). The consequences of this interesting statement are beyond the scope of the theory.

14. R. Schatten, *Norm Ideals of Completely Continuous Operators* (Springer-Verlag, Berlin, 1970).
15. E. Størmer, *Lecture Notes in Physics* **29** (Springer-Verlag, Berlin, 1974), pp. 85–106.
16. V. Paulsen, *Completely Bounded Maps and Operator Algebras* (Cambridge University Press, Cambridge, UK, 2003).
17. B. Russo and H. A. Dye, *Duke Math. J.* **33**, 413 (1966).
18. K. Yosida, *Proc. Imp. Acad. Tokyo* **14**, 292 (1938).
19. E. R. Lorch, *Bull. Amer. Math. Soc.* **45**, 945 (1939).
20. V. P. Fonf, M. Lin, and P. Wojtaszczyk, *Israel J. Math.* **179**, 479 (2010).
21. A. Brunel, *Lecture notes*, Springer-Verlag, t. 160, 7-17 (1970).
22. M. Falkowitz, *PAMS* **38**, 553 (1973).
23. K. Yosida and S. Kakutani, *Ann. Math.* **42**, 188 (1941).
24. B. Sz-Nagy and C. Foiaş, *Acta Sci. Math. Szeged* **21**, 251 (1960).
25. J. Nagler, arXiv: 1706.00663.

¹DEPARTMENT OF PHYSICS, UNIVERSITY OF MALTA, MSIDA MSD 2080, MALTA
E-mail address: `zsolt.bernad@um.edu.mt`