

Optimal robust estimators for families of distributions on the integers*

Ricardo A. Maronna^{1†} and Victor J. Yohai²

¹Universidad Nacional de La Plata

²Universidad de Buenos Aires and CONICET

Abstract

Let F_θ be a family of distributions with support on the set of nonnegative integers Z_0 . In this paper we derive the M-estimators with smallest gross error sensitivity (GES). We start by defining the uniform median of a distribution F with support on Z_0 ($\text{umed}(F)$) as the median of $x + u$, where x and u are independent variables with distributions F and uniform in $[-0.5, 0.5]$ respectively. Under some general conditions we prove that the estimator with smallest GES satisfies $\text{umed}(F_n) = \text{umed}(F_\theta)$, where F_n is the empirical distribution. The asymptotic distribution of these estimators is found. This distribution is normal except when there is a positive integer k so that $F_\theta(k) = 0.5$. In this last case, the asymptotic distribution behaves as normal at each side of 0, but with different variances. A simulation Monte Carlo study compares, for the Poisson distribution, the efficiency and robustness for finite sample sizes of this estimator with those of other robust estimators.

Keywords: Gross-error sensitivity, uniform median, contamination bias.

1 Introduction

Consider a one-parameter family of distributions F_θ . An important problem in the theory of robust estimation is the study of estimators which in some sense optimize their bias under contamination. The *gross-error sensitivity* (GES) is defined as the maximum of the absolute values of the influence function. It gives an approximation to the maximum bias produced by an outlier contamination of rate ε , when ε is "small".

***Acknowledgement.** This research was partially supported by Grants X-094 and 20020170100022BA from Universidad de Buenos Aires, PID 5505 from CONICET and PAV 120 and PICT 21407 from ANPCYT, Argentina.

[†]Corresponding author. E-mail rmaronna@retina.ar

Hampel (1974) dealt with M-estimators defined as solutions of equations of the form

$$\sum_{i=1}^n \psi(x_i, \theta) = 0,$$

and considered the problem of minimizing the asymptotic variance among Fisher-consistent M-estimators which satisfy a bound on the GES. Alternatively, this problem can be stated as minimizing the GES under a bound on the asymptotic variance. Details are given in Section 3.1.

In this paper we consider minimizing the GES without any restrictions on the asymptotic variance. Let F_θ be a family of continuous distributions with densities $p(x, \theta)$ and score function

$$\psi_0(x, \theta) = \frac{\partial \log p(x, \theta)}{\partial \theta}. \quad (1)$$

Maronna et al. (2019, p. 68) show that if $\psi_0(x, \theta)$ is strictly monotone on x , the M-estimator with smallest GES is obtained by solving

$$\text{med}(F_n) = \text{med}(F_{\hat{\theta}}), \quad (2)$$

where “med” stands for the median and F_n denotes the empirical distribution.

This result does not hold when F_θ has support on a discrete set, such as the integers, since $\text{med}(F_\theta)$ is in general not uniquely defined and therefore (2) does not identify θ .

To overcome this problem, we introduce in Section 2 the concept of the *uniform median* of a distribution F with support on the set of nonnegative integers Z_0 . In Section 3 we prove that the estimator with smallest GES can be obtained by solving an equation similar to (2) but replacing the median by the uniform median. Section 4 deals with the asymptotic distribution of this estimator. Section 5 shows an application to the family of Poisson distributions. Finally Section 6 is an appendix containing proofs of the main results.

2 The uniform median

We shall deal with distributions concentrated on a finite or infinite interval of the integers, such as the Poisson distribution. To avoid notational complications it will henceforth be assumed that this interval is the set Z_0 of the nonnegative integers. Let F be a distribution with support on Z_0 and call p the corresponding probability density. The *uniform median* of F ($\text{umed}(F)$) is defined as the median of the distribution that distributes the mass $p(k)$ uniformly on the interval $[k - 0.5, k + 0.5]$. This is equivalent to define $\text{umed}(F)$ as the median of $x + u$ where x and u are independent with distributions F and uniform in $[-0.5, 0.5]$ respectively. To give an explicit formula for $\text{umed}(F)$, define

$$k_0(F) = \min\{k : F(k) \geq 0.5\}. \quad (3)$$

It is easy to verify that

$$\text{umed}(F) = k_0(F) - 0.5 + \frac{0.5 - F(k_0(F) - 1)}{p_0}, \quad (4)$$

where

$$p_0 = p(k_0(F)). \quad (5)$$

Note that by definition $p_0 > 0$, and therefore $\text{umed}(F)$ is well defined.

Ma et al. (2011) define quantiles of discrete distributions based on what they call “mid-distribution functions”. In the case of the median, their definition is similar to that of the umedian but is not exactly equal, nor can any of the two medians be expressed as a function of the other one.

The following property is immediate

$$k_0(F) - 0.5 < \text{umed}(F) \leq k_0(F) + 0.5.$$

Therefore for two distributions F_1 and F_2 we have

$$\text{umed}(F_1) = \text{umed}(F_2) \Rightarrow k_0(F_1) = k_0(F_2). \quad (6)$$

Recall that the median of F is the solution μ of

$$\mathbb{E}_F \text{sign}(x - \mu) = 0.$$

Similarly $\text{umed}(F)$ can be defined as the solution μ of

$$\mathbb{E} \psi_{0.5}^H(x - \mu) = 0,$$

where ψ_m^H is the Huber family of score functions given by

$$\psi_m^H(x) = \max(\min(x, m), -m), \quad (7)$$

Note that $\psi_{0.5}^H(x) = 0.5 \text{sign}(x)$ for $|x| \geq 0.5$ and $\psi_{0.5}^H(x) = x$ for $|x| < 0.5$.

3 Estimators with smallest GES

3.1 The Hampel approach

Before dealing with the unrestricted minimization of the GES we need to consider Hampel’s (1974) approach to robust optimality. These optimal estimators are defined by minimizing the asymptotic variance among the class of M-estimators which are Fisher-consistent and have their GES bounded by a given constant.

If x_1, \dots, x_n is a random sample from F_θ , the Hampel-optimal estimator $\hat{\theta}$ is defined as the solution of the estimating equation

$$\sum_{i=1}^n \psi_m^H[\psi_0(x_i, \hat{\theta}) - c(m, \hat{\theta})] = 0, \quad (8)$$

where m depends on the given bound on the GES, ψ_0 and ψ_m^H are defined in (1) and (7), respectively, and $c(m, \theta)$ is defined by

$$E_{\theta} \psi_m^H(\psi_0(x_i, \theta) - c(m, \theta)) = 0, \quad (9)$$

where F_{θ} denotes the expectation with respect to F_{θ} .

The *dual* Hampel problem consists of minimizing the GES under a bound V on the asymptotic variance. It is known that the solution has again the form (8), where m is a decreasing function of V .

3.2 The optimal estimator

Let F_{θ} be a family of distribution functions with $\theta \in \Theta \subset R$, with support on Z_0 and probability densities $p(x, \theta)$. Given a random sample x_1, \dots, x_n of F_{θ} , denote by F_n the corresponding empirical distribution function, with density

$$p_n(k) = \frac{\#\{i : x_i = k\}}{n}.$$

Then we have:

Theorem 1 *Assume that $\psi_0(x, \theta)$ is continuous and strictly monotone in x and θ . Then the estimator with smallest GES is $\hat{\theta}_n$ defined by*

$$umed(F_n) = umed(F_{\hat{\theta}_n}). \quad (10)$$

The proof is given in Section 6.1

Before showing the existence and uniqueness of the solution to (10) we state some properties of the uniform median.

It will henceforth be assumed that

A1 $F_{\theta}(x)$ is a decreasing function of θ for all x

A2 $F_{\theta}(x)$ is a continuous function of θ for all x

A3 Call Θ the range of θ (e.g., $[0, \infty)$ for the Poisson distribution) and let $\theta_1 = \inf(\Theta)$ and $\theta_2 = \sup(\Theta)$. Then $\lim_{\theta \rightarrow \theta_1} F_{\theta}(x) = 0$ for $x \neq 0$ and $\lim_{\theta \rightarrow \theta_2} F_{\theta}(x) = 0$ for all x

These assumptions are satisfied by the standard discrete families such as the Poisson family. A1 implies that the family is “stochastically increasing” in the sense that $\theta_1 > \theta_2$ implies that F_{θ_1} is strictly stochastically larger than F_{θ_2} .

Lemma 1 *Let $F^{(n)}$, $n \geq 1$ be a sequence of distributions with support on Z_0 . Then $F^{(n)} \rightarrow_w F$ (where \rightarrow_w denotes weak convergence) implies that $umed(F^{(n)}) \rightarrow umed(F)$.*

The proof is given in Section 6.2.

Corollary 1 *Let x_1, \dots, x_n be i.i.d random variables with distribution F with support at Z_0 and call F_n the empirical distribution. Then $\text{umed}(F_n) \rightarrow \text{umed}(F)$ a.s..*

Proof: The result follows from Lemma 1 and the fact that $F_n \rightarrow_w F$ by the Glivenko-Cantelli Theorem.

Lemma 2 *Put for brevity $g(\theta) = \text{umed}(F_\theta)$. If A1-A2-A3 hold, then $g(\theta)$ is a continuous increasing function of θ , and $\lim_{\theta \rightarrow \theta_1} g(\theta) = 0$ and $\lim_{\theta \rightarrow \theta_2} g(\theta) = \infty$.*

Proof: A1 implies that g is increasing; its continuity follows from A2 and Lemma 1, and A3 implies the last statement.

Corollary 2 *(10) has a unique solution.*

4 Asymptotics

4.1 Consistency.

The following result proves the strong consistency of the optimal estimator $\hat{\theta}_n$ defined by (10).

Theorem 2 *Let F_θ satisfy A1-A2-A3 and let x_1, \dots, x_n be i.i.d. random variables with distribution F_θ . Then $\hat{\theta}_n \rightarrow \theta$ a.s.*

Proof. Call F_n the empirical distribution. and put $g(\theta) = \text{umed}(F_\theta)$. Since $\hat{\theta}_n = g^{-1}(\text{umed}(F_n))$, and g^{-1} is continuous, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \hat{\theta}_n &= g^{-1}(\lim_{n \rightarrow \infty} \text{umed}(F_n)) = g^{-1}(\lim_{n \rightarrow \infty} \text{umed}(F_\theta)). \\ &= g^{-1}(g(\theta)) = \theta \text{ a.s..} \end{aligned}$$

This proves the Theorem.

4.2 Asymptotic distribution

The following Theorem states that when $F(k_0(F)) > 0.5$, $\text{umed}(F_n)$ is asymptotically normal, while when $F(k_0(F)) = 0.5$, its left and right tails are asymptotically normal but with different variances.

Theorem 3 *Let x_1, \dots, x_n be i.i.d. random variables with distribution F with support at Z_0 , p its probability density and F_n the empirical distribution. Put for brevity*

$$K = k_0(F), \quad p_0 = p(K), \quad F^1 = F(K-1) \quad \text{and} \quad Z_n = n^{1/2}(\text{umed}(F_n) - \text{umed}(F)).$$

Then,

(a) If $F(K) > 0.5$, then

$$Z_n \rightarrow_d N(0, \sigma^2) \quad (11)$$

where \rightarrow_d stands for convergence in distribution and

$$\sigma^2 = \frac{0.25}{p_0^3} (4F^1 (F^1 - 1 + p_0) - p_0 + 1). \quad (12)$$

(b) If $F(K) = 0.5$, then

$$Z_n \rightarrow_d H$$

with

$$H(t) = \begin{cases} \Phi(2tp(K)) & \text{if } a \leq 0 \\ \Phi(2tp(K+1)) & \text{if } a > 0 \end{cases}, \quad (13)$$

where Φ is the standard normal distribution function.

The proof is given in Section 6.3.

Remark: Note that the phenomenon of an asymptotic distribution with two different normal tails also occurs with the ordinary median of samples from a continuous distribution if the density has different side derivatives at the population median.

The following Theorem deals with the asymptotic distribution of the GES-optimal estimator.

Theorem 4 *Let x_1, \dots, x_n be a random sample of F_θ . Assume that the density $p(x, \theta)$ is continuously differentiable in θ . Let $\hat{\theta}_n$ be the estimator defined by (10). Let*

$$g(\theta) = k_0(F_\theta) - 0.5 + \frac{0.5 - F_\theta(k_0(F_\theta) - 1)}{p(k_0(F_\theta), \theta)}. \quad (14)$$

(a) If $F_\theta(k_0(F_\theta)) > 0.5$, then

$$n^{1/2}(\hat{\theta}_n - \theta) \rightarrow_d N\left(0, \frac{\sigma^2}{g'(\theta)^2}\right)$$

with σ defined in (12).

(b) If $F(k_0(F)) = 0.5$ then.

$$n^{1/2}(\hat{\theta}_n - \theta) \rightarrow_d G,$$

where

$$G(t) = \begin{cases} \Phi(2g'_- tp(K, \theta)) & \text{if } a \leq 0 \\ \Phi(2g'_+ tp(K+1, \theta)) & \text{if } a > 0 \end{cases},$$

with

$$g'_+ = \frac{\partial}{\partial t} \left[\frac{0.5 - F_t(K)}{p(K+1, \theta)} \right]_{t=\theta+} \quad \text{and} \quad g'_- = \frac{\partial}{\partial t} \left[\frac{0.5 - F_t(K-1)}{p(K, \theta)} \right]_{t=\theta-}. \quad (15)$$

The proof is given in Section 6.4.

Table 1: Maximum asymptotic biases for the Poisson family

ε	λ	Optimal	MT	Q-L
0.1	5	0.329	0.421	0.409
	10	0.511	0.608	0.627
	20	0.823	0.985	0.959
0.2	5	0.805	1.087	1.071
	10	1.052	1.413	1.402
	20	1.569	2.057	2.071

Table 2: Asymptotic efficiencies of estimators

λ	Optimal	MT	Q-L
5	0.72	0.89	0.93
10	0.69	0.93	0.96
20	0.67	0.93	0.97

5 Application to the Poisson distribution

In this section we compute the maximum asymptotic bias of the GES-optimal estimator and compare it to those of two robust estimators: the MT estimator of (Valdora and Yohai 1974) and the Quasi-Likelihood (Q-L) estimator of (Cantoni and Ronchetti 2001). Note that the GES is only a rough measure of the bias, and therefore minimizing the GES does not ensure any optimal properties of the actual bias.

For the Poisson distribution with parameter λ we contaminate the data with a point mass with probability ε located at x_0 . For each estimator $\hat{\lambda}$ the absolute bias $|\hat{\lambda} - \lambda|$ is computed, and the maximum over all x_0 is reported in Table 1.

Table 2 gives the asymptotic efficiencies of the three estimators.

These results show that the optimal estimator has a comparatively good bias behavior not only for “small” ε . At the same time, the price for such a low bias is a relatively low efficiency.

To study the small sample behavior of the estimator a simulation with $n = 20$ and 50 was run, with 500 replications. Table 3 shows the estimator’s efficiencies, which are seen to differ little from the asymptotic ones.

To study the estimators’ robustness, in each sample a proportion ε of the values was replaced by a value x_0 , and the estimators’ means squared error (MSE) was computed. This was done for $x_0 = 0, 1, \dots, 3\lambda$. Table 4 displays the maximum MSE of each estimator over all x_0 ’s, for $\varepsilon = 0.1$ and 0.2.

Table 3: Finite sample efficiencies of estimators

n	λ	Optimal	MT	Q-L
20	5	0.80	0.95	0.98
	10	0.64	0.87	0.93
	20	0.56	0.83	0.89
50	5	0.72	0.89	0.94
	10	0.71	0.92	0.96
	20	0.66	0.91	0.94

Table 4: Maximum MSEs of estimators for contamination ϵ

n	ϵ	λ	Optimal	MT	Q-L
20	0.1	5	0.67	0.54	0.52
		10	1.14	1.10	1.04
		20	2.53	2.22	2.32
	0.2	5	1.22	1.72	1.61
		10	2.26	2.81	2.82
		20	4.63	5.81	5.85
	0.1	5	0.30	0.35	0.34
		10	0.68	0.64	0.70
		20	1.40	1.27	1.40
50	0.2	5	0.84	1.29	1.27
		10	1.61	2.29	2.33
		20	3.36	4.92	4.79

The MSEs of the Optimal estimator are slightly higher than the other two for $\varepsilon = 0.1$, and lower for $\varepsilon = 0.2$. This can be explained by the fact that for small ε , the bias is less important than the variability, and viceversa.

6 Appendix: Proofs of results

6.1 Proof of Theorem 1

For the purposes of this proof it will be more convenient to state Hampel's problem in its equivalent *dual* form, namely, to minimize the GES under a bound K on the asymptotic variance. It is known that the solution is again given by (8), where now m is a decreasing function of the bound K .

For given m call $\hat{\theta}_m$ the Hampel-optimal estimator given by (8). In this case (9) takes on the form

$$\sum_{k=0}^{\infty} \psi_m^H(\psi_0(k, \hat{\theta}_m) - c(m, \hat{\theta}_m)) p(k, \hat{\theta}_m) = 0. \quad (16)$$

When the bound K tends to infinity, $m \rightarrow 0$ and the GES of $\hat{\theta}_m$ tends to its lower bound. Then to prove the Theorem it is enough to show that there exists m_0 such that for $m \leq m_0$, this estimator coincides with the estimator given by (10).

We will suppose that $\psi_0(k, \theta)$ is strictly increasing in k . The proof when it is strictly decreasing is similar. Put $k_0^*(\theta) = k_0(F_\theta)$ and let

$$m_0 = \frac{1}{2} \min(\psi_0(k_0^*(\theta), \theta) - \psi_0(k_0^*(\theta) - 1, \theta), \psi_0(k_0^*(\theta) + 1, \theta) - \psi_0(k_0^*(\theta), \theta)). \quad (17)$$

It will be shown that if $m \leq m_0$ then

$$\psi_0(k_0^*(\theta), \theta) - m_0 \leq c(m, \theta) \leq \psi_0(k_0^*(\theta), \theta) + m_0. \quad (18)$$

Suppose that $c(m, \theta) < \psi_0(k_0^*(\theta), \theta) - m_0$. Then we have $\psi_0(k, \theta) - c(m, \theta) > m_0$ for all $k \geq k_0^*(\theta)$, and hence

$$\sum_{k=k_0^*(\theta)}^{\infty} \psi_m^H(\psi_0(k, \theta) - c(m, \theta)) p(k, \theta) = m(1 - F_0(k_0^*(\theta) - 1)) \geq \frac{m}{2}.$$

We also have

$$\sum_{k=0}^{k_0^*-1} \psi_m^H(\psi_0(k, \theta) - c(m, \theta)) p(k, \theta) \geq -m_0 F_0(k_0^*(\theta) - 1) > -m/2,$$

which implies

$$\sum_{k=0}^{\infty} \psi_m^H(\psi_0(k, \theta) - c(m, \theta))p(k, \theta) > 0,$$

contradicting (16), Similarly it can be proved that we can not have $c(m, \theta) > \psi_0(k_0^*(\theta), \theta) - m_0$.

Then from (18) and (17) we get that $\psi_m^H(\psi_0(k, \theta) - c(m, \theta)) \leq -m$ for $k < k_0^*(\theta)$ and $\psi_m^H(\psi_0(k, \theta) - c(m, \theta)) \geq m$ for $k > k_0^*(\theta)$. Then it follows from (16) that

$$-mF(k_0^*(\theta) - 1, \theta) + m(1 - F(k_0^*(\theta), \theta)) + (\psi_0(k_0^*(\theta), \theta) - c(m, \theta))p(k_0^*(\theta), \theta) = 0, \quad (19)$$

or similarly

$$m \frac{(1 - 2F(k_0^*(\theta), \theta)) + p(k_0^*(\theta), \theta)}{p(k_0^*(\theta), \theta)} = \psi_m^H(\psi_0(k_0^*(\theta), \theta) - c(m, \theta)). \quad (20)$$

From (20) we derive

$$k_0^*(\theta) - 0.5 + \frac{(0.5 - F(k_0^*(\theta), \theta))}{p(k_0^*(\theta), \theta)} = k_0^*(\theta) - 1 + \frac{1}{2m} \psi_m^H(\psi_0(k_0^*(\theta), \theta) - c(m, \theta)). \quad (21)$$

Define

$$G_m(\kappa, \theta) = \kappa - 1 + \frac{2}{2m} \psi_m^H(\psi_0(\kappa, \theta) - c(m, \theta)),$$

and note that according to (21) for $m \leq m_0$, G_m does not depend on m . Then (21) and (4) imply that for $m \leq m_0$

$$\text{umed}(F_\theta) = G_m(k_0^*(\theta), \theta). \quad (22)$$

If $m \leq m_0$, (8) is equivalent to

$$-mF_n(k_0^*(\hat{\theta}_m) - 1) + m(1 - F_n(k_0^*(\hat{\theta}_m))) + (\psi_0(k_0^*(\hat{\theta}_m), \hat{\theta}_m) - c(m, \hat{\theta}_m))p_n(k_0^*(\hat{\theta}_m)) = 0, \quad (23)$$

and using the same arguments that lead to (22), we can prove that (23) is equivalent to

$$\text{umed}(F_n) = G_m(k_0^*(\hat{\theta}_m), \hat{\theta}_m). \quad (24)$$

Consider the estimator $\hat{\theta}$ defined by $\text{umed}(\hat{\theta}) = \text{umed}(F_n)$. By (6) we have $k_0(F_n) = k_0^*(\hat{\theta})$ and by (22) we get

$$\text{umed}(F_n) = \text{umed}(F_{\hat{\theta}}) = G_m(k_0^*(\hat{\theta}), \hat{\theta}).$$

Then (24) holds and this implies that (8) holds too. This proves the Theorem.

6.2 Proof of Lemma 1

Let $X_n \sim F^{(n)}$ and $X \sim F$, and call $G^{(n)}$ and G the distributions of $X_n + u$ and $X + u$, respectively, where u has a uniform distribution on $[-0.5, 0.5]$ independent of X_n or of X . Then $G^{(n)}$ and G have a positive density, and $G^{(n)} \rightarrow_w G$. Since $\text{umed}(F^{(n)}) = \text{med}(G^{(n)})$ and $\text{umed}(F) = \text{med}(G)$, and the median is a weakly continuous functional, the result is shown.

6.3 Proof of Theorem 3

(a) If $F(K) > 0.5$, then then for large n we have $k_0(F_n) = K$, and therefore

$$Z_n = n^{1/2} \left(\frac{0.5 - F_n(K-1)}{p_n(K)} - \frac{0.5 - F(K-1)}{p_0} \right).$$

To derive the asymptotic distribution of Z_n we need first to calculate that of the vector

$$\mathbf{d}_n = n^{1/2} [(F_n(K-1) - F^1), (p_n(K) - p_0)]'.$$

The vector d_n converges in distribution to a bivariate normal distribution with mean $(0, 0)$ and covariance matrix

$$\begin{bmatrix} a & c \\ c & b \end{bmatrix}$$

where

$$a = F^1(1 - F^1), \quad b = p_0(1 - p_0), \quad c = -F^1 p_0.$$

Then since for large n

$$Z_n \simeq -n^{1/2} \frac{F_n(K-1) - F^1}{p_0} - n^{1/2} \frac{(0.5 - F^1)(p_n(K) - p_0)}{p_0^2(K)},$$

the delta method yields that $Z_n \rightarrow^D N(0, \sigma^2)$ where

$$\begin{aligned} \sigma^2 &= \frac{a}{p_0^2} + \frac{b(0.5 - F^1)^2}{p_0^4} + \frac{2c(0.5 - F^1)}{p_0^3} \\ &= \frac{F^1(1 - F^1)}{p_0^2} + \frac{(1 - p_0)(0.5 - F^1)^2}{p_0^3} - \frac{2F^1(0.5 - F^1)}{p_0^2}, \end{aligned}$$

and a straightforward calculation yields (12).

(b) If $F(K) = 0.5$ it is easy to verify that $\text{umed}(F) = K$. Therefore for large n we have

$$k_0(F_n) = \begin{cases} K & \text{if } F_n(K) \geq 0.5 \\ K + 1 & \text{if } F_n(K) < 0.5 \end{cases}$$

We are going to calculate Z_n in both cases. If $F_n(K) \geq 0.5$ we have

$$\begin{aligned} \text{umed}(F_n) &= K - 0.5 + \frac{0.5 - F_n(K-1)}{p_n(K)} \\ &= K + 0.5 + \frac{0.5 - F_n(K)}{p_n(K)}, \end{aligned}$$

and therefore

$$Z_n = \frac{n^{1/2}(0.5 - F_n(K))}{p_n(K)} \leq 0. \quad (25)$$

If $F_n(K) < 0.5$ it follows in the same way that

$$Z_n = \frac{n^{1/2}(0.5 - F_n(K))}{p_n(K+1)} > 0.$$

Note that, conversely, $Z_n \leq 0$ implies $F_n(K) \geq 0.5$ and $Z_n > 0$ implies $F_n(K) < 0.5$. Since $n^{1/2}(0.5 - F_n(K)) \rightarrow_d N(0, 0.25)$, the Central Limit Theorem and Slutsky's Lemma yield (13).

6.4 Proof of Theorem 4

Recall that $\hat{\theta}_n = g^{-1}(\text{umed}(F_n))$ and $\theta = g^{-1}(\text{umed}(F_\theta))$, Put for brevity $K = k_0(F_\theta)$.

(a) If $F_\theta(K) > 0.5$ there exists an interval I containing θ such that $t \in I$ implies that $F_t(K) > 0.5$ and $F_t(K-1) < 0.5$, and therefore $k_0(F_t) = K$. Therefore g is differentiable at θ , and the result follows from Theorem 2, part (a), Theorem 3, and Slutsky's Lemma.

(b) Assume now $F_\theta(K) = 0.5$. Then $t < \theta$ implies that $F_t(K) > 0.5$, and therefore for sufficiently small δ we have $k_0(F_{\theta-\delta}) = K$ and $k_0(F_{\theta+\delta}) = K+1$. Then the left- and right side derivatives of g at θ are g'_- and g'_+ given by (15), and therefore the left- and right side derivatives of g^{-1} are $1/g'_-$ and $1/g'_+$, respectively.

We have

$$n^{1/2}(\hat{\theta}_n - \theta) = n^{1/2}(g^{-1}(\text{umed}(F_n)) - g^{-1}(\text{umed}(F_\theta))). \quad (26)$$

Note that for $\tau < 0$ we have $g(t) - g(\theta) = (t - \theta)g'_- + o(t - \theta)$, and that

$$P\left(n^{1/2}(\text{umed}(F_n) - \text{umed}(F_\theta)) < t\right) \rightarrow H(t),$$

with H defined in (13). The result follows by applying the delta method.

References

- Cantoni, E. and Ronchetti, E. (2001). Robust inference for generalized linear models. *Journal of the American Statistical Association*, **96**, 1022-1030.
- Hampel, F. R. (1974) The influence curve and its role in robust estimation. *Journal of the American Statistical Association*, **69**, 383-394.

Ma, Y., Genton, M. and Parzen, E. (2011). Asymptotic properties of sample quantiles of discrete distributions. *Annals of the Institute of Statistical Mathematics*, **63**, 227–243.

Maronna, R.A., Martin, R.D., Yohai, V.J. and Salibián-Barrera, M. (2019) *Robust Statistics: Theory and Methods (with R), Second Edition*, Wiley, Chichester.

Valdora, M. and Yohai, V.J. (1974). Robust estimators for generalized linear models. *Journal of Statistical Planning and Inference*, **146**, 31-48.