

# Neutron Star as Mirror for Gravitational Waves

Hao Wei,\* Da-Chun Qiang, Zhong-Xi Yu, and Hua-Kai Deng  
*School of Physics, Beijing Institute of Technology, Beijing 100081, China*

Gravitational Wave (GW) has become one of the most active fields in physics and astronomy since the first direct detection of GW event in 2015. As is well known, multiple images of GW events are possible through the gravitational lenses. Here, we propose a novel imaging mechanism for GW events different from the gravitational lens. In this work, we predict that there are two types of GW mirror phenomena caused by the neutron star located in Milky Way or the same host galaxy of GW source, which might be detected within a life period of man (namely the time delay  $\Delta t$  can be a few years to a few tens of years). It is expected to witness this predicted GW mirror phenomenon in the near future. In the long term, the observations of this novel GW mirror phenomenon might help us to find the numerous neutron stars unseen by other means, and learn more about the complicated internal structures of neutron stars, as well as their equation of state.

PACS numbers: 04.30.-w, 04.30.Tv, 97.60.Jd

Gravitational Wave (GW) is a long-standing prediction of general relativity since 1916 [1–4]. The debate about the physical reality of GW was mainly settled during the Chapel Hill conference in 1957 [4, 5]. In 1974, Hulse and Taylor discovered the first indirect evidence for the existence of GW from the binary pulsar PSR B1913+16 [6–8], and hence they earned the 1993 Nobel Prize. In 2015, the LIGO/Virgo Collaboration made the first direct observation of GW event (GW150914) from a binary black hole merger [9, 10], and then earned the 2017 Nobel Prize. From the first multimessenger observations of a binary neutron star merger (GW170817) [11, 12], the speed of GW was confirmed to be the speed of light  $c$  [13]. Nowadays, GW becomes one of the most active fields in physics and astronomy.

As is well known, most objects (e.g. earth) are nearly transparent to GWs. On the other hand, the gravitational lenses (e.g. massive galaxies) can deflect GW as well as light [14, 15]. Therefore, multiple images of GW events are possible through the gravitational lenses. Here, we propose a novel imaging mechanism for GW events different from the gravitational lens. We note that in [16–19] the superconducting film was predicted to be highly reflective mirror for GWs (see also e.g. [20–22]). Due to the historical reasons, mainly the GWs at microwave frequencies were considered in [16–22] far before the first direct GW detection by LIGO. However, the arguments in [16–22] are also applicable for GWs at much lower frequencies (e.g. 1 Hz to 100 Hz GWs detectable for LIGO). Due to the so-called “Heisenberg-Coulomb effect”, it was argued that for the incident GWs at angular frequencies  $\omega \ll 10^{16}$  rad/s (equivalently  $f \ll 10^{15}$  Hz), the




FIG. 1. A simple and naive case of neutron star as mirror for GWs. The plot is not to scale. See the text for details.

reflectivity  $\mathcal{R}_G$  is extremely close to 100% [16–19]. So, the specular reflection of GW from the superconducting film is possible. Unfortunately, such kind of experiments on earth for laboratory-scale superconducting mirrors of GWs are very difficult to achieve in fact.

Let us turn eyes to the sky. It is well known that neutron stars exhibit superconductivity and superfluidity (see e.g. [23–27]). Due to the huge luminosity distances of GW sources (namely  $\mathcal{O}(10^2)$  Mpc, or,  $\mathcal{O}(10^8)$  light years, see e.g. [9–12, 28, 29]) and the small size of neutron stars (namely  $\mathcal{O}(10)$  kilometers, see e.g. [23, 24]), it is safe to ignore the spherical shapes of neutron stars and their complicated internal structures, and hence one can simply regard them as plane mirrors (superconducting films) for GWs. On the other hand, we can assume that the reflectivity  $\mathcal{R}_G$  is 100% for GWs at




FIG. 2. The cases of neutron star as mirror for GWs with luminosity distances  $d_{ON} \ll d_{SN}$ , namely the neutron star N is much closer to the observer O on earth. The plots are not to scale. See the text for details.

frequencies  $f \ll 10^{15}$  Hz [16] as mentioned above. In principle, the positions of neutron stars will be perturbed when GWs go to them. However, such kind of perturbations can also be safely ignored with respect to the huge luminosity distances of GW sources.

By estimating the number of stars that have undergone supernova explosions, there are thought to be around  $10^8$  neutron stars in Milky Way [30]. But most of them are old, cold and hence almost undetectable. It is easy to imagine that numerous neutron stars are distributed over the universe. Although most of them cannot be seen by other means, neutron stars might manifest themselves in the specular reflection of GWs.

At first, let us consider a simple and naive case given in Fig. 1 (not to scale). We suppose a transient GW event “S” is directly detected by the observer “O” on earth, while GW goes through the luminosity distance  $d_{OS}$ . A neutron star “N” as a plane mirror for GWs located on the vertex of the isosceles triangle SNO can reflect the GW from S going to it, and hence the observer O will detect a second GW signal later. Thus, O sees a mirror image “S’” of the GW source S. The luminosity distances satisfy  $d_{OS'} \cos \theta = d_{OS}$ . The intensity of the image S’ divided by the intensity of S equals to  $(d_{OS'}/d_{OS})^{-2} = (\cos \theta)^2$ . For example,  $(\cos \theta)^2 = 3/4$  and  $1/2$  even for  $\theta = 30^\circ$  and  $45^\circ$ , respectively. Thus, the second GW signal is still detectable even if the angle  $\theta$  is fairly large. Note that the allowed large angle  $\theta$  between S and S’ makes the mirror phenomenon different from the gravitational lens whose Einstein angle is typically small. It is of interest to calculate the interval be-

tween the arrival times of GW signals S and S’, which is given by  $\Delta t = t_{S'} - t_S = (t_{S'}/t_S - 1) t_S = (d_{OS'}/d_{OS} - 1) t_S = ((\cos \theta)^{-1} - 1) t_S$ . Unfortunately, for a not so small angle  $\theta$ , the arrival time interval  $\Delta t$  is on the order of  $t_S \sim \mathcal{O}(10^8)$  years. This is far beyond the patience. If we hope to detect both GW signals S and S’ within a life period of man (namely a few tens of years), the angle  $\theta$  must be very small. Noting that  $\Delta t \simeq \theta^2 t_S/2$  for  $\theta \rightarrow 0$ , we find that  $\theta \lesssim 10^{-3}$  or  $10^{-4}$  is required by  $\Delta t \lesssim 10^2$  years or 10 years, respectively. This means that the host galaxy of neutron star N lies in the middle of the line of sight from O to S. It will play the role of gravitational lens. So, the mirror phenomenon will be destroyed by the host galaxy of neutron star N in this simple and naive case.

Clearly, in the previous case it is too special that S, N, O form an isosceles triangle. Noting that any observer O on the extension line of S’N can always see the mirror image S’, we can consider the cases with  $d_{ON} \ll d_{SN}$ , i.e. the neutron star N is much closer to O, as shown in Fig. 2 (not to scale). In fact, N can be located in the same host galaxy of O (namely Milky Way). As is well known, the diameter of Milky Way is  $1.5 \sim 2 \times 10^5$  light years [31], and hence  $d_{ON} \lesssim 2 \times 10^5$  light years. However, the neutron star N cannot be too close to earth (O), otherwise the humankind and other lives on earth will all be killed by it. To our best knowledge, the nearest neutron star to earth is claimed to be Calvera (1RXS J141256.0+792204) at the distance around  $250 \sim 1000$  light years [32]. Other two closest known neutron stars are RX J1856.5-3754 and PSR J0108-1431, which are both about 400 light years from earth [33]. So, we take  $d_{ON} \gtrsim 250$




FIG. 3. The cases of neutron star as mirror for GWs with luminosity distances  $d_{\text{SN}} \ll d_{\text{ON}}$ , namely the neutron star N is much closer to the GW source S. The plots are not to scale. See the text for details.

light years. There are mainly three types of configurations for O and N, as shown in Fig. 2. The case (2b) is fairly special, and it becomes possible because earth is nearly transparent to GWs. The mirror image  $S'$  is in the opposite direction of the GW source S. In this case, the interval between the arrival times of GW signals S and  $S'$  is  $\Delta t = 2d_{\text{ON}}/c \gtrsim 500$  years. We turn to the case (2c) with the angle  $\theta \geq 90^\circ$ . In this case,  $\Delta t = (d_{\text{SN}} + d_{\text{ON}} - d_{\text{OS}})/c > d_{\text{ON}}/c \gtrsim 250$  years. Thus, both the arrival time intervals  $\Delta t$  of cases (2b) and (2c) are far beyond a life period of man (namely a few tens of years). The last hope lies in the case (2a) with the angle  $\theta < 90^\circ$ . Obviously,  $\Delta t = (d_{\text{SN}} + d_{\text{ON}} - d_{\text{OS}})/c \rightarrow 0$  if  $\theta \rightarrow 0$ . However, we should avoid  $\theta \rightarrow 0$ , otherwise the GW from S might be reflected before it could reach the observer O. In fact, a large  $\theta$  is possible in the case (2a). Using the law of cosines to the triangle SON, we have  $\cos \theta = (d_{\text{OS}}^2 + d_{\text{ON}}^2 - d_{\text{SN}}^2)/(2d_{\text{OS}}d_{\text{ON}}) = (d_{\text{OS}}^2 + d_{\text{ON}}^2 - (d_{\text{OS}} - d_{\text{ON}} + c\Delta t)^2)/(2d_{\text{OS}}d_{\text{ON}}) \simeq 1 - c\Delta t/d_{\text{ON}}$  for  $d_{\text{ON}} \ll d_{\text{OS}}$  and  $c\Delta t \ll d_{\text{OS}}$ . So, we obtain  $\Delta t \simeq (1 - \cos \theta)d_{\text{ON}}/c$ . If  $d_{\text{ON}} \simeq 250$  light years,  $\Delta t \simeq 0.04, 0.95, 3.8, 8.5, 15.1, 33.5, 47.7$  and  $73.2$  years for  $\theta = 1^\circ, 5^\circ, 10^\circ, 15^\circ, 20^\circ, 30^\circ, 36^\circ$  and  $45^\circ$ , respectively. If  $d_{\text{ON}} \simeq 1000$  light years,  $\Delta t \simeq 0.15, 3.8, 15.2, 34.1$  and  $60.3$  years for  $\theta = 1^\circ, 5^\circ, 10^\circ, 15^\circ$  and  $20^\circ$ , respectively. Thus, we can see the mirror image  $S'$  within a life period of man (namely the time delay  $\Delta t$  can be a few years to a few tens of years) while the angle  $\theta$  can still be fairly large. On the other hand, the intensity of the image  $S'$  divided by the intensity of S equals to  $(d_{\text{OS}'}/d_{\text{OS}})^{-2} = (1 + c\Delta t/d_{\text{OS}})^{-2} \simeq 1$  for  $c\Delta t \ll d_{\text{OS}}$ . So, we will see two GW signals S and

$S'$  with almost same intensities and wave forms.

Similarly, we can also consider the cases with  $d_{\text{SN}} \ll d_{\text{ON}}$ , namely the neutron star N is much closer to the GW source S, as shown in Fig. 3 (not to scale). In fact, N can be located in the same host galaxy of S. Noting that the remnants of most GW events are black holes which can also swallow GWs, in Fig. 3 there is no counterpart to the case (2b). Since we need not to care the alien lives, the neutron star N can be very close to S, unlike the cases in Fig. 2. In fact,  $d_{\text{SN}}$  can be a few light years, or a few tens of light years. Because  $d_{\text{SN}} \ll d_{\text{ON}} \sim d_{\text{OS}}$ , it is easy to see that the angle  $\theta$  must be very close to 0 (on the order of  $d_{\text{SN}}/d_{\text{OS}} \sim 10^{-8}$  to  $10^{-7}$ ). Thus, from the viewpoint of the observer O on earth, the mirror image  $S'$  is almost in the same direction of the GW source S. But this does not mean that the neutron star N must lie in the line OS or its extension line, from the local viewpoints of S and N, as will be shown below. In the case (3a) with the angle  $\phi < 90^\circ$ , following the similar derivations in the previous case (2a), we have  $\cos \phi \simeq 1 - c\Delta t/d_{\text{SN}}$ . Clearly, the angle  $\phi$  can be large if  $c\Delta t$  is comparable to  $d_{\text{SN}}$ . So, the neutron star N does not lie in the line OS actually (although  $\theta \rightarrow 0$  due to the huge  $d_{\text{OS}}$ ). Because  $c\Delta t \simeq (1 - \cos \phi)d_{\text{SN}}$ , we can see the mirror image  $S'$  within a life period of man (namely  $\Delta t$  can be a few years to a few tens of years) if the neutron star N is close enough to the GW source S while the angle  $\phi$  can still be fairly large. Note that even if N is not so close to S,  $\Delta t$  can still be a few years to a few tens of years for a suitable and large  $\theta$  (see the case (2a)). Similarly, in the case (3b) with the angle  $\phi \geq 90^\circ$  (and hence the angle  $\psi < 90^\circ$ ), we

have  $c\Delta t \simeq (1 + \cos\psi) d_{\text{SN}}$ . The angle  $\psi$  can be large, so that the neutron star N does not lie in the extension line of OS actually (although  $\theta \rightarrow 0$  due to the huge  $d_{\text{OS}}$ ). Again, we can see the mirror image  $S'$  within a life period of man (namely  $\Delta t$  can be a few years to a few tens of years) if the neutron star N is close enough to the GW source S while the angle  $\psi$  can still be fairly large. On the other hand, in both cases (3a) and (3b), the intensity of the image  $S'$  divided by the intensity of S equals to  $(d_{\text{OS}'}/d_{\text{OS}})^{-2} = (1 + c\Delta t/d_{\text{OS}})^{-2} \simeq 1$  for  $c\Delta t \ll d_{\text{OS}}$ . So, we will see two GW signals S and  $S'$  with almost same intensities and wave forms.

Besides the three cases shown in Figs. 1–3 (not to scale), the rest is the case with  $d_{\text{SN}}$  comparable to  $d_{\text{ON}}$ . It is easy to see that this case is quite similar to the simple and naive case in Fig. 1, and hence is also inviable. It is worth noting that the main reason for the failure of the cases with  $d_{\text{SN}}$  comparable to  $d_{\text{ON}}$  (including the simple and naive case in Fig. 1) is either  $\Delta t \sim t_{\text{S}} \sim \mathcal{O}(10^8)$  years if the angle  $\theta$  is not so small, or the host galaxy of neutron star N must play the role of gravitational lens to destroy the mirror phenomenon if the angle  $\theta$  is very close to 0. In fact, the latter might be evaded for an isolated neutron star N without a host galaxy, which might be a wandering star in the intergalactic space been kicked out of its original position due to some unknown reasons. However, this must be extremely rare, and hence we do not consider this possibility here.

In summary, we predict that there are two types of GW mirror phenomena caused by the neutron star located in Milky Way (case (2a)) or the same host galaxy of GW source (cases (3a) and (3b)), which might be detected within a life period of man (namely the time delay  $\Delta t$  can be a few years to a few tens of years). In both types of GW mirror phenomena, we will see two GW signals S and  $S'$  with almost same intensities and wave forms. The separate angle  $\theta$  between the GW source S and its mirror image  $S'$  can be fairly large in case (2a), while  $\theta$  should be fairly close to 0 in cases (3a) and (3b). Noting that GWs became directly detectable since 2015, we hope to witness this predicted GW mirror phenomenon in the near future.

In fact, the GW mirror phenomenon predicted here is different from the well-known gravitational lenses [14, 15] which can also image GWs. In the mirror phenomenon, we will see the signal from the GW source S directly, and after a time delay  $\Delta t$  we will see a second GW signal from the unique mirror image  $S'$ . On the contrary, in the case of

gravitational lens, one will see more than (or equal to) two images, or even rings and arcs. In the mirror phenomenon might be detected within a life period of man, we will see two GW signals S and  $S'$  with almost same intensities and wave forms. On the contrary, in the case of gravitational lens, the images should be magnified (and/or distorted). Also, in the case of gravitational lens, the separate angle  $\theta$  (Einstein angle) between images is very small (usually a few arcseconds, namely  $\theta \sim 10^{-6}$ ). But in the mirror phenomenon, the separate angle  $\theta$  between the GW source S and its mirror image  $S'$  can be fairly large in case (2a). Finally, the time delay  $\Delta t$  is usually short (a few days to a few years) in the case of gravitational lens, while the time delay  $\Delta t$  could be relatively long (a few years to  $\mathcal{O}(10^8)$  years) in the mirror phenomenon.

Clearly, the discussions in this work are very preliminary. In fact, many simplifications have been made. For example, we ignored the spherical shape of neutron star and its complicated internal structures. Also, we have not calculated the realistic GW reflectivity  $\mathcal{R}_{\text{G}}$  for an impure superconducting solid sphere. On the other hand, the event rate of the GW mirror phenomenon was not estimated. However, this preliminary work clearly showed the key idea of the novel GW mirror phenomenon, and made clear predictions which might be detected in the near future. This is very important.

We hope to construct some full and realistic models for this GW mirror phenomenon caused by the neutron star in the future works. In the long term, the observations of this novel GW mirror phenomenon might help us to find the numerous neutron stars unseen by other means, and learn more about the complicated internal structures of neutron stars, as well as their equation of state.

We are grateful to Rong-Gen Cai, Hongwei Yu, Puxun Wu, Zhoujian Cao, Wen Zhao, Bin Hu, Hongsheng Zhang for useful discussions. H.W. thank University of Jinan for the kind hospitality. This work was supported in part by NSFC under Grants No. 11975046 and No. 11575022.

---

\* Corresponding author; email: haowei@bit.edu.cn

- [1] A. Einstein, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) **1916**, 688 (1916).
- [2] A. Einstein, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) **1918**, 154 (1918).

- [3] A. Einstein and N. Rosen, *J. Franklin Inst.* **223**, 43 (1937).
- [4] J. L. Cervantes-Cota *et al.*, *Universe* **2**, no. 3, 22 (2016) [arXiv:1609.09400].
- [5] P. R. Saulson, *Gen. Rel. Grav.* **43**, 3289 (2011).
- [6] J. H. Taylor, L. A. Fowler and P. M. McCulloch, *Nature* **277**, 437 (1979).
- [7] J. H. Taylor and J. M. Weisberg, *Astrophys. J.* **253**, 908 (1982).
- [8] D. R. Lorimer, *Living Rev. Rel.* **11**, 8 (2008) [arXiv:0811.0762].
- [9] B. P. Abbott *et al.*, *Phys. Rev. Lett.* **116**, no. 6, 061102 (2016) [arXiv:1602.03837].
- [10] B. P. Abbott *et al.*, *Phys. Rev. Lett.* **116**, no. 22, 221101 (2016) [arXiv:1602.03841].
- [11] B. P. Abbott *et al.*, *Phys. Rev. Lett.* **119**, no. 16, 161101 (2017) [arXiv:1710.05832].
- [12] B. P. Abbott *et al.*, *Astrophys. J.* **848**, no. 2, L12 (2017) [arXiv:1710.05833].
- [13] B. P. Abbott *et al.*, *Astrophys. J.* **848**, no. 2, L13 (2017) [arXiv:1710.05834].
- [14] P. Schneider, J. Ehlers, E. E. Falco, *Gravitational Lenses*, Springer-Verlag, New York (1992).
- [15] P. Schneider, C. Kochanek and J. Wambsganss, *Gravitational Lensing: Strong, Weak and Micro*, Springer-Verlag, Berlin (2006).
- [16] S. J. Minter, K. Wegter-McNelly and R. Y. Chiao, *Physica E* **42**, 234 (2010) [arXiv:0903.0661].
- [17] R. Y. Chiao, S. Minter and K. Wegter-McNelly, arXiv:0903.3280 [gr-qc].
- [18] R. Y. Chiao, arXiv:0710.1378 [gr-qc].
- [19] R. Y. Chiao *et al.*, *J. Br. Interplanet. Soc.* **70**, 405 (2017) [arXiv:1712.08680].
- [20] J. Q. Quach, *Phys. Rev. Lett.* **114**, no. 8, 081104 (2015) [arXiv:1502.07429].
- [21] N. A. Inan, J. J. Thompson and R. Y. Chiao, *Fortsch. Phys.* **65**, no. 6-8, 1600066 (2017).
- [22] N. A. Inan, *Int. J. Mod. Phys. D* **26**, no. 12, 1743031 (2017).
- [23] J. M. Lattimer and M. Prakash, *Science* **304**, 536 (2004) [astro-ph/0405262].
- [24] G. Baym and C. Pethick, *Ann. Rev. Astron. Astrophys.* **17**, 415 (1979).
- [25] F. Weber, *Prog. Part. Nucl. Phys.* **54**, 193 (2005) [astro-ph/0407155].
- [26] D. Page *et al.*, *Phys. Rev. Lett.* **106**, 081101 (2011) [arXiv:1011.6142].
- [27] M. Baldo *et al.*, *Phys. Lett. B* **562**, 153 (2003) [nucl-th/0212096].
- [28] <https://www.ligo.org/detections.php>
- [29] B. P. Abbott *et al.*, *Phys. Rev. X* **9**, no. 3, 031040 (2019) [arXiv:1811.12907].
- [30] M. Camenzind, *Compact Objects in Astrophysics: White Dwarfs, Neutron Stars and Black Holes*, Springer-Verlag, Berlin (2007).
- [31] M. Lopez-Corredoira *et al.*, *Astron. Astrophys.* **612**, L8 (2018) [arXiv:1804.03064].
- [32] R. E. Rutledge, D. B. Fox and A. H. Shevchuk, *Astrophys. J.* **672**, 1137 (2008) [arXiv:0705.1011].
- [33] B. Posselt, R. Neuhauser and F. Haberl, *Astron. Astrophys.* **496**, 533 (2009) [arXiv:0811.0398].