

A polynomial kernel for 3-leaf power deletion

Jungho Ahn^{2,1}, Eduard Eiben³, O-joung Kwon^{4,1}, and Sang-il Oum^{1,2}

¹Discrete Mathematics Group, Institute for Basic Science (IBS),
Daejeon, South Korea

²Department of Mathematical Sciences, KAIST,
Daejeon, South Korea

³Department of Informatics, University of Bergen, Bergen, Norway

⁴Department of Mathematics, Incheon National University,
Incheon, South Korea

Email addresses: junghoahn@kaist.ac.kr, Eduard.Eiben@uib.no,
ojoungkwon@gmail.com, sangil@ibs.re.kr

February 6, 2022

Abstract

A graph G is an ℓ -leaf power of a tree T if $V(G)$ is equal to the set of leaves of T , and distinct vertices v and w of G are adjacent if and only if the distance between v and w in T is at most ℓ . Given a graph G , 3-LEAF POWER DELETION asks whether there is a set $S \subseteq V(G)$ of size at most k such that $G \setminus S$ is a 3-leaf power of some tree T . We provide a polynomial kernel for this problem. More specifically, we present a polynomial-time algorithm for an input instance (G, k) to output an equivalent instance (G', k') such that $k' \leq k$ and G' has at most $O(k^{14} \log^{12} k)$ vertices.

1 Introduction

Nishimura, Ragde, and Thilikos [26] introduced an ℓ -leaf power of a tree to understand the structure of phylogenetic trees in computational biology. A graph G is an ℓ -leaf power of a tree T if $V(G)$ is equal to the set of leaves of

¹JA, OK, and SO are supported by IBS-R029-C1. OK is also supported by the National Research Foundation of Korea (NRF) grant funded by the Ministry of Education (No. NRF-2018R1D1A1B07050294).

T , and distinct vertices v and w of G are adjacent if and only if the distance between v and w in T is at most ℓ . We say that G is an ℓ -leaf power if G is an ℓ -leaf power of some tree. Note that an ℓ -leaf power could have more than one component. For instance, an ℓ -leaf power of a path of length at least $\ell + 1$ has two distinct trivial components. We are interested in the following *vertex deletion* problem.

3-LEAF POWER DELETION

Input : A graph G and a non-negative integer k .

Question : Is there a set $S \subseteq V(G)$ with $|S| \leq k$ such that $G \setminus S$ is a 3-leaf power?

Vertex deletion problems include some of the best studied NP-hard problems in theoretical computer science, including VERTEX COVER and FEEDBACK VERTEX SET. In general, the problem asks whether it is possible to delete at most k vertices from an input graph so that the resulting graph belongs to a specified graph class. Lewis and Yannakakis [23] showed that every vertex deletion problem to a non-trivial hereditary (i.e., closed under taking induced subgraphs) graph class is NP-hard. Since the class of 3-leaf powers is non-trivial and hereditary, it follows that 3-LEAF POWER DELETION is NP-hard.

We study 3-LEAF POWER DELETION through the parameterized complexity paradigm [12, 15], which measures the performance of algorithms not only with respect to the input size but also with respect to an additional numerical parameter. The notion of vertex deletion allows a highly natural choice of the parameter, specifically the size of the deletion set k . From the characterization by Dom et al. [13], it follows that every 3-leaf power is *chordal* and *distance-hereditary*. CHORDAL DELETION is a problem of deciding whether a graph has a vertex set of size at most k whose deletion makes it chordal. Marx [25] showed that CHORDAL DELETION is fixed-parameter tractable by presenting an algorithm with running time $f(k) \cdot n^{O(1)}$ where n is the number of vertices of an input graph, and Cao and Marx [8] improved this result by presenting an algorithm with running time $2^{O(k \log k)} \cdot n^{O(1)}$. DISTANCE-HEREDITARY DELETION is a problem of deciding whether a graph has a vertex set of size at most k whose deletion makes it distance-hereditary. Eiben, Ganian, and Kwon [17] presented a single-exponential fixed-parameter tractable algorithm for DISTANCE-HEREDITARY DELETION, that is an algorithm with running time $O(c^k \cdot n^{O(1)})$ for input size n and some constant c . To obtain our first result, we observe that if an input graph G does not contain a *small obstruction*, that is a minimal induced subgraph of size at most 5 that is not a 3-leaf power, then G is a

3-leaf power if and only if G is distance-hereditary. Hence, after branching on small obstructions, we can use the algorithm by Eiben, Ganian, and Kwon [17] as a black-box. This immediately gives us the following theorem:

Theorem 1.1. *Given an instance (G, k) , we can correctly solve 3-LEAF POWER DELETION in time $O(37^k |V(G)|^7 (|V(G)| + |E(G)|))$.*

After we establish the fixed-parameter tractability of 3-LEAF POWER DELETION, one of the most natural follow-up questions in parameterized complexity is whether the problem admits a polynomial kernel. A *kernel* is basically a polynomial-time preprocessing algorithm that transforms the given instance of the problem into an equivalent instance whose size is bounded above by some function $f(k)$ of the parameter. The function $f(k)$ is usually referred to as the *size* of the kernel. A *polynomial kernel* is then a kernel with size bounded above by some polynomial in k . The existence of polynomial kernels for vertex deletion problems has been widely investigated [4, 14, 19, 24]. Jansen and Pilipczuk [20] presented a kernel with $O(k^{161} \log^{58} k)$ vertices for CHORDAL DELETION, and Agrawal et al. [2] improved this result by presenting a kernel with $O(k^{12} \log^{10} k)$ vertices. Kim and Kwon [21] presented a kernel with $O(k^{30} \log^5 k)$ vertices for DISTANCE-HEREDITARY DELETION. This leads us to the main result of our paper:

Theorem 1.2. *3-LEAF POWER DELETION admits a kernel with $O(k^{14} \log^{12} k)$ vertices.*

The first step of our kernel is to find a “good” approximate solution, that is a set $S \subseteq V(G)$ of size $O(k^2 \log^2 k)$ such that $G \setminus (S \setminus \{v\})$ is a 3-leaf power for each vertex v in S . To construct such a set S in polynomial time, we use Theorem 1.1 in Kim and Kwon [22], Lemma 1.3 in Jansen and Pilipczuk [20], and Theorem 2 in Agrawal et al. [1]. Afterward, we design a series of reduction rules that allows us to bound the number and size of components of $G \setminus S$. We remark that Bessy, Paul, and Perez [5] presented a kernel with $O(k^3)$ vertices for 3-leaf power edge modification problems including editing, completion, and edge-deletion.

We organize this paper as follows. In Section 2, we summarize some terminologies in graph theory and parameterized complexity, and introduce 3-leaf powers, distance-hereditary graphs, and a relation between them. In Section 3, we present a single-exponential fixed-parameter tractable algorithm for 3-LEAF POWER DELETION. In Section 4, we introduce a good modulator of a graph, and present an algorithm that either confirms that an input instance (G, k) is a no-instance, or constructs a small good modulator of G . In Sections 5 and 6, we design a series of reduction rules that

allows us to bound the number of vertices outside of a good modulator of a graph. In Section 7, we combine the above steps to prove our main result. In Section 8, we conclude this paper with some open problems.

2 Preliminaries

In this paper, all graphs are finite and simple. For a vertex v and a set X of vertices of a graph G , let $N_G(v)$ be the set of neighbors of v in $V(G)$, $N_G(X)$ be the set of vertices not in X that are adjacent to some vertices in X , and $N_G[X] := N_G(X) \cup X$. We may omit the subscripts of these notations if it is clear from the context. For disjoint sets X and Y of vertices of G , we say that X is *complete* to Y if each vertex in X is adjacent to all vertices in Y , and X is *anti-complete* to Y if each vertex in X is non-adjacent to all vertices in Y . Let $G \setminus X$ be a graph obtained from G by removing all vertices in X and all edges incident with some vertices in X , and $G[X] := G \setminus (V(G) \setminus X)$. We may write $G \setminus v$ instead of $G \setminus \{v\}$ for each vertex v of G . For a set T of edges of G , let $G \setminus T$ be a graph obtained from G by removing all edges in T .

A graph G is *trivial* if $|V(G)| = 1$, and *non-trivial*, otherwise. A *clique* is a set of pairwise adjacent vertices. A graph is *complete* if every pair of two distinct vertices is adjacent, and *incomplete*, otherwise. An *independent set* is a set of pairwise non-adjacent vertices. Distinct vertices v and w of G are *twins* in G if $N_G(v) \setminus \{w\} = N_G(w) \setminus \{v\}$. Twins v and w in G are *true* if v and w are adjacent, and *false* if v and w are non-adjacent. A *twin-set* in G is a set of pairwise twins in G . A twin-set is *true* if it is a clique, and *false* if it is an independent set.

A vertex of a graph is *isolated* if it has no neighbors. A node of a tree is a *leaf* if it has exactly one neighbor, and is *branching* if it has at least three neighbors. For graphs G_1, \dots, G_m , a graph G is (G_1, \dots, G_m) -*free* if G has no induced subgraphs isomorphic to one of G_1, \dots, G_m .

We say that a reduction rule is *safe* if each input instance is equivalent to the resulting instance obtained from the input instance by applying the rule.

2.1 Parameterized problems and kernels

For a fixed finite set Σ of alphabets, an *instance* is an element in $\Sigma^* \times \mathbb{N}$. For an instance (I, k) , k is called a *parameter*. A *parameterized problem* is a set $L \subseteq \Sigma^* \times \mathbb{N}$. A parameterized problem Π is *fixed-parameter tractable* if there is an algorithm, called a *fixed-parameter tractable algorithm* for Π , that correctly decides whether an input instance $(I, k) \in \Pi$ in time $O(f(k) \cdot n^c)$



Figure 1: A bull, a dart, a gem, a house, and a domino.

for a computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ and a constant c where n is the size of I . A fixed-parameter tractable algorithm for a parameterized problem is *single-exponential* if it takes $O(\alpha^k \cdot n^c)$ time for some constants $\alpha > 1$ and c .

An instance (I, k) is a *yes-instance* for a parameterized problem Π if $(I, k) \in \Pi$, and a *no-instance*, otherwise. Instances (I, k) and (I', k') are *equivalent* with respect to Π if (I, k) is a yes-instance for Π if and only if (I', k') is a yes-instance for Π . A *kernel* for Π is a polynomial-time algorithm that given an instance (I, k) , outputs an instance (I', k') equivalent to (I, k) with respect to Π such that $|I'| + k' \leq g(k)$ for some computable function $g : \mathbb{N} \rightarrow \mathbb{N}$. Such a function $g(k)$ is the *size* of the kernel. A *polynomial kernel* for Π is a kernel for Π with the size as some polynomial in k . We may omit the term “for Π ” and “with respect to Π ” of all these definitions if it is clear from the context. There is a relationship between the fixed-parameter tractability and the existence of a kernel for parameterized problems.

Theorem 2.1 (See Downey and Fellows [15]). *A parameterized problem Π is fixed-parameter tractable if and only if Π admits a kernel.*

2.2 Characterizations of 3-leaf powers

Brandstädt and Le [6] presented a linear-time algorithm to recognize 3-leaf powers.

Theorem 2.2 (Brandstädt and Le [6, Theorem 15]). *Given a graph G , we can either confirm that G is not a 3-leaf power, or find a tree of which G is a 3-leaf power in linear time.*

Figure 1 shows three graphs called a *bull*, a *dart*, and a *gem*. A *hole* is an induced cycle of length at least 4. A graph is *chordal* if it has no holes. Dom, Guo, Hüffner, and Niedermeier [13] presented the following characterization of 3-leaf powers.

Theorem 2.3 (Dom, Guo, Hüffner, and Niedermeier [13, Theorem 1]). *A graph G is a 3-leaf power if and only if G is (bull, dart, gem)-free and chordal.*

We say that a graph H is an *obstruction* if H is not a 3-leaf power, and every proper induced subgraph of H is a 3-leaf power. An obstruction H is *small* if $|V(H)| \leq 5$. We see the following six observations about obstructions.

Observation 1 (O1). *No obstructions have true twins.*

Observation 2 (O2). *No obstructions have independent sets of size at least 4.*

Observation 3 (O3). *No obstructions have K_4 or $K_{2,3}$ as a subgraph.*

Observation 4 (O4). *If an obstruction H has false twins v and w , then both v and w have degree 2 in H .*

Observation 5 (O5). *If a vertex v of an obstruction H has exactly one neighbor w in $V(H)$, then w has degree at least 3 in H .*

Observation 6 (O6). *A graph H is an obstruction having three distinct vertices of degree 2 in H if and only if H is a hole.*

Brandstädt and Le [6] showed that a graph G is a 3-leaf power if and only if G is obtained from some forest F by replacing each node u of F with a non-empty clique B_u of arbitrary size, and each edge vw of F with the edges whose one end is in B_v , and the other end is in B_w . We rephrase this characterization by using the following definition.

A *tree-clique decomposition* of a graph G is a pair $(F, \{B_u : u \in V(F)\})$ of a forest F and a family $\{B_u : u \in V(F)\}$ of non-empty subsets of $V(G)$ satisfying the following two conditions.

- (1) The family $\{B_u : u \in V(F)\}$ is a partition of $V(G)$.
- (2) Distinct vertices x and y of G are adjacent if and only if F has either a node u such that $\{x, y\} \subseteq B_u$, or an edge vw such that $x \in B_v$ and $y \in B_w$.

We call B_u a *bag* of u for each node u of F . We say that B is a *bag* of G if B is a bag of some node of F . We remark that each bag is a clique by (2).

Theorem 2.4 (Brandstädt and Le [6, Theorem 9]). *A graph G is a 3-leaf power if and only if G has a tree-clique decomposition. Moreover, if G is connected and incomplete, then G has a unique tree-clique decomposition.*

We remark that every connected incomplete 3-leaf power has at least three bags. Brandstädt and Le [6] showed that for a connected incomplete 3-leaf power G , distinct vertices v and w of G are in the same bag of G if and only if v and w are true twins in G . Thus, for such a graph G , B is a bag of G if and only if B is a maximal true twin-set in G .

2.3 Characterizations of distance-hereditary graphs

A graph G is *distance-hereditary* if for every connected induced subgraph H of G and vertices v and w of H , the distance between v and w in H is equal to the distance between v and w in G . Figure 1 shows two graphs called a *house* and a *domino*. Bandelt and Mulder [3] presented the following characterization of distance-hereditary graphs.

Theorem 2.5 (Bandelt and Mulder [3, Theorem 2]). *A graph G is distance-hereditary if and only if G is (house, domino, gem)-free, and has no holes of length at least 5.*

Since both the house and the domino have a hole of length 4, every 3-leaf power is distance-hereditary by Theorems 2.3 and 2.5. The following lemma shows when graphs are not distance-hereditary. A proof of the following lemma is readily derived from the definition of a distance-hereditary graph.

Lemma 2.6. *Let P be an induced path of length at least 3 in a graph G . If G has a vertex v adjacent to both ends of P , then $G[V(P) \cup \{v\}]$ is not distance-hereditary.*

3 Single-exponential fixed-parameter tractable algorithms

A set S of vertices of a graph G is a *modulator* of G if $G \setminus S$ is a 3-leaf power. If G has a modulator of size at most k , then it has rank-width at most $k+1$ because every 3-leaf power is distance-hereditary, which has rank-width at most 1 [27]. This already allows us to construct a fixed-parameter tractable algorithm for 3-LEAF POWER DELETION by using an algorithm for graphs of bounded rank-width solving monadic second-order logic [10]. We improve this further by showing that a single-exponential fixed-parameter tractable algorithm exists for 3-LEAF POWER DELETION. To do so, we use the following theorem of Eiben, Ganian, and Kwon [17].

Theorem 3.1 (Eiben, Ganian, and Kwon [17, Theorem 1.1]). *For a graph G and a non-negative integer k , we can decide whether there is a set $S \subseteq V(G)$ with $|S| \leq k$ such that $G \setminus S$ is distance-hereditary in time $O(37^k |V(G)|^7 (|V(G)| + |E(G)|))$.*

Proof of Theorem 1.1. We proceed by induction on k . If $k = 0$, then 3-LEAF POWER DELETION is solved in linear time by Theorem 2.2. Thus, we may

assume that $k > 0$. We investigate every 5-element subset of $V(G)$ to find a small obstruction in time $O(|V(G)|^5)$. If we find a small obstruction H in G , then we branch on each vertex of H to be included in the solution, and solve each of the at most five instances in time $O(37^{k-1}|V(G)|^7(|V(G)| + |E(G)|))$ by the induction hypothesis. Otherwise, we apply Theorem 3.1 for (G, k) . Then since G has no small obstructions, an induced subgraph of G is a 3-leaf power if and only if it is distance-hereditary, and therefore the answer for 3-LEAF POWER DELETION is equal to the answer obtained from G and k by Theorem 3.1. This can be done in time $O(37^k|V(G)|^7(|V(G)| + |E(G)|))$. \square

4 Good modulators

A modulator S of a graph G is *good* if $G \setminus (S \setminus \{v\})$ is a 3-leaf power for each vertex v in S . Note that if G has a modulator S , then for every induced subgraph G' of G , $S \cap V(G')$ is a modulator of G' . This means that if (G, k) is a yes-instance, then so is (G', k) . We remark that if G has an obstruction H and a good modulator S , then H has at least two vertices in S . To find a small good modulator, we first find a modulator by combining a maximal packing of small obstructions with outcomes of the following approximation algorithms for CHORDAL DELETION by Kim and Kwon [22], and by Agrawal et al. [1]:

Theorem 4.1 (Kim and Kwon [22, Theorem 1.1]). *Given a graph G and a positive integer k , we can either find $k + 1$ vertex-disjoint holes in G , or find a set $S \subseteq V(G)$ of size $O(k^2 \log k)$ such that $G \setminus S$ is chordal in time bounded above by a polynomial in $|V(G)|$.*

Theorem 4.2 (Agrawal, Lokshtanov, Misra, Saurabh, and Zehavi [1, Theorem 2]). *For a graph G and a non-negative integer k , we can either confirm that there is no set $S \subseteq V(G)$ with $|S| \leq k$ such that $G \setminus S$ is chordal, or find a set $S \subseteq V(G)$ of size $O(k \log^2 |V(G)|)$ such that $G \setminus S$ is chordal in time bounded above by a polynomial in $|V(G)|$.*

Corollary 4.3. *Given an instance (G, k) with $k > 0$, we can either confirm that G has no modulators of size at most k , or find a modulator of G having size $O(\min(k^2 \log k, k \log^2 |V(G)|))$ in time bounded above by a polynomial in $|V(G)|$.*

Proof. We can find a maximal packing H_1, \dots, H_m of vertex-disjoint small obstructions in G in time $O(|V(G)|^6)$. If $m \geq k + 1$, then we confirm that G has no modulators of size at most k . Thus, we may assume that $m \leq k$.

Let $X := \bigcup_{i \in \{1, \dots, m\}} V(H_i)$. We apply Theorems 4.1 and 4.2 for $G \setminus X$ and k . Note that $|X| \leq 5k$, and $G \setminus X$ has no small obstructions.

If any of the algorithms in Theorems 4.1 and 4.2 confirms that $G \setminus X$ has no set S of size at most k such that $G \setminus (X \cup S)$ is chordal, then we confirm that G has no modulators of size at most k . Thus, let S_1 be the output of the algorithm in Theorem 4.1 having size $O(k^2 \log k)$, and S_2 be the output of the algorithm in Theorem 4.2 having size $O(k \log^2 |V(G)|)$. Then we choose S as one of S_1 and S_2 so that $|S| = \min(|S_1|, |S_2|)$. Then $X \cup S$ is a modulator of G , and therefore $|X \cup S| = |X| + |S| \leq 5k + O(\min(k^2 \log k, k \log^2 |V(G)|)) = O(\min(k^2 \log k, k \log^2 |V(G)|))$. \square

With a modulator of size $O(\min(k^2 \log k, k \log^2 |V(G)|))$ at hand, we are ready to find a small good modulator. We note that, in principle, a small good modulator might not exist, but if that is the case, we are able to identify a vertex that has to be in every modulator of size at most k . Then we can remove it from the input graph, and decrease the parameter k by 1.

Reduction Rule 1 (R1). *Given an instance (G, k) with $k > 0$, if G has $k+1$ obstructions H_1, \dots, H_{k+1} and a vertex v of G such that $V(H_i) \cap V(H_j) = \{v\}$ for every distinct i and j in $\{1, \dots, k+1\}$, then replace (G, k) with $(G \setminus v, k-1)$.*

Proof of Safeness. It suffices to show that if G has a modulator S of size at most k , then S contains v . Suppose not. Then S contains at least one vertex of $H_i \setminus v$ for each $i \in \{1, \dots, k+1\}$. Therefore, $|S| \geq k+1$, a contradiction. \square

To find the obstructions H_1, \dots, H_{k+1} , we make use of the following lemma, which we slightly rephrase to better fit our application.

Lemma 4.4 (Jansen and Pilipczuk [20, Lemma 1.3]). *Given a graph G , a non-negative integer k , and a vertex v , if $G \setminus v$ is chordal, then we can either find holes H_1, \dots, H_{k+1} in G such that $V(H_i) \cap V(H_j) = \{v\}$ for every distinct i and j in $\{1, \dots, k+1\}$, or find a set $S \subseteq V(G) \setminus \{v\}$ of size at most $12k$ such that $G \setminus S$ is chordal in time bounded above by a polynomial in $|V(G)|$.*

Lemma 4.5. *Given an instance (G, k) with $k > 0$, we can find an equivalent instance (G', k') such that $|V(G')| \leq |V(G)|$ and $k' \leq k$, and a good modulator of G' having size $O(\min(k^3 \log k, k^2 \log^2 |V(G)|))$ in time bounded above by a polynomial in $|V(G)|$.*

Proof. We first try to find a modulator S of G having size $O(\min(k^2 \log k, k \log^2 |V(G)|))$ by using Corollary 4.3. If it fails, then (G, k) is a no-instance, and therefore we take $(K_{2,2}, 0)$ as (G', k') and $V(K_{2,2})$ as a good modulator of G' . Otherwise, for each vertex v in S , let $G_v := G \setminus (S \setminus \{v\})$, and $F_1^v, \dots, F_{m(v)}^v$ be a maximal packing of small obstructions in G_v such that $V(F_i^v) \cap V(F_j^v) = \{v\}$ for every distinct i and j in $\{1, \dots, m(v)\}$. Finally, let $G'_v := G_v \setminus ((V(F_1^v) \cup \dots \cup V(F_{m(v)}^v)) \setminus \{v\})$. If $m(v) \geq k + 1$ for some vertex $v \in S$, then we apply our algorithm recursively for $(G \setminus v, k - 1)$. This is safe, because (R1) is safe. Therefore, we may assume that $m(v) \leq k$ for every vertex $v \in S$.

By Lemma 4.4 for G'_v , $k - m(v)$, and v , we can either

- (1) find $k - m(v) + 1$ holes $H_1^v, \dots, H_{k-m(v)+1}^v$ in G'_v such that $V(H_i^v) \cap V(H_j^v) = \{v\}$ for every distinct i and j in $\{1, \dots, k - m(v) + 1\}$, or
- (2) find a set $S'_v \subseteq V(G'_v) \setminus \{v\}$ of size at most $12(k - m(v))$ such that $G'_v \setminus S'_v$ is chordal.

If (1) holds, then we apply our algorithm recursively for $(G \setminus v, k - 1)$. This is safe, because (R1) is safe. Therefore, we may assume that (2) holds for every vertex v in S . Then let $S_v := (V(F_1^v) \cup \dots \cup V(F_{m(v)}^v) \cup S'_v) \setminus \{v\}$. Note that $|S_v| \leq 4m(v) + 12(k - m(v)) \leq 12k$ and $G_v \setminus S_v$ is a 3-leaf power.

We take (G, k) as (G', k') and $X := S \cup \bigcup_{v \in S} S_v$ as a good modulator of G . Clearly, $|X| \leq |S| + 12k|S| = O(\min(k^3 \log k, k^2 \log^2 |V(G)|))$. It remains to argue that X is a good modulator of G . Suppose that H is an obstruction in G . Since S is a modulator of G , H has a vertex $v \in S$. If $|V(H) \cap S| = 1$, then H is an induced subgraph of G_v , and therefore H has at least one vertex in S_v . Since S_v and S are disjoint, H has at least two vertices in X . Therefore, X is a good modulator of G . \square

5 Bounding the number of components outside of a good modulator

Let S be a good modulator of a graph G . In this section, we bound the number of non-trivial components of $G \setminus S$.

In Subsection 5.1, we introduce a complete split of a graph, and present two lemmas observing obstructions with a complete split of a graph. Then we define a blocking pair for a set of vertices, and present a characterization of a complete split of a graph and a lemma observing obstructions with a common blocking pair for two sets of vertices. All lemmas introduced in

this subsection will be used in the next subsection to bound the number of non-trivial components of $G \setminus S$.

In Subsection 5.2, we partition S into S^+ and S^- , and bound the number of components of $G \setminus S$ having neighbors of S^- . Afterward, we design a reduction rule to bound the number of non-trivial components of $G \setminus S$ having no neighbors of S^- .

In Subsection 5.3, we bound the number of isolated vertices of $G \setminus S$.

5.1 Complete splits and blocking pairs

Cunningham [11] introduced a split of a graph. A *split* of a graph G is a partition (A, B) of $V(G)$ such that $|A| \geq 2$, $|B| \geq 2$, and $N(A)$ is complete to $N(B)$. We say that a split (A, B) of G is *complete* if $N(A) \cup N(B)$ is a clique. The following two lemmas observe obstructions from the view of a complete split of a graph.

Lemma 5.1. *Let (A, B) be a complete split of a graph G . If G has a hole H , then $V(H) \cap A = \emptyset$ or $V(H) \cap B = \emptyset$.*

Proof. Suppose not. Since $N(A) \cup N(B)$ is a clique, H has at most two vertices in $N(A) \cup N(B)$, because otherwise H has K_3 as a subgraph. Since both $V(H) \cap A$ and $V(H) \cap B$ are non-empty, and H is connected, H has at least two vertices in $N(A) \cup N(B)$. Therefore, H has exactly two vertices x_1 and x_2 in $N(A) \cup N(B)$, a contradiction, because $H \setminus x_1 x_2$ is disconnected. \square

Lemma 5.2. *Let (A, B) be a complete split of a graph G . If G has an obstruction H having exactly two vertices a_1 and a_2 in A , then a_1 is adjacent to a_2 , one of a_1 and a_2 has degree 1 in H , and the other vertex has degree 3 in H .*

Proof. Suppose that both a_1 and a_2 have neighbors in B . Since $N(A) \cup N(B)$ is a clique, a_1 and a_2 are adjacent, and have the same set of neighbors in B . Then a_1 and a_2 are true twins in H , a contradiction by (O1). Therefore, either a_1 or a_2 , say a_1 , has no neighbors in B . Since H is connected, a_1 is adjacent to a_2 . Thus, a_1 has degree 1 in H . By (O3), a_2 has at most three neighbors in H . By (O5), a_2 has at least three neighbors in H . Therefore, a_2 has degree 3 in H . \square

We remark that for a complete split (A, B) of a graph G , if G has an obstruction H having exactly two vertices in A , then H is isomorphic to the bull.

Now, we define a blocking pair for a set $X \subseteq V(G)$. A *blocking pair* for X is an unordered pair $\{v, w\}$ of distinct vertices in $N(X)$ such that if v and w are adjacent, and $N(v) \cap X = N(w) \cap X$, then $N(v) \cap X$ is not a clique. Note that if $v, w \in N(X)$ are not adjacent, or $N(v) \cap X \neq N(w) \cap X$, then $\{v, w\}$ is a blocking pair for X . We say that X is *blocked by* $\{v, w\}$ if $\{v, w\}$ is a blocking pair for X . We remark that if $N(X)$ has a blocking pair $\{v, w\}$ for some subset of X , then X is blocked by $\{v, w\}$ as well. This definition is motivated by the following lemma.

Lemma 5.3. *Let (A, B) be a partition of the vertex set of a graph G such that $|A| \geq 2$ and $|B| \geq 2$. Then (A, B) is a complete split of G if and only if $N(B)$ is a clique, and B has no blocking pairs for A .*

Proof. It is clear that if (A, B) is a complete split of G , then $N(B)$ is a clique, and B has no blocking pairs for A .

Conversely, suppose that $N(B)$ is a clique, and B has no blocking pairs for A . We may assume that $|N(A)| \geq 2$, because otherwise $N(A) \cup N(B)$ is a clique, and (A, B) is a complete split of G . Since B has no blocking pairs for A , $N(A)$ is a clique, because if $N(A)$ has a non-edge vw , then $\{v, w\}$ is a blocking pair for A . Moreover, $N(v) \cap A = N(w) \cap A$ for all vertices v and w in $N(A)$, because otherwise $\{v, w\}$ is a blocking pair for A . This means that $N(A)$ is complete to $N(B)$. Therefore, $N(A) \cup N(B)$ is a clique, and (A, B) is a complete split of G . \square

The following lemma shows that a blocking pair $\{v, w\}$ for a set $X \subseteq V(G)$ tells us not only that $(X, V(G) \setminus X)$ is not a complete split of G , but also that G is not a 3-leaf power if $G[X]$ has two distinct components whose vertex sets are blocked by $\{v, w\}$.

Lemma 5.4. *Let (A, B) be a partition of the vertex set of a graph G such that $|A| \geq 2$ and $|B| \geq 2$. If $G[A]$ has distinct components C_1 and C_2 such that both $V(C_1)$ and $V(C_2)$ are blocked by $\{v, w\}$ of vertices in B , then $G[V(C_1) \cup V(C_2) \cup \{v, w\}]$ is not a 3-leaf power.*

Proof. If $N_G(v) \cap V(C_1) \neq N_G(w) \cap V(C_1)$, then we may assume that C_1 has a vertex u_1 adjacent to v and non-adjacent to w , because otherwise we can swap v and w . Let u_2 be a neighbor of v in $V(C_2)$, and P be an induced path in $G[V(C_1) \cup V(C_2) \cup \{w\}]$ from u_1 to u_2 . Note that the length of P is at least 3, because P must intersect w that is non-adjacent to u_1 . Since v is adjacent to both ends of P , $G[V(P) \cup \{v\}]$ is not distance-hereditary by Lemma 2.6. Therefore, $G[V(C_1) \cup V(C_2) \cup \{v, w\}]$ is not a 3-leaf power.

Therefore, $N_G(v) \cap V(C_i) = N_G(w) \cap V(C_i)$ for $i = 1, 2$. If v and w are non-adjacent, then for a neighbor u_1 of v in $V(C_1)$ and a neighbor u_2 of v in $V(C_2)$, $G[\{v, w, u_1, u_2\}]$ is a hole. Therefore, we may assume that v and w are adjacent. Then since $\{v, w\}$ is a blocking pair for $V(C_1)$, $N_G(v) \cap V(C_1)$ has a non-edge $u_1 u_2$. Let P be an induced path in C_1 from u_1 to u_2 . Since v is adjacent to both ends of P , we may assume that the length of P is exactly 2 by Lemma 2.6. Let u_3 be a common neighbor of u_1 and u_2 in $V(P)$, and u_4 be a neighbor of v in $V(C_2)$. Then $G[\{v, u_1, u_2, u_3, u_4\}]$ is isomorphic to the dart if u_3 is adjacent to v , and has a hole of length 4 if u_3 is non-adjacent to v . Therefore, $G[V(C_1) \cup V(C_2) \cup \{v, w\}]$ is not a 3-leaf power. \square

5.2 The number of non-trivial components

Let S^+ be the set of vertices v in S such that for each component C of $G \setminus S$, $N_G(v) \cap V(C)$ is a true twin-set in C , and $S^- := S \setminus S^+$. The following proposition shows that $G \setminus S$ has at most $|S^-|$ components having neighbors of S^- .

Proposition 5.5. *Let S be a good modulator of a graph G , v be a vertex in S , and C be a component of $G \setminus S$. If $N_G(v) \cap V(C)$ contains distinct vertices w_1 and w_2 that are not true twins in C , then no components of $G \setminus S$ different from C have neighbors of v .*

Proof. Suppose that there is a component of $G \setminus S$ different from C having a neighbor w of v . If w_1 and w_2 are adjacent, then we may assume that C has a vertex w_3 adjacent to exactly one of w_1 and w_2 , because w_1 and w_2 are not true twins in C . Then $G[\{v, w, w_1, w_2, w_3\}]$ is isomorphic to the dart if w_3 is adjacent to v , and the bull if w_3 is non-adjacent to v , a contradiction, because it has exactly one vertex v in S , which is a good modulator of G .

Therefore, w_1 and w_2 are non-adjacent. Let P be an induced path in C from w_1 to w_2 . Since v is adjacent to both ends of P , and S is a good modulator of G , we may assume that the length of P is exactly 2 by Lemma 2.6. Let w_3 be a common neighbor of w_1 and w_2 in $V(P)$. Then $G[\{v, w, w_1, w_2, w_3\}]$ is isomorphic to the dart if w_3 is adjacent to v , and has a hole of length 4 if w_3 is non-adjacent to v , a contradiction, because it has exactly one vertex v in S . \square

We present a reduction rule to bound the number of non-trivial components of $G \setminus S$ having no neighbors of S^- . We will use the following definition to design such a reduction rule.

Let X be a set of vertices of a graph Q . For a non-negative integer ℓ , a set $M \subseteq E(Q)$ is an (X, ℓ) -*matching* of Q if every vertex in X is incident with at most ℓ edges in M , and every vertex in $V(Q) \setminus X$ is incident with at most one edge in M .

Reduction Rule 2 (R2). *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , let S^+ be the set of vertices u in S such that for each component C of $G \setminus S$, $N_G(u) \cap V(C)$ is a true twin-set in C , X be the set of 2-element subsets of S^+ , and Y be the set of non-trivial components of $G \setminus S$ having no neighbors of $S \setminus S^+$. Let Q be a bipartite graph on $(X \times \{1, 2, 3\}, Y)$ such that the following three statements are true.*

- (1) *Elements $(\{v, w\}, 1) \in X \times \{1\}$ and $C \in Y$ are adjacent in Q if and only if $V(C)$ is blocked by $\{v, w\}$.*
- (2) *Elements $(\{v, w\}, 2) \in X \times \{2\}$ and $C \in Y$ are adjacent in Q if and only if C has a vertex adjacent to both v and w .*
- (3) *Elements $(\{v, w\}, 3) \in X \times \{3\}$ and $C \in Y$ are adjacent in Q if and only if C has an edge xy such that x is adjacent to both v and w , and y is non-adjacent to both v and w .*

If Q has a maximal $(X \times \{1, 2, 3\}, k+2)$ -matching M avoiding some element U in Y , then replace (G, k) with $(G \setminus E(U), k)$.

Proof of Safeness. Let $G' := G \setminus E(U)$. We need to show that (G, k) is a yes-instance if and only if (G', k) is a yes-instance.

Suppose that G has a modulator S' of size at most k , and $G' \setminus S'$ has an obstruction H . Since $G \setminus S'$ is a 3-leaf power, H has vertices b_1 and b_2 such that $b_1 b_2 \in E(U \setminus S')$. Thus, $|V(U) \setminus S'| \geq 2$.

Claim 1. We claim that $(V(U) \setminus S', V(G) \setminus (V(U) \cup S'))$ is a split of $G' \setminus S'$.

We first show that $|V(G) \setminus (V(U) \cup S')| \geq 2$. If H is a hole of length 4, then H has at most two vertices of $U \setminus S'$, because no holes of length 4 have independent sets of size at least 3, and $V(U) \setminus S'$ is an independent set of $G' \setminus S'$. Therefore, H has at least two vertices of $G \setminus (V(U) \cup S')$.

If H is not a hole of length 4, then $|V(H)| \geq 5$. By (O2), H has at most three vertices of $U \setminus S'$, because $V(U) \setminus S'$ is an independent set of $G' \setminus S'$. Therefore, H has at least two vertices of $G \setminus (V(U) \cup S')$, and $|V(G) \setminus (V(U) \cup S')| \geq 2$.

Now, suppose that $(V(U) \setminus S', V(G) \setminus (V(U) \cup S'))$ is not a split of $G' \setminus S'$. Then $G \setminus (V(U) \cup S')$ has vertices v and w such that both v and w have neighbors in $V(U) \setminus S'$, and $N_G(v) \cap (V(U) \setminus S') \neq N_G(w) \cap (V(U) \setminus S')$. Thus, $\{v, w\}$

is a blocking pair for $V(U) \setminus S'$, so for $V(U)$. Then U is adjacent to $(\{v, w\}, 1)$ in Q . Since M is maximal, Y has distinct elements C_1, \dots, C_{k+2} different from U such that $V(C_i)$ is blocked by $\{v, w\}$ for each $i \in \{1, \dots, k+2\}$. Since $|S'| \leq k$, two of them, say C_1 and C_2 , have no vertices in S' . Then $G[V(C_1) \cup V(C_2) \cup \{v, w\}]$ is not a 3-leaf power by Lemma 5.4, a contradiction, because it is an induced subgraph of $G \setminus S'$, and this proves the claim.

Since $V(U) \setminus S'$ is an independent set of $G' \setminus S'$, and H is connected, both b_1 and b_2 have neighbors in $V(G) \setminus (V(U) \cup S')$. Then by Claim 1, b_1 and b_2 are false twins in $G' \setminus S'$. By (O4), both b_1 and b_2 have degree 2 in H . Let z_1 and z_2 be the neighbors of b_1 in $V(H) \cap S$. Then U is adjacent to $(\{z_1, z_2\}, 2)$ in Q . Since M is maximal, Y has distinct elements C'_1, \dots, C'_{k+2} different from U such that C'_i has a vertex adjacent to both z_1 and z_2 for each $i \in \{1, \dots, k+2\}$. Since $|S'| \leq k$, two of them, say C'_1 and C'_2 , have no vertices in S' . Note that S' has no vertices of H , because H is an induced subgraph of $G' \setminus S'$.

If z_1 and z_2 are non-adjacent, then $G[V(C'_1) \cup V(C'_2) \cup \{z_1, z_2\}]$ has a hole of length 4, a contradiction, because it is an induced subgraph of $G \setminus S'$. Therefore, z_1 and z_2 are adjacent. Since $G[\{b_1, z_1, z_2\}]$ is isomorphic to K_3 , H is not a hole, and therefore $|V(H)| = 5$. Let a be a vertex of H different from b_1, b_2, z_1 , and z_2 . We may assume that a is not in $V(C'_1)$, because otherwise we may swap C'_1 and C'_2 . Let c be a vertex of C'_1 adjacent to both z_1 and z_2 . Note that $G[\{b_1, b_2, z_1, z_2\}]$ is isomorphic to $K_4 \setminus b_1 b_2$.

Since the dart and a hole of length 4 are the only obstructions having false twins, H is isomorphic to the dart. Thus, we may assume that $N_H(a) = \{z_1\}$. Then $G[\{a, b_1, c, z_1, z_2\}]$ is isomorphic to the gem if c is adjacent to a , and the dart if c is non-adjacent to a , a contradiction, because it is an induced subgraph of $G \setminus S'$. Therefore, if (G, k) is a yes-instance, then so is (G', k) .

Conversely, suppose that G' has a modulator S' of size at most k , and $G \setminus S'$ has an obstruction H . Since $G' \setminus S'$ is a 3-leaf power, H has an edge of $U \setminus S'$. Thus, $|V(U) \setminus S'| \geq 2$. Since S is a good modulator of G , H has at least two vertices in $S \setminus S'$. Then $|V(G) \setminus (V(U) \cup S')| \geq 2$, because $S \setminus S' \subseteq V(G) \setminus (V(U) \cup S')$.

Claim 2. We claim that $(V(U) \setminus S', V(G) \setminus (V(U) \cup S'))$ is a complete split of $G \setminus S'$.

Suppose not. We first show that $V(G) \setminus (V(U) \cup S')$ has a blocking pair for $V(U) \setminus S'$. Since U is a component of $G \setminus S$, and has no neighbors of $S \setminus S^+$, it suffices to show that $S^+ \setminus S'$ has a blocking pair for $V(U) \setminus S'$. We may assume

that for all vertices v and w in $S^+ \setminus S'$ having neighbors in $V(U) \setminus S'$, v and w are adjacent, and have the same set of neighbors in $V(U) \setminus S'$, because otherwise $\{v, w\}$ is a blocking pair for $V(U) \setminus S'$. For each vertex v in $S^+ \setminus S'$ having neighbors in $V(U) \setminus S'$, the set of neighbors of v in $V(U) \setminus S'$ is a true twin-set in $U \setminus S'$, that is, a clique. Therefore, $N_G(S^+ \setminus S') \cap (V(U) \setminus S')$ is a clique of $U \setminus S'$. Thus, by Lemma 5.3, $S^+ \setminus S'$ has a blocking pair $\{v, w\}$ for $V(U) \setminus S'$, so for $V(U)$.

Since $V(U)$ is blocked by $\{v, w\}$, U is adjacent to $(\{v, w\}, 1)$ in Q . Since M is maximal, Y has distinct elements C_1, \dots, C_{k+2} different from U such that $V(C_i)$ is blocked by $\{v, w\}$ for each $i \in \{1, \dots, k+2\}$. Since $|S'| \leq k$, two of them, say C_1 and C_2 , have no vertices in S' . Then $G[V(C_1) \cup V(C_2) \cup \{v, w\}]$ is not a 3-leaf power by Lemma 5.4, a contradiction, because it is an induced subgraph of $G' \setminus S'$, and this proves the claim.

Since both $U \setminus S'$ and $G \setminus (V(U) \cup S')$ have vertices of H , H is not a hole by Lemma 5.1 and Claim 2, and therefore $|V(H)| = 5$. Let t_1, \dots, t_p be the vertices of H in $V(U) \setminus S'$, and s_1, \dots, s_q be the vertices of H in $V(G) \setminus (V(U) \cup S')$. Note that both p and q are at least 2. Since $|V(H)| = 5$, $(p, q) = (3, 2)$ or $(p, q) = (2, 3)$.

If $(p, q) = (3, 2)$, then we may assume that $N_H(s_1) = \{s_2\}$ and $N_H(s_2) = \{s_1, t_1, t_2\}$ by Lemma 5.2 and Claim 2. Since U has no neighbors of $S \setminus S^+$, s_2 is in S^+ . Thus, t_1 and t_2 are true twins in $U \setminus S'$, a contradiction by (O1).

Therefore, $(p, q) = (2, 3)$. By Lemma 5.2 and Claim 2, we may assume that $N_H(t_1) = \{t_2\}$ and $N_H(t_2) = \{t_1, s_1, s_2\}$. Note that s_1 and s_2 are in $S \setminus S'$. Then U is adjacent to $(\{s_1, s_2\}, 3)$ in Q . Since M is maximal, Y has distinct elements C_1'', \dots, C_{k+2}'' different from U such that C_i'' has an edge $x_i y_i$ such that x_i is adjacent to both s_1 and s_2 , and y_i is non-adjacent to both s_1 and s_2 for each $i \in \{1, \dots, k+2\}$. Since $|S'| \leq k$, two of them, say C_1'' and C_2'' , have no vertices in S' . We may assume that s_3 is not in $V(C_1'')$, because otherwise we may swap C_1'' and C_2'' . We remark that the bull is the only possible graph to which H is isomorphic. Thus, s_1 and s_2 are adjacent, and s_3 is adjacent to exactly one of s_1 and s_2 in H . Then $G[\{x_1, y_1, s_1, s_2, s_3\}]$ is isomorphic to the gem if both x_1 and y_1 are adjacent to s_3 , the bull if both x_1 and y_1 are non-adjacent to s_3 , and the dart if x_1 is adjacent to s_3 and y_1 is non-adjacent to s_3 , and has a hole of length 4 if x_1 is non-adjacent to s_3 and y_1 is adjacent to s_3 , a contradiction, because it is an induced subgraph of $G' \setminus S'$. Therefore, if (G', k) is a yes-instance, then so is (G, k) . \square

Proposition 5.6. *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , if (R2) is not applicable to (G, k) , then $G \setminus S$ has at*

most $2(k+2)|S|^2$ non-trivial components.

Proof. Let S^+ be the set of vertices u in S such that for each component C of $G \setminus S$, $N_G(u) \cap V(C)$ is a true twin-set in C , and $S^- := S \setminus S^+$. By Proposition 5.5, each vertex in S^- is adjacent to at most one component of $G \setminus S$. Therefore, $G \setminus S$ has at most $|S^-|$ non-trivial components having neighbors of S^- .

Let Q and M be defined as in (R2). Since (R2) is not applicable to (G, k) , each non-trivial component of $G \setminus S$ having no neighbors of S^- is incident with exactly one edge in M . Since each edge in M is incident with some element in $X \times \{1, 2, 3\}$, and each element in $X \times \{1, 2, 3\}$ is incident with at most $k+2$ edges, $|M| \leq (k+2) \cdot |X \times \{1, 2, 3\}| \leq (k+2) \cdot 3 \binom{|S^+|}{2} \leq 3(k+2)|S|^2/2$. Then $|S^-| + |M| \leq |S| + 3(k+2)|S|^2/2 \leq (k+2)|S|^2/2 + 3(k+2)|S|^2/2 = 2(k+2)|S|^2$, and therefore $G \setminus S$ has at most $2(k+2)|S|^2$ non-trivial components. \square

5.3 The number of isolated vertices

We present a reduction rule to bound the number of isolated vertices of $G \setminus S$.

Reduction Rule 3 (R3). *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , let \mathcal{A} be the set of ordered pairs (A_1, A_2) of disjoint subsets of S such that $2 \leq |A_1| + |A_2| \leq 4$, and X be the set of isolated vertices of $G \setminus S$. For each element (A_1, A_2) in \mathcal{A} , let X_{A_1, A_2} be a maximal set of vertices v in X such that $N_G(v) \cap (A_1 \cup A_2) = A_1$ and $|X_{A_1, A_2}| \leq k+3$. If X has a vertex u not in $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$, then replace (G, k) with $(G \setminus u, k)$.*

Proof of Safeness. We need to show that if $(G \setminus u, k)$ is a yes-instance, then so is (G, k) . Suppose that $G \setminus u$ has a modulator S' of size at most k , and $G \setminus S'$ has an obstruction H . Then $u \in V(H)$, because $G \setminus (S' \cup \{u\})$ is a 3-leaf power.

If H is a hole, then u has exactly two neighbors v_1 and v_2 in $V(H) \cap S$ such that v_1 is non-adjacent to v_2 . By the construction of $X_{\{v_1, v_2\}, \emptyset}$, $X_{\{v_1, v_2\}, \emptyset}$ contains distinct vertices u_1, \dots, u_{k+3} different from u . Note that H has at most one of u_1, \dots, u_{k+3} , because v_1 and v_2 have at most two common neighbors in $V(H)$ including u . Then since $|S'| \leq k$, two of them, say u_1 and u_2 , are not in $S' \cup V(H)$. Thus, $G[\{v_1, v_2, u_1, u_2\}]$ is a hole, a contradiction, because it is an induced subgraph of $G \setminus (S' \cup \{u\})$.

Suppose that H is isomorphic to the bull, the dart, or the gem. Then $2 \leq |S \cap V(H)| \leq 4$, because H has exactly five vertices including u , and

S is a good modulator of G . Let $B_1 := (S \cap V(H)) \cap N_G(u)$, and $B_2 := (S \cap V(H)) \setminus N_G(u)$. By the construction of X_{B_1, B_2} , X_{B_1, B_2} contains distinct vertices u_1, \dots, u_{k+3} different from u . Since $|V(H)| = 5$ and $2 \leq |S \cap V(H)| \leq 4$, H has at most three vertices in X including u . Thus, H has at most two of u_1, \dots, u_{k+3} . Then since $|S'| \leq k$, one of them, say u_1 , is not in $S' \cup V(H)$. Thus, $G[(V(H) \setminus \{u\}) \cup \{u_1\}]$ is isomorphic to H , a contradiction, because it is an induced subgraph of $G \setminus (S' \cup \{u\})$. \square

Proposition 5.7. *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , if (R3) is not applicable to (G, k) , then $G \setminus S$ has at most $2(k+3)|S|^4/3$ isolated vertices.*

Proof. Let \mathcal{A} , X , and X_{A_1, A_2} be defined as in (R3). If $|S| \leq 1$, then $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$ is empty, and therefore X is empty. Thus, we may assume that $|S| \geq 2$. Let $s := |S|$. For each m -element subset T of S with $2 \leq m \leq 4$, \mathcal{A} contains exactly 2^m elements (A_1, A_2) such that $T = A_1 \cup A_2$. Therefore, $|\mathcal{A}|$ is at most

$$\begin{aligned} 2^4 \cdot \binom{s}{4} + 2^3 \cdot \binom{s}{3} + 2^2 \cdot \binom{s}{2} &\leq \frac{2}{3}(s-1)^4 + \frac{4}{3}(s-1)^3 + 2s(s-1) \\ &= \frac{2}{3}(s-1)^2((s-1)^2 + 2(s-1)) + 2s(s-1) \\ &\leq \frac{2}{3}(s-1)^2 s^2 + 2s(s-1) \\ &= 2s(s-1)(s^2 - s + 3)/3 \\ &\leq 2s(s-1)(s^2 + s)/3 = 2s^2(s^2 - 1)/3 \leq 2s^4/3. \end{aligned}$$

For each element (A_1, A_2) in \mathcal{A} , $|X_{A_1, A_2}| \leq k+3$. Therefore, $|\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}| \leq 2(k+3)|S|^4/3$. Since (R3) is not applicable to (G, k) , every isolated vertex of $G \setminus S$ is in $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$. Therefore, $G \setminus S$ has at most $2(k+3)|S|^4/3$ isolated vertices. \square

6 Bounding the size of components outside of a good modulator

Let S be a good modulator of a graph G . In this section, we bound the size of each component of $G \setminus S$. Subsection 6.1 is about complete components of $G \setminus S$, and Subsection 6.2 is about incomplete components of $G \setminus S$.

6.1 The size of each complete component

We present a reduction rule to bound the size of each complete component of $G \setminus S$.

Reduction Rule 4 (R4). *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , let \mathcal{A} be the set of ordered pairs (A_1, A_2) of disjoint subsets of S such that $2 \leq |A_1| + |A_2| \leq 4$, and C be a complete component of $G \setminus S$. For each element (A_1, A_2) in \mathcal{A} , let X_{A_1, A_2} be a maximal set of vertices v of C such that $N_G(v) \cap (A_1 \cup A_2) = A_1$ and $|X_{A_1, A_2}| \leq k + 3$. If C has a vertex u not in $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$, then replace (G, k) with $(G \setminus u, k)$.*

Proof of Safeness. We need to show that if $(G \setminus u, k)$ is a yes-instance, then so is (G, k) . Suppose that $G \setminus u$ has a modulator S' of size at most k , and $G \setminus S'$ has an obstruction H . Then $u \in V(H)$, because $G \setminus (S' \cup \{u\})$ is a 3-leaf power.

Suppose that H is a small obstruction. Since H has at most five vertices including u , and S is a good modulator of G , $2 \leq |S \cap V(H)| \leq 4$. Let $B_1 := (S \cap V(H)) \cap N_G(u)$, and $B_2 := (S \cap V(H)) \setminus N_G(u)$. By the construction of X_{B_1, B_2} , X_{B_1, B_2} contains distinct vertices u_1, \dots, u_{k+3} different from u . Since $|V(H)| \leq 5$ and $2 \leq |S \cap V(H)| \leq 4$, H has at most three vertices of C including u . Thus, H has at most two of u_1, \dots, u_{k+3} . Then since $|S'| \leq k$, one of them, say u_1 , is not in $S' \cup V(H)$. Thus, $G[(V(H) \setminus \{u\}) \cup \{u_1\}]$ is isomorphic to H , a contradiction, because it is an induced subgraph of $G \setminus (S' \cup \{u\})$.

Therefore, H is a hole of length at least 6. Note that H has at most two vertices of C , because C is complete. Suppose that H has exactly one vertex u of C . In this case, u is adjacent to distinct vertices v_1 and v_2 in $V(H) \cap S$. Then $H \setminus u$ is an induced path of length at least 4 from v_1 to v_2 . By the construction of $X_{\{v_1, v_2\}, \emptyset}$, $X_{\{v_1, v_2\}, \emptyset}$ contains distinct vertices u_1, \dots, u_{k+3} different from u . Since $|S'| \leq k$, one of them, say u_1 , is not in S' . Then $G[(V(H) \setminus \{u\}) \cup \{u_1\}]$ is not distance-hereditary by Lemma 2.6, a contradiction, because it is an induced subgraph of $G \setminus (S' \cup \{u\})$.

Therefore, H has exactly two vertices u and u' of C . In this case, u is adjacent to a vertex v_1 in $V(H) \cap S$, and u' is adjacent to another vertex v_2 in $V(H) \cap S$. Note that u' is non-adjacent to v_1 . Then $H \setminus u$ is an induced path of length at least 4 from v_1 to u' . By the construction of $X_{\{v_1\}, \{v_2\}}$, $X_{\{v_1\}, \{v_2\}}$ contains distinct vertices u_1, \dots, u_{k+3} different from u . Since $|S'| \leq k$, one of them, say u_1 , is not in S' . Then $G[(V(H) \setminus \{u\}) \cup \{u_1\}]$ is not distance-hereditary by Lemma 2.6, a contradiction, because it is an induced subgraph

of $G \setminus (S' \cup \{u\})$. Therefore, if $(G \setminus u, k)$ is a yes-instance, then so is (G, k) . \square

Proposition 6.1. *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , if (R4) is not applicable to (G, k) , then every complete component of $G \setminus S$ has at most $2(k + 3)|S|^4/3$ vertices.*

Proof. Let \mathcal{A} , C , and X_{A_1, A_2} be defined as in (R4). Since (R4) is not applicable to (G, k) , every vertex of C is in $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$. If $|S| \leq 1$, then $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$ is empty, contradicting the assumption that every vertex of C is in $\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}$. Thus, $|S| \geq 2$. For each m -element subset T of S with $2 \leq m \leq 4$, \mathcal{A} contains exactly 2^m elements (A_1, A_2) such that $T = A_1 \cup A_2$. Therefore, $|\mathcal{A}| \leq 2^4 \cdot \binom{|S|}{4} + 2^3 \cdot \binom{|S|}{3} + 2^2 \cdot \binom{|S|}{2} \leq 2|S|^4/3$, as in the proof of Proposition 5.7. For each element (A_1, A_2) in \mathcal{A} , $|X_{A_1, A_2}| \leq k + 3$. Therefore, $|\bigcup_{(A_1, A_2) \in \mathcal{A}} X_{A_1, A_2}| \leq 2(k + 3)|S|^4/3$, and C has at most $2(k + 3)|S|^4/3$ vertices. \square

6.2 The size of each incomplete component

We present a reduction rule to bound the size of each true twin-set in G .

Reduction Rule 5 (R5). *Given an instance (G, k) with $k > 0$, if G has a true twin-set X such that $v \in X$ and $|X| \geq k + 2$, then replace (G, k) with $(G \setminus v, k)$.*

Proof of Safeness. We need to show that if $(G \setminus v, k)$ is a yes-instance, then so is (G, k) . Suppose that $G \setminus v$ has a modulator S of size at most k , and $G \setminus S$ has an obstruction H . Then $v \in V(H)$, because $G \setminus (S \cup \{v\})$ is a 3-leaf power. By (O1), $V(H) \cap X = \{v\}$. Since $|S| \leq k$, X contains a vertex w not in $S \cup \{v\}$. Then $G[(V(H) \setminus \{v\}) \cup \{w\}]$ is isomorphic to H , a contradiction, because it is an induced subgraph of $G \setminus (S \cup \{v\})$. \square

We present a reduction rule to remove some bags of $G \setminus S$ anti-complete to S .

Reduction Rule 6 (R6). *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , let B be a non-empty true twin-set in $G \setminus S$. If $G \setminus (S \cup B)$ has a component D having no neighbors of S such that $V(D) \setminus N_G(B)$ is non-empty, then replace (G, k) with $(G \setminus (V(D) \setminus N_G(B)), k)$.*

Proof of Safeness. Let $G' := G \setminus (V(D) \setminus N_G(B))$. We need to show that if (G', k) is a yes-instance, then so is (G, k) . Suppose that G' has a modulator S' of size at most k , and $G' \setminus S'$ has an obstruction H . Since $G' \setminus S'$ is a 3-leaf power, H has at least one vertex of $D \setminus N_G(B)$. Since H is connected, and D

has no neighbors of S , H has at least one vertex in $V(D) \cap N_G(B)$. Thus, H has at least two vertices of D . Since $V(H) \cap S \neq \emptyset$, $V(H) \cap B$ is a clique cut-set of H and therefore H is not a hole. (A *clique cut-set* of a connected graph is a clique whose deletion makes disconnected.) Thus, $|V(H)| = 5$. Since S is a good modulator of G , $|V(H) \cap S| = 2$, $|V(H) \cap B| = 1$, and $|V(H) \cap V(D)| = 2$, a contradiction, because no obstruction has a cut vertex partitioning its vertex set into two sets both having size 2. \square

We present two reduction rules to reduce the number of bags of $G \setminus S$.

Reduction Rule 7 (R7). *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , let B be a non-empty true twin-set in $G \setminus S$. If $G \setminus (S \cup B)$ has distinct components D_1, \dots, D_{k+4} such that $N_G(V(D_1)) = \dots = N_G(V(D_{k+4}))$, and either $V(D_1) \cup \dots \cup V(D_{k+4}) \subseteq N_G(B)$, or $\emptyset \neq V(D_i) \cap N_G(B) \neq V(D_i)$ for each $i \in \{1, \dots, k+4\}$, then replace (G, k) with $(G \setminus V(D_1), k)$.*

To show that (R7) is safe, we will use the following two lemmas. Lemma 6.2 will be useful because it implies that for a good modulator S of G , a subset B of $V(G) \setminus S$ is a true twin-set in $G \setminus S$ if and only if it is a true twin-set in G .

Lemma 6.2. *Let G be a 3-leaf power having a vertex v such that $G \setminus v$ is connected and incomplete. Then vertices t_1 and t_2 in $V(G) \setminus \{v\}$ are true twins in G if and only if t_1 and t_2 are true twins in $G \setminus v$.*

Proof. It is clear that if t_1 and t_2 are true twins in G , then so are in $G \setminus v$.

Conversely, suppose that t_1 and t_2 are true twins in $G \setminus v$, and v is adjacent to t_1 , and non-adjacent to t_2 . Note that $|N_G(t_2)| \geq 2$, because otherwise $G \setminus v$ is isomorphic to K_2 .

If $N_G(t_2)$ is a clique, then $G \setminus v$ has at least one vertex not in $N_G(t_2)$, because otherwise $G \setminus v$ is complete. Thus, G has an edge xy such that x is adjacent to both t_1 and t_2 , and y is non-adjacent to both t_1 and t_2 , because $G \setminus v$ is connected. Then $G[\{v, x, y, t_1, t_2\}]$ is isomorphic to the gem if both x and y are adjacent to v , the bull if both x and y are non-adjacent to v , and the dart if x is adjacent to v and y is non-adjacent to v , and has a hole of length 4 if x is non-adjacent to v and y is adjacent to v , a contradiction, because it is an induced subgraph of G .

Therefore, t_2 has distinct neighbors x and y such that x is non-adjacent to y . Then $G[\{v, x, y, t_1, t_2\}]$ has a hole of length 4 if both x and y are adjacent to v , and is isomorphic to the gem if exactly one of x and y is adjacent to v ,

and the dart if both x and y are non-adjacent to v , a contradiction, because it is an induced subgraph of G . \square

Lemma 6.3. *Let (A, B) be a complete split of a graph G , and S be a non-empty good modulator of G . If G has an obstruction H , and $S \subseteq B \setminus N(A)$, then H has at most one vertex in A .*

Proof. Suppose not. Since S is a good modulator of G , H has at least two vertices in S . Thus, H has vertices in both A and B . Since (A, B) is a complete split of G , H is not a hole by Lemma 5.1, and therefore $|V(H)| = 5$. Then $|V(H) \cap N(A)| \leq 5 - |V(H) \cap A| - |V(H) \cap S| \leq 5 - 2 - 2$, a contradiction, because no obstruction has a cut vertex partitioning its vertex set into two sets both having size 2. \square

Proof of Safeness of (R7). We need to show that if $(G \setminus V(D_1), k)$ is a yes-instance, then so is (G, k) . Suppose that $G \setminus V(D_1)$ has a modulator S' of size at most k , and $G \setminus S'$ has an obstruction H . Since $G \setminus (V(D_1) \cup S')$ is a 3-leaf power, H has at least one vertex of D_1 . Since S is a good modulator of G , $G \setminus (S \setminus \{v\})$ is a 3-leaf power for each vertex v in S . Thus, if v has a neighbor in a true twin-set X in $G \setminus S$, then $\{v\}$ is complete to X by Lemma 6.2. This means that every true twin-set in $G \setminus S$ is a true twin-set in G as well.

We claim that for each $i \in \{1, \dots, k+4\}$, $V(D_i) \cap N_G(B)$ is a true twin-set in $G \setminus S$. Suppose that $V(D_i) \cap N_G(B)$ contains two vertices x and y such that x is non-adjacent to y . Let P be an induced path in D_i from x to y . We may assume that P has length exactly 2 by Lemma 2.6. Let z be a common neighbor of x and y in $V(P)$. We may assume that $z \in N_G(B)$, because otherwise $V(P)$ with a vertex in B induces a hole of length 4. Then for a vertex v in B , and v' in $V(D_j) \cap N_G(B)$ for some $j \in \{1, \dots, k+4\} \setminus \{i\}$, $G[\{v, v', x, y, z\}]$ is isomorphic to the dart, contradicting the assumption that S is a modulator of G . Therefore, $V(D_i) \cap N_G(B)$ is a clique. Now, suppose that $G \setminus S$ has a vertex w adjacent to a vertex $t_1 \in V(D_i) \cap N_G(B)$ and non-adjacent to a vertex $t_2 \in V(D_i) \cap N_G(B)$. Note that w is a vertex of $D_i \setminus N_G(B)$. Then for a vertex v in B and a vertex v' of $V(D_j) \cap N_G(B)$ for some $j \in \{1, \dots, k+4\} \setminus \{i\}$, $G[\{v, v', w, t_1, t_2\}]$ is isomorphic to the bull, a contradiction, and this proves the claim.

Suppose that $V(D_1) \cup \dots \cup V(D_{k+4}) \subseteq N_G(B)$. By (O1), for each $i \in \{1, \dots, k+4\}$, D_i has at most one vertex of H . By (O2), at most three of D_1, \dots, D_{k+4} have vertices of H . Since $|S'| \leq k$, one of D_2, \dots, D_{k+4} , say D_j , has no vertices in $S' \cup V(H)$. Let t be a vertex in D_j . Since $N_G(V(D_1)) = N_G(V(D_j))$, s and t have the same set of neighbors in $V(H)$.

Then $G[(V(H) \setminus \{s\}) \cup \{t\}]$ is isomorphic to H , a contradiction, because it is an induced subgraph of $G \setminus (V(D_1) \cup S')$.

Therefore, $\emptyset \neq V(D_i) \cap N_G(B) \neq V(D_i)$ for each $i \in \{1, \dots, k+4\}$. We first show that $D_i \setminus N_G(B)$ has no neighbors of S . Suppose that $D_i \setminus N_G(B)$ has a neighbor p_i of some vertex v in S . Let $j \in \{1, \dots, k+4\} \setminus \{i\}$. Since $N_G(V(D_i)) = N_G(V(D_j))$, D_j has a neighbor p_j of v . Since some vertex in B has neighbors in both D_i and D_j , $G \setminus S$ has a path P from p_i to p_j . Note that the length of P is at least 3, because p_i is not in $N_G(B)$. Since v is adjacent to both ends of P , $G[V(P) \cup \{v\}]$ is not distance-hereditary by Lemma 2.6, a contradiction, because it is an induced subgraph of $G \setminus (S \setminus \{v\})$, and this proves the claim.

For each $i \in \{1, \dots, k+4\}$, since $V(D_i) \cap N_G(B)$ is a true twin-set in G , H has at most one vertex in $V(D_i) \cap N_G(B)$ by (O1). Let $D_{i,1}, \dots, D_{i,m(i)}$ be the components of $D_i \setminus N_G(B)$ for each $i \in \{1, \dots, k+4\}$. We claim that for each $j \in \{1, \dots, m(i)\}$, if $|V(D_{i,j})| \geq 2$, then $(V(D_{i,j}), V(G) \setminus V(D_{i,j}))$ is a complete split of G . Since $V(D_i) \cap N_G(B)$ is a true twin-set, and $D_i \setminus N_G(B)$ has no neighbors of S , it suffices to show that $N_G(N_G(B)) \cap V(D_{i,j})$ is a clique. Suppose that $N_G(N_G(B)) \cap V(D_{i,j})$ contains vertices x and y such that x and y are non-adjacent. Let P be an induced path in $D_{i,j}$ from x to y . We may assume that P has length exactly 2 by Lemma 2.6. Let z be a common neighbor of x and y in $V(P)$. We may assume that $z \in N_G(N_G(B))$, because otherwise P with a vertex v in $N_G(B) \cap V(D_i)$ induces a hole of length 4. Then for a vertex v' in B , $G[\{v, v', x, y, z\}]$ is isomorphic to the dart, a contradiction, and this proves the claim.

Therefore, each component of $D_i \setminus N_G(B)$ has at most one vertex of H by Lemma 6.3. Each $V(D_i) \cap N_G(B)$ has at most one vertex of H , because $V(D_i) \cap N_G(B)$ is a true twin-set. Therefore, at most one component of $D_i \setminus N_G(B)$ has a vertex of H , because H cannot have false twins of degree at most 1. By (O2), at most three of D_1, \dots, D_{k+4} have vertices of H . Since $|S'| \leq k$, one of D_2, \dots, D_{k+4} , say D_i , has no vertices in $S' \cup V(H)$. Note that H has a vertex s_1 in $V(D_1) \cap N_G(B)$, because $D_1 \setminus N_G(B)$ has no neighbors of S , H is connected, and has vertices in both S and $V(D_1)$. Let $t_1 t_2$ be an edge of D_i such that $t_1 \in V(D_i) \cap N_G(B)$ and $t_2 \in V(D_i) \setminus N_G(B)$. Since $N_G(V(D_1)) = N_G(V(D_i))$, and both $V(D_1) \cap N_G(B)$ and $V(D_i) \cap N_G(B)$ are true twin-sets, s_1 and t_1 have the same set of neighbors in $V(H) \setminus V(D_1)$. If H has a vertex s_2 in $V(D_1) \setminus N_G(B)$, then $V(D_1) \cap V(H) = \{s_1, s_2\}$, because both $V(D_1) \cap N_G(B)$ and $V(D_1) \setminus N_G(B)$ have at most one vertex of H . Then $G[(V(H) \setminus \{s_1, s_2\}) \cup \{t_1, t_2\}]$ is isomorphic to H , a contradiction, because it is an induced subgraph of $G \setminus (V(D_1) \cup S')$. Therefore, H has no vertices in $V(D_1) \setminus N_G(B)$. Then $G[(V(H) \setminus \{s_1\}) \cup \{t_1\}]$ is isomorphic to H ,

a contradiction, because it is an induced subgraph of $G \setminus (V(D_1) \cup S')$. \square

Reduction Rule 8 (R8). *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , let B_1, \dots, B_m be pairwise disjoint non-empty true twin-sets in $G \setminus S$ for $m \geq 6$ such that $N_G(B_i) = B_{i-1} \cup B_{i+1}$ for each $i \in \{2, \dots, m-1\}$. Let ℓ be an integer in $\{3, \dots, m-2\}$ such that $|B_\ell| \leq |B_i|$ for each $i \in \{3, \dots, m-2\}$, and G' be a graph obtained from $G \setminus ((B_3 \cup \dots \cup B_{m-2}) \setminus B_\ell)$ by making every vertex in B_ℓ adjacent to all vertices in $B_2 \cup B_{m-1}$. Then replace (G, k) with (G', k) .*

To show that (R8) is safe, we will use the following two lemmas.

Lemma 6.4. *Let G be a graph with disjoint true twin-sets B_1, \dots, B_m for $m \geq 5$ such that $N(B_i) = B_{i-1} \cup B_{i+1}$ for each $i \in \{2, \dots, m-1\}$. Then G is a 3-leaf power if and only if $G \setminus (B_3 \cup \dots \cup B_{m-2})$ is a 3-leaf power, and has no paths from a vertex in B_2 to a vertex in B_{m-1} .*

Proof. It is clear that if G is a 3-leaf power, then $G \setminus (B_3 \cup \dots \cup B_{m-2})$ is a 3-leaf power, and has no paths from a vertex in B_2 to a vertex in B_{m-1} , because otherwise G has a hole.

Conversely, suppose that $G \setminus (B_3 \cup \dots \cup B_{m-2})$ is a 3-leaf power, and has no paths from a vertex in B_2 to a vertex in B_{m-1} , and G has an obstruction H . Since $G \setminus (B_3 \cup \dots \cup B_{m-2})$ is a 3-leaf power, H has at least one vertex in $B_3 \cup \dots \cup B_{m-2}$. For each $i \in \{1, \dots, m\}$, since B_i is a true twin-set in G , H has at most one vertex in B_i by (O1). Then every vertex of H in $B_2 \cup \dots \cup B_{m-1}$ has degree at most 2 in H . If H has a vertex v in B_j for some $j \in \{3, \dots, m-2\}$, then both B_{j-1} and B_{j+1} have vertices of H by (O5). This means that B_i contains exactly one vertex of H for each $i \in \{2, \dots, m-1\}$. Then H has vertices in each B_1 and B_m as well by (O5). Thus, $V(H) \cap (B_2 \cup \dots \cup B_{m-1})$ contains at least three vertices of degree 2 in H . Then H is a hole by (O6), a contradiction, because $H \setminus (B_3 \cup \dots \cup B_{m-2})$ is a path in $G \setminus (B_3 \cup \dots \cup B_{m-2})$ from a vertex in B_2 to a vertex in B_{m-1} . Therefore, G is a 3-leaf power. \square

Lemma 6.5. *If G has a modulator S and a true twin-set X such that $X \setminus S$ is non-empty, then $S \setminus X$ is a modulator of G .*

Proof. We may assume that $S \cap X$ is non-empty. Suppose that $G \setminus (S \setminus X)$ has an obstruction H . Since $G \setminus S$ is a 3-leaf power, H has at least one vertex in $S \cap X$. Then H has exactly one vertex v in $S \cap X$ by (O1). Let w be a vertex in $X \setminus S$. Then $G[(V(H) \setminus \{v\}) \cup \{w\}]$ is isomorphic to H , a contradiction, because it is an induced subgraph of $G \setminus S$. \square

Proof of Safeness of (R8). First, let us show that if (G, k) is a yes-instance, then so is (G', k) . Suppose that G has a minimal modulator S' of size at most k . Since S' is minimal, $S' \cap B_i = \emptyset$ or $S' \cap B_i = B_i$ for each $i \in \{1, \dots, m\}$ by Lemma 6.5. We claim that if $S' \cap (B_1 \cup \dots \cup B_m)$ is empty, then S' is a modulator of G' . Since $G \setminus S'$ is a 3-leaf power, $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$ is a 3-leaf power, and has no paths from a vertex in B_2 to a vertex in B_{m-1} by Lemma 6.4. Since $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$ is isomorphic to $G' \setminus (B_\ell \cup S')$, $G' \setminus S'$ is a 3-leaf power by Lemma 6.4, and this proves the claim.

We claim that if $S' \cap (B_1 \cup B_2 \cup B_{m-1} \cup B_m)$ is non-empty, then $S' \cap V(G')$ is a modulator of G' . Suppose that $G' \setminus (S' \cap V(G'))$ has an obstruction H . Since $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$ is a 3-leaf power, and is isomorphic to $G' \setminus (B_\ell \cup (S' \cap V(G')))$, $G' \setminus (B_\ell \cup (S' \cap V(G')))$ is a 3-leaf power. Therefore, H has at least one vertex in B_ℓ . For each $i \in \{1, 2, \ell, m-1, m\}$, since B_i is a true twin-set in G' , H has at most one vertex in B_i by (O1). Then every vertex of H in $B_2 \cup B_\ell \cup B_{m-1}$ has degree at most 2 in H . Thus, for each $i \in \{1, 2, \ell, m-1, m\}$, B_i contains exactly one vertex of H by (O5). Then $S' \cap V(G')$ contains at least one vertex of H , a contradiction, because H is an induced subgraph of $G' \setminus (S' \cap V(G'))$, and this proves the claim.

Thus, we may assume that $S' \cap (B_1 \cup B_2 \cup B_{m-1} \cup B_m)$ is empty, and $S' \cap (B_3 \cup \dots \cup B_{m-2})$ is non-empty. Let $T := (S' \setminus (B_3 \cup \dots \cup B_{m-2})) \cup B_\ell$. Since $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$ is a 3-leaf power, and is isomorphic to $G' \setminus T$, $G' \setminus T$ is a 3-leaf power. Then $B_i \subseteq S'$ for some $i \in \{3, \dots, m-2\}$, and therefore $|T| = |T \setminus (B_3 \cup \dots \cup B_{m-2})| + |B_\ell| \leq |S' \setminus (B_3 \cup \dots \cup B_{m-2})| + |B_i| \leq |S'| \leq k$. Therefore, if (G, k) is a yes-instance, then so is (G', k) .

Secondly, we will show that if (G', k) is a yes-instance, then so is (G, k) . Suppose that G' has a minimal modulator S' of size at most k . Since S' is minimal, $S' \cap B_i = \emptyset$ or $S' \cap B_i = B_i$ for each $i \in \{1, 2, \ell, m-1, m\}$ by Lemma 6.5. We claim that S' is a modulator of G .

Since $G' \setminus S'$ is a 3-leaf power, if $S' \cap (B_1 \cup B_2 \cup B_{m-1} \cup B_m)$ is empty, then $G' \setminus (B_\ell \cup S')$ is a 3-leaf power, and has no paths from a vertex in B_2 to a vertex in B_{m-1} by Lemma 6.4. Since $G' \setminus (B_\ell \cup S')$ is isomorphic to $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$, $G \setminus S'$ is a 3-leaf power by Lemma 6.4.

Thus, we may assume that $S' \cap (B_1 \cup B_2 \cup B_{m-1} \cup B_m)$ is non-empty, and $G \setminus S'$ has an obstruction H . Since $G' \setminus (B_\ell \cup S')$ is a 3-leaf power, and is isomorphic to $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$, $G \setminus (B_3 \cup \dots \cup B_{m-2} \cup S')$ is a 3-leaf power. Therefore, H has at least one vertex in $B_3 \cup \dots \cup B_{m-2}$. For each $i \in \{1, \dots, m\}$, since B_i is a true twin-set in G , H has at most one vertex in B_i by (O1). Then every vertex of H in $B_2 \cup \dots \cup B_{m-1}$ has degree at most 2 in H . If H has a vertex v in B_j for some $j \in \{3, \dots, m-2\}$, then both

B_{j-1} and B_{j+1} have vertices of H by (O5). This means that B_i contains exactly one vertex of H for each $i \in \{2, \dots, m-1\}$ by (O5), and both B_1 and B_m have vertices of H as well by (O5). Thus, S' contains at least one vertex of H , a contradiction, and this proves the claim. Therefore, if (G', k) is a yes-instance, then so is (G, k) . \square

Now, we are ready to prove that after applying some reduction rules exhaustively to (G, k) with a good modulator S of G , each incomplete component of $G \setminus S$ has bounded size.

Proposition 6.6. *Given an instance (G, k) with $k > 0$ and a non-empty good modulator S of G , if (R2), (R5), (R6), (R7), and (R8) are not applicable to (G, k) , then each incomplete component of $G \setminus S$ has at most $(k+1)(k+4)|S|(|S|+2k+14)$ vertices.*

To prove Proposition 6.6, we will use the following lemma and its corollary.

Lemma 6.7. *Let G be a 3-leaf power. If G has a vertex v of degree at least 1 such that $G \setminus v$ is a tree, then $G \setminus v$ has a vertex u such that $N_G(v) = \{u\}$ or $N_G(v) = N_G[u] \setminus \{v\}$.*

Proof. We may assume that v has at least two neighbors, and $G \setminus v$ has at least three vertices, because otherwise the statement clearly holds.

If v has exactly two neighbors u_1 and u_2 , then u_1 and u_2 are adjacent, because otherwise G has a hole. Since $G \setminus v$ has at least three vertices, one of u_1 and u_2 , say u_2 , is not a leaf of $G \setminus v$. If u_1 is not a leaf of $G \setminus v$, then for a neighbor u'_1 of u_1 different from u_2 and a neighbor u'_2 of u_2 different from u_1 , $G[\{v, u_1, u'_1, u_2, u'_2\}]$ is isomorphic to the bull, a contradiction. Therefore, u_1 is a leaf of $G \setminus v$, and $N_G(v) = N_G[u_1] \setminus \{v\}$.

If v has at least three neighbors, then $G[N_G(v)]$ is connected, because otherwise G has a hole. If v has distinct neighbors u_1, u_2, u_3 , and u_4 inducing a path, then $G[\{v, u_1, u_2, u_3, u_4\}]$ is isomorphic to the gem, a contradiction. Therefore, $G \setminus v$ has a vertex u such that $N_G(v) \subseteq N_G(u)$. If u has a neighbor u_1 that is non-adjacent to v , then for distinct neighbors u_2 and u_3 of v different from u , $G[\{v, u, u_1, u_2, u_3\}]$ is isomorphic to the dart, a contradiction. Therefore, every neighbor of u different from v is adjacent to v , and $N_G(v) = N_G[u] \setminus \{v\}$. \square

Corollary 6.8. *Let G be a 3-leaf power. If G has a vertex v of degree at least 1 such that $G \setminus v$ is connected, then $G \setminus v$ has a true twin-set B such that $N_G(v) = B$ or $N_G(v) = N_G[B] \setminus \{v\}$.*

Proof. We may assume that $G \setminus v$ is incomplete. Then every true twin-set in $G \setminus v$ is also a true twin-set in G by Lemma 6.2. We proceed by induction on $|G|$. Suppose that $G \setminus v$ has true twins u and u' . Since u and u' are true twins, $G \setminus \{v, u\}$ is connected, and v has a neighbor in $V(G) \setminus \{u\}$. Then by applying the induction hypothesis, $G \setminus \{u, v\}$ has a true twin-set B' such that $N_G(v) \setminus \{u\}$ is equal to B' or $N_G[B'] \setminus \{u, v\}$. Let

$$B = \begin{cases} B' \cup \{u\} & \text{if } u' \in B', \\ B' & \text{otherwise.} \end{cases}$$

Since u and u' are true twins in $G \setminus v$, B is a true twin-set of $G \setminus v$. It is easy to see that if $N_G(v) \setminus \{u\} = B'$, then $N_G(v) = B$, because u and u' are true twins in G . If $N_G(v) \setminus \{u\} = N_G[B'] \setminus \{u, v\}$, then $N_G[B'] \setminus \{v\} = N_G[B] \setminus \{v\}$ by the same reason.

Thus, we may assume that $G \setminus v$ has no true twins. By Theorem 2.4, since $G \setminus v$ is connected, $G \setminus v$ is a tree. Then by Lemma 6.7, the statement holds. \square

Proof of Proposition 6.6. Let C be an incomplete component of $G \setminus S$ with a tree-clique decomposition $(F, \{B_u : u \in V(F)\})$. Since S is a good modulator of G , $G[V(C) \cup \{v\}]$ is a 3-leaf power for each vertex v in S . Thus, if S has a vertex w having a neighbor in a bag B of C , then $\{w\}$ is complete to B by Lemma 6.2. This means that every bag of C is a true twin-set in G . Since (R5) is not applicable to (G, k) , each bag of C contains at most $k + 1$ vertices. Therefore, in the remaining of this proof, we are going to bound the number of bags of C .

Claim 1. We claim that the maximum degree of F is at most $|S| + 2k + 6$.

Suppose that F has a node u of degree at least $|S| + 2k + 7$ in F . For each vertex w in S , if at least two components of $C \setminus B_u$ have neighbors of w , then all components of $C \setminus B_u$ have neighbors of w by Corollary 6.8. Thus, for each vertex w in S , we can choose a component of $C \setminus B_u$ such that either all other components of $C \setminus B_u$ have neighbors of w , or no other components of $C \setminus B_u$ have neighbors of w . Since $C \setminus B_u$ has at least $|S| + 2k + 7$ components, $C \setminus B_u$ has distinct components D_1, \dots, D_{2k+7} such that for each vertex w in S , either all or none of them have neighbors of w . Thus, $N_G(V(D_1)) = \dots = N_G(V(D_{2k+7}))$. By the pigeonhole principle, $V(D_i) \subseteq N_G(B_u)$ or $\emptyset \neq V(D_i) \cap N_G(B_u) \neq V(D_i)$ is satisfied by at least $k + 4$ values of i , contradicting the assumption that (R7) is not applicable to (G, k) , and this proves the claim.

Let X be the set of leaves of F whose bags are anti-complete to S .

Claim 2. We claim that if u is a node of $F \setminus X$ having degree at most 1 in $F \setminus X$, then B_u contains a neighbor of S .

If $N_F(u) \subseteq X$, then B_u contains a neighbor of S , because otherwise C has no neighbors of S , and (R2) is applicable to (G, k) . If $N_F(u) \setminus X$ is non-empty, then $N_F(u) \setminus X$ contains exactly one node u_1 , because u has degree at most 1 in $F \setminus X$. If B_u contains no neighbors of S , then (R6) is applicable to (G, k) by taking B_{u_1} as B . Therefore, B_u contains a neighbor of S , and this proves the claim.

For each vertex v in S , let X_v be the set of nodes of $F \setminus X$ whose bags contain neighbors of v , S_1 be the set of vertices v in S such that X_v contains some leaf of $F \setminus X$, and $S_2 := S \setminus S_1$. Let F' be a tree obtained from $F \setminus X$ by contracting all edges in $F[X_v]$ for each vertex v in S . Note that F' has at most $|S_1|$ leaves, and therefore it has at most $\max(|S_1| - 2, 0)$ branching nodes. Let Y be the set of nodes of F' which come from X_v for some vertex $v \in S$, and Z be the set of branching nodes of F' . Then $|Y \cup Z| \leq |Y| + |Z| \leq |S| + \max(|S_1| - 2, 0) \leq 2|S|$. Since (R8) is not applicable to (G, k) , each component of $F' \setminus (Y \cup Z)$ has at most three nodes. Therefore, $|V(F' \setminus (Y \cup Z))| \leq 6|S|$. Then $|V(F \setminus X)|$ is at most

$$\begin{aligned} |Y|(|S| + 2k + 7) + |Z| + |V(F' \setminus (Y \cup Z))| &\leq |S|(|S| + 2k + 7) + |S| + 6|S| \\ &= |S|(|S| + 2k + 14). \end{aligned}$$

Since (R7) is not applicable to (G, k) , each node of $F \setminus X$ is adjacent to at most $k + 3$ nodes in X . Thus, $|V(F)| \leq (k + 4)|S|(|S| + 2k + 14)$. By (R5), each bag of C has at most $k + 1$ nodes. Therefore, $|V(C)| \leq (k + 1)(k + 4)|S|(|S| + 2k + 14)$. \square

7 The proof of the main theorem

In this section, we prove Theorem 1.2.

Proof of Theorem 1.2. By Theorem 2.2, we can find all maximal true twin-sets in a 3-leaf power in linear time. Thus, we can apply (R2), \dots , (R8) in polynomial time to an input instance (G, k) with a good modulator S of G by investigating small subsets of $V(G)$ or true twin-sets in $G \setminus S$. Therefore, Algorithm 1 is a polynomial-time algorithm.

We claim that for the instance (G, k) obtained in Line 10, G has at most $O(k^2|S|^6)$ vertices. Note that $|S| \geq k + 1$. By Proposition 5.6, $G \setminus S$ has at most $2(k + 2)|S|^2$ non-trivial components. By Proposition 6.1, each complete component of $G \setminus S$ has at most $2(k + 3)|S|^4/3$ vertices. By Proposition 6.6,

Algorithm 1 Kernelization for 3-LEAF POWER DELETION

```

1: function COMPRESS( $G, k$ )
2:   if  $k = 0$  then
3:     if  $G$  is a 3-leaf power then return  $(K_1, 0)$ .
4:     else return  $(K_{2,2}, 0)$ .
5:   end if
6:   else Find an instance  $(G', k')$  equivalent to  $(G, k)$ , and a good modu-
    lator  $S$  of  $G'$  having size  $O(\min(k^3 \log k, k^2 \log^2 |V(G)|))$  by Lemma 4.5.
7:     if  $|S| \leq k$  then return  $(K_1, 0)$ .
8:     else if  $(G', k') \neq (G, k)$  then return COMPRESS( $G', k'$ ).
9:     else if Rule (Ri) for some  $i \in \{2, \dots, 8\}$  is applicable to  $(G, k)$ 
    then return COMPRESS( $G'', k''$ ) where  $(G'', k'')$  is the resulting instance
    obtained from  $(G, k)$  with  $S$  by applying the rule (Ri).
10:    else return  $(G, k)$ .
11:    end if
12:  end if
13: end function

```

each incomplete component of $G \setminus S$ has at most $(k+1)(k+4)|S|(|S|+2k+14)$ vertices. Therefore, each non-trivial component of $G \setminus S$ has at most $O(k|S|^4)$ vertices. Then the union of all non-trivial components of $G \setminus S$ has at most $2(k+2)|S|^2 \cdot O(k|S|^4) = O(k^2|S|^6)$ vertices. By Proposition 5.7, $G \setminus S$ has at most $2(k+3)|S|^4/3$ isolated vertices. Therefore, $|V(G)| \leq |S| + 2(k+3)|S|^4/3 + O(k^2|S|^6) = O(k^2|S|^6)$, and this proves the claim.

For an instance (G, k) obtained from Line 10, $|S|$ is at most

$$\begin{aligned}
O(\min(k^3 \log k, k^2 \log^2 |V(G)|)) &\leq O(\min(k^3 \log k, k^2 \log^2 (k^2 |S|^6))) \\
&\leq O(\min(k^3 \log k, k^2 \log^2 (k^2 (k^3 \log k)^6))) \\
&\leq O(\min(k^3 \log k, k^2 \log^2 k)) = O(k^2 \log^2 k)
\end{aligned}$$

by Lemma 4.5 and the claim. Therefore, G has at most $O(k^{14} \log^{12} k)$ vertices. \square

8 Conclusions

In this paper, we show that 3-LEAF POWER DELETION admits a fixed-parameter tractable algorithm with running time as $O(37^k |V(G)|^7 (|V(G)| + |E(G)|))$, and a kernel with $O(k^{14} \log^{12} k)$ vertices. It would be an interesting

problem to significantly reduce the size of the kernel for 3-LEAF POWER DELETION.

For an integer $\ell \geq 4$, one may investigate about ℓ -LEAF POWER DELETION, that is a problem of deciding whether deleting at most k vertices makes a graph an ℓ -leaf power. There are linear-time algorithms to recognize 4-, 5-, or 6-leaf powers [7, 9, 16].

Gurski and Wanke [18] stated that for each ℓ , ℓ -leaf powers have bounded clique-width. Rautenbach [28] presented a characterization of 4-leaf powers with no true twins as chordal graphs with ten forbidden induced subgraphs. This can be used to express, in monadic second-order logic, whether a graph is a 4-leaf power and whether there is a vertex set of size at most k whose deletion makes the graph a 4-leaf power. Therefore, by using the algorithm in [10], we deduce that 4-LEAF POWER DELETION is fixed-parameter tractable when parameterized by k , and therefore it admits a kernel. It is natural to ask whether 4-LEAF POWER DELETION admits a polynomial kernel. For $\ell \geq 5$, we do not know whether we can express ℓ -leaf powers in monadic second-order logic. If it is true for some ℓ , then not only ℓ -LEAF POWER DELETION is fixed-parameter tractable, but also ℓ -LEAF POWER RECOGNITION can be solved in polynomial time, which is still open for $\ell \geq 7$.

References

- [1] Akanksha Agrawal, Daniel Lokshtanov, Pranabendu Misra, Saket Saurabh, and Meirav Zehavi, *Polylogarithmic approximation algorithms for weighted- \mathcal{F} -deletion problems*, Approximation, randomization, and combinatorial optimization. Algorithms and techniques, LIPIcs. Leibniz Int. Proc. Inform., vol. 116, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2018, pp. Art. No. 1, 15. MR 3857239
- [2] ———, *Feedback vertex set inspired kernel for chordal vertex deletion*, ACM Trans. Algorithms **15** (2019), no. 1, Art. 11, 28. MR 3914442
- [3] Hans-Jürgen Bandelt and Henry Martyn Mulder, *Distance-hereditary graphs*, J. Combin. Theory Ser. B **41** (1986), no. 2, 182–208. MR 859310
- [4] Stéphane Bessy, Fedor V. Fomin, Serge Gaspers, Christophe Paul, Anthony Perez, Saket Saurabh, and Stéphan Thomassé, *Kernels for feedback arc set in tournaments*, J. Comput. System Sci. **77** (2011), no. 6, 1071–1078. MR 2858010

- [5] Stéphane Bessy, Christophe Paul, and Anthony Perez, *Polynomial kernels for 3-leaf power graph modification problems*, Discrete Appl. Math. **158** (2010), no. 16, 1732–1744. MR 2679772
- [6] Andreas Brandstädt and Van Bang Le, *Structure and linear time recognition of 3-leaf powers*, Inform. Process. Lett. **98** (2006), no. 4, 133–138. MR 2211095
- [7] Andreas Brandstädt, Van Bang Le, and R. Sritharan, *Structure and linear-time recognition of 4-leaf powers*, ACM Trans. Algorithms **5** (2009), no. 1, Art. 11, 22. MR 2479182
- [8] Yixin Cao and Dániel Marx, *Chordal editing is fixed-parameter tractable*, Algorithmica **75** (2016), no. 1, 118–137. MR 3492058
- [9] Maw-Shang Chang and Ming-Tat Ko, *The 3-Steiner root problem*, Graph-theoretic concepts in computer science, Lecture Notes in Comput. Sci., vol. 4769, Springer, Berlin, 2007, pp. 109–120. MR 2428569
- [10] Bruno Courcelle, Johann A. Makowsky, and Udi Rotics, *Linear time solvable optimization problems on graphs of bounded clique-width*, Theory Comput. Syst. **33** (2000), no. 2, 125–150. MR 2001e:68146
- [11] William H. Cunningham, *Decomposition of directed graphs*, SIAM J. Algebraic Discrete Methods **3** (1982), no. 2, 214–228. MR 655562
- [12] Marek Cygan, Fedor V. Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh, *Parameterized algorithms*, Springer, Cham, 2015. MR 3380745
- [13] Michael Dom, Jiong Guo, Falk Hüffner, and Rolf Niedermeier, *Error compensation in leaf power problems*, Algorithmica **44** (2006), no. 4, 363–381.
- [14] Michael Dom, Jiong Guo, Falk Hüffner, Rolf Niedermeier, and Anke Truss, *Fixed-parameter tractability results for feedback set problems in tournaments*, J. Discrete Algorithms **8** (2010), no. 1, 76–86. MR 2558881
- [15] Rodney G. Downey and Michael R. Fellows, *Fundamentals of parameterized complexity*, Texts in Computer Science, Springer, London, 2013. MR 3154461

- [16] Guillaume Ducoffe, *The 4-Steiner root problem*, Graph-theoretic concepts in computer science, Lecture Notes in Comput. Sci., vol. 11789, Springer, Berlin, 2019, pp. 14–26.
- [17] Eduard Eiben, Robert Ganian, and O-joung Kwon, *A single-exponential fixed-parameter algorithm for distance-hereditary vertex deletion*, J. Comput. System Sci. **97** (2018), 121–146. MR 3832000
- [18] Frank Gurski and Egon Wanke, *The clique-width of tree-power and leaf-power graphs (extended abstract)*, Graph-theoretic concepts in computer science, Lecture Notes in Comput. Sci., vol. 4769, Springer, Berlin, 2007, pp. 76–85. MR 2428566
- [19] Bart M. P. Jansen and Hans L. Bodlaender, *Vertex cover kernelization revisited: upper and lower bounds for a refined parameter*, Theory Comput. Syst. **53** (2013), no. 2, 263–299. MR 3063926
- [20] Bart M. P. Jansen and Marcin Pilipczuk, *Approximation and kernelization for chordal vertex deletion*, SIAM J. Discrete Math. **32** (2018), no. 3, 2258–2301. MR 3852710
- [21] Eun Jung Kim and O-joung Kwon, *A polynomial kernel for distance-hereditary vertex deletion*, Algorithms and data structures, Lecture Notes in Comput. Sci., vol. 10389, Springer, Cham, 2017, pp. 509–520. MR 3690913
- [22] ———, *Erdős-Pósa property of chordless cycles and its applications*, Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, SIAM, Philadelphia, PA, 2018, pp. 1665–1684. MR 3775897
- [23] John M. Lewis and Mihalis Yannakakis, *The node-deletion problem for hereditary properties is NP-complete*, J. Comput. System Sci. **20** (1980), no. 2, 219–230, ACM-SIGACT Symposium on the Theory of Computing (San Diego, Calif., 1978). MR 574592
- [24] Diptapriyo Majumdar, Venkatesh Raman, and Saket Saurabh, *Kernels for structural parameterizations of vertex cover—case of small degree modulators*, 10th International Symposium on Parameterized and Exact Computation, LIPIcs. Leibniz Int. Proc. Inform., vol. 43, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2015, pp. 331–342. MR 3452432

- [25] Dániel Marx, *Chordal deletion is fixed-parameter tractable*, Algorithmica **57** (2010), no. 4, 747–768. MR 2629495
- [26] Naomi Nishimura, Prabhakar Ragde, and Dimitrios M. Thilikos, *On graph powers for leaf-labeled trees*, J. Algorithms **42** (2002), no. 1, 69–108. MR 1874637
- [27] Sang-il Oum, *Rank-width and vertex-minors*, J. Combin. Theory Ser. B **95** (2005), no. 1, 79–100. MR 2156341
- [28] Dieter Rautenbach, *Some remarks about leaf roots*, Discrete Math. **306** (2006), no. 13, 1456–1461. MR 2237730