

Modules of infinite projective dimension[★]

Panyue Zhou and Xingjia Zhou

Abstract

We characterize the modules of infinite projective dimension over the endomorphism algebras of Opperman-Thomas cluster tilting objects X in $(n+2)$ -angulated categories $(\mathcal{C}, \Sigma^n, \Theta)$. We define in this article the ideal I_M of $\text{End}_{\mathcal{C}}(\Sigma^n X)$ given by all endomorphisms that factor through M , and show that the $\text{End}_{\mathcal{C}}(X)$ -module $\text{Hom}_{\mathcal{C}}(X, M)$ has infinite projective dimension precisely when I_M is non-zero. As an application, we generalize a recent result by Beaudet-Brüstle-Todorov for cluster-tilted algebras.

Key words: $(n+2)$ -angulated categories; cluster tilting objects; projective dimension.

2010 Mathematics Subject Classification: 18E30; 16D90.

1 Introduction

Cluster-tilting theory sprouted from the categorification of Fomin-Zelevinsky's cluster algebras. It is used to construct abelian categories from some triangulated categories. By Buan-Marsh-Reiten [BMR, Theorem 2.2] in cluster categories, by Keller-Reiten [KR, Proposition 2.1] in the 2-Calabi-Yau case, then by Koenig-Zhu [KZ, Theorem 3.3] and Iyama-Yoshino [IY, Corollary 6.5] in the general case, one can pass from triangulated categories to abelian categories by factoring out any cluster tilting subcategory.

Recall that the notion of cluster-tilting subcategory which is due to Iyama [I]. Let \mathcal{C} be a triangulated category with shift functor Σ . A subcategory \mathcal{X} of \mathcal{C} is called a *cluster-tilting subcategory* of \mathcal{C} , if it satisfies the following conditions:

- (1) \mathcal{X} is contravariantly finite and covariantly finite in \mathcal{C} .
- (2) $X \in \mathcal{X}$ if and only if $\text{Hom}_{\mathcal{C}}(X, \Sigma \mathcal{X}) = 0$, i.e. $\text{Hom}_{\mathcal{C}}(X, \Sigma M) = 0$, for any $M \in \mathcal{X}$.
- (3) $X \in \mathcal{X}$ if and only if $\text{Hom}_{\mathcal{C}}(\mathcal{X}, \Sigma X) = 0$, i.e. $\text{Hom}_{\mathcal{C}}(M, \Sigma X) = 0$, for any $M \in \mathcal{X}$.

An object X is called *cluster-tilting*, if $\text{add} X$ is a cluster-tilting, where $\text{add} X$ is the subcategory of \mathcal{C} consisting of direct summands of direct sum of finitely many copies of X .

In fact, Koenig and Zhu [KZ, Lemma 3.2] show that \mathcal{X} is cluster-tilting if and only if $\text{Hom}_{\mathcal{C}}(\mathcal{X}, \Sigma \mathcal{X}) = 0$, i.e. $\text{Hom}_{\mathcal{C}}(X, \Sigma Y) = 0$, for any $X, Y \in \mathcal{X}$, and for any object $C \in \mathcal{C}$, there exists a triangle $X_0 \rightarrow X_1 \rightarrow C \rightarrow \Sigma X_0$ where $X_0, X_1 \in \mathcal{X}$.

The notion of cluster tilting objects can be generalised to $(n+2)$ -angulated categories, which is due to Oppermann and Thomas [OT, Definition 5.3].

[★]Panyue Zhou is supported by the National Natural Science Foundation of China (Grants No. 11901190 and 11671221) and the Hunan Provincial Natural Science Foundation of China (Grants No. 2018JJ3205).

Definition 1.1. [OT, Definition 5.3] Let \mathcal{C} be an $(n + 2)$ -angulated category with an n -suspension functor Σ^n . An object $X \in \mathcal{C}$ is called *Opperman-Thomas cluster-tilting* if

- (1) $\text{Hom}_{\mathcal{C}}(X, \Sigma^n X) = 0$.
- (2) For any object $C \in \mathcal{C}$, there exists an $(n + 2)$ -angle

$$X_0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_n \rightarrow C \rightarrow \Sigma^n X_0$$

where $X_0, X_1, \dots, X_n \in \text{add} X$.

Let \mathcal{C} be a triangulated category with a shift functor Σ , and $X \in \mathcal{C}$ be a cluster tilting object. For any object $M \in \mathcal{C}$, we denote by I_M the ideal of $\text{End}_{\mathcal{C}}(\Sigma X)$ given by all endomorphisms that factor through M and call it *factorization ideal* of M . Beaudet-Brüstle-Todorov showed the following.

Theorem 1.2. [BBT, Theorem 1.1] *Let \mathcal{C} be a triangulated category with a shift functor Σ and $X \in \mathcal{C}$ be a cluster tilting object. Let M be an indecomposable object in \mathcal{C} which does not belong to $\text{add} \Sigma X$. Then the $\text{End}_{\mathcal{C}}(X)$ -module $\text{Hom}_{\mathcal{C}}(X, M)$ is of infinite projective dimension if and only if the factorization ideal I_M is non-zero.*

Recently, Geiss, Keller and Oppermann introduced in [GKO] a new type of categories, called $(n+2)$ -angulated categories, which generalize triangulated categories: the classical triangulated categories are the special case $n = 1$. These categories appear for instance when considering certain n -cluster tilting subcategories of triangulated categories.

Let \mathcal{C} be an $(n + 2)$ -angulated category with an n -suspension functor Σ^n , and $X \in \mathcal{C}$ be an Opperman-Thomas cluster tilting object. For any object $M \in \mathcal{C}$, we denote by I_M the ideal of $\text{End}_{\mathcal{C}}(\Sigma^n X)$ given by all endomorphisms that factor through M and call it *factorization ideal* of M . Our main result is the following, which is a generalization of Beaudet-Brüstle-Todorov result.

Theorem 1.3. (See Theorem 3.1 for details) *Let \mathcal{C} be an $(n + 2)$ -angulated category with an n -suspension functor Σ^n and $X \in \mathcal{C}$ be an Opperman-Thomas cluster tilting object. Let M be an indecomposable object in \mathcal{C} which does not belong to $\text{add} \Sigma^n X$. Then the $\text{End}_{\mathcal{C}}(X)$ -module $\text{Hom}_{\mathcal{C}}(X, M)$ is of infinite projective dimension if and only if the factorization ideal I_M is non-zero.*

This article is organised as follows: In Section 2, we review some elementary definitions and properties on $(n + 2)$ -angulated categories. In Section 3, we prove our main result.

2 Preliminaries

In this section, we briefly recall the definition and basic properties of $(n + 2)$ -angulated categories from [GKO]. Let \mathcal{C} be an additive category with an automorphism $\Sigma^n : \mathcal{C} \rightarrow \mathcal{C}$, and an integer n greater than or equal to one.

An $(n + 2)$ - Σ^n -sequence in \mathcal{C} is a sequence of objects and morphisms

$$A_0 \xrightarrow{f_0} A_1 \xrightarrow{f_1} A_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_n \xrightarrow{f_n} A_{n+1} \xrightarrow{f_{n+1}} \Sigma^n A_0.$$

Its *left rotation* is the $(n + 2)$ - Σ^n -sequence

$$A_1 \xrightarrow{f_1} A_2 \xrightarrow{f_2} A_3 \xrightarrow{f_3} \cdots \xrightarrow{f_n} A_{n+1} \xrightarrow{f_{n+1}} \Sigma^n A_0 \xrightarrow{(-1)^n \Sigma^n f_0} \Sigma^n A_1.$$

A *morphism* of $(n + 2)$ - Σ^n -sequences is a sequence of morphisms $\varphi = (\varphi_0, \varphi_1, \dots, \varphi_{n+1})$ such that the following diagram commutes

$$\begin{array}{ccccccccccc} A_0 & \xrightarrow{f_0} & A_1 & \xrightarrow{f_1} & A_2 & \xrightarrow{f_2} & \cdots & \xrightarrow{f_n} & A_{n+1} & \xrightarrow{f_{n+1}} & \Sigma^n A_0 \\ \downarrow \varphi_0 & & \downarrow \varphi_1 & & \downarrow \varphi_2 & & & & \downarrow \varphi_{n+1} & & \downarrow \Sigma^n \varphi_0 \\ B_0 & \xrightarrow{g_0} & B_1 & \xrightarrow{g_1} & B_2 & \xrightarrow{g_2} & \cdots & \xrightarrow{g_n} & B_{n+1} & \xrightarrow{g_{n+1}} & \Sigma^n B_0 \end{array}$$

where each row is an $(n + 2)$ - Σ^n -sequence. It is an *isomorphism* if $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_{n+1}$ are all isomorphisms in \mathcal{C} .

Definition 2.1. [GKO, Definition 2.1] An *n-angulated category* is a triple $(\mathcal{C}, \Sigma^n, \Theta)$, where \mathcal{C} is an additive category, Σ^n is an automorphism of \mathcal{C} (Σ^n is called the *n-suspension functor*), and Θ is a class of $(n + 2)$ - Σ^n -sequences (whose elements are called $(n + 2)$ -angles), which satisfies the following axioms:

- (N1) (a) The class Θ is closed under isomorphisms, direct sums and direct summands.
 (b) For each object $A \in \mathcal{C}$ the trivial sequence

$$A \xrightarrow{1_A} A \rightarrow 0 \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \Sigma^n A$$

belongs to Θ .

- (c) Each morphism $f_0: A_0 \rightarrow A_1$ in \mathcal{C} can be extended to $(n + 2)$ - Σ^n -sequence:

$$A_0 \xrightarrow{f_0} A_1 \xrightarrow{f_1} A_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_n \xrightarrow{f_n} A_{n+1} \xrightarrow{f_{n+1}} \Sigma^n A_0.$$

- (N2) An $(n + 2)$ - Σ^n -sequence belongs to Θ if and only if its left rotation belongs to Θ .

- (N3) Each solid commutative diagram

$$\begin{array}{ccccccccccc} A_0 & \xrightarrow{f_0} & A_1 & \xrightarrow{f_1} & A_2 & \xrightarrow{f_2} & \cdots & \xrightarrow{f_n} & A_{n+1} & \xrightarrow{f_{n+1}} & \Sigma^n A_0 \\ \downarrow \varphi_0 & & \downarrow \varphi_1 & & \downarrow \varphi_2 & & & & \downarrow \varphi_{n+1} & & \downarrow \Sigma^n \varphi_0 \\ B_0 & \xrightarrow{g_0} & B_1 & \xrightarrow{g_1} & B_2 & \xrightarrow{g_2} & \cdots & \xrightarrow{g_n} & B_{n+1} & \xrightarrow{g_{n+1}} & \Sigma^n B_0 \end{array}$$

with rows in Θ , the dotted morphisms exist and give a morphism of $(n + 2)$ - Σ^n -sequences.

- (N4) In the situation of (N3), the morphisms $\varphi_2, \varphi_3, \dots, \varphi_{n+1}$ can be chosen such that the

mapping cone

$$A_1 \oplus B_0 \xrightarrow{\begin{pmatrix} -f_1 & 0 \\ \varphi_1 & g_0 \end{pmatrix}} A_2 \oplus B_1 \xrightarrow{\begin{pmatrix} -f_2 & 0 \\ \varphi_2 & g_1 \end{pmatrix}} \cdots \xrightarrow{\begin{pmatrix} -f_{n+1} & 0 \\ \varphi_{n+1} & g_n \end{pmatrix}} \Sigma^n A_0 \oplus B_{n+1} \xrightarrow{\begin{pmatrix} -\Sigma^n f_0 & 0 \\ \Sigma^n \varphi_1 & g_{n+1} \end{pmatrix}} \Sigma^n A_1 \oplus \Sigma^n B_0$$

belongs to Θ .

Now we give an example of $(n+2)$ -angulated categories.

Example 2.2. We recall the standard construction of $(n+2)$ -angulated categories given by Geiss-Keller-Oppermann [GKO, Theorem 1]. Let \mathcal{C} be a triangulated category and \mathcal{T} an n -cluster tilting subcategory which is closed under Σ^n , where Σ is the shift functor of \mathcal{C} . Then $(\mathcal{T}, \Sigma^n, \Theta)$ is an $(n+2)$ -angulated category, where Θ is the class of all sequences

$$A_0 \xrightarrow{f_0} A_1 \xrightarrow{f_1} A_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_n \xrightarrow{f_n} A_{n+1} \xrightarrow{f_{n+1}} \Sigma^n A_0$$

such that there exists a diagram

$$\begin{array}{ccccccc} & & A_1 & \xrightarrow{f_1} & A_2 & & \cdots & & A_n & & \\ & f_0 \nearrow & & & & \nwarrow f_1 & & & & \nwarrow f_n & \\ & & A_0 & \longleftarrow & A_{1.5} & \longleftarrow & A_{2.5} & \cdots & A_{n-1.5} & \longleftarrow & A_{n+1} \end{array}$$

with $A_i \in \mathcal{T}$ for all $i \in \mathbb{Z}$, such that all oriented triangles are triangles in \mathcal{C} , all non-oriented triangles commute, and f_{n+1} is the composition along the lower edge of the diagram.

From now on to the end of this section, we assume that \mathcal{C} is an $(n+2)$ -angulated category with an n -suspension functor Σ^n , $X \in \mathcal{C}$ is an Opperman-Thomas cluster tilting object and $\Gamma := \text{End}_{\mathcal{C}}(X)$ is the endomorphism algebra of X . Recall the following result.

Theorem 2.3. [JJ1, Theorem 0.5] and [ZZ, Theorem 3.8] *Consider the essential image \mathcal{D} of the functor $\mathcal{C}(X, -) : \mathcal{C} \rightarrow \text{mod } \Gamma$. Then \mathcal{D} is an n -cluster tilting subcategory of $\text{mod } \Gamma$ where $\text{mod } \Gamma$ is the category of finite dimensional right Γ -modules. There exists a commutative diagram, as shown below, where the vertical arrow is the quotient functor and the diagonal arrow is an equivalence of categories:*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\mathcal{C}(X, -)} & \mathcal{D} \\ \downarrow & \searrow \cong & \\ \mathcal{C}/\text{add } \Sigma^n X & & \end{array}$$

Remark 2.4. [JJ1, Lemma 2.1] The classic add-proj-correspondence holds, as the functor $\mathcal{C}(X, -)$ restricts to an equivalence $\text{add } X \rightarrow \text{proj } \Gamma$ where $\text{proj } \Gamma$ is the category of projective finite dimensional right Γ -modules.

Lemma 2.5. [JJ2, Lemma 2.2] *If $M \in \mathcal{C}$ has no direct summands in $\text{add}\Sigma^n X$, then there exists an $(n + 2)$ -angle*

$$X_0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_n \rightarrow M \rightarrow \Sigma^n X_0$$

in \mathcal{C} with the following properties: Each X_i is in $\text{add}X$, and applying the functor $\mathcal{C}(X, -)$ gives a complex

$$\mathcal{C}(X, X_0) \rightarrow \mathcal{C}(X, X_1) \rightarrow \cdots \rightarrow \mathcal{C}(X, X_n) \rightarrow \mathcal{C}(X, M) \rightarrow 0$$

which is the start of the augmented minimal projective resolution of $\mathcal{C}(X, M)$.

3 Proof of the Main Theorem

Now we begin to prove the main result of this article.

Theorem 3.1. *Let \mathcal{C} be an $(n+2)$ -angulated category with an Opperman-Thomas cluster-tilting object X , and Γ be the endomorphism algebra of X . Let M be an indecomposable object in \mathcal{C} which does not belong to $\text{add}\Sigma^n X$. Then the Γ -module $\text{Hom}_{\mathcal{C}}(X, M)$ is of infinite projective dimension if and only if the factorization ideal I_M is non-zero.*

Proof. Since M is an indecomposable object in \mathcal{C} which does not belong to $\text{add}\Sigma^n X$, then there exists an $(n + 2)$ -angle

$$X_0 \xrightarrow{\alpha_0} X_1 \xrightarrow{\alpha_1} X_2 \rightarrow \cdots \rightarrow X_n \xrightarrow{\theta} M \xrightarrow{\beta} \Sigma^n X_0 \quad (3.1)$$

which satisfies the properties of Lemma 2.4.

First, we assume Γ -module $\text{Hom}_{\mathcal{C}}(X, M)$ has infinite projective dimension. Applying the functor $\text{Hom}_{\mathcal{C}}(X, -)$ to the $(n + 2)$ -angle (3.1), we have the following exact sequence in $\text{mod}\Gamma$:

$$(X, \Sigma^{-n} M) \xrightarrow{(X, (-1)^n \Sigma^{-n} \beta)} (X, X_0) \rightarrow (X, X_1) \rightarrow \cdots \rightarrow (X, X_n) \rightarrow (X, M) \rightarrow 0$$

where we omitted $\text{Hom}_{\mathcal{C}}$ because of lack of space. It follows that the morphism

$$\text{Hom}_{\mathcal{C}}(X, (-1)^n \Sigma^{-n} \beta) \neq 0,$$

since otherwise the projective dimension of $\text{Hom}_{\mathcal{C}}(X, M)$ would be at most n . Choose a morphism $(-1)^n \Sigma^{-n} \gamma$ in $\text{Hom}_{\mathcal{C}}(X, \Sigma^{-n} M)$ whose image under $\text{Hom}_{\mathcal{C}}(X, (-1)^n \Sigma^{-n} \beta)$ is non-zero, that is, the composition

$$X \xrightarrow{(-1)^n \Sigma^{-n} \gamma} \Sigma^{-n} M \xrightarrow{(-1)^n \Sigma^{-n} \beta} X$$

is non-zero. This yields the non-zero composition

$$\Sigma^n X \xrightarrow{\gamma} M \xrightarrow{\beta} \Sigma^n X.$$

Hence there exists a non-zero element $\beta\gamma$ in the factorization ideal I_M .

Conversely, if the factorization ideal I_M is non-zero, we prove that the Γ -module $\text{Hom}_{\mathcal{C}}(X, M)$ is not of projective dimension at most n . Otherwise, Applying the functor $\text{Hom}_{\mathcal{C}}(X, -)$ to the $(n+2)$ -angle (3.1), we have the following exact sequence in $\text{mod}\Gamma$:

$$0 \rightarrow \text{Hom}_{\mathcal{C}}(X, X_0) \xrightarrow{\text{Hom}_{\mathcal{C}}(X, \alpha_0)} \text{Hom}_{\mathcal{C}}(X, X_1) \rightarrow \cdots \rightarrow \text{Hom}_{\mathcal{C}}(X, X_n) \rightarrow \text{Hom}_{\mathcal{C}}(X, M) \rightarrow 0.$$

Let $ba: \Sigma^n X \xrightarrow{a} M \xrightarrow{b} \Sigma^n X$ be any element in I_M . Since $\text{Hom}_{\mathcal{C}}(X, \Sigma^n X) = 0$, we obtain that the morphism $b\theta \in \text{Hom}_{\mathcal{C}}(X_n, \Sigma^n X)$ is zero. Hence we have the following commutative diagram:

$$\begin{array}{ccc} 0 & \longrightarrow & \Sigma^n X \\ \downarrow & & \downarrow a \\ X_n & \xrightarrow{\theta} & M \\ \downarrow & & \downarrow b \\ 0 & \longrightarrow & \Sigma^n X \end{array}$$

By (N3), there exist morphisms $c: X \rightarrow X_0$ and $d: X_0 \rightarrow X$ which make the following diagram commutative:

$$\begin{array}{ccccccccccccccc} X & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \Sigma^n X & \xrightarrow{(-1)^n} & \Sigma^n X \\ \downarrow c & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow a & & \downarrow \Sigma^n c \\ X_0 & \xrightarrow{\alpha_0} & X_1 & \xrightarrow{\alpha_1} & X_2 & \longrightarrow & \cdots & \longrightarrow & X_{n-1} & \longrightarrow & X_n & \xrightarrow{\theta} & M & \xrightarrow{\beta} & \Sigma^n X_0 \\ \downarrow d & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow b & & \downarrow \Sigma^n d \\ X & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \Sigma^n X & \xrightarrow{(-1)^n} & \Sigma^n X \end{array}$$

Thus $\alpha_0 c = 0$ and then $c = 0$ since $\text{Hom}_{\mathcal{C}}(X, \alpha_0)$ is a monomorphism. This implies $\Sigma^n c = 0$ and, therefore $ba = \Sigma^n d \circ \Sigma^n c = 0$. It follows that $I_M = 0$, which is a contradiction to our assumption. This finishes the proof. \square

As a special case when $n = 1$, we have the following.

Corollary 3.2. [BBT, Theorem 1.1] *Let \mathcal{C} be a triangulated category with a cluster-tilting object X , and Γ be the endomorphism algebra of X . Let M be an indecomposable object in \mathcal{C} which does not belong to $\text{add}\Sigma X$. Then the Γ -module $\text{Hom}_{\mathcal{C}}(X, M)$ is of infinite projective dimension if and only if the factorization ideal I_M is non-zero.*

References

- [BBT] L. Beaudet, T. Brüstle, G. Todorov. Projective dimension of modules over cluster-tilted algebras. *Algebr. Represent. Theory* 17(6): 1797-1807, 2014.
- [BMR] A. B. Buan, R. Marsh, I. Reiten. Cluster-tilted algebra. *Trans. Amer. Math. Soc.* 359(1): 323-332, 2007.
- [GKO] C. Geiss, B. Keller and S. Oppermann. n -angulated categories. *J. Reine Angew. Math.* 675: 101-120, 2013.

- [I] O. Iyama. Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories. *Adv. Math.* 210(1): 22-50, 2007.
- [IY] O. Iyama and Y. Yoshino. Mutation in triangulated categories and rigid Cohen-Macaulay modules. *Invent. Math.* 172(1): 117-168, 2008.
- [JJ1] K. Jacobsen, P. Jørgensen. d -abelian quotients of $(d+2)$ -angulated categories. *J. Algebra*, 521: 114-136, 2019.
- [JJ2] K. Jacobsen, P. Jørgensen. Maximal τ_d -rigid pairs. arXiv: 1812.04871, 2018.
- [KR] B. Keller, I. Reiten. Cluster-tilted algebras are Gorenstein and stably Calabi-Yau. *Adv. Math.* 211: 123-151, 2007.
- [KZ] S. Koenig, B. Zhu. From triangulated categories to abelian categories: cluster tilting in a general framework. *Math. Z.* 258: 143-160, 2008.
- [OT] S. Oppermann, H. Thomas. Higher-dimensional cluster combinatorics and representation theory. *J. Eur. Math. Soc.* 14(6): 1679-1737, 2012.
- [ZZ] P. Zhou, B. Zhu. n -Abelian quotient categories. *J. Algebra*, 527: 264-279, 2019.

Panyue Zhou and Xingjia Zhou

College of Mathematics, Hunan Institute of Science and Technology, Yueyang, Hunan, 414006, People's Republic of China.

E-mail: panyuezhou@163.com and 1149077341@qq.com