

# On the Existence of Heterotic-String and Type-II-Superstring Field Theory Vertices

Seyed Faroogh Moosavian<sup>a,b</sup> Yehao Zhou<sup>a,b</sup>

<sup>a</sup>*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada N2L 2Y5*

<sup>b</sup>*University of Waterloo, Waterloo, ON, Canada*

*E-mail:* [sfmoosavian@pitp.ca](mailto:sfmoosavian@pitp.ca), [yzhou3@pitp.ca](mailto:yzhou3@pitp.ca)

**ABSTRACT:** We consider two problems associated to the space of Riemann surfaces that are closely related to string theory. We first consider the problem of existence of heterotic-string and type-II-superstring field theory vertices in the product of spaces of bordered surfaces parameterizing left- and right-moving sectors of these theories. It turns out that this problem can be solved by proving the existence of a solution to the BV quantum master equation in moduli spaces of bordered spin-Riemann surfaces. We first prove that for arbitrary genus  $g$ ,  $n_{\text{NS}}$  Neveu-Schwarz boundary components, and  $n_{\text{R}}$  Ramond boundary components such solutions exist. We also prove that these solutions are unique up to homotopies in the category of BV algebras. Furthermore, we prove that there exists a map in this category under which these solutions are mapped to fundamental classes of Deligne-Mumford stacks of associated punctured spin curves. These results generalize the work of Costello on the existence of a solution to the BV quantum master equations in moduli spaces of bordered ordinary Riemann surfaces which, through the work of Sen and Zwiebach, are related to the existence of bosonic-string vertices, and their relation to fundamental classes of Deligne-Mumford stacks of associated punctured ordinary curves. Using the existence of solutions to the BV quantum master equation in moduli spaces of spin curves, we prove that heterotic-string and type-II-superstring field theory vertices, for arbitrary genus  $g$  and arbitrary number of any type of boundary components, exist. Furthermore, we prove the existence of a solution to the BV quantum master equation in spaces of bordered  $\mathcal{N} = 1$  super-Riemann surfaces for arbitrary genus  $g$ ,  $n_{\text{NS}}$  Neveu-Schwarz boundary components, and  $n_{\text{R}}$  Ramond boundary components. Secondly, we turn to the moduli problem of stable  $\mathcal{N} = 1$  SUSY curves. Using Vaintrob's deformation theory of complex superspaces and Deligne's model for the compactification of the space of  $\mathcal{N} = 1$  SUSY curves, we prove the representability of this moduli problem by a proper smooth Deligne-Mumford superstack. This is a generalization of the work of Deligne and Mumford on the representability of the moduli problem of stable ordinary curves by a Deligne-Mumford stack. Along the way, we also elaborate some possibly-familiar aspects of the formulation of superstring perturbation theory using spin curves. In particular, we explain a subtlety in the gluing of Ramond punctures on spin curves.

---

## Contents

|          |                                                                                                                                            |           |
|----------|--------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                                                                                        | <b>2</b>  |
| 1.1      | Statement of the Problems                                                                                                                  | 3         |
| 1.1.1    | The Existence of String Vertices                                                                                                           | 5         |
| 1.1.2    | The Moduli Problem of Stable $\mathcal{N} = 1$ SUSY Curves                                                                                 | 6         |
| 1.2      | Contributions of the Paper                                                                                                                 | 6         |
| 1.3      | Notations and Remarks                                                                                                                      | 11        |
| <b>2</b> | <b>Spin Curves, Punctures and their Gluing</b>                                                                                             | <b>15</b> |
| 2.1      | Spin Curves over $\mathbb{C}$                                                                                                              | 15        |
| 2.2      | Punctures on Spin and Super-Riemann Surfaces                                                                                               | 18        |
| 2.3      | Gluing of Punctures on Spin and Super-Riemann Surfaces                                                                                     | 19        |
| <b>3</b> | <b>Nodal Spin Curves and Torsion-Free Sheaves</b>                                                                                          | <b>22</b> |
| 3.1      | Degenerations of Spin Curves                                                                                                               | 22        |
| 3.1.1    | Neveu-Schwarz Degenerations                                                                                                                | 23        |
| 3.1.2    | Ramond Degenerations                                                                                                                       | 24        |
| 3.2      | PCOs and Gluing of Ramond Punctures on Spin Curves                                                                                         | 25        |
| <b>4</b> | <b>Existence and Uniqueness of Solution to the BV QME in <math>\mathcal{S}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})</math></b>   | <b>27</b> |
| 4.1      | The Strategy of Proof                                                                                                                      | 28        |
| 4.2      | Gluing of Boundary Components                                                                                                              | 34        |
| 4.3      | Connected Components of $\partial\overline{\mathcal{S}}_{\mathfrak{g}, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$                 | 35        |
| 4.4      | An Alternative Model for $\mathcal{S}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$                                                  | 41        |
| 4.5      | The Proof of Existence and Uniqueness                                                                                                      | 42        |
| 4.6      | Relation of Solutions to Fundamental Classes of $\overline{\mathcal{S}}_{\mathfrak{g}, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ | 48        |
| <b>5</b> | <b>Existence of Solutions to the BV QME in <math>\mathcal{SM}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})</math></b>                | <b>49</b> |
| 5.1      | Obstructions of Splitness and Projectedness                                                                                                | 50        |
| 5.1.1    | Lifting Space of the Bosonic Truncation of a Supermanifold                                                                                 | 53        |
| 5.2      | Singular Homology on Superspaces                                                                                                           | 56        |
| 5.3      | Proof of the Existence of Solution                                                                                                         | 58        |
| <b>6</b> | <b>Existence of Heterotic-String and Type-II-Superstring Vertices</b>                                                                      | <b>59</b> |
| 6.1      | Introduction                                                                                                                               | 60        |
| 6.2      | The Case of Heterotic-String Field Theory Vertices                                                                                         | 62        |
| 6.3      | The Case of Type-II-Superstring Field Theory Vertices                                                                                      | 66        |

|          |                                                                                               |            |
|----------|-----------------------------------------------------------------------------------------------|------------|
| <b>7</b> | <b>On the Moduli Problem of Stable <math>\mathcal{N} = 1</math> SUSY Curves</b>               | <b>69</b>  |
| 7.1      | Definitions, Elementary Results and Super Analogs of Classical Theorems                       | 70         |
| 7.1.1    | Superschemes over Inductive Limit of Superrings                                               | 77         |
| 7.1.2    | Affineness Criteria                                                                           | 79         |
| 7.1.3    | Chow's Lemma for Superschemes                                                                 | 80         |
| 7.1.4    | Pushforward and Flatness                                                                      | 81         |
| 7.1.5    | Super Analog of Grothendieck's Comparison and Existence Theorems                              | 82         |
| 7.1.6    | Super Analog of Castelnuovo-Mumford Regularity                                                | 84         |
| 7.1.7    | Flattening Stratification                                                                     | 85         |
| 7.1.8    | The Quot Superscheme                                                                          | 87         |
| 7.1.9    | Deformation Theory and Super Analog of Schlessinger's Theorem                                 | 89         |
| 7.2      | Stable $\mathcal{N} = 1$ SUSY Curves                                                          | 91         |
| 7.2.1    | Étale Descent for $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$                  | 94         |
| 7.2.2    | Local Structures of Stable SUSY Curves                                                        | 96         |
| 7.2.3    | Infinitesimal Automorphisms of Stable SUSY Curves                                             | 103        |
| 7.2.4    | Isomorphisms of Stable SUSY Curves                                                            | 105        |
| 7.2.5    | Étale Local Structures of Stable SUSY Curves                                                  | 105        |
| 7.2.6    | Absence of Obstruction for Lifting Stable SUSY Curves                                         | 107        |
| 7.3      | Moduli of Stable $\mathcal{N} = 1$ SUSY Curves                                                | 110        |
| 7.3.1    | Gluing of Punctures                                                                           | 117        |
| <b>8</b> | <b>Discussion, and Future Directions</b>                                                      | <b>119</b> |
| <b>A</b> | <b>The Picture-Changing Operation</b>                                                         | <b>120</b> |
| A.1      | The Canonical Framework: Friedan-Shenker-Martinec Approach                                    | 120        |
| A.2      | The Path Integral Framework: Verlinde-Verlinde Approach                                       | 123        |
| A.3      | The Supergeometry Framework: Belopolsky Approach                                              | 126        |
| A.4      | The Unified Perspective: From Supermanifolds to Superstrings                                  | 129        |
| <b>B</b> | <b>Moduli Stacks of Stable Spin Curves</b>                                                    | <b>131</b> |
| B.1      | Definitions and Examples                                                                      | 131        |
| B.2      | Rank-One Torsion-Free Sheaves                                                                 | 136        |
| B.3      | Cornalba's Definition of Spin Curves                                                          | 138        |
| B.4      | Stack of Prespin Structures and Representability of $\overline{\mathcal{S}}_{g,n}^{\text{m}}$ | 139        |
| B.5      | Diagonal of $\overline{\mathcal{S}}_{g,n}^{\text{m}}$                                         | 141        |
| B.6      | Deformation Theory of $\overline{\mathcal{S}}_{g,n}^{\text{m}}$                               | 142        |
| B.7      | Properness of $\overline{\mathcal{S}}_{g,n}^{\text{m}}$                                       | 144        |
| B.8      | Connected Components of $\overline{\mathcal{S}}_{g,n}^{\text{m}}$                             | 144        |
| B.9      | Gluing Punctures                                                                              | 147        |

---

# 1 Introduction

The best-understood formulation of superstring theory is the superstring perturbation theory. Much of the mathematical difficulty of the superstring perturbation theory in the current formulation comes from the intricate mathematics of spin-Riemann or super-Riemann surfaces and stacks<sup>1</sup> or superstacks thereof. The appearance of these spaces is resulted from two different approaches to superstring perturbation theory in the RNS formalism,

1. The natural setting for the formulation of the RNS superstring theory is the language of supergeometry. The worldsheet theory is written on an  $\mathcal{N} = 1$  super-Riemann surface and one needs to deal with superstacks of these surfaces to compute the scattering amplitudes [10–12]. The concept of super-Riemann surface, as an appropriate generalization of ordinary Riemann surfaces for doing superstring perturbation theory, was introduced by Baranov and Schwarz [13] and independently by Friedan [14].
2. In the second formulation, the so-called *picture-changing formalism*, superstring amplitudes are defined as integrals over the moduli stack of spin-Riemann surfaces where the string measure contains the so-called picture-changing operators (PCOs) [15]. Since the worldsheet theory contains fermions, the Riemann surfaces are equipped with a choice of spin structure, i.e. a line bundle  $\mathcal{E}$  over the surface such that  $\mathcal{E}^{\otimes 2} \equiv \mathcal{E} \otimes \mathcal{E} \simeq \omega$ , where  $\omega$  is the canonical line bundle over the surface<sup>2</sup>. In physical terms, a choice of spin structure corresponds to the phase shift of the fermions after parallel transport along a closed 1-cycle of the surface. There are two possible boundary conditions by going around a closed 1-cycle: periodic or antiperiodic boundary conditions<sup>3</sup>. The fermions with periodic boundary condition on the plane (or anti-periodic boundary condition on the cylinder) form the so-called Neveu-Schwarz (NS) sector of the Hilbert space, and the fermions with antiperiodic boundary condition on the plane (or periodic boundary condition on the cylinder) form the so-called Ramond (R) sector of the Hilbert space. The antiperiodic boundary condition on R states is one of the main sources of difficulties in superstring theory.

---

<sup>1</sup> To have a uniform and mathematically-precise language throughout the paper, we use the term stack or superstack for spaces of (possibly *punctured*) ordinary, spin- or  $\mathcal{N} = 1$  SUSY curves. Stack and superstack in this context are synonyms for orbifold and superorbifold respectively, which are what these spaces are due to the existence of singularities [1–5]. These terms can be replaced with the possibly-more-familiar terms moduli or supermoduli space. A concise and accessible discussion of the stacky nature of these spaces can be found in section 3.2.1 of [6]. We also refer to sections 2 and 3 of [7], and also [8], for relevant definitions of superstacks and coarse superspaces. A thorough treatment of stacks, especially stacks of curves, Deligne-Mumford stacks, and coarse moduli spaces can be found in [9]. As far as we aware, it is not known that what the spaces of *bordered* ordinary, spin- or super-Riemann surfaces are. As we prove in this paper, the moduli of *punctured*  $\mathcal{N} = 1$  SUSY curves is a Deligne-Mumford superstack.

<sup>2</sup>The canonical line bundle  $\omega$  is isomorphic to the holomorphic cotangent bundle over the surface.

<sup>3</sup>The boundary conditions we are referring to are in the radial quantization, i.e. after mapping the cylinder, with coordinates  $(\sigma, \tau)$ , to the complex  $z$ -plane via  $z = \exp(\tau + i\sigma)$ .

As a result there are two types of punctures in superstring theory, depending on whether the corresponding vertex operator belongs to the NS or R sector. We will call the corresponding punctures NS or R punctures. It has been argued in [16] and has been further elaborated in [17, 18] that the two formulations are equivalent. In this paper, we use both formalism and it will be hopefully clear which framework we are working within.

### 1.1 Statement of the Problems

One of the surprising feature of the superstring theory is that the modular invariance removes the UV divergences of the theory. However, the theory suffers from IR divergences [12, 19]. The IR divergences come from the points at infinity<sup>4</sup> of the superstack or equivalently the moduli stack of spin curves. In the picture-changing formalism, on one hand, one has to deal with moduli stack of punctured spin curves and on the other hand, as was mentioned, the IR divergences of the theory come from the points at infinity of the compactified moduli stack. One thus needs to deal with the compactification of the moduli stack of punctured spin curves. However, unlike the case of bosonic-string theory, where the amplitudes are obtained by an integration over the moduli stack of ordinary Riemann surface, there is no natural moduli stack to be integrated over in the superstring theories, in either the supergeometry or the picture-changing formulation. The reason is as follows; 1) for heterotic-string theory the fields of right-moving sector, having  $\mathcal{N} = 1$  supersymmetry, are parameterized by  $(z|\theta)$  while the fields of left-moving sector, having no supersymmetry, are parametrized by  $\tilde{z}$ . Therefore, there is no notion of complex conjugate variable for  $\theta$  and therefore we can not set  $\tilde{z} = \bar{z}$  since the transition functions may depend on  $\theta$  on different patches; 2) For type-II theories, the fields of right-moving sector, having  $\mathcal{N} = 1$  supersymmetry, are parameterized by  $(z|\theta)$  while the fields of left-moving sector, again having  $\mathcal{N} = 1$ , are parametrized by  $(\tilde{z}|\tilde{\theta})$ . Again, there is no natural notion of complex conjugation between  $(z|\theta)$  and  $(\tilde{z}|\tilde{\theta})$  since  $\theta$  and  $\tilde{\theta}$  can have different spin structures. Therefore, in the supergeometry formulation, one needs to integrate over an appropriate integration cycle with appropriate behaviors at infinity inside an appropriate product of left and right stack and/or superstacks, where the appropriate moduli or superstacks are determined by the worldsheet of the theory [10, 11]. In the picture-changing formalism, the construction of a proper integration cycle involves the delicate procedure of vertical integration [17, 18, 21, 22]. The integrands associated to the amplitudes are invariant under the *spin- $\mathcal{E}$  mapping-class group*, the subgroup of the mapping-class group of the surface that preserves the spin structure  $\mathcal{E}$ , and thus naturally can be integrated over the moduli stack of spin curves with appropriate signature. One also need to sum over all spin structures for at

---

<sup>4</sup>The compactification of the moduli stack of punctured Riemann surfaces without boundary corresponds to the addition of complex co-dimension one divisor to the uncompactified moduli stack. The resulting space, the Deligne-Mumford space [20], is a compact space without boundary. The divisor points thus should be thought of as points at infinity. There is a subtle difference between this case and the case of open-closed superstring theories, where the superstack has boundaries, i.e. the compactification locus consists of real codimension one divisors. The presence of these boundaries is related to the presence of various anomalies. For more on this see section 9.2 of [12].

least two crucial reasons: 1) we need to integrate over  $\overline{\mathcal{M}}_{g,n}$ , the compactified moduli stack of genus- $g$  surfaces with  $n$  punctures, and as such, we thus need a mapping-class-group-invariant integrand. Since the action of mapping-class group generically changes the spin structure from  $\mathcal{E}$  to say  $\mathcal{E}'$ , the summation over all spin structures guarantees that the resulting expression is mapping-class-group-invariant. 2) the summation over all spin structures is equivalent to imposing GSO projection in the path integral [23].

The compactified moduli stack of punctured spin curves can be viewed as a covering of the compactified moduli stack of ordinary curves, where the covering map corresponds to the forgetting of spin structures. Therefore, the region near the compactification divisors in the moduli stack of spin curves is mapped into the region near the compactification divisors in the moduli stack of ordinary curves. A good set of coordinate near the compactification divisor can be constructed by the so-called *plumbing-fixture construction* [24, 25]. In this construction, the local coordinate around two punctures on a single surface or disjoint surfaces are glued together using appropriate gluing relations<sup>5</sup>. The parameter(s) of the gluing together with the moduli of the component surfaces form a coordinate chart near the compactification divisor of the moduli stack. To compactify the stack of spin curve, one is thus interested in the following question

**Question 1:** *Consider two punctures of the same type on a single surface equipped with a choice of spin structure or on two separate surfaces, each of them equipped with a choice of spin structure. Glue them using the appropriate gluing relation. What is the spin structure of the resulting surface and what does happen to it in the limit that the resulting surface develops a node<sup>6</sup>?*

There is another subtlety in the gluing of punctures in superstring theory. Since the total number of PCOs<sup>7</sup> on a genus- $g$  spin curve with  $n \equiv (n_{\text{NS}}, n_{\text{R}})$  punctures, as we will explain later, is equal to the fermionic dimension of the corresponding superstack, we need to insert  $2g - 2 + n_{\text{NS}} + \frac{1}{2}n_{\text{R}}$  PCOs. From this relation, it turns out that the gluing of two NS punctures reproduces the correct number of PCOs in the resulting surface. However, the total picture number of the surface resulting from the gluing of two R punctures is one less than the one it should be. One thus has to *add* a PCO by hand on the surface with a suitable prescription and check that it reproduces the desired results [17]. Therefore, one is interested in the following question

**Question 2:** *Consider two R punctures on a single surface equipped with a spin structure, or on two separate surfaces each of which is equipped with a choice of spin structure. Glue them using the appropriate gluing relation. What is the reason for the addition of the extra picture-changing operator from the supergeometry point of view?*

---

<sup>5</sup>We refer to the gluing of the local coordinates around the punctures as gluing of punctures.

<sup>6</sup>A node or an ordinary double point on an ordinary, spin curve, or super-Riemann surface is a point whose neighborhood is homeomorphic to two disk joined at a single point, the node.

<sup>7</sup>We exclude the cases where the surface admits conformal-Killing spinors. These cases can be treated separately.

The aim is to show that there is a purely geometric explanation for this missing PCO, otherwise it is obvious that one has to include a PCO in the definition of kinetic term of the Ramond sector.

Although these questions are trivial for experts of the field, we felt that it might be useful for some readers.

However, the main objectives of this work are

1. the proof of the existence of heterotic-string and type-II-superstring field theory vertices;
2. the proof of the representability of the moduli problem of  $\mathcal{N} = 1$  (possibly punctured) stable SUSY curves.

Let us briefly explain each of these problems.

### 1.1.1 The Existence of String Vertices

In the formulation of any string field theory, two choices have to be made: 1) a choice of the worldsheet theory, and 2) a choice of string vertices<sup>8</sup>. These two choices provide us with a way to describe the interaction vertices of the associated string field theory. String vertices are some regions inside the product of moduli spaces parameterizing left- and right-moving sectors of the relevant worldsheet theory. If we denote the moduli space parameterizing left- and right-moving sectors by  $\mathcal{M}_L$  and  $\mathcal{M}_R$ , respectively, then we have the following

- For heterotic-string theory  $\mathcal{M}_L^H = \mathcal{M}_g(n_{NS} + n_R)$ <sup>9</sup>, the moduli space of genus- $g$  ordinary Riemann surfaces with  $n_{NS} + n_R$  borders, and  $\mathcal{M}_R^H = \mathcal{SM}_g(n_{NS}, n_R)$ , the moduli space of genus- $g$  super-Riemann surfaces with  $n_{NS}$  NS boundary components and  $n_R$  R boundary components. Let us denote a choice heterotic-string vertices corresponding to genus- $g$  vertices with  $n_{NS}$  NS boundary components, and  $n_R$  R boundary components by  $\mathcal{V}_g^H(n_{NS}, n_R)$ . Then, the genus- $g$  heterotic-string field theory interaction vertices  $\mathcal{I}_g^H(n_{NS}, n_R)$  is given by

$$\mathcal{I}_g^H(n_{NS}, n_R) = \int_{\mathcal{V}_g^H(n_{NS}, n_R)} \Omega_{\mathcal{T}^H}, \quad (1.1)$$

where  $\mathcal{V}_g^H(n_{NS}, n_R) \subset \mathcal{M}_L^H \times \mathcal{M}_R^H$ , and  $\Omega_{\mathcal{T}^H}$  is some correlation function computed in the worldsheet theory  $\mathcal{T}^H$  of heterotic-string theory.

- For type-II-superstring theory  $\mathcal{M}_L^H = \mathcal{SM}_g(n_{NS}, n_R) = \mathcal{M}_R^H$ . Let us denote a choice type-II-superstring vertices corresponding to genus- $g$  vertices with  $n_{NSNS}$  NSNS,  $n_{RNS}$  RNS,  $n_{NSR}$  NSR, and  $n_{RR}$  RR boundary components by  $\mathcal{V}_g^H(n_{NSNS}, n_{RNS}, n_{NSR}, n_{RR})$ . Then, the

---

<sup>8</sup>For more on these choices see Section 6.1.

<sup>9</sup>Here, by  $n_{NS} + n_R$ , we mean that we ignore all the complications associated to these borders on spin- or super-Riemann surfaces, and we think of them as borders on an ordinary Riemann surface.

genus- $g$  type-II-superstring field theory interaction vertices  $\mathcal{I}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RR}})$  is given by

$$\mathcal{I}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RR}}) = \int_{\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RR}})} \Omega_{\mathbf{T}^{\text{II}}}, \quad (1.2)$$

where  $\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RR}}) \subset \mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$ , and  $\Omega_{\mathbf{T}^{\text{II}}}$  is some correlation function computed in the worldsheet theory  $\mathbf{T}^{\text{II}}$  of type-II-superstring theory.

The gauge-invariance of the string field theory action is guaranteed if the string vertices satisfy the BV quantum master equation (BV QME). However, it is not *a priori* guaranteed that such subspaces exist. In this paper, we show that  $\mathcal{V}_g^{\text{H}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  and  $\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RR}})$  *always* exist.

### 1.1.2 The Moduli Problem of Stable $\mathcal{N} = 1$ SUSY Curves

One problem of interest in super algebraic geometry is the moduli problem of stable  $\mathcal{N} = 1$  (possibly punctured) SUSY curves, and one can ask the representability of the associated moduli functor. By this in the case of the moduli problem of stable  $\mathcal{N} = 1$  (possibly punctured) SUSY curves, we mean a fibered functor  $\mathcal{F}$  which maps a superscheme to the category of  $\mathcal{N} = 1$  (possibly punctured) SUSY curves. The representability requires the existence of a superscheme  $\mathcal{SM}$  such that  $\mathcal{F}$  is isomorphic to the functor of points of  $\mathcal{SM}$ . If it happened to be the case, then  $\mathcal{SM}$  is the fine moduli space for this moduli problem.

There is a catch as it is usual with the moduli problem of stable curves [20]. Since stable  $\mathcal{N} = 1$  SUSY curves have nontrivial automorphisms, there is no fine moduli spaces for the representability of the moduli functor, i.e. there is no superscheme  $\mathcal{SM}$  that classifies all such curves. One thus need to find an alternative solution. One such solution is to extend the category of superschemes to the *category of superstacks*. Then, the moduli functor might be representable by a superstack. So the question is the following: *is the moduli functor associated to the moduli problem of stable  $\mathcal{N} = 1$  (possibly punctured) SUSY curves representable by a stack?* As we will show the moduli functor is more rigid and similar to the case of (possibly punctured) ordinary and spin curves can be represented by a Deligne-Mumford (DM) superstack.

## 1.2 Contributions of the Paper

Although we elaborate on many points, but the main results of this paper can be summarized as follows

### 1) The Existence and Uniqueness of Solution to the BV QME in $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$

Let  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  denote the moduli space of possibly disconnected spin-Riemann surfaces with  $\mathbf{n}_{\text{NS}}$  NS boundary components and  $\mathbf{n}_{\text{R}}$  R boundary components. We denote the moduli stack



of spin curves of connected genus- $g$  spin-Riemann surfaces with  $n_{\text{NS}}$  NS boundary components and  $n_{\text{R}}$  boundary components by  $\mathcal{S}_g(n_{\text{NS}}, n_{\text{R}})$ . We associate a complex to such moduli spaces

$$\mathcal{F}(\mathcal{S}) \equiv \bigoplus_{n_{\text{NS}}, n_{\text{R}}} \mathcal{C}_* \left( \frac{\mathcal{S}(n_{\text{NS}}, n_{\text{R}})}{S^1 \wr S_{n_{\text{NS}}, n_{\text{R}}}} \right).$$

$\mathcal{C}_*$  is the functor of normalized singular chains with coefficient in any field containing  $\mathbb{Q}$ ,  $S^1 \wr S_{n_{\text{NS}}, n_{\text{R}}}$  is the wreath product  $(S^1)^{n_{\text{NS}}+n_{\text{R}}} \rtimes S_{n_{\text{NS}}, n_{\text{R}}}$ .  $S_{n_{\text{NS}}, n_{\text{R}}}$  is the group that permutes NS boundaries and R boundaries separately among each other. The quotient by the wreath product turns  $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$  to the space of bordered spin-Riemann surfaces with unparametrized boundary. We also denote the contribution from connected genus- $g$  surfaces by  $\mathcal{F}_{g, n_{\text{NS}}, n_{\text{R}}}(\mathcal{S})$ .  $\mathcal{F}(\mathcal{S})$  carries the structure of a commutative differential-graded algebra (CDGA)<sup>10</sup>, where differential  $d$  is the boundary map of chain complexes and the product comes from disjoint union of surfaces. Furthermore, we define the following gluing operation: let  $\Delta^{\text{NS}}$  ( $\Delta^{\text{R}}$ ) denotes the operation of gluing of two NS (respectively R) boundaries. We define

$$\Delta \equiv \Delta^{\text{NS}} + \Delta^{\text{R}}.$$

This turns  $\mathcal{F}(\mathcal{S})$  into a BV algebra<sup>11</sup>. We can now state one of the results of this paper

**Theorem** (The Existence and Uniqueness of Solution to the BV QME in  $\mathcal{S}_g(n_{\text{NS}}, n_{\text{R}})$ ). *For each triple  $(g, n_{\text{NS}}, n_{\text{R}})$  with  $2g-2+n_{\text{NS}}+n_{\text{R}} > 0$ , there exists an element  $\mathcal{V}_g(n_{\text{NS}}, n_{\text{R}}) \in \mathcal{F}_{g, n_{\text{NS}}, n_{\text{R}}}(\mathcal{S})$  of homological degree  $6g-6+2n_{\text{NS}}+2n_{\text{R}}$ , with the following properties*

1.  $\mathcal{V}_0(3, 0)$  is the fundamental cycle of  $\mathcal{S}_0(3, 0)/S_1 \wr S_{3, 0}$ , i.e. 0-chain of coefficient  $1/3!$ .
2.  $\mathcal{V}_0(1, 2)$  is the fundamental cycle of  $\mathcal{S}_0(1, 2)/S_1 \wr S_{1, 2}$ , i.e. 0-chain of coefficient  $1/2!$ .
3. The generating function

$$\mathcal{V} \equiv \sum_{2g-2+n_{\text{NS}}+n_{\text{R}} > 0} \hbar^{2g-2+n_{\text{NS}}+n_{\text{R}}} \mathcal{V}_g(n_{\text{NS}}, n_{\text{R}}) \in \hbar \mathcal{F}(\mathcal{S})[[\hbar]], \quad (1.3)$$

---

<sup>10</sup>A differential-graded algebra  $\mathcal{A}$  is a graded algebra equipped with a map  $d : \mathcal{A} \rightarrow \mathcal{A}$  which has degree  $-1$  (chain-complex convention) or degree  $+1$  (cochain-complex convention) with product  $\cdot : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$  that satisfies the following two conditions:

1.  $d \circ d = 0$ : this means that  $d$  turns  $\mathcal{A}$  to a chain or cochain complex depending on its degree.
2. the graded Leibniz rule: For any two elements  $x, y \in \mathcal{A}$ , we have

$$d(x \cdot y) = dx \cdot y + (-1)^{\deg(x)} x \cdot dy,$$

where  $\deg(x)$  is the degree of the element  $x$  in the graded algebra  $\mathcal{A}$ .

<sup>11</sup>Recall that a BV algebra  $\mathcal{A}$  is defined as an algebra with the following structures

1.  $\mathcal{A}$  is a commutative differential-graded algebra with differential  $d$ .
2. it is equipped with an odd differential  $\delta$ , i.e.  $\delta^2 = 0$ .
3.  $\delta$  satisfies  $[\delta, d] = 0$ , where  $[\cdot, \cdot]$  is a graded commutator.

satisfies the BV QME

$$(d + \hbar\Delta) \exp\left(\frac{\mathcal{V}}{\hbar}\right) = 0.$$

4.  $\mathcal{V}$  is unique up to homotopy in the category of BV algebras.

For the proof of this result see Section 4.5 and especially Theorem 4.2. This result is the generalization of Proposition 10.1.1<sup>12</sup> of [26]. The physical interpretation of the result of Costello is the existence and uniqueness of bosonic-string field theory vertices. However, we cannot make a similar claim in the case of heterotic-string and type-II-superstring theories. We need to work more to establish the existence of vertices in these theories, as we explain below.

One might ask *what is the importance of this result for string field theory?* As we argue below, the existence of heterotic-string and type-II superstring field theory vertices can be deduced from this result. The reason is as follows

1. We can associate BV algebras to complexes associated to spaces parameterizing world-sheets of heterotic-string and type-II superstring field theory. These BV algebras are quasi-isomorphic to BV algebra  $\mathcal{F}(\mathcal{S})$  which means that their corresponding homology groups are isomorphic.
2. On the other hand, if two differential-graded algebras are quasi-isomorphic, their sets of homotopy classes of solutions of the Maurer-Cartan equation are isomorphic. The BV QME can be considered as the Maurer-Cartan equation of the corresponding BV algebra (see Lemma 5.2.1 of [26]). Therefore, if two BV algebras are quasi-isomorphic, their sets of homotopy classes of solutions of the BV QME are isomorphic (see Lemma 5.3.1 and Definition 5.4.1 of [26]).
3. Putting 1 and 2 together and noting that heterotic-string and type-II superstring field theory vertices must satisfy the BV QME, we conclude that they exist.

## 2) Solutions to the BV QME in $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$ and Fundamental Classes of DM Stacks

Let  $\overline{\mathcal{S}}_{n_{\text{NS}}, n_{\text{R}}}$  denote the moduli stack of stable spin curves with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  Ramond punctures, and  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  denote the contribution coming from connected genus- $g$  surfaces. We associate a chain complex to these moduli spaces

$$\mathcal{F}(\overline{\mathcal{S}}) \equiv \bigoplus_{n_{\text{NS}}, n_{\text{R}}} \mathcal{C}_* \left( \frac{\overline{\mathcal{S}}_{n_{\text{NS}}, n_{\text{R}}}}{\mathcal{S}_{n_{\text{NS}}, n_{\text{R}}}} \right). \quad (1.4)$$

$\mathcal{S}_{n_{\text{NS}}, n_{\text{R}}}$  permutes the set of NS and R punctures among each other separately. After taking the homology, and setting  $\Delta = 0$ ,  $\mathcal{F}(\overline{\mathcal{S}})$  turns to a BV algebra. We then define the following

---

<sup>12</sup>The numbers of theorem, proposition, etc of [26] that we are referring to are the ones that appear in the published version not the Arxiv preprint. We explicitly mention it if we are referring to the ArXiv version.

formal sum

$$[\overline{\mathcal{S}}] \equiv \sum_{\substack{g, n_{\text{NS}}, n_{\text{R}} \\ 2g-2+n_{\text{NS}}+n_{\text{R}} \geq 0}} \hbar^{2g-2+n_{\text{NS}}+n_{\text{R}}} [\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}] \in \hbar H_*(\mathcal{F}(\overline{\mathcal{S}}))[[\hbar]], \quad (1.5)$$

where  $[\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}]$  is the fundamental class of  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}$ . One can then establish the following Theorem which provides a link between  $\mathcal{V}$ , the solution of the BV quantum master equation in the BV algebra  $\mathcal{F}(\mathcal{S})$  defined in (1.3), and  $[\overline{\mathcal{S}}]$ .

**Theorem** (Solutions to the BV QME in  $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$  and Fundamental Classes of DM Stacks). *There is a map  $\mathcal{F}(\mathcal{S}) \rightarrow H_*(\mathcal{F}(\overline{\mathcal{S}}))$  in the homotopy category of BV algebras that sends  $\mathcal{V} \rightarrow [\overline{\mathcal{S}}]$ . This means that in the homotopy category of BV algebras  $\mathcal{V}_{g, n_{\text{NS}}, n_{\text{R}}}$  is homotopy-equivalent to the orbifold fundamental class  $[\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}]$  of the moduli stack of connected stable genus- $g$  spin curves with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  R punctures, i.e. it is homotopy-equivalent to the whole moduli stack.*

For the proof of this result see Theorem 4.3 in Section 4.6.

### 3) The Existence of Solution to the BV QME in $\mathcal{SM}(n_{\text{NS}}, n_{\text{R}})$

Let  $\mathcal{SM}(n_{\text{NS}}, n_{\text{R}})$  denote the space of bordered  $\mathcal{N} = 1$  super-Riemann surfaces with  $n_{\text{NS}}$  NS and  $n_{\text{R}}$  R boundary components. We can associate a singular superchain<sup>13</sup> complex to these spaces

$$\mathcal{F}(\mathcal{SM}) \equiv \bigoplus_{n_{\text{NS}}, n_{\text{R}}} \mathcal{SC}_* \left( \frac{\mathcal{SM}(n_{\text{NS}}, n_{\text{R}})}{S^1 \wr S_{n_{\text{NS}}, n_{\text{R}}}} \right) \quad (1.6)$$

where  $\mathcal{SC}_*$  is the functor of normalized singular simplicial superchains of top fermionic dimension with coefficients in any field containing  $\mathbb{Q}$ . This is a differential-graded algebra, where the derivative is the boundary operation, as defined in Definition 5.1, acting on singular superchains. We can define the gluing operation as follows

$$\Delta_{\mathfrak{s}} \equiv \Delta_{\mathfrak{s}}^{\text{NS}} + \Delta_{\mathfrak{s}}^{\text{R}}, \quad (1.7)$$

where  $\Delta_{\mathfrak{s}}^{\text{NS}}$  denotes the gluing of two NS boundaries and  $\Delta_{\mathfrak{s}}^{\text{R}}$  is the gluing operation of two R boundaries. This turns  $\mathcal{F}(\mathcal{SM})$  to a BV algebra. We then have the following

**Theorem** (The Existence of Solution to BV QME in  $\mathcal{SM}(n_{\text{NS}}, n_{\text{R}})$ ). *There exists a solution to the BV QME in  $\mathcal{F}(\mathcal{SM})$ .*

For the proof of this Theorem see Section 5.3.

We might need to stress one point here. We have constructed a solution to the BV QME in the space of bordered spin-Riemann surfaces. In principle, we should be able to *thicken* this solution in fermionic direction to find a solution to the BV QME in the moduli space of bordered  $\mathcal{N} = 1$  super-Riemann surfaces. However, life is not that simple. As we will

---

<sup>13</sup>For the definition of singular superchains on superspaces see Section 5.2.

rigorously prove in Section 5.1.1, to construct a supermanifold  $M$ , we can start from its reduced space  $M_{\text{red}}$  and thicken it in the fermionic direction. However, this defines  $M$  up to *homology*<sup>14</sup>. This means that any two different thickenings in the fermionic directions can be connected by a homotopy, which is intuitively due to the fact that fermionic directions are infinitesimal. Therefore, by considering the solutions to the BV QME in  $\mathcal{S}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  *as a smooth subspace* and thickening it in all fermionic directions, we get a subspace of  $\mathcal{SM}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  which *does not necessarily satisfies the BV QME*. Therefore, we need to come-up with another way to prove the existence of such solutions in  $\mathcal{SM}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$ , as we do in Section 5.3.

Perhaps we must stress that the idea of thickening does not work in the realm of complex supermanifolds. It is established that there are always obstructions in extending a holomorphic thickening to a complex supermanifold [28]. However, we are only interested in integration over these spaces, and for that reason, the results of integration over different thickenings are the same, and do not depend on the choice of thickening in the fermionic direction [10].

#### 4) The Existence of Heterotic-String and Type-II-Superstring Vertices

The next result of the paper is the proof of existence of heterotic-string and type-II superstring field theory vertices. As we stated above, the proof of this result depends on the first result, i.e. the existence of solution to the BV QME in  $\mathcal{S}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$ .

To state the result for the heterotic-string field theory, we define the following complexes

$$\mathcal{F}((\mathcal{M}_L^{\text{H}} \times \mathcal{M}_R^{\text{H}})^{\mathbf{D}}) \equiv \bigoplus_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} \mathcal{SC}_* \left( \frac{(\mathcal{M}_L^{\text{H}}(\mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}}) \times \mathcal{M}_R^{\text{H}}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}))^{\mathbf{D}}}{S_L^{\text{H}} \times S_R^{\text{H}}} \right), \quad (1.8)$$

where superscript  $\mathbf{D}$  denotes diagonal and subscript red means that we consider the reduced spaces, and

$$S_L^{\text{H}} \equiv S_1 \wr S_{\mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}}}, \quad S_R^{\text{H}} \equiv S_1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}, \quad (1.9)$$

where  $S_1 \wr S_{\mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}}} = (S_1)^{\mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}}} \rtimes S_{\mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}}}$  and  $S_1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} = (S_1)^{\mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}}} \rtimes S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ . We then have the following result

**Theorem (The Existence of Heterotic-String Vertices).** *There exists a solution to the BV QME in  $\mathcal{F}((\mathcal{M}_L^{\text{H}} \times \mathcal{M}_R^{\text{H}})^{\mathbf{D}})$ . The genus- $g$  part is a genus- $g$  heterotic-string field theory vertex with  $\mathfrak{n}_{\text{NS}}$  NS and  $\mathfrak{n}_{\text{R}}$  R boundary components.*

For the proof of this Theorem see Section 6.2.

To state the result for the case of type-II-superstring field theory, we define the following complexes

$$\mathcal{F}((\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}) \equiv \bigoplus_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} \mathcal{SC}_* \left( \frac{(\mathcal{M}_L^{\text{II}}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}) \times \mathcal{M}_R^{\text{II}}(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}))^{\mathbf{D}}}{S_L^{\text{II}} \times S_R^{\text{II}}} \right). \quad (1.10)$$

---

<sup>14</sup>This can also be seen from the result of [27] where it is proven that any *smooth* thickening can be extended to a smooth supermanifold, and conversely any smooth supermanifold may be obtained in this way.

where

$$S_L^{\text{II}} \equiv S_1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} = S_R^{\text{II}}. \quad (1.11)$$

We then have the following result

**Theorem (The Existence of Type-II-Superstring Vertices).** *There exists a solution to the BV QME in  $\mathcal{F}((\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}})$ . The genus- $g$  part is a genus- $g$  type-II-string field theory vertex with  $\mathfrak{n}_{\text{NSNS}}$  NSNS,  $\mathfrak{n}_{\text{RNS}}$  RNS,  $\mathfrak{n}_{\text{NSR}}$  NSR, and  $\mathfrak{n}_{\text{RR}}$  RR boundary components.*

For the proof of this Theorem see Section 6.3.

### 5) The Representability of the Moduli Problem of Stable $\mathcal{N} = 1$ SUSY Curves

There is a model for the compactification of the space of smooth genus- $g$  possibly punctured  $\mathcal{N} = 1$  SUSY curves introduced by Deligne [29]. Let  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  denotes the space of stable genus- $g$   $\mathcal{N} = 1$  SUSY curve with  $\mathfrak{n}_{\text{NS}}$  NS punctures and  $\mathfrak{n}_{\text{R}}$  Ramond punctures. Then, we have the following result

**Theorem (The Representability of the Moduli Problem of Stable  $\mathcal{N} = 1$  SUSY Curves).** *Assume that a triple of natural numbers  $(g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  satisfies  $\mathfrak{n}_{\text{R}} \in 2\mathbb{Z}_{\geq 0}$ , and  $2g - 2 + \mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}} > 0$ .*

- (1)  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  is a smooth proper Deligne-Mumford superstack of dimension

$$\left(3g - 3 + \mathfrak{n}_{\text{NS}} + \mathfrak{n}_{\text{R}} \mid 2g - 2 + \mathfrak{n}_{\text{NS}} + \frac{\mathfrak{n}_{\text{R}}}{2}\right).$$

Moreover its bosonic truncation  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}; \text{bos}}$  is the moduli stack  $\overline{\mathcal{S}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  of punctured stable spin curves.

- (2)  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  has a coarse moduli superspace  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  which is actually bosonic and is also the coarse moduli space for the bosonic quotient  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}; \text{ev}}$ . Moreover its reduced subspace  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}; \text{red}}$  is the coarse moduli space  $\overline{\mathcal{S}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  of punctured stable spin curves.

For the proof of this Theorem see Section 7.3.

There is an alternative model for the compactification of the space of stable  $\mathcal{N} = 1$  SUSY curve different from the one proposed by Deligne based on studying the spaces of stable log twisted SUSY curves [30]. This model will not be pursued here. Recently, the moduli spaces of *smooth* SUSY curves with  $R$  punctures have also been constructed in the literature [31, 32].

### 1.3 Notations and Remarks

For the convenience of reader, we collect some notations used in this paper

- $\mathcal{M}_{g, \mathfrak{n}}$ : the moduli stack of genus- $g$  Riemann surfaces with  $\mathfrak{n}$  punctures.
- $\overline{\mathcal{M}}_{g, \mathfrak{n}}$ : the moduli stack of stable genus- $g$  Riemann surfaces with  $\mathfrak{n}$  punctures.

- $\partial\overline{\mathcal{M}}_{g,n}$ : the boundary (i.e. points at infinity) of  $\overline{\mathcal{M}}_{g,n}$ . It is isomorphic to  $\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}$ .
- $\mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}}$ : the moduli stack of genus- $g$  spin curves with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures. The moduli stack of genus- $g$  spin curves with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures with an even spin structure is denoted as  $\mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}}^{\text{ev}}$  for the even spin structure, and  $\mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}}^{\text{od}}$  for an odd spin structure.
- $\overline{\mathcal{S}}_{n_{\text{NS}},n_{\text{R}}}$ : the stable-curve compactification of the moduli stack of possible disconnected spin curves with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures
- $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}$ : the compactified moduli stack of connected genus- $g$  spin curves with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures. This moduli stack is the component of  $\overline{\mathcal{S}}_{n_{\text{NS}},n_{\text{R}}}$  coming from the connected genus- $g$  surfaces. The moduli stack of genus- $g$  spin curves with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures with a specific spin structure is denoted as  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}^{\text{ev}}$  for the even spin structure, and  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}^{\text{od}}$  for the odd spin structure.
- $\partial\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}$ : the boundary (i.e. points at infinity) of the compactified moduli stack of genus- $g$  spin curves with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures. It is isomorphic to  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}} \setminus \mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}}$ .
- $\mathcal{S}_{g,n}^{\mathbf{m}}$ : The moduli stack of genus- $g$  spin curve with  $n$  punctures such that the vector  $\mathbf{m} \equiv (m_1, \dots, m_n)$  is a vector whose components  $m_a$  are the order of vanishing at punctures  $\mathbf{p}_a$ . Let us make this a bit more precise. As we will explain in section 2, there is a distinguished subbundle  $\mathcal{D}$  inside the tangent bundle  $T\mathcal{R}$  of an  $\mathcal{N} = 1$  super-Riemann surface  $\mathcal{R}$  which is nonintegrable. This means that there exists a short exact sequence

$$0 \longrightarrow \mathcal{D} \longrightarrow T\mathcal{R} \longrightarrow T\mathcal{R}/\mathcal{D} \longrightarrow 0. \quad (1.12)$$

The bundle  $T\mathcal{R}/\mathcal{D}$  is generated by  $\partial_z$ . When there is no punctures, we will see that the bundle  $\mathcal{D}^2$  is also generated by  $\partial_z$ . Once punctures are introduced, the bundle  $T\mathcal{R}/\mathcal{D}$  is still generated by  $\partial_z$ . However, it is not necessarily true that  $\mathcal{D}^2$  is generated by  $\partial_z$ . For example, we will see in section 2 that in the presence of Ramond puncture  $\mathcal{D}^2$  is actually generated by  $P(z)\partial_z$ , where  $P(z)$  has a simple zero at the location of the Ramond puncture  $\mathbf{p}_a$ , i.e.  $P(z) = z - z_a$ . We thus have  $\mathcal{D}^2 \simeq T\mathcal{R}/\mathcal{D} \otimes \mathcal{O}(-\mathcal{D})$ , where

$$\mathcal{D} \equiv \sum_{a=1}^{n_{\text{R}}} \mathcal{D}_a = \sum_{a=1}^{n_{\text{R}}} \mathbf{p}_a, \quad (1.13)$$

is the divisor defined by the locations of Ramond punctures  $\mathbf{p}_a$  at  $z = z_a$ . This means that a section of  $\mathcal{D}^2$  is a section of  $T\mathcal{R}/\mathcal{D}$  which vanishes along the divisor  $\mathcal{D}$ . In general,  $\mathcal{D}^2$  is generated by  $P(z)\partial_z$ , where now  $P(z)$  has higher-order zeros at the location of punctures. If near the puncture  $\mathbf{p}_a$ ,  $P(z)$  has zero of order  $m_a$ , then  $\mathcal{D}^2 \simeq$

$T\mathcal{R}/\mathcal{D} \otimes \mathcal{O}(-\mathcal{D})$  where now

$$\mathcal{D} = \sum_{a=1}^n \mathcal{D}_a = \sum_{a=1}^n m_a \mathfrak{p}_a. \quad (1.14)$$

This means that a section of  $\mathcal{D}^2$  is a section of  $T\mathcal{R}/\mathcal{D}$  which vanishes along the divisor  $\mathcal{D}$  with zeros of order  $m_a$  at the puncture  $\mathfrak{p}_a$ , i.e.  $P(z) = \prod_a (z - z_a)^{m_a}$ . Once we reduce from an  $\mathcal{N} = 1$  super-Riemann surface to the underlying spin curve,  $\mathcal{D}$  is essentially a choice of spin structure. Therefore,  $\mathcal{S}_{g,n}^{\mathbf{m}}$  is the moduli stack of genus- $g$  spin curve with  $n$  punctures such that the vector  $\mathbf{m}$  is the vector of the order of vanishing at punctures. For the cases we are interested in, the order of vanishing are as follows:  $m_a = 0$  for NS punctures and  $m_a = 1$  for Ramond punctures.

- $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$ : the moduli space of possibly disconnected spin-Riemann surfaces with  $n_{\text{NS}}$  Neveu-Schwarz boundary components (see Definition 4.2) and  $n_{\text{R}}$  Ramond boundary components (see Definition 4.4).
- $\mathcal{S}_g(n_{\text{NS}}, n_{\text{R}})$ : the moduli space of connected genus- $g$  spin-Riemann surfaces with  $n_{\text{NS}}$  Neveu-Schwarz boundary components and  $n_{\text{R}}$  Ramond boundary components.
- $\tilde{\mathcal{S}}(n_{\text{NS}}, n_{\text{R}})$ : the moduli space of stable spin-Riemann surfaces, decorated at each Neveu-Schwarz puncture with a ray in the tangent space, at each Ramond punctures with a ray in the spinor bundle, at each Neveu-Schwarz node with a ray in the tensor product of rays of the tangent spaces at each side, and at each Ramond node with a ray in the spinor bundle. We emphasize that a surface in  $\tilde{\mathcal{S}}(n_{\text{NS}}, n_{\text{R}})$  does not have boundary components.  $\tilde{\mathcal{S}}(n_{\text{NS}}, n_{\text{R}})$  is however homotopy-equivalent to  $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$ . As we will explain, there is a crucial fact about the gluing of Ramond punctures in  $\tilde{\mathcal{S}}(n_{\text{NS}}, n_{\text{R}})$  which makes it more suitable than  $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$  to work with.
- $\mathcal{SM}_{g,n_{\text{NS}},n_{\text{R}}}$ : the moduli space of genus- $g$   $\mathcal{N} = 1$  super-Riemann surfaces with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures.
- $\overline{\mathcal{SM}}_{g,n_{\text{NS}},n_{\text{R}}}$ : the compactified superstack of genus- $g$   $\mathcal{N} = 1$  super-Riemann surfaces with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures.
- $\mathcal{SM}_g(n_{\text{NS}}, n_{\text{R}})$ : the moduli space of genus- $g$   $\mathcal{N} = 1$  super-Riemann surfaces with  $n_{\text{NS}}$  Neveu-Schwarz boundary components and  $n_{\text{R}}$  Ramond boundary components.

**Remark 1:** When we say *puncture*, we do not mean removing the point from the surface. The surfaces are still closed with marked points on them. There is a delicate difference between marked points and punctures in the presence of spin structure. We only use this terminology to cope with the literatures of superstring theory. We also would like to mention that that since a Ramond puncture is part of the defining data of a superconformal structure, it defines

a divisor (we make this statement precise in section 2.2). Therefore, Ramond divisor is a better terminology, as has been used in [11, 12].

**Remark 2:** We use the notation  $(z|\theta)$  to denote the local coordinates on a  $\mathcal{N} = 1$  super-Riemann surface. We also use the notation  $(p|q)$  to denote the dimension of supermanifolds, supervector spaces, etc with  $p$  even directions and  $q$  fermionic directions. An exception is when we talk about pseudoforms. In this case,  $p$  is related to the form degree of pseudoforms and  $-q$  is the so-called picture number of pseudoforms, as we shall explain in Appendix A.

**Remark 3:** A surface equipped with a line bundle  $\mathcal{E}$  whose  $r^{\text{th}}$  tensor power is isomorphic to the canonical line bundle,  $\mathcal{E}^{\otimes r} \simeq \omega$ , is called an  $r$ -spin curve and  $\mathcal{E}$  is called a generalized spin structure. We are interested in the case that  $r = 2$  corresponding to the ordinary spin structure, and as such it should be called a 2-spin curve. To avoid cluttering, we call them spin curves. A discussion of spin curves defined over a base scheme can be found in Appendix B.

**Remark 4:** We denote the *signature* of a genus- $g$  spin curve or  $\mathcal{N} = 1$  super-Riemann surface with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  R punctures as  $(g; n_{\text{NS}}, n_{\text{R}})$ .

**Remark 5:** For *punctured surfaces*, we use ordinary, spin- or SUSY curves (the language of algebraic geometry where we can work with curves over an algebraically-closed field  $\mathbb{F}^{15}$ ) and ordinary, spin- or super-Riemann surfaces (the language of analytic geometry where  $\mathbb{F} = \mathbb{C}$ ) interchangeably throughout the text. We use the terms *bordered* ordinary, spin- or super-Riemann surfaces for surfaces with geodesic boundary components. For example, We call a genus- $g$  Riemann surface with  $n_{\text{NS}}$  Neveu-Schwarz punctures and  $n_{\text{R}}$  Ramond punctures equipped with a spinor bundle  $\mathcal{E}$ , a genus- $g$  punctured spin curve<sup>16</sup>. Sometimes we omit  $\mathcal{N} = 1$  but we always mean  $\mathcal{N} = 1$  (smooth or stable) SUSY curve.

**Remark 6:** When it is needed to describe more than one surface, we denote the signature of the  $a^{\text{th}}$  surface by  $(g_a; n_{\text{NS}}^a, n_{\text{R}}^a)$ . Here,  $n_{\text{NS}}^a$  ( $n_{\text{R}}^a$ ) could denote either the  $a^{\text{th}}$  NS (respectively R) puncture or NS (respectively R) boundary.

The organization of the paper is as follows. In Section 2, we define spin curves, punctures on them, and the gluing of punctures. In Section 3, we investigate the relation between gluing of punctures on spin curves and the compactification of the moduli stack of spin curves, and also a subtlety in the gluing of Ramond punctures. We then establish the existence and uniqueness of a solution to the BV quantum master equation in the moduli space of bordered spin-Riemann surfaces, and their relation to fundamental classes of DM spaces in Section 4. In Section 5, we prove the existence of solution to the BV QME in the moduli space of bordered  $\mathcal{N} = 1$  super-Riemann surfaces. Section 6 is devoted to the proof of the existence of heterotic-string and type-II-superstring vertices. Finally, we turn to the moduli problem of  $\mathcal{N} = 1$

<sup>15</sup>One can define curves over fields which are not algebraically-closed. We are not considering such curves in this work.

<sup>16</sup>There are generalized  $r$ -spin curves, the curves equipped with a line bundle  $\mathcal{E}$  whose  $r^{\text{th}}$  tensor power is isomorphic to  $\omega$ .



SUSY curves in Section 7. In the Appendix A, we explain the picture-changing operation from canonical, path-integral and supergeometry point of view, and their equivalence. The compactification of the moduli stack of spin curves is the subject of Appendix B.

## 2 Spin Curves, Punctures and their Gluing

In this section, we define spin curves and various punctures on them. We then explain the gluing of punctures which can be used to construct new surfaces.

### 2.1 Spin Curves over $\mathbb{C}$

The gauge-fixed worldsheet action of the RNS-superstring theory has superconformal (gauge) invariance and, as such, the worldsheet theory should only be sensitive to the superconformal structure of the worldsheet. This means that the worldsheet cannot be a general  $(1|1)$  complex supermanifold whose transition functions are arbitrary superholomorphic functions. Instead, it should be a  $(1|1)$  complex supermanifold with superconformal transition functions, i.e. an  $\mathcal{N} = 1$  super-Riemann surface.

An  $\mathcal{N} = 1$  super-Riemann surface is a complex supermanifold of dimension  $(1|1)$  with the following properties

- 1) It can be locally mapped to  $\mathbb{C}^{1|1}$ . Each local chart is parametrized by  $(z|\theta)$ , where  $z$  is a local complex commuting coordinate and  $\theta$  is a complex anti-commuting coordinate. Since we are interested in super-Riemann surfaces appearing in the heterotic-string theory, the complex conjugate of  $\theta$  does not appear, and as such there is no reality condition on  $\theta$  [33]. A similar argument holds for the case of surfaces appearing in the type-II superstring theories.
- 2) The transition function between the overlapping charts  $(z|\theta)$  and  $(z'|\theta')$  is given by superconformal coordinate transformations, i.e. those coordinate transformations that satisfy the following relations

$$D_\theta z' = \theta' D_\theta \theta', \quad (2.1)$$

where  $D_\theta$  is the spinor derivative

$$D_\theta \equiv \frac{\partial}{\partial \theta} + \theta \frac{\partial}{\partial z}. \quad (2.2)$$

A local coordinate  $(z|\theta)$  in which  $D_\theta$  takes the form (2.2) is called a *superconformal coordinate system*. On any super-Riemann surface there is a local superconformal coordinate  $(z|\theta)$  such that  $D_\theta$  has the form (2.2). For the proof of this fact, see Lemma 1.2 of [4] or Lemma 3.1 of [6].  $D_\theta$  generates a  $(0|1)$ -dimensional subbundle  $\mathcal{D}$  of the tangent bundle such that if  $s$  is a nonzero local section of  $\mathcal{D}$ , then  $s$  and  $s^2 \equiv \frac{1}{2}\{s, s\}$  are linearly independent<sup>17</sup>, i.e. the section is *completely nonintegrable*.

---

<sup>17</sup>As we explain later, this statement fails to hold in the presence of Ramond punctures.

Another useful and equivalent way to define an  $\mathcal{N} = 1$  super-Riemann surface which is closer to the supergravity description is as follows. A  $(1|1)$ -dimensional supermanifold  $\mathcal{R}$  is locally parametrized by  $\mathbf{z} \equiv (x_1, x_2; \theta_1, \theta_2)$ , where  $x_a$  and  $\theta_a$  for  $a = 1, 2$  are real. The geometry of the worldsheet is described by a supermetric. Dealing with supermetric is easier if we use the language of frame fields. As such we introduce the vierbein  $\mathbf{E}_A(\mathbf{z}) = E_A{}^B(\mathbf{z})\partial_B$  on each chart  $(\mathbf{z}_\alpha, \mathbf{u}_\alpha)$ , subject to the torsion constraints required to remove the redundant components [34]. Here  $A, B = x_1, x_2, \theta_1, \theta_2$ . There is always a choice of supercoordinate  $\mathbf{z} \equiv (z, \bar{z}; \theta, \bar{\theta})$ , where  $\bar{z}$  and  $\bar{\theta}$  are the complex conjugate of  $z$  and  $\theta$  respectively, such that the complex superfield  $\mathbf{E}(\mathbf{z}) = E^A(\mathbf{z})\partial_A$  is proportional to the spinor derivative (2.2)

$$\mathbf{E}(\mathbf{z}) = \rho(\mathbf{z})D_\theta, \quad (2.3)$$

for some function  $\rho(\mathbf{z})$ .  $\{\mathbf{E}, \bar{\mathbf{E}}; \mathbf{E}^2, \bar{\mathbf{E}}^2\}$  form a basis for the tangent space at each point  $\mathbf{z}$ . (2.3) is invariant precisely under coordinate transformations (2.1). If we choose to work with superconformal coordinate transformations in the overlap of patches  $\mathbf{u}_\alpha$  and  $\mathbf{u}_\beta$ , we end-up with an atlas which defines a superconformal structure on  $\mathcal{R}$ . A  $(1|1)$ -dimensional supermanifold endowed with such a superconformal structure is called an  $\mathcal{N} = 1$  super-Riemann surface. For a mathematically-precise definition of a family of *smooth*  $\mathcal{N} = 1$  super-Riemann surfaces, see Definition 7.10.

All super-Riemann surfaces we are dealing with in heterotic-string and type-II-superstring theories have trivial topology in the fermionic direction. This restriction is implemented by endowing the surface with the DeWitt topology. Although nontrivial topology in fermionic direction is also possible [35, 36], it can be argued that they are not contribute to the path integrals in string theory [37].

In the case of ordinary Riemann surface, the operator  $\partial_z$  transforms as  $\left(\frac{\partial z'}{\partial z}\right)^{-1} \partial_z$ . Then, the Dolbeault operator  $\partial \equiv dz \otimes \partial_z$  is globally-defined. Similarly, on an  $\mathcal{N} = 1$  super-Riemann surface, the spinor derivative transforms as

$$D_{\theta'} = D_{\theta'} \theta D_\theta. \quad (2.4)$$

We can define an operator

$$D \equiv [dz|d\theta] \otimes D_\theta, \quad (2.5)$$

which can be thought of as the super version of the usual Dolbeault operator.  $[dz|d\theta]$  denotes a section of the canonical (or Berezinian) line bundle (similar to the ordinary case that  $dz$  is a section of the usual canonical bundle). Under a coordinate transformation  $(z|\theta) \longrightarrow (z'|\theta')$ , we have

$$\begin{aligned} D' &= [dz'|d\theta'] \otimes D_{\theta'} = \text{Ber} \left( \frac{\partial(z'|\theta')}{\partial(z|\theta)} \right) \cdot (D_{\theta'} \theta) [dz|d\theta] \otimes D_\theta \\ &= (D_\theta \theta') \cdot (D_{\theta'} \theta) [dz|d\theta] \otimes D_\theta = [dz|d\theta] \otimes D_\theta = D, \end{aligned} \quad (2.6)$$

where  $\text{Ber}(\cdots)$  denotes the Berezinian or superdeterminant<sup>18</sup>. We thus conclude that  $D$  is a globally-defined operator and  $D_\theta$  is a section of the line bundle dual to the canonical line bundle.

The generic solutions of (2.1) are given by [38]

$$\begin{aligned} z' &= f(z) + \theta(\partial_z f(z))\varepsilon(z), \\ \theta' &= (\partial_z f(z))^{\frac{1}{2}} \left( \theta + \varepsilon(z) + \frac{1}{2}\theta\varepsilon(z)\partial_z\varepsilon(z) \right), \end{aligned} \quad (2.7)$$

where  $f(z)$  and  $\varepsilon(z)$  are commuting and anticommuting functions and both of which are holomorphic functions of  $z$ . In the absence of odd moduli for the super-Riemann surface, i.e.  $\varepsilon(z) = 0$ , the transition functions (2.7) reduce to

$$\begin{aligned} z' &= f(z), \\ \theta' &= (\partial_z f(z))^{\frac{1}{2}} \theta, \end{aligned} \quad (2.8)$$

on the intersection of two superconformal patches. The first equation is the transition function between two overlapping charts on an ordinary Riemann surface  $\mathcal{R}$ , i.e. a conformal transformation, and the second is the coordinate transformation of the fibers of a fiber bundle on  $\mathcal{R}$ . Together, they are transition functions of the total space of a spinor bundle  $\mathcal{E}$  over  $\mathcal{R}$ . The choice of square root in the transition function of  $\theta$  corresponds to the choice of spin structure. For a mathematically-precise definition of spin curves see Definitions B.1 and B.5.

By going around each 1-cycle of  $\mathcal{R}$ ,  $\theta$  can be either periodic or anti-periodic. For a genus- $g$  surface  $\mathcal{R}$ , since  $\dim(H_1(\mathcal{R}, \mathbb{Z})) = 2g$ , there are  $2g$  possible non-trivial 1-cycles, and therefore  $2^{2g}$  possible choices of signs for the transition functions of  $\theta$ , i.e. there are  $2^{2g}$  spin structures. We thus conclude that *a genus- $g$  super-Riemann surface with vanishing odd moduli is an ordinary genus- $g$  Riemann surface. Since the transition function for  $\theta$  depends on the square root of  $f(z)$ , we get a genus- $g$  spin curve.* This discussion shows that  $\mathcal{S}_g$ , the moduli stack of genus- $g$  spin curves is the reduced space<sup>19</sup> of  $\mathcal{SM}_g$ , the superstack of genus- $g$   $\mathcal{N} = 1$  super-Riemann surfaces. It is also a  $2^{2g}$ -fold covering of  $\mathcal{M}_g$ , the moduli stack of ordinary genus- $g$  Riemann surfaces. We thus have the following

$$\mathcal{SM}_g \xleftarrow{\iota_S} \mathcal{S}_g \xrightarrow{\pi_M} \mathcal{M}_g, \quad (2.9)$$

where  $\iota_S$  is a closed embedding [6], and  $\pi_M$  is a covering map of degree  $2^{2g}$ . There is a similar diagram for the punctured spin curves and compactified moduli stacks. In the construction of the compactification divisor of  $\overline{\mathcal{S}}_g$  (or  $\overline{\mathcal{S}}_{g, \text{nNS}, \text{nR}}$  in the presence of NS and R punctures) using plumbing of surfaces, one needs to deal with *gluing of spin bundles* on the original surface(s).

<sup>18</sup>Note that  $\cdot$  in the above expression does not mean anything special and we inserted it for clarity of expressions.

<sup>19</sup>The reduced space of a supermanifold is an ordinary manifold which can be obtained by putting equal to zero all of the odd coordinates and all (possible) odd moduli of the supermanifold. For more on this see [10].

The question that arises is the proper way to glue spin bundles to get a well-defined notion of spin structure. It turns out that this is not a trivial task [39, 40]. We clarify some aspects of this construction in Section 2.3, and give a detailed account of the construction in Appendix B.

## 2.2 Punctures on Spin and Super-Riemann Surfaces

In heterotic-string theory, one has to specify the boundary condition for the worldsheet fermions, which could be either periodic or anti-periodic. The inclusion of both types of boundary conditions is necessary for the unitarity of loop amplitudes [41]. Since these two types of boundary conditions cannot be continuously deformed into each other, the total Hilbert space  $\mathcal{H}$  consists of two sectors with their own vacua: the Neveu-Schwarz (NS) sector  $\mathcal{H}_{\text{NS}}$  corresponding to the periodic boundary condition and the Ramond (R) sector  $\mathcal{H}_{\text{R}}$  corresponding to the antiperiodic boundary condition

$$\mathcal{H} = \mathcal{H}_{\text{NS}} \oplus \mathcal{H}_{\text{R}}. \quad (2.10)$$

The external states can be from either of these two sectors. From the state-operator correspondence, there are thus two types of punctures associated to the external states [42]

- **Neveu-Schwarz or Super Punctures:** This type of puncture is just a marked point on the surface. In a local coordinate  $(z|\theta)$ , any point  $(z|\theta) = (z_0|\theta_0)$  can be the location of an NS puncture. Any NS puncture defines a divisor  $\Delta_\alpha$  given by  $z = z_0 + \alpha\theta$  and  $\theta = \theta_0 + \alpha$ , for an anti-commuting variable  $\alpha$ . This divisor is the orbit through the NS puncture located at  $(z_0|\theta_0)$  and is generated by  $D_\theta$ <sup>20</sup> [11]. There can be any number of NS puncture on an  $\mathcal{N} = 1$  super-Riemann surface. Therefore, inserting an NS puncture on a super-Riemann surface increases the complex dimension of its superstack by  $(1|1)$ ;
- **Ramond or Spin Punctures:** This type of puncture is actually a singularity in the superconformal structure (and not the surface itself), and so is part of the defining data of a superconformal structure. What do we mean by this is the following. Away from an R puncture, there is a local superconformal coordinate system  $(z|\theta)$  such that

$$D = [dz|d\theta] \otimes D_\theta = [dz|d\theta] \otimes \left( \frac{\partial}{\partial \theta} + \theta \frac{\partial}{\partial z} \right). \quad (2.11)$$

The local expression of the spinor derivative satisfies the following relation

$$D_\theta^2 = \frac{1}{2} \{D_\theta, D_\theta\} = \frac{\partial}{\partial z}. \quad (2.12)$$

In particular,  $D_\theta^2$  is linearly independent of  $D_\theta$  and non-zero everywhere. However, near an R puncture, there is a local coordinate  $(w|\eta)$  such that

$$D = [dw|d\eta] \otimes D_\eta = [dw|d\eta] \otimes \left( \frac{\partial}{\partial \eta} + \eta(w - w_0) \frac{\partial}{\partial w} \right), \quad (2.13)$$

---

<sup>20</sup>Consider a point  $p$  on a super-Riemann surface and an odd tangent vector  $v$ . Then, there is a subspace of dimension  $(0|1)$  that passes through  $p$  and has  $v$  as the tangent vector.

which satisfies

$$D_\eta^2 = \frac{1}{2}\{D_\eta, D_\eta\} = (w - w_0)\frac{\partial}{\partial w}, \quad (2.14)$$

where  $w_0$  is the location of insertion of the R puncture. This expression shows that  $D_\theta^2$  vanishes at  $w - w_0 = 0$ , i.e. along the divisor defined by the location of the R puncture<sup>21</sup>. A possible change of coordinates is

$$(w|\eta) \longrightarrow (z|\theta) = (w - w_0|(w - w_0)^{1/2}\eta), \quad (2.15)$$

where  $D$  retain its usual form as (2.5). However, this change of coordinate introduces cuts between pairs of R punctures. Since by going around an R punctures  $\theta \longrightarrow -\theta$ , there can not be a single R puncture, i.e. R punctures come in pairs<sup>22</sup>. Each pair of R punctures contributes two even and two odd moduli to the complex dimension of superstack. However, one of the odd moduli of a pair of R punctures can be removed by the conformal-Killing spinor of a disk with two R punctures [43]. Therefore, inserting a pair of R punctures on a super-Riemann surface increases the complex dimension of its superstack by (2|1).

On a spin curve, the punctures are given by a point on the surface and a choice of  $D_\theta$ . Just as the case of super-Riemann surfaces, the choice of  $D_\theta$  depends on whether the puncture is of NS or R type. We now turn to the gluing of punctures on a spin- or an  $\mathcal{N} = 1$  super-Riemann surface.

### 2.3 Gluing of Punctures on Spin and Super-Riemann Surfaces

In this section, we describe the gluing of two punctures on a spin or an  $\mathcal{N} = 1$  super-Riemann surface. Intuitively, the gluing corresponds to propagation of states between the glued states, just like a Feynman diagram. The direct sum structure of the decomposition of the total Hilbert space (2.10) shows that there can not be propagation between a state from NS sector and a state from R sector, since otherwise there can be mixing of states. We can thus only glue two punctures associated to states from the same sector of the theory, i.e. we can glue either two NS punctures or two R punctures. On the other hand, the gluing can be done either on a single surface or on two separate surfaces. If we consider the underlying spin surfaces of the component super-Riemann surfaces, the surfaces are endowed with spin structures, the gluing relations are different for the two types of punctures. When two punctures are glued, the corresponding spin bundles are glued as well. For a spin curve, we thus need a gluing relation that describes the gluing of surfaces as well as gluing of spinor bundles.

Consider two punctures on a single or two disconnected  $\mathcal{N} = 1$  super-Riemann surfaces. The neighborhoods of these punctures can be described by local superconformal coordinates  $(z|\theta)$

---

<sup>21</sup>In general,  $D_\eta^2 = (w - w_0)^n \frac{\partial}{\partial w}$  for  $n \geq 1$  near a puncture. Such a surface is called a super-Riemann surface with level- $n$  parabolic structure. The case of  $n = 1$  corresponds to the superconformal structure in the presence of Ramond puncture.

<sup>22</sup>For another explanation see section 4.2.2 of [11].

and  $(w|\eta)$ . The punctures are located at  $(z_0|\theta_0)$  and  $(w_0|\eta_0)$ . In terms of these coordinates, the gluing relations can be described as follows

- **Gluing of two NS Punctures:** To describe the gluing relation, we define the following coordinates

$$\begin{aligned} x_1 &\equiv z - z_0 - \theta\theta_0, & \widehat{x}_1 &\equiv \theta - \theta_0, \\ x_2 &\equiv w - w_0 - \eta\eta_0, & \widehat{x}_2 &\equiv \eta - \eta_0. \end{aligned} \quad (2.16)$$

Two NS punctures can then be glued via the following superconformal mapping [33]

$$\begin{aligned} x_1 x_2 + t^2 &= 0, \\ x_1 \widehat{x}_2 - t \widehat{x}_1 &= 0, \\ x_2 \widehat{x}_1 + t \widehat{x}_2 &= 0, \\ \widehat{x}_1 \widehat{x}_2 &= 0. \end{aligned} \quad (2.17)$$

$t$  is a complex parameter. For non-zero  $t$ , the two middle equations are equivalent which can be seen by multiplying the second equation by  $x_2$  (or the third equation by  $x_1$ ) and using the first equation. However, these equations are independent for  $t = 0$ . One usually is interested in the case that the punctures are located at  $(z_0|\theta_0) = (0|0) = (w_0|\eta_0)$  in the respective local coordinates. In this case, the gluing relations can be simply written as

$$\begin{aligned} zw + t^2 &= 0, \\ z\eta - t\theta &= 0, \\ w\theta + t\eta &= 0, \\ \theta\eta &= 0. \end{aligned} \quad (2.18)$$

To connect with the usual notation in the literature, one defines  $q_{\text{NS}} \equiv -t^2$ . These relations reduce to the plumbing of punctures on ordinary Riemann surfaces by setting  $\theta = \eta = 0$  and  $q \equiv q_{\text{NS}}$ . The one-parameter family of glued surfaces are  $\mathcal{N} = 1$  super-Riemann surfaces parametrized by the even and odd moduli of the component surface(s) and the extra complex parameter  $t$ . It is easy to check that the resulting surfaces have correct number of even and odd moduli.

- **Gluing of two R Punctures:** Two R punctures can be glued using the following relations [11]

$$\begin{aligned} (z - z_0)(w - w_0) + t^2 &= 0, \\ (\theta - \theta_0) + \alpha \mp i(\eta - \eta_0) &= 0, \end{aligned} \quad (2.19)$$

where  $\alpha$  is an odd gluing parameter. The origin of this parameter is an extra symmetry  $\theta \longrightarrow \theta + \alpha$  of the divisor defined by an R puncture. We will explain the idea behind this

extra odd symmetry in section 3.1.2. Putting the punctures at  $(z_0|\theta_0) = (0|0) = (w_0|\eta_0)$  in the respective local coordinates, these relations become

$$\begin{aligned} zw + t^2 &= 0, \\ \theta + \alpha \mp i\eta &= 0. \end{aligned} \tag{2.20}$$

One can connect these to the notation in the literature by defining  $q_R \equiv -t^2$ . Again, these relations reduce to the plumbing of punctures on ordinary Riemann surfaces by setting  $\theta = \eta = 0$  and  $q \equiv q_R$ . The two-parameter family of glued surfaces are  $\mathcal{N} = 1$  super-Riemann surfaces parametrized by the even and odd moduli of the component surface(s) and the extra complex parameters  $t$  and  $\alpha$ <sup>23</sup>. It is easy to check that the resulting surfaces have correct number of even and odd moduli.

**Remark:** The gluing of two R punctures can also be described as follows [33]. We first define the following coordinates

$$y_1^2 \equiv -(z - z_0), \quad y_2^2 \equiv +(w - w_0). \tag{2.21}$$

Two R punctures can then be glued via the following superconformal mapping

$$\begin{aligned} \left(y_1 \mp \frac{1}{4}\theta\hat{t}\right) \left(y_2 - \frac{1}{4}\eta\hat{t}\right) - t &= 0, \\ \mp\theta y_2 + \eta y_1 + t\hat{t} &= 0. \end{aligned} \tag{2.22}$$

$\hat{t}$  is an odd gluing parameter. In this paper, we only use the gluing relation (2.20) for the gluing of two R punctures.

The gluing relations of both NS punctures (2.18) and R punctures (2.20) are two-valued. This is related to the imposition of GSO projection by summing over different spin structures. For more explanation see section 6.2.3 of [11].

We were interested in the gluing of two R punctures on spin-Riemann surfaces. *How do the gluing relations change when we project super-Riemann surfaces to their underlying spin curves?* The gluing relations of NS punctures remains the same. However, the gluing relations of R punctures will change a bit. The reason is that the symmetry of the divisor defined by an R punctures is lost by projecting to the underlying spin curve, i.e. there is no parameter

---

<sup>23</sup>As we mentioned, the odd gluing parameter  $\alpha$  enters the gluing relations due to an extra symmetry of the divisors defined by the glued R punctures, as we discuss in section 3.2. However, there are special cases that  $\alpha$  cannot be considered as a modulus. Essentially, these are the cases that the component surfaces have superconformal automorphisms that acts on the divisors associated to the glued punctures by  $\theta \rightarrow \theta + \alpha$  (or  $\eta \rightarrow \eta + \alpha$ ). These automorphisms can then be used to remove the extra symmetry of the R punctures. In such cases, there is no odd gluing parameter in the gluing relations of R punctures and can be studied separately.

$\alpha$ . Therefore, we use the following gluing relations of two R punctures on an spin-Riemann surfaces become

$$\begin{aligned} zw + t^2 &= 0, \\ \theta \pm i\eta &= 0. \end{aligned} \tag{2.23}$$

As usual, the gluing of even coordinates is the usual plumbing fixture of ordinary Riemann surfaces. The gluing of odd coordinates is interpreted as the gluing of the corresponding spin bundles which induces the spin bundle, i.e. the choice of spin structure, on the resulting surface. For a mathematically-precise description of gluing of punctures on spin curves see Section B.9.

### 3 Nodal Spin Curves and Torsion-Free Sheaves

In this section, we consider the gluing of two punctures of the same kind on a single or two disconnected spin curves using plumbing prescription given in section 2.3. We describe how the gluing provides a proper compactification of the moduli stack of smooth spin curves. This means that starting from the component surface(s)<sup>24</sup> equipped with spin structures, one can construct a family of degenerating surfaces equipped with spin structures such that 1) spin structures are induced from the spin structures of the component surface(s), 2) their moduli can be described in terms of the moduli of component surface(s), and 3) they properly compactify the moduli stack of irreducible spin curves as constructed in [39, 40, 44, 45].

#### 3.1 Degenerations of Spin Curves

The plumbing fixture construction of a family of degenerating spin curves can be applied to either a single surface or two (and more) surfaces. To be concrete, we consider only the gluing of two surfaces. All considerations are applicable for the other types of gluing.

Consider two split surfaces  $\mathcal{R}(\mathbf{m}, \mathbf{0})$  and  $\mathcal{R}'(\mathbf{m}', \mathbf{0})$ , where  $\mathbf{m}$  and  $\mathbf{m}'$  denote the even moduli of the surfaces.  $\mathcal{R}$  has signature  $(g; \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  and is equipped with a spin structure  $\mathcal{E}$  and  $\mathcal{R}'$  has signature  $(g'; \mathbf{n}'_{\text{NS}}, \mathbf{n}'_{\text{R}})$  and is equipped with a spin structure  $\mathcal{E}'$ . To glue punctures, we consider the local coordinate around the glued punctures on  $\mathcal{R}$  and  $\mathcal{R}'$  to be  $z$  and  $z'$ , respectively. Since the surfaces are equipped with spin structure, we also consider the coordinates in the fiber of the respective spinor bundles to be  $\theta$  and  $\theta'$ . These coordinate define the following spinor derivatives acting on the respective spinor bundles

$$D_\theta = \frac{\partial}{\partial \theta} + \theta \frac{\partial}{\partial z}, \quad D_{\theta'} = \frac{\partial}{\partial \theta'} + \theta' \frac{\partial}{\partial z'}. \tag{3.1}$$

We would like to understand what is the effect of plumbing fixture on the spinor derivative. These will tell us the spinor bundle on the resulting family of surfaces parametrized by  $\mathbf{m}$ ,  $\mathbf{m}'$ , and the gluing parameters.

---

<sup>24</sup>We write surface(s) to include the situation that the glued punctures are located on a single surface.



### 3.1.1 Neveu-Schwarz Degenerations

Consider the NS punctures  $\mathfrak{p}$  and  $\mathfrak{p}'$  on  $\mathcal{R}$  and  $\mathcal{R}'$ , and the gluing relation (2.18) for the NS puncture. For  $t \neq 0$ , the independent equations are

$$\begin{aligned} zz' + t^2 &= 0, \\ z\theta' - t\theta &= 0. \end{aligned} \tag{3.2}$$

By putting the first equation into the second one, we get  $z'\theta + t\theta' = 0$ , which is the third equation of (2.18). Using these relations, we have

$$\begin{aligned} \frac{\partial}{\partial \theta} &= \frac{\partial z'}{\partial \theta} \frac{\partial}{\partial z'} + \frac{\partial \theta'}{\partial \theta} \frac{\partial}{\partial \theta'} = -\frac{z'}{t} \frac{\partial}{\partial \theta'}, \\ \theta \frac{\partial}{\partial z} &= -\frac{t\theta'}{z'} \left( \frac{\partial z'}{\partial z} \frac{\partial}{\partial z'} + \frac{\partial \theta'}{\partial z} \frac{\partial}{\partial \theta'} \right) = -\frac{t\theta'}{z'} \left( \frac{z'^2}{t^2} \frac{\partial}{\partial z'} + \frac{z'\theta'}{t^2} \frac{\partial}{\partial \theta'} \right) = -\frac{z'\theta'}{t} \frac{\partial}{\partial z'}. \end{aligned} \tag{3.3}$$

Note that we have used the fact that  $\frac{\partial z'}{\partial \theta} = 0$ . This is because we use one of the last two equations of (2.18), and as such,  $z'$  is independent of  $\theta$ . Thus, the spinor derivative  $D_\theta$  transforms as


$$D_\theta = -\frac{z'}{t} D_{\theta'}. \tag{3.4}$$

It is clear from this form that the spinor derivative blows up as  $t \rightarrow 0$ . What does this blow-up mean in terms of the compactification of the moduli stack of spin curves as described in appendix B? When  $t \neq 0$ , the spin bundles  $\mathcal{E}$  and  $\mathcal{E}'$  are glued together in a natural way to form a spin bundle defined over the whole glued surface. However, when  $t = 0$ , the second and third equations of (2.18) become independent, and (3.4) does not make sense. We thus have the following gluing relations at  $t = 0$

$$zz' = 0, \quad z\theta' = 0, \quad z'\theta = 0. \tag{3.5}$$

There are two possible cases


1.  $z = 0$  or  $z' = 0$ : since the equations (3.5) are symmetric, let us assume that  $z' = 0$ . The third equation of (3.5) is trivially satisfied. Therefore, the odd parameter  $\theta$  is left as a free parameter. The corresponding spinor derivative  $D_\theta$  generates a rank-1 bundle with odd fibers, i.e. a spin structure. Therefore, a spinor bundle is defined over the whole glued surface.
2.  $z = z' = 0$ : This case is when, by definition, a node is formed, see figure 1. In this case, all the equations are trivially satisfied. We thus left with two odd parameters  $\theta$  and  $\theta'$  whose corresponding spinor derivative  $D_\theta$  and  $D_{\theta'}$  are independent and generate a two-dimensional vector space over the node. Therefore, formation of an NS degeneration is characterized by formation of a 2-dimensional vector space over the node. In this sense, the NS degenerations do not appear naturally in the compactification of the moduli stack of spin curves.




**Figure 1:** In the gluing of two NS punctures, a node  $\mathfrak{n}_{\text{NS}}$  is formed when  $z = z' = 0$ .

Therefore the question is that: *what is the proper compactification of the moduli stack of spin curves in which NS degenerations appear naturally?* As has been explained in Appendix B, the divisor points of a proper compactification of the moduli stack of spin curves consist of triples  $(\widehat{\mathcal{R}}, \mathcal{E}, \psi)$ .  $\widehat{\mathcal{R}}$  is a semi-stable curve with copies of  $\mathbb{P}^1$  intersecting the rest of the curve in at most two points<sup>25</sup>, and  $\mathcal{E}$  is a line bundle which has degree one on each  $\mathbb{P}^1$ . The contraction of all  $\mathbb{P}^1$  described by the blowup map  $\rho : \widehat{\mathcal{R}} \rightarrow \mathcal{R}^\bullet$  turns  $\widehat{\mathcal{R}}$  to a stable curve  $\mathcal{R}^\bullet$ , see figure 2. The pushforward  $\rho_* \widehat{\mathcal{E}}$  is a rank-one torsion-free sheaf.  $\psi : (\rho_* \widehat{\mathcal{E}})^{\otimes 2} \rightarrow \omega_{\mathcal{R}^\bullet}$ , where  $\omega_{\mathcal{R}^\bullet}$  is the canonical sheaf of  $\mathcal{R}^\bullet$ , is a homomorphism between the tensor product of  $\rho_* \widehat{\mathcal{E}}$  and the canonical bundle over  $\mathcal{R}^\bullet$  with the following properties, 1) it is an isomorphism where  $\rho_* \widehat{\mathcal{E}}$  is locally-free, and 2) its cokernel, i.e.  $\omega_{\mathcal{R}^\bullet} / \text{Im} \psi$ , has length one where  $\rho_* \widehat{\mathcal{E}}$  is failed to be locally-free, i.e. the singular points of  $\mathcal{R}^\bullet$ .

In the case of an NS node, we thus need to consider the blowup geometry. This means that we turn an NS node to a  $\mathbb{P}^1$  over which a line bundle  $\widehat{\mathcal{E}}$  lives. The pushforward of this line bundle by the blowup map  $\mathfrak{b} : \mathcal{R}^{\text{b}} \rightarrow \mathcal{R}^\bullet$ , where  $\mathcal{R}^{\text{b}}$  denotes the blow-up geometry, is a rank-one torsion-free sheaf over the surface with node which, according to [40, 44, 45], define a spin structure over the nodal curve.



(a) The formation of an NS node.



(b) The blowup geometry of an NS node.

**Figure 2:** The figure (a) shows the formation of an NS node when  $z = z' = 0$ .  $\theta$  and  $\theta'$  are two free parameters whose corresponding spinor derivatives  $D_\theta$  and  $D_{\theta'}$  generate a rank-2 vector space over the node. The figure (b) shows the blowup geometry where a  $\mathbb{P}^1$  intersect the rest of the curve in two points.

### 3.1.2 Ramond Degenerations

Consider R punctures  $\mathfrak{q}$  and  $\mathfrak{q}'$  on  $\mathcal{R}$  and  $\mathcal{R}'$ , and the gluing relation (2.23) for R punctures

$$\begin{aligned} zz' + t^2 &= 0, \\ \theta \mp i\theta' &= 0. \end{aligned} \tag{3.6}$$

<sup>25</sup>Such irreducible components of a semi-stable curve are called *exceptional curves*.

Let us repeat the same exercise as above. We have

$$\begin{aligned}\frac{\partial}{\partial\theta} &= \mp i \frac{\partial}{\partial\theta'}, \\ \theta z \frac{\partial}{\partial z} &= \pm i \theta' \left(-\frac{t^2}{z'}\right) \left(\frac{z'^2}{t^2}\right) \frac{\partial}{\partial z'} = \mp i \theta' z' \frac{\partial}{\partial z'}.\end{aligned}\tag{3.7}$$

We thus have

$$D_\theta = \mp i D_{\theta'}.\tag{3.8}$$

The relation shows that the bundles generated by  $D_\theta$  and  $D_{\theta'}$  are isomorphic. Therefore, irrespective of the values of  $z$  and  $z'$ , there is a rank-1 bundle over the glued curve which equips the surface with spin structure. Unlike the NS-type degenerations, there is no need to consider the blowup geometry of the surface with node. This fact shows that R-type degenerations appear in the natural compactification of the moduli stack of spin curves.

We can thus provide the answer to the [Question 1](#) posed in section 1.

In the gluing of NS punctures, there are two separate situation. In the moduli stack of spin curves and away from the compactification divisor, the spin structure of the component surfaces are glued together using the gluing relations (3.2) for non-zero  $t$  and (3.5) for  $t = 0$ . The spin bundle over the gluing tube is generated by either  $D_\theta$  or  $D_{\theta'}$ , related by (3.4). However, (3.5) also shows that exactly at the node, i.e.  $z = z' = 0$ , there are free parameters  $\theta$  and  $\theta'$  whose corresponding spinor derivatives generate a rank-2 vector space over the node which cannot be naturally thought of as a spin bundle. The resolution is to consider the blowup geometry of the node on which a rank-one bundle is defined. The pushforward of this bundle under the blowup map is a rank-1 torsion-free sheaf over the node. This procedure provides a proper compactification of the moduli stack of spin curves [39].

In the gluing of R punctures, there is no ambiguity and the gluing relations gives a rank-one bundle over the glued surface. As we mentioned above, the gluing of R punctures on spin curves rather than the parent  $\mathcal{N} = 1$  super-Riemann surface is accompanied by a missing odd parameter. This has a consequence when one defines the superstring amplitudes in terms of spin curves and the corresponding moduli stack of such surfaces. We turn to this consequence in the next section.

### 3.2 PCOs and Gluing of Ramond Punctures on Spin Curves

In this section, we consider the gluing of two R punctures on the spin curves  $\mathcal{R}$  and  $\mathcal{R}'$  equipped with spin structures  $\mathcal{E}$  and  $\mathcal{E}'$ . Equation (3.8) shows that the parameter  $\alpha$  is not visible from the point of view of the spin curves. The question is then *what is the consequence of losing the symmetry whose parameter is  $\alpha$  by studying the gluing on the underlying spin curve(s) rather than the parent  $\mathcal{N} = 1$  super-Riemann surfaces?* To answer this question, we first need to understand the answer to the following question.

*What is the role of parameter  $\alpha$  from the point of view of the gluing of super-Riemann surfaces?* As we explained in section 2.2, the location of an R puncture defines a divisor on

a super-Riemann surface. A generic even superconformal vector field which preserves the superconformal structure in the presence of an R puncture has the following form<sup>26</sup>

$$v_{\mathfrak{e}} = F(z) \left( f(z) \frac{\partial}{\partial z} + \frac{f'(z)}{2} \theta \frac{\partial}{\partial \theta} \right), \quad (3.9)$$

where  $f(z)$  is some holomorphic function of  $z$  and  $F(z)$  vanishes along the divisor defined by the R puncture. It is thus clear that this vector field vanishes along that divisor. However, a generic odd superconformal vector field which also preserves the superconformal structure in the presence of an R puncture has the following form

$$v_{\mathfrak{o}} = g(z) \left( \frac{\partial}{\partial \theta} - F(z) \theta \frac{\partial}{\partial z} \right), \quad (3.10)$$

where  $g(z)$  is some holomorphic function of  $z$ . This vector field is in general non-vanishing along the divisor defined by the R puncture and is proportional to  $\partial_{\theta}$ . Such a vector field generates the transformation  $\theta \longrightarrow \theta + \alpha$  which is a symmetry of the divisor. This means that when we are gluing two R punctures, the gluing is not unique (the odd coordinate of each of the punctures is defined up to addition by an odd parameter) and the gluing relation is defined up to addition by an odd parameter  $\alpha$ . Therefore, the divisors in the superstack defined by an R-type degeneration are fiber bundles over  $\mathcal{SM}_{g-1; \mathfrak{n}_{\text{NS}} \mathfrak{n}_{\text{R}}+2}$  (for nonseparating R-type degenerations where a single genus- $g$  surface with  $\mathfrak{n}_{\text{NS}}$  NS puncture and  $\mathfrak{n}_{\text{R}}$  R punctures degenerates into a surface whose normalization has signature  $(g-1, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}+2)$ ) and  $\mathcal{SM}_{g_1, \mathfrak{n}_{\text{NS}}^1, \mathfrak{n}_{\text{R}}^1} \cup_{\text{R}} \mathcal{SM}_{g_2, \mathfrak{n}_{\text{NS}}^2, \mathfrak{n}_{\text{R}}^2}$  (for separating R-type degenerations where a genus  $g = g_1 + g_2$  surface with  $\mathfrak{n}_{\text{NS}} = \mathfrak{n}_{\text{NS}}^1 + \mathfrak{n}_{\text{NS}}^2$  NS punctures and  $\mathfrak{n}_{\text{R}} = \mathfrak{n}_{\text{R}}^1 + \mathfrak{n}_{\text{R}}^2 - 2$  R punctures degenerates into two surfaces with signatures  $(g_1; \mathfrak{n}_{\text{NS}}^1, \mathfrak{n}_{\text{R}}^1)$  and  $(g_2; \mathfrak{n}_{\text{NS}}^2, \mathfrak{n}_{\text{R}}^2)$ ). The fibers of these fibration are isomorphic to  $\mathbb{C}^{0|1}$  parameterized by an odd parameter. On the other hand, the integration over  $\alpha$ , which together with the other even and odd parameters provide a coordinate chart near the divisor of  $\mathcal{SM}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ , is necessary to produce the correct propagator in the Ramond sector [11, 46].

If one starts from the supergeometry definition of the superstring amplitudes, and tries to reduce the computation to the underlying moduli stack of spin curves, one has to integrate over the odd moduli of the superstack. As we explained in section A.2, the effect of integration over complex odd moduli of superstack is the introduction of PCOs to the string measure on the moduli stack of spin curves. Since  $\alpha$  is one of the odd moduli of the resulting surface, it has to be integrated over. The parameter  $\alpha$  enters the gluing relations of two R punctures due to a freedom in the gluing and it is not associated to intrinsic odd moduli of the component surfaces. Therefore, the resulting PCO has to be added on the gluing tube using some prescription that produces the correct results. However, if one uses the picture-changing formalism as a basic definition to define the string amplitudes, the parameter  $\alpha$  does not enter the computation since the symmetry of the R puncture does not exist anymore. The

---

<sup>26</sup>For more details see section 4.2 of [11].

effect of losing this symmetry by reducing everything on the underlying spin curve is as follows. If we integrate all of the odd moduli, there will be the  $2g - 2 + n_{\text{NS}} + \frac{1}{2}n_{\text{R}}$  PCOs in the string measure. Let us glue two R punctures on spin curves with signatures  $(g_1; n_{\text{NS}}^1, n_{\text{R}}^1)$  and  $(g_2; n_{\text{NS}}^2, n_{\text{R}}^2)$ . A simple counting shows that the number of PCOs on  $\mathcal{R} \cup_{\text{R}} \mathcal{R}'$ , the resulting glued surface with signature  $(g_1 + g_2; n_{\text{NS}}^1 + n_{\text{NS}}^2, n_{\text{R}}^1 + n_{\text{R}}^2 - 2)$ , is

$$\# \text{ of PCOs on } \mathcal{R} \cup_{\text{R}} \mathcal{R}' = 2(g_1 + g_2) - 2 + (n_{\text{NS}}^1 + n_{\text{NS}}^2) + \frac{n_{\text{R}}^1 + n_{\text{R}}^2 - 2}{2} - 1. \quad (3.11)$$

This is one less than the number it should be. On the other hand, to deal with IR divergences, it is required that the off-shell amplitudes to be defined in such a way that the choice of local coordinates around the punctures and the choice of PCOs be gluing-compatible on the separating-type degenerations [47, 48]. For PCOs, this essentially means that the choice of PCOs on the glued surface must be induced from the choice of PCOs on the component surfaces. However, (3.11) shows that the choice of PCOs with picture number  $-\frac{1}{2}$  is not gluing-compatible, i.e. the number of PCOs on the resulting surface is not equal to the sum of the number of PCOs on the component surfaces. The resolution is as follows, 1) for computing the two-point function, we choose one of the R states that are being glued to have picture number  $-\frac{3}{2}$ , and 2) for other situations, we introduce a wighted average of PCOs inserted on a cycle on the tube joining the two glued R punctures. For more details, see sections 6.2 and 6.3 of [17]. In this way, the number of PCOs on the glued surface match with the one that is induced from the component surfaces.

We can thus provide the answer to the [Question 2](#) posed in section 1.

On an  $\mathcal{N} = 1$  super-Riemann surface with R punctures, each of the R punctures defines a divisor with a symmetry  $\theta \rightarrow \theta + \alpha$ . This symmetry is lost by projecting onto the underlying spin curve. Therefore, there is a missing odd parameter. On the other hand, the number of PCOs is equal to the fermionic dimension of the corresponding superstack. As such, there is a missing PCO when we glue two R punctures on spin curves. We need to compensate the missing PCO by adding an extra one on the gluing tube as prescribed in [17, 49].

## 4 Existence and Uniqueness of Solution to the BV QME in $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$

In this section, we discuss on of the main results of this paper, i.e. 1) the existence and uniqueness of a solution to the BV QME in  $\mathcal{S}(n_{\text{NS}}, n_{\text{R}})$ , and 2) the relation of these solutions to the fundamental class of DM stack of corresponding spin curves. For our main purposes, we introduce an alternative model for the moduli space of bordered spin-Riemann surfaces. This model is a natural generalization of the model for the moduli space of bordered ordinary Riemann surfaces introduced in [50]. After setting the stage, we prove the existence and uniqueness of a solution to the BV quantum master equation, and find that it can be mapped to the fundamental classes of DM spaces.

#### 4.1 The Strategy of Proof

Since the procedure for proving the results of this section is long, we provide a summary of the strategy of proof in this subsection. We first describe the problems in the context of bordered ordinary Riemann surfaces, and the results obtained in [26]. We then briefly explain the generalization of these results to the case of bordered spin-Riemann surfaces considered in this paper.

Let us first define the Neveu-Schwarz and Ramond boundary components. For this purpose, we define a more general notion of a superconformal cycle on an  $\mathcal{N} = 1$  super-Riemann surface [51].

**Definition 4.1 (Neveu-Schwarz Cycles).** *A cycle<sup>27</sup> on an  $\mathcal{N} = 1$  super-Riemann surface is called a Neveu-Schwarz cycle if it is isomorphic to  $S^{1|1}$ . If we denote the coordinates along  $S^{1|1}$  by  $(x|\theta)$ , then they are identified as  $(x + 2\pi|\theta) \sim (x|\theta)$  on an NS cycle.*

This definition shows that on an spin curve, a cycle with antiperiodic boundary condition is an NS cycle. Similarly, we can define NS boundary components

**Definition 4.2 (Neveu-Schwarz Boundary Component).** *An NS boundary component of an  $\mathcal{N} = 1$  super-Riemann surface is a cs sub supermanifolds isomorphic to  $S^{1|1} \otimes_{\mathbb{R}} \mathbb{C}$  of dimension<sup>28</sup>  $(1|1)$ .*

**Definition 4.3 (Ramond Cycles).** *A cycle on an  $\mathcal{N} = 1$  super-Riemann surface is called a Ramond cycle if it is isomorphic to  $S^1 \times \mathbb{R}^{0|1}$ . If we denote the coordinates along  $S^1 \times \mathbb{R}^{0|1}$  by  $(x|\theta)$ , then they are identified as  $(x + 2\pi|\theta) \sim (x|\theta)$  on an R cycle.*

This definition shows that on an spin curve, a cycle with periodic boundary condition is an R cycle. Similarly, we can define R boundary components

**Definition 4.4 (Ramond Boundary Component).** *An R boundary component of an  $\mathcal{N} = 1$  super-Riemann surface is a cs sub supermanifolds isomorphic to  $S^1 \times \mathbb{R}^{0|1} \otimes_{\mathbb{R}} \mathbb{C}$  of dimension  $(1|1)$ .*

One of the main result of [26] is the proof of existence and uniqueness, in the sense we explain below, of the solution of the BV quantum master equation for the BV algebra associated to the moduli space of bordered ordinary Riemann surfaces. From the work of Sen and Zwiebach [52, 53], this is the problem of existence and uniqueness of the closed bosonic-string vertices. Let  $\mathcal{C}_*$  be the functor of normalized singular simplicial chains with coefficients in any field  $\mathbb{F}$  containing  $\mathbb{Q}$ , and  $\mathcal{M}(\mathfrak{n})$  be the moduli space of Riemann surfaces with  $\mathfrak{n}$  boundaries. Associated to these spaces, we can define the following complex

$$\mathcal{F}(\mathcal{M}) \equiv \bigoplus_{\mathfrak{n}} \mathcal{C}_* \left( \frac{\mathcal{M}(\mathfrak{n})}{S^1 \wr S_{\mathfrak{n}}} \right), \quad (4.1)$$

<sup>27</sup>By *cycle* we mean a closed  $C^\infty$  sub-supermanifold of real codimension  $(1|1)$ .

<sup>28</sup>By dimension  $(m|n)$  of a cs supermanifold, we mean even real dimension  $m$  and odd complex dimension  $n$ .

where  $S^1$  is a circle action,  $S_n$  denotes the permutation group, and  $S^1 \wr S_n \equiv (S^1)^n \ltimes S_n$  is the wreath product group. The action of wreath product on  $\mathcal{M}(n)$  turns it into the moduli space of Riemann surfaces with unlabeled boundaries.  $\mathcal{F}(\mathcal{M})$  has the structure of a BV algebra by considering  $d$  to be the boundary operator acting on chains and  $\delta$  to be  $\Delta$ , the operation of sewing two boundaries on the same surface [26, 53]. It is clear that there is an  $S^1$ -worth possible ways of gluing. This is called the Sen-Zwiebach BV algebra [53]. We denote the part of  $\mathcal{F}(\mathcal{M})$  coming from genus- $g$  surfaces with  $n$  boundaries as  $\mathcal{F}_{g,n}(\mathcal{M})$ . To prove the existence and uniqueness of a solution to the BV QME, Costello considered a model for  $\mathcal{M}(n)$  introduced by Kimura, Stasheff, and Voronov [50]. We denote these spaces as  $\tilde{\mathcal{M}}(n)$ <sup>29</sup>. This is the moduli space of Riemann surfaces in  $\overline{\mathcal{M}}_n$ , the stable-curve compactification of the moduli stack of Riemann surfaces with  $n$  punctures, decorated at each puncture with a ray in the tangent space and decorated at each node by a ray in the tensor product of tangent spaces in the two sides.  $\tilde{\mathcal{M}}(n)$  is homotopy-equivalent to  $\mathcal{M}(n)$ . The reason is clear: 1) each puncture is equipped with a ray, and the phase of this ray parametrizes a circle which can be thought of as a boundary, and 2) each node is equipped with a ray which can be used to open the node. Similar to  $\mathcal{M}(n)$ , we can associate a complex to  $\tilde{\mathcal{M}}(n)$  as follows

$$\mathcal{F}(\tilde{\mathcal{M}}) \equiv \bigoplus_n \mathcal{C}_* \left( \frac{\tilde{\mathcal{M}}(n)}{S^1 \wr S_n} \right). \quad (4.2)$$

The quotient by the wreath product turns  $\tilde{\mathcal{M}}(n)/S^1 \wr S_n$  to the space of Riemann surfaces possibly with nodes, unordered marked points (without any ray in the tangent space), unparametrized boundary, but with a ray in the tensor product of tangent spaces at each side for each node. If  $d$  is the boundary operation, and  $\Delta$  is the operation of taking two marked points and gluing them (we give a more precise definition in Section 4.4),  $\mathcal{F}(\tilde{\mathcal{M}})$  turns into a BV algebra. It is clear that the gluing can be done in  $S^1$  possible ways. For brevity, let us define

$$\mathbb{X}(n) \equiv \frac{\tilde{\mathcal{M}}(n)}{S^1 \wr S_n}, \quad (4.3)$$

and denote its fundamental chain by  $[\mathbb{X}(n)]$ . After passing to homology, these classes satisfy [26]

$$d[\mathbb{X}(n)] + \Delta[\mathbb{X}(n+2)] = 0. \quad (4.4)$$

If  $\mathbb{X}_g(n)$  denotes the subspace of connected genus- $g$  surfaces, and  $[\mathbb{X}_g(n)]$ , after passing to homology, its fundamental class, we can define the following formal sum

$$[\mathbb{X}] \equiv \sum_{\substack{g,n=0 \\ 2g-2+n>0}}^{\infty} \hbar^{2g-2+n} [\mathbb{X}_g(n)]. \quad (4.5)$$

---

<sup>29</sup>The model in [50] is really for the space of surfaces all whose connected components have negative Euler number.

Using this and (4.4), we have

$$(d + \hbar\Delta) \exp\left(\frac{[\mathbb{X}]}{\hbar}\right) = 0, \quad (4.6)$$

i.e.  $[\mathbb{X}]$  satisfies the BV quantum master equation. A more transparent way of writing this equation is

$$d[\mathbb{X}_g(\mathbf{n})] + \Delta[\mathbb{X}_{g-1}(\mathbf{n} + 2)] + \frac{1}{2} \sum_{\substack{g_1 + g_2 = g \\ \mathbf{n}_1 + \mathbf{n}_2 = \mathbf{n} - 2}} \{[\mathbb{X}_{g_1}(\mathbf{n}_1)], [\mathbb{X}_{g_2}(\mathbf{n}_2)]\} = 0. \quad (4.7)$$

Using this result, we have (Proposition 10.1.1<sup>30</sup> of [26])

**Proposition (The Existence and Uniqueness of Closed Bosonic-String Vertices).** *For any  $g$  and  $\mathbf{n}$  such that  $2g - 2 + \mathbf{n} > 0$ , there exist elements  $\mathcal{V}_g(\mathbf{n}) \in \mathcal{F}_{g,\mathbf{n}}(\mathcal{M})$ , i.e. which can be thought of as genus- $g$  bosonic-string vertices with  $\mathbf{n}$  punctures, of homological degree  $6g - 6 + 2\mathbf{n}$  such that*

1.  $\mathcal{V}_0(3)$  is the fundamental cycle of  $\mathcal{M}_0(3)/S_3$ , i.e. a 0-chain of degree  $\frac{1}{3!}$ .
2. Consider the following formal power series

$$\mathcal{V} \equiv \sum_{\substack{g, \mathbf{n}=0 \\ 2g-2+\mathbf{n}>0}}^{\infty} \hbar^{2g-2+\mathbf{n}} \mathcal{V}_g(\mathbf{n}) \in \hbar\mathcal{F}(\mathcal{M})[[\hbar]]. \quad (4.8)$$

$\mathcal{V}$  satisfies the BV quantum master equation

$$(d + \hbar\Delta) \exp\left(\frac{\mathcal{V}}{\hbar}\right) = 0.$$

3.  $\mathcal{V}$  is unique up to homotopy in the category of BV algebras.

At this point, one might ask why do we need to use  $\tilde{\mathcal{M}}(\mathbf{n})$  and its associated complex rather than  $\mathcal{M}(\mathbf{n})$  and its associated complex? The reason is that the above proposition is proven for  $\mathcal{F}(\tilde{\mathcal{M}})$ . However, the BV algebras  $\mathcal{F}(\mathcal{M})$  and  $\mathcal{F}(\tilde{\mathcal{M}})$  are quasi-isomorphic (see Lemma 10.4.2 of [26]) which means that their corresponding homology groups are isomorphic [26]. On the other hand, if two differential-graded algebras are quasi-isomorphic, their sets of homotopy classes of solutions of the Maurer-Cartan equation are isomorphic [26]. The BV quantum master equation can be considered as the Maurer-Cartan equation of the corresponding BV algebra (see Lemma 5.2.1 of [26]). Therefore, if two BV algebras are quasi-isomorphic, their sets of homotopy classes of solutions of the BV quantum master equation are isomorphic (see Lemma 5.3.1 and Definition 5.4.1 of [26]). We can then say that once the existence and

---

<sup>30</sup>The numbers of theorem, proposition, etc of [26] that we are referring to are the ones that appear in the published version not the Arxiv preprint.



uniqueness of the solution of the BV quantum master equation in the BV algebra  $\mathcal{F}(\tilde{\mathcal{M}})$  is proven, we can use the quasi-isomorphism of  $\mathcal{F}(\tilde{\mathcal{M}})$  and  $\mathcal{F}(\mathcal{M})$ , and the above considerations, to conclude the existence and uniqueness of the solution of the BV quantum master equation, which we interpret as closed bosonic-string vertices, in the BV algebra  $\mathcal{F}(\mathcal{M})$ .

Let us now describe the analogous result for the case of bordered spin-Riemann surfaces which has been proven in Section 4.5. Analogous to  $\tilde{\mathcal{M}}(\mathbf{n})$  which is a model for  $\mathcal{M}(\mathbf{n})$ , we introduce the space  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  which is a nice model for  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . It is the moduli space of stable spin curves, decorated at each NS puncture with a ray in the tangent space, at each R puncture with a ray in the spinor bundle, at each NS node with a ray in the tensor product of rays of the tangent spaces at each side, and at each R node with a ray in the spinor line bundle.  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is again homotopy-equivalent to  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . The reason is similar to the one explained in the case of  $\mathcal{F}(\mathcal{M})$  and  $\mathcal{F}(\tilde{\mathcal{M}})$ . Associate to these spaces, and completely analogous to  $\mathcal{F}(\mathcal{M})$  and  $\mathcal{F}(\tilde{\mathcal{M}})$ , we define two complexes

$$\begin{aligned}\mathcal{F}(\mathcal{S}) &\equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{C}_* \left( \frac{\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})}{S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}} \right), \\ \mathcal{F}(\tilde{\mathcal{S}}) &\equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{C}_* \left( \frac{\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})}{S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}} \right).\end{aligned}\tag{4.9}$$

$S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  is the wreath product  $(S^1)^{\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}} \rtimes S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ .  $S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  is the group that permutes NS boundaries and R boundaries separately among each other. The quotient by the wreath product turns  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  to the space of bordered spin-Riemann surfaces possibly with nodes, unordered marked points (without any ray in either the tangent space or the spinor bundle), unordered boundary, but with a ray in the tensor product of tangent spaces at each side for each NS node and a ray in the spinor bundle at each R node. Let  $\mathcal{F}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}(\mathcal{S})$  be the part coming from connected surfaces of genus  $g$  with  $\mathbf{n}_{\text{NS}}$  NS boundaries and  $\mathbf{n}_{\text{R}}$  R boundaries.  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\tilde{\mathcal{S}})$  naturally carry structures of commutative differential graded algebra, where differential is the boundary map of chain complexes and the product comes from disjoint union of surfaces. Since  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is homotopy-equivalent to  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ ,  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\tilde{\mathcal{S}})$  are quasi-isomorphic. Furthermore, we define the following gluing operation<sup>31</sup> acting on surfaces in  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$

$$\begin{aligned}\Delta &\equiv \Delta^{\text{NS}} + \Delta^{\text{R}+} + \Delta^{\text{R}-}, \\ \{\cdot, \cdot\} &\equiv \{\cdot, \cdot\}_{\text{NS}} + \{\cdot, \cdot\}_{\text{R}}.\end{aligned}$$

$\Delta^{\text{NS}}$  is the operation of gluing two NS punctures on a surface, and  $\Delta^{\text{R}\pm}$  is the operation of gluing of two R punctures on a surface. We note that surfaces in  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  are equipped with a ray in the spinor bundle at each R puncture. Therefore, unlike  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , there are two canonical choices for the gluing of R punctures on a single surface in  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .  $\Delta^{\text{R}+}$  and

---

<sup>31</sup>We have used the same notation of  $\Delta$  for gluing operation on the spin curves as well as ordinary Riemann surface. In this paper, we exclusively work with spin curves.

$\Delta^{R-}$  denote these two possible choices, i.e. two rays associated to the glued punctures are either of the same phase  $\Delta^{R+}$  or of the opposite phase  $\Delta^{R-}$ . The operation  $d + \hbar\Delta$ , where  $d$  is as usual the boundary map on chain complexes, turns  $\mathcal{F}(\tilde{\mathcal{S}})$  into a BV algebra. For brevity, let us define

$$\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \equiv \frac{\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})}{S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}}, \quad (4.10)$$

and denote its fundamental chain by  $[\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$ . After passing to homology, these classes satisfy (see the proof of Lemma 4.1 in Section 4.5)

$$d[\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})] + \Delta^{\text{NS}}[\mathbb{X}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}})] + \Delta^{R+}[\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)] + \Delta^{R-}[\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)] = 0. \quad (4.11)$$

If  $\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  denotes the subspace of connected genus- $g$  surfaces, and  $[\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$  its fundamental class, we can define the following formal sum<sup>32</sup>

$$[\mathbb{X}] \equiv \sum_{\substack{g, \mathbf{n}=0 \\ 2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}}>0}}^{\infty} \hbar^{2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}}} [\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]. \quad (4.12)$$

Using this and (4.11), we have (see Lemma 4.1 in section 4.5)

$$(d + \hbar\Delta) \exp\left(\frac{[\mathbb{X}]}{\hbar}\right) = 0, \quad (4.13)$$

i.e.  $[\mathbb{X}]$  satisfies the BV quantum master equation. A more transparent way to write this equation is

$$\begin{aligned} & d[\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})] + \Delta^{\text{NS}}[\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}})] + \Delta^{R+}[\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)] + \Delta^{R-}[\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)] \\ & + \frac{1}{2} \sum_{\substack{g_1+g_2=g \\ \mathbf{n}_{\text{NS}}^1+\mathbf{n}_{\text{NS}}^2=\mathbf{n}_{\text{NS}}+2 \\ \mathbf{n}_{\text{R}}^1+\mathbf{n}_{\text{R}}^2=\mathbf{n}_{\text{R}}}} \{[\mathbb{X}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1)], [\mathbb{X}_{g_2}(\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2)]\}_{\text{NS}} \\ & + \frac{1}{2} \sum_{\substack{g_1+g_2=g \\ \mathbf{n}_{\text{NS}}^1+\mathbf{n}_{\text{NS}}^2=\mathbf{n}_{\text{NS}} \\ \mathbf{n}_{\text{R}}^1+\mathbf{n}_{\text{R}}^2=\mathbf{n}_{\text{R}}+2}} \{[\mathbb{X}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1)], [\mathbb{X}_{g_2}(\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2)]\}_{\text{R}} = 0. \end{aligned} \quad (4.14)$$

Using this result, we can show one of the main results of this paper (Theorem 4.2 in Section 4.5)

**Theorem (The Existence and Uniqueness of Closed Superstring Vertices).** *For each triple  $(g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  with  $2g - 2 + \mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}} > 0$ , there exists an element  $\mathcal{V}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \in \mathcal{F}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}(\mathcal{S})$ , of homological degree  $6g - 6 + 2\mathbf{n}_{\text{NS}} + 2\mathbf{n}_{\text{R}}$ , with the following properties*

1.  $\mathcal{V}_0(3, 0)$  is the fundamental cycle of  $\mathcal{S}_0(3, 0)/S_{3,0}$ , i.e. 0-chain of coefficient  $1/3!$ .

---

<sup>32</sup>We have used the same notation for the formal sum of  $[\mathbb{X}]$  for formal power series of  $[\mathbb{X}_g(\mathbf{n})]$  as well. In this paper however  $[\mathbb{X}]$  exclusively mean the formal sum (4.12).

2.  $\mathcal{V}_0(1,2)$  is the fundamental cycle of  $\mathcal{S}_0(1,2)/S_{1,2}$ , i.e. 0-chain of coefficient  $1/2!$ .

3. The generating function

$$\mathcal{V} \equiv \sum_{2g-2+n_{\text{NS}}+n_{\text{R}}>0} \hbar^{2g-2+n_{\text{NS}}+n_{\text{R}}} \mathcal{V}_g(n_{\text{NS}}, n_{\text{R}}) \in \hbar\mathcal{F}(\mathcal{S})[[\hbar]], \quad (4.15)$$

satisfies the BV quantum master equation

$$(d + \hbar\Delta) \exp\left(\frac{\mathcal{V}}{\hbar}\right) = 0.$$

4.  $\mathcal{V}$  is unique up to homotopy in the category of BV algebras.

Once again the quasi-isomorphism of  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\tilde{\mathcal{S}})$  make it possible to conclude the results for  $\mathcal{F}(\mathcal{S})$ .

Another main result of [26], which again we generalize to the case of bordered spin-Riemann surfaces in Section 4.6, is the relation between the solution of the BV quantum master equation in the BV algebra  $\mathcal{F}_{g,n}(\mathcal{M})$  and the fundamental class of the DM stack of stable ordinary Riemann surfaces. Let  $\overline{\mathcal{M}}_n$  denote the moduli stack of stable Riemann surfaces with  $n$  punctures, and  $\overline{\mathcal{M}}_{g,n}$  denote the moduli stack of stable connected genus- $g$  Riemann surfaces with  $n$  punctures. One can define a commutative differential-graded algebra as follows

$$\mathcal{F}(\overline{\mathcal{M}}) = \bigoplus_n \mathbb{C}_* \left( \frac{\overline{\mathcal{M}}_n}{S_n} \right), \quad (4.16)$$

and turn it into a BV algebra by setting  $\Delta = 0$ . We then define the following formal sum

$$[\overline{\mathcal{M}}] \equiv \sum_{\substack{g,n \\ 2g-2+n>0}} \hbar^{2g-2+n} [\overline{\mathcal{M}}_{g,n}/S_n] \in \hbar\mathcal{F}(\overline{\mathcal{M}})[[\hbar]], \quad (4.17)$$

where  $[\overline{\mathcal{M}}_{g,n}/S_n]$  is the fundamental chain of  $\overline{\mathcal{M}}_{g,n}/S_n$ . The following theorem provides the link between  $\mathcal{V}$ , the solution of the BV quantum master equation in the BV algebra  $\mathcal{F}(\mathcal{M})$  defined in (4.8), and  $[\overline{\mathcal{M}}]$  (Theorem 10.4.1 of [26])

**Theorem (Bosonic-String Vertices and Fundamental Classes of DM Stacks).** *There is a map in the homotopy category of BV algebras that sends  $\mathcal{V} \longrightarrow [\overline{\mathcal{M}}]$ . This means that the bosonic-string vertices  $\mathcal{V}_g(n)$  is homotopy-equivalent in the homotopy category of BV algebras to an orbifold fundamental chain  $[\overline{\mathcal{M}}_{g,n}/S_n]$  of the compactified moduli stack of connected genus- $g$  Riemann surfaces with  $n$  punctures, i.e. it is homotopy-equivalent to the whole compactified moduli stack.*

Let us present the analogous result for the case of bordered spin-Riemann surfaces. Let  $\overline{\mathcal{S}}_{n_{\text{NS}}, n_{\text{R}}}$  denote the moduli stack of stable spin curves with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  Ramond punctures, and  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  denote the moduli stack of stable connected spin curves with genus

$g$ ,  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  Ramond punctures. One again can define the following chain complex

$$\mathcal{F}(\bar{\mathcal{S}}) \equiv \bigoplus_{n_{\text{NS}}, n_{\text{R}}} \mathcal{C}_* \left( \frac{\bar{\mathcal{S}}_{n_{\text{NS}}, n_{\text{R}}}}{S_{n_{\text{NS}}, n_{\text{R}}}} \right). \quad (4.18)$$

$S_{n_{\text{NS}}, n_{\text{R}}}$  permutes the set of NS and R punctures among each other separately. After taking the homology, and setting  $\Delta = 0$ , we have a BV algebra. We then define the following formal sum

$$[\bar{\mathcal{S}}] \equiv \sum_{\substack{g, n_{\text{NS}}, n_{\text{R}} \\ 2g-2+n_{\text{NS}}+n_{\text{R}} \geq 0}} \hbar^{2g-2+n_{\text{NS}}+n_{\text{R}}} [\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}] \in \hbar H_*(\mathcal{F}(\bar{\mathcal{S}}))[[\hbar]], \quad (4.19)$$

where  $[\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}]$  is the fundamental class of  $\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}$ . One can then establish the following theorem which provides a link between  $\mathcal{V}$ , the solution of the BV quantum master equation in the BV algebra  $\mathcal{F}(\mathcal{S})$  defined in (4.15), and  $[\bar{\mathcal{S}}]$  (Proposition 4.5.1 in Section 4.6)

**Theorem (Superstring Vertices and Fundamental Classes of DM Stacks).** *There is a map in the homotopy category of BV algebras that sends  $\mathcal{V} \longrightarrow [\bar{\mathcal{S}}]$ . This means that  $\mathcal{V}_{g, n_{\text{NS}}, n_{\text{R}}}$  is homotopy-equivalent in the homotopy category of BV algebras to the orbifold fundamental class  $[\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} / S_{n_{\text{NS}}, n_{\text{R}}}]$  of the compactified moduli stack of connected genus- $g$  spin curves with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  R punctures, i.e. it is homotopy-equivalent to the whole compactified moduli stack.*

This ends our brief summary of the results in this section.

## 4.2 Gluing of Boundary Components

In this section, we explain the gluing of boundary components on spin-Riemann surfaces. We take two bordered spin-Riemann surfaces  $\mathcal{C}_1$  and  $\mathcal{C}_2$ . Suppose that  $n_{\text{NS}}^1$  and  $n_{\text{NS}}^2$ , the number of NS boundary components on  $\mathcal{C}_1$  and  $\mathcal{C}_2$  respectively, are nonzero. We can then pick one NS boundary component from each curve, say that the  $i_1$ th NS boundary component of  $\mathcal{C}_1$  and  $i_2$ th NS boundary component of  $\mathcal{C}_2$ , and glue them to get a new spin curve  $\mathcal{C}_1 \cup \mathcal{C}_2$ , and the spin structure  $\mathcal{E}$  on  $\mathcal{C}_1 \cup \mathcal{C}_2$  is the direct sum of pushforward of spin structures  $\mathcal{E}_1$  and  $\mathcal{E}_2$  via the gluing map  $\mathcal{C}_1 \sqcup \mathcal{C}_2 \rightarrow \mathcal{C}_1 \cup \mathcal{C}_2$ . This can be easily generalized to families of bordered spin-Riemann surfaces hence giving rise to an operation on corresponding stacks

$$\mathbf{m}_{i_1, i_2}^{\text{NS}} : \mathcal{S}_{g_1}(n_{\text{NS}}^1, n_{\text{R}}^1) \times \mathcal{S}_{g_2}(n_{\text{NS}}^2, n_{\text{R}}^2) \rightarrow \partial \mathcal{S}_{g_1+g_2}(n_{\text{NS}}^1 + n_{\text{NS}}^2 - 2, n_{\text{R}}^1 + n_{\text{R}}^2), \quad (4.20)$$

where  $n_{\text{NS}}^i$  and  $n_{\text{R}}^i$  for  $i = 1, 2$  are the number of NS and R boundary components on  $\mathcal{C}_i$ . This can be defined for a bordered spin-Riemann surfaces with at least two NS boundary components, say the  $i$ th and the  $j$ th ones, as well

$$\mathbf{g}_{i,j}^{\text{NS}} : \mathcal{S}_g(n_{\text{NS}}, n_{\text{R}}) \rightarrow \partial \mathcal{S}_{g+1}(n_{\text{NS}} - 2, n_{\text{R}}). \quad (4.21)$$

Note that these maps depends on the choice of  $i_1$ ,  $i_2$ ,  $i$ , and  $j$ , in an equivariant way. For example, if  $\sigma_{i_1, i'_1} \in S_{n_1}$  switches  $i_1$  and  $i'_1$  (it acts on  $\mathcal{S}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1)$  naturally), then we have  $\mathbf{m}_{i'_1, i_2}^{\text{NS}} = \mathbf{m}_{i_1, i_2}^{\text{NS}} \circ \sigma_{i_1, i'_1}$ .

We can also define the gluing operation for a pair of R boundary components belong to two bordered spin-Riemann surfaces: the gluing of spin structures is simply choosing a square root of gluing data for the twisted canonical sheaf (since the map  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega$  is non-degenerate at R-nodes, as we explained in section 3.1). Note that the choice of the sign of square root does not matter because changing the sign corresponds to a  $\mathbb{Z}/2$ -automorphism of the spinor bundle on one of the curve (fix the spinor bundle on the other one), but this  $\mathbb{Z}/2$ -automorphism has been quotient out in the definition of the space of bordered spin-Riemann surfaces, so the resulting spin structure is canonically isomorphic to each other. The gluing of borders give rises to an operation on the corresponding moduli spaces

$$\mathbf{m}_{j_1, j_2}^{\text{R}} : \mathcal{S}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1) \times \mathcal{S}_{g_2}(\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2) \rightarrow \partial \mathcal{S}_{g_1+g_2}(\mathbf{n}_{\text{NS}}^1 + \mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^1 + \mathbf{n}_{\text{R}}^2 - 2). \quad (4.22)$$

So far there was no difference with the classical gluing property of the moduli space of bordered ordinary Riemann surfaces  $\mathcal{M}_g(\mathbf{n})$ . However, it turns out that there is no self-gluing operation of R boundary components<sup>33</sup>: the reason is that the  $\mathbb{Z}/2$  ambiguity can not be removed by doing an automorphism (there is only one curve, an automorphism affects two boundary components simultaneously), i.e. there is no canonical choice of gluing data. Nevertheless, we will see in Section 4.4 that this ambiguity can be cured by introducing a new model for  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .

### 4.3 Connected Components of $\partial \bar{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$

A crucial step for the proof of the existence of bosonic-string vertices used in [26] is to establish a bound on the homological dimension of  $\mathcal{M}_{g, \mathbf{n}}/S_{\mathbf{n}}$  for  $(g, \mathbf{n}) \neq (0, 3)$  (see the proof of Proposition 10.1.1 in [26])


$$H_i(\mathcal{M}_{g, \mathbf{n}}/S_{\mathbf{n}}, \mathbb{C}) = 0, \quad i \geq 6g - 7 + 2\mathbf{n}. \quad (4.23)$$

We will prove a similar result for stacks of spin curves. We find that a similar result holds in general other than a few exceptional cases. It turns out that to establish the result, we need two know the number of connected components of  $\partial \bar{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ . Therefore, we describe the components of  $\partial \bar{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  in this section. The main result is

**Theorem 4.1** (Connected Components of  $\partial \bar{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ ). *For each connected component of  $\bar{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ , its boundary is connected, except for  $\bar{\mathcal{S}}_{0,4}^{\mathbf{m}}$ <sup>34</sup> and  $\bar{\mathcal{S}}_{1,1,0}^{ev}$ , in which cases the number of boundary components are 3 and 2, respectively.*

<sup>33</sup>This means the gluing of two R boundary components on the same surface.

<sup>34</sup>Notice that here we have four punctures whose puncturing pattern is encoded in the vector  $\mathbf{m} = (m_1, m_2, m_3, m_4)$ , as we explained in Section 1.3. They can be either four NS punctures, two NS punctures and two R punctures, or four R punctures.



**Figure 3:** A surface with a node whose connected components are an elliptic curve with a single R puncture, and a genus- $(g-1)$  curve with  $n_R$  R punctures and  $n_{NS}$  NS punctures. The red dots denote NS punctures and blue dots denote R punctures.

*Proof.* We divide the proof into different cases.

**The Case  $g = 0$ :** Since  $\bar{\mathcal{S}}_{0,n_{NS},n_R} \cong \bar{\mathcal{M}}_{0,n_{NS}+n_R}$ , and the result for the latter is classical [20].


For other cases, notice that the classical result of connectedness of  $\partial\bar{\mathcal{M}}_{g,n}$ , together with the finiteness and flatness of the morphism  $f : \bar{\mathcal{S}}_{g,n_{NS},n_R} \longrightarrow \bar{\mathcal{M}}_{g,n_{NS}+n_R}$ , imply that every connected component of  $\partial\bar{\mathcal{S}}_{g,n_{NS},n_R}$  maps surjectively to  $\partial\bar{\mathcal{M}}_{g,n_{NS}+n_R}$ . We make use of this fact in the following way: if we can show that for some connected substack  $\mathcal{S} \subset \partial\bar{\mathcal{M}}_{g,n_{NS}+n_R}$ , the preimage  $f^{-1}(\mathcal{S})$  lies in the same connected component of  $\partial\bar{\mathcal{S}}_{g,n_{NS},n_R}$ , then it follows that  $\partial\bar{\mathcal{S}}_{g,n_{NS},n_R}$  is connected.

**The Case  $n_R \neq 0$ :** This separates into 2 subcases:


- $g \geq 2$ : Consider the locus  $\mathcal{S}$  of  $\partial\bar{\mathcal{M}}_{g,n_{NS}+n_R}$  consisting of gluing an elliptic curve with a genus  $g-1$  curve at a node, with one punctures on the elliptic curve and all other punctures on the other component. We claim that every connected component of the preimage  $f^{-1}(\mathcal{S})$  can be connected to the component representing the configuration shown in Figure 3, i.e. one R puncture on the elliptic curve and all other punctures are on the other component. This can be seen from the degeneration pattern of Figure 4.
- $g = 1$  and  $n_R \geq 2$ : Consider the locus  $\mathcal{S}$  of  $\partial\bar{\mathcal{M}}_{1,n_{NS}+n_R}$  consisting of gluing  $n_{NS} + n_R$  projective lines at  $n_{NS} + n_R$  node, with a punctures at each component (there is a unique such curve up to isomorphism). We claim that every connected component of the preimage  $f^{-1}(\mathcal{S})$  can be connected to the component representing consecutively  $n_R$  components with R punctures on them. This can be seen from the degeneration pattern of figure 5.

**The Case  $n_R = 0$ :** This separates into 6 sub-cases:

1.  $g \geq 3$ : Consider the locus  $\mathcal{S}$  of  $\partial\bar{\mathcal{M}}_{g,n_{NS}}$  consisting of gluing an elliptic curve with a genus- $(g-1)$  curve at a node, with all punctures on the elliptic curve. For the even spin structure component, the preimage  $f^{-1}(\mathcal{S})$  consists of two connected components arising from gluing spin structures of the same parity. They are connected via degenerations




**Figure 4:** The degeneration pattern of a surface with a node consists of two connected components, a genus- $(g-1)$  curve with a number of NS and R punctures, and a genus-1 surface with a single NS or R puncture. The red dots denote the NS punctures and blue dots denote the R punctures.




**Figure 5:** The degeneration pattern of a surface consists of multiple connected components joined at several nodes. The red dot denotes the NS puncture, and blue dots denote the R punctures.

shown in Figure 6. Similarly for odd spin structure component, the preimage  $f^{-1}(\mathcal{S})$  consists of two connected components arising from gluing spin structures of opposite parity. They are connected via degenerations shown in Figure 7

2.  $g \geq 1$  and  $n_{\text{NS}} \geq 2$ : Consider the locus  $\mathcal{S}$  of  $\partial\overline{\mathcal{M}}_{g,n_{\text{NS}}}$  consisting of gluing a projective line with a genus- $g$  curve at a node, with all punctures on the projective line. It is easy to see that the preimage  $f^{-1}(\mathcal{S})$  restricted to each parity component is connected, since the parity of spin structure on the genus- $g$  curve is the same as the one of the nodal curve.
3.  $g = 1$  and  $n_{\text{NS}} = 1$  with odd spin structure: The boundary of  $\partial\overline{\mathcal{M}}_{1,1}$  is the unique genus-1 nodal curve with a puncture. The preimage  $\partial\overline{\mathcal{M}}_{1,1}$  is connected since the only odd spin structure comes from gluing the spin structure of  $\mathbb{P}^1$  with one NS puncture and



**Figure 6:** The degeneration patterns of a spin curve with a node and even spin structure. One of the connected components is a genus- $(g-1)$  curve with  $n_{\text{NS}}$  NS punctures, and the other component is an elliptic curve. The red dots denote NS punctures. In the figure,  $n_{\text{NS}} = 3$ .




**Figure 7:** The degeneration patterns of a spin curve with a node and odd spin structure. One of the connected components is a genus- $(g-1)$  curve with  $n_{\text{NS}}$  NS punctures, and the other component is an elliptic curve. The red dots denote NS punctures. In the figure,  $n_{\text{NS}} = 3$ .

two R punctures (note that the spin structure  $\mathcal{E}$  is isomorphic to the structure sheaf) in a way that the global section of the spinor bundle  $\mathcal{E}$  (which is one dimensional) is preserved.



4.  $g = 2$  and  $n_{\text{NS}} = 1$  with odd spin structure: Consider the locus  $\mathcal{S}$  of  $\partial\overline{\mathcal{M}}_{2,1}$  consisting of gluing two elliptic curves at a node. Let us label them by  $E_1$  and  $E_2$ . We choose the only punctures to be on  $E_2$ . The preimage  $f^{-1}(\mathcal{S})$  consists of two connected components arising from gluing spin structures of opposite parity. They are connected via degeneration to the same boundary:




**Figure 8:** The degeneration pattern of a genus-2 spin curve with a node and odd spin structure. The component surfaces are elliptic curves and there is an NS puncture, the red dot, on one of them.

5.  $g = 2$  and  $n_{\text{NS}} = 0$  with odd spin structure: Consider the locus  $\mathcal{S}$  of  $\partial\overline{\mathcal{M}}_{2,0}$  consisting of gluing two *isomorphic* elliptic curves  $E_1$  and  $E_2$  at a node. The preimage  $f^{-1}(\mathcal{S})$  is connected since spin structures on two elliptic curves must have opposite parity, i.e. either even on  $E_1$  and odd on  $E_2$  or the other way around, and they are isomorphic via pullback along the isomorphism of switching two components.
6.  $g = 2$  and  $n_{\text{NS}} = 0$  or 1 with even spin structure: This is the hardest case, we first consider the locus  $\mathcal{S}_1$  of  $\partial\overline{\mathcal{M}}_{2,n_{\text{NS}}}$  consisting of gluing two elliptic curves at a node. The preimage  $f^{-1}(\mathcal{S}_1)$  has two connected components since spin structures on two elliptic curves must have the same parity, i.e. either even-even or odd-odd, denoted by  $f^{-1}(\mathcal{S}_1)_e$  and  $f^{-1}(\mathcal{S}_1)_o$ . For a general point  $p$  in  $\mathcal{S}_1$ , its preimage consists of 10 points: nine of which lies in  $f^{-1}(\mathcal{S}_1)_e$  and one of which lies in  $f^{-1}(\mathcal{S}_1)_o$ . To show that  $\partial\overline{\mathcal{S}}_{2,n_{\text{NS}},0}$  is connected, it suffices to show that the unique point lying in  $f^{-1}(\mathcal{S}_1)_o$  can be connected one of points lying in  $f^{-1}(\mathcal{S}_1)_e$ .

We temporarily switch to another locus  $\mathcal{S}_2$  of  $\partial\overline{\mathcal{M}}_{2,n_{\text{NS}}}$  consisting of irreducible curves with one node. Its preimage  $f^{-1}(\mathcal{S}_1)$  has two connected components: the one with an NS node and another one with an R node<sup>35</sup>. Let us focus on the component with an R node. We conclude the proof of this case by the following degeneration procedure.

It remains to deal with the exceptional cases:  $\overline{\mathcal{S}}_{1,1,0}^{\text{ev}}$ . It consists of two connected components, corresponding to two even spin structures associated to the unique genus-1 nodal curve with a puncture. One of these spin structures comes from the gluing of a pair of NS punctures on a  $\mathbb{P}^1$  with three NS punctures, and the other one comes from the gluing of the spin structure

<sup>35</sup>As we explained in section 4.2, there is no self contraction  $\overline{\mathcal{S}}_{1,n_{\text{NS}},2} \rightarrow \partial\overline{\mathcal{S}}_{2,n_{\text{NS}},0}$ . In fact, the connectedness of the second component follows from the construction in 4.4.



**Figure 9:** The degeneration pattern of genus-2 curves with NS or R nodes. The red dots denote NS nodes and blue dots denote R nodes.

of  $\mathbb{P}^1$  with one NS puncture and two R punctures in a way that the global section of  $\mathcal{E}$  is not preserved. ■

We can state the following corollary to the Theorem 4.1

**Corollary 4.1.** *Let permutation groups  $S_{n_{\text{NS}}}$  and  $S_{n_{\text{R}}}$  act on NS and R punctures naturally, and define  $S_{n_{\text{NS}}, n_{\text{R}}} \equiv S_{n_{\text{NS}}} \times S_{n_{\text{R}}}$ . Then for  $(g, n_{\text{NS}} + n_{\text{R}}) \neq (3, 0)$  we have*

$$H_{6g-6+2(n_{\text{NS}}+n_{\text{R}})}(\mathcal{S}_{g, n_{\text{NS}}, n_{\text{R}}}/S_{n_{\text{NS}}, n_{\text{R}}}, \mathbb{C}) = 0,$$

and

$$H_{6g-7+2(n_{\text{NS}}+n_{\text{R}})}(\mathcal{S}_{g, n_{\text{NS}}, n_{\text{R}}}/S_{n_{\text{NS}}, n_{\text{R}}}, \mathbb{C}) = \begin{cases} \mathbb{C}, & (g, n_{\text{NS}}, n_{\text{R}}) = (0, 2, 2) \text{ or } (1, 1, 0)^{ev}, \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* The top homology is trivial since  $\mathcal{S}_{g, n_{\text{NS}}, n_{\text{R}}}$  is noncompact. For the other homology groups, we first compute the group without quotient, using the Poincaré duality

$$H_{6g-7+2(n_{\text{NS}}+n_{\text{R}})}(\mathcal{S}_{g, n_{\text{NS}}, n_{\text{R}}}, \mathbb{C}) \cong H^1(\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \partial \bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \mathbb{C}), \quad (4.24)$$

Observe that  $\pi_1(\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}})$  is trivial since the spin mapping-class group is generated by Dehn twists, and the boundary of  $\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  trivializes those Dehn twists. It follows from the long exact sequence of cohomologies for the pair  $(\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \partial \bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}})$  that

$$H^1(\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \partial \bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \mathbb{C}) \cong H^0(\partial \bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \mathbb{C})/H^0(\bar{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}, \mathbb{C}). \quad (4.25)$$

This identification shows that  $H_{6g-7+2(n_{\text{NS}}+n_{\text{R}})}(\mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}},\mathbb{C})$  is canonically identified with  $\mathbb{C}^{\#\text{Boundary Components}-1}$ . This proves the case when  $(g,n_{\text{NS}},n_{\text{R}})$  is not one of  $(0,4,0)$ ,  $(0,2,2)$ ,  $(0,0,4)$ , or  $(1,1,0)^{\text{ev}}$ , since there is only one boundary component. For  $(g,n_{\text{NS}},n_{\text{R}}) = (1,1,0)^{\text{ev}}$ , it is easy to see that the permutation group  $S_{n_{\text{NS}},n_{\text{R}}}$  preserves those boundary components, hence taking quotient does not affect the homology, i.e. they are still one dimensional. The action of  $S_4$  on the boundary of  $\overline{\mathcal{M}}_{0,4}$  (which consists of 3 points) is transitive, hence after taking quotient the homology becomes trivial. It remains to examine the action of  $S_2 \times S_2$  on the the boundary of  $\overline{\mathcal{M}}_{0,4}$ . The order of  $S_2 \times S_2$  is 4, its image in  $S_3$  is either trivial or  $S_2$ . Since we know that the action must be nontrivial: there is a  $S_2$  component corresponding to switching two NS punctures. Hence the image of  $S_2 \times S_2$  in  $S_3$  is  $S_2$ , after taking the quotient, and the homology is one dimensional.  $\blacksquare$

**Remark 4.1.** *It follows from the Corollary 4.1 that the stack  $\partial\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}/S_{n_{\text{NS}},n_{\text{R}}}$  has at most two connected components. The intersection pairing between  $H_{6g-7+2(n_{\text{NS}}+n_{\text{R}})}(\mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}}/S_{n_{\text{NS}},n_{\text{R}}},\mathbb{C})$  and  $H_1(\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}/S_{n_{\text{NS}},n_{\text{R}}},\partial\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}/S_{n_{\text{NS}},n_{\text{R}}},\mathbb{C})$  (when it is not zero) is given by connecting two boundary components with a generic smooth arc and counting the intersection number with a  $(6g-7+2(n_{\text{NS}}+n_{\text{R}}))$ -cycle.*

#### 4.4 An Alternative Model for $\mathcal{S}(n_{\text{NS}},n_{\text{R}})$

As we discussed in Section 2.3, there is an ambiguity of gluing R boundary components on a single spin-Riemann surfaces due to a  $\mathbb{Z}/2$ -automorphism acting on the spinor bundle over the surface. This motivates the following question: *can we define a classifying space for spin curves such that there is no such ambiguity?* The answer turns out to be positive. We can indeed define such a space by natural generalization of the space considered in [50] as a model for the space of bordered ordinary Riemann surfaces. In this section, we elaborate the generalization of this construction for the space of bordered spin-Riemann surfaces.

$\mathcal{S}_g(n_{\text{NS}},n_{\text{R}})$  has a nice model  $\widetilde{\mathcal{S}}_g(n_{\text{NS}},n_{\text{R}})$ , which is the moduli stack of stable Riemann surfaces, i.e. the surfaces belong to  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}$ , decorated at each NS puncture with a ray in the tangent space, at each R puncture with a ray in the spinor line bundle  $\mathcal{E}$ , at each NS node with a ray in the tensor product of rays of the tangent spaces at each side, and at each R node with a ray in the spinor line bundle  $\mathcal{E}$ . This is an orbifold with corners whose interior consists of smooth spin curves, so it is homotopy-equivalent to  $\mathcal{S}_g(n_{\text{NS}},n_{\text{R}})$ . According to Section 2.3, we have the natural gluing operation which give rises to an operation on the corresponding moduli stacks

$$\begin{aligned} \mathbf{m}_{i_1,i_2}^{\text{NS}} : \widetilde{\mathcal{S}}_{g_1}(n_{\text{NS}}^1,n_{\text{R}}^1) \times \widetilde{\mathcal{S}}_{g_2}(n_{\text{NS}}^2,n_{\text{R}}^2) &\rightarrow \widetilde{\mathcal{S}}_{g_1+g_2}(n_{\text{NS}}^1+n_{\text{NS}}^2-2,n_{\text{R}}^1+n_{\text{R}}^2), \\ \mathbf{m}_{i_1,i_2}^{\text{R}} : \widetilde{\mathcal{S}}_{g_1}(n_{\text{NS}}^1,n_{\text{R}}^1) \times \widetilde{\mathcal{S}}_{g_2}(n_{\text{NS}}^2,n_{\text{R}}^2) &\rightarrow \widetilde{\mathcal{S}}_{g_1+g_2}(n_{\text{NS}}^1+n_{\text{NS}}^2,n_{\text{R}}^1+n_{\text{R}}^2-2), \end{aligned} \quad (4.26)$$

as well as self-gluing for NS punctures

$$\mathbf{g}_{i,j}^{\text{NS}} : \widetilde{\mathcal{S}}_g(n_{\text{NS}},n_{\text{R}}) \rightarrow \widetilde{\mathcal{S}}_{g+1}(n_{\text{NS}}-2,n_{\text{R}}). \quad (4.27)$$

Note that the gluing operation takes two rays at two NS points to their tensor product.

On the other hand, the ambiguity of gluing R boundaries for  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  disappears due to the fact that there are two canonical choices of gluing R punctures for surfaces in  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ : two rays are either of the same phase or of the opposite phase. We denote the corresponding self-gluing operation  $\mathfrak{g}_{i,j}^{\text{R}+}$  and  $\mathfrak{g}_{i,j}^{\text{R}-}$ :

$$\mathfrak{g}_{i,j}^{\text{R}\pm} : \tilde{\mathcal{S}}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \rightarrow \tilde{\mathcal{S}}_{g+1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} - 2). \quad (4.28)$$

Note that all of these gluing operations are equivariant under the permutation group action. Therefore, they can be defined on the quotient space without referring to an explicit choice of punctures, i.e. we have

$$\begin{aligned} \mathbf{m}^{\text{NS}} : \tilde{\mathcal{S}}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1) / S_{\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1} &\times \tilde{\mathcal{S}}_{g_2}(\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2) / S_{\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2} \rightarrow \tilde{\mathcal{S}}_{g_1+g_2}(\mathbf{n}_{\text{NS}}^1 + \mathbf{n}_{\text{NS}}^2 - 2, \mathbf{n}_{\text{R}}^1 + \mathbf{n}_{\text{R}}^2) / S_{\mathbf{n}_{\text{NS}}^1 + \mathbf{n}_{\text{NS}}^2 - 2, \mathbf{n}_{\text{R}}^1 + \mathbf{n}_{\text{R}}^2}, \\ \mathbf{m}^{\text{R}} : \tilde{\mathcal{S}}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1) / S_{\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1} &\times \tilde{\mathcal{S}}_{g_2}(\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2) / S_{\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2} \rightarrow \tilde{\mathcal{S}}_{g_1+g_2}(\mathbf{n}_{\text{NS}}^1 + \mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^1 + \mathbf{n}_{\text{R}}^2 - 2) / S_{\mathbf{n}_{\text{NS}}^1 + \mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^1 + \mathbf{n}_{\text{R}}^2 - 2}, \\ \mathfrak{g}^{\text{NS}} : \tilde{\mathcal{S}}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) / S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} &\rightarrow \tilde{\mathcal{S}}_{g+1}(\mathbf{n}_{\text{NS}} - 2, \mathbf{n}_{\text{R}}) / S_{\mathbf{n}_{\text{NS}} - 2, \mathbf{n}_{\text{R}}}, \\ \mathfrak{g}^{\text{R}\pm} : \tilde{\mathcal{S}}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) / S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} &\rightarrow \tilde{\mathcal{S}}_{g+1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} - 2) / S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} - 2}. \end{aligned} \quad (4.29)$$

#### 4.5 The Proof of Existence and Uniqueness

We are finally in a position to present one of the main results of this work, i.e. the existence and uniqueness, in the sense we explain, of a solution to the BV QME. Let  $\mathcal{C}_*$  be the functor of normalized singular simplicial chains with coefficients in any field  $\mathbb{F}$  containing  $\mathbb{Q}$ . We define two complexes

$$\begin{aligned} \mathcal{F}(\mathcal{S}) &\equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{C}_*(\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) / S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}), \\ \mathcal{F}(\tilde{\mathcal{S}}) &\equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{C}_*(\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) / S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}), \end{aligned} \quad (4.30)$$

where  $S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  is the wreath product  $(S^1)^{\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}} \rtimes S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ . Let  $\mathcal{F}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}(\mathcal{S})$  (respectively  $\mathcal{F}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}(\tilde{\mathcal{S}})$ ) be the part coming from connected surfaces of with  $g$  handles,  $\mathbf{n}_{\text{NS}}$  NS boundary components (respectively NS punctures) and  $\mathbf{n}_{\text{R}}$  R boundary components (respectively R punctures).  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\tilde{\mathcal{S}})$  naturally carry structures of commutative differential graded algebra, where the differential is the boundary map of chain complexes and the product comes from disjoint union of surfaces. Since  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is homotopy-equivalent to  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ <sup>36</sup>,  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\tilde{\mathcal{S}})$  are quasi-isomorphic. Let us define the following spaces

$$\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \equiv \tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) / S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}. \quad (4.31)$$

$\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is the moduli stack of spin curves with unordered punctures, together with at each NS node, a ray in the tensor product of the tangent spaces at each side, and at each R node,

---

<sup>36</sup>Remember that 1) the phase of ray at a puncture parametrizes a circle, which can be thought of as a boundary component, and 2) the ray at a node can be used to open it.

a ray in the spinor bundle, and unparametrized boundaries since there is no ray on punctures anymore. The boundary of  $\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , i.e. the locus of nodal curves, has real codimension one inside  $\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .

Consider the moduli stack of a curve  $\mathcal{C} \in \mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}})$  together with a chosen pair of NS punctures and a choice of gluing at these two punctures. The choice does not matter because the permutation symmetry has been quotient out. There is an  $S^1$  possible ways of gluing rays. Note that it can be identified with an irreducible component of  $\partial\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . Let us denote this component by  $\mathbb{Y}_g^{\text{NS}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . Similarly, we can define the space  $\mathbb{Y}_g^{\text{R}+}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  and  $\mathbb{Y}_g^{\text{R}-}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , they are differed by a choice of gluing of spin structures at punctures. There is a sequence of maps:

$$\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}}) \longleftarrow \mathbb{Y}_g^{\text{NS}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \longrightarrow \mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \quad (4.32)$$

then define the operator  $\Delta^{\text{NS}} : \mathcal{C}_*(\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}})) \rightarrow \mathcal{C}_*(\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))$  to be the minus of pulling back to  $\mathbb{Y}_g^{\text{NS}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  followed by pushing forward to  $\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . Similarly for  $\Delta^{\text{R}\pm} : \mathcal{C}_*(\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)) \rightarrow \mathcal{C}_*(\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))$ . A similar procedure applies to a pair of punctures coming from different connected curves, which gives rise to brackets  $\{\cdot, \cdot\}_{\text{NS}}$  and  $\{\cdot, \cdot\}_{\text{R}}$ . Note that the pulling back procedure increases the degree of chain by one, since  $\mathbb{Y}_g^\bullet \rightarrow \mathbb{X}_{g-1}$ <sup>37</sup> is an  $S^1$  fibration.

**Definition 4.5.** Let  $\mathcal{F}_k(\tilde{\mathcal{S}})$  denote the subspace spanned by chains on the space of surfaces with at least  $k$  modes of either type. We define the following operators

$$\begin{aligned} \Delta &\equiv \Delta^{\text{NS}} + \Delta^{\text{R}+} + \Delta^{\text{R}-} : \mathcal{F}_k(\tilde{\mathcal{S}}) \rightarrow \mathcal{F}_{k+1}(\tilde{\mathcal{S}}), \\ \{\cdot, \cdot\} &\equiv \{\cdot, \cdot\}_{\text{NS}} + \{\cdot, \cdot\}_{\text{R}} : \mathcal{F}_k(\tilde{\mathcal{S}}) \otimes \mathcal{F}_l(\tilde{\mathcal{S}}) \rightarrow \mathcal{F}_{k+l+1}(\tilde{\mathcal{S}}). \end{aligned} \quad (4.33)$$

The operator  $\hat{d} \equiv d + \hbar \Delta$  makes  $\mathcal{F}(\tilde{\mathcal{S}})$  into a BV algebra.

One can then prove the following lemma which helps us in proving the solution to the BV QME in  $\tilde{\mathcal{S}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S_{S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}}$ .

**Lemma 4.1.** Define the subspace  $\partial_i \mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  to be the locus of spin curves with at least  $i$  nodes, and consider the following decreasing filtration on  $\mathcal{F}(\tilde{\mathcal{S}})$

$$F^i \mathcal{F}(\tilde{\mathcal{S}}) \equiv \text{span} \left\{ \alpha \in \mathcal{F}_*(\tilde{\mathcal{S}}) \mid \alpha \text{ is supported in } \partial_i \mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \right\}. \quad (4.34)$$

Let  $[\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$  be the fundamental class of  $(\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))$ , then it satisfies the BV QME:

$$\begin{aligned} d[\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})] &+ \Delta^{\text{NS}}[\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}})] + \Delta^{\text{R}+}[\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)] + \Delta^{\text{R}-}[\mathbb{X}_{g-1}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)] \\ &+ \frac{1}{2} \sum_{\substack{g_1 + g_2 = g \\ \mathbf{n}_{\text{NS}}^1 + \mathbf{n}_{\text{NS}}^2 = \mathbf{n}_{\text{NS}} + 2 \\ \mathbf{n}_{\text{R}}^1 + \mathbf{n}_{\text{R}}^2 = \mathbf{n}_{\text{R}}}} \{ [\mathbb{X}_{g_1}(\mathbf{n}_{\text{NS}}^1, \mathbf{n}_{\text{R}}^1)], [\mathbb{X}_{g_2}(\mathbf{n}_{\text{NS}}^2, \mathbf{n}_{\text{R}}^2)] \}_{\text{NS}} \end{aligned}$$

---

<sup>37</sup>• denotes either NS or R.

$$+ \frac{1}{2} \sum_{\substack{g_1 + g_2 = g \\ n_{\text{NS}}^1 + n_{\text{NS}}^2 = n_{\text{NS}} \\ n_{\text{R}}^1 + n_{\text{R}}^2 = n_{\text{R}} + 2}} \{ [\mathbb{X}_{g_1}(n_{\text{NS}}^1, n_{\text{R}}^1)], [\mathbb{X}_{g_2}(n_{\text{NS}}^2, n_{\text{R}}^2)] \}_{\text{R}} = 0,$$

in the BV algebra  $(H_*(\mathcal{F}(\tilde{\mathcal{S}})/F^2\mathcal{F}(\tilde{\mathcal{S}})), d + \hbar\Delta)$ .

*Proof.* By definition of  $\Delta$  and  $\{\cdot, \cdot\}$ , they are of degree one with respect to the filtration, i.e.  $\Delta(F^k) \subset F^{k+1}$  and  $\{F^k, F^l\} \subset F^{k+l+1}$ . It follows that  $\text{Gr}_F^i \mathcal{F}(\tilde{\mathcal{S}})$ , where  $\text{Gr}_F^i \mathcal{F}(\tilde{\mathcal{S}})$  is the  $i$ th graded component of the graded algebra associated to the BV algebra  $\mathcal{F}(\tilde{\mathcal{S}})$  and the grading is induced by the filtration  $F^{38}$ , is the relative chain complex  $(\mathcal{C}_*(\partial_i \mathbb{X})/\mathcal{C}_*(\partial_{i+1} \mathbb{X}), d)$ , hence

$$H_*(\text{Gr}_F^i \mathcal{F}(\tilde{\mathcal{S}})) \cong \bigoplus_{n_{\text{NS}}, n_{\text{R}}} H_*(\partial_i \mathbb{X}(n_{\text{NS}}, n_{\text{R}}), \partial_{i+1} \mathbb{X}(n_{\text{NS}}, n_{\text{R}})). \quad (4.35)$$

$\Delta^{\text{NS}}$  acts on the homology class:  $H_k(\text{Gr}_F^i \mathcal{F}(\tilde{\mathcal{S}})) \rightarrow H_{k+1}(\text{Gr}_F^{i+1} \mathcal{F}(\tilde{\mathcal{S}}))$ , i.e. it is a map

$$H_k(\partial_i \mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}}), \partial_{i+1} \mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}})) \rightarrow H_{k+1}(\partial_{i+1} \mathbb{X}(n_{\text{NS}}, n_{\text{R}}), \partial_{i+2} \mathbb{X}(n_{\text{NS}}, n_{\text{R}})). \quad (4.36)$$

Similar statement holds for  $\Delta^{\text{R}\pm}$ .

We look closer to the case  $i = 0$ : there is a fundamental class  $[\mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}})] \in H_*(\mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}}), \partial \mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}}))$ , its image

$$\Delta^{\text{NS}}[\mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}})] \in H_{*+1}(\partial \mathbb{X}(n_{\text{NS}}, n_{\text{R}}), \partial_2 \mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}})), \quad (4.37)$$

is by construction  $-[\mathbb{Y}^{\text{NS}}(n_{\text{NS}} + 2, n_{\text{R}})]$ , i.e. minus the fundamental class of the corresponding irreducible component of  $\partial \mathbb{X}(n_{\text{NS}}, n_{\text{R}})$ . Adding the contribution of  $\Delta^{\text{R}\pm}$ , we see that

$$\Delta^{\text{NS}}[\mathbb{X}(n_{\text{NS}} + 2, n_{\text{R}})] + \Delta^{\text{R}+}[\mathbb{X}(n_{\text{NS}}, n_{\text{R}} + 2)] + \Delta^{\text{R}-}[\mathbb{X}(n_{\text{NS}}, n_{\text{R}} + 2)] = -[\partial \mathbb{X}(n_{\text{NS}}, n_{\text{R}})]. \quad (4.38)$$

On the other hand, the  $\mathbb{Q}$ -orientation of  $\partial \mathbb{X}(n_{\text{NS}}, n_{\text{R}})$  is induced from the  $\mathbb{Q}$ -orientation of  $\mathbb{X}(n_{\text{NS}}, n_{\text{R}})$ , i.e. the fundamental class of the boundary  $\partial \mathbb{X}(n_{\text{NS}}, n_{\text{R}})$  is induced from the

---

<sup>38</sup>Let  $(F_i^M)_{i \in \mathbb{Z}}$  be a decreasing filtration of a module  $M$  over a commutative ring  $R$ . We can define

$$M_i \equiv F_i^M / \sum_{j > i} F_j^M, \quad i, j \in \mathbb{Z}.$$

Then, we can define

$$\text{Gr}_F M \equiv \bigoplus_{i \in \mathbb{Z}} M_i.$$

$\text{Gr}_F M$  is the associated graded  $R$ -module of  $M$ . Assuming that  $M$  has an algebra structure  $*$ :  $M \times M \rightarrow M$  such that  $F_i^M \times F_j^M$  is sent to  $F_{i+j}^M$ , we can define an algebra structure on  $\text{Gr}_F M$  as follows. Consider the canonical projections  $\pi_i : F_i^M \rightarrow M_i$ . For two elements  $a_i \in M_i$  and  $a_j \in M_j$ , we can define the multiplication operation  $\cdot$ :  $\text{Gr}_F M \times \text{Gr}_F M \rightarrow \text{Gr}_F M$  on  $\text{Gr}_F M$

$$a_i \cdot a_j \equiv \pi_{i+j}(a'_i * a'_j),$$

where  $a'_i = \pi_i^{-1}(a_i)$  and  $a'_j = \pi_j^{-1}(a_j)$  are lifts of  $a_i$  and  $a_j$ . The filtered algebra  $\text{Gr}_F M$  endowed with this multiplication is called the associated graded  $R$ -algebra of  $M$ .

boundary of fundamental class of  $\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . Note that the map is exactly the boundary map in the following exact sequence of homologies:

$$\begin{aligned} 0 &= H_{6g-6+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial_2 \mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})) \longrightarrow H_{6g-6+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial \mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})) \\ &\xrightarrow{d} H_{6g-7+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\partial \mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial_2 \mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})) \longrightarrow H_{6g-7+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial_2 \mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})). \end{aligned}$$

It then follows that

$$d[\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})] = [\partial \mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})], \quad (4.39)$$

in the BV algebra  $(H_*(\mathcal{F}(\tilde{\mathcal{S}})/F^2\mathcal{F}(\tilde{\mathcal{S}})), d + \Delta)$ , whence

$$d([\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]) + \Delta^{\text{NS}}([\mathbb{X}(\mathbf{n}_{\text{NS}} + 2, \mathbf{n}_{\text{R}})]) + \Delta^{\text{R}+}([\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)]) + \Delta^{\text{R}-}([\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} + 2)]) = 0. \quad (4.40)$$

We can consider the following formal sum

$$\sum_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} [\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})] = \exp \left( \sum_{\mathbf{g}, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \hbar^{2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}}} [\mathbb{X}_{\mathbf{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})] \right) \equiv \exp([\mathbb{X}]). \quad (4.41)$$

Using this sum and (4.40), we have

$$(d + \hbar \Delta) \exp \left( \frac{[\mathbb{X}]}{\hbar} \right) = 0,$$

i.e. the class  $[\mathbb{X}]$  satisfies the BV QME. ■

Using this result, we have the following Theorem

**Theorem 4.2** (The Existence and Uniqueness of Solution to the BV QME in  $\mathcal{F}(\mathcal{S})$ ). *For each triple  $(g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  with  $2g - 2 + \mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}} > 0$ , there exists an element  $\mathcal{V}_{\mathbf{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \in \mathcal{F}_{\mathbf{g}, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}(\tilde{\mathcal{S}})$ , of homological degree  $6g - 6 + 2\mathbf{n}_{\text{NS}} + 2\mathbf{n}_{\text{R}}$ , with the following properties.*

1.  $\mathcal{V}_0(3, 0)$  is the fundamental cycle of  $\mathbb{X}_0(3, 0)$ , i.e. 0-chain of coefficient  $1/6$ ;
2.  $\mathcal{V}_0(1, 2)$  is the fundamental cycle of  $\mathbb{X}_0(2, 1)$ , i.e. 0-chain of coefficient  $1/2$ ;
3. The generating function

$$\mathcal{V} \equiv \sum_{\substack{\mathbf{g}, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} \\ 2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}} > 0}} \hbar^{2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}}} \mathcal{V}_{\mathbf{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \in \hbar \mathcal{F}(\tilde{\mathcal{S}})[[\hbar]], \quad (4.42)$$

satisfies the BV quantum master equation

$$(d + \hbar \Delta) \exp \left( \frac{\mathcal{V}}{\hbar} \right) = 0. \quad (4.43)$$

4. Such solution  $\mathcal{V}$  is unique up to homotopy through such elements.

*Proof.* Define a dg Lie algebra  $\mathfrak{g} \equiv \oplus_i \mathfrak{g}_i$  where  $\mathfrak{g}_i$  is the set of

$$\mathcal{V} \equiv \sum_{\substack{\mathfrak{g}, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}} \\ 2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}} > 0}} \hbar^{2g-2+\mathbf{n}_{\text{NS}}+\mathbf{n}_{\text{R}}} \mathcal{V}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \in \hbar \mathcal{F}(\tilde{\mathcal{S}})[[\hbar]], \quad (4.44)$$

such that  $\mathcal{V}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \in \mathcal{F}_{\mathfrak{g}, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}(\tilde{\mathcal{S}})$  and  $\deg \mathcal{V}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = 6g - 5 + 2\mathbf{n}_{\text{NS}} + 2\mathbf{n}_{\text{R}} + i$ . Consider the filtration (4.34),  $\mathfrak{g} = F^0 \mathfrak{g} \supset F^1 \mathfrak{g} \supset \dots$ . We define  $\mathfrak{g}'$  to be  $\mathfrak{g}/F^2 \mathfrak{g}$ . It follows that  $\forall k \geq 0$  and  $\forall i \geq 0$ ,  $H_i(F^k \mathfrak{g}/F^{k+1} \mathfrak{g}) = 0$ , for there is no element of non-negative degree. Obviously, the quotient map  $\mathfrak{g} \rightarrow \mathfrak{g}'$  induces isomorphisms

$$H_i(F^k \mathfrak{g}/F^{k+1} \mathfrak{g}) \cong H_i(F^k \mathfrak{g}'/F^{k+1} \mathfrak{g}'), \quad (4.45)$$

when  $k = 0$  or  $k = 1$ . Moreover, when  $k \geq 2$ , there is no chain of degree  $\geq -2$ , so  $H_i(F^k \mathfrak{g}/F^{k+1} \mathfrak{g}) = 0$  when  $i \geq -2$ . To sum up, the quotient map  $\mathfrak{g} \rightarrow \mathfrak{g}'$  induces isomorphisms

$$H_i(\text{Gr}_F \mathfrak{g}) \cong H_i(\text{Gr}_F \mathfrak{g}') \quad i = 0, -1, -2, \quad (4.46)$$

where  $\text{Gr}_F \mathfrak{g}$  is the graded algebra associated to the Lie algebra  $\mathfrak{g}$  and the grading is induced by the filtration  $F$ . Let  $\text{MC}(\mathfrak{g})$  be the set of elements of  $\mathfrak{g}$  satisfying the Maurer-Cartan equation, and  $\pi_0(\text{MC}(\mathfrak{g}))$  be the set of homotopy-equivalence classes of solutions of the Maurer-Cartan equation in  $\mathfrak{g}$  (see Definition 5.1.2 and Lemma 5.2.1 of [26]). It follows from the Lemma 5.3.1 of [26] that the map

$$\pi_0(\text{MC}(\mathfrak{g})) \rightarrow \pi_0(\text{MC}(\mathfrak{g}')), \quad (4.47)$$

is an isomorphism. Since the set of homotopy-equivalence classes of solutions of the Maurer-Cartan equation in the Lie algebra  $\mathfrak{g}$ , defined using (4.44), can be identified with the set of homotopy-equivalence classes of solutions  $\mathcal{V}$  of the BV QME in  $\mathcal{F}(\tilde{\mathcal{S}})$ , the existence of  $\mathcal{V}$  follows immediately from the Lemma 4.1, as the quotient map sends  $\mathcal{V}_0(3, 0)$  to  $[\mathbb{X}_0(3, 0)]$ , and we have seen that the classes  $[\mathbb{X}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$  satisfy the BV QME. It is thus suffices to show that  $[\mathbb{X}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$  is the unique solution to the BV QME in the BV algebra  $H_*(\mathcal{F}(\tilde{\mathcal{S}})/F^2 \mathcal{F}(\tilde{\mathcal{S}}))$  such that the degree  $(\mathfrak{g}, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = (0, 3, 0)$  part is the fundamental class. In fact, suppose that there is another system of solutions

$$\mathcal{V}'_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \in H_{6g-6+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\mathbb{X}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial \mathbb{X}_{\mathfrak{g}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})),$$

then it must satisfy

$$d\mathcal{V}'_1{}^{\text{ev}}(1, 0) + \Delta^{\text{NS}}[\mathbb{X}_0(3, 0)] + \Delta^{\text{R-}}\mathcal{V}'_0(1, 2) = 0.$$

Note that there is no  $\Delta^{\text{R+}}$  term<sup>39</sup>. It then follows from the exact sequence of homologies

$$0 = H_2(\mathbb{X}_1(1, 0)^{\text{ev}}) \longrightarrow H_2(X_1(1, 0)^{\text{ev}}, \partial \mathbb{X}_1(1, 0)^{\text{ev}}) \xrightarrow{d} H_1(\partial \mathbb{X}_1(1, 0)^{\text{ev}}) \longrightarrow H_1(\mathbb{X}_1(1, 0)^{\text{ev}}),$$

---

<sup>39</sup>Since the operator  $\Delta^{\text{R+}}$  preserves the global section of the spin structure  $\mathcal{E}$  on  $\mathbb{P}^1$  with one NS puncture and two R punctures, hence producing an odd spin structure.



that the image of  $\Delta^{\text{NS}}[\mathbb{X}_0(3, 0)] + \Delta^{\text{R-}}\mathcal{V}'_0(1, 2)$  in  $H_1(\mathbb{X}_{1,1,0}^{\text{ev}})$  is trivial. On the other hand, we already know that  $[\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$  is a solution, so the same argument implies that the image of  $\Delta^{\text{NS}}[\mathbb{X}_0(3, 0)] + \Delta^{\text{R-}}[\mathbb{X}_0(1, 2)]$  in  $H_1(\mathbb{X}_1(1, 0)^{\text{ev}})$  is trivial. Comparing these equations we find that

$$\Delta^{\text{R-}}(\mathcal{V}'_0(1, 2) - [\mathbb{X}_0(1, 2)]) = 0,$$

in  $H_1(\mathbb{X}_1(1, 0)^{\text{ev}})$ . However, we know that  $\mathbb{X}_1(1, 0)^{\text{ev}}$ , after reducing to the coarse moduli, is just a sphere with a real blow-up at two points (i.e. the boundary of  $\tilde{\mathbb{S}}_1^{\text{ev}}(1, 0)$ ), and  $\Delta^{\text{R-}}$  sends a zero cycle in  $\tilde{\mathbb{S}}_0(1, 2)$  to the circle fundamental class then embed it isomorphically to one of the boundary circle of  $\mathbb{X}_1(1, 0)^{\text{ev}}$ <sup>40</sup>. In particular,  $\Delta^{\text{R-}}$  is injective. As a consequence, we see that

$$\mathcal{V}'_0(1, 2) - [\mathbb{X}_0(1, 2)] = 0.$$

Now the uniqueness of solution follows from induction and the fact that

$$d : H_{6g-6+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial\mathbb{X}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})) \longrightarrow H_{6g-7+2\mathbf{n}_{\text{NS}}+2\mathbf{n}_{\text{R}}}(\partial\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \partial_2\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})),$$

is injective. This means that lower-degree terms completely determine higher-degree terms via the BV QME. ■

*Another Proof.* Alternatively, one can prove this theorem using the filtration in Proposition 10.1.1 of [26], but the argument there need to be modified since the vanishing of top minus one degree homologies is not always true for the moduli of spin curves. Nevertheless we can take the filtration  $\mathfrak{g} = F^1\mathfrak{g} \supset F^2\mathfrak{g} \supset F^3\mathfrak{g} \cdots$  by letting  $F^k\mathfrak{g}$  generated by the set of  $\mathcal{V}$  such that  $\mathcal{V}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is zero for  $2g - 2 + \mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}} < k$ , when  $k \geq 3$ , and  $F^2\mathfrak{g}$  generated by the set of  $\mathcal{V}$  such that  $\mathcal{V}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is zero for  $2g - 2 + \mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}} < k$  except for  $(g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = (0, 2, 2)$ . The point is that

$$H_i(F^k\mathfrak{g}/F^{k+1}\mathfrak{g}) = 0, \quad \text{for } k \geq 2, \quad \text{and } i = 0, -1, -2,$$

which follows from Corollary 4.1. It follows from the Lemma 5.3.1 of [26] that the map

$$\pi_0(\text{MC}(\mathfrak{g})) \rightarrow \pi_0(\text{MC}(\mathfrak{g}')),$$

is isomorphism, where  $\mathfrak{g}' = \mathfrak{g}/F^2\mathfrak{g}$ . It remains to solve the QME in  $\mathfrak{g}'$ , and this is a consequence of Lemma 4.1 since  $\partial_i\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is empty for

$$(g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = (0, 3, 0), (0, 1, 2), (0, 2, 2), (1, 1, 0).$$

■

---

<sup>40</sup> $\mathbb{X}_1(1, 0)^{\text{ev}}$  is a copy of  $\mathbb{P}^1$  which two boundaries at north and south poles of  $\mathbb{P}^1$ . The real blow-up of these boundaries are circles. We are referring to one of these circles as the boundary circle.

This theorem shows that there exists certain subspaces  $\mathcal{V}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  inside  $\widetilde{\mathcal{S}}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  that satisfy the BV QME. However, we are interested in finding a solution to the BV QME in  $\mathcal{F}(\mathcal{S})$ , and hence find a certain subspace inside  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ . To show that there indeed exists such a subspace, we proceed as follows. As we have explained,  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\widetilde{\mathcal{S}})$  naturally carry structures of commutative differential-graded algebra, where the differential is the boundary map of chain complexes and the product comes from disjoint union of surfaces. We can also define a BV algebra structure on  $\mathcal{F}(\mathcal{S})$  in a similar way to the definition of  $\Delta$  operator of  $\mathcal{F}(\widetilde{\mathcal{S}})$ . Then the Lemma 10.4.2 of [26] can be applied to this case, and we have a homotopy-equivalence of BV algebras:

$$\mathcal{F}(\mathcal{S}) \cong \mathcal{F}(\widetilde{\mathcal{S}}). \quad (4.48)$$

Therefore, all statements in Theorem 4.2 remain true for  $\mathcal{F}(\mathcal{S})$ , i.e. there exists an element (unique up to homotopy)<sup>41</sup>

$$\mathcal{V} \equiv \sum_{\substack{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}} \\ 2g-2+\mathfrak{n}_{\text{NS}}+\mathfrak{n}_{\text{R}} > 0}} \hbar^{2g-2+\mathfrak{n}_{\text{NS}}+\mathfrak{n}_{\text{R}}} \mathcal{V}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}) \in \hbar \mathcal{F}(\mathcal{S})[[\hbar]], \quad (4.49)$$

and  $\mathcal{V}_0(3, 0) = [\mathcal{S}_0(3, 0)/S_{3,0}]$  and  $\mathcal{V}_0(1, 2) = [\mathcal{S}_0(1, 2)/S_{1,2}]$ , satisfying the BV QME

$$(d + \hbar \Delta) \exp \left( \frac{\mathcal{V}}{\hbar} \right) = 0. \quad (4.50)$$

We have thus established that there exist certain subspace  $\mathcal{V}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  in  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  which satisfy the BV QME.

#### 4.6 Relation of Solutions to Fundamental Classes of $\overline{\mathcal{S}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$

In this section, we prove that there is a map in the homotopy category of BV algebras that sends the solution  $\mathcal{V}$  of the BV quantum master equation in the BV algebra  $\mathcal{F}(\widetilde{\mathcal{S}})$  to the fundamental class of the Deligne-Mumford stacks of associated *punctured* spin curves. Consider the following complex

$$\mathcal{F}(\overline{\mathcal{S}}) \equiv \bigoplus_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} \mathcal{C}_*(\overline{\mathcal{S}}_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}/S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}). \quad (4.51)$$

We pass to homology of this complex by taking the boundary operation on chains. We further turn the homology groups to a BV algebra by setting  $\Delta = 0$  [26]. Projecting down to the part coming from connected genus- $g$  surfaces, i.e. to  $\overline{\mathcal{S}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}/S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ , we have a morphism of linear spaces

$$\pi : \mathcal{F}(\widetilde{\mathcal{S}}) \rightarrow H_*(\mathcal{F}(\overline{\mathcal{S}})).$$

In fact, we have

---

<sup>41</sup>By abuse of notation, we use the same notation  $\widehat{\mathcal{V}}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  for subspace of  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ .

**Theorem 4.3** (Relation of Solutions to the BV QME and Fundamental Classes of DM Stacks).  *$\pi$  is a homomorphism of BV algebras, and  $\pi(\widehat{\mathcal{V}}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))$  is the fundamental class  $[\overline{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}/S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}]$ .*

*Proof.* The first statement follows from similar argument for Lemma 10.4.3 in [26]. In the proof of Theorem 4.2, we have seen that the quotient map

$$\mathcal{F}(\widetilde{\mathcal{S}}) \longrightarrow H_*(\mathcal{F}(\widetilde{\mathcal{S}})/F^2\mathcal{F}(\widetilde{\mathcal{S}})),$$

takes  $\mathcal{V}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  to the fundamental class of  $\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . Composing this map with a further quotient

$$H_*(\mathcal{F}(\widetilde{\mathcal{S}})/F^2\mathcal{F}(\widetilde{\mathcal{S}})) \longrightarrow H_*(\mathcal{F}(\widetilde{\mathcal{S}})/F\mathcal{F}(\widetilde{\mathcal{S}})),$$

and then projecting down to  $\overline{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}/S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  sends  $\mathcal{V}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  to the image of  $[\mathbb{X}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})]$  in  $H_*(\mathcal{F}(\widetilde{\mathcal{S}})/F\mathcal{F}(\widetilde{\mathcal{S}}))$ , which is  $[\overline{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}/S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}]$ .  $\blacksquare$

On the other hand, as we explained, there is a homotopy-equivalence (4.48) between the BV algebras  $\mathcal{F}(\mathcal{S})$  and  $\mathcal{F}(\widetilde{\mathcal{S}})$ . Therefore, this fact together with Theorem 4.3 implies that there is a homomorphism in the homotopy category of BV algebras

$$\pi : \mathcal{F}(\mathcal{S}) \rightarrow H_*(\mathcal{F}(\widetilde{\mathcal{S}})),$$

which takes  $\mathcal{V}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , as certain subspaces of  $\mathcal{S}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  which appear in (4.49), to the fundamental class  $[\overline{\mathcal{S}}_{g, \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}/S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}]$ . This establishes the result we were looking for.

## 5 Existence of Solutions to the BV QME in $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$

The main aim of this section is to give a proof for the existence of the solution to the BV QME in  $\mathcal{SM}$  (or more precisely  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ ). The basic idea that we use is the one we have used in previous sections, i.e. we give  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  the structure of a BV algebra, and then show that there is a quasi-isomorphisms between this BV algebra and the BV algebra associated to the underlying moduli space of bordered spin-Riemann surfaces.

One might ask *why we do not try the seemingly simpler method of lifting the solution to the BV quantum master equation in the space of bordered spin-Riemann surfaces to the corresponding space of bordered  $N = 1$  super-Riemann surfaces?* As we show in Section 5.1.1, the solution to the BV QME in  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  can always be lifted to  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ . However, this lifting is unique up to homotopy. As such, the lifting of a solution in  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  to  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  does not necessary give us a solution to BV QME in  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ . One thus need to prove the existence of solution to BV QME in  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$  by a different method. One of these methods is using the quasi-isomorphism of BV algebras and the isomorphism of the set of homotopy-equivalent solutions to the BV QME of such BV algebras. There are other methods for such proof. We mention one of them in the end of Section 5.1.1. The background for this section is provided in Section 7.1.

### 5.1 Obstructions of Splitness and Projectedness

It is interesting to ask when a superscheme  $(X, \mathcal{O}_X)$  is projected or split. To simplify the life, we restrict to the case of smooth superschemes over  $\mathbb{C}$ . Since  $\mathbb{A}^{p|q}$  is automatically split, the definition of smoothness together with Lemma 7.11 implies that smooth superschemes over  $\mathbb{C}$  splits locally. So the question becomes whether local projections (resp. splittings) glue to a global projection (resp. splitting). This is similar to the situation of principal  $G$ -bundles, where bundles are locally trivializable but not necessarily globally trivial, and the obstruction is exactly captured by the cohomology class  $H^1(X, G)$ . Here the gauge group  $G$  is replaced by the group of automorphisms of local models, i.e.  $\text{Aut}(\wedge^\bullet \mathcal{V})$  for a free  $\mathcal{O}_{X_{\text{bos}}}$ -sheaf  $\mathcal{V}$ .

To analyse this problem in detail, we adopt the usual set-up for lifting problems: let  $X_n$  be the superscheme  $(X, \mathcal{O}_X/\mathfrak{n}_X^n)$ , and assume that  $X_n$  splits<sup>42</sup>, i.e.  $\mathcal{O}_{X_n} = \bigoplus_{i=0}^{n-1} \wedge^i \mathcal{V}$ . Let  $G_n$  be the subgroup of  $\text{Aut}(\wedge^\bullet \mathcal{V})$  whose image in  $\text{Aut}(\bigoplus_{i=0}^{n-1} \wedge^i \mathcal{V})$  is the identity, and let  $H_n \equiv G_n/G_{n+1}$ . Then we fix a cover  $\{U_i\}$  of  $X$  such that  $\mathcal{O}_{X_{n+1}}$  splits on  $U_i$ . On overlap  $U_{ij} \equiv U_i \cap U_j$ , splittings on  $U_i$  and  $U_j$  are related by an isomorphism of  $\bigoplus_{i=0}^{n+1} \wedge^i \mathcal{V}$  which maps to identity in  $\text{Aut}(\bigoplus_{i=0}^{n-1} \wedge^i \mathcal{V})$ , i.e. an element  $h_{ij}$  in  $H_n$ ; moreover on  $U_{ijk} \equiv U_i \cap U_j \cap U_k$ ,  $h_{ij}h_{jk} = h_{ik}$ , hence  $\{h_{ij}\}$  defines a class in  $H^1(X, H_n)$ , and is trivial if and only if there is a collection  $\{h_i \in \Gamma(U_i, H_n)\}$ , such that  $h_{ij} = h_i h_j^{-1}$ , whence is equivalent to a splitting of  $X_{n+1}$  over  $X_n$ .

On the other hand, it is easy to see that

$$H_n \cong T_X^{(-)n} \otimes_{\mathcal{O}_{X_{\text{bos}}}} \bigwedge^n \mathcal{V}, \quad (5.1)$$

where  $T_X^+ \equiv T_{X_{\text{bos}}}$  and  $T_X^- \equiv \mathcal{V}^\vee$ . To sum up, we have

**Proposition 5.1.** *Assume  $X_n$  splits, then there is an element*

$$\text{ob}(X_n) \in H^1 \left( X, T_X^{(-)n} \otimes \bigwedge^n \mathcal{V} \right), \quad (5.2)$$

*whose vanishing is necessary and sufficient for the splitting of  $X_{n+1}$ . If it vanishes, then the space of splittings of  $X_{n+1}$  is a torsor<sup>43</sup> under*

$$H^0 \left( X, T_X^{(-)n} \otimes \bigwedge^n \mathcal{V} \right). \quad (5.3)$$

**Corollary 5.1.** *If  $H^1 \left( X, T_X^{(-)n} \otimes \wedge^n \mathcal{V} \right)$  vanishes for all  $n \geq 2$ , then  $X$  splits, and the space of splittings is a torsor under*

$$H^0 \left( X, \bigoplus_{n \geq 1} T_X^{(-)n} \otimes \bigwedge^n \mathcal{V} \right). \quad (5.4)$$

<sup>42</sup> $X_1$  is just  $X_{\text{bos}}$  and  $X_2$  always splits.

<sup>43</sup>For the definition of torsor see [54].

For projectedness, the analysis is analogous. Thanks to Lemma 7.2, we can forget about the odd grading part and analyze the projection problem for embedding  $X_{\text{bos}} \hookrightarrow X_{\text{ev}}$  of ordinary schemes. Let  $X_n^0$  be the scheme  $(X, \mathcal{O}_X^0 / (\mathcal{O}_X^1)^{2n})$ , and assume that  $X_n^0$  splits, i.e.  $\mathcal{O}_{X_n^0} = \bigoplus_{i=0}^{n-1} \wedge^{2i} \mathcal{V}$ . An argument similar to the splitness case that we will skip shows the following:

**Proposition 5.2.** *Assume  $X_n^0$  splits, then there is an element*

$$\text{ob}(X_n^0) \in H^1 \left( X, T_{X_{\text{bos}}} \otimes \bigwedge^{2n} \mathcal{V} \right), \quad (5.5)$$

*whose vanishing is necessary and sufficient for the projectedness of  $X_{n+1}^0$ . If it vanishes, then the space of projection of  $X_{n+1}^0$  onto  $X_n^0$  is a torsor under*

$$H^0 \left( X, T_{X_{\text{bos}}} \otimes \bigwedge^{2n} \mathcal{V} \right). \quad (5.6)$$

**Corollary 5.2.** *If  $H^1 \left( X, T_{X_{\text{bos}}} \otimes \bigwedge^{2n} \mathcal{V} \right)$  vanishes for all  $n \geq 1$ , then  $X$  is projected, and the space of projections is a torsor under*

$$H^0 \left( X, \bigoplus_{n \geq 1} T_{X_{\text{bos}}} \otimes \bigwedge^{2n} \mathcal{V} \right). \quad (5.7)$$

**Remark 5.1.** *A supermanifold is defined to be a superspace  $X$  such that  $X_{\text{bos}}$  is a  $C^\infty$ -manifold and locally on  $X$ ,  $\mathcal{O}_X$  is isomorphic to  $\wedge^\bullet V$  for a vector bundle  $V$  on  $X_{\text{bos}}$ <sup>44</sup>. The above arguments also apply when  $X$  is a supermanifold. In fact,  $H^1 \left( X, T_X^{(-)^n} \otimes \wedge^n V \right)$  vanishes automatically since sections of vector bundles form a fine sheaf. Hence supermanifolds are always split.*

In general it is hard to compute the obstruction class  $\text{ob}(X_n)$  or  $\text{ob}(X_n^0)$  except for small dimensions and small  $n$ . In [6], an interesting example is constructed as follows: assume that  $g > 1$ , then there is a morphism  $f : \mathcal{SM}_{g,1,0} \rightarrow \mathcal{SM}_{g,0,0}$  which is defined by the projection from the universal super-Riemann surfaces (SRS) to the base. Choose a SRS  $\mathcal{C}$  whose spin structure is denoted by  $\mathcal{E}$ , and an element  $\eta \in H^1(C, \mathcal{E}^\vee)$ , then  $\eta$  is an odd tangent vector at the point  $[\mathcal{C}] \in \mathcal{SM}_{g,0,0}(\mathbb{C})$ , which can be identified with a superaffine space  $\mathbb{A}^{0|1}$ , denoted by  $\mathbb{C}_\eta$ . Consider  $\mathcal{C}_\eta \equiv f^{-1}\mathcal{C}_\eta$ . It is a smooth superscheme of dimension  $(1|2)$  and is a first order odd deformation of  $\mathcal{C}$ . Now the obstruction class  $\text{ob}(C_{\eta,2})$  lives in the cohomology

$$H^1(\mathcal{C}, T_{\mathcal{C}} \otimes \wedge^2(\mathcal{E} \oplus \mathcal{O}_{\mathcal{C}})) \cong H^1(\mathcal{C}, \mathcal{E}^\vee). \quad (5.8)$$

An explicit computation in Lemma 3.5 of [6] shows that  $\text{ob}(\mathcal{C}_{\eta,2})$  is identified with  $\eta$  via this isomorphism, in particular, it is nontrivial if  $\eta \neq 0$ .

---

<sup>44</sup>What is called supermanifold in [6] is a special class of superspace defined in this work, and  $C^\infty$ -supermanifold in the sense of [6] is simply called supermanifold here.

In [6], Donagi and Witten used this example and a comparison between obstruction classes of an ambient superscheme and its smooth closed super subscheme (*loc. cit.* Corollary 2.10) to conclude that for  $g \geq 2$ , the even spin component  $\mathcal{SM}_{g,1,0}^{\text{ev}}$  is non-projected. Moreover, for  $g \geq 5$ , they construct a class of non-split super subschemes of supermoduli  $\mathcal{SM}_{g,0,0}$  to conclude that the supermoduli in these cases are non-projected. This example has an interesting consequence which is slightly stronger than non-projectedness.

**Proposition 5.3.** *Assume  $g \geq 2$ , then there is no projection  $\pi : \mathcal{SM}_{g,1,0}^{\text{ev}} \rightarrow \mathcal{M}_{g,1}$  such that the following diagram commutes*

$$\begin{array}{ccc} \mathcal{S}_{g,1,0}^{\text{ev}} & \xleftarrow{i} & \mathcal{SM}_{g,1,0}^{\text{ev}} \\ & \searrow f & \downarrow \pi \\ & & \mathcal{M}_{g,1}. \end{array}$$

*Proof.* Assume that there is such projection  $\pi : \mathcal{SM}_{g,1,0}^{\text{ev}} \rightarrow \mathcal{M}_{g,1}$  making the diagram commutative. Choose a spin curve  $\mathcal{C}$  of genus  $g$  with even spin structure  $\mathcal{E}$  such that  $H^1(\mathcal{C}, \mathcal{E}) = 0$ . Note that a generic choice of  $[\mathcal{C}]$  in  $\mathcal{S}_{g,0,0}^{\text{ev}}(\mathbb{C})$ <sup>45</sup> will satisfy  $H^1(\mathcal{C}, \mathcal{E}) = 0$  since the locus in  $\mathcal{S}_{g,0,0}^{\text{ev}}$  where  $H^1(\mathcal{C}, \mathcal{E}) = 0$  is open by semicontinuity of cohomology, and nonempty by explicit construction on a hyperelliptic curve<sup>46</sup>. Now consider the commutative diagram

$$\begin{array}{ccccc} \mathcal{S}_{g,1,0}^{\text{ev}} & & \xrightarrow{\quad} & & \mathcal{SM}_{g,1,0}^{\text{ev}} \\ & \searrow f & & \swarrow \pi & \downarrow \\ & & \mathcal{M}_{g,1} & & \\ & \downarrow & \downarrow & \downarrow & \downarrow \\ \mathcal{S}_{g,0,0}^{\text{ev}} & \xleftarrow{\quad} & & \xrightarrow{\quad} & \mathcal{SM}_{g,0,0}^{\text{ev}} \\ & \searrow f' & & & \downarrow \\ & & \mathcal{M}_{g,0}. & & \end{array}$$

Note that  $f$  and  $f'$  are étale coverings. Take an étale neighborhood  $U$  of  $[\mathcal{C}] \in \mathcal{M}_{g,0}(\mathbb{C})$ <sup>47</sup> such that  $f'^{-1}(U)$  is isomorphic to a disjoint union of  $U$ . Take the component of  $f'^{-1}(U)$  which contains the chosen spin structure, and call it  $U_1$ . Denote the base change of  $U$  (resp.  $U_1$ ) to  $\mathcal{M}_{g,1}$  (resp.  $\mathcal{S}_{g,1,0}^{\text{ev}}$ ) by  $V$  (resp.  $V_1$ ). Then, there is a commutative diagram

$$\begin{array}{ccc} V_1 & \xleftarrow{i} & \pi^{-1}(V) \\ & \searrow f|_{V_1} & \downarrow \pi \\ & & V. \end{array}$$

Since  $f|_{V_1}$  is an isomorphism, this diagram says that  $\pi^{-1}(V)$  is projected. Note that given a non-zero  $\eta \in H^1(\mathcal{C}, \mathcal{E}^\vee)$ , the superscheme  $\mathcal{C}_\eta$  is still a super subscheme of  $\pi^{-1}(V)$ , hence

<sup>45</sup> $\mathcal{S}_{g,0,0}^{\text{ev}}(\mathbb{C})$  denotes complex points of the stack  $\mathcal{S}_{g,0,0}^{\text{ev}}$ , i.e.  $\mathcal{C}$  is a spin curve over complex numbers.

<sup>46</sup>The divisor  $\mathfrak{b}_S^{(-1)}$  in the section 4 of [55], for example.

<sup>47</sup> $\mathcal{M}_{g,0}(\mathbb{C})$  denotes the complex points of the stack  $\mathcal{M}_{g,0}$ , i.e.  $\mathcal{C}$  is a genus  $g$  curve over complex numbers.

the same argument in Proposition 4.1 of [6] shows that  $\pi^{-1}(V)$  is not projected, which is a contradiction. This concludes the proof.  $\blacksquare$

### 5.1.1 Lifting Space of the Bosonic Truncation of a Supermanifold

As we have seen so far, we could find certain subspaces  $\mathcal{V}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  of the moduli stack of bordered spin-Riemann surfaces  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  (or more precisely  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ ) with the following properties

1.  $\mathcal{V}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  satisfies the BV QME;
2. Under a map in the homotopy category of BV algebras between chain complexes defined in (4.30),  $\mathcal{V}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})$  is mapped to the fundamental class of the stable-curve compactification of the moduli stack of punctured spin curves;

The question that we are interested to address is whether we can lift such solutions to  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  and then interpret the lifting as a solution to the BV QME in  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ ? We thus need to show that the lifting is unique. We investigate this question in the following.

In a precise language, we would like to understand the following question: given a smooth superscheme  $(X, \mathcal{O}_X)$ , and a smooth closed subscheme  $Y_{\text{bos}} \subset X_{\text{bos}}$ <sup>48</sup>, what is the deformation-obstruction of lifting  $Y_{\text{bos}}$  to a closed super subscheme  $Y \subset X$  of codimension  $(n|0)$ ? To simplify our discussion, we assume that  $X$  is a scheme.

Let us start with assuming that  $X_{\text{bos}}$  is affine and  $X$  is split, i.e. there is an isomorphism

$$\mathcal{O}_X \cong \mathcal{O}_{X_{\text{bos}}} \oplus \bigoplus_{i \geq 1}^m \bigwedge^i \mathcal{V}, \quad (5.9)$$

for a locally-free sheaf  $\mathcal{V}$  of rank  $m$ , and  $Y_{\text{bos}}$  is generated by  $\{f_1, f_2, \dots, f_n\} \subset \Gamma(X, \mathcal{O}_{X_{\text{bos}}})$ , a regular sequence of length  $n$ . Then there is a canonical lifting of  $Y_{\text{bos}}$ , namely by pulling back  $Y_{\text{bos}}$  along the projection  $X \rightarrow X_{\text{bos}}$ . This is equivalent to setting the ideal defining  $Y$  to be generated by the same set of generators  $\{f_1, f_2, \dots, f_n\}$ . Suppose that there is another lifting  $Y'$  defined by  $n$  even functions  $\{f'_1, f'_2, \dots, f'_n\} \subset \Gamma(X, \bigoplus_{i \geq 0} \bigwedge^{2i} \mathcal{V})$  such that  $f'_i - f_i \in \Gamma(X, \bigoplus_{i \geq 1} \bigwedge^{2i} \mathcal{V})$ . Moreover, two sets  $\{f'_1, f'_2, \dots, f'_n\}$  and  $\{f''_1, f''_2, \dots, f''_n\}$  generates the same ideal if and only if  $f''_i - f'_i \in \{f_1, f_2, \dots, f_n\} \cdot \Gamma(X, \bigoplus_{i \geq 1} \bigwedge^{2i} \mathcal{V})$ . Hence the liftings have one to one correspondence to the Hom set

$$\text{Hom}_{\mathcal{O}_{X_{\text{bos}}}} \left( \mathcal{I}_{Y_{\text{bos}}}, \bigoplus_{i \geq 1}^m \bigwedge^{2i} \mathcal{V}|_{Y_{\text{bos}}} \right) \quad (5.10)$$

where  $\mathcal{I}_{Y_{\text{bos}}}$  is the ideal defining  $Y_{\text{bos}}$ .

---

<sup>48</sup>From the definition 7.2,  $X_{\text{bos}}$  is the locally-ringed topological space  $(X, \mathcal{O}_X/\mathfrak{n}_X)$  of codimension  $n$ , where  $\mathfrak{n}_X$  is the ideal generated by the odd part  $\mathcal{O}_X^1$  of the sheaf  $\mathcal{O}_X$ .

If  $X_{\text{bos}}$  is not necessarily affine, then we need to lift  $Y_{\text{bos}}$  order by order, and we possibly encounter obstructions. Let  $X_k$  be the commutative space  $(X, \mathcal{O}_X^+ / (\mathcal{O}_X^-)^{2k+2})$ . From our previous discussion, we know that to lift  $Y_{\text{bos}}$  to a subspace of  $X$ , it suffices to lift it to a subspace of  $X_{\text{ev}}$ <sup>49</sup>, by lifting it to  $X_k$  recursively. Suppose that there is a lifting  $Y_k \subset X_k$ , we want to know the deformation-obstruction space of lifting it to  $X_{k+1}$ , this is completely classical, in fact we have:

**Proposition 5.4.** *There is an element*

$$\text{ob}(Y_k) \in H^1 \left( Y_{\text{bos}}, \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right), \quad (5.11)$$

whose vanishing is necessary and sufficient for the existence of lifting  $Y_{k+1}$  of  $Y_k$ .  $\mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}}$  is the normal bundle of  $Y_{\text{bos}}$  in  $X_{\text{bos}}$ . If it vanishes, then the space of lifting  $Y_{k+1}$  is a torsor under

$$\text{Hom}_{\mathcal{O}_{X_{\text{bos}}}} \left( \mathcal{I}_{Y_{\text{bos}}}, \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right) = H^0 \left( Y_{\text{bos}}, \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right). \quad (5.12)$$

*Proof.* This is essentially Theorem 6.4.5 of [56] with modifications. We will basically follow *loc. cit.*

Step (a): construction of  $\text{ob}(Y_k)$ . We have following diagram of coherent sheaves on  $X_{k+1}$  with exact row and columns

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \mathcal{I}_{Y_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V} & & \mathcal{I}_{Y_k} & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \bigwedge^{2k+2} \mathcal{V} & \longrightarrow & \mathcal{O}_{X_{k+1}} & \longrightarrow & \mathcal{O}_{X_k} \longrightarrow 0, \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} & & \mathcal{O}_{Y_k} & & \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

Consider the induced maps  $\alpha : \mathcal{I}_{Y_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V} \rightarrow \mathcal{O}_{X_{k+1}}$  and  $\beta : \mathcal{O}_{X_{k+1}} \rightarrow \mathcal{O}_{Y_k}$ , and we also have  $\text{im}(\alpha) \subset \ker(\beta)$ . Define  $\tilde{\mathcal{O}}_{X_{k+1}} := \ker(\beta) / \text{im}(\alpha)$  and we have short exact sequence of coherent sheaves on  $X_{k+1}$

$$0 \longrightarrow \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \longrightarrow \tilde{\mathcal{O}}_{X_{k+1}} \longrightarrow \mathcal{I}_{Y_k} \longrightarrow 0,$$

---

<sup>49</sup>The locally-ringed topological space  $(X, \mathcal{O}_X^0)$  is the **bosonic quotient** of  $X$ , denoted by  $X_{\text{ev}}$ . For more on this, see Section 7.1.



Similar to *loc. cit.* we need to check that this is in fact a sequence of  $\mathcal{O}_{X_k}$  sheaves. Since this is a local property and we can assume that  $X$  splits and  $\mathcal{V}$  is free. In this case  $X_k$  is isomorphic to  $X_{\text{bos}} \times \text{Spec } A_k$  for an Artinian ring  $A_k$  and this goes back to the situation of *loc. cit.* hence concludes the proof.

Step (b). It is enough to prove that extensions are in a natural bijection with splittings of the above short exact sequence. Assume we are given a splitting  $\xi : \mathcal{I}_{Y_k} \rightarrow \tilde{\mathcal{O}}_{X_{k+1}}$ , and let  $\mathcal{I}_{Y_{k+1}}$  be the preimage of  $\xi(\mathcal{I}_{Y_k})$  in  $\ker(\beta) \subset \mathcal{O}_{X_{k+1}}$ . We can then complete the H-shaped diagram to a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathcal{I}_{Y_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V} & \longrightarrow & \mathcal{I}_{Y_{k+1}} & \longrightarrow & \mathcal{I}_{Y_k} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \bigwedge^{2k+2} \mathcal{V} & \longrightarrow & \mathcal{O}_{X_{k+1}} & \longrightarrow & \mathcal{O}_{X_k} \longrightarrow 0, \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} & \longrightarrow & \mathcal{O}_{Y_{k+1}} & \longrightarrow & \mathcal{O}_{Y_k} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

and  $\mathcal{O}_{Y_{k+1}}$  is the required lifting. To show that it's generated by a regular sequence of length  $n$ , we can again assume that  $X$  splits and  $\mathcal{V}$  is free. In this case  $Y_{k+1}$  is flat over  $A_{k+1}$  (see *loc. cit.*) hence the statement follows from *loc. cit.* Lemma 9.3.4<sup>50</sup>. Conversely, given a lifting  $\mathcal{O}_{Y_{k+1}}$  we can build such commutative diagram as above by exactness and the subsheaf  $\mathcal{I}_{Y_{k+1}}/\text{im}(\alpha) \subset \ker(\beta)/\text{im}(\alpha)$  maps isomorphically to  $\mathcal{I}_{Y_k}$  hence gives a splitting of  $\tilde{\mathcal{O}}_{X_{k+1}} \rightarrow \mathcal{I}_{Y_k}$ .

Step (c). We have only shown that  $\text{ob}(Y_k) \in \text{Ext}_{\mathcal{O}_{X_{\text{bos}}}}^1(\mathcal{I}_{Y_{\text{bos}}}, \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}})$ . To proceed further we notice that there is a spectral sequence

$$\text{Ext}_{\mathcal{O}_{Y_{\text{bos}}}}^q \left( \text{Tor}_p^{\mathcal{O}_{X_{\text{bos}}}}(\mathcal{I}_{Y_{\text{bos}}}, \mathcal{O}_{Y_{\text{bos}}}), \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right) \implies_p \text{Ext}_{\mathcal{O}_{X_{\text{bos}}}}^{p+q} \left( \mathcal{I}_{Y_{\text{bos}}}, \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right),$$

which gives rise to a short exact sequence

$$\begin{aligned}
0 &\longrightarrow \text{H}^1 \left( Y_{\text{bos}}, \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right) \longrightarrow \text{Ext}_{\mathcal{O}_{X_{\text{bos}}}}^1 \left( \mathcal{I}_{Y_{\text{bos}}}, \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right) \\
&\longrightarrow \text{Hom}_{\mathcal{O}_{Y_{\text{bos}}}} \left( \text{Tor}_1^{\mathcal{O}_{X_{\text{bos}}}}(\mathcal{I}_{Y_{\text{bos}}}, \mathcal{O}_{Y_{\text{bos}}}), \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right) = \Gamma \left( Y_{\text{bos}}, \bigwedge^2 \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \bigwedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right).
\end{aligned}$$

Take an open covering  $\{U_i\} \rightarrow X_{\text{bos}}$  such that  $U_i$  are affine and split. From our previous discussions, we see that the obstruction vanishes on  $U_i$ , in particular the image of  $\text{ob}(Y_k)$  in

<sup>50</sup>There  $n$  is set to 1 but the proof presented there works for general  $n$ .

$\Gamma(Y_{\text{bos}}, \wedge^2 \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \wedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}})$ , denoted by  $\overline{\text{ob}(Y_k)}$ , vanishes on  $U_i$ . Since  $U_i$  is a covering, this implies that  $\text{ob}(Y_k)$  is trivial, hence  $\text{ob}(Y_k)$  lives in the subspace

$$H^1 \left( Y_{\text{bos}}, \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \wedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}} \right). \quad (5.13)$$

This concludes the proof.  $\blacksquare$

**Corollary 5.3.** *If  $H^1(Y_{\text{bos}}, \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \wedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}})$  vanishes for all  $k$ , then the set of lifting of  $Y_{\text{bos}}$  is a torsor under*

$$\text{Hom}_{\mathcal{O}_{X_{\text{bos}}}} \left( \mathcal{I}_{Y_{\text{bos}}}, \bigoplus_{i \geq 1}^m \wedge^{2i} \mathcal{V}|_{Y_{\text{bos}}} \right). \quad (5.14)$$

The case we are interested is when  $X$  is a supermanifold, in the sense of Remark 5.1. Above arguments also apply when  $X$  is a supermanifold, the case that we are interested in. In that case, the cohomology group  $H^1(Y_{\text{bos}}, \mathcal{N}_{Y_{\text{bos}}/X_{\text{bos}}} \otimes \wedge^{2k+2} \mathcal{V}|_{Y_{\text{bos}}})$  vanishes automatically since sections of vector bundles form a fine sheaf. So the space of liftings has the structure of an affine space, hence contractible. As a consequence, any two liftings can be connected by a homotopy.

We thus end-up with the conclusion we were looking: We have shown that

1. We can always find a solution to the BV QME in  $\mathcal{S}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  to  $\mathcal{SM}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ . The lifting is unique up to homology;
2. Precisely because the lifted solution is defined up to homology, we cannot consider the lifted solution as a solution to the BV QME in  $\mathcal{SM}_g(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}})/S^1 \wr S_{\mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$ .

## 5.2 Singular Homology on Superspaces

In this section, we discuss singular homology on superspaces. We mainly follow [57] and [58]. From now on we focus on cs-manifolds<sup>51</sup> (possibly with boundaries and corners). Consider the  $n$ -dimensional simplex  $\Delta^n$  and embed it into a cs-manifold  $\Delta^{n|m}$  defined by the trivial  $m$ -dimensional bundle on  $\Delta^n$  as its first order part.

**Definition 5.1** (Singular Superchains on Superspaces). *Fix a cs-manifold  $M$  of dimension  $(p|q)$  such that  $\mathcal{V} \equiv \mathcal{O}_M[1]$  is a complex vector bundle of dimension  $q$ . A singular supersimplex  $\sigma$  of dimension  $(n|m)$  on  $M$  is a morphism of superspaces  $\sigma : \Delta^{n|m} \rightarrow M$  such that the associated bundle map  $\underline{\mathbb{C}}^m \rightarrow \sigma^* \mathcal{V}$  is **injective**. Define the singular superchain complex valued in a ring  $R$ , denoted by  $C_{n|m}(M; R)$ , to be the free  $R$  module with basis singular supersimplices on  $M$  of dimension  $(n|m)$ . And also define the boundary map  $\partial_{n|m} : C_{n|m}(M; R) \rightarrow C_{n-1|m}(M; R)$  by*

$$\partial_{n|m} \sigma := \sum_{i=0}^n (-1)^i \partial_i \sigma$$

---

<sup>51</sup>Here 'cs' means complex-super but be aware that this is by definition a real manifold with a complex structure sheaf. Dimension  $(p|q)$  means real even dimension  $p$  and complex odd dimension  $q$ .

where  $\partial_i \sigma$  is the singular supersimplex induced by restriction of  $\sigma$  to sub-superspace  $\Delta^{n-1|m}$  of  $\Delta^{n|m}$ , which is topologically the  $i$ 'th boundary of  $\Delta^n$  with the same orientation as in the definition of singular chain complex.

The singular homology of  $M$  of fermionic degree  $m$  with coefficient ring  $R$ , denoted by  $H_{\bullet|m}(M; R)$  is defined to be the homology of the complex  $(C_{\bullet|m}(M; R), \partial_{\bullet|m})$ .

We then have the following result

**Proposition 5.5.** *Fix a cs-manifold  $M$  of dimension  $(p|q)$  such that  $\mathcal{V} \equiv \mathcal{O}_M[1]$  is a complex vector bundle of dimension  $q$ . We have isomorphism:*

$$H_{n|m}(M; R) \cong H_n(\text{Grass}_m(\mathcal{V}); R),$$

in particular,  $H_{n|0}(M; R) \cong H_{n|q}(M; R) \cong H_n(M_{\text{bos}}; R)$ . Here  $\text{Grass}_m(\mathcal{V})$  means the Grassmannian bundle of  $m$  dimensional subspaces of  $\mathcal{V}$ .

*Proof.* The proof goes similarly to the proof of Theorem 3 and 4 of [57]. We apply Corollary 1 of [57] and see that the chain complex  $(C_{\bullet|m}(M; R), \partial_{\bullet|m})$  is quasi-isomorphic to the chain complex  $(C'_{n,m}, d_n)$  of free  $R$  modules generated by  $n$ -simplices on  $M_{\text{bos}}$ , together with a choice of  $m$ -dimensional subbundle of pull-back of  $\mathcal{V}$  to the simplex, and  $d_n$  is alternating sum of restriction to boundaries. It is easy to see that this is exactly  $(C_{\bullet}(\text{Grass}_m(\mathcal{V}), R), \partial_{\bullet})$ . ■

**Remark 5.2.** *The definition of homology does not guarantee that for any morphism between cs-manifolds  $f : M \rightarrow N$ , there is a natural morphism of homologies. However, if the associated bundle map  $\mathcal{V}_M \rightarrow f^* \mathcal{V}_N$  is **injective**, then  $f$  indeed induces a natural morphism of homologies  $f_* : H_{n|m}(M; R) \rightarrow H_{n|m}(N; R)$ . In fact, in this case the isomorphism in Proposition 5.5 is natural, i.e. there is a commutative diagram*

$$\begin{array}{ccc} H_{n|m}(M; R) & \xrightarrow{\cong} & H_n(\text{Grass}_m(\mathcal{V}_M); R) \\ f_* \downarrow & & \downarrow f_* \\ H_{n|m}(N; R) & \xrightarrow{\cong} & H_n(\text{Grass}_m(\mathcal{V}_N); R). \end{array}$$

Using Proposition 5.5, we can prove the existence of solution to the BV QME in the moduli space of bordered  $N = 1$  super-Riemann surfaces. For that purpose, we can define a further generalization of the Kimura et al spaces for the moduli space of bordered  $N = 1$  super-Riemann surface. As asserted in Proposition 5.5, the homology group with top fermionic dimensions for these spaces are isomorphic to their reduced space, which is the generalization of the Kimura et al spaces for bordered spin-Riemann surface we introduced in Section 4.4, the proof for the existence of solution to the BV QME in the moduli space of bordered  $N = 1$  super-Riemann surfaces goes through without change. We will not attempt to reproduce that proof here. Instead, we present a shorter and more elegant proof for the existence of such solutions.

### 5.3 Proof of the Existence of Solution

In this section we give the proof for the existence of solution to the BV QME in  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ . Consider the following complex

$$\mathcal{F}(\mathcal{SM}) \equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{SC}_* \left( \frac{\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})}{S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}} \right), \quad (5.15)$$

where  $\mathcal{SC}_*$  is the functor of normalized singular simplicial superchains of top fermionic dimension with coefficients in any field containing  $\mathbb{Q}$ . This complex has the structure of a dg algebra. As usual the product comes from the disjoint union of surfaces, which gives  $\mathcal{F}(\mathcal{SM})$  an algebra structure, and the differential comes from the boundary operation on singular superchains, as defined in Definition 5.1. We would like to give  $\mathcal{F}(\mathcal{SM})$  the structure of a BV algebra. For that purpose, we define the following gluing operations of boundaries: let  $\tilde{\Delta}_s^{\text{NS}}$  (respectively  $\tilde{\Delta}_s^{\text{R}}$ ) be the operation of gluing two NS (respectively, R) boundaries with a full  $S^1$  twist. There is one additional complication in defining  $\tilde{\Delta}_s^{\text{R}}$ , namely we let it increase the odd dimension of the superchain by 1. The question is that why we can do that for R boundaries while we do not have this freedom for NS boundaries? From Definition 4.4 of R boundaries, they are isomorphic to  $S^1 \times \mathbb{R}^{0|1} \otimes_{\mathbb{R}} \mathbb{C}$ , and as such there is a translation symmetry along the only complex odd direction of  $\mathbb{R}^{0|1} \otimes_{\mathbb{R}} \mathbb{C}$ . This also makes sure that while we glue R boundaries, the resulting surface does have correct number of global fermionic parameters. From Definition 4.2 of NS boundaries there is no such freedom.

We would like to consider the operation of gluing as an operation acting on singular superchains. For this purpose, consider a singular superchain associated to  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \equiv \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ , which we denote by  $C_{m|d_f}(\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}); \mathbb{F})$ . Note that  $d_f = 2g - 2 + \mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}/2$  is the top fermionic dimension of the part of  $\mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  coming from connected genus- $g$  surfaces. This singular superchain is generated by the  $(m|d_f)$ -singular supersimplices  $\sigma_s^{m|d_f} : \Delta^{m|d_f} \longrightarrow \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , i.e. a morphism of superspaces  $\sigma_s^{m|d_f} : \Delta^{m|d_f} \longrightarrow \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  with properties given in Definition 5.1. We would like to define the operations  $\tilde{\Delta}_s^{\text{NS}}$  and  $\tilde{\Delta}_s^{\text{R}}$  on the singular superchains. We thus consider the operations corresponding to  $\tilde{\Delta}_s^{\text{NS}}$  and  $\tilde{\Delta}_s^{\text{R}}$  for all values of  $\mathbf{n}_{\text{NS}}$  and  $\mathbf{n}_{\text{R}}$  which are acting on singular superchains and define<sup>52</sup>

$$\begin{aligned} \Delta_s^{\text{NS}} : \mathcal{F}^{\mathfrak{b}|\mathfrak{f}}(\mathcal{SM}) &\longrightarrow \mathcal{F}^{\mathfrak{b}+1|\mathfrak{f}}(\mathcal{SM}), \\ \Delta_s^{\text{R}} : \mathcal{F}^{\mathfrak{b}|\mathfrak{f}}(\mathcal{SM}) &\longrightarrow \mathcal{F}^{\mathfrak{b}+1|\mathfrak{f}+1}(\mathcal{SM}), \end{aligned} \quad (5.16)$$

where superscript  $\mathfrak{b}|\mathfrak{f}$  denotes the  $\mathbb{Z}_2$ -graded homological degree of the respective superchains. It is clear that this operation is compatible with gluing, i.e.  $[\Delta_s, d_s] = 0$ , where  $\Delta_s \equiv \Delta_s^{\text{NS}} + \Delta_s^{\text{R}}$ , and  $d_s \equiv \partial$  is the boundary map acting on singular superchains, as defined in Definition 5.1. We denote this BV algebra as  $\mathbf{BV}(\mathcal{SM}) \equiv (\mathcal{F}(\mathcal{SM}); d_s, \Delta_s)$ .

<sup>52</sup>These operations are not canonically defined. See Footnote 6 on page 5 of [26].

$$\begin{array}{ccc}
\mathbb{SM} & \xrightarrow{\mathcal{SC}_*} & \mathbf{BV}(\mathbb{SM}) \\
\downarrow \pi & & \downarrow \varphi \\
\mathbb{S} & \xrightarrow{\mathcal{C}_*} & \mathbf{BV}(\mathbb{S}).
\end{array}$$

**Figure 10:** The commutative diagram associated to the map  $\varphi$  between the BV algebras  $\mathbf{BV}(\mathbb{SM})$  and  $\mathbf{BV}(\mathbb{S})$ .  $\mathcal{SC}_*$  associates singular superchains to  $\mathbb{SM}$ ,  $\mathcal{C}_*$  associates singular chains to  $\mathbb{S}$ , and  $\pi$  is the map to the underlying bosonic space that sends  $\mathbb{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  to  $\mathbb{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ . The commutative diagram basically tells us that projecting  $\mathbb{SM}$  to  $\mathbb{S}$ , and then taking singular chains is the same as associating singular superchains with top fermionic degree to  $\mathbb{SM}$ , and then projecting onto underlying singular chains.

Our next task is to define a map between the BV algebras  $\mathbf{BV}(\mathbb{SM})$  and  $\mathbf{BV}(\mathbb{S}) \equiv (\mathcal{F}(\mathbb{S}); d, \Delta)$ , the BV associated to the moduli space of bordered spin-Riemann surfaces, and show that it is a quasi-isomorphism. We define the map  $\varphi : \mathbf{BV}(\mathbb{SM}) \longrightarrow \mathbf{BV}(\mathbb{S})$  by sending any singular superchain in  $\mathcal{F}(\mathbb{SM})$  to its underlying singular chain,  $d_{\mathbb{S}}$  to  $d$ , and  $\Delta_{\mathbb{S}}$  to  $\Delta$ . Proposition 5.5 asserts that the  $\mathbb{Z}_2$ -graded homology groups of the singular supercomplex associated to  $\mathbb{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  with top fermionic degree are isomorphic to the singular homology groups associated to  $\mathbb{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \equiv \mathbb{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})/S^1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}}$ . This shows that the associated singular chain complexes are quasi-isomorphic. Since this holds for all values of  $\mathbf{n}_{\text{NS}}$  and  $\mathbf{n}_{\text{R}}$ , the BV algebras  $\mathbf{BV}(\mathbb{SM})$  and  $\mathbf{BV}(\mathbb{S})$  are quasi-isomorphic. If we define

$$\mathbb{S} = \bigcup_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathbb{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \quad \mathbb{SM} = \bigcup_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathbb{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \quad (5.17)$$

there is a commutative diagram which is shown in Figure 10. We thus have the following result

**Proposition 5.6.** *The BV algebras  $\mathbf{BV}(\mathbb{SM})$  and  $\mathbf{BV}(\mathbb{S})$  associated to moduli spaces of bordered spin-Riemann surfaces and moduli spaces of bordered  $\mathcal{N} = 1$  super-Riemann surfaces are quasi-isomorphic.*

We already mentioned that the set of homotopy-equivalent solutions of the BV QME for quasi-isomorphic BV algebras are isomorphic. Combining this result together with Proposition 5.6, we conclude that

**Corollary 5.4** (The Existence of Solution to the BV QME in  $\mathbf{BV}(\mathbb{SM})$ ). *There exists a solution to the BV QME in  $\mathbf{BV}(\mathbb{SM})$ .*

With this result at hand, we conclude this section.

## 6 Existence of Heterotic-String and Type-II-Superstring Vertices

In this section, we will prove the existence of heterotic-string and type-II-superstring field theory vertices, i.e. we prove the existence of subspaces in the product of stacks  $\mathcal{M}_L$  and  $\mathcal{M}_R$

that respectively parameterize left-moving and right-moving modes of the theory. The worldsheet  $\mathcal{R}$  of heterotic-string or type-II-superstring theories can be described by an embedding of the following form<sup>53</sup>

$$\mathcal{R} \hookrightarrow \mathcal{R}_L \times \mathcal{R}_R, \quad (6.1)$$

where  $\mathcal{R}_L$  is either an ordinary Riemann surface, in the case of heterotic-string theory, or an  $\mathcal{N} = 1$  super-Riemann surface, in the case of type-II-superstring theories, and  $\mathcal{R}_R$  is an  $\mathcal{N} = 1$  super-Riemann surface. In the following, we denote by  $\pi$  the map that forgets the spin structure, by  $X_{\text{red}}$  the reduced space of the supermanifold  $X$ , and by  $X_{\overline{\text{red}}} \equiv X_{\pi \circ \text{red}}$  the reduced space of the supermanifold together with the forgetting of spin structure of either  $X$  or the geometric objects that  $X$  parametrizes. For an ordinary surface  $\mathcal{R}$ , we denote a surface which has the complex-conjugate complex structure by  $\overline{\mathcal{R}}$ .

## 6.1 Introduction

In the current understanding of superstring field theory, one must make two choices to be able to formulate such a theory

1. **a choice of worldsheet superconformal field theory:** this choice is essential to be able to define interaction vertices of superstring field theory. This choice begs one of the most important question in the formulation of superstring field theory, and string theory in general, i.e. the question of *background independence*. Since the results of this paper do not have any implication for this choice, the reader can directly go to [a choice of string vertex](#). We however briefly describe the problem. There are two versions of this question.

- **Background-Independence (Weak Version):** In this version, the question is the following: *consider two arbitrary 2d superconformal field theories which live in the same connected component of the space of 2d superconformal field theories with appropriate central charges and supersymmetry in the space of all such superconformal field theories. If we use these two theories to formulate two superstring field theories, to which extent the physical quantities computed using these two theories are sensitive to the choice of 2d superconformal field theory?* Since this is a kind of gauge choice, for the consistency of the whole theory, the physical quantities must be independent of this choice. For the case of closed bosonic-string field theory is has been shown that two string field theories formulated using two infinitesimally-different conformal field theories related to each other by some exactly marginal perturbation are related to each other by a field redefinition [52, 59]. In the case of closed superstring field theories formulated in the picture-changing formalism, a similar result has been established recently [60]. There exist stronger results for

---

<sup>53</sup>For more on this viewpoint of worldsheets in string theory, see section 5.5 of [10].

open bosonic-string field theory, i.e. the field redefinition for even two finitely-separated backgrounds belonging to the same connected component can be constructed [61, 62]. Some other references on the issue of background-independence in string theory are [63–70]

- **Background-Independence (Strong Version):** In general, superconformal field theories are classical backgrounds for the formulation of superstring theory. However, the full quantum theory should be formulated using an arbitrary background. By this we mean more than just starting from a string field theory constructed on a conformal background and then constructing string field theory around a new background by giving VEV to original string fields. Such a theory is a gauge-invariant theory but it explicitly depends on the original conformal background and as such cannot be the solution to the problem of formulating theory on a generic background. String field theory suggest that such a background can be constructed by starting with a conformal background, and then consider the space of all, not necessarily marginal, deformations of this conformal background. This suggest that the space of quantum string background is the space of all  $2d$  supersymmetric field theories possibly with some yet unknown features [71]. If this is the case, then the strong version of the concept of background-independence can be phrased as follows: *consider two arbitrary  $2d$  supersymmetric field theories with some specific properties. If we use these two theories to formulate two superstring field theories, to which extent the physical quantities computed using these two theories are sensitive to the choice of  $2d$  supersymmetric field theory?* As far as we know, this question and more generally the question of string theory on nonconformal backgrounds have not been touched at all. The only attempt in this direction that we are aware of is the work of Zwiebach on the closed bosonic-string field theory on nonconformal backgrounds [71].

We also note that there is more to the concept of background-independence in string theory. This motivated the definition of the concept of *string background* rather than a conformal background. An example of a deformation of string background without deforming the underlying conformal field theory is provided by the zero-momentum ghost-dilaton state, the component of the dilaton state constructed by the action of operators from the ghost sector [72–74]. It can be shown that the deformation of a CFT by a zero-momentum ghost-dilaton state is a trivial deformation of the CFT, i.e. it is the old CFT represented in a new basis of states. Despite this fact, such deformation provides a new string background in the sense defined in section 7.2 of [72]. This shows that the space of deformation of string background is larger than the space of all  $2d$  conformal field theories with appropriate properties. Therefore, the independence of the physical quantities from such deformation must also be part of the proof of the weak version of the background-independence. For the case of zero-momentum ghost-dilaton, and also the matter dilaton and the true dilaton, constructed by a combination



of matter and ghost dilatons<sup>54</sup>, in bosonic-string field theory, it is shown that such shifts change the dimensionless string coupling constant, and thus the string background, but the resulting bosonic-string field theories are related by a field redefinition [74]. Also, as far as we know, the other possible deformations of the quantum string background, similar to the ghost-dilaton deformation of the classical string background, has not been explored in the literature.

2. **a choice of string vertex:** string vertices are regions inside the moduli space of surfaces on which the relevant theories are defined. They must satisfy the BV QME. This guarantees the gauge-invariance of the action. Then, the remaining region of the moduli space is constructed by gluing of surfaces with lower genus and/or number of punctures, which can be interpreted as Feynman diagrams in string field theory. This region together with the region corresponding to the string vertices must provide a single cover of the moduli space. This essential requirement, related to the modular invariance, is demanded by the unitarity of the theory [12, 75–77]. Regarding this choice, there is a similar question. This question can be phrased as follows. *Let us choose two string vertices. If we formulate two string field theory using these two choices, to which extent the physical quantities computed in these two theories are sensitive to the choice of string vertex?* Again, the consistency of the theory demands that the physical results must be independent of this choice. It has been shown in [78] and it has been further elaborated in [52, 53] that two closed bosonic-string field theories constructed using two infinitesimally-different vertices give the same gauge-fixed action for the bosonic-string field theory upon using two different gauge-fixing conditions. Similar results for finitely-different closed bosonic-string vertices is considered in [79]. As far as we aware, similar results for heterotic-string and type-II-superstring vertices are not considered in the literature.

To construct string vertices, we can either work with *punctured* surfaces and consider local (super)conformal coordinates, or we can work with *bordered* surfaces. The gluing of local (super)conformal coordinates around the punctures correspond to gluing of boundaries. In this paper, we work with *bordered* surfaces to characterize string vertices. We consider the two cases of heterotic-string and type-II superstring theory separately in Sections 6.2 and 6.3, respectively.

## 6.2 The Case of Heterotic-String Field Theory Vertices

Heterotic-string theories are described by a worldsheet theory with  $\mathcal{N} = (0, 1)$  supersymmetry. The notation means that the right-moving sector has one right-moving supersymmetry, and the left-moving sector is similar to the left-moving sector of the bosonic-string theory.  $\mathcal{R}$  is a smooth submanifold inside  $\mathcal{R}_L^H \times \mathcal{R}_R^H$ , whose dimension is  $2|1$ , and whose reduced space  $\mathcal{R}_{\text{red}}$  is taken to be such that the bosonic moduli of  $\mathcal{R}_{R;\text{red}}^H$  is the complex conjugate of the moduli

---

<sup>54</sup>For more on the definition of these states see section 5 of [74].



of  $\mathcal{R}_L^H$  (note that  $\mathcal{R}_L$  does not have fermionic moduli), and there is no condition on the spin structure. This means that  $\mathcal{R}_{\text{red}}$  is diagonal in  $\mathcal{R}_L^H \times \mathcal{R}_{R;\text{red}}^H$ . Worldsheets that appear in the definition of  $g$ -loop heterotic-string vertices with  $n_{\text{NS}}$  NS boundary components and  $n_R$  R boundary components are parametrized by a cycle  $\Gamma^H$  inside  $\mathcal{M}_L^H \times \mathcal{M}_R^H$ .  $\mathcal{M}_L^H$  is the moduli space of genus- $g$  bordered ordinary Riemann surfaces with  $n_{\text{NS}} + n_R$  boundary components and  $\mathcal{M}_L^H$  is the moduli space of genus- $g$   $\mathcal{N} = 1$  bordered super-Riemann surfaces with  $n_{\text{NS}}$  NS boundary components and  $n_R$  R boundary components. From what we explained above, the choice of  $\Gamma^H$  is such that  $\Gamma_{\text{red}}^H$  is diagonal in  $(\mathcal{M}_L^H \times \mathcal{M}_R^H)_{\text{red}} = \mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H$ .  $\Gamma^H$  can be constructed as follows. One first consider a diagonal space  $\Gamma_{\text{red}}^H$  in  $\mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H$  such that the complex structure of  $\mathcal{R}_L^H$  parametrized by  $\mathcal{M}_L^H$  and the complex structure of  $\mathcal{R}_{R;\text{red}}^H$  parameterized by  $\mathcal{M}_{R;\text{red}}^H$  are complex conjugate of each other. One then thicken  $\Gamma_{\text{red}}^H$  in the fermionic directions to get  $\Gamma^H$ . Since the thickening is arbitrary, the choice of  $\Gamma^H$  is unique up to homolog, as we have explained in Section 5.1.1. As it has been explained in section 5.6 of [10], the result of the integration over  $\Gamma^H$  is independent of  $\Gamma^H$ , and it is only sensitive to  $\Gamma_{\text{red}}^H$ <sup>55</sup>. However, the definition of genus- $g$  heterotic-string vertices with some number of boundary components requires more elaboration since  $\Gamma^H$  must furthermore satisfies the BV QME. If we were dealing with  $\mathcal{M}_L^H \times \mathcal{M}_R^H$ , i.e. we were letting left and right moduli vary independently, then we could construct a solution of the BV quantum master equation by considering the diagonal product of the solutions of the BV QME in  $\mathcal{M}_L$  and  $\mathcal{M}_R$ . On the other hand, it is not always clear that if we start from a diagonal solution of the BV QME in  $\mathcal{M}_L \times \mathcal{M}_{R;\text{red}}$  and thicken that solution in the fermionic directions, we land on a solution of the BV QME in the product  $\mathcal{M}_L \times \mathcal{M}_R$  since the thickening is defined up to homology. We thus need to consider the problem of the existence of the solution of the BV quantum master equation in  $\mathcal{M}_L^H \times \mathcal{M}_R^H$  such that once we reduce that solution to  $(\mathcal{M}_L^H \times \mathcal{M}_R^H)_{\text{red}}$ , the reduced solution parametrizes irreducible surfaces  $\mathcal{R}_{\text{red}}^H$  which are embedded diagonally inside  $(\mathcal{R}_L^H \times \mathcal{R}_R^H)_{\text{red}}$ , i.e. those surfaces in  $(\mathcal{R}_L^H \times \mathcal{R}_R^H)_{\text{red}}$  whose left-moving complex structure is complex conjugate of the right-moving complex structure. Thus we call  $\mathcal{V}_g^H(n_{\text{NS}}, n_R) \equiv \Gamma^H$  the genus- $g$  heterotic-string field theory with  $n_{\text{NS}}$  NS boundary components and  $n_R$  R boundary components if it satisfies the following requirements

1. The complex dimension of  $\mathcal{V}_g^H(n_{\text{NS}}, n_R)$  is  $(3g - 3 + n_{\text{NS}} + n_R)|2g - 2 + n_{\text{NS}} + n_R/2)$ ;
2.  $\mathcal{V}_g^H(n_{\text{NS}}, n_R)$ , as a subspace of  $\mathcal{M}_L \times \mathcal{M}_R$ , satisfies the BV quantum master equation;
3.  $(\mathcal{V}_g^H(n_{\text{NS}}, n_R))_{\text{red}}$  is embedded diagonally in  $(\mathcal{M}_L \times \mathcal{M}_R)_{\text{red}}$ , where by diagonal embedding, we mean all points in  $(\mathcal{M}_L \times \mathcal{M}_R)_{\text{red}}$  of the form  $(\mathcal{R}, \overline{\mathcal{R}})$ .

---

<sup>55</sup>To be more precise, suppose that the ideal sheaf of  $\mathcal{M}_L^H \times \mathcal{M}_R^H$  generated by fermionic directions is  $\mathcal{I}$ , then deformation theory implies that the space of all possible lifting of  $\Gamma_n \subset \mathcal{M}_L^H \times \mathcal{M}_R^H / \mathcal{I}^n$  to  $\Gamma_{n+1} \subset \mathcal{M}_L^H \times \mathcal{M}_R^H / \mathcal{I}^{n+1}$  is an affine space modeled on some linear space, hence contractible. This means that for any two liftings  $\Gamma$  and  $\Gamma'$ , there is a homotopy connecting them.

To construct such a solution, we proceed as follows<sup>56</sup>. We introduce the maps  $f_L^H : \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \longrightarrow \mathcal{M}_L^H$  defined by forgetting the spin structure, i.e.

$$f_L^H(\mathcal{R}^\varepsilon) \equiv \mathcal{R}, \quad \forall \mathcal{R}^\varepsilon \in \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}), \quad (6.2)$$

where  $\mathcal{R}$  is the underlying bosonic surface, and  $f_R^H \equiv J : \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \longrightarrow \mathcal{M}_{R;\text{red}}^H$ , where  $J$  is the operation of anti-holomorphic involution which sends a spin-Riemann surface  $\mathcal{R}^\varepsilon$  with a specific complex structure to a spin-Riemann surface with complex-conjugate complex structure  $\overline{\mathcal{R}}^\varepsilon$ . Note that the complex-conjugate spin structure  $\bar{\varepsilon}$  is defined such that  $\bar{\varepsilon}^{\otimes 2} \cong \bar{\omega}$ , where  $\bar{\omega}$  can be identified with the antiholomorphic cotangent bundle of  $\mathcal{R}^\varepsilon$ . Furthermore, these maps can be restricted to  $\mathcal{S}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .

Using these maps, we can define

$$f^H \equiv f_L^H \times f_R^H : \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \longrightarrow \mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H. \quad (6.3)$$

We then have

$$(\pi \circ f^H)(\mathcal{R}^\varepsilon) = \pi(f^H(\mathcal{R}^\varepsilon)) = \pi(\mathcal{R}, \overline{\mathcal{R}}^\varepsilon) = (\mathcal{R}, \overline{\mathcal{R}}). \quad (6.4)$$

This shows that the image of a spin Riemann surface  $\mathcal{R}^\varepsilon$  under  $f$  after reduction and forgetting of the spin structure is embedded diagonally in  $(\mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H)$  such that the left and right complex structures are complex conjugate of each other. To proceed further, we define the following space

$$(\mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H)^{\mathbf{D}} \equiv f^H(\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})) = \left\{ (\mathcal{R}, \overline{\mathcal{R}}^\varepsilon) \in (\mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H) \right\}, \quad (6.5)$$

i.e. all pairs  $(\mathcal{R}, \overline{\mathcal{R}}^\varepsilon)$  of a surface  $\mathcal{R}$  with a specific complex structures, and the surface  $\overline{\mathcal{R}}^\varepsilon$  whose complex structure is complex-conjugate of  $\mathcal{R}$  together with a choice of spin structure compatible with the complex-conjugate complex structure. The superscript  $\mathbf{D}$  stands for diagonal. The dimension of this space, for genus- $g$  surfaces with  $\mathbf{n}_{\text{NS}}$  NS boundary components, and  $\mathbf{n}_{\text{R}}$  R boundary components, is  $6g - 6 + 2\mathbf{n}_{\text{NS}} + 2\mathbf{n}_{\text{R}}$ , the same is as the space of spin-Riemann surfaces with the same signature. We also consider the thickening of this space in all fermionic dimensions, and denote the resulting space by  $(\mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H)^{\mathbf{D}}$ . Note that

$$\mathcal{M}_L^H = \mathcal{M}(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}) = \bigcup_{g=0}^{\infty} \mathcal{M}_g(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}), \quad \mathcal{M}_{R;\text{red}}^H = \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = \bigcup_{g=0}^{\infty} \mathcal{S}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}),$$

and the thickening is done for all values of  $(g; \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  in  $2g - 2 + \mathbf{n}_{\text{NS}} + \frac{\mathbf{n}_{\text{R}}}{2}$  fermionic directions. We can thus define the following chain complex

$$\mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_{R;\text{red}}^H)^{\mathbf{D}}) \equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{C}_* \left( \frac{(\mathcal{M}_L^H(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}) \times \mathcal{M}_{R;\text{red}}^H(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))^{\mathbf{D}}}{S_L^H \times S_R^H} \right), \quad (6.6)$$

---

<sup>56</sup>Note that for heterotic-string theories  $\mathcal{M}_L^H = \mathcal{M}(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}})$ ,  $\mathcal{M}_R^H = \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , and  $\mathcal{M}_{R;\text{red}}^H = \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .

where

$$S_L^H \equiv S_1 \wr S_{n_{NS}+n_R}, \quad S_R^H \equiv S_1^{n_{NS}+n_R} \wr S_{n_{NS}, n_R}, \quad (6.7)$$

$\mathcal{M}_L^H(n_{NS} + n_R) \equiv \mathcal{M}(n_{NS} + n_R)$ , and  $\mathcal{M}_R^H(n_{NS}, n_R) \equiv \mathcal{SM}(n_{NS}, n_R)$ .  $S_L^H$ , respectively  $S_R^H$ , acts on  $\mathcal{M}_R^H$ , respectively  $\mathcal{M}_L^H$ , trivially. We also associate a superchain complex to  $(\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}$ , defined by thickening  $(\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}_{\text{red}}$  in all fermionic directions,

$$\mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}) \equiv \bigoplus_{n_{NS}, n_R} \mathcal{SC}_* \left( \frac{(\mathcal{M}_L^H(n_{NS} + n_R) \times \mathcal{M}_R^H(n_{NS}, n_R))^{\mathbf{D}}}{S_L^H \times S_R^H} \right), \quad (6.8)$$

where  $\mathcal{SC}_*$  is the functor of normalized singular simplicial superchains with coefficients in any field containing  $\mathbb{Q}$  which associates singular simplicial superchains with top fermionic dimension.

The crucial property of the map  $f^H$  defined in (6.3) is that it is compatible with the gluing operation, i.e. the operation of gluing and mapping with  $f^H$  commute and there is a commutative diagram. We can thus define the following induced map

$$F_{f^H}^H : \mathcal{F}(\mathcal{S}) \longrightarrow \mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}_{\text{red}}). \quad (6.9)$$

If  $\mathcal{F}(\mathcal{M}_L^H \times \mathcal{M}_R^H)$  is the complex associated to the product  $\mathcal{M}_L^H \times \mathcal{M}_R^H$ , it has naturally the structure of a differential-graded algebra where the differential  $d^H \equiv d_L^H \oplus d_R^H$  is the left  $d_L^H$  and right  $d_R^H$  boundary operations on left and right singular chains, and the product is the disjoint union of left and right surfaces.  $\mathcal{F}(\mathcal{M}_L^H \times \mathcal{M}_R^H)$  can be turned into a BV algebra by the usual twist-swing operation  $\Delta_{LR}^H \equiv \Delta_L^H \oplus \Delta_R^H$ , where  $\Delta_L^H(\Delta_R^H)$  is the gluing operation defined on the complex  $\mathcal{F}(\mathcal{M}_L^H)$  ( $\mathcal{F}(\mathcal{M}_R^H)$ ).  $\mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}_{\text{red}})$  inherits this BV-algebra structure. By construction,  $F_{f^H}^H$  is a homomorphism of BV algebras, and thus maps solutions of the BV QME onto each other. Therefore, having a solution in  $\mathcal{F}(\mathcal{S})$  whose existence we have proven, we have a solution in  $\mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}_{\text{red}})$ . We have thus constructed a subspace of  $(\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}_{\text{red}}$  that satisfies the BV QME. To construct the heterotic-string vertices, we consider the following maps of BV algebras

$$\widehat{F}_{f^H}^H : \mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}_{\text{red}}) \longrightarrow \mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}}). \quad (6.10)$$

Since the BV algebra associated to singular superchains with top fermionic dimensions is quasi-isomorphic to the BV algebra of associated bosonic chains,  $\widehat{F}_{f^H}^H$  is a quasi-isomorphism of BV algebras. Again using the fact that quasi-isomorphic BV algebras have the same set of homotopy-classes of solutions of the BV QME, we conclude that there is a subspace of  $(\mathcal{M}_L^H(n_{NS} + n_R) \times \mathcal{M}_R^H(n_{NS}, n_R))^{\mathbf{D}}$ , whose genus- $g$  contribution we denote by  $\mathcal{V}_g^H(n_{NS}, n_R)$ , which satisfies the BV QME.  $\mathcal{V}_g^H(n_{NS}, n_R)$  is a genus- $g$  heterotic-string field theory vertex with  $n_{NS}$  boundary components and  $n_R$  boundary components since

1. its complex dimension is by construction  $(3g - 3 + n_{NS} + n_R|2g - 2 + n_{NS} + n_R/2)$ ;

2.  $\mathcal{V}_g^H(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_R)$ , as a subspace of  $\mathcal{M}_L^H \times \mathcal{M}_R^H$ , satisfies the BV QME;
3. by construction,  $(\mathcal{V}_g^H(\mathfrak{n}_{\text{NS}}, \mathfrak{n}_R))_{\text{red}}^{\overline{\phantom{x}}}$  is embedded diagonally in  $(\mathcal{M}_L \times \mathcal{M}_R)_{\text{red}}^{\overline{\phantom{x}}}$ .

We thus have the following

**Theorem 6.1** (The Existence of Heterotic-String Vertices). *There exists a solution to the BV QME in  $\mathcal{F}((\mathcal{M}_L^H \times \mathcal{M}_R^H)^{\mathbf{D}})$ . The genus- $g$  part is a genus- $g$  heterotic-string field theory vertex with  $\mathfrak{n}_{\text{NS}}$  NS and  $\mathfrak{n}_R$  R boundary components.*

### 6.3 The Case of Type-II-Superstring Field Theory Vertices

We now turn to the case of type-II-superstring theories. The worldsheet theory has  $\mathcal{N} = (1, 1)$  supersymmetry. The notation means that the right-moving sector has one right-moving supersymmetry, and the left-moving sector has one left-moving supersymmetry. The worldsheet  $\mathcal{R}$  is a smooth submanifold inside  $\mathcal{R}_L^{\text{II}} \times \mathcal{R}_R^{\text{II}}$ , whose dimension is  $2|2$ , and whose reduced space  $\mathcal{R}_{\text{red}}$  is taken to be such that the bosonic moduli of  $\mathcal{R}_{R;\text{red}}^{\text{II}}$  is the complex conjugate of the bosonic moduli of  $\mathcal{R}_{L;\text{red}}^{\text{II}}$ , and we do not assume any relation between fermionic moduli. This means that  $\mathcal{R}_{\text{red}}^{\text{II}}$  is diagonal in  $\mathcal{R}_{L;\text{red}}^{\text{II}} \times \mathcal{R}_{R;\text{red}}^{\text{II}}$ . Worldsheets that appear in the definition of  $g$ -loop type-II-superstring vertices with  $\mathfrak{n}_{\text{SNS}}$  SNS boundary components,  $\mathfrak{n}_{\text{NSR}}$  NSR boundary components,  $\mathfrak{n}_{\text{RNS}}$  boundary components, and  $\mathfrak{n}_{\text{RR}}$  RR boundary components are parametrized by a cycle  $\Gamma^{\text{II}}$  inside  $\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$ .  $\mathcal{M}_L^{\text{II}}$  and  $\mathcal{M}_R^{\text{II}}$  are the moduli space of genus- $g$   $\mathcal{N} = 1$  bordered super-Riemann surfaces with  $\mathfrak{n}_{\text{NS}}$  NS boundary components and  $\mathfrak{n}_R$  R boundary components. Again from what we explained so far, the choice of  $\Gamma^{\text{II}}$  is such that  $\Gamma_{\text{red}}^{\text{II}}$  is diagonal in  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})_{\text{red}}^{\overline{\phantom{x}}} = \mathcal{M}_{L;\text{red}}^{\text{II}} \times \mathcal{M}_{R;\text{red}}^{\text{II}}$ .  $\Gamma^{\text{II}}$  can be constructed as follows. One first consider a diagonal space  $\Gamma_{\text{red}}^{\text{II}}$  in  $\mathcal{M}_{L;\text{red}}^{\text{II}} \times \mathcal{M}_{R;\text{red}}^{\text{II}}$  such that the complex structure of  $\mathcal{R}_L^{\text{II}}$  parametrized by  $\mathcal{M}_{L;\text{red}}^{\text{II}}$  and the complex structure of  $\mathcal{R}_R^{\text{II}}$  parameterized by  $\mathcal{M}_{R;\text{red}}^{\text{II}}$  are complex conjugate of each other. One then thicken  $\Gamma_{\text{red}}^{\text{II}}$  in the fermionic directions to get  $\Gamma^{\text{II}}$ . Thickening is arbitrary, the choice of  $\Gamma^{\text{II}}$  is unique up to homology, the result of the integration over  $\Gamma^{\text{II}}$  is independent of  $\Gamma^{\text{II}}$ , and it is only sensitive to  $\Gamma_{\text{red}}^{\text{II}}$ . Again, the definition of genus- $g$  type-II-superstring vertices with some number of boundary components requires more elaboration since  $\Gamma^{\text{II}}$  must furthermore satisfies the BV QME. If we were dealing with  $\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$ , i.e. we were letting left and right moduli vary independently, then we could construct a solution of the BV QME by considering the diagonal product of the solutions of the BV QME in  $\mathcal{M}_L^{\text{II}}$  and  $\mathcal{M}_R^{\text{II}}$ . Again, it is not always clear that if we start from a diagonal solution of the BV QME in  $\mathcal{M}_{L;\text{red}}^{\text{II}} \times \mathcal{M}_{R;\text{red}}^{\text{II}}$  and thicken that solution in the fermionic directions, we land on a solution of the BV quantum master equation in the product  $\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$  since the thickening is defined up to homology. We thus need to consider the problem of existence of the solution of the BV quantum master equation in  $\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$  such that once we reduce that solution to  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})_{\text{red}}^{\overline{\phantom{x}}}$ , the reduced solution parametrizes irreducible surfaces  $\mathcal{R}_{\text{red}}^{\text{II}}$  which are embedded diagonally inside  $(\mathcal{R}_L^{\text{II}} \times \mathcal{R}_R^{\text{II}})_{\text{red}}^{\overline{\phantom{x}}}$ , i.e. those surfaces in  $(\mathcal{R}_L^{\text{II}} \times \mathcal{R}_R^{\text{II}})_{\text{red}}^{\overline{\phantom{x}}}$  whose left-moving complex structure is complex conjugate of the right-moving complex structure. Thus

we call  $\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}}) \equiv \Gamma^{\text{II}}$  the genus- $g$  type-II-superstring field theory vertex with  $\mathbf{n}_{\text{NSNS}}$  NSNS,  $\mathbf{n}_{\text{NSR}}$  RNS,  $\mathbf{n}_{\text{RNS}}$  NSR, and  $\mathbf{n}_{\text{RR}}$  RR boundary components if it satisfies the following requirements

1. The dimension of  $\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})$  is

$$\begin{aligned} \dim_{\mathbb{C}}^e(\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})) &= 3g - 3 + \mathbf{n}_{\text{NSNS}} + \mathbf{n}_{\text{NSR}} + \mathbf{n}_{\text{RNS}} + \mathbf{n}_{\text{RR}}, \\ \dim_{\mathbb{C}}^o(\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})) &= 4g - 4 + 2\mathbf{n}_{\text{NSNS}} + \frac{3}{2}(\mathbf{n}_{\text{NSR}} + \mathbf{n}_{\text{RNS}}) + \mathbf{n}_{\text{RR}}, \end{aligned} \quad (6.11)$$

where  $\dim_{\mathbb{C}}^e$  and  $\dim_{\mathbb{C}}^o$  are bosonic and fermionic dimensions, respectively.

2.  $\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})$ , as a subspace of  $\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$ , satisfies the BV QME;
3.  $(\mathcal{V}_g^{\text{II}}(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}}))_{\overline{\text{red}}}$  is embedded diagonally in  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})_{\overline{\text{red}}}$ , where by diagonal embedding, we mean all points in  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})_{\overline{\text{red}}}$  of the form  $(\mathcal{R}, \overline{\mathcal{R}})$ .

To construct such a solution, we proceed as follows<sup>57</sup>. We consider the fiber product with the twisted diagonal of  $\mathcal{M}(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}})$ :

$$\begin{array}{ccc} \mathcal{S}'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) & \xrightarrow{f^{\text{II}}} & \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \times \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \\ \pi \downarrow & & \downarrow \pi \\ \mathcal{M}(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}) & \xrightarrow{\mathbf{D}^t} & \mathcal{M}(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}) \times \mathcal{M}(\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}), \end{array}$$

where  $\mathbf{D}^t$  is the twisted diagonal defined by sending a surface  $\mathcal{R}$  to  $(\mathcal{R}, \overline{\mathcal{R}})$ . Note that the space  $\mathcal{S}'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  represents Riemann surfaces with  $\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}$  borders and two choices of spin structures, one with respect to a complex structure and the other one with respect to its complex conjugate. Therefore,  $\mathcal{S}'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  can be obtained by distributing  $\mathbf{n}_{\text{NS}}$  and  $\mathbf{n}_{\text{R}}$  such that  $\mathbf{n}_{\text{NS}} + \mathbf{n}_{\text{R}}$  is fixed. To each of such distribution of boundary components, one can assign  $2^{2g}$  choices of spin structures (and also its complex conjugate). Therefore,  $\mathcal{S}'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is a disjoint union of  $\mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .

Using this construction, we have

$$f^{\text{II}} : \mathcal{S}'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \longrightarrow \mathcal{M}_{L;\text{red}}^{\text{II}} \times \mathcal{M}_{R;\text{red}}^{\text{II}}. \quad (6.12)$$

We then have

$$(\pi \circ f^{\text{II}})(\mathcal{R}^{\mathcal{E}}) = \pi(f^{\text{II}}(\mathcal{R}^{\mathcal{E}})) = \pi(\mathcal{R}^{\mathcal{E}_L}, \overline{\mathcal{R}}^{\mathcal{E}_R}) = (\mathcal{R}, \overline{\mathcal{R}}), \quad (6.13)$$

where  $\mathcal{E}_L$  and  $\mathcal{E}_R$  are the spin structures of surfaces left- and right-moving sectors. *A priori, there is no relation between these spin structures.* This shows that the image of a spin-Riemann surface  $\mathcal{R}^{\mathcal{E}}$  under  $f^{\text{II}}$  after forgetting of the spin structure is embedded diagonally

---

<sup>57</sup>Note that for type-II-superstring theories  $\mathcal{M}_L^{\text{II}} = \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ ,  $\mathcal{M}_R^{\text{II}} = \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ ,  $\mathcal{M}_{L;\text{red}}^{\text{II}} = \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , and  $\mathcal{M}_{R;\text{red}}^{\text{II}} = \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .

in  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})_{\text{red}}^{\overline{\phantom{x}}}$  such that the left and right complex structures are complex conjugate of each other. To proceed further, we define the following space

$$(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}_{\text{red}} \equiv f^{\text{II}}(\mathcal{S}'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})) = \left\{ (\mathcal{R}^{\mathcal{E}_L}, \overline{\mathcal{R}}^{\mathcal{E}_R}) \in (\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})_{\text{red}} \right\}, \quad (6.14)$$

i.e. all pairs  $(\mathcal{R}^{\mathcal{E}_L}, \overline{\mathcal{R}}^{\mathcal{E}_R})$  of a surface  $\mathcal{R}^{\mathcal{E}_L}$  with a specific complex structures equipped with an arbitrary spin structure, and the surface  $\overline{\mathcal{R}}^{\mathcal{E}_R}$  whose complex structure is complex-conjugate of  $\mathcal{R}$  together with a choice of spin structure compatible with the complex-conjugate complex structure. As we mentioned above, there is no relation between spin structures  $\mathcal{E}_L$  and  $\mathcal{E}_R$ . As usual, the superscript  $\mathbf{D}$  stands for diagonal. The dimension of this space, for genus  $g$  surfaces with  $\mathbf{n}_{\text{NSNS}}$  NSNS,  $\mathbf{n}_{\text{NSR}}$  NSR,  $\mathbf{n}_{\text{RNS}}$  RNS, and  $\mathbf{n}_{\text{RR}}$  RR boundary components is  $6g - 6 + 2\mathbf{n}_{\text{NSNS}} + 2\mathbf{n}_{\text{NSR}} + 2\mathbf{n}_{\text{RNS}} + 2\mathbf{n}_{\text{RR}}$ , the same is as the space of spin-Riemann surfaces with the same signature. We also consider the thickening of this space in all fermionic dimensions, and denote the resulting space by  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}$ . Note that

$$\mathcal{M}_{L;\text{red}}^{\text{II}} = \mathcal{S}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = \bigcup_{g=0}^{\infty} \mathcal{S}_g(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) = \mathcal{M}_{R;\text{red}}^{\text{II}},$$

and the thickening is done for all values of  $(g; \mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  in  $2g - 2 + \mathbf{n}_{\text{NS}} + \frac{\mathbf{n}_{\text{R}}}{2}$  fermionic directions. We can thus define the following chain complex

$$\mathcal{F}((\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}_{\text{red}}) \equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{C}_* \left( \frac{(\mathcal{M}_L^{\text{II}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \times \mathcal{M}_R^{\text{II}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))^{\mathbf{D}}_{\text{red}}}{S_L^{\text{II}} \times S_R^{\text{II}}} \right), \quad (6.15)$$

where

$$S_L^{\text{II}} \equiv S_1 \wr S_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} = S_R^{\text{II}}, \quad (6.16)$$

$\mathcal{M}_L^{\text{II}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \equiv \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , and  $\mathcal{M}_R^{\text{II}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \equiv \mathcal{SM}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ .  $S_L^{\text{II}}$ , respectively  $S_R^{\text{II}}$ , acts on  $\mathcal{M}_L^{\text{II}}$ , respectively  $\mathcal{M}_R^{\text{II}}$ , trivially. We also define the superchain complex associated to  $(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}$

$$\mathcal{F}^{\text{II}}((\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}) \equiv \bigoplus_{\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}} \mathcal{SC}_* \left( \frac{(\mathcal{M}_L^{\text{II}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \times \mathcal{M}_R^{\text{II}}(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))^{\mathbf{D}}}{S_L^{\text{II}} \times S_R^{\text{II}}} \right). \quad (6.17)$$

The crucial property of the map  $f^{\text{II}}$  defined in (6.12) is that it is compatible with the gluing operation, i.e. the operation of gluing and mapping with  $f^{\text{II}}$  commute and there is a commutative diagram. We can thus define the following induced map

$$F_{f^{\text{II}}}^{\text{II}} : \mathcal{F}(\mathcal{S}') \longrightarrow \mathcal{F}((\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})^{\mathbf{D}}_{\text{red}}). \quad (6.18)$$

If  $\mathcal{F}(\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}})$  is the complex associated to the stack  $\mathcal{M}_L^{\text{II}} \times \mathcal{M}_R^{\text{II}}$ , it has naturally the structure of a differential-graded algebra where the differential  $d^{\text{II}} \equiv d_L^{\text{II}} \oplus d_R^{\text{II}}$  is the left  $d_L^{\text{II}}$  and right  $d_R^{\text{II}}$

boundary operations on left and right singular chains, and the product is the disjoint union of left and right surfaces.  $\mathcal{F}(\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)$  can be turned into a BV algebra by the usual twist-swing operation  $\Delta_{LR}^\Pi \equiv \Delta_L^\Pi \oplus \Delta_R^\Pi$ , where  $\Delta_L^\Pi(\Delta_R^\Pi)$  is the gluing operation defined on the complex  $\mathcal{F}(\mathcal{M}_L^\Pi)$  ( $\mathcal{F}(\mathcal{M}_R^\Pi)$ ).  $\mathcal{F}((\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)_{\text{red}}^\mathbf{D})$  inherits this BV-algebra structure. By construction,  $F_{f^\Pi}^\Pi$  is a homomorphism of BV algebras, and thus maps solutions of the BV QME onto each other. Therefore, having a solution in  $\mathcal{F}(S')$  whose existence we have proven<sup>58</sup>, we have a solution in  $\mathcal{F}((\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)_{\text{red}}^\mathbf{D})$ . We have thus constructed a subspace of  $(\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)_{\text{red}}^\mathbf{D}$  that satisfies the BV QME. To construct the type-II-superstring vertices, we consider the following maps of BV algebras

$$\widehat{F}_{f^\Pi}^\Pi : \mathcal{F}((\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)_{\text{red}}^\mathbf{D}) \longrightarrow \mathcal{F}((\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)^\mathbf{D}). \quad (6.19)$$

Since the BV algebra associated to singular superchains with top fermionic dimensions is quasi-isomorphic to the BV algebra of associated bosonic chains,  $\widehat{F}_{f^\Pi}^\Pi$  is a quasi-isomorphism of BV algebras. Again using the fact that quasi-isomorphic BV algebras have the same set of homotopy-classes of solutions of the BV QME, we conclude that there is a subspace of  $(\mathcal{M}_L^\Pi(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}) \times \mathcal{M}_R^\Pi(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}}))^\mathbf{D}$ , whose genus- $g$  contribution we denote by  $\mathcal{V}_g^\Pi(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})$ , which satisfies the BV QME.  $\mathcal{V}_g^\Pi(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})$  is the genus- $g$  type-II-superstring field theory vertex with  $\mathbf{n}_{\text{NSNS}}$  NSNS,  $\mathbf{n}_{\text{NSR}}$  NSR,  $\mathbf{n}_{\text{RNS}}$  RNS, and  $\mathbf{n}_{\text{RR}}$  RR boundary components since

1. its complex dimension is by construction the one given in (6.11);
2.  $\mathcal{V}_g^\Pi(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}})$ , as a subspace of  $\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi$ , satisfies the BV quantum master equation;
3. by construction  $(\mathcal{V}_g^\Pi(\mathbf{n}_{\text{NSNS}}, \mathbf{n}_{\text{NSR}}, \mathbf{n}_{\text{RNS}}, \mathbf{n}_{\text{RR}}))_{\text{red}}^\Pi$  is embedded diagonally in  $(\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)_{\text{red}}^\Pi$ .

We thus have proven the following

**Theorem 6.2 (The Existence of Type-II-Superstring Vertices).** *There exists a solution to the BV QME in  $\mathcal{F}((\mathcal{M}_L^\Pi \times \mathcal{M}_R^\Pi)^\mathbf{D})$ . The genus- $g$  part is a genus- $g$  type-II-string field theory vertex with  $\mathbf{n}_{\text{NSNS}}$  NSNS,  $\mathbf{n}_{\text{RNS}}$  RNS,  $\mathbf{n}_{\text{NSR}}$  NSR, and  $\mathbf{n}_{\text{RR}}$  RR boundary components.*

## 7 On the Moduli Problem of Stable $\mathcal{N} = 1$ SUSY Curves

This section is devoted to the representability of the moduli problem of stable  $\mathcal{N} = 1$  SUSY curves. We review some basics of superspace. We will basically follow [6, 7] for various definitions and splitness of superspaces, and follow [57, 58] for homology on superspaces. Unless specified, all rings discussed in this section are of characteristic 0, i.e. contains  $\mathbb{Q}$ . For early development on moduli problem and deformation theory see [80–83]. Deformation theory in the context of supergeometry is developed in [84, 85].

---

<sup>58</sup>Note that  $S'(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$  is a disjoint union of  $S(\mathbf{n}_{\text{NS}}, \mathbf{n}_{\text{R}})$ , and as such, the existence of the solution of the BV QME in  $\mathcal{F}(S')$  follows at once from the existence of solution in  $\mathcal{F}(S)$ .



## 7.1 Definitions, Elementary Results and Super Analogs of Classical Theorems

**Definition 7.1** (Superrings, Ideals and Modules). A superring is a  $\mathbb{Z}_2$ -graded ring  $R = R^+ \oplus R^-$  such that for homogeneous elements  $x, y \in R$ ,  $x \cdot y = (-1)^{|x||y|} y \cdot x$ . An ideal of  $R$  is a  $\mathbb{Z}_2$ -graded ideal of  $R$  (considered as an ordinary ring). A module of  $R$  is a  $\mathbb{Z}_2$ -graded  $R^+$ -module  $M = M^+ \oplus M^-$ , with an  $R^+$ -homomorphism  $a : R^- \otimes_{R^+} M^\pm \rightarrow M^\mp$  such that following diagram

$$\begin{array}{ccc} R^- \otimes_{R^+} R^- \otimes_{R^+} M^\pm & \xrightarrow{\text{Id} \otimes a} & R^- \otimes_{R^+} M^\mp \\ & \searrow \mu \otimes \text{Id} & \downarrow a \\ & & M^\pm \end{array}$$

commutes, where  $\mu : R^- \otimes_{R^+} R^- \rightarrow R^+$  is the multiplication map.

Unless specified, we only consider ideals and modules which are defined above, instead of ideals or modules in the ordinary sense.

**Definition 7.2** (Superspace). A superspace is a locally-superringed topological space  $(X, \mathcal{O}_X)$ , i.e. there is a  $\mathbb{Z}_2$ -grading

$$\mathcal{O}_X = \mathcal{O}_X^+ \oplus \mathcal{O}_X^-. \quad (7.1)$$

We also define the ideal generated by  $\mathcal{O}_X^-$  to be  $\mathfrak{n}_X$ , and define the locally-ringed topological space  $(X, \mathcal{O}_X/\mathfrak{n}_X)$  to be the **bosonic truncation** of  $X$ , denoted by  $X_{\text{bos}}$ . The locally-ringed topological space  $(X, \mathcal{O}_X^+)$  will be called the **bosonic quotient** of  $X$ , denoted by  $X_{\text{ev}}$ .

A morphism between superspaces  $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  is a morphism between locally-ringed topological space  $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  such that the induced homomorphism  $f^{-1}\mathcal{O}_Y \rightarrow \mathcal{O}_X$  preserves the  $\mathbb{Z}_2$ -grading. The category of superspaces will be denoted by  $\text{SupSp}$ , and the category of locally-commutative-ringed topological space which we call commutative space will be denoted by  $\text{ComSp}$ .

**Remark 7.1.** Given a superspace  $(X, \mathcal{O}_X)$ , there is a natural  $\mathbb{Z}_2$  action on it, namely, let  $\mathbb{Z}_2$  acts on  $\mathcal{O}_X^i$  by  $(-1)^i$ , hence the  $\mathbb{Z}_2$ -invariant is exactly  $\mathcal{O}_X^+$ . This is where the name bosonic quotient comes from. Moreover this defines a functor:

$$C : \text{SupSp} \rightarrow \text{ComSp}.$$

**Remark 7.2.**  $X_{\text{bos}} \equiv (X, \mathcal{O}_{X_{\text{bos}}})$  has a natural superspace structure, namely let the odd part to be trivial. It has a natural closed embedding into  $X$  in the category of superspaces. In fact, for any locally-commutative-ringed topological space  $(X, \mathcal{O}_X)$ , there is a canonical superspace  $(X, \mathcal{O}_X \oplus 0)$  associated to it, and this is a functor:

$$S : \text{ComSp} \rightarrow \text{SupSp}.$$

**Lemma 7.1.**  $C$  is left adjoint to  $S$ , and  $S$  is left adjoint to the functor  $B$  which takes a superspace to its bosonic truncation.



*Proof.* Suppose that  $X \in \text{SupSp}$  and  $Y \in \text{ComSp}$ , and given a morphism  $f : X \rightarrow S(Y)$ , since the induced homomorphism  $f^{-1}\mathcal{O}_Y \rightarrow \mathcal{O}_X$  preserves  $\mathbb{Z}_2$ -grading and the odd part of  $f^{-1}\mathcal{O}_Y$  is trivial, this map factors through the  $\mathbb{Z}_2$ -invariant of  $\mathcal{O}_X$ , hence there is morphism  $S(X) \rightarrow Y$  given by the same map on topological spaces as  $f$  but send  $f^{-1}\mathcal{O}_Y$  to  $\mathcal{O}_X^+$ . The converse is obvious. Hence we see that  $C$  is left adjoint to  $S$ .

Suppose that there is a morphism  $g : S(Y) \rightarrow X$ , since the induced homomorphism  $g^{-1}\mathcal{O}_X \rightarrow \mathcal{O}_Y$  preserves  $\mathbb{Z}_2$  grading and the odd part of  $\mathcal{O}_Y$  is trivial, hence  $\mathfrak{n}_Y$  is sent to zero and  $g$  factors through  $X_{\text{bos}}$ . The converse is composing with canonical morphism  $X_{\text{bos}} \hookrightarrow X$ . Hence we see that  $S$  is left adjoint to  $B$ .  $\blacksquare$

**Definition 7.3 (Split and Projected Superspace).** A superspace is called **split** if  $\mathcal{O}_X$  is  $\mathbb{Z}_{\geq 0}$  graded, i.e.

$$\mathcal{O}_X = \bigoplus_{i \geq 0} \mathcal{O}_X[i],$$

such that  $\mathcal{O}_X$  is generated by  $\mathcal{O}_X[0]$  and  $\mathcal{O}_X[1]$ , and the  $\mathbb{Z}_2$ -grading is induced from the  $\mathbb{Z}_{\geq 0}$  grading. Note that this implies that  $\mathfrak{n}_X = \bigoplus_{i > 0} \mathcal{O}_X[i]$  and  $\mathcal{O}_{X_{\text{bos}}} \cong \mathcal{O}_X[0]$ .

A superspace is called **projected** if there is a morphism  $\pi : X \rightarrow X_{\text{bos}}$  of superspaces such that the following diagram commutes:

$$\begin{array}{ccc} X_{\text{bos}} & \xhookrightarrow{i} & X \\ & \searrow Id & \downarrow \pi \\ & & X_{\text{bos}}. \end{array}$$

It is easy to see that a split superspace is projected.

The following is obvious:

**Lemma 7.2.** A superspace  $X$  is projected if and only if the induced closed immersion  $X_{\text{bos}} \hookrightarrow X_{\text{ev}}$  has a retract  $X_{\text{ev}} \rightarrow X_{\text{bos}}$ .

The following lemma gives equivalent definitions of Noetherian superrings.

**Lemma 7.3.** Let  $R = R^+ \oplus R^-$  be a superring, then the following are equivalent:

1.  $R$  is Noetherian as an ordinary ring, i.e. its ideals (including non-homogeneous ideals) satisfies Ascending Chain Condition (ACC).
2.  $R$  satisfies ACC on ideals.
3.  $\text{Gr}_{\mathfrak{n}} R$  satisfies ACC on ideals.
4.  $\mathfrak{n}$  is a finitely-generated ideal and  $R_{\text{bos}} = R/\mathfrak{n}$  is Noetherian.

5.  $R^-$  is finitely-generated as  $R^+$ -module and  $R_{\text{bos}} = R/\mathfrak{n}$  is Noetherian.

6.  $R^-$  is finitely-generated as  $R^+$ -module and  $R^+$  is Noetherian.

*Proof.*  $1 \Rightarrow 2$  and  $6 \Rightarrow 5 \Rightarrow 4$  are obvious.  $6 \Rightarrow 1$ :  $R$  is a finitely-generated  $R^+$ -module so every left ideal in  $R$  is a finitely-generated  $R^+$ -module, same for right ideals.  $2 \Rightarrow 6$ : if  $I_i \subset I_{i+1} \subset \dots$  is an ascending chain of ideals of  $R^+$ , then  $I_i R = I_i \oplus I_i R^-$  is an ascending chain of homogeneous ideals of  $R$  hence stable, so  $\{I_i\}$  is stable and  $R^+$  is Noetherian; on the other hand,  $\mathfrak{n} = R^- \oplus (R^-)^2$  is a finitely-generated  $R$ -ideal so its odd part  $R^-$  is finitely-generated as an  $R^+$ -module.  $4 \Rightarrow 3$ : 4 implies that  $\text{Gr}_{\mathfrak{n}} R$  satisfies 6 hence  $\text{Gr}_{\mathfrak{n}} R$  is Noetherian whence 3 follows.  $3 \Rightarrow 6$ : 3 implies that  $\text{Gr}_{\mathfrak{n}} R$  satisfies 2 so by what we have shown  $\text{Gr}_{(R^-)^2} R^+$  is Noetherian and  $\text{Gr}_{(R^-)^2} R^-$  is a finitely-generated  $\text{Gr}_{(R^-)^2} R^+$ -module, so  $(R^-)^{2N-1}/(R^-)^{2N+1} = 0$  for some  $N \geq 1$ , thus  $(R^-)^{2N-1} = 0$  by Nakayama Lemma (since elements in  $(R^-)^2$  are nilpotent), whence  $R^+$  is Noetherian<sup>59</sup> and  $R^-$  is a finitely-generated  $R^+$ -module<sup>60</sup>.  $\blacksquare$

Analogous to classical case, there is a notion of depth of modules in terms of Ext groups:

**Definition 7.4 (Depth of Module).** Let  $A$  be a local Noetherian superring with residue field  $k$ ,  $M$  is a finitely-generated  $A$  module, define  $\text{depth}_A(M)$  to be smallest  $n \in \mathbb{N}$  such that  $\text{Ext}_A^n(k, M)$  is nontrivial.

In fact this equals to the depth of  $M$  regarded as an  $A^+$ -module

**Lemma 7.4.**  $\text{depth}_{A^+}(M) = \text{depth}_A(M)$ . Consequently, the depth of  $M$  equals to the length of maximal regular sequence of  $M$ .

*Proof.* First of all, for any  $A$ -module  $N$  with finite length,  $\text{depth}_A(M)$  is the smallest integer  $n$  such that  $\text{Ext}_A^n(N, M)$  is nontrivial. This can be seen from induction on the length of  $N$  and the long exact sequence of Ext groups. Note that  $\text{Tor}_p^{A^+}(k, A)$  has finite length for all  $p$ . Then it follows immediately from the spectral sequence

$$\text{Ext}_A^q \left( \text{Tor}_p^{A^+}(k, A), M \right) \implies_p \text{Ext}_{A^+}^{p+q}(k, M),$$

that  $\text{depth}_A(M) = \text{depth}_{A^+}(M)$ .  $\blacksquare$

**Definition 7.5 (Cohen-Macaulay Module).** Let  $A$  be a local Noetherian superring,  $M$  is a finitely-generated  $A$  module.  $M$  is called Cohen-Macaulay if  $\text{depth}_A(M)$  equals to the dimension of  $\text{Supp}(M)$ <sup>61</sup>.

<sup>59</sup>For a commutative ring  $S$  with an ideal  $I$  such that  $I^2 = 0$ , if  $\text{Gr}_I S = S/I \oplus I$  is Noetherian then so is  $S$ , since for every ideal  $J$  of  $S$ ,  $J/J \cap I \oplus J \cap I$  is finitely-generated  $\text{Gr}_I S$ -module, so  $J/J \cap I$  and  $J \cap I$  are finitely-generated  $S$  modules, thus  $J$  is a finitely-generated ideal.

<sup>60</sup>Similarly, for a commutative ring  $S$  with an ideal  $I$  such that  $I^2 = 0$ , if  $S$ -module  $M$  satisfies that  $\text{Gr}_I M = M/IM \oplus IM$  is a finitely-generated  $\text{Gr}_I S$ -module, then  $M$  is a finitely-generated  $S$ -module.

<sup>61</sup> $\text{Supp}(M)$  denote the support of the module  $M$ .

Obviously we have the inequality:

$$\begin{aligned} \text{depth}_A(M) &= \min\{\text{depth}_{A^+}(M^+), \text{depth}_{A^+}(M^-)\} \\ &\leq \max\{\dim \text{Supp}(M^+), \dim \text{Supp}(M^-)\} = \dim \text{Supp}(M), \end{aligned}$$

which implies the following

**Lemma 7.5.** *M is Cohen-Macaulay if and only if both  $M^+$  and  $M^-$  are Cohen-Macaulay  $A^+$  modules and either they have the same dimension of supports or one of them is zero. In particular, ring A itself is Cohen-Macaulay if and only if  $A^+$  is a Cohen-Macaulay ring and either  $A^-$  is a Cohen-Macaulay module for  $A^+$  with  $\dim \text{Supp}(A^-) = \dim \text{Spec } A$  or  $A^-$  is trivial.*

**Definition 7.6 (Superscheme).** *A superscheme is a superspace  $(X, \mathcal{O}_X)$  such that  $(X, \mathcal{O}_X^+)$  is a scheme and  $\mathcal{O}_X^-$  is a quasi-coherent  $\mathcal{O}_X^+$ -module. This implies that  $\mathfrak{n}_X/\mathfrak{n}_X^2$ , which is isomorphic to  $\mathcal{O}_X^-/(\mathcal{O}_X^-)^3$ , is a quasi-coherent  $\mathcal{O}_{X_{\text{bos}}}$ -sheaf. A superscheme is called Noetherian if  $(X, \mathcal{O}_X^+)$  is Noetherian and  $\mathcal{O}_X^-$  is a coherent  $\mathcal{O}_X^+$  sheaf. This implies that  $(\mathcal{O}_X^-)^N$  and  $\mathfrak{n}_X^N$  vanish for large N. We say that a superscheme  $(X, \mathcal{O}_X)$  is affine if its bosonic quotient  $(X, \mathcal{O}_X^+)$  is affine*

**Example 7.1.** *The basic example is the super affine space  $\mathbb{A}^{p|q}$ , which has the same topological space with  $\mathbb{A}^p$ , but the structure sheaf*

$$\mathcal{O}_{\mathbb{A}^{p|q}} = \bigoplus_{i=0}^q \bigwedge^i \mathcal{O}_{\mathbb{A}^p}^q,$$

*endowed with the natural  $\mathbb{Z}_2$ -grading.*

**Definition 7.7 (Quasicoherent Sheaves).** *An  $\mathcal{O}_X$ -module  $\mathcal{F}$  on a superscheme X is said to be quasicoherent if there is an open covering of X such that for each open U in the covering family,  $\mathcal{F}$  is the cokernel of an **even** homomorphism*

$$\mathcal{O}_U^{K|L} \rightarrow \mathcal{O}_U^{I|J},$$

*where  $I, J, K, L$  are index sets, and  $\mathcal{O}_U^{I|J} := \mathcal{O}_U^I \oplus \prod \mathcal{O}_U^J$ . Moreover  $\mathcal{F}$  is called finitely-generated (respectively finite presented) if I and J are finite (respectively all index sets are finite) for all U. If X is Noetherian, then we say that  $\mathcal{F}$  is coherent if it is finitely-generated.*

Similar to the classical case, we have the following lemma on the extension of coherent sheaves:

**Lemma 7.6.** *Let S be a Noetherian superscheme,  $U \subset X$  is open and  $E \in \text{Coh}(\mathcal{O}_U)$ , then  $\exists F \in \text{Coh}(\mathcal{O}_S)$  such that  $E = F|_U$ .*

---

<sup>62</sup>By Chevalley's Criterion, this is equivalent to its bosonic truncation  $(X, \mathcal{O}_X/\mathfrak{n}_X)$  being affine.

*Proof.* First consider the case where  $S = \operatorname{Spec} A$  is affine, and let  $j : U \rightarrow S$  denote the inclusion. The quasi-coherent sheaf  $j_*(E)$  corresponds to the  $A$ -module  $M = \Gamma(U, E)$ . Given any  $u \in U$ , there exist finitely many elements  $e_1, \dots, e_n \in M$  which generate the fiber  $E(u)$  regarded as a super-vector space over the residue field  $k(u)$ . By Nakayama Lemma these elements will generate the stalks of  $E$  in an open neighbourhood of  $u$  in  $U$ . Therefore by the Noetherian hypothesis, there exist finitely many elements  $f_1, \dots, f_N \in M$  which generate the stalk of  $E$  at each point of  $U$ . If  $N \subset M$  is the submodule generated by these elements, then  $F = N^\sim$  is a coherent prolongation of  $E$  to  $S$ , proving the result in the affine case. In the general case, we can use the Noetherian hypothesis to assume that there is a maximal  $V$  on which  $E$  can be extended, if  $V \neq S$  then take an open affine  $V'$  not contained in  $V$  and extend  $F_V$  to  $V \cup V'$ , contradicting the maximality of  $V$ . ■

**Definition 7.8.** Given a morphism  $f : X \rightarrow Y$  between Noetherian superschemes, we say that

1.  $f$  has property **P** if  $f_{\text{bos}} : X_{\text{bos}} \rightarrow Y_{\text{bos}}$  has property **P**, where **P** can be finite type, separated, proper, finite.
2.  $f$  is flat if  $\mathcal{O}_X$  is a flat  $f^{-1}\mathcal{O}_Y$  module.
3.  $f$  is unramified if  $f$  is of finite type and for any geometric point  $y$  of  $Y$ , the fiber  $X_y$  is bosonic and étale.
4.  $f$  is étale if  $f$  is flat and unramified.
5.  $f$  is smooth of relative dimension  $(p|q)$  if locally on  $X$  there is an étale morphism to  $Y \times \mathbb{A}^{p|q}$  whose composition with projection  $Y \times \mathbb{A}^{p|q} \rightarrow Y$  is  $f$ .

**Lemma 7.7 (Local Criteria of Flatness).** Let  $A$  be a superring,  $I$  an ideal of  $A$  and  $M$  an  $A$ -module. Assume that either

- (a)  $I$  is nilpotent, or
- (b)  $A$  is local Noetherian,  $B$  is a local Noetherian  $A$ -superalgebra, and  $M$  is a finitely-generated  $B$ -module.

Then the following are equivalent

- (1)  $M$  is  $A$ -flat.
- (2)  $\operatorname{Tor}_1^A(N, M) = 0$  for all  $A/I$ -module  $N$ .
- (3)  $M/IM$  is  $A/I$ -flat and  $I \otimes M \xrightarrow{\sim} IM$  by natural map.
- (3')  $M/IM$  is  $A/I$ -flat and  $\operatorname{Tor}_1^A(A/I, M) = 0$ .
- (4) The natural map

$$\operatorname{Gr}_I A \otimes_{A/I} M/IM \rightarrow \operatorname{Gr}_I M,$$

is isomorphism.

(5)  $M/I^n M$  is  $A/I^n$ -flat, for all  $n \geq 0$ .

The proof is the same as in the classical case, see 20.C of [86], except that the classical Artin-Rees Lemma used in the proof of *loc.cit.* should be enhanced to its super analog proven below.

**Lemma 7.8 (Artin-Rees).** *Let  $I$  be an ideal of Noetherian superring  $A$ ,  $M$  a finitely-generated  $A$ -module and  $N$  a submodule of  $M$ . Then there exists a  $k \geq 1$  such that*

$$I^n M \cap N = I^{n-k}(I^k M \cap N),$$

for all  $n \geq k$ .

*Proof.* Apparently  $I^n M \cap N \supset I^{n-k}(I^k M \cap N)$ , so it suffices to prove the reverse inclusion. Define

$$\widetilde{M}^+ \equiv I^+ M^+ + I^- M^-, \quad \widetilde{M}^- \equiv I^+ M^- + I^- M^+,$$

then  $I^n M$  decomposes into even part  $(I^+)^{n-1} \widetilde{M}^+$  and odd part  $(I^+)^{n-1} \widetilde{M}^-$ , so we have  $(I^n M \cap N)^+ = (I^+)^{n-1} \widetilde{M}^+ \cap N^+$ . Using the classical Artin-Rees Lemma, there exists a  $k$  such that

$$(I^+)^{n-1} \widetilde{M}^+ \cap N^+ = (I^+)^{n-k}((I^+)^{k-1} \widetilde{M}^+ \cap N^+) = (I^+)^{n-k}((I^k M)^+ \cap N^+),$$

for all  $n \geq k$ . The last term is contained in  $I^{n-k}(I^k M \cap N)$ , whence the reverse inclusion holds.  $\blacksquare$

For the rest of this section, we will only consider Noetherian superschemes unless otherwise specified.

**Lemma 7.9.** *Suppose that  $f : X \rightarrow Y$  is flat and  $f^{-1} \mathfrak{n}_Y$  generates  $\mathfrak{n}_X$ , then the commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X_{\text{ev}} & \xrightarrow{f_{\text{ev}}} & Y_{\text{ev}}. \end{array}$$

is Cartesian, and  $f_{\text{ev}}$  is flat.

*Proof.* Since  $f^{-1} \mathfrak{n}_Y$  generates  $\mathfrak{n}_X$ , the local flatness implies that the natural map is isomorphism:

$$\mathcal{O}_{X_{\text{bos}}} \otimes_{f^{-1} \mathcal{O}_{Y_{\text{bos}}}} f^{-1} \text{Gr}_{\mathfrak{n}_Y} \mathcal{O}_Y \cong \text{Gr}_{\mathfrak{n}_X} \mathcal{O}_X.$$

Hence the homomorphism between bosonic graded algebras

$$\mathcal{O}_{X_{\text{bos}}} \otimes_{f^{-1} \mathcal{O}_{Y_{\text{bos}}}} \bigoplus_{i \geq 0} f^{-1} \text{Gr}_{\mathfrak{n}_Y}^{2i} \mathcal{O}_Y \rightarrow \bigoplus_{i \geq 0} \text{Gr}_{\mathfrak{n}_X}^{2i} \mathcal{O}_X,$$

is an isomorphism. So  $f_{\text{ev}}$  is flat by local flatness again. It is straightforward from the isomorphism between graded algebras that the induced homomorphism  $f_{\text{ev}}^* \mathcal{O}_Y \rightarrow \mathcal{O}_X$  is an isomorphism of sheaves, i.e.  $X \cong X_{\text{ev}} \times_{Y_{\text{ev}}} Y$ .  $\blacksquare$

**Lemma 7.10.**  $f : X \rightarrow Y$  is étale if and only if  $f_{\text{ev}} : X_{\text{ev}} \rightarrow Y_{\text{ev}}$  is étale and the commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X_{\text{ev}} & \xrightarrow{f_{\text{ev}}} & Y_{\text{ev}}. \end{array}$$

is Cartesian.

*Proof.* If  $f$  is étale, then  $X \times_Y Y_{\text{bos}}$  is bosonic, since its nilpotent ideal  $\mathfrak{n}$  generated by fermionic elements is zero after restriction to fibers, thus it is identically zero by Nakayama Lemma. So we see that  $f^{-1}\mathfrak{n}_Y$  generates  $\mathfrak{n}_X$ . By Lemma 7.9,  $f_{\text{ev}}$  is flat and the diagram is Cartesian. It follows that  $f_{\text{ev}}$  and  $f$  have the same fibers hence  $f_{\text{ev}}$  is étale. The converse is obvious. ■

**Lemma 7.11.** Suppose that  $f : X \rightarrow Y$  is étale. If  $Y$  is split (resp. projected), then so is  $X$ .

*Proof.* We use the characterization of étale morphism in Lemma 7.10. First of all,  $f$  being étale implies that  $\mathcal{O}_X^-$  is generated by  $f^{-1}\mathcal{O}_Y^-$ , hence there is an isomorphism

$$X_{\text{bos}} \cong X_{\text{ev}} \times_{Y_{\text{ev}}} Y_{\text{bos}}.$$

Assume that  $Y$  is projected, then there is a retract  $\pi : Y_{\text{ev}} \rightarrow Y_{\text{bos}}$ . To show that  $X$  is projected, it suffices to show that there is a retract  $X_{\text{ev}} \rightarrow X_{\text{bos}}$ . This follows from  $f_{\text{ev}}$  being étale hence there is a unique dotted arrow making following diagram commutes

$$\begin{array}{ccc} X_{\text{bos}} & \xlongequal{\quad} & X_{\text{bos}} \\ f_{\text{bos}} \downarrow & \nearrow \exists! & \downarrow \\ Y_{\text{bos}} & \xleftarrow{\pi \circ f_{\text{ev}}} & X_{\text{ev}}. \end{array}$$

By uniqueness, the composition of embedding  $X_{\text{bos}} \hookrightarrow X_{\text{ev}}$  with the dotted arrow is identity, hence  $X$  is projected.

Assume that  $Y$  is split, then  $Y$  is projected hence  $X$  is also projected and the previous argument shows that  $\mathcal{O}_{X_{\text{ev}}} \cong f_{\text{bos}}^* \mathcal{O}_{Y_{\text{ev}}}$  as a sheaf of  $\mathcal{O}_{X_{\text{bos}}}$ -algebras. This in turn implies that  $\mathcal{O}_X \cong f_{\text{bos}}^* \mathcal{O}_Y$  whence the splitness of  $X$  follows. ■

**Lemma 7.12.** Suppose that  $f : X \rightarrow Y$  is étale. If  $Y \hookrightarrow Y'$  is a nilpotent thickening, then there exists a unique lifting of étale morphism  $f' : X' \rightarrow Y'$  such that its restriction to  $Y$  is  $f$ .

*Proof.* Consider bosonic quotient of  $X, Y, Y'$  and fit them into

$$\begin{array}{ccc} X_{\text{ev}} & & \\ \downarrow f_{\text{ev}} & & \\ Y_{\text{ev}} & \hookrightarrow & Y'_{\text{ev}}. \end{array}$$

Now the classical nilpotent lifting result for étale morphisms shows that there exists a unique  $f'_{\text{ev}} : X'_{\text{ev}} \rightarrow Y'_{\text{ev}}$  which makes the diagram Cartesian

$$\begin{array}{ccc} X_{\text{ev}} & \longrightarrow & X'_{\text{ev}} \\ \downarrow f_{\text{ev}} & & \downarrow f'_{\text{ev}} \\ Y_{\text{ev}} & \longrightarrow & Y'_{\text{ev}}. \end{array}$$

Then  $X' = X'_{\text{ev}} \times_{Y'_{\text{ev}}} Y'$  is a lifting and the uniqueness is obvious.  $\blacksquare$

**Remark 7.3.** *There is a notion of (Artin, Deligne-Mumford) superstack which is defined in a similar flavor of usual stack [7], but the exact definition is not important here and we will skip it. Note that all statements remain true if we relax the conditions to allow DM superstacks [7].*

### 7.1.1 Superschemes over Inductive Limit of Superrings

Here we include some results for superschemes/sheaves/morphisms under the inverse limit of base superscheme. The setup is similar to the classical algebraic geometry: Let  $(I, \leq)$  be a directed set, with a distinguished element  $0 \in I$ , suppose that  $\{A_i\}_{i \in I}$  is an directed system of superrings, with inductive limit  $A$ .

**Lemma 7.13.** *If  $X_0, Y_0$  are finite presented surperschemes over  $\text{Spec } A_0$ , let  $X_i, Y_i$  be the base change of  $X_0, Y_0$  to  $\text{Spec } A_i$ , and  $X, Y$  be the base change of  $X_0, Y_0$  to  $\text{Spec } A$ , then the natural map*

$$\varphi : \varinjlim_i \text{Map}_{\text{Spec } A_i}(X_i, Y_i) \rightarrow \text{Map}_{\text{Spec } A}(X, Y),$$

*is an equivalence of sets.*

*Proof.*  **$\varphi$  is injective:** suppose that two morphisms  $f_0, g_0 : X_0 \rightarrow Y_0$  has the same limit  $f = g : X \rightarrow Y$ , consider the kernel of two morphisms  $j_0 : \text{Ker}(f_0, g_0) \rightarrow X_0$ , which is a finite presented immersion, becomes an isomorphism after passing to the limit, hence  $j_i : \text{Ker}(f_i, g_i) \rightarrow X_i$  is a homeomorphism for some index  $i \in I$ . Since the ideal defining  $j_i$  is finitely-generated, so locally we can take finitely many generators and they vanishes after taking limit, thus vanishing for some index  $i' \in I$ , consequently  $j_{i'} : \text{Ker}(f_{i'}, g_{i'}) \rightarrow X_{i'}$  is an isomorphism and  $f_{i'} = g_{i'}$ .

**$\varphi$  is surjective:** suppose that  $f : X \rightarrow Y$  is a morphism, we cover  $Y_0$  by finite many affine open sub superschemes  $\{U_{\lambda 0} \rightarrow Y_0\}$ , and let  $U_\lambda$  be the base change of  $U_{\lambda 0}$  to  $\text{Spec } A$ , then cover  $X$  by finite many open affine sub superschemes  $\{V_{\lambda \mu} \rightarrow X\}$  such that  $f(V_{\lambda \mu}) \subset U_\lambda$ . According to Theorem 10.57 of [87], we can assume that there is an index  $i \in I$  and an open covering  $\{V_{\lambda \mu i} \rightarrow X_i\}$  whose base change to  $X$  is  $\{V_{\lambda \mu} \rightarrow X\}$ . Refine  $V_{\lambda \mu i}$  such that each open sub superscheme is affine. Let us still use the same notation to denote this refined cover, then  $\mathcal{O}_X(V_{\lambda \mu})$  is a finite presented  $\mathcal{O}_Y(U_\lambda)$  superalgebra, then there exists an index  $i'$  such that  $\mathcal{O}_X(V_{\lambda \mu i'})$  is a finite presented  $\mathcal{O}_Y(U_{\lambda i'})$  superalgebra, i.e. we found  $f_{\lambda \mu i'} : V_{\lambda \mu i'} \rightarrow Y_{i'}$

whose base change to  $Y$  is  $f$  restricted on  $V_{\lambda\mu}$ . Finally, increase the index  $i'$  such that  $f_{\lambda\mu i'}$  agrees on the overlap (because of the injectivity of  $\varphi$ ). Hence  $f_{\lambda\mu i'}$  glues to a morphism  $f_{i'} : X_{i'} \rightarrow Y_{i'}$   $\blacksquare$

**Lemma 7.14.** *If  $X$  is a finite presented surperscheme over  $\operatorname{Spec} A$ , then there exists a finite presented superscheme  $X_i$  over  $\operatorname{Spec} A_i$  such that its base change to  $\operatorname{Spec} A$  is isomorphic to  $X$ .*

*Proof.* Cover  $X$  by open affine sub superschemes  $\{U_\lambda \rightarrow X\}$ , then each  $\mathcal{O}_X(U_\lambda)$  is a finite presented  $A$  superalgebra, thus  $\exists i \in I$  such that there exists finite presented  $A_i$  superalgebras  $B_{\lambda i}$  whose base change to  $A$  is  $\mathcal{O}_X(U_\lambda)$ . Let  $U_\lambda$  be  $\operatorname{Spec} B_{\lambda i}$ . Now increase the index  $i$  such that there exists open sub superschemes  $V_{\lambda\mu i} \subset U_\lambda$  whose base change to  $U_\lambda$  is  $U_\mu \cap U_\lambda$ , together with isomorphism  $\psi_{\lambda\mu i} : V_{\lambda\mu i} \cong V_{\mu\lambda i}$  satisfying cocycle conditions whose base change to  $\operatorname{Spec} A$  is the gluing morphism for the cover  $U_\lambda$ . The existence of such open sub superschemes follows from Theorem 10.57 of [87] and the existence of such isomorphisms follows from Lemma 7.13. Hence  $V_{\lambda\mu i}$  glues to a superscheme  $X_i$  over  $\operatorname{Spec} A_i$ , whose base change to  $\operatorname{Spec} A$  is isomorphic to  $X$ .  $\blacksquare$

In summary, the natural functor between categories of finite presented schemes

$$\varinjlim_i \operatorname{Sch}_{\text{fp}}(S_i) \rightarrow \operatorname{Sch}_{\text{fp}}(S),$$

is an equivalence of categories. We also have similar result for finite presented sheaves, whose proof is similar to the bosonic case, which will be omitted:

**Lemma 7.15.** *Let  $(I, \leq)$  be a directed set, suppose that  $\{S_i\}_{i \in I}$  is an directed system of qcqs<sup>63</sup> superschemes with affine transition morphisms, with inverse limit  $S$ , then the natural functor between categories of finite presented sheaves*

$$\varinjlim_i \operatorname{QCoh}_{\text{fp}}(\mathcal{O}_{S_i}) \rightarrow \operatorname{QCoh}_{\text{fp}}(\mathcal{O}_S),$$

*is an equivalence of categories.*

For later application, we also need a result about flatness:

**Proposition 7.1.** *If  $X$  is a finite presented surperscheme over  $\operatorname{Spec} A$ ,  $\mathcal{F} \in \operatorname{QCoh}_{\text{fp}}(\mathcal{O}_X)$ ,  $X$  is the inverse limit of a system of finite presented superscheme  $X_i$  over  $\operatorname{Spec} A_i$  and  $\mathcal{F}$  is the inverse limit of  $\mathcal{F}_i \in \operatorname{QCoh}_{\text{fp}}(\mathcal{O}_{X_i})$ , then  $\mathcal{F}$  is flat over  $\operatorname{Spec} A$  if and only if  $\exists i \in I$  such that  $\mathcal{F}_i$  is flat over  $\operatorname{Spec} A_i$ .*

*Proof.* This is the super analog of Théorème 11.2.6 of [88]. Let's explain why the proof of *loc.cit.* Théorème 11.2.6 works for superschemes as well. In the step I, the key ingredient is to find a Noetherian sub superrings of  $A_0$ , denoted by  $A'_0$  such that there exists  $(X'_0, \mathcal{F}'_0)$

---

<sup>63</sup>This stands for quasi-compact quasi-separated.



on  $\text{Spec } A'_0$  whose base change to  $\text{Spec } A_0$  is  $(X_0, \mathcal{F}_0)$ , this can be done because we can finite many even elements  $\{x_i\}_{i=1}^m$  in  $A_0^+$  and finite many odd elements  $\{\theta_j\}_{j=1}^n$  in  $A_0^-$ , then we define a homomorphism

$$\mathbb{Z}[X_i|\Theta_j] \rightarrow A_0,$$

by sending  $X_i$  to  $x_i$  and  $\Theta_j$  to  $\theta_j$ , its image in  $A_0$  is a Noetherian superring. Increase the size of this set and we can apply Lemma 7.14 and 7.15. The step II is a pure topological statement so it works for superschemes as well. The key ingredient of step III is showing that  $\text{Tor}_1^{A_0}(A_0/\mathfrak{m}_0, M_0)$  is finitely-generated  $B_0$ -module in the setup that  $A_0$  is a Noetherian superring,  $B_0$  is a finite type  $A_0$  superalgebra and  $M_0$  is a finitely-generated  $B_0$ -module. This is because we can take a resolution of  $A_0/\mathfrak{m}_0$  by finite free  $A_0$  modules, with even connecting homomorphisms, by the Noetherian property of  $A_0$ . Now the same argument in step III shows that the canonical homomorphism

$$\text{Tor}_1^{A_0}(A_0/\mathfrak{m}_0, M_0) \rightarrow \text{Tor}_1^{A_i}(A_i/\mathfrak{m}_0 A_i, M_i),$$

is zero for some  $i \in I$ , then the flatness of  $M_i$  over  $B_i$  follows from the super analog of *loc.cit.* Lemme 11.2.4 (stated in the Lemma 7.16) and the local criterion for flatness.  $\blacksquare$

**Lemma 7.16.** *Let  $A$  be a superring,  $C$  is an  $A$  superalgebra,  $M$  is an  $A$  module,  $N$  is a  $C$  module and  $A'$  is an  $A$  algebra. Let  $C' = C \otimes_A A'$ ,  $M' = M \otimes_A A'$  and  $N' = N \otimes_A A'$ , then the canonical homomorphism*

$$\text{Tor}_1^A(M, N) \otimes_A A' \rightarrow \text{Tor}_1^{A'}(M', N'),$$

*is surjective.*

The proof is formally the same as the proof of Lemme 11.2.4 of [88], except that the resolutions  $L_1 \rightarrow L_0 \rightarrow M \rightarrow 0$  and  $L'_1 \rightarrow L'_0 \rightarrow M' \rightarrow 0$  in the *loc.cit.* are required to have even connecting homomorphisms.

### 7.1.2 Affineness Criteria

Here are super analog of Serre's and Chevalley's Criteria for affineness.

**Proposition 7.2 (Serre's Criterion).** *Let  $X$  be a quasi-compact superscheme. Then the following statements are equivalent:*

- (1)  $X$  is affine.
- (2) *There exists a family of even elements  $f_i \in A \equiv \Gamma(X, \mathcal{O}_X)$  such that the open sub superscheme  $X_{f_i} := \{x \in X | f_i(x) \neq 0\}$  of  $X$  is affine for all  $i$  and the ideal generated by the  $f_i$  in  $A$  is the unit ideal.*
- (3)  $H^1(X, \mathcal{F}) = 0$  for all  $\mathcal{F} \in \text{QCoh}(\mathcal{O}_X)$ .
- (4)  $H^1(X, \mathcal{I}) = 0$  for all quasi-coherent ideal  $\mathcal{I}$ .

*Proof.* Since statement (2) only involves even elements  $f_i$  and  $f_i$  generate unit ideal in  $A$  if and only if they generate unit ideal in  $A^+$ , the equivalence between (1) and (2) follows from classical Serre's criterion.

(1)  $\implies$  (3) : Write  $\mathcal{F} = \mathcal{F}^+ \oplus \mathcal{F}^-$  be its even and odd parts, considered as  $\mathcal{O}_X^+$  module, then  $(X, \mathcal{O}_X^+)$  being affine implies that  $H^1(X, \mathcal{F}^\pm) = 0$ .

(3)  $\implies$  (4) is trivial, so let us prove (4)  $\implies$  (1) : For any quasi-coherent ideal  $\mathcal{J}$  of  $\mathcal{O}_X^+$ , consider  $\mathcal{I} = \mathcal{J} \oplus \mathcal{J}\mathcal{O}_X^- \subset \mathcal{O}_X$ , which is a quasi-coherent  $\mathcal{O}_X$ -ideal, so  $H^1(X, \mathcal{I}) = H^1(X, \mathcal{J}) \oplus H^1(X, \mathcal{J}\mathcal{O}_X^-) = 0$ , hence  $H^1(X, \mathcal{J}) = 0$  and by classical Serre's criterion,  $X_{\text{ev}}$  is affine.  $\blacksquare$

**Proposition 7.3 (Chevalley's Criterion).** *Suppose that  $f : X \rightarrow Y$  is a finite morphism between superschemes, then  $X$  is affine if and only if  $Y$  is affine.*

*Proof.* Since  $f_{\text{bos}} : X_{\text{bos}} \rightarrow Y_{\text{bos}}$  is finite, then the result follows from classical Chevalley's criterion.  $\blacksquare$

### 7.1.3 Chow's Lemma for Superschemes

**Theorem 7.1.** *Let  $S$  be a Noetherian superscheme,  $f : X \rightarrow S$  is of finite type and separated. Then there exists a dense open  $U \subset X$  and a projective morphism  $p : X' \rightarrow X$ , such that  $p|_{p^{-1}(U)}$  is an isomorphism, and  $X'$  is quasi-projective over  $S$ .*

*Proof.* First we consider the bosonic quotient of  $X$  and  $S$ , denoted by  $X_{\text{ev}}$  and  $S_{\text{ev}}$ , and fit them into a commutative diagram:

$$\begin{array}{ccccc}
 X & & & & \\
 \searrow \pi & \searrow f & & & \\
 & X_{\text{ev}} \times_{S_{\text{ev}}} S & \xrightarrow{f'} & S & \\
 & \downarrow & & \downarrow & \\
 & X_{\text{ev}} & \xrightarrow{f_{\text{ev}}} & S_{\text{ev}} & 
 \end{array}$$

Both  $f_{\text{ev}}$  and  $f'$  are of finite type, since  $X_{\text{bos}} \rightarrow S_{\text{ev}}$  is of finite type and nilpotent ideal defining  $X_{\text{bos}} \subset X_{\text{ev}}$  is finitely-generated.  $\pi$  is finite, since  $\pi_{\text{bos}} : X_{\text{bos}} \rightarrow (X_{\text{ev}} \times_{S_{\text{ev}}} S)_{\text{bos}} = X_{\text{bos}}$  is identity, and nilpotent ideal defining  $X_{\text{bos}} \subset X$  is finitely-generated.

By applying the classical Chow's Lemma to  $f_{\text{ev}}$ , we get a dense open  $U_{\text{ev}} \subset X_{\text{ev}}$  and a projective morphism  $q : Y \rightarrow X_{\text{ev}}$ , such that  $q|_{q^{-1}(U_{\text{ev}})}$  is an isomorphism, and  $Y$  is a quasi-projective  $S_{\text{ev}}$ -scheme. So the natural morphism  $Y \times_{S_{\text{ev}}} S \rightarrow X_{\text{ev}} \times_{S_{\text{ev}}} S$  is projective, and  $Y \times_{S_{\text{ev}}} S$  is quasi-projective over  $S$ . Since  $\pi$  is finite, pullback of  $Y \times_{S_{\text{ev}}} S$  along  $\pi$  is also quasi-projective over  $S$ , and call it  $X'$ . The projection  $p : X' \rightarrow X$  is an isomorphism on  $p^{-1}(U)$  so it satisfies all conditions.  $\blacksquare$

### 7.1.4 Pushforward and Flatness

The following is the super analog of Grothendieck's Complex, whose proof is also analogous to the classical case.

**Theorem 7.2.** *Let  $S = \operatorname{Spec} R$  be a Noetherian affine superscheme,  $f : X \rightarrow S$  is proper,  $\mathcal{F} \in \operatorname{Coh}(\mathcal{O}_X)$  and is flat over  $S$ , then  $Rf_*\mathcal{F}$  is quasi-isomorphic to a perfect complex with cohomological amplitudes in  $[0, d]$ , where  $d$  is the maximal dimension of  $\operatorname{Supp}(\mathcal{F}_s)$ , and its formation commutes with arbitrary base change.*

In the same way as the classical situation, Grothendieck's Complex implies the base-change theorem for flat sheaves:

**Theorem 7.3** (Theorem 5.10 of [56]). *Let  $\pi : X \rightarrow S$  be a proper morphism of Noetherian superschemes, and let  $\mathcal{F}$  be a coherent  $\mathcal{O}_X$ -module which is flat over  $S$ . Then the following statements hold:*

1. *For any integer  $i$  the functions  $s \mapsto \dim_{\kappa(s)} H^i(X_s, \mathcal{F}_s)_+$  and  $s \mapsto \dim_{\kappa(s)} H^i(X_s, \mathcal{F}_s)_-$  are upper semi-continuous on  $S$ .*
2. *The functions  $s \mapsto \dim_{\kappa(s)} \chi(X_s, \mathcal{F}_s)_+$  and  $s \mapsto \dim_{\kappa(s)} \chi(X_s, \mathcal{F}_s)_-$  are locally constant on  $S$ .*
3. *Suppose that  $S$  is a reduced scheme, and if for some integer  $i$ , there is a pair of integers  $d_{\pm} \geq 0$  such that for all  $s \in S$  we have  $\dim_{\kappa(s)} H^i(X_s, \mathcal{F}_s)_{\pm} = d_{\pm}$ , then  $R^i\pi_*\mathcal{F}$  is locally free of rank  $(d_+|d_-)$  and  $(R^i\pi_*\mathcal{F})_s \rightarrow H^i(X_s, \mathcal{F}_s)$  is an isomorphism for all  $s \in S$ .*
4. *If for some integer  $i$  and point  $s \in S$  the map  $(R^i\pi_*\mathcal{F})_s \rightarrow H^i(X_s, \mathcal{F}_s)$  is surjective, then there exists an open subscheme  $U \subset S$  such that both  $\tau_{\leq i} R\pi_*\mathcal{F}|_U$  and  $\tau_{\geq i+1} R\pi_*\mathcal{F}|_U$  are quasi-isomorphic to perfect complex on  $U$ .*

**Corollary 7.1** (Hom Superscheme). *Let  $\pi : X \rightarrow S$  be quasi-projective morphism between Noetherian superschemes, and let  $G$  and  $F$  be coherent sheaves on  $X$ . Suppose that  $F$  is flat over  $S$  and  $\operatorname{Supp}(F)$  is proper over  $S$ . Then the functor  $\mathfrak{H}\operatorname{om}_S(G, F)_+ : T \mapsto \operatorname{Hom}_{X_T}(G_T, F_T)_+$  is represented by a linear superscheme  $\mathbb{V}(\mathcal{Q})$  for a coherent sheaf  $\mathcal{Q}$  on  $S$ .*

*Proof.* Since  $\pi$  is quasi-projective, there exists finite locally free sheaves  $E_1, E_2$  on  $X$  and a map  $\phi : E_2 \rightarrow E_1$  with  $\operatorname{coker} \phi = G$ . It follows that

$$\operatorname{Hom}_{X_T}(G_T, F_T) \cong H^0 R\pi_{T*} \underline{R}\operatorname{Hom}_{X_T}([E_{2T} \rightarrow E_{1T}], F_T).$$

Theorem 7.2 implies that  $R\pi_* \underline{R}\operatorname{Hom}_X([E_2 \rightarrow E_1], F)$  is a perfect complex of non-negative cohomological degrees, whose formation commutes with arbitrary base change. Denote this perfect complex by  $\mathcal{K}^\bullet$ .  $(\mathcal{K}^\bullet)^\vee$  is of non-positive cohomological degrees, whence

$$H^0(\mathcal{K}_T^\bullet) = \operatorname{Hom}_{\mathcal{O}_T}(H^0((\mathcal{K}^\bullet)_T^\vee), \mathcal{O}_T).$$

So  $\mathfrak{H}\operatorname{om}_S(G, F)$  is represented by  $\mathbb{V}(\mathcal{Q})$  where  $\mathcal{Q} = H^0((\mathcal{K}^\bullet)^\vee)$ . ■

**Lemma 7.17.** *Let  $S = \operatorname{Spec} R$  be a Noetherian affine superscheme,  $X = \mathbb{P}_S^{n|m}$ ,  $\mathcal{F} \in \operatorname{Coh}(\mathcal{O}_X)$ , then  $\mathcal{F}$  is flat over  $S$  if and only if  $\exists N$  such that  $\forall i > N$ ,  $\pi_* \mathcal{F}(i)$  is flat on  $S$ .*

*Proof.* This is the super analog of Lemma 5.5 of [56]. The key point is that

$$\mathcal{F} = \Gamma_*(\mathbb{P}_S^{n|m}, \mathcal{F})^\sim.$$

The reason is that there is a canonical homomorphism  $\Gamma_*(\mathbb{P}_S^{n|m}, \mathcal{F})^\sim \rightarrow \mathcal{F}$  and we know that there is a projection  $\mathbb{P}^{n|m} \rightarrow \mathbb{P}^n$ , so  $\Gamma_*(\mathbb{P}_S^{n|m}, \mathcal{F})$  is identified with  $\Gamma_*(\mathbb{P}_S^n, \mathcal{F})$  under this projection, where the latter is considered as a  $\mathcal{O}_{\mathbb{P}^n}$  module. Now the isomorphism follows from its classical counterpart. The rest of proof is the same as *loc. cit.* ■

We also have the super analog of the Formal Function Theorem:

**Theorem 7.4 (Super Version of Formal Function Theorem).** *Let  $\pi : X \rightarrow S$  be a proper morphism of Noetherian superschemes, and let  $\mathcal{F}$  be a coherent  $\mathcal{O}_X$ -module, let  $s \in S$ .  $X_s^{(n)}$  is the  $n$ 'th formal neighborhood of  $X_s$  and  $(R^i \pi_* \mathcal{F})_s^\wedge$  be the completion of  $R^i \pi_* \mathcal{F}$  at  $s$ . Then there is an isomorphism*

$$(R^i \pi_* \mathcal{F})_s^\wedge \cong \varprojlim_n H^i(X_s^{(n)}, \mathcal{F}_s^{(n)}).$$

*Proof.* Localize  $S$  and we can assume that  $S = \operatorname{Spec} R$  for a local superring  $R = R^+ \oplus R^-$ , whose closed point is  $s$ . Let the maximal ideal of  $R^+$  be  $\mathfrak{m}_0$ , and let the ideal of  $R$  generated by  $R^-$  be  $\mathfrak{n}$ . Then the maximal ideal of  $R$  defining  $s$  is  $\mathfrak{m}_0 R + \mathfrak{n}$ . Note that  $(\mathfrak{m}_0 R + \mathfrak{n})^N \subset \mathfrak{m}_0^N R + \mathfrak{n}^N$ , so  $(\mathfrak{m}_0 R + \mathfrak{n})^N = \mathfrak{m}_0^N R$  for large  $N$ . We first replace  $\pi$  by  $\pi' : X \rightarrow S_{\text{ev}} = \operatorname{Spec} R^+$ , and the classical Formal Function Theorem states that

$$(R^i \pi'_* \mathcal{F})_s^\wedge \cong \varprojlim_n H^i(X_s^{(n)}, \mathcal{F}_s^{(n)}).$$

Note that  $R^i \pi'_* \mathcal{F}$  is the same as  $R^i \pi_* \mathcal{F}$  considered as a  $R^+$ -module. Now the theorem follows from the fact that  $\mathfrak{m}_0 R$  and  $\mathfrak{m}_0 R + \mathfrak{n}$  generate the same topology on  $R$  hence two limit systems are cofinal to each other. ■

### 7.1.5 Super Analog of Grothendieck's Comparison and Existence Theorems

Let  $A$  be an adic Noetherian superring,  $I$  an homogeneous ideal of definition of  $A$ ,  $Y = \operatorname{Spec} A$ ,  $Y_n = \operatorname{Spec} A/I^n$ , and

$$\widehat{Y} = \varprojlim_n Y_n = \operatorname{Spf} A.$$

**Theorem 7.5 (Grothendieck's Comparison Theorem).** *Let  $f : X \rightarrow Y$  be a finite type morphism between superschemes, and let  $\widehat{X}$  be the  $I$ -adic completion of  $X$ . Suppose that  $\mathcal{F} \in \operatorname{Coh}(\mathcal{O}_X)$  has proper support over  $Y$ , then the base change homomorphisms*

$$\begin{aligned} (R^q f_* \mathcal{F})^\wedge &\rightarrow R^q \widehat{f}_* \widehat{\mathcal{F}}. \\ R^q \widehat{f}_* \widehat{\mathcal{F}} &\rightarrow \varprojlim_n R^q (f_n)_* \mathcal{F}_n, \end{aligned} \tag{7.2}$$

*are isomorphisms.*

*Proof.* Write  $A = A^+ \oplus A^-$  be the decomposition of  $A$  into even and odd parts and also  $I = I^+ \oplus I^-$ , where  $I^+$  is even and  $I^-$  is odd. Since  $I^-$  is odd thus is nilpotent, the topology on  $\widehat{X}$  and  $\widehat{Y}$  can also be generated by  $(I^+)^n$ . Now we regard every thing to be superscheme over the bosonic adic Noetherian ring  $A^+$  with  $I^+$ -topology. Write  $\mathcal{F} = \mathcal{F}_+ \oplus \mathcal{F}_-$  be decomposition into even and odd parts. By classical Grothendieck's Comparison Theorem, there are natural isomorphisms of  $A^+$  modules

$$\begin{aligned} (R^q f_* \mathcal{F}_\pm)^\wedge &\rightarrow R^q \widehat{f}_* \widehat{\mathcal{F}}_\pm, \\ R^q \widehat{f}_* \widehat{\mathcal{F}}_\pm &\rightarrow \varprojlim_n R^q (f_n)_* (\mathcal{F}_\pm)_n. \end{aligned}$$

The super analog comes from the direct sum of even and odd parts. ■

**Corollary 7.2.** *In the same setup as Theorem 7.5, let  $\mathcal{F}, \mathcal{G} \in \text{Coh}(\mathcal{O}_X)$  whose supports have an intersection which is proper over  $Y$ . Then the natural map below is an isomorphism*

$$\text{Ext}^r(\mathcal{F}, \mathcal{G})^\wedge \xrightarrow{\sim} \text{Ext}^r(\widehat{\mathcal{F}}, \widehat{\mathcal{G}}). \quad (7.3)$$

The proof is formally the same as Corollary 8.2.9 in [56].

**Theorem 7.6 (Grothendieck's Existence Theorem).** *Let  $f : X \rightarrow Y$  be a separated finite type morphism between superschemes, and let  $\widehat{X}$  be the  $I$ -adic completion of  $X$ . Then the functor  $\mathcal{F} \mapsto \widehat{\mathcal{F}}$  where*

$$\widehat{\mathcal{F}} := \varprojlim_n \mathcal{F}/I^n \mathcal{F},$$

*from the category of coherent sheaves on  $X$  whose support is proper over  $Y$  to the category of coherent sheaves on  $\widehat{X}$  whose support is proper over  $Y$  is an equivalence.*

*Proof.* The fully faithfulness follows from Corollary 7.2. Let us prove essential surjectivity. Again we regard everything over the bosonic adic Noetherian ring  $A^+$  with  $I^+$ -topology. Suppose that  $F \in \text{Coh}(\mathcal{O}_{\widehat{X}})$  with proper support over  $\widehat{Y}$ , we want to show that there exists  $\mathcal{F} \in \text{Coh}(\mathcal{O}_X)$  with proper support over  $Y$  such that  $F = \widehat{\mathcal{F}}$ . Write  $F = F_+ \oplus F_-$  be decomposition into even and odd parts. By the classical Grothendieck's Existence Theorem, there exists coherent  $\mathcal{O}_X^+$  sheaves  $\mathcal{F}_+$  and  $\mathcal{F}_-$  whose supports are proper over  $Y$ , such that  $F_\pm = \widehat{\mathcal{F}}_\pm$ . Now an  $\mathcal{O}_X$ -module structure on  $\mathcal{F}_+ \oplus \mathcal{F}_-$  is a pair of homomorphisms

$$a : \mathcal{O}_X^- \otimes_{\mathcal{O}_X^+} \mathcal{F}_\pm \rightarrow \mathcal{F}_\mp,$$

such that the diagram

$$\begin{array}{ccc} \mathcal{O}_X^- \otimes_{\mathcal{O}_X^+} \mathcal{O}_X^- \otimes_{\mathcal{O}_X^+} \mathcal{F}_\pm & \xrightarrow{\text{Id} \otimes a} & \mathcal{O}_X^- \otimes_{\mathcal{O}_X^+} \mathcal{F}_\mp \\ & \searrow \mu \otimes \text{Id} & \downarrow a \\ & & \mathcal{F}_\pm \end{array},$$

commutes. Now every sheaf in this diagram has proper support, hence the existence of the homomorphisms  $a$  and the comutativity of this diagram follows from the classical Grothendieck's Existence Theorem. ■

### 7.1.6 Super Analog of Castelnuovo-Mumford Regularity

A coherent sheaf  $\mathcal{F}$  on  $\mathbb{P}^{n|s}$  is called  $m$ -regular if

$$H^i(\mathbb{P}^{n|s}, \mathcal{F}(m-i)) = 0, \quad \forall i \geq 1.$$

An important trick we will use is that  $\mathbb{P}^{n|s}$  is split, in fact  $\mathbb{P}^{n|s} = \wedge^\bullet \mathcal{O}_{\mathbb{P}^n}(-1)^{\oplus s}$ , so coherent sheaf  $\mathcal{F}$  on  $\mathbb{P}^{n|s}$  can be regarded as a pair of coherent sheaves  $(\mathcal{F}_+ | \mathcal{F}_-)$  on  $\mathbb{P}^n$  with a homomorphism  $a : \mathcal{O}_{\mathbb{P}^n}(-1)^s \otimes_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{F}_\pm \rightarrow \mathcal{F}_\mp$  such that the sum of two homomorphisms

$$\psi = a \circ (\text{Id} \otimes a) \text{ and } \phi = \psi \circ (\sigma \otimes \text{Id}) : \mathcal{F}_\pm(-2)^{\binom{s}{2}} \rightarrow \mathcal{F}_\pm, \quad (7.4)$$

is zero, where

$$\sigma \in \text{Aut}(\mathcal{O}_{\mathbb{P}^n}(-1)^s \otimes_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{O}_{\mathbb{P}^n}(-1)^s), \quad (7.5)$$

is the swap of two components. Under this identification, there is an isomorphism

$$H^i(\mathbb{P}^{n|s}, \mathcal{F}(r)) \cong H^i(\mathbb{P}^n, \mathcal{F}_+(r)) \oplus \prod H^i(\mathbb{P}^n, \mathcal{F}_-(r)).$$

Hence  $\mathcal{F}$  being  $m$ -regular is equivalent to that  $\mathcal{F}_+$  and  $\mathcal{F}_-$  are both  $m$ -regular (in the classical sense). With the help of this trick, we can formulate the Castelnuovo-Mumford  $m$ -regularity theory on super projective space purely in terms of  $m$ -regularity theory on classical projective space, namely we have

**Lemma 7.18** (Lemma 5.1 of [56]). *If  $\mathcal{F}$  is a  $m$ -regular sheaf on  $\mathbb{P}^{n|s}$ , then the following statements hold:*

1. *The canonical map  $H^0(\mathbb{P}^{n|s}, \mathcal{O}_{\mathbb{P}^n}(1)) \otimes H^0(\mathbb{P}^{n|s}, \mathcal{F}(r)) \rightarrow H^0(\mathbb{P}^{n|s}, \mathcal{F}(r+1))$  is surjective whenever  $r \geq m$ .*
2. *We have  $H^i(\mathbb{P}^{n|s}, \mathcal{F}(r)) = 0$  whenever  $i > 1$  and  $r > m|i$ . In other words, if  $\mathcal{F}$  is  $m$ -regular, then it is  $m'$ -regular for all  $m' > m$ .*
3. *The sheaf  $\mathcal{F}(r)$  is generated by its global sections, and all its higher cohomologies vanish, whenever  $r > m$ .*

**Theorem 7.7** (Theorem 5.3 of [56]). *For any non-negative integers  $p$ ,  $n$  and  $s$ , there exists a polynomial  $F_{p,n,s}$  in  $n+1$  variables with integral coefficients, which has the following property: Let  $k$  be any field, and let  $\mathcal{F}$  be any coherent sheaf on  $\mathbb{P}^{n|s}$ , which is isomorphic to a subsheaf of  $\mathcal{O}_{\mathbb{P}^{n|s}}^{\oplus p}$ . Let the Hilbert polynomial of  $\mathcal{F}$  be written in terms of binomial coefficients as*

$$\chi(\mathcal{F}(r))_+ = \sum_{i=0}^n a_i \binom{n}{i}, \quad \chi(\mathcal{F}(r))_- = \sum_{i=0}^n b_i \binom{n}{i},$$

where  $a_0, \dots, a_n \in \mathbb{Z}$ , and  $\chi(\mathcal{F}(r))_+$  (resp.  $\chi(\mathcal{F}(r))_-$ ) are even (resp. odd) Euler characteristics. Then  $\mathcal{F}$  is  $m$ -regular, where  $m = \max\{F_{p,n,s}(a_0, \dots, a_n), F_{p,n,s}(b_0, \dots, b_n)\}$ .

### 7.1.7 Flattening Stratification

Let's first recall the definition of flattening stratification. Suppose that  $f : X \rightarrow S$  is a morphism of superschemes and  $\mathcal{F} \in \text{QCoh}(\mathcal{O}_X)$ , a flattening stratification of  $\mathcal{F}$  over  $S$  (if exists) is a set of locally closed sub superschemes  $\{S_i\}_{i \in I}$  of  $S$ , such that

1. As a set,  $|S| = \coprod_{i \in I} |S_i|$ .
2. For any morphism of superschemes  $g : T \rightarrow S$ ,  $\mathcal{F}_T$  is flat over  $T$  if and only if there exists  $i \in I$  and  $g$  factors through  $S_i \hookrightarrow S$ .

An elementary case is when  $f$  is identity map so  $\mathcal{F}$  is a quasi-coherent sheaf on  $S$ . To study the existence of flattening stratification, we start from the affine case. Let  $R = R^+ \oplus R^-$  be a superring (not necessarily Noetherian) and  $M = \text{coker}(f : R^{p|q} \rightarrow R^{m|n})$  be a finite presented module. We write  $f$  in the matrix form

$$f = \begin{bmatrix} A & B \\ C & D \end{bmatrix}.$$

**Proposition 7.4.** *There exists flattening stratification for  $M$  on  $\text{Spec } R$ .*

*Proof.* It is well-known that  $M_{\text{bos}}$  on  $R_{\text{bos}} = R/\mathfrak{n}$  has flattening stratification

$$\coprod_{\substack{0 \leq k \leq m \\ 0 \leq l \leq n}} U'_{k|l} \rightarrow \text{Spec } R_{\text{bos}},$$

such that  $M_{S_{\text{bos}}}$  is flat of rank  $(k|l)$  if and only if  $\text{Spec } S_{\text{bos}} \rightarrow \text{Spec } R_{\text{bos}}$  factors through stratum  $U'_{k|l}$ . So if there is a flattening stratification for  $M$  on  $R$ , then it must have the same underlying topological space as the  $U'_{k|l}$ 's. Take any point  $p \in U'_{k|l}$ , defined by prime ideal  $\mathfrak{p}$ . Passing to residue field  $k(p)$ ,  $f$  is similar to a matrix of the form

$$\left[ \begin{array}{cc|cc} \text{Id}_{(m-k) \times (m-k)} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \text{Id}_{(n-l) \times (n-l)} \end{array} \right].$$

Hence on  $R_{\mathfrak{p}}$  (the localization of  $R$  at  $p$ ),  $f$  is similar to a matrix of the form

$$\left[ \begin{array}{cc|cc} \text{Id}_{(m-k) \times (m-k)} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A' & B' & \mathbf{0} \\ \hline \mathbf{0} & C' & D' & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \text{Id}_{(n-l) \times (n-l)} \end{array} \right],$$

where  $A', B', C', D'$  are matrices with input from prime ideal  $\mathfrak{p}$ , by Gauss elimination algorithm. The same is true if  $R_{\mathfrak{p}}$  is replaced by  $R_a$  for some  $a \in \mathfrak{p}_0$ . Then  $M_a$  is the cokernel of

$$f' = \begin{bmatrix} A' & B' \\ C' & D' \end{bmatrix} : R_a^{k+p-m|l+q-n} \rightarrow R_a^{k|l}.$$

Hence the pullback of  $M_a$  to a  $R_a$ -superscheme  $S$  is locally free of rank  $(k|l)$  if and only if pullback of  $A', B', C', D'$  are identically zero, i.e.  $S \rightarrow \operatorname{Spec} R_a$  factors through the closed super subscheme defined by the ideal generated by entries of  $A', B', C', D'$ . Let  $p$  runs through  $U'_{k|l}$  and we get a open super subcheme  $V \subset \operatorname{Spec} R$  and a closed super subscheme  $U_{k|l} \subset V$  by gluing. This is the flattening stratum for  $(k|l)$  piece.  $\blacksquare$

**Remark 7.4.** *Since  $U_{k|l}$  has the same topological space as  $U'_{k|l}$ , by the lower semicontinuity of the rank of matrix in the bosonic case, we see that the closure of  $U_{k|l}$  is contained in the union of all  $U_{k'|l'}$  where  $k' \geq k, l' \geq l$ .*

By a gluing argument, Proposition 7.4 generalizes to

**Corollary 7.3.** *There exists flattening stratification for finite presented sheaves on superschemes.*

A more general situation is when  $f : X \rightarrow S$  is projective, and  $\mathcal{F}$  is finite presented, in this case the Hilbert polynomials are constant for any flat strata (if exists) by Theorem 7.3. Analogous to the classical case, we can add one more condition to the flattening stratification for projective morphisms:

- The index set  $I$  for the flattening stratification is the set of Hilbert polynomials  $(f_+|f_-)$ , and  $I$  has a partial ordering, defined by putting  $(f_+|f_-) \leq (g_+|g_-)$  whenever  $f_+(n) \leq g_+(n)$  and  $f_-(n) \leq g_-(n)$  for all  $n \gg 0$ . Then the closure in  $S$  of the subset  $S_{(f_+|f_-)}$  is contained in the union of all  $S_{(g_+|g_-)}$  where  $(f_+|f_-) \leq (g_+|g_-)$ .

**Theorem 7.8** (Theorem 5.13 of [56]). *Let  $\mathcal{V}$  be a locally-free sheaf on a superscheme  $S$ , and  $\mathcal{F}$  be a finite presented sheaf on  $\mathbb{P}(\mathcal{V})$ , then there exists flattening stratification for  $\mathcal{F}$  over  $S$ .*

*Proof.* We first assume that  $S$  is affine. By Noetherian approximation, we can assume that  $S$  is also Noetherian and  $\mathcal{F}$  is coherent. If we can prove the proposition for affine Noetherian  $S$ , then the result automatically holds for general  $S$  by functorial property of flattening stratification.

Let us explain how to modify the arguments in the proof of Theorem 5.13 of *loc.cit.* so that it can be applied to its super analog: The special case corresponds to Corollary 7.3; Generic Flatness (*loc.cit.* 5.12) is still true, since if the base is an integral scheme, then we can project  $X$  down to its bosonic quotient  $X_{\text{ev}}$ , and apply the generic flatness to  $\mathcal{F}$  which is considered as a coherent sheaf on  $X_{\text{ev}}$ ; *loc.cit.* 5.5 is replaced by its super analog (Lemma 7.17). Now the remaining step is to repeat the proof of *loc.cit.* 5.13, which will be omitted here.  $\blacksquare$

**Corollary 7.4 (Incidence Superscheme).** *Let  $S$  be a Noetherian superscheme and  $\pi : X \rightarrow S$  is a projective morphism,  $Y, Z \subset X$  are two closed sub superschemes, and  $Y$  is flat over  $S$ . Then there is a closed sub superscheme  $S'$  of  $S$ , such that for any morphism of superschemes  $g : T \rightarrow S$ ,  $Y_T$  is a closed sub superscheme of  $Z_T$  if and only if  $g$  factors through  $S'$ .*



*Proof.* Let us first prove a weaker version: there is a set of disjoint locally closed sub superschemes  $\{S_i\}$  of  $S$ , such that for any morphism of superschemes  $g : T \rightarrow S$ ,  $Y_T$  is a closed sub superscheme of  $Z_T$  if and only if  $g$  factors through some  $S_i$ .

Let  $g : T \rightarrow S$  be a morphism such that  $Y_T$  is a closed sub superscheme of  $Z_T$ , then  $(Y \times_X Z)_T = Y_T \times_{X_T} Z_T = Y_T$  is flat over  $T$ , so  $g$  factors through one of the flattening strata  $S'_i$  for the coherent sheaf  $\mathcal{O}_{Y \times_X Z}$ . On the other hand, on the strata  $S'_i$ ,  $Y \times_X Z$  is flat over  $S'_i$  and is a closed sub superscheme of  $Y$ , defined by an ideal  $J_i$  which is flat over  $S'_i$ , then pull-back of  $Y \times_X Z$  to  $T$  is  $Y_T$  exactly when the pull back of the ideal  $J_i$  vanishes, i.e.  $g$  factor through  $S'_i \setminus \pi(\text{Supp } J_i)$ , the latter is open since  $\text{Supp } J_i$  and  $\pi$  is proper. Now we take  $S_i = S'_i \setminus \pi(\text{Supp } J_i)$  and it satisfies the condition.

Next we show that those  $S_i$  are closed. By Valuative Criterion, it is enough to assume that  $S = \text{Spec } R$  where  $R$  is a DVR<sup>64</sup>, and  $(Y \times_X Z)_\eta = Y_\eta$  where  $\eta$  is the generic point of  $S$ . Then the ideal of  $\mathcal{O}_Y$  defining the closed sub superscheme  $Y \times_X Z$  is supported on the special fiber, hence it must be trivial because  $Y$  is flat over  $S$ . Finally we take  $S' = \cup_i S_i$  and it satisfies the condition.  $\blacksquare$

### 7.1.8 The Quot Superscheme

Let  $S$  be a Noetherian superscheme,  $\pi : X \rightarrow S$  is a quasi-projective morphism,  $\mathcal{L}$  is relatively very ample line bundle of rank  $(1|0)$  on  $X$ . Then for any  $E \in \text{Coh}(\mathcal{O}_X)$  and any pair of polynomials  $\Phi(t), \Psi(t) \in \mathbb{Q}[t]$ , the functor  $\mathbf{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}$  on the category of superschemes over  $S$  is defined by

$$(f : T \rightarrow S) \mapsto \{E_T \twoheadrightarrow \mathcal{F} | \mathcal{F} \text{ is flat over } T \text{ with Hilber polynomials } (\Phi|\Psi)\}.$$

**Theorem 7.9.** *The functor  $\mathbf{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}$  is represented by a quasi-projective  $S$ -superscheme  $\text{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}$ . Moreover, if  $\pi : X \rightarrow S$  is projective then  $\text{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}$  is projective.*

The proof will be carried out in several steps.

**Lemma 7.19.** *Let  $X_1 \hookrightarrow X_2$  be an open immersion between quasi-projective  $S$ -superschemes,  $E \in \text{Coh}(\mathcal{O}_{X_2})$ ,  $\mathcal{L}$  is relatively very ample line bundle of rank  $(1|0)$  on  $X_2$ . Then the natural transform  $\mathbf{Quot}_{E/X_1/S}^{(\Phi|\Psi), \mathcal{L}} \rightarrow \mathbf{Quot}_{E/X_2/S}^{(\Phi|\Psi), \mathcal{L}}$  is an open immersion.*

*Proof.* Given an  $S$ -superscheme  $T$  and any point

$$(\phi : E \rightarrow F) \in \mathbf{Quot}_{E/X_2/S}^{(\Phi|\Psi), \mathcal{L}}(T).$$

The pullback of  $F$  to  $X_2 \times_S U$  for any  $T$ -superscheme  $U$  lies inside  $X_1 \times_S U$  if and only if the image of  $U$  in  $T$  does not intersect with  $K = \pi(\text{Supp}(F) \setminus X_1)$ . Hence the  $T$ -value of  $\mathbf{Quot}_{E/X_1/S}^{(\Phi|\Psi), \mathcal{L}} \rightarrow \mathbf{Quot}_{E/X_2/S}^{(\Phi|\Psi), \mathcal{L}}$  is represented by the complement of  $K$  in  $T$ , which is open.  $\blacksquare$

<sup>64</sup>DVR stands for discrete valuation ring.

**Lemma 7.20.** *Let  $\psi : E \rightarrow F$  be a surjective map between coherent sheaves on  $X$ . Then the natural transform  $\mathbf{Quot}_{F/X/S}^{(\Phi|\Psi), \mathcal{L}} \rightarrow \mathbf{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}$  is a closed immersion.*

*Proof.* Given an  $S$ -superscheme  $T$  and any point

$$(\phi : E \rightarrow G) \in \mathbf{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}(T).$$

Let  $K = \ker \psi$  and  $\eta : K \rightarrow G$  be the composition of inclusion of  $K$  in  $E$  and projection  $E \rightarrow G$ . The pullback of  $\phi$  to  $X \times_S U$  for any  $T$ -superscheme  $U$  factors through  $F$  if and only if the pullback of  $\eta$  to  $X \times_S U$  is zero. Corollary 7.1 says that  $\eta$  gives rise to a section  $\iota$  of a linear superscheme  $\mathbb{V}(\mathcal{Q})$  on  $T$ , and pullback to  $X \times_S U$  is zero if and only if  $U \rightarrow \mathbb{V}(\mathcal{Q})$  lies in the zero section  $\mathbf{0}$ , i.e.  $U \rightarrow T$  lies in the closed sub superscheme  $\iota^{-1}(\mathbf{0})$ . Hence the  $T$ -value of  $\mathbf{Quot}_{F/X/S}^{(\Phi|\Psi), \mathcal{L}} \rightarrow \mathbf{Quot}_{E/X/S}^{(\Phi|\Psi), \mathcal{L}}$  is represented by  $\iota^{-1}(\mathbf{0})$ .  $\blacksquare$

Representability of a Zariski functor can be proven Zariski locally on  $S$ , so we can assume that  $S$  is affine and there is an open embedding  $X \hookrightarrow Y$  followed by closed embedding  $Y \hookrightarrow \mathbb{P}^{n|m}$ , such that  $\mathcal{L}^{\otimes a}$  is the pullback of  $\mathcal{O}(1)$  on  $\mathbb{P}^{n|m}$ . Lemma 7.19 implies the existence of prolongation of  $E$  to a coherent sheaf  $F$  on  $Y$ , and Lemma 7.20 implies that we can replace  $F$  by  $\pi^*(\pi_* F(N))(-N)$  for large enough  $N$ , in fact  $\pi_* F(N)$  can be replaced by a free sheaf  $\mathcal{O}_S^{p|q}$  which surjects onto it. This step reduces the original question to showing the representability of

$$\mathbf{Quot}_{\mathcal{O}^{p|q}/\mathbb{P}_S^{n|m}/S}^{(\tilde{\Phi}|\tilde{\Psi}), \mathcal{O}(1)},$$

where  $\tilde{\Phi}(t) = \Phi(a(t + N))$  and  $\tilde{\Psi}(t) = \Psi(a(t + N))$ . For simplicity, we keep using  $\Phi$  and  $\Psi$  instead of  $\tilde{\Phi}$  or  $\tilde{\Psi}$ .

By  $m$ -regularity (Theorem 7.7), there exists  $m$  such that for any geometric point  $s$  of  $S$ , and any quotient  $\phi : \mathcal{O}^{p|q} \rightarrow \mathcal{F}$  on  $\mathbb{P}_s^{n|m}$ , the sheaves  $\mathcal{O}^{p|q}(r)$ ,  $\mathcal{F}(r)$ ,  $\mathcal{G}(r)$  (where  $\mathcal{G}(r) = \ker \phi(r)$ ) are generated by their global sections and have trivial  $H^i$  for  $i > 0$ , whenever  $r > m$ . Then for any  $S$ -superscheme  $T$  and

$$(\phi : \mathcal{O}^{p|q} \rightarrow \mathcal{F}) \in \mathbf{Quot}_{\mathcal{O}^{p|q}/\mathbb{P}_S^{n|m}/S}^{(\Phi|\Psi), \mathcal{O}(1)}(T),$$

there is a short exact sequence

$$0 \rightarrow \pi_* \mathcal{G}(r) \rightarrow \pi_* \mathcal{O}^{p|q}(r) \rightarrow \pi_* \mathcal{F}(r) \rightarrow 0,$$

which gives rise to a point in  $\text{Grass}(\Phi(r)|\Psi(r), \pi_* \mathcal{O}^{p|q}(r))$ .

**Lemma 7.21.** *The natural transform  $i : \mathbf{Quot}_{\mathcal{O}^{p|q}/\mathbb{P}_S^{n|m}/S}^{(\Phi|\Psi), \mathcal{O}(1)} \rightarrow \text{Grass}(\Phi(r)|\Psi(r), \pi_* \mathcal{O}^{p|q}(r))$  is represented by locally closed sub superscheme of  $\text{Grass}(\Phi(r)|\Psi(r), \pi_* \mathcal{O}^{p|q}(r))$ .*

*Proof.* Let the kernel of the canonical map  $\pi^* \pi_* \mathcal{O}^{p|q}(r) \rightarrow \mathcal{O}^{p|q}(r)$  be  $\mathcal{K}$ , then the image of  $\mathcal{K}$  in  $\pi^* \pi_* \mathcal{F}(r)$  is the kernel of canonical map  $\pi^* \pi_* \mathcal{F}(r) \rightarrow \mathcal{F}(r)$ , hence  $\mathcal{F}(r)$  is uniquely determined by the quotient  $\pi_* \mathcal{O}^{p|q}(r) \rightarrow \pi_* \mathcal{F}(r)$ . In other words,  $i$  is a monomorphism.

Let  $\mathcal{Q}$  be the universal quotient of  $\pi_*\mathcal{O}^{p|q}(r)$  of rank  $(\Phi(r)|\Psi(r))$ , let  $\mathcal{N}$  be the cokernel of the canonical map  $\mathcal{K} \rightarrow \pi^*\mathcal{Q}$ , then the formation of  $\mathcal{N}$  commutes with base change and comes from a  $\mathcal{F}(r)$  if and only if it is flat over the base and has characteristic polynomial  $(\Phi(t+r)|\Psi(t+r))$ . It follows that the lemma is a consequence of the flattening stratification (Theorem 7.8).  $\blacksquare$

Finally it remains to show that if  $\pi$  is projective then  $\text{Quot}_{E/X/S}^{(\Phi|\Psi),\mathcal{L}}$  is proper over  $S$ . This follows from the lemma below.

**Lemma 7.22 (Valuative Criterion).** *Let  $A$  be a discrete valuation ring, with fractional field  $K$ . Let  $X$  be a proper superscheme on  $\text{Spec } A$  and  $E \in \text{Coh}(\mathcal{O}_X)$ . If  $E_K \rightarrow F$  is a quotient of  $E_K$  on the generic fiber  $X_K$ , then there exists a quotient  $E \rightarrow G$  on  $X$  such that  $G_K = F$  and  $G$  is flat over  $A$ .*

*Proof.* Take  $G$  to be the image of  $E$  in  $j_*F$  under the canonical map  $E \rightarrow j_*F$  where  $j : X_K \rightarrow X$  is the open immersion. Clearly  $G$  restricts to  $F$ .  $G$  is torsion free as an  $A$ -module since it is a submodule of a torsion free  $A$ -module  $j_*F$ , so  $G$  is flat over  $A$ .  $\blacksquare$

The followings are corollaries to the existence of  $\text{Quot}$  superschemes, which are analogous to the classical case, and the proofs are formally identical to those in the *loc.cit.*, which will be omitted here.

**Theorem 7.10 (Theorem 5.23 of [56]).** *Let  $S$  be a Noetherian superscheme, let  $X$  be a projective superscheme over  $S$ , and let  $Y$  be quasi-projective superscheme over  $S$ . Assume moreover that  $X$  is flat over  $S$ . Then the functor*

$$\mathfrak{Map}_S(X, Y) : (f : T \rightarrow S) \mapsto \text{Map}_T(X_T, Y_T),$$

*is represented by an open sub superscheme  $\text{Map}_S(X, Y)$  of  $\underline{\text{Hilb}}_{X \times_S Y/S}$ .*

**Theorem 7.11 (Theorem 5.25 of [56]).** *Let  $S$  be a Noetherian superscheme, and let  $X \rightarrow S$  be a quasi-projective morphism. Let  $f : R \rightarrow X \times_S X$  be schematic equivalence relation on  $X$  over  $S$ , such that the projections  $f_1, f_2 : R \rightrightarrows X$  are proper and flat. Then a schematic quotient  $X \rightarrow Q$  exists over  $S$ . Moreover,  $Q$  is quasi-projective over  $S$ , the morphism  $X \rightarrow Q$  is faithfully flat and projective, and the induced morphism  $(f_1, f_2) : R \rightarrow X \times_Q X$  is an isomorphism.*

### 7.1.9 Deformation Theory and Super Analog of Schlessinger's Theorem

Let  $k$  be a fixed algebraically-closed ground field, and let the category of local Artinian  $k$ -superalgebras be  $\text{sArt}_k$ , and let the category of local complete Noetherian  $k$ -superalgebras be  $\text{sLoc}_k$ . We consider a covariant functor  $F : \text{sArt}_k \rightarrow \text{Sets}$ . One such example is taking an  $R \in \text{sLoc}_k$  and letting  $h_R : A \rightarrow \text{Hom}_{\text{sLoc}_k}(R, A)$ . By Yoneda Lemma, a natural transform  $\varphi : h_R \rightarrow F$  is given by a compatible family  $\{\xi_n \in F(R/\mathfrak{m}_R^n)\}$ .

**Definition 7.9.** Suppose that  $F : \mathbf{sArt}_k \rightarrow \mathbf{Sets}$  is a covariant functor,  $R \in \mathbf{sLoc}_k$  and  $\varphi : h_R \rightarrow F$  is a natural transform, represented by  $\xi := \{\xi_n \in F(R/\mathfrak{m}_R^n)\}$ . Then

- $(R, \xi)$  is called a **versal** family if  $\varphi : h_R \rightarrow F$  is strongly surjective, i.e.  $\forall A \in \mathbf{sLoc}_k$ ,  $h_R(A) \rightarrow F(A)$  is surjective, and for every surjection  $B \rightarrow A$  in  $\mathbf{sLoc}_k$ ,  $h_R(B) \rightarrow h_R(A) \times_{F(A)} F(B)$  is surjective.
- $(R, \xi)$  is called a **miniversal** family if it's versal, and maps

$$h_R(k[\theta]) \rightarrow F(k[\theta]), \quad h_R(k[t]/(t^2)) \rightarrow F(k[t]/(t^2)),$$

are bijective.

- $(R, \xi)$  is called a **universal** family if  $\varphi$  is an isomorphism.

Note that analogous to the classical case, if the the functor  $F : \mathbf{sArt}_k \rightarrow \mathbf{Sets}$  satisfies that  $F(k)$  has just one element and

$$F(A \times_k k[\theta]) \rightarrow F(A) \times F(k[\theta]), \quad F(A \times_k k[t]/(t^2)) \rightarrow F(A) \times F(k[t]/(t^2)), \quad (7.6)$$

are bijective for all  $A \in \mathbf{sArt}_k$ , then  $F(k[t, \theta]/(t^2, t\theta))$  has a natural structure of  $k$ -super vector space, which shall be denoted by  $T_F = T_F^+ \oplus T_F^-$ , where  $T_F^+ = F(k[t]/(t^2))$ ,  $T_F^- = F(k[\theta])$ .

**Theorem 7.12 (Schlessinger's Theorem).** The functor  $F : \mathbf{sArt}_k \rightarrow \mathbf{Sets}$  has a miniversal family if and only if

(S<sub>0</sub>)  $F(k)$  has just one element.

(S<sub>1</sub>)  $F(A' \times_A A'') \rightarrow F(A') \times_{F(A)} F(A'')$  is surjective for every small <sup>65</sup> extension  $A'' \rightarrow A$ .

(S<sub>2</sub>) The map in (S<sub>1</sub>) is bijective when  $A = k$  and  $A'' = k[\theta]$  or  $k[t]/(t^2)$ .

(S<sub>3</sub>)  $T_F$  is a finite-dimensional  $k$ -super vector space.

Moreover,  $F$  has a universal family if and only if in addition

(S<sub>4</sub>) For every small extension  $p : A'' \rightarrow A$  with even kernel (resp. odd kernel) and for every  $\eta \in F(A)$  which  $p^{-1}(\eta)$  is nonempty, the group action of  $T_F^+$  (resp.  $T_F^-$ ) on  $p^{-1}(\eta)$  is bijective.

The proof is almost the same as in the classical case (e.g. Theorem 16.2 of [89]), except that all ideals in the original proof should be  $\mathbb{Z}_2$ -graded in the super case, and the proof in the *loc.cit.* works formally.

---

<sup>65</sup>An extension  $0 \rightarrow I \rightarrow A'' \rightarrow A \rightarrow 0$  is called small if  $I$  is one dimensional.

## 7.2 Stable $\mathcal{N} = 1$ SUSY Curves

Assume that a triple of natural numbers  $(g, n_{\text{NS}}, n_{\text{R}})$  satisfies  $n_{\text{R}} \in 2\mathbb{Z}_{\geq 0}$ , and

$$2g - 2 + n_{\text{NS}} + n_{\text{R}} > 0.$$

**Definition 7.10 (Stable SUSY Curves).** *Given a superscheme  $S$ , a family of stable SUSY curves of genus  $g$  with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  R punctures over  $S$ , is a finite presented proper flat relatively Cohen-Macaulay  $S$ -superscheme  $X$ , such that the smooth open locus  $X^{\text{sm}} \subset X$  is of relative dimension  $(1|1)$  and is dense on each geometric fiber, together with*

1. *Sections  $E = E_1 \sqcup \cdots \sqcup E_{n_{\text{NS}}}$  inside  $X^{\text{sm}}$ .  $E_i$  will be called the  $i$ 'th NS punctures;*
2. *A closed sub superscheme  $\mathcal{D} = \mathcal{D}_1 \sqcup \cdots \sqcup \mathcal{D}_{n_{\text{R}}}$  inside  $X^{\text{sm}}$ , proper and flat over  $S$ , of codimension  $(1|0)$ , and  $\mathcal{D}_{i,s}$  is a point for every geometric point  $s \in S$ .  $\mathcal{D}_i$  will be called the  $i$ 'th R puncture;*
3. *An  $\mathcal{O}_S$ -linear derivation*

$$\delta : \mathcal{O}_X \rightarrow \omega_{X/S}(\mathcal{D}), \tag{7.7}$$

*or equivalently a homomorphism  $\delta : \Omega_{X/S}^1 \rightarrow \omega_{X/S}(\mathcal{D})$ .  $\delta$  is sometimes called the supersymmetric structure, or the SUSY structure;*

*such that*

- A. *For every geometric fiber  $X_s$ ,  $(X_s, \mathcal{O}_{X_s}^+, E_s, \mathcal{D}_s)$  is a punctured stable curve of genus  $g$ , and  $\delta^-$  induces isomorphism of  $\mathcal{O}_{X_s}^+$  modules*

$$\delta^- : \mathcal{O}_{X_s}^- \cong \omega_{X_s/s}^-(\mathcal{D}_s).$$

*Consequently the homomorphism  $\delta : \Omega_{X/S}^1 \rightarrow \omega_{X/S}(\mathcal{D})$  is surjective on the smooth locus of  $X$ .*

- B. *Restricted on  $X^{\text{sm}}$ , the dual of the surjective homomorphism  $\delta : \Omega_{X/S}^1 \rightarrow \omega_{X/S}(\mathcal{D})$ , write*

$$0 \rightarrow D \rightarrow T_{X/S} \rightarrow T_{X/S}/D \rightarrow 0,$$

*where  $D = \omega_{X/S}^\vee(-\mathcal{D})$ , induces an injective homomorphism  $D^{\otimes 2} \rightarrow T_{X/S}/D$  via  $D_1 \otimes D_2 \mapsto \frac{1}{2}[D_1, D_2]$  and the image is  $T_{X/S}/D(-\mathcal{D})$ .*

**Definition 7.11.** *An isomorphism between stable SUSY curves  $(X, E, \mathcal{D}, \delta)$  and  $(X', E', \mathcal{D}', \delta')$  over  $S$  is an  $S$ -isomorphism of superschemes  $f : X \rightarrow X'$  such that  $f^{-1}E' = E$ ,  $f^{-1}\mathcal{D}' = \mathcal{D}$ , and  $f^*\delta' = \delta$ .*

**Definition 7.12.** *The category of stable SUSY curves is the fibered groupoid over the category of superschemes, where the fiber over a superscheme  $S$  is the groupoid of stable SUSY curve of genus  $g$  with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  R punctures. Denote this category by  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$ .*

The following lemma is a direct consequence of the definition of stable SUSY curves:

**Lemma 7.23.** *Let  $(I, \leq)$  be a directed set, suppose that  $\{A_i\}_{i \in I}$  is an directed system of superrings, with inductive limit  $A$ , then the natural functor*

$$\varinjlim_i \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}(\text{Spec } A_i) \rightarrow \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}(\text{Spec } A),$$

*is an equivalence of categories.*

*Proof.* **Fully faithful:** first forgetting the derivation, then the natural map

$$\varinjlim_i \text{Map}_{\text{Spec } A_i}(X_i, Y_i) \rightarrow \text{Map}_{\text{Spec } A}(X, Y),$$

is an equivalence of sets, by Lemma 7.13. Now a morphism  $f$  preserves the derivation if and only if  $f^*\delta_Y - \delta_X : \Omega_{X/S}^1 \rightarrow \omega_{X/S}(\mathcal{D})$  is zero, then the result follows from Lemma 7.15.

**Essentially surjective:** the functor is essentially surjective if we forget the conditions A and B in the Definition 7.10, this follows from the Lemmas 7.13, 7.14, 7.15, and Proposition 7.1. Note that  $X \rightarrow S$  is proper if and only if  $X_{\text{bos}} \rightarrow S_{\text{bos}}$  is proper, so we can apply Theorem 8.10.5 of [88] to ensure that there exists  $i \in I$  such that  $X_i \rightarrow S_i$  is proper.

Now if the derivation  $\delta$  on  $X$  satisfies condition A, then the derivation  $\delta_i$  on  $X_i$  automatically satisfies condition A, since a homomorphism of sheaves on a variety over a field is an isomorphism if and only if it is an isomorphism under some field extension. If the derivation  $\delta$  on  $X$  satisfies condition B, then it follows that the natural map  $D_i^{\otimes 2} \rightarrow T_{X_i/S_i}/D_i$  is injective on  $X_i^{\text{sm}}$  since it is a map between locally free  $\mathcal{O}_{X_i}$  sheaves and is injective on geometric fibers  $X_{i,s}^{\text{sm}}$  for all geometric point  $s$ . Moreover after increasing  $i$ , the composition  $D_i^{\otimes 2} \rightarrow T_{X_i/S_i}/D_i \rightarrow T_{X_i/S_i}/D_i \otimes \mathcal{O}_{\mathcal{D}_i}$  becomes zero, since it is zero after taking limit, so the image of  $D_i^{\otimes 2}$  in  $T_{X_i/S_i}/D_i$  is a subsheaf of  $T_{X_i/S_i}/D_i(-\mathcal{D}_i)$ , thus it must be  $T_{X_i/S_i}/D_i(-\mathcal{D}_i)$  since the induced map  $D_i^{\otimes 2} \rightarrow T_{X_i/S_i}/D_i(-\mathcal{D}_i)$  is an isomorphism on geometric fibers  $X_{i,s}^{\text{sm}}$ . This concludes that  $\delta_i$  satisfies condition B.  $\blacksquare$

By usual argument (see e.g. Lemma 3.1 of [6]), the condition B in Definition 7.10 tells that away from an R puncture, locally there is a coordinate system  $(z|\theta)$ , such that  $\delta$  can be written as

$$\delta : f \mapsto (\partial_\theta f + \theta \partial_z f)[dz|d\theta],$$

and nearby a R puncture there is a local coordinate system  $(z|\theta)$ , such that  $z = 0$  is the location of R puncture and  $\delta$  can be written as

$$\delta : f \mapsto (\partial_\theta f + \theta z \partial_z f) \left[ \frac{dz}{z} | d\theta \right].$$

On the other hand, suppose that  $S$  is a bosonic scheme, then condition A in the Definition 7.10 tells that  $(X, \mathcal{O}_X^+, E, \mathcal{D})$  is a pointed stable curve of genus  $g$ . The Cohen-Macaulay

condition implies that  $\mathcal{O}_X^-$  is a relatively torsion free  $\mathcal{O}_X^+$  sheaf, and it is of rank one because it is a line bundle on  $U$ . This implies that

$$\mathcal{O}_X^- \cdot \mathcal{O}_X^- = 0, \quad (7.8)$$

since it is generically zero and  $\mathcal{O}_X^+$  is relatively Cohen-Macaulay. Moreover the odd part of  $\delta$  induces an isomorphism

$$\delta^- : \mathcal{O}_X^- \cong \omega_{X/S}^-(\mathcal{D}),$$

since  $\delta_s^-$  is an isomorphism on each fiber  $X_s$ . Note that the relative dualizing complex (in fact a sheaf) is

$$\omega_{X/S} = R\mathbf{Hom}_{\mathcal{O}_X^+}(\mathcal{O}_X, \omega_{X_{\text{bos}}/S}) = \mathbf{Hom}_{\mathcal{O}_X^+}(\mathcal{O}_X, \omega_{X_{\text{bos}}/S}),$$

and it decomposes as  $\mathcal{O}_X^+$ -module into

$$\omega_{X/S}^+ = \omega_{X_{\text{bos}}/S}, \quad \omega_{X/S}^- \cong \mathbf{Hom}_{\mathcal{O}_X^+}(\mathcal{O}_X^-, \omega_{X_{\text{bos}}/S}).$$

This realizes  $(\mathcal{O}_X^-, \delta^-)$  as a spin structure on the punctured stable curve  $(X, \mathcal{O}_X^+, E, \mathcal{D})$ . Finally, the condition B Definition 7.10 uniquely determines the even part of  $\delta$ , in fact  $\delta^+ : \mathcal{O}_X^+ \rightarrow \omega_{X/S}^+(\mathcal{D})$  is the composition of exterior differential  $d : \mathcal{O}_X^+ \rightarrow \omega_{X/S}^+$  with natural embedding  $\omega_{X/S}^+ \hookrightarrow \omega_{X/S}^+(\mathcal{D})$ , because this is the case on the open  $S$ -dense locus  $U$  (see the coordinate presentation of  $\delta$  above), and  $\mathcal{O}_X \rightarrow j_*\mathcal{O}_U$  is injective where  $j : U \hookrightarrow X$  is the open embedding. As a summary,

**Proposition 7.5.** *Given a bosonic scheme  $S$ , a family of stable  $\mathcal{N} = 1$  SUSY curves of genus  $g$  with  $n_{\text{NS}}$  NS punctures and  $n_{\text{R}}$  R punctures over  $S$ , is a superscheme  $(X, \mathcal{O}_X)$ , flat over  $S$ , such that  $\mathcal{O}_X = \mathcal{O}_X^+ \oplus \mathcal{O}_X^-$ ,  $\mathcal{O}_X^-$  has zero multiplication with itself,  $X_{\text{bos}} := (X, \mathcal{O}_X^+)$  is a relative stable curve with puncturing divisors  $E \sqcup \mathcal{D}$ , with  $\deg E_s = n_{\text{NS}}$ ,  $\deg \mathcal{D}_s = n_{\text{R}}$ , and  $\mathcal{O}_X^-$  is a rank one relatively torsion free sheaf over  $X_{\text{bos}}$  with an isomorphism*

$$a : \mathcal{O}_X^- \rightarrow \mathbf{Hom}_{\mathcal{O}_X^+}(\mathcal{O}_X^-, \omega_{X_{\text{bos}}/S}(\mathcal{D})). \quad (7.9)$$

**Remark 7.5.** *From the proposition, it is straightforward to see that the data of an stable  $\mathcal{N} = 1$  punctured SUSY curve over a scheme  $S$  is equivalent to the data of a relative punctured spin curve over  $S$  (Definition B.5).*

Recall that on a stable spin curve  $(\mathcal{C}, E, \mathcal{D}, \mathcal{E})$  over a bosonic base scheme  $S$ , the line bundle  $\mathcal{L} = \omega_{\mathcal{C}/S}(E + \mathcal{D})$  is relatively ample and  $\mathcal{L}^{\otimes 3}$  is relatively very ample. Consider a stable SUSY curve  $X$  over  $S$  with underlying spin curve  $(\mathcal{C}, E, \mathcal{D}, \mathcal{E})$ , since  $p : X \rightarrow \mathcal{C}$  is finite, we see that  $p^*\mathcal{L}$  is a relatively ample line bundle of rank  $(1|0)$  on  $X$ . Consequently, we have

**Lemma 7.24.** *Let  $S$  be a Noetherian affine superscheme and  $X \rightarrow S$  is a stable SUSY curve, then  $X$  is projective over  $S$ .*

*Proof.* Suppose that  $\mathfrak{n}_S^{N+1} = 0$  and the restriction of  $X$  on  $(S, \mathcal{O}_S/\mathfrak{n}_S^N)$  has an ample line bundle  $\mathcal{L}$ , we first lift  $\mathcal{L}$  to  $X$ : This can be done locally on an affine covering  $U_i$ , since  $\mathcal{L}$  is locally isomorphic to the structure sheaf; from local to global we need to lift the transition function  $\varphi_{ij}$  from  $\Gamma(U_{ij}, \mathcal{O}_X/\mathfrak{n}_S^N \mathcal{O}_X)_+$  to  $\Gamma(U_{ij}, \mathcal{O}_X)_+$ , we make an arbitrary choice  $\tilde{\varphi}_{ij}$ ; then

$$\tilde{\varphi}_{ij} \circ \tilde{\varphi}_{jk} \circ \tilde{\varphi}_{ki} - 1 \in \Gamma(U_{ijk}, \mathfrak{n}_S^N \mathcal{O}_X)_+,$$

defines a Čech cocycle valued in the coherent sheaf  $\mathfrak{n}_S^N \mathcal{O}_X$ , but  $S$  is affine and the dimensions of fibers  $X_s$  are 1 hence  $H^2$  of a coherent sheaf vanishes; so we can modify  $\tilde{\varphi}_{ij}$  by an element in  $1 + \Gamma(U_{ij}, \mathfrak{n}_S^N \mathcal{O}_X)_+$  to force it satisfying cocycle conditions, hence local liftings glue to a global one. Now the lift  $\tilde{\mathcal{L}}$  is ample since  $X \rightarrow S$  is proper and  $\mathcal{L}_s$  is ample for all geometric point  $s \in S$ .  $\blacksquare$

### 7.2.1 Étale Descent for $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$

Suppose that  $B$  is a Noetherian superring,  $S = \text{Spec } B$ ,  $B_1$  is an étale  $B$  superalgebra, let  $B_2 = B_1 \otimes_B B_1$ ,  $B_3 = B_1 \otimes_B B_1 \otimes_B B_1$ ,  $S_i = \text{Spec } B_i$ . Let  $p_1, p_2$  be two canonical projections from  $S_2$  to  $S_1$ .

**Lemma 7.25.** *Suppose that  $M, M'$  are  $B$ -modules, let  $M_i = M \otimes_B B_i$ ,  $M'_i = M' \otimes_B B_i$ , then the sequence*

$$0 \rightarrow \text{Hom}_B(M, M') \rightarrow \text{Hom}_{B_1}(M_1, M'_1) \rightrightarrows \text{Hom}_{B_1}(M_2, M'_2), \quad (7.10)$$

*is exact.*

*Proof.* Let  $B = B^+ \oplus B^-$  where  $B^+$  and  $B^-$  are its even and odd parts. Considering similar decomposition for  $B_i$ , then  $B^+ \rightarrow B_1^+$  is étale,  $B_2^+ = B_1^+ \otimes_{B^+} B_1^+$ , and  $B_i^- = B_i^+ \otimes_{B^+} B^-$ , by Lemma 7.10. If a  $B$ -homomorphism  $\varphi : M \rightarrow M'$  is zero after base change to  $B_1$ , then it is zero as a  $B^+$ -homomorphism, by classical descent theory, thus  $\varphi = 0$ . If there is a  $B_1$ -homomorphism  $\psi : M_1 \rightarrow M'_1$  whose base changes to  $B_2$  in two ways agree, then there exists a  $B^+$ -homomorphism  $\rho : M \rightarrow M'$  whose base change to  $B_1^+$  is  $\psi$  (as  $B_1^+$ -homomorphism). Note that a  $B^+$ -homomorphism is a  $B$ -homomorphism if it commutes with the  $B^-$  action on  $M$  and  $M'$ . Now the base change of  $\rho$  to  $B_1^+$  commutes with the action of  $B_1^-$ , so  $\rho$  commutes with the  $B^-$  action, by classical descent theory. Hence we get a  $B$ -homomorphism  $\rho : M \rightarrow M'$  whose base change to  $B_1$  is  $\psi$ .  $\blacksquare$

**Lemma 7.26.** *Suppose that  $N$  is a  $B_1$  module with an isomorphism*

$$\phi : p_1^* N \xrightarrow{\sim} p_2^* N,$$

*satisfying cocycle condition. Then there is a  $B$  module  $M$  with isomorphism  $\psi : M \otimes_B B_1 \xrightarrow{\sim} N$  such that  $\phi$  is the canonical isomorphism  $p_2^* \psi \circ p_1^* \psi^{-1} : p_1^* N \xrightarrow{\sim} p_2^* N$ . Moreover,  $N$  is a flat (resp. finite type)  $B_1$ -module if and only if  $M$  is a flat (resp. finite type)  $B$ -module.*



*Proof.* Let  $B = B^+ \oplus B^-$  where  $B^+$  and  $B^-$  are its even and odd parts, similar for  $B_i$ , then  $B^+ \rightarrow B_1^+$  is étale,  $B_2^+ = B_1^+ \otimes_{B^+} B_1^+$ , and  $B_i^- = B_i^+ \otimes_{B^+} B^-$ , by Lemma 7.10. Then it follows that  $p_i^* N \cong p_{i,\text{ev}}^* N$  when  $N$  is considered as a  $\mathbb{Z}_2$ -graded  $B_1^+$ -module, and  $p_{i,\text{ev}} : S_{2,\text{ev}} \rightarrow S_{1,\text{ev}}$  are two natural projections. Moreover, the isomorphism  $\phi$  preserves  $\mathbb{Z}_2$ -grading. By classical descent theory, we have a  $B^+$ -module  $M = M^+ \oplus M^-$  with isomorphism  $\psi : M^\pm \otimes_{B^+} B_1^+ \xrightarrow{\sim} N^\pm$  such that  $\phi$  is the canonical isomorphism  $p_{2,\text{ev}}^* \psi \circ p_{1,\text{ev}}^* \psi^{-1} : p_{1,\text{ev}}^* N^\pm \xrightarrow{\sim} p_{2,\text{ev}}^* N^\pm$ . Note that  $\mathbb{Z}_2$ -graded  $B^+$  module  $M = M^+ \oplus M^-$  has a  $B$ -module structure if and only if there is a pair of  $B^+$ -homomorphism  $a : B^- \otimes_{B^+} M^\pm \rightarrow M^\mp$  such that the diagram

$$\begin{array}{ccc} B^- \otimes_{B^+} B^- \otimes_{B^+} M^\pm & \xrightarrow{\text{Id} \otimes a} & B^- \otimes_{B^+} M^\mp \\ & \searrow \mu \otimes \text{Id} & \downarrow a \\ & & M^\pm \end{array},$$

commutes. Now the existence of homomorphism  $a$  comes from the classical descent theory, and the commutativity of the diagram follows as well. The statement of flatness comes from the fact that  $B \rightarrow B_1$  is faithfully flat, and statement of finiteness comes from reducing  $B$  to  $B/\mathfrak{n}_B$  and use the classical descent theory.  $\blacksquare$

**Remark 7.6.** *In fact we can relax the condition  $B \rightarrow B_1$  being étale to that  $B \rightarrow B_1$  is flat and  $\mathfrak{n}_B$  generates  $\mathfrak{n}_{B_1}$ , then the trick of decomposing  $B = B^+ \oplus B^-$  still applies and Lemmas 7.25 and 7.26 is still true under this condition.*

Consider a stable SUSY curve  $X_1$  over  $S_1$ , and an isomorphism

$$\varphi : X_1 \times_{S_1, p_1} S_2 \xrightarrow{\sim} X_1 \times_{S_1, p_2} S_2,$$

satisfying the cocycle condition.

**Lemma 7.27.** *There exists an open affine cover  $\{U_\alpha \rightarrow X_1\}$ , such that  $\varphi$  maps  $U_\alpha \times_{S_1, p_1} S_2$  to  $U_\alpha \times_{S_1, p_2} S_2$ .*

*Proof.* According to the affineness criterion Proposition 7.3, it suffices to prove this lemma under the additional assumption that  $B$  is purely bosonic<sup>66</sup>, then  $X_1$  is a spin curve  $(\mathcal{C}_1, E, \mathcal{D}, \mathcal{E})$  together with the derivation induced from the structure map of the spinor sheaf  $\mathcal{E}$ . By descent theory for classical schemes, we get a stable SUSY curve  $X$  on  $S$  with isomorphism  $\psi : X \times_S S_1 \rightarrow X_1$  such that  $\varphi$  is the canonical isomorphism  $p_2^* \psi \circ p_1^* \psi^{-1} : X_1 \times_{S_1, p_1} S_2 \rightarrow X_1 \times_{S_1, p_2} S_2$ . Take an open affine cover  $\{V_\alpha\}$  of  $X$ , then the base change of  $V_\alpha$  is what we want.  $\blacksquare$

**Proposition 7.6.** *There exists a stable SUSY curve  $X$  over  $S$  with an isomorphism  $\psi : X \times_S S_1 \rightarrow X_1$  such that  $\varphi$  is the canonical isomorphism  $p_2^* \psi \circ p_1^* \psi^{-1} : X_1 \times_{S_1, p_1} S_2 \xrightarrow{\sim} X_1 \times_{S_1, p_2} S_2$ .*

<sup>66</sup>Then it follows from the definition of étale morphism that  $B_i$  are also bosonic.

*Proof.* We take the cover  $\{U_\alpha \rightarrow X_1\}$  in the Lemma 7.27, then  $\mathcal{O}_X(U_\alpha)$  is a flat  $B_1$ -superalgebra with a gluing isomorphism when base changed to  $B_2$ . By Lemmas 7.25 and 7.26, there exists a flat and finite type  $B$ -superalgebra  $C_\alpha$  with isomorphism  $\psi_\alpha : C_\alpha \otimes_B B_1 \xrightarrow{\sim} \mathcal{O}_X(U_\alpha)$  such that  $\varphi|_{U_\alpha \times_{S_1, p_1} S_2}$  is  $p_{2, \alpha}^* \psi \circ p_{1, \alpha}^* \psi^{-1}$ . Moreover  $C_\alpha$  is relatively Cohen-Macaulay since its geometric fibers are Cohen-Macaulay. By Lemma 7.25 again, we obtain a Zariski gluing data for  $\{\text{Spec } C_\alpha\}$ , thus there exists an  $S$ -superscheme  $X$  with an isomorphism  $\psi : X \times_S S_1 \rightarrow X_1$  and  $\varphi = p_2^* \psi \circ p_1^* \psi^{-1}$ .  $X$  is proper and its smooth locus  $X^{\text{sm}}$  is dense in every geometric fiber, since  $X \times_S S_{\text{bos}}$  is the descent for  $X_1 \times_{S_1} S_{1, \text{bos}}$  and classical descent theory is applied. The puncture divisors  $E, \mathcal{D}$  are determined by their ideal of definition, so they also descent to  $X$ , by Lemma 7.26. We also have a homomorphism  $\delta : \Omega_{X/S}^1 \rightarrow \omega_{X/S}(\mathcal{D})$  by Lemma 7.25.  $\delta$  satisfies Condition A in Definition 7.10 since it is a condition on geometric fibers;  $\delta$  satisfies Condition B as well, since  $D^{\otimes 2} \rightarrow T_{X/S}/D$  is injective after base change to  $S_1$ , and the image is  $T_{X/S}/D(-\mathcal{D})$  because it is after base change to  $S_1$ .  $\blacksquare$

### 7.2.2 Local Structures of Stable SUSY Curves

Before giving the characterization of the local structures of stable SUSY curves, let us define some canonical local models. Take  $S = \text{Spec } A$ , where  $A = A^+ \oplus A^-$  is a superring

1. Smooth Canonical Model:  $X = \text{Spec } A[z|\theta]$ , with derivation

$$\delta : f \mapsto (\partial_\theta f + \theta \partial_z f)[dz|d\theta].$$

2. R Puncture Canonical Model:  $X = \text{Spec } A[z|\theta]$ , with derivation

$$\delta : f \mapsto (\partial_\theta f + \theta z \partial_z f) \left[ \frac{dz}{z} \middle| d\theta \right].$$

3. R Node Canonical Model:  $X = \text{Spec } A[z_1, z_2|\theta]/(z_1 z_2 + t)$  where  $t \in A^+, \alpha \in A^-$ , with derivation

$$\delta : f \mapsto \begin{cases} (\partial_\theta f + \theta z_1 \partial_{z_1} f) \left[ \frac{dz_1}{z_1} \middle| d\theta \right] & \text{on } D(z_1) \\ (\partial_\theta f - \theta z_2 \partial_{z_2} f) \left[ -\frac{dz_2}{z_2} \middle| d\theta \right] & \text{on } D(z_2) \end{cases}.$$

4. NS Node Canonical Model:  $X = \text{Spec } A[z_1, z_2|\theta_1, \theta_2]/(z_1 z_2 + t^2, z_1 \theta_2 - t \theta_1, z_2 \theta_1 + t \theta_2, \theta_1 \theta_2)$  where  $t \in A^+$ , with derivation

$$\delta : f \mapsto \begin{cases} (\partial_{\theta_1} f + \theta_1 \partial_{z_1} f)[dz_1|d\theta_1] & \text{on } D(z_1) \\ (\partial_{\theta_2} f + \theta_2 \partial_{z_2} f)[dz_2|d\theta_2] & \text{on } D(z_2) \end{cases}.$$

Let  $S$  be a bosonic scheme and  $X/S$  is a stable SUSY curve. Suppose that  $S = \text{Spec } A$  where  $A$  is a complete local Noetherian ring with algebraically-closed residue field  $\kappa$ . Take a point  $p$  in the special fiber of  $X$ , and take completion of  $\mathcal{O}_X$  at  $p$ , then according to the local structure of spin curves (Proposition B.7), there are three possibilities:

- a.  $p$  is a smooth point:  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z|\theta]]$ .
- b.  $p$  is an R node:  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z_1, z_2|\theta]]/(z_1 z_2 + t)$  where  $t \in A$ .
- c.  $p$  is an NS node:  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z_1, z_2|\theta_1, \theta_2]]/(z_1 z_2 + t^2, z_1 \theta_2 + t \theta_1, z_2 \theta_1 - t \theta_2, \theta_1 \theta_2)$  where  $t \in A$ .

To figure out the derivation, let us first write down a set of generators of  $\omega_{X/S}$  as  $\widehat{\mathcal{O}}_{X,p}^+$  module in each case:

- a.  $p$  is a smooth point:  $\omega_{X/S}$  is generated by  $\mathfrak{b}^- = [dz|d\theta]$  and  $\mathfrak{b}^+ = \theta[dz|d\theta]$ .
- b.  $p$  is a R node:  $\omega_{X/S}$  is generated by

$$\mathfrak{b}^- = \begin{cases} \left[ \frac{dz_1}{z_1} \middle| d\theta \right] & \text{on } D(z_1) \\ \left[ -\frac{dz_2}{z_2} \middle| d\theta \right] & \text{on } D(z_2) \end{cases}, \quad \mathfrak{b}^+ = \begin{cases} \theta \left[ \frac{dz_1}{z_1} \middle| d\theta \right] & \text{on } D(z_1) \\ \theta \left[ -\frac{dz_2}{z_2} \middle| d\theta \right] & \text{on } D(z_2) \end{cases}.$$

- c.  $p$  is a NS node:  $\omega_{X/S}$  is generated by

$$\mathfrak{b}_1^- = \begin{cases} [dz_1|d\theta_1] & \text{on } D(z_1) \\ \frac{t}{z_2} [dz_2|d\theta_2] & \text{on } D(z_2) \end{cases}, \quad \mathfrak{b}_2^- = \begin{cases} -\frac{t}{z_1} [dz_1|d\theta_1] & \text{on } D(z_1) \\ [dz_2|d\theta_2] & \text{on } D(z_2) \end{cases}.$$

and

$$\mathfrak{b}^+ = \begin{cases} \frac{\theta_1}{z_1} [dz_1|d\theta_1] & \text{on } D(z_1) \\ -\frac{\theta_2}{z_2} [dz_2|d\theta_2] & \text{on } D(z_2) \end{cases}.$$

By construction, if  $p$  is a smooth point but not an R node, then  $\delta$  maps  $\theta$  to  $\mathfrak{b}^-$  and maps  $z$  to  $\mathfrak{b}^+$ ; otherwise  $\delta$  maps  $\theta_i$  to  $\mathfrak{b}_i^-$  and maps  $z_i$  to  $\pm z_i \mathfrak{b}^+$ . It's easy to check that this is exactly the derivation in the canonical models. Hence we have

**Lemma 7.28.** *Let  $S = \text{Spec } A$  where  $A$  is a complete local Noetherian ring and  $X/S$  is a stable SUSY curve. Take a point  $p$  in the special fiber of  $X$ , then the completion of  $\mathcal{O}_X$  at  $p$  together with derivation  $\delta$  is isomorphic to the completion of one of the canonical models at the origin of the special fiber.*

The next goal is to extend this result to a more general situation where  $A$  can be a complete local Noetherian superring. The idea is to first prove it for local Artinian  $A$ , then take the inverse limit. For local Artinian  $A$ , use the induction on the length of  $A$ , more precisely, suppose that there is an extension

$$0 \rightarrow \mathcal{J} \rightarrow A \rightarrow A/\mathcal{J} \rightarrow 0,$$

where  $\mathcal{J}$  is an ideal of  $A$ , purely bosonic or purely fermionic, such that the maximal ideal  $\mathfrak{m}_A$  annihilates it. If Lemma 7.28 is true for  $A/\mathcal{J}$ , then we want to show that Lemma 7.28 is true for  $A$ . Denote the residue field of  $A$  by  $\kappa$ .

To begin with, let us temporarily forget the derivation and try to figure out the underlying scheme structure of  $\widehat{\mathcal{O}}_{X,p}$ , assuming that  $\widehat{\mathcal{O}}_{X,p}/\mathcal{J}\widehat{\mathcal{O}}_{X,p}$  is isomorphic to the completion of one of the canonical models at the origin of the special fiber. The following list of possibilities is a direct consequence of flatness:

- a.  $p$  is a smooth point:  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z|\theta]]$ .
- b.  $p$  is an R node:  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z_1, z_2|\theta]]/(z_1z_2 + t' + \eta\theta)$  where  $t' \in A^+$ ,  $\eta \in \mathcal{J}$  such that  $t' \equiv t \pmod{\mathcal{J}}$ .
- c.  $p$  is an NS node:  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z_1, z_2|\theta_1, \theta_2]]/(z_1z_2 + t_1t_2 + a, z_1\theta_2 + t_1\theta_1 + b, z_2\theta_1 - t_2\theta_2 + c, \theta_1\theta_2 + d)$  where  $t_i \in A^+$  such that  $t_i \equiv t \pmod{\mathcal{J}}$  and

$$a, b, c, d \in B := \mathcal{J} \otimes_{\kappa} \widehat{\mathcal{O}}_{X,p}/\mathfrak{m}_A \widehat{\mathcal{O}}_{X,p} = \mathcal{J} \widehat{\mathcal{O}}_{X,p}.$$

In fact, suppose that  $Y$  is stable SUSY curve over  $\text{Spec } A/\mathcal{J}$ , then for any such  $p \in Y$ , take an open affine neighborhood  $W_p$  of  $p$  such that  $W_p$  contains no other node (if  $p$  is a node) or puncture divisor, then by Theorem 2.1.2 of [84], the obstruction of lifting  $W_p$  to a flat superscheme over  $\text{Spec } A$  is an element in  $H^2(T_{W_{p,s}}^\bullet)$  where  $W_{p,s}$  is the special fiber of  $W_p$ . Note that  $H^i(T_{W_{p,s}}^\bullet)$  is supported at  $p$  for  $i > 0$ , since  $W_p \setminus \{p\}$  is smooth. If the obstruction vanishes, then the set of deformation modulo infinitesimal automorphism  $\text{Def}_A(W_p, \mathcal{J})$  is a torsor under  $H^1(T_{W_{p,s}}^\bullet)$ , and we also have

$$\text{Def}_A(W_p, \mathcal{J}) = \text{Def}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J}). \quad (7.11)$$

In fact, the obstruction always vanishes. For the situation a, i.e.  $p$  is a smooth point,  $H^i(T_{W_{p,s}}^\bullet)$  vanishes for  $i > 0$ ; for situation b, i.e.  $p$  is an R node,  $H^2(T_{W_{p,s}}^\bullet)$  vanishes since the relation defining  $\widehat{\mathcal{O}}_{Y,p}$  in  $A[[z_1, z_2|\theta]]$  is  $z_1z_2 + t$  which is not a zero-divisor, i.e. there is no syzygy; for the situation c, i.e.  $p$  is an NS node, the obstruction of lifting  $W_p$  is identified with the obstruction of lifting  $\text{Spec } \widehat{\mathcal{O}}_{Y,p}$ , which vanishes because there exists a lifting:  $A[[z_1, z_2|\theta_1, \theta_2]]/(z_1z_2 + t_1t_2, z_1\theta_2 + t_1\theta_1, z_2\theta_1 - t_2\theta_2, \theta_1\theta_2)$ <sup>67</sup>.

Following [29], we can put more constraints on the elements  $a, b, c, d$  in the case c. Namely,  $a, b, c, d$  are components of an element in the  $H^1$  of the chain complex

$$B\partial_{z_i} \oplus B\partial_{\theta_i} \xrightarrow{\frac{\partial f_j}{\partial z_i} \oplus \frac{\partial f_j}{\partial \theta_i}} B\partial_{f_j} \xrightarrow{h_{kj}} B\partial_{S_k},$$

where  $f_j$  are relations of  $\widehat{\mathcal{O}}_{X,p}/\mathfrak{m}_A \widehat{\mathcal{O}}_{X,p}$  and  $S_k = \sum_j h_{kj} f_j$  are the syzygies, namely

$$\begin{cases} f_1 = z_1z_2 \\ f_2 = z_1\theta_2 \\ f_3 = z_2\theta_1 \\ f_4 = \theta_1\theta_2 \end{cases}, \quad \begin{cases} S_1 = \theta_1f_1 - z_1f_3 \\ S_2 = \theta_2f_1 - z_2f_2 \\ S_3 = z_1f_4 - \theta_1f_2 \\ S_4 = z_2f_4 + \theta_2f_3 \end{cases}.$$

<sup>67</sup>Notice that this is actually a germ of a nodal curve over  $A$  defined by  $A[[z_1, z_2]]/(z_1z_2 + t_1t_2)$ , together with a rank one torsion-free sheaf  $\mathcal{E}$  on it generated by  $\theta_1, \theta_2$  with relation  $z_1\theta_2 + t_1\theta_1 = 0, z_2\theta_1 - t_2\theta_2 = 0$ , and obvious relation  $\theta_1\theta_2 = 0$  to ensure that  $\mathcal{E} \cdot \mathcal{E} = 0$ .

So we get four equations:

$$\theta_1 a = z_1 c, \theta_2 a = z_2 b, z_1 d = \theta_1 b, z_2 d = -\theta_2 c,$$

modulo the image of the matrix

$$M = \begin{bmatrix} z_2 & z_1 & 0 & 0 \\ \theta_2 & 0 & 0 & z_1 \\ 0 & \theta_1 & z_2 & 0 \\ 0 & 0 & \theta_2 & -\theta_1 \end{bmatrix},$$

which corresponds to coordinate transformation. If  $N$  is odd, then  $d \in (\theta_1, \theta_2)$  and is even, so it can be written as  $d = \lambda_1 \theta_1 + \lambda_2 \theta_2$ , and a coordinate transform  $\theta_1 \mapsto \theta_1 - \lambda_2, \theta_2 \mapsto \theta_2 - \lambda_1$  will set  $d$  to zero; if  $N$  is even, then  $b, c \in (\theta_1, \theta_2)$ , so the last two equations become  $z_1 d = z_2 d = 0$ , hence  $d = 0$ . So in any case we can set  $d = 0$ . In the following discussion, we only consider coordinate transform whose image is in  $B\partial_{f_1} \oplus B\partial_{f_2} \oplus B\partial_{f_3}$ , for example the image of the matrix

$$\begin{bmatrix} z_2 & z_1 & 0 & 0 \\ \theta_2 & 0 & 0 & -z_1 \theta_1 \\ 0 & \theta_1 & z_2 \theta_2 & 0 \end{bmatrix}.$$

Using this coordinate transformation, together with the observation that any constant  $\times \theta_1$  term of  $b$  can be absorbed by  $t_1 \theta_1$  and any constant  $\times \theta_2$  term of  $c$  can be absorbed by  $t_2 \theta_2$ , we can transform  $b$  and  $c$  to Laurent polynomials in  $z_1, z_2$  simultaneously, i.e.

$$b = f_1(z_1) + z_2 f_2(z_2), \quad c = z_1 g_1(z_1) + g_2(z_2).$$

Moreover, the equations  $\theta_1 b = \theta_2 c = 0$  spells out that

$$f_1(z_1) \theta_1 + t f_2(z_2) \theta_2 = g_2(z_2) \theta_2 - t g_1(z_1) \theta_1 = 0,$$

hence

$$f_1(z_1) = t f_2(z_2) = g_2(z_2) = t g_1(z_1) = 0.$$

Let us turn to consider equations

$$\theta_1 a = z_1 c, \quad \theta_2 a = z_2 b.$$

Since  $b, c$  has no  $\theta_i$  dependence, the equations become

$$\theta_1 a = z_1 c = \theta_2 a = z_2 b = 0.$$

We thus see that  $a \in (\theta_1, \theta_2)$  and  $f_2(z_2) = g_1(z_1) = 0$ , i.e.  $b = c = 0$ . Now we can use coordinate transforms with images in  $B\partial_{f_1}$  to modify  $a$ , for example we can choose  $\theta_i$  times

the first and the second rows of the matrix  $M$  hence all terms in  $a$  involving  $z_i\theta_j$  can be removed, leaving

$$a = \eta_1\theta_1 + \eta_2\theta_2,$$

where  $\eta_i \in \mathcal{J}$ . To sum up, we have that

- When  $p$  is an NS node,  $\widehat{\mathcal{O}}_{X,p} \cong \text{Spec } A[[z_1, z_2 | \theta_1, \theta_2]] / (z_1z_2 + t_1t_2 + \eta_1\theta_1 + \eta_2\theta_2, z_1\theta_2 + t_1\theta_1, z_2\theta_1 - t_2\theta_2, \theta_1\theta_2)$  where  $t_i \in A^+$  such that  $t_i \equiv t \pmod{\mathcal{J}}$  and  $\eta_i \in \mathcal{J}$ .

Next we take the derivation  $\delta$  into account, in fact we have

**Lemma 7.29.**  *$\widehat{\mathcal{O}}_{X,p}$  described above has a SUSY structure derivation  $\delta$  extending the derivation on  $\widehat{\mathcal{O}}_{X,p}/\mathcal{J}\widehat{\mathcal{O}}_{X,p}$  if and only if*

- If  $p$  is a smooth point, there is no condition.*
- If  $p$  is a R node, then  $\eta = 0$ .*
- If  $p$  is a NS node, then  $\eta_i = 0$  and  $t_1 = t_2$ .*

In other words, derivations extend exactly when  $\widehat{\mathcal{O}}_{X,p}$  is isomorphic to the completion of one of the canonical models at the origin of the special fiber. Moreover, the choice of extension is unique in all three cases, in the sense that there exists coordinate transform under which derivations are transformed to that of the corresponding canonical models.

Assume this lemma for now, then it concludes the induction step and we have the following generalization of Lemma 7.28.

**Proposition 7.7.** *Let  $S = \text{Spec } A$  where  $A$  is a complete local Noetherian superring and  $X/S$  is a stable SUSY curve. Take a point  $p$  in the special fiber of  $X$ , then the completion of  $\mathcal{O}_X$  at  $p$  together with derivation  $\delta$  is isomorphic to the completion of one of the canonical models at the origin of the special fiber.*

*Proof of Lemma 7.29.* We consider each cases separately

**If  $p$  is smooth,**  $\widehat{\mathcal{O}}_{X,p}$  is already isomorphic to that of canonical models so we can always choose the derivation in the smooth or R puncture canonical models, and the uniqueness follows from the condition B of Definition 7.10.

**If  $p$  is a R node,** write

$$\delta = \delta_0 + \delta',$$

where  $\delta_0$  is

$$f \mapsto \begin{cases} (\partial_\theta f + \theta z_1 \partial_{z_1} f) \left[ \frac{dz_1}{z_1} \Big| d\theta \right], & \text{on } D(z_1) \\ (\partial_\theta f - \theta z_2 \partial_{z_2} f) \left[ -\frac{dz_2}{z_2} \Big| d\theta \right], & \text{on } D(z_2) \end{cases},$$

and

$$\delta' \in \text{Der}_\kappa (\mathcal{O}_U, \mathcal{J} \otimes_\kappa \omega_{U/\text{Spec } \kappa}),$$

where  $U = \text{Spec}(\widehat{\mathcal{O}}_{X,p}/\mathfrak{m}_A \widehat{\mathcal{O}}_{X,p}) \setminus \{p\}$  is a disjoint union of  $\text{Spec } \kappa((z_1))$  and  $\text{Spec } \kappa((z_2))$ . So we can write down  $\delta'$  as

$$\delta' f = (xDf + yD^2f)\mathfrak{b}^-, \quad D = \begin{cases} (\partial_\theta f + \theta z_1 \partial_{z_1} f) & \text{on } D(z_1) \\ (\partial_\theta f - \theta z_2 \partial_{z_2} f) & \text{on } D(z_2) \end{cases},$$

where  $x, y \in \mathcal{J} \otimes_\kappa \mathcal{O}_U$  such that  $x$  is even and  $y$  is odd. Using Lemma 1.1 of [29], we have that  $x + Dy = 0$ , so  $\delta'$  can be written as

$$\delta' : f \mapsto -D(y \cdot Df)\mathfrak{b}^-.$$

Consider  $\delta' \theta = (1 - Dy)\mathfrak{b}^- \in \omega_{X/S}$ , this implies that  $y \in \mathcal{J}[[z_1]] \times \mathcal{J}[[z_2]]$  or  $\theta \mathcal{J}[[z_1]] \times \theta \mathcal{J}[[z_2]]$  depending on the parity of  $\mathcal{J}$ . Now consider

$$\delta z_1 = \begin{cases} (\theta + y)z_1 \mathfrak{b}^- & \text{on } D(z_1) \\ \frac{\eta - \theta t'}{z_2} \mathfrak{b}^- & \text{on } D(z_2) \end{cases}, \quad \delta z_2 = \begin{cases} \frac{\eta + \theta t'}{z_1} \mathfrak{b}^- & \text{on } D(z_1) \\ -(\theta + y)z_2 \mathfrak{b}^- & \text{on } D(z_2) \end{cases}$$

So  $\delta z_1 - (\theta + y)z_1 \mathfrak{b}^-$  is an element in  $\omega_{X/S}$  which is zero on  $D(z_1)$  and is  $\eta/z_2 \mathfrak{b}^-$  on  $D(z_2)$ , and  $\delta z_2 + (\theta + y)z_2$  is zero on  $D(z_2)$ , and is  $\eta/z_1 \mathfrak{b}^-$  on  $D(z_1)$ . However,  $\eta/z_1$  is not an element of  $\widehat{\mathcal{O}}_{X,p}$ , unless  $\eta = 0$ . Hence it must be that  $\eta = 0$ .

Assume that  $\mathcal{J}$  is fermionic then  $t' = t$  and  $y = (f_1(z_1), f_2(z_2))$ , where  $f_1, f_2$  are Laurent polynomials in  $z_1, z_2$  valued in  $\mathcal{J}$  respectively, then we can write action of  $\delta$  on generators by

$$\begin{aligned} \theta &\mapsto (1 - \theta z_1 f'_1(z_1) - \theta z_2 f'_2(z_2))\mathfrak{b}^-, \\ z_1 &\mapsto (\theta + f_1(z_1))z_1 \mathfrak{b}^-, \\ z_2 &\mapsto -(\theta + f_2(z_2))z_2 \mathfrak{b}^-. \end{aligned}$$

Take coordinate transformation  $\theta \mapsto \theta + f_1(z_1) + f_2(z_2) - f_2(0) \equiv \theta'$  and  $z_2 \mapsto z_2(1 + (f_2(0) - f_1(0))\theta) \equiv z'_2$ , and note that this does not change  $\mathfrak{b}^-$ . Under the new coordinate  $(z_1, z'_2, \theta')$ ,  $\delta$  acts on generators by

$$\theta' \mapsto \mathfrak{b}^-, \quad z_1 \mapsto \theta' z_1 \mathfrak{b}^-, \quad z'_2 \mapsto -\theta' z'_2 \mathfrak{b}^-.$$

It is thus the derivation of the R node canonical model.

Now assume that  $\mathcal{J}$  is bosonic then  $y = \theta(f_1(z_1), f_2(z_2))$ , where  $f_1, f_2$  are Laurent polynomials in  $z_1, z_2$  valued in  $\mathcal{J}$  respectively, then

$$\delta \theta = \begin{cases} (1 - f_1(z_1))\mathfrak{b}^- & \text{on } D(z_1) \\ (1 - f_2(z_2))\mathfrak{b}^- & \text{on } D(z_2) \end{cases}.$$

So we must have  $f_1(0) = f_2(0)$  in order that  $\delta\theta \in \omega_{X/S}$ , thus  $y \in \mathcal{J}\widehat{\mathcal{O}}_{X,p}$ . we can write action of  $\delta$  on generators by

$$\begin{aligned}\theta &\mapsto (1 - \partial_\theta y)\mathfrak{b}^-, \\ z_1 &\mapsto (\theta + y)z_1\mathfrak{b}^-, \\ z_2 &\mapsto -(\theta + y)z_2\mathfrak{b}^-.\end{aligned}$$

Take the coordinate transformation  $\theta \mapsto \theta + y =: \theta'$ , note that this does not change  $\mathfrak{b}^-$ . Under the new coordinate  $(z_1, z_2, \theta')$ ,  $\delta$  acts on generators by

$$\theta' \mapsto \mathfrak{b}^-, \quad z_1 \mapsto \theta' z_1 \mathfrak{b}^-, \quad z_2 \mapsto -\theta' z_2 \mathfrak{b}^-.$$

It is thus the derivation of the R node canonical model. This concludes the situation b.

**If  $p$  is a NS node**, write

$$\delta = \delta_0 + \delta',$$

where  $\delta_0$  is

$$\delta_0 : f \mapsto \begin{cases} (\partial_{\theta_1} f + \theta_1 \partial_{z_1} f)[dz_1|d\theta_1], & \text{on } D(z_1), \\ (\partial_{\theta_2} f + \theta_2 \partial_{z_2} f)[dz_2|d\theta_2], & \text{on } D(z_2) \end{cases},$$

and

$$\delta' \in \text{Der}_\kappa(\mathcal{O}_U, \mathcal{J} \otimes_\kappa \omega_{U/\text{Spec } \kappa}),$$

where  $U = \text{Spec}(\widehat{\mathcal{O}}_{X,p}/\mathfrak{m}_A \widehat{\mathcal{O}}_{X,p}) \setminus \{p\}$  is a disjoint union of  $\text{Spec } \kappa((z_1))$  and  $\text{Spec } \kappa((z_2))$ . The same argument in the situation b shows that  $x + Dy = 0$ ,  $y \in \mathcal{J}[[z_1]] \times \mathcal{J}[[z_2]]$  or  $\theta \mathcal{J}[[z_1]] \times \theta \mathcal{J}[[z_2]]$  depending on the parity of  $\mathcal{J}$ , and  $\delta'$  can be written as

$$\delta' : f \mapsto \begin{cases} -D(y \cdot Df)[dz_1|d\theta_2], & \text{on } D(z_1) \\ -D(y \cdot Df)[dz_2|d\theta_2], & \text{on } D(z_2) \end{cases}. \quad (7.12)$$

Consequently  $\delta'f$  has nonnegative power of  $z_1$  (respectively nonnegative power of  $z_2$ ) on  $D(z_1)$  (respectively on  $D(z_2)$ ), for all  $f \in \widehat{\mathcal{O}}_{X,p}/\mathfrak{m}_A \widehat{\mathcal{O}}_{X,p}$ . Now assume that  $\mathcal{J}$  is fermionic, then  $t_i = t$ . Consider

$$\delta_0 z_1 = \begin{cases} \theta_1 [dz_1|d\theta_1], & \text{on } D(z_1) \\ \frac{\eta_2 - z_1 \theta_2}{z_2} [dz_2|d\theta_2], & \text{on } D(z_2) \end{cases}$$

So  $\delta z_1 - z_1 \mathfrak{b}^+ - \delta' z_1$  is zero on  $D(z_1)$  and is  $\eta_2/z_2$  on  $D(z_2)$ . Since  $\delta' z_1$  has nonnegative power of  $z_2$  on  $D(z_2)$ , thus  $\eta_2$  must be zero. Similarly  $\eta_1 = 0$ . Assume that  $\mathcal{J}$  is bosonic, then  $\eta_i = 0$  automatically. Consider

$$\delta_0 \theta_1 = \begin{cases} [dz_1|d\theta_1], & \text{on } D(z_1) \\ \frac{t_2}{z_2} [dz_2|d\theta_2], & \text{on } D(z_2) \end{cases}.$$



Note that a set of generators of  $\omega_{X/S}$  is

$$\begin{aligned} \mathfrak{b}_1^- &= \begin{cases} [dz_1|d\theta_1], & \text{on } D(z_1) \\ \frac{t_1}{z_2} [dz_2|d\theta_2], & \text{on } D(z_2) \end{cases}, \\ \mathfrak{b}_2^- &= \begin{cases} -\frac{t_2}{z_1} [dz_1|d\theta_1], & \text{on } D(z_1) \\ [dz_2|d\theta_2], & \text{on } D(z_2) \end{cases}, \\ \mathfrak{b}^+ &= \begin{cases} \frac{\theta_1}{z_1} [dz_1|d\theta_1], & \text{on } D(z_1) \\ -\frac{\theta_2}{z_2} [dz_2|d\theta_2], & \text{on } D(z_2) \end{cases}. \end{aligned}$$

So  $\delta_0\theta_1 - \mathfrak{b}_1^-$  is zero on  $D(z_1)$  and is  $\frac{t_2-t_1}{z_2}$  on  $D(z_2)$ . Since  $\delta'\theta_1$  has nonnegative power of  $z_2$  on  $D(z_2)$ , this concludes that  $t_1 - t_2$  must be zero. Then we can write down the action of  $\delta$  on generators:

$$\delta\theta_i = (1 - Dy)\mathfrak{b}_i^-, \quad \delta z_i = (z_i\theta_i + y)\mathfrak{b}_i^-.$$

Suppose that  $y = (f_1(z_1), f_2(z_2))$ , where  $f_1, f_2$  are Laurent polynomials in  $z_1, z_2$  respectively, then the constant term of  $\delta z_1$  is  $f_1(0)$ , so  $f_1(0) = 0$  in order that  $\delta z_1 \in \omega_{X/S}$ . Similarly  $f_2(0) = 0$ . Thus  $y = (z_1\tilde{f}_1(z_1), z_2\tilde{f}_2(z_2))$ , and  $\delta z_i = (\theta_i + \tilde{f}_i(z_i))z_i\mathfrak{b}_i^-$ . Now take the coordinate transform  $\theta_i \mapsto \theta_i + \tilde{f}_i(z_i) \equiv \theta'_i$ , note that this does not change  $\mathfrak{b}_i^-$ . Under the new coordinate  $(z_1, z_2, \theta'_1, \theta'_2)$ ,  $\delta$  acts on generators by

$$\theta'_i \mapsto \mathfrak{b}_i^-, \quad z_i \mapsto \pm z_i \mathfrak{b}^+.$$

It is thus the derivation of the NS node canonical model. This concludes the situation c.  $\blacksquare$

### 7.2.3 Infinitesimal Automorphisms of Stable SUSY Curves

Let the sheaf of automorphism group of  $X$  which induces identity modulo  $\mathcal{J}$  be  $\mathcal{A}$ , then if  $X$  is smooth then it is a classical fact that

$$\mathcal{A} = \left( T_{X_s}^+(-E - \mathcal{D}) \oplus \Pi T_{X_s}^+(-E - \mathcal{D}) \otimes_{\mathcal{O}_{X_s}^+} \mathcal{E} \right) \otimes_{\kappa} \mathcal{J},$$

where  $T_{X_s}^+$  is the even tangent sheaf of the special fiber  $X_s$ , and  $\mathcal{E}$  is the spinor sheaf. If  $X$  is a general stable SUSY curve, then we need to take R and NS nodes into account. In fact, suppose that an automorphism at a formal neighborhood of a node comes from a derivation  $x\mathcal{D} + y\mathcal{D}^2$  where  $x$  is odd and  $y$  is even, then it must preserves the superconformal structure, i.e.  $[D, x\mathcal{D} + y\mathcal{D}^2] \equiv 0 \pmod{D}$ , hence  $x = \frac{1}{2}Dy$ . In local coordinate we can write it as

- If  $p$  is a R node: when  $\mathcal{J}$  is bosonic

$$x\mathcal{D} + y\mathcal{D}^2 = z_1 f_1(z_1) \partial_{z_1} + z_2 f_2(z_2) \partial_{z_2} + \frac{z_1 f'_1(z_1) + z_2 f'_2(z_2)}{2} \theta \partial_\theta,$$

where  $f_i(z_i) \in \mathcal{J}[[z_i]]$ . When  $\mathcal{J}$  is fermionic

$$x\mathcal{D} + y\mathcal{D}^2 = \theta g(z_1, z_2) (z_1 \partial_{z_1} + z_2 \partial_{z_2}) + g(z_1, z_2) \partial_\theta,$$

where  $g(z_1, z_2) \in \mathcal{J}[[z_1, z_2]]/(z_1 z_2)$ . Note that  $z_1 \partial_{z_1} + z_2 \partial_{z_2}$  is a generator of  $(\omega_{X_s, p}^+)^{-1}$ , and  $\partial_\theta$  is a generator of the dual of spin structure  $\mathcal{E}_p$ .

- If  $p$  is a NS node: when  $\mathcal{J}$  is bosonic

$$xD + yD^2 = f_1(z_1)\partial_{z_1} + f_2(z_2)\partial_{z_2} + \frac{1}{2}f'_1(z_1)\theta_1\partial_{\theta_1} + \frac{1}{2}f'_2(z_2)\theta_2\partial_{\theta_2},$$

where  $f_i(z_i) \in z_i\mathcal{J}[[z_i]]$ . When  $\mathcal{J}$  is fermionic

$$xD + yD^2 = g_1(z_1)(\partial_{\theta_1} - \theta_1\partial_{z_1}) + g_2(z_2)(\partial_{\theta_2} - \theta_2\partial_{z_2}),$$

where  $g_i(z_i) \in \mathcal{J}[[z_i]]$ . Note that  $\partial_{\theta_i} - \theta_i\partial_{z_i}$  is a generator of the dual of spin structure on the  $i$ 'th component of the normalization at  $p$ , and the direct sum of their pushforward along the normalization map is the dual of spin structure  $\mathcal{E}_p$ .

so we see that

$$\widehat{\mathcal{A}}_p = \left(T_{X_{s,p}}^+ \oplus \Pi(\omega_{X_{s,p}}^+)^{-1} \otimes \mathcal{E}_p\right) \otimes_{\kappa} \mathcal{J} \cong \left(T_{X_{s,p}}^+ \oplus \Pi\mathcal{E}_p^{\vee}\right) \otimes_{\kappa} \mathcal{J},$$

if  $p$  is a R node or NS node. Combine the smooth case and the nodes by formal gluing, we get

**Proposition 7.8.** *The sheaf of infinitesimal automorphism group  $\mathcal{A}$  is isomorphic to*

$$\mathcal{A} = (T_{X_s}^+(-E - \mathcal{D}) \oplus \Pi\mathcal{E}^{\vee}) \otimes_{\kappa} \mathcal{J}, \quad (7.13)$$

**Corollary 7.5.** *There is no global infinitesimal automorphism, neither even nor odd.*

*Proof.* Let  $\pi : C \rightarrow X_{s,\text{bos}}$  be the normalization map of the bosonic truncation of  $X_s$ , then

$$T_{X_s}^+(-E - \mathcal{D}) = \pi_* (\omega_{\mathcal{C}}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N})),$$

where  $\mathcal{R}$  (resp.  $\mathcal{N}$ ) is the preimage of R nodes (resp. NS nodes) in  $\mathcal{C}$ . Since  $X_{s,\text{bos}}$  is a stable curve,  $\omega_{\mathcal{C}}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N})$  has negative degree on every connected component, hence

$$\Gamma(X, T_{X_s}^+(-E - \mathcal{D})) = \Gamma(\mathcal{C}, \omega_{\mathcal{C}}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N})) = 0.$$

On the other hand, there is a short exact sequence

$$0 \rightarrow (\omega_{X_{s,p}}^+)^{-1}(-E - \mathcal{D}) \otimes_{\mathcal{O}_{X_s}^+} \mathcal{E} \rightarrow \pi_* (\omega_{\mathcal{C}}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N}) \otimes \mathcal{E}_{\mathcal{C}}) \rightarrow \bigoplus_{p \in \text{R Node}} \kappa \rightarrow 0,$$

where  $\mathcal{E}_{\mathcal{C}}$  is a square root of  $\omega_{\mathcal{C}}(\mathcal{D} + \mathcal{R})$ . On each connected component  $\mathcal{C}_i$ ,

$$\deg \mathcal{E}_{\mathcal{C}} = \frac{1}{2} \deg \omega_{\mathcal{C}_i}(\mathcal{D} + \mathcal{R}) \leq \frac{1}{2} \deg \omega_{\mathcal{C}_i}(E + \mathcal{D} + \mathcal{R} + \mathcal{N}),$$

consequently

$$\deg \omega_{\mathcal{C}_i}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N}) \otimes \mathcal{E}_{\mathcal{C}_i} \leq -\frac{1}{2} \deg \omega_{\mathcal{C}_i}(E + \mathcal{D} + \mathcal{R} + \mathcal{N}) < 0.$$

Then it follows that

$$\Gamma(X, (\omega_{X_{s,p}}^+)^{-1}(-E - \mathcal{D}) \otimes_{\mathcal{O}_{X_s}^+} \mathcal{E}) \subset \Gamma(\mathcal{C}, \omega_{\mathcal{C}}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N}) \otimes \mathcal{E}_{\mathcal{C}}) = 0.$$

■

### 7.2.4 Isomorphisms of Stable SUSY Curves

**Proposition 7.9.** *Suppose that  $S$  is a Noetherian superscheme and  $X, Y$  are two stable SUSY curves over  $S$ , then the functor*

$$\mathfrak{Isom}_S(X, Y) : (f : T \rightarrow S) \mapsto \text{Isom}_T(X_T, Y_T),$$

*is represented by a finite unramified superscheme over  $S$ .*

*Proof.* Since representability can be proven Zariski locally, so we may additionally assume that  $S$  is affine, so  $X$  is projective over  $S$  according to Lemma 7.24. First consider the natural transform  $\mathfrak{Isom}_S(X, Y) \rightarrow \mathfrak{Map}_S(X, Y)$  (see Theorem 7.10). According to *loc.cit.*  $\mathfrak{Map}_S(X, Y)$  is represented by a locally finite type  $S$ -superscheme  $\text{Map}_S(X, Y)$ , we claim that  $\mathfrak{Isom}_S(X, Y)$  is represented by a locally closed sub superscheme of  $M \equiv \text{Map}_S(X, Y)$ . Take the universal map  $\Phi : X_M \rightarrow Y_M$ . The locus  $U$  on  $M$  where  $\Phi$  is an isomorphism of superschemes is open. Restricting on  $U$ , and let the puncture of  $X$  (resp.  $Y$ ) be  $E, \mathcal{D}$  (resp.  $E', \mathcal{D}'$ ), then the incidence relations  $\Phi^{-1}E'_i \subset E_i$ ,  $E_i \subset \Phi^{-1}E'_i$ ,  $\Phi^{-1}\mathcal{D}'_i \subset \mathcal{D}_i$  and  $\mathcal{D}_i \subset \Phi^{-1}\mathcal{D}'_i$  are represented by closed sub superschemes  $U_\alpha$  of  $U$  (Corollary 7.4). Take intersections of these  $U_\alpha$  and we obtain a closed sub superscheme  $V$  of  $U$  which represents isomorphisms between  $X$  and  $Y$  as superschemes as preserves punctures. Finally the preservation of derivation can be interpreted as  $\Phi^*\delta' - \delta = 0$ , so it is represented by the intersection of the section induced by  $\Phi^*\delta' - \delta$  and the zero section in the Hom superscheme  $\mathfrak{Hom}_V(\Omega_{X_V/V}^1, \omega_{X_V/V})$ . Note that moduli spaces showing up in these steps are locally finite presented, whence  $\mathfrak{Isom}_S(X, Y)$  is represented by a locally finite type  $S$ -superscheme  $\text{Isom}_S(X, Y)$ .

Next, if  $T$  is a bosonic scheme, then isomorphisms between  $X_T$  and  $Y_T$  are one to one corresponds to isomorphisms between their underlying spin curves. By Yoneda Lemma, this implies that

$$\text{Isom}_S(X, Y)_{\text{bos}} \cong \text{Isom}_{S_{\text{bos}}}(\mathcal{C}, \mathcal{C}'),$$

where  $\mathcal{C}, \mathcal{C}'$  are underlying spin curves of  $X \times_S S_{\text{bos}}, Y \times_S S_{\text{bos}}$ . Consequently  $\text{Isom}_S(X, Y)_{\text{bos}}$  is finite over  $S_{\text{bos}}$  hence  $\text{Isom}_S(X, Y)$  is finite over  $S$  by Nakayama Lemma.

Finally, Corollary 7.5 implies that for any geometric point  $s$  of  $S$  and any  $\phi \in \text{Isom}_s(X_s, Y_s)$ , the tangent space of  $\text{Isom}_S(X, Y)_s$  at  $\phi$  is trivial, hence  $\text{Isom}_S(X, Y)$  is unramified over  $S$ . This concludes the proof.  $\blacksquare$

### 7.2.5 Étale Local Structures of Stable SUSY Curves

Let us discuss the étale local structures of stable SUSY curves.

**Lemma 7.30.** *Suppose that  $A$  is a strictly Henselian local Noetherian superring, such that  $A/\mathfrak{n}_A$  is a  $G$ -ring. Then there exists an inductive system  $\{B_\lambda\}_{\lambda \in L}$  of smooth  $A$  superalgebras, such that*

$$\widehat{A} = \varinjlim_{\lambda} B_\lambda.$$

*Proof.* Let  $A = A^+ \oplus A^-$  be its bosonic and fermionic parts.  $A/\mathfrak{n}_A$  being a G-ring implies that  $A^+$  is also a G-ring, so  $\widehat{A^+} = \widehat{A^+}$  is a regular- $A^+$  algebra. By Popescu's Theorem, there exists an inductive system  $\{B'_\lambda\}_{\lambda \in L}$  of smooth  $A^+$  algebras, such that

$$\widehat{A^+} = \varinjlim_{\lambda} B'_\lambda.$$

Then  $B_\lambda = B'_\lambda \otimes_{A^+} A$  is what we want. ■

**Proposition 7.10.** *Suppose that  $S$  is a Noetherian superscheme such that  $S_{\text{bos}}$  is a G-scheme. Let  $X \rightarrow S$  be a stable SUSY curve, then there exists an étale cover  $\{V_i \rightarrow S\}_{i \in I}$  and for each  $i \in I$  there exists a finite set  $J_i$  and an étale cover  $\{\rho_{ij} : U_{ij} \rightarrow X_i\}_{j \in J_i}$  where  $X_i = X \times_S V_i$ , such that*

1. *Every  $U_{ij}$  has an étale morphism  $\varphi_{ij} : U_{ij} \rightarrow Y$  where  $Y$  is one of the canonical models such that  $\rho_{ij}^* \delta_X = \varphi_{ij}^* \delta_{\text{can}}$ .*
2. *For every pair of indices  $j, j' \in J_i$ ,  $j \neq j'$ ,  $U_{ij} \times_{X_i} U_{ij'}$  is smooth over  $V_i$  and contains no marking divisors.*

*Proof.* By Lemma 7.23, it suffices to prove the proposition with additional assumption that  $S = \text{Spec } A$  where  $A$  is a strict Henselian local Noetherian superring. Since  $S_{\text{bos}}$  is a G-scheme, we can apply Lemma 7.30 to furthermore assume that  $S = \text{Spec } A$  where  $A$  is a complete local Noetherian superring with algebraically closed residue field<sup>68</sup>. Let the set of nodes and marking divisors of the special fiber  $X_s$  be  $J_0$ , then for every  $p \in J_0$ , there exists an isomorphism

$$\varphi_p : \text{Spec } \widehat{\mathcal{O}}_{X,p} \cong \text{Spec } \widehat{\mathcal{O}}_{Y,0},$$

where  $Y$  is one of the canonical models and 0 is the origin of its special fiber over  $\text{Spec } A$ , by Proposition 7.7. Choose an affine neighborhood of  $p$ , denoted by  $W_p$ , such that it does not contain any point in the set  $J_0 \setminus \{p\}$ . Now  $X_{\text{bos}}$  is a G-scheme, so Lemma 7.30 implies that there exists an directed system  $\{B_\lambda\}_{\lambda \in L}$  of smooth  $\mathcal{O}_X(W_p)$  superalgebras such that

$$\widehat{\mathcal{O}}_{X,p} = \varinjlim_{\lambda} B_\lambda.$$

Denote the morphism  $\text{Spec } B_\lambda \rightarrow W_p$  by  $\rho_{p,\lambda}$ . Next we prove the NS node case and the rest are similar. There exists  $\lambda \in L$  such that  $\exists z_1, z_2, \theta_1, \theta_2 \in B_\lambda$  whose images in  $\widehat{\mathcal{O}}_{X,p}$  are the corresponding generators of the maximal ideal and satisfies the relations in the NS node canonical model, this defines a morphism of  $A$  superschemes  $\varphi_{p,\lambda} : \text{Spec } B_\lambda \rightarrow Y$ . Moreover, after increasing  $\lambda$ , there is a set of generators  $\mathfrak{b}_1^-, \mathfrak{b}_2^-, \mathfrak{b}^+ \in \rho_{p,\lambda}^* \omega_{X/S}$ , whose images

---

<sup>68</sup>  $B_i$  in the Lemma 7.30 are only smooth, not étale, but we can always choose a closed point  $p$  on the special fiber of  $\text{Spec } B_i \rightarrow \text{Spec } A$  and lift a system of parameters of  $p$  on the special fiber to the germ of  $B_i$  at  $p$ , after taking quotient of these elements, we get a local superring which is étale over  $A$  hence isomorphic to  $A$  since  $A$  is strictly henselian.

in  $\omega_{\text{Spec } \widehat{\mathcal{O}}_{X,p}/S}$  are the corresponding generators in the NS node canonical model, such that  $\rho_{p,\lambda}^* \delta_X : \rho_{p,\lambda}^* \Omega_{X/S}^1 \rightarrow \rho_{p,\lambda}^* \omega_{X/S}$  acts on generators by

$$\begin{aligned} dz_1 &\rightarrow z_1 \mathbf{b}^+, & dz_2 &\rightarrow -z_2 \mathbf{b}^+, \\ d\theta_1 &\rightarrow \mathbf{b}_1^-, & d\theta_2 &\rightarrow \mathbf{b}_2^-. \end{aligned}$$

Use the same notation  $p$  to denote the image of the closed point of  $\text{Spec } \widehat{\mathcal{O}}_{X,p}$  in  $\text{Spec } B_\lambda$ , then lift a system of homogeneous parameters of the fiber of  $\rho_{p,\lambda}$  at point  $p$  to an open affine neighborhood  $U'_{p,\lambda}$  and take the closed sub superscheme of  $U'_{p,\lambda}$  defined by these elements, denoted by  $U''_{p,\lambda}$ , then  $U''_{p,\lambda}$  is étale over  $W_p$  at  $p$ . We keep using the notation  $\varphi_{p,\lambda}$  to denote its restriction to  $U''_{p,\lambda}$ . Then taking completion at  $p$  induces isomorphism  $\widehat{\mathcal{O}}_{U''_{p,\lambda},p} \cong \widehat{\mathcal{O}}_{Y,0}$ , hence  $\varphi_{p,\lambda}$  is étale at  $p$ . Shrinking  $U''_{p,\lambda}$  if needed, we have two morphisms  $\rho_{p,\lambda} : U''_{p,\lambda} \rightarrow U_p$  and  $\varphi_{p,\lambda} : U''_{p,\lambda} \rightarrow Y$  such that  $\rho_{p,\lambda}^* \delta_X = \varphi_{p,\lambda}^* \delta_{\text{can}}$  since they agree on generators of  $\Omega_{U''_{p,\lambda}/S}^1$ .

Applying the same procedure to other elements in the set  $J_0$ , we found a finite set of superschemes  $\{U_p\}_{p \in J_0}$  with étale morphisms  $\rho_p : U_p \rightarrow X, \varphi_p : U_p \rightarrow Y$  such that  $\rho_p^* \delta_X = \varphi_p^* \delta_{\text{can}}$ . Moreover  $U_p \times_X U_{p'}$  is smooth over  $S$  and contains no puncturing divisors if  $p \neq p'$ , this follows from the construction that  $U_p$  are superschemes over  $W_p$  for some open affine neighborhood of  $p$  which contains no element  $p' \in J_0 \setminus \{p\}$ , so the special fiber of  $U_p \times_X U_{p'}$  is smooth and contains no marking divisor. The remaining issue is that  $\bigcup_{p \in J_0} U_p \rightarrow X$  might not be a covering, this can be cured by enlarging  $J_0$  to include all points in  $X_s \setminus \bigcup_{p \in J_0} W_p$ , note that this is a finite set since  $X_s$  is one dimensional.  $\blacksquare$

## 7.2.6 Absence of Obstruction for Lifting Stable SUSY Curves

Recall the setup for discussing local models of stable SUSY curves: suppose that there is an extension

$$0 \rightarrow \mathcal{J} \rightarrow A \rightarrow A/\mathcal{J} \rightarrow 0,$$

of local Artinian superrings, where  $\mathcal{J}$  is an ideal of  $A$ , purely bosonic or purely fermionic, such that the maximal ideal  $\mathfrak{m}_A$  annihilates it. We also assume that the residue field of  $A$ , denote by  $\kappa$ , is algebraically closed.

**Proposition 7.11.** *Suppose that there is a stable SUSY curve  $Y$  over  $\text{Spec } A/\mathcal{J}$ , then there exists a stable SUSY curve  $X$  over  $\text{Spec } A$  such that its base change to  $\text{Spec } A/\mathcal{J}$  is isomorphic to  $Y$ .*

*Proof. Step 1.* We prove that Zariski-locally on  $Y$  there always exists such lifting. Let the deformation space of  $\mathcal{O}_{Y,p}$  (see the notation in equation (7.11)) as a germ of SUSY curve be  $\text{sDef}_A(\mathcal{O}_{Y,p}, \mathcal{J})$ , we want to show that this is not empty. According to Equation 7.11, we have a commutative diagram of maps between sets

$$\begin{array}{ccc} \text{sDef}_A(\mathcal{O}_{Y,p}, \mathcal{J}) & \longrightarrow & \text{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J}) \\ \downarrow & & \downarrow \\ \text{Def}_A(\mathcal{O}_{Y,p}, \mathcal{J}) & \xlongequal{\quad} & \text{Def}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J}). \end{array}$$

We claim that the map  $\mathrm{sDef}_A(\mathcal{O}_{Y,p}, \mathcal{J}) \rightarrow \mathrm{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J})$  is bijective.

- **Injectivity:** Suppose that there are two deformations whose completions are isomorphic, then the underlying deformation of superscheme structures are isomorphic, denoted by  $\mathcal{O}'_{Y,p}$ , by the identification  $\mathrm{Def}_A(\mathcal{O}_{Y,p}, \mathcal{J}) = \mathrm{Def}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J})$ . Let the derivations be  $\delta_1, \delta_2$ , using Lemma 7.30, there exists an étale morphism  $\mathcal{O}'_{Y,p} \rightarrow C$  such that  $C$  is local, and  $\pi^* \delta_1, \pi^* \delta_2$  are related by infinitesimal automorphism  $\alpha$  on  $\mathrm{Spec} C$ , then  $\pi_1^* \alpha = \pi_2^* \alpha$  because they both transform  $q^* \delta_1$  to  $q^* \delta_2$ , see notations below

$$\begin{array}{ccc} \mathrm{Spec} C \otimes_{\mathcal{O}'_{Y,p}} C & \xrightarrow{\pi_1} & \mathrm{Spec} C \\ \downarrow \pi_2 & \searrow q & \downarrow \pi \\ \mathrm{Spec} C & \xrightarrow{\pi} & \mathrm{Spec} \mathcal{O}'_{Y,p} \end{array},$$

Hence  $\alpha$  descends to an infinitesimal automorphism on  $\mathrm{Spec} \mathcal{O}'_{Y,p}$  and transform  $\delta_1$  to  $\delta_2$ .

- **Surjectivity:** Suppose that  $(\widehat{\mathcal{O}}_{Y,p}, \delta')$  is a deformation of  $(\widehat{\mathcal{O}}_{Y,p}, \delta)$  whose underlying superscheme is the completion of a deformation of  $\mathcal{O}_{Y,p}$ , denoted by  $\mathcal{O}'_{Y,p}$ . Using Lemma 7.30, there exists an étale morphism  $\mathcal{O}'_{Y,p} \rightarrow C$  such that  $C$  is local and  $\delta'$  descends to a SUSY derivation  $\delta_C$  on  $\mathrm{Spec} C$ . Then  $\pi_1^* \delta_C, \pi_2^* \delta_C$  are isomorphic derivations because their completions are isomorphic, hence they are related by an infinitesimal automorphism  $\alpha$ , see notations below

$$\begin{array}{ccc} \mathrm{Spec} C \otimes_{\mathcal{O}'_{Y,p}} C & \xrightarrow{\pi_1} & \mathrm{Spec} C \\ \downarrow \pi_2 & & \downarrow \pi \\ \mathrm{Spec} C & \xrightarrow{\pi} & \mathrm{Spec} \mathcal{O}'_{Y,p} \end{array},$$

Apparently  $\alpha$  satisfies cocycle condition when pulled back to  $\mathrm{Spec} C \otimes_{\mathcal{O}'_{Y,p}} C \otimes_{\mathcal{O}'_{Y,p}} C$ , hence defines an element in the cohomology group

$$H^1((\mathrm{Spec} \mathcal{O}'_{Y,p})_{\mathrm{ét}}, \mathcal{A})_+,$$

which is trivial since  $\mathcal{A}$  is coherent and  $\mathrm{Spec} \mathcal{O}'_{Y,p}$  is affine. So there exists an étale morphism  $C \rightarrow D$  and a SUSY derivation  $\delta_D$  which is isomorphic to the pull back of  $\delta_C$  to  $\mathrm{Spec} D$ , such that  $f_1^* \delta_D = f_2^* \delta_D$  where  $f_1, f_2$  are two projections from  $\mathrm{Spec} D \otimes_{\mathcal{O}'_{Y,p}} D$  to  $\mathrm{Spec} C$ . Hence  $\delta_D$  descends to  $\mathrm{Spec} \mathcal{O}'_{Y,p}$ .

Now we take an element of  $\mathrm{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J})$  (see Proposition 7.7), which induces a deformation of  $W_p$  as a superscheme (see equation (7.11) and notations there), denote the deformation of  $W_p$  by  $W'_p$ , and by the identification  $\mathrm{sDef}_A(\mathcal{O}_{Y,p}, \mathcal{J}) \xrightarrow{\sim} \mathrm{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J})$  there exists a SUSY derivation  $\delta_p$  on  $\mathrm{Spec} \mathcal{O}_{W'_p, p}$ . Extend  $\delta_p$  to a Zariski open neighborhood of  $p$  and we are done.

**Step 2.** For smooth open affine  $W \subset Y$ ,  $W$  deforms uniquely, since it deforms uniquely as superscheme with puncturing divisor<sup>69</sup>, and the derivation is also unique up to infinitesimal automorphism, since for any point  $p \in W$ ,  $\text{Spec } \widehat{\mathcal{O}}_{W,p}$  deforms uniquely as germ of SUSY curves.

**Step 3.** Take an open affine cover  $\{W_j\}_{j \in J}$  of  $Y$  such that  $W_j$  contains at most one node or puncture divisor, and  $W_j \cap W_k$  is smooth and contains no node or puncture divisor, whenever  $j \neq k$ . Shrinking  $W_j$  if necessary, we assume that there exists deformations  $W'_j$ .  $W_j \cap W_k$  is smooth hence has a unique deformation. Thus  $W'_j \cap W_k$  and  $W'_k \cap W_j$  are related by an isomorphism  $\phi_{jk}$ . Consider

$$\phi_{jk} \circ \phi_{kl} \circ \phi_{lj} \in \Gamma(W_j \cap W_k \cap W_l, \mathcal{A})_+,$$

which apparently satisfies cocycle condition, thus it defines an element in  $H^2(Y, \mathcal{A})_+$  which is trivial because  $\dim Y = 1$ . This means that after a refinement of  $\{W_j\}_{j \in J}$ , we can modify  $\phi_{jk}$  by elements in  $\Gamma(W_j \cap W_k, \mathcal{A})_+$  to force it satisfying cocycle condition, thus  $\{W'_j\}$  glue to a deformation of  $Y$  as a stable SUSY curve.  $\blacksquare$

In fact, the choice of gluing data in the last step of proof is in one-to-one correspondence with  $H^1(Y, \mathcal{A})_+ = H^1(Y_{s,\text{bos}}, \mathcal{A})_+$ , so we have

**Corollary 7.6.** *There is a fiber sequence of sets*

$$H^1(Y_{s,\text{bos}}, \mathcal{A})_+ \rightarrow \text{sDef}_A(Y, \mathcal{J}) \rightarrow \prod_{p \in \text{NS Nodes}} \text{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J}) \times \prod_{p \in \text{R Nodes}} \text{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J}). \quad (7.14)$$

When  $A = B \times_{\kappa} \kappa[\theta]$  or  $A = B \times_{\kappa} \kappa[t]/(t^2)$  and  $\mathcal{J}$  is the ideal  $(\theta)$  or  $(t)$ , there is a canonical  $\kappa$ -vector space structure on  $\text{sDef}_A(Y, \mathcal{J})$  with zero being trivial extension and the fiber sequence (7.14) becomes a short exact sequence of vector spaces

$$0 \rightarrow H^1(Y_{s,\text{bos}}, \mathcal{A})_+ \rightarrow \text{sDef}_A(Y, \mathcal{J}) \rightarrow \bigoplus_{p \in \text{NS Nodes}} (\mathcal{J}_p)_+ \oplus \bigoplus_{p \in \text{R Nodes}} (\mathcal{J}_p)_+ \rightarrow 0.$$

Therefore, we can use it to compute the dimension of  $\text{sDef}_A(Y, \mathcal{J})$ :

- $A = B \times_{\kappa} \kappa[t]/(t^2)$ ,  $H^1(Y_{s,\text{bos}}, \mathcal{A})_+ = H^1(Y_{s,\text{bos}}, T_{Y_s}^+(-E - \mathcal{D}))$ , use the normalization in the proof of Corollary 7.5, there is isomorphism

$$H^1(Y_{s,\text{bos}}, T_{Y_s}^+(-E - \mathcal{D})) \cong \bigoplus_i H^1(C_i, \omega_{C_i}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N})).$$

So the dimension of  $H^1(Y_{s,\text{bos}}, \mathcal{A})_+$  can be computed

$$\dim_{\kappa} H^1(Y_{s,\text{bos}}, \mathcal{A})_+ = \sum_i (3g_i - 3) + n_{\text{NS}} + n_{\text{R}} + 2\#(\text{NS Node}) + 2\#(\text{R Node})$$

---

<sup>69</sup>The deformation space of smooth superscheme  $W$  is  $H^1(W, T_{W_s})$ , by Theorem 2.1.2 of [84], which is trivial when  $W$  is affine.

$$= 3g - 3 + n_{\text{NS}} + n_{\text{R}} - \#(\text{NS Node}) - \#(\text{R Node}).$$

The spaces  $(\mathcal{J}_p)_+$  are one dimensional so the dimension of even deformations can be computed

$$\dim_{\kappa} \text{sDef}_A(Y, \mathcal{J}) = 3g - 3 + n_{\text{NS}} + n_{\text{R}}. \quad (7.15)$$

- $A = B \times_{\kappa} \kappa[\theta]$ ,  $H^1(Y_{s,\text{bos}}, \mathcal{A})_+ = H^1(Y_{s,\text{bos}}, (\omega_{X_{s,p}}^+)^{-1}(-E - \mathcal{D}) \otimes \mathcal{E})$ , using the short exact sequence

$$\begin{aligned} 0 \rightarrow \bigoplus_{p \in \text{R Node}} \kappa \rightarrow H^1(Y_{s,\text{bos}}, (\omega_{X_{s,p}}^+)^{-1}(-E - \mathcal{D}) \otimes \mathcal{E}) \\ \rightarrow \bigoplus_i H^1(C_i, \omega_{C_i}^{-1}(-E - \mathcal{D} - \mathcal{R} - \mathcal{N}) \otimes \mathcal{E}_{C_i}) \rightarrow 0, \end{aligned}$$

we see that the dimension of  $H^1(Y_{s,\text{bos}}, \mathcal{A})_+$  is

$$\begin{aligned} \dim_{\kappa} H^1(Y_{s,\text{bos}}, \mathcal{A})_+ &= \sum_i (2g_i - 2) + n_{\text{NS}} + \frac{n_{\text{R}}}{2} + 2\#(\text{NS Node}) + 2\#(\text{R Node}) \\ &= 2g - 2 + n_{\text{NS}} + \frac{n_{\text{R}}}{2} \end{aligned} \quad (7.16)$$

The spaces  $(\mathcal{J}_p)_+$  are trivial so the dimension of odd deformations is

$$\dim_{\kappa} \text{sDef}_A(Y, \mathcal{J}) = 2g - 2 + n_{\text{NS}} + \frac{n_{\text{R}}}{2}.$$

In these two situations, the restriction maps are actually isomorphisms

$$\begin{aligned} \text{sDef}_{B \times_{\kappa} \kappa[t]/(t^2)}(Y, (t)) &\cong \text{sDef}_{\kappa[t]/(t^2)}(Y_s, (t)), \\ \text{sDef}_{B \times_{\kappa} \kappa[\theta]}(Y, (\theta)) &\cong \text{sDef}_{\kappa[\theta]}(Y_s, (\theta)). \end{aligned} \quad (7.17)$$

Since the restriction of local deformation parameters

$$\text{sDef}_{B \times_{\kappa} \kappa[t]/(t^2)}(\hat{\mathcal{O}}_{Y,p}, (t)) \rightarrow \text{sDef}_{\kappa[t]/(t^2)}(\hat{\mathcal{O}}_{Y_s,p}, (t))$$

is an isomorphism as can be seen from the Proposition 7.7 (same for the other case), and the Čech cocycles restricts isomorphically, so the isomorphisms (7.17) follows from the Five Lemma.

### 7.3 Moduli of Stable $\mathcal{N} = 1$ SUSY Curves

For simplicity, we will work over base field  $\mathbb{C}$ . Analogous to the non-supersymmetric case,  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is an algebraic superstack, more precisely

**Theorem 7.13.** *Assume that a triple of natural numbers  $(g, n_{\text{NS}}, n_{\text{R}})$  satisfies  $n_{\text{R}} \in 2\mathbb{Z}_{\geq 0}$ , and  $2g - 2 + n_{\text{NS}} + n_{\text{R}} > 0$ .*



(1)  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is a smooth proper Deligne-Mumford superstack of dimension

$$\left(3g - 3 + n_{\text{NS}} + n_{\text{R}} \mid 2g - 2 + n_{\text{NS}} + \frac{n_{\text{R}}}{2}\right).$$

Moreover its bosonic truncation  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}; \text{bos}}$  is the moduli stack  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  of punctured stable spin curves.

(2)  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  has a coarse moduli superspace  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  which is actually bosonic and is also the coarse moduli space for the bosonic quotient  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}; \text{ev}}$ . Moreover its reduced subspace  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}; \text{red}}$  is the coarse moduli space  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  of punctured stable spin curves.

*Proof. (1):* Proposition 7.6 shows that  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is an étale superstack, and Lemma 7.23 implies that  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is locally finite presented. Moreover, Proposition 7.9 says that the diagonal of  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is representable and is finite and unramified. Assuming that  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  has an étale atlas, then  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is a Deligne-Mumford superstack. Smoothness follows from Proposition 7.11, and the computation of tangent space is done in equations (7.15) and (7.16). Remark 7.5 implies that for any Noetherian scheme  $T$  over  $\mathbb{C}$ , there is an isomorphism  $\text{Hom}_{\text{ComSp}}(T, \overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}) \cong \text{Hom}_{\text{SupSp}}(S(T), \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}})$  which is functorial in  $T$ , then it follows from Lemma 7.1 and Yoneda Lemma that

$$\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}; \text{bos}} = \overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}.$$

Hence the properness of  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}; \text{bos}}$  follows from the properness of  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  (Theorem B.1).

(2): For a stable SUSY curve  $\pi : X \rightarrow S$ , consider automorphism  $\Gamma_X : X \rightarrow X$  acting by 1 on  $\mathcal{O}_X^+$  and  $-1$  on  $\mathcal{O}_X^-$ , similarly  $\Gamma_S : S \rightarrow S$  is defined. They are compatible in the sense that the diagram

$$\begin{array}{ccc} X & \xrightarrow{\Gamma_X} & X \\ \downarrow \pi & & \downarrow \pi \\ C & \xrightarrow{\Gamma_C} & C \end{array},$$

commutes.  $\Gamma_X$  preserves punctures  $E_i$ , since  $E_i$  is locally in coordinate system  $(z|\theta)$  defined by the ideal  $(z, \theta)$  which is preserved by  $\Gamma_X$ . Similarly  $\Gamma_X$  preserves punctures  $\mathcal{D}_i$ .  $\Gamma_X$  commutes with derivation  $\delta$ , since  $\delta$  in the smooth canonical model can be written as  $f \mapsto (\partial_\theta f + \theta \partial_z f)[dz|d\theta]$  which is invariant under  $\theta \mapsto -\theta$ , similar for other three canonical models. Hence we have an embedding  $\text{Id}_S \times \Gamma_S : S \rightarrow \text{Isom}_{S \times S}(p_1^* X, p_2^* X)$  where  $p_1, p_2 : S \times S \rightarrow S$  are two projections. The the proof of (2) follows from the same argument for the moduli of smooth SUSY curves (Proposition 5.4 and Corollary 5.5 of [7]).  $\blacksquare$

It remains to construct an étale atlas of  $\overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$ , which will be done in the next two sections.

## Universal Deformation

For a stable SUSY curve  $X$  over  $\text{Spec } \mathbb{C}$ , define a functor  $F_X : \text{sArt}_{\mathbb{C}} \rightarrow \text{Sets}$  by sending  $A \in \text{sArt}_{\mathbb{C}}$  to the set of equivalence class of  $(Y, \phi)$  where  $Y$  is a stable SUSY curve over  $A$  and  $\phi : Y_{\mathbb{C}} \rightarrow X$  is an isomorphism, with the obvious equivalence relation. From the definition it is easy to see that  $F_X$  satisfies the Schlessinger's condition  $(S_0)$ ; Proposition 7.11 implies that  $F_X$  satisfies  $(S_1)$  and isomorphism (7.17) is the condition  $(S_1)$ . The tangent space of  $F_X$  is of dimension

$$\left(3g - 3 + n_{\text{NS}} + n_{\text{R}} | 2g - 2 + n_{\text{NS}} + \frac{n_{\text{R}}}{2}\right).$$

So  $(S_3)$  does also hold. We claim that condition  $(S_4)$  is true for  $F_X$  as well. Suppose that  $p : A \rightarrow A/\mathcal{I}$  is an *even* small extension, and  $(Y, \phi) \in F_X(A/\mathcal{I})$ , then  $p^{-1}(Y, \phi)$  is the affine space  $\text{sDef}_A(Y, \mathcal{I})$  with a transitive action of  $T_{F_X}^+$ <sup>70</sup>. The action is actually faithful, since the affine space

$$\Lambda := \prod_{p \in \text{NS Nodes}} \text{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{I}) \times \prod_{p \in \text{R Nodes}} \text{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{I}),$$

has dimension  $\#(\text{R Nodes}) + \#(\text{NS Nodes})$ , so the stabilizer of a given point  $\lambda \in \Lambda$  under the induced action of  $T_{F_X}^+$  has dimension  $\dim_{\mathbb{C}} T_{F_X}^+ - \#(\text{R Nodes}) - \#(\text{NS Nodes})$ , which is exactly  $\dim_{\mathbb{C}} H^1(X, \mathcal{A})_+$ . The action of  $\text{Stab}_{\lambda} T_{F_X}^+$  on  $H^1(X, \mathcal{A})_+$  is thus faithful, and as such the action of  $T_{F_X}^+$  on  $\text{sDef}_A(Y, \mathcal{I})$  is also faithful. The odd small extension case is straightforward, since the fiber sequence (7.14) degenerates to first two terms thus realizing  $\text{sDef}_A(Y, \mathcal{I})$  as an affine space modeled on  $H^1(X, \mathcal{A})_+ = T_{F_X}^-$ . To summarize, we have the following result

**Proposition 7.12.** *Functor  $F_X : \text{sArt}_{\mathbb{C}} \rightarrow \text{Sets}$  is isomorphic to the functor  $h_R$  where*

$$\begin{aligned} R &= \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]], \\ a &= 3g - 3 + n_{\text{NS}} + n_{\text{R}}, \\ b &= 2g - 2 + n_{\text{NS}} + \frac{n_{\text{R}}}{2}. \end{aligned} \tag{7.18}$$

*The isomorphism is realized by a compatible family  $\{(Y_n, \phi_n) \in F_X(R/\mathfrak{m}_R^n)\}$ .*

*Proof.*  $F_X$  has a universal family by applying Schlessinger's Theorem 7.12 to what we have explained.  $R$  is formally smooth because the functor  $F_X$  is formally smooth (Proposition 7.11), hence  $R$  is the formal power series  $\mathbb{C}$ -superalgebra generated by the dual of  $T_{F_X}$ . ■

**Corollary 7.7.** *There exists a morphism  $\text{Spec } \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]] \rightarrow \overline{\mathfrak{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  whose reduction modulo  $\mathfrak{m}_R^n$  is equivalent to  $(Y_n, \phi_n)$ .*

*Proof.* Since  $X$  is projective over  $\text{Spec } \mathbb{C}$  (Lemma 7.24) and the ample line bundle lifts to  $Y_n$  in a compatible way for all  $n \geq 1$ , so we get a compatible system of closed embeddings  $Y_n \hookrightarrow \mathbb{P}_{R/\mathfrak{m}_R^n}^{M|N}$  for some integers  $M, N$ . By Grothendieck's Existence Theorem 7.6, there exists a

---

<sup>70</sup>Transitivity comes from the condition  $(S_1)$ .

proper  $R$ -superscheme  $Y$  with punctures  $E, \mathcal{D}$  and map  $\delta : \Omega_{Y/R}^1 \rightarrow \omega_{Y/R}$  and an isomorphism  $\phi : Y_{\mathbb{C}} \cong X$ , and the reduction of  $(Y, \phi)$  to  $R/\mathfrak{m}_R^n$  is equivalent to  $(Y_n, \phi_n)$ . The flatness of  $Y$  is a consequence of local flatness criterion.  $\blacksquare$

Note that the restriction of the universal family  $\{(Y_n, \phi_n) \in F_X(R/\mathfrak{m}_R^n)\}$  to  $\text{Art}_{\mathbb{C}}$  (category of Artinian bosonic  $\mathbb{C}$ -algebras) is a universal family for the deformations of  $X$  as spin curves, so the restriction of the morphism  $\text{Spec } \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]] \rightarrow \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  to  $\text{Spec } \mathbb{C}[[t_1, \dots, t_a]]$  factors through some affine étale chart  $U_0$  of  $\overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$ , which induces an isomorphism between  $\mathbb{C}[[t_1, \dots, t_a]]$  and the completion of  $\mathcal{O}_{U_0}$  at a closed point  $p$ . Let  $U$  be the superscheme  $(U_0, \wedge^\bullet \mathcal{O}_{U_0}^b)$ , i.e. formally adding independent odd variables  $\{\theta_1, \dots, \theta_n\}$  to the coordinate ring of  $U_0$ , so there is an isomorphism  $\widehat{\mathcal{O}}_{U,p} \cong \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]]$ . Combining Lemma 7.30 and Lemma 7.23, we obtain a superscheme  $V$  smooth over  $U$  with a morphism  $\text{Spec } \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]] \rightarrow V$  and a stable SUSY curve  $Y_V$  over  $V$  such that the universal family  $Y$  is isomorphic to the pull back of  $Y_V$ . Denote the image of the close point of  $\text{Spec } \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]]$  in  $V$  by  $q$ , consider the the embeddings

$$\begin{array}{ccccc} \text{Spec } \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]]/\mathfrak{m}^2 & \longrightarrow & \text{Spec } \mathcal{O}_V(V)/\mathfrak{m}_q^2 & \longrightarrow & \text{Spec } \mathcal{O}_U(U)/\mathfrak{m}_p^2 \\ \downarrow & & \downarrow & \nearrow & \downarrow \\ \text{Spec } \mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]] & \longrightarrow & V & \xrightarrow{\quad} & U \end{array},$$

defined by modulo  $\mathfrak{m}^2$ . Since the composition in the top row is an isomorphism, this gives an embedding of  $\text{Spec } \mathcal{O}_U(U)/\mathfrak{m}_p^2$  into  $\text{Spec } \mathcal{O}_V(V)/\mathfrak{m}_q^2$ . Extending the image of this embedding to a neighborhood of  $q$ , we obtain an locally closed sub superscheme  $W \subset V$  and  $W$  is étale over  $U$ . The restriction of  $Y_V$  to  $W$  induces a homomorphism  $\mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]] \rightarrow \widehat{\mathcal{O}}_{W,q}$  which is an isomorphism modulo  $\mathfrak{m}^2$ , thus it is an isomorphism because any automorphism of  $\mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]]$  which induces isomorphism on tangent space is an isomorphism. To summarize, we have the following

**Proposition 7.13.** *For every  $p \in \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}(\mathbb{C})$ , there exists an affine smooth  $\mathbb{C}$ -variety  $W$  with an element  $P \in \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}(W)$ , and a closed point  $q \in W$  such that  $P|_q$  is isomorphic to  $p$ . Furthermore,*

- (1) *The induced map  $P_{\text{bos}} : W_{\text{bos}} \rightarrow \overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is étale.*
- (2) *The induced map  $\widehat{P} : \widehat{\mathcal{O}}_{W,q} \rightarrow \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  identifies  $\widehat{\mathcal{O}}_{W,q}$  with the base of universal deformation of  $p$ .*

The next goal is to prove that  $P : W \rightarrow \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}}$  is étale (shrink  $W$  if necessary), for which we will need more ingredients, discussed in the next section.

### The Odd Tangent Space

Suppose that  $S$  is a affine smooth  $\mathbb{C}$ -variety of positive dimension and  $\pi : X \rightarrow S$  is a stable SUSY curve and is smooth on a open subscheme  $S_0 \subset S$ . Denote the underlying spin curve

of  $X$  by  $(\mathcal{C}, E, \mathcal{D}, \mathcal{E})$ . Let  $\mathcal{J}$  be a finite free sheaf on  $S$  and let the trivial extension of  $S$  by  $\mathcal{J}$  be  $S'$ . Assume that stable SUSY curve  $\pi' : X' \rightarrow S'$  is a lift of  $X \rightarrow S$ , then we have

**Lemma 7.31.** *The sheaf  $\mathcal{A}$  of automorphisms of  $X' \rightarrow S'$  which induce identity on  $X \rightarrow S$  is*

$$\omega_{\mathcal{C}/S}^{-1}(-E - \mathcal{D}) \otimes \mathcal{E} \otimes \pi^* \mathcal{J} \cong \mathcal{E}^\vee \otimes \pi^* \mathcal{J}.$$

*Proof.* On the smooth locus  $X^{\text{sm}}$ , it is well-known that (see Sections 4.2.1 and 4.2.3 of [11])  $\mathcal{A} = T_{\mathcal{C}/S}(-E - \mathcal{D}) \otimes \mathcal{E} \otimes \pi^* \mathcal{J} \cong \omega_{\mathcal{C}/S}^{-1}(-E - \mathcal{D}) \otimes \mathcal{E} \otimes \pi^* \mathcal{J}$ . It remains to determine the local structure of  $\mathcal{A}$  at nodes. Let the closed subscheme of R node (resp. NS node) be  $Z_R$  (resp.  $Z_{NS}$ ). Suppose that  $p \in X$  is closed and is a R node, then by the same argument in the proof of Lemma 7.29, we see that in an open affine neighborhood  $W_p$  of  $p$ , an element in  $\Gamma(W_p, \mathcal{A})$  is determined by derivation of the form  $yD^2$ , which is a rational section of  $T_{\mathcal{C}/S} \otimes \mathcal{E} \otimes \pi^* \mathcal{J}$  on the smooth locus  $W_p \setminus Z_R$ . Notice that  $W_p$  is normal,  $Z_R$  has codimension at least 2, and  $\omega_{\mathcal{C}/S}^{-1} \otimes \mathcal{E} \otimes \pi^* \mathcal{J}$  is reflexive and its restriction to  $W_p \setminus Z_R$  is isomorphic to  $T_{\mathcal{C}/S} \otimes \mathcal{E} \otimes \pi^* \mathcal{J}$ , so

$$\Gamma(W_p, \mathcal{A}) = \Gamma(W_p, \omega_{\mathcal{C}/S}^{-1} \otimes \mathcal{E} \otimes \pi^* \mathcal{J}),$$

by algebraic Hartogs's Theorem. The similar statement is also true for an open neighborhood of a NS node, whence the result follows by gluing.  $\blacksquare$

The object we will concern about is the space of deformations  $\text{sDef}_{S'}(X, \mathcal{J})$ . Note that if  $\mathcal{J} \rightarrow \mathcal{K}$  is a homomorphism of free  $\mathcal{O}_S$  sheaves, then there is a canonical map  $\text{sDef}_{S'}(X, \mathcal{J}) \rightarrow \text{sDef}_{S'}(X, \mathcal{K})$  defined by pull back of liftings of  $X$ . This endows  $\text{sDef}_{S'}(X, \mathcal{J})$  with a canonical  $\mathcal{O}_S$ -module structure via

- Addition: Induced by the addition map  $\mathcal{J} \times \mathcal{J} \rightarrow \mathcal{J}, (x, y) \mapsto x + y$ .
- Zero element: Induced by the trivial map  $0 \rightarrow \mathcal{J}$ .
- Inverse: Induced by inverse map  $\mathcal{J} \rightarrow \mathcal{J}, x \mapsto -x$ .
- $\mathcal{O}_S$ -action: For any element  $f \in \mathcal{O}_S(S)$ , the action of  $f$  on  $\text{sDef}_{S'}(X, \mathcal{J})$  is induced by  $\mathcal{J} \rightarrow \mathcal{J}, x \mapsto f \cdot x$ .

The goal of this section is to determine the  $\mathcal{O}_S$ -module structure of  $\text{sDef}_{S'}(X, \mathcal{J})$ . Indeed, we have

**Proposition 7.14.**  *$\text{sDef}_{S'}(X, \mathcal{J})$  is isomorphic to*

$$H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{J}.$$

*Moreover, the isomorphism is functorial in  $\mathcal{J}$  in the sense that for a homomorphism  $\mathcal{J} \rightarrow \mathcal{K}$  of free  $\mathcal{O}_S$  sheaves, the diagram*

$$\begin{array}{ccc}
\mathrm{sDef}_{S'}(X, \mathcal{J}) & \longrightarrow & \mathrm{sDef}_{S'}(X, \mathcal{K}) \\
\downarrow & & \downarrow \\
H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{J} & \longrightarrow & H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{K}
\end{array},$$

commutes.

*Proof.* The proof basically follows the same idea of Proposition 7.11. Again we have  $\mathrm{sDef}_A(\mathcal{O}_{Y,p}, \mathcal{J}) \xrightarrow{\sim} \mathrm{sDef}_A(\widehat{\mathcal{O}}_{Y,p}, \mathcal{J})$  and the latter is a one element set, since the odd deformation is trivial for all canonical models. So the same gluing argument shows that  $\mathrm{sDef}_{S'}(X, \mathcal{J})$  is a torsor under the action of  $H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{J}$ . It has a distinguished element which is the trivial deformation, so we have a canonical isomorphism of sets  $H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{J} \xrightarrow{\sim} \mathrm{sDef}_{S'}(X, \mathcal{J})$ . It is easy to see that this isomorphism commutes with the Abelian group structure and  $\mathcal{O}_S$ -action, since the deformations are characterized by a Čech cochain valued in  $\mathcal{E}^\vee \otimes \pi^* \mathcal{J}$ , whose groups structures and  $\mathcal{O}_S$ -action are exactly given by the description above. Now the functoriality in  $\mathcal{J}$  comes from the functoriality of Čech cochains. ■

Observe that  $\mathcal{E}^\vee$  is flat over  $S$  and its restrictions to fibers have cohomology concentrated in degree 1, so  $H^1(\mathcal{C}, \mathcal{E}^\vee)$  is a locally-free  $\mathcal{O}_S$ -module. From the Čech cochain description of  $\mathrm{sDef}_{S'}(X, \mathcal{J})$ , we see that

**Corollary 7.8.** *The isomorphism  $\mathrm{sDef}_{S'}(X, \mathcal{J}) \xrightarrow{\sim} H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{J}$  commutes with restriction to fiber map, i.e. the diagram*

$$\begin{array}{ccc}
\mathrm{sDef}_{S'}(X, \mathcal{J}) & \xrightarrow{\mathrm{res}} & \mathrm{sDef}_{s'}(X_s, \mathcal{J}(s)) \\
\downarrow & & \downarrow \\
H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_S} \mathcal{J} & \xrightarrow{\mathrm{res}} & H^1(\mathcal{C}_s, \mathcal{E}_s^\vee) \otimes_{\mathbb{C}} \mathcal{J}(s)
\end{array},$$

commutes, where  $s \in S$  is a closed point,  $s'$  is the superscheme constructed from  $s$  trivially extended by the fiber  $\mathcal{J}(s)$ .

### An Étale Atlas of $\overline{\mathcal{SM}}_{g, n_{\mathrm{NS}}, n_{\mathrm{R}}}$

**Lemma 7.32.** *Keep using the same notation as Proposition 7.13, then there exists an open affine neighborhood  $S$  of  $q$  in  $W$  such that  $\forall s \in S$ , the completion of  $P$  at  $s$  is a universal deformation family of  $P|_s$ .*

*Proof.* From the construction of  $W$ , we already know that the pullback of  $P$  to  $\widehat{\mathcal{O}}_{W,s,\mathrm{bos}}$  is a universal deformation family of  $P|_s$  as spin curves (since  $W \rightarrow \overline{\mathcal{S}}_{g, n_{\mathrm{NS}}, n_{\mathrm{R}}}$  is étale). To show that the pullback of  $P$  to  $\widehat{\mathcal{O}}_{W,s}$  is a universal deformation family of  $P|_s$  as stable SUSY curves, it suffices to show that the homomorphism  $\mathbb{C}[[t_1, \dots, t_a | \theta_1, \dots, \theta_b]] \rightarrow \widehat{\mathcal{O}}_{W,s}$  induced by  $\widehat{P}_s$  is an isomorphism on tangent spaces. The even tangent spaces are obviously isomorphic, so it remains to show that this induces isomorphism on odd tangent spaces.

Consider the restriction of  $P$  to  $W_1 \equiv \text{Spec } \mathcal{O}_W(W)/\mathfrak{n}_W^2$ , it gives rise to an element

$$[P] \in \text{sDef}_{W_1}(P_{W_0}, \mathcal{O}_{W_0}^{\oplus b}),$$

where  $W_0 = W_{\text{bos}}$  is the bosonic truncation. Hence by Proposition 7.14, it corresponds to an element in  $H^1(\mathcal{C}, \mathcal{E}^\vee) \otimes_{\mathcal{O}_{W_0}} \mathcal{O}_{W_0}^{\oplus b}$ , or equivalently

$$[P] : H^1(\mathcal{C}, \mathcal{E}^\vee)^\vee \rightarrow \mathcal{O}_{W_0}^{\oplus b},$$

where  $\mathcal{C}$  is the spin curve underlying  $P|_{W_0}$  with spinor sheaf  $\mathcal{E}$ . Because the restriction commutes with the identification between  $\text{sDef}$  and  $H^1$  (Corollary 7.8), the restriction of the homomorphism  $[P]$  to the fiber  $q$  is the homomorphism

$$[P_q] : H^1(\mathcal{C}_q, \mathcal{E}_q^\vee)^\vee \rightarrow \mathbb{C}^{\oplus b},$$

where the vector space  $\mathbb{C}^{\oplus b}$  is spanned by  $\{\theta_i\}$ , i.e. the cotangent space of the universal deformation base of  $P_q$ , thus  $[P_q]$  is isomorphism. It follows that there exists an open affine neighborhood  $S$  of  $q$  in  $W$  such that  $[P]$  is an isomorphism on  $S_{\text{bos}}$ . Then  $\forall s \in S$ , the homomorphism  $[P_s] : H^1(\mathcal{C}_s, \mathcal{E}_s^\vee)^\vee \rightarrow \mathbb{C}^{\oplus b}$  is isomorphism, whence  $\hat{P}_s$  induces an isomorphism between the universal deformation base of  $P|_s$  and  $\hat{\mathcal{O}}_{S,s}$ .  $\blacksquare$

**Proposition 7.15.** *The morphism  $S \rightarrow \overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  in Lemma 7.32 is representable and étale.*

*Proof.* Representability of  $S \rightarrow \overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  is a consequence of representability of the diagonal of  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  (Proposition 7.9). To show that it is étale is equivalent to show that for every small extension  $p : A \rightarrow A/\mathcal{J}$  of local Artinian  $\mathbb{C}$ -superalgebras, and for every commutative diagram

$$\begin{array}{ccc} \text{Spec } A/\mathcal{J} & \xrightarrow{f} & S \\ \downarrow i & & \downarrow \\ \text{Spec } A & \xrightarrow{g} & \overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} \end{array},$$

there exists a unique  $\tilde{f} : \text{Spec } A \rightarrow S$  making the diagram still commutative. Let the image of the closed point of  $\text{Spec } A/\mathcal{J}$  under  $f$  be  $s \in S$ , then the pull back of stable SUSY curve  $P$  over  $S$  to  $\text{Spec } A/\mathcal{J}$ , denoted by  $P_{A/\mathcal{J}}$ , extends to a stable SUSY curve  $P_A$  over  $\text{Spec } A$ , where  $P_A$  is defined by the morphism  $g$ . Since  $\hat{P}_s$  over  $\hat{\mathcal{O}}_{S,s}$  is a universal deformation, there exists a unique homomorphism  $\tilde{f} : \text{Spec } A \rightarrow \text{Spec } \hat{\mathcal{O}}_{S,s}$  which extends  $f : \text{Spec } A/\mathcal{J} \rightarrow \text{Spec } \hat{\mathcal{O}}_{S,s}$  such that  $\tilde{f}^*P \cong P_A$ .  $\blacksquare$

Now for every object  $p \in \overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}(\mathbb{C})$ , we can construct such  $S \rightarrow \overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  in the Proposition 7.15 that  $p$  is equivalent to  $q \in S(\mathbb{C})$ , thus we obtained an étale covering of  $\overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}}$  and concludes the proof of Theorem 7.13.

### 7.3.1 Gluing of Punctures

Similar to the spin curves, we can define a set of gluing operations functorially for stable SUSY curves. For NS punctures, this is easy, since NS punctures are sections of the base superscheme, so we can use this to define a gluing directly. More precisely, suppose that  $X_1$  and  $X_2$  are stable SUSY curves over a base superscheme  $S$ , and that  $\mathfrak{n}_{\text{NS}}^1$  and  $\mathfrak{n}_{\text{NS}}^2$  are nonzero, then we can pick one NS puncture from each curve, say that the  $i_1$ 'th NS puncture  $E_{i_1}$  on  $X_1$  and  $i_2$ 'th NS puncture  $E_{i_2}$  on  $X_2$ . The projection  $E_{i_1} \rightarrow S$  is an isomorphism, same for  $E_{i_2}$ , so we can use these isomorphisms to define a canonical identification between  $\varphi : E_{i_1} \cong E_{i_2}$ , and glue them to get a new superscheme  $X = X_1 \cup_{\varphi} X_2$ . Note that there are two short exact

sequence of sheaves

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X & \longrightarrow & p_{1*}\mathcal{O}_{X_1} \oplus p_{2*}\mathcal{O}_{X_2} & \longrightarrow & \mathcal{O}_S \longrightarrow 0 \\ & & & & \downarrow \delta_1 \oplus \delta_2 & & \\ 0 & \longrightarrow & p_{1*}\omega_{X_1/S} \oplus p_{2*}\omega_{X_2/S} & \longrightarrow & \omega_{X/S} & \longrightarrow & \mathcal{O}_S \longrightarrow 0 \end{array},$$

where  $p_i : X_i \rightarrow X$  are natural projections. So we can define the derivation  $\delta$  on  $X$  as the composition

$$\mathcal{O}_X \rightarrow p_{1*}\mathcal{O}_{X_1} \oplus p_{2*}\mathcal{O}_{X_2} \xrightarrow{\delta_1 \oplus \delta_2} p_{1*}\omega_{X_1/S} \oplus p_{2*}\omega_{X_2/S} \rightarrow \omega_{X/S}.$$

This construction is functorial hence giving rise to an operation on corresponding moduli spaces:

$$\mathfrak{m}_{i_1, i_2}^{\text{NS}} : \overline{\mathcal{SM}}_{g_1, \mathfrak{n}_{\text{NS}}^1, \mathfrak{n}_{\text{R}}^1} \times \overline{\mathcal{SM}}_{g_2, \mathfrak{n}_{\text{NS}}^2, \mathfrak{n}_{\text{R}}^2} \rightarrow \partial \overline{\mathcal{SM}}_{g_1+g_2, \mathfrak{n}_{\text{NS}}^1+\mathfrak{n}_{\text{NS}}^2-2, \mathfrak{n}_{\text{R}}^1+\mathfrak{n}_{\text{R}}^2}.$$

This can be defined for a single stable SUSY curve with at least two NS punctures as well:

$$\mathfrak{g}_{i,j}^{\text{NS}} : \overline{\mathcal{SM}}_{g, \mathfrak{n}_{\text{NS}}, \mathfrak{n}_{\text{R}}} \rightarrow \partial \overline{\mathcal{SM}}_{g+1, \mathfrak{n}_{\text{NS}}-2, \mathfrak{n}_{\text{R}}}.$$

Note that these maps depends on the choice of  $i_1$ ,  $i_2$ ,  $i$ , and  $j$ , in an equivariant way. For example, if  $\sigma_{i_1, i'_1} \in S_{\mathfrak{n}_{\text{NS}}^1}$  switches  $i_1$  and  $i'_1$  (it acts  $\overline{\mathcal{SM}}_{g_1, \mathfrak{n}_{\text{NS}}^1, \mathfrak{n}_{\text{R}}^1}$  naturally), then we have  $\mathfrak{m}_{i'_1, i_2}^{\text{NS}} = \mathfrak{m}_{i_1, i_2}^{\text{NS}} \circ \sigma_{i_1, i'_1}$ .

For gluing R divisors, the  $\mathbb{Z}/2$ -ambiguity is lifted to the ambiguity of choosing an isomorphism between the structure sheaf of two R divisors. The solution is also similar to the spin curve case: we add a canonical isomorphism *by hand*. Suppose that  $X_1$  and  $X_2$  are stable SUSY curves over a base superscheme  $S$ , and that  $\mathfrak{n}_{\text{R}}^1$  and  $\mathfrak{n}_{\text{R}}^2$  are nonzero, we can pick one R divisor from each curve,  $\iota_1 : \mathcal{D}_{i_1} \subset X_1$  and  $\iota_2 : \mathcal{D}_{i_2} \subset X_2$ . Note that  $\mathcal{D}_i$  are smooth over  $S$  of relative dimension  $(0|1)$ . The derivations  $\delta_1, \delta_2$  induce  $\mathcal{O}_S$ -linear derivations

$$\begin{aligned} d_1 &: \mathcal{O}_{\mathcal{D}_{i_1}} \rightarrow \omega_{\mathcal{D}_{i_1}/S}, \\ d_2 &: \mathcal{O}_{\mathcal{D}_{i_2}} \rightarrow \omega_{\mathcal{D}_{i_2}/S}, \end{aligned}$$

as can be easily seen from a local computation. Consider the superscheme

$$G_{i_1, i_2} \subset \text{Isom}_S(\mathcal{D}_{i_1}, \mathcal{D}_{i_2}),$$

parametrizing isomorphisms from  $\mathcal{D}_{i_1}$  to  $\mathcal{D}_{i_2}$  which commutes with  $d_i$ . Locally on  $S$ , after choosing coordinate system of  $\mathcal{D}_{i_1}, \mathcal{D}_{i_2}$ , denoted by  $\theta_1, \theta_2$ , we can assume that  $d_1, d_2$  can be written as

$$d_i : \theta_i \mapsto [1/d\theta_i], \quad 1 \mapsto 0,$$

so for any  $S$ -superscheme  $T$ ,  $G_{i_1, i_2}(T)$  is locally generated by homomorphisms of form

$$\theta_2 \mapsto \{\pm\theta_1 + a | a \in \mathcal{O}_T^-\}.$$

Therefore,  $G_{i_1, i_2}$  is locally on  $S$  isomorphic to two copies of  $\mathbb{A}^{0|1}$ . Note that  $G_{i_1, i_2, \text{bos}}$  on  $S_{\text{bos}}$  is exactly the  $G_{i_1, i_2}$  we constructed in B.9.

Base change to the superscheme  $G_{i_1, i_2}$ , there is a universal isomorphism  $\alpha : \mathcal{D}_{i_1} \cong \mathcal{D}_{i_2}$  which commutes with  $d_i$ , so we can take the gluing of  $X_1 \times_S G_{i_1, i_2}$  and  $X_2 \times_S G_{i_1, i_2}$  to be  $X = X_1 \cup_{\alpha} X_2$ . Note that there are two short exact sequence of sheaves connected by derivations:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X & \longrightarrow & p_{1*}\mathcal{O}_{X_1} \oplus p_{2*}\mathcal{O}_{X_2} & \xrightarrow{\alpha + \text{Id}} & \mathcal{O}_{\mathcal{D}_{i_2}} \longrightarrow 0 \\ & & & & \downarrow \delta_1 \oplus \delta_2 & & \downarrow d_2 \\ 0 & \longrightarrow & \omega_{X/S} & \longrightarrow & p_{1*}\omega_{X_1/S} \oplus p_{2*}\omega_{X_2/S} & \xrightarrow{\alpha + \text{Id}} & \omega_{\mathcal{D}_{i_2}/S} \longrightarrow 0 \end{array},$$

such that the diagram commutes. So there is a derivation  $\delta : \mathcal{O}_X \rightarrow \omega_{X/S}$  and this gives  $X$  a structure of stable SUSY curve. The construction is also functorial, hence giving rise to an operation on corresponding moduli spaces:

$$\begin{array}{ccc} \mathcal{G}_{i_1, i_2} & \xrightarrow{\mathfrak{m}_{i_1, i_2}^R} & \partial \overline{\mathcal{SM}}_{g_1+g_2, n_{\text{NS}}^1+n_{\text{NS}}^2, n_{\text{R}}^1+n_{\text{R}}^2-2} \\ \pi_{i_1, i_2} \downarrow & & \\ \overline{\mathcal{SM}}_{g_1, n_{\text{NS}}^1, n_{\text{R}}^1} \times \overline{\mathcal{SM}}_{g_2, n_{\text{NS}}^2, n_{\text{R}}^2} & & \end{array}.$$

This can be defined for a single punctured SUSY curve with at least two NS puncturing divisors as well:

$$\begin{array}{ccc} \mathcal{G}_{i, j} & \xrightarrow{\mathfrak{g}_{i, j}^R} & \partial \overline{\mathcal{SM}}_{g+1, n_{\text{NS}}, n_{\text{R}}-2} \\ \pi_{i, j} \downarrow & & \\ \overline{\mathcal{SM}}_{g, n_{\text{NS}}, n_{\text{R}}} & & \end{array}.$$

Note that  $\pi_{i_1, i_2}$  and  $\pi_{i, j}$  are locally isomorphic to a  $\mathbb{Z}/2 \ltimes \mathbb{A}^{0|1}$ -bundle. Similar to the gluing of NS punctures, these maps depends on the choice of  $i_1, i_2$  and  $i, j$  in an equivariant way.



## 8 Discussion, and Future Directions

In this last section, we would like to point out few interesting research directions

- As we have shown that genus- $g$  heterotic-string and type-II-superstring field theory vertices exist for any type of boundary components.

It would be interesting to explicitly construct these vertices. An attempt to construct type-II-superstring field theory vertices using a generalization of the minimal-area problem of ordinary Riemann surfaces [90–98] to the case of  $\mathcal{N} = 1$  super-Riemann surface is considered in<sup>71</sup> [99].

- We also proved that the solution to the BV QME in  $\mathcal{F}(\mathcal{S})$  are unique up to homotopies in the homotopy category of BV algebras.

It is shown by Costello that the space of homotopy-equivalent solutions to the BV QME in the moduli of bordered ordinary Riemann surfaces is a point [26]. This together with [78] shows that the gauge-fixed action of bosonic-string field theory is unique. For another investigation in proving the uniqueness of bosonic-string field theory see [100].

As we have explained in [a choice of string vertex](#), the physics of string field theory should not be dependent on the choice of string vertices. On the other hand, we have proven the existence of string vertices by defining a quasi-isomorphism to the BV algebra of the complexes associated to the bordered spin-Riemann surfaces, and the solution to the BV QME in the latter BV algebra is unique up to homotopy. It would be very interesting to find the space of homotopy-equivalent solutions to the BV QME that give rise to string vertices. This sheds some light on the question of dependence of resulting string field theory on the choice of string vertex. One possibility is the following. Assuming that the space of homotopy-equivalent solutions of the BV QME that give rise to string vertices has a trivial mapping-class group (the group of large diffeomorphisms), then any two solutions of the BV quantum master equation can be related by a sequence of small deformations of one of them. Then, a result similar to [78] for the heterotic-string and type-II superstring field theories would show that the two gauge-fixed actions constructed using the two string vertices are the same upon using two different gauge-fixing conditions.

- We have proven the existence of type-II-superstring vertices. Two related questions are as follows
  1. In the presence of D-branes, we need to deal with surfaces with boundaries with both open-string punctures on the boundaries and closed-string punctures on the interior of the surface [101, 102]. One way to deal with such surfaces is to consider

---

<sup>71</sup>We would like to thank Ted Erler who pointed us to this paper.

the double of the surface. We refer to section 7 of [11] for details. So, one might be able to prove the existence of type-II-superstring vertices in the presence of D-branes by considering the doubled surface. As we have shown, there is always a solution to the BV QME in the BV algebra of complexes associated to moduli space of such or corresponding bordered surfaces.

2. Another interesting and relevant direction is the case of unoriented string field theory [102, 103]. Since this theory is the orientifold of type-IIB-superstring theory, and the proof of the existence of the string vertex for the latter has been given here, it would be interesting to prove the existence of string vertices for unoriented string field theory as well.

We leave these and related questions for future works.

**Acknowledgments** It is our pleasure to thank Kevin Costello whose works and ideas shaped the main motivation behind this work. We also thank him for discussion and guidance on this work. A special gratitude goes to Ted Erler who provided detailed and critical comments on various parts of an earlier version of this manuscript. We also thank Ivo Sachs for useful discussions. SFM is grateful to Davide Gaiotto for his support during his stay at Perimeter Institute. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research and Innovation.

## A The Picture-Changing Operation

In this appendix, we define the picture-changing operation and explain its various incarnations. We also explain its relation to the natural operation defined on space of form on a supermanifold. Following [16, 51, 104], we then argue that all these incarnations are different manifestations of the same concept.

### A.1 The Canonical Framework: Friedan-Shenker-Martinec Approach

In this section, we explain the picture-changing operation in the Friedan-Shenker-Martinec (FSM) formulation of covariant superstring perturbation theory [15].

The fixing of superconformal invariance of the worldsheet theory of RNS-superstring theory will give rise to two sets of antighost-ghost systems

1. The fermionic reparametrization ghosts corresponding to a  $bc$  system with fields having conformal dimensions  $\Delta_b = 2$  and  $\Delta_c = -1$ .
2. The bosonic local supersymmetry ghosts corresponding to a  $\beta\gamma$  system with fields having conformal dimensions  $\Delta_\beta = 3/2$  and  $\Delta_\gamma = -1/2$ .

The  $\beta\gamma$  system has a first-order action and as such there is no ground state for the system<sup>72</sup>. The resolution of the problem was found in [15]. The idea is that instead of  $\beta\gamma$  system, one considers its bosonization in terms of the bosonic field  $\phi$  and the fermionic fields  $\eta$  and  $\xi$  using the following identifications

$$\beta(z) \mapsto \partial_z \xi(z) e^{-\phi(z)}, \quad \gamma(z) \mapsto \eta(z) e^{\phi(z)}. \quad (\text{A.1})$$

One can then define a family of states parametrized by  $\phi$ -charge  $n$  of the corresponding state

$$|n\rangle \equiv \lim_{z \rightarrow 0} e^{n\phi(z)} |0\rangle, \quad (\text{A.2})$$

where  $|0\rangle$  is the  $\text{SL}(2, \mathbb{C})$ -invariant vacuum.  $n$  is called the *picture number* [105, 106]. The action of  $\beta$  and  $\gamma$  does not change  $n$ . This means that  $\{|n\rangle\}_{n \in \mathbb{Z}}$  can be used to construct representations of the algebra of modes of  $\beta$  and  $\gamma$  fields, since they have picture number zero. Representations constructed using different  $n$  are inequivalent. This is a crucial property of the bosonic nature of the  $\beta\gamma$  system. The choice of a picture is thus a choice of vacuum for the  $\beta\gamma$  system [15].

One can expand these fields as

$$\beta(z) = \sum_k \frac{\beta_k}{z^{k+\frac{3}{2}}}, \quad \gamma(z) = \sum_k \frac{\gamma_k}{z^{k-\frac{1}{2}}}, \quad (\text{A.3})$$

where  $k \in \frac{1}{2} + \mathbb{Z}$  (for NS sector) or  $k \in \mathbb{Z}$  (for R sector). From these, we conclude that

$$\beta_k = \oint \frac{dz}{2\pi i} z^{k+\frac{1}{2}} \beta(z), \quad \gamma_k = \oint \frac{dz}{2\pi i} z^{k-\frac{3}{2}} \gamma(z). \quad (\text{A.4})$$

These modes satisfy the commutation relation

$$[\gamma_m, \beta_n] = \delta_{m, -n}. \quad (\text{A.5})$$

The other commutators are zero. There is a ghost current  $\mathcal{Q}_{\text{gh}} \equiv \sum_n : \beta_n \gamma_{-n} :$ , where  $: \cdot :$  denotes the normal ordering. The ghost-charges of modes are

$$[\mathcal{Q}_{\text{gh}}, \beta_n] = +n\beta_n, \quad [\mathcal{Q}_{\text{gh}}, \gamma_n] = -n\gamma_n. \quad (\text{A.6})$$

We can also introduce the operators that interpolate between the ghost-see levels. These operators are formally written as  $\delta(\beta_n)$  and  $\delta(\gamma_n)$ . The action of these operators on a vacuum with picture number  $n$  is defined via [107]

$$\delta(\beta_{-n-\frac{3}{2}})|n\rangle = |n+1\rangle, \quad \delta(\gamma_{n+\frac{1}{2}})|n\rangle = |n-1\rangle. \quad (\text{A.7})$$

---

<sup>72</sup>This can be seen as follows. The action of the mode  $\gamma_{\frac{1}{2}}$  of  $\gamma(z)$  field on the  $\text{SL}(2, \mathbb{C})$ -invariant vacuum  $|0\rangle$  is not zero. Since  $\gamma_{\frac{1}{2}}$  is a bosonic mode, it can be applied to  $|0\rangle$  arbitrarily. However, this action reduced the  $L_0$  eigenvalue, i.e. the energy, of the resulting state by  $\frac{1}{2}$ . In this way one can construct states with arbitrary negative energy and as a result there is no ground state for the system. To cure this problem, one has to construct a state that is annihilated by all of the positive modes. For more details, see section 5.2 of [15].

We can thus assign picture-numbers  $+1$  to  $\delta(\beta_n)$  and  $-1$  to  $\delta(\gamma_n)$ .

To determine the picture number of the NS and R vacua, we consider the state (A.2) and act with the modes of  $\beta$  and  $\gamma$  fields. The action of the mode  $\beta_k$  on the state (A.2) gives

$$\begin{aligned}\beta_k|n\rangle &\sim \oint \frac{dz}{2\pi i} z^{k+\frac{1}{2}} \beta(z) e^{n\phi(0)} = \oint \frac{dz}{2\pi i} z^{k+\frac{1}{2}} \partial_z \xi(z) e^{-\phi(z)} e^{n\phi(0)} \\ &= \oint \frac{dz}{2\pi i} z^{k+\frac{1}{2}+n} : \partial_z \xi(z) e^{-\phi(z)} e^{n\phi(0)} : .\end{aligned}\tag{A.8}$$

This vanishes only if  $k + \frac{1}{2} + n \geq 0$ . Similarly, the action of the mode  $\gamma_k$  on the state (A.2) gives

$$\begin{aligned}\gamma_k|n\rangle &\sim \oint \frac{dz}{2\pi i} z^{k-\frac{3}{2}} \gamma(z) e^{n\phi(0)} = \oint \frac{dz}{2\pi i} z^{k-\frac{3}{2}} \eta(z) e^{\phi(z)} e^{n\phi(0)} \\ &= \oint \frac{dz}{2\pi i} z^{k-\frac{3}{2}-n} : \eta(z) e^{\phi(z)} e^{n\phi(0)} : .\end{aligned}\tag{A.9}$$

This vanishes only if  $k - \frac{3}{2} - n \geq 0$ . We want to choose  $n$  such that  $\beta_k|n\rangle = \gamma_k|n\rangle = 0$  for  $k > 0$ . From the possible values of  $k$  for the NS and R sectors, it is clear that  $n$  is integer for the NS sector and half-integer for the R sector. The only picture numbers that satisfy these inequalities are  $n = -1$  for states in the NS sector, and  $n = -\frac{1}{2}, -\frac{3}{2}$  for states in the R sector. The R vacua are obtained by the action of spin fields

$$|\Sigma_{\pm}\rangle \equiv \lim_{z \rightarrow 0} e^{\pm \frac{1}{2} \phi(z)} |0\rangle \tag{A.10}$$

on the NS ground state with picture number  $-1$ , i.e. spin fields interpolate between NS and R ground states. The picture numbers  $-1$  and  $-\frac{1}{2}$  are called canonical picture numbers<sup>73</sup>. It turns out that R states with picture-number  $-\frac{3}{2}$  are useful for formulation of off-shell amplitudes in superstring theory [17] and also for the formulation of heterotic and type-II closed-superstring field theories [108]. The main reason is that a formulation using the states from R sector with picture number  $-\frac{3}{2}$  avoids the introduction of the so-called *inverse picture-changing operators*.

Once a particular vacuum  $|n\rangle$  is chosen, one can build the spectrum of the ghost system upon it. Therefore, the question is that whether these inequivalent description (i.e. inequivalent representation of the algebra of modes) of the spectrum build upon different vacua with different picture numbers are related. It is clear from the formulas (A.1) that the superstring BRST complex does not contain states involving the zero-mode  $\xi_0$  of the  $\xi$  field, states like  $\xi_0|\psi\rangle$ . Consider the state  $|\psi'\rangle \equiv \mathcal{Q}\xi_0|\psi\rangle$ , for a BRST-closed state  $|\psi\rangle$ , where  $\mathcal{Q}$  is the

---

<sup>73</sup>They give rise to the canonical fermionic dimension for the superstack. Doing superstring theory on other negative pictures corresponds to the modification of the fermionic dimension of superstack. For more details see section 4.3 of [11].

superstring BRST operator.  $|\psi'\rangle$  has three properties, 1) it does not contain the zero-mode of  $\xi(z)$  field, which can be seen by using the explicit form of  $\mathcal{Q}$ , 2) it is obviously BRST-closed, i.e. it is an element of the BRST cohomology and as such it is a physical state, 3) it is not a representative of the trivial class of the superstring BRST cohomology since  $\xi_0|\psi\rangle$  is not an element of the superstring BRST complex (it does contain zero-mode of  $\xi(z)$  field and, as such, it is not part of the spectrum of the  $\beta\gamma$  system). One can thus define the so-called *picture-changing operator* (PCO) <sup>74</sup>. If  $\mathcal{V}_n(z)$  denotes a vertex operator built on a vacuum with picture number  $n$ , then

$$\mathcal{V}_{n+1}(z) = \mathcal{X}[\mathcal{V}_n(z)] \equiv \{\mathcal{Q}, \xi(z)\mathcal{V}_n(z)\}, \quad (\text{A.11})$$

where  $\mathcal{X}$  is the PCO (for explicit examples of the construction of vertex operators in two different pictures using the PCO, see section 5.3 of [15]). The explicit form of PCO is thus

$$\mathcal{X}(z) = \{\mathcal{Q}, \xi(z)\}. \quad (\text{A.12})$$

Thus, the answer to the above question is that the vertex operators associated to the same physical state built upon vacua with different picture number are related by the picture-changing operation [15]. In other words, if we think of physical states with a particular picture number as elements of the BRST cohomology of the BRST complex defined by that picture number (picture numbers are part of the grading of the complex [51]), the PCO maps the BRST cohomologies of the BRST complexes with different picture numbers into each other. It can be shown that the computation of *on-shell* superstring scattering amplitudes using different picture numbers gives the same result [15].

It turns out that picture-changing operators naturally enters the superstring path integral by integration over the odd moduli of superstack [16]. We turn to this point in the next section.

## A.2 The Path Integral Framework: Verlinde-Verlinde Approach

In this section, we review the results of [16] due to Verlinde and Verlinde (VV). It is argued there that the picture-changing operation arises in the string measure as the integration over odd moduli of superstack. We will be brief and refer to [16] or section 3.6.1 and 3.6.2 of [12] for more details.

The simplest amplitude is the genus- $g$  superstring vacuum amplitude given by [109]

$$\mathcal{Z} = \int_{\mathcal{SM}_g} \mathcal{Z}(\mathbf{m}, \widehat{\mathbf{m}}) = \int_{\mathcal{SM}_g} \mathcal{Z}(X, \mathcal{G}; \mathbf{m}, \widehat{\mathbf{m}}) |\mathcal{Z}(B, C; \mathbf{m}, \widehat{\mathbf{m}})|^2, \quad (\text{A.13})$$

where  $\mathcal{SM}_g$  is the superstack of genus- $g$  super-Riemann surfaces  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$ , and  $\mathbf{m}$  and  $\widehat{\mathbf{m}}$  collectively denote the even and odd moduli of  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$ , and

$$\mathcal{Z}(X, \mathcal{G}; \mathbf{m}, \widehat{\mathbf{m}}) \equiv \int [\mathcal{D}X] e^{-S[X, \mathcal{G}]},$$

---

<sup>74</sup>Here, we are not dealing with the more complicated issue of vertical integration which involves the introduction of a modified picture-changing operator [49]. See also [22] for an alternative approach to the vertical integration procedure from the point of view of large Hilbert space.

$$\mathcal{Z}(B, C; \mathbf{m}, \widehat{\mathbf{m}}) \equiv \int [\mathcal{D}B, \mathcal{D}C] e^{-S[B, C]} \left( \prod_{i=1}^{3g-3} \langle \mu_i, B \rangle \prod_{a=1}^{2g-2} \delta(\langle \widehat{\mu}_a, B \rangle) \right). \quad (\text{A.14})$$

Here  $\mathcal{G}$  denotes the gravitino field. The fields  $X$ ,  $B$ , and  $C$  have the following expansions

$$X^\mu(z|\theta) = x^\mu(z) + \theta\psi^\mu(z), \quad B(z|\theta) = \beta(z) + \theta b(z), \quad C(z|\theta) = c(z) + \theta\gamma(z). \quad (\text{A.15})$$

$x^\mu(z)$  are space-time coordinates,  $\psi^\mu(z)$  are space-time fermions,  $b(z)$  and  $c(z)$  are fields of the diffeomorphism ghost system, and  $\beta(z)$  and  $\gamma(z)$  are fields of the superconformal ghosts system.  $\{\mu_i, \widehat{\mu}_a\}$  is a basis of super-Beltrami differentials and the respective pairings  $\langle \mu_i, B \rangle$  and  $\langle \widehat{\mu}_a, B \rangle$  are defined by integration over  $\mathcal{R}$ .

The integration over the odd coordinates of the superstack can be done as follows. It turns out that the dependence of the action on the odd moduli of super-Riemann surfaces  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$  is contained in the gravitino field  $\mathcal{G}$ . Let us explain the reason. Consider a surface  $\mathcal{R}(\mathbf{m}, \mathbf{0})$ , i.e. a surface that does not have any odd moduli. These moduli can be turned on by doing a quasi-superconformal transformation generated by an odd vector field  $\widehat{v}(z, \bar{z})$  [107]

$$\begin{aligned} z &\longrightarrow z' = z + \theta \widehat{v}(z, \bar{z}), \\ \theta &\longrightarrow \theta' = \theta + \widehat{v}(z, \bar{z}) + \frac{1}{2} \theta \widehat{v}(z, \bar{z}) \partial_z \widehat{v}(z, \bar{z}). \end{aligned} \quad (\text{A.16})$$

The fields on the surface  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$ , obtained from  $\mathcal{R}(\mathbf{m}, \mathbf{0})$  by a quasi-superconformal transformation like (A.16), can be written in terms of the field on  $\mathcal{R}(\mathbf{m}, \mathbf{0})$  by a field-redefinition depending on  $\widehat{v}(z, \bar{z})$ . Once the matter action  $S[X, \mathcal{G}]$  on  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$  is written, it turns out that it can be identified with the Brink-Di Vecchia-Howe action for massless fields  $x^\mu(z)$  and  $\psi^\mu(z)$  coupled to  $\mathcal{N} = 1$  supergravity in the Wess-Zumino gauge [110], in which gravitino field can be identified with the odd vector field  $\widehat{v}(z, \bar{z})$  as follows. Since  $\widehat{v}(z, \bar{z})$  is a  $(-\frac{1}{2}, 0)$ -differential, on a genus- $g$   $\mathcal{N} = 1$  super-Riemann surface, it can be expanded in a basis of odd moduli  $\{\widehat{m}^a\}$  as follows

$$\widehat{v}(z, \bar{z}) = \sum_{a=1}^{2g-2} \widehat{m}^a v_a(z, \bar{z}). \quad (\text{A.17})$$

On the other hand, the gravitino field<sup>75</sup> can be similarly expanded

$$\mathcal{G} = \sum_{a=1}^{2g-2} \widehat{m}^a \mathcal{G}_a(z, \bar{z}), \quad (\text{A.18})$$

---

<sup>75</sup>Consider a super-Riemann surface  $\mathcal{R}$  with  $\mathfrak{n}_{\text{NS}}$  NS punctures located at  $p_1, \dots, p_{\mathfrak{n}_{\text{NS}}}$  and  $\mathfrak{n}_{\text{R}}$  R punctures located at  $q_1, \dots, q_{\mathfrak{n}_{\text{R}}}$ . As has been explained in section 5.4 of [12], the gravitino field  $\mathcal{G}$  is an element of  $H^1(\mathcal{R}_{\text{red}}, \widehat{L})$ , where

$$\widehat{L} \simeq L \otimes \mathcal{O} \left( - \sum_{a=1}^{\mathfrak{n}_{\text{NS}}} p_a \right), \quad L^2 \simeq T\mathcal{R}_{\text{red}} \otimes \mathcal{O} \left( - \sum_{a=1}^{\mathfrak{n}_{\text{R}}} q_a \right),$$

up to  $\widehat{L}$ -valued gauge transformations.  $\mathcal{R}_{\text{red}}$  is the reduced space of the super-Riemann surface  $\mathcal{R}$ . The dimension of  $H^1(\mathcal{R}_{\text{red}}, \widehat{L})$  is  $2g - 2 + \mathfrak{n}_{\text{NS}} + \frac{1}{2}\mathfrak{n}_{\text{R}}$ , i.e. the fermionic dimension of the superstack. In the above computation,  $\mathfrak{n}_{\text{NS}} = 0 = \mathfrak{n}_{\text{R}}$ , and  $\{\widehat{m}^a\}$  is a basis for  $H^1(\mathcal{R}_{\text{red}}, T\mathcal{R}_{\text{red}}^{\frac{1}{2}})$ . See also section 2.2 of [11].

where the differentials  $\mathcal{G}_a(z, \bar{z})$  can be chosen to be independent of both even and odd moduli of  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$ . One can then identify

$$\mathcal{G}_a(z, \bar{z}) = 2\bar{\partial}v_a(z, \bar{z}). \quad (\text{A.19})$$

Therefore,  $\mathcal{G}$  is a  $(-\frac{1}{2}, 1)$ -differential. In the basis chosen for the gravitino modes  $\mathcal{G}_a(z, \bar{z})$ ,  $\langle \mu_i, B \rangle_{\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})} = \langle \mu_i, b \rangle_{\mathcal{R}(\mathbf{m}, \mathbf{0})}$ , and  $\langle \widehat{\mu}_a, B \rangle_{\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})} = \langle \mathcal{G}_a, \beta \rangle_{\mathcal{R}(\mathbf{m}, \mathbf{0})}$ <sup>76</sup>. Using this choice, the worldsheet action  $S[X, \mathcal{G}; B, C]_{\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})}$  on  $\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})$  can be expanded as

$$S[X, \mathcal{G}; B, C]_{\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})} = S[X; B, C]_{\mathcal{R}(\mathbf{m}, \mathbf{0})} + \sum_{a=1}^{2g-2} \widehat{m}^a \langle \mathcal{G}_a, T_F \rangle, \quad (\text{A.20})$$

where  $T_F$  is the worldsheet supercurrent, the fermionic counterpart of the usual energy-momentum tensor  $T_B$ <sup>77</sup>, of the total superconformal field theories of matter and ghosts whose explicit form is not important for us. Expanding the exponential, we get

$$\mathcal{Z}(\mathbf{m}, \widehat{\mathbf{m}}) = \int \mathcal{D}[X; B, C] \left[ \prod_{i=1}^{3g-3} \langle \mu_i, b \rangle \prod_{a=1}^{2g-2} (1 + \widehat{m}^a \langle \mathcal{G}_a, T_F \rangle) \delta(\langle \mathcal{G}_a, \beta \rangle) \right] e^{-S[X; B, C]_{\mathcal{R}(\mathbf{m}, \mathbf{0})}}. \quad (\text{A.21})$$

In this final form, the integration over  $\{\widehat{m}^a\}$  can be easily done. The result is

$$\mathcal{Z} = \int_{\mathbb{S}_g} \prod_{a=1}^{2g-2} d\widehat{m}^a \mathcal{Z}(\mathbf{m}, \widehat{\mathbf{m}}) = \int_{\mathbb{S}_g} \int \mathcal{D}[X; B, C] \left[ \prod_{i=1}^{3g-3} \langle \mu_i, b \rangle_{\mathcal{R}(\mathbf{m}, \mathbf{0})} \prod_{a=1}^{2g-2} \mathcal{X}(\mathcal{G}_a) \right] e^{-S[X; B, C]_{\mathcal{R}(\mathbf{m}, \mathbf{0})}}, \quad (\text{A.22})$$

where  $\mathbb{S}_g$  is the moduli space of split surfaces  $\mathcal{R}(\mathbf{m}, \mathbf{0})$ , and  $\mathcal{X}$  is the *picture-changing operator* given by

$$\mathcal{X}(\mathcal{G}_a) \equiv \langle \mathcal{G}_a, T_F \rangle \delta(\langle \mathcal{G}_a, \beta \rangle) \Big|_{\mathcal{R}(\mathbf{m}, \mathbf{0})}. \quad (\text{A.23})$$

Therefore, the picture-changing operation is the operation of absorbing a zero-mode of  $\beta$  ( $\delta(\langle \mathcal{G}_a, \beta \rangle)$ ) together with the integration over the corresponding odd moduli of superstack.

---

<sup>76</sup>In general, we have

$$\langle \mu_i, B \rangle_{\mathcal{R}(\mathbf{m}, \widehat{\mathbf{m}})} = \langle \mu_i, b \rangle_{\mathcal{R}(\mathbf{m}, \mathbf{0})} + \sum_{a=1}^{2g-2} \widehat{m}^a \langle \nabla_i \mathcal{G}_a, \beta \rangle_{\mathcal{R}(\mathbf{m}, \mathbf{0})},$$

where  $\nabla_i \mathcal{G}_a$  denotes the covariant derivative of the gravitino mode  $\mathcal{G}_a$  with respect to the even moduli of the surface. The explicit form of  $\nabla_i \mathcal{G}_a$  is not important for the following conclusion. Since we have chosen  $\mathcal{G}_a$ s to be independent of even moduli  $\{m^a\}$ , and hence  $\nabla_i \mathcal{G}_a$  vanishes identically.

<sup>77</sup>This means that the OPEs of  $T_B$  and  $T_F$ , i.e.  $T_B(z)T_B(w)$ ,  $T_B(z)T_F(w)$ , and  $T_F(z)T_F(w)$ , generates a copy of the  $\mathcal{N} = 1$  superconformal algebra. They have the following mode-expansion

$$T_B(z) = \sum_n \frac{L_n}{z^{n+2}}, \quad T_F(z) = \sum_r \frac{G_r}{z^{n+\frac{3}{2}}}.$$

$r \in \mathbb{Z} + \frac{1}{2}$  (for NS sector) or  $r \in \mathbb{Z}$  (for R sector). The modes satisfy the NS or R extensions of the Virasoro algebra.

This analysis can be repeated for the scattering amplitudes [16]. A special choice of the gravitino modes  $\mathcal{G}_a(z, \bar{z})$  is the delta-function support at points  $z_a$

$$\mathcal{G}_a(z, \bar{z}) = \delta^{(2)}(z - z_a; \bar{z} - \bar{z}_a) \quad a = 1, \dots, 2g - 2. \quad (\text{A.24})$$

Using this choice, (A.23) can be written as

$$\mathcal{X}(z_a) = \delta(\beta(z_a)) T_F(z_a) = \delta(\beta(z_a)) \{\mathcal{Q}, \beta(z_a)\}, \quad (\text{A.25})$$

where we have used the fact that  $\{\mathcal{Q}, \beta(z)\} = T_F(z)$ . From the relation  $\frac{d\Theta(\beta(z))}{d\beta(z)} = \delta(\beta(z))$ , this expression can be formally written as

$$\mathcal{X}(z_a) = \{\mathcal{Q}, \Theta(\beta(z_a))\}. \quad (\text{A.26})$$

The locations of the support of the gravitino modes  $\mathcal{G}_a(z, \bar{z})$  are the locations of insertion of PCOs. Considering the fact that  $\Theta(\beta(z))$  can be identified with  $\xi(z)$ <sup>78</sup> [16], this form matches with (A.12), the FSM form of the picture-changing operator.

One is also interested in understanding the meaning of the picture-changing operation in the superstack of  $\mathcal{N} = 1$  Super-Riemann surfaces. It turns out that the correct interpretation of the picture-changing operation can be understood by a proper generalization of the concept of differential forms on ordinary manifolds to the so-called *pseudoforms* on supermanifolds. The picture-changing operation can then be understood as a natural operation on pseudoforms [51, 104]. We will explain this viewpoint in the next section.

### A.3 The Supergeometry Framework: Belopolsky Approach

In this section, we explain the geometrical meaning of the picture-changing operation. For the full exposition, we will refer to the original papers [51, 104] in which the relation between FSM's interpretation of the picture-changing operation and the operation of forms on supermanifolds has been elucidated.

In the theory of supermanifolds, the concept of differential forms is not enough to define the integration theory. The reason is that for odd coordinates  $\theta_i$  and  $\theta_j$ , the differential  $d\theta_i \wedge d\theta_j$  is even under  $i \longleftrightarrow j$ . This means that there is no differential form of top degree on a supermanifold. To properly define the integration on supermanifold, one has to generalize the concept of differential and top-degree forms.

---

<sup>78</sup>One way to determine this is as follows. By considering the integral representation of the delta function, we have

$$\beta(z)\delta(\beta(w)) \sim (z - w)\partial_w\beta(w)\delta(\beta(w)) = (z - w)\partial_w\Theta(w),$$

where we have used the fact that  $d\Theta(\beta(z)) = d\beta(z)\delta(\beta(z))$ , which implies that  $\partial_z\Theta(\beta(z)) = \partial_z\beta(z)\delta(\beta(z))$ . On the other hand, using the bosonization formula  $\delta(\beta(z)) \mapsto \exp(\phi(z))$  and (A.1), we find that

$$\beta(z)\exp(\phi(w)) \sim (z - w)\partial_w\xi(w).$$

Comparing the above equations, we can identify  $\xi(z) = \Theta(\beta(z))$ .



The simplest integrand for the integration over a supermanifold is the volume form. To define a volume form on a supermanifold, one tries to generalize the idea of integration over the bosonic manifolds. A volume form on an ordinary manifold is a non-degenerate nowhere-vanishing section of the canonical sheaf of the manifold. The idea can be generalized to supermanifold: a natural object that can be integrated over a supermanifold is a section of the canonical (or Berezinian) sheaf of the supermanifold. It turns out that defining such a section requires a generalization of the concept of differential form on an ordinary manifold to the so-called *pseudoforms* on supermanifolds [111]. If  $(x|\theta) \equiv (x^1 \cdots x^m | \theta^1 \cdots \theta^n)$  denotes the local coordinate on a supermanifold, then an  $(p|q)$ -pseudoform  $\Omega$  can be written locally as

$$\Omega = \Omega(x|\theta) dx^1 \cdots dx^r d\theta^1 \cdots d\theta^s \delta^{(n_1)}(d\theta^1) \cdots \delta^{(n_q)}(d\theta^q), \quad (\text{A.27})$$

where  $\delta^{(n_i)}(d\theta^i)$  denotes the  $n_i^{\text{th}}$  derivative of the delta function with respect to  $d\theta^i$ , and  $p \equiv r + s - \sum_{a=1}^q n_a$ . There are two degrees associated to a pseudo-form:  $p$  is called *the form degree*, corresponding to the usual degree of a differential form, and  $-q$ , minus of the number of fermionic dimension of the supermanifold, is called *the picture number*, which does not have an analog in the theory of differential forms on ordinary manifolds.

We would like to have an object that can be integrated over a supermanifold. On an ordinary manifold, a natural object that can be integrated over is a section of the canonical sheaf of the manifold<sup>79</sup>. These objects transform as a density on the manifold, i.e. under a coordinate transformations a density is multiplied by the Jacobian determinant of the coordinate transformation. Therefore, to be able to define the analogue of a density on a supermanifold, we need a supergeometric analog of the Jacobian determinant. This notion is called *Berezinian* of a coordinate transformation<sup>80</sup>. It thus turns out that proper objects that can be integrated over an  $(m|n)$ -dimensional supermanifold are sections of the canonical (or Berezinian) sheaf of the supermanifold, the so-called *integral forms* of top degree [111, 113]. These are special cases of pseudoforms (A.27) with the restriction  $s = 0$ ,  $n_i = 0$ ,  $r = m$  and  $q = n$ . A top-degree integral form  $\omega$  can thus locally be written as

$$\omega = \omega(x|\theta) dx^1 \cdots dx^m \delta(d\theta^1) \cdots \delta(d\theta^n). \quad (\text{A.28})$$

It has form number  $m - n$  and picture number  $-n$ . We note that  $\delta(dx) = dx$  has picture number zero since  $dx$  is a *fermionic* variable. The fundamental properties of such forms are that 1)  $\omega dx^i = \omega d\theta^j = 0$  for  $i = 1, \dots, m$  and  $j = 1, \dots, n$ , and 2)  $\delta(d\theta^a)\delta(d\theta^b) = -\delta(d\theta^b)\delta(d\theta^a)$  for  $a, b = 1, \dots, q$ <sup>81</sup>, therefore, top-degree integral forms are indeed the objects that can be integrated over a supermanifold of dimension  $(m|n)$ .

*What is the significance of integral forms in the superstring theory?* In superstring theory, the supermanifolds we are interested in is some integration cycle  $\Gamma$  inside the product  $\mathcal{M}_L \times \mathcal{M}_R$  of

<sup>79</sup>Here, we consider real manifold. For a complex manifold, a section of the canonical sheaf provides the holomorphic volume form.

<sup>80</sup>For more on the notion of density on a supermanifold see section 7 of chapter 4 of [112].

<sup>81</sup>We refer to section 3.3.2 of [10] for the details.

spaces  $\mathcal{M}_L$  and  $\mathcal{M}_R$  that parameterize the left- and right-moving sectors of relevant theories, as we have explained in Sections 6.2 and 6.3. The fermionic dimension of the superstack of genus- $g$  super-Riemann surfaces is equals to the space of meromorphic and multivalued quadratic  $\frac{3}{2}$ -superdifferentials associated to the variation of odd moduli [114]. The space of such differentials has dimension  $2g - 2$ . On the other hand, each vertex operator from the NS sector has picture number  $-1$ , i.e. it comes with a factor of  $\delta(\gamma)$ , and each vertex operator from the R sector has picture number  $-\frac{1}{2}$  since the spin field, i.e. the vacuum of the  $\beta\gamma$  system, has picture number  $-\frac{1}{2}$ . Therefore, the fermionic dimension of the superstack is  $2g - 2 + n_{\text{NS}} + \frac{1}{2}n_{\text{R}}$ , i.e. the superstring measure must be a top-degree integral form with picture number  $-(2g - 2 + n_{\text{NS}} + \frac{1}{2}n_{\text{R}})$ . The superstring measure associated to an arbitrary scattering process involving  $\mathbf{n} = (n_{\text{NS}}, n_{\text{R}})$  external states from either sectors, given by the path integral over the worldsheet superfields, naturally produces a top-degree integral form on  $\Gamma$  [11, 12, 42]

$$\langle\langle \mathcal{V}_1 \cdots \mathcal{V}_{\mathbf{n}} \rangle\rangle_{\mathcal{R}} \equiv \int \mathcal{D}(X, \mathcal{G}; B, C) \mathcal{V}_1 \cdots \mathcal{V}_{\mathbf{n}} e^{-S}, \quad (\text{A.29})$$

where  $\mathcal{R}$  is a fixed genus- $g$  surface with  $\mathbf{n} = (n_{\text{NS}}, n_{\text{R}})$  punctures,  $X, B, C$  are superfields of the matter and ghost superconformal field theories respectively,  $\mathcal{V}_i$ s are appropriate vertex operators associated to the external states, and  $S$  is the action describing the worldsheet theory (the gauge-fixed version of the RNS-superstring action or any other matter-ghost superconformal field theory with appropriate supersymmetry and central charges). Therefore,  $\langle\langle \mathcal{V}_1 \cdots \mathcal{V}_{\mathbf{n}} \rangle\rangle_{\mathcal{R}}$  can be integrated over  $\Gamma$  to give  $\mathcal{A}_{\text{String}}(1, \cdots, \mathbf{n})$ , the genus- $g$  stringy contribution to the scattering amplitude of  $\mathbf{n}$  arbitrary external states

$$\mathcal{A}_{\text{String}}(1, \cdots, \mathbf{n}) = \int_{\Gamma} \langle\langle \mathcal{V}_1 \cdots \mathcal{V}_{\mathbf{n}} \rangle\rangle_{\mathcal{R}}. \quad (\text{A.30})$$

After defining the pseudoforms, the next step is to define operations on them. Some of these operations may change the form degree or the picture number of pseudoform. In particular, one is interested to define an operator that changes the picture number of a generic pseudoform. To define such an operator, consider an odd vector field  $\widehat{V}$ , even tangent vectors  $\{v_1, \cdots, v_p\}$  and odd tangent vectors  $\{\widehat{v}_1, \cdots, \widehat{v}_q\}$ , which collectively denoted as  $\mathbf{v}$ , on a  $(m|n)$ -dimensional supermanifold. We can define the analog of the Lie derivative along  $\widehat{V}$  on a  $(p|q)$  form  $\omega$

$$[\mathcal{L}_{\widehat{V}} \omega](v_1, \cdots, v_p | \widehat{v}_1, \cdots, \widehat{v}_q) \equiv \widehat{V}^I \frac{\partial \omega(\mathbf{v})}{\partial x^I} + (-1)^{[A]} v_A^I \frac{\partial \widehat{V}^J}{\partial x^I} \frac{\partial \omega(\mathbf{v})}{\partial v_A^J}, \quad (\text{A.31})$$

where  $A = 1, \cdots, (p|q)$  and  $I, J = 1, \cdots, (m|n)$ , the dimension of the supermanifold.  $[\cdot]$  denotes the statistics of the quantity inside it. This operation does not change the form degree or the picture number of pseudoforms. Next, we define an operation that changes the form degree of a pseudoform but preserves its picture number. It can thus be interpreted as

the supergeometry analog of the de Rham differential. It is given by

$$[\mathbf{d}\omega](v_1, \dots, v_{p+1}|\widehat{v}_1, \dots, \widehat{v}_q) \equiv (-1)^p v_{p+1}^I \frac{\partial \omega(\mathbf{v})}{\partial x^I} + (-1)^{[I][A]} v_A^J \frac{\partial^2 \omega(\mathbf{v})}{\partial x^J \partial v_A^I}, \quad (\text{A.32})$$

$\mathbf{d}$  maps  $(p|q)$ -forms to  $(p+1|q)$ -forms. Finally, we define an operation that preserves the form number of pseudoforms but changes their picture number. It is given by

$$[\delta(\mathbf{i}_{\widehat{V}})\omega](v_1, \dots, v_p|\widehat{v}_1, \dots, \widehat{v}_{q-1}) \equiv (-1)^p \omega(v_1, \dots, v_p|\widehat{V}, \widehat{v}_1, \dots, \widehat{v}_{q-1}). \quad (\text{A.33})$$

$\delta(\mathbf{i}_{\widehat{V}})$  maps a  $(p|q)$  form to a  $(p|q-1)$  form. Note that  $\mathbf{i}_{\widehat{V}}$  is the usual interior product with the vector  $\widehat{V}$ . It might be tempting to identify  $\delta(\mathbf{i}_{\widehat{V}})$  with the picture-changing operation in the superstring theory. However, it turns out that the relevant operation is a little more subtle than this operation. One defines the so-called *picture-changing operator* on the space of pseudoforms as follows

$$\mathcal{X}_{\widehat{V}} \equiv \frac{1}{2} (\delta(\mathbf{i}_{\widehat{V}})\mathcal{L}_{\widehat{V}} + \mathcal{L}_{\widehat{V}}\delta(\mathbf{i}_{\widehat{V}})) = \{\mathbf{d}, \Theta(\mathbf{i}_{\widehat{V}})\}, \quad (\text{A.34})$$

where  $\Theta(x)$  is the step function. In the last identity, the fact that  $\mathcal{L}_{\widehat{V}} = \{\mathbf{d}, \mathbf{i}_{\widehat{V}}\}$  has been used. It is clear from the first equality that  $\mathcal{X}_{\widehat{V}}$  maps  $(p|q)$  forms to  $(p|q-1)$  forms, and indeed changes the picture number of a form on a supermanifold. Considering these operations, we can give the complex of  $(p|q)$ -forms on a supermanifold in figure 11.

From (A.34), and the fact that de Rham differential  $\mathbf{d}$  on the superstack can be interpreted as the superstring BRST operator<sup>82</sup>, and the fact that  $\Theta(\beta(z))$  can be identified with  $\xi(z)$ , it seems reasonable to guess that this operator is the same as the superstring picture-changing operation. This turns out to be the case [104]. We will explain this fact in the next section.

#### A.4 The Unified Perspective: From Supermanifolds to Superstrings

In this section, we explain the relation between the Belopolsky's picture-changing operator (A.34) and the superstring picture-changing operator. We will not explain the details of this relationship since it requires the explanation of the relation between semi-infinite forms defined over a suitable algebra and the ghost sector of the superstring theory. This section is just for the completion of discussion and we just quote the results from [51, 104] where the relation between FSM-VV approaches and the semi-infinite forms has been elucidated.

It turns out that the space of semi-infinite forms on the algebra of superconformal vector fields on the supercircle  $S^{1|1}$  can be identified with the ghost sector of superstring theory.

---

<sup>82</sup>The precise statement is as follows. Let  $\mathbf{d}$  denotes the de Rham differential acting on the superstack, and  $\Omega(\Psi_1, \dots, \Psi_n)$  is the off-shell superstring measure on the superstack for scattering of  $n$  external off-shell states  $\{\Psi_1, \dots, \Psi_n\}$ . Then,

$$\mathbf{d}\Omega_{p-1}(\Psi_1, \dots, \Psi_n) = \sum_{a=1}^n \Omega_p(\Psi_1, \dots, \Omega\Psi_a, \dots, \Psi_n).$$

$p$  and  $p-1$  denotes the form degree of the integral form  $\Omega$ . For more details see section 3 of [115].

$$\begin{array}{ccccccc}
0 & \xrightarrow{\mathbf{d}} & \Omega^{(0|0)} & \xrightarrow{\mathbf{d}} & \dots & \xrightarrow{\mathbf{d}} & \Omega^{(m|0)} & \xrightarrow{\mathbf{d}} & \Omega^{(m+1|0)} & \xrightarrow{\mathbf{d}} & \dots \\
\Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \\
\vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots \\
\Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \\
\vdots & \xrightarrow{\mathbf{d}} & \Omega^{(-1|s)} & \xrightarrow{\mathbf{d}} & \Omega^{(0|s)} & \xrightarrow{\mathbf{d}} & \dots & \xrightarrow{\mathbf{d}} & \Omega^{(m|s)} & \xrightarrow{\mathbf{d}} & \Omega^{(m+1|s)} & \xrightarrow{\mathbf{d}} & \dots \\
\Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \\
\vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots & \xrightarrow{\mathbf{d}} & \vdots \\
\Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \Xi \updownarrow \chi & & \\
\vdots & \xrightarrow{\mathbf{d}} & \Omega^{(-1|n)} & \xrightarrow{\mathbf{d}} & \Omega^{(0|n)} & \xrightarrow{\mathbf{d}} & \dots & \xrightarrow{\mathbf{d}} & \Omega^{(m|n)} & \xrightarrow{\mathbf{d}} & 0.
\end{array}$$

**Figure 11:** The complex of forms on a  $(m|n)$ -dimensional supermanifold. The first row is the complex of superforms. the intermediate complexes are complexes of pseudoforms with various picture numbers. The last row is the complex of integral forms, which as it is clear, there is a notion of top-degree integral forms. We have denoted two natural operations on forms on a supermanifold: 1) the picture-changing operation  $\chi$ , which acts from bottom to top and decreases the picture-number of forms by one, and 2) the picture-changing operation  $\Xi$ , which acts from top to bottom and increases the picture-number of forms by one. The space  $\Omega^{(m|n)}$  is the space of top-degree integral forms. The elements of this space can naturally be integrated over an  $(m|n)$ -dimensional supermanifold.

Let us denote the even and odd vector fields concentrated at  $z$  by  $l(z)$  and  $g(z)$ . Then, superconformal ghosts can be identified as follows

$$\begin{aligned}
b(z) &\mapsto \mathbf{i}_{l(z)}, & c(z) &\mapsto \mathbf{e}_{l^\vee(z)}, \\
\beta(z) &\mapsto \mathbf{i}_{g(z)}, & \gamma(z) &\mapsto \mathbf{e}_{g^\vee(z)}.
\end{aligned} \tag{A.35}$$

$v^\vee$  is the covector dual to the tangent vector  $v$ .  $\mathbf{e}_{v^\vee}$  is the operation of exterior product defined using  $v^\vee$ . For a  $(p|q)$ -form  $\omega$ , it is defined as follows

$$[\mathbf{e}_{v^\vee} \omega](v_1, \dots, v_{p+1} | \widehat{v}_1, \dots, \widehat{v}_q) \equiv (-1)^p \left[ v^\vee(v_{p+1}) - (-1)^{[v^\vee][A]} v^\vee(v_A) v_{p+1}^I \frac{\partial}{\partial v_A^I} \right] \omega. \tag{A.36}$$

Comparing (A.34) and (A.35) shows that we can identify  $\delta(\mathbf{i}_{\widehat{V}})$  with  $\delta(\beta(z))$ . The odd vector fields that appears in the FSM approach is thus  $g(z)$ , the odd generator of the super-Virasoro algebra. To complete the identification, we need to identify  $\mathcal{L}_{\widehat{V}}$ , the operation of Lie derivative along the odd vector field  $\widehat{V} = g(z)$ , with some conformal field in the ghost sector of superstring theory. It turns out that the Lie derivative  $\mathcal{L}_{g(z)}$  can be identified with  $T_F(z)$ . Therefore, the picture-changing operator (A.34) can be identified with the following conformal field

$$\mathcal{X}_{\widehat{V}} \mapsto \mathcal{X}_{g(z)} \equiv \frac{1}{2} (\delta(\beta(z)) T_F(z) + T_F(z) \delta(\beta(z))) = \{\mathcal{Q}, \Theta(\beta(z))\}. \tag{A.37}$$

Since  $\delta(\beta(z))$  is a conformal field with conformal dimension  $-\frac{3}{2}$ <sup>83</sup>,  $\mathcal{X}_{g(z)}$  has conformal dimension zero. The final identity is the FSM (A.12) and VV (A.26) expressions for the picture-changing operator. Note that this is a formal identification. The picture-changing operator (A.12) acting on a vertex operator *increases* the picture-number of the associated vertex operator. However, (A.34) acts on forms on a supermanifold and *decreases* their picture number.

In summary, we found that various interpretations of the picture-changing operation, the FSM's interpretation (i.e. an operation that changes the picture number of the vertex operator and maps the superstring BRST cohomology with a particular picture number to another superstring BRST cohomology with a different picture number), VV's interpretation (i.e. the combined operation of the absorption of a zero-mode of the  $\beta(z)$  field and the integration over the corresponding odd moduli of the superstack of  $\mathcal{N} = 1$  super-Riemann surfaces), and the Belopolsky's interpretation (i.e. an operation that acts on the space of pseudofields and changes their picture number) match. This concludes our discussion of the picture-changing operation.

## B Moduli Stacks of Stable Spin Curves

In this appendix, we review some details on the construction of the moduli stack of punctured spin curves over  $\text{Spec } \mathbb{Z}[1/2]$ , following [39, 44, 45]. In [44, 45], Jarvis discusses higher-spin structures and they are more complicated than  $r = 2$  case, i.e. ordinary spin structures. For convenience, we focus on  $r = 2$  since only ordinary spin structures are relevant for string theory. In fact, all four versions of spin structures in [44, 45], i.e. quasi-spin  $\text{QSpin}_{r,g}$ , spin  $\overline{\text{Spin}}_{r,g}$ , pure spin  $\text{Pure}_{r,g}$ , and finally the  $\overline{\mathcal{S}}_g^{1/r}$ , coincide when  $r = 2$ :

$$\text{QSpin}_{2,g} = \overline{\text{Spin}}_{2,g} = \text{Pure}_{2,g} = \overline{\mathcal{S}}_g^{1/2}.$$

### B.1 Definitions and Examples

We begin by recalling some definition. We begin with the definition of spin curve with and without punctures.

**Definition B.1 (Spin Curve).** A spin curve of genus  $g$  is a triple  $(\mathcal{C}, \mathcal{E}, b)$ , where  $\mathcal{C}$  is a stable curve of genus  $g$ ,  $\mathcal{E}$  is a rank one torsion-free sheaf, a coherent  $\mathcal{O}_X$ -module  $\mathcal{F}$  is called a torsion-free or relatively-torsion-free sheaf on a family of stable or semi-stable curves  $f : X \rightarrow S$  over a scheme  $S$  if it is a flat module over  $S$  in such a way that on  $\mathcal{C}$  of degree  $g - 1$  with a homomorphism  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}}$  which is an isomorphism on the smooth locus of  $\mathcal{C}$ .  $\omega_{\mathcal{C}}$  is the dualizing sheaf on  $\mathcal{C}$ .

**Definition B.2 (Family of Spin Curves).** A family of spin curves of genus  $g$  over the base scheme  $T$  is a triple  $(\mathcal{C}, \mathcal{E}, b)$ , where  $\mathcal{C} \rightarrow T$  is a relative stable curve,  $\mathcal{E}$  is a finite-presented

---

<sup>83</sup>Note that  $\beta(z)$  is a conformal field with conformal dimension  $+\frac{3}{2}$ .

sheaf on  $\mathcal{C}$  which is **flat over**  $T$ , with a morphism  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/T}$ , such that for every geometric point  $t$  of  $T$ ,  $(\mathcal{C}_t, \mathcal{E}_t, b_t)$  is a stable curve.  $\omega_{\mathcal{C}/T}$  is the relative dualizing sheaf on  $\mathcal{C}$ .

**Remark B.1.** Since relative stable curves are relative Gorenstein, dualizing complex  $\omega_{\mathcal{C}/T}^\bullet$  is quasi-isomorphic to a line bundle  $\omega_{\mathcal{C}/T}$ , we call it the relative dualizing sheaf.

**Remark B.2.** Let  $X$  be a topological space and  $S$  be a scheme. Suppose that  $X \rightarrow S$  is a locally finite-presented morphism between schemes. We call a finite-presented sheaf  $\mathcal{F}$  on  $X$  **relatively torsion-free** if  $\mathcal{F}$  is flat over  $S$  and  $\mathcal{F}_s$  is torsion-free on  $X_s$  for every geometric point  $s$  of  $S$ , i.e. the associated primes of the induced  $\mathcal{F}_s$  on each fiber  $X_s \equiv X \times_S \text{Spec } \mathbb{K}(s)$ , for some algebraically-closed field  $\mathbb{K}$  and all  $s \in S$ , do have height zero<sup>84</sup>. In Definition B.2,  $\mathcal{E}$  is relatively torsion-free.

A useful fact is the following:

**Lemma B.1.** Suppose that  $X \rightarrow T$  is a relative stable curve. If  $\mathcal{E} \in \text{QCoh}(\mathcal{O}_X)$ , where  $\text{QCoh}(\mathcal{O}_X)$  denotes the category of quasi-coherent sheaves on  $X$ , whose structure sheaf is  $\mathcal{O}_X$ <sup>85</sup>, is relatively torsion free, and  $\mathcal{F} \in \text{QCoh}(\mathcal{O}_X)$  is locally-free of finite rank, then  $\text{RHom}^\bullet(\mathcal{E}, \mathcal{F})$ <sup>86</sup>, is quasi-isomorphic to  $\text{Hom}(\mathcal{E}, \mathcal{F})$ <sup>87</sup>, which is relatively torsion free and its construction commutes with arbitrary base change.

*Proof.* There is a canonical morphism  $\text{Hom}(\mathcal{E}, \mathcal{F}) \rightarrow \text{RHom}^\bullet(\mathcal{E}, \mathcal{F})$  defined by truncation, and it's a quasi-isomorphism if it is locally on  $X$ , so we can assume that both  $X$  and  $T$  are affine. Our strategy is to prove it first for Noetherian  $T$  then show that it remains true after base change.

**When  $T$  is a spectrum of a field**, i.e.  $T = \text{Spec } \mathbb{K}$  for some field  $\mathbb{K}$ , then  $X$  is a stable curve over  $\mathbb{K}$ . Since  $X$  is Gorenstein of dimension 1 and  $\mathcal{F}$  is locally free, injective dimension of  $\mathcal{F}$  is 1, hence  $\text{RHom}^\bullet(\mathcal{E}, \mathcal{F})$  is concentrated in degree 0 and 1. Moreover,  $\mathcal{E}$  is torsion free thus torsionless, so locally  $\mathcal{E}$  can be embedded into a free sheaf

$$\mathcal{E}|_U \hookrightarrow \mathcal{O}_U^{\oplus n},$$

which implies that there is a surjective morphism

$$\text{Ext}^1(\mathcal{O}_U^{\oplus n}, \mathcal{F}|_U) \twoheadrightarrow \text{Ext}^1(\mathcal{E}|_U, \mathcal{F}|_U).$$

<sup>84</sup>Let  $M$  be an  $R$ -module over a commutative ring  $R$ . An associated prime ideal of  $M$  is a prime ideal  $I_p$  that is the annihilator of a nonzero element  $\mathfrak{m} \in M$ . The height of a prime ideal is the maximal length of a chain of prime ideals contained in  $I_p$ , which is also called the Krull dimension of the localization  $R_{I_p}$ .

<sup>85</sup>A local-ringed space  $(X, \mathcal{O}_X)$  is a topological space  $X$  together with a sheaf of rings  $\mathcal{O}_X$ , called the structure sheaf, on  $X$ .

<sup>86</sup>Here,  $\text{RHom}^\bullet$  denotes the derived Hom functor and the  $\text{RHom}$  means it is an object in the derived category of sheaves, not just literally taking the  $\text{Ext}^\bullet$ .

<sup>87</sup>Here,  $\text{Hom}$  denotes the Hom functor and  $\text{Hom}$  denotes the sheaf of Homs, not just the global Hom between two sheaves.

Hence  $\text{Ext}^1(\mathcal{E}|_U, \mathcal{F}|_U) = 0$  and

$$\underline{\text{Hom}}(\mathcal{E}, \mathcal{F}) \cong \underline{\text{RHom}}^\bullet(\mathcal{E}, \mathcal{F}).$$

Moreover  $\underline{\text{Hom}}(\mathcal{E}, \mathcal{F})$  is torsion free (in fact reflexive).

**For general Noetherian  $T$ ,** take any geometric point  $t$  of  $T$ , denote the natural morphism  $X_t \rightarrow X$  by  $i_t$ . For any integer  $N > 1$ , consider a complex of finite free  $\mathcal{O}_X$  modules

$$\cdots \rightarrow 0 \rightarrow \mathcal{E}^{-N} \rightarrow \mathcal{E}^{-N+1} \rightarrow \cdots \rightarrow \mathcal{E}^0$$

with trivial cohomologies at degree  $\{-N+1, \dots, -1\}$  and  $H^0$  being  $\mathcal{E}$ . Recall that we assume that  $X$  is affine, hence there exists such complex. Note that there is a distinguished triangle in the derived category:

$$\mathcal{K}^{-N}[N] \longrightarrow \mathcal{E}^\bullet \longrightarrow \mathcal{E} \xrightarrow{+1},$$

where  $\mathcal{K}^{-N}$  is the kernel of  $\mathcal{E}^{-N} \rightarrow \mathcal{E}^{-N+1}$ . Hence we have a quasi-isomorphism

$$\tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}, \mathcal{F}) \cong \tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}^\bullet, \mathcal{F}), \quad (\text{B.1})$$

where  $\tau_{\leq N-1}$  is the truncation functor. Moreover this quasi-isomorphism is stable under arbitrary base change  $T' \rightarrow T$ , since derived base change of  $\mathcal{K}^{-N}[N]$  has nonzero cohomologies only in degree below  $-N+1$ . Now take the base change  $i_t : X_t \rightarrow X$ , we have

$$Li_t^* \underline{\text{RHom}}^\bullet(\mathcal{E}^\bullet, \mathcal{F}) \cong \underline{\text{RHom}}^\bullet(i_t^* \mathcal{E}^\bullet, i_t^* \mathcal{F}), \quad (\text{B.2})$$

since both  $\mathcal{E}^\bullet$ ,  $\mathcal{F}$ , and  $\underline{\text{RHom}}^\bullet(\mathcal{E}^\bullet, \mathcal{F})$  are represented by complex of free modules. The case of  $T$  being  $\text{Spec } \mathbb{K}(t)$  implies that  $\tau_{\leq N-1} \underline{\text{RHom}}^\bullet(i_t^* \mathcal{E}^\bullet, i_t^* \mathcal{F})$  is concentrated in degree zero and torsion free. Since  $t$  can be arbitrary, we see that the truncation of perfect complex,  $\tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}^\bullet, \mathcal{F})$ , is again perfect, concentrated in degree zero, and whose degree zero cohomology is relatively torsion free. Hence the same is true for  $\tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}, \mathcal{F})$ . Take  $N \rightarrow \infty$  and we conclude the proof for arbitrary Noetherian  $T$ .

**For general scheme  $T$ ,** Note that we can take Noetherian approximation and assume that there is a Noetherian affine  $X_0 \rightarrow T_0$  with  $\mathcal{E}_0$  and  $\mathcal{F}_0$  satisfying conditions of this lemma, such that  $X$  and  $\mathcal{E}$  and  $\mathcal{F}$  are pull-back of a morphism  $\phi : T \rightarrow T_0$ . We have shown a quasi-isomorphism

$$\tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}_0, \mathcal{F}_0) \cong \tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}_0^\bullet, \mathcal{F}_0)$$

in step 2 which is stable under arbitrary base change. We also know that the perfect complex  $i_t^* \underline{\text{RHom}}^\bullet(\mathcal{E}_0, \mathcal{F}_0)$  has non-trivial cohomologies only in degree 0 and  $N$ , for arbitrary geometric point  $t$ , hence  $\underline{\text{Ext}}^N(\mathcal{E}_0, \mathcal{F}_0)$  is also flat over  $T_0$ . Hence we see that

$$\tau_{\leq N-1} \underline{\text{RHom}}^\bullet(L\phi^* \mathcal{E}_0^\bullet, L\phi^* \mathcal{F}_0) \cong L\phi^* \tau_{\leq N-1} \underline{\text{RHom}}^\bullet(\mathcal{E}_0^\bullet, \mathcal{F}_0)$$

is concentrated in degree zero and whose degree zero cohomology is relatively torsion free. Take  $N \rightarrow \infty$  and we conclude the proof for arbitrary scheme  $T$ .  $\blacksquare$

**Definition B.3 (Isomorphism of Spin Curves).** An isomorphism between two families of spin curves over  $T$  denoted by  $(\mathcal{C}, \mathcal{E}, b)$  and  $(\mathcal{C}', \mathcal{E}', b')$  is a pair  $(\phi, \psi)$ , where  $\phi : \mathcal{C}' \rightarrow \mathcal{C}$  is an isomorphism between spin curves,  $\psi$  is an isomorphism between  $\mathcal{E}'$  and  $\phi^*\mathcal{E}$  which is compatible with  $b$ .

**Example B.1.** For any family of spin curve such that 2 is invertible in the base scheme, there always exists an automorphism  $(\mathbb{I}, -1)$ , for the identity map  $\mathbb{I}$ .

**Definition B.4 (Stack of Spin Curves).** Fix a scheme  $S \equiv \text{Spec } \mathbb{Z}[1/2]$  and an integer  $g \geq 2$ , define the following prestack in  $(\text{Sch}_S)_{\text{ét}}$ <sup>88</sup>

$$(T \in \text{Sch}_S) \mapsto \begin{cases} \mathbf{Obj} : \text{Families of Spin curves of genus-}g \text{ over } T, \\ \mathbf{Mor} : \text{Isomorphisms between families of spin curves.} \end{cases}$$

It's in fact a stack: every stable curve  $\mathcal{C} \rightarrow S$  is canonically polarized by dualizing sheaf  $\omega_{\mathcal{C}/S}$ . We denote it by  $\overline{\mathcal{S}}_g$ .

We also need the following generalization:

**Definition B.5 (Marked Spin Curve).** A family of punctured spin curves of genus  $g$  over the base scheme  $T$ , of puncturing type  $\mathbf{m} \equiv (m_1, \dots, m_n)$  is a quadruple  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$ , where  $\mathcal{C} \rightarrow T$  is a relative semi-stable curve,  $\mathcal{D} \equiv (\mathcal{D}_1, \dots, \mathcal{D}_n)$  are  $n$  sections of  $\mathcal{C} \rightarrow T$  with images lying in the smooth locus and making  $(\mathcal{C}, \mathcal{D})$  into an  $n$ -punctured stable curve,  $\mathcal{E}$  is a rank-1 relatively torsion-free sheaf on  $\mathcal{C}$ , with a morphism  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i)$ <sup>89</sup>, such that for every geometric point  $t$  of  $T$

1.  $\deg \mathcal{E}_t = g - 1 + \frac{1}{2} \sum_{i=1}^n m_i$ ,
2.  $b_t$  is isomorphism on the smooth locus of  $\mathcal{C}_t$ .

It turns out that there is an equivalent definition which is sometimes more convenient:

**Lemma B.2.** Suppose that  $(\mathcal{C}, \mathcal{D})/T$  is a relative  $n$ -punctured stable curve.  $\mathcal{E}$  is a rank-1 relatively torsion free sheaf on  $\mathcal{C}$ , of degree  $g - 1 + \frac{1}{2} \sum_i m_i$ , then there is a one-to-one correspondence between

1. Homomorphisms  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i)$  such that  $b_t$  is isomorphism on the smooth locus of  $\mathcal{C}_t$  for any geometric point  $t$  of  $T$ ;
2. Isomorphisms  $a : \mathcal{E} \rightarrow \underline{\text{Hom}}(\mathcal{E}, \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i))$ .

<sup>88</sup> $(\text{Sch}_S)_{\text{ét}}$  denotes the category of schemes over  $S$  endowed with the étal topology.

<sup>89</sup> $\omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i)$  means that sections of the canonical sheaf have poles of order  $m_i$  along the divisors  $\mathcal{D}_i$  defined by the puncture  $\mathbf{p}_i$ . For more on this see [the definition of the notation  \$\mathcal{S}\_{g,n}^{\mathbf{m}}\$](#)  in the Introduction.



*Proof.* First of all, according to Lemma B.1,  $\underline{\text{Hom}}(\mathcal{E}, \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i))$  is a rank-1 relatively torsion-free sheaf on  $\mathcal{C}$ , and we have

$$\deg \underline{\text{Hom}} \left( \mathcal{E}_t, \omega_{\mathcal{C}_t/t} \left( \sum_i m_i \mathcal{D}_i \right) \right) = \deg \omega_{\mathcal{C}_t/t} \left( \sum_i m_i \mathcal{D}_i \right) - \deg \mathcal{E}_t = g-1 + \frac{1}{2} \sum_{i=1}^n m_i = \deg \mathcal{E}_t.$$

By adjunction, there is a one to one correspondence between

1. Homomorphisms  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i)$  such that  $b_t$  is isomorphism on the smooth locus of  $\mathcal{C}_t$  for any geometric point  $t$  of  $T$ ;
2. Homomorphisms  $c : \mathcal{E} \rightarrow \underline{\text{Hom}}(\mathcal{E}, \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i))$  such that  $c_t$  is isomorphism on the smooth locus of  $\mathcal{C}_t$  for any geometric point  $t$  of  $T$ .

Since  $c_t$  is an isomorphism on the smooth locus of  $\mathcal{C}_t$ , we see that  $\ker c_t = 0$ , this follows from the fact that a subsheaf of torsion free sheaf is torsion free, hence if  $\ker c_t$  is not empty,  $\text{Supp}(\ker c_t)$  has dimension 1 so it must have nontrivial intersection with the smooth locus, which is absurd. The equation  $\deg \underline{\text{Hom}}(\mathcal{E}_t, \omega_{\mathcal{C}_t/t}(\sum_i m_i \mathcal{D}_i)) = \deg \mathcal{E}_t$  implies that any homomorphism  $c_t$  which is isomorphism on smooth locus is actually an isomorphism, otherwise  $\text{coker}(c_t) \neq 0$  and

$$\begin{aligned} \deg \underline{\text{Hom}} \left( \mathcal{E}_t, \omega_{\mathcal{C}_t/t} \left( \sum_i m_i \mathcal{D}_i \right) \right) &= \chi \left( \underline{\text{Hom}} \left( \mathcal{E}_t, \omega_{\mathcal{C}_t/t} \left( \sum_i m_i \mathcal{D}_i \right) \right) \right) - \chi(\mathcal{O}_{\mathcal{C}_t}) \\ &= \chi(\mathcal{E}_t) - \chi(\mathcal{O}_{\mathcal{C}_t}) + \dim_{k(t)} \text{coker}(c_t) > \deg \mathcal{E}_t, \end{aligned}$$

where  $\chi$  denotes the Euler number. This is a contradiction. Hence  $c$  is forced to be an isomorphism.  $\blacksquare$

**Definition B.6 (Stack of Punctured Spin Curves).** Fix a scheme  $S \equiv \text{Spec } \mathbb{Z}[1/2]$ , a pair of natural numbers  $(g, n)$  such that  $2g - 2 + n > 0$ , and an  $n$ -tuple  $\mathbf{m} = (m_1, m_2, \dots, m_n)$  such that  $\sum_i m_i$  is even, define the following prestack in  $(\text{Sch}_S)_{\text{ét}}$ :

$$(T \in \text{Sch}_S) \mapsto \begin{cases} \text{Obj} : \text{Families of spin curves of genus } g \text{ over } T \text{ of puncturing type } \mathbf{m}, \\ \text{Mor} : \text{Isomorphisms between families of maked spin curves.} \end{cases}$$

It's in fact a stack: every marked stable curve  $\mathcal{C} \rightarrow S$  is canonically polarized by  $\omega_{\mathcal{C}/S}(\sum_i D_i)$ . We denote it by  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$ .

**Remark B.3.** There is an obvious relation between these  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$ : shifting  $\mathcal{E}$  to  $\mathcal{E} \otimes \mathcal{O}_{\mathcal{C}}(\mathcal{D}_i)$ , we get an isomorphism between stacks:

$$\overline{\mathcal{S}}_{g,n}^{\mathbf{m}} \cong \overline{\mathcal{S}}_{g,n}^{\mathbf{m}'},$$

where  $\mathbf{m}' \equiv \mathbf{m} + (0, \dots, 0, 2, 0, \dots, 0)$ .

**Remark B.4.** There is an obvious morphism from  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  to  $\overline{\mathcal{M}}_{g,n}$  by forgetting the spin structure.

The main result of this appendix is the following

**Theorem B.1.** *Suppose that  $2g - 2 + n > 0$ . Then,  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  is a smooth proper Deligne-Mumford stack over  $\mathrm{Spec} \mathbb{Z}[1/2]$  and contains an open dense substack  $\mathcal{S}_{g,n}^{\mathbf{m}}$ .  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  is flat and quasi-finite over  $\overline{\mathcal{M}}_{g,n}$ . Moreover the number of connected components of geometric fibers of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}} \rightarrow \mathrm{Spec} \mathbb{Z}[1/2]$  is  $\gcd(2, m_1, m_2, \dots, m_n)$ .*

In the following sections, we discuss the proof of this Theorem. The different parts of the Theorem is proven in Corollary B.2, Corollary B.3, and Proposition B.9.

## B.2 Rank-One Torsion-Free Sheaves

Recall the classical fact of local classification of torsion-free sheaves on semistable curves, due to Faltings:

**Theorem B.2 (Theorem 3.5 of [116], The Rank-One Case).** *Let  $R$  be a complete local Noetherian ring with the maximal ideal  $\mathfrak{m}$ , then any rank-one relative torsion-free sheaf which is not free over the ring  $A = R[[x, y]]/(xy - \pi)$ ,  $\pi \in \mathfrak{m}$  is isomorphic to  $E(p, q)$ , for  $p, q \in \mathfrak{m}$  and  $pq = \pi$ , defined as the image of*

$$\alpha = \begin{bmatrix} x & q \\ p & y \end{bmatrix} : A^{\oplus 2} \rightarrow A^{\oplus 2}.$$

Moreover  $E(p, q) \cong E(p', q')$  if and only if  $\exists u \in R^\times$  such that  $p' = up$ ,  $q' = q/u$ .

**Remark B.5.** *This theorem is also true for  $R$  being a Henselian local  $G$ -ring, by Artin's Approximation Theorem.*

In fact the module  $E(p, q)$  in the Theorem B.2 has a nice geometric description. Consider the projective scheme

$$X = \mathrm{Proj} A[z, w]/(xz - pw, qz - yw). \quad (\text{B.3})$$

We can check that  $X$  is flat over  $R$ : on the chart  $D_+(w)$ ,  $X$  is given by

$$\mathrm{Spec} A[T]/(qT - y, p - xT) = \mathrm{Spec} R[x, y, T]^\wedge/(qT - y, p - xT).$$

Since  $\{xT, y\}$  is a regular sequence on the special fiber,  $R[x, y, T]/(qT - y, p - xT)$  is flat over  $R$  by local criterion for flatness, so its completion is also flat over  $R$ . Note that the special fiber of  $X \rightarrow \mathrm{Spec} R$  is a  $\mathbb{P}^1$ -arc connecting closed points of  $\mathrm{Spec} \mathbb{K}[[x]]$  and  $\mathrm{Spec} \mathbb{K}[[y]]$  with nodal singularity, where  $\mathbb{K}$  is the residue field of  $R$ , while  $f : X \rightarrow \mathrm{Spec} A$  on the open locus (possibly empty)  $\mathrm{Spec} R - V(p, q) = D(p) \cup D(q)$  is an isomorphism.

Now we can push forward the line bundle  $\mathcal{O}(1)$  on  $X$  to  $\mathrm{Spec} A$ , which can be computed explicitly:

$$f_*\mathcal{O}(1) = \ker \left( A[T]/(qT - y, p - xT) \oplus A[S]/(q - yS, pS - x) \rightarrow A[T, T^{-1}]/(qT - y, p - xT) \right),$$

where the map is given by embedding the first direct summand naturally and send  $S$  to  $T^{-1}$  meanwhile multiply the second direct summand by  $T$ . Notice that  $A[T]/(qT - y, p - xT)$  has a free resolution as  $A[T]$ -module:

$$\cdots \longrightarrow A[T]^{\oplus 2} \xrightarrow{\alpha} A[T]^{\oplus 2} \xrightarrow{\beta} A[T]^{\oplus 2} \xrightarrow{\alpha} A[T]^{\oplus 2} \xrightarrow{\beta_1} A[T],$$

where

$$\alpha = \begin{bmatrix} x & q \\ p & y \end{bmatrix}, \quad \beta = \begin{bmatrix} y & -q \\ -p & x \end{bmatrix}, \quad \beta_1 = \begin{bmatrix} qT - y & p - xT \end{bmatrix}.$$

This is a resolution since it is a resolution modulo  $\mathfrak{m}$  and  $A[T]$  is flat over  $R$ . For the same reason,  $A[S]/(q - yS, pS - x)$  has a free resolution as  $A[S]$ -module:

$$\cdots \longrightarrow A[S]^{\oplus 2} \xrightarrow{\alpha} A[S]^{\oplus 2} \xrightarrow{\beta} A[S]^{\oplus 2} \xrightarrow{\alpha} A[S]^{\oplus 2} \xrightarrow{\beta_2} A[S],$$

where

$$\beta_2 = \begin{bmatrix} q - yS & pS - x \end{bmatrix},$$

and  $A[T, T^{-1}]/(qT - y, p - xT)$  has a free resolution as  $A[T, T^{-1}]$ -module:

$$\cdots \longrightarrow A[T, T^{-1}]^{\oplus 2} \xrightarrow{\beta} A[T, T^{-1}]^{\oplus 2} \xrightarrow{\alpha} A[T, T^{-1}]^{\oplus 2} \xrightarrow{\beta_3} A[T, T^{-1}],$$

where

$$\beta_3 = \begin{bmatrix} qT - y & p - xT \end{bmatrix},$$

It's easy to see that the original map lifts to a map between complexes:

$$\begin{array}{ccccccc} \cdots & \longrightarrow & A[T]^{\oplus 2} \oplus A[S]^{\oplus 2} & \xrightarrow{\alpha \oplus \alpha} & A[T]^{\oplus 2} \oplus A[S]^{\oplus 2} & \xrightarrow{\beta_1 \oplus \beta_2} & A[T] \oplus A[S] \\ & & \downarrow (\text{Id}, \text{Id}) & & \downarrow (\text{Id}, \text{Id}) & & \downarrow (\text{Id}, T) \\ \cdots & \longrightarrow & A[T, T^{-1}]^{\oplus 2} & \xrightarrow{\alpha} & A[T, T^{-1}]^{\oplus 2} & \xrightarrow{\beta_3} & A[T, T^{-1}]. \end{array}$$

Obviously, vertical arrows are surjective, so we end up with a chain of kernels:

$$\cdots \longrightarrow A^{\oplus 2} \xrightarrow{\beta} A^{\oplus 2} \xrightarrow{\alpha} A^{\oplus 2} \xrightarrow{\beta} A^{\oplus 2}.$$

It is easy to see that this chain is exact except for the terminal place, and it gives rise to an exact sequence

$$A^{\oplus 2} \xrightarrow{\beta} A^{\oplus 2} \longrightarrow f_*\mathcal{O}(1) \longrightarrow 0.$$

As a result,  $f_*\mathcal{O}(1) \cong \text{im}(\alpha) = E(p, q)$ .

**Remark B.6.** The identification  $f_*\mathcal{O}(1) \cong \text{coker}(\beta)$  implies the following isomorphism of  $A$ -algebras:

$$\bigoplus_{n \geq 0} \text{Sym}^n(E(p, q)) \cong \bigoplus_{n \geq 0} \text{Sym}^n(A^{\oplus 2})/\text{relations} = A[z, w]/(xz - pw, qz - yw),$$

via identifying two generators of  $A^{\oplus 2}$  with  $z, w$ , hence the relations are exactly  $xz - pw = 0$  and  $qz - yw = 0$ . Consequently

$$X \cong \mathbb{P}(E(p, q)).$$

### B.3 Cornalba's Definition of Spin Curves

Above local characterization of rank one torsion free sheaves indicates that the rank one torsion free sheaf defining the spin structure should be a pushforward of a locally free sheaf on a certain "blow-up" of the family of curves. Indeed, suppose that  $\pi : \mathcal{C} \rightarrow T$  is a family of spin curves with puncturing divisors  $\{\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_n\}$  and spin structure  $\mathcal{E}$ , then  $\tilde{\pi} : \tilde{\mathcal{C}} := \mathbb{P}(\mathcal{E}) \rightarrow T$  is a flat family of nodal curves (not necessarily stable) over  $T$ , together with a distinct line bundle  $\mathcal{L} = \mathcal{O}(1)$  such that

$$f_*\mathcal{L} \cong \mathcal{E},$$

where  $f : \tilde{\mathcal{C}} \rightarrow \mathcal{C}$  is the natural projection. Moreover the structure map  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/T}(\sum_i m_i \mathcal{D}_i)$  induces a homomorphism by adjunction

$$\alpha : \mathcal{L}^{\otimes 2} \rightarrow \omega_{\tilde{\mathcal{C}}/T} \left( \sum_i m_i \mathcal{D}_i \right),$$

which is an isomorphism outside of the locus of exceptional curves.

In [39], Cornalba defines a spin curve over  $T$  as a *semi-stable* curve  $\mathcal{C}$  over  $T$  with a line bundle  $\mathcal{L}$  and a homomorphism  $\alpha : \mathcal{L}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/T}$  such that for all geometric point  $t$  of  $T$ ,  $b$  is an isomorphism outside of the rational components of  $\mathcal{C}_t$ , and  $\mathcal{L}_t$  has degree one on each rational component. An isomorphism between spin curves  $\mathcal{C}, \mathcal{C}'$  is defined to be a pair of isomorphisms  $\sigma : \mathcal{C} \rightarrow \mathcal{C}'$  and  $\varphi : \sigma^*\mathcal{L}' \rightarrow \mathcal{L}$  which is compatible with the canonical isomorphism of  $\sigma^*\omega_{\mathcal{C}'/T} \rightarrow \omega_{\mathcal{C}/T}$ . In this way we can define the stack of spin curves over a base scheme  $S$  by

$$(T \in \text{Sch}_S) \mapsto \begin{cases} \mathbf{Obj} : \text{Spin curves of genus-}g \text{ in Cornalba's sense over } T, \\ \mathbf{Mor} : \text{Isomorphisms between families of spin curves.} \end{cases}$$

It is in fact a stack: every spin curve  $\mathcal{C}$  over  $T$  with spinor bundle  $\mathcal{L}$  is canonically polarized by  $\omega_{\mathcal{C}/S} \otimes \mathcal{L}$ . We denote it by  $\overline{\mathcal{S}}'_g$ . By what we discussed above, there is a canonical morphism  $p : \overline{\mathcal{S}}_g \rightarrow \overline{\mathcal{S}}'_g$  defined by

$$(\mathcal{C}, \mathcal{E}) \mapsto (\mathbb{P}(\mathcal{E}), \mathcal{O}(1)).$$

It is easy to see from the above *blowing-up* construction that over an algebraic closed field  $\mathbb{K}$ ,  $p$  is an equivalence between categories. In fact,  $p$  is an isomorphism between stacks, proven below.

**Lemma B.3.**  $p : \overline{\mathcal{S}}_g \rightarrow \overline{\mathcal{S}}'_g$  is fully faithful.

*Proof.* Fix a scheme  $T$  and let  $(\mathcal{C}_1, \mathcal{E}_1)$  and  $(\mathcal{C}_2, \mathcal{E}_2)$  be two objects in  $\overline{\mathcal{S}}_g(T)$ , and let  $(\tilde{\mathcal{C}}_i, \mathcal{L}_i)$  be  $(\mathbb{P}(\mathcal{E}_i), \mathcal{O}(1))$ . We ought to show that  $\text{Isom}_T((\mathcal{C}_1, \mathcal{E}_1), (\mathcal{C}_2, \mathcal{E}_2)) \rightarrow \text{Isom}_T((\tilde{\mathcal{C}}_1, \mathcal{L}_1), (\tilde{\mathcal{C}}_2, \mathcal{L}_2))$  is bijective.

**Injectivity:** Suppose that there are two isomorphisms  $\phi, \psi : (\mathcal{C}_1, \mathcal{E}_1) \rightarrow (\mathcal{C}_2, \mathcal{E}_2)$  which give rise to the same isomorphism  $\varphi : (\tilde{\mathcal{C}}_1, \mathcal{L}_1) \rightarrow (\tilde{\mathcal{C}}_2, \mathcal{L}_2)$ , then the  $\phi$  and  $\psi$  induces the same map for underlying topological space, because  $\tilde{\mathcal{C}}_1 \rightarrow \mathcal{C}_1$  is surjective. Furthermore,  $\phi^\# : \phi^{-1}\mathcal{O}_{\mathcal{C}_2} \rightarrow \mathcal{O}_{\mathcal{C}_1}$  is the same as  $\pi_{1,*}(\varphi^\#) : \pi_{1,*}\varphi^{-1}\mathcal{O}_{\tilde{\mathcal{C}}_2} \rightarrow \pi_{1,*}\mathcal{O}_{\tilde{\mathcal{C}}_1}$ , and so is  $\psi^\#$ , thus  $\phi^\# = \psi^\#$ , i.e.  $\phi$  and  $\psi$  are the same for the underlying stable curve. Similar argument for spin structure shows that  $\phi$  agrees with  $\psi$  on spinor sheaves, hence  $\phi = \psi$ .

**Surjectivity:** Suppose that there is an isomorphism  $\varphi : (\tilde{\mathcal{C}}_1, \mathcal{L}_1) \rightarrow (\tilde{\mathcal{C}}_2, \mathcal{L}_2)$ . Then exceptional curves of  $\tilde{\mathcal{C}}_1$  are identified with exceptional curves of  $\tilde{\mathcal{C}}_2$  under  $\varphi$ , so they are contracted to single points by the map  $\pi_2 \circ \varphi$ , thus there is a continuous map  $\psi$  between topological spaces  $\mathcal{C}_1$  and  $\mathcal{C}_2$  (because  $\pi_1$  is submersive) such that  $\psi \circ \pi_1 = \pi_2 \circ \varphi$ , hence  $\psi$  is a homeomorphism. It follows that  $\psi^{-1}\mathcal{O}_{\mathcal{C}_2} = \pi_{1,*}\varphi^{-1}\mathcal{O}_{\tilde{\mathcal{C}}_2}$ . Define  $\psi^\# : \psi^{-1}\mathcal{O}_{\mathcal{C}_2} \rightarrow \mathcal{O}_{\mathcal{C}_1}$  by  $\pi_{1,*}(\varphi^\#)$ , which is an isomorphism between sheaves of rings and also  $\mathcal{O}_T$ -linear, thus  $\psi$  is a isomorphism between  $T$ -schemes. Finally, define the isomorphism between spinor sheaves by  $\pi_{1,*}$  of the isomorphism between  $\varphi^*\mathcal{L}_2$  and  $\mathcal{L}_1$ . ■

**Lemma B.4.** The diagonal of  $\overline{\mathcal{S}}'_g$  is represented by separated and locally quasi-finite scheme.

*Proof.* Since spin curves are projective,  $\text{Isom}_T(\mathcal{C}_1, \mathcal{C}_2)$  is a locally closed subscheme of  $\text{Hilb}_{\mathcal{C}_1 \times_T \mathcal{C}_2 / T}$ , which is separated and locally finite type over  $T$ . For each geometric point  $t$  of  $T$ ,  $\text{Isom}_t(\mathcal{C}_{1,t}, \mathcal{C}_{2,t})$  has finite many points (Lemma 2.2 of [39]), so  $\text{Isom}_T(\mathcal{C}_1, \mathcal{C}_2)$  is locally quasi-finite. ■

**Proposition B.1.**  $p : \overline{\mathcal{S}}_g \rightarrow \overline{\mathcal{S}}'_g$  is isomorphism.

*Proof.* By Lemma B.3 and B.4, for every scheme  $T$  and a  $T$ -point  $g : T \rightarrow \overline{\mathcal{S}}'_g$ ,  $p^{-1}T$  is represented by an algebraic space, which is separated and locally finite type over  $T$ . For a geometric point  $s$  of  $\overline{\mathcal{S}}'_g$ ,  $p^{-1}(s)$  has only one point, thus  $p$  is universally injective. Furthermore, comparing Proposition B.7 and the construction of universal deformation in [39], we see that  $p$  induces isomorphism between universal deformation basis, thus  $p$  is étale. Applying Zariski's Main Theorem [117] to the separated, universally injective, and étale morphism  $p$ , we see that  $p$  is an isomorphism. ■

#### B.4 Stack of Prespin Structures and Representability of $\overline{\mathcal{S}}_{g,n}^m$

In this section, we describe the stack of prespin curves. We begin by giving the definition of the relevant concepts.

**Definition B.7 (Prespin Curve).** A family of prespin curves of genus  $g$  with  $n$  punctures over the base scheme  $T$ , of puncturing number  $N \in 2\mathbb{Z}$ , is a triple  $(\mathcal{C}, \mathcal{D}, \mathcal{E})$ , where  $(\mathcal{C}, \mathcal{D}) \rightarrow T$

is an  $n$ -punctured stable curve,  $\mathcal{E}$  is a rank-1 relatively torsion-free sheaf on  $C$ , such that for every geometric point  $t$  of  $T$

$$\deg \mathcal{E}_t = g - 1 + \frac{1}{2}N.$$

**Definition B.8 (Stack of Prespin Curves).** Fix a scheme  $S \equiv \operatorname{Spec} \mathbb{Z}[1/2]$ , a pair of natural numbers  $(g, n)$  such that  $2g - 2 + n > 0$ , define the following prestack in  $(\operatorname{Sch}_S)_{\text{ét}}$ :

$$(T \in \operatorname{Sch}_S) \mapsto \begin{cases} \text{Obj} : \text{Families of } (g, n) \text{ prespin curves over } T \text{ of marking number } N, \\ \text{Mor} : \text{Isomorphisms between families of prespin curves}/(\mathbb{I}, -1). \end{cases}$$

Note that we do not distinguish two isomorphisms if they are related by  $(\mathbb{I}, -1)$ . We denote the stackification of the above prestack by  $\mathcal{PS}_{g,n}^N$ . There is a morphism  $\pi : \mathcal{PS}_{g,n}^N \rightarrow \overline{\mathcal{M}}_{g,n}$  forgetting the sheaf  $\mathcal{E}$ , and there is a morphism  $p : \overline{\mathcal{S}}_{g,n}^m \rightarrow \overline{\mathcal{S}}_{g,n}^N$  forgetting  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{C/T}(\sum_i m_i \mathcal{D}_i)$ , where  $N = \sum_i m_i$ .

**Proposition B.2.**  $\pi : \mathcal{PS}_{g,n}^N \rightarrow \overline{\mathcal{M}}_{g,n}$  is relatively represented by finite-type quasi-separated Artin stacks with separated diagonal.

*Proof.* For any  $T$ -point of  $\overline{\mathcal{M}}_{g,n}$ , i.e. an  $n$ -punctured stable curve  $(C, \mathcal{D})/T$ ,  $\mathcal{PS}_{g,n}^N \times_{\overline{\mathcal{M}}_{g,n}} T$  is the stack of rank-1 relatively torsion-free sheaves of degree  $g - 1 + N/2$  on  $(\operatorname{Sch}_T)_{\text{ét}}$ . Since rank-1 relatively torsion free-sheaves are obviously semistable, so  $\mathcal{PS}_{g,n}^N \times_{\overline{\mathcal{M}}_{g,n}} T$  is a substack of  $\mathcal{M}_{C/T}^{1, g-1+N/2}$ , i.e. stack of rank-1 semistable sheaves of degree  $g - 1 + N/2$ . It is well-known that  $\mathcal{M}_{C/T}^{1, g-1+N/2}$  is a finite-type quasi-separated Artin stack with separated diagonal. Since being relatively torsion-free is an open condition on the base,  $\mathcal{PS}_{g,n}^N \times_{\overline{\mathcal{M}}_{g,n}} T$  is a finite-type quasi-separated Artin stack with separated diagonal. ■

**Proposition B.3.**  $p : \overline{\mathcal{S}}_{g,n}^m \rightarrow \mathcal{PS}_{g,n}^N$  is relatively represented by finite type quasi-affine schemes.

*Proof.* For any  $T$ -point of  $\mathcal{PS}_{g,n}^N$ , i.e. a  $n$ -pointed prespin curve  $(C, \mathcal{D}, \mathcal{E})/T$ ,  $\overline{\mathcal{S}}_{g,n}^m \times_{\mathcal{PS}_{g,n}^N} T$  is the functor of isomorphisms  $a : \mathcal{E} \rightarrow \underline{\operatorname{Hom}}(\mathcal{E}, \omega_{C/T}(\sum_i m_i \mathcal{D}_i))$ . The next lemma says that  $\overline{\mathcal{S}}_{g,n}^m \times_{\mathcal{PS}_{g,n}^N} T$  is represented by a finite type quasi-affine scheme, hence  $p : \overline{\mathcal{S}}_{g,n}^m \rightarrow \mathcal{PS}_{g,n}^N$  is relatively represented by finite type quasi-affine schemes. ■

**Lemma B.5.** Suppose that  $X$  is proper over a Noetherian scheme  $S$ ,  $\mathcal{F}$  and  $\mathcal{G}$  are coherent sheaves on  $X$  and flat over  $S$ , then  $\underline{\operatorname{Isom}}(\mathcal{F}, \mathcal{G})$  is represented by a scheme finite type and quasi-affine over  $S$ .

*Proof.* The functor  $T \mapsto \operatorname{Hom}_T(\mathcal{F}_T, \mathcal{G}_T)$  is represented by a scheme finite type and affine over  $S$  (see section 7.7.8, 7.7.9 of [118]), denoted by  $M$ . The locus in  $M$  where  $\mathcal{F} \rightarrow \mathcal{G}$  is surjective is open since it's the complement of the image of  $\operatorname{Supp}(\operatorname{coker}(\mathcal{F} \rightarrow \mathcal{G}))$  in  $M$ , the latter is closed since  $X$  is proper over  $S$ . Now in the locus  $U \subset M$  where  $\mathcal{F} \rightarrow \mathcal{G}$  is surjective, the kernel of  $\mathcal{F} \rightarrow \mathcal{G}$ , denoted by  $\mathcal{K}$ , is flat over  $S$ , since both  $\mathcal{F}$  and  $\mathcal{G}$  are flat over  $S$ . Hence  $\underline{\operatorname{Isom}}(\mathcal{F}, \mathcal{G})$  is represented by the vanishing locus of  $\mathcal{K}$ , i.e. the complement of the image of  $\operatorname{Supp}(\mathcal{K})$  in  $U$ , which is open by the properness of  $X$ . ■

Combining propositions B.2 and B.3, we conclude that

**Corollary B.1.**  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  is represented by a finite-type quasi-separated Artin stack over  $\mathrm{Spec} \mathbb{Z}[1/2]$  with separated diagonal.

### B.5 Diagonal of $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$

We are going to describe the diagonal  $\Delta_f$  of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  by investigating the diagonal of morphism  $f : \overline{\mathcal{S}}_{g,n}^{\mathbf{m}} \rightarrow \overline{\mathcal{M}}_{g,n}$ .

**Proposition B.4.** Suppose that  $\mathbb{K}$  is an algebraically-closed field,  $p$  is a  $\mathbb{K}$ -point of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$ , represented by a punctured spin curve  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  over  $\mathbb{K}$ . Let  $\{N_1, \dots, N_r\}$  be the set of nodes where  $\mathcal{E}$  is not locally-free,  $\tilde{\mathcal{C}}$  be the curve  $\mathcal{C}$  normalized at  $\{N_1, \dots, N_r\}$ , then the group of automorphisms of  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  which induce identity on  $(\mathcal{C}, \mathcal{D})$  is

$$(\mathbb{Z}/2\mathbb{Z})^{\oplus s},$$

where  $s$  is the number of connected components of  $\tilde{\mathcal{C}}$ .

*Proof.* It is easy to see that there is a one-to-one correspondence between automorphisms of  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  and automorphisms of  $(\tilde{\mathcal{C}}, \mathcal{D}, \pi^*\mathcal{E}, \pi^*b)$ , where  $\pi : \tilde{\mathcal{C}} \rightarrow \mathcal{C}$  is the normalization morphism at NS punctures. Since  $\pi^*\mathcal{E}$  is locally free, automorphism of  $\pi^*\mathcal{E}$  is given by  $\mathbb{K}^{\times s}$ , hence those automorphisms compatible with  $\pi^*b$  are square roots of  $(1, \dots, 1) \in \mathbb{K}^{\times s}$ , i.e.  $(\mathbb{Z}/2\mathbb{Z})^{\oplus s}$ . ■

**Proposition B.5.**  $\Delta_f$  is unramified.

*Proof.* Suppose that  $\mathbb{K}$  is an algebraically-closed field,  $p$  is a  $\mathbb{K}$ -point of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$ , represented by a punctured spin curve  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  over  $\mathbb{K}$ . We want to show that the automorphism group scheme of  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  which fixes  $(\mathcal{C}, \mathcal{D})$  is étale, or equivalently, the tangent space at identity is trivial.

Let  $D_\epsilon \equiv \mathrm{Spec} \mathbb{K}[\epsilon]/\epsilon^2$ . Consider an automorphism  $\alpha$  of  $\mathcal{E} \times D_\epsilon$  on  $\mathcal{C} \times D_\epsilon$  which reduces to identity on the special fiber. Then  $\alpha$  induces an automorphism  $\tilde{\alpha}$  of  $\pi^*\mathcal{E} \times D_\epsilon$  on  $\tilde{\mathcal{C}} \times D_\epsilon$ , and  $\tilde{\alpha}$  reduces to identity on the special fiber, so

$$\tilde{\alpha} \in (1 + \mathbb{K}\epsilon)^{\oplus s}.$$

Moreover  $\tilde{\alpha}^2 = \mathbb{I}$ , which implies that  $\tilde{\alpha} = \mathbb{I}$ , since 2 is invertible in  $\mathbb{K}$ . As a result we see that  $\alpha = \mathbb{I}$ , i.e. the tangent space at identity is trivial. ■

**Proposition B.6.**  $\Delta_f$  is proper.

*Proof.* Let  $R$  be a complete discrete valuation ring with algebraically-closed residue field  $\mathbb{K}$  and fraction field  $\mathbb{K}_\eta$ ,  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  is a relative punctured spin curve on  $\mathrm{Spec} R$ , and  $\alpha_\eta$  is an automorphism of  $\mathcal{E}$  on generic fiber which is compatible with  $b$ . Using the valuative criterion,

we need to show that  $\alpha_\eta$  extends to an automorphism  $\alpha$  on the whole  $\mathcal{C}$  and is still compatible with  $b$ .

Suppose that  $\{N_1, \dots, N_r\}$  is the set of nodes of generic fiber  $\mathcal{C}_\eta$  where  $\mathcal{E}_\eta$  is not locally free. We can assume that  $N_i$  is a geometric point of  $\mathbb{K}_\eta$  otherwise we can always take a finite extension of  $R$  to make it a geometric point. Note that the valuative criterion still holds for a finite extension of  $R$ . So  $N_i$  extends to a section of  $R$ , which is still denoted by  $N_i$ . Completion of the local ring of  $\mathcal{C}$  at the closed point of  $N_i$  is  $\llbracket R[[x, y]]/(xy) \rrbracket$ . According to the Faltings' classification theorem (Theorem B.2), the completion of  $\mathcal{E}$  at the closed point of  $N_i$  is of type  $(0, 0)$ . Let  $\pi : \tilde{\mathcal{C}} \rightarrow \mathcal{C}$  be the normalization of  $\mathcal{C}$  at sections  $\{N_1, \dots, N_r\}$ , then the adjunction  $\mathcal{E} \rightarrow \pi_* \pi^* \mathcal{E}$  is an isomorphism. Thus, it is enough to extend  $\pi^* \alpha_\eta$  to an automorphism of  $\pi^* \mathcal{E}$ . According to Proposition B.4,  $\pi^* \alpha_\eta$  is a direct product of  $\pm 1$  on connected components of  $\tilde{\mathcal{C}}_\eta$ , so we can extend it tautologically.  $\blacksquare$

As a corollary, we have the following

**Corollary B.2.**  $\bar{\mathcal{S}}_{g,n}^{\mathbf{m}}$  is a separated Deligne-Mumford stack.

## B.6 Deformation Theory of $\bar{\mathcal{S}}_{g,n}^{\mathbf{m}}$

In this section, we discuss the deformation theory of  $\bar{\mathcal{S}}_{g,n}^{\mathbf{m}}$ . We begin with the following proposition

**Proposition B.7.** Suppose that  $\mathbb{K}$  is an algebraically-closed field,  $p$  is a  $\mathbb{K}$ -point of  $\bar{\mathcal{S}}_{g,n}^{\mathbf{m}}$ , represented by a punctured spin curve  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  over  $\mathbb{K}$ . Let  $\{N_1, \dots, N_r\}$  be the set of nodes where  $\mathcal{E}$  is not locally free, then the universal deformation of  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  is the pull-back of the universal deformation of punctured stable curve  $(\mathcal{C}, \mathcal{D})$  under homomorphism

$$\mathfrak{o}_{\mathbb{K}}[[t_1, \dots, t_r, t_{r+1}, \dots, t_{3g-3+n}]] \rightarrow \mathfrak{o}_{\mathbb{K}}[[\tau_1, \dots, \tau_r, t_{r+1}, \dots, t_{3g-3+n}]]], \quad (\text{B.4})$$

$\mathfrak{o}_{\mathbb{K}}$  is the Cohen ring with residue field  $\mathbb{K}$  [20], by sending  $t_i$  to  $\tau_i^2$  for  $i \in \{1, 2, \dots, r\}$ .

*Proof.* Recall that it is shown in [20] that the universal deformation of  $(\mathcal{C}, \mathcal{D})$  is

$$R \equiv \mathfrak{o}_k[[t_1, \dots, t_{3g-3+n}]].$$

At a node of  $\mathcal{C}$ , the complete local ring of the universal deformation of  $\mathcal{C}$  is

$$R[[x, y]]/(xy - t_j),$$

for some index  $j$ . According to the Faltings' classification theorem (Theorem B.2), any deformation of the completion of  $\mathcal{E}$  at  $N_i$  is uniquely determined by a parameter  $\tau_i$  such that  $\tau_i^2 = t_i$ . On the other hand, for any Artinian local ring  $A$  with residue field  $\mathbb{K}$ , and choose a deformation of  $(\mathcal{C}, \mathcal{D})$  with base scheme  $\text{Spec } A$ , denote it by  $(\mathcal{C}_A, \mathcal{D}_A)$ , we claim that  $\mathcal{E}|_{\mathcal{C} - \{N_1, \dots, N_r\}}$  has a unique deformation to  $\mathcal{C}_A - \{N_1, \dots, N_r\}$  as a square root of  $\omega_{\mathcal{C}_A}(\sum_i m_i \mathcal{D}_{A,i})$ . This can be proven as follows. First we notice that there is an exact sequence<sup>90</sup>

<sup>90</sup>Note that  $U_{\text{ét}}$  means the we equip  $U$  with étale topology, and cohomologies are taken with respect to the étale topology of  $U$ .  $\text{Pic}(U)$  is the Picard group of  $U$ .



$$H^1(U_{\text{ét}}, \mu_2) \longrightarrow \text{Pic}(U) \xrightarrow{2} \text{Pic}(U) \longrightarrow H^2(U_{\text{ét}}, \mu_2),$$

where  $U = \mathbb{C} - \{N_1, \dots, N_r\}$ ,  $\mu_2$  is the sheaf of the 2<sup>nd</sup> roots of unity, and the first and the last terms are 2-torsion groups. There is an exact sequence on  $U_A$ , the open subscheme of  $\mathbb{C}_A$  with underlying topological space  $U$ , as well. They are related by restriction map:

$$\begin{array}{ccccccc} H^1(U_{\text{ét}}, \mu_2) & \longrightarrow & \text{Pic}(U_A) & \xrightarrow{2} & \text{Pic}(U_A) & \longrightarrow & H^2(U_{\text{ét}}, \mu_2) \\ \parallel & & \downarrow \text{res} & & \downarrow \text{res} & & \parallel \\ H^1(U_{\text{ét}}, \mu_2) & \longrightarrow & \text{Pic}(U) & \xrightarrow{2} & \text{Pic}(U) & \longrightarrow & H^2(U_{\text{ét}}, \mu_2). \end{array}$$

Notice that the kernel of restriction map,  $\ker(\text{res})$  has a composition series of quotient of  $\mathbb{K}$ -linear spaces where 2 is invertible. Since  $\omega_{\mathbb{C}}(\sum_i m_i \mathcal{D}_i)$  has a square root, from the commutative diagram above, we see that  $\omega_{\mathbb{C}_A}(\sum_i m_i \mathcal{D}_{A,i})$  also has a square root, we want to show that there is a one-to-one correspondence between square roots of  $\omega_{\mathbb{C}_A}(\sum_i m_i \mathcal{D}_{A,i})$  and square roots of  $\omega_{\mathbb{C}}(\sum_i m_i \mathcal{D}_i)$ . Suppose that there are two line bundles on  $\mathbb{C}_A$  denoted by  $\mathcal{L}$  and  $\mathcal{L}'$ , both square to  $\omega_{\mathbb{C}_A}(\sum_i m_i \mathcal{D}_{A,i})$ , and their restrictions to  $\mathbb{C}$  are isomorphic, then  $\mathcal{L}^{-1} \otimes \mathcal{L}' \in \ker(\text{res})$ . However,  $\mathcal{L}^{-1} \otimes \mathcal{L}'$  is in the image of  $H^1(U_{\text{ét}}, \mu_2)$  which is 2-torsion, so  $\mathcal{L}^{-2} \otimes \mathcal{L}'^2 \cong \mathcal{O}_{U_A}$ . This implies that  $\mathcal{L}^{-1} \otimes \mathcal{L}' \cong \mathcal{O}_{U_A}$  since 2 is invertible in  $\ker(\text{res})$ . Thus we see that there is a unique deformation of  $\mathcal{E}|_U$ .

Finally, we are going to show that given any set of deformations of the completion of  $\mathcal{E}$  at  $\{N_1, \dots, N_r\}$ , there exists a unique deformation of  $\mathcal{E}$  which gives rise to deformations of the completion of  $\mathcal{E}$  at  $\{N_1, \dots, N_r\}$ . This is achieved by observing that all we need is the formal gluing at generic points of  $\widehat{\mathcal{O}}_{N_i, \mathbb{C}_A}$ , the completion of  $\mathcal{O}_{N_i, \mathbb{C}_A}$  at the maximal ideal. Note that  $\omega_{\mathbb{C}_A}(\sum_i m_i \mathcal{D}_{A,i})$  naturally carries a formal gluing data, which is a set  $\{a_1, b_1, \dots, a_r, b_r\} \in A((x))^{\times 2r}$ . Since we already have formal gluing data coming from  $\mathcal{E}$  when restrict to  $\mathbb{C}$ , i.e. a set of square roots of  $\{\bar{a}_1, \bar{b}_1, \dots, \bar{a}_r, \bar{b}_r\} \in \mathbb{K}((x))^{\times 2r}$  where  $\bar{a}_i$  (resp.  $\bar{b}_i$ ) is  $a_i$  (resp.  $b_i$ ) modulo the maximal ideal of  $A$ . Since 2 is invertible in  $A$ , square roots lift uniquely in  $A((x))$ , hence there exists a unique deformation of formal gluing data.

To sum up, we have shown that formal deformation of  $\mathcal{E}$  is uniquely determined by deformations of completions  $\widehat{\mathcal{E}}_{N_i}$ , which is determined by a square root of  $t_i$ . Hence the universal deformation of  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  is the pull-back of the universal deformation of punctured stable curve  $(\mathcal{C}, \mathcal{D})$  under homomorphism

$$\mathfrak{o}_{\mathbb{K}}[[t_1, \dots, t_r, t_{r+1}, \dots, t_{3g-3+n}]] \rightarrow \mathfrak{o}_{\mathbb{K}}[[\tau_1, \dots, \tau_r, t_{r+1}, \dots, t_{3g-3+n}]], \quad (\text{B.5})$$

by sending  $t_i$  to  $\tau_i^2$  for  $i \in \{1, 2, \dots, r\}$ . ■

We denote the preimage of  $\mathcal{M}_{g,n}$  under the natural projection  $f : \overline{\mathcal{S}}_{g,n}^{\mathbf{m}} \rightarrow \overline{\mathcal{M}}_{g,n}$  by  $\mathcal{S}_{g,n}^{\mathbf{m}}$ , then we can easily see from the proposition B.7 that

**Corollary B.3.**  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  is a smooth stack, and contains  $\mathcal{S}_{g,n}^{\mathbf{m}}$  as an open dense substack. Moreover  $f : \overline{\mathcal{S}}_{g,n}^{\mathbf{m}} \rightarrow \overline{\mathcal{M}}_{g,n}$  is flat and quasi-finite.

## B.7 Properness of $\overline{\mathcal{S}}_{g,n}^m$

In this section, we prove the following proposition.

**Proposition B.8.** *The projection  $f : \overline{\mathcal{S}}_{g,n}^m \rightarrow \overline{\mathcal{M}}_{g,n}$  is proper.*

*Proof.* Consider the following data: a complete discrete valuation ring (DVR)  $(R, \pi)$  with algebraically-closed residue field  $\mathbb{K}$  and generic point  $\text{Spec } \mathbb{K}(\eta)$ , a relative punctured curve  $(\mathcal{C}, \mathcal{D})$  on  $\text{Spec } R$  such that  $\mathcal{C}_\eta$  is smooth, and a relative punctured spin curve  $(\mathcal{C}_\eta, \mathcal{D}_\eta, \mathcal{E}_\eta, b_\eta)$  on  $\text{Spec } \mathbb{K}(\eta)$ . We are about to show that there exists a pair  $(\mathcal{E}, b)$  extending  $(\mathcal{E}_\eta, b_\eta)$  on the generic fiber which makes  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  a relative punctured spin curve.

Denote the set of generic points of the special fiber  $\mathcal{C}_s$  by  $\{\xi_1, \dots, \xi_e\}$ , and the local ring of  $\mathcal{C}$  at  $\{\xi_1, \dots, \xi_e\}$  is  $\{R_1, \dots, R_e\}$ . Note that  $R_i$  are DVRs. Since  $\mathcal{E}_\eta$  is torsion-free, its localization at  $R_{i\eta}$  extends naturally to a free module of rank one on  $R_i$ , and can be furthermore extended to an open neighborhood of  $\text{Spec } R_i$ . Assume that those open neighborhoods do not intersect with each other. Let  $j : U \hookrightarrow \mathcal{C}$  be the union of  $\mathcal{C}_\eta$  with those neighborhoods, and the extension of  $\mathcal{E}_\eta$  be  $\mathcal{E}_U$ . Since  $\mathcal{C}$  is Cohen-Macaulay<sup>91</sup>,  $\mathcal{E}_U$  is torsion-free, and  $\mathcal{C} - U$  is of codimension 2, so  $j_*\mathcal{E}_U$  is coherent, and satisfies the Serre's  $S_2$  condition. Define  $\mathcal{E} := j_*\mathcal{E}_U$ . It is torsion-free over  $R$  hence flat over  $R$ . As a result,  $\mathcal{E}_s$  satisfies the Serre's  $S_1$  condition, i.e. it is torsion-free. So  $\mathcal{E}$  is relatively torsion-free of rank one, and has the same degree as  $\mathcal{E}_\eta$ .

For the morphism  $b$ , it is equivalent to find an extension of the isomorphism  $a_\eta : \mathcal{E}_\eta \cong \underline{\text{Hom}}(\mathcal{E}_\eta, \omega_{\mathcal{C}_\eta}(\sum_i m_i \mathcal{D}_i))$  (Lemma B.2). Notice that  $a_\eta$  extends to  $U$  (up to a shrinking of  $U$ ), and up to multiplication by a power of uniformizer  $\pi$  on each neighborhood  $U_i$ , we can assume that restriction of  $a$  to  $R_i$  is isomorphism. By adjunction

$$\text{Hom}\left(\mathcal{E}, j_*\underline{\text{Hom}}\left(\mathcal{E}_U, \omega_{\mathcal{C}_U}\left(\sum_i m_i \mathcal{D}_i\right)\right)\right) \cong \text{Hom}\left(\mathcal{E}_U, \underline{\text{Hom}}\left(\mathcal{E}_U, \omega_{\mathcal{C}_U}\left(\sum_i m_i \mathcal{D}_i\right)\right)\right).$$

There is a globally-defined homomorphism  $a : \mathcal{E} \rightarrow \underline{\text{Hom}}(\mathcal{E}, \omega_{\mathcal{C}}(\sum_i m_i \mathcal{D}_i))$  which is isomorphism on  $R_i$ . The same argument in Lemma B.2 shows that  $a$  is an isomorphism. This completes the proof.  $\blacksquare$

## B.8 Connected Components of $\overline{\mathcal{S}}_{g,n}^m$

We are going to prove the last statement of Theorem B.1: the number of connected components of geometric fibers of  $\overline{\mathcal{S}}_{g,n}^m \rightarrow \text{Spec } \mathbb{Z}[1/2]$  is

$$\gcd(2, m_1, m_2, \dots, m_n).$$

Since  $\overline{\mathcal{S}}_{g,n}^m$  is smooth and proper over  $\text{Spec } \mathbb{Z}[1/2]$ , the number of connected components of its geometric fibers is locally constant, so it is enough to prove the statement for the base scheme being  $\text{Spec } \mathbb{C}$ . Using Remark B.3, we can assume that  $m_i$  are either zero or one, so the statement is equivalent to:

<sup>91</sup>This basically means that the depth (maximal length of regular sequence) of every local ring is equal to its Krull dimension.

**Proposition B.9.** *The number of connected components of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  over  $\text{Spec } \mathbb{C}$  is 2 if all  $m_i$  are zero, and 1 if some of  $m_i$  is 1.*

**Remark B.7.** *When all  $m_i$  are zero, the number of connected components of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$  is at least 2, since  $\mathcal{E} \mapsto \dim(H^0(\mathcal{E})) \bmod 2$  is locally-constant on  $\mathcal{S}_{g,n}^{\mathbf{m}}$  and both 1 and 0 show up as value of this function [55]. In fact it is not hard to show that  $\mathcal{E} \mapsto \dim(H^0(\mathcal{E})) \bmod 2$  is locally-constant on  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$ , we give a sketch here:*

Assume that  $(\mathcal{C}, \mathcal{D}, \mathcal{E}, b)$  is a punctured spin curve over algebraically-closed field  $\mathbb{K}$ . Consider a divisor  $\mathbf{a} = \sum_{j=1}^N P_j$  consisting of  $N$  distinct smooth points, such that each irreducible component has at least  $g$  points, where  $g$  is the genus of the curve. Then an argument similar to the one in [55] shows that,  $W_1 \equiv \Gamma(\mathcal{E}(\mathbf{a}))^{92}$  and  $W_2 \equiv \Gamma(\mathcal{E}/\mathcal{E}(-\mathbf{a}))$  are subspaces of  $V \equiv \Gamma(\mathcal{E}(\mathbf{a})/\mathcal{E}(-\mathbf{a}))$ , and  $W_1 \cap W_2 = \Gamma(\mathcal{E})$ . Moreover

$$\dim W_1 = \dim W_2 = \frac{1}{2} \dim V = N.$$

Define the quadratic form  $q$  on  $V$  by

$$q(\mathbf{s}) = \sum_{j=1}^N \text{Res}_{P_j} Q(s_j),$$

where  $\mathbf{s} = (s_1, s_2, \dots, s_N)$  is a collection of sections of  $\mathcal{E}(\mathbf{a})_{P_j}/\mathcal{E}(-\mathbf{a})_{P_j}$ , and  $Q : \mathcal{E}(\mathbf{a}) \rightarrow \omega_{\mathcal{C}}(2\mathbf{a})$  is the quadratic morphism induced by  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}}$ . The same argument in [55] shows that  $W_1$  and  $W_2$  are isotropic and of half the dimension of  $V$ , and hence the dimension modulo 2 of their intersection is locally constant, for the same reason in [55].

The spin structure  $\mathcal{E}$  with  $h^0(\mathcal{E}) \bmod 2 = 0$  (resp.  $h^0(\mathcal{E}) \bmod 2 = 1$ ) is called even spin structure (resp. odd spin structure). Recall that for smooth curve of genus  $g$ , there are  $2^{g-1}(2^g + 1)$  even spin structures and  $2^{g-1}(2^g - 1)$  odd spin structures [55].

The strategy of the proof of Proposition B.9 is the following: Utilizing the fact that  $\overline{\mathcal{M}}_{g,n}$  (equivalently  $\mathcal{M}_{g,n}$ ) is connected over  $\text{Spec } \mathbb{C}$  [20], we investigate the monodromy action of  $\pi_1(\mathcal{M}_{g,n})$  (the mapping-class group of a genus- $g$  Riemann surface with  $n$  punctures) on the fiber of  $\mathcal{S}_{g,n}^{\mathbf{m}} \rightarrow \mathcal{M}_{g,n}$ , and prove by induction on  $g$  to reduce to  $g = 1$  and  $g = 0$ , of which the proof is straightforward. We deal with the induction step first.

**Lemma B.6.** *Suppose that  $g > 1$ . If Proposition B.9 is true for genus from 1 to  $g - 1$ , then Proposition B.9 is true for genus  $g$ .*

*Proof.* The idea is reducing to the boundary of  $\overline{\mathcal{M}}_{g,n}$ , and degenerate the curve into components with lower genus.

**Case  $\mathbf{m} = \mathbf{0}$ :** We want to show that for a generic smooth curve of genus  $g$  with  $n$  punctures, its even spin structures (resp. odd spin structures) lie in the same connected component of  $\overline{\mathcal{S}}_{g,n}$ . Consider two kinds of degenerations of this curve:

---

<sup>92</sup> $\Gamma$  is used to denote the space of global sections, which is the same as  $H^0$ .

1.  $\mathcal{C} = E \cup \mathcal{C}'$  where  $E$  is an elliptic curve, and  $\mathcal{C}'$  is smooth of genus  $g - 1$ ;
2.  $\mathcal{C}$  is irreducible with one node.

For a degeneration of the first kind, it is easy to see that every spin structure  $\mathcal{E}$  comes from pushing forward spin structures on  $E \amalg \mathcal{C}'$ , the disjoint union of the elliptic curve  $E$  and the curve  $\mathcal{C}'$ . The set of even spin structures is thus the disjoint union of two subsets of size  $3 \cdot 2^{g-2}(2^{g-1} + 1)$  and  $2^{g-2}(2^{g-1} - 1)$ , coming from pushing forward of even-even and odd-odd spin structures. We denote them by  $A_1$  and  $A_2$ . Assuming that Proposition 7.8 holds for genus from 1 to  $g - 1$ , we see that each one of these subsets lies in a connected component of  $\overline{\mathcal{S}}_{g,n}$ . For a degeneration of the second kind, it is easy to see that every spin structure  $\mathcal{E}$  comes from either gluing square root of  $\omega_{\tilde{\mathcal{C}}}(r + s)$  or pushing forward spin structure for  $\omega_{\tilde{\mathcal{C}}}$ , where  $\tilde{\mathcal{C}}$  is the normalization of  $\mathcal{C}$ ,  $r$  and  $s$  are preimages of the node of  $\mathcal{C}$ . As a result, the set of even spin structures is disjoint union of two subsets of size  $2^{2g-2}$  and  $2^{g-2}(2^{g-1} + 1)$ . We denote them by  $B_1$  and  $B_2$ . Note that  $B_2$  is the degeneration of twice as many spin structures of a generic smooth curve, since Proposition B.7 shows that universal deformation of spin structure in  $B_2$  is a double covering of universal deformation of  $\mathcal{C}$ , and Proposition B.4 show that the automorphism is trivial, so every spin structure from  $B_2$  is the degeneration of a pair of spin structures. Using the assumption, we see that there are two subsets of even spin structures of a generic smooth curve of size  $2^{2g-2}$  and  $2^{g-1}(2^{g-1} + 1)$ , each lies in a connected component of  $\overline{\mathcal{S}}_{g,n}$ . We denote them by  $B_1$  and  $B'_2$ . Combining results from two kinds of degenerations, we see that  $A_1$  must intersects both  $B_1$  and  $B'_2$ , hence even spin structures of a generic smooth curve lie in a connected component of  $\overline{\mathcal{S}}_{g,n}$ . For odd spin structures, it is essentially the same. The difference is that now  $A_1$  has size  $3 \cdot 2^{g-2}(2^{g-1} - 1)$ ,  $A_2$  has size  $2^{g-2}(2^{g-1} + 1)$ , corresponding to pushing forward of even-odd, and odd-even spin structures;  $B_1$  has size  $2^{2g-2}$ , and  $B_2$  has size  $2^{g-2}(2^{g-1} - 1)$ , so  $B'_2$  has size  $2^{g-1}(2^{g-1} - 1)$ ; the rest of the arguments is the same.

**Case  $\mathbf{m} \neq \mathbf{0}$ :** Consider the following degeneration of a generic smooth punctured curve:

- $\mathcal{C} = E \cup \mathcal{C}'$  where  $E$  is an elliptic curve with 1 punctured point  $\mathcal{D}_i$  such that  $m_i = 1$ , and  $\mathcal{C}'$  is smooth of genus  $g - 1$  with  $n - 1$  punctures.

It is easy to see that every spin structure of  $\mathcal{C}$  is a line bundle and is the gluing of square roots of  $\omega_E(\mathcal{D}_i + r)$  and  $\omega_{\mathcal{C}'}(s + \sum_{j \neq i} m_j \mathcal{D}_j)$ , where  $r$  and  $s$  are preimage of the node under the projection  $E \amalg \mathcal{C}' \rightarrow \mathcal{C}$ . Using the assumption, we see that all spin structures of  $\mathcal{C}$  lie in the same connected component of  $\overline{\mathcal{S}}_{g,n}^{\mathbf{m}}$ . ■

**Lemma B.7.** *Proposition B.9 is true for  $g = 0$ .*

*Proof.* Since for smooth curves of genus zero, the spin structure is unique, so  $\mathcal{S}_{0,n}^{\mathbf{m}} \rightarrow \mathcal{M}_{0,n}$  is an isomorphism and the connectedness is obvious. ■

**Lemma B.8.** *Proposition B.9 is true for  $g = 1$ .*

*Proof.* We fix an elliptic curve  $E$  with  $n$  punctures, the set of spin structures is an  $\mathbb{F}_2$ -affine space modeled on  $H^1(E_{\text{ét}}, \mu_2)$ , where  $H^1(E_{\text{ét}}, \mu_2)$  is the first étale cohomology group of  $E_{\text{ét}}$  with coefficients in the sheaf of the  $2^{\text{nd}}$  roots of unity. Then the monodromy action of  $\pi_1(\mathcal{M}_{1,n}) = \text{MCG}_{1,n}$  on the set of spin structures quotient by transportation is a subgroup of  $\text{SL}_2(\mathbb{F}_2) = \text{Sp}_2(\mathbb{F}_2)$ . Note that the homomorphism  $\text{MCG}_{1,n} \rightarrow \text{Sp}_2(\mathbb{F}_2)$  factors through  $\text{Sp}_2(\mathbb{Z})$ , and this is well-known to be surjective [119]. Since  $\text{Sp}_2(\mathbb{Z}) \rightarrow \text{Sp}_2(\mathbb{F}_2)$  is surjective, we see that  $\text{MCG}_{1,n} \rightarrow \text{Sp}_2(\mathbb{F}_2)$  is surjective.

**Case  $\mathbf{m} = \mathbf{0}$ :** The space of spin structures is exactly  $H^1(E_{\text{ét}}, \mu_2)$ , hence  $\text{MCG}_{1,n}$  acts on by  $\text{SL}_2(\mathbb{F}_2)$  and it is easy to see that the only odd spin structure is the trivial one and even spin structures are permuted under  $\text{SL}_2(\mathbb{F}_2)$  action. As a result, even spin structures lie in the same connected component of  $\mathcal{S}_{1,n}$ .

**Case  $\mathbf{m} \neq \mathbf{0}$ :** We want to show that  $\text{MCG}_{1,n}$  acts on the set of spin structures transitively. We first show that the action has no fixed point, in other word, for any spin structure  $\mathcal{E}$ , there exists another spin structure  $\mathcal{E}'$  such that they are in the same connected component of  $\mathcal{S}_{1,n}^{\mathbf{m}}$ . Consider the the degeneration  $\mathcal{C} = F_1 \cup F_2$  where  $F_i \cong \mathbb{P}^1$ , they intersect at two nodes  $r_1$  and  $r_2$ , there is one puncturing point  $\mathcal{D}_1 \in F_1(\mathbb{C})$  with  $m_1 = 1$  and all other punctures are on the  $F_2$ . It is easy to see that there are two spin structures of  $\mathcal{C}$ ,  $\mathcal{E}_1$  and  $\mathcal{E}_2$ , such that  $\mathcal{E}_i$  is the push-forward of the spin structure for  $\omega_{\mathcal{C}_i}(\sum_j m_j \mathcal{D}_j)$  on curve  $\mathcal{C}_i$ , where  $\mathcal{C}_i$  is the normalization of  $\mathcal{C}$  at the node  $r_i$ . Using Proposition B.4, we see that  $\mathcal{E}_i$  has trivial automorphism, hence according to Proposition B.7,  $\mathcal{E}_i$  is the degeneration of two spin structures of a generic smooth curve. As a result, every spin structure of a generic smooth curve share the connected component of  $\mathcal{S}_{1,n}^{\mathbf{m}}$  with another spin structure. Now, if the action of  $\text{MCG}_{1,n}$  on the set of spin structures is not transitive, then the only possibility is that there are two pairs of spin structures, each pair is stabilized and permuted by  $\text{MCG}_{1,n}$ . However, this implies that the image of  $\text{MCG}_{1,n}$  is of order at most 4, which is absurd because a further quotient by transportation subgroup has order 6. This concludes the proof of the case  $\mathbf{m} \neq \mathbf{0}$ . ■

## B.9 Gluing Punctures

To simplify notation, we will use  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}$  to denote the moduli space of spin curves which is used to be called  $\overline{\mathcal{S}}_{g,n+|\mathbf{m}|}^{\mathbf{m}}$ , i.e. curves of genus  $g$  with  $n$  NS marked points and  $n_{\text{R}}$  R marked points. Similarly for uncompactified version,  $\mathcal{S}_{g,n,n_{\text{R}}} := \mathcal{S}_{g,n_{\text{NS}}+|\mathbf{m}|}^{\mathbf{m}}$ . And we define the boundary of  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}$  to be  $\overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}} \setminus \mathcal{S}_{g,n_{\text{NS}},n_{\text{R}}}$ , with notation  $\partial \overline{\mathcal{S}}_{g,n_{\text{NS}},n_{\text{R}}}$ . Suppose that  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are punctured spin curves over a base  $S$ , and that  $n_{\text{NS}}^1$  and  $n_{\text{NS}}^2$  are nonzero, then we can pick one NS divisor from each curve, say that the  $i_1$ 'th NS divisor  $\mathcal{D}_{i_1}$  on  $\mathcal{C}_1$  and  $i_2$ 'th NS divisor  $\mathcal{D}_{i_2}$  on  $\mathcal{C}_2$ . The projection  $\mathcal{D}_{i_1} \rightarrow S$  is an isomorphism, same for  $\mathcal{D}_{i_2}$ , so we can use these isomorphisms to define a canonical identification between  $\varphi : \mathcal{D}_{i_1} \cong \mathcal{D}_{i_2}$ , and glue them to get a new curve  $\mathcal{C} = \mathcal{C}_1 \cup_{\varphi} \mathcal{C}_2$ , and the spin structure  $\mathcal{E}$  on  $\mathcal{C}$  is the direct sum of pushforward of spin structures  $\mathcal{E}_1$  and  $\mathcal{E}_2$  via the gluing map  $\mathcal{C}_1 \sqcup \mathcal{C}_2 \rightarrow \mathcal{C}$ . This

construction is functorial hence giving rise to an operation on corresponding moduli spaces:

$$\mathbf{m}_{i_1, i_2}^{\text{NS}} : \overline{\mathcal{S}}_{g_1, n_{\text{NS}}^1, n_{\text{R}}^1} \times \overline{\mathcal{S}}_{g_2, n_{\text{NS}}^2, n_{\text{R}}^2} \rightarrow \partial \overline{\mathcal{S}}_{g_1+g_2, n_{\text{NS}}^1+n_{\text{NS}}^2-2, n_{\text{R}}^1+n_{\text{R}}^2}.$$

This can be defined for a single punctured spin curve with at least two NS puncturing divisors as well:

$$\mathfrak{g}_{i,j}^{\text{NS}} : \overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} \rightarrow \partial \overline{\mathcal{S}}_{g+1, n_{\text{NS}}-2, n_{\text{R}}}.$$

Note that these maps depends on the choice of  $i_1$ ,  $i_2$ ,  $i$ , and  $j$ , in an equivariant way. For example, if  $\sigma_{i_1, i'_1} \in S_{n_{\text{NS}}^1}$  switches  $i_1$  and  $i'_1$  (it acts  $\overline{\mathcal{S}}_{g_1, n_{\text{NS}}^1, n_{\text{NS}}^1}$  naturally), then we have  $\mathbf{m}_{i'_1, i_2}^{\text{NS}} = \mathbf{m}_{i_1, i_2}^{\text{NS}} \circ \sigma_{i_1, i'_1}$ .

We move on to gluing R divisors: suppose that  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are punctured spin curves over a base  $S$ , and that  $m_1$  and  $m_2$  are nonzero, we can pick one R puncture from each curve,  $\iota_1 : S \xrightarrow{\sim} \mathcal{D}_{i_1} \subset C_1$  and  $\iota_2 : S \xrightarrow{\sim} \mathcal{D}_{i_2} \subset C_2$ , then  $\iota_1^* \mathcal{E}_1$  and  $\iota_2^* \mathcal{E}_2$  are two line bundles on  $S$  with isomorphisms  $b_1 : (\iota_1^* \mathcal{E}_1)^{\otimes 2} \cong \mathcal{O}_S, b_2 : (\iota_2^* \mathcal{E}_2)^{\otimes 2} \cong \mathcal{O}_S$ . If we try to glue  $\mathcal{C}_1$  and  $\mathcal{C}_2$ , surely we can still get a curve as in the NS case:  $\mathcal{C} = \mathcal{C}_1 \cup_{\varphi}^{\mathcal{D}_{i_1}} \mathcal{C}_2$ . However, there is no canonical isomorphism between  $\iota_1^* \mathcal{E}_1$  and  $\iota_2^* \mathcal{E}_2$ , every isomorphism can be multiplied by  $-1$  to get another isomorphism. A way out is to adding a choice *by hand*, namely, consider the line bundle  $\mathcal{L}_{12} = \iota_1^* \mathcal{E}_1^\vee \otimes \iota_2^* \mathcal{E}_2$  with the isomorphism  $b_{12} = b_1^{-1} \otimes b_2 : \mathcal{L}_{12}^{\otimes 2} \cong \mathcal{O}_S$ , then we have a morphism between schemes  $f : \mathbb{V}(\mathcal{L}_{12}^\vee) \rightarrow \mathbb{A}_S^1$  by sending a section of  $\mathcal{L}_{12}$  to the image of its square under  $b_{12}$ . The scheme

$$G_{i_1, i_2} \equiv f^{-1}(\mathbf{1}),$$

parametrizes isomorphisms from  $\iota_1^* \mathcal{E}_1$  to  $\iota_2^* \mathcal{E}_2$  which commutes with  $b_i$ . Locally on  $S$ , after choosing trivializations of  $\iota_1^* \mathcal{E}_1, \iota_2^* \mathcal{E}_2$ ,  $f$  becomes the morphisms  $\mathbb{A}^1 \rightarrow \mathbb{A}^1$  by sending a section to its square, hence locally  $G_{i_1, i_2}$  is disjoint union of two copies of the base scheme, and globally there is a  $\mathbb{Z}/2$ -action on  $G_{i_1, i_2}$  via multiplication by  $-1$ , thus  $G_{i_1, i_2}$  is a  $\mathbb{Z}/2$ -torsor.

By base change to the scheme  $G_{i_1, i_2}$ , there is a universal isomorphism  $\alpha : \iota_1^* \mathcal{E}_1 \cong \iota_2^* \mathcal{E}_2$  which commutes with  $b_i$ , so we can define a gluing of  $\mathcal{C}_1 \times_S G_{i_1, i_2}$  and  $\mathcal{C}_2 \times_S G_{i_1, i_2}$  by taking the spinor sheaf  $\mathcal{E}$  on  $\mathcal{C} = \mathcal{C}_1 \cup_{\varphi}^{\mathcal{D}_{i_1}} \mathcal{C}_2$  to be the kernel of

$$p_{1*} \mathcal{E}_1 \oplus p_{2*} \mathcal{E}_2 \rightarrow \iota_1^* \mathcal{E}_1 \oplus \iota_2^* \mathcal{E}_2 \xrightarrow{(\alpha, \text{Id})} \iota_2^* \mathcal{E}_2,$$

where  $p_i : \mathcal{C}_i \rightarrow \mathcal{C}$  is the canonical projection. It is easy to see that  $\mathcal{E}$  is locally-free in a neighborhood of the R node formed by gluing, and

$$\deg \mathcal{E} = \deg \mathcal{E}_1 + \deg \mathcal{E}_2 - 1.$$

Furthermore, it follows from the definition of  $\mathcal{E}$  and  $\alpha$  commuting with  $b_i$  that the composition

$$\mathcal{E}^{\otimes 2} \rightarrow (p_{1*} \mathcal{E}_1 \oplus p_{2*} \mathcal{E}_2)^{\otimes 2} \rightarrow p_{1*} \mathcal{E}_1^{\otimes 2} \oplus p_{2*} \mathcal{E}_2^{\otimes 2}$$

$$\xrightarrow{b_1 \oplus b_2} p_{1*} \omega_{\mathcal{C}_1/S}(\mathcal{D}_{i_1} + E) \oplus p_{2*} \omega_{\mathcal{C}_2/S}(\mathcal{D}_{i_2} + H) \rightarrow \iota_2^* \omega_{\mathcal{C}_2/S}(\mathcal{D}_{i_2} + H),$$

is zero, where  $E$  and  $H$  are other puncturing divisors on  $\mathcal{C}_1$  and  $\mathcal{C}_2$ . The kernel of  $p_{1*} \omega_{\mathcal{C}_1/S}(\mathcal{D}_{i_1} + E) \oplus p_{2*} \omega_{\mathcal{C}_2/S}(\mathcal{D}_{i_2} + H) \rightarrow \iota_2^* \omega_{\mathcal{C}_2/S}(\mathcal{D}_{i_2} + H)$  is  $\omega_{\mathcal{C}/S}(E + H)$ . We thus have a homomorphism  $b : \mathcal{E}^{\otimes 2} \rightarrow \omega_{\mathcal{C}/S}(E + H)$  which is an isomorphism on the smooth locus. This confirms that we obtain a spin structure on  $\mathcal{C}$ , and the construction is also functorial. Consequently,  $G_{i_1, i_2}$  forms a relative scheme  $\mathcal{G}_{i_1, i_2}$  over the product moduli space  $\overline{\mathcal{S}}_{g_1, n_{\text{NS}}^1, n_{\text{R}}^1} \times \overline{\mathcal{S}}_{g_2, n_{\text{NS}}^2, n_{\text{R}}^2}$ . Hence giving rise to an operation on corresponding moduli spaces:

$$\begin{array}{ccc} \mathcal{G}_{i_1, i_2} & \xrightarrow{m_{i_1, i_2}^{\text{R}}} & \partial \overline{\mathcal{S}}_{g_1 + g_2, n_{\text{NS}}^1 + n_{\text{NS}}^2, n_{\text{R}}^1 + n_{\text{R}}^2 - 2} \\ \pi_{i_1, i_2} \downarrow & & \\ \overline{\mathcal{S}}_{g_1, n_{\text{NS}}^1, n_{\text{R}}^1} \times \overline{\mathcal{S}}_{g_2, n_{\text{NS}}^2, n_{\text{R}}^2} & & \end{array}$$

This can be defined for a single punctured spin curve with at least two NS marking divisors as well:

$$\begin{array}{ccc} \mathcal{G}_{i, j} & \xrightarrow{g_{i, j}^{\text{R}}} & \partial \overline{\mathcal{S}}_{g+1, n_{\text{NS}}, n_{\text{R}} - 2} \\ \pi_{i, j} \downarrow & & \\ \overline{\mathcal{S}}_{g, n_{\text{NS}}, n_{\text{R}}} & & \end{array}$$

Note that  $\pi_{i_1, i_2}$  and  $\pi_{i, j}$  are  $\mathbb{Z}/2$ -torsors. Similar to the gluing of NS divisors case, these maps depends on the choice of  $i_1, i_2$  and  $i, j$  in an equivariant way.

## References

- [1] J. Harris and D. Mumford, *On the Kodaira Dimension of the Moduli Space of Curves*, *Invent. Math.* **67** (1982) 23.
- [2] J. L. Harer, *The Cohomology of the Moduli Space of Curves*, in *Theory of Moduli*, pp. 138–221. Springer Berlin Heidelberg, 1988.
- [3] K. Ludwig, *On the Geometry of the Moduli Space of Spin Curves*, *J. Algebraic. Geom.* **19** (2010) 133 [[arXiv:0707.1831](#)].
- [4] C. LeBrun and M. Rothstein, *Moduli of Super-Riemann Surfaces*, *Comm. Math. Phys.* **117** (1988) 159.
- [5] L. Hodgkin, *The Complex Structure of Supermoduli Space*, *Class. Quantum Grav.* **12** (1995) 2135.
- [6] R. Donagi and E. Witten, *Supermoduli Space is not Projected*, in *Proc. Symp. Pure Math*, vol. 90, p. 19, 2015, [1304.7798](#).
- [7] G. Codogni and F. Viviani, *Moduli and Periods of Supersymmetric Curves*, *arXiv e-prints* (2017) [[1706.04910](#)].

- [8] J. D. Pérez, D. H. Ruipérez and C. S. de Salas, *Global Structures for the Moduli of (Punctured) Super-Riemann Surfaces*, *J. Geom. Phys.* **21** (1997) 199.
- [9] M. Olsson, *Algebraic Spaces and Stacks*. American Mathematical Society, 2016.
- [10] E. Witten, *Notes on Supermanifolds and Integration*, *arXiv e-prints* (2012) [[1209.2199](#)].
- [11] E. Witten, *Notes on Super Riemann Surfaces and their Moduli*, *arXiv e-prints* (2012) [[1209.2459](#)].
- [12] E. Witten, *Superstring Perturbation Theory Revisited*, *arXiv e-prints* (2012) [[1209.5461](#)].
- [13] M. A. Baranov and A. Schwarz, *Multiloop Contribution to String Theory*, *JETP Lett.* **42** (1985) 419.
- [14] D. Friedan, *Notes on String Theory and Two-Dimensional Conformal Field Theory*, in *Unified String Theories*, pp. 162–213, World Scientific, 1986.
- [15] D. Friedan, E. J. Martinec and S. Shenker, *Conformal Invariance, Supersymmetry and String Theory*, *Nucl. Phys.* **B271** (1986) 93.
- [16] E. Verlinde and H. Verlinde, *Multiloop Calculations in Covariant Superstring Theory*, *Phys. Lett.* **192** (B1987) 95.
- [17] A. Sen, *Off-Shell Amplitudes in Superstring Theory*, *Fortsch. Phys.* **63** (2015) 149 [[1408.0571](#)].
- [18] A. Sen and E. Witten, *Filling the Gaps with PCOs*, *J. High Energ. Phys.* **2015** (2015) [[1504.00609](#)].
- [19] A. Sen, *Supersymmetry Restoration in Superstring Perturbation Theory*, *J. High Energ. Phys.* **12** (2015) 75 [[1508.02481](#)].
- [20] P. Deligne and D. Mumford, *The Irreducibility of the Space of Curves of Given Genus*, *Publications Mathématiques de IHÉS* **36** (1969) 75.
- [21] T. Erler, S. Konopka and I. Sachs, *Resolving Witten’s Superstring Field Theory*, *Journal of High Energy Physics* **4** (2014) 150 [[1312.2948](#)].
- [22] T. Erler and S. Konopka, *Vertical Integration from the Large Hilbert Space*, *J. High Energ. Phys.* **2017** (2017) [[arXiv:1710.07232](#)].
- [23] N. Seiberg and E. Witten, *Spin Structures in String Theory*, *Nucl. Phys.* **B276** (1986) 272.
- [24] H. Masur, *The Extension of the Weil-Petersson Metric to the Boundary of Teichmüller Space*, *Duke Math. J.* **43** (1976) 623.
- [25] S. A. Wolpert, *Asymptotics of the Spectrum and the Selberg Zeta-Function on the Space of Riemann Surfaces*, *Comm. Math. Phys.* **112** (1987) 283.
- [26] K. J. Costello, *The Partition Function of a Topological Field Theory*, *J. Topol.* **2** (2009) 779 [[math/0509264](#)].
- [27] M. Batchelor, *The Structure of Supermanifolds*, *Trans. Am. Math. Soc.* **253** (1979) 329.
- [28] M. Eastwood and C. LeBrun, *Thickening and Supersymmetric Extensions of Complex Manifolds*, *Am. J. Math.* **108** (1986) 1177.
- [29] P. Deligne, *Letter to Y. I. Manin*, 25 September (1987) .
- [30] Y. Wakabayashi, *Moduli of Log-Twisted  $N = 1$  SUSY Curves*, *arXiv e-prints* (2016)



- [1602.07059].
- [31] N. Ott and A. Voronov, *The Supermoduli Space of Genus-Zero Super-Riemann Surfaces with Ramond Punctures*, *arXiv e-prints* (2019) [[1910.05655](#)].
  - [32] U. Bruzzo and D. Hernández Ruipérez, *The Supermoduli of SUSY Curves with Ramond Punctures*, *arXiv e-prints* (2019) [[1910.12236](#)].
  - [33] J. D. Cohn, *Modular Geometry of Superconformal Field Theory*, *Nucl. Phys.* **B306** (1988) 239.
  - [34] P. S. Howe, *Super-Weyl Transformations in Two Dimensions*, *J. Phys.* **A12** (1979) 393.
  - [35] L. Crane and J. M. Rabin, *Global Properties of Supermanifolds*, *Comm. Math. Phys.* **100** (1985) 141.
  - [36] L. Crane and J. M. Rabin, *How Different are the Supermanifolds of Rogers and DeWitt?*, *Comm. Math. Phys.* **102** (1985) 123.
  - [37] L. Crane and J. M. Rabin, *Super-Riemann Surfaces: Uniformization and Teichmüller Theory*, *Comm. Math. Phys.* **113** (1988) 601.
  - [38] A. A. Rosly, A. S. Schwarz and A. A. Voronov, *Geometry of Superconformal Manifolds*, *Comm. Math. Phys.* **119** (1988) 129.
  - [39] M. Cornalba, *Moduli of Curves and Theta-Characteristics*, in *Lectures on Riemann Surfaces*, pp. 560–589. World Scientific, 1989.
  - [40] T. J. Jarvis, *Compactification of the Moduli Space of Generalized Spin Curves*, Ph.D. Thesis, Princeton University, 1994.
  - [41] D. J. Gross, J. A. Harvey, E. J. Martinec and R. Rohm, *Heterotic-String Theory I. The Free Heterotic String*, *Nucl. Phys.* **B256** (1985) 253.
  - [42] L. Alvarez-Gaumé, P. Nelson, C. Gomez, G. Sierra and C. Vafa, *Fermionic Strings in the Operator Formalism*, *Nucl. Phys.* **B311** (1988) 333.
  - [43] S. B. Giddings, *Punctures on Super-Riemann Surfaces*, *Comm. Math. Phys.* **143** (1992) 355.
  - [44] T. J. Jarvis, *Torsion-Free Sheaves and Moduli of Generalized Spin Curves*, *Compositio Mathematica* **110** (1998) 291–333 [[alg-geom/9502022](#)].
  - [45] T. J. Jarvis, *Geometry of the Moduli of Higher-Spin Curve*, *International Journal of Mathematics* **11** (2000) 637 [[math/9809138](#)].
  - [46] P. Ramond, *Dual Theory for Free Fermions*, *Phys. Rev.* **D3** (1971) 2415.
  - [47] R. Pius, A. Rudra and A. Sen, *Mass Renormalization in String Theory: Special States*, *J. High Energ. Phys.* **2014** (2014) [[arXiv:1311.1257](#)].
  - [48] R. Pius, A. Rudra and A. Sen, *Mass Renormalization in String Theory: General States*, *J. High Energ. Phys.* **2014** (2014) [[arXiv:1401.7014](#)].
  - [49] A. Sen, *Gauge-Invariant 1PI Effective Superstring Field Theory: Inclusion of the Ramond Sector*, *J. High Energ. Phys.* **2015** (2015) [[arXiv:1501.00988](#)].
  - [50] T. Kimura, J. Stasheff and A. A. Voronov, *On Operad Structures of Moduli Spaces and String Theory*, *Comm. Math. Phys.* **171** (1995) 1 [[hep-th/9307114](#)].
  - [51] A. Belopolsky, *New Geometrical Approach to Superstrings*, *arXiv e-prints* (1997)

- [[hep-th/9703183](#)].
- [52] A. Sen and B. Zwiebach, *Quantum Background-Independence of Closed-String Field Theory*, *Nucl. Phys.* **B423** (1994) 580 [[hep-th/9311009](#)].
  - [53] A. Sen and B. Zwiebach, *Background-Independent Algebraic Structures in Closed-String Field Theory*, *Comm. Math. Phys.* **177** (1996) 305 [[hep-th/9408053](#)].
  - [54] The Stacks Project Authors, *First Cohomology and Torsors*, 2019.
  - [55] D. Mumford, *Theta Characteristics of an Algebraic Curve*, *Annales Scientifiques de l'École Normale Supérieure* **4** (1971) 181.
  - [56] B. Fantechi, L. Illusie and L. Göttsche, *Fundamental Algebraic Geometry: Grothendieck's FGA Explained*, vol. 123 of *Mathematical Surveys and Monographs*. American Mathematical Soc., 2005.
  - [57] F. F. Voronov and A. V. Zorich, *Bordism Theory and Homotopy Properties of Supermanifolds*, *Funct. Anal. Appl.* **21** (1987) 237.
  - [58] T. Voronov, *Geometric Integration Theory on Supermanifolds*, vol. 1. CRC Press, 1991.
  - [59] A. Sen and B. Zwiebach, *A Proof of Local Background-Independence of Classical Closed-String Field Theory*, *Nucl. Phys.* **B414** (1994) 649 [[hep-th/9307088](#)].
  - [60] A. Sen, *Background-Independence of Closed-Superstring Field Theory*, *J. High Energy. Phys.* **2** (2018) 155 [[1711.08468](#)].
  - [61] T. Erler and C. Maccaferri, *String Field Theory Solution for Any Open String Background*, *J. High Energy. Phys.* **10** (2014) 29 [[1406.3021](#)].
  - [62] T. Erler and C. Maccaferri, *String Field Theory Solution for Any Open String Background II*, *arXiv e-prints* (2019) [[1909.11675](#)].
  - [63] A. Sen, *On the Background-Independence of String Field Theory*, *Nucl. Phys.* **B345** (1990) 551.
  - [64] A. Sen, *On the Background-Independence of String Field Theory II. Analysis of On-Shell S-Matrix Elements*, *Nucl. Phys.* **B347** (1990) 270.
  - [65] A. Sen, *On the Background-Independence of String Field Theory III. Explicit Field Redefinitions*, *Nucl. Phys.* **B391** (1993) 550 [[hep-th/9201041](#)].
  - [66] E. Witten, *On Background-Independent Open-String Field Theory*, *Phys. Rev.* **46** (1992) 5467 [[hep-th/9208027](#)].
  - [67] E. Witten, *Some Computations in Background-Independent Off-Shell String Theory*, *Phys. Rev.* **D47** (1993) 3405 [[hep-th/9210065](#)].
  - [68] E. Witten, *Quantum Background Independence In String Theory*, *arXiv High Energy Physics - Theory e-prints* (1993) [[hep-th/9306122](#)].
  - [69] S. L. Shatashvili, *Comment on the Background-Independent Open-String Theory*, *Phys. Lett.* **B311** (1993) 83 [[hep-th/9303143](#)].
  - [70] S. L. Shatashvili, *On Background-Independence in String Theory*, in *Integrable Models and Strings*, pp. 221–234. Springer Berlin Heidelberg, 1994. [[hep-th/9311177](#)].

- [71] B. Zwiebach, *Building String Field Theory around Non-Conformal Backgrounds*, *Nucl. Phys.* **B480** (1996) 541 [[hep-th/9606153](#)].
- [72] O. Bergman and B. Zwiebach, *The Dilaton Theorem and Closed-String Backgrounds*, *Nucl. Phys.* **B441** (1995) 76 [[hep-th/9411047](#)].
- [73] S. Rahman and B. Zwiebach, *Vacuum Vertices and the Ghost-Dilaton*, *Nucl. Phys.* **B471** (1996) 233 [[hep-th/9507038](#)].
- [74] A. Belopolsky and B. Zwiebach, *Who Changes the String Coupling?*, *Nucl. Phys.* **B472** (1996) 109 [[hep-th/9511077](#)].
- [75] S. Mandelstam, *Interacting-String Picture of Dual-Resonance Models*, *Nucl. Phys.* **B64** (1973) 205.
- [76] S. Mandelstam, *Interacting-String Picture of the Fermionic String*, *Prog. Theoretical Phys. Supplement* **86** (1986) 163.
- [77] S. B. Giddings and E. Martinec, *Conformal Geometry and String Field Theory*, *Nucl. Phys.* **B278** (1986) 91.
- [78] H. Hata and B. Zwiebach, *Developing the Covariant Batalin-Vilkovisky Approach to String Theory*, *Annals of Physics* **229** (1994) 177 [[hep-th/9301097](#)].
- [79] K. Costello and B. Zwiebach, *Hyperbolic String Vertices*, *arXiv e-prints* (2019) [[1909.00033](#)].
- [80] M. Artin, *Algebraization of Formal Moduli I*, in *Global Analysis (Papers in Honor of K. Kodaira)*, pp. 21–72. Univ. Tokyo Press, 1969.
- [81] M. Artin, *Algebraization of Formal Moduli II: Existence of Modifications*, *Ann. Math.* **91** (1970) 88.
- [82] M. Artin, *Versal Deformations and Algebraic Stacks*, *Invent. Math.* **27** (1974) 165.
- [83] M. Schlessinger, *Functors of Artin Rings*, *Trans. Am. Math. Soc.* **130** (1968) 208.
- [84] A. Y. Vaintrob, *Deformation of Complex Superspaces and Coherent Sheaves on Them*, *J. Sov. Math.* **51** (1990) 2140.
- [85] H. Flenner and D. Sundararaman, *Analytic Geometry of Complex Superspaces*, *Trans. Am. Math. Soc.* **330** (1992) 1.
- [86] H. Matsumura, *Commutative Algebra*, Mathematics Lecture Note Series, vol. 56. Benjamin/Cummings Publishing Co., 1989.
- [87] U. Görtz and T. Wedhorn, *Algebraic Geometry I: Schemes with Examples and Exercises*, Advanced Lectures in Mathematics. Vieweg+Teubner Verlag, 2010.
- [88] A. Grothendieck, *Éléments de Géométrie Algébrique IV. Étude Locale des Schémas et des Morphismes de Schémas, Quatrième Partie*, *Publications Mathématiques de l'IHÉS* **32** (1967) 5.
- [89] R. Hartshorne, *Deformation Theory*. Springer New York, 2010.
- [90] B. Zwiebach, *Consistency of Closed-String Polyhedra from Minimal Area*, *Phys. Lett.* **B241** (1990) 343.
- [91] B. Zwiebach, *Quantum Closed-Strings from Minimal Area*, *Mod. Phys. Lett.* **A05** (1990) 2753.

- [92] B. Zwiebach, *Quantum Open-String Theory with Manifest Closed-String Factorization*, *Phys. Lett.* **B256** (1991) 22.
- [93] B. Zwiebach, *How Covariant Closed-String Theory Solves a Minimal-Area Problem*, *Comm. Math. Phys.* **136** (1991) 83.
- [94] B. Zwiebach, *Minimal-Area Problems and Quantum Open Strings*, *Comm. Math. Phys.* **141** (1991) 577.
- [95] B. Zwiebach, *Closed-String Field Theory: Quantum Action and the BV Master Equation*, *Nucl. Phys.* **B390** (1993) 33 [[hep-th/9206084](#)].
- [96] M. Headrick and B. Zwiebach, *Convex Programs for Minimal-Area Problems*, *arXiv e-prints* (2018) [[1806.00449](#)].
- [97] M. Headrick and B. Zwiebach, *Minimal-Area Metrics on the Swiss Cross and Punctured Torus*, *arXiv e-prints* (2018) [[1806.00450](#)].
- [98] U. Naseer and B. Zwiebach, *Extremal Isosystolic Metrics with Multiple Bands of Crossing Geodesics*, *arXiv e-prints* (2019) [[1903.11755](#)].
- [99] B. Jurčo and K. Münster, *Type-II Superstring Field Theory: Geometric Approach and Operadic Description*, *J. High Energ. Phys.* **4** (2013) 126 [[1303.2323](#)].
- [100] K. Münster and I. Sachs, *Homotopy Classification of Bosonic String Field Theory*, *Comm. Math. Phys.* **330** (2014) 1227 [[1208.5626](#)].
- [101] B. Zwiebach, *Oriented Open-Closed String Theory Revisited*, *Ann. Phys.* **267** (1998) 193 [[hep-th/9705241](#)].
- [102] S. F. Moosavian, A. Sen and M. Verma, *Superstring Field Theory with Open and Closed Strings*, *arXiv e-prints* (2019) [[1907.10632](#)].
- [103] O. DeWolfe, *Minimal Area Nonorientable String Diagrams*, *arXiv e-prints* (1997) [[hep-th/9708084](#)].
- [104] A. Belopolsky, *Picture-Changing Operators in Supergeometry and Superstring Theory*, *arXiv e-prints* (1997) [[hep-th/9706033](#)].
- [105] A. Neveu, J. Schwarz and C. Thorn, *Reformulation of the Dual Pion Model*, *Phys. Lett.* **B35** (1971) 529.
- [106] L. Brink, D. Olive, C. Rebbi and J. Scherk, *The Missing Gauge Conditions for the Dual Fermion Emission Vertex and their Consequences*, *Phys. Lett.* **B45** (1973) 379.
- [107] E. Verlinde and H. Verlinde, *Lectures on String Perturbation Theory*, in *Superstrings '88* (M. Green, M. Grisaru, R. Iengo, E. Sezgin and A. Strominger, eds.), pp. 189–250, World Scientific Singapore, 1989.
- [108] A. Sen, *Master Action for Heterotic and Type-II String Field Theories*, *J. High Energ. Phys.* **2** (2016) 87 [[1508.05387](#)].
- [109] E. Martinec, *Conformal Field Theory on a (Super-)Riemann Surface*, *Nucl. Phys.* **B281** (1987) 157.
- [110] L. Brink, P. Di Vecchia and P. Howe, *A Locally-Supersymmetric and Reparametrization-Invariant Action for the Spinning String*, *Phys. Lett.* **B65** (1976) 471.

- [111] I. N. Bernshtein and D. A. Leites, *Integration of Differential Forms on Supermanifolds*, *Funct. Anal. Appl.* **11** (1978) 219.
- [112] Y. I. Manin, *Gauge Field Theory and Complex Geometry*. Springer Berlin Heidelberg, 1997.
- [113] I. N. Bernshtein and D. A. Leites, *Integral Forms and the Stokes Formula on Supermanifolds*, *Funct. Anal. Appl.* **11** (1977) 45.
- [114] V. G. Knizhnik, *Covariant Superstring Fermion Amplitudes from the Sum over Fermionic Surfaces*, *Phys. Lett.* **B178** (1986) 21.
- [115] A. Belopolsky, *De Rham Cohomology of the Supermanifolds and Superstring BRST Cohomology*, *Phys. Lett.* **B403** (1997) 47 [[hep-th/9609220](#)].
- [116] G. Faltings, *Moduli-Stacks for Bundles on Semistable Curves*, *Math. Ann.* **304** (1996) 489.
- [117] The Stacks Project Authors, *Zariski's Main Theorem*, 2019.
- [118] A. Grothendieck, *Éléments de Géométrie Algébrique : III. Étude Cohomologique des Faisceaux Cohérents, Seconde Partie*, *Publications Mathématiques de l'IHÉS* **17** (1963) 5.
- [119] W. H. Meeks and J. Patrusky, *Representing Homology Classes by Embedded Circles on a Compact Surface*, *Illinois J. Math* **22** (1978) 262.