

Towards Gallai's path decomposition conjecture

F. Botler¹ M. Sambinelli²

¹Programa de Engenharia de Sistemas e Computação
Universidade Federal do Rio de Janeiro

²Centro de Matemática, Computação e Cognição
Universidade Federal do ABC

Abstract

A path decomposition of a graph G is a collection of edge-disjoint paths of G that covers the edge set of G . Gallai (1968) conjectured that every connected graph on n vertices admits a path decomposition of cardinality at most $\lceil n/2 \rceil$. Seminal results towards its verification consider the graph obtained from G by removing its vertices of odd degree, which is called the *E-subgraph* of G . Lovász (1968) verified Gallai's Conjecture for graphs whose E-subgraphs consist of at most one vertex, and Pyber (1996) verified it for graphs whose E-subgraphs are forests. In 2005, Fan verified Gallai's Conjecture for graphs in which each block of their E-subgraph is triangle-free and has maximum degree at most 3. Let \mathcal{G} be the family of graphs for which (i) each block has maximum degree at most 3; and (ii) each component either has maximum degree at most 3 or has at most one block that contains triangles. In this paper, we generalize Fan's result by verifying Gallai's Conjecture for graphs whose E-subgraphs are subgraphs of graphs in \mathcal{G} . This allows the components of the E-subgraphs to contain any number of blocks with triangles as long as they are subgraphs of graphs in \mathcal{G} .

Keywords: Graph, path, decomposition, Gallai's Conjecture, E-subgraph, maximum degree.

1 Introduction

In this paper, all graphs considered are finite and simple, i.e., contain a finite number of vertices and edges and have neither loops nor multiple edges. A *path decomposition* \mathcal{D} of a graph G is a collection of edge-disjoint paths of G that covers all the edges of G . A path decomposition \mathcal{D} of a graph G is *minimum* if for every path decomposition \mathcal{D}' of G we have $|\mathcal{D}| \leq |\mathcal{D}'|$, and the cardinality of such a minimum path decomposition, denoted by $\text{pn}(G)$, is called the *path number* of G . Gallai proposed the following conjecture (see [2, 13]).

*This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. F. Botler is partially supported by CNPq (Grant 423395/2018-1). M. Sambinelli is supported by FAPESP (Grant 2017/23623-4) and CNPq (Grant 423833/2018-9). E-mails: fbotler@cos.ufrj.br (F. Botler), m.sambinelli@ufabc.edu.br (M. Sambinelli).

Conjecture 1 (Gallai, 1968). *If G is a connected simple graph on n vertices, then $\text{pn}(G) \leq \lfloor n/2 \rfloor$.*

Lovász [13] verified Conjecture 1 for graphs that have at most one vertex with even degree. Pyber [14] extended Lovász's result by proving that Conjecture 1 holds for graphs in which each cycle contains at least one vertex with odd degree. In 2005, Fan [9] strengthened these results by extending Lovász's technique (see Section 2). Given a graph G , the *even subgraph* of G (*E-subgraph*, for short), denoted by $EV(G)$, is the graph obtained from G by removing its vertices with odd degree, or, equivalently, the subgraph of G induced by its even degree vertices. A *block* in a graph G is a maximal 2-connected subgraph of G , and a *leaf block* of G is a block that contains at most one cut-vertex of G . Thus, the results above may be restated as follows.

Theorem 2 (Lovász, 1968; Pyber, 1996; Fan, 2005). *Let G be a connected graph on n vertices.*

- (a) *If $EV(G)$ contains at most one vertex, then $\text{pn}(G) \leq \lfloor n/2 \rfloor$;*
- (b) *If $EV(G)$ is a forest, then $\text{pn}(G) \leq \lfloor n/2 \rfloor$;*
- (c) *If each block of $EV(G)$ is triangle-free and has maximum degree at most 3, then $\text{pn}(G) \leq \lfloor n/2 \rfloor$.*

Given a vertex u with even degree in G , the degree $d_{EV(G)}(u)$ is often referred as the *E-degree* of u . Consider a graph G for which each block of $EV(G)$ has maximum degree at most 3 (and that may contain triangles), and that has a minimum number of edges. One can show that each leaf block of $EV(G)$ must be a triangle, this is a key idea in the proof of Theorem 2(c). In this paper we extend Theorem 2(c) by presenting a strategy for dealing with these remaining triangles in the following special case. Let \mathcal{G} denote the family of graphs for which (i) each block has maximum degree at most 3; and (ii) each component either has maximum degree at most 3 or has at most one block that contains triangles. Note that \mathcal{G} contains (properly) the E-subgraphs of the graphs considered in [9]. We verify Conjecture 1 for graphs whose E-subgraphs are subgraphs of graphs in \mathcal{G} .

Theorem 3. *If G is a connected graph on n vertices such that $EV(G)$ is a subgraph of a graph in \mathcal{G} , then $\text{pn}(G) \leq \lfloor n/2 \rfloor$.*

We remark that Theorem 3 extends Theorem 2(c) by allowing some blocks to contain triangles. More specifically, a component of the E-subgraph may contain any number of blocks that contain triangles as long as it is a subgraph of graph in \mathcal{G} (see Figure 1).

Conjecture 1 has been deeply explored, and the literature indicating its correctness include results for Eulerian graphs with maximum degree at most 4 [10]; a family of regular graphs [5]; a family of triangle-free graphs [12]; and maximal outerplanar graphs and 2-connected outerplanar graphs [11]. Recent results were obtained by Bonamy and Perrett [1] who verified Conjecture 1 for graphs with maximum degree at most 5.

Note that the results in Theorem 2 give a bound of $\lfloor n/2 \rfloor$ for the graphs studied, which is slightly different from the bound of $\lceil n/2 \rceil$ proposed by Gallai. A straightforward condition for a graph on n vertices not to admit the former bound is to have sufficiently many edges. More precisely, if $|E(G)| > \lfloor n/2 \rfloor(n-1)$, then we have $\text{pn}(G) \geq \lfloor n/2 \rfloor$. In this case, n must be an odd integer. Such graphs are known as *odd semi-cliques* [1]. This motivates the following strengthening of Conjecture 1, which was considered in [6, 7].




Figure 1: Examples of graphs in which more than one block contain triangles, and that may be completed (by using the dotted lines) to graphs in \mathcal{G} .

Conjecture 4. *If G is a connected simple graph on n vertices, then either $\text{pn}(G) \leq \lfloor n/2 \rfloor$, or $\text{pn}(G) = \lceil n/2 \rceil$ and G is an odd semi-clique.*

Graphs G for which $\text{pn}(G) \leq \lfloor n/2 \rfloor$ are called *Gallai graphs*. Botler, Coelho, Lee, and Sambinelli [7] verified Conjecture 4 for graphs with treewidth at most 3 by proving that a partial 3-tree are either Gallai graphs, or one of the two odd semi-cliques that are partial 3-trees (K_3 and $K_5 - e$). They also prove [4] that a graph with maximum degree at most 4 is either a Gallai graph, or one of the three odd semi-cliques with maximum degree at most 4 (K_3 , $K_5 - e$, and K_5). More recently, Botler, Jiménez, and Sambinelli [6] verified Conjecture 4 for triangle-free planar graphs by proving that every such graph is a Gallai graph. In this paper, we explore an intermediate statement between Conjectures 1 and 4. We prove that, for the classes of graphs studied, all graphs are Gallai graphs except for a special family \mathcal{S} . The family \mathcal{S} , which we call the *SET graphs* (see Section 3), and for which we check Conjecture 1, differs from previous families of exceptions for two reasons. First, \mathcal{S} contains an infinite number of odd semi-cliques. Second, apart from containing odd semi-cliques, \mathcal{S} also contains non odd semi-cliques which we cannot guarantee the bound of $\lfloor n/2 \rfloor$. We remark that this is the first result regarding E-subgraphs to handle odd semi-cliques. Our result can be more specifically stated as follows.

Theorem 5. *If G is a connected graph such that $EV(G)$ is a subgraph of a graph in \mathcal{G} , then G is a Gallai graph or $G \in \mathcal{S}$.*

This work is organized as follows. In Section 2 we present some technical lemmas. In Section 3, we verify Conjecture 1 for the special case of graphs whose E-subgraphs have maximum degree at most 3; In Section 4, we verify Conjecture 1 for graphs G for which $EV(G)$ is a subgraph of a graph in \mathcal{G} ; and in Section 5, we present some concluding remarks.

Notation. The basic terminology and notation used in this paper are standard (see, e.g. [3]). Given a graph G , we denote its vertex set by $V(G)$ and its edge set by $E(G)$. The set of neighbors of a vertex u in a graph G is denoted by $N_G(u)$ and its degree by $d_G(u)$. When G is clear from the context, we simply write $N(u)$ and $d(u)$. Since G is simple, we always have $d_G(u) = |N_G(u)|$.

A graph H is a *subgraph* of a graph G , denoted by $H \subseteq G$, if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. Given a set of vertices $X \subseteq V(G)$, we say that H is the subgraph of

G induced by X , denoted by $G[X]$, if $V(H) = X$ and $E(H) = \{xy \in E(G) : x, y \in X\}$. Given a set of edges $Y \subseteq E(G)$, we say that H is the subgraph of G induced by Y , denoted by $G[Y]$, if $E(H) = Y$ and $V(H) = \{x \in V(G) : xy \in Y\}$. For ease of notation, when convenient, we write simply Y to refer to the graph $G[Y]$. Given $X \subseteq V(G)$, we define $G - X = G[V(G) \setminus X]$. In the case that $X = \{u\}$, we simply write $G - u$. Given a set Y of edges, we define the graphs $G \setminus Y = (V(G), E(G) \setminus Y)$. As before, in the case that $Y = \{e\}$, we simply write $G \setminus e$.

A *path* P in a graph G is a sequence $u_0u_1 \cdots u_\ell$ of distinct vertices in $V(G)$ such that $u_iu_{i+1} \in E(G)$, for $i = 0, \dots, \ell - 1$. We say that u_0 and u_ℓ are the *end vertices* of P , and that P *joins* u_0 and u_ℓ . When convenient, we consider a path as the subgraph of G induced by the set of edges $\{u_iu_{i+1} : i = 0, \dots, \ell - 1\}$. A *shortest* path joining two vertices u and v is a path that joins u and v with a minimum number of edges.

Given a vertex u of a graph G , we say that u is an odd (resp. even) vertex if its degree is odd (resp. even). Analogously, we say that a neighbor v of u is an odd (resp. even) neighbor of u if v has odd (resp. even) degree. Given a path decomposition \mathcal{D} of a graph G and a vertex $u \in V(G)$, we denote by $\mathcal{D}(u)$ the number of paths in \mathcal{D} that have u as an end vertex. It is not hard to check that $\mathcal{D}(u) \equiv d(u) \pmod{2}$. In particular, if u is an odd vertex, we have $\mathcal{D}(u) \geq 1$, for any path decomposition \mathcal{D} of G .

2 Technical Lemmas

In this section we present some technical results used throughout our proof. Following the strategy presented by Fan [9], our technique relies in the following definition.

Definition 1. Let u be a vertex in a graph G and let B be a set of edges incident to u . Let $G' = G \setminus B$, let \mathcal{D}' be a path decomposition of G' , and let $A = \{ux_i : 1 \leq i \leq k\}$ be a subset of B . We say that A is *addible towards* u (resp. *addible outwards* u) *with respect to* \mathcal{D}' if $G' + A$ admits a path decomposition \mathcal{D} such that

- (i) $|\mathcal{D}| = |\mathcal{D}'|$;
- (ii) $\mathcal{D}(u) = \mathcal{D}'(u) + |A|$ and $\mathcal{D}(x_i) = \mathcal{D}'(x_i) - 1$, for $1 \leq i \leq k$
(resp. $\mathcal{D}(u) = \mathcal{D}'(u) - |A|$ and $\mathcal{D}(x_i) = \mathcal{D}'(x_i) + 1$, for $1 \leq i \leq k$);
- (iii) $\mathcal{D}(v) = \mathcal{D}'(v)$ for each $v \in V(G) \setminus \{u, x_1, \dots, x_k\}$.

In these cases, we say that \mathcal{D} is an *A-transformation* of \mathcal{D}' *towards* (resp. *outwards*) u . For simplifying the notation, when $k = 1$, we write ux_1 -transformation instead of $\{ux_1\}$ -transformation.

Definition 1 is an extension of the definition of addible [9, Definition 3.2]. In fact, the present definition of *addible towards* precisely matches the definition of *addible at* given by Fan [9]. The next observation is used frequently in our proof.

Remark 1. Let B be a set of edges incident to a vertex u of a graph G . Let \mathcal{D}' be a path decomposition of $G' = G \setminus B$, and $A \subset B$ be an addible set towards (resp. outwards) u with respect to \mathcal{D}' , and let \mathcal{D}'' be an *A-transformation* of \mathcal{D}' . If $A' \subset B \setminus A$ is addible towards (resp. outwards) u with respect to \mathcal{D}'' , then $A \cup A'$ is addible towards (resp. outwards) u with respect to \mathcal{D}' .

The next two lemmas are results of Fan [9, Lemmas 3.4 and 3.6].

Lemma 6. *Let G be a graph and $uv \in E(G)$. Suppose that \mathcal{D}' is a path decomposition of $G' = G \setminus uv$. If $\mathcal{D}'(v) > |\{w \in N_{G'}(u) : \mathcal{D}'(w) = 0\}|$, then uv is addible towards u with respect to \mathcal{D}' .*

Lemma 6 motivates the following definitions. Let \mathcal{D} be a path decomposition of a graph G and let u be a vertex of G . We say that a vertex u is *passing* in \mathcal{D} if $\mathcal{D}(u) = 0$. Thus, we say that vertices in $\{v \in N_{G'}(u) : \mathcal{D}'(v) = 0\}$ are the *passing neighbors* of u in \mathcal{D}' . Lemma 6 then says that if in a path decomposition \mathcal{D}' of $G \setminus uv$ there are more paths having v as end vertex than passing neighbors of u , then uv is addible towards u with respect to \mathcal{D}' .

Lemma 7. *Let u be a vertex in a graph G and let $G' = G \setminus \{ux_1, \dots, ux_h\}$, where $x_i \in N_G(u)$. Suppose that \mathcal{D}' is a path decomposition of G' with $\mathcal{D}'(v) \geq 1$ for every $v \in N_{G'}(u)$. Then, for any $x \in \{x_1, \dots, x_h\}$, there is $A \subseteq \{ux_1, \dots, ux_h\}$ such that*

- $ux \in A$ and $|A| \geq \lceil \frac{h}{2} \rceil$; and
- A is addible towards u with respect to \mathcal{D}'

Given a graph G and a set M of edges of G , we denote by $G[M]$ the subgraph of G induced by the vertices incident to edges of M . We say that a set of edges $M \subseteq E(G)$ is an *induced matching* if the set edges of $G[M]$ is precisely M . Note that every subset of an induced matching is also an induced matching. The next two lemmas are used in the proof of Claim 5.

Lemma 8. *Let G be a graph and let $M = \{e_1, \dots, e_k\}$ be an induced matching in G , where $e_i = u_i v_i$ for $i = 1, \dots, k$. Let \mathcal{D}' be a path decomposition of $G' = G \setminus M$. If, for $i = 1, \dots, k$, the vertex u_i has no passing neighbor in \mathcal{D}' and $\mathcal{D}'(v_i) \geq 1$, then there is a path decomposition \mathcal{D} of G such that*

- $|\mathcal{D}| = |\mathcal{D}'|$;
- for $i = 1, \dots, k$, we have $\mathcal{D}(u_i) = \mathcal{D}'(u_i) + 1$ and $\mathcal{D}(v_i) = \mathcal{D}'(v_i) - 1$; and
- $\mathcal{D}(w) = \mathcal{D}'(w)$ for every $w \in V(G) \setminus V(M)$.

Proof. The proof follows by induction on the size of M . If $k = 0$, then $M = \emptyset$, and the statement holds with $\mathcal{D} = \mathcal{D}'$. Thus, assume $k \geq 1$. Since u_1 has no passing neighbor in \mathcal{D}' , and $\mathcal{D}'(v_1) \geq 1$, by Lemma 6, e_1 is addible towards u_1 with respect to \mathcal{D}' . Let $G'' = G' + e_1$, and let \mathcal{D}'' be the e_1 -transformation of \mathcal{D}' towards u_1 . Note that $\mathcal{D}''(u_1) = \mathcal{D}'(u_1) + 1$, $\mathcal{D}''(v_1) = \mathcal{D}'(v_1) - 1$, and $\mathcal{D}''(w) = \mathcal{D}'(w)$, for every $w \in V(G) \setminus \{u_1, v_1\}$. Thus, v_1 is the only vertex that possibly became a passing vertex through the transformation from \mathcal{D}' to \mathcal{D}'' . Now, consider $M' = M \setminus \{e_1\} = \{e_2, \dots, e_k\}$. Clearly, M' is an induced matching and $G'' = G \setminus M'$. Since $\{e_1, \dots, e_k\}$ is an induced matching, v_1 is not adjacent to u_2, \dots, u_k in G'' . Thus u_i has no passing neighbor in \mathcal{D}'' and $\mathcal{D}''(v_i) \geq 1$, for $i = 2, \dots, k$, and hence, by induction hypothesis, there is a path decomposition \mathcal{D} of G such that $|\mathcal{D}| = |\mathcal{D}''| = |\mathcal{D}'|$; for $i = 2, \dots, k$, we have $\mathcal{D}(u_i) = \mathcal{D}''(u_i) + 1 = \mathcal{D}'(u_i) + 1$ and $\mathcal{D}(v_i) = \mathcal{D}''(v_i) - 1 = \mathcal{D}'(v_i) - 1$; and $\mathcal{D}(w) = \mathcal{D}''(w) = \mathcal{D}'(w)$ for every $w \in V(G) \setminus \{u_i, v_i : i = 2, \dots, k\}$. Note that $\mathcal{D}(u_1) = \mathcal{D}''(u_1) = \mathcal{D}'(u_1) + 1$ and $\mathcal{D}(v_1) = \mathcal{D}''(v_1) = \mathcal{D}'(v_1) - 1$, and $\mathcal{D}(w) = \mathcal{D}''(w) = \mathcal{D}'(w)$ for every $w \notin \{u_i, v_i : i = 1, \dots, k\}$, as desired. \square

It is not hard to check that the condition that $\{e_1, \dots, e_k\}$ is an induced matching in Lemma 8 may be easily replaced by the condition that $N(u_i) \cap \{v_j: j = 1, \dots, k\} = \{v_j\}$, i.e., u_i 's may be adjacent to other u_i 's and v_i 's may be adjacent to other v_i 's, but each u_i is adjacent to precisely one v_j , namely, v_i .

In what follows, given a graph G and a vertex u , we denote by $E_G(u)$ the set of edges of G that are incident to u . The next lemma says, roughly, that after removing a set $E' \subseteq E_G(u)$ of edges incident to an odd vertex u , and applying Lemma 7, any further transformation outward u leaves more paths having u as an end vertex, than the number of edges that remain to add.

Lemma 9. *Let u be an odd vertex in a graph G , $B \subseteq E_G(u)$, and \mathcal{D}_0 be a path decomposition of $G \setminus B$. Let $A_1 \subseteq B$ be an addible set towards u with respect to \mathcal{D}_0 , and let $A_2 \subseteq B \setminus A_1$ be an addible set outwards u with respect to the A_1 -transformation \mathcal{D}_1 of \mathcal{D}_0 . If \mathcal{D}_2 is the A_2 -transformation of \mathcal{D}_1 and $|A_1| \geq \lceil |B|/2 \rceil$, then $\mathcal{D}_2(u) > |B| - |A_1 \cup A_2|$.*

Proof. We claim that $\mathcal{D}_1(u) \geq 1 + \lfloor |B|/2 \rfloor$. Indeed, since $d_G(u)$ is odd, if $|B|$ is even, then $\mathcal{D}_0(u) \geq 1$, and $\mathcal{D}_1(u) \geq \mathcal{D}_0(u) + |B|/2 \geq 1 + |B|/2 = 1 + \lfloor |B|/2 \rfloor$; if $|B|$ is odd, then $\mathcal{D}_1(u) \geq \lceil |B|/2 \rceil = 1 + \lfloor |B|/2 \rfloor$. Note that $\mathcal{D}_2(u) = \mathcal{D}_1(u) - |A_2|$. Since $|B| = \lfloor |B|/2 \rfloor + \lceil |B|/2 \rceil$, we have

$$\begin{aligned} \mathcal{D}_2(u) &= \mathcal{D}_1(u) - |A_2| \\ &\geq 1 + \lfloor |B|/2 \rfloor - |A_2| \\ &= 1 + |B| - \lceil |B|/2 \rceil - |A_2| \\ &\geq 1 + |B| - |A_1| - |A_2|. \end{aligned}$$

□

2.1 Absorbing lemmas

In this section, we present some lemmas that allow Gallai graphs to absorb special subgraphs while keeping its Gallai property. The next remark is used often in our proofs.

Remark 2. *If every component of a graph G is a Gallai graph, then G is a Gallai graph.*

Proof. Let H_1, \dots, H_k be the components of G . Thus $\text{pn}(G) = \sum_{i=1}^k \text{pn}(H_i) \leq \sum_{i=1}^k \lfloor |V(H_i)|/2 \rfloor \leq \lfloor |V(G)|/2 \rfloor$. Therefore G is a Gallai graph. □

We say that a graph G is a *single even triangle graph (SET graph)* if $EV(G)$ is a triangle and every odd vertex of G has at least two even neighbors. Since a SET graph has three even vertices, it must have odd order. Single even triangle graphs are special cases of our proof. Note that the graphs obtained from a complete graph on $2k+1$ vertices by removing a matching of size $k-1$ is a SET graph that is an odd semi-clique. Although we were not able to fully characterize which SET graphs are Gallai graphs, checking the validity of Conjecture 1 for them is a straightforward task (see Lemma 10). In what follows, we extend the definition of SET graphs, and indicate some of their vertices to be *connection vertices*. We say that a graph K is an *extended single even triangle graph (ESET graph)* if one of the following hold.

- i) K is a SET graph. In this case, every vertex of K is a *connection vertex*;

- ii) K is obtained from a SET graph K^- by adding a new vertex z adjacent to an odd and an even vertex of K^- . In this case, z is the *connection* vertex of K .

We say that an ESET is of *type i* or *ii*, according to the items above. Throughout the proof we are required to absorb ESET graphs without increasing the size of the path decomposition. The next lemma provide, for ESET graphs, path decompositions that contain two paths that have a fixed connection vertex as end vertex.

Lemma 10. *Let K be an ESET graph on n vertices. If u is a connection vertex of K , then K admits a path decomposition \mathcal{D} of K such that $\mathcal{D}(u) \geq 2$ and $|\mathcal{D}| \leq \lceil n/2 \rceil$.*

Proof. Let K , n , and u be as in the statement. We divide the proof according to the type of K .

Type i. By the definition of ESET graph of type *i*, K is a SET graph and u is any of its vertices. Let S be the set of edges that join u to the even vertices of K , and let $K' = K \setminus S$. Note that K' has precisely one even vertex, and hence, by Theorem 2(a), it follows that $\text{pn}(K') \leq \lfloor n/2 \rfloor$. Let \mathcal{D}' be a minimum path decomposition of K' . Note that every neighbor of u in K has odd degree in K' . Thus, by Lemma 7, there is $B \subseteq S$, with $|B| \geq |S|/2$, which is addible with respect to \mathcal{D}' . Let \mathcal{D}'' be the B -transformation of \mathcal{D}' towards u . Note that $S \setminus B$ contains at most one edge. If $S \setminus B = \{e\}$, then let P be the path containing only e , and put $\mathcal{D} = \mathcal{D}'' \cup \{P\}$, otherwise, put $\mathcal{D} = \mathcal{D}''$. It is not hard to check that $\mathcal{D}(u) \geq 2$ and $|\mathcal{D}| \leq \lceil n/2 \rceil$, as desired.

Type ii. By the definition of ESET graph of type *ii*, K is obtained from a SET graph K^- by adding a new vertex u adjacent to an odd vertex x and an even vertex y of K^- . Since K^- is an ESET of type *i*, K^- admits a path decomposition \mathcal{D}^- such that $\mathcal{D}^-(y) \geq 2$ and $|\mathcal{D}^-| \leq \lceil (n-1)/2 \rceil$. Since x is an odd vertex of K^- , we have $\mathcal{D}^-(x) \geq 1$. Let P_x^- be a path in \mathcal{D}^- that has x as end vertex. Since $\mathcal{D}^-(y) \geq 2$, there are two paths in \mathcal{D}^- that have y as end vertex, and hence there is at least one such path, say P_y^- , that is different from P_x^- . Let $P_x = P_x^- + ux$ and $P_y = P_y^- + yu$, and hence $(\mathcal{D}^- \setminus \{P_x^-, P_y^-\}) \cup \{P_x, P_y\}$ is the desired decomposition. \square

As a direct application of Lemma 10, we have the following result.

Lemma 11. *Let G be a graph that can be decomposed into an ESET graph K on n vertices with a connection vertex $u \in V(K)$, and a path P such that $V(P) \cap V(K) = \{u\}$. Then $\text{pn}(G) \leq \lceil n/2 \rceil$.*

Proof. Let G , K , P , n , and u be as in the statement. By Lemma 10, K admits a path decomposition \mathcal{D}_K such that $\mathcal{D}_K(u) \geq 2$ and $|\mathcal{D}_K| \leq \lceil n/2 \rceil$. Let Q_1 and Q_2 be two paths in \mathcal{D}_K having u as end vertex. Decompose P into two paths P_1 and P_2 having u as end vertex. Let $R_1 = P_1 \cup Q_1$ and $R_2 = P_2 \cup Q_2$, and let $\mathcal{D} = (\mathcal{D}_K \setminus \{Q_1, Q_2\}) \cup \{R_1, R_2\}$. Clearly, \mathcal{D} is a path decomposition of G with $|\mathcal{D}| \leq \lceil n/2 \rceil$. Therefore, $\text{pn}(G) \leq \lceil n/2 \rceil$. \square

Let K , G' , and $G = K \cup G'$ be graphs. We say that K is a *hanging ESET subgraph* of G if K is an ESET graph, $V(K) \cap V(G') = \{u\}$, and u is a connection vertex of K .

Lemma 12. *Let G be a graph that contains a hanging ESET subgraph K , and let G' be such that $G = K \cup G'$ and $V(K) \cap V(G') = \{u\}$. Then $\text{pn}(G) \leq \lceil |V(K)|/2 \rceil + \text{pn}(G') - 1$.*

Proof. Let G , K , G' , and u be as in the statement. Let \mathcal{D}' be a path decomposition of G' . Let P be an element of \mathcal{D}' that contains u and let $H = K \cup P$. By Lemma 11, we have $\text{pn}(H) \leq \lceil |V(K)|/2 \rceil$. Let \mathcal{D}_H be a minimum path decomposition of H , and note that $\mathcal{D}_H \cup (\mathcal{D}' \setminus \{P\})$ is a path decomposition of G such that $|\mathcal{D}_H \cup (\mathcal{D}' \setminus \{P\})| \leq \lceil |V(K)|/2 \rceil + \text{pn}(G') - 1$, as desired. \square

3 Graphs with maximum E-degree at most 3

The strategy of the proof of the main theorem of this section is to show that the even subgraph of a minimal counterexample consists of vertex-disjoint triangles, and then proving that these graphs satisfy Gallai's Conjecture.

Theorem 13. *If G is a connected simple graph on n vertices such that $\Delta(EV(G)) \leq 3$, then either G is a Gallai graph or G is a SET graph.*

Proof. Suppose that the statement does not hold, and let G be a counterexample on n vertices with $\Delta(EV(G)) \leq 3$ and which minimizes $|E(G)|$. In what follows, we prove a few claims regarding G . First, we prove that every hanging ESET subgraph of G must be connected at a special vertex.

Claim 1. *Let K be a hanging ESET subgraph of G , and let G' be such that $G = K \cup G'$, and $V(K) \cap V(G') = \{u\}$. Then u is an odd vertex of G and an even vertex of G' .*

Proof. Let K , G' , and u be as in the statement. Suppose, for a contradiction, that u has even degree in G or odd degree in G' . Then $EV(G') \subseteq EV(G)$, and hence $\Delta(EV(G')) \leq 3$. By the minimality of G , the graph G' is either a Gallai graph or a SET graph. First, suppose that G' is a Gallai graph, i.e., $\text{pn}(G') \leq \lfloor |V(G')|/2 \rfloor$. By Lemma 12, it follows that

$$\begin{aligned} \text{pn}(G) &\leq \lceil |V(K)|/2 \rceil + \lfloor |V(G')|/2 \rfloor - 1 \\ &\leq (|V(K)| + 1)/2 + |V(G')|/2 - 1 \\ &= (|V(K)| + |V(G')| - 1)/2 \\ &= |V(G)|/2. \end{aligned}$$

Therefore, $\text{pn}(G) \leq \lfloor |V(G)|/2 \rfloor$, and G is a Gallai graph, a contradiction.

Thus, we may assume that G' is a SET graph. By Lemma 10, K (resp. G') admits a path decomposition \mathcal{D}_K (resp. \mathcal{D}') such that $\mathcal{D}_K(u) \geq 2$ and $|\mathcal{D}_K| \leq \lceil |V(K)|/2 \rceil$ (resp. $\mathcal{D}'(u) \geq 2$ and $|\mathcal{D}'| \leq \lceil |V(G')|/2 \rceil$). Let P_1 and P_2 be paths in \mathcal{D}_K having u as end vertices, and Q_1 and Q_2 be paths in \mathcal{D}' having u as end vertices. Put $R_1 = P_1 \cup Q_1$ and $R_2 = P_2 \cup Q_2$, and note that $\mathcal{D} = (\mathcal{D}_K \setminus \{P_1, P_2\}) \cup (\mathcal{D}' \setminus \{Q_1, Q_2\}) \cup \{R_1, R_2\}$ is a path

decomposition of G with cardinality

$$\begin{aligned}
|\mathcal{D}| &\leq \lceil |V(K)|/2 \rceil + \lceil |V(G')|/2 \rceil - 2 \\
&\leq (|V(K)| + 1)/2 + (|V(G')| + 1)/2 - 2 \\
&\leq (|V(K)| + |V(G')| - 2)/2 \\
&< (|V(K)| + |V(G')| - 1)/2 \\
&= |V(G)|/2.
\end{aligned}$$

Therefore, $\text{pn}(G) \leq \lfloor |V(G)|/2 \rfloor$, and G is a Gallai graph, a contradiction. \square

Now we use Fan's techniques to prove that $EV(G)$ consists of vertex-disjoint triangles. First, we prove that no vertex of G has a unique even neighbor.

Claim 2. *No vertex of G has exactly one even neighbor.*

Proof. Suppose, for a contradiction, that G contains a vertex u that has precisely one even neighbor, say v , and let $G' = G \setminus uv$. Note that v has odd degree in G' and u has no even neighbor in G' . Therefore, $\Delta(EV(G')) \leq 3$. We claim that G' is a Gallai graph. By Remark 2, it is enough to prove that no component of G' is a SET graph. Indeed, G' has at most two components, say G'_u and G'_v , that contain, respectively, u and v . If u is an odd vertex of G'_u and G'_u is a SET graph, then u must be adjacent to at least two even vertices of G'_u , say x, y . But, in this case, x and y have even degree in G , and hence u has at least three even neighbors in G , namely v, x , and y , a contradiction. Thus, we may assume that u is an even vertex of G'_u and an odd vertex of G , and hence u is an isolated even vertex of G' . Thus, $EV(G'_u)$ is not a triangle, and hence G'_u is not a SET graph. Thus, if G' is connected, i.e., $G'_u = G'_v$, then G' is a Gallai graph, as desired. Thus, we may assume $G'_u \neq G'_v$. In this case, note that, if G'_v is a SET graph, then G'_v is a hanging ESET subgraph of G connected at v , which is an even vertex of G , a contradiction to Claim 1. Thus, G'_u and G'_v are Gallai graphs as desired.

Let \mathcal{D}' be a minimum path decomposition of G' . Since v has odd degree in G' , it follows $\mathcal{D}'(v) \geq 1$, and since u has no even neighbor in G' , we have $|\{x \in N_{G'}(u) : \mathcal{D}'(x) = 0\}| = 0$. Thus, by Lemma 6, uv is addible towards u with respect to \mathcal{D}' . Thus, $\text{pn}(G) \leq |\mathcal{D}'| = \text{pn}(G')$, and hence G is a Gallai graph, a contradiction. \square

Definition 2. Let G be a graph, and let F be a non-empty subgraph of G with components F_1, F_2, \dots, F_ℓ . We say that F is a *Fan subgraph* if, for $j = 2, \dots, \ell$, the graph F_j consists of a single edge joining even vertices of G , and F_1 is either the null graph (graph with an empty set of vertices) or the following hold.

- (i) F_1 is a star with center at a vertex u and at least two leaves v_1, v_2, \dots, v_k and, for $i \geq 2$, the vertex v_i is even in G ;
- (ii) u has no even neighbor in $G \setminus E(F)$; and
- (iii) if v_1 is odd in G , then u is odd in G and each component of $EV(G)$ is a triangle.

Claim 3. *Let F be a Fan subgraph of G and let $G' = G \setminus E(F)$. If $\Delta(EV(G')) \leq 3$, then G' is a Gallai graph.*

Proof. Let F and G' be as in the statement. By Remark 2, it is enough to prove that no component of G' is a SET graph. Thus, let H' be a component of G' , and suppose, for a contradiction, that H' is a SET graph. Let $E = \{x_1, x_2, x_3\}$ and $O = V(H') \setminus E = \{y_1, \dots, y_t\}$ be, respectively, the set of even and odd vertices of H' . Since G is connected, it follows that $V(H') \cap V(F) \neq \emptyset$.

Let F_1, F_2, \dots, F_ℓ be the components of F as in Definition 2 where F_1 is either the null graph or a star. First, suppose that F_1 is the null graph. Since, by Definition 2, for $j = 2, \dots, \ell$, F_j consists of a single edge joining even vertices of G , every vertex of F has odd degree in G' , which implies that $V(F) \cap V(H') \subseteq O$. Thus, x_1, x_2, x_3 have even degree in G . Suppose that there are two vertices, say y and y' , in $V(F) \cap V(H')$. By the definition of SET graph, y and y' are each adjacent to at least two vertices in E . Therefore, y and y' have a common neighbor, say x_1 , in E . But then x_2, x_3, y, y' are four even neighbors of x_1 in G , a contradiction to $\Delta(EV(G)) \leq 3$. Thus, we may assume that there is precisely one vertex, say y , in $V(F) \cap V(H')$. In this case, H' is a hanging ESET (of type i) connected at y , which is an even vertex of G , a contradiction to Claim 1.

Thus, we may assume that F_1 is a star, and let u be its center and v_1, v_2, \dots, v_k be its leaves. By Definition 2(i), we may assume that v_i is even in G for $i \geq 2$. Also, by Definition 2, we have $k \geq 2$ and, for $j \geq 2$, F_j consists of a single edge joining even vertices of G . This implies that the only vertices of F that may have even degree in G' are u and v_1 , and hence $V(F) \cap E \subseteq \{u, v_1\}$. We may assume that $u \notin O$, otherwise, by the definition of SET graph, u would have at least two even neighbors in G' , which contradicts Definition 2(ii). In what follows, we divide the proof into three cases, depending on whether $\{u, v_1\} \cap E = \emptyset$, $u \in E$, or $v_1 \in E$. Since $uv_1 \in E(F)$, we have $uv_1 \notin E(H')$, and hence the conditions $u \in E$ and $v_1 \in E$ are disjoint because $EV(H')$ is a triangle. Let L be the set of leaves of F , i.e., $L = V(F) \setminus \{u\}$, and let $L_E = \{v \in L: v \text{ has even degree in } G\}$. Moreover, note that

$$d_{G'}(w) = d_G(w) - 1 \text{ for every vertex } w \in L, \quad (1)$$

and that every vertex in L_E has odd degree in G' .

Suppose that $\{u, v_1\} \cap E = \emptyset$, and hence every vertex in E has even degree in G . Analogously to the case F_1 is the null graph, if there are two distinct vertices in $L \cap O$, then there is a vertex, say x_1 , in E with E-degree at least 4 in G , a contradiction to $\Delta(EV(G)) \leq 3$. Thus we may assume that $|L \cap O| \leq 1$. Thus $V(H') \cap V(F) = \{y_i\} \subset O$, for some $i = 1, \dots, t$, and hence H' is a hanging ESET subgraph (of type i) connected at y_i , which is an even vertex of G , a contradiction do Claim 1. Hence, we may assume that $\{u, v_1\} \cap E \neq \emptyset$.

Suppose that $u \in E$, and suppose, without loss of generality, that $u = x_1$. Since $v_1 \notin E$, x_2 and x_3 have even degree in G , a contradiction to Definition 2(ii). Thus, we may assume that $u \notin E$.

Now, suppose that $v_1 \in E$ and suppose, without loss of generality, that $v_1 = x_1$. As noted above, $u \notin E$, and hence x_2 and x_3 have even degree in G . Moreover, v_1 has odd degree in G , and hence by Definition 2(iii), u is odd in G and each component of $EV(G)$ is a triangle. If $L \cap O = \emptyset$, then H' is a hanging ESET subgraph (of type i) connected at v_1 , which is an odd vertex of G . Let G'' be such that $G = H' \cup G''$ and $V(H') \cap V(G'') = \{v_1\}$. Since $v_1 \in E$, the vertex v_1 has odd degree in G'' , a contradiction do Claim 1. Thus, we may assume that there exists a vertex $y \in L \cap O$, and hence by (1),

the vertex y have even degree in G . Since every vertex in O has at least two neighbors in E , we may assume, without loss of generality, that $yx_2 \in E(G)$. Therefore, $yx_3 \in E(G)$ because x_2, x_3, y are even in G , $\{x_2x_3, yx_2\} \subseteq E(G)$, and each component of $EV(G)$ is a triangle. In particular, x_2 and x_3 are the only even neighbors of y in G . Note that every vertex in O is adjacent to x_2 or x_3 . Thus, if there is a vertex in $y' \in L \cap O$ such that $y' \neq y$, then y' would have even degree in G and x_2, x_3, y, y' would belong to the same component of $EV(G)$, a contradiction. Therefore $V(H') \cap V(F) = \{x_1, y\}$. Suppose that $y \in V(F_j)$, for some $j \geq 2$, and let $V(F_j) = \{y, y'\}$. Then y' is an even neighbor of y in G , and hence $y' \in \{x_2, x_3\} \subseteq E$, a contradiction. Thus, we have $y = v_i$, for some $i \geq 2$, and $uy \in E(F_1)$. Note that $\{ux_1, uy\}$ is an edge cut of G , where H' is a component of $G - x_1v - yv$. In other words, $H'' = H' + ux_1 + uy$ is a hanging ESET subgraph (of type ii) connected at u , which is an odd vertex of G . Let G'' be such that $G = H'' \cup G''$ and $V(H'') \cap V(G'') = \{u\}$. Since $d_{H''}(u) = 2$, the vertex u has odd degree in G'' , a contradiction do Claim 1. \square

Claim 4. *Every component of $EV(G)$ is a triangle or an isolated vertex.*

Proof. Suppose, for a contradiction, that $EV(G)$ contains a component H which is neither a triangle nor an isolated vertex.

Subclaim 4.1. *Every vertex in H has degree 2.*

Proof. By Claim 2, no vertex of H has degree 1. Thus, we prove that H has maximum degree 2. Suppose, for a contradiction, that H has a vertex u with degree 3, and let $\{v_1, v_2, v_3\} \subseteq N_H(u)$ be three even neighbors of u . Let F be the subgraph of G induced by the edges uv_1, uv_2, uv_3 , and let $G_1 = G \setminus E(F)$. Note that every vertex in $V(F)$ has odd degree in G_1 and, hence, that $EV(G_1) \subseteq EV(G)$. Thus, it follows that $\Delta(EV(G_1)) \leq 3$. Note that F is a star with center at u and with at least two leaves, in which every leaf has even degree in G . Moreover, u has no odd neighbor in G_1 . Thus, F is a Fan subgraph. By Claim 3, we have $\text{pn}(G_1) \leq \lfloor n/2 \rfloor$. Let \mathcal{D}_1 be a minimum path decomposition of G_1 . Since the vertices of F have odd degree in G_1 , it follows that $\mathcal{D}_1(v) \geq 1$ for every $v \in V(F)$. By Lemma 7, there is a set $B \subseteq \{uv_1, uv_2, uv_3\}$ addible towards u with respect to \mathcal{D}_1 and containing at least two edges. Let \mathcal{D}_2 be the B -transformation of \mathcal{D}_1 towards u , and let $G_2 = G_1 + B$. If $|B| = 3$, then \mathcal{D}_2 is a path decomposition of G such that $|\mathcal{D}_2| = |\mathcal{D}_1| \leq \lfloor n/2 \rfloor$, a contradiction. Thus, we may assume $|B| = 2$. Suppose, without loss of generality that $uv_3 \notin B$. Note that $\mathcal{D}_2(u) \geq 3$, and that, since $d_{EV(G)}(v_3) \leq 3$, we have $d_{EV(G_2)}(v_3) \leq 2$, because u is not a neighbor of v_3 in G_1 . Thus, by Lemma 6, uv_3 is addible towards v_3 with respect to \mathcal{D}_2 , but the uv_3 -transformation \mathcal{D} of \mathcal{D}_2 towards v_3 is a path decomposition of G such that $|\mathcal{D}_3| \leq \lfloor n/2 \rfloor$, a contradiction. \square

Suppose that H is not a triangle, and let u be a vertex in H . Let v_1 and v_2 be the even neighbors of u . Since H is not a triangle, v_1 and v_2 are not adjacent. Let w be an even neighbor of v_2 different from u . Let F_1 be the null graph, $F_2 = G[\{u, v_1\}]$ and $F_3 = G[\{v_2, w\}]$, and let $F = F_1 \cup F_2 \cup F_3$. Let $G' = G \setminus E(F)$, and note that every vertex in $V(F)$ has odd degree in G' , and hence $EV(G') \subseteq EV(G)$. Thus $\Delta(EV(G')) \leq 3$. Since F_2 and F_3 consist of single edges joining even vertices of G , F is a Fan subgraph. By Claim 3, we have $\text{pn}(G') \leq \lfloor n/2 \rfloor$. Let \mathcal{D}' be a minimum path decomposition of G' . Note that v_2 has no even neighbors in G' . Thus, by Lemma 6, v_2w is addible towards v_2 with

respect to \mathcal{D}' . Let \mathcal{D}'' be the v_2w -transformation of \mathcal{D}' towards v_2 . Note that v_2 is the only even neighbor of u in $G' + v_2w$, but $\mathcal{D}''(v_2) \geq 2$. Thus, by Lemma 6, uv_1 is addible towards u with respect to \mathcal{D}'' , but the uv_1 -transformation \mathcal{D} of \mathcal{D}'' towards u is a path decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$, a contradiction. \square

In what follows, if x is a vertex of odd degree, and $T \subseteq EV(G)$ is a triangle containing a neighbor of x , we say that T is a *triangle neighbor* of x . Given a vertex v and a triangle T , we denote by $E_G(u, T)$ the set of edges in G joining v to a vertex in T . First, we prove that each odd vertex of G has even neighbors in at most one component of $EV(G)$, which implies that every odd vertex of G has at most one triangle neighbor. This implies that two vertices of even degree have a common neighbor if and only if they belong to the same (triangle) component of $EV(G)$. The proof of the next claim consists in applying Fan's technique on vertices of odd degree. For that, let T_1, \dots, T_s be the triangles in $EV(G)$.

Claim 5. *If $d_G(u)$ is odd, then u has even neighbors in at most one component of $EV(G)$.*

Proof. Suppose, for a contradiction, that u has even neighbors in at least two components of $EV(G)$. Let \mathcal{I} be the set of even neighbors of u which are isolated vertices in $EV(G)$. For $k = 1, 2, 3$, let \mathcal{T}_k be indexes of the triangle neighbors of u containing k neighbors of u and let $\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3$. For each $k \in \{1, 2, 3\}$ and $j \in \mathcal{T}_k$, let $V(T_j) = \{x_1^j, x_2^j, x_3^j\}$, where $x_i^j \in N_G(u)$ for $i \leq k$. In what follows, we remove a set of edges incident to u and of some triangles neighbors of u . For that, for $v \in \mathcal{I}$, let $\varphi(v) = \{uv\}$; for $j \in \mathcal{T}_1 \cup \mathcal{T}_2$, let $\varphi(T_j) = \{ux_1^j, x_2^j x_3^j\}$; and for $j \in \mathcal{T}_3$, let $\varphi(T_j) = \{ux_1^j, ux_2^j, ux_3^j\}$. Let $B_0 = \left(\bigcup_{v \in \mathcal{I}} \varphi(v) \right) \cup \left(\bigcup_{j \in \mathcal{T}} \varphi(T_j) \right)$ (see Figure 2) and let $B_0^u = E_G(u) \cap B_0$.

Subclaim 5.1. $\mathcal{T}_2 \neq \emptyset$ and $|B_0^u|$ is odd.

Proof. Suppose, for a contradiction, that $\mathcal{T}_2 = \emptyset$ or $|B_0^u|$ is even. Let $G_0 = G \setminus B_0$. Note that every vertex of $B_0 - u$ has odd degree in G_0 , and hence if $d_{G_0}(u)$ is even, then u is an isolated vertex in $EV(G_0)$. Thus, $\Delta(EV(G_0)) \leq 3$. Let H_1, H_2, \dots, H_ℓ be the components of B_0 , where H_1 is the subgraph of G edge-induced by B_0^u . By definition of B_0 , the graphs H_2, \dots, H_ℓ consist of single edges joining even vertices. Moreover, H_1 is a star with center at u with at least two leaves, since u has even neighbors in at least two components of $EV(G)$. In addition, the leaves of H_1 are even vertices of G , and every neighbor of u in G_0 is odd. Thus, B_0 is a Fan subgraph of G , and by Claim 3, $\text{pn}(G_0) \leq \lfloor n/2 \rfloor$. Let \mathcal{D}_0 be a minimum path decomposition of G_0 .




Figure 2: Illustration of the Fan subgraph B_0 in the proof of Subclaim 5.1. The edges of B_0 are highlighted in red.

Let $A_0 = \{x_2^j x_3^j : j \in \mathcal{T}_1 \cup \mathcal{T}_2\}$ and let $G_1 = G_0 + A_0$. Since every vertex in $B_0 - u \supset V(A_0)$ has odd degree in G_0 , it follows that $\mathcal{D}_0(v) \geq 1$ for every $v \in V(B_0) \setminus \{u\}$. Note that if $j \in \mathcal{T}_1$, then x_2^j and x_3^j are not adjacent to u (that may possibly have even degree in G_0), and hence x_2^j and x_3^j have no passing neighbor in \mathcal{D}_0 . Moreover, if $j \in \mathcal{T}_2$, then $\mathcal{T}_2 \neq \emptyset$ and, by hypothesis, $|B_0^u|$ is even; hence u has odd degree in G_0 . Thus, x_2^j and x_3^j have no passing neighbors in \mathcal{D}_0 , for $j \in \mathcal{T}_1 \cup \mathcal{T}_2$. Since A_0 is an induced matching in G_1 , it follows by Lemma 8 that there is a path decomposition \mathcal{D}_1 of G_1 such that $|\mathcal{D}_1| = |\mathcal{D}_0|$; for every $j \in \mathcal{T}_1 \cup \mathcal{T}_2$, we have $\mathcal{D}_1(x_2^j) = \mathcal{D}_0(x_2^j) + 1$ and $\mathcal{D}_1(x_3^j) = \mathcal{D}_0(x_3^j) - 1$; and $\mathcal{D}_1(v) = \mathcal{D}_0(v)$ for every $v \notin V(A_0)$.

For $v \in (V(B_0) \setminus \{u\}) \setminus V(A_0)$, it follows that $\mathcal{D}_1(v) = \mathcal{D}_0(v) \geq 1$. Moreover, for $j \in \mathcal{T}_1 \cup \mathcal{T}_2$, we have $\mathcal{D}_1(x_2^j) \geq 2$ and x_3^j is not a neighbor of u in G . Thus no neighbor of u in G is a passing vertex in \mathcal{D}_1 . Let $B_1 = E(G) \setminus E(G_1)$, and note that $B_1 = B_0^u$. By Lemma 7, there is an addible set $A_1 \subseteq B_1$ towards u with respect to \mathcal{D}_1 such that $|A_1| \geq \lceil |B_1|/2 \rceil$. Let $G_2 = G_1 + A_1$, \mathcal{D}_2 be the A_1 -transformation of \mathcal{D}_1 towards u , and $B_2 = E(G) \setminus E(G_2)$. Note that

$$\mathcal{D}_2(u) \geq \mathcal{D}_1(u) + \lceil |B_1|/2 \rceil. \quad (2)$$

Let $B_2^* = \{uv_j \in B_2 \cap E_G(u, T_j) : j \in \mathcal{T}_3 \text{ and } |E_{G_2}(u, T_j)| = 2\}$. Note that $\mathcal{D}_2(u) \geq 2|B_2^*|$. Let $A_2 \subseteq B_2^*$ be a maximal addible set outwards u with respect to \mathcal{D}_2 , $G_3 = G_2 + A_2$, \mathcal{D}_3 be the A_2 -transformation of \mathcal{D}_2 outwards u , and $B_3 = E(G) \setminus E(G_3)$. We show that $A_2 = B_2^*$. Suppose, for a contradiction, that $A_2 \subsetneq B_2^*$ (hence $B_2^* \setminus A_2 \neq \emptyset$ and $B_2^* \neq \emptyset$). Since $B_2^* \neq \emptyset$, it follows that $\mathcal{T}_3 \neq \emptyset$, and since u has neighbors in at least two components of $EV(G)$ in G , it follows that $|B_1| = |B_0^u| \geq 4$. We claim that $\mathcal{D}_2(u) \geq 3$. This is clear from Equation (2) if $|B_1|$ is odd. If $|B_1|$ is even, then u has odd degree in G_1 , and hence $\mathcal{D}_1(u) \geq 1$. By Equation (2), we have $\mathcal{D}_2(u) \geq 3$. Let $uv_j \in B_2^* \setminus A_2$. First, we show that $|A_2| \geq 1$. For that note that the only possible passing neighbors of v_j in \mathcal{D}_2 are the vertices in $V(T_j) \setminus \{v_j\}$. Since $\mathcal{D}_2(u) \geq 3$, it follows by Lemma 6 that $\{uv_j\}$ is an addible set towards v (i.e. outwards u) in \mathcal{D}_2 and, therefore, $|A_2| \geq 1$. Now, since $A_2 \subsetneq B_2^*$, we have, $|B_2^* \setminus A_2| \geq 1$, and hence,

$$\mathcal{D}_3(u) = \mathcal{D}_2(u) - |A_2| \geq 2|B_2^*| - |A_2| = 2(|A_2| + |B_2^* \setminus A_2|) - |A_2| \geq 3.$$

By Lemma 6, uv_j is addible towards v_j (i.e. outwards u) with respect to \mathcal{D}_3 , and hence, by Remark 1, the set $A_2 \cup \{uv_j\} \subseteq B_2^*$ is an addible set outwards u with respect to \mathcal{D}_2 , a contradiction to the maximality of A_2 . Therefore, $A_2 = B_2^*$.

Let $A_3 \subseteq B_3$ be a maximum addible set outwards u with respect to \mathcal{D}_3 , $G_4 = G_3 + A_3$, \mathcal{D}_4 be the A_3 -transformation outwards u with respect to \mathcal{D}_3 , and $B_4 = E(G) \setminus E(G_4)$. In what follows, we show that $B_4 = \emptyset$, and hence $G = G_4$. Suppose, for a contradiction, that $B_4 \neq \emptyset$. By Lemma 9, $\mathcal{D}_4(u) > |B_0^u| - |A_1| - |A_2| - |A_3| = |B_4|$, and since $|B_4| \geq 1$, it follows that $\mathcal{D}_4(u) \geq 2$. Let $uv \in B_4$, and note that v has at most one passing neighbor with respect to \mathcal{D}_4 in G_4 :

- if $v \in \mathcal{I}$, then every neighbor of v in G_4 has odd degree, and hence v has no passing neighbor with respect to \mathcal{D}_4 ;
- if $v = x_i^j$ for some $j \in \mathcal{T}_1 \cup \mathcal{T}_2$, then $i \leq 2$, and the only possible passing neighbor of x_i^j is x_3^j ; and

- if $v = x_i^j$ for some $j \in \mathcal{T}_3$, then $|E_{G_2}(u, T_j)| \leq 1$, otherwise $uv \in A_2$. Thus there is at most one passing vertex in T_j with respect to \mathcal{D}_4 and, therefore, x_i^j has at most one passing neighbor with respect to \mathcal{D}_4 .

Since $\mathcal{D}_4(u) \geq 2$, by Lemma 6, uv is addible towards v (i.e. outwards u) with respect to \mathcal{D}_4 , and hence by Remark 1, $A_3 \cup \{uv\}$ is an addible set outwards u with respect to \mathcal{D}_3 , a contradiction to the choice of A_3 . Therefore, $B_4 = \emptyset$ and $G_4 = G$. \square

By Subclaim 5.1, we have $\mathcal{T}_2 \neq \emptyset$ and $|B_0^u|$ is odd. Choose $z \in \mathcal{T}_2$, and define $\varphi'(T_z) = \{ux_1^z, ux_2^z\}$. Let $R_0 = \left(\bigcup_{v \in \mathcal{I}} \varphi(v)\right) \cup \left(\bigcup_{j \in \mathcal{T} \setminus \{z\}} \varphi(T_j)\right) \cup \varphi'(T_z)$ (see Figure 3), and let $R_0^u = R_0 \cap E_G(u)$. Note that $|R_0^u| = |B_0^u| + 1$, and hence $|R_0^u|$ is even. Let $G_0 = G \setminus R_0$, and note that every vertex in $V(R_0) \setminus \{x_3^z\}$ has odd degree in G_0 . Since no odd vertex of G is an even vertex of G_0 , we have $EV(G_0) \subseteq EV(G)$, and hence $\Delta(EV(G_0)) \leq 3$. Let H_1, \dots, H_ℓ be the components of R_0 , where H_1 is the subgraph of G induced by the edges in R_0^u . Note that H_1 is a star with center at u and at least four leaves, i.e., $|R_0^u| \geq 4$, since u has even neighbors in at least two components of $EV(G)$, $\mathcal{T}_2 \neq \emptyset$, and $|R_0^u|$ is even. Also, all the leaves of R_0^u are even in G , and u has no even neighbors in G_0 . Moreover, note that H_2, \dots, H_ℓ consist of single edges joining even vertices. Therefore, R_0 is a Fan subgraph, and by Claim 3, $\text{pn}(G_0) \leq \lfloor n/2 \rfloor$. Let \mathcal{D}_0 be a minimum path decomposition of G_0 .




Figure 3: Illustration of the Fan subgraph R_0 in the proof of Claim 5. The edges of R_0 are highlighted in red.

Note that every vertex $v \in V(R_0)$ has odd degree in G_0 , and hence $\mathcal{D}_0(v) \geq 1$. In particular, $\mathcal{D}_0(u) \geq 1$. Also, note that every vertex $v \in V(R_0) \setminus \{x_1^z, x_2^z\}$ has no neighbor with even degree in G_0 , and hence no passing neighbor in \mathcal{D}_0 . On the other hand, x_1^z and x_2^z has only one possible passing neighbor in \mathcal{D}_0 , namely the vertex x_3^z which has even degree in G_0 . Let $A_0 = \{x_2^z x_3^z : j \in \mathcal{T}_1 \cup \mathcal{T}_2 \setminus \{z\}\}$, $G_1 = G_0 + A_0$, and $R_1 = E(G) \setminus E(G_1)$ (remark that $R_1 = R_0^u$). Note that A_0 is an induced matching in G_1 . If $A_0 \neq \emptyset$, then let \mathcal{D}_1 be the path decomposition given by Lemma 8; otherwise let $\mathcal{D}_1 = \mathcal{D}_0$. In either case, it follows that \mathcal{D}_1 is a path decomposition of G_1 such that

- $|\mathcal{D}_1| = |\mathcal{D}_0|$;
- for all $j \in \mathcal{T}_1 \cup \mathcal{T}_2 \setminus \{z\}$, we have $\mathcal{D}_1(x_2^j) = \mathcal{D}_0(x_2^j) + 1$ and $\mathcal{D}_1(x_3^j) = \mathcal{D}_0(x_3^j) - 1$; and
- $\mathcal{D}_1(v) = \mathcal{D}_0(v)$ for every $v \notin V(A_0)$.

Note that, for $j \in \mathcal{T}_1 \cup \mathcal{T}_2 \setminus \{z\}$, we have $\mathcal{D}_1(x_2^j) \geq 2$, and for $j \in \mathcal{T}_1 \cup \mathcal{T}_2$, the vertex x_3^j is not adjacent to u in G . Therefore, no neighbor of u in G is a passing vertex in \mathcal{D}_1 . By Lemma 7, there is an addible set $A_1 \subseteq R_1$ towards u with respect to \mathcal{D}_1 such that $|A_1| \geq \lceil |R_1|/2 \rceil$. Let $G_2 = G_1 + A_1$, \mathcal{D}_2 be the A_1 -transformation of \mathcal{D}_1 towards u , and $R_2 = E(G) \setminus E(G_2)$. If $R_2 = \emptyset$, then $G = G_2$, and \mathcal{D}_2 is a path decomposition of G such that $|\mathcal{D}_2| = |\mathcal{D}_1| \leq \lfloor n/2 \rfloor$, a contradiction to the choice of G . Thus, we may assume that $R_2 \neq \emptyset$.

By the construction of \mathcal{D}_2 , it follows that

$$\mathcal{D}_2(u) = \mathcal{D}_1(u) + |A_1|. \quad (3)$$

Let R_2^* be the set of edges $uv_j \in R_2$, where $j \in \mathcal{T}_3 \cup \{z\}$, such that uv_j is the only edge in $E(G)$ joining u to a vertex of T_j which does not belong to G_2 , i.e.,

$$R_2^* = \{uv_j \in R_2 \cap E_G(u, T_j) : j \in \mathcal{T}_3 \cup \{z\} \text{ and } |R_2 \cap E_G(u, T_j)| = 1\}.$$

For every edge $uv_j \in R_2^*$, it follows that the vertex v_j has at most two passing neighbors in \mathcal{D}_2 , namely, the vertices in $V(T_j) \setminus \{v_j\}$, which have even degree in G_2 . Let $A_2 \subseteq R_2^*$ be a maximum addible set outwards u with respect to \mathcal{D}_2 . Let $G_3 = G_2 + A_2$, \mathcal{D}_3 be the A_2 -transformation outwards u with respect to \mathcal{D}_2 , and $R_3 = E(G) \setminus E(G_3)$.

Now we show that $A_2 = R_2^*$. Suppose, for a contradiction, that $A_2 \subsetneq R_2^*$, and let $uv_t \in R_2^* \setminus A_2$. Note that $\mathcal{D}_2(u) = \mathcal{D}_1(u) + |A_1| \geq 1 + \lceil |R_1|/2 \rceil = 1 + \lceil |R_0^u|/2 \rceil \geq 3$. Thus, by Lemma 6, $\{uv_t\}$ is an addible set outwards u with respect to \mathcal{D}_2 , and hence $|A_2| \geq 1$. By the construction of \mathcal{D}_3 , it follows that $\mathcal{D}_3(u) = \mathcal{D}_2(u) - |A_2|$. Now we show that $\mathcal{D}_2(u) \geq 2|R_2^*|$. This is clear if $uv_z \notin R_2^*$; thus suppose that $uv_z \in R_2^*$, and hence

$$\mathcal{D}_2(u) = \mathcal{D}_1(u) + |A_1| \geq 1 + 2(|R_2^*| - 1) + 1 = 2|R_2^*|.$$

Therefore,

$$\mathcal{D}_3(u) = \mathcal{D}_2(u) - |A_2| \geq 2|R_2^*| - |A_2| = 2(|A_2| + |R_2^* \setminus A_2|) - |A_2| = |A_2| + 2|R_2^* \setminus A_2| \geq 3.$$

By Remark 1, $A_2 \cup \{uv_t\}$ is an addible set outwards u with respect to \mathcal{D}_2 , a contradiction to the choice of A_2 .

If $R_3 = \emptyset$, then $G = G_3$ and \mathcal{D}_3 is a path decomposition of G such that $|\mathcal{D}_3| = |\mathcal{D}_2| \leq \lfloor n/2 \rfloor$, a contradiction to the choice of G . Thus, we may assume that $R_3 \neq \emptyset$. Let $A_3 \subseteq R_3$ be a maximum addible set outwards u with respect to \mathcal{D}_3 . Let $G_4 = G_3 + A_3$, \mathcal{D}_4 be the A_3 -transformation outwards u with respect to \mathcal{D}_3 , and $R_4 = E(G) \setminus E(G_4)$. Now, we show that $A_3 = R_3$. Suppose, for a contradiction, that $A_3 \subsetneq R_3$, and let $uv_t \in R_3 \setminus A_3$. By Lemma 9, we have $\mathcal{D}_4(u) > |R_3 \setminus A_3| \geq 1$, and hence $\mathcal{D}_4(u) \geq 2$. We claim that v_t has at most one passing neighbor with respect to \mathcal{D}_4 . Indeed, if $v_t \in \mathcal{I}$, then every neighbor of v_t in G_4 has odd degree, and hence v_t has no passing neighbor; if $v_t = x_1^j$ for some $j \in \mathcal{T}_1 \cup \mathcal{T}_2 \setminus \{z\}$, then the only passing neighbor of x_1^j is possibly x_3^j ; if $v_t = x_i^j$ for some $j \in \mathcal{T}_3$ and $i \in \{1, 2, 3\}$, then, $|E_{G_2}(u, T_t)| \leq 1$, otherwise $uv_t \in A_2$. Thus there is at most one passing vertex in T_t with respect to \mathcal{D}_4 and, therefore, v_t has at most one passing neighbor with respect to \mathcal{D}_4 ; finally, if $v_t = x_i^z$ for some $i \in \{1, 2\}$, then we must have $|E_{G_2}(u, T_z)| = 0$, otherwise $uv_t \in A_2$. Thus, the unique passing vertex in T_z is x_3^z , and hence v_t has at most one passing neighbor in \mathcal{D}_4 . Since $\mathcal{D}_4(u) \geq 2$, by

Lemma 6, uv_t is addible towards v_t with respect to \mathcal{D}_4 , hence by Remark 1, $A_3 \cup \{uv_t\}$ is an addible set outwards u with respect to \mathcal{D}_3 , a contradiction to the choice of A_3 . Thus, $A_3 = R_3 = E(G) \setminus E(G_3)$ and $G = G_4$. Therefore, \mathcal{D}_4 is a path decomposition of G such that $|\mathcal{D}_4| = |\mathcal{D}_3| \leq \lfloor n/2 \rfloor$, a contradiction to the choice of G . \square

Now, we can prove that $EV(G)$ contains no isolated vertex. Indeed, let x be an isolated vertex of $EV(G)$, and let y be any neighbor of x . Since x is isolated, y has odd degree. By Claim 5, x is the unique even neighbor of y , a contradiction to Claim 2.

Given a vertex u of G that has a triangle neighbor T , we say that u is a *full vertex* if every vertex of T is a neighbor of u .

Claim 6. *Let u be a vertex of G that has a triangle neighbor. If u has an odd neighbor that has no even neighbor, then u is a full vertex.*

Proof. Let u be as in the statement, let T be its triangle neighbor, and let v be an odd neighbor of u that has no even neighbor. Let $V(T) = \{x, y, z\}$. Suppose, for a contradiction, that u has at most two neighbors in T . Thus, we may suppose, without loss of generality, that $z \notin N(u)$. Note that, by Claims 2 and 5, u must be adjacent to x and y . Let F_1 be the subgraph of G induced by the edges uv and ux , and F_2 be the subgraph of G induced by yz , and let $F = F_1 \cup F_2$. Let $G_0 = G \setminus E(F)$ and note that $d_{G_0}(w)$ is odd for every $w \in V(T) \cup \{u\}$, and since v has no even neighbors in G , v is an isolated vertex of $EV(G_0)$. Therefore, $EV(G_0) - v \subseteq EV(G)$, and hence $\Delta(EV(G_0)) \leq 3$. We claim that F is a Fan subgraph. Indeed, F_2 consists of a single edge joining even vertices of G . Moreover, F_1 is a star with center at u and two leaves $v_1 = v$ and $v_2 = x$, where $d_G(v_2)$ is even, and hence Definition 2(i) holds. Also, every neighbor of u has odd degree in G_0 , and hence Definition 2(ii) holds. Finally, v_1 is an odd vertex in G , and u has odd degree in G , and every component of $EV(G)$ is a triangle, which verifies Definition 2(iii). By Claim 3, $\text{pn}(G_0) \leq \lfloor n/2 \rfloor$. Let \mathcal{D}_0 be a minimum path decomposition of G_0 . In what follows, we restore the edges yz , xu , and uv , in this order (see Figure 4). Note that every neighbor of y has odd degree in G_0 , thus, by Lemma 6, yz is addible towards y with respect to \mathcal{D}_0 . Let \mathcal{D}_1 be the yz -transformation of \mathcal{D}_0 towards y , and note that every neighbor of u in $G_1 = G_0 + yz$ has odd degree, except for y , but we have $\mathcal{D}_1(y) \geq 2$. Thus, by Lemma 6, xu is addible towards u with respect to \mathcal{D}_1 . Let \mathcal{D}_2 be the xu -transformation of \mathcal{D}_1 towards u , and note that every neighbor of v in $G_2 = G_1 + xu$ has odd degree (note that u is not a neighbor of v in G_2). Again, by Lemma 6, uv is addible towards v with respect to \mathcal{D}_2 . Let \mathcal{D} be the uv -transformation of \mathcal{D}_2 towards v . Therefore, $|\mathcal{D}| = |\mathcal{D}_2| = |\mathcal{D}_1| = |\mathcal{D}_0| \leq \lfloor n/2 \rfloor$, a contradiction. \square




Figure 4: Illustration of the Fan subgraph F in the proof of Claim 6. The edges of F are highlighted in red. A directed edge \vec{ab} indicates the use of Lemma 6 adding the edge ab towards b , following the ordering yz, xu, uv .

Claim 7. *Let u be a vertex of G that has a triangle neighbor. Then every odd neighbor u has an even neighbor.*

Proof. Let u be as in the statement, let T be its triangle neighbor. Suppose, for a contradiction, that u has an odd neighbor v that has no even neighbor. By Claim 6, every vertex of T is a neighbor of u . Let $V(T) = \{x, y, z\}$ and let $S = \{ux, uy, uz\}$. Let $G_0 = G - uv - S$, and note that $d_{G_0}(w)$ is odd for every $w \in V(T) \cup \{u\}$, and since v has no even neighbors in G , v is an isolated vertex of $EV(G_0)$. Therefore, $EV(G_0) - v \subseteq EV(G)$, and hence $\Delta(EV(G_0)) \leq 3$. Let $F = F_1$ be the subgraph of G induced by the edges in $S \cup \{uv\}$. We claim that F is a Fan subgraph. Indeed, F_1 is a star with center at u and four leaves $v_1 = v$ and $v_2 = x, v_3 = y, v_4 = z$ where $d_G(v_i)$ is even, for $i \in \{2, 3, 4\}$, and hence Definition 2(i) holds. Also, every neighbor of u has odd degree in G_0 , and hence Definition 2(ii) holds. Finally, v_1 is an odd vertex in G , and u has odd degree in G , and every component of $EV(G)$ is a triangle, which verifies Definition 2(iii). By Claim 3, $\text{pn}(G_0) \leq \lfloor n/2 \rfloor$. Let \mathcal{D}_0 be a minimum path decomposition of G_0 . Note that no neighbor of u in $G_0 + S = G - uv$ is a passing vertex in \mathcal{D}_0 . By Lemma 7, there is a $B \subseteq S$ such that $|B| \geq \lceil |S|/2 \rceil$ and B is addible towards u with respect to \mathcal{D}_0 . Let \mathcal{D}_1 be the B -transformation of \mathcal{D}_0 towards u . We have $\mathcal{D}_1(u) \geq 1 + \lceil |S|/2 \rceil \geq 3$. Note that $S \setminus B$ contains at most one edge. In what follows, we obtain a decomposition \mathcal{D}_2 of $G_2 = G_0 + S = G - uv$ such that $\mathcal{D}_2(u) \geq 2$. If $S \setminus B = \emptyset$, then $\mathcal{D}_2 = \mathcal{D}_1$ is the desired decomposition (see Figure 5a). If $S \setminus B \neq \emptyset$, then suppose $uz \in S \setminus B$ and put $G_1 = G_0 + B$. Note that the only passing neighbors of z are possibly x and y , and hence, by Lemma 6, uz is addible towards z with respect to \mathcal{D}_1 . Then, the uv -transformation \mathcal{D}_2 of \mathcal{D}_1 towards z is the desired decomposition (see Figure 5b). Finally, note that every neighbor of v in G_2 has odd degree, and hence, by Lemma 6, uv is addible towards v with respect to \mathcal{D}_2 . Then, the uv -transformation \mathcal{D} of \mathcal{D}_2 towards v is a decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$, a contradiction. \square




Figure 5: Illustration of the Fan subgraph F in the proof of Claim 7. The edges of F are highlighted in red. A directed edge \vec{ab} indicates the use of Lemmas 6 and 7 adding the edge ab towards b .

Claim 8. *Every odd vertex of G has a triangle neighbor.*

Proof. Let v be an odd vertex, and let P a shortest path joining v to an even vertex, say x . Let u be the neighbor of x in P , and let w be the neighbor of u different from x in P . By Claim 7, w has an even neighbor, say y . If $w = v$, then the claim follow; and if $w \neq v$, the path $(P \setminus \{xu, uw\}) \cup \{wy\}$ is a path joining v to an even vertex and that is shorter than P , a contradiction. \square

Claim 9. *Let F be a non-empty subgraph of G and let $G' = G \setminus E(F)$. Suppose that for every odd vertex u of G in $V(F)$ the set $N_F(u)$ contains an even vertex of G . Then $\text{pn}(G') \leq \lfloor n/2 \rfloor$.*

Proof. By Remark 2, it is enough to prove that no component of G' is a SET graph. Indeed, suppose that some component, say H' of G' is a SET graph, and let E and O be, respectively, the sets of even and odd vertices of H' . Also, let E_F and O_F be, respectively, the even and odd vertices of G in $V(F)$. Since every vertex of F has odd degree in G' , we have $EV(G') \subseteq EV(G)$, and hence $\Delta(EV(G')) \leq 3$. Moreover, the vertices of F are not in E . Thus, the vertices in E have even degree in G . Since every vertex of O has a neighbor in E , if $O \cap E_F \neq \emptyset$, then $EV(G)$ contains a component different from a triangle. Now, let $u \in O \cap O_F$. By hypothesis, if $u \in O_F$, then F contains an edge uv such that v is an even vertex of G . Thus, u is an odd vertex of G that has neighbors in more than one component of $EV(G)$, a contradiction to Claim 5. \square

Claim 10. *If u and v are two adjacent odd vertices having neighbors in two distinct even components, then u and v are a full vertices.*

Proof. Let u and v be as in the statement, and let T_u and T_v be the triangle neighbors of u and v , respectively. Let $V(T_u) = \{a, b, c\}$ and $V(T_v) = \{x, y, z\}$. Suppose, for a contradiction, that v has at most two neighbors in T_v . Thus, we may suppose, without loss of generality, that $z \notin N(v)$. By Claims 2 and 5, $x, y \in N(v)$. If $|N(u) \cap V(T_u)| = 2$, then suppose, without loss of generality, that $c \notin N(u)$. In this case, put $S = \{ua, bc\}$. If $|N(u) \cap V(T_u)| \neq 2$, by Claim 2, we have $|N(u) \cap V(T_u)| = 3$, then let $S = \{ua, ub, uc\}$. Let $G_0 = G - \{uv, xv, yz\} - S$. Note that $d_{G_0}(w) = d_G(w) - 1$ for every $w \in \{a, b, c, x, y, z\}$, $d_{G_0}(v) = d_G(v) - 2$, and $d_{G_0}(u) \in \{d_G(u) - 2, d_G(u) - 4\}$, and hence $d_{G_0}(w)$ is odd for every $w \in \{a, b, c, x, y, z, u, v\}$. By Claim 9, $\text{pn}(G_0) \leq \lfloor n/2 \rfloor$.

Let \mathcal{D}_0 be a minimum path decomposition of G_0 . In what follows, we restore the edges yz, xv , and uv , in this order. Note that every neighbor of y has odd degree in G_0 , thus, by Lemma 6, yz is addible towards y with respect to \mathcal{D}_0 . Let \mathcal{D}_1 be the yz -transformation of \mathcal{D}_0 towards y , and note that every neighbor of v in $G_1 = G_0 + yz$ has odd degree, except for y , but we have $\mathcal{D}_1(y) \geq 2$. Thus, by Lemma 6, xv is addible towards v with respect to \mathcal{D}_1 . Let \mathcal{D}_2 be the xv -transformation of \mathcal{D}_1 towards v , and note that every neighbor of u in $G_2 = G_1 + xv$ has odd degree. Again, by Lemma 6, uv is addible towards u with respect to \mathcal{D}_2 . Let \mathcal{D}_3 be the uv -transformation of \mathcal{D}_2 towards u , and note that every neighbor of u in $G_3 = G_2 + uv = G \setminus S$ has odd degree, and $\mathcal{D}_3(u) \geq 2$. In what follows, we divide the proof on whether $|N(u) \cap V(T_u)| = 2$ or $|N(u) \cap V(T_u)| \neq 2$.

First, suppose that $|N(u) \cap V(T_u)| = 2$. Note that a has no even neighbors in G_3 , and hence, by Lemma 6, ua is addible towards a with respect to \mathcal{D}_3 . Let \mathcal{D}_4 be the ua -transformation of \mathcal{D}_3 towards a , and note that every neighbor of c in $G_3 + ua$ is odd, except for a , but $\mathcal{D}_4(a) \geq 2$, and hence c has no passing neighbor in \mathcal{D}_4 . Thus, by Lemma 6, bc is addible towards c with respect to \mathcal{D}_4 . But the bc -transformation \mathcal{D} of \mathcal{D}_4 towards c is a path decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$, a contradiction (see Figure 6a).

Thus, we may assume $|N(u) \cap V(T_u)| \neq 2$. Note that no neighbor of u in $G_3 = G \setminus S$ is a passing vertex in \mathcal{D}_3 . By Lemma 7, there is $B \subseteq S$ such that $|B| \geq \lceil |S|/2 \rceil = 2$, and B is addible towards u with respect to \mathcal{D}_3 . Let \mathcal{D}_4 be the B -transformation of \mathcal{D}_3 towards u , we have $\mathcal{D}_4(u) \geq 1 + \lceil |S|/2 \rceil \geq 3$. Note that $S \setminus B$ contains at most one edge. If

$S \setminus B = \emptyset$, then \mathcal{D}_4 is a path decomposition of G such that $|\mathcal{D}_4| \leq \lfloor n/2 \rfloor$, a contradiction. Thus, we may assume $S \setminus B \neq \emptyset$. Suppose, without loss of generality, that $S \setminus B = \{uc\}$ and let $G_4 = G_3 + B$. Note that the only possible passing neighbors of c in \mathcal{D}_4 are a and b . By Lemma 6, uc is addible towards c with respect to \mathcal{D}_4 . Then, the uc -transformation \mathcal{D} of \mathcal{D}_4 towards c is a decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$, a contradiction (see Figure 6b). \square




Figure 6: Illustration of the proof of Claim 10. The edges in $\{uv, xv, yz\} \cup S$ are highlighted in red. A directed edge \vec{ab} indicates the use of Lemmas 6 and 7 adding the edge ab towards b .

Recall that T_1, \dots, T_s are the triangle components of $EV(G)$. Note that, given a graph G , if $s = 1$, then $EV(G)$ consists of one triangle. By Claim 8, every odd vertex of G has an even neighbor, and by Claim 2, every odd vertex of G has at least two even neighbors. Thus, G is a SET graph, a contradiction to the choice of G . Thus, we may assume $s \geq 2$.

Now, let P be a shortest path joining vertices of two different components of $EV(G)$. By Claims 4, 5, and 8, P contains precisely two internal vertices, say u and v . Suppose, without loss of generality, that T_1 is the triangle neighbor of u , and T_2 is the triangle neighbor of v . Let $V(T_1) = \{a, b, c\}$ and $V(T_2) = \{x, y, z\}$. By Claim 10, the vertices u and v are full vertices. Let $S_u = \{ua, ub, uc\}$ and $S_v = \{vx, vy, vz\}$, and let $G_0 = G - uv - S_u - S_v$. Clearly, every vertex in $\{a, b, c, x, y, z, u, v\}$ has odd degree in G_0 . By Claim 9, $\text{pn}(G_0) \leq \lfloor n/2 \rfloor$. Let \mathcal{D}_0 be a minimum path decomposition of G_0 .

In what follows, we obtain a path decomposition \mathcal{D}_3 of $G_3 = G_0 + uv + S_u = G - S_v$ such that $\mathcal{D}_3(u), \mathcal{D}_3(v) \geq 1$, and then we extend it to a path decomposition of G . First, we obtain a path decomposition \mathcal{D}_2 of $G_2 = G_0 + S_u$ such that $\mathcal{D}_2(u) \geq 2$. By Lemma 7, there is a $B_u \subseteq S_u$ such that $|B_u| \geq \lceil |S_u|/2 \rceil \geq 2$ and B_u is addible towards u with respect to \mathcal{D}_0 . Let \mathcal{D}_1 be the B_u -transformation of \mathcal{D}_0 towards u . We have $\mathcal{D}_1(u) \geq 1 + \lceil |S_u|/2 \rceil \geq 3$. Note that $S_u \setminus B_u$ contains at most one edge. If $S_u \setminus B_u = \emptyset$, then put $\mathcal{D}_2 = \mathcal{D}_1$ is the desired decomposition. If $S_u \setminus B_u \neq \emptyset$, then suppose $uc \in S_u \setminus B_u$ and put $G_1 = G_0 + B_u$. Note that the only possible passing neighbors of c are a and b . By Lemma 6, uc is addible towards c with respect to \mathcal{D}_1 . Then, the uc -transformation \mathcal{D}_2 of \mathcal{D}_1 towards c is the desired decomposition. Now, note that every neighbor of v in $G_2 = G_1 + uc = G_0 + S_u$ has odd degree, and hence, by Lemma 6, uv is addible towards v with respect to \mathcal{D}_2 . Then, the uv -transformation \mathcal{D}_3 of \mathcal{D}_2 towards v is a path decomposition of $G_3 = G_0 + uv + S_u = G - S_v$ such that $\mathcal{D}_3(u) \geq 1$ and $\mathcal{D}_3(v) \geq 2$.

By Lemma 7, there is a $B_v \subseteq S_v$ such that $|B_v| \geq \lceil |S_v|/2 \rceil \geq 2$ and B_v is addible towards v with respect to \mathcal{D}_3 . Let \mathcal{D}_4 be the B_v -transformation of \mathcal{D}_3 towards v . We have $\mathcal{D}_4(v) \geq 1 + \lceil |S_v|/2 \rceil \geq 3$. Note that $S_v \setminus B_v$ contains at most one edge. If $S_v \setminus B_v = \emptyset$, then $\mathcal{D} = \mathcal{D}_4$ is a path decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$. If $S_v \setminus B_v \neq \emptyset$, then

suppose $vz \in S_v \setminus B_v$. Note that the only passing neighbors of z in $G_4 = G_3 + B_v$ are x and y . By Lemma 6, vz is addible towards z with respect to \mathcal{D}_4 . Then, the vz -transformation \mathcal{D} of \mathcal{D}_4 towards z is a decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$. \square

The next corollary then is a straightforward application of Lemma 10 and Theorem 13.

Corollary 14. *If G is a connected graph with $\Delta(EV(G)) \leq 3$, then $\text{pn}(G) \leq \lfloor |V(G)|/2 \rfloor$.*

4 Towards Gallai's Conjecture

In this section, we use Theorem 13 to prove Theorem 5, which extends Fan's result [9]. Recall that \mathcal{G} denotes the family of graphs for which (i) each block has maximum degree at most 3; and (ii) each component either has maximum degree at most 3 or has at most one block that contains triangles.

Theorem 15. *If G is a connected simple graph on n vertices such that $EV(G)$ is a subgraph of a graph in \mathcal{G} , then G is a Gallai graph or G is a SET graph.*

Proof. Suppose that the statement does not hold, and let G be a graph on n vertices such that $EV(G)$ is a subgraph of a graph in \mathcal{G} . Suppose that G is a counterexample that minimizes $|E(G)|$. In what follows, we prove three claims regarding G . The proof of Claims 11 and 12 are analogous to the proofs of Claims 1 and 2, respectively. We present their proof for completeness. First, we prove that every hanging ESET subgraph of G must be connected at a special vertex.

Claim 11. *Let K be a hanging ESET subgraph of G , and let G' be such that $G = K \cup G'$, and $V(K) \cap V(G') = \{u\}$. Then u is an odd vertex of G .*

Proof. Let K , G' , and u be as in the statement. Suppose, for a contradiction, that u has even degree in G . Then $EV(G') \subseteq EV(G)$, and hence $EV(G')$ is a subgraph of a graph in \mathcal{G} . By the minimality of G , the graph G' is either a Gallai graph or a SET graph. First, suppose that G' is a Gallai graph, i.e., $\text{pn}(G') \leq \lfloor |V(G')|/2 \rfloor$. By Lemma 12, it follows that

$$\begin{aligned} \text{pn}(G) &\leq \left\lceil |V(K)|/2 \right\rceil + \left\lceil |V(G')|/2 \right\rceil - 1 \\ &\leq (|V(K)| + 1)/2 + |V(G')|/2 - 1 \\ &= (|V(K)| + |V(G')| - 1)/2 \\ &= |V(G)|/2. \end{aligned}$$

Therefore, $\text{pn}(G) \leq \lfloor |V(G)|/2 \rfloor$, and G is a Gallai graph, a contradiction.

Thus, we may assume that G' is a SET graph. By Lemma 10, K (resp. G') admits a path decomposition \mathcal{D}_K (resp. \mathcal{D}') such that $\mathcal{D}_K(u) \geq 2$ and $|\mathcal{D}_K| \leq \lceil |V(K)|/2 \rceil$ (resp. $\mathcal{D}'(u) \geq 2$ and $|\mathcal{D}'| \leq \lceil |V(G')|/2 \rceil$). Let P_1 and P_2 be paths in \mathcal{D}_K having u as end vertex, and Q_1 and Q_2 be paths in \mathcal{D}' having u as end vertex. Put $R_1 = P_1 \cup Q_1$ and $R_2 = P_2 \cup Q_2$,

and note that $\mathcal{D} = (\mathcal{D}_K \setminus \{P_1, P_2\}) \cup (\mathcal{D}' \setminus \{Q_1, Q_2\}) \cup \{R_1, R_2\}$ is a path decomposition of G with cardinality

$$\begin{aligned} |\mathcal{D}| &\leq \lceil |V(K)|/2 \rceil + \lceil |V(G')|/2 \rceil - 2 \\ &\leq (|V(K)| + 1)/2 + (|V(G')| + 1)/2 - 2 \\ &\leq (|V(K)| + |V(G')| - 2)/2 \\ &< (|V(K)| + |V(G')| - 1)/2 \\ &= |V(G)|/2. \end{aligned}$$

Therefore, $\text{pn}(G) \leq \lfloor |V(G)|/2 \rfloor$, and G is a Gallai graph, a contradiction. \square

Now use Fan's techniques to prove that $EV(G)$ consists of vertex-disjoint triangles. First, we prove that no vertex of G has a unique even neighbor.

Claim 12. *No vertex of G has exactly one even neighbor.*

Proof. Suppose, for a contradiction, that G contains a vertex u that has precisely one even neighbor, say v , and let $G' = G \setminus uv$. Note that v has odd degree in G' and u has no even neighbor in G' . Therefore, $EV(G') - u \subseteq EV(G)$ and if $d_G(u)$ is odd, then u is an isolated vertex in $EV(G')$. Thus, $EV(G')$ is a subgraph of a graph in \mathcal{G} . We claim that G' is a Gallai graph. By Remark 2, it is enough to prove that no component of G' is a SET graph. Indeed, G' has at most two components, say G'_u and G'_v , that contain, respectively, u and v . Since u has no even neighbors in G' , G'_u is not a SET graph. Thus, if G' is connected, i.e., $G'_u = G'_v$, then G' is a Gallai graph, as desired. Thus, we may assume $G'_u \neq G'_v$. In this case, note that, if G'_v is a SET graph, then G'_v is a hanging ESET subgraph of G connected at v , which is an even vertex of G , a contradiction to Claim 11. Thus, G'_u and G'_v are Gallai graphs as desired.

Let \mathcal{D}' be a minimum path decomposition of G' . Since v has odd degree in G' , it follows $\mathcal{D}'(v) \geq 1$, and since u has no even neighbor in G' , we have $|\{x \in N_{G'}(u) : \mathcal{D}'(x) = 0\}| = 0$. Thus, by Lemma 6, uv is addible towards u with respect to \mathcal{D}' , a contradiction. \square

The next claim follows the steps introduced by Fan [9].

Claim 13. *Every leaf block of $EV(G)$ is a triangle or an isolated vertex.*

Proof. Suppose, for a contradiction, that $EV(G)$ contains a component C which is neither a triangle nor an isolated vertex, and let H be a leaf block of C . If $H \neq C$, then let w be the cut vertex of C in H , otherwise, let w be any vertex of H .

Subclaim 13.1. *H is a cycle.*

Proof. By Claim 12, no vertex of H has E-degree 1. We prove that $d_H(u) \leq 2$ for every $u \in V(H) \setminus \{w\}$. Suppose, for a contradiction, that H has a vertex $u \neq w$ with degree 3, and let $\{v_1, v_2, v_3\} \subseteq N_H(u)$ be three even neighbors of u . Let F be the subgraph of G induced by the edges uv_1, uv_2, uv_3 , and let $G' = G \setminus E(F)$. Note that every vertex in $V(F)$ has odd degree in G' and, hence, that $EV(G') \subset EV(G)$. Thus, it follows that $EV(G')$ is a subgraph of a graph in \mathcal{G} . We claim that no component of G' is a SET graph. Indeed, suppose that some component C' of G' is a SET graph. Let x, y, z be even vertices of C' .

Suppose that C' contains at least two vertices in $\{u, v_1, v_2, v_3\}$, say a and b . Note that x , y , and z are vertices of H . Since C' is a SET graph, a and b have a common neighbor in x, y, z , say x , but this implies that x has degree at least 4 in H , a contradiction. Thus, C' contains at most one vertex in $\{u, v_1, v_2, v_3\}$, say a . This implies that C' is a hanging ESET of G , but a has even degree in G , a contradiction to Claim 11. By Remark 2, G' is a Gallai graph. Let \mathcal{D}_1 be a minimum path decomposition of G' . Since the vertices of F have odd degree in G' , it follows that $\mathcal{D}_1(v) \geq 1$ for every $v \in V(F)$. By Lemma 7, there is a set $B \subseteq \{uv_1, uv_2, uv_3\}$ addible towards u with respect to \mathcal{D}_1 and containing at least two edges. Moreover, if w is a neighbor of u , say $w = v_1$, then Lemma 7 guarantees that we can choose B such that $uv_1 \in B$. Let \mathcal{D}_2 be the B -transformation of \mathcal{D}_1 towards u . If $|B| = 3$, then \mathcal{D}_2 is a path decomposition of G such that $|\mathcal{D}_2| = |\mathcal{D}_1| \leq \lfloor n/2 \rfloor$, a contradiction. Thus, we may assume $|B| = 2$. Suppose, without loss of generality that $uv_3 \notin B$. Note that $\mathcal{D}_2(u) \geq 3$, and that, since $d_{EV(G)}(v_3) \leq 3$, we have $d_{EV(G'+B)}(v_3) \leq 2$, because u is not a neighbor of v_3 in G' . Thus, by Lemma 6, uv_3 is addible towards v_3 with respect to \mathcal{D}_2 , a contradiction. Thus, every vertex in H different from w has degree 2. \square

Suppose that H is not a triangle, and let u be a neighbor of w in H . Let v_1 and v_2 be the even neighbors of u , where $w = v_1$. Since H is not a triangle, v_1 and v_2 are not adjacent. Let w' be an even neighbor of v_2 different from u . Let $G' = G \setminus \{uv_1, v_2w'\}$, and note that every vertex in $\{u, v_1, v_2, w'\}$ has odd degree in G' , and hence $EV(G') \subseteq EV(G)$. Thus, $EV(G')$ is a subgraph of a graph in \mathcal{G} . We claim that no component of G' is a SET graph. Indeed, suppose that some component C' of G' is a SET graph. Let x, y, z be even vertices of C' . Suppose that C' contains at least two vertices in $\{u, v_1, v_2, v_3\}$, say a and b . Note that x, y , and z are vertices of H . Since C' is a SET graph, a and b have a common neighbor in x, y, z , say x , but this implies that x has degree at least 4 in B' , a contradiction. Thus, C' contains at most one vertex in $\{u, v_1, v_2, w'\}$, say a . This implies that C' is a hanging ESET of G , but a has even degree in G , a contradiction to Claim 11. By Remark 2, G' is a Gallai graph. Let \mathcal{D}_1 be a minimum path decomposition of G' . Note that v_2 has no even neighbors in G' . Thus, by Lemma 6, v_2w' is addible towards v_2 with respect to \mathcal{D}_1 . Let \mathcal{D}_2 be the v_2w' -transformation of \mathcal{D}_1 towards v_2 . Note that v_2 is the only even neighbor of u in $G' + v_2w'$, but $\mathcal{D}_2(v_2) \geq 2$. Thus, by Lemma 6, uv_1 is addible towards u with respect to \mathcal{D}_2 . Therefore, the uv_1 -transformation \mathcal{D} of \mathcal{D}_2 towards u is a path decomposition of G such that $|\mathcal{D}| \leq \lfloor n/2 \rfloor$, a contradiction. \square

Now, if every component of $EV(G)$ has maximum degree at most 3, the statement follows by Theorem 13. Thus, we may assume that $EV(G)$ contains a vertex u of degree at least 4. By hypothesis, $EV(G)$ is a subgraph of a graph H in which each block has maximum degree at most 3; and each component either has maximum degree at most 3 or has at most one block that contains triangles. Let H_u be the component of H that contains u . Since $d_H(u) \geq 4$, u must be a cut vertex of H (and hence, a cut vertex of H_u). Let H'_1, \dots, H'_k be the components of $H_u - u$, and put $H_i = H[V(H'_i) \cup \{u\}]$ for $i = 1, \dots, k$. Clearly, H_1, \dots, H_k decompose H_u . Since $d_{EV(G)}(u) \geq 4$, at least two of these graphs, say H_i and H_j , contain edge of $EV(G)$ incident to u . Let H''_i and H''_j be the subgraphs of $EV(G)$ contained in H_i and H_j , respectively. By Claim 13, every leaf block of $EV(G)$ in H''_i and in H''_j is a triangle. This implies that each H_i and H_j contain a block that is not a triangle-free graph, and hence H_u contains at least two blocks which are not triangle-free graphs, a contradiction. \square

5 Concluding remarks

In this paper we give a step towards verifying Conjecture 1. In fact, Theorems 13 and 15 present statements which are in between Conjecture 1 and Conjecture 4. This indicates that an intermediate statement may be easier to deal than Conjectures 1 and 4. In order to strengthen the results in this paper for fitting Conjecture 4, one need only to verify Conjecture 4 for SET graphs. In fact, knowing that the only non-Gallai graphs with E-degree at most 3 are the odd semi-cliques would simplify the proof of Claim 3, which was introduced to deal with SET subgraphs. By using the Integer Linear Formulation presented in [8], we were able to check this fact for SET graphs up to eleven vertices.

This work benefited greatly from the techniques introduced by Fan [9], and there are two directions that we believe to be natural for extending the results presented in this paper, i.e., two graph classes for which Conjectures 1 and 4 are worth exploring with the techniques introduced here and in [9].

1. Graphs with E-degree at most 4;
2. Graphs in which each block of $EV(G)$ has maximum degree at most 3.

References

- [1] M. Bonamy and T. J. Perrett, *Gallai's path decomposition conjecture for graphs of small maximum degree*, Discrete Math. **342** (2019), no. 5, 1293–1299. MR3905200 ↑2
- [2] A. Bondy, *Beautiful conjectures in graph theory*, European J. Combin. **37** (2014), 4–23. MR3138588 ↑1
- [3] J. A. Bondy and U. S. R. Murty, *Graph theory*, Graduate texts in mathematics, Springer, New York, London, 2008. ↑3
- [4] F. Botler, M. Sambinelli, R. S. Coelho, and O. Lee, *On Gallai's and Hajós' conjectures for graphs with treewidth at most 3*, ArXiv e-prints (June 2017), available at [arXiv:1706.04334](https://arxiv.org/abs/1706.04334). submitted. ↑3
- [5] F. Botler and A. Jiménez, *On path decompositions of $2k$ -regular graphs*, Discrete Math. **340** (2017), no. 6, 1405–1411. MR3624626 ↑2
- [6] F. Botler, A. Jiménez, and M. Sambinelli, *Gallai's path decomposition conjecture for triangle-free planar graphs*, arXiv preprint arXiv:1803.06768 (2018). ↑2, 3
- [7] F. Botler, M. Sambinelli, R. S Coelho, and O. Lee, *Gallai's path decomposition conjecture for graphs with treewidth at most 3*, Journal of Graph Theory (2019). ↑2, 3
- [8] F. Botler, R. Cano, and M. Sambinelli, *On computing the path number of a graph*, X latin and american algorithms, graphs and optimization symposium (LAGOS), 2019. to appear. ↑23
- [9] G. Fan, *Path decompositions and Gallai's conjecture*, J. Combin. Theory Ser. B **93** (2005), no. 2, 117–125. MR2117933 (2005m:05125) ↑2, 4, 5, 20, 21, 23
- [10] O. Favaron and M. Kouider, *Path partitions and cycle partitions of Eulerian graphs of maximum degree 4*, Studia Sci. Math. Hungar. **23** (1988), no. 1-2, 237–244. MR962453 ↑2
- [11] X. Geng, M. Fang, and D. Li, *Gallai's conjecture for outerplanar graphs*, Journal of Interdisciplinary Mathematics **18** (2015), no. 5, 593–598. ↑2
- [12] A. Jiménez and Y. Wakabayashi, *On path-cycle decompositions of triangle-free graphs*, Discrete Math. Theor. Comput. Sci. **19** (2017), no. 3, Paper No. 7, 21. MR3717034 ↑2
- [13] L. Lovász, *On covering of graphs*, Theory of Graphs (Proc. Colloq., Tihany, 1966), 1968, pp. 231–236. MR0233723 (38 #2044) ↑1, 2
- [14] L. Pyber, *Covering the edges of a connected graph by paths*, J. Combin. Theory Ser. B **66** (1996), no. 1, 152–159. MR1368522 (96i:05101) ↑2