

H.E.S.S. detection of very-high-energy γ -ray emission from the quasar PKS 0736+017

H.E.S.S. Collaboration, H. Abdalla¹, R. Adam², F. Aharonian^{3,4,5}, F. Ait Benkhali³, E.O. Angüner⁶, M. Arakawa⁷, C. Arcaro¹, C. Armand⁸, H. Ashkar⁹, M. Backes^{10,1}, V. Barbosa Martins¹¹, M. Barnard¹, Y. Becherini¹², D. Berge¹¹, K. Bernlöhr³, R. Blackwell¹³, M. Böttcher¹, C. Boisson¹⁴, J. Bolmont¹⁵, S. Bonnefoy¹¹, J. Bregeon¹⁶, M. Breuhaus³, F. Brun⁹, P. Brun⁹, M. Bryan¹⁷, M. Büchele¹⁸, T. Bulik¹⁹, T. Bylund¹², M. Capasso²⁰, S. Caroff¹⁵, A. Carosi⁸, S. Casanova^{21,3}, M. Cerruti^{15,42}, T. Chand¹, S. Chandra¹, A. Chen²², S. Colafrancesco^{†22}, M. Curyło²³, I.D. Davids¹⁰, C. Deil³, J. Devin²⁴, P. deWilt¹³, L. Dirson²⁵, A. Djannati-Atai²⁶, A. Dmytriiev¹⁴, A. Donath³, V. Doroshenko²⁰, L.O’C. Drury⁴, J. Dyks²⁷, K. Egberts²⁸, G. Emery¹⁵, J.-P. Ernenwein⁶, S. Eschbach¹⁸, K. Feijen¹³, S. Fegan², A. Fiasson⁸, G. Fontaine², S. Funk¹⁸, M. Füßling¹¹, S. Gabici²⁶, Y.A. Gallant¹⁶, F. Gaté⁸, G. Giavitto¹¹, D. Glawion²⁹, J.F. Glicenstein⁹, D. Gottschall²⁰, M.-H. Grondin²⁴, J. Hahn³, M. Haupt¹¹, G. Heinzlmann²⁵, G. Henri³⁰, G. Hermann³, J.A. Hinton³, W. Hofmann³, C. Hoischen²⁸, T. L. Holch³¹, M. Holler³², D. Horns²⁵, D. Huber³², H. Iwasaki⁷, M. Jamroz²³, D. Jankowsky¹⁸, F. Jankowsky²⁹, A. Jardin-Blicq³, I. Jung-Richardt¹⁸, M.A. Kastendieck²⁵, K. Katarzyński³³, M. Katsuragawa³⁴, U. Katz¹⁸, D. Khangulyan⁷, B. Khélifi²⁶, J. King²⁹, S. Klepser¹¹, W. Kluźniak²⁷, Nu. Komin²², K. Kosack⁹, D. Kostunin¹¹, M. Kraus¹⁸, G. Lamanna⁸, J. Lau¹³, A. Lemièrre²⁶, M. Lemoine-Goumard²⁴, J.-P. Lenain¹⁵, E. Leser^{28,11}, C. Levy¹⁵, T. Lohse³¹, I. Lypova¹¹, J. Mackey⁴, J. Majumdar¹¹, D. Malyshev²⁰, V. Marandon³, A. Marcowith¹⁶, A. Mares²⁴, C. Mariaud², G. Martí-Devesa³², R. Marx³, G. Maurin⁸, P.J. Meintjes³⁵, A.M.W. Mitchell^{3,43}, R. Moderski²⁷, M. Mohamed²⁹, L. Mohrmann¹⁸, J. Muller², C. Moore³⁶, E. Moulin⁹, T. Murach¹¹, S. Nakashima³⁷, M. de Naurois², H. Ndiyavala¹, F. Niederwanger³², J. Niemiec²¹, L. Oakes³¹, P. O’Brien³⁶, H. Odaka³⁸, S. Ohm¹¹, E. de Oa Wilhelmi¹¹, M. Ostrowski²³, I. Oya¹¹, M. Panter³, R.D. Parsons³, C. Perennes¹⁵, P.-O. Petrucci³⁰, B. Peyaud⁹, Q. Piel⁸, S. Pita²⁶, V. Poireau⁸, A. Priyana Noel²³, D.A. Prokhorov²², H. Prokoph¹¹, G. Pühlhofer²⁰, M. Punch^{26,12}, A. Quirrenbach²⁹, S. Raab¹⁸, R. Rauth³², A. Reimer³², O. Reimer³², Q. Remy¹⁶, M. Renaud¹⁶, F. Rieger³, L. Rinchiuso⁹, C. Romoli³, G. Rowell¹³, B. Rudak²⁷, E. Ruiz-Velasco³, V. Sahakian³⁹, S. Saito⁷, D.A. Sanchez⁸, A. Santangelo²⁰, M. Sasaki¹⁸, R. Schlickeiser⁴⁰, F. Schüssler⁹, A. Schulz¹¹, H. Schutte¹, U. Schwanke³¹, S. Schwemmer²⁹, M. Seglar-Arroyo⁹, M. Senniappan¹², A.S. Seyffert¹, N. Shafi²², K. Shiningayamwe¹⁰, R. Simoni¹⁷, A. Sinha²⁶, H. Sol¹⁴, A. Specovius¹⁸, M. Spir-Jacob²⁶, Ł. Stawarz²³, R. Steenkamp¹⁰, C. Stegmann^{28,11}, C. Steppa²⁸, T. Takahashi³⁴, T. Tavernier⁹, A.M. Taylor¹¹, R. Terrier²⁶, D. Tiziani¹⁸, M. Tluczykont²⁵, C. Trichard², M. Tsirou¹⁶, N. Tsuji⁷, R. Tuffs³, Y. Uchiyama⁷, D.J. van der Walt¹, C. van Eldik¹⁸, C. van Rensburg¹, B. van Soelen³⁵, G. Vasileiadis¹⁶, J. Veh¹⁸, C. Venter¹, P. Vincent¹⁵, J. Vink¹⁷, F. Voisin¹³, H.J. Völk³, T. Vuillaume⁸, Z. Wadiasingh¹, S.J. Wagner²⁹, R. White³, A. Wiercholska^{21,29}, R. Yang³, H. Yoneda³⁴, M. Zacharias¹, R. Zanin³, A.A. Zdziarski²⁷, A. Zech¹⁴, A. Ziegler¹⁸, J. Zorn³, N. Żywucka¹, and P. S. Smith⁴¹

(Affiliations can be found after the references)

November 13, 2019

ABSTRACT

Context. Flat-spectrum radio-quasars (FSRQs) are rarely detected at very-high-energies (VHE; $E \geq 100$ GeV) due to their low-frequency-peaked spectral energy distributions. At present, only six FSRQs are known to emit VHE photons, representing only 7% of the VHE extragalactic catalog, which is largely dominated by high-frequency-peaked BL Lacertae objects.

Aims. Following the detection of MeV-GeV γ -ray flaring activity from the FSRQ PKS 0736+017 ($z=0.189$) with *Fermi*-LAT, the H.E.S.S. array of Cherenkov telescopes triggered target-of-opportunity (ToO) observations on February 18, 2015, with the goal of studying the γ -ray emission in the VHE band.

Methods. H.E.S.S. ToO observations were carried out during the nights of February 18, 19, 21, and 24, 2015. Together with *Fermi*-LAT the multi-wavelength coverage of the flare includes *Swift* observations in soft-X-rays and optical/UV, and optical monitoring (photometry and spectro-polarimetry) by the Steward Observatory, the ATOM, the KAIT and the ASAS-SN telescope.

Results. VHE emission from PKS 0736+017 was detected with H.E.S.S. during the night of February 19, 2015, only. *Fermi*-LAT data indicate the presence of a γ -ray flare, peaking at the time of the H.E.S.S. detection, with a flux doubling time-scale of around six hours. The γ -ray flare was accompanied by at least a 1 mag brightening of the non-thermal optical continuum. No simultaneous observations at longer wavelengths are available for the night of the H.E.S.S. detection. The γ -ray observations with H.E.S.S. and *Fermi*-LAT are used to put constraints on the location of the γ -ray emitting region during the flare: it is constrained to be just outside the radius of the broad-line-region r_{BLR} with a bulk Lorentz factor $\Gamma \simeq 20$, or at the level of the radius of the dusty torus r_{torus} with $\Gamma \simeq 60$.

Conclusions. PKS 0736+017 is the seventh FSRQ known to emit VHE photons and, at $z=0.189$, is the nearest so far. The location of the γ -ray emitting region during the flare can be tightly constrained thanks to opacity, variability, and collimation arguments.

Key words. Astroparticle physics; Relativistic processes; Gamma-rays: general; Galaxies: quasars: individual: PKS 0736+017

1. Introduction

The very-high-energy (VHE; $E \geq 100$ GeV) window on the Universe was opened with the discovery of VHE emission from the Crab Nebula (Weekes et al. 1989) using the *Whipple* 10-m Imaging Atmospheric Cherenkov Telescope (IACT). Soon after, the first extragalactic VHE source, the blazar Markarian 421, was discovered by Punch et al. (1992). A few decades later, thanks to the current generation of IACTs (H.E.S.S., MAGIC and VERITAS), the number of known VHE extragalactic sources has grown to 82¹. The majority of them (around 90%) are blazars.

Within the current unification model of active galactic nuclei (AGNs), blazars are interpreted as radio-loud AGNs whose relativistic jet points in the direction of the observer (see Blandford & Rees 1978). From an observational point of view, two sub-classes of blazars exist: flat-spectrum radio-quasars (FSRQs) and BL Lacertae objects, according to the equivalent width of emission lines from the Broad-Line Region (BLR), which is $> 5 \text{ \AA}$ in FSRQs (see e.g. Urry & Padovani 1995). All blazars are characterized by a similar spectral energy distribution (SED), which consists of two distinct components peaking in the infrared-to-X-ray band and the MeV-to-TeV band, respectively (see e.g. Abdo et al. 2010). While FSRQs are in general characterized by a relatively low frequency (in the infrared) of the low-energy SED peak, BL Lac objects are further classified by the peak frequency of their first SED component into low / intermediate / high frequency-peaked BL Lac objects (LBLs / IBLs / HBLs, Padovani & Giommi 1995; Sambruna et al. 1996). Hence, observations in a narrow frequency window unavoidably preselect a particular blazar sub-class, and it is not a surprise that observations at VHE γ rays detect more likely HBLs whose overall spectrum can peak in the VHE band at energies up to few TeVs (such as the extreme HBL 1ES 0229+200, see Aharonian et al. 2007; Aliu et al. 2014), and represent about 70% of extragalactic VHE sources. In addition, FSRQs and BL Lac objects have different redshift distributions, with the former being located on average at larger distances (see Padovani 1992). VHE astronomy has the important property of being limited in redshift, due to the absorption of VHE photons via pair-production over the extragalactic background light (EBL, see Salamon & Stecker 1998). Thus, both intrinsic source properties and propagation effects make FSRQs difficult to be observed with IACTs. So far only six FSRQs have been detected by IACTs: 3C 279 (MAGIC Collaboration et al. 2008; H.E.S.S. Collaboration et al. 2019), PKS 1222+216 (Aleksić et al. 2011; Cerruti 2015), PKS 1510-089 (H.E.S.S. Collaboration et al. 2013; Aleksić et al. 2014), PKS 1441+252 (Abeysekara et al. 2015; Ahnen et al. 2015), S3 0218+35 (Ahnen et al. 2016), which is currently the most distant source of VHE photons ever observed ($z=0.944$), and Ton 599 (Mirzoyan 2017; Mukherjee & VERITAS Collaboration 2017). With the notable exception of PKS 1510-089 (MAGIC Collaboration et al. 2018), which at $z=0.361$ is the nearest among them, all other FSRQs have been detected at VHE only during bright flaring

activity, and are all characterized by very soft VHE spectra.

The radiation mechanism responsible for the low-energy SED component of blazars is thought to be synchrotron emission by a non-thermal population of leptons (electrons and positrons) in the jet. The radiation mechanism responsible for the γ -ray emission is thought to be inverse-Compton scattering off low-energy photons by the same leptons which produce the synchrotron SED component. For FSRQs, the low-energy target photons are thought to be thermal photons from the accretion disc, or from the dusty torus, or emission lines produced in the BLR (see Dermer & Schlickeiser 1993; Sikora et al. 1994; Błażejowski et al. 2000). This type of scenario, called External-Inverse-Compton (EIC), proved to be able to successfully reproduce the broad-band SED of γ -ray FSRQs (see e.g. Ghisellini et al. 2010; Meyer et al. 2012; Böttcher et al. 2013). Alternative hadronic emission scenarios, although capable of modeling the SED of FSRQs (see e.g. Böttcher et al. 2013), encounter difficulties due to the high power required to reproduce the photon emission (Sikora et al. 2009; Reimer 2012; Petropoulou & Dimitrakoudis 2015; Zdziarski & Böttcher 2015).

Even though there is general consensus on the EIC as the emission mechanism, several questions remain open. Among them is the uncertainty as to where the γ -ray emission region is located within the relativistic jet. Does the emission come from the base of the jet, near the supermassive black hole (SMBH) powering the quasar, or is it rather produced downstream in the jet? The answer to this question, which arose with the detection of the first γ -ray AGNs by EGRET (see e.g., Dermer et al. 1992; Dermer & Schlickeiser 1994; Becker & Kafatos 1995; Blandford & Levinson 1995; Marcowith et al. 1995; Jorstad et al. 2001), is still a major active research topic in blazar physics and has been addressed by several authors in the last years (see e.g., Ghisellini & Tavecchio 2009; Finke & Dermer 2010; Poutanen & Stern 2010; Tavecchio et al. 2010; Agudo et al. 2011a,b; Hayashida et al. 2012; Yan et al. 2012; Cao & Wang 2013; Rani et al. 2013; Brown 2013; Dermer et al. 2014; Max-Moerbeck et al. 2014; Nalewajko et al. 2014; Dotson et al. 2015; Ramakrishnan et al. 2015; Coogan et al. 2016; Finke 2016). In this paper it is shown that, making use only of γ -ray observations of the quasar PKS 0736+017, it is possible to put a tight constraint on the location of the γ -ray emitting region within the jet.

The quasar PKS 0736+017 was first detected as a radio-source with the Parkes telescope (Day et al. 1966). Its radio morphology is typical of a blazar, with a compact core and a single-sided, parsec-scale jet (Lister & Homan 2005; Lister et al. 2009). The optical-UV spectrum is characterized by broad emission lines and a big-blue bump, associated with thermal emission from the SMBH accretion disk (Baldwin 1975; Malkan & Moore 1986). Thanks to the emission lines, the redshift of PKS 0736+017 is well determined: the most recent measurement is $z = 0.189$ (Ho & Kim 2009). As is typical for blazars, a giant elliptical galaxy hosts the AGN (Kotilainen et al. 1998; Wright et al. 1998; McLure et al. 1999). During January 2002, PKS 0736+017 exhibited an extremely bright and fast optical flare (0.6 magnitudes/hour, see Clements et al. 2003), which classifies the source also as an Optically Violently Variable (OVV) quasar.

Send offprint requests to: H.E.S.S. collaboration,
e-mail: contact.hess@hess-experiment.eu;

* Corresponding authors

† Deceased

¹ In October 2019. For an up-to-date list of VHE sources see <http://tevcat.uchicago.edu>, Wakely & Horan (2008)

In high-energy γ rays (HE; $100 \text{ MeV} \leq E \leq 100 \text{ GeV}$), PKS 0736+017 is detected by *Fermi*-LAT, and is included in the most recent *Fermi*-LAT catalog (3FGL, Acero et al. 2015) under the name 3FGL J0739.4+0137. Since the beginning of the *Fermi* mission in 2008, the source remained relatively quiescent until November 2014, when a γ -ray flare was detected (D’Ammando & Orienti 2014). The source remained active in HE γ rays for the following months, and reached the maximum γ -ray flux in February 2015. The flaring state in HE γ rays triggered VHE observations with the High Energy Stereoscopic System (H.E.S.S.), resulting in the first detection of VHE photons from PKS 0736+017, which is reported in this paper.

The paper is organized as follows: in Section 2 are presented the observations of PKS 0736+017 and the data analysis, for H.E.S.S. and for lower-energy instruments; in Section 3 are discussed the implications of the VHE detection, in particular in terms of the location of the γ -ray emitting region, and is presented the SED of the source during the flare; conclusions are in Section 4.

2. Observations and Data Analysis

2.1. H.E.S.S.

H.E.S.S. is an array of IACTs located in the Khomas Highland of Namibia ($23^\circ 16' 18'' \text{ S}$, $16^\circ 30' 00'' \text{ E}$), at an altitude of about 1800 m above sea level. As all other IACTs, H.E.S.S. images the Cherenkov light emitted by particle showers triggered by interaction of γ rays with the Earth’s atmosphere. The study of the shower images makes it possible to reconstruct the incoming direction of a γ ray, its energy and its arrival time. The array consists of four 12-m diameter reflectors arranged in a square of 120 m side length and, since 2012, one additional 28-m diameter telescope (in the following called CT5) in the middle of the array (H.E.S.S. Collaboration et al. 2017). This hybrid configuration of the array allows data to be taken in different modes by triggering on events either detected by CT5 only (monoscopic mode) or by any combination of two or more telescopes (stereoscopic mode). The standard observation mode is to collect both monoscopic and stereoscopic events during the same observation to allow for a low-energy analysis threshold (below 100 GeV in the monoscopic mode) as well as a good spatial and spectral reconstruction based on the excellent background rejection power in the stereoscopic mode.

Target of Opportunity (ToO) observations of PKS 0736+017 with H.E.S.S. were triggered on February 18, 2015, following the detection of flaring activity in *Fermi*-LAT data (see next Section). H.E.S.S. observations were carried out with the full array in the same night. Follow-up observations were performed on February 19, 21, and 24 (with the full array for the first two nights, while data were recorded with CT5 only during the last night). All observations were taken in “wobble mode” where the source position is offset by 0.5° from the camera center to allow for simultaneous background estimation (Berge et al. 2007). The details of H.E.S.S. observations are reported in Table 1.

Data were analyzed with the ImPACT analysis chain (Parsons & Hinton 2014; Parsons et al. 2015) using both monoscopic and stereoscopic reconstruction and were crosschecked with an independent analysis chain (De Naurois & Rolland 2009; Holler et al. 2015) yielding consistent results. With the




Fig. 1. Gamma-ray emission from PKS 0736+017 centred on the night of February 19, 2015. The red bow-tie represents the spectrum measured with H.E.S.S. using the monoscopic (bold line) and the stereoscopic analysis (dashed line). The green and blue bow-ties represent the spectra measured with *Fermi*-LAT strictly simultaneous with the H.E.S.S. detection, and integrating over an exposure of 24 hours (MJD 57072.5–57073.5) around the H.E.S.S. detection, respectively; the extrapolation of *Fermi*-LAT spectra to higher energies (green and blue long-dashed lines) takes into account the absorption on the EBL (Franceschini et al. 2008).

loose configuration of the monoscopic analysis, an excess of 473 γ -like events with a signal-to-background ratio of 0.17 is found at the source position in the overall good-quality 7.2 hour dataset, corresponding to a detection significance of 8.5σ using Eq. 17 of Li & Ma (1983). The overall signal is dominated by the strong excess coming from the second night of observations (February 19, 2015). The monoscopic (respectively, stereoscopic) analysis of this 1.8 h live time data set yields an excess of 364 (49) γ -like events, corresponding to an 11.1σ (5.3σ) post-trial² significance, while the source is not detected with a significance greater than 5σ in any other night.

To determine the position of the VHE γ -ray emission, a two-dimensional fit to the γ -ray excess (for the night of the VHE detection only) is performed using the stereoscopic data set, yielding a shape consistent with a point-like source. The best-fit position is found at RA(J2000) = $07^{\text{h}} 39^{\text{m}} 17^{\text{s}} \pm 3^{\text{s}}$, DEC(J2000) = $01^\circ 36' 29'' \pm 56''$, and is positionally consistent with the radio position of PKS 0736+017 (Lanyi et al. 2010).

The differential energy spectrum of the γ -ray emission is derived by performing a spectral fit, again for the night of the detection only. Both monoscopic and stereoscopic spectra are consistent with a power-law model of the form $dN/dE = N_0(E/E_0)^{-\Gamma}$. The photon index is estimated to be $\Gamma = 3.1 \pm 0.3_{\text{stat}} \pm 0.2_{\text{sys}}$ for the monoscopic analysis and $\Gamma = 4.2 \pm 0.8_{\text{stat}} \pm 0.2_{\text{sys}}$ for the stereoscopic one, and the flux normalization at 200 GeV is found to be $N_0 = (1.0 \pm 0.2_{\text{stat}} \pm 0.3_{\text{sys}}) \times 10^{-10} \text{ cm}^{-2} \text{ s}^{-1} \text{ TeV}^{-1}$ and

² Four trials are considered for the computation of significances: two for the analyses (monoscopic and stereoscopic), and two for the time selection (the full data-set, and the night of February 19, 2015).

Table 1. Details of the H.E.S.S. observations of PKS 0736+017.

Array configuration	Starting time [UTC]	Exposure [h]	Flux ($E > 100$ GeV) [10^{-11} cm $^{-2}$ s $^{-1}$]
CT1-5	Feb 18, 2015, 21:21	1.8	< 3.6
CT1-5	Feb 19, 2015, 18:53	1.8	5.7 ± 1.1
CT1-5	Feb 21, 2015, 20:31	2.7	< 2.0
CT5	Feb 24, 2015, 21:43	0.9	< 3.1

$N_0 = (1.1 \pm 0.3_{\text{stat}} \pm 0.2_{\text{syst}}) \times 10^{-10}$ cm $^{-2}$ s $^{-1}$ TeV $^{-1}$, respectively. It is important to underline that the two analyses have different energy thresholds, equal to 80 and 150 GeV, respectively. The spectral results obtained are presented in Fig. 1.

The night-by-night light curve for the monoscopic reconstruction, provided as an integral flux above 100 GeV, assuming a photon index of 3.1, is shown in Fig. 2, panel “a”. For the nights of no detection with H.E.S.S., upper limits on the VHE emission (estimated following Rolke et al. (2005) at the 95% confidence level) are also shown. No evidence for intra-night variability is found in the H.E.S.S. data during the night of February 19, 2015: a fit of the 28-minutes-binned light curve with a constant function results in a χ^2 value of 1.9 for three degrees of freedom.

2.2. Fermi-LAT

Detecting γ -ray photons with energies between 20 MeV and above 300 GeV, the LAT instrument (Atwood et al. 2009) onboard the *Fermi* satellite monitors the high-energy γ -ray sky every three hours. This instrument is thus ideal to reveal active states in AGNs, which could be used to trigger observations with other facilities as ToO observations. On February 18, 2015, such an active state was detected in PKS 0736+017 using the public monitored source list³ as well as the dedicated FLAapLUC aperture-photometry pipeline (Lenain 2018). Following this early flare detection, a ToO campaign was launched with H.E.S.S., as reported in the previous Section.

The *Fermi*-LAT data are analyzed with the public ScienceTools v10r0p5⁴, and events are selected within a circular region of interest of 10° in radius centred on the nominal position of 3FGLJ0739.4+0137, in order to perform a binned analysis as implemented in the gtlite tool. To encompass the entire active state studied here, data between February 1, 2015 and April 1, 2015 are considered, in the 100 MeV–500 GeV energy range. The P8R3_SOURCE_V6 instrument response functions were used, together with a zenith angle cut of 90° to avoid contamination by the γ -ray bright Earth limb emission. The model of the region of interest was built based on the 3FGL catalog (Acero et al. 2015), and it has been checked *a posteriori* that no significant residual remains, which could have hinted for new sources not referenced in the 3FGL catalogue. The Galactic diffuse emission has been modeled using the file gll_iem_v06.fits (Acero et al. 2016) and the isotropic background using iso_P8R3_SOURCE_V6_v06.txt.

For the time window considered here, the spectrum of PKS 0736+017 is best described using a log-parabolic shape.⁵ A comparison with a power-law spectrum yielded a log-likelihood ratio of 27.5 in favor of the log-parabola. Under this spectral hypothesis, PKS 0736+017 is detected with a test statistic (TS; Mattox et al. 1996) of 2061, i.e. $\sim 45\sigma$, with a flux $F_{(100\text{MeV}-500\text{GeV})} = (5.42 \pm 0.27) \times 10^{-7}$ cm $^{-2}$ s $^{-1}$, index $a = 2.22 \pm 0.06$, curvature index $b = 0.037 \pm 0.025$, and reference energy $E_0 = 327$ MeV, averaged over the 2 months of data considered here. In Fig. 2, panel “b”, the *Fermi*-LAT light curve of PKS 0736+017 is reported (starting from February 13, 2015), using a time binning of 6 h around the γ -ray flare (MJD 57070–57074) and 12 h elsewhere, and a power-law spectrum with both the flux and photon index of PKS 0736+017 free to vary. In the light curve, flux upper-limits at the 68% level are provided when $TS < 9$. The evolution in time of the best-fit photon index is shown in panel “c”.

To better compare the *Fermi*-LAT data of PKS 0736+017 with H.E.S.S. data during the night it was detected at VHE, as well as to retain sufficient statistics, a *Fermi*-LAT data subset was analyzed for the time window MJD 57072.5–57073.5. The source is detected in this subset with a TS=339 ($\sim 18\sigma$), a flux of $F_{(100\text{MeV}-500\text{GeV})} = (2.08 \pm 0.26) \times 10^{-6}$ cm $^{-2}$ s $^{-1}$, and a photon index of $\Gamma = 2.15 \pm 0.10$. In this case, no evidence of curvature is found (log-likelihood ratio of -0.12). The *Fermi*-LAT highest energy for a photon associated with PKS 0736+017 (at the 95% confidence level) is 106 GeV (detected on MJD 57072, at 14:42:20 UTC). This is also the highest photon energy when considering the larger two-months interval described above. Given the high significance of the detection, a *Fermi*-LAT data analysis has also been performed in a time window strictly simultaneous with the H.E.S.S. detection (MJD 57072, 18:53–20:55 UTC). PKS 0736+017 is detected in this dataset with a TS= 37 ($\sim 6\sigma$), a flux of $F_{(100\text{MeV}-500\text{GeV})} = (4.57 \pm 1.52) \times 10^{-6}$ cm $^{-2}$ s $^{-1}$, and a photon index of $\Gamma = 2.43 \pm 0.33$. The highest-energy LAT photon simultaneous with the H.E.S.S. detection has $E = 1.71$ GeV (detected on MJD 57072, at 20:40:16 UTC). Fig. 1 shows the *Fermi*-LAT spectra of PKS 0736+017 for the two time windows described here.

To estimate the variability in the HE γ -ray band, the fractional variability $F_{\text{var,HE}}$ of the 12-hours binned light curve

⁵ The best-fit γ -ray spectrum of 3FGLJ0739.4+0137 as provided in the 3FGL catalog is a log-parabola with index $a = 2.25 \pm 0.05$, curvature $b = 0.17 \pm 0.03$, reference energy $E_0 = 327.1$ MeV and $EF_{(1\text{GeV}-100\text{GeV})} = (3.66 \pm 0.13) \times 10^{-11}$ erg cm $^{-2}$ s $^{-1}$. The *Fermi*-LAT Collaboration has recently released a new γ -ray catalog, the 4FGL (the *Fermi*-LAT Collaboration 2019). The 4FGL associated source for PKS 0736+017, 4FGLJ0739.2+0137, is again best described by a log-parabolic spectrum, with $a = 2.33 \pm 0.02$, curvature $b = 0.09 \pm 0.01$, reference energy $E_0 = 503.3$ MeV and $EF_{(100\text{MeV}-100\text{GeV})} = (6.12 \pm 0.14) \times 10^{-11}$ erg cm $^{-2}$ s $^{-1}$.

³ <https://fermi.gsfc.nasa.gov/ssc/data/access/lat/msl/lc/>

⁴ See <http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation>.




Fig. 2. Multi-wavelength light curve of PKS 0736+017 during February 2015. Panels from a to g: H.E.S.S. integral flux above 100 GeV from the monoscopic analysis (in gray, the flux for the nights with no significant detection); *Fermi*-LAT integral flux above 100 MeV; *Fermi*-LAT photon index; *Swift*-XRT integral flux between 0.3 and 10 keV, corrected for absorption; *Swift*-UVOT (orange, light-blue, blue, light-violet, violet and dark-violet squares for the V,B,U,W1,M2 and W2 filter, respectively), ATOM (red squares, for the R filter), Steward Observatory (orange and red triangles, for the V and R filter, respectively), ASAS-SN (orange open circles, for the V filter), and KAIT (red open circles, unfiltered) fluxes, de-reddened; Steward Observatory broad-band polarization fraction; and Steward Observatory polarization position angle (both averaged over 5000-7000 Å). The vertical dashed red line indicates the time of the H.E.S.S. detection.

Table 2. Details of the *Swift* observations of PKS 0736+017.

Obs. ID	Starting time [UTC]	Exposure [ks]	Power-law Index	Flux 0.3-10 keV [10^{-12} erg cm $^{-2}$ s $^{-1}$]	χ^2/DOF	UVOT filters
00033535009	Feb 20, 2015, 19:26	3.4	1.62 ± 0.08	8.4 ± 0.5	11.2 / 14	V,B,U,W1,W2,M2
00033535010	Feb 22, 2015, 01:46	2.5	1.49 ± 0.09	9.2 ± 0.6	13.7 / 10	V,B,U,W1,W2,M2
00033535011	Feb 22, 2015, 21:25	2.6	1.46 ± 0.13	7.3 ± 0.7	5.4 / 6	W1
00033535012	Feb 27, 2015, 08:01	3.8	1.34 ± 0.09	6.9 ± 0.6	9.4 / 11	U

was computed following [Vaughan et al. \(2003\)](#). This results in $F_{var,HE} = (62 \pm 7)\%$, clearly supporting the presence of variability of the HE γ -ray flux. This finding does not depend on the binning of the light curve: for a 6-hours binned light curve, $F_{var,HE} = (48 \pm 2)\%$, while for a 24-hours binned light curve, $F_{var,HE} = (57 \pm 4)\%$. An important property of a flare is its variability timescale, which is defined here as the flux-doubling timescale (see e.g. [Aliu et al. 2016](#)):

$$F(t) = \frac{a}{2^{-(t-t_0)/b} + 2^{(t-t_0)/c}} \quad (1)$$

where t_0 is the time of the peak flux in the light curve, a fits the peak flux, and b and c denote the rising and falling flux-doubling time-scales, respectively. The evolution of the γ -ray flare in the HE band shows a slow flux increase starting on February 13, 2015, with a first flare on February 18, 2015, at an integral flux level of around 3×10^{-6} cm $^{-2}$ s $^{-1}$. The flux then dropped by a factor of around four, before brightening again on February 19, 2015, coincidentally with the H.E.S.S. detection. The flux doubling time-scale for this second flare centered on the H.E.S.S. detection is fitted using the 6-hours binned light curve as 6 ± 4 hours for the rising and 4 ± 2 hours for the falling part of the flare.

2.3. *Swift*-XRT

Following the detection of the γ -ray flaring activity in PKS 0736+017, ToO observations were requested to the *Neil Gehrels Swift* satellite ([Gehrels et al. 2005](#)) in order to measure the flux of PKS 0736+017 in soft-X-rays and optical/UV. The ToO resulted in four observations with *Swift* starting from February 20, 2015, for a total exposure time of around 12.3 ks. The details are provided in Table 2.

Data from *Swift*-XRT ([Burrows et al. 2005](#)) are analyzed using *Heasoft* version 6.18. All observations are taken in the standard *Photon Counting* observing mode. The event files are cleaned using standard screening criteria. Images, spectra and light curves are extracted using a circular region with radius equal to 20 pixels. The background is extracted from an annular region with inner radius equal to 50 pixels, and outer radius equal to 160 pixels. The count rates are around 0.11-0.16 s $^{-1}$, and are low enough that no pile-up affects the analysis.

Spectral analyses of *Swift*-XRT data are performed using *XSpec* version 12.9. The ancillary response files are recomputed using *xrtmkarf*, while the redistribution matrix files provided by the *Swift* team are used. Data are rebinned using *grppha* imposing a minimum of 30 counts per bin, and data below 0.3 keV are excluded.

The X-ray spectrum from each of the four *Swift*-XRT observations is fitted with an absorbed powerlaw model (*phabs*cflux*powerlaw* within *Xspec*, to access the unabsorbed flux in the 0.3-10 keV band), to take into account the absorption by neutral material in the Milky Way. The value of N_H in the direction of PKS 0736+017 is fixed to 7.8×10^{20} cm $^{-2}$ as estimated by [Dickey & Lockman \(1990\)](#). The results of the fit are provided in Table 2. The *Swift*-XRT light curve is shown in Fig. 2, panel “d”. No significant variability is detected in the X-ray data: the four *Swift*-XRT observations are consistent with a constant flux ($\chi^2/DOF = 9/3$, which corresponds to a chance probability of about 3%), with an average $F_{0.3-10 \text{ keV}} = (8.1 \pm 0.3) 10^{-12}$ erg cm $^{-2}$ s $^{-1}$.

2.4. *Swift*-UVOT

Simultaneously with *Swift*-XRT, the UVOT telescope ([Roming et al. 2005](#)) also observed PKS 0736+017, guaranteeing a coverage in the optical and UV band. During the first two observations all UVOT filters were available, while the two other observations were taken only with the W1 and U filter, respectively (see Table 2). Fluxes are calculated using *uvotmaghist*, integrating over a 5''-radius circular region for the source, and a 20''-radius circular region for the background. Data are de-reddened following [Roming et al. \(2009\)](#) using $E_{B-V} = 0.121$ as value for the Galactic extinction ([Schlafly & Finkbeiner 2011](#)). The *Swift*-UVOT light curve is presented in Fig. 2, panel “e”. There is clear variability in the optical band, showing a decrease from February 20, 2015 to February 27, 2015.

2.5. ATOM

The ATOM telescope ([Hauser et al. 2004](#)), located on the H.E.S.S. site, regularly observes PKS 0736+017 as part of its blazar monitoring program, providing a long term light curve of the source in the R band. Fluxes from ATOM observations are extracted using a 4''-radius aperture, and have been de-reddened using the same value of Galactic extinction $E_{B-V} = 0.121$ as for the UVOT data analysis. There are no ATOM observations during the night of February 19, 2015, when H.E.S.S. detected VHE emission from PKS 0736+017. On the other hand, ATOM observations on the night of February 20, 2015, are contemporaneous with the UVOT ones. The light curve from ATOM data is provided in Fig. 2, panel “e”.

2.6. Steward Observatory

PKS 0736+017 is among the sources covered by the Steward Observatory blazar monitoring program ([Smith et al. 2009](#);




Fig. 3. Optical spectra (4000–7500 Å) of PKS 0736+017 obtained with SPOL on three consecutive nights of 2015: February 18 (in red), February 19 (in green), and February 20 (in blue). Due to absence of photometric conditions on February 19, this spectrum is provided in arbitrary units. For reference, it is shown in black a spectrum from a low-state of the source (February 9, 2016). The emission lines and telluric features are identified in the spectrum from February 20, 2015.

Carnerero et al. 2015). The program provides optical monitoring for bright *Fermi*-LAT blazars, including photometry, spectroscopy and spectro-polarimetry. During February 2015, PKS 0736+017 was observed on nine occasions from February 12 to February 20, using the 1.5-m Kuiper Telescope located on Mt. Bigelow, Arizona and the SPOL CCD spectropolarimeter (Schmidt et al. 1992). The SPOL apertures range were $5.1'' \times 10''$ with the first number being the width of the slit and the second number being the width of the spectroscopic extraction aperture. The extraction aperture is in the east-west direction on the sky. For the SPOL differential photometry measurements, the slit width used was $12.7''$ and the extraction aperture was either $10''$, $11''$, or $12''$, depending on the seeing at the time of the measurement. Photometry measurements are provided in the V (Johnson) and R (Kron-Cousins) bands, and have been de-reddened using $E_{B-V} = 0.121$ as for the UVOT and ATOM data. The light curve from the Steward Observatory monitoring of PKS 0736+017 is shown in Fig. 2, and complements UVOT and ATOM measurements to provide optical coverage of the γ -ray flaring activity in PKS 0736+017. Observations using SPOL were made on the morning of February 20, 2015 UTC, 8.4 hours after the H.E.S.S. detection, and show a clear optical flare. There was an increase in luminosity of about one magnitude compared to the measurements taken 48 hours previously with

SPOL and with the Swift-UVOT and ATOM observations made 17 hours afterwards. Measurements of the equivalent widths of the $H\beta$ and $H\gamma$ broad-emission lines of PKS 0736+017 using the flux spectra obtained with SPOL confirm the rapid brightening of the blazar on February 20. The decrease in the equivalent widths of the emission lines from February 19 to February 20 is consistent with a 1-mag increase in the continuum brightness over the period of about 24 hours. The onset of the optical flare appears to be as rapid as its decay (see panel “e” in Fig. 2), and the coincidence of the optical, HE, and VHE outbursts is strong evidence that they are closely related physically.

Measurements of the optical polarization fraction and position angle are shown in panels “f” and “g” of Fig. 2. Uncertainties on both quantities are smaller than the symbols used: the uncertainty on the polarization fraction is $\leq 0.2\%$, while the uncertainty on the angle is $\leq 1^\circ$. The polarization fraction reported has been corrected by subtracting the assumed unpolarized contributions to the spectrum by the BLR and big blue bump. This is best done by subtracting an optical spectrum of PKS0736+017 obtained during a faint period when the blazar shows little polarization. This is when the BLR and thermal optical continuum are most dominant in the spectrum. After subtracting the unpolarized flux, the measured polarization fraction is a better approximation to the intrinsic polarization of the synchrotron continuum responsible for the rapid optical flux variations. Given that the spectrum of PKS0736+017 still includes some polarization from the non-thermal continuum even when it is optically faint, the values of the polarization fraction plotted in Figure 2 are the highest possible intrinsic polarization levels of the synchrotron emission. The highest polarization is observed not for the night of the optical flare, but for the night before, at $(12.5 \pm 0.1)\%$. The polarization position angle is extremely variable during the week leading up to the γ -ray flare and is observed throughout the full 180° range. It is interesting to note that the two measurements taken on February 20 show a variation of the polarization angle of $(44.9 \pm 0.6)^\circ$ in three hours, which is a very rapid rotation even for a blazar (see Blinov et al. 2015, 2016; Lyutikov & Kravchenko 2017).

Spectro-polarimetry of PKS 0736+017 yields the spectral index α (defined as $f_\lambda \propto \lambda^{-\alpha}$) of the optical synchrotron continuum during the flare, which is key to constraining radiative models (see Section 3.2). The spectral index is derived directly from the polarized flux spectrum, assuming that the polarization of the synchrotron continuum is constant with wavelength. This may not be the case, but observations of BL Lac objects, where the synchrotron emission dominates the optical flux, do not often show strong wavelength-dependent polarization (Sitko et al. 1985; Jannuzi et al. 1994; Smith 1996). Also, PKS0736+017 shows no evidence for a strong variation in polarization position angle with wavelength in 2015 February. The de-reddened polarized flux spectrum of PKS 0736+017 is consistent with a featureless power law and indicates that the emission lines are unpolarized. During the morning of Feb 20, 2015, the spectral index is found to be 1.0 ± 0.1 , significantly flatter than for the nights before the flare, when it is observed to be between 1.4 and 1.8, indicating a shift of the synchrotron peak frequency to higher energies.

2.7. ASAS-SN and KAIT

PKS 0736+017 is also regularly monitored in the optical by the ASAS-SN project (Shappee et al. 2014) and the KAIT telescope (Cohen et al. 2014). Their public data have been retrieved for the observations around the February 2015 γ -ray flare. ASAS-SN V-magnitudes, and KAIT unfiltered magnitudes (which are close to the R ones, see Li et al. 2003) have been converted to fluxes and de-reddened using $E_{B-V} = 0.121$ as for the other optical observations. ASAS-SN and KAIT observed PKS 0736+017 during the morning (UTC) of February 19, respectively 11.5 hours and 14 hours before the H.E.S.S. VHE detection, and complement the optical light curve from UVOT, ATOM, and Steward Observatory. The contribution of the host-galaxy to the optical measurements has not been investigated. The emission in the optical band is dominated by the non-thermal continuum from the quasar, as it is evident from the SED shown in Fig. 5.

3. Discussion

3.1. On the location of the γ -ray emission region

As shown in the multi-wavelength light curve in Fig. 2, the only instrument which was observing PKS 0736+017 together with H.E.S.S. during the night of February 19, 2015, was *Fermi*-LAT. Observations by *Swift* covered only the post-flare period, and there are no optical observations available within 8 hours of the H.E.S.S. detection. It is thus not possible to follow the usual approach of modeling the simultaneous SED to access blazar physics. In particular, there is no coverage of the behavior of the synchrotron component of PKS 0736+017 simultaneously with the γ -ray flare. On the other hand, constraints can be put on the location of the emitting region r (defined as the distance from the SMBH) within the relativistic jet, under some hypotheses. In the following, it is assumed that⁶:

- the main radiation mechanism responsible for the γ -ray emission is inverse-Compton scattering by a population of electrons/positrons in the jet, off a low-energy photon field represented by the radiation from the accretion disc, the BLR and the dusty torus;
- the γ -ray emission during the flare is produced from a single region within the relativistic jet;
- the jet of PKS 0736+017 is closely aligned to the line of sight, and the Doppler factor δ of the emitting region equals its bulk Lorentz factor Γ , that is the angle to the line of sight ϑ_{los} is equal to $1/\Gamma$;
- the emitting region is approximated by a spherical plasmoid in the jet, characterized by its radius R' , which is related to the variability time-scale τ_{var} via $R' \simeq c \tau_{var} \frac{\Gamma}{1+z}$, where c is the speed of light in vacuum, and z is the redshift of the source; the most constraining estimate for τ_{var} comes from the falling part of the *Fermi*-LAT flare as $\tau_{var} = (4 \pm 2)$ hours; the two extreme values of 2 and 6 hours are used in the following;
- the emitting region fills the entire cross-section of the relativistic jet and thus the location of the emitting region r and its size R are simply related by $R/r = \tan \vartheta_{open}$, where ϑ_{open} is the jet opening angle;
- the location of the BLR, r_{BLR} , is derived from the luminosity of the H β line, which is measured from Steward

⁶ Here and in the following, quantities in the co-moving jet frame are indicated by a prime.




Fig. 4. Constraint on the location of the emitting region r in cm, defined as distance from the SMBH, as a function of the bulk Lorentz factor Γ . The red exclusion region represents the opacity constraint, the blue exclusion region represents the collimation constraint, while the orange exclusion region represents the cooling constraint, the last two computed for $\tau_{var} = 6$ hours. Dashed lines show how the constraints change assuming $\tau_{var} = 2$ hours. The white regions indicate the allowed part of the parameter space. The vertical black lines represent the estimated locations of the BLR r_{BLR} and of the dust torus r_{torus} . The green star identifies the parameters of the EIC solution discussed in Sec. 3.2 and plotted in Fig. 5.

Observatory data to be $L_{H\beta} = 4.20 \times 10^{42} \text{ erg s}^{-1}$. Following Greene & Ho (2005), r_{BLR} is then estimated as 1.45×10^{17} cm; the total BLR luminosity is similarly estimated from $L_{H\beta}$ (see Finke 2016) as $L_{BLR} = 1.24 \times 10^{44} \text{ erg s}^{-1}$, and $L_{disk} \simeq 10 L_{BLR} = 1.24 \times 10^{45} \text{ erg s}^{-1}$. The location of the dusty torus r_{torus} is assumed to scale as $r_{torus} \simeq 2.5 \times 10^{18} \sqrt{L_{disk}/10^{45} \text{ erg s}^{-1}} = 2.85 \times 10^{18}$ cm (see e.g. Sikora et al. 2009; Hayashida et al. 2012). The values for the luminosity of the accretion disc and the SMBH mass of PKS 0736+017 vary largely in the literature: estimates for L_{disk} range from $10^{44.6}$ to $10^{45.7} \text{ erg s}^{-1}$, while estimates for M_{\bullet} range from 10^8 to $10^{8.7} M_{\odot}$ (Wandel 1991; McLure & Dunlop 2001; Woo & Urry 2002; Marchesini et al. 2004; Dai et al. 2007). The adopted value of $L_{disk} \simeq 1.24 \times 10^{45} \text{ erg s}^{-1}$ is consistent with these previous estimates;

- the BLR is modeled as a spherical shell centered at r_{BLR} , with lower boundary $r_{in} = 0.9 * r_{BLR}$ and outer boundary $r_{out} = 1.1 * r_{BLR}$; as discussed in Böttcher & Els (2016), the choice of the boundaries have negligible effects on the BLR opacity; the opening angle of the dusty torus is assumed to be $\pi/4$ as in Nalewajko et al. (2014).

The first constraint on the location of the γ -ray emitting region comes from opacity to $\gamma - \gamma$ pair-production. The VHE photons can pair-produce over the bright environment of the SMBH and, as shown by several authors (see e.g. Donea & Protheroe 2003; Liu & Bai 2006; Reimer 2007; Tavecchio & Ghisellini 2012; Böttcher & Els 2016; Finke 2016), the absorption can be so severe that the simple detection of VHE photons can be used to exclude an emitting region located at the base of the jet. The γ -ray spectra presented in

Fig. 1 show that the extrapolated *Fermi*-LAT spectrum, once the absorption on the EBL is taken into account, is consistent with the H.E.S.S. detection. This translates into an internal opacity equal to zero, constraining the location of the emitting region well beyond r_{BLR} . However, given the large uncertainty in the simultaneous *Fermi*-LAT spectrum, the most conservative opacity constraint has to be computed using the upper-end of the *Fermi*-LAT extrapolated bow-tie. In this case a break exists between the *Fermi*-LAT and H.E.S.S. energy band. This break can be intrinsic (for example due to a break in the lepton population, or due to the transition from the Thomson to the Klein-Nishina regime of the inverse-Compton scattering), or due to additional pair-production absorption at the source. In particular, while pair-production on the infrared photons from the dusty torus is expected to produce a spectral break at a few TeV, absorption on Ly α photons can produce a spectral break at around 100 GeV. Given that it is not possible to discriminate between an intrinsic cut-off in the γ -ray emission and an absorption effect, the opacity argument can only be used to put a lower limit on the location of the emitting region r_{min} : if the emitting region is located further in, closer to the SMBH, the spectral break between the *Fermi*-LAT and H.E.S.S. observations would have been stronger. To quantitatively estimate r_{min} , the model described in Böttcher & Els (2016) is used, together with the values of L_{disk} and r_{BLR} provided above. The calculation of r_{min} is performed by extrapolating into the VHE band the upper-end of the *Fermi*-LAT bow-tie, including absorption by internal and EBL photons, and varying r until the extrapolated spectrum matches the H.E.S.S. measurement. This estimate results in $r_{min} = 1.6 \times 10^{17}$ cm, or 1.1 r_{BLR} . This limit is shown in Fig. 4 by the red exclusion region.

The second constraint comes from the collimation of the relativistic jet. As shown by radio observations, relativistic jets from SMBHs are highly collimated and in particular $\Gamma \vartheta_{open} < 1$. In the following, the reformulation by Nalewajko et al. (2014) is used, which translates this inequality into a limit in the $\Gamma - r$ plane, under the assumption that $R' \simeq c \tau_{var} \frac{\Gamma}{1+z}$, and that $R'/r = \tan \vartheta_{open}$. This constraint is shown in Fig. 4 by the blue exclusion region.

The last constraint comes from the cooling of the leptons due to inverse Compton scattering. In particular, the cooling time-scale τ_{cool} is required to be shorter than the observed variability time-scale τ_{var} , under the assumption again that $R' \simeq c \tau_{var} \frac{\Gamma}{1+z}$, and that the cooling time-scale is dominated by the inverse Compton scattering on the external photon fields. To translate this condition into an exclusion region in the $\Gamma - r$ plane, it is used a modified version of the equations described in Nalewajko et al. (2014): instead of considering a BLR opening angle of $\pi/4$, as assumed in the original paper, the equations have been recomputed for a spherical BLR, to be consistent with the opacity constraint. The impact of this change is to open up the parameter space, allowing a larger range of Γ values when the emitting region is located close to r_{BLR} . The cooling constraint is shown in Fig. 4 by the orange exclusion region. The cooling time-scale depends on the energy density of the target photons, and thus depends on r : the shape of the exclusion region in Fig. 4 is due to the changes in the photon field seen by the γ -ray emitting region when approaching the BLR radius r_{BLR} and torus radius r_{torus} .

The three aforementioned constraints limit significantly the $\Gamma - r$ plane. A location of the emitting region close to the SMBH, where the dominant photon field is the thermal radiation from the accretion disc, is excluded due to the opacity constraint. Two scenarios are, however, allowed: an emitting region located at around r_{BLR} , with $\Gamma \simeq 10 - 20$, or an emitting region close to r_{torus} , with $\Gamma \simeq 60$. In the first scenario, the dominant external photon field is the emission from the BLR, while in the second case it is the thermal emission from the dusty torus. The first solution implies lower values of Γ , in line with radio observations of γ -ray FSRQs and of PKS 0736+017 in particular, for which an estimate of $\Gamma = 16.5 - 17.0$ is provided by Pushkarev et al. (2009) and Hovatta et al. (2009), respectively, from direct measurements of the jet speed. Changing the assumed value of τ_{var} results in a translation of the constraints towards larger values of Γ , as shown in Fig. 4.

3.2. Spectral Energy Distribution

The SED of PKS 0736+017 during the February 2015 flaring activity, using the data described in the previous section and together with archival observations⁷, is presented in Fig. 5. As discussed previously, it is not possible to fit the overall SED during the night of the H.E.S.S. detection due to the absence of simultaneous information on the behavior of the synchrotron component. It is however interesting to use the constraints on Γ and r computed in the previous section as input parameters for a leptonic model fitting the γ -ray SED. The goal is first to see if it is possible to find a model which remains consistent with the observations during the flare, and for what parameter values (other than Γ and r), and then discuss the implications for the behaviour of the synchrotron component during the flare. For this purpose, the γ -ray emitting region is assumed to be at $r = 1.7 \times 10^{17}$ cm, and located at the limit of the collimation constraint, i.e. $\Gamma \vartheta_{open} = 1$ (the solution is identified as a green star in Fig. 4). The Lorentz factor Γ is equal to 17.7, and the size of the emitting region R is estimated via $\tau_{var} = 6$ hours to be 9.6×10^{15} cm. The ratio of R/r gives a jet opening angle ϑ_{open} of 3.2° (which by construction respects $\Gamma \vartheta_{open} = 1$). The estimate of the jet opening angle of PKS 0736+017 from radio observations is smaller, $\vartheta_{open} = 1.8^\circ \pm 0.3^\circ$ (Pushkarev et al. 2017). This value refers however to the jet opening angle as measured at much larger (kpc) distances from the SMBH. The fact that the intrinsic opening angle gets smaller downstream is consistent with the findings of Pushkarev et al. (2017) from an analysis of 65 AGNs. The leptons in the emitting region scatter primarily BLR photons: the BLR photon energy density at r is estimated following Nalewajko et al. (2014) resulting in $u_{BLR} = 3.95$ erg cm⁻³. The energy density of torus photons is similarly estimated as $u_{torus} = 0.005$ erg cm⁻³, clearly negligible with respect to the BLR one. The numerical code used to compute synchrotron and EIC radiation is described in Cerruti et al. (2013). The electron distribution in the emitting region is modeled by a broken power-law function with exponential cut-off

$$N'_e(\gamma') = \begin{cases} N'_0 \gamma'^{-n_1} & \text{if } \gamma'_{min} \leq \gamma' \leq \gamma'_{break} \\ N'_0 \gamma'^{(n_2-n_1)} \gamma'^{-n_2} e^{-\gamma'/\gamma'_{Max}} & \text{if } \gamma'_{break} < \gamma' \end{cases} \quad (2)$$

with indices n_1 and n_2 below and above the break Lorentz factor γ'_{break} , cut-off Lorentz factor γ'_{Max} , and minimum Lorentz factor $\gamma'_{min} = 10$. A complex BLR line spectrum is assumed,

⁷ from the SSDC SED builder, <https://tools.ssdsc.asi.it>.




Fig. 5. Spectral energy distribution of PKS 0736+017 during the multi-wavelength campaign of February 2015, together with archival data (in grey). The black and red bow-ties represent the H.E.S.S. spectra during the night of February 19, 2015, for the monoscopic and stereoscopic reconstruction, respectively; the blue bow-tie represents the *Fermi*-LAT spectrum simultaneous with the H.E.S.S. detection; the orange bow-ties represents the four *Swift*-XRT spectra, acquired after the H.E.S.S. detection; the red points represent all optical/UV measurements before and after the flare, from ATOM, Steward Observatory and *Swift*-UVOT. The EIC model components, from low to high energies are: synchrotron emission by leptons (in red), synchrotron-self-Compton emission (in green), EIC emission over $\text{Ly}\alpha$ line (in blue, emission from EIC over other BLR lines is not plotted for clarity purposes), EIC emission over the dust torus (in magenta). The violet dotted lines represent the thermal emission from the dusty torus, and the accretion disk, respectively.

using the seven most prominent emission lines from the average quasar model by Telfer et al. (2002). The absorption on BLR photons is calculated following Dermer et al. (2009), under the assumption that the most relevant absorption is the one due to the $\text{Ly}\alpha$ line. Absorption by the EBL is computed using the model by Franceschini et al. (2008). The remaining free parameters are adjusted to reproduce the γ -ray data from *Fermi*-LAT and H.E.S.S., a peak of the synchrotron component $\nu_{\text{sync}} = 8 \times 10^{14}$ Hz, as constrained by the flat optical spectrum measured by Steward Observatory observations 8 hours after the H.E.S.S. detection, and a flux of the synchrotron peak similar to the maximum observed optical flux. A good description of the SED can be obtained assuming $n_1 = 2.7$, $n_2 = 3.4$, $\gamma_{\text{break}} = 2 \times 10^3$, $\gamma_{\text{Max}} = 1 \times 10^5$, $N'_0 = 1 \times 10^6 \text{ cm}^{-3}$, and a magnetic field $B' = 1.6$ G. The corresponding value of the equipartition factor (the ratio of the electron energy density to the magnetic energy density) u'_e/u'_B is 2.3. The resulting modeling is shown in Fig. 5. The peak of the EIC component of the model is in the GeV range, resulting in a flat spectrum in the LAT energy band. The hard HE spectrum is needed in order to fit the VHE spectrum, as both the transition to the Klein-Nishina regime and the absorption on BLR photons lead to an under-prediction of the VHE spectrum for softer HE spectra. The model indicates that an X-ray flare could have occurred simultaneously with the VHE γ -ray flare, with significant softening of the spectrum, due to the emergence of the synchrotron component, similar to the one observed in PKS 1441+25 (Abeysekara et al. 2015; Ahnen et al. 2015). Although the model predicts a soft-X-ray flare a factor of about

ten brighter at 0.1 keV compared to the *Swift*-XRT observations of PKS 0736+017, it is important to note that the electron synchrotron cooling time-scale at these energies is very short: for electrons with Lorentz factor $\gamma = 10^4$ in a magnetic field $B' = 1.6$ G, and moving towards the observer with bulk Lorentz factor $\Gamma = 17.7$, the synchrotron cooling time-scale is only 35 minutes, to be compared with the 24 hours delay of the *Swift*-XRT observations with respect to the time of the H.E.S.S. detection. This prediction of an unseen soft-X-ray flare is solid with respect to the free parameters considered: having fixed Γ , R , and u_{BLR} , the normalization of the particle distribution N'_0 and the magnetic field B' are adjusted to reproduce the peak flux of the synchrotron and EIC components, and γ'_{break} is adjusted to have a synchrotron peak in the optical band. The high value of γ'_{Max} , which is the parameter that implies the occurrence of a simultaneous X-ray flare from the source, is required for the model to reach VHE γ -rays.

4. Conclusions

H.E.S.S. observations of the FSRQ PKS 0736+017, triggered on the basis of a γ -ray flare detected with *Fermi*-LAT, resulted in the discovery of VHE emission from this quasar during the night of February 19, 2015. PKS 0736+017 is the seventh member of the elusive population of FSRQs known to emit VHE photons, the nearest so far. *Fermi*-LAT and H.E.S.S. high-energy flares were accompanied by at least a 1 mag brightening of the non-thermal optical continuum. *Fermi*-LAT observations show

the presence of a relatively fast γ -ray flare, with flux-doubling time-scale of around six hours. No optical nor X-ray observations were performed strictly simultaneously with the H.E.S.S. detection. Nonetheless, both temporal and spectral measurements in the γ -ray band can be used to put model-dependent constraints on the location of the γ -ray emitting region in the jet, and on its Lorentz factor. The *Fermi*-LAT and the H.E.S.S. spectra collectively constrain the location of the emitting region to be located beyond $r = 1.1 r_{BLR} = 1.6 \times 10^{17}$ cm to avoid absorption due to γ - γ pair-production with BLR photons. A location of the γ -ray emitting region just outside the BLR with a Lorentz factor $\Gamma \simeq 10 - 20$ is thus a viable solution which satisfies both temporal and spectral constraints. Alternatively, the emitting region may be located much farther away, at around $r_{torus} = 2.85 \times 10^{18}$ cm, with a higher value of $\Gamma \simeq 60$, with electrons in the jet in this case scattering thermal photons from the dusty torus.

The main limitation of this study is the absence of strictly simultaneous observations of the behavior of the synchrotron component during the H.E.S.S. detection. With simultaneous optical, UV, and X-ray data, it will be possible to uniquely constrain the electron energy distribution, and the opacity and variability constraints could be coupled with a full SED fitting. In addition with tighter constraints on the location of the γ -ray emitting region, it will be possible to study in more details the opacity in the VHE band, potentially putting constraints on the geometry of the BLR itself (its aperture angle and its profile). This kind of study is more easily done using nearby quasars, for which the internal absorption is not hidden by the EBL absorption. In this context, PKS 0736+017 represents the ideal FSRQ, being located much closer than all other VHE FSRQs. Its location near the celestial equator makes it also a perfect target for all current ground-based γ -ray telescopes, with only marginal zenith-angle effects. The next γ -ray flare from PKS 0736+017 has thus the potential to be an extremely interesting event for the study of γ -ray quasars, and future multi-wavelength coordinated observing campaigns are strongly encouraged.

Acknowledgements. The support of the Namibian authorities and of the University of Namibia in facilitating the construction and operation of H.E.S.S. is gratefully acknowledged, as is the support by the German Ministry for Education and Research (BMBF), the Max Planck Society, the German Research Foundation (DFG), the Helmholtz Association, the Alexander von Humboldt Foundation, the French Ministry of Higher Education, Research and Innovation, the Centre National de la Recherche Scientifique (CNRS/IN2P3 and CNRS/INSU), the Commissariat Inergie atomique et auxnergies alternatives (CEA), the U.K. Science and Technology Facilities Council (STFC), the Knut and Alice Wallenberg Foundation, the National Science Centre, Poland grant no. 2016/22/M/ST9/00382, the South African Department of Science and Technology and National Research Foundation, the University of Namibia, the National Commission on Research, Science & Technology of Namibia (NCRST), the Austrian Federal Ministry of Education, Science and Research and the Austrian Science Fund (FWF), the Australian Research Council (ARC), the Japan Society for the Promotion of Science and by the University of Amsterdam. We appreciate the excellent work of the technical support staff in Berlin, Zeuthen, Heidelberg, Palaiseau, Paris, Saclay, Tbingen and in Namibia in the construction and operation of the equipment. This work benefited from services provided by the H.E.S.S. Virtual Organisation, supported by the national resource providers of the EGI Federation. The CC-IN2P3 (cc.in2p3.fr) is gratefully acknowledged for providing a significant amount of the computing resources and services needed for this work. Part of this work is based on archival data, software or online services provided by the Space Science Data Center - ASI. Blazar observations at Steward Observatory are funded though NASA/Fermi Guest Investigator Program grant NNX15AU81G. Matteo Ceruti has received financial support through the Postdoctoral Junior Leader Fellowship Programme from la Caixa Banking Foundation (LCF/BQ/LI18/11630012). The

authors would like to thank the anonymous referee, whose constructive comments significantly improved the manuscript.

References

- Abdo, A. A., Ackermann, M., Agudo, I., et al. 2010, *ApJ*, 716, 30
 Abeysekara, A. U., Archambault, S., Archer, A., et al. 2015, *ApJ*, 815, L22
 Acero, F., Ackermann, M., Ajello, M., et al. 2015, *ApJS*, 218, 23
 Acero, F., Ackermann, M., Ajello, M., et al. 2016, *ApJS*, 223, 26
 Agudo, I., Jorstad, S. G., Marscher, A. P., et al. 2011a, *ApJ*, 726, L13
 Agudo, I., Marscher, A. P., Jorstad, S. G., et al. 2011b, *ApJ*, 735, L10
 Aharonian, F., Akhperjanian, A. G., Barres de Almeida, U., et al. 2007, *A&A*, 475, L9
 Ahnen, M. L., Ansoldi, S., Antonelli, L. A., et al. 2016, *A&A*, 595, A98
 Ahnen, M. L., Ansoldi, S., Antonelli, L. A., et al. 2015, *ApJ*, 815, L23
 Aleksić, J., Ansoldi, S., Antonelli, L. A., et al. 2014, *A&A*, 569, A46
 Aleksić, J., Antonelli, L. A., Antonz, P., et al. 2011, *ApJ*, 730, L8
 Aliu, E., Archambault, S., Archer, A., et al. 2016, *A&A*, 594, A76
 Aliu, E., Archambault, S., Arlen, T., et al. 2014, *ApJ*, 782, 13
 Atwood, W. B., Abdo, A. A., Ackermann, M., et al. 2009, *ApJ*, 697, 1071
 Baldwin, J. A. 1975, *ApJ*, 201, 26
 Becker, P. A. & Kafatos, M. 1995, *ApJ*, 453, 83
 Berge, D., Funk, S., & Hinton, J. 2007, *A&A*, 466, 1219
 Blandford, R. D. & Levinson, A. 1995, *ApJ*, 441, 79
 Blandford, R. D. & Rees, M. J. 1978, *Phys. Scr*, 17, 265
 Błażejowski, M., Sikora, M., Moderski, R., & Madejski, G. M. 2000, *ApJ*, 545, 107
 Blinov, D., Pavlidou, V., Papadakis, I., et al. 2015, *MNRAS*, 453, 1669
 Blinov, D., Pavlidou, V., Papadakis, I. E., et al. 2016, *MNRAS*, 457, 2252
 Böttcher, M. & Els, P. 2016, *ApJ*, 821, 102
 Böttcher, M., Reimer, A., Sweeney, K., & Prakash, A. 2013, *ApJ*, 768, 54
 Brown, A. M. 2013, *MNRAS*, 431, 824
 Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
 Cao, G. & Wang, J.-C. 2013, *MNRAS*, 436, 2170
 Carnerero, M. I., Raiteri, C. M., Villata, M., et al. 2015, *MNRAS*, 450, 2677
 Ceruti, M. 2015, *ArXiv e-prints* 1501.03554
 Ceruti, M., Dermer, C. D., Lott, B., Boisson, C., & Zech, A. 2013, *ApJ*, 771, L4
 Clements, S. D., Jenks, A., & Torres, Y. 2003, *AJ*, 126, 37
 Cohen, D. P., Romani, R. W., Filippenko, A. V., et al. 2014, *ApJ*, 797, 137
 Coogan, R. T., Brown, A. M., & Chadwick, P. M. 2016, *MNRAS*, 458, 354
 Dai, H., Xie, G. Z., Zhou, S. B., et al. 2007, *AJ*, 133, 2187
 D'Ammando, F. & Orienti, M. 2014, *The Astronomer's Telegram*, 6731
 Day, G. A., Shimmins, A. J., Ekers, R. D., & Cole, D. J. 1966, *Australian Journal of Physics*, 19, 35
 De Naurois, M. & Rolland, L. 2009, *Astroparticle Physics*, 32, 231
 Dermer, C. D., Ceruti, M., Lott, B., Boisson, C., & Zech, A. 2014, *ApJ*, 782, 82
 Dermer, C. D., Finke, J. D., Krug, H., & Böttcher, M. 2009, *ApJ*, 692, 32
 Dermer, C. D. & Schlickeiser, R. 1993, *ApJ*, 416, 458
 Dermer, C. D. & Schlickeiser, R. 1994, *ApJS*, 90, 945
 Dermer, C. D., Schlickeiser, R., & Maticchiadis, A. 1992, *A&A*, 256, L27
 Dickey, J. M. & Lockman, F. J. 1990, *ARA&A*, 28, 215
 Donea, A.-C. & Protheroe, R. J. 2003, *Astroparticle Physics*, 18, 377
 Dotson, A., Georganopoulos, M., Meyer, E. T., & McCann, K. 2015, *ApJ*, 809, 164
 Finke, J. D. 2016, *ApJ*, 830, 94
 Finke, J. D. & Dermer, C. D. 2010, *ApJ*, 714, L303
 Franceschini, A., Rodighiero, G., & Vaccari, M. 2008, *A&A*, 487, 837
 Gehrels, N., Chincarini, G., Giommi, P., et al. 2005, *ApJ*, 621, 558
 Ghisellini, G. & Tavecchio, F. 2009, *MNRAS*, 397, 985
 Ghisellini, G., Tavecchio, F., Foschini, L., et al. 2010, *MNRAS*, 402, 497
 Greene, J. E. & Ho, L. C. 2005, *ApJ*, 630, 122
 Hauser, M., Möllenhoff, C., Pühlhofer, G., et al. 2004, *Astronomische Nachrichten*, 325, 659
 Hayashida, M., Madejski, G. M., Nalewajko, K., et al. 2012, *ApJ*, 754, 114
 H.E.S.S. Collaboration, Abdalla, H., Abramowski, A., et al. 2017, *A&A*, 600, A89
 H.E.S.S. Collaboration, Abdalla, H., Adam, R., et al. 2019, *A&A*, 627, A159
 H.E.S.S. Collaboration, Abramowski, A., Acero, F., et al. 2013, *A&A*, 554, A107
 Ho, L. C. & Kim, M. 2009, *ApJS*, 184, 398
 Holler, M., Balzer, A., Chalmé-Calvet, R., et al. 2015, *ArXiv e-prints* 1509.02896
 Hovatta, T., Valtaoja, E., Tornikoski, M., & Lähteenmäki, A. 2009, *A&A*, 494, 527
 Jannuzi, B. T., Smith, P. S., & Elston, R. 1994, *ApJ*, 428, 130
 Jorstad, S. G., Marscher, A. P., Mattox, J. R., et al. 2001, *ApJ*, 556, 738
 Kotilainen, J. K., Falomo, R., & Scarpa, R. 1998, *A&A*, 332, 503
 Lanyi, G. E., Boboltz, D. A., Charlot, P., et al. 2010, *AJ*, 139, 1695

- Lenain, J.-P. 2018, *Astronomy and Computing*, 22, 9
- Li, T.-P. & Ma, Y.-Q. 1983, *ApJ*, 272, 317
- Li, W., Filippenko, A. V., Chornock, R., & Jha, S. 2003, *PASP*, 115, 844
- Lister, M. L., Aller, H. D., Aller, M. F., et al. 2009, *AJ*, 137, 3718
- Lister, M. L. & Homan, D. C. 2005, *AJ*, 130, 1389
- Liu, H. T. & Bai, J. M. 2006, *ApJ*, 653, 1089
- Lyutikov, M. & Kravchenko, E. V. 2017, *MNRAS*, 467, 3876
- MAGIC Collaboration, Acciari, V. A., Ansoldi, S., et al. 2018, *A&A*, 619, A159
- MAGIC Collaboration, Albert, J., Aliu, E., et al. 2008, *Science*, 320, 1752
- Malkan, M. A. & Moore, R. L. 1986, *ApJ*, 300, 216
- Marchesini, D., Celotti, A., & Ferrarese, L. 2004, *MNRAS*, 351, 733
- Marcowith, A., Henri, G., & Pelletier, G. 1995, *MNRAS*, 277, 681
- Mattox, J. R., Bertsch, D. L., Chiang, J., et al. 1996, *ApJ*, 461, 396
- Max-Moerbeck, W., Hovatta, T., Richards, J. L., et al. 2014, *MNRAS*, 445, 428
- McLure, R. J. & Dunlop, J. S. 2001, *MNRAS*, 327, 199
- McLure, R. J., Kukula, M. J., Dunlop, J. S., et al. 1999, *MNRAS*, 308, 377
- Meyer, E. T., Fossati, G., Georganopoulos, M., & Lister, M. L. 2012, *ApJ*, 752, L4
- Mirzoyan, R. 2017, *The Astronomer's Telegram*, 11061
- Mukherjee, R. & VERITAS Collaboration. 2017, *The Astronomer's Telegram*, 11075
- Nalewajko, K., Begelman, M. C., & Sikora, M. 2014, *ApJ*, 789, 161
- Padovani, P. 1992, *MNRAS*, 257, 404
- Padovani, P. & Giommi, P. 1995, *ApJ*, 444, 567
- Parsons, R., Murach, T., & Gajdus, M. 2015, in *International Cosmic Ray Conference*, Vol. 34, 34th International Cosmic Ray Conference (ICRC2015), 826
- Parsons, R. D. & Hinton, J. A. 2014, *Astroparticle Physics*, 56, 26
- Petropoulou, M. & Dimitrakoudis, S. 2015, *MNRAS*, 452, 1303
- Poutanen, J. & Stern, B. 2010, *ApJ*, 717, L118
- Punch, M., Akerlof, C. W., Cawley, M. F., et al. 1992, *Nature*, 358, 477
- Pushkarev, A. B., Kovalev, Y. Y., Lister, M. L., & Savolainen, T. 2009, *A&A*, 507, L33
- Pushkarev, A. B., Kovalev, Y. Y., Lister, M. L., & Savolainen, T. 2017, *MNRAS*, 468, 4992
- Ramakrishnan, V., Hovatta, T., Nieppola, E., et al. 2015, *MNRAS*, 452, 1280
- Rani, B., Lott, B., Krichbaum, T. P., Fuhrmann, L., & Zensus, J. A. 2013, *A&A*, 557, A71
- Reimer, A. 2007, *ApJ*, 665, 1023
- Reimer, A. 2012, in *Journal of Physics Conference Series*, Vol. 355, *Journal of Physics Conference Series*, 012011
- Rolke, W. A., López, A. M., & Conrad, J. 2005, *Nuclear Instruments and Methods in Physics Research A*, 551, 493
- Roming, P. W. A., Kennedy, T. E., Mason, K. O., et al. 2005, *Space Sci. Rev.*, 120, 95
- Roming, P. W. A., Koch, T. S., Oates, S. R., et al. 2009, *ApJ*, 690, 163
- Salamon, M. H. & Stecker, F. W. 1998, *ApJ*, 493, 547
- Sambruna, R. M., Maraschi, L., & Urry, C. M. 1996, *ApJ*, 463, 444
- Schlafly, E. F. & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Schmidt, G. D., Stockman, H. S., & Smith, P. S. 1992, *ApJ*, 398, L57
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48
- Sikora, M., Begelman, M. C., & Rees, M. J. 1994, *ApJ*, 421, 153
- Sikora, M., Stawarz, Ł., Moderski, R., Nalewajko, K., & Madejski, G. M. 2009, *ApJ*, 704, 38
- Sitko, M. L., Schmidt, G. D., & Stein, W. A. 1985, *ApJS*, 59, 323
- Smith, P. S. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 110, *Blazar Continuum Variability*, ed. H. R. Miller, J. R. Webb, & J. C. Noble, 135
- Smith, P. S., Montiel, E., Rightley, S., et al. 2009, *ArXiv e-prints* 0912.3621
- Tavecchio, F. & Ghisellini, G. 2012, *ArXiv e-prints* 1209.2291
- Tavecchio, F., Ghisellini, G., Bonnoli, G., & Ghirlanda, G. 2010, *MNRAS*, 405, L94
- Telfer, R. C., Zheng, W., Kriss, G. A., & Davidsen, A. F. 2002, *ApJ*, 565, 773
- the Fermi-LAT Collaboration. 2019, *ArXiv e-prints* 1902.10045
- Urry, C. M. & Padovani, P. 1995, *PASP*, 107, 803
- Vaughan, S., Edelson, R., Warwick, R. S., & Uttley, P. 2003, *MNRAS*, 345, 1271
- Wakely, S. P. & Horan, D. 2008, *International Cosmic Ray Conference*, 3, 1341
- Wandel, A. 1991, *A&A*, 241, 5
- Weekes, T. C., Cawley, M. F., Fegan, D. J., et al. 1989, *ApJ*, 342, 379
- Woo, J.-H. & Urry, C. M. 2002, *ApJ*, 579, 530
- Wright, S. C., McHardy, I. M., & Abraham, R. G. 1998, *MNRAS*, 295, 799
- Yan, D., Zeng, H., & Zhang, L. 2012, *PASJ*, 64, 80
- Zdziarski, A. A. & Böttcher, M. 2015, *MNRAS*, 450, L21
-
- 1 Centre for Space Research, North-West University, Potchefstroom 2520, South Africa
 - 2 Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France
 - 3 Max-Planck-Institut für Kernphysik, P.O. Box 103980, D 69029 Heidelberg, Germany
 - 4 Dublin Institute for Advanced Studies, 31 Fitzwilliam Place, Dublin 2, Ireland
 - 5 High Energy Astrophysics Laboratory, RAU, 123 Hovsep Emin St Yerevan 0051, Armenia
 - 6 Aix Marseille Université, CNRS/IN2P3, CPPM, Marseille, France
 - 7 Department of Physics, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan
 - 8 Laboratoire d'Annecy de Physique des Particules, Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LAPP, 74000 Annecy, France
 - 9 IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France
 - 10 University of Namibia, Department of Physics, Private Bag 13301, Windhoek, Namibia, 12010
 - 11 DESY, D-15738 Zeuthen, Germany
 - 12 Department of Physics and Electrical Engineering, Linnaeus University, 351 95 Växjö, Sweden
 - 13 School of Physical Sciences, University of Adelaide, Adelaide 5005, Australia
 - 14 LUTH, Observatoire de Paris, PSL Research University, CNRS, Université Paris Diderot, 5 Place Jules Janssen, 92190 Meudon, France
 - 15 Sorbonne Université, Université Paris Diderot, Sorbonne Paris Cité, CNRS/IN2P3, Laboratoire de Physique Nucléaire et de Hautes Energies, LPNHE, 4 Place Jussieu, F-75252 Paris, France
 - 16 Laboratoire Univers et Particules de Montpellier, Université Montpellier, CNRS/IN2P3, CC 72, Place Eugène Bataillon, F-34095 Montpellier Cedex 5, France
 - 17 GRAPPA, Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands
 - 18 Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen Centre for Astroparticle Physics, Erwin-Rommel-Str. 1, D 91058 Erlangen, Germany
 - 19 Astronomical Observatory, The University of Warsaw, Al. Ujazdowskie 4, 00-478 Warsaw, Poland
 - 20 Institut für Astronomie und Astrophysik, Universität Tübingen, Sand 1, D 72076 Tübingen, Germany
 - 21 Instytut Fizyki Jądrowej PAN, ul. Radzikowskiego 152, 31-342 Kraków, Poland
 - 22 School of Physics, University of the Witwatersrand, 1 Jan Smuts Avenue, Braamfontein, Johannesburg, 2050 South Africa
 - 23 Obserwatorium Astronomiczne, Uniwersytet Jagielloński, ul. Orla 171, 30-244 Kraków, Poland
 - 24 Université Bordeaux, CNRS/IN2P3, Centre d'Études Nucléaires de Bordeaux Gradignan, 33175 Gradignan, France
 - 25 Universität Hamburg, Institut für Experimentalphysik, Luruper Chaussee 149, D 22761 Hamburg, Germany
 - 26 APC, AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/Irfu, Observatoire de Paris, Sorbonne Paris

- Cité, 10, rue Alice Domon et Léonie Duquet, 75205 Paris Cedex 13, France
- ²⁷ Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warsaw, Poland
- ²⁸ Institut für Physik und Astronomie, Universität Potsdam, Karl-Liebknecht-Strasse 24/25, D 14476 Potsdam, Germany
- ²⁹ Landessternwarte, Universität Heidelberg, Königstuhl, D 69117 Heidelberg, Germany
- ³⁰ Univ. Grenoble Alpes, CNRS, IPAG, F-38000 Grenoble, France
- ³¹ Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, D 12489 Berlin, Germany
- ³² Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität Innsbruck, A-6020 Innsbruck, Austria
- ³³ Centre for Astronomy, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5, 87-100 Torun, Poland
- ³⁴ Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), The University of Tokyo Institutes for Advanced Study (UTIAS), The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba, 277-8583, Japan
- ³⁵ Department of Physics, University of the Free State, PO Box 339, Bloemfontein 9300, South Africa
- ³⁶ Department of Physics and Astronomy, The University of Leicester, University Road, Leicester, LE1 7RH, United Kingdom
- ³⁷ RIKEN, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan
- ³⁸ Department of Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
- ³⁹ Yerevan Physics Institute, 2 Alikhanian Brothers St., 375036 Yerevan, Armenia
- ⁴⁰ Institut für Theoretische Physik, Lehrstuhl IV: Weltraum und Astrophysik, Ruhr-Universität Bochum, D 44780 Bochum, Germany
- ⁴¹ Steward Observatory, University of Arizona, Tucson, AZ 85721, USA
- ⁴² Now at Institut de Ciències del Cosmos (ICC UB), Universitat de Barcelona (IEEC-UB), Martí Franquès 1, E08028 Barcelona, Spain
- ⁴³ Now at Physik Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland