
Systematic Analysis of Cluster Similarity Indices: Towards Bias-free Cluster Validation

Martijn Gösgens¹ Liudmila Prokhorenkova^{2,3} Alexey Tikhonov⁴

Abstract

There are many cluster similarity indices used to evaluate clustering algorithms and choosing the best one for a particular task is usually an open problem. In this paper, we perform a thorough analysis of this problem: we develop a list of desirable properties (requirements) and theoretically verify which indices satisfy them. In particular, we investigate dozens of pair-counting indices and prove that none of them satisfies all the requirements. Based on our analysis, we propose using the arccosine of the correlation coefficient as a similarity measure and prove that it satisfies almost all requirements (except for one, which is still satisfied asymptotically). This new measure can be thought of as an angle between partitions.

1. Introduction

Clustering is an unsupervised machine learning problem, where the task is to group together objects which are similar to each other. In network analysis, a related problem is called *community detection*, where grouping is based on relations between items (links) and the obtained clusters are expected to be densely interconnected. Clustering is used across various applications including text mining, online advertisement, anomaly detection and many others (Allahyari et al., 2017; Xu & Tian, 2015).

To measure the quality of a clustering algorithm, one usually compares two partitions: predicted and ground truth (the latter one can be obtained, e.g., by human assessors). Nowadays, there are many cluster similarity indices proposed for that, but which one is the best is still a subject of debates (Lei et al., 2017). Depending on an application, different properties of a similarity index could be desirable. In this paper, we formally define requirements that are desirable across different applications, discuss their importance and formally analyze which similarity indices satisfy them.

While many of the ideas discussed in the paper can be applied to all similarity indices, we particularly focus on pair-counting ones (Rand and Jaccard are the most well-known examples). We formally prove that among dozens of known indices only two satisfy all the properties except for being a distance: correlation coefficient and Sokal & Sneath’s first index (Lei et al., 2017). Surprisingly, both indices are rarely used for cluster evaluation. The correlation coefficient has an additional advantage of being easily convertible to a distance measure via the arccosine function. The obtained measure, which can be thought of as an *angle between partitions*, satisfies all the requirements except the *constant baseline*, which is still satisfied asymptotically.

The *constant baseline* requirement is a particular focus of the current research. Informally, a good index should not prefer one candidate partition over another just because it has too large or too small clusters. To the best of our knowledge, we are the first to formalize this requirement. For all indices under consideration we proved whether they satisfy this requirement. For pair-counting indices, we carried some further analysis and defined several particular types of biases. In this aspect, our work improves the results from the recent work by Lei et al. (2017).

2. Related work

Several attempts to the comparative analysis of cluster similarity indices have been made in the literature, both in machine learning and complex networks communities. In particular, the problem of indices favoring clusterings with smaller or larger clusters has been identified in numerous works (Strehl, 2002; Albatineh et al., 2006; Vinh et al., 2009; 2010; Lei et al., 2017). Some remedies to these biases have been proposed: in one attempt (Amelio & Pizzuti, 2015), the similarity index is multiplied by a penalty factor that is small when the difference in the number of clusters is large. In another attempt (Romano et al., 2014), the index is standardized with respect to the number of clusters.

A paper closely related to the current research (Amigó et al., 2009) formulates several constraints (axioms) for cluster similarity indices. Some of the properties are particular cases of those discussed in the

¹Department of Mathematics and Computer Science, Eindhoven University of Technology, Eindhoven, Netherlands
²Yandex, Moscow, Russia ³Moscow Institute of Physics and Technology, Dolgoprudny, Russia ⁴Yandex, Berlin, Germany. Correspondence to: Gösgens M. <research@martijngosgens.nl>.

current paper, while others seem to be strongly application dependent. In the current paper, we give a more comprehensive list of constraints and focus on those which are desirable in a wide range of applications.

While we focus on cluster similarity indices (partition-partition comparisons), some work has been done for graph-partition similarity indices, often referred to as goodness or quality measures. Such indices quantify how well a community structure (given by a partition) fits a graph. The most well-known example of such measure is *modularity* (Newman & Girvan, 2004). Axioms that these measures ought to satisfy are given in (Ben-David & Ackerman, 2009; Van Laarhoven & Marchiori, 2014). Note that all pair-counting indices discussed in this paper can also be used for graph-partition similarity. Furthermore, these indices can also be used for graph-graph comparison. For example, (Donnat & Holmes, 2018) discusses measuring the Hamming distance and Jaccard distance between the edge-sets of graphs. Here, the Hamming distance corresponds to the Mirkin metric (defined in the next section).

3. Background and notation

We assume that there is a set of elements V with size $n = |V|$. A clustering is a partition of V into disjoint subsets. Capital letters A, B, C will be used to name the clusterings and we will represent them as $A = \{A_1, \dots, A_{k_A}\}$ (similarly for B, C), where k_A is the number of clusters in A and A_i , for $i \in [k_A]$, is the set of elements belonging to i -th cluster. If a pair of elements $v, w \in V$ lie in the same cluster in A , we refer to them as an *intra-cluster pair* of A , while *inter-cluster pair* will be used otherwise. The total number of pairs is denoted by $N = \binom{n}{2}$.

The value that an index I assigns to the similarity between the clusterings A and B will be denoted by $I(A, B)$. Similarity indices used throughout the literature generally fall into four categories (Amigó et al., 2009): 1) indices based on set matching such as Purity, Inverse Purity and their harmonic mean (F-measure); 2) indices based on entropy such as Variation of Information (Meilă, 2007), Normalized Mutual Information (Strehl, 2002) and Standardized Mutual Information (Romano et al., 2014); 3) indices based on edit distance; and 4) pair-counting indices such as the well-known Rand (Rand, 1971) and Jaccard (Jaccard, 1912) indices. Some indices are mixtures of the above categories such as BCubed (Amigó et al., 2009). In Appendix A.1 and A.2, we define the indices that will be discussed throughout the current work.

In this paper, a special attention will be given to pair-counting indices. For this, it will be convenient to use an alternative representation for clusterings. Let \vec{A} be the N -dimensional vector indexed by the set of element-pairs, so

that the entry corresponding to (v, w) equals 1 if (v, w) is an intra-cluster pair and 0 otherwise. Note that this representation has some redundancy: whenever u, v and v, w form intra-cluster pairs, we know that u, w must also be an intra-cluster pair. Hence, not every binary vector of length N represents a clustering. The class of N -dimensional binary vectors is, however, isomorphic to the class of undirected graphs on n vertices.

Each index I can be represented as a function of the two vectors \vec{A}, \vec{B} , or, equivalently, of the $N \times 2$ matrix M_{AB} that results from concatenating the two (column-) vectors. Each row of M_{AB} is either 11, 10, 01 or 00. Let the pair-counts $N_{11}, N_{10}, N_{01}, N_{00}$ denote the number of occurrences for each of these rows in M_{AB} . Pair-counting indices are defined as follows.

Definition 1. A similarity index is a pair-counting index if it can be expressed as a function of the four pair-counts $N_{11}, N_{10}, N_{01}, N_{00}$.

Note that even though not every binary vector corresponds to a clustering, pair-counts can be computed for any two binary vectors of equal length. For example, given an undirected graph $G = (V, E)$ and the corresponding incidence vector $\vec{G} = (\mathbf{1}\{\{v, w\} \in E\})_{v, w \in V}$, pair-counting indices can be used to measure similarity between two graphs. Similarly, such indices can be used to measure similarity between a graph and a clustering (a clustering A corresponds to a graph $G(A)$ where each connected component of $G(A)$ is a clique corresponding to a cluster of A). So, one may see a connection between graph and cluster similarity indices. For example, the Mirkin metric is a pair-counting index that coincides with the Hamming distance between the edge-sets of two graphs (Donnat & Holmes, 2018). Another example is the Jaccard graph distance, which turns out to be more appropriate for comparing sparse graphs (Donnat & Holmes, 2018). To sum up, all pair-counting indices and their properties discussed in the current paper can also be applied to graph-graph and graph-partition similarities.

In Appendix A.3 we prove that pair-counting similarity indices can be uniquely characterized by the property of being pair-symmetric.

In (Lei et al., 2017), 26 known pair-counting indices are listed. We have extended this list to Table 4 in Appendix and will only mention indices of special interest throughout the main text. An even more extensive list is given by (Choi et al., 2009). Some of the pair-counting indices have slight variants that are essentially the same. For example, the Hubert Index (Hubert, 1977) can be expressed as a linear transformation of the Rand index as $H(A, B) = 2R(A, B) - 1$. Similarly, the two Wallace indices are related as $W_1(A, B) = W_2(B, A)$. As all the requirements

that are defined in this paper are invariant under linear transformations and interchanging A and B , requirements do not have to be checked for such variants. Therefore, we define the following linear equivalence relation on similarity indices and check the requirements for at most one representative of each equivalence class.

Definition 2. *Two similarity indices I_1, I_2 are linearly equivalent if there exists a nonconstant linear function f such that either $I_1(A, B) = f(I_2(A, B))$ or $I_1(A, B) = f(I_2(B, A))$.*

This definition is symmetric, reflexive and transitive, thus it indeed defines an equivalence relation. Furthermore, it allows us to conveniently restrict to indices for which higher numerical values indicate higher similarity. Table 3 in Appendix lists the equivalence classes containing more than one known index along with their representative indices. Note that our linear equivalence differs from the less restrictive monotonous equivalence given in (Batagelj & Bren, 1995). In the current work, we have to restrict to linear equivalence as the constant baseline requirement is not invariant to non-linear transformations.

4. Requirements for cluster similarity indices

In this section, we motivate and formally define requirements which are desirable for cluster similarity indices. In Table 1, several indices of special interest are listed along with the requirements that are satisfied by them. In Table 2 a similar list is given for pair-counting indices, along with some additional pair-counting-specific requirements. In Appendix B, we give proofs for each entry of the two tables.

Requirement 1. Maximal agreement. The numerical value that an index assigns to a similarity must be easily interpretable. In particular, it should be easy to see whether the candidate clustering is maximally similar to (i.e., coincides with) the ground truth clustering. Formally, we require that $I(A, A)$ is constant and either a strict upper or a strict lower bound of $I(A, B)$ for all $A \neq B$. Note that the equivalence defined in Definition 2 allows us to further assume that $I(A, A)$ is a maximum w.l.o.g. This requirement is easy to check and it is satisfied by almost all indices, except for SMI and Wallace.

Requirement 2. Symmetry. Similarity is intuitively understood as a symmetric concept. Therefore, in general, a good similarity index ought to be symmetric, i.e., $I(A, B) = I(B, A)$ for all partitions A, B .¹

¹However, in some specific applications A and B may have different roles (e.g., ground truth and candidate partitions) and an asymmetric index may be appropriate if there are different consequences of making false positives or false negatives.

As shown in Tables 1 and 2, most indices are symmetric. Examples of asymmetric indices are precision and recall (Wallace). FNMI (Amelio & Pizzuti, 2015) is asymmetric as it is a product of the symmetric NMI with the asymmetric penalty factor $e^{-|k_A - k_B|/k_A}$.

Requirement 3. Monotonicity. When one clustering is changed such that it resembles the other clustering more, the similarity score ought to improve. Hence, we require an index to be monotone w.r.t. changes that increase the similarity.

Definition 3. *For two clusterings A and B , we say that B' is an A -consistent improvement upon B iff $B \neq B'$ and every pair of elements that is agreed on by A and B , is also agreed on by A and B' .*

This leads to the following monotonicity requirement.

Definition 4. *An index I satisfies the monotonicity requirement if for every two clusterings A, B and any B' that is an A -consistent improvement upon B , it holds that $I(A, B') > I(A, B)$.*

Here we again use Definition 2 which allows us to assume that I gives higher numerical values for higher similarity.

Let us now discuss two particular improvements to which this monotonicity applies. Recall that a clustering is defined as a set of clusters (e.g., $B = \{B_1, \dots, B_{k_B}\}$).

- **Perfect split:** We say that B' is a perfect split of B (w.r.t. A) if B' is obtained from B by splitting a single cluster B_1 into two clusters B'_1, B'_2 such that no two elements of the same cluster of A are in different parts of this split, i.e., for all i , $A_i \cap B_1$ is a subset of either B'_1 or B'_2 .
- **Perfect merger:** We say that B' is a perfect merger of B (w.r.t. A) if there exists some A_i and $B_1, B_2 \subset A_i$ such that B' is obtained by merging B_1, B_2 into B'_1 .

It is easily verified that if B' is obtained from B by a sequence of perfect splits and mergers, then B' is an A -consistent improvement of B , since at each step we only create new agreeing pairs between A and B' . The following theorem states that the opposite also holds.

Theorem 1. *B' is an A -consistent improvement upon B iff B' can be obtained from B by a sequence of perfect splits and perfect mergers.*

Proof. Let B' be an A -consistent improvement upon B and define

$$B \otimes B' = \{B_j \cap B'_{j'} \mid B_j \in B, B'_{j'} \in B', B_j \cap B'_{j'} \neq \emptyset\}.$$

We will prove that $B \otimes B'$ can be obtained from B by a sequence of perfect splits, while B' can be obtained

Table 1. Requirements for non-pair-counting similarity indices

	Max. agreement	Symmetry	Monotonicity	Distance	Const. baseline
NMI	✓	✓	✓	✗	✗
NMI _{max}	✓	✓	✗	✓	✗
FNMI	✓	✗	✗	✗	✗
VI	✓	✓	✓	✓	✗
SMI	✗	✓	✗	✗	✓
FMeasure	✓	✓	✓	✗	✗
BCubed	✓	✓	✓	✗	✗

from $B \otimes B'$ by a sequence of perfect mergers. Indeed, the assumption that B' does not introduce new disagreeing pairs guarantees that any $B_j \in B$ can be split into $B_j \cap B'_1, \dots, B_j \cap B'_{k_{B'}}$ without splitting over an intra-cluster pair of A . Now we will prove that B' can be obtained from $B \otimes B'$ by perfect mergers. Suppose there are two $B'_1, B'_2 \in B \otimes B'$ such that both are subsets of some B'_j . Assume this merger is not perfect, then there must be $v \in B'_1, w \in B'_2$ such v, w are in different clusters of A . As v, w are in the same cluster of B' , it follows from the definition of $B \otimes B'$ that v, w must be in different clusters of B . Hence, v, w is an inter-cluster pair in both A and B while it is an intra-cluster pair of B' , contradicting the assumption that B' is an A -consistent improvement of B . This concludes the proof. \square

Note that this monotonicity is a stronger form of the first two constraints defined in (Amigó et al., 2009): *Cluster Homogeneity* is a weaker form of our monotonicity w.r.t. perfect splits while *Cluster Equivalence* is equivalent to our monotonicity w.r.t. perfect mergers. The authors prove that BCubed satisfies their constraints.

Monotonicity is a critical requirement for cluster similarity indices. However, not all indices satisfy this: we have found counterexamples that prove that SMI, FNMI and Wallace do not satisfy our monotonicity requirement. Furthermore, for NMI, whether the monotonicity is satisfied depends on the normalization: the common normalization by the average of the entropies satisfies monotonicity while normalization by the maximum of the entropies does not. See Appendix B.2 for the proofs.

Requirement 4. Distance. For some applications, a distance-interpretation of dissimilarity may be desirable: whenever A is similar to B and B is similar to C , then

²All indices of Table 4 that are excluded from this table, either do not satisfy constant baseline, symmetry or maximal agreement.

 Table 2. Requirements for pair-counting indices. ACB stands for Asymptotic Constant Baseline.²

	Max. agreement	Min. agreement	Symmetry	Monotonicity	Strong monotonicity	Distance	Const. baseline	ACB
R	✓	✓	✓	✓	✓	✓	✗	✗
AR	✓	✗	✓	✓	✓	✗	✓	✓
J	✓	✗	✓	✓	✗	✗	✓	✗
W	✗	✗	✗	✗	✗	✗	✗	✗
D	✓	✗	✓	✓	✗	✗	✗	✗
CC	✓	✓	✓	✓	✓	✗	✓	✓
S&S1	✓	✓	✓	✓	✓	✗	✓	✓
CD	✓	✓	✓	✓	✓	✓	✗	✓

A should also be somewhat similar to C . A function d is a distance metric if it satisfies the three distance axioms: 1) symmetry ($d(A, B) = d(B, A)$); 2) positive-definiteness ($d(A, B) \geq 0$ with equality iff $A = B$); 3) the triangle inequality ($d(A, C) \leq d(A, B) + d(B, C)$). We say that I is linearly transformable to a distance metric if there exists a linearly equivalent index that satisfies these three distance axioms. Note that all three axioms are invariant under re-scaling of d . We have already imposed the symmetry as a separate requirement while the positive-definiteness is equivalent to the maximal agreement requirement. Therefore, whenever I is symmetric and satisfies the maximal agreement requirement, then it satisfies the distance requirement iff $d(A, B) = c_{\max} - I(A, B)$ satisfies the triangle inequality.

Examples of popular similarity distances are Variation of Information and Mirkin metric. In (Vinh et al., 2010) it is proven that when Mutual Information is normalized by the maximum of entropies, the resulting NMI is equivalent to a distance metric. A proof that Jaccard index is equivalent to a distance (Jaccard distance) is given in (Kosub, 2019). See Appendix B.3 for all proofs.

Requirement 5. Constant baseline. Obviously, a good similarity index should not give a preference to a candidate clustering B over another clustering C just because B has very small or very large clusters. This intuition can be formalized using random partitions: assume that we have some ground truth clustering A and two random partitions B and C . While intuitively both random guesses are equally bad approximations of A , it has been known throughout the literature (Albatineh et al., 2006; Vinh et al., 2010; Romano et al., 2014) that some indices tend to give higher scores for random guesses with a larger number of clusters. Ideally, we want the similarity value of a random candidate w.r.t. the ground truth to have a fixed ex-

pected value c_{base} (independent of A). We will formalize this in the following way: Let $S(B)$ denote the specification of the cluster sizes of the clustering B , i.e., $S(B) := [|B_1|, \dots, |B_{k_B}|]$. Here we use the notation $[\dots]$ to denote that it is a multiset, since the sizes and their multiplicities matter but not their order. For a cluster sizes specification s , let $\mathcal{C}(s)$ be the uniform distribution over clusterings B with $S(B) = s$.

Definition 5. *An index I satisfies the constant baseline requirement whenever there exists a constant c_{base} so that for any cluster-sizes specification s and clustering A with $1 < k_A < n$, it holds that $\mathbf{E}_{B \sim \mathcal{C}(s)}[I(A, B)] = c_{\text{base}}$.*

In this definition, we excluded the cases where A is a trivial clustering consisting of either 1 or n clusters. If we would include these cases to the definition, then we would run into problems for $s = S(A)$, as $\mathcal{C}(s)$ would be a constant distribution, surely returning A and any sensible index should have $I(A, A) \neq c_{\text{base}}$.

We will now show that this definition of the constant baseline applies to all symmetric distributions over clusterings.

Definition 6. *We say that a distribution over clusterings \mathcal{B} is element-symmetric if for every two clusterings B and B' that have the same cluster-sizes, \mathcal{B} returns B and B' with equal probabilities.*

In particular, we note that the distribution that assigns each element to one of k clusters uniformly at random is an element-symmetric distribution.

Lemma 1. *Let I be an index with a constant baseline as defined in Definition 5, let A be a clustering with $1 < k_A < n$ and let \mathcal{B} be an element-symmetric distribution. Then $\mathbf{E}_{B \sim \mathcal{B}}[I(A, B)] = c_{\text{base}}$.*

Proof. We write

$$\begin{aligned} & \mathbf{E}_{B \sim \mathcal{B}}[I(A, B)] \\ &= \sum_s \mathbf{P}_{B \sim \mathcal{B}}(S(B) = s) \mathbf{E}_{B \sim \mathcal{B}}[I(A, B) | S(B) = s] \\ &= \sum_s \mathbf{P}_{B \sim \mathcal{B}}(S(B) = s) \mathbf{E}_{B \sim \mathcal{C}(s)}[I(A, B)] \\ &= \sum_s \mathbf{P}_{B \sim \mathcal{B}}(S(B) = s) c_{\text{base}} = c_{\text{base}}, \end{aligned}$$

where the sum ranges over all cluster-sizes of n elements. \square

Note that the Definition 5 may be challenging to verify, as the contingency-distribution for fixed cluster-sizes is combinatorically complex. For pair-counting indices, we developed some additional tools that help proving the constant baseline requirement (see Section 5).

Constant baseline requirement is extremely important in many practical applications: if an index violates this requirement, its optimization may lead to undesirably biased results. Examples of indices that satisfy the constant baseline requirement with $c_{\text{base}} = 0$ are Adjusted Rand index, correlation coefficient and SMI. Sokal&Sneath-1 has a constant baseline at $c_{\text{base}} = \frac{1}{2}$. However, all other considered indices including such popular ones as NMI, Rand and Jaccard do not satisfy this requirement. See Appendix B.4 for the proofs.

5. Requirements for pair-counting indices

In this section, we discuss additional requirements for pair-counting indices. These three requirements can be seen as variants of Maximal agreement, Monotonicity and Constant baseline respectively. Throughout this section, we interchangeably use the notation $I(A, B)$ and $I(N_{11}, N_{10}, N_{01}, N_{00})$.

Requirement 1'. Minimal agreement. The maximal agreement requirement gives a numerical value to high agreement, which helps make the upper range of the index more interpretable. Similarly, a numerical value for low agreement would make the lower range of the index interpretable. For general partitions, minimal agreement is not well defined: it is not clear which partition would be most dissimilar to a given partition. However, referring to Theorem 2, pair-counting indices form a subclass of graph similarity indices. For a given graph $G = (V, E)$, it is clear that the graph most dissimilar to G is its complement $G^C = (V, E^C)$. Comparing a graph to its complement would result in pair-counts $N_{11} = N_{00} = 0$ and $N_{10} + N_{01} = N$.³ This motivates the following definition:

Definition 7. *We define a pair-counting index I to satisfy the minimal agreement requirement if there exists a constant c_{min} so that $I(N_{11}, N_{10}, N_{01}, N_{00}) \geq c_{\text{min}}$ with equality if and only if $N_{11} = N_{00} = 0$.*

Clearly, this requirement is satisfied by Rand, Correlation Coefficient and Sokal&Sneath-1, while it is not satisfied by Jaccard, Wallace and Dice. Less obvious is the fact that Adjusted Rand does not satisfy this requirement. Substituting $N_{11} = N_{00} = 0$ gives the non-constant

$$AR(0, N_{10}, N_{01}, 0) = -\frac{N_{10}N_{01}}{\frac{1}{2}N^2 - N_{10}N_{01}}.$$

Requirement 3'. Strong monotonicity. Note that a perfect split increases N_{00} and decreases N_{01} while a perfect merger increases N_{11} and decreases N_{10} . Therefore, it can

³Note, however, that for a general partition A , there exists no partition B that disagrees on every pair; this is only possible if A consists of either 1 or n clusters.

easily be seen that any pair-counting index that increases when incrementing either N_{11} or N_{00} while decrementing N_{10} or N_{01} , satisfies the monotonicity requirement.

However, we will give a stronger definition of pair-counting monotonicity whose domain is not constrained to the four pair-counting variables summing to N .

Definition 8. A pair-counting index I satisfies strong monotonicity if it increases with N_{11}, N_{00} and decreases with N_{10}, N_{01} .

This stronger monotonicity requirement allows for comparing similarities across different settings. For example, we could compare the similarity between two clusterings A_1, B_1 on n_1 elements with the similarity between A_2, B_2 on n_2 elements, even when $n_1 \neq n_2$. This ability to compare similarity scores across different numbers of elements is similar to the *Few data points* property of SMI (Romano et al., 2014) that allows its scale to have a similar interpretation across different settings.

Examples of indices that satisfy Requirement 3 while not satisfying Requirement 3' are Jaccard and Dice indices, as these are constant w.r.t. N_{00} . In Table 2 we list both monotonicity and strong monotonicity requirements. The proofs can be found in Appendix B.1.

Requirement 5'. Asymptotic Constant Baseline (ACB).

For pair-counting indices, some further analysis of the expected value of the index is possible. Recall constant baseline from Definition 5. Let $m_A = N_{11} + N_{10}, m_B = N_{11} + N_{01}$ be the number of intra-cluster pairs of A and B , respectively, and note that these are constant as A is constant and $B \sim \mathcal{C}(s)$, so that its cluster-sizes are constant. Furthermore, the pair-counts N_{10}, N_{01}, N_{00} are functions of N, m_A, m_B, N_{11} . Hence, to find the expected value of the index, we only need to inspect it as a function of a single random variable N_{11} . For a given pair, the probability that it is an intra-cluster pair of both clusterings is given by $m_A m_B / N^2$, so the expected values of the pair-counts are

$$\begin{aligned} \overline{N_{11}} &:= \frac{m_A m_B}{N}, & \overline{N_{10}} &:= m_A - \overline{N_{11}} \\ \overline{N_{10}} &:= m_B - \overline{N_{11}}, & \overline{N_{00}} &:= N - m_A - m_B + \overline{N_{11}} \end{aligned} \quad (1)$$

To the best of our knowledge, all pair-counting indices that satisfy the constant baseline requirement are linear functions of N_{11} when expressed in terms of N_{11}, m_A, m_B, N . This may be explained by the fact that only the first moment of N_{11} can be expressed in terms of the pair-counting variables while all higher moments depend on the specific cluster-sizes of both clusterings. For these N_{11} -linear indices, substituting the expected pair-counts gives the expected value of the index. This motivates the following relaxation of the constant baseline requirement.

Definition 9. A pair-counting index I satisfies the Asymptotic Constant Baseline requirement if there exists a constant c_{base} so that for all A with $1 < k_A < n$ and for all cluster-size specifications s we have $I(\overline{N_{11}}, \overline{N_{10}}, \overline{N_{01}}, \overline{N_{00}}) = c_{base}$, where the arguments are the expected values of the pair-counts for $B \sim \mathcal{C}(s)$.

Note that the above definition coincides with the earlier constant baseline from Definition 5 if I is a linear function in N_{11} for fixed m_A, m_B, N . The name of the above requirement is justified by the next result, for which we need to make a mild assumption.

Definition 10. An index I is said to be scale-invariant, if it can be expressed as a continuous function of the three variables $p_A := m_A/N, p_B := m_B/N$ and $p_{AB} := N_{11}/N$.

This assumption holds for all indices in Table 2. For such indices, we will write $I^{(p)}(p_{AB}, p_A, p_B)$. Note that when $B \sim \mathcal{C}(s)$ for some s , the values p_A, p_B are constants while p_{AB} is a random variable. Therefore, we may write P_{AB} to stress that this is a random variable.

Theorem 2. Let I be a scale-invariant pair-counting index, let $n \rightarrow \infty$ and consider a sequence of clusterings $A^{(n)}$ and cluster-size specifications $s^{(n)}$. Let $N_{11}^{(n)}, N_{10}^{(n)}, N_{01}^{(n)}, N_{00}^{(n)}$ be the corresponding pair-counts. Then for any $\varepsilon > 0$

$$\mathbf{P} \left(\left| I \left(N_{11}^{(n)}, N_{10}^{(n)}, N_{01}^{(n)}, N_{00}^{(n)} \right) - I \left(\overline{N_{11}^{(n)}}, \overline{N_{10}^{(n)}}, \overline{N_{01}^{(n)}}, \overline{N_{00}^{(n)}} \right) \right| > \varepsilon \right) \rightarrow 0.$$

Proof. Let $P_{AB}^{(n)}, p_A^{(n)}, p_B^{(n)}$ be the corresponding fractions as defined in Definition 10. We prove the equivalent statement

$$I^{(p)} \left(P_{AB}^{(n)}, p_A^{(n)}, p_B^{(n)} \right) - I^{(p)} \left(p_A^{(n)}, p_B^{(n)}, p_A^{(n)}, p_B^{(n)} \right) \xrightarrow{P} 0$$

We first prove that $P_{AB}^{(n)} - p_A^{(n)} p_B^{(n)} \xrightarrow{P} 0$ so that the above follows from the continuous mapping theorem. Chebyshev's inequality gives

$$\mathbf{P} \left(\left| P_{AB}^{(n)} - p_A^{(n)} p_B^{(n)} \right| > \varepsilon \right) \leq \frac{1}{\binom{n}{2} \varepsilon^2} \text{Var} \left(N_{11}^{(n)} \right) \rightarrow 0.$$

The last step follows from the fact that $\text{Var}(N_{11}) = o(n^4)$, as proven in Appendix C.1. \square

This result justifies the usage of the name *Asymptotic Constant Baseline requirement* as indices satisfying it, will converge in probability to c_{base} for $n \rightarrow \infty$. For smaller n , by computing the second order Taylor expansion of the index around $P_{AB} = p_A p_B$, one can approximate the difference from the expected value to its limiting value in terms of the variance of P_{AB} .

Biases of cluster similarity indices. In (Lei et al., 2017), three biases for cluster similarity indices are described: *NCinc* — the average value of a random guess increases monotonously with the Number of Clusters (NC) in the candidate; *NCdec* — the average value of a random guess decreases monotonously with the number of clusters and *ground truth bias* — the direction of the monotonicity depends on the specific Ground Truth (GT). In particular, the authors conclude from numerical experiments that Jaccard suffers from *NCdec* and analytically prove that Rand suffers from *GTbias*, where the direction of the bias depends on the quadratic entropy of the ground truth clustering. We will first argue that these biases are not well defined, then we will redefine them and show how our analysis allows to easily test indices on these biases.

First, we argue that the quantity of interest should not be the number of clusters, but the number of intra-cluster pairs of the candidate. Theorem 2 shows that the asymptotic value of the index depends on the number of intra-cluster pairs of both clusterings. Although in general clusterings with more clusters tend to have less intra-cluster pairs, one can easily construct clusterings that both have many clusters and intra-cluster pairs. For instance, let B be a random clustering consisting of three clusters, each of size $n/3$. Consider constructing B' from B by merging the first two clusters and splitting the third cluster into $n/3$ clusters of size 1. The number of clusters increases by $n/3 - 2$ while the number of intra-cluster pairs increases by $(n/3)^2 - \binom{n/3}{2}$. Hence, the expected value of the Jaccard index increases with respect to any ground truth A . In contrast, (Lei et al., 2017) classifies Jaccard as an *NCdec* index, so that expected value should increase, contradicting the definition of *NCdec*.

Therefore, we rename *NCinc* to *NPdec* and *NCdec* to *NPinc*, where NP stands for Number of Pairs. We will give the following definitions for these biases.

Definition 11. Let I be a scale-invariant pair-counting index whose corresponding $I^{(p)}$ is differentiable w.r.t. p_B and p_{AB} . Then we define the following biases

- i I suffers from *NPinc* bias if there are $p_A, p_B \in (0, 1)$ such that $\frac{d}{dp_B}[I^{(p)}(p_{APB}, p_A, p_B)] > 0$;
- ii I suffers from *NPdec* bias if there are $p_A, p_B \in (0, 1)$ such that $\frac{d}{dp_B}[I^{(p)}(p_{APB}, p_A, p_B)] < 0$;
- iii I suffers from ground truth bias if it suffers from both *NPinc* and *NPdec* biases.

Applying the previous definition to Jaccard $J^{(p)}(p_{APB}, p_A, p_B) = \frac{p_{APB}}{p_A + p_B - p_{APB}}$ and Rand $R^{(p)}(p_{APB}, p_A, p_B) = 1 - p_A - p_B + 2p_{APB}$ immediately shows that Jaccard suffers from *NPinc* bias and Rand suffers from ground truth bias, confirming the

findings of (Lei et al., 2017). Furthermore, the direction of the monotonicity for the ground truth bias of Rand is now determined by the condition $p_A > \frac{1}{2}$ instead of the more complicated but equivalent condition on the quadratic entropy of A that is given in (Lei et al., 2017). Note that an index satisfying the Asymptotic Constant Baseline requirement will not have any of these biases as $I^{(p)}(p_{APB}, p_A, p_B) = c_{\text{base}}$.

6. Correlation Distance

Although there are some works (Lei et al., 2017; Choi et al., 2009) listing Pearson correlation as a cluster similarity index, it has not received attention that our results suggest it deserves. The correlation coefficient satisfies all requirements except being a distance. In this section, we will show how a monotone transformation of the correlation coefficient results in an index that may be even more suitable in applications where the distance requirements is important. When taking the arccosine of the coefficient, the resulting index is a distance metric, at the cost of not satisfying the exact constant baseline. It does, however, still satisfy an asymptotic constant baseline and we will prove that its expectation is very close to being constant. To the best of our knowledge, this Correlation Distance has never before been used as similarity index for comparing clusterings throughout the literature.

We define Correlation Distance (CD) as

$$\text{CD}(A, B) := \frac{1}{\pi} \arccos \text{CC}(A, B),$$

where CC is the Pearson correlation coefficient, which has scale-invariant representation

$$\text{CC}^{(p)}(P_{AB}, p_A, p_B) = \frac{P_{AB} - p_A p_B}{\sqrt{p_A(1-p_A)p_B(1-p_B)}}.$$

The factor $\frac{1}{\pi}$ scales the index to $[0, 1]$.

This monotone arccosine transformation only affects the (exact) constant baseline and distance requirements, the rest of the requirements it inherits from the correlation coefficient. We first verify that this index is indeed a distance metric:

Theorem 3. *The Correlation Distance is indeed a distance.*

Proof. First we map each partition A to an N -dimensional vector on the unit sphere by

$$\vec{u}(A) := \begin{cases} \frac{1}{\sqrt{N}} \mathbf{1} & \text{if } k_A = 1, \\ \frac{A - p_A \mathbf{1}}{\|A - p_A \mathbf{1}\|} & \text{if } 1 < k_A < n, \\ -\frac{1}{\sqrt{N}} \mathbf{1} & \text{if } k_A = n, \end{cases}$$

where $\mathbf{1}$ is the N -dimensional all-one vector and \vec{A} is the binary vector representation of a partition introduced in Section 3. Straightforward computation gives

$$\|\vec{A} - p_A \mathbf{1}\| = \sqrt{N p_A (1 - p_A)},$$

and standard inner product

$$\langle \vec{A} - p_A \mathbf{1}, \vec{B} - p_B \mathbf{1} \rangle = p_{AB} - p_A p_B,$$

so that indeed

$$\frac{\langle \vec{A} - p_A \mathbf{1}, \vec{B} - p_B \mathbf{1} \rangle}{\|\vec{A} - p_A \mathbf{1}\| \|\vec{B} - p_B \mathbf{1}\|} = CC^{(p)}(p_{AB}, p_A, p_B).$$

It is a well-known fact that the inner product of two vectors of unit length corresponds to the cosine of their angle. Hence, taking the arccosine gives us the angle. The angle between unit vectors corresponds to the distance along the unit sphere. As \vec{u} is an injection from the set of partitions to points on the unit sphere, we may conclude that this index is indeed a distance on the set of partitions. \square

As the correlation has a constant baseline at 0, CD has an asymptotic constant baseline at $\frac{1}{2}$. The expected deviation from this baseline is given by the following theorem.

Theorem 4. *Given ground truth A with a number of clusters $1 < k_A < n$, a cluster-size specification s and a random partition $B \sim \mathcal{C}(s)$, the expected difference between Correlation Distance and its baseline is given by*

$$\begin{aligned} & \mathbf{E}_{B \sim \mathcal{C}(s)}[\text{CD}(A, B)] - \frac{1}{2} \\ &= -\frac{1}{\pi} \sum_{k=1}^{\infty} \frac{(2k)!}{2^{2k} (k!)^2} \frac{\mathbf{E}_{B \sim \mathcal{C}(s)}[\text{CC}(A, B)^{2k+1}]}{2k+1}. \end{aligned}$$

Proof. We take the Taylor expansion of the arccosine around $\text{CC}(A, B) = 0$ and get

$$\text{CD}(A, B) = \frac{1}{2} - \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{2^{2k} (k!)^2} \frac{\text{CC}(A, B)^{2k+1}}{2k+1}.$$

We take the expectation of both sides and note that the first moment of CC equals zero so that the starting index changes to $k = 1$. Rearranging terms gives the desired result. \square

This result explains that in practice, the mean tends to be very close to the asymptotic baseline. To the best of our knowledge, there does not exist a cluster similarity index that is a distance while having an exact constant baseline. For applications where both these properties are needed, CD seems to be the most suitable index.

7. Conclusion

In this paper, we have formally defined several requirements for cluster validation indices. The most important and non-trivial ones are monotonicity and constant baseline. While monotonicity reflects the core idea of similarity indices (it requires better values for more similar partitions), constant baseline is less trivial, but equally important, since violating it in practice can lead to unexpectedly biased algorithms. In Table 1, it can be seen that among the non-pair-counting indices considered, Standardized Mutual Information is the only index that has a constant baseline, yet it does not satisfy monotonicity. Among pair-counting indices, these two requirements are satisfied by the Adjusted Rand Index (AR), Sokal&Sneath-1 (S&S1) and the Correlation Coefficient (CC). With respect to the proposed requirements, AR is dominated by S&S-1 and CC. The only requirement that is not satisfied by the latter two is distance. CC has the advantage of having an easier interpretation than S&S1. We have shown that CC can be monotonously transformed into a Correlation Distance (CD) that satisfies all requirements except constant baseline, which is still satisfied asymptotically. Furthermore, the difference between the expected value of CD and its asymptotic value is shown to be negligible. We conclude that in applications where a distance is desirable, we advise using the Correlation Distance while in situations where this is not necessary, we advise using the Correlation Coefficient.

References

- Albatineh, A. N., Niewiadomska-Bugaj, M., and Michalko, D. On similarity indices and correction for chance agreement. *Journal of Classification*, 23(2): 301–313, Sep 2006. ISSN 1432-1343. doi: 10.1007/s00357-006-0017-z.
- Allahyari, M., Pouriyeh, S., Assefi, M., Safaei, S., Trippe, E. D., Gutierrez, J. B., and Kochut, K. A brief survey of text mining: Classification, clustering and extraction techniques. *arXiv preprint arXiv:1707.02919*, 2017.
- Amelio, A. and Pizzuti, C. Is normalized mutual information a fair measure for comparing community detection methods? In *Proceedings of the 2015 IEEE/ACM International Conference on Advances in Social Networks Analysis and Mining 2015*, pp. 1584–1585. ACM, 2015.
- Amigó, E., Gonzalo, J., Artiles, J., and Verdejo, F. A comparison of extrinsic clustering evaluation metrics based on formal constraints. *Information retrieval*, 12(4):461–486, 2009.
- Batagelj, V. and Bren, M. Comparing resemblance mea-

- asures. *Journal of Classification*, 12(1):73–90, Mar 1995. ISSN 1432-1343. doi: 10.1007/BF01202268.
- Ben-David, S. and Ackerman, M. Measures of clustering quality: A working set of axioms for clustering. In *Advances in neural information processing systems*, pp. 121–128, 2009.
- Choi, S., Cha, S.-H., and Tappert, C. A survey of binary similarity and distance measures. *J. Syst. Cybern. Inf.*, 8, 11 2009.
- Donnat, C. and Holmes, S. Tracking network dynamics: a survey of distances and similarity metrics, 2018.
- Hubert, L. Nominal scale response agreement as a generalized correlation. *British Journal of Mathematical and Statistical Psychology*, 30(1):98–103, 1977. doi: 10.1111/j.2044-8317.1977.tb00728.x.
- Jaccard, P. The distribution of the flora in the alpine zone.1. *New Phytologist*, 11(2):37–50, 1912. doi: 10.1111/j.1469-8137.1912.tb05611.x.
- Kosub, S. A note on the triangle inequality for the jaccard distance. *Pattern Recognition Letters*, 120:36–38, 2019. ISSN 0167-8655. doi: https://doi.org/10.1016/j.patrec.2018.12.007.
- Lei, Y., Bezdek, J. C., Romano, S., Vinh, N. X., Chan, J., and Bailey, J. Ground truth bias in external cluster validity indices. *Pattern Recognition*, 65:58–70, 2017. ISSN 0031-3203. doi: https://doi.org/10.1016/j.patcog.2016.12.003.
- Meilä, M. Comparing clusterings an information based distance. *Journal of multivariate analysis*, 98(5):873–895, 2007.
- Newman, M. E. and Girvan, M. Finding and evaluating community structure in networks. *Physical review E*, 69(2):026113, 2004.
- Rand, W. M. Objective criteria for the evaluation of clustering methods. *Journal of the American Statistical Association*, 66(336):846–850, 1971. doi: 10.1080/01621459.1971.10482356.
- Romano, S., Bailey, J., the vinh, N., and Verspoor, K. Standardized mutual information for clustering comparisons: One step further in adjustment for chance. *31st International Conference on Machine Learning, ICML 2014*, 4: 2873–2882, 01 2014.
- Strehl, A. *Relationship-based clustering and cluster ensembles for high-dimensional data mining*. PhD thesis, 2002.
- Van Laarhoven, T. and Marchiori, E. Axioms for graph clustering quality functions. *The Journal of Machine Learning Research*, 15(1):193–215, 2014.
- Vinh, N. X., Epps, J., and Bailey, J. Information theoretic measures for clusterings comparison: is a correction for chance necessary? In *Proceedings of the 26th annual international conference on machine learning*, pp. 1073–1080. ACM, 2009.
- Vinh, N. X., Epps, J., and Bailey, J. Information theoretic measures for clusterings comparison: Variants, properties, normalization and correction for chance. *Journal of Machine Learning Research*, 11(Oct):2837–2854, 2010.
- Xu, D. and Tian, Y. A comprehensive survey of clustering algorithms. *Annals of Data Science*, 2(2):165–193, 2015.

A. Cluster similarity indices

A.1. General indices

Here we give the definitions of the indices listed in Table 1. We define the contingency variables as $n_{ij} = |A_i \cap B_j|$. We note that all indices discussed in this paper can be expressed as functions of these contingency variables.

The *F-Measure* is defined as the harmonic mean of recall and precision. Recall is defined as

$$r(A, B) = \frac{1}{n} \sum_{i=1}^{k_A} \max_{j \in [k_B]} \{n_{ij}\},$$

and precision is its symmetric counterpart $r(B, A)$.

In (Amigó et al., 2009), recall is redefined as

$$r'(A, B) = \frac{1}{n} \sum_{i=1}^{k_A} \frac{\max_{j \in [k_B]} \{n_{ij}^2\}}{|A_i|},$$

and *BCubed* is defined as the harmonic mean of $r'(A, B)$ and $r'(B, A)$.

The remainder of the indices are information-theoretic and require some additional definitions. Let p_1, \dots, p_ℓ be a discrete distribution (i.e., all values are nonnegative and sum to 1). The Shannon entropy is then defined as

$$H(p_1, \dots, p_\ell) := - \sum_{i=1}^{\ell} p_i \log(p_i).$$

The entropy of a clustering is defined as the entropy of the cluster-label distribution of a random item, i.e.,

$$H(A) := H(|A_1|/n, \dots, |A_{k_A}|/n),$$

and similarly for $H(B)$. The joint entropy $H(A, B)$ is then defined as the entropy of the distribution with probabilities $(p_{ij})_{i \in [k_A], j \in [k_B]}$, where $p_{ij} = n_{ij}/n$.

Variation of Information (Meilä, 2007) is defined as

$$VI(A, B) = 2H(A, B) - H(A) - H(B).$$

Mutual information is defined as

$$M(A, B) = H(A) + H(B) - H(A, B).$$

The mutual information between A and B is upper-bounded by $H(A)$ and $H(B)$, which gives multiple possibilities to normalize the mutual information. In this paper, we discuss two normalizations: normalization by the average of the entropies $\frac{1}{2}[H(A) + H(B)]$, and normalization by the maximum of entropies $\max\{H(A), H(B)\}$. We will refer to the corresponding indices as NMI and NMI_{\max} , respectively:

$$\begin{aligned} \text{NMI}(A, B) &= \frac{2M(A, B)}{H(A) + H(B)}, \\ \text{NMI}_{\max}(A, B) &= \frac{M(A, B)}{\max\{H(A), H(B)\}}. \end{aligned}$$

Fair NMI is a variant of NMI that includes a factor that penalizes large differences in the number of clusters (Amelio & Pizzuti, 2015). It is given by

$$\text{FNMI}(A, B) = e^{|k_A - k_B|/k_A} \text{NMI}(A, B).$$

In this definition, NMI may be normalized in various ways. We note that a different normalization would not result in more requirements being satisfied.

Standardized Mutual Information standardizes the mutual information w.r.t. random permutations of the items (Romano et al., 2014), i.e.,

$$\text{SMI}(A, B) = \frac{M(A, B) - \mathbf{E}_{B' \sim \mathcal{C}(S(B))}[M(A, B')]}{\sigma_{B' \sim \mathcal{C}(S(B))}(M(A, B'))},$$

where σ denotes the standard deviation. Calculating the expected value and standard deviation of the mutual information is nontrivial and requires significantly more computation power than other indices. For this, we refer to the original paper (Romano et al., 2014).

A.2. Pair-counting indices and their equivalences

Pair-counting similarity indices are defined in Table 4. Table 3 lists linearly equivalent indices (see Definition 2).

Table 3. Equivalent pair-counting indices

Representative Index	Equivalent indices
Rand	Adjusted Mirkin Metric, Hubert
Jaccard	Jaccard Distance
Wallace1	Wallace2
Kulczynski	McConnaughey

A.3. Defining subclass of pair-counting indices

In this section, we show that a subclass of pair-counting similarity indices can be uniquely defined by the property of being pair-symmetric.

For two graph G_1, G_2 let $M_{G_1 G_2}$ denote the $N \times 2$ matrix that is obtained by concatenating the two adjacency vectors. Let us write $I_M^{(G)}(M_{G_1 G_2})$ for the similarity between two graphs G_1, G_2 according to some graph similarity index $I^{(G)}$. We will now characterize pair-counting similarity indices as a subclass of the class of similarity indices between undirected graphs.

Definition 12. We define a graph similarity index $I_M^{(G)}(M_{G_1 G_2})$ to be pair-symmetric if interchanging two rows of M_{G_1, G_2} leaves the index unchanged.

We give the following result.

Lemma 2. The class of pair-symmetric graph similarity indices coincides with the class of pair-counting cluster similarity indices.

Proof. A matrix is an ordered list of its rows. An unordered list is a multiset. Hence, when we disregard the ordering of the matrix M_{AB} , we get a multiset of the rows. This multiset contains at most four distinct elements, each corresponding to one of the pair-counts. Therefore, each $I_M^{(G)}(M_{AB})$ that is symmetric w.r.t. interchanging rows is equivalently a function of the pair-counts of A and B . \square

B. Checking requirements for indices

Here all non-trivial requirements will be checked for all indices. The requirements of symmetry, maximal/minimal agreement and asymptotic constant baseline can trivially be tested by simply checking $I(B, A) = I(A, B)$, $I(A, A) = c_{\max}$, $I(0, N_{10}, N_{01}, 0) = c_{\min}$ and $I^{(p)}(p_A p_B, p_A, p_B)$ respectively.

B.1. Strong monotonicity

B.1.1. POSITIVE CASES

Correlation Coefficient and Sokal&Sneath-1. These two indices have the property that inverting one of the bi-

nary vectors results in the index flipping sign. Therefore, we only need to prove that these indices are increasing in N_{11} . For Sokal&Sneath-1, this can easily be seen by the fact that two of the four terms are increasing in N_{11} while the other two terms are constant w.r.t. N_{11} . For the Correlation coefficient, it is less obvious. We will take the derivative but first omit the factor $[(N_{00} + N_{10})(N_{00} + N_{01})]^{-\frac{1}{2}}$. We get

$$\begin{aligned} & \frac{\partial}{\partial N_{11}} \frac{N_{11}N_{00} - N_{10}N_{01}}{\sqrt{(N_{11} + N_{10})(N_{11} + N_{01})}} \\ &= \frac{N_{00}}{\sqrt{(N_{11} + N_{10})(N_{11} + N_{01})}} \\ & \quad - \frac{(N_{11}N_{00} - N_{10}N_{01}) \cdot \frac{1}{2}(2N_{11} + N_{10} + N_{01})}{[(N_{11} + N_{10})(N_{11} + N_{01})]^{1.5}} \\ &= \frac{\frac{1}{2}N_{11}N_{00}(N_{10} + N_{01}) + N_{00}N_{10}N_{01}}{[(N_{11} + N_{10})(N_{11} + N_{01})]^{1.5}} \\ & \quad + \frac{\frac{1}{2}N_{10}N_{01}(2N_{11} + N_{10} + N_{01})}{[(N_{11} + N_{10})(N_{11} + N_{01})]^{1.5}} > 0. \end{aligned}$$

Correlation Distance. The correlation distance satisfies the strong monotonicity as it is a monotone transformation of the correlation coefficient, which satisfies the requirement.

Rand Index. For the Rand index, it can be seen that the derivatives w.r.t. N_{11}, N_{00} are positive, while for N_{10}, N_{01} they are negative. Hence, strong monotonicity is satisfied.

Adjusted Rand. For this index, the analysis can be simplified by noting that the index is given by

$$\frac{f}{f + \frac{N_{10} + N_{01}}{2}},$$

where

$$f = N_{11} - \frac{(N_{11} + N_{10})(N_{11} + N_{01})}{N_{11} + N_{10} + N_{01} + N_{00}}.$$

Therefore, for the index to be increasing in N_{11}, N_{00} , we only need f to be increasing in these variables. Increasingness in N_{00} follows from the fact that this variable only appears in the denominator. For N_{11} , we have

$$\begin{aligned} & \frac{\partial}{\partial N_{11}} f \\ &= 1 - \frac{N_{11} + N_{01}}{N_{11} + N_{10} + N_{01} + N_{00}} - \frac{N_{11} + N_{10}}{N_{11} + N_{10} + N_{01} + N_{00}} \\ & \quad + \frac{(N_{11} + N_{10})(N_{11} + N_{01})}{(N_{11} + N_{10} + N_{01} + N_{00})^2} \\ &= \frac{(N_{00} + N_{10})(N_{00} + N_{01})}{(N_{11} + N_{10} + N_{01} + N_{00})^2} > 0. \end{aligned}$$

We now analyse the index w.r.t. N_{01}

$$\begin{aligned} & \frac{\partial}{\partial N_{01}} \frac{f}{f + \frac{N_{10} + N_{01}}{2}} \\ &= \frac{\left(\frac{\partial}{\partial N_{01}} f\right) \left(f + \frac{N_{10} + N_{01}}{2}\right) - f \left(\frac{\partial}{\partial N_{01}} f + \frac{1}{2}\right)}{\left(f + \frac{N_{10} + N_{01}}{2}\right)^2} \\ &= \left(f + \frac{N_{10} + N_{01}}{2}\right)^{-2} \left(\frac{N_{10} + N_{01}}{2} \frac{\partial}{\partial N_{01}} f - \frac{1}{2}f\right). \end{aligned}$$

Decreasingness in N_{01} will follow from $\frac{\partial}{\partial N_{01}} f < 0$

$$\begin{aligned} \frac{\partial}{\partial N_{01}} f &= -\frac{N_{11} + N_{10}}{N_{11} + N_{10} + N_{01} + N_{00}} \\ & \quad + \frac{(N_{11} + N_{10})(N_{11} + N_{01})}{(N_{11} + N_{10} + N_{01} + N_{00})^2} \\ &= -\frac{(N_{11} + N_{10})(N_{00} + N_{10})}{(N_{11} + N_{10} + N_{01} + N_{00})^2} < 0. \end{aligned}$$

Decreasingness in N_{10} follows from symmetry.

B.1.2. NEGATIVE CASES

Jaccard, Wallace and Dice. All these three indices are constant w.r.t. N_{00} . Therefore, these indices do not satisfy the strong monotonicity.

B.2. Monotonicity

B.2.1. POSITIVE CASES

Rand, Adjusted Rand, Correlation coefficient, Sokal&Sneath-1, Correlation Distance. Strong monotonicity implies monotonicity. Therefore, these pair-counting indices satisfy the monotonicity requirement.

Jaccard and Dice. It can easily be seen that these indices are increasing in N_{11} while decreasing in N_{10}, N_{01} . For increasingness in N_{00} we note that whenever N_{00} gets increased, either N_{10} or N_{01} must decrease, resulting in an increase in the index. Therefore these indices satisfy monotonicity.

NMI and VI. Let B' be a obtained by a perfect split of a cluster B_1 into B'_1, B'_2 . Note that this increases the entropy of the candidate while keeping the joint entropy constant. Let us denote this increase in the candidate entropy by $H(B'|B) = H(B') - H(B) > 0$. Now, for NMI, the numerator will increase by $H(B'|B)$ while the denominator will increase by at most $H(B'|B)$ (dependent on $H(A)$ and the specific normalization that is used). Therefore, NMI will increase. Similarly, VI will decrease by $H(B'|B)$. Concluding, both NMI and VI are monotonous w.r.t. perfect splits. Now let B'' be obtained by a perfect merger of B_1, B_2 into B''_1 . This results in a difference of the entropy

of the candidate $H(B'') - H(B) = -H(B|B'') < 0$. The joint entropy will decrease by the same amount so that the mutual information remains unchanged. Therefore, the numerator of NMI will remain unchanged while the denominator may or may not change, depending on the normalization. For min- or max-normalization, it may remain unchanged while for any other average it will increase. Hence, NMI does not satisfy monotonicity w.r.t. perfect mergers for min- and max-normalization but does satisfy this for average-normalization. For VI, the distance will decrease by $H(B|B'')$ so that it indeed satisfies monotonicity w.r.t. perfect mergers.

FMeasure BCubed. Note that a perfect merger increases recall while leaving precision unchanged and that a perfect split increases precision while leaving recall unchanged. Idem for BCubed recall and BCubed precision. The harmonic mean will increase all of these cases.

B.2.2. NEGATIVE CASES

FNMI We will give the following numerical counter-example: Consider $A = \{\{0, 1\}, \{2\}, \{3\}\}$, $B = \{\{0\}, \{1\}, \{2, 3\}\}$ and merge the first two clusters to obtain $B' = \{\{0, 1\}, \{2, 3\}\}$. This results in

$$FNMI(A, B) \approx 0.67 > 0.57 \approx FNMI(A, B').$$

This non-monotonicity is caused by the severe penalty factor that equals 1 on the l.h.s. while it equals $\exp(-1/3) \approx 0.72$ on the r.h.s..

SMI. For this numerical counter-example we rely on the Matlab-implementation of the index by its original authors (Romano et al., 2014). Let $A = \{\{0, \dots, 4\}, \{5\}\}$, $B = \{\{0, 1\}, \{2, 3\}, \{4\}, \{5\}\}$ and consider merging the two clusters resulting in $B' = \{\{0, 1, 2, 3\}, \{4\}, \{5\}\}$. The index remains unchanged and equals 2 before and after the merger.

Wallace. Let $k_A = n$ and let $k_B < n$. Then any split of B will be a perfect split yet $W(A, B) = 1$ so that no increase will occur.

B.3. Distance

B.3.1. POSITIVE CASES

NMI and VI. In (Vinh et al., 2010) it is proven that for max-normalization, $1 - NMI$ is a distance while in (Meilä, 2007) it is proven that VI is a distance.

Rand. The Mirkin metric $1 - R$ corresponds to a rescaled version of the size of the symmetric difference between the sets of intra-cluster pairs. The symmetric difference is known to be a distance metric.

Jaccard. In (Kosub, 2019) it is proven that the Jaccard distance $1 - J$ is indeed a distance.

Correlation distance. In Theorem 3 of the main text it has been proven that the correlation distance is indeed a distance.

B.3.2. NEGATIVE CASES

FNMI and Wallace. These indices cannot be transformed to distances as they are not symmetric.

SMI. SMI does not satisfy the maximal agreement requirement and can therefore not be transformed to a metric.

FMeasure and BCubed. We will use a simple counter-example where $|V| = 3$, $k_A = 1$, $k_B = 2$, $k_C = 3$. In this case

$$1 - FM(A, C) = 1 - 0.5 > (1 - 0.8) + (1 - 0.8) \\ = (1 - FM(A, B)) + (1 - FM(B, C)),$$

and

$$1 - BC(A, C) = 1 - 0.5 > (1 - 0.71) + (1 - 0.8) \\ \approx (1 - BC(A, B)) + (1 - BC(B, C)),$$

so that both indices violate the triangle inequality in this case.

Adjusted Rand, Dice, Correlation Coefficient and Sokal&Sneath-1. For these indices, we use the following counter-example: Let $A = \{\{0, 1\}, \{2\}, \{3\}\}$, $B = \{\{0, 1\}, \{2, 3\}\}$, $C = \{\{0\}, \{1\}, \{2, 3\}\}$. Then $p_{AB} = p_{BC} = 1/6$ and $p_{AC} = 0$ while $p_A = p_C = 1/6$ and $p_B = 1/3$. By substituting these variables, one can see that

$$1 - I^{(p)}(p_{AC}, p_A, p_C) \\ > (1 - I^{(p)}(p_{AB}, p_A, p_B)) + (1 - I^{(p)}(p_{BC}, p_B, p_C)),$$

holds for each of these indices, contradicting the triangle inequality.

B.4. Constant baseline

B.4.1. POSITIVE CASES

SMI. As SMI is standardized, it satisfies the constant baseline requirement by construction.

Adjusted Rand, Correlation Coefficient and Sokal&Sneath-1. These indices all satisfy ACM while being P_{AB} -linear for fixed p_A, p_B . Therefore, the expected value equals the asymptotic constant.

B.4.2. NEGATIVE CASES

For all the following indices, we will analyse the following counter-example. Let $|V| = n, k_A = k_B = n - 1$. For each index, we will compute the expected value and show that it is not constant. All of these indices satisfy the maximal agreement requirement and maximal agreement is achieved with probability $1/N$ (the probability that the single intra-pair of A coincides with the single intra-pair of B). Furthermore, each case where the intra-pairs do not coincide will result in the same contingency variables and hence the same value of the index. We will refer to this value as $c_n(I)$. Therefore, the expected value will only have to be taken over two values and will be given by

$$\mathbf{E}[I(A, B)] = \frac{1}{N}c_{\max} + \frac{N-1}{N}c_n(I).$$

For each of these indices we will conclude that this is a non-constant function of n so that the index does not satisfy the constant baseline requirement.

Jaccard and Dice. For both these indices we have $c_{\max} = 1$ and $c_n(I) = 0$ (as $N_{11} = 0$ whenever the intra-pairs do not coincide). Hence, $\mathbf{E}[I(A, B)] = \frac{1}{N}$, which is not constant.

Rand and Wallace. As both functions are linear in P_{AB} , we can compute the expected value by simply substituting $P_{AB} = p_A p_B$. This will result in expected values $1 - 2/N + 2/N^2$ and $1/N$ for Rand and Wallace respectively, which are both non-constant.

Correlation distance. Here $c_{\max} = 0$ and

$$c_n(CD) = \frac{1}{\pi} \arccos \left(\frac{0 - 1/N^2}{(N-1)/N^2} \right),$$

so that the expected value will be given by

$$\mathbf{E}[CD(A, B)] = \frac{N-1}{N\pi} \arccos \left(-\frac{1}{N-1} \right).$$

This is non-constant (it evaluates to 0.44, 0.47 for $n = 3, 4$ respectively). Note that this expected value converges to $\frac{1}{2}$ for $n \rightarrow \infty$, which is indeed the asymptotic baseline of the index.

FNMI and NMI. Note that in this case $k_A = k_B$ so that the penalty term of FNMI will equal 1 and FNMI will coincide with NMI. Again $c_{\max} = 1$. For the case where the intra-pairs do not coincide, the joint entropy will equal $H(A, B) = \ln(n)$ while each of the marginal entropies will equal

$$H(A) = H(B) = \frac{n-2}{n} \ln(n) + \frac{2}{n} \ln(n/2) = \ln(n) - \frac{2}{n} \ln(2).$$

This results in

$$c_n(NMI) = \frac{2H(A) - H(A, B)}{H(A)} = 1 - \frac{2 \ln(n)}{n \ln(n) - 2 \ln(2)},$$

and the expected value will be given by the non-constant

$$\mathbf{E}[NMI(A, B)] = 1 - \frac{N-1}{N} \frac{2 \ln(n)}{n \ln(n) - 2 \ln(2)}.$$

Note that as $H(A) = H(B)$, all normalizations of MI will be equal so that this counter-example proves that none of the variants of (F)NMI satisfy the constant baseline requirement.

VI. In this case $c_{\max} = 0$. We will use the entropies from the NMI-computations to conclude that

$$\begin{aligned} \mathbf{E}[VI(A, B)] &= \frac{N-1}{N} (2H(A, B) - H(A) - H(B)) \\ &= \frac{N-1}{N} \frac{4}{n} \ln(2), \end{aligned}$$

which is again non-constant.

F-measure. Here $c_{\max} = 1$. In the case where the intra-pairs do not coincide, all contingency variables will be either one or zero so that both recall and precision will equal $1 - 1/n$ so that $c_n(FM) = 1 - 1/n$. This results in the following non-constant expected value

$$\mathbf{E}[FM(A, B)] = 1 - \frac{N-1}{N} \frac{1}{n}.$$

Note that because recall equals precision in both cases, this counter-example also works for other averages than the harmonic average.

BCubed. Again $c_{\max} = 1$. In the other case, the recall and precision will again be equal. Because for BCubed, the contribution of cluster i is given by $\frac{1}{n} \max\{n_{ij}^2\}/|A_i|$, the contributions of the one- and two-clusters will be given by $\frac{1}{n}, \frac{1}{2n}$ respectively. Hence, $c_n(BC) = \frac{n-2}{n} + \frac{1}{2n} = 1 - \frac{3}{2n}$ and we get the non-constant

$$\mathbf{E}[BC(A, B)] = 1 - \frac{N-1}{N} \frac{3}{2n}.$$

We note that again, this counter-example can be extended to non-harmonic averages of the BCubed recall and precision.

C. Additional proofs

C.1. Theorem 2

We will show that $\text{Var}(N_{11}) = o(n^4)$. To compute the variance, we first inspect the second moment. Let $A(S)$ denote the indicator function of the event that all elements of

$S \subset V$ are in the same cluster in A . Define $B(S)$ similarly and let $AB(S) = A(S)B(S)$. Let e, e_1, e_2 range over subsets of V of size 2. We write

$$\begin{aligned} N_{11}^2 &= \left(\sum_e AB(e) \right)^2 \\ &= \sum_{e_1, e_2} AB(e_1)AB(e_2) \\ &= \sum_{|e_1 \cap e_2|=2} AB(e_1)AB(e_2) \\ &\quad + \sum_{|e_1 \cap e_2|=1} AB(e_1)AB(e_2) \\ &\quad + \sum_{|e_1 \cap e_2|=0} AB(e_1)AB(e_2) \\ &= N_{11} + \sum_{|e_1 \cap e_2|=1} AB(e_1 \cup e_2) \\ &\quad + \sum_{e_1 \cap e_2 = \emptyset} AB(e_1)AB(e_2). \end{aligned}$$

We take the expectation

$$\begin{aligned} \mathbf{E}[N_{11}^2] &= \mathbf{E}[N_{11}] + 6 \binom{n}{3} \mathbf{E}[AB(\{v_1, v_2, v_3\})] \\ &\quad + \binom{n}{2} \binom{n-2}{2} \mathbf{E}[AB(e_1)AB(e_2)], \end{aligned}$$

where $v_1, v_2, v_3 \in V$ distinct and $e_1 \cap e_2 = \emptyset$. The first two terms are obviously $o(n^4)$. We inspect the last term

$$\begin{aligned} &\binom{n}{2} \binom{n-2}{2} \mathbf{E}[AB(e_1)AB(e_2)] \\ &= \binom{n}{2} \sum_{i,j} \mathbf{P}(e_1 \subset A_i \cap B_j) \\ &\quad \times \binom{n-2}{2} \mathbf{E}[AB(e_2)|e_1 \subset A_i \cap B_j] \quad (2) \end{aligned}$$

Now we rewrite $\mathbf{E}[N_{11}]^2$ to

$$\mathbf{E}[N_{11}]^2 = \binom{n}{2} \sum_{i,j} \mathbf{P}(e_1 \subset A_i \cap B_j) \binom{n}{2} \mathbf{E}[AB(e_2)].$$

Note that $\binom{n}{2} \mathbf{E}[AB(e_2)] > \binom{n-2}{2} \mathbf{E}[AB(e_2)]$ so that the difference between (2) and $\mathbf{E}[N_{11}]^2$ can be bounded by

$$\begin{aligned} &\binom{n}{2} \binom{n-2}{2} \sum_{i,j} \mathbf{P}(e_1 \subset A_i \cap B_j) \\ &\cdot (\mathbf{E}[AB(e_2)|e_1 \subset A_i \cap B_j] - \mathbf{E}[AB(e_2)]). \end{aligned}$$

As $\binom{n}{2} \binom{n-2}{2} = O(n^4)$, what remains to be proven is

$$\begin{aligned} &\sum_{i,j} \mathbf{P}(e_1 \subset A_i \cap B_j) \\ &\quad \times (\mathbf{E}[AB(e_2)|e_1 \subset A_i \cap B_j] - \mathbf{E}[AB(e_2)]) \\ &= o(1). \end{aligned}$$

Note that it is sufficient to prove that

$$\mathbf{E}[AB(e_2)|e_1 \subset A_i \cap B_j] - \mathbf{E}[AB(e_2)] = o(1),$$

for all i, j . To this end, we will write

$$\begin{aligned} &\mathbf{E}[AB(e_2)|e_1 \subset A_i \cap B_j] \\ &= \mathbf{E}[A(e_2)|e_1 \subset A_i] \mathbf{E}[B(e_2)|e_1 \subset B_j] \\ &= \mathbf{E}[A'(e_2)] \mathbf{E}[B'(e_2)], \end{aligned}$$

where A' is the clustering over $V \setminus e_1$ obtained by removing e_1 from A_i and B' is defined analogously. We have

$$\begin{aligned} \mathbf{E}[A'(e_2)] &= \frac{m_A - (2a_i - 3)}{N - (2n - 3)}, \\ \mathbf{E}[B'(e_2)] &= \frac{m_B - (2b_j - 3)}{N - (2n - 3)}. \end{aligned}$$

This allows us to compute

$$\begin{aligned} &\mathbf{E}[A'B'(e_2)] - \mathbf{E}[AB(e_2)] \\ &= \frac{(m_A - (2a_i - 3))(m_B - (2b_j - 3))}{(N - (2n - 3))^2} - \frac{m_A m_B}{N^2} \\ &= \frac{N^2(m_A - (2a_i - 3))(m_B - (2b_j - 3))}{N^2(N - (2n - 3))^2} \\ &\quad - \frac{(N - (2n - 3))^2 m_A m_B}{N^2(N - (2n - 3))^2} \\ &= \frac{N^2((2a_i - 3)(2b_j - 3) - m_A(2b_j - 3) - m_B(2a_i - 3))}{N^2(N - (2n - 3))^2} \\ &\quad + \frac{m_A m_B (2N(2n - 3) - (2n - 3)^2)}{N^2(N - (2n - 3))^2} \\ &= \frac{((2a_i - 3)(2b_j - 3) - m_A(2b_j - 3) - m_B(2a_i - 3))}{(N - (2n - 3))^2} \\ &\quad + \frac{m_A m_B (2N(2n - 3) - (2n - 3)^2)}{N^2(N - (2n - 3))^2} \\ &= \frac{O(n^3)}{(N - (2n - 3))^2} + \frac{m_A m_B}{N^2} \frac{O(n^3)}{N^2(N - (2n - 3))^2} \\ &= o(1). \end{aligned}$$

Which completes the proof.

⁴Throughout the literature, the Mirkin metric is defined as $2(N_{10} + N_{01})$ but we use this variant as it satisfies scale-invariance as defined in Definition 10.

Table 4. A selection of pair-counting indices. Most of these indices are taken from (Lei et al., 2017).

Index (Abbreviation)	Expression
Rand (R)	$\frac{N_{11}+N_{00}}{N_{11}+N_{10}+N_{01}+N_{00}}$
Adjusted Rand (AR)	$\frac{N_{11} - \frac{(N_{11}+N_{10})(N_{11}+N_{01})}{N_{11}+N_{10}+N_{01}+N_{00}}}{\frac{(N_{11}+N_{10})+(N_{11}+N_{01})}{2} - \frac{(N_{11}+N_{10})(N_{11}+N_{01})}{N_{11}+N_{10}+N_{01}+N_{00}}}$
Jaccard (J)	$\frac{N_{11}}{N_{11}+N_{10}+N_{01}}$
Jaccard Distance (JD)	$\frac{N_{10}+N_{01}}{N_{11}+N_{10}+N_{01}}$
Wallace1 (W)	$\frac{N_{11}}{N_{11}+N_{10}}$
Wallace2	$\frac{N_{11}}{N_{11}+N_{01}}$
Dice	$\frac{2N_{11}}{2N_{11}+N_{10}+N_{01}}$
Correlation Coefficient (CC)	$\frac{N_{11}N_{00}-N_{10}N_{01}}{\sqrt{(N_{11}+N_{10})(N_{11}+N_{01})(N_{00}+N_{10})(N_{00}+N_{01})}}$
Correlation Distance (CD)	$\frac{1}{\pi} \arccos \left(\frac{N_{11}N_{00}-N_{10}N_{01}}{\sqrt{(N_{11}+N_{10})(N_{11}+N_{01})(N_{00}+N_{10})(N_{00}+N_{01})}} \right)$
Sokal&Sneath-I ($S\&S_1$)	$\frac{1}{4} \left(\frac{N_{11}}{N_{11}+N_{10}} + \frac{N_{11}}{N_{11}+N_{01}} + \frac{N_{00}}{N_{00}+N_{10}} + \frac{N_{00}}{N_{00}+N_{01}} \right)$
Minkowski	$\sqrt{\frac{N_{10}+N_{01}}{N_{11}+N_{10}}}$
Hubert (H)	$\frac{N_{11}+N_{00}-N_{10}-N_{01}}{N_{11}+N_{10}+N_{01}+N_{00}}$
Folkes&Mallow	$\frac{N_{11}}{\sqrt{(N_{11}+N_{10})(N_{11}+N_{01})}}$
Sokal&Sneath-II	$\frac{\frac{1}{2}N_{11}}{\frac{1}{2}N_{11}+N_{10}+N_{01}}$
Adjusted Mirkin ⁴	$\frac{N_{10}+N_{01}}{N_{11}+N_{10}+N_{01}+N_{00}}$
Kulczynski	$\frac{1}{2} \left(\frac{N_{11}}{N_{11}+N_{10}} + \frac{N_{11}}{N_{11}+N_{01}} \right)$
McConnaughey	$\frac{N_{11}^2-N_{10}N_{01}}{(N_{11}+N_{10})(N_{11}+N_{01})}$
Yule	$\frac{N_{11}N_{00}-N_{10}N_{01}}{N_{11}N_{10}+N_{01}N_{00}}$
Baulieu-I	$\frac{(N_{11}+N_{10}+N_{01}+N_{00})(N_{11}+N_{00})+(N_{10}-N_{01})^2}{(N_{11}+N_{10}+N_{01}+N_{00})^2}$
Russell&Rao	$\frac{N_{11}}{N_{11}+N_{10}+N_{01}+N_{00}}$
Fager&McGowan	$\frac{N_{11}}{\sqrt{(N_{11}+N_{10})(N_{11}+N_{01})}} - \frac{1}{2\sqrt{N_{11}+N_{10}}}$
Peirce	$\frac{N_{11}N_{00}-N_{10}N_{01}}{(N_{11}+N_{01})(N_{00}+N_{10})}$
Baulieu-II	$\frac{N_{11}N_{00}-N_{10}N_{01}}{(N_{11}+N_{10}+N_{01}+N_{00})^2}$
Sokal&Sneath-III	$\frac{N_{11}N_{00}}{\sqrt{(N_{11}+N_{10})(N_{11}+N_{01})(N_{00}+N_{10})(N_{00}+N_{01})}}$
Gower&Legendre	$\frac{N_{11}+N_{00}}{N_{11}+\frac{1}{2}(N_{10}+N_{01})+N_{00}}$
Rogers&Tanimoto	$\frac{N_{11}+N_{00}}{N_{11}+2(N_{10}+N_{01})+N_{00}}$
Goodman&Kruskal	$\frac{N_{11}N_{00}-N_{10}N_{01}}{N_{11}N_{00}+N_{10}N_{01}}$