FURTHER EQUIDISTRIBUTION OF SET-VALUED STATISTICS ON PERMUTATIONS

JIANXI MAO AND JIANG ZENG

ABSTRACT. We construct bijections to show that two pairs of sextuple set-valued statistics of permutations are equidistributed on symmetric groups. This extends a recent result of Sokal and the second author valid for integer-valued statistics as well as a previous result of Foata and Han for bivariable set-valued statistics.

1. INTRODUCTION

Equidistribution problems of set-valued statistics on permutations have attracted much attention in recent literature, see [BV17, KL18, PO14, FH09]. A decade ago, answering a conjecture of Foata and Han, Cori [CO09] proved that the number of permutations in the symmetric group \mathfrak{S}_n with p cycles and q left-to-right maxima is equal to the number of permutations in \mathfrak{S}_n with q cycles and p left-to-right maxima. In a follow-up [FH09] Foata and Han showed that Cori's result can be further extended to set-valued statistics by using two simple permutation codings called the A-code and the B-code. Recently Sokal and the second author [SZ19] have extended Cori's result on integer-valued bi-statistics to integervalued multiple statistics. The purpose of this paper is to show that the latter has also a setvalued analogue using a classical *Laguerre history* encoding of permutations [DV94, CSZ97] and two new bijections from Laguerre histories onto themselves. For a permutation $\sigma = \sigma(1) \cdots \sigma(n)$ of $12 \dots n$, the pair $(i, \sigma(i))$ $(1 \le i \le n)$ is called

- a record of σ if $\sigma(j) < \sigma(i)$ for all j < i;
- an *antirecord* of σ if $\sigma(j) > \sigma(i)$ for all j > i;
- an *exclusive record* of σ if it is a record but not an antirecord;
- a record-antirecord of σ if it is both a record and an antirecord;
- an excedance if $\sigma(i) > i$.

The corresponding numbers of the above five statistics are denoted by $\operatorname{rec} \sigma$, $\operatorname{arec} \sigma$, $\operatorname{erec} \sigma$, $\operatorname{rar} \sigma$ and $\operatorname{exc} \sigma$. Moreover, the indices *i* and $\sigma(i)$ are called *position* and *letter* of the corresponding statistic. The ten sets of corresponding positions and letters of the above five statistics are denoted, respectively, by $\operatorname{Recp} \sigma$, $\operatorname{Recl} \sigma$, $\operatorname{Arecp} \sigma$, $\operatorname{Arecl} \sigma$, $\operatorname{Erecp} \sigma$, $\operatorname{Erecl} \sigma$, $\operatorname{Rarp} \sigma$, $\operatorname{Rarl} \sigma$, $\operatorname{Excp} \sigma$, and $\operatorname{Excl} \sigma$. For convenience we intoduce the four bi-set-valued statistics

 $\operatorname{Rec} \sigma = (\operatorname{Recp} \sigma, \operatorname{Recl} \sigma), \quad \operatorname{Arec} \sigma = (\operatorname{Arecp} \sigma, \operatorname{Arecl} \sigma)$ $\operatorname{Erec} \sigma = (\operatorname{Erecp} \sigma, \operatorname{Erecl} \sigma), \quad \operatorname{Exc} \sigma = (\operatorname{Excp} \sigma, \operatorname{Excl} \sigma).$

Date: March 1, 2022.

Since the position and letter of a record-antirecord must be equal, we have $\operatorname{Rar} \sigma := \operatorname{Rarp} \sigma = \operatorname{Rarl} \sigma$. Furthermore, if the bijection $i \mapsto \sigma(i)$ $(1 \leq i \leq n)$ has r disjoint cycles, whose maximum elements are c_1, \ldots, c_r , we set $\operatorname{Cyc} \sigma := \{c_1, c_2, \ldots, c_r\}$, and an index $i \in [n] := \{1, \ldots, n\}$ is called a

- cycle peak of σ if $\sigma^{-1}(i) < i > \sigma(i)$;
- cycle valley of σ if $\sigma^{-1}(i) > i < \sigma(i)$;
- cycle double rise of σ if $\sigma^{-1}(i) < i < \sigma(i)$;
- cycle double fall of σ if $\sigma^{-1}(i) > i > \sigma(i)$;
- fixed point of σ if $\sigma(i) = i$.

The corresponding sets (resp. numbers) of the above statistics are denoted by Cpeak σ , Cval σ , Cdrise σ , Cdfall σ and Fix σ (resp. cpeak σ , cval σ , cdrise σ , cdfall σ and fix σ), respectively. The following is our running example in this paper.

Example. Consider the permutation $\sigma \in \mathfrak{S}_{17}$ given by

Then

 $\operatorname{Rec} \sigma = (\{1, 2, 4, 12, 13, 14\}, \{4, 9, 11, 12, 16, 17\})$ $\operatorname{Arec} \sigma = (\{7, 8, 9, 11, 12, 15, 16, 17\}, \{1, 3, 6, 7, 12, 13, 14, 15\})$ $\operatorname{Erec} \sigma = (\{1, 2, 4, 13, 14\}, \{4, 9, 11, 16, 17\})$ $\operatorname{Exc} \sigma = (\{1, 2, 4, 6, 13, 14\}, \{4, 9, 10, 11, 16, 17\})$

and $\operatorname{Rar} \sigma = \{12\}$. The cycle decomposition of σ reads

$$\sigma = (1, 4, 11, 7) (2, 9, 6, 10, 8, 3) (5) (12) (13, 16, 14, 17, 15),$$

thus $\operatorname{Cyc} \sigma = \{5, 10, 11, 12, 17\}$, $\operatorname{Cpeak} \sigma = \{9, 10, 11, 16, 17\}$, $\operatorname{Cval} \sigma = \{1, 2, 6, 13, 14\}$, $\operatorname{Cdrise} \sigma = \{4\}$, $\operatorname{Cdfall} \sigma = \{3, 7, 8, 15\}$, and $\operatorname{Fix} \sigma = \{5, 12\}$.

In a recent work [SZ19] Sokal and the second author studied the polynomials $\mu_n(x, y, u, v)$ in the Taylor expansion of the Stieltjes continued fraction

$$\sum_{n\geq 0} \mu_n(x, y, u, v) t^n = \frac{1}{1 - \frac{xt}{1 - \frac{yt}{1 - \frac{yt}{1 - \frac{(x+u)t}{1 - \frac{(y+v)t}{\dots}}}}},$$
(1.1)

where the coefficients α_k $(k \ge 1)$ are defined by

$$\alpha_{2k-1} = x + (k-1)u, \quad \alpha_{2k} = y + (k-1)v.$$

They [SZ19, Theorem 2.2] showed that the polynomial μ_n has the two interpretations

$$\mu_n(x, y, u, v) = \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{arec}(\sigma)} y^{\operatorname{erec}(\sigma)} u^{n - \operatorname{exc}(\sigma) - \operatorname{arec}(\sigma)} v^{\operatorname{exc}(\sigma) - \operatorname{erec}(\sigma)}$$
(1.2)

$$= \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{cyc}(\sigma)} y^{\operatorname{erec}(\sigma)} u^{n - \operatorname{exc}(\sigma) - \operatorname{cyc}(\sigma)} v^{\operatorname{exc}(\sigma) - \operatorname{erec}(\sigma)}, \qquad (1.3)$$

where $\operatorname{cyc} \sigma$ is the cycle number of σ . Besides, they [SZ19, Theorem 2.4] proved the Jacobi continued fraction expansion

$$\frac{\sum_{n=0}^{\infty} \left(\sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{cyc}(\sigma)} y^{\operatorname{arec}(\sigma)} z^{\operatorname{exc}(\sigma)} w_0^{\operatorname{rar}\sigma}\right) t^n}{1} =$$
(1.4)

$$1 - xyw_0 t - \frac{xyz t}{1 - (x + y + z) t - \frac{(x + 1)(y + 1)z t}{1 - \dots}}$$

where the coefficients γ_k $(k \ge 0)$ and β_k $(k \ge 1)$ are defined by $\gamma_0 = xyw_0$,

$$\gamma_n = x + y + n - 1 + nz, \quad \beta_n = (x + n - 1)(y + n - 1) \text{ for } n \ge 1.$$

It follows from (1.2) and (1.3) that the two triple integer-valued statistics

$$(\text{erec}, \text{cyc}, \text{exc}) \quad \text{and} \quad (\text{erec}, \text{arec}, \text{exc})$$
 (1.5)

are equidistributed on \mathfrak{S}_n . As the right-hand side of (1.4) is symmetric under $x \leftrightarrow y$, the two triple integer-valued statistics

$$(cyc, arec, exc)$$
 and $(arec, cyc, exc)$ (1.6)

are also equidistributed on \mathfrak{S}_n . Note that the symmetry of the integer valued statistics (arec, cyc) is due to Cori [CO09]. Motivated by the set-valued analogue of Cori's result [FH09], we shall prove the following set-valued analogue of (1.5) and (1.6).

Theorem 1. The two sextuple set-valued statistics

$$(Cyc, Erec, Exc, Rar)$$
 and $(Arecp, Erec, Exc, Rar)$

are equidistibuted on \mathfrak{S}_n .

Theorem 2. The two quintuple set-valued statistics

are equidistributed on \mathfrak{S}_n .

As $\text{Rec} = (\text{Erecp} \cup \text{Rar}, \text{Erecl} \cup \text{Rar})$, we derive immediately that the following two pairs of triple set-statistics

(Cyc, Rec, Exc) and (Arecp, Rec, Exc), (1.7)

(Cyc, Arecp, Exc) and (Arecp, Cyc, Exc) (1.8)

are equidistributed on \mathfrak{S}_n . We note that (1.7) is a stronger version of Han's conjecture [Han19], i.e., (cyc, Recp, Excp) and (arec, Recp, Excp) are equidistributed on \mathfrak{S}_n and the bijections in [CO09, FH09] do not keep track of excedances.

A 3-Motzkin word of length n is a word $s := s_1 \dots s_i$ on the alphabet $\{U, D, L_a, L_b, L_c\}$ such that $|s_1 \dots s_n|_U = |s_1 \dots s_n|_D$ and

$$h_i := |s_1 \dots s_i|_{\mathcal{U}} - |s_1 \dots s_i|_{\mathcal{D}} \ge 0 \quad (i = 1, \dots, n),$$
(1.9)

where $|s_1 \dots s_i|_{\mathbf{a}}$ is the number occurences of letter **a** in the word $s_1 \dots s_i$. A Motzkin path of length n is a sequence of points $\omega = (\omega_0, \omega_1, \dots, \omega_{n-1}, \omega_n)$ in $\mathbb{N} \times \mathbb{N}$, starting from $\omega_0 = (0,0)$ and ending at $\omega_n = (n,0)$, such that each step $s_i := (\omega_{i-1}, \omega_i)$ is of type up if $\omega_i - \omega_{i-1} = (1, 1)$, or type down if $\omega_i - \omega_{i-1} = (1, -1)$, or type level if $\omega_i - \omega_{i-1} = (1, 0)$ for $i = 1, \ldots, n$. Clearly we can depict a Motzkin word with a Motzkin path with three types of level steps. A Laguerre history of length n is a pair (s, γ) , where s is a 3-Motzkin word $s := s_1 \dots s_n$ with $h_n = 0$ (i.e., $|s|_U = |s|_D$) and $\gamma = (\gamma_1, \dots, \gamma_n)$ is a sequence satisfying the following:

- $\gamma_i = (\xi_i, \eta_i) = (\Delta, \Delta)$ if $s_i = U$,
- $\gamma_i = (\xi_i, \eta_i) \in \{1, \dots, h_{i-1}\}^2$ if $s_i = D$,
- $\gamma_i = (\xi_i, \eta_i) \in \{1, \dots, h_{i-1}\} \times \{\Delta\}$ if $s_i = L_a$, $\gamma_i = (\xi_i, \eta_i) \in \{\Delta\} \times \{1, \dots, h_{i-1}\}$ if $s_i = L_b$,
- $\gamma_i = (\xi_i, \eta_i) = (\Delta, h_{i-1} + 1)$ if $s_i = L_c$.

Let \mathcal{LH}_n be the set of Laguerre histories of length n. There are several well-known related such bijections between \mathfrak{S}_n and \mathcal{LH}_n , see [Bi93, DV94, CSZ97] and references therein. We shall present a variant of such encoding θ , which is very close to Biane's version [Bi93]. Our strategy is to first encode permutations using Laguerre histories and then build bijections ρ_i on the latter, where ρ_i is the bijection on \mathcal{LH}_n used in the proof of Theorem i (i = 1, 2). In otherwords, we have the following diagram

$$\mathfrak{S}_n \xrightarrow{\theta} \mathcal{LH}_n \xrightarrow{\rho_i} \mathcal{LH}_n \xrightarrow{\theta^{-1}} \mathfrak{S}_n.$$

The paper is organized as follows: in Section 2 we present permutation code, i.e., a bijection θ between \mathfrak{S}_n and \mathcal{LH}_n with the main properties. In Section 3 we construct the bijection ρ_1 from \mathcal{LH}_n onto itself and prove Theorem 1 by composing θ and ρ_1 . In Section 4 we construct the bijection ρ_2 from \mathcal{LH}_n onto itself and prove Theorem 2 by composing θ and ρ_2 .

2. Encoding permutations by Laguerre histories

For $\sigma \in \mathfrak{S}_n$ and $i \in [n]$, the refined *lower-nesting* and *upper-nesting* numbers (see [SZ19]) are defined by

lnest
$$(i, \sigma) = \#\{j \in [n] : \sigma(j) < \sigma(i) \text{ and } i < j\},\$$

unest $(i, \sigma) = \#\{j \in [n] : \sigma^{-1}(j) < \sigma^{-1}(i) \text{ and } i < j\}$

Note that $\operatorname{unest}(i, \sigma) = \operatorname{lnest}(i, \sigma^{-1})$.



FIGURE 1. The Laguerre history of $\sigma = (1, 4, 11, 7)(2, 9, 6, 10, 8, 3)(5)(12)(13, 16, 14, 17, 15)$

Lemma 3. Given $\sigma \in \mathfrak{S}_n$ and $i \in [n]$, we have

(1) $i \in \operatorname{Arecp} \sigma$ if and only if $\operatorname{lnest}(i, \sigma) = 0$, (2) $i \in \operatorname{Recl} \sigma$ if and only if $\operatorname{unest}(i, \sigma) = 0$.

Proof. Statement (1) is clear by definition. For $(i, \sigma) \in [n] \times \mathfrak{S}_n$, as unest $(i, \sigma) = \text{lnest}(i, \sigma^{-1})$ we have

$$(i, \sigma(i)) \in \operatorname{Arec} \sigma \Leftrightarrow (\sigma(i), i) \in \operatorname{Rec} \sigma^{-1}.$$

Hence (2) is equivalent to (1).

Algorithm θ . For $\sigma \in \mathfrak{S}_n$, let $\theta(\sigma) = (s, \gamma) \in \mathcal{LH}_n$ where the pair (s_i, γ_i) for $1 \le i \le n$ is defined as follows:

$$(s_i, \gamma_i) = \begin{cases} (\mathbf{U}, (\Delta, \Delta)) & \text{if } i \in \operatorname{Cval} \sigma; \\ (\mathbf{D}, (\operatorname{unest}(i, \sigma) + 1, \operatorname{lnest}(i, \sigma) + 1)) & \text{if } i \in \operatorname{Cpeak} \sigma; \\ (\mathbf{L}_a, (\operatorname{unest}(i, \sigma) + 1, \Delta)) & \text{if } i \in \operatorname{Cdrise} \sigma; \\ (\mathbf{L}_b, (\Delta, \operatorname{lnest}(i, \sigma) + 1)) & \text{if } i \in \operatorname{Cdfall} \sigma; \\ (\mathbf{L}_c, (\Delta, h_{i-1} + 1)) & \text{if } i \in \operatorname{Fix} \sigma, \end{cases}$$
(2.1)

We can show that θ is a variant of some well-known bijections from permutations to Laguerre hostories; see [Bi93, DV94, CSZ97, SZ19] for more details and other variants.

Lemma 4. Let $\theta(\sigma) = (s, \gamma)$ and $h_0 = 0$. For $i = 1, \ldots, n$,

- if $i \in \operatorname{Cval} \sigma$, then $h_i = h_{i-1} + 1$.
- if $i \in \operatorname{Cpeak} \sigma$, then $h_i = h_{i-1} 1$.
- if $i \in \operatorname{Cdrise} \sigma \cup \operatorname{Cdfall} \sigma \cup \operatorname{Fix} \sigma$, then $h_i = h_{i-1}$,

and $h_i = \#\{j \le i : \sigma(j) > i\} = \#\{j \le i : \sigma^{-1}(j) > i\}.$

A permutation $\sigma \in \mathfrak{S}_n$ can be represented by a *bipartite digraph* such that

- the top row of vertices is labelled by $1, \ldots, n$,
- the bottom row of vertices is labelled by $1', \ldots, n'$,
- there is an edge $i \to j'$ from the top row to the bottom row if and only if $\sigma(i) = j$.



FIGURE 2. The bipartite digraph of $\sigma = (1, 4, 11, 7)(2, 9, 6, 10, 8, 3)(5)(12)(13, 16, 14, 17, 15)$

The bipartite digraph associated to the running example is depicted in Figure 2. We can visualize the statistics $\text{lnest}(i, \sigma)$ and $\text{unest}(i, \sigma)$ in (2.1) by the *i*-th restriction of the bipartite digraph on $\frac{1}{1'} \cdots \frac{i}{i'}$ for $i = 1, \ldots, n$. In other words, we have the following graphical description of the mapping θ .

Algorithm θ (bis). Let $\sigma \in \mathfrak{S}_n$ and $g_0 = \emptyset$. For $i = 1, \ldots, n$, the *i*-th restriction g_i is obtained from g_{i-1} by adding the column $_{i'}^i$ at each time *i* from left to right as follows:

- (i) if $i \in \operatorname{Cval} \sigma$, then $(s_i, \gamma_i) = (U, (\Delta, \Delta))$,
- (ii) if $i \in \text{Cpeak } \sigma$, then $(s_i, \gamma_i) = (L_a, (\xi_i, \eta_i))$, where $\xi_i 1$ is the number of vacant vertices to the left of $\sigma^{-1}(i)$ at the top row of g_i and $\eta_i 1$ is the number of vacant vertices to the left of $\sigma(i)$ at the bottom row of g_i .
- (iii) if $i \in \text{Cdrise } \sigma$, then $(s_i, \gamma_i) = (L_a, (\xi_i, \Delta))$, where $\xi_i 1$ is the number of vacant vertices to the left of $\sigma^{-1}(i)$ at the top row of g_i ,
- (iv) if $i \in \text{Cdfall } \sigma$, then $(s_i, \gamma_i) = (L_b, (\Delta, \eta_i))$, where $\eta_i 1$ is the number of vacant vertices to the left of $\sigma(i)$ at the bottom row of g_i ,
- (v) if $i \in \text{Fix } \sigma$, then $(s_i, \gamma_i) = (L_c, (\Delta, h_{i-1} + 1))$, where h_{i-1} is the number of vacant vertices at the top (or bottom) row of g_{i-1} .

Lemma 5. If g is the bipartite digraph of $\sigma \in \mathfrak{S}_n$ and g_i is the *i*-th restriction of g $(1 \leq i \leq n)$, then the index *i* is a cycle maximum of σ if and only if either $i \rightarrow i'$ is an edge or there are integers i_1, i_2, \ldots, i_k in [*i*] such that $i \rightarrow i'_1, i_1 \rightarrow i'_2, \ldots, i_k \rightarrow i'$ are edges of g_i .

Proof. Given a permutation $\sigma \in \mathfrak{S}_n$, an index i is a cycle maximum if and only if $\sigma^{\ell}(i) \leq i$ and for all $\ell \geq 0$, this means either $i \to i'$ is an edge or there are vertices i_1, i_2, \ldots, i_k between 1 and i at the top row and i'_1, \ldots, i'_k at the bottom row such that $i \to i'_1, i_1 \to i'_2, \ldots, i_k \to i'$ are edges in the bipartite graph g_i .

Remark. Starting from any permutation we can draw the corresponding bipartite digraph or Laguerre history through the above correspondences. By the above lemma, the graphical interpretation enables us to to count the cycle maxima of a permutation in our bijections between Laguerre histories, which seems difficult in the corresponding Laguerre history via the bijection θ .

For each Laguerre history $(s, \gamma) \in \mathcal{LH}_n$ we define the set-valued statistics:

$$\operatorname{Arecp}(s,\gamma) = \{i : s_i \in \{D, L_b, L_c\} \text{ and } \eta_i = 1\}$$

$$(2.2)$$

$$\operatorname{Erecl}(s,\gamma) = \{i : s_i = D \text{ or } L_a, \text{ and } \xi_i = 1\}$$

$$(2.3)$$

$$\operatorname{Erecp}(s,\gamma) = \{i : i \to j' \text{ in g and } j \in \operatorname{Erecl}(s,\gamma)\}$$

$$(2.4)$$

$$\operatorname{Erec}(s,\gamma) = (\operatorname{Erecp}(s,\gamma),\operatorname{Erecl}(s,\gamma))$$
(2.5)

$$\operatorname{Excp}(s,\gamma) = \{i : s_i = \mathrm{U} \text{ or } \mathrm{L}_a\}, \quad \operatorname{Excl}(s,\gamma) = \{i : s_i = \mathrm{D} \text{ or } \mathrm{L}_a\}$$
(2.6)

$$\operatorname{Exc}(s,\gamma) = (\operatorname{Excp}(s,\gamma), \operatorname{Excl}(s,\gamma))$$
(2.7)

$$\operatorname{Rar}(s,\gamma) = \{i : s_i = \mathcal{L}_c \text{ and } \eta_i = 1\}$$

$$(2.8)$$

$$\operatorname{Cyc}(s,\gamma) = \{i : s_i = \operatorname{L}_c \text{ or } s_i = \operatorname{D} \text{ with } i \to i'_1, i_1 \to i'_2, \dots, i_k \to i' \text{ in } g_i\}.$$
(2.9)

Lemma 6. Let $\sigma \in \mathfrak{S}_n$ and $\theta(\sigma) = (s, \gamma)$. Then,

- (i) $\operatorname{Arecp}(s, \gamma) = \operatorname{Arecp} \sigma$.
- (ii) $\operatorname{Erecl}(s, \gamma) = \operatorname{Erecl} \sigma$.
- (iii) $\operatorname{Rar}(s, \gamma) = \operatorname{Rar} \sigma$.
- (iv) $\operatorname{Excp}(s, \gamma) = \operatorname{Excp} \sigma$.
- (v) $\operatorname{Excl}(s, \gamma) = \operatorname{Excl} \sigma$.
- (vi) $\operatorname{Cyc}(s, \gamma) = \operatorname{Cyc} \sigma$.

Proof. Clearly, if $i \in \operatorname{Arecp} \sigma$, then $\sigma(i) \leq i$, i.e., $i \in \operatorname{Cdfall} \sigma \cup \operatorname{Cpeak} \sigma \cup \operatorname{Fix} \sigma$, and if $i \in \operatorname{Erecl} \sigma$, then $i > \sigma^{-1}(i)$, i.e., $i \in \operatorname{Cdrise} \sigma \cup \operatorname{Cpeak} \sigma$. By Lemma 3 and (2.1) we get (i) and (ii). Next, an integer $i \in \operatorname{Rar} \sigma \Leftrightarrow i \in \operatorname{Fix} \sigma \cap \operatorname{Arecp} \sigma$, so by (i) and (2.1) we get (iii). Since $\operatorname{Excp} = \operatorname{Cval} \cup \operatorname{Cdrise}$ and $\operatorname{Excl} = \operatorname{Cpeak} \cup \operatorname{Cdrise}$, by (2.1), we get (iv) and (v). By Lemma 5 we get (vi).

3. PROOF OF THEOREM 1

3.1. Algorithm ρ_1 . For $(s, \gamma) \in \mathcal{LH}_n$, we define $\rho_1(s, \gamma) = (s', \gamma')$ through the corresponding bipartite digraphs g'_i for $1 \le i \le n$. Set $g'_0 = \emptyset$. The graph g'_i is obtained from g'_{i-1} by first adding the column $\frac{i}{i'}$ with possibles edges as follows:

(i)-(ii) If $s_i = U$ or L_a , then $(s'_i, \gamma') = (s_i, \gamma_i)$. (iii) If $s_i = L_b$ and $\gamma_i = (\Delta, \eta_i)$, then

$$(s'_{i}, (\xi'_{i}, \eta'_{i})) = \begin{cases} (\mathcal{L}_{c}, (\Delta, h_{i-1} + 1)) & \text{if } \eta_{i} = 1\\ (\mathcal{L}_{b}, (\Delta, \eta_{i})) & \text{if } \eta_{i} > 1. \end{cases}$$
(3.1)

Note that $\eta'_i > 1$ because $h_{i-1} \ge 1$. (iv) If $s_i = L_c$ and $\gamma_i = (\Delta, h_{i-1} + 1)$, then

$$(s'_{i}, (\xi'_{i}, \eta'_{i})) = \begin{cases} (\mathbf{L}_{c}, (\Delta, 1)) & \text{if } \eta_{i} = 1\\ (\mathbf{L}_{b}, (\Delta, 1)) & \text{if } \eta_{i} > 1. \end{cases}$$
(3.2)

(v) If $s_i = D$ and $\gamma_i = (\xi_i, \eta_i)$, let $s'_i = s_i$ and $\xi'_i = \xi_i$. Thus the vertex i' is connected to the ξ_i -th vacant vertex v_1 at the top row of g'_i and there are edges $i' \to v_1, v'_1 \to v_1$.

 $v_2, \ldots, v'_{r-1} \to v_r$ such that v'_r is vacant. Assume that v'_r is the η^*_i -th vacant vertex at the bottom row from left to right. Then

$$\eta_{i}' = \begin{cases} \eta_{i}^{*} & \text{if } \eta_{i} = 1\\ \eta_{i} & \text{if } \eta_{i} > \eta_{i}^{*}\\ \eta_{i} - 1 & \text{if } 1 < \eta_{i} \le \eta_{i}^{*}. \end{cases}$$
(3.3)

Set $\rho_1(s,\gamma) = (s',\gamma')$, where $s' = (s'_1, \ldots, s'_n)$ and $\gamma' = (\gamma'_1, \ldots, \gamma'_n)$.

Lemma 7. The mapping $\rho_1 : \mathcal{LH}_n \to \mathcal{LH}_n$ is a bijection.

Proof. We construct the inverse mapping ρ_1^{-1} . Let $(s', \gamma') = \rho_1(s, \gamma)$ and $g''_0 = \emptyset$. For $1 \leq i \leq n$, if $s'_i \in \{U, L_a, L_b, L_c\}$, we define (s''_i, γ''_i) in the same way as ρ_1 ; if $s'_i = D$ and $\gamma'_i = (\xi'_i, \eta'_i)$, let $s''_i = D$ and $\xi''_i = \xi'_i$. Let v_r be the unique vertex such that $i' \leftarrow v_1, v'_1 \leftarrow v_2, \ldots, v'_{r-1} \leftarrow v_r$ and $v'_r \leftarrow i$ or v'_r is vacant in g'_i . Assume that there are $\tilde{\eta}^*_i - 1$ vacant vertices at the left of v'_r at the bottom row of g'_i . Then

$$\eta_i'' = \begin{cases} 1 & \text{if } \eta_i' = \tilde{\eta}_i^* \\ \eta_i' & \text{if } \eta_i' > \tilde{\eta}_i^* \\ \eta_i' + 1 & \text{if } 1 \le \eta_i' < \tilde{\eta}_i^*. \end{cases}$$
(3.4)

Set $\rho_1^{-1}(s',\gamma') = (s'',\gamma'')$, where $s'' = (s''_1,\ldots,s''_n)$ and $\gamma'' = (\gamma''_1,\ldots,\gamma''_n)$. We show that $(s''_i,\gamma''_i) = (s_i,\gamma_i)$ by induction on *i*. If i = 1, then $(s_1,\gamma_1) = (U, (\Delta, \Delta))$ or $(L_c, (1,1))$. By definition of ρ_1 (see (i) and (iv) (a)) and ρ_1^{-1} , for both two cases, we get $(\bar{s}_1,\bar{\gamma}_1) = (s_1,\gamma_1)$. For $i \geq 2$ assume $(s''_k,\gamma''_k) = (s_k,\gamma_k)$ for $1 \leq k \leq i-1$. We have to verify the validity for k = i for the five types of s_i in § 3.1. This is trivial for the cases (i)-(iv). For case (v), we have $s_i = s'_i = s''_i = D$ and $\xi''_i = \xi'_i = \xi_i$. By (v), the mapping ρ_1 provides the index η^*_i , which is equal to $\tilde{\eta}^*_i$. There are three cases :

- $\eta_i = 1$. Then $\eta'_i = \eta^*_i$ by (3.3), this means $\eta''_i = 1$ by (3.4).
- $\eta_i > \eta_i^*$. Then $\eta_i' = \eta_i$, and $\bar{\eta}_i = \eta_i' = \eta_i$ by (3.3)and (3.4).
- $1 < \eta_i \le \eta_i^*$. Then $\eta_i' = \eta_i 1$ by (3.3) and $\bar{\eta}_i = \eta_i' + 1 = \eta_i$ by (3.4).

Summarizing the above five cases, we have completed the proof.

Remark. Using the graph g'_i we determine η^*_i by (3.3) and characterize $i \in \text{Cyc}(s', \gamma')$ by the equation $\eta'_i = \eta^*_i$. Next, using g'_i we get the index $\tilde{\eta}^*_i$ by (3.4) and identify $i \in \text{Cyc}(s', \gamma')$ by the equation $\eta'_i = \tilde{\eta}^*_i$. Thus $\eta^*_i = \tilde{\eta}^*_i$.

Lemma 8. For $(s, \gamma) \in \mathcal{LH}_n$, if $\rho_1(s, \gamma) = (s', \gamma')$, then

 $(Cyc, Exc, Erecl, Rar)(s', \gamma') = (Arecp, Exc, Erecl, Rar)(s, \gamma).$

Proof. By definition of ρ_1 in § 3.1, if $\rho_1(s_i, \gamma_i) = (s'_i, \gamma'_i)$ for $1 \le i \le n$, then $s_i = s'_i$ for $s_i \in \{U, D, L_a\}$ and $\{i : s_i = L_b \text{ or } L_c\} = \{i : s'_i = L_b \text{ or } L_c\}$.

- (1) As $\operatorname{Excp}(s,\gamma) = \{i : s_i = U \text{ or } L_a\}$ and $\operatorname{Excl}(s,\gamma) = \{i : s_i = D \text{ or } L_a\}$, see (2.2)-(2.9), we have $\operatorname{Exc}(s',\gamma') = \operatorname{Exc}(s,\gamma)$.
- (2) Recall that $\operatorname{Arecp}(s, \gamma) := \{i : s_i \in \{D, L_b, L_c\} \text{ and } \eta_i = 1\}$ and $\operatorname{Cyc}(s', \gamma') = \{i : s'_i = L_c \text{ or } s'_i = D \text{ with } i \to i'_1, i_1 \to i'_2, \dots, i_k \to i' \text{ in } g'_i\}$ (cf. (2.2) and (2.9)).



FIGURE 3. The graph of $\varphi(\sigma) = (1, 4, 11, 6, 10, 5)(7)(8)(2, 9, 3)(12)(13, 16)(14, 17)(15)$

- $(s_i, \gamma_i) = (L_b, (\Delta, 1))$ or $(L_c, (\Delta, 1))$ if and only if $(s'_i, \gamma'_i) = (L_c, (\Delta, h_{i-1} + 1))$, in other word there is an edge $i \to i'$ in g'_i by (iii) and (iv).
- $(s_i, \gamma_i) = (D, (\xi_i, 1))$ if and only if $s'_i = D$, $\xi'_i = \xi_i$, and in g'_i there are edges $i \to i'_1, i_1 \to i'_2, \ldots, i_k \to i'$ by § 3.1 (v).

Thus $\operatorname{Cyc}(s', \gamma') = \operatorname{Arecp}(s, \gamma).$

- (3) Recall that $\operatorname{Erecl}(s,\gamma) := \{i : (s_i,\xi_i) = (D,1) \text{ or } (L_a,1)\}$. By § 3.1 (ii)(v) the latter is equivalent to $(s'_i,\xi'_i) = (D,1) \text{ or } (L_a,1)\}$. So $\operatorname{Erecl}(s',\gamma') = \operatorname{Erecl}(s,\gamma)$.
- (4) By § 3.1 (iv), $(s_i, \gamma_i) = (L_c, (\Delta, 1))$ if and only if $(s'_i, \gamma'_i) = (L_c, (\Delta, 1))$. Thus $\operatorname{Rar}(s', \gamma') = \operatorname{Rar}(s, \gamma)$.

Lemma 9. For
$$(s, \gamma) \in \mathcal{LH}_n$$
, if $\rho_1(s, \gamma) = (s', \gamma')$, then $\operatorname{Erecp}(s', \gamma') = \operatorname{Erecp}(s, \gamma)$.

Proof. Let g and g' denote the bipartite digraphs corresponding to (s, γ) and (s', γ') . The construction of ρ_1 shows that $\xi'_i = \xi_i$ for $1 \le i \le n$. Let (cf. Lemma 8)

$$P := \operatorname{Excp}(s', \gamma') = \operatorname{Excp}(s, \gamma) = \{u_1, \dots, u_t\}_{<}$$
(3.5)

$$L := \operatorname{Excl}(s', \gamma') = \operatorname{Excl}(s, \gamma) = \{v_1, \dots, v_t\}_{<}.$$
(3.6)

Since $\operatorname{Erecl}(s,\gamma) \subset L$ (see (2.3) and (2.6)) and $\operatorname{Erecl}(s',\gamma') = \operatorname{Erecl}(s,\gamma)$ (see Lemma 8), it suffices to prove that $u_j \to v'_i$ is an edge in g with $(u_j, v_i) \in P \times L$ if and only if $u_j \to v'_i$ is an edge in g'. Let E_i (resp. E'_i) be the set of vacant vertices at the top row of g_{v_i-1} (resp. g'_{v_i-1}). We show that $E_i = E'_i$ for all $v_i \in L$ for $i = 1, \ldots, t$.

- For i = 1, a vertex v at the top row of g_{v_1-1} (resp. g'_{v_1-1}) is vacant if and only if $v \in P$ and $v < v_1$. Thus $E_1 = E'_1$ by (3.5).
- Assume $E_{i-1} = E'_{i-1}$. If $u_j \to v'_{i-1}$ is an edge in $g_{v_{i-1}}$, then it is also in g' because $\xi_{v_{i-1}} = \xi'_{v_{i-1}}$. Thus $E_i = (E_{i-1} \setminus \{u_j\}) \cup \{v \in P : v_{i-1} \le v \le v_i 1\}$. So $E_i = E'_i$.

Since $\xi_{v_i} = \xi'_{v_i}$ for $v_i \in L$, each v'_i is connected to the same vertex in both g and g'. \Box

By Lemma 8 and 9, for $(s, \gamma) \in \mathcal{LH}_n$, if $\rho_1(s, \gamma) = (s', \gamma')$, then

 $(Cyc, Exc, Erec, Rar)(s', \gamma') = (Arecp, Exc, Erec, Rar)(s, \gamma).$

Let $\varphi := \theta^{-1} \circ \rho_1 \circ \theta$. By Lemmas 6, 8 and 9, we see that φ is the desired bijection from \mathfrak{S}_n onto itself for Theorem 1.



FIGURE 4. The Laguerre history of $\varphi(\sigma)$

Example. For our running example σ , we get $\omega := \varphi(\sigma)$ with

$\omega =$	(1)	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
	$\begin{pmatrix} 4 \end{pmatrix}$	9	2	11	1	10	7	8	3	5	6	12	16	17	15	13	14)

We have

 $\operatorname{Rec} \omega = (\{1, 2, 4, 12, 13, 14\}, \{4, 9, 11, 12, 16, 17\})$ $\operatorname{Arec} \omega = (\{5, 9, 10, 11, 12, 16, 17\}, \{1, 3, 5, 6, 12, 13, 14\})$ $\operatorname{Erec} \omega = (\{1, 2, 4, 13, 14\}, \{4, 9, 11, 16, 17\})$ $\operatorname{Exc} \omega = (\{1, 2, 4, 6, 13, 14\}, \{4, 9, 11, 10, 16, 17\})$

and Rar $\omega = \{12\}$. Also, the cycle decomposition of ω is

$$\omega = (1, 4, 11, 6, 10, 5) (7) (8) (2, 9, 3) (12) (13, 16) (14, 17) (15),$$

so, Cyc $\omega = \{7, 8, 9, 11, 12, 15, 16, 17\}$, Cpeak $\omega = \{9, 10, 11, 16, 17\}$, Cval $\omega = \{1, 2, 6, 13, 14\}$, Cdrise $\omega = \{4\}$, Cdfall $\omega = \{3, 5\}$, and Fix $\omega = \{7, 8, 12, 15\}$.

4. Proof of Theorem 2

4.1. Algorithm ρ_2 . For $(s, \gamma) \in \mathcal{LH}_n$, we define $\rho_2(s, \gamma) = (s', \gamma')$ through the corresponding bipartite digraphs (g_i, g'_i) for $1 \le i \le n$ as follows.

(i)-(ii) If $s_i = U$ or L_a , then $(s'_i, \gamma') = (s_i, \gamma_i)$.

(iii) If $s_i = L_b$ and $\gamma_i = (\Delta, \eta_i)$, then

$$(s'_{i}, (\xi'_{i}, \eta'_{i})) = \begin{cases} (\mathcal{L}_{c}, (\Delta, h_{i-1} + 1)) & \text{if } \eta_{i} = 1\\ (\mathcal{L}_{b}, (\Delta, \eta_{i})) & \text{if } \eta_{i} > 1. \end{cases}$$
(4.1)

Note that $\eta'_i > 1$ because $h_{i-1} \ge 1$.

(iv) If $s_i = L_c$ and $\gamma_i = (\Delta, h_{i-1} + 1)$, then

$$(s'_{i}, (\xi'_{i}, \eta'_{i})) = \begin{cases} (\mathbf{L}_{c}, (\Delta, 1)) & \text{if } \eta_{i} = 1\\ (\mathbf{L}_{b}, (\Delta, 1)) & \text{if } \eta_{i} > 1. \end{cases}$$
(4.2)

- (v) If $s_i = D$ and $\gamma_i = (\xi_i, \eta_i)$, then $s'_i = D$. We use the corresponding bipartite diagraphs g_i and g'_i to define η'_i and ξ'_i .
 - Definition of η'_i . There are edges $i \to v'_1, v_1 \to v'_2, \ldots, v_{s-1} \to v'_s$ in g_i such that v_s is vacant or $v_s \to i'$. Assume that there are $\xi^*_i 1$ vacant vertices at the left of v_s at the top row of g_i . Let

$$\eta_i' = \begin{cases} 1 & \text{if } \xi_i = \xi_i^* \\ \xi_i & \text{if } \xi_i > \xi_i^* \\ \xi_i + 1 & \text{if } 1 \le \xi_i < \xi_i^*. \end{cases}$$
(4.3)

- Definition of ξ'_i . Connecting vertex *i* to the η'_i -th vacant vertex p'_1 at the bottom row of g'_{i-1} yields edges $i \to p'_1, p_1 \to p'_2, \ldots, p_{t-1} \to p'_t$ such that p_t is vacant in g'_{i-1} . Assume that p_t is the η^*_i -th vacant vertex at the top row of g'_{i-1} . Let

$$\xi_{i}' = \begin{cases} \eta_{i}^{*} & \text{if } \eta_{i} = 1\\ \eta_{i} & \text{if } \eta_{i} > \eta_{i}^{*}\\ \eta_{i} - 1 & \text{if } 1 < \eta_{i} \le \eta_{i}^{*}. \end{cases}$$
(4.4)

Set $\rho_2(s,\gamma) = (s',\gamma')$, where $s' = (s'_1,\ldots,s'_n)$ and $\gamma' = (\gamma'_1,\ldots,\gamma'_n)$.

Lemma 10. The mapping $\rho_2 : \mathcal{LH}_n \to \mathcal{LH}_n$ is an involution.

Proof. For $(s,\gamma) \in \mathcal{LH}_n$ let $\rho_2(s,\gamma) = (s',\gamma')$ and $\rho_2(s',\gamma') = (s'',\gamma'')$. We show that $(s''_i,\gamma''_i) = (s_i,\gamma_i)$ by induction on i with $1 \leq i \leq n$. For i = 1 we have $(s_1,\gamma_1) = (\mathbf{U}, (\Delta, \Delta))$ or $(\mathbf{L}_c, (\Delta, 1))$. It is clear from (1)-(iv) we have $(s''_1,\gamma''_1) = (s_1,\gamma_1)$. Let $i \geq 2$ and assume that $((s''_1,\gamma''_1),\ldots,(s''_{i-1},\gamma''_{i-1}) = ((s_1,\gamma_1),\ldots,(s_{i-1},\gamma_{i-1}))$. For the cases (i)-(iv) it is easy to see that $(s''_i,\gamma''_i) = (s_i,\gamma_i)$. Here we just verify the case (v) with $s_i = D$ and $\gamma_i = (\mathbf{D}, (\xi_i,\eta_i))$. Then $s''_i = s'_i = D$, and the mapping ρ_2 (resp. $\rho_2 \circ \rho_2$) provides the indices ξ^*_i and η^*_i (resp. $\tilde{\xi}^*_i$ and $\tilde{\eta}^*_i$) for the computation of η'_i and ξ'_i (resp. η''_i and ξ''_i) in (4.3) and (4.6). We show that $\eta^*_i = \tilde{\xi}^*_i, \xi^*_i = \tilde{\eta}^*_i, \text{ and } \gamma''_i = \gamma_i \ (i \geq 2)$ in the following way. Using g_i and g'_i we determine ξ^*_i and η^*_i by (4.3) and (4.4), and then characterize $i \in I$.

Using g_i and g'_i we determine ξ^*_i and η^*_i by (4.3) and (4.4), and then characterize $i \in \operatorname{Cyc}(s,\gamma)$ and $i \in \operatorname{Cyc}(s',\gamma')$ by the equation $\xi_i = \xi^*_i$ and $\xi'_i = \eta^*_i$. This yields $\gamma'_i = (\xi'_i,\eta'_i)$ by (4.3) and (4.4). On the other hand, using g'_i we get the index $\tilde{\xi}^*_i$ by (4.3) and identify $i \in \operatorname{Cyc}(s',\gamma')$ by the equation $\xi'_i = \tilde{\xi}^*_i$. Thus $\eta^*_i = \tilde{\xi}^*_i$.

Next we define η_i'' by (4.3) using the index $\tilde{\xi}_i^* = \eta_i^*$.

$$\eta_i'' = \begin{cases} 1 & \text{if } \xi_i' = \tilde{\xi}_i^* \\ \xi_i' & \text{if } \xi_i' > \tilde{\xi}_i^* \\ \xi_i' + 1 & \text{if } 1 \le \xi_i' < \tilde{\xi}_i^*. \end{cases}$$
(4.5)

Comparing (4.5) with (4.4), we have $\eta''_i = \eta_i$. Now, we know that $i \in \operatorname{Cyc}(s'', \gamma'')$ in g''_i if and only if $\xi''_i = \tilde{\eta}^*_i$, and with η''_i and $\gamma''_1, \ldots, \gamma''_{i-1}$ we can construct the edges $i \to v'_1, v_1 \to v'_2, \ldots, v_{s-1} \to v'_s$ in g''_i . By induction hypothesis $\gamma''_j = \gamma_j$ for $1 \le j \le i-1$ and $\eta''_i = \eta_i$. So



FIGURE 5. The graph of $\Phi(\sigma) = (1, 4, 9, 5)(7)(8)(2, 11, 6, 10, 3)(12)(13, 17)(14, 16)(15)$

 $\tilde{\eta}_i^* = \xi_i^*$. Finally

$$\xi_i'' = \begin{cases} \tilde{\eta}_i^* & \text{if } \eta_i' = 1\\ \eta_i' & \text{if } \eta_i' > \tilde{\eta}_i^*\\ \eta_i' - 1 & \text{if } 1 < \eta_i' \le \tilde{\eta}_i^*. \end{cases}$$
(4.6)

Comparing (4.6) with (4.3), we have $\xi_i'' = \xi_i$. Thus we complete the proof.

Lemma 11. For $(s, \gamma) \in \mathcal{LH}_n$, if $\rho_2(s, \gamma) = (s', \gamma')$, then

 $(Cyc, Arecp, Exc, Rar)(s', \gamma') = (Arecp, Cyc, Exc, Rar)(s, \gamma).$

Proof. By definition of ρ_2 in § 4.1, if $\rho_2(s_i, \gamma_i) = (s'_i, \gamma'_i)$ for $1 \le i \le n$, then $s_i = s'_i$ for $s_i \in \{U, D, L_a\}$ and $\{i : s_i = L_b \text{ or } L_c\} = \{i : s'_i = L_b \text{ or } L_c\}$.

- (1) As $\operatorname{Excp}(s,\gamma) = \{i : s_i = U \text{ or } L_a\}$ and $\operatorname{Excl}(s,\gamma) = \{i : s_i = D \text{ or } L_a\}$, see (2.6), we have $\operatorname{Exc}(s',\gamma') = \operatorname{Exc}(s,\gamma)$.
- (2) Recall that $\operatorname{Cyc}(s,\gamma) = \{i : s_i = \mathcal{L}_c \text{ or } s_i = \mathcal{D} \text{ with } i \to i'_1, i_1 \to i'_2, \dots, i_k \to i' \text{ in } g_i\}$ and $\operatorname{Arecp}(s',\gamma') = \{i : s'_i \in \{\mathcal{D}, \mathcal{L}_b, \mathcal{L}_c\} \text{ and } \eta'_i = 1\}$, see (2.2)-(2.9). Now,
 - the case $(s_i, \gamma_i) = (L_c, (\Delta, h_{i-1} + 1))$ means there is an edge $i \to i'$ in g_i , which is equivalent to $(s'_i, \gamma'_i) = (L_b, (\Delta, 1))$ or $(L_c, (\Delta, 1))$ by § 4.1 (iv).
 - the case $s_i = D$ with $i \to v'_1, v_1 \to v'_2, \dots, v_{s-1} \to v'_s, v_s \to i'$ in g_i is equivalent to $s'_i = D, \eta'_i = 1$ by § 4.1 (v).

Thus $\operatorname{Arecp}(s', \gamma') = \operatorname{Cyc}(s, \gamma)$. As ρ_2 is an involution from \mathcal{LH}_n onto itself, we derive immediately that $\operatorname{Cyc}(s', \gamma') = \operatorname{Arecp}(s, \gamma)$.

(3) Recall that $\operatorname{Rar}(s, \gamma) := \{i : s_i = L_c \text{ and } \eta_i = 1\}$. By § 4.1 (iv), $(s_i, \gamma_i) = (L_c, (\Delta, 1))$ if and only if $(s'_i, \gamma'_i) = (L_c, (\Delta, 1))$. Hence $\operatorname{Rar}(s', \gamma') = \operatorname{Rar}(s, \gamma)$.

Let $\Phi := \theta^{-1} \circ \rho_1 \circ \theta$. By Lemmas 6 and 11 we see that Φ is the desired bijection from \mathfrak{S}_n onto itself for Theorem 2.

Example. For our running example σ , we get $\tau := \Phi(\sigma)$ with

Thus

$$\operatorname{Rec} \tau = (\{1, 2, 6, 12, 13\}, \{4, 10, 11, 12, 17\})$$



FIGURE 6. The Laguerre history of $\Phi(\sigma)$

Arec
$$\tau = (\{5, 10, 11, 12, 17\}, \{1, 3, 6, 12, 13\})$$

Erec $\tau = (\{1, 2, 6, 13\}, \{4, 10, 11, 17\})$
Exc $\tau = (\{1, 2, 4, 6, 13, 14\}, \{4, 9, 10, 11, 16, 17\})$

and Rar $\tau = \{12\}$. Also, the cycle decomposition of τ is

$$\tau = (1, 4, 9, 5) (2, 11, 6, 10, 3) (7) (8) (12) (13, 17) (14, 16) (15).$$

Thus Fix $\tau = \{7, 8, 12, 15\}$, Cyc $\tau = \{7, 8, 9, 11, 12, 15, 16, 17\}$, Cpeak $\tau = \{9, 10, 11, 16, 17\}$, Cval $\tau = \{1, 2, 6, 13, 14\}$, Cdrise $\tau = \{4\}$, and Cdfall $\tau = \{3, 5\}$.

5. Acknowledgements

The first author was supported by the *China Scholarship Council*. This work was done during his visit at Université Claude Bernard Lyon 1 in 2018-2019.

References

- [BV17] J.-L. Baril, V. Vajnovszki, A permutation code preserving a double Eulerian bistatistic, Discrete Appl. Math. 224 (2017) 9–15.
- [Bi93] P. Biane, Permutations suivant le type d'excédance et le nombre d'inversions et interprétation combinatoire d'une fraction continue de Heine, European J. Combin. 14 (1993), no. 4, 277–284.
- [CSZ97] R. J. Clarke, E. Steingrímsson, and J. Zeng, New Euler-Mahonian statistics on permutations and words, Adv. in Appl. Math. 18 (1997), no. 3, 237–270.
- [CO09] R. Cori, Indecomposable permutations, hypermaps and labeled Dyck paths, J. Combin. Theory Ser. A 116 (2009), no. 8, 1326–1343.
- [DV94] A. de Médicis, G. X. Viennot, Moments des q-polynômes de Laguerre et la bijection de Foata-Zeilberger. Adv. in Appl. Math. 15 (1994), no. 3, 262–304.
- [FH09] D. Foata and Guo-Niu, Han, New permutation coding and equidistribution of set-valued statistics. Theoret. Comput. Sci. 410 (2009), no. 38–40, 3743–3750.
- [Han19] G. N. Han, private communication, January 2019.
- [KL18] D.-S., Kim and Z. Lin, A sextuple equidistribution arising in pattern avoidance. J. Combin. Theory Ser. A 155 (2018), 267–286.
- [PO14] S. Poznanović, The sorting index and equidistribution of set-valued statistics over restricted permutations. J. Combin. Theory Ser. A 125 (2014), 254–272.

JIANXI MAO AND JIANG ZENG

[SZ19] A. Sokal and J. Zeng, Some multivariate master polynomials for permutations, set partitions, and perfect matchings, and their continued fractions, in preparation, 2019.

(Jianxi Mao) School of Mathematic Sciences, Dalian University of Technology, Dalian 116024, P. R. China

E-mail address: maojianxi@hotmail.com

(Jiang Zeng) UNIV LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, CNRS UMR 5208, INSTITUT CAMILLE JORDAN, 43 BLVD. DU 11 NOVEMBRE 1918, F-69622 VILLEURBANNE CEDEX, FRANCE *E-mail address*: zeng@math.univ-lyon1.fr

14