

Some remarks on relative framed motives

Mingazov A. A.

November 13, 2019

1. Introduction

The purpose of this paper is to prove two statements which are key in the construction of motivic fibration replacement for suspension spectrum concerned with motivic space $\frac{X}{X-Z}$, where X is smooth variety and Z is its smooth subvariety. In fact this generalizes the main result of the paper [5]. The main results are formulated in theorems 5.2 and 6.2.

2. Basic notions

Definition 2.1. Let Y be a k -smooth scheme, $S \subset Y$ be a closed subset and $U \in Sm/k$. A framed correspondence of level m from U to $Y/(Y-S)$ is the set of data

$$(Z, W, \phi_1, \dots, \phi_m; g: W \rightarrow Y),$$

where $Z \subset U \times \mathbb{A}^m$ is closed and finite over U , W is étale neighborhood of Z in $U \times \mathbb{A}^m$, ϕ_1, \dots, ϕ_m are regular functions on W , g is a regular map such as $Z = g^{-1}(S) \cap V(\phi_1, \dots, \phi_m)$.

Two correspondences are equivalent if they coincide on the common étale neighborhood.

Notation 2.1. 1) $Fr_m(U, Y/(Y-S))$ is the set of framed correspondences from U to $Y/(Y-S)$ up to equivalence.

2) $F_m(U, Y/(Y-S))$ is the set of framed correspondences from U to $Y/(Y-S)$ up to equivalence with connected support Z .

$$3) Fr_*(U, Y/(Y-S)) = \bigsqcup_{m \geq 0} Fr_m(U, Y/(Y-S)).$$

4) $Fr(U, Y/(Y-S))$ is the set stabilized up to the map $\Sigma: \Phi \mapsto \Phi \boxtimes \sigma$, where $\sigma = (\{0\}, \mathbb{A}^1, id: \mathbb{A}^1 \rightarrow \mathbb{A}^1, const: \mathbb{A}^1 \rightarrow pt) \in Fr_1(pt, pt)$.

5) $\mathbb{Z}Fr_m(U, Y/(Y - S))$ is the free abelian group generated by Fr_m .

6) $\mathbb{Z}F_*(U, Y/(Y - S))$ is the factor of $\mathbb{Z}Fr_m(U, Y/(Y - S))$ up to the relation:

$$(Z \sqcup Z', W, (\phi_1, \dots, \phi_m); g: W \rightarrow Y) = \\ = (Z, W \setminus Z', (\phi_1, \dots, \phi_m)|_{W \setminus Z'}; g|_{W \setminus Z'}) + (Z, W \setminus Z, (\phi_1, \dots, \phi_m)|_{W \setminus Z}; g|_{W \setminus Z})$$

Note that $F_m(U, Y/(Y - S))$ is a basis of the group $\mathbb{Z}F_m(U, Y/(Y - S))$.

Definition 2.2. Presheaf \mathcal{F} with $\mathbb{Z}F_*$ -transfers is a functor $\mathcal{F}: \mathbb{Z}F_*^{op} \mathring{A}b$. The category of Nisnevich shaeves with $\mathbb{Z}F_*$ -transfers will be denoted as $NSZF_*$.

Lemma 2.1. $\mathbb{Z}F_*(-, X)$ is a Nisnevich sheaf.

3. Some statements on cohomologies of sheaves with $\mathbb{Z}F_*$ -transfers

The statements of this section are modifications of similar theorems of [1].

Lemma 3.1. Let $f: Y \rightarrow X$ be a Nisnevich covering of variety X . Then the sequence of the sheaves

$$0 \leftarrow \mathbb{Z}F_n(X) \xleftarrow{f_*} \mathbb{Z}F_n(Y) \xleftarrow{(p_2)_* - (p_1)_*} \mathbb{Z}F_n(Y \times_X Y) \leftarrow \dots$$

is exact.

Proof. Let U be a local Henselian ring. We need to prove the exactness of the complex

$$A_*(U) = \left(0 \leftarrow \mathbb{Z}F_n(U, X) \xleftarrow{f_*} \mathbb{Z}F_n(U, Y) \xleftarrow{(p_2)_* - (p_1)_*} \mathbb{Z}F_n(U, Y \times_X Y) \leftarrow \dots \right).$$

Fix framed correspondence α

$$\begin{array}{ccccc} & & W & \longrightarrow & X \times \mathbb{A}^n \\ & \swarrow & \uparrow & & \uparrow \\ U \times \mathbb{A}^n & \longleftarrow & Z & \longrightarrow & X \times 0, \\ & & \downarrow & & \\ & & U & & \end{array}$$

where scheme Z is connected.

Let $A_k^\alpha(U)$ be a subgroup of $A_k(U)$ generated by correspondences β , which after composition with $\text{map } Y \times_X Y \times \dots \times_X Y \rightarrow Y$ are divisible by α . Note that $A_*^\alpha(U)$ is subcomplex in $A_*(U)$ and $A_* = \bigoplus_{\alpha} A_*^\alpha(U)$. In particular, the support of all elements of $A_*^\alpha(U)$ is Z . It is enough to prove that the complex $A_*^\alpha(U)$ is contractible.

As Z is connected and finite over local Henselian ring U , it is local and Henselian. Consider two Cartesian squares

$$\begin{array}{ccccc} Z \times_X Y & \longrightarrow & W \times_X Y & \longrightarrow & Y \\ \downarrow & & \downarrow & & \downarrow f \\ Z & \longrightarrow & W & \longrightarrow & X. \end{array}$$

The map $Z \times_X Y \rightarrow Z$ is etale as the pullback of the etale map $f: Y \rightarrow X$, so it has a section $s: Z \rightarrow Z \times_X Y$ as Z is Henselian. Let fix it. As W is Henselian too, we will suppose that there exist some underline diagramm $Z \hookrightarrow W \xrightarrow{g} Y$.

Now we construct a contracting homotopy $h_k^\alpha: A_k^\alpha(U) \rightarrow A_{k+1}^\alpha(U)$. Let us take $\beta \in A_k^\alpha(U)$

$$\begin{array}{ccccc} & & W & \xrightarrow{q} & Y \times_X \dots \times_X Y \times \mathbb{A}^n \\ & \swarrow & \uparrow & & \uparrow \\ U \times \mathbb{A}^n & \longleftarrow & Z & \longrightarrow & Y \times_X \dots \times_X Y \times 0. \\ & \searrow & \downarrow & & \downarrow \\ & & U & & \end{array}$$

Then $h(\beta)$ can be defined as

$$\begin{array}{ccccc} & & W & \xrightarrow{q \times_X g} & Y \times_X \dots \times_X Y \times_X Y \times \mathbb{A}^n \\ & \swarrow & \uparrow & & \uparrow \\ U \times \mathbb{A}^n & \longleftarrow & Z & \longrightarrow & Y \times_X \dots \times_X Y \times_X Y \times 0. \\ & \searrow & \downarrow & & \downarrow \\ & & U & & \end{array}$$

The properties of contracting homotopy can be verified straightly. \square

Statement 3.1. *Let X be a smooth variety. Then presheaves $H^i(C^*ZF(X))$ are quasistable.*

Corollary 3.1. *Let I be an injective Nisnevich sheaf with $\mathbb{Z}F_*$ -transfers. Then for any $X \in \text{Sm}/k$*

$$H_{\text{Nis}}^i(X, I) = 0.$$

Corollary 3.2. *For any Nisnevich sheaf \mathcal{F} with $\mathbb{Z}F_*$ -transfers*

$$\text{Ext}_{\text{NSZF}_*}^i(\mathbb{Z}F_*(X), \mathcal{F}) = H_{\text{Nis}}^i(X, \mathcal{F}).$$

Proposition 3.1. *Let $A^* \in D^-(\text{NSZF}_*)$ be a bounded complex of Nisnevich sheaves with $\mathbb{Z}F_*$ -transfers. Then for any $X \in \text{Sm}/k$*

$$H_{\text{Nis}}^i(X, A^*) = \text{Hom}_{D^-(\text{NSZF}_*)}(\mathbb{Z}F_*(X), A^*[i]).$$

Definition 3.1. Presheaf is called contractible if there exist the presheaf morphism $\phi: \mathcal{F}(-) \rightarrow \mathcal{F}(\Delta^1 \times -)$ such as $\partial_0\phi = 0$, $\partial_1\phi = 1_{\mathcal{F}}$.

Proposition 3.2. *Let \mathcal{G}, \mathcal{F} be quasistable Nisnevich sheaves with $\mathbb{Z}F_*$ -transfers. Suppose that \mathcal{G} is contractible, \mathcal{F} is strongly homotopy invariant. Then $\text{Ext}_{\text{NSZF}_*}^i(\mathcal{G}, \mathcal{F}) = 0$ for all $i \geq 0$.*

Corollary 3.3. *Let \mathcal{G}, \mathcal{F} be quasistable Nisnevich sheaves with $\mathbb{Z}F_*$ -transfers. Suppose that \mathcal{F} is strongly homotopy invariant and \mathcal{G} has a finite resolution*

$$0 \rightarrow \mathcal{G} \rightarrow \mathcal{G}^0 \rightarrow \dots \rightarrow \mathcal{G}^n \rightarrow 0,$$

where \mathcal{G}^i is contractible and quasistable Nisnevich sheaf with $\mathbb{Z}F_*$ -transfers for all i . Then $\text{Ext}_{\text{NSZF}_*}^i(\mathcal{G}, \mathcal{F}) = 0$.

Corollary 3.4. *Let A^* be a bounded complex of quasistable Nisnevich sheaves with $\mathbb{Z}F_*$ -transfers, A^i is contractible for all i , cohomology presheaves $\mathcal{H}^i = H^i(A^*)$ are homotopy invariant and quasistable. Then the complex A^* is acyclic.*

Corollary 3.5. *Let A^* be a complex of Nisnevich sheaves with $\mathbb{Z}F_*$ -transfers which satisfies the following conditions:*

- 1) $A_{\text{Nis}}^* = 0$;
 - 2) presheaves $H^i(C^*(A^*))$ are quasistable.
- Then $C^*(A^*)$ is locally acyclic.

4. Mayer-Vietoris sequence and etale excision for framed motives

Statement 4.1. *Let $0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_3 \rightarrow 0$ be an exact sequence of Nisnevich sheaves with $\mathbb{Z}F_*$ -transfers and cohomology presheaves $H^i(C^*(\mathcal{F}_r))$ are quasistable for all i, r . Then*

$$C^*(\mathcal{F}_1) \rightarrow C^*(\mathcal{F}_2) \rightarrow C^*(\mathcal{F}_3) \rightarrow C^*(\mathcal{F}_1)[1]$$

is distinguished triangle in $D^-(NSZF_*)$.

Statement 4.2. Let A, B, C, D be sets. Consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & B \\ \downarrow \beta & & \downarrow \delta \\ C & \xrightarrow{\gamma} & D \end{array}$$

and the sequence of abelian groups

$$0 \rightarrow \mathbb{Z}[A] \xrightarrow{\begin{pmatrix} \alpha \\ -\beta \end{pmatrix}} \begin{array}{c} \mathbb{Z}[B] \\ \oplus \\ \mathbb{Z}[C] \end{array} \xrightarrow{(\delta \ \gamma)} \mathbb{Z}[D] \rightarrow 0.$$

Then

- 1) the sequence is the complex if and only if the diagram comutes;
- 2) the sequence is exact on the left if and only if the diagram is Cartesian square;
- 3) the sequence is exact on the right if and only if the diagram is pushout square.

Lemma 4.1. Let U be a spectrum of Henselian ring, $X \subset Y$ be an open imbedding. Then the map $F_n(U, X) \rightarrow F_n(U, Y)$ is injective.

Lemma 4.2. Let $X = X_1 \cup X_2$ be a union of its open subsets, $Z \subset X$ be a closed subvariety. Denote $X_{12} = X_1 \cap X_2$, $Z_1 = Z \cap X_1$, $Z_2 = Z \cap X_2$, $Z_{12} = Z \cap X_1 \cap X_2$. Let $v_i: X_{12} \rightarrow X_i$ be inclusion $i = \overline{1}, \overline{2}$, $u_i: X_i \rightarrow X$, $i = \overline{1}, \overline{2}$. Then the sequence of Nisnevich sheaves

$$0 \rightarrow \mathbb{Z}F_n \left(\frac{X_{12}}{X_{12} - Z_{12}} \right) \xrightarrow{\begin{pmatrix} v_{1*} \\ -v_{2*} \end{pmatrix}} \begin{array}{c} \mathbb{Z}F_n \left(\frac{X_1}{X_1 - Z_1} \right) \\ \oplus \\ \mathbb{Z}F_n \left(\frac{X_2}{X_2 - Z_2} \right) \end{array} \xrightarrow{(u_{1*} \ u_{2*})} \mathbb{Z}F_n \left(\frac{X}{X - Z} \right) \rightarrow 0$$

is exact.

Definition 4.1. Let X be a smooth variety and $Z \subset X$ is its clozed subset. Let us denote

$$M_Z^{fact} X = \frac{C^* \mathbb{Z}F(X)}{C^* \mathbb{Z}F(X - Z)},$$

$$M_Z^{geom} X = C^* \mathbb{Z}F \left(\frac{X}{X-Z} \right).$$

Note that there exist the natural map

$$M_Z^{fact}(X) \rightarrow M_Z^{geom}(X).$$

Statement 4.3. *Let the variety $X = X_1 \cup X_2$ be an union of its open subsets, $Z \subset X$ be its clozed subvariety. Denote $X_{12} = X_1 \cap X_2$, $Z_1 = Z \cap X_1$, $Z_2 = Z \cap X_2$, $Z_{12} = Z \cap X_1 \cap X_2$. Then the triangles*

$$1) M_{Z_{12}}^{fact}(X_{12}) \rightarrow M_{Z_1}^{fact}(X_1) \oplus M_{Z_2}^{fact}(X_2) \rightarrow M_Z^{fact} X \rightarrow M_{Z_{12}}^{fact}(X_{12})[1],$$

$$2) M_{Z_{12}}^{geom}(X_{12}) \rightarrow M_{Z_1}^{geom}(X_1) \oplus M_{Z_2}^{geom}(X_2) \rightarrow M_Z^{fact} X \rightarrow M_{Z_{12}}^{geom}(X_{12})[1]$$

are distinguished.

Corollary 4.1. *Let $\pi: L \rightarrow X$ be a vector bundle over X , $X_1, X_2 \subset X$ be open subsets, $X_{12} = X_1 \cap X_2$, $L_i = L|_{X_i}$. Then the triangles*

$$M_{X_{12}}^{fact}(L_{12}) \rightarrow M_{X_1}^{fact}(L_1) \oplus M_{X_2}^{fact}(L_2) \rightarrow M_X^{fact}(L),$$

$$M_{X_{12}}^{geom}(L_{12}) \rightarrow M_{X_1}^{geom}(L_1) \oplus M_{X_2}^{geom}(L_2) \rightarrow M_X^{geom}(L)$$

are distinguished in $D^-(NSZF_*)$.

Theorem 4.1. *Let $p: L \rightarrow X$ be a vector bundle. Then the map*

$$M_X^{fact}(L) \rightarrow M_X^{geom}(L)$$

is the local isomorphism.

Proof. We'll use the induction on the number of open sets covering X on which L is trivial. If L is trivial on X , the statement of the theorem is the main result of [5]. Let $X_1, X_2 \subset X$ be open subsets and $X_1 \cup X_2 = X$. If $L_1 = L|_{X_1}$ and $L_2 = L|_{X_2}$ are trivial, then $L_{12} = L|_{X_{12}}$ is trivial too. Consider the diagramm

$$\begin{array}{ccccc} C^* \mathbb{Z}F \left(\frac{L_{12}}{L_{12} - X_{12}} \right) & \longrightarrow & C^* \mathbb{Z}F \left(\frac{L_1}{L_1 - X_1} \right) \oplus C^* \mathbb{Z}F \left(\frac{L_2}{L_2 - X_2} \right) & \longrightarrow & C^* \mathbb{Z}F \left(\frac{L}{L - X} \right) \\ \uparrow & & \uparrow & & \uparrow \\ \frac{C^* \mathbb{Z}F(L_{12})}{C^* \mathbb{Z}F(L_{12} - X_{12})} & \longrightarrow & \frac{C^* \mathbb{Z}F(L_1)}{C^* \mathbb{Z}F(L_1 - X_1)} \oplus \frac{C^* \mathbb{Z}F(L_2)}{C^* \mathbb{Z}F(L_2 - X_2)} & \longrightarrow & \frac{C^* \mathbb{Z}F(L)}{C^* \mathbb{Z}F(L - X)} \end{array}$$

String are distinguished triangles and two vertical maps are isomorphisms in $D^-(NSZF_*)$. \square

Statement 4.4. Let $f: X' \rightarrow X$ be etale morphism, $Z \subset X$ be a closed subset and $f: f^{-1}(Z) \rightarrow Z$ be an isomorphism. Then

- 1) the map $f_*^{fact}: M_Z^{fact}(X') \rightarrow M_Z^{fact}(X)$ is isomorphism in $D^-(NSZF_*)$,
- 2) the map $f_*^{geom}: M_Z^{geom}(X') \rightarrow M_Z^{geom}(X)$ is isomorphism in $D^-(NSZF_*)$.

5. Deformation to normal bundle

Statement 5.1. Let X be a smooth variety, Z is its closed subset. Denote with $p: X \times \mathbb{A}^1 \rightarrow X$ projection. Then the map

$$p_*: M_{Z \times \mathbb{A}^1}^{geom}(X \times \mathbb{A}^1) \rightarrow M_Z^{geom}(X)$$

is isomorphism in $D^-(NSZF_*)$.

Corollary 5.1. Let $p: E \rightarrow X$ be a vector bundle over X , $Z \subset X$, $S = p^{-1}(Z)$. Then the map

$$p_*: M_S^{geom}(E) \rightarrow M_Z^{geom}(X)$$

is isomorphism in $D^-(NSZF_*)$.

Proof. This follows from the previous statement by applying 4.3. \square

Statement 5.2. Let X be a smooth variety and Z is its closed subset. Denote with $p: X \times \mathbb{A}^1 \rightarrow X$ the projection. Then the map

$$p_*: M_{Z \times \mathbb{A}^1}^{fact}(X \times \mathbb{A}^1) \rightarrow M_Z^{fact}(X)$$

is isomorphism in $D^-(NSZF_*)$.

Proof. This follows from the homotopical invariance of $C^*\mathbb{Z}F(X)$. \square

Corollary 5.2. Let $p: E \rightarrow X$ be a vector bundle over X , $Z \subset X$, $S = p^{-1}(Z)$. Then the map

$$p_*: M_S^{fact}(E) \rightarrow M_Z^{fact}(X)$$

is an isomorphism in $D^-(NSZF_*)$.

Proof. This follows from the previous statement by applying 4.3. \square

Statement 5.3. Let X be a smooth variety, $Z \subset X$ be its closed subset and $N = N_{X/Z}$ be the normal bundle. Denote with X_t deformation to normal bundle. Let U and V be open subsets of X . Then

- a) $U_t \cap V_t = (U \cap V)_t$;

- b) $U_t \cup V_t = (U \cup V)_t$;
c) Let

$$\begin{array}{ccccc} \tilde{X} - \tilde{Z} & \hookrightarrow & \tilde{X} & \longleftarrow & \tilde{Z} \\ \downarrow e|_{\tilde{X}-\tilde{Z}} & & \downarrow e & & \parallel \\ X - Z & \hookrightarrow & X & \longleftarrow & Z \end{array}$$

be an elemental Nisnevich square. Then

$$\begin{array}{ccccc} \tilde{X}_t - \tilde{Z} \times \mathbb{A}^1 & \hookrightarrow & \tilde{X}_t & \longleftarrow & \tilde{Z} \times \mathbb{A}^1 \\ \downarrow & & \downarrow e_t & & \parallel \\ X_t - Z \times \mathbb{A}^1 & \hookrightarrow & X_t & \longleftarrow & Z \times \mathbb{A}^1 \end{array}$$

is also elemental Nisnevich square.

Proof. This follows from construction of X_t described in [6]. \square

Theorem 5.1. Let X be a smooth variety, $Z \subset X$ be its smooth subvariety and $N = N_{X/Z}$ be the normal bundle. Denote with X_t deformation to normal bundle. The maps of pairs

$$(N, N - Z) \xrightarrow{i_0} (X_t, X_t - Z \times \mathbb{A}^1) \xleftarrow{i_1} (X, X - Z)$$

induce the isomorphisms

- 1) $M_Z^{geom}(N) \xrightarrow{\sim} M_{Z \times \mathbb{A}^1}^{geom}(X_t) \xleftarrow{\sim} M_Z^{geom}(X)$,
- 2) $M_Z^{fact}(N) \xrightarrow{\sim} M_{Z \times \mathbb{A}^1}^{fact}(X_t) \xleftarrow{\sim} M_Z^{fact}(X)$.

Proof. The proof is similar to the proof of Theorem 1.2 from [6]. \square

Lemma 5.1. Let $p: E \rightarrow Z$ be a vector bundle over smooth Z , $i: Z \hookrightarrow E$ be a zero section. Then the maps of pairs

$$(N, N - Z) \xrightarrow{i_0} (E_t, E_t - Z \times \mathbb{A}^1) \xleftarrow{i_1} (E, E - Z),$$

where the normal bundle N also coincides with E , induce the isomorphisms

- 1) $M_Z^{geom}(N) \xrightarrow{\sim} M_{Z \times \mathbb{A}^1}^{geom}(E_t) \xleftarrow{\sim} M_Z^{geom}(E)$,
- 2) $M_Z^{fact}(N) \xrightarrow{\sim} M_{Z \times \mathbb{A}^1}^{fact}(E_t) \xleftarrow{\sim} M_Z^{fact}(E)$.

Proof. The proof is similar to the proof of lemma 2.5 in [6]. \square

Theorem 5.2. Let X be a smooth variety and $Z \subset X$ be its smooth subvariety. Then the map

$$M_Z^{fact}(X) \rightarrow M_Z^{geom}(X)$$

is the isomorphism in $D^-(NSZF_*)$.

Proof. Consider the deformation to normal bundle X_t . We have the diagram

$$\begin{array}{ccccc} M_Z^{fact}(X) & \xrightarrow{i_{1*}} & M_{Z \times \mathbb{A}^1}^{fact}(X_t) & \xleftarrow{i_{0*}} & M_Z^{fact}(N) \\ \downarrow & & \downarrow & & \downarrow \\ M_Z^{geom}(X) & \xrightarrow{i_{1*}} & M_{Z \times \mathbb{A}^1}^{geom}(X_t) & \xleftarrow{i_{0*}} & M_Z^{geom}(N). \end{array}$$

By the theorems 5.1 and 4.1 we get the statement. \square

6. Cancelcation theorem

Definition 6.1. Let U, X be smooth varieties over field k , $S \subset X$ be a closed subset. We'll define the set $Fr_n^{qf} \left(U, \frac{X}{X-Z} \right)$ as a set of the following data

$$\begin{array}{ccc} W = (X \times \mathbb{A}^n)_Z^h & \xrightarrow{(g, \phi_1, \dots, \phi_n)} & X \times \mathbb{A}^n \\ \uparrow e & \swarrow i & \uparrow \\ U \times \mathbb{A}^n & \xleftarrow{\quad} Z' = V(\phi_1, \dots, \phi_n) & \xrightarrow{\quad} S \times 0 \\ \downarrow p_U & \downarrow p & \downarrow \\ U & & U \end{array}$$

where $e : W = (X \times \mathbb{A}^n)_Z^h \rightarrow X \times \mathbb{A}^n$ is Henselian neighborhood of Z , Z' is a set-theoretical type pullback of $S \times 0$ in W , besides the composition $p_U \circ e \circ i : Z' \rightarrow U$ is quasifinite.

Denote with $F_n^{qf} \left(U, \frac{X}{X-S} \right)$ the subset of $Fr_n^{qf} \left(U, \frac{X}{X-S} \right)$ with connected supports Z .

Lemma 6.1. *Let V be an affine scheme, $Z \subset V$ be a connected closed subset, $can = can_{V,Z} : V_Z^h \rightarrow V$ be a Henselisation of V in Z , $s : Z \rightarrow V_Z^h$ be a section can over Z . Let U be a regular local Henselian scheme, $q : V \rightarrow U$ be a smooth morphism such that $q|_Z : Z \rightarrow U$ is finite. Suppose that $Y \subset V_Z^h$ is closed subset contained $s(Z)$ and quasifinite over U . Then $can : Y \rightarrow V$ is closed imbedding, $can(Y)$ contains Z and $can^{-1}(can(Y)) = Y$.*

Proof. This lemma proved in [5]. \square

Lemma 6.2. *Let X be a smooth affine variety over field k , $Z \subset Y \subset X$ be closed subsets, $g: X_Z^h \rightarrow X$ be a henselization of X in Z . Suppose that there exist the inclusion $j: Y \hookrightarrow X_Z^h$ such that the diagram*

$$\begin{array}{ccc} X_Z^h & & \\ g \downarrow & \nearrow j & \\ X & \xleftarrow{i} & Y \end{array}$$

conutes and $g^{-1}(Y) = j(Y)$. Then the canonical map $X_Y^h \rightarrow X_Z^h$ is the isomorphism of schemes.

Remark 6.1. The canonical map $p: F_n(U, X) \rightarrow F_n\left(U, \frac{X}{X-S}\right)$ induces the map $p: F_n(U, X) \rightarrow F_n^{qf}\left(U, \frac{X}{X-S}\right)$.

Theorem 6.1. *Let U be a Henselian scheme, X be a smooth variety, $S \subset X$ be a closed subset. Then the next square is cocartesian*

$$\begin{array}{ccc} F_n(U, X-S) & \xrightarrow{i} & F_n(U, X) \\ \downarrow & & \downarrow p \\ * & \hookrightarrow & F_n^{qf}\left(U, \frac{X}{X-S}\right). \end{array}$$

Corollary 6.1. *Let X be a smooth variety over field k , $S \subset X$ be closed subset. Then there exist the following exact sequence*

$$0 \rightarrow \mathbb{Z}F(X-S) \rightarrow \mathbb{Z}F(X) \rightarrow \mathbb{Z}F^{qf}\left(\frac{X}{X-S}\right) \rightarrow 0.$$

Corollary 6.2. *Let X be a smooth variety over field k , $S \subset X$ be closed subset. Then the following triangle*

$$\mathbb{Z}F(X-S) \rightarrow \mathbb{Z}F(X) \rightarrow \mathbb{Z}F^{qf}\left(\frac{X}{X-S}\right) \rightarrow \mathbb{Z}F(X-S)[1]$$

is distinguished in $D^-(NSZF_)$.*

Lemma 6.3. *Let \mathcal{F} be a homotopy invariati quasistable presheave with $\mathbb{Z}F_*$ -transfers, X be a local Henselian scheme over field k . Then*

$$H_{Nis}^i(X \times \mathbb{G}_m, \mathcal{F}) = \begin{cases} \mathcal{F}(X \times \mathbb{G}_m) & \text{for } i = 0, \\ 0 & \text{for } i \neq 0. \end{cases}$$

Theorem 6.2. *Let $X = \text{Spec } \mathcal{O}$, where \mathcal{O} is Henselian local ring over field k , Y be a smooth variety, $S \subset Y$ be its closed subset. Then the map*

$$-\boxtimes(\text{id}_{\mathbb{G}_m} - e_1): \mathbb{Z}F^{qf} \left(\Delta^\bullet \times X, \frac{Y}{Y-S} \right) \rightarrow \mathbb{Z}F^{qf} \left(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1} \right)$$

is quasiisomorphism of complexes.

Proof. Consider two distinguished triangles in $D^-(NSZF_*)$:

$$\begin{aligned} 1) \quad & \mathbb{Z}F(\Delta^\bullet \times -, Y-S) \rightarrow \mathbb{Z}F(\Delta^\bullet \times -, Y) \rightarrow \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S}), \\ 2) \quad & \mathbb{Z}F(\Delta^\bullet \times -, (Y-S) \times \mathbb{G}_m^{\wedge 1}) \rightarrow \mathbb{Z}F(\Delta^\bullet \times -, Y \times \mathbb{G}_m^{\wedge 1}) \rightarrow \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}). \end{aligned}$$

From 1) we get the following exact sequence for scheme X

$$0 \rightarrow \mathbb{Z}F(\Delta^\bullet \times X, Y-S) \rightarrow \mathbb{Z}F(\Delta^\bullet \times X, Y) \rightarrow \mathbb{Z}F^{qf}(\Delta^\bullet \times X, \frac{Y}{Y-S}) \rightarrow 0$$

From 2) we get for $X \times \mathbb{G}_m$:

$$\begin{aligned} 0 \rightarrow & H_{Nis}^0(X \times \mathbb{G}_m, \mathbb{Z}F(\Delta^\bullet \times -, (Y-S) \times \mathbb{G}_m^{\wedge 1})) \rightarrow \\ & \rightarrow H_{Nis}^0(X \times \mathbb{G}_m, \mathbb{Z}F(\Delta^\bullet \times -, Y \times \mathbb{G}_m^{\wedge 1})) \rightarrow \\ & \rightarrow H_{Nis}^0(X \times \mathbb{G}_m, \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})) \rightarrow \\ & \rightarrow H_{Nis}^1(X \times \mathbb{G}_m, \mathbb{Z}F(\Delta^\bullet \times -, (Y-S) \times \mathbb{G}_m^{\wedge 1})) \rightarrow \dots \end{aligned}$$

We'll calculate the hypercohomologies in this sequence. We restrict ourselves to calculating cohomology $H_{Nis}^i \left(X \times \mathbb{G}_m, \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}) \right)$. The remaining cohomologies can be calculated in the same way.

Consider the spectral sequence

$$\begin{aligned} H_{Nis}^p \left(X \times \mathbb{G}_m, \left(\underline{h}^q(\mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})) \right)_{Nis} \right) & \Longrightarrow \\ \Longrightarrow H_{Nis}^{p+q} \left(X \times \mathbb{G}_m, \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}) \right), & \end{aligned}$$

where $\underline{h}^q(\mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}))$ — are preasheave cohomologies of the complex $\mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})$. On the left there are the usual

cohomology of sheaves, on the right there are the hypercohomology of the sheaves complex.

Denote the sheaf $\mathbb{Z}F^{qf}(-, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})$ with \mathcal{F} . Note that it is quasistable and is a sheaf with $\mathbb{Z}F_*$ -transfers. Then by the theorem from [4] presheaves $h^q(\mathcal{F}(\Delta^\bullet \times -))$ are homotopy invariant, quasistable and has $\mathbb{Z}F_*$ -transfers. Then by the theorem from [4] sheaves $\underline{h}^q(\mathcal{F}(\Delta^\bullet \times -))$ are also homotopy invariant, quasistable and has $\mathbb{Z}F_*$ -transfers. Then by lemma 6.3 we get

$$\begin{aligned} H_{Nis}^p \left(X \times \mathbb{G}_m, \left(\underline{h}^q(\mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})) \right)_{Nis} \right) &= \\ = \begin{cases} \underline{h}^q(\mathbb{Z}F^{qf}(\Delta^\bullet \times X \times \mathbb{G}_m, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})) & \text{for } i = 0, \\ 0 & \text{for } i \neq 0. \end{cases} \end{aligned}$$

That is, the spectral sequence degenerates at the first step, and exactly one nonzero group falls into the corresponding dimension of the limit. Then

$$H_{Nis}^n \left(X \times \mathbb{G}_m, \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}) \right) = \underline{h}^q \left(\mathbb{Z}F^{qf}(\Delta^\bullet \times X \times \mathbb{G}_m, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}) \right).$$

Considering the direct summands on the left and on the right we get

$$H_{Nis}^n \left(X \times \mathbb{G}_m^{\wedge 1}, \mathbb{Z}F^{qf}(\Delta^\bullet \times -, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}) \right) = \underline{h}^q \left(\mathbb{Z}F^{qf}(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1}) \right).$$

Similar calculation is true for sheaves $\mathbb{Z}F(\Delta^\bullet \times -, (Y-S) \times \mathbb{G}_m^{\wedge 1})$, $\mathbb{Z}F(\Delta^\bullet \times -, Y \times \mathbb{G}_m^{\wedge 1})$, so there is a long exact sequence

$$\begin{aligned} 0 \rightarrow \underline{h}^0(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, (Y-S) \times \mathbb{G}_m^{\wedge 1})) &\rightarrow \\ \rightarrow \underline{h}^0(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, Y \times \mathbb{G}_m^{\wedge 1})) &\rightarrow \\ \rightarrow \underline{h}^0(\mathbb{Z}F^{qf}(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})) &\rightarrow \\ \rightarrow \underline{h}^1(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, (Y-S) \times \mathbb{G}_m^{\wedge 1})) &\rightarrow \dots \end{aligned}$$

We'll show that the map

$$-\boxtimes(id_{\mathbb{G}_m} - e_1): \mathbb{Z}F^{qf}(\Delta^\bullet \times X, \frac{Y}{Y-S}) \rightarrow \mathbb{Z}F^{qf}(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, \frac{Y}{Y-S} \times \mathbb{G}_m^{\wedge 1})$$

induces the isomorphisms of complexes. The map $-\boxtimes (id_{\mathbb{G}_m} - e_1)$ induces the triangle morphism, and so induces a map of long exact sequences:

$$\begin{array}{ccc}
\cdots & & \cdots \\
\downarrow & & \downarrow \\
\underline{h}^n(\mathbb{Z}F(\Delta^\bullet \times X, (Y - S))) & \longrightarrow & \underline{h}^n(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, (Y - S) \times \mathbb{G}_m^{\wedge 1})) \\
\downarrow & & \downarrow \\
\underline{h}^n(\mathbb{Z}F(\Delta^\bullet \times X, Y)) & \longrightarrow & \underline{h}^n(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, Y \times \mathbb{G}_m^{\wedge 1})) \\
\downarrow & & \downarrow \\
\underline{h}^n\left(\mathbb{Z}F^{qf}\left(\Delta^\bullet \times X, \frac{Y}{Y - S}\right)\right) & \longrightarrow & \underline{h}^n\left(\mathbb{Z}F^{qf}\left(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, \frac{Y}{Y - S} \times \mathbb{G}_m^{\wedge 1}\right)\right) \\
\downarrow & & \downarrow \\
\underline{h}^{n+1}(\mathbb{Z}F(\Delta^\bullet \times X, (Y - S))) & \longrightarrow & \underline{h}^{n+1}(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, (Y - S) \times \mathbb{G}_m^{\wedge 1})) \\
\downarrow & & \downarrow \\
\underline{h}^{n+1}(\mathbb{Z}F(\Delta^\bullet \times X, (Y - S))) & \longrightarrow & \underline{h}^{n+1}(\mathbb{Z}F(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, (Y - S) \times \mathbb{G}_m^{\wedge 1})) \\
\downarrow & & \downarrow \\
\cdots & & \cdots
\end{array}$$

Horizontal maps except maybe

$$\underline{h}^n\left(\mathbb{Z}F^{qf}\left(\Delta^\bullet \times X, \frac{Y}{Y - S}\right)\right) \rightarrow \underline{h}^n\left(\mathbb{Z}F^{qf}\left(\Delta^\bullet \times X \times \mathbb{G}_m^{\wedge 1}, \frac{Y}{Y - S} \times \mathbb{G}_m^{\wedge 1}\right)\right)$$

are isomorphisms by theorem C from [2]. Therefore, the map of our interest is also an isomorphism. \square

References

- [1] Suslin A., Voevodsky V. Bloch–Kato Conjecture and Motivic Cohomology with Finite Coefficients. // The arithmetics and geometry of algebraic cycles, NATO Sci. Ser. C Math. Phys. Sci. 548, Kluwer Acad. Publ., Dordrecht, 2000, 117–189.
- [2] A. Ananyevskiy, G. Garkusha, I. Panin, Cancellation theorem for framed motives of algebraic varieties, preprint arXiv:1601.06642.
- [3] G. Garkusha, I. Panin, Framed motives of algebraic varieties (after V. Voevodsky), preprint arXiv:1409.4372v4.

- [4] G. Garkusha, I. Panin, Homotopy invariant presheaves with framed transfers, preprint arXiv:1504.00884.
- [5] G. Garkusha, A. Neshitov, I. Panin, Framed motives of relative motivic spheres, preprint arXiv:1604.02732.
- [6] Panin I., Smirnov A. Oriented cohomologies of algebraic varieties.