

## A NOTE ON A QUANTIZATION VIA THE RAN SPACE.

ALEXEY KALUGIN

ABSTRACT. In a present note we give a new proof of Etingof-Kazhdan quantization theorem.

## 1. INTRODUCTION

1.1. **Main result.** A quantization of Lie bialgebras is a functorial way to associate with a conilpotent<sup>1</sup> Lie bialgebra  $\mathfrak{g}$  a certain conilpotent bialgebra  $Q(\mathfrak{g})$ , called a *quantization* of  $\mathfrak{g}$ , in a such way that a quantization induces an equivalence between the corresponding categories (an inverse functor is called *dequantization*). After the original construction by Etingof and Kazhdan [EK] many others proofs appeared latter (for example [Tam]). In a preset paper we are going to give an another "motivic" construction of Lie bialgebras quantization.

1.2. **Technique.** There are three main ingredients in our construction of a quantization (resp. a dequantization) of Lie bialgebras (resp. bialgebras):

**The first one** is the equivalence between category of *factorization algebras* on  $Ran(\mathbb{A}^1)$  in the sense of Beilinson and Drinfeld and conilpotent Lie bialgebras, which can be though as a geometric instance of *Quillen duality*:

$$\Lambda: LieB_c \xrightarrow{\sim} FA_{un}(Ran(\mathbb{A})), \quad \mathfrak{g} \mapsto \Lambda(\mathfrak{g}) := \{\Lambda(\mathfrak{g})_I\}_{I \in Fin}.$$

Here  $\Lambda(\mathfrak{g})_I$  is a unipotent  $\mathcal{D}$ -module on  $\mathbb{A}^I$  and  $Fin$  is category of finite set with surjections as morphisms. This equivalence is achieved by using a *quiver description* of a category of unipotent  $\mathcal{D}$ -modules on  $\mathbb{A}^I$  following [Kho]. For example when  $|I| = 2$  out of Lie bialgebra  $\mathfrak{g}$  we can define the following double quiver:

$$A^-: (\mathfrak{g}_1 \otimes \mathfrak{g}_1 \oplus \mathfrak{g}_1 \otimes \mathfrak{g}_1)^{-\Sigma_2} \longleftrightarrow \mathfrak{g}_2: A^+,$$

where  $-\Sigma_2$  indicates skew invariants with respect to a symmetric group and operators are defined from a bracket and a cobracket:

$$A^+ := \delta \oplus \delta, \quad A^- := \frac{1}{2}[-, -] \oplus [-, -].$$

The quasi-inverse functor to  $\Lambda$  is given by computing *vanishing cycles* of  $\mathcal{D}$ -modules.

**The second one** is the equivalence between category of *factorizable sheaves* on  $Ran(\mathbb{A}^1)$  (topological incarnations of factorization algebras) and conilpotent bialgebras, which can be though as a topological instance of *Koszul duality*:

$$\Omega: AlgB_c \xrightarrow{\sim} FS_{un}(Ran(\mathbb{A})), \quad A \mapsto \Omega(A) := \{\Omega(A)_I\}_{I \in Fin}.$$

A perverse sheaf  $\Omega(A)_I$  on  $\mathbb{A}^I$  is defined using *Cousin resolutions*. Note that our construction is essentially parallel to the recent result of Kapranov and Schechtman

<sup>1</sup>For a bialgebra  $A$  with a reduced comultiplication  $\overline{\Delta}$  (resp. a Lie bialgebra  $\mathfrak{g}$  with a cobracket  $\delta$ ) a canonical filtration  $A_k := \{x \in A^+ \mid \overline{\Delta}^k(x) = 0\}$  (resp.  $\mathfrak{g}_k := \{x \in \mathfrak{g} \mid \delta(x) = 0\}$ ) is exhaustive.

[KS2], where instead of conilpotent bialgebras they have considered graded ones. As it was explained in *ibid* Cousin resolution approach is essentially equivalent to an elementary quiver description of categories of perverse sheaves on  $Sym^n(\mathbb{A}^1)^2$  for small values of  $n$ .

For an example when  $|I| = 2$  using the classical quiver description of perverse sheaves on  $Sym^2(\mathbb{A})$  [Bei] we have a perverse sheaf  $\mathcal{E}_2$  associated with a quiver:

$$\overline{\Delta}: A_2 \longleftarrow A_1 \otimes A_1: m,$$

where  $m$  is a multiplication in bialgebra  $A$ . Then we define  $\Omega(A)_I := p^! \mathcal{E}_2(A)$ , where  $p: \mathbb{A}^I \rightarrow Sym^{|I|}(\mathbb{A})$  is a natural projection.

Inverse functor to  $\Omega$  is defined by computing cohomology of perverse sheaves with support in real dimensional strata i.e. *hyperbolic stalks* in the sense of [KS1].

**The third one** is given by the *Riemann-Hilbert correspondence* which relates factorization algebras to factorizable sheaves:

$$RH: FA_{un}(Ran(\mathbb{A})) \longrightarrow FS_{un}(Ran(\mathbb{A})).$$

Then we define a *quantization functor*  $Q$  and a *dequantization functor*  $DQ$ :

$$Q: LieB_c \longleftarrow AlgB_c: DQ,$$

by the rule:

$$Q := \Omega^{-1} \circ RH \circ \Lambda, \quad DQ := \Lambda^{-1} \circ RH^{-1} \circ \Omega.$$

**1.3. Content.** In the first section, we recall some basic facts about conilpotent Lie bialgebras and collect all necessary information about unipotent  $\mathcal{D}$ -modules which we will need.

In the second section, we recall some basic facts about conilpotent bialgebras and collect all necessary facts about perverse sheaves on symmetric powers that we need. In our exposition about perverse sheaves, we will mostly follow the work [KS2]. We omit proofs and refer to *ibid* for details.

In the third subsection, we define a quantization and a dequantization functors and give some insights about how one can see associators in our setting.

**1.4. Acknowledgments.** We wish to thank Boris Feigin, Sergei Merkulov and Nikita Markarian for fruitful discussions.

Special thanks are due to Sasha Beilinson for his interest which has encouraged the author to write this note and Sergei Merkulov for his constant help and support during the preparation of this note.

This work was supported by the FNR project number PRIDE15/10949314/GSM and partially supported by the Simons Foundation

**1.5. Notation.** For a natural number  $n$  by  $[n]$  we denote a closed interval of natural numbers from 1 to  $n$ . Category of all finite sets and surjective maps between them will be denoted by  $Fin$ . By  $Vect$  we denote category of  $\mathbb{C}$ -vector spaces with the corresponding subcategory of finite-dimensional vector spaces  $Vect^f$ . Let  $I \in Fin$  be a finite set and  $\{V\}_{i \in I}$  be a collection of vector spaces indexed by  $I$  with the corresponding tensor product  $\otimes_I V_i$ . For a permutation  $\pi \in \Sigma_{|I|}$  we denote by  $\sigma_\pi$  a standard twisting operator  $\sigma_\pi: \otimes_I V_i \rightarrow \otimes_I V_i$ .

---

<sup>2</sup>A quotient of  $\mathbb{A}^n$  with respect to a natural symmetric group action

## 2. QUILLEN DUALITY

**2.1. Lie bialgebras.** Recall that a *Lie bialgebra* is a vector space  $\mathfrak{g}$  with skew symmetric operations  $[-, -]: \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$  (bracket) and  $\delta: \mathfrak{g} \rightarrow \mathfrak{g} \otimes \mathfrak{g}$  (cobracket) which satisfies Jacobi and coJacobi identity. Moreover a bracket and a cobracket are compatible in the following way:  $\delta$  is a 1-cocycle of  $\mathfrak{g}$  with values in  $\mathfrak{g} \otimes \mathfrak{g}$  [Dri]. We say that a Lie bialgebra  $\mathfrak{g}$  is *conilpotent* if a canonical filtration  $\mathfrak{g}_k := \{x \in \mathfrak{g} \mid \delta^k(x) = 0\}$  is exhaustive and for every  $k$  a component  $\mathfrak{g}_k$  is finite dimensional. An important property of this filtration is that a bracket and a cobracket respect this filtration:  $\delta(\mathfrak{g}_k) \subset \oplus_{i+j=k} \mathfrak{g}_i \otimes \mathfrak{g}_j$ ,  $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j-1}$ . For an element  $x \in \mathfrak{g}$  we will use notation  $\delta(x)_{ij}$  for a component of  $\delta(x)$ , which lies in the  $\mathfrak{g}_i \otimes \mathfrak{g}_j$ . The category of conilpotent Lie bialgebras will be denoted by  $LieB_c$ .

**2.2.  $\mathcal{D}$ -modules.** For every finite set  $I \in Fin$  we consider the corresponding complex affine space  $\mathbb{A}^I$ , equipped with a *diagonal stratification*  $\mathcal{S}_\emptyset = \{\Delta_{ij}\}$ , where  $\Delta_{ij} = z_i - z_j$ . The unique minimal closed stratum of  $\mathcal{S}_\emptyset$  will be denoted by  $\Delta$  and the unique maximal open stratum will be denoted by  $U^I$ . We say that a stratum  $S$  is adjusted to a stratum  $S'$  if  $S \subset \overline{S'}$ . Strata of dimension  $k$  can be identified with a set of equivalence classes of surjections  $[\pi]$ , where we say that two surjections  $\pi: I \rightarrow [k]$  and  $\pi': I \rightarrow [k]$  are equivalent if  $\pi' = \pi \circ \tau$ , where  $\tau$  is a bijection. Set of equivalence classes of surjections will be denoted by  $Q(I)$  and has a structure of a poset where we say that  $[\pi]: I \rightarrow T$  is less or equal to  $[\pi']: I \rightarrow T'$  if  $T$  is a quotient of  $T'$ . Set of equivalence classes of surjections of a cardinality  $k$  will be denoted by  $Q(I, k)$ . For an equivalence class  $[\pi]$  the corresponding stratum will be denoted by  $S_{[\pi]}$ . We will freely switch between these two notations. For  $[\pi]: I \rightarrow T \in Q(I)$  and  $i \neq j$  we associate another surjection  $[\pi_{ij}]: I \rightarrow T_{ij}$ , where  $T_{ij} = T - \{ij\} \sqcup *$ , by the following rule let  $\gamma_{ij}: T \rightarrow T_{ij}$  be a map sending  $i$  and  $j$  to  $*$  and every  $k$  not equal to  $i$  or  $j$  to  $k$ , then we define  $[\pi_{ij}] = \gamma_{ij} \circ [\pi]$ . In this way we get all codimension one adjunctions.

By  $\mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_\emptyset)$ , we denote a category of  $\mathcal{D}$ -modules on  $\mathbb{A}^I$  which are generated by delta functions with a support on diagonals. This is a subcategory of the category of holonomic  $\mathcal{D}$ -modules smooth along stratification  $\mathcal{S}_\emptyset$ . Objects of the category  $\mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_\emptyset)$  are called *unipotent  $\mathcal{D}$ -modules*.

For every finite set  $I$  with an affine space  $\mathbb{A}^I$  we can associate a quiver  $\Gamma_I$  whose vertices correspond to elements of  $\mathcal{S}_\emptyset$  and arrows correspond to an adjusted strata of codimension one:

$$[\alpha] \rightarrow [\beta] \Leftrightarrow S_{[\beta]} \subset \overline{S_{[\alpha]}}, \quad \text{codim}_{\overline{S_{[\alpha]}}} S_{[\beta]} = 1.$$

We will use a notation  $[\alpha_1] \circ \dots \circ [\alpha_n]$  for a minimal path in a quiver which passes through vertices  $[\alpha_i]$ , such that it starts at  $[\alpha_1]$  and ends at  $[\alpha_n]$ .

Following [Kho] with a quiver  $\Gamma_I$  we can associate a category  $Rep_{un}(\Gamma_I)$  of *double representations of quiver*  $\Gamma_I$  with certain relations. An object  $V$  of this category is by the definition consists of a vector space  $V_{[\beta]}$ , where  $[\beta]$  is a vertex of  $\Gamma_I$  and a pair of linear operators  $A_{[\lambda][\beta]}^-: V_{[\beta]} \leftarrow V_{[\lambda]}: A_{[\beta][\lambda]}^+$  for every arrow  $[\beta] \rightarrow [\lambda]$ . These linear operators must satisfy relations which we omit (See *ibid*). The main result of *ibid* (Theorem 2.1) gives an *explicit equivalence* between category  $Rep_{un}(\Gamma_I)$  and the category  $\mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_\emptyset)$ . For a path  $[\tau] = [\alpha_1] \circ \dots \circ [\alpha_n]$  we will use notation  $A_\tau^+$  for a composition  $A_{[\alpha_1],[\alpha_2]}^+ \circ \dots \circ A_{[\alpha_{n-1}],[\alpha_n]}^+$  of operators.

**Remark 2.2.1.** Consider a *specialization functor*:

$$Sp_{\Delta}: \mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_{\emptyset}) \longrightarrow \mathcal{M}_{un}^{dR}(T_{\Delta}\mathbb{A}^I, \mathcal{S}_{\emptyset})$$

along minimal diagonal  $\Delta \subset \mathbb{A}^I$ . Then we define an exact functor from a category unipotent  $\mathcal{D}$ -modules to a category finite dimensional vector spaces by the rule:

$$\omega_I^{dR}: \mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_{\emptyset}) \longrightarrow Vect^f, \quad \omega^{dR} := H_{dR}^0(T_{\Delta}\mathbb{A}^I, -)$$

Under the equivalence from *ibid* we have the identical isomorphism between a functor  $\omega^{dR}$  and a forgetful functor from  $Rep_{un}(\Gamma_I)$  to vector spaces. Thus a quiver description of category  $\mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_{\emptyset})$  can be considered as an instance of *Tannakian duality*. Note that by a construction  $\omega_I^{dR}$  carries a natural  $\mathbb{C}^{\times}$ -action i.e. grading. We will denote the first grading component by  $\Phi_I$ , which corresponds to the vector space associated with the smallest diagonal in the quiver description. It is easy to see that this construction extends to an exact functor:

$$\Phi_I: \mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_{\emptyset}) \longrightarrow Vect^f$$

For every surjection  $\pi: J \twoheadrightarrow I$  of finite sets we have the corresponding diagonal embedding  $\Delta^{(\pi)}: \mathbb{A}^I \hookrightarrow \mathbb{A}^J$ , which induces a pair of adjoint functors for  $\mathcal{D}$ -modules:

$$\Delta_*^{(\pi)}: \mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_{\emptyset}) \longleftarrow \mathcal{M}_{un}^{dR}(\mathbb{A}^J, \mathcal{S}_{\emptyset}): \Delta^{(\pi)!},$$

where a pushforward functor is exact and a  $!$ -pullback functor is left exact. Let us recall following [KV] a definition of a *quiver pushforward functor*  $\Delta_*^{(\pi)}$ :

For an element  $M \in Rep_{un}(\Gamma_I)$  we define a new quiver representation  $\Delta_*^{(\pi)}M \in Rep_{un}(\Gamma_J)$  by the following rule. For every stratum  $[\gamma]: I \twoheadrightarrow T$  of  $\mathbb{A}^I$  we define a stratum  $[\gamma_{\pi}]$  of  $\mathbb{A}^J$  by taking a composition with  $\pi$ . Thus we set  $\Delta_*^{(\pi)}M_{[\gamma_{\pi}]} := M_{[\gamma]}$ . For all other strata we define  $\Delta_*^{(\pi)}M$  to be zero. For a pair of non zero vector spaces we take corresponding operators from  $M$  and for all others we set operators to be zero. It is easy to see that  $\Delta_*^{(\pi)}M \in Rep_{un}(\Gamma_J)$  and the under equivalence between quiver representations and  $\mathcal{D}$ -modules quiver pushforward corresponds to  $\mathcal{D}$ -modules pushforward. Now we will define a *quiver  $!$ -pullback functor*:

**Definition 2.2.2.** For every  $\pi: J \twoheadrightarrow I$  we define quiver  $\Delta^{(\pi)!}M$  by the following rule. If  $\pi$  is a bijection we set  $(\Delta^{(\pi)!}M)_{[\alpha]} := M_{[\alpha_{\pi}]}$  with induced operators. In the case when  $\pi$  is not a bijection we set:

For  $[\alpha] \in \Gamma_I$  we define a vector space:

$$(\Delta^{(\pi)!}M)_{[\alpha]} := \bigcap_{\tau} \text{Ker}(A_{\tau}^+)$$

where intersection is taken over all *admissible paths*  $\tau := [\gamma] \circ [\alpha_j] \circ \dots \circ [\alpha_1] \circ [\alpha_{\pi}]$ , that are paths from a stratum  $[\gamma]$  of dimension  $|J| - \dim[\alpha]$  to a stratum  $[\alpha_{\pi}]$  in  $\Gamma_J$  of minimal length such that  $[\alpha_i]$  can not be factored into  $J \twoheadrightarrow I \twoheadrightarrow T$ . Operators  $A^+$  and  $A^-$  are induced from operators in  $M$ .


**Proposition 2.2.3.** The construction above defines functor  $\Delta^{(\pi)!}$  which is the right adjoint to a quiver pushforward

$$\Delta_*^{(\pi)}: Rep_{un}(\Gamma_I) \longleftarrow Rep_{un}(\Gamma_J): \Delta^{(\pi)!},$$

and thus under the equivalence between double quiver representations and unipotent  $\mathcal{D}$ -modules a quiver  $!$ -pullback functor corresponds to a  $!$ -pullback functor for  $\mathcal{D}$ -modules.

*Proof.* First we will show that operators in a quiver representation  $M$  induce will defined operators in  $\Delta^{(\pi)!}M$ . In the case when  $\pi$  is a bijection there is nothing to prove. Let us consider a case when  $\pi$  is not a bijection:

Without loss of a generality we can consider a case when  $|J| \geq 3$  (in the case when  $|J| < 3$  there is nothing to prove). Let  $[\pi] \rightarrow [\pi']$  be an adjusted pair of strata in  $\Gamma_I$ . Operator  $A_{[\pi],[\pi']}^+ : M_{[\pi']} \rightarrow M_{[\pi]}$  descends to an operator  $\Delta^{(\pi)!}M_{[\pi']} \rightarrow \Delta^{(\pi)!}M_{[\pi]}$ , because we have the condition one from [Kho]. More precisely let  $\tau = [\gamma] \circ \dots \circ [\alpha_1] \circ [\pi]$  be an arbitrary admissible path to the stratum  $[\pi]$  then we have an admissible path  $\tau_1$  (resp.  $\tau_2$ ) to the stratum  $[\pi']$  which passes through  $[\alpha_1]$  and equal to  $\tau$  everywhere except vertex  $[\beta]$  (resp.  $[\delta]$ ) (these strata have the same dimension as  $[\alpha_1]$ ). We have a picture like this:




Let  $x \in (\Delta^{(\pi)!}M)_{[\pi']}$ , then in particular  $x \in \text{Ker}A_{\tau_1}^+$  and  $x \in \text{Ker}A_{\tau_2}^+$ . Then by the first condition from *ibid* we have:

$$A_{[\alpha_1],[\pi]}^+ \circ A_{[\pi],[\pi']}^+ = -A_{[\alpha_1],[\beta]}^+ \circ A_{[\beta],[\pi']}^+ - A_{[\alpha_1],[\delta]}^+ \circ A_{[\delta],[\pi']}^+$$

and thus  $A_{\tau_1}^+ \circ A_{[\pi],[\pi']}^+(x) = -(A_{\tau_1}^+ + A_{\tau_2}^+)(x) = 0$ .

An operator  $A_{[\pi'],[\pi]}^- : M_{[\pi]} \rightarrow M_{[\pi']}$  descends to an operator  $\Delta^{(\pi)!}M_{[\pi]} \rightarrow \Delta^{(\pi)!}M_{[\pi']}$ , because we have the last condition from *ibid*. More precisely let  $\tau = [\gamma] \circ \dots \circ [\alpha_1][\pi']$  be an arbitrary admissible path to the stratum  $[\pi']$ . Then we have an adjusted strata  $[\pi_1] \rightarrow [\pi]$ . We have a picture like this:



Now due to the last condition from *ibid* we have the following:

$$A_{[\alpha_1],[\pi']}^+ \circ A_{[\pi],[\pi]}^- = -A_{[\alpha_1],[\pi_1]}^- \circ A_{[\pi],[\pi_1]}^+$$

Let  $x \in (\delta^{(\pi)!}M)_{[\pi]}$  if  $A_{[\alpha_1],[\pi_1]}^- \circ A_{[\pi],[\pi_1]}^+$  does not vanish we pick up an adjusted to  $[\alpha_1]$  vertex  $[\alpha_2]$  in a path  $\tau$  and consider another quadruple of strata  $[\alpha_1], [\pi_1], [\alpha_2]$  and  $[\pi_2]$ , where  $[\alpha_2] \rightarrow [\pi_2] \rightarrow [\pi_1] \rightarrow [\alpha_1]$  and repeat the procedure. At some point we will reach the strata of dimension  $|J| - \dim[\alpha]$  and thus obtain the admissible path  $\tau'$  to a stratum  $[\pi]$ . Thus an operator  $A_{\tau}^+$  must vanish on  $x$  and by the last condition from *ibid* we have  $A_{[\pi'],[\pi]}^-(x) \in \Delta^{(\pi)!}M_{[\pi']}$ . All conditions from *ibid* for  $A^+$  operators and  $A^-$  operators are satisfied because they descend from a double quiver

representation  $M$ . Moreover it is easy to see that  $\Delta^{(\pi)!}$  extends to a functor which is the right adjoint to a quiver direct image functor  $\Delta_*^{(\pi)}$  by the construction. Thus by the uniqueness of the right adjoint functor we obtain that under the equivalence with  $\mathcal{D}$ -modules a  $!$ -quiver pullback  $\Delta^{(\pi)!}$  corresponds to a  $!$ -pullback functor for  $\mathcal{D}$ -modules.  $\square$

With every surjection  $\pi: J \twoheadrightarrow I$  we can associate a space  $U^{(\pi)} := \{(x_j) \mid x_i \neq x_{i'} \text{ if } \pi(i) \neq \pi(i')\}$  with a corresponding open inclusion  $j^{(\pi)}: U^{(\pi)} \hookrightarrow \mathbb{A}^J$ . A space  $U^{(\pi)}$  inherits a diagonal stratification from  $\mathbb{A}^J$ : strata of  $U^{(\pi)}$  are given by equivalence classes of surjections  $[\gamma]: J \twoheadrightarrow T$  which are greater or equal to  $[\pi]: J \twoheadrightarrow I$ . We have an exact  $*$ -pullback functor along map  $j^{(\pi)}$  for  $\mathcal{D}$ -module:

$$j^{(\pi)*}: \mathcal{M}_{un}^{dR}(\mathbb{A}^J, \mathcal{S}_\emptyset) \longrightarrow \mathcal{M}_{un}^{dR}(U^{(\pi)}, \mathcal{S}_\emptyset),$$

Since  $U^{(\pi)}$  is a stratified subspace of  $\mathbb{A}^J$  methods of [Kho] can be used to study category of unipotent  $\mathcal{D}$ -modules on it. Let us recall following [KV] a definition of a *quiver  $*$ -pullback functor*  $j^{(\pi)*}$ :

For an element  $M \in \text{Rep}_{un}(\Gamma_J)$  we define a new quiver representation  $j^{(\pi)*}M \in \text{Rep}_{un}(\Gamma_{(\pi)})$  by the following rule. For every stratum  $[\gamma]: J \twoheadrightarrow T$  of  $U^{(\pi)}$  we define  $j^{(\pi)*}M_{[\gamma]} := M_{[\gamma]}$ . Operators are induced from operators in  $M$ .

Let  $\pi: J \twoheadrightarrow I$  be a surjection. For unipotent  $\mathcal{D}$ -modules we have a standard operation of the exterior tensor product, which is a functor

$$\boxtimes_I: \times_I \mathcal{M}_{un}^{dR}(\mathbb{A}^{J_i}, \mathcal{S}_\emptyset) \longrightarrow \mathcal{M}_{un}^{dR}(\mathbb{A}^J, \mathcal{S}_\emptyset), \quad J_i := \pi^{-1}(i).$$

Following [KS3] we define a *quiver exterior tensor product*:

Let  $[\gamma]: J \twoheadrightarrow T$  be a stratum, then if  $\pi$  can be factored as  $\pi = \gamma \circ \beta$ , where  $\beta: T \twoheadrightarrow I$  we define

$$(\boxtimes_I M_i)_{[\gamma]} := \otimes_I M_{[\gamma_i]},$$

where  $\gamma_i: J_i \twoheadrightarrow T_i$ , and  $T_i = \beta^{-1}(i)$ . Otherwise we define  $(\boxtimes_I M_i)_{[\gamma]}$  to be zero. Operators in a quiver representation are defined by the following rule: if  $[\gamma]$  and  $[\gamma']$  is a pair of adjusted strata which satisfy a property above we define  $A^+$  operators and  $A^-$  operators as a tensor product of appropriate operators acting between vector spaces assigned to a strata  $[\gamma_i]$  and  $[\gamma'_i]$ , otherwise operators are defined to be zero. It is easy to see that  $\boxtimes_I M_i$  is an element of  $\text{Rep}_{un}(\Gamma_J)$  and moreover  $\boxtimes_I$  extends to a functor and under the equivalence between quiver representations and  $\mathcal{D}$ -modules corresponds to  $\mathcal{D}$ -module exterior tensor product.

**2.3. Factorization algebra  $\Lambda(\mathfrak{g})$ .** In this subsection out of conilpotent Lie bialgebra we built a certain factorization algebra.

**Proposition 2.3.1.** With every conilpotent Lie bialgebra  $\mathfrak{g}$  we associate unipotent  $\mathcal{D}$ -module  $\Lambda(\mathfrak{g})_I$  on  $\mathbb{A}^I$ .

*Proof.* We will use quiver description of a category  $\mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_\emptyset)$  from [Kho]:

Let  $[\pi] \in Q(I, k)$  be a  $k$ -dimensional strata, then with an element of the equivalence class  $\pi: I \twoheadrightarrow [k] \in [\pi]$  we can associate a vector space  $\mathfrak{g}_\pi = \mathfrak{g}_{i_1} \otimes \cdots \otimes \mathfrak{g}_{i_k}$ , where  $i_j = |\pi^{-1}(j)|$ . Hence for a stratum  $[\pi]$  we define the corresponding vector space by the rule

$$\mathfrak{g}_{[\pi]} := \bigoplus_{\pi: I \twoheadrightarrow [k] \in [\pi]} \mathfrak{g}_\pi^{-\Sigma_k}.$$

For an adjusted pair of strata  $[\pi] \twoheadrightarrow [\pi_{pq}]$  we choose a representative  $I \twoheadrightarrow [k] \in [\pi]$ , such that  $p$  will be the first coordinate and  $q$  will be the second one. Then we can define a map  $A_{\pi_{pq}, \pi}^-: \mathfrak{g}_\pi \longrightarrow \mathfrak{g}_{\pi_{pq}}$  by the rule  $([-, -] \otimes id \otimes \cdots \otimes id)$ . Thus we define a map  $\mathfrak{g}_\pi \longrightarrow \mathfrak{g}_{[\pi_{pq}]}$  as:

$$A_{[\pi_{pq}], \pi}^- = \bigoplus_{\sigma \in \Sigma_{k-1}} \tau_\sigma \circ A_{\pi_{pq}, \pi}^-.$$

And finally we define a map  $A_{[\pi_{pq}], T}^-: \mathfrak{g}_{[\pi]} \longrightarrow \mathfrak{g}_{[\pi_{pq}]}$  by the rule:

$$A_{[\pi_{pq}], [\pi]}^- := \frac{1}{k!} \bigoplus_{\sigma \in \Sigma_k} A_{\pi_{pq}, \pi}^- \circ \tau_\sigma.$$

Analogically we define a map  $A_{[\pi], [\pi_{pq}]}^+: \mathfrak{g}_{[\pi_{pq}]} \longrightarrow \mathfrak{g}_{[\pi]}$  but instead of a bracket we use a cobracket. Thus we set  $A_{\pi, \pi_{pq}}^+ := \delta_{i_1, i_2} \otimes id \otimes \cdots \otimes id$  and then:

$$A_{[\pi], \pi_{pq}}^+ := \bigoplus_{\sigma \in \Sigma_k} \tau_\sigma \circ A_{\pi, \pi_{pq}}^+,$$

and thus:

$$A_{[\pi], [\pi_{pq}]}^+ := \frac{1}{(k-1)!} \bigoplus_{\sigma \in \Sigma_{k-1}} A_{[\pi], \pi_{pq}}^+ \circ \tau_\sigma.$$

We need to check conditions from *ibid*:

The first two identities follow from Jacobi and coJacobi identities for a Lie bialgebra  $\mathfrak{g}$ . Let us consider a case of a map  $A_{[\pi_{ij}], \pi}^-$ . Choice a representative  $\pi: I \twoheadrightarrow [k]$  such that  $i, j, k \in [k]$  will be a first three coordinates. Then  $A_{[\pi_{jk}], \pi}^-$  acts on  $\pi$  component as  $A_{[\pi_{ij}], \pi}^- \circ \tau_{(ijk)}$  and  $A_{[\pi_{ki}], \pi}^-$  acts on  $\pi$  component as  $A_{[\pi_{ij}], \pi}^- \circ \tau_{(ijk)^2}$ . Hence from Jacobi identity we obtain that:

$$A_{[\pi_{jk}], [\pi_{jk}]}^- \circ A_{[\pi_{jk}], [\pi]}^- + A_{[\pi_{ki}], [\pi_{ki}]}^- \circ A_{[\pi_{ki}], [\pi]}^- + A_{[\pi_{ij}], [\pi_{ij}]}^- \circ A_{[\pi_{ij}], [\pi]}^- = 0.$$

Case of  $+$ -map can be treated analogically and follows from coJacobi identity for a Lie cobracket.

The third condition from *ibid* follows from 1-cocycle property for a Lie bialgebra. Choice a representative  $\pi: I \twoheadrightarrow [k]$  such that  $i, j, k \in [k]$  will be first three coordinates. Then  $A_{\pi_{jk}, \pi_{ij}}^+ \circ A_{\pi_{ij}, \pi_{ij}}^-$  acts between  $\pi_{ij}$  and  $\pi_{jk}$  component as  $\tau_{(ij)} \circ A_{\pi_{ij}, \pi_{ij}}^+ \circ A_{\pi_{ij}, \pi_{ij}}^-$  and  $A_{\pi, \pi_{jk}}^- \circ A_{\pi, \pi_{ij}}^+$  acts as  $A_{\pi, \pi_{ij}}^- \circ \tau_{(ijk)} \circ A_{\pi, \pi_{ij}}^+$ . Thus by 1-cocycle property of a Lie bialgebra we obtain that:

$$A_{[\pi_{jk}], [\pi_{ij}]}^+ \circ A_{[\pi_{ij}], [\pi_{ij}]}^- + A_{[\pi], [\pi_{jk}]}^- \circ A_{[\pi], [\pi_{ij}]}^+ = 0.$$

It remains to show that monodromy operators are nilpotent *ibid*. This obviously follows from a Lie property of a canonical filtration.  $\square$

**Proposition 2.3.2.** For every  $\pi: J \twoheadrightarrow I$  conilpotent filtration induces a natural isomorphism:

$$\theta(\pi): \Lambda(\mathfrak{g})_I \xrightarrow{\sim} \Delta^{(\pi)!} \Lambda(\mathfrak{g})_J,$$

compatible with compositions of surjections.

*Proof.* We will use a description of quiver  $!$ -pullback from Proposition 2.2.3. Let us first consider a case when  $\pi: I \xrightarrow{\sim} I$  is a bijection. For every stratum  $[\gamma]: I \twoheadrightarrow T$  we have a new stratum  $[\gamma_\pi]$ . Then we define an operator:

$$\rho: \mathfrak{g}_{[\gamma]} \longrightarrow \mathfrak{g}_{[\gamma_\pi]},$$

for a representative  $\gamma \in [\gamma]$  by taking a signed twist morphism  $\tau_\pi$  associated with a permutation  $\pi$  :

$$\tau_\pi: \mathfrak{g}_{i_1} \otimes \cdots \otimes \mathfrak{g}_{i_k} \longrightarrow \mathfrak{g}_{i_1} \otimes \cdots \otimes \mathfrak{g}_{i_k}.$$

Due to skew symmetry of a bracket and a cobracket isomorphisms  $\rho$  extend to an isomorphism of quivers  $\Lambda(\mathfrak{g})_I \xrightarrow{\sim} \Delta^{(\pi)!} \Lambda(\mathfrak{g})_I$ . It is easy to see that defined isomorphisms are compatible with compositions of bijections, i.e.  $\mathcal{D}$ -module  $\Lambda(\mathfrak{g})_I$  is equivariant.

Now we need to consider a case of a non bijective surjection  $\pi: J \twoheadrightarrow I$ . For a given stratum  $[\alpha]$  in  $\mathbb{A}^I$  and any stratum  $[\gamma]$  of dimension  $|J| - \dim[\alpha]$  in  $\mathbb{A}^J$  choose a representatives  $\alpha$  and  $\gamma$ . Denote by  $\tau$  a sequence of elementary surjections from  $\gamma$  to  $\alpha_\pi$ . Denote by  $A_\tau^+$  a component of an operator  $A_{[\tau]}^+$  which acts between  $\mathfrak{g}_{\alpha_\pi}$  and  $\mathfrak{g}_\gamma$ . Operator  $A_\tau^+$  acts as a composition of appropriate components of  $\delta$  and signed twisting maps. We have a canonical colimit map  $can: \mathfrak{g}_\alpha \longrightarrow \mathfrak{g}_{[\alpha]}$ . Then it is easy to see that we have an isomorphism  $can: \text{Ker} A_\tau^+ \xrightarrow{\sim} \text{Ker} A_{[\tau]}^+$ . Since we have proved that  $\mathcal{D}$ -module  $\Lambda(\mathfrak{g})_J$  is  $\Sigma_{|J|}$ -equivariant it is enough to consider equivalence classes of admissible paths with respect to a natural symmetric group action on  $\mathbb{A}^J$ . Thus by a skew symmetry of a cobracket and remarks above for a stratum  $[\alpha]$  in  $\mathbb{A}^I$  we obtain:

$$\bigcap_{[\tau]/\Sigma_{|J|}} \text{Ker} A_{[\tau]}^+ \xrightarrow{\sim} \mathfrak{g}_{[\alpha]}.$$

It is easy to see that constructed isomorphisms extends to an isomorphism of quivers which are compatible with compositions of surjections.  $\square$

**Remark 2.3.3.** Following Beilinson and Drinfeld [BD] it is natural to call  $\Lambda(\mathfrak{g}) := \{\Lambda(\mathfrak{g})_I, \theta(\pi)\}_{Fin^\circ}$  an *unipotent left  $\mathcal{D}$ -module on Ran prestack* of  $\mathbb{A}^1$ . These objects with obvious morphisms form a category which will be denoted by  $\mathcal{M}_{un}^{dR}(\text{Ran}(\mathbb{A}^1))$ .

**Proposition 2.3.4.** Unipotent  $\mathcal{D}$ -module  $\Lambda(\mathfrak{g})$  on  $\text{Ran}(\mathbb{A})$  has the following additional structure:

- For every surjection  $\pi: J \twoheadrightarrow I$ , we have a natural isomorphism:

$$c_\pi: j^{(\pi)*} \Lambda(\mathfrak{g})_J \xrightarrow{\sim} j^{(\pi)*} \boxtimes_I \Lambda(\mathfrak{g})_{J_i}, \quad J_i := \pi^{-1}(i).$$

- Compatibilities between  $c$ 's and compositions of surjections of finite sets.
- Compatibilities between maps  $c$  and  $\theta$ .

*Proof.* Let  $\pi: J \twoheadrightarrow I$  be a surjection and let  $[\gamma]: J \twoheadrightarrow T$  be a stratum in  $\mathbb{A}^J$ , then if  $\pi$  can be factored as  $\pi = \beta \circ \gamma$ , where  $\beta: T \twoheadrightarrow I$  we have

$$(\boxtimes_I \Lambda(\mathfrak{g})_{J_i})_{[\gamma]} := \otimes_I \mathfrak{g}_{[\gamma_i]} = \otimes_I \left( \bigoplus_{J_i \rightarrow T_i} \otimes_{t \in T_i} \mathfrak{g}_{(J_i)_t} \right)^{-\Sigma},$$

where  $\gamma_i: J_i \twoheadrightarrow T_i$ , and  $T_i = \beta^{-1}(i)$ . On another hand we have:

$$(\Lambda(\mathfrak{g})_J)_{[\gamma]} = \left( \bigoplus_{\gamma: J \rightarrow T \in [\gamma]} \otimes_T \mathfrak{g}_{J_t} \right)^{-\Sigma_{|T|}}.$$

Note that a diagram which underlies  $(\boxtimes_I \Lambda(\mathfrak{g})_{J_i})_{[\gamma]}$  is naturally a subdiagram of a diagram which underlies  $(\Lambda(\mathfrak{g})_J)_{[\gamma]}$  thus we obtain a canonical map:

$$can: (\Lambda(\mathfrak{g})_J)_{[\gamma]} \xrightarrow{\sim} (\boxtimes_I \Lambda(\mathfrak{g})_{J_i})_{[\gamma]},$$



which is an isomorphism obviously. According to the definition of a quiver pullback  $j^{(\pi)}$  it is easy to see that canonical maps define an isomorphism of quivers:

$$c_\pi: j^{(\pi)*} \Lambda(\mathfrak{g})_J \xrightarrow{\sim} j^{(\pi)*} \boxtimes_I \Lambda(\mathfrak{g})_{J_i},$$

which satisfies all properties listed above.  $\square$

**Remark 2.3.5.** Following Beilinson and Drinfeld [BD] we say that an object  $\mathcal{N}$  in  $\mathcal{M}_{un}^{dR}(\text{Ran}(\mathbb{A}), \mathcal{S}_\emptyset)$  is an *unipotent factorization algebra* on  $\mathbb{A}^1$  if conditions above are satisfied. Factorization algebras naturally organize into a category denoted by  $FA_{un}(\text{Ran}(\mathbb{A}^1))$ . It is obvious that the construction defined above is functorial i.e. we have a functor:

$$\Lambda: LieB_c \longrightarrow FA_{un}(\text{Ran}(\mathbb{A})), \quad \Lambda: \mathfrak{g} \longmapsto \Lambda(\mathfrak{g}).$$

**2.4. Vanishing cycles.** In this subsection we will prove that functor  $\Lambda$  induces equivalence between conilpotent Lie bialgebras and unipotent factorization algebras on  $\mathbb{A}$ :

**Proposition 2.4.1.** We have the following functor:

$$(1) \quad \Phi_{Ran}: FA_{un}(\text{Ran}(\mathbb{A})) \longrightarrow LieB_c, \quad \Phi_{Ran} := \lim_{Fin} \Phi_I.$$

Such that  $\Phi_{Ran}$  is a quasi inverse a functor to  $\Lambda$ .

*Proof.* Let  $\mathcal{K} \in FA_{un}(\text{Ran}(\mathbb{A}))$  then for every  $I \in Fin$  we define a vector space  $\Phi_{Ran}(\mathcal{K})_{|I|} := \Phi_I(\mathcal{K}_I)^{\Sigma_{|I|}}$ . Note that for every  $\pi: J \rightarrow I$  we have an inclusion  $\Delta_*^{(\pi)} \mathcal{K}_I \hookrightarrow \mathcal{K}_J$ , which comes as an adjoint map to an isomorphism from Proposition 2.3.2. These maps descends to an injection:  $\Phi_{Ran}(\mathcal{K})_{|I|} \hookrightarrow \Phi_{Ran}(\mathcal{K})_{|J|}$ . Thus we have:

$$\Phi_{Ran}(\mathcal{K})_1 \subset \cdots \subset \Phi_{Ran}(\mathcal{K})_n \subset \cdots \subset \Phi_{Ran}(\mathcal{K}).$$

Note that  $A^+$  operators together with factorization isomorphisms from Proposition 2.3.4 define the following map:

$$\delta_{|I'|, |I''|}: \Phi_{Ran}(\mathcal{K})_{|I|} \longrightarrow \Phi_{Ran}(\mathcal{K})_{|I'|} \otimes \Phi_{Ran}(\mathcal{K})_{|I''|}, \quad |I'| + |I''| = |I|.$$

From the quiver description of a category of unipotent  $\mathcal{D}$ -modules, we obtain that  $\delta$  satisfies coJacobi identity. Also from the Koszul sign rule for the exterior tensor product  $\boxtimes$  of  $\mathcal{D}$ -modules we obtain that  $\delta$  is skew-symmetric.

Analogically using  $A^-$  operators and factorization isomorphisms we define the following map:

$$[-, -]_{|I'|, |I''|}: \Phi_{Ran}(\mathcal{K})_{|I'|} \otimes \Phi_{Ran}(\mathcal{K})_{|I''|} \longrightarrow \Phi_{Ran}(\mathcal{K})_{|I|}, \quad |I'| + |I''| = |I|.$$

Which is skew-symmetric and satisfies Jacobi identity. Operators  $\delta$  and  $[-, -]$  are compatible in the sense of Drinfeld ( $\delta$  is 1-cocycle for the bracket  $[-, -]$ ) because of the last condition of quiver description of category of unipotent  $\mathcal{D}$ -modules.

Thus we have defined a structure of conilpotent Lie bialgebra on  $\Phi_{Ran}(\mathcal{K})$ . It is obvious that  $\Phi_{Ran}$  is a quasi-inverse functor to  $\Lambda$ .  $\square$

**Remark 2.4.2.** Analogically to Remark (2.2.1) we can define functor  $\omega^{dR}$  on a category of  $\mathcal{D}$ -modules on a Ran prestack:

$$(2) \quad \omega_{Ran}^{dR}: \mathcal{M}_{un}^{dR}(\text{Ran}(\mathbb{A}), \mathcal{S}_\emptyset) \longrightarrow Vect, \quad \omega_{Ran}^{dR} := \lim_{Fin} \omega_I^{dR}.$$

Denote by  $AlgCB_c$  is a category of *conilpotent coPoisson bialgebras*. We have a primitive elements functor  $Prim: AlgCB_c \rightarrow LieB_c$ , defined by taking the underlying Lie algebra i.e.  $Prim(A) := \{x \in A^+ \mid \overline{\Delta}(x) = 0\}$ . One can show that a functor (2) makes the following diagram into the commutative:

$$\begin{array}{ccc}
 FA_{un}(Ran(\mathbb{A})) & \xrightarrow{\omega_{Ran}^{dR}} & AlgCB_c \\
 & \searrow \Phi_{Ran} & \downarrow Prim \\
 & & LieB_c
 \end{array}$$

$\sim$  (under  $\Phi_{Ran}$ )       $\sim$  (under  $Prim$ )

Note that a primitive elements functor  $Prim$  defines an equivalences of categories. This is corollary from the Milnor-Moore equivalence [Dri].

### 3. KOSZUL DUALITY

**3.1. Hopf algebras.** For a bialgebra  $A$  we denote by  $\Delta: A \rightarrow A \otimes A$  the corresponding comultiplication with the corresponding reduced comultiplication  $\overline{\Delta}(x) = \Delta(x) - x \otimes 1 - 1 \otimes x$  and by  $m: A \otimes A \rightarrow A$  we denote the corresponding multiplication and  $\varepsilon: A \rightarrow \mathbb{k}$  the counit. By  $A^+ := \text{Ker}(\varepsilon)$  we denote an augmentation ideal. An operator  $\overline{\Delta}$  defines a coalgebra structure on  $A^+$ . Let  $A$  be a *conilpotent bialgebra*, that is a bialgebra such that a canonical filtration  $A_k := \{x \in A^+ \mid \overline{\Delta}^k(x) = 0\}$  is exhaustive and  $\dim A_k < \infty$ . Note that  $m$  and  $\overline{\Delta}$  respect a canonical filtration i.e.  $m(A_i, A_j) \subset A_{i+j}$  and  $\overline{\Delta}(A_k) \subset \oplus_{i+j=k} A_i \otimes A_j$  and by  $\overline{\Delta}_{ij}$  we denote a component which lies in  $A_i \otimes A_j$ . Conilpotent bialgebras form a category  $AlgB_c$  in the obvious way which is equivalent to the category of conilpotent Hopf algebras  $AlgH_c$ . Further we will not distinguish between conilpotent bialgebras and conilpotent Hopf algebras.

**3.2. Perverse sheaves.** With every  $I \in Fin$  we associate complex affine space  $\mathbb{A}^I$  (in analytic topology) equipped with a diagonal stratification  $\mathcal{S}_\emptyset$ . Let  $\mathcal{M}_{un}^B(\mathbb{A}^I, \mathcal{S}_\emptyset)$  be a category of unipotent perverse sheaves on  $\mathbb{A}^I$  smooth with respect to a diagonal stratification.

For every  $I$  we will also consider  $\mathbb{A}_{\mathbb{R}}^I$  with a real diagonal stratification denoted by  $\mathcal{S}_{\emptyset, \mathbb{R}}$ . Real strata of dimension  $|T|$  correspond to a surjections  $\rho: I \rightarrow T$ . We denote by  $i: \mathbb{A}_{\mathbb{R}}^I \rightarrow \mathbb{A}^I$  embedding of  $\mathbb{A}_{\mathbb{R}}^I$  as  $x$ -axes. A category of constructible sheaves smooth with respect to  $\mathcal{S}_{\emptyset, \mathbb{R}}$  will be denoted by  $Const(\mathbb{A}_{\mathbb{R}}^I, \mathcal{S}_{\emptyset, \mathbb{R}})$ . In [KS1] it was proved that a functor  $i^!: \mathcal{M}_{un}^B(\mathbb{A}^I, \mathcal{S}_\emptyset) \rightarrow Const(\mathbb{A}_{\mathbb{R}}^I, \mathcal{S}_{\emptyset, \mathbb{R}})$  is an exact functor. Moreover using this functor *explicit description* of a category  $\mathcal{M}_{un}^B(\mathbb{A}^I, \mathcal{S}_\emptyset)$  was given in *ibid*. A category of perverse sheaves was identified with a category of double representations with relations of a certain quiver  $\Gamma_{I, \mathbb{R}}$  (which is defined as a quiver  $\Gamma_I$  *mutatis mutandis* with respect to a real diagonal stratification of  $\mathbb{A}_{\mathbb{R}}^I$ ).

**Remark 3.2.1.** Denote by  $\Delta_{\mathbb{R}}$  the unique closed minimal stratum of  $\mathcal{S}_{\emptyset, \mathbb{R}}$ . For every  $I$  we define the following functor:

$$\omega_I^B: \mathcal{M}_{un}^B(\mathbb{A}^I, \mathcal{S}_\emptyset) \rightarrow Vect^f, \quad \omega_I^B := \Gamma(\Delta_{\mathbb{R}}, i^!-).$$

The equivalence from *ibid* can be though as an instance of *Tannakian duality* with respect to a functor  $\omega_I^B$ .

By  $Sym^{|I|}(\mathbb{A})$  we denote a geometric quotient of  $\mathbb{A}^I$  for the natural symmetric group action. Geometric quotient  $Sym^{|I|}(\mathbb{A})$  can be naturally identified with a space  $Div^{|I|}(\mathbb{A})$  of *effective Cartier divisors* on  $\mathbb{A}$  of degree  $|I|$ . The diagonal stratification on  $\mathbb{A}^I$  induces a stratification on  $Sym^{|I|}(\mathbb{A})$ , which by the abuse of notation will be denoted by the same symbol  $\mathcal{S}_\emptyset$ . From point of view of Cartier divisors elements in  $\mathcal{S}_\emptyset$  can be identified with *partitions*. By  $\mathcal{M}^B(Sym^{|I|}(\mathbb{A}), \mathcal{S}_\emptyset)$  we denote a category of perverse sheaves on  $Sym^{|I|}(\mathbb{A})$  smooth with respect to a diagonal stratification.

We also consider a space  $Sym^n(\mathbb{A}_{\mathbb{R}}) := \mathbb{A}_{\mathbb{R}}^n / \Sigma_n$ . This space also has a real diagonal stratification denoted by  $\mathcal{S}_{\emptyset, \mathbb{R}}$ . Elements of this stratification can be identified with *ordered partitions*. Following [KS2] a poset of ordered partitions, equipped with a partial order of reverse inclusion of subsets will be denoted by  $\mathbf{2}^{n-1}$ . The corresponding category of combinatorial (constructible) sheaves on  $Sym^n(\mathbb{A}_{\mathbb{R}})$  will be denoted by  $Const(Sym^n(\mathbb{A}_{\mathbb{R}}), \mathcal{S}_{\emptyset, \mathbb{R}})$ . By  $Rep(\mathbf{2}^{n-1})$  we will denote a category of functors from  $\mathbf{2}^{n-1}$  to the category  $Vect^f$ . Note [KS1] that a category  $Const(Sym^n(\mathbb{A}_{\mathbb{R}}))$  is equivalent to a category  $Rep(\mathbf{2}^{n-1})$ .

For every ordered partition  $(n_1, \dots, n_k)$  of  $n$  we have the corresponding embedding of strata:  $\Delta^{(n_1, \dots, n_k)}: Sym^k(\mathbb{A}_{\mathbb{R}}) \rightarrow Sym^n(\mathbb{A}_{\mathbb{R}})$  and the corresponding pair of adjoint functors:

$$(3) \quad \Delta_*^{(n_1, \dots, n_k)}: Const(Sym^k(\mathbb{A}_{\mathbb{R}}), \mathcal{S}_{\emptyset, \mathbb{R}}) \longleftrightarrow Const(Sym^n(\mathbb{A}_{\mathbb{R}}), \mathcal{S}_{\emptyset, \mathbb{R}}): \Delta^{(n_1, \dots, n_k)!}$$

between constructible sheaves. We have the corresponding functors between combinatorial data:

$$(4) \quad \Delta_*^{(n_1, \dots, n_k)}: Rep(\mathbf{2}^{k-1}) \longleftrightarrow Rep(\mathbf{2}^{n-1}): \Delta^{(n_1, \dots, n_k)!}$$

Note that we have the following commutative square:

$$(5) \quad \begin{array}{ccc} \mathbb{A}_{\mathbb{R}}^I & \xrightarrow{i} & \mathbb{A}^I \\ \downarrow p_{\mathbb{R}, I} & & \downarrow p_I \\ Sym^{|I|}(\mathbb{A}_{\mathbb{R}}) & \xrightarrow{i_\sigma} & Sym^{|I|}(\mathbb{A}) \end{array}$$

Where  $p_I$  and  $p_{\mathbb{R}, I}$  are projection maps and  $i_\sigma$  is the induced map between quotient. Note that map  $p_I$  is finite flat and surjective.

Let  $\pi: J \rightarrow I$  be a surjection with a corresponding ordered partition  $|\pi| := (|\pi^{-1}(i)|)_I$ . Then we have the following commutative square:

$$(6) \quad \begin{array}{ccc} \mathbb{A}_{\mathbb{R}}^I & \xrightarrow{\Delta^{(\pi)}} & \mathbb{A}_{\mathbb{R}}^J \\ \downarrow p_I & & \downarrow p_J \\ Sym^{|I|}(\mathbb{A}_{\mathbb{R}}) & \xrightarrow{\Delta^{|\pi|}} & Sym^{|J|}(\mathbb{A}_{\mathbb{R}}) \end{array}$$

It is easy to see that this square is fibered.

**3.3. Factorizable sheaf  $\Omega(A)$ .** In this subsection out of a conilpotent bialgebra we built a certain factorizable sheaf.

**Proposition 3.3.1.** With every conilpotent bialgebra  $A$  and for every  $n$  we associate perverse sheaf  $\mathcal{E}(A)_n$  on  $Sym^n(\mathbb{A})$ .

*Proof.* We essentially follow a construction from [KS2] which goes *mutatis mutandis* in a conilpotent case. This construction consists of the following steps:

We have a natural map  $Sym^n(\mathbb{A}) \rightarrow Sym^n(\mathbb{A}_{\mathbb{R}})$ , which sends a complex tuple to its imaginary part  $\mathcal{J}: Sym^n(\mathbb{A}) \rightarrow Sym^n(\mathbb{A}_{\mathbb{R}})$ . Following *ibid* we will call a preimage of a real diagonal stratification an *imaginary stratification*. We denote by  $S_{\alpha}^{\mathcal{J}} := \mathcal{J}^{-1}(K_{\alpha})$ , and  $j_{\alpha}: S_{\alpha}^{\mathcal{J}} \hookrightarrow Sym^n(\mathbb{A})$ . Then for every  $n$  and  $\alpha \in \mathbf{2}^{n-1}$  we have a sheaf  $\tilde{\mathcal{E}}^{\alpha}(A)$  on  $S_{\alpha}^{\mathcal{J}}$ . Stalk of this sheaf at the point  $x = \Sigma \lambda_i x_i$  is identified with  $A_{\lambda_1} \otimes \cdots \otimes A_{\lambda_k}$  and generalization maps are given by the appropriate component of  $\tilde{\Delta}$ . We omit proof and refer reader to *ibid*. We call  $\mathcal{E}^{\alpha}(A) := Rj_{\alpha*} \tilde{\mathcal{E}}^{\alpha}(A)$  the  $\alpha$ th *Cousin sheaf* of  $A$ .

For  $\alpha, \beta \in \mathbf{2}^{n-1}$  we define the morphism of sheaves on  $S_{\beta}^{\mathcal{J}}$ :

$$\delta'_{\alpha, \beta}: j_{\beta}^* \mathcal{E}^{\alpha}(A) = j_{\beta}^* Rj_{\alpha*} \tilde{\mathcal{E}}^{\alpha}(A) \rightarrow \tilde{\mathcal{E}}^{\beta}(A),$$

using multiplication in a bialgebra  $A$ , we again refer to *ibid* for details. By an adjunction this map gives a morphism of sheaves on  $Sym^n(\mathbb{A})$ :

$$\delta_{\alpha, \beta}: \mathcal{E}^{\alpha}(A) \rightarrow \mathcal{E}^{\beta}(A).$$

Thus have the following complex of sheaves:

$$\mathcal{E}_n^*(A) = \{\mathcal{E}^{1^n}(A) \rightarrow \bigoplus_{l(\alpha)=n-1} \mathcal{E}^{\alpha}(A) \rightarrow \cdots \rightarrow \mathcal{E}^n(A)\},$$

where  $l(\alpha)$  is a length of an ordered partition.

The last step is to show that  $\mathcal{E}_n^*(A)$  is a perverse sheaf on  $Sym^n(\mathbb{A})$ , constructible with respect to a diagonal stratification  $\mathcal{S}_{diag}$ . This is Proposition 4.2.16 from *ibid*.  $\square$

For every  $I$  we put  $\Omega(A)_I := p_I^! \mathcal{E}(A)_{|I|}$ . Since  $p$  is a finite flat and surjective map a sheaf  $\Omega(A)_I$  will be perverse. Note that because  $p^!$  is a submersion we have  $p^! \cong p^*$ .

**Proposition 3.3.2.** For every  $\pi: J \twoheadrightarrow I$  conilpotent filtration induces natural isomorphism:

$$\varphi(\pi): \Omega(A)_I \xrightarrow{\sim} \Delta^{(\pi)!} \Omega(A)_J,$$

compatible with compositions of surjections.

*Proof.* By the definition of a quiver  $!$ -pullback (4) and the definition of a conilpotent filtration for every surjection  $\pi: J \twoheadrightarrow I$  we have an isomorphism:

$$i_{\sigma}^! \mathcal{E}(A)_{|I|} \xrightarrow{\sim} \Delta^{|\pi|!} i_{\sigma}^! \mathcal{E}(A)_{|J|}$$

Moreover this map lifts to an isomorphism:

$$(7) \quad \phi(\pi): i^! \Omega(A)_{|I|} \xrightarrow{\sim} \Delta^{(\pi)!} i^! \Omega(A)_{|J|},$$

compatible with compositions of surjections. This follows from the standard diagram chasing:

$$(8) \quad \Delta^{(\pi)!} i^! \Omega(A)_{|J|} := \Delta^{(\pi)!} i^! p_J^! \mathcal{E}(A)_{|J|} \cong \Delta^{(\pi)!} p_J^! i_\sigma^! \mathcal{E}(A), \quad (\text{square (5)}),$$

$$(9) \quad \Delta^{(\pi)!} p_J^! i_\sigma^! \mathcal{E}(A) \cong p_I^! \Delta^{|\pi|!} i_\sigma^! \mathcal{E}(A), \quad (\text{square (6)}).$$

For every  $\pi: J \rightarrow I$  we also have the following morphism:

$$(10) \quad \Delta_*^{|\pi|} i_\sigma^* \mathcal{E}(A)_{|I|} \longrightarrow i_\sigma^* \mathcal{E}(A)_{|J|}.$$

This morphism follows from the fact that a multiplication respects the filtration on  $A$ . More precisely a functor  $i_\sigma^*$  sends perverse sheaf  $\mathcal{E}(A)$  to a complex of sheaves (in a  $t$ -structure dual to the standard one). The later complex is equivalent to a datum of cosheaf on  $Sym^n(\mathbb{A}_{\mathbb{R}})$  associated with an algebra underlying  $A$  (See [KS2]). Then map (10) lifts to a morphism:

$$\Delta_*^{(\pi)} i^* \Omega(A)_{|I|} \longrightarrow i^* \Omega(A)_{|J|}$$

Once again this follows from the standard diagram chasing:

$$\Delta_*^{(\pi)} i^* \Omega(A)_{|I|} := \Delta_*^{(\pi)} i^* p_I^! \mathcal{E}(A) \cong \Delta_*^{(\pi)} i^* p_I^* \mathcal{E}(A), \quad (p_I \text{ is submersion}),$$

$$\Delta_*^{(\pi)} i^* p_I^* \mathcal{E}(A) \cong \Delta_*^{(\pi)} p_{\mathbb{R}, I}^* i_\sigma^* \mathcal{E}(A), \quad (\text{square (5)})$$

$$\Delta_*^{(\pi)} p_{\mathbb{R}, I}^* i_\sigma^* \mathcal{E}(A) \cong p_{\mathbb{R}, J}^* \Delta_*^{|\pi|} i_\sigma^* \mathcal{E}(A), \quad (\text{base change (6)}).$$

Hence from the quiver description of a category  $\mathcal{M}^B(\mathbb{A}^I, \mathcal{S}_\emptyset)$  from [KS1] isomorphism (7) extends to an isomorphism of perverse sheaves.  $\square$

Note that for every  $I \in Fin$  sheaf  $\Omega(A)_I$  belongs to the subcategory of unipotent perverse sheaves.

**Remark 3.3.3.** Just like in the case of  $\mathcal{D}$ -modules it is natural to call  $\Omega(A) := \{\Omega(A)_I, \phi(\pi)\}_{Fin}$  an *unipotent perverse on Ran space* of  $\mathbb{A}^1$ . These objects with obvious morphisms form a category denoted  $\mathcal{M}_{un}^B(Ran(\mathbb{A}^1))$ .

**Proposition 3.3.4.** Unipotent perverse sheaf  $\Omega(A)$  over  $Ran(\mathbb{A})$  has the following additional structure:

- For every surjection  $\pi: J \rightarrow I$ , bialgebra structure induces an isomorphism:

$$v_\pi: j^{(\pi)*} \Omega(A)_J \xrightarrow{\sim} j^{(\pi)*} \boxtimes_{i \in I} \Omega(A)_{J_i}.$$

- Compatibilities between  $v$ 's and compositions of surjections.
- Compatibilities between maps  $\phi$  and  $v$ .

*Proof.* For every  $n$  and an ordered partition  $(n_1, \dots, n_k)$  of  $n$  we have an *addition map*:

$$a_{(n_1, \dots, n_k)}: Sym^{n_1}(\mathbb{A}) \times \dots \times Sym^{n_k}(\mathbb{A}) \longrightarrow Sym^n(\mathbb{A}).$$

Denote by  $Sym^{(n_1, \dots, n_k)}(\mathbb{A})_{disj} \hookrightarrow Sym^{n_1}(\mathbb{A}) \times \dots \times Sym^{n_k}(\mathbb{A})$  a subspace which consists  $k$ -tuples of divisors with a disjoint support. Then following [KS2] one can show that we have natural isomorphisms:

$$v_{(n_1, \dots, n_k)}: j^{(n_1, \dots, n_k)*} a_{(n_1, \dots, n_k)}^* \mathcal{E}(A)_n \xrightarrow{\sim} j^{(n_1, \dots, n_k)*} \mathcal{E}(A)_{n_1} \boxtimes \dots \boxtimes \mathcal{E}(A)_{n_k}$$

which is associative in a natural case.

For very  $\pi: J \twoheadrightarrow I$  we have the following commutative diagram:

$$(11) \quad \begin{array}{ccccc} U(\pi) & \xrightarrow{j^{(\pi)}} & \times_{i \in I} \mathbb{A}^{J_i} & \xrightarrow{id} & \mathbb{A}^J \\ \downarrow p_{disj} & & \downarrow \times_I(p_{J_i}) & & \downarrow p_J \\ Sym^{|\pi|}(\mathbb{A})_{disj} & \xrightarrow{j^{|\pi|}} & \times_I Sym^{|\pi^{-1}(i)|}(\mathbb{A}) & \xrightarrow{a_{|\pi|}} & Sym^{|J|}(\mathbb{A}) \end{array}$$

Then a standard diagram chasing gives us:

$$(12) \quad j^{(\pi)*} \Omega(A)_J \cong j^{(\pi)*} p_J^* \mathcal{E}(A)_{|J|} \cong p_{disj}^* j^{|\pi|*} a_{|\pi|}^* \mathcal{E}(A)_{|J|},$$

$$(13) \quad p_{disj}^* j^{|\pi|*} a_{|\pi|}^* \mathcal{E}(A)_{|J|} \cong p_{disj}^* j^{|\pi|} \boxtimes_I \mathcal{E}(A)_{|\pi^{-1}(i)|} \quad (\text{morphism } c_{|\pi|}),$$

$$(14) \quad p_{disj}^* j^{|\pi|*} \boxtimes_I \mathcal{E}(A)_{|\pi^{-1}(i)|} \cong p_{disj}^* j^{(\pi)*} \boxtimes_I p_{J_i}^* \mathcal{E}(A)_{|\pi^{-1}(i)|},$$

$$(15) \quad \cong j^{(\pi)*} \boxtimes_{i \in I} \Omega(A)_{J_i}.$$

Compatibilities between  $v_\pi$  and  $\phi(\pi)$  are obvious ones.  $\square$

**Remark 3.3.5.** We say that  $\mathcal{K}$  in  $\mathcal{M}_{un}^B(Ran(\mathbb{A}), \mathcal{S}_\emptyset)$  is an *unipotent factorizable sheaf* on  $\mathbb{A}^1$  if conditions above are satisfied. The corresponding category will be denoted by  $FS_{un}(Ran(\mathbb{A}^1))$ . It obvious that we have a functor:

$$\Omega: AlgB_c \longrightarrow FS_{un}(Ran(\mathbb{A})), \quad \Omega: A \longmapsto \Omega(A).$$

**3.4. Hyperbolic stalk.** In this subsection we will prove that the functor  $\Omega$  is an equivalence.

**Proposition 3.4.1.** We have a functor: <sup>3</sup>

$$(16) \quad \omega_{Ran}^B: FS_{un}(Ran(\mathbb{A})) \longrightarrow AlgB_c, \quad \omega_{Ran}^B := \lim_{Fin} \omega_I^B.$$

which is the quasi inverse to the functor  $\Omega$ .

*Proof.* Let  $\mathcal{K} \in FS_{un}(Ran(\mathbb{A}))$  then for every  $I \in Fin$  we define a space  $\omega_{Ran}^B(\mathcal{K})_{|I|} := \omega_I^B(\mathcal{K}_I)^{\Sigma_{|I|}}$ . Note that for every  $\pi: J \twoheadrightarrow I$  we have an inclusion  $\Delta_*^{(\pi)} \mathcal{K}_I \hookrightarrow \mathcal{K}_J$  which descends to an injection:  $\omega_{Ran}^B(\mathcal{K})_{|I|} \hookrightarrow \omega_{Ran}^B(\mathcal{K})_{|J|}$  and thus we have:

$$\omega_{Ran}^B(\mathcal{K})_1 \subset \dots \subset \omega_{Ran}^B(\mathcal{K})_n \subset \dots \subset \omega_{Ran}^B(\mathcal{K}).$$

Note that we have:  $i_\sigma^! \mathcal{E}(A) = (p_{I*} i^! p_I^* \mathcal{E}(A))^\Sigma$ , hence analogically to [KS2] generalization maps define a comultiplication:

$$\overline{\Delta}_{|I'|, |I''|}: \omega_{Ran}^B(\mathcal{K})_{|I|} \longrightarrow \omega_{Ran}^B(\mathcal{K})_{|I'|} \otimes \omega_{Ran}^B(\mathcal{K})_{|I''|}, \quad |I| = |I'| + |I''|,$$

which is coassociative.

Using *ibid* we also define a map:

$$m_{|I'|, |I''|}: \omega_{Ran}^B(\mathcal{K})_{|I'|} \otimes \omega_{Ran}^B(\mathcal{K})_{|I''|} \longrightarrow \omega_{Ran}^B(\mathcal{K})_{|I|}, \quad |I| = |I'| + |I''|,$$

which is associative. Once again by *ibid* operators  $m_{|I'|, |I''|}$  and  $\overline{\Delta}_{|I'|, |I''|}$  are compatible, therefore  $\omega_{Ran}^B(\mathcal{K})$  is a conilpotent bialgebra with a unit.  $\square$

<sup>3</sup>Strictly speaking functor  $\omega_{Ran}^B$  takes values in the category of bialgebras without a unit, but since the latter category is equivalent to  $AlgB_c$  we will not distinguish between them.

## 4. QUANTIZATION AND COMPLEMENTS

**4.1. Quantization.** Recall that we have a de Rham functor which induces *Riemann-Hilbert equivalence*  $RH: \mathcal{M}_{un}^{dR}(\mathbb{A}^I, \mathcal{S}_\emptyset) \xrightarrow{\sim} \mathcal{M}_{un}^B(\mathbb{A}^I, \mathcal{S}_\emptyset)$ . Important property of this equivalence that  $RH$  commutes with a six functor formalism and therefore induces an equivalence between unipotent  $\mathcal{D}$ -modules and unipotent perverse sheaves on  $Ran(\mathbb{A}^1)$  and also an equivalence between categories of unipotent factorization algebras and unipotent factorizable sheaves. Now we are ready to give a construction of Lie bialgebras (resp. Hopf algebras) quantization (resp. dequantization):

**Definition 4.1.1.** We define a **quantization functor**  $Q$  over  $\mathbb{C}$  as a functor which makes the following diagram into commutative:

$$\begin{array}{ccc} LieB_c & \xrightarrow[\sim]{Q} & AlgB_c \\ \Lambda \downarrow \sim & & \Omega \downarrow \sim \\ FA_{un}(Ran(\mathbb{A}^1)) & \xrightarrow[\sim]{RH} & FS_{un}(Ran(\mathbb{A}^1)) \end{array}$$

**Remark 4.1.2** (Associators). Recall that all proofs of Etingof-Kazhdan quantization (for example [EK] [Tam]) require a choice of *associator*. Here we will briefly sketch how associators appear in our construction. To define a connection with Drinfeld associators let us note that categories  $\mathcal{M}_{un}^B(Ran(\mathbb{A}^1))$  and  $\mathcal{M}_{un}^{dR}(Ran(\mathbb{A}^1))$  can be equipped with certain *natural tensor structures*. These tensor structures come from *fusion tensor structure* on category of perverse sheaves (resp.  $\mathcal{D}$ -modules) on  $\mathbb{A}^I$ . The definition of the fusion tensor product in the case of perverse sheaves is the following:

Denote by  $\Delta$  a flat in a product stratification of  $\mathbb{A}_{\mathbb{R}}^I \times \mathbb{A}_{\mathbb{R}}^I$ , which corresponds to a  $\mathbb{A}_{\mathbb{R}}^I$ . The inclusion of a complex diagonal  $\Delta_{\mathbb{C}}$  can be factored as:

$$(17) \quad k: \Delta_{\mathbb{C}} = \Delta + i\Delta \hookrightarrow (\mathbb{A}_{\mathbb{R}}^I \times \mathbb{A}_{\mathbb{R}}^I) + i\Delta, \quad j_{\Delta}: (\mathbb{A}_{\mathbb{R}}^I \times \mathbb{A}_{\mathbb{R}}^I) + i\Delta \hookrightarrow \mathbb{A}^I \times \mathbb{A}^I.$$

Then for two perverse sheaves  $\mathcal{K}_1$  and  $\mathcal{K}_2$  on  $\mathbb{A}^I$  the corresponding exterior tensor product  $\mathcal{K}_1 \boxtimes \mathcal{K}_2$  is naturally a perverse sheaf on  $\mathbb{A}^I \times \mathbb{A}^I$  smooth with respect to a product stratification. We define a *new perverse sheaf*  $\mathcal{K}_1 \otimes^{*!} \mathcal{K}_2$  by the rule:

$$(18) \quad \mathcal{K}_1 \otimes^{*!} \mathcal{K}_2 := k^* j_{\Delta}^! \mathcal{K}_1 \boxtimes \mathcal{K}_2[|I|],$$

This operation naturally extends to the symmetric unital product on  $\mathcal{M}_{un}^B(\mathbb{A}^I, \mathcal{S}_\emptyset)$ .

Categories  $\mathcal{M}_{un}^B(Ran(\mathbb{A}^1))$  and  $\mathcal{M}_{un}^{dR}(Ran(\mathbb{A}^1))$  are equipped with "fiber" functors  $\omega_{Ran}^B$  and  $\omega_{Ran}^{dR}$  to vector spaces. Composing the functor  $\omega_{Ran}^B$  with the Riemann-Hilbert functor we obtain two "fiber" functors from category of  $\mathcal{D}$ -modules on  $Ran(\mathbb{A})$ . Then we have the following:

**Conjecture 1.** Denote by  $\text{Isom}^{\otimes}(\omega_{Ran}^{dR}, \omega_{Ran}^B RH)$  scheme of tensor isomorphisms between "fiber" functors, then we have an isomorphism of schemes:

$$\text{Isom}^{\otimes}(\omega_{Ran}^{dR}, \omega_{Ran}^B RH) \cong \text{Ass}(\mathbb{C})$$

Note that equivalences  $\Lambda$  and  $\Omega$  can be easily defined over  $\mathbb{Q}$ . It seems that together with methods from [Dre] this allows us to extend the above conjecture to the case of rational associators.

## REFERENCES

- [BD] Alexander Beilinson and Vladimir Drinfeld. *Chiral Algebras*, volume 51 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [Bei] Alexander Beilinson. How to glue perverse sheaves. In Yuri Manin, I., editor, *K-Theory, Arithmetic and Geometry*, volume 1289 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1987.
- [Dre] Brad Drew. *Réalisations tannakiennes des motifs mixtes triangulés*. PhD thesis, Univ. Paris 13, 2013.
- [Dri] Vladimir Drinfeld. Quantum groups. In *Proceedings of the International Congress of Mathematicians 1986*.
- [EK] Pavel Etingof and David Kazhdan. Quantization of Lie bialgebras. *Selecta math.*, 2:1–41, 1996.
- [Kho] Sergey Khoroshkin.  $D$ -modules over the arrangements of hyperplanes. *Communications in Algebra*, 23:3481–3504, 1995.
- [KS1] Mikhail Kapranov and Vadim Schechtman. Perverse sheaves over real hyperplane arrangements. *Ann. Math*, 183:619–679, 2016.
- [KS2] Mikhail Kapranov and Vadim Schechtman. Shuffle algebras and perverse sheaves. <https://arxiv.org/pdf/1904.09325.pdf>, 2019.
- [KS3] Sergey Khoroshkin and Vadim Schechtman. Factorizable  $\mathcal{D}$ -modules. *Theory and Applications of Categories*, 11:107–131, 2003.
- [KV] Sergey Khoroshkin and Alexander Varchenko. Quiver  $D$ -modules and homology of local systems over an arrangement of hyperplanes. *International Mathematics Research Papers*, pages 1–116, 2006.
- [Tam] Dmitry Tamarkin. Quantization of Lie bialgebras via the formality of the operad of little disks. *GAGA Geometric And Functional Analysis*, 17:537–604, 2007.

MATHEMATICS RESEARCH UNIT, LUXEMBOURG UNIVERSITY, GRAND DUCHY OF LUXEMBOURG  
 AND INDEPENDENT UNIVERSITY OF MOSCOW, B. VLASIEVSKY PER., 11, MOSCOW 121002, RUSSIA  
*E-mail address:* alexey.kalugin@uni.lu