Finite-dimensional Hopf algebras over the Hopf algebra $H_{b:1}$ of Kashina

Yiwei Zheng ^a, Yun Gao ^{a,b}, Naihong Hu ^{c, *}

^aDepartment of Mathematics, Shanghai University, Shanghai 200444, China

^bDepartment of Mathematics and Statistics, York University, Toronto M3J 1P3, Canada

^cSchool of Mathematical Sciences, Shanghai Key Laboratory of PMMP,

East China Normal University, Shanghai 200241, China

Abstract

Let H be the 16-dimensional nontrivial (namely, noncommutative and noncocommutative) semisimple Hopf algebra $H_{b:1}$ appeared in Kashina's work [28]. We obtain all simple Yetter-Drinfeld modules over H and then determine all finite-dimensional Nichols algebras satisfying $\mathcal{B}(V) \cong \bigotimes_{i \in I} \mathcal{B}(V_i)$, where $V = \bigoplus_{i \in I} V_i$, each V_i is a simple object in ${}^{H}_{H}\mathcal{YD}$. Finally, we describe all liftings of those $\mathcal{B}(V)$ over H.

1 Introduction

Let k be an algebraically closed field of characteristic zero. In 1975, Kaplansky [27] posed ten conjectures including the number of iso-classes of finite-dimensional Hopf algebras over k. Until now, although there are many classification results, there are no standard methods, except that in 1998, Andruskiewitsch and Schneider [10] put forward the lifting method which later on plays a very important role in the classification of finite-dimensional (co-)pointed Hopf algebras.

Actually, the lifting method works well for those classes of finite-dimensional Hopf algebras with Radford biproduct structure or their deformations by certain Hopf 2-cocycles. Let Abe a Hopf algebra. If the coradical A_0 is a subalgebra, then the coradical filtration $\{A_n\}_{n\geq 0}$ is a Hopf algebra filtration, and the associated graded coalgebra $grA = \bigoplus_{n>0} A_n/A_{n-1}$ is

^{*}E-mails: nhhu@math.ecnu.edu.cn, ygao@yorku.ca, zhengyiwei12@foxmail.com

also a Hopf algebra, where $A_{-1} = 0$. Let $\pi : grA \to A_0$ be the homogeneous projection. By a theorem of Radford [31], there exists a unique connected graded braided Hopf algebra $R = \bigoplus_{n>0} R(n)$ in the monoidal category $A_0 \mathcal{YD}$ such that $grA \cong R \sharp A_0$. We call R or R(1)the diagram or infinitesimal braiding of A, respectively. Moreover, R is strictly graded, that is, R(0) = k, $R(1) = \mathcal{P}(R)$ (see Definition 1.13 [12]).

The lifting method has been applied to classify some finite-dimensional pointed Hopf algebras such as [3], [4], [5], [6], [9], [13], [17], [20], etc., and copointed Hopf algebras [14], [26], etc. Nevertheless, there are a few classification results on finite-dimensional Hopf algebras whose coradical is neither a group algebra nor the dual of a group algebra, for instance, [7], [16], [19], [24], [25], [33], [34], etc. In fact, Shi began a program in [33] to classify the objects of finite-dimensional growth from a given nontrivial semisimple Hopf algebra $A_0 = H_8$ via some relevant Nichols algebras $\mathcal{B}(V)$ derived from its semisimple Yetter-Drinfeld modules $V \in {}^{A_0}_{A_0}\mathcal{YD}.$

The classification of finite-dimensional Hopf algebras A whose coradical A_0 is a (Hopf) subalgebra needs the following main steps:

(a) Determine those V in ${}^{A_0}_{A_0}\mathcal{YD}$ such that the Nichols algebra $\mathcal{B}(V)$ is finite-dimensional and present $\mathcal{B}(V)$ by generators and relations.

(b) If $R = \bigoplus_{n \ge 0} R(n)$ is a finite-dimensional Hopf algebra in ${}^{A_0}_{A_0} \mathcal{YD}$ with V = R(1), decide if $R \cong \mathcal{B}(V)$.

(c) Given V as in (a), classify all A such that $grA \cong \mathcal{B}(V) \sharp A_0$. We call A a lifting of $\mathcal{B}(V)$ over A_0 .

In this paper, we fix a 16-dimensional non-trivial semisimple Hopf algebra $H = H_{b:1}$, which appeared in [28], and study these questions. For the definition of Hopf algebra H, see Definition 3.1. We first determine the Drinfeld double $D := D(H^{cop})$ of H^{cop} and describe the simple D-modules. In fact, we prove in Theorem 3.6 that there are 32 one-dimensional simple objects $\mathbb{k}_{\chi_{i,j,k,l}}$ with $0 \le i, k, l < 2, 0 \le j < 4$, and 56 two-dimensional simple objects $V_{i,j,k,l,m,n}$ with $(i,j,k,l,m,n) \in \Omega$, $W_{i,j,k,l}$ with $(i,j,k,l) \in \Lambda^1$, $U_{i,j,k,l}$ with $(i,j,k,l) \in \Lambda^2$. Using the fact that the baided categories ${}_{D}\mathcal{M}$ and ${}_{H}^{H}\mathcal{YD}$ are monoidally isomorphic (see [29]), we actually get the simple objects in ${}^{H}_{H}\mathcal{YD}$. Furthermore, we get all the possible finitedimensional Nichols algebras of semisimple objects satisfying $\mathcal{B}(V) \cong \bigotimes_{i \in I} \mathcal{B}(V_i)$. Here is our first main result.

Theorem A Let $V = \bigoplus_{i \in I} V_i$, each V_i be a simple object in ${}^H_H \mathcal{YD}$. Then the Nichols algebra $\mathcal{B}(V)$ satisfying $\mathcal{B}(V) \cong \bigotimes_{i \in I} \mathcal{B}(V_i)$ is finite-dimensional if and only if V is isomorphic to one of the following Yetter-Drinfeld modules

- (1) $\Omega_1(n_1, n_2, n_3, n_4, n_5, n_6, n_7, n_8) \triangleq \bigoplus_{j=1}^8 V_j \text{ with } \sum_{j=1}^8 n_j \ge 1.$ (2) $\Omega_2(n_1, n_2, n_3, n_4) \triangleq V_3^{\oplus n_1} \oplus V_4^{\oplus n_2} \oplus V_7^{\oplus n_3} \oplus V_8^{\oplus n_4} \oplus M_1 \text{ with } \sum_{j=1}^4 n_j \ge 0.$ (3) $\Omega_3(n_1, n_2, n_3, n_4) \triangleq V_1^{\oplus n_1} \oplus V_2^{\oplus n_2} \oplus V_5^{\oplus n_3} \oplus V_6^{\oplus n_4} \oplus M_2 \text{ with } \sum_{j=1}^4 n_j \ge 0.$
- (4) $\Omega_4(n_1, n_2, n_3, n_4) \triangleq V_1^{\oplus n_1} \oplus V_3^{\oplus n_2} \oplus V_5^{\oplus n_3} \oplus V_7^{\oplus n_4} \oplus M_3 \text{ with } \sum_{i=1}^4 n_i \geq 0.$

 $\begin{array}{l} (5) \ \Omega_{5}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{1}^{\oplus n_{1}} \oplus V_{4}^{\oplus n_{2}} \oplus V_{5}^{\oplus n_{3}} \oplus V_{8}^{\oplus n_{4}} \oplus M_{4} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (6) \ \Omega_{6}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{2}^{\oplus n_{1}} \oplus V_{4}^{\oplus n_{2}} \oplus V_{6}^{\oplus n_{3}} \oplus V_{8}^{\oplus n_{4}} \oplus M_{5} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (7) \ \Omega_{7}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{2}^{\oplus n_{1}} \oplus V_{3}^{\oplus n_{2}} \oplus V_{6}^{\oplus n_{3}} \oplus V_{7}^{\oplus n_{4}} \oplus M_{6} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (8) \ \Omega_{8}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{1}^{\oplus n_{1}} \oplus V_{2}^{\oplus n_{2}} \oplus V_{5}^{\oplus n_{3}} \oplus V_{6}^{\oplus n_{4}} \oplus M_{7} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (9) \ \Omega_{9}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{3}^{\oplus n_{1}} \oplus V_{4}^{\oplus n_{2}} \oplus V_{7}^{\oplus n_{3}} \oplus V_{8}^{\oplus n_{4}} \oplus M_{8} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (10) \ \Omega_{10}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{2}^{\oplus n_{1}} \oplus V_{4}^{\oplus n_{2}} \oplus V_{6}^{\oplus n_{3}} \oplus V_{8}^{\oplus n_{4}} \oplus M_{9} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (11) \ \Omega_{11}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{2}^{\oplus n_{1}} \oplus V_{4}^{\oplus n_{2}} \oplus V_{6}^{\oplus n_{3}} \oplus V_{7}^{\oplus n_{4}} \oplus M_{9} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (12) \ \Omega_{12}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{1}^{\oplus n_{1}} \oplus V_{3}^{\oplus n_{2}} \oplus V_{6}^{\oplus n_{3}} \oplus V_{7}^{\oplus n_{4}} \oplus M_{10} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (13) \ \Omega_{13}(n_{1},n_{2},n_{3},n_{4}) \triangleq V_{1}^{\oplus n_{1}} \oplus V_{3}^{\oplus n_{2}} \oplus V_{5}^{\oplus n_{3}} \oplus V_{7}^{\oplus n_{4}} \oplus M_{10} \ with \sum_{j=1}^{4} n_{j} \geq 0. \\ (14) \ \Omega_{14} \triangleq M_{1} \oplus M_{1}, \ \Omega_{15} \triangleq M_{1} \oplus M_{2}, \ \Omega_{16} \triangleq M_{1} \oplus M_{7}, \ \Omega_{17} \triangleq M_{3} \oplus M_{3}, \ \Omega_{18} \triangleq M_{3} \oplus M_{5}, \\ \Omega_{19} \triangleq M_{3} \oplus M_{9}, \ \Omega_{20} \triangleq M_{4} \oplus M_{4}, \ \Omega_{21} \triangleq M_{4} \oplus M_{6}, \ \Omega_{22} \triangleq M_{7} \oplus M_{7}, \ \Omega_{23} \triangleq M_{7} \oplus M_{1}, \\ \Omega_{24} \triangleq M_{13} \oplus M_{13}, \ \Omega_{25} \triangleq M_{13} \oplus M_{14}, \ \Omega_{26} \triangleq M_{15} \oplus M_{15}, \ \Omega_{27} \triangleq M_{15} \oplus M_{16}, \ \Omega_{28} \triangleq M_{17} \oplus M_{17}, \\ \Omega_{29} \triangleq M_{17} \oplus M_{18}. \end{array}$

 $(15) \ \Omega_{30} \triangleq M_2 \oplus M_2, \ \Omega_{31} \triangleq M_2 \oplus M_8, \ \Omega_{32} \triangleq M_4 \oplus M_{10}, \ \Omega_{33} \triangleq M_5 \oplus M_5, \ \Omega_{34} \triangleq M_5 \oplus M_{11}, \\ \Omega_{35} \triangleq M_6 \oplus M_6, \ \Omega_{36} \triangleq M_6 \oplus M_{12}, \ \Omega_{37} \triangleq M_8 \oplus M_8, \ \Omega_{38} \triangleq M_9 \oplus M_9, \ \Omega_{39} \triangleq M_9 \oplus M_{11}, \\ \Omega_{40} \triangleq M_{10} \oplus M_{10}, \ \Omega_{41} \triangleq M_{10} \oplus M_{12}, \ \Omega_{42} \triangleq M_{11} \oplus M_{11}, \ \Omega_{43} \triangleq M_{12} \oplus M_{12}, \ \Omega_{44} \triangleq M_{14} \oplus M_{14}, \\ \Omega_{45} \triangleq M_{16} \oplus M_{16}, \ \Omega_{46} \triangleq M_{18} \oplus M_{18}, \ \Omega_{47} \triangleq M_{19} \oplus M_{19}, \ \Omega_{48} \triangleq M_{19} \oplus M_{20}, \ \Omega_{49} \triangleq M_{20} \oplus M_{20}.$

Remark 1.1 We point out which of the Yetter-Drinfeld modules have principal realizations and which not, since the liftings are known when there exist principal realizations [1, Subsection 2.2]. Let (h) and (δ_h) be dual bases of H and H^{*}. Define

$$\begin{aligned} \chi := & \delta_1 + \delta_{x^2} + (-1)^i (\delta_x + \delta_{x^3}) + (-1)^j (\delta_y + \delta_{x^2y}) + (-1)^{i+j} (\delta_{xy} + \delta_{x^3y}) \\ &+ (-1)^k (\delta_t + \delta_{x^2t}) + (-1)^{i+k} (\delta_{xt} + \delta_{x^3t}) + (-1)^{j+k} (\delta_{yt} + \delta_{x^2yt}) \\ &+ (-1)^{i+j+k} (\delta_{xyt} + \delta_{x^3yt}) \in Alg(H, \Bbbk), \end{aligned}$$

then $(x^j y^l, \chi)$ is a YD-pair [2] if and only if $\mathbb{K}^{\chi}_{x^j y^l}$ is a one-dimensional Yetter-Drinfeld module. So only $\Omega_1(n_1, ..., n_8)$ has a principal realization.

Based on the principle of the lifting method, we classify the finite-dimensional Hopf algebras over H such that their infinitesimal braidings are those Yetter-Drinfeld modules listed in Theorem A. Here is the second main result.

Theorem B Let A be a finite-dimensional Hopf algebra over H such that its infinitesimal braiding is isomorphic to one of the Yetter-Drinfeld modules in Theorem A, then A is isomorphic either to

- (1) $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$, see Definition 6.4;
- (2) $\mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2)$, see Definition 6.7;
- (3) $\mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4)$, see Definition 6.11;
- (4) $\mathfrak{U}_5(n_1, n_2, n_3, n_4; I_5)$, see Definition 6.12;

- (5) $\mathfrak{U}_8(n_1, n_2, n_3, n_4; I_8)$, see Definition 6.9;
- (6) \$\mathbf{U}_{14}(I_{14})\$, see Definition 6.14; \$\mathbf{U}_{15}(\lambda, \mu)\$, see Definition 6.15;
 \$\mathbf{U}_{16}(\lambda, \mu)\$, see Definition 6.16; \$\mathbf{U}_{17}(I_{17})\$, see Definition 6.17;
 \$\mathbf{U}_{18}(I_{18})\$, see Definition 6.18; \$\mathbf{U}_{19}(\lambda, \mu)\$, see Definition 6.19;
 \$\mathbf{U}_{20}(I_{20})\$, see Definition 6.21; \$\mathbf{U}_{21}(\lambda, \mu)\$, see Definition 6.22;
 \$\mathbf{U}_{22}(I_{22})\$, see Definition 6.23; \$\mathbf{U}_{23}(I_{23})\$, see Definition 6.24;
 \$\mathbf{U}_{24}(\lambda)\$, see Definition 6.26; \$\mathbf{U}_{26}(I_{26})\$, see Definition 6.27;
 \$\mathbf{U}_{27}(\lambda, \mu)\$, see Definition 6.28; \$\mathbf{U}_{28}(I_{28})\$, see Definition 6.31;
 \$\mathbf{U}_{29}(\lambda, \mu)\$, see Definition 6.32.
- (7) $\mathcal{B}(\Omega_{25})$ $\sharp H$, see Proposition 6.30.

Except for the case (7), the remaining families of Hopf algebras contain non-trivial lifting relations.

Remark 1.2 Note that (1) dim $\mathfrak{U}_1(n_1, n_2, \dots, n_8; I_1) = 2^{4 + \sum_{i=1}^8 n_i};$ (2) For $k \in \{2, 4, 5, 8\}$, dim $\mathfrak{U}_k(n_1, n_2, n_3, n_4; I_k) = 2^{6 + \sum_{i=1}^4 n_i};$ (3) For $i \in \{14, 17, 18, 20, 22, 23, 26, 28\}$, dim $\mathfrak{U}_i(I_i) = 256;$ (4) For $i \in \{15, 16, 19, 21, 27, 29\}$, dim $\mathfrak{U}_i(\lambda, \mu) = \dim \mathfrak{U}_{24}(\lambda) = 256.$

The paper is organized as follows. In section 2, we recall some basics and notations of Yetter-Drinfeld modules, Nichols algebras, Radford biproduct and Drinfeld double. In section 3, we describe the structures of H and present the Drinfeld double $D = D(H^{cop})$ by generators and relations. We also determine the simple D-modules. In section 4, we describe the simple objects of ${}^{H}_{H}\mathcal{YD}$ by using the monoidal isomorphism of braided categories ${}^{H}_{H}\mathcal{YD} \cong {}^{D}\mathcal{M}$. In section 5, we obtain all the possible finite-dimensional Nichols algebras of semisimple modules satisfying $\mathcal{B}(V) \cong \bigotimes_{i \in I} \mathcal{B}(V_i)$. In section 6, based on the principle of the lifting method, we classify the finite-dimensional Hopf algebras over H such that their infinitesimal braidings are those Yetter-Drinfeld modules listed in Theorem A. Then we get Theorem B.

2 Preliminaries

Conventions. Throughout the paper, the ground field \Bbbk is an algebraically closed field of characteristic zero and we denote by ξ a primitive 4-th root of unity. For the references of Hopf algebra theory, one can consult [30], [29], [32], [35], etc.

If H is a Hopf algebra over \Bbbk , then \triangle , ε , S denote the comultiplication, the counit and the antipode, respectively. We use Sweedler's notation for the comultiplication and coaction, e.g., $\triangle(h) = h_{(1)} \otimes h_{(2)}$ for $h \in H$. We denote by H^{op} the Hopf algebra with the opposite multiplication, by H^{cop} the Hopf algebra with the opposite comultiplication, and by H^{bop} the Hopf algebra $H^{op\ cop}$. Denote by G(H) the set of group-like elements of H.

$$\mathcal{P}_{g,h}(H) = \{ x \in H \mid \triangle(x) = x \otimes g + h \otimes x \}, \quad \forall \ g, \ h \in G(H).$$

In particular, the linear space $\mathcal{P}(H) = \mathcal{P}_{1,1}(H)$ is called the set of primitive elements.

If V is a k-vector space, $v \in V$, $f \in V^*$, we use either f(v), $\langle f, v \rangle$ or $\langle v, f \rangle$ to denote the evaluation. Given $n \geq 0$, we denote $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ and $\mathbb{I}_{0,n} = \{0, 1, \dots, n\}$. It is emphasized that the operations ij and $i \pm j$ are considered modulo n+1 for $i, j \in \mathbb{I}_{0,n}$ when not specified.

2.1 Yetter-Drinfeld modules and Nichols algebras

Let *H* be a Hopf algebra. A left Yetter-Drinfeld module *V* over *H* is a left *H*-module (V, \cdot) and a left *H*-comodule (V, δ) with $\delta(v) = v_{(-1)} \otimes v_{(0)} \in H \otimes V$ for all $v \in V$, satisfying

$$\delta(h \cdot v) = h_{(1)} v_{(-1)} S(h_{(3)}) \otimes h_{(2)} \cdot v_{(0)}, \qquad \forall \ v \in V, \ h \in H.$$

We denote by ${}^{H}_{H}\mathcal{YD}$ the category of finite-dimensional left Yetter-Drinfeld modules over H. It is a braided monoidal category: for $V, W \in {}^{H}_{H}\mathcal{YD}$, the braiding $c_{V,W} : V \otimes W \to W \otimes V$ is given by

$$c_{V,W}(v \otimes w) = v_{(-1)} \cdot w \otimes v_{(0)}, \quad \forall \ v \in V, \ w \in W.$$

$$(2.1)$$

In particular, $(V, c_{V,V})$ is a braided vector space, that is, $c := c_{V,V}$ is a linear isomorphism satisfying the braid equation

$$(c \otimes id)(id \otimes c)(c \otimes id) = (id \otimes c)(c \otimes id)(id \otimes c).$$

Moreover, ${}^{H}_{H}\mathcal{YD}$ is rigid. Denote by V^{*} the left dual defined by

$$\langle h \cdot f, v \rangle = \langle f, S(h)v \rangle, \qquad f_{(-1)} \langle f_{(0)}, v \rangle = S^{-1}(v_{(-1)}) \langle f, v_{(0)} \rangle.$$

Definition 2.1 [12, Definition 2.1] Let H be a Hopf algebra and V a Yetter-Drinfeld module over H. A braided \mathbb{N} -graded Hopf algebra $R = \bigoplus_{n\geq 0} R(n)$ in ${}^{H}_{H}\mathcal{YD}$ is called a Nichols algebra of V if

$$\mathbb{k} \simeq R(0), \ V \simeq R(1), \ R(1) = \mathcal{P}(R), \ R \ is \ generated \ as \ an \ algebra \ by \ R(1)$$

For any $V \in {}^{H}_{H}\mathcal{YD}$ there is a Nichols algebra $\mathcal{B}(V)$ associated to it. It is the quotient of the tensor algebra T(V) by the largest homogeneous two-sided ideal I satisfying:

- I is generated by homogeneous elements of degree ≥ 2 .
- $\triangle(I) \subseteq I \otimes T(V) + T(V) \otimes I$, i.e., it is also a coideal.

In such case, $\mathcal{B}(V) = T(V)/I$. See [12, Section 2.1] for details.

Remark 2.2 As well known, the Nichols algebra as an algebra and a coalgebra is completely determined by the braided space. If $W \subseteq V$ is a subspace such that $c(W \otimes W) \subseteq W \otimes W$, then dim $\mathcal{B}(V) = \infty$ if dim $\mathcal{B}(W) = \infty$. In particular, if V contains a non-zero element v such that $c(v \otimes v) = v \otimes v$, then dim $\mathcal{B}(V) = \infty$.

Lemma 2.3 [21, Theorem 2.2] Let (V, c) be a braided vector space, $V = \bigoplus_{i=1}^{n} V_i$, such that each $\mathcal{B}(V_i)$ is finite-dimensional. Then dim $\mathcal{B}(V) \ge \prod_{i=1}^{n} \dim \mathcal{B}(V_i)$. Furthermore, the equality holds if and only if $b_{ij} = b_{ji}^{-1}$ for all $i \ne j$ ($b_{ij} = c \mid_{V_i \otimes V_j}$).

2.2 Bosonization and Hopf algebras with a projection

Let R be a Hopf algebra in ${}^{H}_{H}\mathcal{YD}$ and denote the coproduct by $\Delta_{R}(r) = r^{(1)} \otimes r^{(2)}$ for $r \in R$. We define the Radford biproduct or bosonization $R \sharp H$ as follows: as a vector space, $R \sharp H = R \otimes H$, and the multiplication and comultiplication are given by the smash product and smash-coproduct, respectively:

$$(r\sharp g)(s\sharp h) = r(g_{(1)} \cdot s) \sharp g_{(2)}h, \quad \triangle(r\sharp g) = r^{(1)} \sharp (r^{(2)})_{(-1)} g_{(1)} \otimes (r^{(2)})_{(0)} \sharp g_{(2)}.$$

Clearly, the map $\iota: H \to R \sharp H, h \mapsto 1 \sharp h, \forall h \in H$ is injective and the map

$$\pi: R \sharp H \to H, \ r \sharp h \mapsto \varepsilon_R(r)h, \ \forall \ r \in R, \ h \in H$$

is surjective such that $\pi \circ \iota = id_H$. Moreover, it holds that

$$R = (R \sharp H)^{coH} = \{ x \in R \sharp H \mid (id \otimes \pi) \triangle (x) = x \otimes 1 \}.$$

Conversely, if A is a Hopf algebra with bijective antipode and $\pi : A \to H$ is a Hopf algebra epimorphism admitting a Hopf algebra section $\iota : H \to A$ such that $\pi \circ \iota = id_H$. Then $R = A^{co\pi}$ is a braided Hopf algebra in ${}^{H}_{H}\mathcal{YD}$ and $A \cong R \sharp H$ as Hopf algebras. See [31] and [32] for more details.

2.3 The Drinfeld double

Let H be a finite-dimensional Hopf algebra over \Bbbk . The Drinfeld double $D(H) = H^{*cop} \otimes H$ is a Hopf algebra with the tensor product coalgebra structure and algebra structure defined by

$$(p \otimes a)(q \otimes b) = p\langle q_{(3)}, a_{(1)} \rangle q_{(2)} \otimes a_{(2)} \langle q_{(1)}, S^{-1}(a_{(3)}) \rangle b, \quad \forall \ p, \ q \in H^*, \ a, \ b \in H.$$

Proposition 2.4 [30, Proposition 10.6.16] Let H be a finite-dimensional Hopf algebra. Then the Yetter-Drinfeld category ${}^{H}_{H}\mathcal{YD}$ can be identified with the category ${}_{D(H^{cop})}\mathcal{M}$ of left modules over the Drinfeld double $D(H^{cop})$.

3 The Hopf algebra *H* and its Drinfeld double

In this section, we will recall the structure of the 16-dimensional non-trivial semisimple Hopf algebra $H = H_{b:1}$ in [28] and present the Drinfeld Double $D = D(H^{cop})$ by generators and relations. Then we will determine all simple left *D*-modules.

Definition 3.1 As an algebra H is generated by x, y, t satisfying the relations

$$x^{4} = 1, \quad y^{2} = 1, \quad t^{2} = 1, \quad xy = yx, \quad tx = x^{-1}t, \quad ty = yt,$$
 (3.2)

and its coalgebra is given by

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y, \quad \varepsilon(x) = \varepsilon(y) = 1, \tag{3.3}$$

$$\Delta(t) = \frac{1}{2} [(1+y)t \otimes t + (1-y)t \otimes x^2 t], \qquad \varepsilon(t) = 1, \tag{3.4}$$

and its antipode is given by

$$S(x) = x^{-1}, \quad S(y) = y, \quad S(t) = \frac{1}{2}[(1+y)t + (1-y)x^2t]$$

Remark 3.2 1. $G(H) = \langle x \rangle \times \langle y \rangle$, $\mathcal{P}_{1,g}(H) = \mathbb{k}\{1-g\}$ for $g \in G(H)$ and a linear basis of H is given by $\{x^i, x^iy, x^it, x^iyt, i \in \mathbb{I}_{0,3}\}$.

2. Denote by $\{(x^i)^*, (x^iy)^*, (x^it)^*, (x^iyt)^*, i \in \mathbb{I}_{0,3}\}$ the the dual basis of Hopf algebra H^* . Let

$$a = [(1 - x + x^{2} - x^{3})(1 + y)(1 + t)]^{*}, \quad b = [(1 + x + x^{2} + x^{3})(1 - y)(1 + t)]^{*},$$

$$c = [(1 + x + x^{2} + x^{3})(1 - y)(1 - t)]^{*}, \quad d = [(1 + \xi x - x^{2} - \xi x^{3})(1 + y)(1 + t)]^{*}.$$

Then using the multiplication table induced by the relations of H, it follows that

$$a^{2} = b^{2} = c^{2} = 1, \quad ab = ba, \quad ac = ca, \quad bc = cb,$$

$$d^{2} = a, \quad da = ad, \quad db = cd, \quad dc = bd,$$

$$\triangle(a) = a \otimes a, \quad \triangle(b) = b \otimes b, \quad \triangle(c) = c \otimes c,$$

$$\triangle(d) = \frac{1}{2}(d + bcd) \otimes d + \frac{1}{2}(d - bcd) \otimes ad.$$

3. The automorphisms of H are given in Table 1.

	x	y	t		x	y	t		x	y	t		x	y	t
$ au_1$	x	y	t	$ au_9$	x^3	y	t	$ au_{17}$	xy	y	t	$ au_{25}$	x^3y	y	t
$ au_2$	x	y	xt	$ au_{10}$	x^3	y	xt	$ au_{18}$	xy	y	xt	$ au_{26}$	x^3y	y	xt
$ au_3$	x	y	x^2t	$ au_{11}$	x^3	y	x^2t	$ au_{19}$	xy	y	x^2t	$ au_{27}$	x^3y	y	x^2t
$ au_4$	x	y	x^3t	$ au_{12}$	x^3	y	x^3t	$ au_{20}$	xy	y	x^3t	$ au_{28}$	x^3y	y	x^3t
$ au_5$	x	y	yt	$ au_{13}$	x^3	y	yt	$ au_{21}$	xy	y	yt	$ au_{29}$	x^3y	y	yt
$ au_6$	x	y	xyt	$ au_{14}$	x^3	y	xyt	$ au_{22}$	xy	y	xyt	$ au_{30}$	x^3y	y	xyt
$ au_7$	x	y	x^2yt	$ au_{15}$	x^3	y	x^2yt	$ au_{23}$	xy	y	x^2yt	$ au_{31}$	x^3y	y	x^2yt
$ au_8$	x	y	x^3yt	$ au_{16}$	x^3	y	x^3yt	$ au_{24}$	xy	y	x^3yt	$ au_{32}$	x^3y	y	x^3yt

Table 1. Automorphisms of H

Now we describe the Drinfeld Double $D(H^{cop})$ of H^{cop} .

Proposition 3.3 $D := D(H^{cop})$ as a coalgebra is isomorphic to tensor coalgebra $H^{*bop} \otimes H^{cop}$, and as an algebra is generated by the elements a, b, c, d, x, y, t such that x, y, t satisfying the relations of H^{cop} , a, b, c, d satisfying the relations of H^{*bop} and

$$\begin{array}{ll} xa=ax, & xb=bx, & xc=cx, & xd=bcdx, \\ ya=ay, & yb=by, & yc=cy, & yd=dy, \\ ta=at, & tb=bx^2t, & tc=cx^2t, & td=adyt. \end{array}$$

Proof. After a direct computation, we have that

$$\begin{split} & \triangle_{H^{cop}}^2(x) = x \otimes x \otimes x, \quad \triangle_{H^{cop}}^2(y) = y \otimes y \otimes y, \\ & \triangle_{H^{*bop}}^2(a) = a \otimes a \otimes a, \quad \triangle_{H^{*bop}}^2(b) = b \otimes b \otimes b, \quad \triangle_{H^{*bop}}^2(c) = c \otimes c \otimes c, \\ & \triangle_{H^{cop}}^2(t) = \frac{1}{4} [t \otimes (1+y)t \otimes (1+y)t + x^2t \otimes (1-y)t \otimes (1+y)t \\ & \quad + x^2t \otimes x^2(1+y)t \otimes (1-y)t + t \otimes x^2(1-y)t \otimes (1-y)t], \\ & \triangle_{H^{*bop}}^2(d) = \frac{1}{4} [d \otimes (d+bcd) \otimes (d+bcd) + ad \otimes (d-bcd) \otimes (d+bcd) \\ & \quad + ad \otimes a(d+bcd) \otimes (d-bcd) + d \otimes a(d-bcd) \otimes (d-bcd)]. \end{split}$$

It follows that

$$\begin{split} xa = \langle a, x \rangle ax \langle a, S(x) \rangle &= ax, \quad ya = \langle a, y \rangle ay \langle a, S(y) \rangle = ay, \\ xb = \langle b, x \rangle bx \langle b, S(x) \rangle &= bx, \quad yb = \langle b, y \rangle by \langle b, S(y) \rangle = by, \\ xc = \langle c, x \rangle cx \langle c, S(x) \rangle &= cx, \quad yc = \langle c, y \rangle cy \langle c, S(y) \rangle = cy, \\ xd = \frac{1}{4} [\langle d, x \rangle (d + bcd) x \langle d + bcd, S(x) \rangle + \langle ad, x \rangle (d - bcd) x \langle d + bcd, S(x) \rangle] = bcdx, \\ yd = \frac{1}{4} [\langle d, y \rangle (d + bcd) y \langle d + bcd, S(y) \rangle + \langle ad, y \rangle (d - bcd) y \langle d + bcd, S(y) \rangle] = dy, \\ ta = \frac{1}{4} [\langle a, t \rangle a(1 + y) t \langle a, S(t + yt) \rangle + \langle a, x^2 t \rangle a(1 - y) t \langle a, S(t + yt) \rangle] = at, \\ tb = \frac{1}{4} [\langle b, x^2 t \rangle bx^2 (1 + y) t \langle b, S(t - yt) \rangle + \langle b, t \rangle bx^2 (1 - y) t \langle b, S(t - yt) \rangle] = bx^2 t, \\ tc = \frac{1}{4} [\langle c, x^2 t \rangle cx^2 (1 + y) t \langle c, S(t - yt) \rangle + \langle c, t \rangle cx^2 (1 - y) t \langle c, S(t - yt) \rangle] = cx^2 t, \\ td = \frac{1}{16} [\langle ad, t \rangle a(d + bcd) (1 + y) t \langle d - bcd, S(t + yt) \rangle \\ + \langle ad, x^2 t \rangle a(d + bcd) (1 - y) t \langle d - bcd, S(t + yt) \rangle \\ + \langle d, t \rangle a(d - bcd) (1 + y) t \langle d - bcd, S(t + yt) \rangle \end{split}$$

$$+ \langle d, x^2 t \rangle a(d - bcd)(1 - y)t \langle d - bcd, S(t + yt) \rangle = adyt.$$

This completes the proof.

We begin by describing the one-dimensional *D*-modules.

Lemma 3.4 There are 32 non-isomorphic one-dimensional simple modules $\mathbb{k}_{\chi_{i,j,k,l}}$ given by the characters $\chi_{i,j,k,l}$, $0 \leq i$, k, l < 2, $0 \leq j < 4$, where

$$\begin{aligned} \chi_{i,j,k,l}(x) &= (-1)^i, \quad \chi_{i,j,k,l}(y) = (-1)^j, \quad \chi_{i,j,k,l}(t) = (-1)^k, \\ \chi_{i,j,k,l}(a) &= (-1)^j, \quad \chi_{i,j,k,l}(b) = (-1)^l, \quad \chi_{i,j,k,l}(c) = (-1)^l, \quad \chi_{i,j,k,l}(d) = \xi^j. \end{aligned}$$

Moreover, any one-dimensional D-module is isomorphic to $\mathbb{k}_{\chi_{i,j,k,l}}$ for some $0 \leq i, k, l < 2, 0 \leq j < 4$.

Proof. Let $\lambda: D \to \mathbb{k}$ be a character and write

$$\lambda(x) = \lambda_1, \quad \lambda(y) = \lambda_2, \quad \lambda(t) = \lambda_3, \quad \lambda(a) = \lambda_4, \quad \lambda(b) = \lambda_5, \quad \lambda(c) = \lambda_6, \quad \lambda(d) = \lambda_7.$$

Since $x^4 = 1$, $y^2 = t^2 = a^2 = b^2 = c^2 = 1$, it follows that

$$\lambda_1^4 = 1, \quad \lambda_2^2 = \lambda_3^2 = \lambda_4^2 = \lambda_5^2 = \lambda_6^2 = 1.$$

By $tx = x^{-1}t$, we have $\lambda_1 = \lambda_1^3$, that is $\lambda_1^2 = 1$. As db = cd, dc = bd, then $\lambda_5 = \lambda_6$. Because td = adyt, it follows that $\lambda_2\lambda_4 = 1$, i.e., $\lambda_2 = \lambda_4$. From $d^2 = a$, we get $\lambda_7^2 = \lambda_4$. Hence, λ is completely determined by $\lambda(x)$, $\lambda(y)$, $\lambda(t)$, $\lambda(b)$. Let

$$\lambda(x) = (-1)^i, \quad \lambda(y) = (-1)^j, \quad \lambda(t) = (-1)^k, \quad \lambda(b) = (-1)^l.$$

It is clear that these modules are pairwise non-isomorphic and any one-dimensional *D*-module is isomorphic to some $\mathbb{K}_{\chi_{i,j,k,l}}$, where $0 \leq i, k, l < 2, 0 \leq j < 4$.

Next we describe the simple D-modules of dimension two. For convenience, let

$$\begin{aligned} \Omega^1 &= \{ (0, j, k, l, m, n) \mid j, k, l, m, n \in \mathbb{Z}_2, m + n = 1 \}, \\ \Omega^2 &= \{ (i, j, 0, l, m, n) \mid i, j, l, m, n \in \mathbb{Z}_2, m + n = 0, j + l = 1 \}, \quad \Omega = \Omega^1 \cup \Omega^2, \\ \Lambda^1 &= \{ (1, j, k, l) \mid k \in \mathbb{Z}_4, j, l \in \mathbb{Z}_2 \}, \quad \Lambda^2 = \{ (1, j, k, l) \mid j \in \mathbb{Z}_4, k, l \in \mathbb{Z}_2 \}. \end{aligned}$$

Clearly, $|\Omega| = 24$, $|\Lambda^1| = |\Lambda^2| = 16$.

Lemma 3.5 For any 6-tuple $(i, j, k, l, m, n) \in \Omega$, there exists a simple left *D*-module $V_{i,j,k,l,m,n}$ of dimension 2 with the action on a fixed basis given by

$$[x] = \begin{pmatrix} (-1)^i & 0\\ 0 & (-1)^{i+m+n} \end{pmatrix}, \quad [y] = \begin{pmatrix} (-1)^j & 0\\ 0 & (-1)^j \end{pmatrix},$$

$$\begin{split} [t] &= \begin{pmatrix} (-1)^k & 0\\ 0 & (-1)^{j+l+k} \end{pmatrix}, \quad [a] = \begin{pmatrix} (-1)^l & 0\\ 0 & (-1)^l \end{pmatrix}, \\ [b] &= \begin{pmatrix} (-1)^m & 0\\ 0 & (-1)^n \end{pmatrix}, \quad [c] = \begin{pmatrix} (-1)^n & 0\\ 0 & (-1)^m \end{pmatrix}, \quad [d] = \begin{pmatrix} 0 & 1\\ (-1)^l & 0 \end{pmatrix}. \end{split}$$

For any 4-tuple $(i, j, k, l) \in \Lambda^1$, there exists a simple left D-module $W_{i,j,k,l}$ of dimension 2 with the action on a fixed basis given by

$$[x] = \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix}, \quad [y] = \begin{pmatrix} (-1)^j & 0 \\ 0 & (-1)^j \end{pmatrix}, \quad [t] = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$
$$[a] = \begin{pmatrix} (-1)^k & 0 \\ 0 & (-1)^k \end{pmatrix}, \qquad [b] = \begin{pmatrix} (-1)^l & 0 \\ 0 & (-1)^{l+1} \end{pmatrix},$$
$$[c] = \begin{pmatrix} (-1)^l & 0 \\ 0 & (-1)^{l+1} \end{pmatrix}, \qquad [d] = \begin{pmatrix} \xi^k & 0 \\ 0 & (-1)^{j+k} \xi^k \end{pmatrix}.$$

For any 4-tuple $(i, j, k, l) \in \Lambda^2$, there exists a simple left D-module $U_{i,j,k,l}$ of dimension 2 with the action on a fixed basis given by

$$\begin{split} [x] &= \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix}, \quad [y] = \begin{pmatrix} (-1)^j & 0 \\ 0 & (-1)^j \end{pmatrix}, \quad [t] = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ [a] &= \begin{pmatrix} (-1)^k & 0 \\ 0 & (-1)^k \end{pmatrix}, \qquad [b] = \begin{pmatrix} (-1)^l & 0 \\ 0 & (-1)^{l+1} \end{pmatrix}, \\ [c] &= \begin{pmatrix} (-1)^{l+1} & 0 \\ 0 & (-1)^l \end{pmatrix}, \qquad [d] = \begin{pmatrix} 0 & \xi^j \\ (-1)^{j+k}\xi^j & 0 \end{pmatrix}. \end{split}$$

Moreover, if V is a simple D-module of dimension 2, then V is isomorphic to $V_{i,j,k,l,m,n}$ for some $(i, j, k, l, m, n) \in \Omega$ or $W_{i,j,k,l}$ for some $(i, j, k, l) \in \Lambda_1$ or $U_{i,j,k,l}$ for some $(i, j, k, l) \in \Lambda_2$ and $V_{i,j,k,l,m,n} \cong V_{p,q,r,\kappa,\mu,\nu}$ if and only if $(i, j, k, l, m, n) = (p, q, r, \kappa, \mu, \nu)$, $W_{i,j,k,l} \cong W_{p,q,r,\kappa}$ if and only if $(i, j, k, l) = (p, q, r, \kappa)$, $U_{i,j,k,l} \cong U_{p,q,r,\kappa}$ if and only if $(i, j, k, l) = (p, q, r, \kappa)$.

Proof. Since the elements x, y, a, b, c commute each other and $x^4 = 1, y^2 = a^2 = b^2 = c^2 = 1$, the matrices defining *D*-action on *V* can be of the form

$$[x] = \begin{pmatrix} x_1 & 0 \\ 0 & x_2 \end{pmatrix}, \quad [y] = \begin{pmatrix} y_1 & 0 \\ 0 & y_2 \end{pmatrix}, \quad [t] = \begin{pmatrix} t_1 & t_2 \\ t_3 & t_4 \end{pmatrix},$$

$$[a] = \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix}, \quad [b] = \begin{pmatrix} b_1 & 0 \\ 0 & b_2 \end{pmatrix}, \quad [c] = \begin{pmatrix} c_1 & 0 \\ 0 & c_2 \end{pmatrix}, \quad [d] = \begin{pmatrix} d_1 & d_2 \\ d_3 & d_4 \end{pmatrix},$$

where $x_1^4 = x_2^4 = y_1^2 = y_2^2 = a_1^2 = a_2^2 = b_1^2 = b_2^2 = c_1^2 = c_2^2 = 1$. Since ty = yt, dy = yd, it follows that

$$t_2(y_1 - y_2) = 0,$$
 $t_3(y_1 - y_2) = 0,$ $d_2(y_1 - y_2) = 0,$ $d_3(y_1 - y_2) = 0.$

If $y_1 \neq y_2$, then $t_2 = t_3 = d_2 = d_3 = 0$ and whence V is not simple D-module, a contradiction. Thus we have $y_1 = y_2$. Similarly, the relations ta = at, da = ad give $a_1 = a_2$.

Since $t^2 = 1$, $tx = x^{-1}t$, it follows that

$$t_1^2 + t_2 t_3 = 1, \quad t_4^2 + t_2 t_3 = 1, \quad t_2(t_1 + t_4) = 0, \quad t_3(t_1 + t_4) = 0,$$
 (3.5)

$$t_1 x_1 (1 - x_1^2) = 0, \quad t_4 x_2 (1 - x_2^2) = 0, \quad t_2 (x_2 - x_1^3) = 0, \quad t_3 (x_1 - x_2^3) = 0.$$
 (3.6)

From the relations $tb = bx^2t$, $tc = cx^2t$, td = adyt, we have the relations

$$t_1b_1(1-x_1^2) = 0, \quad t_4b_2(1-x_2^2) = 0, \quad t_2(b_2-b_1x_1^2) = 0, \quad t_3(b_1-b_2x_2^2) = 0, \quad (3.7)$$

$$t_1c_1(1-x_1^2) = 0, \quad t_4c_2(1-x_2^2) = 0, \quad t_2(c_2-c_1x_1^2) = 0, \quad t_3(c_1-c_2x_2^2) = 0, \quad (3.8)$$

$$t_1c_1(1-x_1) = 0, \quad t_4c_2(1-x_2) = 0, \quad t_2(c_2-c_1x_1) = 0, \quad t_3(c_1-c_2x_2) = 0, \quad (3.8)$$

$$t_1d_1 + t_2d_3 = a_1y_1(t_1d_1 + t_3d_2), t_1d_2 + t_2d_4 = a_1y_1(t_2d_1 + t_4d_2), (3.9)$$

$$t_3d_1 + t_4d_3 = a_1y_1(t_1d_3 + t_3d_4), \qquad t_3d_2 + t_4d_4 = a_1y_1(t_2d_3 + t_4d_4).$$
(3.10)

By the relations $d^2 = a$, db = cd, dc = bd, xd = bcdx, it follows that

$$d_1^2 + d_2 d_3 = a_1, \quad d_4^2 + d_2 d_3 = a_1, \quad d_2(d_1 + d_4) = 0, \quad d_3(d_1 + d_4) = 0,$$
 (3.11)

$$d_1(b_1 - c_1) = 0, \quad d_2(b_2 - c_1) = 0, \quad d_3(b_1 - c_2) = 0, \quad d_4(b_2 - c_2) = 0,$$
 (3.12)

$$d_1(b_1 - c_1) = 0, \quad d_2(c_2 - b_1) = 0, \quad d_3(c_1 - b_2) = 0, \quad d_4(b_2 - c_2) = 0,$$
 (3.13)

$$x_1d_1(1-b_1c_1) = 0, \ x_2d_4(1-b_2c_2) = 0, \ d_2(x_1-b_1c_1x_2) = 0, \ d_3(x_2-b_2c_2x_1) = 0.$$
(3.14)

If $x_1^2 = 1$, $x_2^2 \neq 1$, then by (3.6) we have that $t_4 = 0$ and whence from (3.5) we get that $t_2t_3 = 1$. Thus by relation (3.6), it follows that $x_2 = x_1$, contradiction. Similarly, the case $x_1^2 \neq 1$, $x_2^2 = 1$ doesn't appear.

If $x_1^2 \neq 1$, $x_2^2 \neq 1$, then by (3.6) we have that $t_1 = t_4 = 0$ and whence by (3.5) we get $t_2t_3 = 1$. From the relations (3.6) – (3.10), it follows that

$$x_2 = x_1^3$$
, $b_2 = -b_1$, $c_2 = -c_1$, $d_4 = a_1 y_1 d_1$, $t_2 d_3 = a_1 y_1 t_3 d_2$.

If $d_2 = d_3 = 0$, then by relations (3.11) - (3.14), we get that

$$d_1^2 = a_1 = d_4^2, \qquad b_1 = c_1, \qquad b_2 = c_2.$$

Whence the matrices defining the action on V are of the form $W_{i,j,k,l}$, where $i \in \{1,3\}, 0 \le k < 4, 0 \le j, l < 2$. If $d_1 = d_4 = 0$, then

$$d_2 d_3 = a_1, \qquad b_2 = c_1, \qquad b_1 = c_2.$$

Whence the module is isomorphic to $U_{i,j,k,l}$, where $i \in \{1,3\}$, $0 \le j < 4$, $0 \le k$, l < 2. If $d_1d_4 \ne 0$, $d_2d_3 \ne 0$, then $b_1 = b_2$, a contradiction. Other conditions are contained in the above conditions. A direct computation shows that

when j + k = 0, $W_{i,j,k,l} \cong W_{-i,j,k,l+1}$, $U_{i,j,k,l} \cong U_{-i,j,k,l+1}$; when j + k = 1, $W_{i,j,k,l} \cong W_{-i,j,k+2,l+1}$, $U_{i,j,k,l} \cong U_{-i,j+2,k,l+1}$.

If $x_1^2 = 1$, $x_2^2 = 1$, suppose that $t_1 t_4 \neq 0$, $t_2 = t_3 = 0$, then it is clear that V is simple if and only if $d_2d_3 \neq 0$, whence by (3.11) we have that $d_1 + d_4 = 0$. By (3.5), (3.9), (3.10), (3.12), (3.14) we have that

$$t_1^2 = t_4^2 = 1, \qquad (1 - a_1 y_1) t_1 d_1 = 0, \qquad (1 - a_1 y_1) t_4 d_4 = 0,$$

$$t_4 = a_1 y_1 t_1, \qquad b_2 = c_1, \qquad b_1 = c_2, \qquad x_2 = b_2 c_2 x_1.$$

If $d_1 = d_4 = 0$, then the matrices defining the action on V are of the form $V_{i,j,k,l,m,n}$, where i, j, k, l, m, $n \in \{0,1\}$; If $d_1 \neq 0$, $d_4 \neq 0$, the module is not simple. Suppose that $t_1 = t_4 = 0$, then by (3.5)-(3.14), we get that

$$t_2t_3 = 1$$
, $x_1 = x_2$, $b_1 = b_2$, $c_1 = c_2$, $t_2d_3 = a_1y_1t_3d_2$, $d_4 = a_1y_1d_1$.

Whence the module is isomorphic to $V_{i,j,k,l,m,n}$. Similarly, for the case

$$t_1 + t_4 = 0, \quad t_1 \neq 0, \quad t_4 \neq 0, \quad t_2 \neq 0, \quad t_3 \neq 0,$$

the module is isomorphic to $V_{i,j,k,l,m,n}$. It is clear that $V_{i,j,k,l,m,n}$ is simple if and only if $m + n \neq 0$ or $j + l \neq 0$. A direct computation shows that

when m + n = 0, j + l = 1, $V_{i,j,k,l,m,n} \cong V_{i,j,k+1,l,n,m}$; when m + n = 1, j + l = 0, $V_{i,j,k,l,m,n} \cong V_{-i,j,k,l,n,m}$;

when m + n = 1, j + l = 1, $V_{i,j,k,l,m,n} \cong V_{-i,j,k+1,l,n,m}$.

Obviously, $V_{i,j,k,l,m,n}$ with $(i, j, k, l, m, n) \in \Omega$ is not isomorphic to the above two modules and $W_{i,j,k,l} \ncong U_{i',j',k',l'}$ for $(i,j,k,l) \in \Lambda^1$, $(i',j',k',l') \in \Lambda^2$.

We claim that $V_{i,j,k,l,m,n} \cong V_{i',j',k',l',m',n'}$ if and only if (i, j, k, l, m, n) = (i', j', k', l', m', n')in Ω . Assume that $\Phi: V_{i,j,k,l,m,n} \to V_{i',j',k',l',m',n'}$ is an isomorphism of *D*-modules. Denote by $[\Phi] = (p_{i,j})_{i,j=1,2}$ the matrix of Φ with respect to the given basis. Since

$$[y][\Phi] = [\Phi][y], \quad [a][\Phi] = [\Phi][a], \quad [d][\Phi] = [\Phi][d],$$

we have that j = j', l = l', $p_1 = p_4$, $p_3 = (-1)^l p_2$. Since

$$[x][\Phi] = [\Phi][x], \quad [t][\Phi] = [\Phi][t], \quad [b][\Phi] = [\Phi][b],$$

we have that

$$[(-1)^{i'} - (-1)^{i}]p_1 = 0, \qquad [(-1)^{i'} - (-1)^{i+m+n}]p_2 = 0, \qquad (3.15)$$
$$[(-1)^{i'+m'+n'} - (-1)^{i}]p_2 = 0, \qquad [(-1)^{i'+m'+n'} - (-1)^{i+m+n}]p_1 = 0, \qquad (3.16)$$

$$^{(+n)} - (-1)^{i}]p_2 = 0, \qquad [(-1)^{i'+m'+n'} - (-1)^{i+m+n}]p_1 = 0, \qquad (3.16)$$

$$[(-1)^{k'} - (-1)^{k}]p_1 = 0, \qquad [(-1)^{k'} - (-1)^{j+l+k}]p_2 = 0, \qquad (3.17)$$

$$[(-1)^{m'} - (-1)^m]p_1 = 0, \qquad [(-1)^{m'} - (-1)^n]p_2 = 0, \qquad (3.18)$$

$$[(-1)^{n'} - (-1)^m]p_2 = 0, \qquad [(-1)^{n'} - (-1)^n]p_1 = 0.$$
(3.19)

If m + n = 0, then k = k' = 0 and by (3.15), (3.16) we have that i = i', m' + n' = 0. Whence by (3.18), (3.19), it follows that m' = m, n' = n. If m + n = 1, suppose m' + n' = 0, then by (3.15), (3.16) we have that i = -i, contradiction. So m' + n' = 1. Whence i = i' = 0, $p_2 = 0$. Then by (3.17)-(3.19), k = k', m = m', n = n'. Thus the claim follows.

Similarly, we have that $W_{i,j,k,l} \cong W_{i',j',k',l'}$ if and only if (i, j, k, l) = (i', j', k', l') in Λ^1 ; $U_{i,j,k,l} \cong U_{i',j',k',l'}$ if and only if (i, j, k, l) = (i', j', k', l') in Λ^2 .

Theorem 3.6 There are 88 simple left D-modules up to isomorphism, among which 32 onedimensional objects given by Lemma 3.4 and 56 two-dimensional objects given by Lemma 3.5.

Proof. Assume there is a simple module of dimension d > 2 and let n denote the number of simple d-dimensional modules pairwise non-isomorphic. By Lemmas 3.4 & 3.5, we have

$$32 \cdot 1^2 + 56 \cdot 2^2 + nd^2 = 256 + nd^2 \le \dim(D^*)_0 = \dim D^* = 256.$$

Then n = 0.

4 The category ${}^{H}_{H}\mathcal{YD}$

In this section, by using the monoidal isomorphism ${}^{H}_{H}\mathcal{YD} \cong {}_{D}\mathcal{M}$, we will describe the simple objects in ${}^{H}_{H}\mathcal{YD}$ and determine their braidings.

Proposition 4.1 Let $\mathbb{k}_{\chi_{i,j,k,l}} = \mathbb{k}\{v\}$ be a one-dimensional *D*-module with $i, k, l \in \mathbb{I}_{0,1}, j \in \mathbb{I}_{0,3}$. Then $\mathbb{k}_{\chi_{i,j,k,l}} \in {}^{H}_{H}\mathcal{YD}$ with its module and comodule structure given by

$$x \cdot v = (-1)^i v, \quad y \cdot v = (-1)^j v, \quad t \cdot v = (-1)^k v, \quad \delta(v) = x^j y^l \otimes v.$$

Proof. The action is given by the restriction of the action given in Lemma 3.4. Since $\mathbb{k}_{\chi_{i,j,k,l}}$ is one-dimensional, we must have that $\delta(v) = h \otimes v$ with $h \in G(H) = \langle x \rangle \times \langle y \rangle$. As $f \cdot v = \langle f, h \rangle v$ for all $f \in H^*$, it follows that $\delta(v) = x^j y^l \otimes v$.

Proposition 4.2 The braiding of the one-dimensional Yetter-Drinfeld module $\mathbb{k}_{\chi_{i,j,k,l}}$ is $c(v \otimes v) = (-1)^{(i+l)j} v \otimes v.$

Proof. By formula (2.1) and Proposition 4.1, we have that

$$c(v \otimes v) = x^{j}y^{l} \cdot v \otimes v = (-1)^{(i+l)j}v \otimes v.$$

This completes the proof.

Proposition 4.3 Let $V_{i,j,k,l,m,n} = \mathbb{k}\{v_1, v_2\}$ be a two-dimensional simple *D*-module with $(i, j, k, l, m, n) \in \Omega$, then $V_{i,j,k,l,m,n} \in {}^{H}_{H}\mathcal{YD}$ with its action given by

$$\begin{aligned} x \cdot v_1 &= (-1)^i v_1, \qquad y \cdot v_1 &= (-1)^j v_1, \qquad t \cdot v_1 &= (-1)^k v_1, \\ x \cdot v_2 &= (-1)^{i+m+n} v_2, \qquad y \cdot v_2 &= (-1)^j v_2, \qquad t \cdot v_2 &= (-1)^{j+l+k} v_2, \end{aligned}$$

and its coaction by

$$\begin{aligned} 1. \ for \ l &= m = n = 0, \\ \delta(v_1) &= \frac{1}{2}(1+x^2) \otimes v_1 + \frac{1}{2}(1-x^2) \otimes v_2, \quad \delta(v_2) = \frac{1}{2}(1+x^2) \otimes v_2 + \frac{1}{2}(1-x^2) \otimes v_1; \\ 2. \ for \ l &= 0, m = n = 1, \\ \delta(v_1) &= \frac{1}{2}(1+x^2)y \otimes v_1 + \frac{1}{2}(1-x^2)y \otimes v_2, \ \delta(v_2) &= \frac{1}{2}(1+x^2)y \otimes v_2 + \frac{1}{2}(1-x^2)y \otimes v_1; \\ 3. \ for \ l &= m = 0, n = 1, \\ \delta(v_1) &= \frac{1}{2}(1+x^2)t \otimes v_1 + \frac{1}{2}(1-x^2)yt \otimes v_2, \ \delta(v_2) &= \frac{1}{2}(1+x^2)yt \otimes v_2 + \frac{1}{2}(1-x^2)t \otimes v_1; \\ 4. \ for \ l &= 0, m = 1, n = 0, \\ \delta(v_1) &= \frac{1}{2}(1+x^2)yt \otimes v_1 + \frac{1}{2}(1-x^2)t \otimes v_2, \ \delta(v_2) &= \frac{1}{2}(1+x^2)t \otimes v_2 + \frac{1}{2}(1-x^2)yt \otimes v_1; \\ 5. \ for \ l &= 1, m = n = 0, \\ \delta(v_1) &= \frac{1}{2}x(1+x^2) \otimes v_1 + \frac{1}{2}\xi x(1-x^2) \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}x(1+x^2) \otimes v_2 - \frac{1}{2}\xi x(1-x^2) \otimes v_1; \\ 6. \ for \ l &= m = n = 1, \\ \delta(v_1) &= \frac{1}{2}xy(1+x^2) \otimes v_1 + \frac{1}{2}\xi xy(1-x^2) \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}xy(1+x^2) \otimes v_2 - \frac{1}{2}\xi xy(1-x^2) \otimes v_1; \\ 7. \ for \ l &= 1, m = 0, n = 1, \\ \delta(v_1) &= \frac{1}{2}xt(1+x^2) \otimes v_1 + \frac{1}{2}\xi xyt(1-x^2) \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}xyt(1+x^2) \otimes v_2 - \frac{1}{2}\xi xy(1-x^2) \otimes v_1; \\ 8. \ for \ l &= m = 1, n = 0, \\ \delta(v_1) &= \frac{1}{2}xyt(1+x^2) \otimes v_1 + \frac{1}{2}\xi xt(1-x^2) \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}xyt(1+x^2) \otimes v_2 - \frac{1}{2}\xi xyt(1-x^2) \otimes v_1; \\ 8. \ for \ l &= m = 1, n = 0, \\ \delta(v_1) &= \frac{1}{2}xyt(1+x^2) \otimes v_1 + \frac{1}{2}\xi xt(1-x^2) \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}xyt(1+x^2) \otimes v_2 - \frac{1}{2}\xi xyt(1-x^2) \otimes v_1; \\ 8. \ for \ l &= m = 1, n = 0, \\ \delta(v_1) &= \frac{1}{2}xyt(1+x^2) \otimes v_1 + \frac{1}{2}\xi xt(1-x^2) \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}xt(1+x^2) \otimes v_2 - \frac{1}{2}\xi xyt(1-x^2) \otimes v_1. \end{aligned}$$

Proof. We just need to describe the coaction. Denote by $\{h_i\}_{1 \le i \le 16}$ and $\{h^i\}_{1 \le i \le 16}$ a basis of H and its dual basis, respectively. Then the comodule structure is given by $\delta(v) = \sum_{i=1}^{16} h_i \otimes h^i \cdot v$, it follows that

$$\begin{split} \delta(v_1) &= \sum_{i=0}^{1} \sum_{j=0}^{1} \sum_{k=0}^{1} \sum_{l=0}^{1} (a^i b^j c^k d^l)^* \otimes a^i b^j c^k d^l \cdot v_1 \\ &= [1^* + (-1)^{la}^* + (-1)^m b^* + (-1)^n c^* + (-1)^{m+l} (ab)^* + (-1)^{l+n} (ac)^* + (-1)^{m+n} (bc)^* \\ &+ (-1)^{l+m+n} (abc)^*] \otimes v_1 \\ &+ [(-1)^l d^* + (ad)^* + (-1)^{m+l} (bd)^* + (-1)^{n+l} (cd)^* + (-1)^m (abd)^* + (-1)^n (acd)^* \\ &+ (-1)^{m+n+l} (bcd)^* + (-1)^{m+n} (abcd)^*] \otimes v_2, \\ \delta(v_2) &= \sum_{i=0}^{1} \sum_{j=0}^{1} \sum_{k=0}^{1} \sum_{l=0}^{1} (a^i b^j c^k d^l)^* \otimes a^i b^j c^k d^l \cdot v_2 \\ &= [1^* + (-1)^l a^* + (-1)^n b^* + (-1)^m c^* + (-1)^{n+l} (ab)^* + (-1)^{l+m} (ac)^* + (-1)^{m+n} (bc)^* \\ &+ (-1)^{l+m+n} (abc)^*] \otimes v_2 \\ &+ [d^* + (-1)^l (ad)^* + (-1)^n (bd)^* + (-1)^m (cd)^* + (-1)^{n+l} (abd)^* + (-1)^{m+l} (acd)^* \end{split}$$

$$+(-1)^{m+n}(bcd)^*+(-1)^{m+n+l}(abcd)^*]\otimes v_1$$

By discussing the possible situations, the claim follows.

Using the same method, we have the following propositions.

Proposition 4.4 Let $W_{i,j,k,l} = \mathbb{k}\{v_1, v_2\}$ be a two-dimensional simple *D*-module with $(i, j, k, l) \in \Lambda^1$, then $W_{i,j,k,l} \in {}^H_H \mathcal{YD}$ with its action given by

$$\begin{aligned} x \cdot v_1 &= \xi v_1, \qquad y \cdot v_1 = (-1)^j v_1, \qquad t \cdot v_1 = v_2, \\ x \cdot v_2 &= \xi^{-1} v_2, \qquad y \cdot v_2 = (-1)^j v_2, \qquad t \cdot v_2 = v_1, \end{aligned}$$

and its coaction by $\delta(v_1) &= x^k y^l \otimes v_1,$
1. for $(k, j, l) = (0, 0, 0), \quad \delta(v_2) = y \otimes v_2; \ (k, j, l) = (2, 0, 0), \quad \delta(v_2) = x^2 y \otimes v_2; \end{aligned}$
2. for $(k, j, l) = (0, 1, 0), \quad \delta(v_2) = x^2 y \otimes v_2; \ (k, j, l) = (2, 1, 0), \quad \delta(v_2) = y \otimes v_2; \end{aligned}$
3. for $(k, j, l) = (0, 0, 1), \quad \delta(v_2) = 1 \otimes v_2; \ (k, j, l) = (2, 0, 1), \quad \delta(v_2) = x^2 \otimes v_2; \end{aligned}$
4. for $(k, j, l) = (0, 1, 1), \quad \delta(v_2) = x^2 \otimes v_2; \ (k, j, l) = (2, 1, 1), \quad \delta(v_2) = 1 \otimes v_2; \end{aligned}$
5. for $(k, j, l) = (1, 0, 0), \quad \delta(v_2) = x^3 y \otimes v_2; \ (k, j, l) = (3, 0, 0), \quad \delta(v_2) = xy \otimes v_2; \end{aligned}$
6. for $(k, j, l) = (1, 0, 1), \quad \delta(v_2) = x^3 \otimes v_2; \ (k, j, l) = (3, 0, 1), \quad \delta(v_2) = x \otimes v_2; \end{aligned}$
7. for $(k, j, l) = (1, 0, 1), \quad \delta(v_2) = x \otimes v_2; \ (k, j, l) = (3, 0, 1), \quad \delta(v_2) = x \otimes v_2; \end{aligned}$
8. for $(k, j, l) = (1, 1, 1), \quad \delta(v_2) = x \otimes v_2; \ (k, j, l) = (3, 1, 1), \quad \delta(v_2) = x^3 \otimes v_2. \end{aligned}$

Proposition 4.5 Let $U_{i,j,k,l} = \mathbb{k}\{v_1, v_2\}$ be a two-dimensional simple *D*-module with $(i, j, k, l) \in \Lambda_2$, then $U_{i,j,k,l} \in {}^H_H \mathcal{YD}$ with its action same as $W_{i,j,k,l}$ and its coaction by 1. for (j, l, k) = (0, 0, 0),

 $\delta(v_1) = \frac{1}{2}(1+x^2)t \otimes v_1 + \frac{1}{2}(1-x^2)yt \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)yt \otimes v_2 + \frac{1}{2}(1-x^2)t \otimes v_1;$ for (j, l, k) = (2, 0, 0), $\delta(v_1) = \frac{1}{2}(1+x^2)t \otimes v_1 - \frac{1}{2}(1-x^2)yt \otimes v_2, \quad \delta(v_2) = \frac{1}{2}(1+x^2)yt \otimes v_2 - \frac{1}{2}(1-x^2)t \otimes v_1;$ 2. for (j, l, k) = (0, 1, 0), $\delta(v_1) = \frac{1}{2}(1+x^2)yt \otimes v_1 + \frac{1}{2}(1-x^2)t \otimes v_2, \quad \delta(v_2) = \frac{1}{2}(1+x^2)t \otimes v_2 + \frac{1}{2}(1-x^2)yt \otimes v_1;$ for (j, l, k) = (2, 1, 0), $\delta(v_1) = \frac{1}{2}(1+x^2)yt \otimes v_1 - \frac{1}{2}(1-x^2)t \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)t \otimes v_2 - \frac{1}{2}(1-x^2)yt \otimes v_1;$ 3. for (j, l, k) = (0, 0, 1), $\delta(v_1) = \frac{1}{2}(1+x^2)xt \otimes v_1 + \frac{\xi}{2}(1-x^2)xyt \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)xyt \otimes v_2 - \frac{\xi}{2}(1-x^2)xt \otimes v_1;$ for (j, l, k) = (2, 0, 1). $\delta(v_1) = \frac{1}{2}(1+x^2)xt \otimes v_1 - \frac{\xi}{2}(1-x^2)xyt \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)xyt \otimes v_2 + \frac{\xi}{2}(1-x^2)xt \otimes v_1;$ 4. for (j, l, k) = (0, 1, 1), $\delta(v_1) = \frac{1}{2}(1+x^2)xyt \otimes v_1 + \frac{\xi}{2}(1-x^2)xt \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)xt \otimes v_2 - \frac{\xi}{2}(1-x^2)xyt \otimes v_1;$ for (j, l, k) = (2, 1, 1), $\delta(v_1) = \frac{1}{2}(1+x^2)xyt \otimes v_1 - \frac{\xi}{2}(1-x^2)xt \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)xt \otimes v_2 + \frac{\xi}{2}(1-x^2)xyt \otimes v_1;$ 5. for (j, l, k) = (1, 0, 0). $\delta(v_1) = \frac{1}{2}(1+x^2)t \otimes v_1 - \frac{\xi}{2}(1-x^2)yt \otimes v_2, \\ \delta(v_2) = \frac{1}{2}(1+x^2)yt \otimes v_2 + \frac{\xi}{2}(1-x^2)t \otimes v_1;$

for (j, l, k) = (3, 0, 0),

$$\begin{split} \delta(v_1) &= \frac{1}{2}(1+x^2)t \otimes v_1 + \frac{\xi}{2}(1-x^2)yt \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)yt \otimes v_2 - \frac{\xi}{2}(1-x^2)t \otimes v_1; \\ 6. \ for \ (j,l,k) &= (1,1,0), \\ \delta(v_1) &= \frac{1}{2}(1+x^2)yt \otimes v_1 - \frac{\xi}{2}(1-x^2)t \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)t \otimes v_2 + \frac{\xi}{2}(1-x^2)yt \otimes v_1; \\ for \ (j,l,k) &= (3,1,0), \\ \delta(v_1) &= \frac{1}{2}(1+x^2)yt \otimes v_1 + \frac{\xi}{2}(1-x^2)t \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)t \otimes v_2 - \frac{\xi}{2}(1-x^2)yt \otimes v_1; \\ 7. \ for \ (j,l,k) &= (1,0,1), \\ \delta(v_1) &= \frac{1}{2}(1+x^2)xt \otimes v_1 + \frac{1}{2}(1-x^2)xyt \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)xyt \otimes v_2 + \frac{1}{2}(1-x^2)xt \otimes v_1; \\ for \ (j,l,k) &= (3,0,1), \\ \delta(v_1) &= \frac{1}{2}(1+x^2)xt \otimes v_1 - \frac{1}{2}(1-x^2)xyt \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)xyt \otimes v_2 - \frac{1}{2}(1-x^2)xt \otimes v_1; \\ 8. \ for \ (j,l,k) &= (1,1,1), \\ \delta(v_1) &= \frac{1}{2}(1+x^2)xyt \otimes v_1 + \frac{1}{2}(1-x^2)xt \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)xt \otimes v_2 + \frac{1}{2}(1-x^2)xyt \otimes v_1; \\ for \ (j,l,k) &= (3,1,1), \\ \delta(v_1) &= \frac{1}{2}(1+x^2)xyt \otimes v_1 - \frac{1}{2}(1-x^2)xt \otimes v_2, \\ \delta(v_2) &= \frac{1}{2}(1+x^2)xt \otimes v_2 - \frac{1}{2}(1-x^2)xyt \otimes v_1. \\ \end{cases}$$

5 Nichols algebras in ${}^{H}_{H}\mathcal{YD}$

Let $V = \bigoplus_{i \in I} V_i$, where V_i is a simple object in ${}^{H}_{H}\mathcal{YD}$. In this section we will try to determine all the finite-dimensional Nichols algebras satisfying $\mathcal{B}(V) \cong \bigotimes_{i \in I} \mathcal{B}(V_i)$. We begin by studying the Nichols algebras of simple Yetter-Drinfeld modules.

Lemma 5.1 Let $(i, j, k, l) \in \mathbb{I}_{0,1} \times \mathbb{I}_{0,3} \times \mathbb{I}_{0,1} \times \mathbb{I}_{0,1}$. The Nichols algebras $\mathcal{B}(\mathbb{k}_{\chi_{i,j,k,l}})$ associated to $\mathbb{k}_{\chi_{i,j,k,l}} = \mathbb{k}v$ are

$$\mathcal{B}(\mathbb{k}_{\chi_{i,j,k,l}}) = \begin{cases} \mathbb{k}[v], & \text{if } (i+l)j = 0, \ 2, \\ \mathbb{k}[v]/(v^2) = \bigwedge \mathbb{k}_{\chi_{i,j,k,l}}, & \text{if } (i+l)j = 1, \ 3. \end{cases}$$

Proof. The claim follows by Proposition 4.2.

Remark 5.2 $\mathcal{B}(\mathbb{k}_{\chi_{i,j,k,l}})$ is finite-dimensional if and only if $\mathbb{k}_{\chi_{i,j,k,l}}$ is isomorphic either to

$$\begin{split} V_1 &:= \mathbb{k}_{\chi_{1,1,0,0}}, \quad V_2 &:= \mathbb{k}_{\chi_{1,1,1,0}}, \quad V_3 &:= \mathbb{k}_{\chi_{0,1,0,1}}, \quad V_4 &:= \mathbb{k}_{\chi_{0,1,1,1}}, \\ V_5 &:= \mathbb{k}_{\chi_{1,3,0,0}}, \quad V_6 &:= \mathbb{k}_{\chi_{1,3,1,0}}, \quad V_7 &:= \mathbb{k}_{\chi_{0,3,0,1}}, \quad V_8 &:= \mathbb{k}_{\chi_{0,3,1,1}}. \end{split}$$

For the convenience of our statements, we let

$$\begin{split} M_1 &= V_{0,1,0,0,1,1}, \quad M_2 = V_{1,1,0,0,1,1}, \quad M_3 = V_{0,0,1,0,0,1}, \quad M_4 = V_{0,1,1,0,0,1}, \\ M_5 &= V_{0,0,1,0,1,0}, \quad M_6 = V_{0,1,0,0,1,0}, \quad M_7 = V_{1,0,0,1,0,0}, \quad M_8 = V_{1,0,0,1,1,1}, \\ M_9 &= V_{0,1,1,1,0,1}, \quad M_{10} = V_{0,0,1,1,0,1}, \quad M_{11} = V_{0,1,0,1,1,0}, \quad M_{12} = V_{0,0,1,1,1,0}, \\ M_{13} &= W_{1,1,0,1}, \quad M_{14} = W_{1,1,2,0}, \quad M_{15} = W_{1,0,2,0}, \quad M_{16} = W_{1,0,2,1}, \\ M_{17} &= U_{1,2,0,0}, \quad M_{18} = U_{1,2,0,1}, \quad M_{19} = U_{1,0,1,0}, \quad M_{20} = U_{1,0,1,1}. \end{split}$$

Lemma 5.3 Let $V_{i,j,k,l,m,n} \in {}^{H}_{H}\mathcal{YD}$ with $(i, j, k, l, m, n) \in \Omega$. Then

(1) dim $\mathcal{B}(V_{i,j,k,l,m,n}) = \infty$, for $V_{i,j,k,l,m,n} \notin \{M_1, M_2, \dots, M_{12}\}.$

(2) $\forall V \in \{M_1, M_2, \dots, M_{12}\}, \dim \mathcal{B}(V) = 4 \text{ and relations of } \mathcal{B}(M_4), \mathcal{B}(M_6), \mathcal{B}(M_9), \mathcal{B}(M_{11}) \text{ are given by } v_1^2 = 0, v_2^2 = 0, v_1v_2 - v_2v_1 = 0, \text{ others are given by } v_1^2 = 0, v_2^2 = 0, v_1v_2 + v_2v_1 = 0.$

Proof. If $V_{i,j,k,l,m,n} \notin \{M_1, M_2, \dots, M_{12}\}$, then the braiding has an eigenvector $v_1 \otimes v_1$ of eigenvalue 1, hence (1) follows by Remark 2.2. For (2), the braiding is of type $A_1 \times A_1$, then we can get the relations of Nichols algebras.

Using the same method, we have the following Lemmas.

Lemma 5.4 Let $W_{i,j,k,l} \in {}^{H}_{H}\mathcal{YD}$ with $(i, j, k, l) \in \Lambda^{1}$. Then

(1) dim $\mathcal{B}(W_{i,j,k,l}) = \infty$, for $W_{i,j,k,l,m,n} \notin \{M_{13}, M_{14}, M_{15}, M_{16}\}$.

(2) $\forall V \in \{M_{13}, M_{14}, M_{15}, M_{16}\}, \dim \mathcal{B}(V) = 4 \text{ and the relations are given by } v_1^2 = 0, v_2^2 = 0, v_1v_2 + v_2v_1 = 0.$

Lemma 5.5 Let $U_{i,j,k,l} \in {}^{H}_{H}\mathcal{YD}$ with $(i, j, k, l) \in \Lambda^{2}$. Then

(1) dim $\mathcal{B}(U_{i,j,k,l}) = \infty$, for $U_{i,j,k,l} \notin \{M_{17}, M_{18}, M_{19}, M_{20}\}$.

(2) $\forall V \in \{M_{17}, M_{18}, M_{19}, M_{20}\}$, dim $\mathcal{B}(V) = 4$ and the relations of $\mathcal{B}(M_{17})$, $\mathcal{B}(M_{18})$ are given by $v_1v_2 = 0$, $v_2v_1 = 0$, $v_1^2 + v_2^2 = 0$, the relations of $\mathcal{B}(M_{19})$, $\mathcal{B}(M_{20})$ are given by $v_1v_2 = 0$, $v_2v_1 = 0$, $v_1^2 - v_2^2 = 0$.

Proof. (1) is similar to Proposition 5.3 (1).

(2) For $V \in \{M_{17}, M_{18}, M_{19}, M_{20}\}$, it belongs to the case $\mathfrak{R}_{1,4}$ in [8]. Then the assertion follows by [8, Proposition 3.15].

Proposition 5.6 (1) Let V, W be simple objects in ${}^{H}_{H}\mathcal{YD}$. Finite-dimensional $\mathcal{B}(V \oplus W) \cong \mathcal{B}(V) \otimes \mathcal{B}(W)$ for the following cases

(a) $V, W \in \{V_1, V_2, \dots, V_8\};$ (b) $V \in \{V_3, V_4, V_7, V_8\}, W \in \{M_1, M_8\};$ (c) $V \in \{V_1, V_2, V_5, V_6\}, W \in \{M_2, M_7\};$ (d) $V \in \{V_1, V_3, V_5, V_7\}, W \in \{M_3, M_{11}\};$ (e) $V \in \{V_1, V_4, V_5, V_8\}, W \in \{M_4, M_{12}\};$ (f) $V \in \{V_2, V_4, V_6, V_8\}, W \in \{M_5, M_9\};$ (g) $V \in \{V_2, V_3, V_6, V_7\}, W \in \{M_6, M_{10}\};$ (h) $V = M_1, W \in \{M_2, M_7\}; V = M_3 \text{ or } M_{11}, W \in \{M_5, M_9\};$ (i) $V = M_4 \text{ or } M_{12}, W \in \{M_6, M_{10}\};$ (j) $(V, W) \in \{(M_2, M_8), (M_7, M_8), (M_{13}, M_{14}), (M_{15}, M_{16}), (M_{17}, M_{18}), (M_{19}, M_{20})\};$ (k) $V = W = M_i \text{ for } i \in \mathbb{I}_{1,20}.$ (2) For $W \in \{M_{13}, M_{14}, M_{15}, M_{16}\}, V \in \{V_1, \dots, V_8\}, \dim \mathcal{B}(V \oplus W) = \infty.$ (3) For $W \in \{M_{17}, M_{18}, M_{19}, M_{20}\}, V \in \{V_1, \dots, V_8\}, \dim \mathcal{B}(V \oplus W) = \infty.$ *Proof.* The part (1) is a direct result of Lemma 2.3.

(2) Let $M_3 = \mathbb{k}\{v_1, v_2\}, \mathbb{k}p = \mathbb{k}_{\chi_{i,j,k,l}} \in \{V_1, \dots, V_8\}$, then

$$c(p \otimes v_1) = (-1)^l \xi^j v_1 \otimes p, \qquad c(p \otimes v_2) = -(-1)^l \xi^j v_2 \otimes p,$$

$$c(v_1 \otimes p) = -p \otimes v_1, \qquad c(v_2 \otimes p) = p \otimes v_2.$$

The generalized Dynkin diagram associated to the braiding is given by

$$\underbrace{\begin{array}{ccc} -1 & (-1)^{l+1}\xi^j & -1 & (-1)^{l+1}\xi^j & -1 \\ \widetilde{v_1} & \widetilde{p} & \widetilde{v_2} \end{array}}_{p}$$

Then by [22] we know that the Nichols algebra $\mathcal{B}(\mathbb{k}_{\chi_{i,j,k,l}} \oplus M_{13})$ is infinite-dimensional. The proofs for M_{14} , M_{15} , M_{16} are completely analogous.

(3) Let $M_{17} = \mathbb{k}\{v_1, v_2\}, \mathbb{k}p = \mathbb{k}_{\chi_{i,j,k,l}} \in \{V_1, \dots, V_8\}$, then

$$(id - c^2)(p \otimes v_1) = [1 - (-1)^k \xi^j] p \otimes v_1, \quad (id - c^2)(p \otimes v_2) = [1 - (-1)^k \xi^j] p \otimes v_2.$$

Thus $ad(\mathbb{k}_{\chi_{i,j,k,l}})(M_{17}) = (id - c^2)(\mathbb{k}_{\chi_{i,j,k,l}} \otimes M_{17}) \cong \mathbb{k}_{\chi_{i,j,k,l}} \otimes M_{17}$. Let $u = p \otimes v_1$, $v = p \otimes v_2$, then

$$\begin{aligned} x \cdot u &= (-1)^{i} \xi u, \qquad y \cdot u = -u, \qquad t \cdot u = -(-1)^{k} v, \\ x \cdot v &= -(-1)^{i} \xi v, \qquad y \cdot v = -v, \qquad t \cdot v = -(-1)^{k} u, \\ \delta(u) &= \frac{1}{2} x^{j} y^{l} (1 + x^{2}) t \otimes u - \frac{1}{2} x^{j} y^{l} (1 - x^{2}) y t \otimes v, \\ \delta(v) &= \frac{1}{2} x^{j} y^{l} (1 + x^{2}) y t \otimes v - \frac{1}{2} x^{j} y^{l} (1 - x^{2}) t \otimes u. \end{aligned}$$

Whence by Lemma 5.5, we have that $\dim \mathcal{B}(ad(\Bbbk_{\chi_{i,j,k,l}})(M_{17})) = \infty$. As $ad(\Bbbk_{\chi_{i,j,k,l}})(M_{17})$ is the Yetter-Drinfeld submodule of $\mathcal{B}(\Bbbk_{\chi_{i,j,k,l}} \oplus M_{17})$ [23], then $\dim \mathcal{B}(\Bbbk_{\chi_{i,j,k,l}} \oplus M_{17}) = \infty$. The proofs for M_{18} , M_{19} , M_{20} are completely analogous.

Proof of Theorem A. The claim follows by Remark 5.2, Lemmas 5.3, 5.4, 5.5 and Proposition 5.6.

6 Hopf algebras over *H*

In this section, based on the principle of the lifting method, we will determine finitedimensional Hopf algebras over H such that their infinitesimal braidings are those Yetter-Drinfeld modules listed in Theorem A. We first show that the diagrams of these Hopf algebras are Nichols algebras.

Theorem 6.1 Let A be a finite-dimensional Hopf algebra over H such that its infinitesimal braiding is isomorphic to a Yetter-Drinfeld module V in Theorem A. Then $gr(A) \cong \mathcal{B}(V) \sharp H$.

Proof. Let S be the graded dual of the diagram R of A. By [12, Lemma 2.4], S is generated by W = S(1). Since $R(0) = \mathbb{k}$, $V := R(1) = \mathcal{P}(R)$, there exists an epimorphism $S \to \mathcal{B}(W)$. We know that R is a Nichols algebra if and only if $\mathcal{P}(S) = S(1)$, that is, S is also a Nichols algebra. It is enough to show that the relations of $\mathcal{B}(W)$ also hold in S. If V is a simple object in ${}^{H}_{H}\mathcal{YD}$, then W must be simple as an object in ${}^{H}_{H}\mathcal{YD}$.

Assume $W = \Omega_2(n_1, n_2, n_3, n_4)$. Then W is generated by

$$p_1, p_2, \{C_k\}_{k=1,\dots,n_1}, \{D_\ell\}_{\ell=1,\dots,n_2}, \{G_s\}_{s=1,\dots,n_3}, \{H_r\}_{r=1,\dots,n_4}$$

with $M_1 = \mathbb{k}\{p_1, p_2\}, \mathbb{k}C_k \simeq V_3, \mathbb{k}D_\ell \simeq V_4, \mathbb{k}G_s \simeq V_7, \mathbb{k}H_r \simeq V_8$, and the defining ideal of the Nichols algebra $\mathfrak{B}(W)$ is generated by the elements

$$\begin{array}{ll} p_1^2, & p_2^2, & p_1p_2 + p_2p_1, & \{C_{k_1}C_{k_2} + C_{k_2}C_{k_1}\}_{1 \leq k_1, k_2 \leq n_1}, & \{D_{\ell_1}D_{\ell_2} + D_{\ell_2}D_{\ell_1}\}_{1 \leq \ell_1, \ell_2 \leq n_2}, \\ & \{G_{s_1}G_{s_2} + G_{s_2}G_{s_1}\}_{1 \leq s_1, s_2 \leq n_3}, & \{H_{r_1}H_{r_2} + H_{r_2}H_{r_1}\}_{1 \leq r_1, r_2 \leq n_4}, & C_kD_\ell + D_\ell C_k, \\ & C_kG_s + G_sC_k, & C_kH_r + H_rC_k, & D_\ell G_s + G_sD_\ell, & D_\ell H_r + H_rD_\ell, & G_sH_r + H_rG_s, \\ & p_1C_k + C_kp_1, & p_2C_k + C_kp_2, & p_1D_\ell + D_\ell p_1, & p_2D_\ell + D_\ell p_2, & p_1G_s + G_sp_1, \\ & p_2G_s + G_sp_2, & p_1H_r + H_rp_1, & p_2H_r + H_rp_2. \end{array}$$

Since S is finite-dimensional, it is enough to show that $c(r \otimes r) = r \otimes r$ for all generators given in above for the defining ideal. By [18, Theorem 6], all those generators are primitive elements. As $\delta(C_k) = xy \otimes C_k$, $\delta(D_l) = xy \otimes D_l$,

$$\delta(p_1) = \frac{1}{2}(1+x^2)y \otimes p_1 + \frac{1}{2}(1-x^2)y \otimes p_2,$$

$$\delta(p_2) = \frac{1}{2}(1+x^2)y \otimes p_2 + \frac{1}{2}(1-x^2)y \otimes p_1,$$

we have that

$$\begin{split} \delta(p_1^2) &= \frac{1}{2}(1+x^2) \otimes p_1^2 + \frac{1}{2}(1-x^2) \otimes p_2^2, \\ \delta(p_2^2) &= \frac{1}{2}(1+x^2) \otimes p_2^2 + \frac{1}{2}(1-x^2) \otimes p_1^2, \\ \delta(C_k D_l + D_l C_k) &= x^2 \otimes (C_k D_l + D_l C_k), \\ \delta(p_1 p_2 + p_2 p_1) &= \frac{1}{2}(1+x^2) \otimes (p_1 p_2 + p_2 p_1) + \frac{1}{2}(1-x^2) \otimes (p_1 p_2 + p_2 p_1), \\ \delta(p_1 C_k + C_k p_1) &= \frac{1}{2}x(1+x^2) \otimes (p_1 C_k + C_k p_1) + \frac{1}{2}x(1-x^2) \otimes (p_2 C_k + C_k p_2). \end{split}$$

Then by the definition of the braiding in ${}^{H}_{H}\mathcal{YD}$, the claim follows. We leave the rest to the reader.

Lemma 6.2 [11, Lemma 6.1] Let H be a Hopf algebra, $\psi : H \to H$ an automorphism of Hopf algebra, and V, W Yetter-Drinfeld modules over H.

(1) Let V^{ψ} be the same space as that of the underlying V but with action and coaction

$$h \cdot_{\psi} v = \psi(h) \cdot v, \quad \delta^{\psi}(v) = (\psi^{-1} \otimes id)\delta(v), \quad h \in H, \ v \in V.$$

Then V^{ψ} is also a Yetter-Drinfeld module over H. If $T: V \to W$ is a morphism in ${}^{H}_{H}\mathcal{YD}$, so is $T^{\psi}: V^{\psi} \to W^{\psi}$. Moreover, the braiding $c: V^{\psi} \otimes W^{\psi} \to W^{\psi} \otimes V^{\psi}$ coincides with the braiding $c: V \otimes W \to W \otimes V$.

(2) If R is an algebra (resp., a coalgebra, a Hopf algebra) in ${}^{H}_{H}\mathcal{YD}$, so is R^{ψ} , with the same structural maps.

(3) Let R be a Hopf algebra in ${}^{H}_{H}\mathcal{YD}$. Then the map $\varphi : R^{\psi}\sharp H \to R\sharp H$ given by $\varphi(r\sharp h) = r\sharp\psi(h)$ is an isomorphism of Hopf algebras.

Corollary 6.3 (1) $V_1^{\tau_{17}} \simeq V_3$, $V_2^{\tau_{17}} \simeq V_4$, $V_5^{\tau_{17}} \simeq V_7$, $V_6^{\tau_{17}} \simeq V_8$, $M_2^{\tau_{17}} \simeq M_1$, $M_7^{\tau_{17}} \simeq M_8$. (2) $V_1^{\tau_5} \simeq V_2$, $V_3^{\tau_5} \simeq V_4$, $V_5^{\tau_5} \simeq V_6$, $V_7^{\tau_5} \simeq V_8$, $M_3^{\tau_5} \simeq M_5$, $M_{11}^{\tau_5} \simeq M_9$, $M_4^{\tau_5} \simeq M_6$, $M_{12}^{\tau_5} \simeq M_{10}$, $M_{17}^{\tau_5} \simeq M_{18}$, $M_{13}^{\tau_9} \simeq M_{14}$, $M_{15}^{\tau_9} \simeq M_{16}$.

(3) $M_3^{\tau_2} \simeq M_{10}, \ M_4^{\tau_2} \simeq M_9, \ M_5^{\tau_2} \simeq M_{12}, \ M_6^{\tau_2} \simeq M_{11}, \ M_{17}^{\tau_2} \simeq M_{19}, \ M_{18}^{\tau_2} \simeq M_{20}, \ V_1^{\tau_2} \simeq V_2, V_5^{\tau_2} \simeq V_6, \ V_i^{\tau_2} \simeq V_i \ for \ i \in \{3, 4, 7, 8\}.$

(4) $\mathcal{B}(\Omega_i(n_1, n_2, n_3, n_4)) \sharp H \simeq \mathcal{B}(\Omega_j(n_1, n_2, n_3, n_4)) \sharp H$ for

 $(i, j) \in \{(2, 3), (4, 6), (5, 7), (8, 9), (10, 12), (11, 13), (4, 11), (5, 10)\}.$

(5) $\mathcal{B}(\Omega_i) \sharp H \simeq \mathcal{B}(\Omega_j) \sharp H$ for

$$\begin{aligned} (i,j) \in \{(14,30), (16,31), (19,32), (17,33), (19,34), (20,35), (32,36), \\ (22,37), (20,38), (21,39), (17,40), (18,41), (38,42), (40,43), \\ (24,44), (26,45), (28,46), (28,47), (29,48), (46,49)\}. \end{aligned}$$

Let $i, j, k, \ell, m, q, s, r \in \mathbb{N}^*$, denote

$$\begin{split} & \mathbb{k} A_i \simeq V_1, \quad \mathbb{k} B_j \simeq V_2, \quad \mathbb{k} C_k \simeq V_3, \quad \mathbb{k} D_\ell \simeq V_4, \\ & \mathbb{k} E_m \simeq V_5, \quad \mathbb{k} F_q \simeq V_6, \quad \mathbb{k} G_s \simeq V_7, \quad \mathbb{k} H_r \simeq V_8. \end{split}$$

Definition 6.4 For $n_1, n_2, ..., n_8 \in \mathbb{N}^*$ with $\sum_{i=1}^8 n_i \ge 1$ and

$$I_{1} = \{ (\alpha_{i_{1},i_{2}})_{n_{1}\times n_{1}}, \ (\beta_{j_{1},j_{2}})_{n_{2}\times n_{2}}, \ (\gamma_{k_{1},k_{2}})_{n_{3}\times n_{3}}, \ (\eta_{\ell_{1},\ell_{2}})_{n_{4}\times n_{4}}, \\ (\zeta_{m_{1},m_{2}})_{n_{5}\times n_{5}}, \ (\theta_{q_{1},q_{2}})_{n_{6}\times n_{6}}, \ (\lambda_{s_{1},s_{2}})_{n_{7}\times n_{7}}, \ (\mu_{r_{1},r_{2}})_{n_{8}\times n_{8}} \}$$

with entries in k, let us denote by $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$ the algebra that is generated by x, y, t, $\{A_i\}_{i=1,..,n_1}, \{B_j\}_{j=1,..,n_2}, \{C_k\}_{k=1,..,n_3}, \{D_\ell\}_{\ell=1,..,n_4}, \{E_m\}_{m=1,..,n_5}, \{F_q\}_{q=1,..,n_6}, \{G_s\}_{s=1,..,n_7}, \{H_r\}_{r=1,..,n_8}$ satisfying the relations (3.2), (3.3), (3.4) and

$$xA_i = -A_i x, \quad yA_i = -A_i y, \quad tA_i = A_i x^2 t,$$
 (6.20)

$$xB_j = -B_j x, \quad yB_j = -B_j y, \quad tB_j = -B_j x^2 t,$$
(6.21)

$$xC_k = C_k x, \quad yC_k = -C_k y, \quad tC_k = C_k x^2 t,$$
(6.22)

$$xD_{\ell} = D_{\ell}x, \quad yD_{\ell} = -D_{\ell}y, \quad tD_{\ell} = -D_{\ell}x^{2}t,$$
(6.23)

$$xE_m = -E_m x, \quad yE_m = -E_m y, \quad tE_m = E_m x^2 t,$$
(6.24)

$$xF_q = -F_q x, \quad yF_q = -F_q y, \quad tF_q = -F_q x^2 t,$$
(6.25)

$$xG_s = G_s x, \quad yG_s = -G_s y, \quad tG_s = G_s x^2 t,$$
 (6.26)

$$xH_r = H_r x, \quad yH_r = -H_r y, \quad tH_r = -H_r x^2 t,$$
 (6.27)

$$A_{i_1}A_{i_2} + A_{i_2}A_{i_1} = \alpha_{i_1,i_2}(1-x^2), \quad i_1, i_2 \in \{1, \dots, n_1\},$$
(6.28)

$$B_{j_1}B_{j_2} + B_{j_2}B_{j_1} = \beta_{j_1,j_2}(1-x^2), \quad j_1, j_2 \in \{1, \dots, n_2\},$$
(6.29)

$$C_{k_1}C_{k_2} + C_{k_2}C_{k_1} = \gamma_{k_1,k_2}(1-x^2), \quad k_1, k_2 \in \{1, \dots, n_3\},$$

$$(6.30)$$

$$D_{\ell_1} D_{\ell_2} + D_{\ell_2} D_{\ell_1} = \eta_{\ell_1, \ell_2} (1 - x^2), \quad \ell_1, \ell_2 \in \{1, \dots, n_4\},$$
(6.31)

$$E_{m_1}E_{m_2} + E_{m_2}E_{m_1} = \zeta_{m_1,m_2}(1-x^2), \quad m_1, m_2 \in \{1, \dots, n_5\},$$
(6.32)

$$F_{q_1}F_{q_2} + F_{q_2}F_{q_1} = \theta_{q_1,q_2}(1-x^2), \quad q_1,q_2 \in \{1,\dots,n_6\},$$
(6.33)

$$G_{s_1}G_{s_2} + G_{s_2}G_{s_1} = \lambda_{s_1,s_2}(1-x^2), \quad s_1, s_2 \in \{1, \dots, n_7\},$$
(6.34)

$$H_{r_1}H_{r_2} + H_{r_2}H_{r_1} = \mu_{r_1,r_2}(1-x^2), \quad r_1, r_2 \in \{1, \dots, n_8\},$$
(6.35)

$$A_i B_j + B_j A_i = 0, \quad A_i C_k - C_k A_i = 0, \quad A_i D_\ell - D_\ell A_i = 0, \quad A_i E_m + E_m A_i = 0, \tag{6.36}$$

$$A_i F_q + F_q A_i = 0, \quad A_i G_s - G_s A_i = 0, \quad A_i H_r - H_r A_i = 0, \quad B_j C_k - C_k B_j = 0, \tag{0.37}$$

$$B_{j}D_{\ell} - D_{\ell}B_{j} = 0, \quad B_{j}E_{m} + E_{m}B_{j} = 0, \quad B_{j}F_{q} + F_{q}B_{j} = 0, \quad B_{j}G_{s} - G_{s}B_{j} = 0, \quad (6.38)$$

$$B_{j}H_{j} - H_{j}B_{j} = 0, \quad C_{j}D_{s} + D_{s}C_{j} = 0, \quad C_{j}E_{j} - E_{j}C_{j} = 0, \quad C_{j}E_{j} - E_{j}C_{j} = 0, \quad (6.39)$$

$$D_{j}H_{r} - H_{r}D_{j} = 0, \quad C_{k}D_{\ell} + D_{\ell}C_{k} = 0, \quad C_{k}E_{m} - E_{m}C_{k} = 0, \quad C_{k}F_{q} - F_{q}C_{k} = 0, \quad (0.39)$$

$$C_k G_s + G_s C_m = 0, \quad C_k H_r + H_r C_k = 0, \quad D_\ell E_m - E_m D_\ell = 0, \quad D_\ell F_q - F_q D_\ell = 0, \quad (6.40)$$

$$D_{\ell}G_s + G_sD_{\ell} = 0, \quad D_{\ell}H_r + H_rD_{\ell} = 0, \quad E_mF_q + F_qE_m = 0, \quad E_mG_s - G_sE_m = 0, \quad (6.41)$$

$$E_m H_r - H_r E_m = 0, \quad F_q G_s - G_s F_q = 0, \quad F_q H_r - H_r F_q = 0, \quad G_s H_r + H_r G_s = 0.$$
(6.42)

It is a Hopf algebra with its coalgebra structure determined by

$$\begin{split} & \triangle(A_i) = A_i \otimes 1 + x \otimes A_i, \qquad \triangle(B_j) = B_j \otimes 1 + x \otimes B_j, \\ & \triangle(C_k) = C_k \otimes 1 + xy \otimes C_k, \qquad \triangle(D_\ell) = D_\ell \otimes 1 + xy \otimes D_\ell, \\ & \triangle(E_m) = E_m \otimes 1 + x^3 \otimes E_m, \qquad \triangle(F_q) = F_q \otimes 1 + x^3 \otimes F_q, \\ & \triangle(G_s) = G_s \otimes 1 + x^3y \otimes G_s, \qquad \triangle(H_r) = H_r \otimes 1 + x^3y \otimes H_r. \end{split}$$

Remark 6.5 It is clear that $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; 0) \cong \mathcal{B}(\Omega_1(n_1, n_2, \ldots, n_8)) \sharp H$. Also note that $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1) \cong T(\Omega_1(n_1, n_2, \ldots, n_8))/J(I_1)$, where $J(I_1)$ is a Hopf ideal generated by relations (6.28) - (6.35).

Now we determine the liftings and discuss the isomorphism classes of $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$.

Proposition 6.6 (1) Let A be a finite-dimensional Hopf algebra over H such that its infinitesimal braiding is $\Omega_1(n_1, n_2, \ldots, n_8)$, then $A \cong \mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$.

 $\begin{array}{ll} (2) \ \mathfrak{U}_{1}(n_{1},\ldots,n_{8};I_{1}) \cong \mathfrak{U}_{1}(n_{1},\ldots,n_{8};I_{1}') \ if \ and \ only \ if \ there \ exist \ invertible \ matrices \\ (a_{i_{1}i_{1}})_{n_{1}\times n_{1}}, \ (a_{i_{2}i_{2}})_{n_{1}\times n_{1}}, \ (b_{j_{1}'j_{1}})_{n_{2}\times n_{2}}, \ (b_{j_{2}'j_{2}})_{n_{2}\times n_{2}}, \ (c_{k_{1}'k_{1}})_{n_{3}\times n_{3}}, \ (c_{k_{2}'k_{2}})_{n_{3}\times n_{3}}, \ (d_{\ell_{1}\ell_{1}})_{n_{4}\times n_{4}}, \\ (d_{\ell_{2}\ell_{2}})_{n_{4}\times n_{4}}, \ (e_{m_{1}'m_{1}})_{n_{5}\times n_{5}}, \ (e_{m_{2}'m_{2}})_{n_{5}\times n_{5}}, \ (f_{q_{1}'q_{1}})_{n_{6}\times n_{6}}, \ (f_{q_{2}'q_{2}})_{n_{6}\times n_{6}}, \ (g_{s_{1}'s_{1}})_{n_{7}\times n_{7}}, \ (g_{s_{2}'s_{2}})_{n_{7}\times n_{7}}, \\ (h_{r_{1}'r_{1}})_{n_{8}\times n_{8}}, \ (h_{r_{2}'r_{2}})_{n_{8}\times n_{8}} \ such \ that \end{array}$

$$\sum_{i_{1}=1}^{n_{1}}\sum_{i_{2}'=1}^{n_{1}}a_{i_{1}'i_{1}}a_{i_{2}'i_{2}}\alpha_{i_{1}',i_{2}'}' = \alpha_{i_{1},i_{2}}, \qquad \sum_{j_{1}'=1}^{n_{2}}\sum_{j_{2}'=1}^{n_{2}}b_{j_{1}'j_{1}}b_{j_{2}'j_{2}}\beta_{j_{1}',j_{2}'}' = \beta_{j_{1},j_{2}}, \qquad (6.43)$$

$$\sum_{k_1'=1}^{n_3} \sum_{k_2'=1}^{n_3} c_{k_1'k_1} c_{k_2'k_2} \gamma_{k_1',k_2'}' = \gamma_{k_1,k_2}, \qquad \sum_{\ell_1'=1}^{n_4} \sum_{\ell_2'=1}^{n_4} d_{\ell_1'\ell_1} d_{\ell_2'\ell_2} \eta_{\ell_1',\ell_2'}' = \eta_{\ell_1,\ell_2}, \qquad (6.44)$$

$$\sum_{\substack{m_1'=1\\n_2}}^{n_5} \sum_{\substack{m_2'=1\\n_2}}^{n_5} e_{m_1'm_1} e_{m_2'm_2} \zeta_{m_1',m_2'}' = \zeta_{m_1,m_2}, \qquad \sum_{\substack{q_1'=1\\q_2'=1}}^{n_6} \sum_{\substack{n_6'\\q_1'q_1}}^{n_6} f_{q_1'q_1} f_{q_2'q_2} \theta_{q_1',q_2'}' = \theta_{q_1,q_2}, \tag{6.45}$$

$$\sum_{s_1'=1}^{n_7} \sum_{s_2'=1}^{n_7} g_{s_1's_1} g_{s_2's_2} \lambda_{s_1',s_2'}' = \lambda_{s_1,s_2}, \qquad \sum_{r_1'=1}^{n_8} \sum_{r_2'=1}^{n_8} h_{r_1'r_1} h_{r_2'r_2} \mu_{r_1',r_2'}' = \mu_{r_1,r_2}, \qquad (6.46)$$

or $n_1 = n_2$, $n_5 = n_6$ satisfying (6.44), (6.46) and

$$\sum_{j_{1}'=1}^{n_{1}} \sum_{j_{2}'=1}^{n_{1}} a_{j_{1}'i_{1}} a_{j_{2}'i_{2}} \beta_{j_{1}',j_{2}'}^{'} = \alpha_{i_{1},i_{2}}, \qquad \sum_{i_{1}'=1}^{n_{1}} \sum_{i_{2}'=1}^{n_{1}} b_{i_{1}'j_{1}} b_{i_{2}'j_{2}} \alpha_{i_{1}',i_{2}'}^{'} = \beta_{j_{1},j_{2}},$$

$$\sum_{q_{1}'=1}^{n_{5}} \sum_{q_{2}'=1}^{n_{5}} \sum_{q_{1}'m_{1}}^{n_{5}} e_{q_{1}'m_{1}} e_{q_{2}'m_{2}} \theta_{q_{1}',q_{2}'}^{'} = \zeta_{m_{1},m_{2}}, \qquad \sum_{m_{1}'=1}^{n_{5}} \sum_{m_{2}'=1}^{n_{5}} f_{m_{1}'q_{1}} f_{m_{2}'q_{2}} \zeta_{m_{1}',m_{2}'}^{'} = \theta_{q_{1},q_{2}}, \qquad (6.47)$$

or $n_1 = n_2$, $n_3 = n_4$, $n_5 = n_6$, $n_7 = n_8$ satisfying (6.47) and

$$\sum_{\ell_{1}^{\prime}=1}^{n_{3}} \sum_{\ell_{2}^{\prime}=1}^{n_{3}} c_{\ell_{1}^{\prime}k_{1}} c_{\ell_{2}^{\prime}k_{2}} \eta_{\ell_{1}^{\prime},\ell_{2}^{\prime}}^{\prime} = \gamma_{k_{1},k_{2}}, \qquad \sum_{k_{1}^{\prime}=1}^{n_{3}} \sum_{k_{2}^{\prime}=1}^{n_{3}} d_{k_{1}^{\prime}\ell_{1}} d_{k_{2}^{\prime}\ell_{2}} \gamma_{k_{1}^{\prime},k_{2}^{\prime}}^{\prime} = \eta_{\ell_{1},\ell_{2}}, \\ \sum_{r_{1}^{\prime}=1}^{n_{7}} \sum_{r_{2}^{\prime}=1}^{n_{7}} \sum_{r_{1}^{\prime}s_{1}} g_{r_{1}^{\prime}s_{2}} g_{r_{1}^{\prime}s_{1}} g_{r_{2}^{\prime}s_{2}} \mu_{r_{1}^{\prime},r_{2}^{\prime}}^{\prime} = \lambda_{s_{1},s_{2}}, \qquad \sum_{s_{1}^{\prime}=1}^{n_{7}} \sum_{s_{2}^{\prime}=1}^{n_{7}} h_{s_{1}^{\prime}r_{1}} h_{s_{2}^{\prime}r_{2}} \lambda_{s_{1}^{\prime},s_{2}^{\prime}}^{\prime} = \mu_{r_{1},r_{2}}, \qquad (6.48)$$

or $n_3 = n_4$, $n_7 = n_8$ satisfying (6.43), (6.45) and (6.48) or $n_1 = n_5$, $n_2 = n_6$, $n_3 = n_7$, $n_4 = n_8$ and there exist invertible matrices $(a_{m_1'i_1})_{n_1 \times n_1}$, $(a_{m_2'i_2})_{n_1 \times n_1}$, $(b_{q_1'j_1})_{n_2 \times n_2}$, $(b_{q_2'j_2})_{n_2 \times n_2}$, $(c_{s_1'k_1})_{n_3 \times n_3}$, $(c_{s_2'k_2})_{n_3 \times n_3}$, $(d_{r_1'\ell_1})_{n_4 \times n_4}$, $(d_{r_2'\ell_2})_{n_4 \times n_4}$, $(e_{i_1'm_1})_{n_1 \times n_1}$, $(e_{i_2'm_2})_{n_1 \times n_1}$, $(f_{j_1'q_1})_{n_2 \times n_2}$, $(f_{j_2'q_2})_{n_2 \times n_2}$, $(g_{k_1's_1})_{n_3 \times n_3}$, $(g_{k_2's_2})_{n_3 \times n_3}$, $(h_{\ell_1'r_1})_{n_4 \times n_4}$, $(h_{\ell_2'r_2})_{n_4 \times n_4}$ such that

$$\sum_{m_1'=1}^{n_1} \sum_{m_2'=1}^{n_1} a_{m_1'i_1} a_{m_2'i_2} \zeta_{m_1',m_2'}' = \alpha_{i_1,i_2}, \qquad \sum_{q_1'=1}^{n_2} \sum_{q_2'=1}^{n_2} b_{q_1'j_1} b_{q_2'j_2} \theta_{q_1',q_2'}' = \beta_{j_1,j_2}, \tag{6.49}$$

$$\sum_{s_1'=1}^{n_3} \sum_{s_2'=1}^{n_3} c_{s_1'k_1} c_{s_2'k_2} \lambda_{s_1',s_2'}' = \gamma_{k_1,k_2}, \qquad \sum_{r_1'=1}^{n_4} \sum_{r_2'=1}^{n_4} d_{r_1'\ell_1} d_{r_2'\ell_2} \mu_{r_1',r_2'}' = \eta_{\ell_1,\ell_2}, \tag{6.50}$$

$$\sum_{i_1'=1}^{n_1} \sum_{i_2'=1}^{n_1} e_{i_1'm_1} e_{i_2'm_2} \alpha_{i_1',i_2'}' = \zeta_{m_1,m_2}, \qquad \sum_{j_1'=1}^{n_2} \sum_{j_2'=1}^{n_2} f_{j_1'q_1} f_{j_2'q_2} \beta_{j_1',j_2'}' = \theta_{q_1,q_2}, \tag{6.51}$$

$$\sum_{k_1'=1}^{n_3} \sum_{k_2'=1}^{n_3} g_{k_1's_1} g_{k_2's_2} \gamma_{k_1',k_2'}' = \lambda_{s_1,s_2}, \qquad \sum_{\ell_1'=1}^{n_4} \sum_{\ell_2'=1}^{n_4} h_{\ell_1'r_1} h_{\ell_2'r_2} \eta_{\ell_1',\ell_2'}' = \mu_{r_1,r_2}, \qquad (6.52)$$

or $n_1 = n_6$, $n_2 = n_5$, $n_3 = n_7$, $n_4 = n_8$ satisfying (6.50), (6.52) and

$$\sum_{q_1'=1}^{n_1} \sum_{q_2'=1}^{n_1} a_{q_1'i_1} a_{q_2'i_2} \theta_{q_1',q_2'}' = \alpha_{i_1,i_2}, \qquad \sum_{m_1'=1}^{n_2} \sum_{m_2'=1}^{n_2} b_{m_1'j_1} b_{m_2'j_2} \zeta_{m_1',m_2'}' = \beta_{j_1,j_2},$$

$$\sum_{j_1'=1}^{n_2} \sum_{j_2'=1}^{n_2} e_{j_1'm_1} e_{j_2'm_2} \beta_{j_1',j_2'}' = \zeta_{m_1,m_2}, \qquad \sum_{i_1'=1}^{n_1} \sum_{i_2'=1}^{n_1} f_{i_1'q_1} f_{i_2'q_2} \alpha_{i_1',i_2'}' = \theta_{q_1,q_2},$$
(6.53)

or $n_1 = n_6$, $n_2 = n_5$, $n_3 = n_8$, $n_4 = n_7$ satisfying (6.53) and

$$\sum_{r_{1}^{\prime}=1}^{n_{3}} \sum_{r_{2}^{\prime}=1}^{n_{3}} c_{r_{1}^{\prime}k_{1}} c_{r_{2}^{\prime}k_{2}} \mu_{r_{1}^{\prime},r_{2}^{\prime}}^{\prime} = \gamma_{k_{1},k_{2}}, \qquad \sum_{s_{1}^{\prime}=1}^{n_{4}} \sum_{s_{2}^{\prime}=1}^{n_{4}} d_{s_{1}^{\prime}\ell_{1}} d_{s_{2}^{\prime}\ell_{2}} \lambda_{s_{1}^{\prime},s_{2}^{\prime}}^{\prime} = \eta_{\ell_{1},\ell_{2}}, \qquad \sum_{s_{1}^{\prime}=1}^{n_{4}} \sum_{s_{2}^{\prime}=1}^{n_{4}} d_{s_{1}^{\prime}\ell_{1}} d_{s_{2}^{\prime}\ell_{2}} \lambda_{s_{1}^{\prime},s_{2}^{\prime}}^{\prime} = \eta_{\ell_{1},\ell_{2}}, \qquad (6.54)$$

or $n_1 = n_5$, $n_2 = n_6$, $n_3 = n_8$, $n_4 = n_7$ satisfying (6.49), (6.51) and (6.54) or $n_1 = n_3$, $n_2 = n_4$, $n_5 = n_7$, $n_6 = n_8$ and there exist invertible matrices $(a_{k'_1i_1})_{n_1 \times n_1}$, $(a_{k'_2i_2})_{n_1 \times n_1}$, $(b_{\ell'_1j_1})_{n_2 \times n_2}$, $(b_{\ell'_2j_2})_{n_2 \times n_2}$, $(c_{i'_1k_1})_{n_1 \times n_1}$, $(c_{i'_2k_2})_{n_1 \times n_1}$, $(d_{j'_1\ell_1})_{n_2 \times n_2}$, $(d_{j'_2\ell_2})_{n_2 \times n_2}$, $(e_{s'_1m_1})_{n_5 \times n_5}$, $(e_{s'_2m_2})_{n_5 \times n_5}$, $(f_{r'_1q_1})_{n_6 \times n_6}$, $(f_{r'_2q_2})_{n_6 \times n_6}$, $(g_{m'_1s_1})_{n_5 \times n_5}$, $(g_{m'_2s_2})_{n_5 \times n_5}$, $(h_{q'_1r_1})_{n_6 \times n_6}$, $(h_{q'_2r_2})_{n_6 \times n_6}$ such that

$$\sum_{k_1'=1}^{n_1} \sum_{k_2'=1}^{n_1} a_{k_1'i_1} a_{k_2'i_2} \gamma_{k_1',k_2'}' = \alpha_{i_1,i_2}, \qquad \sum_{\ell_1'=1}^{n_2} \sum_{\ell_2'=1}^{n_2} b_{\ell_1'j_1} b_{\ell_2'j_2} \eta_{\ell_1',\ell_2'}' = \beta_{j_1,j_2}, \tag{6.55}$$

$$\sum_{i_1'=1}^{n_1} \sum_{i_2'=1}^{n_1} c_{i_1'k_1} c_{i_2'k_2} \alpha_{i_1',i_2'} = \gamma_{k_1,k_2}, \qquad \sum_{j_1'=1}^{n_2} \sum_{j_2'=1}^{n_2} d_{j_1'\ell_1} d_{j_2'\ell_2} \beta_{j_1',j_2'} = \eta_{\ell_1,\ell_2}, \tag{6.56}$$

$$\sum_{s_1'=1}^{n_5} \sum_{s_2'=1}^{n_5} e_{s_1'm_1} e_{s_2'm_2} \lambda_{s_1',s_2'}' = \zeta_{m_1,m_2}, \qquad \sum_{r_1'=1}^{n_6} \sum_{r_2'=1}^{n_6} f_{r_1'q_1} f_{r_2'q_2} \mu_{r_1',r_2'}' = \theta_{q_1,q_2}, \tag{6.57}$$

$$\sum_{m_1'=1}^{n_5} \sum_{m_2'=1}^{n_5} g_{m_1's_1} g_{m_2's_2} \zeta_{m_1',m_2'}' = \lambda_{s_1,s_2}, \quad \sum_{q_1'=1}^{n_6} \sum_{q_2'=1}^{n_6} h_{q_1'r_1} h_{q_2'r_2} \theta_{q_1',q_2'}' = \mu_{r_1,r_2}, \tag{6.58}$$

or $n_1 = n_2 = n_3 = n_4$, $n_5 = n_6 = n_7 = n_8$ satisfying (6.55), (6.57) and

$$\sum_{j_{1}^{'}=1}^{n_{1}} \sum_{j_{2}^{'}=1}^{n_{1}} c_{j_{1}^{'}k_{1}} c_{j_{2}^{'}k_{2}} \beta_{j_{1}^{'},j_{2}^{'}}^{'} = \gamma_{k_{1},k_{2}}, \qquad \sum_{i_{1}^{'}=1}^{n_{1}} \sum_{i_{2}^{'}=1}^{n_{1}} d_{i_{1}^{'}\ell_{1}} d_{i_{2}^{'}\ell_{2}} \alpha_{i_{1}^{'},i_{2}^{'}}^{'} = \eta_{\ell_{1},\ell_{2}}, \qquad \sum_{i_{1}^{'}=1}^{n_{5}} \sum_{i_{2}^{'}=1}^{n_{5}} \sum_{j_{2}^{'}=1}^{n_{5}} g_{q_{1}^{'}s_{1}} g_{q_{2}^{'}s_{2}} \theta_{q_{1}^{'},q_{2}^{'}}^{'} = \lambda_{s_{1},s_{2}}, \qquad \sum_{m_{1}^{'}=1}^{n_{5}} \sum_{m_{2}^{'}=1}^{n_{5}} h_{m_{1}^{'}r_{1}} h_{m_{2}^{'}r_{2}} \zeta_{m_{1}^{'},m_{2}^{'}}^{'} = \mu_{r_{1},r_{2}}, \qquad (6.59)$$

or $n_1 = n_4, n_2 = n_3, n_5 = n_8, n_6 = n_7$ satisfying (6.59) and

$$\sum_{\ell_{1}^{\prime}=1}^{n_{1}} \sum_{\ell_{2}^{\prime}=1}^{n_{1}} a_{\ell_{1}^{\prime}i_{1}} a_{\ell_{2}^{\prime}i_{2}} \eta_{\ell_{1}^{\prime},\ell_{2}^{\prime}}^{\prime} = \alpha_{i_{1},i_{2}}, \qquad \sum_{k_{1}^{\prime}=1}^{n_{2}} \sum_{k_{2}^{\prime}=1}^{n_{2}} b_{k_{1}^{\prime}j_{1}} b_{k_{2}^{\prime}j_{2}} \gamma_{k_{1}^{\prime},k_{2}^{\prime}}^{\prime} = \beta_{j_{1},j_{2}}, \qquad \sum_{k_{1}^{\prime}=1}^{n_{2}} \sum_{k_{2}^{\prime}=1}^{n_{2}} b_{k_{1}^{\prime}j_{1}} b_{k_{2}^{\prime}j_{2}} \gamma_{k_{1}^{\prime},k_{2}^{\prime}}^{\prime} = \beta_{j_{1},j_{2}}, \qquad (6.60)$$

or $n_1 = n_2 = n_3 = n_4$, $n_5 = n_6 = n_7 = n_8$ satisfying (6.56), (6.58) and (6.60) or $n_1 = n_7$, $n_2 = n_8$, $n_3 = n_5$, $n_4 = n_6$ and there exist invertible matrices $(a_{s'_1i_1})_{n_1 \times n_1}$, $(a_{s'_2i_2})_{n_1 \times n_1}$, $(b_{r'_1j_1})_{n_2 \times n_2}$, $(b_{r'_2j_2})_{n_2 \times n_2}$, $(c_{m'_1k_1})_{n_3 \times n_3}$, $(c_{m'_2k_2})_{n_3 \times n_3}$, $(d_{q'_1\ell_1})_{n_4 \times n_4}$, $(d_{q'_2\ell_2})_{n_4 \times n_4}$, $(e_{k'_1m_1})_{n_3 \times n_3}$, $(e_{k'_2m_2})_{n_3 \times n_3}$, $(f_{\ell'_1q_1})_{n_4 \times n_4}$, $(f_{\ell'_2q_2})_{n_4 \times n_4}$, $(g_{i'_1s_1})_{n_1 \times n_1}$, $(g_{i'_2s_2})_{n_1 \times n_1}$, $(h_{j'_1r_1})_{n_2 \times n_2}$, $(h_{j'_2r_2})_{n_2 \times n_2}$ such that

$$\sum_{s_{1}^{'}=1}^{n_{1}}\sum_{s_{2}^{'}=1}^{n_{1}}a_{s_{1}^{'}i_{1}}a_{s_{2}^{'}i_{2}}\lambda_{s_{1}^{'},s_{2}^{'}}^{'} = \alpha_{i_{1},i_{2}}, \qquad \sum_{r_{1}^{'}=1}^{n_{2}}\sum_{r_{2}^{'}=1}^{n_{2}}b_{r_{1}^{'}j_{1}}b_{r_{2}^{'}j_{2}}\mu_{r_{1}^{'},r_{2}^{'}}^{'} = \beta_{j_{1},j_{2}}, \qquad (6.61)$$

$$\sum_{m_1'=2}^{n_3} \sum_{m_2'=1}^{n_3} c_{m_1'k_1} c_{m_2'k_2} \zeta_{m_1',m_2'}' = \gamma_{k_1,k_2}, \quad \sum_{q_1'=1}^{n_4} \sum_{q_2'=1}^{n_4} d_{q_1'\ell_1} d_{q_2'\ell_2} \theta_{q_1',q_2'}' = \eta_{\ell_1,\ell_2}, \quad (6.62)$$

$$\sum_{k_1'=1}^{n_3} \sum_{k_2'=1}^{n_3} e_{k_1'm_1} e_{k_2'm_2} \gamma_{k_1',k_2'}' = \zeta_{m_1,m_2}, \qquad \sum_{\ell_1'=1}^{n_4} \sum_{\ell_2'=1}^{n_4} f_{\ell_1'q_1} f_{\ell_2'q_2} \eta_{\ell_1',\ell_2'}' = \theta_{q_1,q_2}, \tag{6.63}$$

$$\sum_{i_1'=1}^{n_1} \sum_{i_2'=1}^{n_1} g_{i_1's_1} g_{i_2's_2} \alpha'_{i_1',i_2'} = \lambda_{s_1,s_2}, \qquad \sum_{j_1'=1}^{n_2} \sum_{j_2'=1}^{n_2} h_{j_1'r_1} h_{j_2'r_2} \beta'_{j_1',j_2'} = \mu_{r_1,r_2}, \qquad (6.64)$$

or $n_1 = n_2 = n_7 = n_8$, $n_3 = n_4 = n_5 = n_6$ satisfying (6.61), (6.63) and

$$\sum_{q_{1}'=1}^{n_{3}} \sum_{q_{2}'=1}^{n_{3}} c_{q_{1}'k_{1}} c_{q_{2}'k_{2}} \theta_{q_{1}',q_{2}'}' = \gamma_{k_{1},k_{2}}, \qquad \sum_{m_{1}'=1}^{n_{3}} \sum_{m_{2}'=1}^{n_{3}} d_{m_{1}'\ell_{1}} d_{m_{2}'\ell_{2}} \zeta_{m_{1}',m_{2}'}' = \eta_{\ell_{1},\ell_{2}}, \sum_{j_{1}'=1}^{n_{1}} \sum_{j_{2}'=1}^{n_{1}} g_{j_{1}'s_{1}} g_{j_{2}'s_{2}} \beta_{j_{1}',j_{2}'}' = \lambda_{s_{1},s_{2}}, \qquad \sum_{i_{1}'=1}^{n_{1}} \sum_{i_{2}'=1}^{n_{1}} h_{i_{1}'r_{1}} h_{i_{2}'r_{2}} \alpha_{i_{1}',i_{2}'}' = \mu_{r_{1},r_{2}},$$

$$(6.65)$$

or $n_1 = n_8$, $n_2 = n_7$, $n_3 = n_6$, $n_4 = n_5$ satisfying (6.65) and

$$\sum_{r_{1}^{'}=1}^{n_{1}} \sum_{r_{2}^{'}=1}^{n_{1}} a_{r_{1}^{'}i_{1}} a_{r_{2}^{'}i_{2}} \mu_{r_{1}^{'},r_{2}^{'}}^{'} = \alpha_{i_{1},i_{2}}, \qquad \sum_{s_{1}^{'}=1}^{n_{2}} \sum_{s_{2}^{'}=1}^{n_{2}} b_{s_{1}^{'}j_{1}} b_{s_{2}^{'}j_{2}} \lambda_{s_{1}^{'},s_{2}^{'}}^{'} = \beta_{j_{1},j_{2}}, \qquad \sum_{s_{1}^{'}=1}^{n_{4}} \sum_{s_{2}^{'}=1}^{n_{4}} \sum_{s_{1}^{'}=1}^{n_{4}} e_{\ell_{1}^{'}m_{1}} e_{\ell_{2}^{'}m_{2}} \eta_{\ell_{1}^{'},\ell_{2}^{'}}^{'} = \zeta_{m_{1},m_{2}}, \qquad \sum_{k_{1}^{'}=1}^{n_{3}} \sum_{k_{2}^{'}=1}^{n_{3}} f_{k_{1}^{'}q_{1}} f_{k_{2}^{'}q_{2}} \gamma_{k_{1}^{'},k_{2}^{'}}^{'} = \theta_{q_{1},q_{2}}, \qquad (6.66)$$

or $n_1 = n_2 = n_7 = n_8$, $n_3 = n_4 = n_5 = n_6$ satisfying (6.62), (6.64) and (6.66).

Proof. (1) By Theorem 6.1, we have that $\operatorname{gr}(A) \cong \mathcal{B}(\Omega_1(n_1, n_2, \dots, n_8)) \sharp H$. We can check the relations listed in Definition 6.4, except (6.28) – (6.42), hold in A from the bosonization $\mathcal{B}(\Omega_1(n_1, n_2, \dots, n_8)) \sharp H$. As $\Delta(A_i) = A_i \otimes 1 + x \otimes A_i, \Delta(B_j) = B_j \otimes 1 + x \otimes B_j$, a direct computation shows that

$$\Delta(A_{i_1}A_{i_2} + A_{i_2}A_{i_1}) = (A_{i_1}A_{i_2} + A_{i_2}A_{i_1}) \otimes 1 + x^2 \otimes (A_{i_1}A_{i_2} + A_{i_2}A_{i_1}),$$

$$\Delta(A_iB_j + B_jA_i) = (A_iB_j + B_jA_i) \otimes 1 + x^2 \otimes (A_iB_j + B_jA_i).$$

Then $A_{i_1}A_{i_2} + A_{i_2}A_{i_1}, A_iB_j + B_jA_i \in \mathcal{P}_{1,x^2}(\mathcal{B}(\Omega_1(n_1, n_2, \dots, n_8)) \sharp H) = \mathcal{P}_{1,x^2}(H) = \Bbbk\{1 - x^2\}.$ Since $t(A_iB_j + B_jA_i) = -(A_iB_j + B_jA_i)t$, it follows that

$$A_{i_1}A_{i_2} + A_{i_2}A_{i_1} = \alpha_{i_1,i_2}(1-x^2), \quad A_iB_j + B_jA_i = 0$$

for some $\alpha_{i_1,i_2} \in \mathbb{k}$. Similarly, the relations (6.29) - (6.42) hold in A. Then there is a surjective Hopf morphism from $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$ to A. We can observe that each element of $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$ can be expressed by a linear combination of

$$\{A_i^{\alpha_i}B_j^{\beta_j}C_k^{\gamma_k}D_\ell^{\eta_\ell}E_m^{\zeta_m}F_q^{\theta_q}G_s^{\lambda_s}H_r^{\mu_r}x^ey^ft^g; \ \alpha_i,\beta_j,\gamma_k,\eta_\ell,\zeta_m,\theta_q,\lambda_s,\mu_r,f,g\in\mathbb{I}_{0,1},e\in\mathbb{I}_{0,3}\}.$$

In fact, according to the Diamond Lemma [15], the set is a basis of $\mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$. Then dim $A = \dim \mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$ and whence $A \cong \mathfrak{U}_1(n_1, n_2, \ldots, n_8; I_1)$.

(2) Suppose that $\Phi : \mathfrak{U}_1(n_1, \ldots, n_8; I_1) \to \mathfrak{U}_1(n_1, \ldots, n_8; I_1')$ is an isomorphism of Hopf algebras, where

$$I_{1}' = \{ (\alpha_{i_{1},i_{2}}')_{n_{1}\times n_{1}}, \ (\beta_{j_{1},j_{2}}')_{n_{2}\times n_{2}}, \ (\gamma_{k_{1},k_{2}}')_{n_{3}\times n_{3}}, \ (\eta_{\ell_{1},\ell_{2}}')_{n_{4}\times n_{4}}, \\ (\zeta_{m_{1},m_{2}}')_{n_{5}\times n_{5}}, \ (\theta_{q_{1},q_{2}}')_{n_{6}\times n_{6}}, \ (\lambda_{s_{1},s_{2}}')_{n_{7}\times n_{7}}, \ (\mu_{r_{1},r_{2}}')_{n_{8}\times n_{8}} \}$$

and $\mathfrak{U}_1(n_1, n_2, \dots, n_8; I'_1)$ is generated by $x, y, t, A'_i, B'_j, C'_k, D'_\ell, E'_m, F'_q, G'_s, H'_r$.

When $\Phi|_H = \tau_1$ or τ_3 , then $\Phi(A_i)$ is (1, x)-skew primitive, so

$$\Phi(A_i) \in \bigoplus_{i'=1}^{n_1} \mathbb{k}A'_{i'} \bigoplus_{j'=1}^{n_2} \mathbb{k}B'_{j'} \bigoplus \mathbb{k}(1-x).$$

 $tA_i = A_i x^2 t$, $tB'_{j'} = -B'_{j'} x^2 t$ imply that $\Phi(A_i)$ doesn't contain the term of $\bigoplus_{j'=1}^{n_2} \mathbb{k}B'_{j'}$. $xA_i = -A_i x$ implies that $\Phi(A_i)$ doesn't contain the term of 1 - x. So there exists an invertible matrix $(a_{i'i})_{n_1 \times n_1}$ such that $\Phi(A_i) = \sum_{i'=1}^{n_1} a_{i'i} A'_{i'}$. Similarly, $\Phi(B_j) = \sum_{j'=1}^{n_2} b_{j'j} B'_{j'}$,

$$\Phi(C_k) = \sum_{k'=1}^{n_3} c_{k'k} C'_{k'}, \quad \Phi(D_\ell) = \sum_{\ell'=1}^{n_4} d_{\ell'\ell} D'_{\ell'}, \quad \Phi(E_m) = \sum_{m'=1}^{n_5} e_{m'm} E'_{m'},$$

$$\Phi(F_q) = \sum_{q'=1}^{n_6} f_{q'q} F'_{q'}, \quad \Phi(G_s) = \sum_{s'=1}^{n_7} g_{s's} G'_{s'}, \quad \Phi(H_r) = \sum_{r'=1}^{n_8} h_{r'r} H'_{r'}.$$

In this case, Φ is an isomorphism of Hopf algebras if and only if the relations (6.43) – (6.46) hold. Similarly, Φ is an isomorphism of Hopf algebras if and only if

when $\Phi|_{H} = \tau_{2}$ or τ_{4} , the relations (6.44), (6.46), (6.47) hold; when $\Phi|_{H} = \tau_{5}$ or τ_{7} , the relations (6.48), (6.47) hold; when $\Phi|_{H} = \tau_{6}$ or τ_{8} , the relations (6.43), (6.45), (6.48) hold; when $\Phi|_{H} = \tau_{9}$ or τ_{11} , the relations (6.49) – (6.52) hold; when $\Phi|_{H} = \tau_{10}$ or τ_{12} , the relations (6.50), (6.52), (6.53) hold; when $\Phi|_{H} = \tau_{13}$ or τ_{15} , the relations (6.53) , (6.54) hold; when $\Phi|_{H} = \tau_{14}$ or τ_{16} , the relations (6.49), (6.51), (6.54) hold; when $\Phi|_{H} = \tau_{17}$ or τ_{19} , the relations (6.55) – (6.58) hold; when $\Phi|_{H} = \tau_{21}$ or τ_{23} , the relations (6.59), (6.60) hold; when $\Phi|_{H} = \tau_{22}$ or τ_{24} , the relations (6.56), (6.58), (6.60) hold; when $\Phi|_{H} = \tau_{25}$ or τ_{27} , the relations (6.61) – (6.64) hold; when $\Phi|_{H} = \tau_{29}$ or τ_{31} , the relations (6.65), (6.66) hold; when $\Phi|_{H} = \tau_{29}$ or τ_{31} , the relations (6.62), (6.64), (6.66) hold;

Definition 6.7 For n_1 , n_2 , n_3 $n_4 \in \mathbb{N}^*$ with $\sum_{i=1}^4 n_i \ge 0$ and

$$I_2 = \{ \gamma_{k_1,k_2}, \ \eta_{\ell_1,\ell_2}, \ \lambda_{s_1,s_2}, \ \mu_{r_1,r_2}, \ \lambda_k, \ \zeta_\ell, \ \iota_s, \ \theta_r, \ \nu \},\$$

where k, k_1 , $k_2 = 1, ..., n_1$, ℓ , ℓ_1 , $\ell_2 = 1, ..., n_2$, s, s_1 , $s_2 = 1, ..., n_3$, r, r_1 , $r_2 = 1, ..., n_4$. Let us denote by $\mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2)$ the algebra that is generated by x, y, t, p_1 , p_2 , $\{C_k\}$, $\{D_\ell\}$, $\{G_s\}$, $\{H_r\}$ satisfying the relations (3.2) - (3.4), (6.22), (6.23), (6.26), (6.27), (6.30), (6.31), (6.34), (6.35) and

$$\begin{aligned} xp_1 &= p_1 x, \ xp_2 = p_2 x, \ yp_1 = -p_1 y, \ yp_2 = -p_2 y, \ tp_1 = p_1 x^2 t, \ tp_2 = -p_2 x^2 t, \\ C_k D_\ell + D_\ell C_k &= 0, \quad C_k G_s + G_s C_m = 0, \quad C_k H_r + H_r C_k = 0, \\ D_\ell G_s + G_s D_\ell &= 0, \quad D_\ell H_r + H_r D_\ell = 0, \quad G_s H_r + H_r G_s = 0, \end{aligned}$$

$$p_1^2 = \nu(1 - x^2), \quad p_2^2 = -\nu(1 - x^2), \quad p_1 p_2 + p_2 p_1 = 0,$$
(6.67)

$$p_1C_k + C_kp_1 = \lambda_k(x + x^3 - 2), \quad p_2C_k + C_kp_2 = \lambda_k(x - x^3),$$
(6.68)

$$p_1 D_\ell + D_\ell p_1 = \zeta_\ell (x - x^3), \qquad p_2 D_\ell + D_\ell p_2 = \zeta_\ell (x + x^3 - 2),$$
(6.69)

$$p_1G_s + G_sp_1 = \iota_s(x + x^3 - 2), \quad p_2G_s + G_sp_2 = \iota_s(x^3 - x),$$
(6.70)

$$p_1H_r + H_rp_1 = \theta_r(x^3 - x), \qquad p_2H_r + H_rp_2 = \theta_r(x + x^3 - 2).$$
 (6.71)

a Hopf algebra with its coalgebra structure determined by

$$\triangle(p_1) = p_1 \otimes 1 + \frac{1}{2}(1+x^2)y \otimes p_1 + \frac{1}{2}(1-x^2)y \otimes p_2, \\ \triangle(p_2) = p_2 \otimes 1 + \frac{1}{2}(1+x^2)y \otimes p_2 + \frac{1}{2}(1-x^2)y \otimes p_1, \\ \triangle(C_k) = C_k \otimes 1 + xy \otimes C_k, \qquad \triangle(D_\ell) = D_\ell \otimes 1 + xy \otimes D_\ell, \\ \triangle(G_s) = G_s \otimes 1 + x^3y \otimes G_s, \qquad \triangle(H_r) = H_r \otimes 1 + x^3y \otimes H_r.$$

It is

Remark 6.8 It is clear that $\mathfrak{U}_2(n_1, n_2, n_3, n_4; 0) \cong \mathcal{B}(\Omega_2(n_1, n_2, n_3, n_4)) \sharp H$. Also note that $\mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2) \cong T(\Omega_2(n_1, n_2, n_3, n_4))/J(I_2)$, where $J(I_2)$ is a Hopf ideal generated by relations (6.30), (6.31), (6.34), (6.35), (6.67) - (6.71).

Definition 6.9 For n_1 , n_2 , n_3 , $n_4 \in \mathbb{N}^*$ with $\sum_{i=1}^4 n_i \ge 0$ and a set

 $I_8 = \{ (\alpha_{i_1,i_2})_{n_1 \times n_1}, \ (\beta_{j_1,j_2})_{n_2 \times n_2}, \ (\zeta_{m_1,m_2})_{n_3 \times n_3}, \ (\theta_{q_1,q_2})_{n_4 \times n_4}, \ \nu \},$

we denote by $\mathfrak{U}_8(n_1, n_2, n_3, n_4; I_8)$ the algebra that is generated by $x, y, t, p_1, p_2, \{A_i\}_{i=1,..,n_1}, \{B_j\}_{j=1,..,n_2}, \{E_m\}_{m=1,...,n_3}, \{F_q\}_{q=1,...,n_4}$, satisfying the relations (3.2) - (3.4), (6.20), (6.21), (6.24), (6.25), (6.28), (6.29), (6.32), (6.33), (6.67) and

$$\begin{aligned} xp_1 &= -p_1 x, \ xp_2 &= -p_2 x, \ yp_1 &= p_1 y, \ yp_2 &= p_2 y, \ tp_1 &= p_1 t, \ tp_2 &= -p_2 t, \\ A_i B_j + B_j A_i &= 0, \quad A_i E_m + E_m A_i &= 0, \quad A_i F_q + F_q A_i &= 0, \\ B_j E_m + E_m B_j &= 0, \quad B_j F_q + F_q B_j &= 0, \quad E_m F_q + F_q E_m &= 0, \\ p_1 A_i + A_i p_1 &= 0, \quad p_2 A_i + A_i p_2 &= 0, \quad p_1 B_j + B_j p_1 &= 0, \quad p_2 B_j + B_j p_2 &= 0, \\ p_1 E_m + E_m p_1 &= 0, \quad p_2 E_m + E_m p_2 &= 0, \quad p_1 F_q + F_q p_1 &= 0, \quad p_2 F_q + F_q p_2 &= 0. \end{aligned}$$
(6.72)

It is a Hopf algebra with its coalgebra structure determined by

$$\begin{split} & \triangle(p_1) = p_1 \otimes 1 + \frac{1}{2}(1+x^2)x \otimes p_1 + \frac{\xi}{2}(1-x^2)x \otimes p_2, \\ & \triangle(p_2) = p_2 \otimes 1 + \frac{1}{2}(1+x^2)x \otimes p_2 - \frac{\xi}{2}(1-x^2)x \otimes p_1, \\ & \triangle(A_i) = A_i \otimes 1 + x \otimes A_i, \qquad \triangle(B_j) = B_j \otimes 1 + x \otimes B_j, \\ & \triangle(E_m) = E_m \otimes 1 + x^3 \otimes E_m, \qquad \triangle(F_q) = F_q \otimes 1 + x^3 \otimes F_q. \end{split}$$

Proposition 6.10 (1) Let A be a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is $\Omega_i(n_1, n_2, n_3, n_4)$ for $i \in \{2, 8\}$, then $A \cong \mathfrak{U}_i(n_1, n_2, n_3, n_4; I_i)$.

(2) $\mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2) \cong \mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2')$ if and only if there exist nonzero parameter z and invertible matrices $(c_{k'k})_{n_1 \times n_1}, (c_{k'_1k_1})_{n_1 \times n_1}, (c_{k'_2k_2})_{n_1 \times n_1}, (d_{\ell'\ell})_{n_2 \times n_2}, (d_{\ell'_1\ell_1})_{n_2 \times n_2}, (d_{\ell'_1\ell_1})_{n$

 $(d_{\ell'_2\ell_2})_{n_2\times n_2}, (g_{s's})_{n_3\times n_3}, (g_{s'_1s_1})_{n_3\times n_3}, (g_{s'_2s_2})_{n_3\times n_3}, (h_{r'r})_{n_4\times n_4}, (h_{r'_1r_1})_{n_4\times n_4}, (h_{r'_2r_2})_{n_4\times n_4} \text{ satis-fying } z^2v' = v, \ (6.44), \ (6.46) \text{ and }$

$$z\sum_{k'=1}^{n_1} c_{k'k} \lambda'_{k'} = \lambda_k, \quad z\sum_{\ell'=1}^{n_2} d_{\ell'\ell} \zeta'_{\ell'} = \zeta_\ell, \quad z\sum_{s'=1}^{n_3} g_{s's} \iota'_{s'} = \iota_s, \quad z\sum_{r'=1}^{n_4} h_{r'r} \theta'_{r'} = \theta_r.$$
(6.74)

or $n_1 = n_2$, $n_3 = n_4$ satisfying $z^2 v' = -v$, (6.48) and

$$z\sum_{\ell'=1}^{n_1} c_{\ell'k}\zeta_{\ell'} = \lambda_k, \quad z\sum_{k'=1}^{n_1} d_{k'\ell}\lambda_{k'} = \zeta_\ell, \quad z\sum_{r'=1}^{n_3} g_{r's}\theta_{r'} = \iota_s, \quad z\sum_{s'=1}^{n_3} h_{s'r}\iota_{s'} = \theta_r.$$
(6.75)

or $n_1 = n_3$, $n_2 = n_4$ satisfying $z^2 v' = v$, (6.50), (6.52) and

$$z\sum_{s'=1}^{n_1} c_{s'k} \iota_{s'}^{'} = \lambda_k, \quad z\sum_{r'=1}^{n_2} d_{r'\ell} \theta_{r'}^{'} = \zeta_\ell, \quad z\sum_{k'=1}^{n_1} g_{k's} \lambda_{k'}^{'} = \iota_s, \quad z\sum_{\ell'=1}^{n_2} h_{\ell'r} \zeta_{\ell'}^{'} = \theta_r.$$
(6.76)

or $n_1 = n_4$, $n_2 = n_3$ satisfying $z^2 v' = -v$, (6.54) and

$$z\sum_{r'=1}^{n_1} c_{r'k} \theta'_{r'} = \lambda_k, \quad z\sum_{s'=1}^{n_2} d_{s'l} \iota'_{s'} = \zeta_\ell, \quad z\sum_{\ell'=1}^{n_2} g_{\ell's} \zeta'_{\ell'} = \iota_s, \quad z\sum_{k'=1}^{n_1} h_{k'r} \lambda'_{k'} = \theta_r.$$
(6.77)

(3) $\mathfrak{U}_8(n_1, n_2, n_3, n_4; I_8) \cong \mathfrak{U}_8(n_1, n_2, n_3, n_4; I'_8)$ if and only if there exist nonzero parameter z and invertible matrices $(a_{i'_{111}})_{n_1 \times n_1}, (a_{i'_{212}})_{n_1 \times n_1}, (b_{j'_{111}})_{n_2 \times n_2}, (b_{j'_{212}})_{n_2 \times n_2}, (e_{m'_{1m_1}})_{n_3 \times n_3}, (e_{m'_{2m_2}})_{n_3 \times n_3}, (f_{q'_{1q_1}})_{n_4 \times n_4}, (f_{q'_{2q_2}})_{n_4 \times n_4}$ satisfying $z^2v' = v$, (6.43), (6.45) or $z^2v' = -v$ or $n_1 = n_2, n_3 = n_4$ satisfying $z^2v' = v$, (6.47) or $z^2v' = -v$ or $n_1 = n_3, n_2 = n_4$ satisfying $z^2v' = v$, (6.49), (6.51) or $z^2v' = -v$ or $n_1 = n_4, n_2 = n_3$ satisfying $z^2v' = v$, (6.53) or $z^2v' = -v$.

Proof. (1) We prove the claim only for $\Omega_2(n_1, n_2, n_3, n_4)$, since the proof for $\Omega_8(n_1, n_2, n_3, n_4)$ follows the same lines. By Theorem 6.1, we have that $gr(A) \cong \mathcal{B}(\Omega_2(n_1, n_2, n_3, n_4)) \sharp H$. Let $M_1 = \Bbbk\{p_1, p_2\}$. From Proposition 6.6 and a direct computation, we get that the relations except (6.67) - (6.71) hold in A. As

it follows that

So we have the relation (6.67) holds in A. Since $\triangle(C_k) = C_k \otimes 1 + xy \otimes C_k$, we get that

$$\Delta(p_1C_k + C_kp_1) = (p_1C_k + C_kp_1) \otimes 1 + \frac{1}{2}(x + x^3) \otimes (p_1C_k + C_kp_1)$$

$$+ \frac{1}{2}(x - x^3) \otimes (p_2C_k + C_kp_2),$$

$$\Delta(p_2C_k + C_kp_2) = (p_2C_k + C_kp_2) \otimes 1 + \frac{1}{2}(x + x^3) \otimes (p_2C_k + C_kp_2)$$

$$+ \frac{1}{2}(x - x^3) \otimes (p_1C_k + C_kp_1).$$

Since

$$t(p_1C_k + C_kp_1) = (p_1C_k + C_kp_1)t, \qquad t(p_2C_k + C_kp_2) = -(p_2C_k + C_kp_2)t,$$

we have the relation (6.68) holds in A.

Similarly, the relations (6.69), (6.70), (6.71) hold in A. Then there is a surjective Hopf homomorphism from $\mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2)$ to A. By the Diamond Lemma, we have that the linear basis of $\mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2)$ is

$$\{C_{k}^{\gamma_{k}}D_{\ell}^{\eta_{\ell}}G_{s}^{\lambda_{s}}H_{r}^{\mu_{r}}p_{1}^{i}p_{2}^{j}x^{e}y^{f}t^{g}; \ \gamma_{k},\eta_{\ell},\lambda_{s},\mu_{r},i,j,f,g\in\mathbb{I}_{0,1},e\in\mathbb{I}_{0,3}\}.$$

Then dim $A = \dim \mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2)$ and whence $A \cong \mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2)$.

(2) Suppose that $\Phi : \mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2) \to \mathfrak{U}_2(n_1, n_2, n_3, n_4; I_2')$ is an isomorphism of Hopf algebras. By Proposition 6.6, we know that $\Phi|_H \in \{\tau_1, \cdots, \tau_{16}\}$. If $\Phi|_H \in \{\tau_1, \tau_2, \tau_3, \tau_4\}$, then

$$\begin{split} \Phi(p_1) &= zp'_1, \quad \Phi(p_2) = zp'_2, \quad \Phi(C_k) = \sum_{k'=1}^{n_1} c_{k'k} C'_{k'}, \\ \Phi(D_\ell) &= \sum_{\ell'=1}^{n_2} d_{\ell'\ell} D'_{\ell'}, \quad \Phi(G_s) = \sum_{s'=1}^{n_3} g_{s's} G'_{s'}, \quad \Phi(H_r) = \sum_{r'=1}^{n_4} h_{r'r} H'_{r'} \end{split}$$

Whence the relations $z^2v' = v$, (6.44), (6.46), (6.74) hold. Similarly, Φ is an isomorphism of Hopf algebras if and only if

when $\Phi|_H \in \{\tau_5, \tau_6, \tau_7, \tau_8\}$, the relations $z^2 v' = -v$, (6.48), (6.75) hold;

when $\Phi|_H \in \{\tau_9, \tau_{10}, \tau_{11}, \tau_{12}\}$, the relations $z^2 v' = v$, (6.50), (6.52), (6.76) hold;

when $\Phi|_H \in \{\tau_{13}, \tau_{14}, \tau_{15}, \tau_{16}\}$, the relations $z^2 v' = -v$, (6.54), (6.77) hold.

The proof of (3) is completely analogous.

Definition 6.11 For n_1 , n_2 , n_3 , $n_4 \in \mathbb{N}^*$ with $\sum_{i=1}^4 n_i \ge 0$ and a set

$$I_4 = \{ (\alpha_{i_1,i_2})_{n_1 \times n_1}, \ (\gamma_{k_1,k_2})_{n_2 \times n_2}, \ (\zeta_{m_1,m_2})_{n_3 \times n_3}, \ (\lambda_{s_1,s_2})_{n_4 \times n_4}, \ \nu \},$$

we denote by $\mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4)$ the algebra that is generated by $x, y, t, p_1, p_2, \{A_i\}_{i=1,..,n_1}, \{C_k\}_{k=1,..,n_2}, \{E_m\}_{m=1,..,n_3}, \{G_s\}_{s=1,..,n_4}$ satisfying the relations (3.2) - (3.4), (6.20), (6.22), (6.24), (6.26), (6.28), (6.30), (6.32), (6.34), (6.67) and

$$xp_{1} = p_{1}x, \ xp_{2} = -p_{2}x, \ yp_{1} = p_{1}y, \ yp_{2} = p_{2}y, \ tp_{1} = -p_{1}t, \ tp_{2} = -p_{2}t,$$

$$A_{i}C_{k} - C_{k}A_{i} = 0, \quad A_{i}E_{m} + E_{m}A_{i} = 0, \quad A_{i}G_{s} - G_{s}A_{i} = 0,$$

$$C_{k}E_{m} - E_{m}C_{m} = 0, \quad C_{k}G_{s} + G_{s}C_{m} = 0, \quad E_{m}G_{s} - G_{s}E_{m} = 0,$$

$$p_{1}A_{i} - A_{i}p_{1} = 0, \quad p_{2}A_{i} + A_{i}p_{2} = 0, \quad p_{1}C_{k} - C_{k}p_{1} = 0, \quad p_{2}C_{k} + C_{k}p_{2} = 0,$$

$$p_{1}E_{m} - E_{m}p_{1} = 0, \quad p_{2}E_{m} + E_{m}p_{2} = 0, \quad p_{1}G_{s} - G_{s}p_{1} = 0, \quad p_{2}G_{s} + G_{s}p_{2} = 0.$$
(6.79)

Definition 6.12 For n_1 , n_2 , n_3 , $n_4 \in \mathbb{N}^*$ with $\sum_{i=1}^4 n_i \ge 0$ and a set

$$I_5 = \{ \alpha_{i_1, i_2}, \ \eta_{\ell_1, \ell_2}, \ \zeta_{m_1, m_2}, \mu_{r_1, r_2}, \ \nu, \ \lambda_i, \ \kappa_\ell, \ \iota_m, \ \theta_r \}$$

with $i, i_1, i_2 \in \{1, ..., n_1\}, \ell, \ell_1, \ell_2 \in \{1, ..., n_2\}, m, m_1, m_2 \in \{1, ..., n_3\}, r, r_1, r_2 \in \{1, ..., n_4\},$ denote by $\mathfrak{U}_5(n_1, n_2, n_3, n_4; I_5)$ the algebra that is generated by $x, y, t, p_1, p_2, \{A_i\}, \{D_\ell\}, \{E_m\}, \{H_r\}$ satisfying the relations (3.2) - (3.4), (6.20), (6.23), (6.24), (6.27), (6.28), (6.31), (6.32), (6.35) and

$$xp_{1} = p_{1}x, \ xp_{2} = -p_{2}x, \ yp_{1} = -p_{1}y, \ yp_{2} = -p_{2}y, \ tp_{1} = -p_{1}x^{2}t, \ tp_{2} = p_{2}x^{2}t,$$

$$A_{i}D_{\ell} - D_{\ell}A_{i} = 0, \quad A_{i}E_{m} + E_{m}A_{i} = 0, \quad A_{i}H_{r} - H_{r}A_{i} = 0,$$

$$D_{\ell}E_{m} - E_{m}D_{\ell} = 0, \quad D_{\ell}H_{r} + H_{r}D_{\ell} = 0, \quad E_{m}H_{r} - H_{r}E_{m} = 0,$$

$$p_{1}^{2} = \nu(1 - x^{2}), \quad p_{2}^{2} = -\nu(1 - x^{2}), \quad p_{1}p_{2} - p_{2}p_{1} = 0,$$

$$p_{1}A_{i} - A_{i}p_{1} = \lambda_{i}(xyt - x^{3}yt), \quad p_{2}A_{i} + A_{i}p_{2} = \lambda_{i}(2 - xyt - x^{3}yt),$$

$$(6.81)$$

$$p_1 D_\ell + D_\ell p_1 = \kappa_\ell (2 - xyt - x^3 yt), \quad p_2 D_\ell - D_\ell p_2 = \kappa_\ell (xyt - x^3 yt), \tag{6.82}$$

$$p_1 E_m - E_m p_1 = \iota_m (xyt - x^3yt), \quad p_2 E_m + E_m p_2 = \iota_m (-2 + xyt + x^3yt), \tag{6.83}$$

$$p_1H_r + H_rp_1 = \theta_r(-2 + xyt + x^3yt), \quad p_2H_r - H_rp_2 = \theta_r(xyt - x^3yt).$$
(6.84)

It is a Hopf algebra with its coalgebra structure determined by

Proposition 6.13 (1) Let A be a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is $\Omega_i(n_1, n_2, n_3, n_4)$ for $i \in \{4, 5\}$, then $A \cong \mathfrak{U}_i(n_1, n_2, n_3, n_4; I_i)$.

(2) $\mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4) \cong \mathfrak{U}_4(n_1, n_2, n_3, n_4; I'_4)$ if and only if there exist nonzero parameter z such that $z^2v' = v$ and invertible matrices $(a_{i'_1i_1})_{n_1 \times n_1}$, $(a_{i'_2i_2})_{n_1 \times n_1}$, $(c_{k'_1k_1})_{n_2 \times n_2}$, $(c_{k'_2k_2})_{n_2 \times n_2}$, $(e_{m'_1m_1})_{n_3 \times n_3}$, $(e_{m'_2m_2})_{n_3 \times n_3}$, $(g_{s'_1s_1})_{n_4 \times n_4}$, $(g_{s'_2s_2})_{n_4 \times n_4}$ satisfying the first equations of (6.43) - (6.46) or $n_1 = n_3$, $n_2 = n_4$ satisfying the first equations of (6.49) - (6.52) or $n_1 = n_2$, $n_3 = n_4$ satisfying the first equations of (6.55) - (6.58) or $n_1 = n_4$, $n_2 = n_3$ satisfying the first equations of (6.61) - (6.64).

(3) $\mathfrak{U}_{5}(n_{1}, n_{2}, n_{3}, n_{4}; I_{5}) \cong \mathfrak{U}_{5}(n_{1}, n_{2}, n_{3}, n_{4}; I'_{5})$ if and only if there exist nonzero parameter z and invertible matrices $(a_{i'i})_{n_{1} \times n_{1}}, (a_{i'_{1}i_{1}})_{n_{1} \times n_{1}}, (a_{i'_{2}i_{2}})_{n_{1} \times n_{1}}, (d_{\ell'\ell})_{n_{2} \times n_{2}}, (d_{\ell'_{1}\ell_{1}})_{n_{2} \times n_{2}}, (d_{\ell'_{1}\ell_{1}})_{n_{3} \times n_{3}}, (e_{m'_{1}m_{1}})_{n_{3} \times n_{3}}, (e_{m'_{2}m_{2}})_{n_{3} \times n_{3}}, (h_{r'r})_{n_{4} \times n_{4}}, (h_{r'_{1}r_{1}})_{n_{4} \times n_{4}}, (h_{r'_{2}r_{2}})_{n_{4} \times n_{4}}$ satisfying $z^{2}v' = v$, the first equations of (6.43), (6.45), the second equations of (6.44), (6.46) and

$$z\sum_{i'=1}^{n_1} a_{i'i}\lambda'_{i'} = \lambda_i, \quad z\sum_{\ell'=1}^{n_2} d_{\ell'\ell}\kappa'_{\ell'} = \kappa_\ell, \quad z\sum_{m'=1}^{n_3} e_{m'm}\iota'_{m'} = \iota_m, \quad z\sum_{r'=1}^{n_4} h_{r'r}\theta'_{r'} = \theta_r.$$
(6.85)

or

$$z\sum_{i'=1}^{n_1} a_{i'i}\lambda'_{i'} = -\lambda_i, \quad z\sum_{\ell'=1}^{n_2} d_{\ell'\ell}\kappa'_{\ell'} = \kappa_\ell, \quad z\sum_{m'=1}^{n_3} e_{m'm}\iota'_{m'} = -\iota_m, \quad z\sum_{r'=1}^{n_4} h_{r'r}\theta'_{r'} = \theta_r. \quad (6.86)$$

or $n_1 = n_3$, $n_2 = n_4$ satisfying $z^2 v' = v$, the first equations of (6.49), (6.51), the second equations of (6.50), (6.52) and

$$z\sum_{m'=1}^{n_1} a_{m'i}\iota'_{m'} = -\lambda_i, \ z\sum_{r'=1}^{n_2} d_{r'\ell}\theta'_{r'} = -\kappa_\ell, \ z\sum_{i'=1}^{n_1} e_{i'm}\lambda'_{i'} = -\iota_m, \ z\sum_{\ell'=1}^{n_2} h_{\ell'r}\kappa'_{\ell'} = -\theta_r.$$
(6.87)

or

$$z\sum_{m'=1}^{n_1} a_{m'i}\iota'_{m'} = \lambda_i, \ z\sum_{r'=1}^{n_2} d_{r'\ell}\theta'_{r'} = -\kappa_\ell, \ z\sum_{i'=1}^{n_1} e_{i'm}\lambda'_{i'} = \iota_m, \ z\sum_{\ell'=1}^{n_2} h_{\ell'r}\kappa'_{\ell'} = -\theta_r.$$
(6.88)

or $n_1 = n_2$, $n_3 = n_4$ satisfying $z^2 v' = -v$, the first equation of (6.60), the second equation of (6.59) and

$$z\sum_{\ell'=1}^{n_1} a_{\ell'i} \kappa_{\ell'}^{'} = \lambda_i, \ z\sum_{i'=1}^{n_1} d_{i'\ell} \lambda_{i'}^{'} = \kappa_\ell, \ z\sum_{r'=1}^{n_3} e_{r'm} \theta_{r'}^{'} = \iota_m, \ z\sum_{m'=1}^{n_3} h_{m'r} \iota_{m'}^{'} = \theta_r.$$
(6.89)

or

$$z\sum_{\ell'=1}^{n_1} a_{\ell'i} \kappa_{\ell'}^{'} = -\lambda_i, \ z\sum_{i'=1}^{n_1} d_{i'\ell} \lambda_{i'}^{'} = \kappa_\ell, \ z\sum_{r'=1}^{n_3} e_{r'm} \theta_{r'}^{'} = -\iota_m, \ z\sum_{m'=1}^{n_3} h_{m'r} \iota_{m'}^{'} = \theta_r.$$
(6.90)

or $n_1 = n_4$, $n_2 = n_3$ satisfying $z^2 v' = -v$, the first equation of (6.66), the second equation of (6.65) and

$$z\sum_{r'=1}^{n_1} a_{r'i} \theta'_{r'} = -\lambda_i, \ z\sum_{m'=1}^{n_2} d_{m'\ell} \iota'_{m'} = -\kappa_\ell, \ z\sum_{\ell'=1}^{n_2} e_{\ell'm} \kappa'_{\ell'} = -\iota_m, \ z\sum_{i'=1}^{n_1} h_{i'r} \lambda'_{i'} = -\theta_r.$$
(6.91)

or

$$z\sum_{r'=1}^{n_1} a_{r'i}\theta_{r'}^{'} = \lambda_i, \ z\sum_{m'=1}^{n_2} d_{m'\ell}\iota_{m'}^{'} = -\kappa_\ell, \ z\sum_{\ell'=1}^{n_2} e_{\ell'm}\kappa_{\ell'}^{'} = \iota_m, \ z\sum_{i'=1}^{n_1} h_{i'r}\lambda_{i'}^{'} = -\theta_r.$$
(6.92)

Proof. (1) We prove the claim only for $\Omega_4(n_1, n_2, n_3, n_4)$, since the proof for $\Omega_5(n_1, n_2, n_3, n_4)$ follows the same lines. By Theorem 6.1, we have that $gr(A) \cong \mathcal{B}(\Omega_4(n_1, n_2, n_3, n_4)) \sharp H$. Let $M_3 = \Bbbk\{p_1, p_2\}$. Similar to Proposition 6.10, we only need to prove that the relations (6.78), (6.79) hold in A. Since $\Delta(A_i) = A_i \otimes 1 + x \otimes A_i$, we have that

$$\Delta(p_1A_i - A_ip_1) = (p_1A_i - A_ip_1) \otimes 1 + \frac{1}{2}(xt + x^3t) \otimes (p_1A_i - A_ip_1)$$

+ $\frac{1}{2}(x^3yt - xyt) \otimes (p_2A_i + A_ip_2),$
$$\Delta(p_2A_i + A_ip_2) = (p_2A_i + A_ip_2) \otimes 1 + \frac{1}{2}(xyt + x^3yt) \otimes (p_2A_i + A_ip_2)$$

+ $\frac{1}{2}(x^3t - xt) \otimes (p_1A_i - A_ip_1).$

If the relations $p_1A_i - A_ip_1 = 0$ admit non-trivial deformations, then $p_1A_i - A_ip_1 \in A_{[1]}$ is a linear combination of

 $\{x^{j}y^{l}t^{r}, A_{i}x^{j}y^{l}t^{r}, C_{k}x^{j}y^{l}t^{r}, E_{m}x^{j}y^{l}t^{r}, G_{s}x^{j}y^{l}t^{r}, p_{1}x^{j}y^{l}t^{r}, p_{2}x^{j}y^{l}t^{r}, j \in \mathbb{I}_{0,3}, l, r \in \mathbb{I}_{0,1}\}.$ Since

 $y(p_1A_i - A_ip_1) = -(p_1A_i - A_ip_1)y, \quad t(p_1A_i - A_ip_1) = -(p_1A_i - A_ip_1)x^2t,$

then $p_1A_i - A_ip_1 = 0$ hold in A. Whence by the above two equations we have that the relations $p_2A_i + A_ip_2 = 0$ hold in A.

Similarly, other relations hold in A. Then there is a surjective Hopf morphism from $\mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4)$ to A. By the Diamond Lemma, the linear basis of $\mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4)$ is $\{A_i^{\alpha_i}C_k^{\gamma_k}E_m^{\zeta_m}G_s^{\lambda_s}p_1^jp_2^lx^ey^ft^g\}$ for all parameters $\alpha_i, \gamma_k, \zeta_m, \lambda_s, j, l, f, g \in \mathbb{I}_{0,1}, e \in \mathbb{I}_{0,3}$. Then dim $A = \dim \mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4)$ and whence $A \cong \mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4)$.

(2) Suppose that $\Phi : \mathfrak{U}_4(n_1, n_2, n_3, n_4; I_4) \to \mathfrak{U}_4(n_1, n_2, n_3, n_4; I'_4)$ is an isomorphism of Hopf algebras. Similar to the proof of Proposition 6.10, $\Phi|_H \in \{\tau_1, \tau_3, \tau_9, \tau_{11}, \tau_{17}, \tau_{19}, \tau_{25}, \tau_{27}\}$. If $\Phi|_H = \tau_1$, then

$$\Phi(p_1) = zp'_1, \quad \Phi(p_2) = zp'_2, \quad \Phi(A_i) = \sum_{i'=1}^{n_1} a_{i'i}A'_{i'},$$

$$\Phi(C_k) = \sum_{k'=1}^{n_2} c_{k'k} C'_{k'}, \ \Phi(E_m) = \sum_{m'=1}^{n_3} e_{m'm} E'_{m'}, \ \Phi(G_s) = \sum_{s'=1}^{n_4} g_{s's} G'_{s'}.$$

Whence the first equations of (6.43) - (6.46) hold. Similarly, Φ is an isomorphism of Hopf algebras if and only if

when $\Phi|_{H} = \tau_{3}$, the first equations of (6.43) - (6.46) hold; when $\Phi|_{H} \in {\tau_{9}, \tau_{11}}$, the first equations of (6.49) - (6.52) hold; when $\Phi|_{H} \in {\tau_{17}, \tau_{19}}$, the first equations of (6.55) - (6.58) hold; when $\Phi|_{H} \in {\tau_{25}, \tau_{27}}$, the first equations of (6.61) - (6.64) hold.

The proof of (3) is completely analogous.

Definition 6.14 For a set of parameters $I_{14} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{14}(I_{14})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4) and

$$xp_1 = p_1x, yp_1 = -p_1y, tp_1 = p_1x^2t, xp_2 = p_2x, yp_2 = -p_2y, tp_2 = -p_2x^2t,$$
 (6.93)

$$xq_1 = q_1x, \quad yq_1 = -q_1y, \quad tq_1 = q_1x^2t, \quad xq_2 = q_2x, \quad yq_2 = -q_2y, \quad tq_2 = -q_2x^2t,$$
 (6.94)

$$p_1^2 = \lambda(1 - x^2), \qquad p_2^2 = -\lambda(1 - x^2), \qquad p_1 p_2 + p_2 p_1 = 0,$$
(6.95)

$$q_1^2 = \mu(1 - x^2), \qquad q_2^2 = -\mu(1 - x^2), \qquad q_1q_2 + q_2q_1 = 0,$$
(6.96)

$$p_1q_1 + q_1p_1 = \alpha(1 - x^2), \ p_2q_2 + q_2p_2 = -\alpha(1 - x^2), \ p_1q_2 + q_2p_1 = p_2q_1 + q_1p_2 = 0.$$
 (6.97)

It is a Hopf algebra with its coalgebra structure determined by

Definition 6.15 Denote by $\mathfrak{U}_{15}(\lambda, \mu)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.93), (6.95), (6.96) and

$$xq_1 = -q_1x, \ yq_1 = -q_1y, \ tq_1 = q_1x^2t, \ xq_2 = -q_2x, \ yq_2 = -q_2y, \ tq_2 = -q_2x^2t,$$
 (6.98)

$$p_1q_1 + q_1p_1 = 0, \quad p_2q_2 + q_2p_2 = 0, \quad p_1q_2 + q_2p_1 = 0, \quad p_2q_1 + q_1p_2 = 0.$$
 (6.99)

It is a Hopf algebra with the same coalgebra structure as $\mathfrak{U}_{14}(I_{14})$.

Definition 6.16 Denote by $\mathfrak{U}_{16}(\lambda, \mu)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.93), (6.95), (6.96) and

$$xq_1 = -q_1x, \quad yq_1 = q_1y, \quad tq_1 = q_1t, \quad xq_2 = -q_2x, \quad yq_2 = q_2y, \quad tq_2 = -q_2t,$$
 (6.100)

$$p_1q_1 - q_1p_1 = 0, \quad p_2q_2 - q_2p_2 = 0, \quad p_1q_2 - q_2p_1 = 0, \quad p_2q_1 - q_1p_2 = 0.$$
 (6.101)

It is a Hopf algebra with its coalgebra structure determined by

Definition 6.17 For a set of parameters $I_{17} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{17}(I_{17})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.95) - (6.97)and

$$xp_1 = p_1x, \quad yp_1 = p_1y, \quad tp_1 = -p_1t, \quad xp_2 = -p_2x, \quad yp_2 = p_2y, \quad tp_2 = -p_2t, \quad (6.102)$$

 $xq_1 = q_1x, \quad yq_1 = q_1y, \quad tq_1 = -q_1t, \quad xq_2 = -q_2x, \quad yq_2 = q_2y, \quad tq_2 = -q_2t.$ (6.103)

It is a Hopf algebra with its coalgebra structure determined by

Definition 6.18 For a set of parameters $I_{18} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{18}(I_{18})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.95), (6.96), (6.102), (6.103) and

$$p_1q_1 + q_1p_1 = \alpha(y + x^2y - 2), \qquad p_2q_2 + q_2p_2 = \alpha(y - x^2y),$$
(6.104)

$$p_1q_2 + q_2p_1 = 0, \qquad p_2q_1 + q_1p_2 = 0.$$
 (6.105)

It is a Hopf algebra with its coalgebra structure determined by

Definition 6.19 Denote by $\mathfrak{U}_{19}(\lambda, \mu)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.95), (6.102) and

$$xq_1 = q_1x, \ yq_1 = -q_1y, \ tq_1 = -q_1x^2t, \ xq_2 = -q_2x, \ yq_2 = -q_2y, \ tq_2 = -q_2x^2t,$$
 (6.106)

$$q_1^2 = \mu(1 - x^2), \qquad q_2^2 = -\mu(1 - x^2), \qquad q_1 q_2 - q_2 q_1 = 0,$$
(6.107)

$$p_1q_1 + q_1p_1 = 0, \quad p_2q_2 - q_2p_2 = 0, \quad p_1q_2 + q_2p_1 = 0, \quad p_2q_1 - q_1p_2 = 0.$$
 (6.108)

It is a Hopf algebra with its coalgebra structure determined by

Proposition 6.20 (1) Suppose A is a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is isomorphic to Ω_i for $i \in \{14, 15, 16, 17, 18, 19\}$, then $A \cong$ $\mathfrak{U}_{14}(I_{14})$ or $\mathfrak{U}_{15}(\lambda, \mu)$ or $\mathfrak{U}_{16}(\lambda, \mu)$ or $\mathfrak{U}_{17}(I_{17})$, $\mathfrak{U}_{18}(I_{18})$, $\mathfrak{U}_{19}(\lambda, \mu)$.

(2) $\mathfrak{U}_{17}(I_{17}) \cong \mathfrak{U}_{17}(I'_{17})$ if and only if there exist nonzero parameters a_1, a_2, b_1, b_2 satisfying (6.109). $\mathfrak{U}_{14}(I_{14}) \cong \mathfrak{U}_{14}(I'_{14})$ if and only if there exist nonzero parameters a_1, a_2, b_1, b_2 such that

$$a_{1}^{2}\lambda^{'} + a_{1}a_{2}\alpha^{'} + a_{2}^{2}\mu^{'} = \lambda, \qquad b_{1}^{2}\lambda^{'} + b_{1}b_{2}\alpha^{'} + b_{2}^{2}\mu^{'} = \mu,$$

$$2a_{1}b_{1}\lambda^{'} + 2a_{2}b_{2}\mu^{'} + (a_{1}b_{2} + a_{2}b_{1})\alpha^{'} = \alpha,$$
(6.109)

or satisfying

$$a_{1}^{2}\lambda^{'} + a_{1}a_{2}\alpha^{'} + a_{2}^{2}\mu^{'} = -\lambda, \qquad b_{1}^{2}\lambda^{'} + b_{1}b_{2}\alpha^{'} + b_{2}^{2}\mu^{'} = -\mu,$$

$$2a_{1}b_{1}\lambda^{'} + 2a_{2}b_{2}\mu^{'} + (a_{1}b_{2} + a_{2}b_{1})\alpha^{'} = -\alpha.$$
(6.110)

(3) $\mathfrak{U}_{18}(I_{18}) \cong \mathfrak{U}_{18}(I'_{18})$ if and only if there exist nonzero parameters a_1, a_2 such that

$$a_1^{\ 2}\lambda' = \lambda, \qquad a_2^{\ 2}\mu' = \mu, \qquad a_1a_2\alpha' = \alpha,$$
 (6.111)

or satisfying

$$a_1^{\ 2}\mu' = \lambda, \qquad a_2^{\ 2}\lambda' = \mu, \qquad a_1a_2\alpha' = \alpha.$$
 (6.112)

(4) Let $i \in \{15, 16, 19\}$. $\mathfrak{U}_i(\lambda, \mu) \cong \mathfrak{U}_i(1, 1)$ for $\lambda, \mu \neq 0$, and $\mathfrak{U}_i(1, 1) \ncong \mathfrak{U}_i(0, 0)$.

Proof. (1) We prove the claim for Ω_{14} . The proofs for others follow the same lines. By Theorem 6.1, we have that $gr(A) \cong \mathcal{B}(\Omega_{14}) \sharp H$. Let $M_1 = \Bbbk\{p_1, p_2\} = \Bbbk\{q_1, q_2\}$. By Proposition 6.10, we only need to prove that (6.97) holds in A. After a direct computation, we have that

$$\Delta(p_1q_1 + q_1p_1) = (p_1q_1 + q_1p_1) \otimes 1 + \frac{1}{2}(1 + x^2) \otimes (p_1q_1 + q_1p_1)$$

$$+ \frac{1}{2}(1 - x^2) \otimes (p_2q_2 + q_2p_2),$$

$$\Delta(p_2q_2 + q_2p_2) = (p_2q_2 + q_2p_2) \otimes 1 + \frac{1}{2}(1 + x^2) \otimes (p_2q_2 + q_2p_2)$$

$$+ \frac{1}{2}(1 - x^2) \otimes (p_1q_1 + q_1p_1).$$

So $p_1q_1 + q_1p_1 = \alpha(1 - x^2)$, $p_2q_2 + q_2p_2 = -\alpha(1 - x^2)$ for some $\alpha \in \mathbb{k}$. Similarly, other relations hold in A. Then there is a surjective Hopf homomorphism from $\mathfrak{U}_{14}(I_{14})$ to A. We can observe that all elements of $\mathfrak{U}_{14}(I_{14})$ can be expressed by linear combinations of

$$\{p_1^i p_2^j q_1^k q_2^l x^e y^f t^g; \ i, j, k, l, f, g \in \mathbb{I}_{0,1}, \ e \in \mathbb{I}_{0,3}\}$$

In fact, according to the Diamond Lemma, the set is a basis of $\mathfrak{U}_{14}(I_{14})$. Then dim $A = \dim \mathfrak{U}_{14}(I_{14})$, whence $A \cong \mathfrak{U}_{14}(I_{14})$.

(2) Suppose $\Phi : \mathfrak{U}_{14}(I_{14}) \to \mathfrak{U}_{14}(I'_{14})$ is an isomorphism of Hopf algebras. Similar to the proof of Proposition 6.10, $\Phi|_H \in \{\tau_1 - \tau_{16}\}$: when $\Phi|_H \in \{\tau_1 - \tau_4, \tau_9 - \tau_{12}\}$,

$$\Phi(p_1) = a_1 p_1' + a_2 q_1', \ \Phi(p_2) = a_1 p_2' + a_2 q_2', \ \Phi(q_1) = b_1 p_1' + b_2 q_1', \ \Phi(q_2) = b_1 p_2' + b_2 q_2'.$$

then the relation (6.109) holds; when $\Phi|_H \in \{\tau_5 - \tau_8, \tau_{13} - \tau_{16}\},\$

$$\Phi(p_1) = a_1 p_2' + a_2 q_2', \ \Phi(p_2) = a_1 p_1' + a_2 q_1', \ \Phi(q_1) = b_1 p_2' + b_2 q_2', \ \Phi(q_2) = b_1 p_1' + b_2 q_1',$$

the relation (6.110) holds.

For $\mathfrak{U}_{17}(I_{17})$, the proof is completely analogous.

(3) Suppose $\Phi : \mathfrak{U}_{18}(I_{18}) \to \mathfrak{U}_{18}(I'_{18})$ is an isomorphism of Hopf algebras. When $\Phi|_H \in \{\tau_1, \tau_9, \tau_{17}, \tau_{25}\},\$

 $\Phi(p_1) = a_1 p'_1, \quad \Phi(p_2) = a_1 p'_2, \quad \Phi(q_1) = a_2 q'_1, \quad \Phi(q_2) = a_2 q'_2;$

when $\Phi|_H \in \{\tau_3, \tau_{11}, \tau_{19}, \tau_{27}, \},\$

$$\Phi(p_1) = a_1 p'_1, \quad \Phi(p_2) = -a_1 p'_2, \quad \Phi(q_1) = a_2 q'_1, \quad \Phi(q_2) = -a_2 q'_2,$$

then the relation (6.111) holds. When $\Phi|_H \in \{\tau_5, \tau_{13}, \tau_{21}, \tau_{29}\},\$

$$\Phi(p_1) = a_1 q'_1, \quad \Phi(p_2) = a_1 q'_2, \quad \Phi(q_1) = a_2 p'_1, \quad \Phi(q_2) = a_2 p'_2;$$

when $\Phi|_H \in \{\tau_7, \tau_{15}, \tau_{23}, \tau_{31}, \},\$

$$\Phi(p_1) = a_1 q'_1, \quad \Phi(p_2) = -a_1 q'_2, \quad \Phi(q_1) = a_2 p'_1, \quad \Phi(q_2) = -a_2 p'_2,$$

then the relation (6.112) holds.

(4) When $\lambda, \ \mu \neq 0, \ \Phi : \mathfrak{U}_{15}(1,1) \to \mathfrak{U}_{15}(\lambda,\mu)$ by $\Phi|_H = id, \ p_i \mapsto \sqrt{\lambda}p_i, \ q_i \mapsto \sqrt{\mu}q_i$ for i = 1, 2. For $\mathfrak{U}_{16}(\lambda,\mu), \ \mathfrak{U}_{19}(\lambda,\mu)$, the proofs are completely analogous. \Box

Definition 6.21 For a set of parameters $I_{20} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{20}(I_{20})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4) and

$$xp_1 = p_1x, \ yp_1 = -p_1y, \ tp_1 = -p_1x^2t, \ xp_2 = -p_2x, \ yp_2 = -p_2y, \ tp_2 = p_2x^2t, \tag{6.113}$$

$$xq_1 = q_1x, \ yq_1 = -q_1y, \ tq_1 = -q_1x^2t, \ xq_2 = -q_2x, \ yq_2 = -q_2y, \ tq_2 = q_2x^2t,$$
 (6.114)

$$p_1^2 = \lambda(1 - x^2), \qquad p_2^2 = -\lambda(1 - x^2), \qquad p_1 p_2 - p_2 p_1 = 0,$$
(6.115)

$$q_1^2 = \mu(1 - x^2), \qquad q_2^2 = -\mu(1 - x^2), \qquad q_1q_2 - q_2q_1 = 0,$$
 (6.116)

$$p_1q_1 + q_1p_1 = \alpha(1 - x^2), \qquad p_2q_2 + q_2p_2 = -\alpha(1 - x^2),$$
(6.117)

$$p_1q_2 - q_2p_1 = 0, \qquad p_2q_1 - q_1p_2 = 0.$$
 (6.118)

It is a Hopf algebra with the same coalgebra structure as $\mathfrak{U}_{17}(I_{17})$.

Definition 6.22 Denote by $\mathfrak{U}_{21}(\lambda, \mu)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.113), (6.115), (6.116) and

$$xq_1 = q_1x, \ yq_1 = -q_1y, \ tq_1 = q_1x^2t, \ xq_2 = -q_2x, \ yq_2 = -q_2y, \ tq_2 = -q_2x^2t,$$
 (6.119)

$$p_1q_1 - q_1p_1 = 0, \quad p_2q_2 - q_2p_2 = 0, \quad p_1q_2 + q_2p_1 = 0, \quad p_2q_1 + q_1p_2 = 0.$$
 (6.120)

It is a Hopf algebra with the same coalgebra structure as $\mathfrak{U}_{18}(I_{18})$.

Definition 6.23 For a set of parameters $I_{22} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{22}(I_{22})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.95), (6.96),(6.100) and

 $xp_{1} = -p_{1}x, \quad yp_{1} = p_{1}y, \quad tp_{1} = p_{1}t, \quad xp_{2} = -p_{2}x, \quad yp_{2} = p_{2}y, \quad tp_{2} = -p_{2}t, \quad (6.121)$ $p_{1}q_{1} + q_{1}p_{1} = p_{2}q_{2} + q_{2}p_{2} = \alpha(1 - x^{2}), \quad p_{1}q_{2} + q_{2}p_{1} = 0, \quad p_{2}q_{1} + q_{1}p_{2} = 0. \quad (6.122)$

It is a Hopf algebra with its coalgebra structure determined by

Definition 6.24 For a set of parameters $I_{23} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{23}(I_{23})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.95), (6.96), (6.100), (6.121) and

$$p_1q_1 + q_1p_1 = \alpha(y + x^2y - 2), \qquad p_2q_2 + q_2p_2 = \alpha(x^2y - y),$$
(6.123)

$$p_1q_2 + q_2p_1 = 0, \qquad p_2q_1 + q_1p_2 = 0.$$
 (6.124)

It is a Hopf algebra with its coalgebra structure determined by

Similar to Proposition 6.20, we have the following

Proposition 6.25 (1) Suppose A is a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is isomorphic to Ω_i for $i \in \{20, 21, 22, 23\}$, then $A \cong \mathfrak{U}_i(I_i)$.

(2) $\mathfrak{U}_{20}(I_{20}) \cong \mathfrak{U}_{20}(I'_{20}), \mathfrak{U}_{22}(I_{22}) \cong \mathfrak{U}_{22}(I'_{22})$ if and only if there exist nonzero parameters a_1, a_2, b_1, b_2 satisfying (6.109) or (6.110).

(3) $\mathfrak{U}_{23}(I_{23}) \cong \mathfrak{U}_{23}(I'_{23})$ if and only if there exist nonzero parameters a_1 , a_2 satisfying (6.111) or (6.112) or satisfying

$$a_1^2 \lambda' = -\lambda, \qquad a_2^2 \mu' = -\mu, \qquad \alpha = 0,$$
 (6.125)

or satisfying

$$a_1^2 \mu' = -\lambda, \qquad a_2^2 \lambda' = -\mu, \qquad \alpha = 0.$$
 (6.126)

(4) $\mathfrak{U}_{21}(\lambda,\mu) \cong \mathfrak{U}_{21}(1,1)$ for $\lambda,\mu \neq 0$, and $\mathfrak{U}_{21}(1,1) \ncong \mathfrak{U}_{21}(0,0)$.

Definition 6.26 Denote by $\mathfrak{U}_{24}(\lambda)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4) and

$$xp_1 = \xi p_1 x, \ yp_1 = -p_1 y, \ tp_1 = p_2 x^2 t, \ xp_2 = -\xi p_2 x, \ yp_2 = -p_2 y, \ tp_2 = p_1 x^2 t,$$
 (6.127)

$$xq_1 = \xi q_1 x, \ yq_1 = -q_1 y, \ tq_1 = q_2 x^2 t, \ xq_2 = -\xi q_2 x, \ yq_2 = -q_2 y, \ tq_2 = q_1 x^2 t,$$
 (6.128)

$$p_1^2 = 0, \quad p_2^2 = 0, \quad p_1 p_2 + p_2 p_1 = 0, \quad q_1^2 = 0, \quad q_2^2 = 0, \quad q_1 q_2 + q_2 q_1 = 0,$$
 (6.129)

$$p_1q_1 + q_1p_1 = 0, \qquad p_2q_2 + q_2p_2 = 0,$$
(6.130)

$$p_1q_2 + q_2p_1 = \lambda(1 - x^2y), \qquad p_2q_1 + q_1p_2 = -\lambda(1 - x^2y).$$
 (6.131)

It is a Hopf algebra with its coalgebra structure determined by

Definition 6.27 For a set of parameters $I_{26} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{26}(I_{26})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4) and

- $xp_1 = \xi p_1 x, \ yp_1 = p_1 y, \ tp_1 = p_2 t, \ xp_2 = -\xi p_2 x, \ yp_2 = p_2 y, \ tp_2 = p_1 t,$ (6.132)
- $xq_1 = \xi q_1 x, \ yq_1 = q_1 y, \ tq_1 = q_2 t, \ xq_2 = -\xi q_2 x, \ yq_2 = q_2 y, \ tq_2 = q_1 t,$ (6.133)

$$p_1^2 = 0, \qquad p_2^2 = 0, \qquad p_1 p_2 + p_2 p_1 = \lambda (1 - y),$$
(6.134)

$$q_1^2 = 0, \qquad q_2^2 = 0, \qquad q_1 q_2 + q_2 q_1 = \mu (1 - y),$$
(6.135)

$$p_1q_1 + q_1p_1 = 0, \quad p_2q_2 + q_2p_2 = 0, \quad p_1q_2 + q_2p_1 = p_2q_1 + q_1p_2 = \alpha(1-y).$$
 (6.136)

It is a Hopf algebra with its coalgebra structure determined by

$$\Delta(p_1) = p_1 \otimes 1 + x^2 \otimes p_1, \qquad \Delta(p_2) = p_2 \otimes 1 + x^2 y \otimes p_2, \Delta(q_1) = q_1 \otimes 1 + x^2 \otimes q_1, \qquad \Delta(q_2) = q_2 \otimes 1 + x^2 y \otimes q_2.$$

Definition 6.28 Denote by $\mathfrak{U}_{27}(\lambda, \mu)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) – (3.4), (6.132), (6.134), (6.135) and

$$xq_1 = -\xi q_1 x, \ yq_1 = q_1 y, \ tq_1 = q_2 t, \ xq_2 = \xi q_2 x, \ yq_2 = q_2 y, \ tq_2 = q_1 t, \tag{6.137}$$

 $p_1q_1 + q_1p_1 = 0, \quad p_2q_2 + q_2p_2 = 0, \quad p_1q_2 + q_2p_1 = 0, \quad p_2q_1 + q_1p_2 = 0.$ (6.138)

It is a Hopf algebra with the same coalgebra structure as $\mathfrak{U}_{26}(I_{26})$.

Proposition 6.29 (1) Suppose A is a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is isomorphic to Ω_i for $i \in \{24, 26, 27\}$, then $A \cong \mathfrak{U}_{24}(\lambda)$ or $\mathfrak{U}_{26}(I_{26})$ or $\mathfrak{U}_{27}(\lambda, \mu)$.

(2) $\mathfrak{U}_{24}(\lambda) \cong \mathfrak{U}_{24}(1), \mathfrak{U}_{27}(\lambda,\mu) \cong \mathfrak{U}_{27}(1,1)$ for $\lambda, \ \mu \neq 0, \ \mathfrak{U}_{24}(1) \not\cong \mathfrak{U}_{24}(0)$ and $\mathfrak{U}_{27}(1,1) \not\cong \mathfrak{U}_{27}(0,0).$

(3) $\mathfrak{U}_{26}(I_{26}) \cong \mathfrak{U}_{26}(I'_{26})$ if and only if there exist nonzero parameters a_1, a_2, b_1, b_2 such that

$$a_{1}^{2}\lambda' + 2a_{1}a_{2}\alpha' + a_{2}^{2}\mu' = \lambda, \quad b_{1}^{2}\lambda' + 2b_{1}b_{2}\alpha' + b_{2}^{2}\mu' = \mu,$$

$$a_{1}b_{1}\lambda' + (a_{1}b_{2} + a_{2}b_{1})\alpha' + a_{2}b_{2}\mu' = \alpha,$$
(6.139)

or satisfying

$$a_{1}^{2}\lambda' + 2a_{1}a_{2}\alpha' + a_{2}^{2}\mu' = -\xi\lambda, \quad b_{1}^{2}\lambda' + 2b_{1}b_{2}\alpha' + b_{2}^{2}\mu' = -\xi\mu, a_{1}b_{1}\lambda' + (a_{1}b_{2} + a_{2}b_{1})\alpha' + a_{2}b_{2}\mu' = -\xi\alpha,$$
(6.140)

or satisfying

$$a_{1}^{2}\lambda' + 2a_{1}a_{2}\alpha' + a_{2}^{2}\mu' = -\lambda, \quad b_{1}^{2}\lambda' + 2b_{1}b_{2}\alpha' + b_{2}^{2}\mu' = -\mu,$$

$$a_{1}b_{1}\lambda' + (a_{1}b_{2} + a_{2}b_{1})\alpha' + a_{2}b_{2}\mu' = -\alpha,$$
(6.141)

or satisfying

$$a_{1}^{2}\lambda' + 2a_{1}a_{2}\alpha' + a_{2}^{2}\mu' = \xi\lambda, \quad b_{1}^{2}\lambda' + 2b_{1}b_{2}\alpha' + b_{2}^{2}\mu' = \xi\mu, a_{1}b_{1}\lambda' + (a_{1}b_{2} + a_{2}b_{1})\alpha' + a_{2}b_{2}\mu' = \xi\alpha.$$
(6.142)

Proof. (1) We prove the claim for Ω_{24} . The proofs for Ω_{26} , Ω_{27} follow the same lines. By Theorem 6.1, we have that $gr(A) \cong \mathcal{B}(\Omega_{24}) \sharp H$. Let $M_{13} = \Bbbk\{p_1, p_2\} = \Bbbk\{q_1, q_2\}$. Similar to Proposition 6.20, we only need to prove that (6.129) – (6.131) in Definition 6.26 hold in A. Since

then p_1^2 , p_2^2 , $p_1q_1 + q_1p_1$, $p_2q_2 + q_2p_2$ are primitive elements and

$$\Delta(p_1p_2 + p_2p_1) = (p_1p_2 + p_2p_1) \otimes 1 + x^2y \otimes (p_1p_2 + p_2p_1), \Delta(p_1q_2 + q_2p_1) = (p_1q_2 + q_2p_1) \otimes 1 + x^2y \otimes (p_1q_2 + q_2p_1), \Delta(p_2q_1 + q_1p_2) = (p_2q_1 + q_1p_2) \otimes 1 + x^2y \otimes (p_2q_1 + q_1p_2).$$

As

$$t(p_1p_2 + p_2p_1) = -(p_1p_2 + p_2p_1)t, \quad t(p_1q_2 + q_2p_1) = -(p_2q_1 + q_1p_2)t,$$

the relations (6.129) – (6.131) hold in A. Then there is a surjective Hopf morphism from $\mathfrak{U}_{24}(I_{24})$ to A. We can observe that each element of $\mathfrak{U}_{24}(I_{24})$ can be expressed by a linear combination of $\{p_1^i p_2^j q_1^k q_2^l x^e y^f t^g\}$ for all parameters $i, j, k, l, f, g \in \mathbb{I}_{0,1}, e \in \mathbb{I}_{0,3}$. In fact, according to the Diamond Lemma, the set is a basis of $\mathfrak{U}_{24}(I_{24})$. Then dim $A = \dim \mathfrak{U}_{24}(I_{24})$, whence $A \cong \mathfrak{U}_{24}(I_{24})$.

(2) The proof is the same as that of Proposition 6.20 (4).

(3) Suppose $\Phi : \mathfrak{U}_{26}(I_{26}) \to \mathfrak{U}_{26}(I'_{26})$ is an isomorphism of Hopf algebras. Similar to Proposition 6.20, $\Phi|_H \in \{\tau_1 - \tau_8, \tau_{17} - \tau_{24}\}$. When $\Phi|_H \in \{\tau_1, \tau_5, \tau_{17}, \tau_{21}\}$,

$$\Phi(p_1) = a_1 p'_1 + a_2 q'_1, \quad \Phi(p_2) = a_1 p'_2 + a_2 q'_2, \quad \Phi(q_1) = b_1 p'_1 + b_2 q'_1, \quad \Phi(q_2) = b_1 p'_2 + b_2 q'_2,$$

then the relation (6.139) holds. When $\Phi|_H \in \{\tau_2, \tau_6, \tau_{18}, \tau_{22}\},\$

$$\Phi(p_1) = a_1 p'_1 + a_2 q'_1, \ \Phi(p_2) = \xi(a_1 p'_2 + a_2 q'_2), \ \Phi(q_1) = b_1 p'_1 + b_2 q'_1, \ \Phi(q_2) = \xi(b_1 p'_2 + b_2 q'_2),$$

then the relation (6.140) holds. When $\Phi|_{H} \in \{\tau_{3}, \tau_{7}, \tau_{19}, \tau_{23}\},\$

$$\Phi(p_1) = a_1 p'_1 + a_2 q'_1, \ \Phi(p_2) = -(a_1 p'_2 + a_2 q'_2), \ \Phi(q_1) = b_1 p'_1 + b_2 q'_1, \ \Phi(q_2) = -(b_1 p'_2 + b_2 q'_2),$$

then the relation (6.141) holds. When $\Phi|_H \in \{\tau_4, \tau_8, \tau_{20}, \tau_{24}\},\$

$$\Phi(p_1) = a_1 p'_1 + a_2 q'_1, \ \Phi(p_2) = -\xi(a_1 p'_2 + a_2 q'_2), \ \Phi(q_1) = b_1 p'_1 + b_2 q'_1, \ \Phi(q_2) = -\xi(b_1 p'_2 + b_2 q'_2),$$

then the relation (6.142) holds.

Proposition 6.30 Suppose A is a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is Ω_{25} , then $A \cong \mathcal{B}(\Omega_{25}) \sharp H$.

Proof. Let $M_{13} = \mathbb{k}\{p_1, p_2\}, M_{14} = \mathbb{k}\{q_1, q_2\}$. By Theorem 6.1, we have that $gr(A) \cong \mathcal{B}(\Omega_{25}) \notin H$. Recall that $\mathcal{B}(\Omega_{25}) \notin H$ is the algebra generated by $p_1, p_2, q_1, q_2, x, y, t$ with p_1, p_2, q_1, q_2 satisfying the relation of $\mathcal{B}(\Omega_{25}), x, y, t$ satisfying the relations of H, and all together satisfying the cross relations:

$$xp_1 = \xi p_1 x, \quad yp_1 = -p_1 y, \quad tp_1 = p_2 x^2 t \quad xp_2 = -\xi p_2 x, \quad yp_2 = -p_2 y, \quad tp_2 = p_1 x^2 t, \\ xq_1 = -\xi q_1 x, \quad yq_1 = -q_1 y, \quad tq_1 = q_2 x^2 t \quad xq_2 = \xi q_2 x, \quad yq_2 = -q_2 y, \quad tq_2 = q_1 x^2 t.$$

By Proposition 6.29 (1), we know that the relations $p_1^2 = 0$, $p_2^2 = 0$, $p_1p_2 + p_2p_1 = 0$ hold in A. Similarly, the relations $q_1^2 = 0$, $q_2^2 = 0$, $q_1q_2 + q_2q_1 = 0$ hold in A. After a direct computation, $p_1q_1 + q_1p_1$, $p_2q_2 + q_2p_2$ are primitive elements and $p_1q_2 + q_2p_1$, $p_2q_1 + q_1p_2$ are $(1, x^2y)$ -skew primitive. As

$$x(p_1q_2+q_2p_1) = -(p_1q_2+q_2p_1)x, \quad x(p_2q_1+q_1p_2) = -(p_2q_1+q_1p_2)x,$$

then the relations

$$p_1q_1 + q_1p_1 = 0$$
, $p_2q_2 + q_2p_2 = 0$, $p_1q_2 + q_2p_1 = 0$, $p_2q_1 + q_1p_2 = 0$

hold in A. Therefore, $A \cong gr(A)$.

Definition 6.31 For a set of parameters $I_{28} = \{\lambda, \mu, \alpha\}$, denote by $\mathfrak{U}_{28}(I_{28})$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.132), (6.133)and

$$p_1^2 + p_2^2 = 0, \quad p_1 p_2 = p_2 p_1 = \lambda (1 - y), \quad q_1^2 + q_2^2 = 0, \quad q_1 q_2 = q_2 q_1 = \mu (1 - y),$$
 (6.143)

 $p_1q_1 + q_1p_1 + p_2q_2 + q_2p_2 = 0, \quad p_1q_2 + q_2p_1 + p_2q_1 + q_1p_2 = \alpha(1-y),$ (6.144)

$$p_1q_1 - q_1p_1 - p_2q_2 + q_2p_2 = 0, \quad p_1q_2 - q_2p_1 - p_2q_1 + q_1p_2 = 0.$$
(6.145)

It is a Hopf algebra with its coalgebra structure determined by

$$\begin{split} & \triangle(p_1) = p_1 \otimes 1 + \frac{1}{2}(1+x^2)t \otimes p_1 - \frac{1}{2}(1-x^2)yt \otimes p_2, \\ & \triangle(p_2) = p_2 \otimes 1 + \frac{1}{2}(1+x^2)yt \otimes p_2 - \frac{1}{2}(1-x^2)t \otimes p_1, \\ & \triangle(q_1) = q_1 \otimes 1 + \frac{1}{2}(1+x^2)t \otimes q_1 - \frac{1}{2}(1-x^2)yt \otimes q_2, \\ & \triangle(q_2) = q_2 \otimes 1 + \frac{1}{2}(1+x^2)yt \otimes q_2 - \frac{1}{2}(1-x^2)t \otimes q_1. \end{split}$$

Definition 6.32 Denote by $\mathfrak{U}_{29}(\lambda, \mu)$ the algebra that is generated by $x, y, t, p_1, p_2, q_1, q_2$ satisfying the relations (3.2) - (3.4), (6.132), (6.137), (6.143), (6.145) and

 $p_1q_1 + q_1p_1 + p_2q_2 + q_2p_2 = 0, \quad p_1q_2 + q_2p_1 + p_2q_1 + q_1p_2 = 0.$ (6.146)

It is a Hopf algebra with the same coalgebra structure as $\mathfrak{U}_{28}(I_{28})$.

Proposition 6.33 (1) Suppose A is a finite-dimensional Hopf algebra with coradical H such that its infinitesimal braiding is isomorphic to Ω_{28} or Ω_{29} , then $A \cong \mathfrak{U}_{28}(I_{28})$ or $\mathfrak{U}_{29}(\lambda,\mu)$.

(2) $\mathfrak{U}_{28}(I_{28}) \cong \mathfrak{U}_{28}(I'_{28})$ if and only if there exist nonzero parameters a_1, a_2, b_1, b_2 such that

$$a_{1}^{2}\lambda' + \frac{1}{2}a_{1}a_{2}\alpha' + a_{2}^{2}\mu' = \lambda, \quad b_{1}^{2}\lambda' + \frac{1}{2}b_{1}b_{2}\alpha' + b_{2}^{2}\mu' = \mu,$$

$$4a_{1}b_{1}\lambda' + (a_{1}b_{2} + a_{2}b_{1})\alpha' + 4a_{2}b_{2}\mu' = \alpha,$$
(6.147)

or satisfying

$$a_{1}^{2}\lambda^{'} + \frac{1}{2}a_{1}a_{2}\alpha^{'} + a_{2}^{2}\mu^{'} = -\lambda, \quad b_{1}^{2}\lambda^{'} + \frac{1}{2}b_{1}b_{2}\alpha^{'} + b_{2}^{2}\mu^{'} = -\mu,$$

$$4a_{1}b_{1}\lambda^{'} + (a_{1}b_{2} + a_{2}b_{1})\alpha^{'} + 4a_{2}b_{2}\mu^{'} = -\alpha.$$
(6.148)

(3) $\mathfrak{U}_{29}(\lambda,\mu) \cong \mathfrak{U}_{29}(1,1)$ for $\lambda,\mu \neq 0$, and $\mathfrak{U}_{29}(1,1) \ncong \mathfrak{U}_{29}(0,0)$.

Proof. (1) We prove the claim for Ω_{28} . The proof for Ω_{29} follows the same lines. By Theorem 6.1, we have that $gr(A) \cong \mathcal{B}(\Omega_{28}) \sharp H$. Let $M_{17} = \Bbbk \{p_1, p_2\} = \Bbbk \{q_1, q_2\}$. Similar to Proposition 6.20, we only need to prove that (6.143) - (6.145) in Definition 6.31 hold in A. Let $u = p_1q_1 - q_1p_1 - p_2q_2 + q_2p_2$, $v = p_1q_2 + q_2p_1 + p_2q_1 + q_1p_2$, $w = p_1q_2 - q_2p_1 - p_2q_1 + q_1p_2$. After a direct computation we have that $p_1^2 + p_2^2$, $q_1^2 + q_2^2$, $p_1q_1 + q_1p_1 + p_2q_2 + q_2p_2$ are primitive elements and

As

$$t(p_1p_2 - p_2p_1) = -(p_1p_2 - p_2p_1)t, \quad tu = -ut, \quad tw = -wt,$$

the relations (6.143) - (6.145) hold in A. Then there is a surjective Hopf homomorphism from $\mathfrak{U}_{28}(I_{28})$ to A. We can observe that all elements of $\mathfrak{U}_{28}(I_{28})$ can be expressed by linear combinations of

$$\{(p_1+p_2)^i(p_1-p_2)^j(q_1+q_2)^k(q_1-q_2)^l x^e y^f t^g; i, j, k, l, f, g \in \mathbb{I}_{0,1}, e \in \mathbb{I}_{0,3}\}$$

In fact, according to the Diamond Lemma, the set is a basis of $\mathfrak{U}_{28}(I_{28})$. Then dim $A = \dim \mathfrak{U}_{28}(I_{28})$, whence $A \cong \mathfrak{U}_{28}(I_{28})$.

(2) Suppose $\Phi : \mathfrak{U}_{28}(I_{28}) \to \mathfrak{U}_{28}(I'_{28})$ is an isomorphism of Hopf algebras. Similar to Proposition 6.20, $\Phi|_H \in \{\tau_1, \tau_3, \tau_{13}, \tau_{15}, \tau_{17}, \tau_{19}, \tau_{29}, \tau_{31}\}$. When $\Phi|_H \in \{\tau_1, \tau_{17}\}$, then

$$\Phi(p_1) = a_1 p'_1 + a_2 q'_1, \quad \Phi(p_2) = a_1 p'_2 + a_2 q'_2, \quad \Phi(q_1) = b_1 p'_1 + b_2 q'_1, \quad \Phi(q_2) = b_1 p'_2 + b_2 q'_2;$$

when $\Phi|_{H} \in \{\tau_{13}, \tau_{29}\}$, then

$$\Phi(p_1) = a_1 p'_2 + a_2 q'_2, \ \Phi(p_2) = a_1 p'_1 + a_2 q'_1, \ \Phi(q_1) = b_1 p'_2 + b_2 q'_2, \ \Phi(q_2) = b_1 p'_1 + b_2 q'_1,$$

thus the relation (6.147) holds. When $\Phi|_H \in {\tau_3, \tau_{19}}$, then

$$\Phi(p_1) = a_1 p'_1 + a_2 q'_1, \ \Phi(p_2) = -a_1 p'_2 - a_2 q'_2, \ \Phi(q_1) = b_1 p'_1 + b_2 q'_1, \ \Phi(q_2) = -b_1 p'_2 - b_2 q'_2;$$

when $\Phi|_{H} \in \{\tau_{15}, \tau_{31}\}$, then

$$\Phi(p_1) = a_1 p'_2 + a_2 q'_2, \ \Phi(p_2) = -a_1 p'_1 - a_2 q'_1, \ \Phi(q_1) = b_1 p'_2 + b_2 q'_2, \ \Phi(q_2) = -b_1 p'_1 - b_2 q'_1,$$

thus the relation (6.148) holds.

(3) The proof is the same as that of Proposition 6.20 (4).

Proof of Theorem B. Let M be one of the Yetter-Drinfeld modules listed in Theorem A. Let A be a finite-dimensional Hopf algebra over H such that its infinitesimal braiding is isomorphic to M. By Theorem 6.1, $\operatorname{gr} A \cong \mathcal{B}(M) \sharp H$. Propositions 6.6, 6.10, 6.13, 6.20, 6.25, 6.29, 6.30, 6.33 finish the proof.

ACKNOWLEDGMENT

The authors would like to thank Dr. Yuxing Shi and Dr. Rongchuan Xiong for their helpful discussions. The first author is indebted to Dr. Jiao Zhang for her seminar. The second author is supported by NSERC of Canada. The third author is supported by the NSFC (Grant No. 11771142) and in part by the Science and Technology Commission of Shanghai Municipality (No. 18dz2271000).

References

- N. Andruskiewitsch, I. Angiono and A.G. Iglesias, Liftings of Nichols algebras of diagonal type I. Cartan type A, Int. Math. Res. Not. IMRN (9) (2017), 2793–2884.
- [2] N. Andruskiewitsch, I. Angiono, A.G. Iglesias, A. Masuoka and C. Vay, Lifting via cocycle deformation, J. Pure Appl. Algebra 218 (4) (2014), 684—703.
- [3] N. Andruskiewitsch, G. Carnovale and G.A. García, Finite-dimensional pointed Hopf algebras over finite simple groups of Lie type I. Non-semisimple classes in $PSL_n(q)$, J. Algebra 442 (2015), 36–65.
- [4] —, Finite-dimensional pointed Hopf algebras over finite simple groups of Lie type II: unipotent classes in symplectic groups, Commun. Contemp. Math. 18 (4) (2016), 1550053, 35.
- [5] N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, *Pointed Hopf algebras over some sporadic simple groups*, C. R. Math. Acad. Sci. Paris 348 (2010), 605–608.
- [6] —, Finite-dimensional pointed Hopf algebras with alternating groups are trivial, Ann. Mat. Pura Appl. (4) 190 (2) (2011), 225—245.
- [7] N. Andruskiewitsch, C. Galindo and M. Müller, Examples of finite-dimensional Hopf algebras with the dual Chevalley property, Publ. Mat., 61 (2) (2017): 445-474.
- [8] N. Andruskiewitsch and J.M.J. Giraldi, Nichols algebras that are quantum planes, Linear and Multilinear Algebra 66 (5) (2018), 961—991.
- [9] N. Andruskiewitsch, I. Heckenberger and H.-J. Schneider, The Nichols algebra of a semisimple Yetter-Drinfeld module, Amer. J. Math. 132 (6) (2010), 1493—1547.
- [10] N. Andruskiewitsch and H.-J. Schneider, Lifting of quantum linear spaces and pointed Hopf algebras of order p³, J. Algebra 209 (2) (1998), 658—691.
- [11] —, Finite quantum groups and Cartan matrices, Adv. Math., 154 (1) (2000): 1–45.
- [12] —, Pointed Hopf algebras, New directions in Hopf algebras, 1-68, Math. Sci. Res. Inst. Publ., 43, Cambridge Univ. Press, Cambridge, 2002.
- [13] —, On the classification of finite-dimensional pointed Hopf algebras, Ann. of Math. 171 (1) (2010), 375—417.

- [14] N. Andruskiewitsch and C. Vay, Finite dimensional Hopf algebras over the dual group algebra of the symmetric group in three letters, Comm. Algebra 39 (12) (2011), 4507-4517.
- [15] G.M. Bergman, The diamond lemma for ring theory, Adv. in Math. 29 (2) (1978), 178—218.
- [16] C. Călinescu, S. Dăscălescu, A. Masuoka and C. Menini, Quantum lines over noncocommutative cosemisimple Hopf algebras, J. Algebra 273 (2) (2004), 53—779.
- [17] F. Fantino and G.A. García, On pointed Hopf algebras over dihedral groups, Pacific J. Math. 252 (1) (2011), 69—91.
- [18] X. Fang, On defining ideals and differential algebras of Nichols algebras, J. Algebra 346 (2011), 299—331.
- [19] G.A. García and J.M.J. Giraldi, On Hopf algebras over quantum subgroups, J. Pure Appl. Algebra 223 (2) (2019), 738—768.
- [20] G.A. García and A.G. Iglesias, *Finite-dimensional pointed Hopf algebras over* S₄, Israel J. Math. 183 (2011), 417-444.
- [21] M. Graña, A freeness theorem for Nichols algebras, J. Algebra 231 (1) (2000), 235— 257.
- [22] I. Heckenberger, Classification of arithmetic root systems, Adv. Math. 220 (1) (2009), 59-124.
- [23] I. Heckenberger and H.-J. Schneider, Nichols algebras over groups with finite root system of rank two I, J. Algebra, 324 (11) (2010): 3090-3114.
- [24] N.H. Hu and R.C. Xiong, Some Hopf algebras of dimension 72 without the Chevalley property, arXiv:1612.0498.
- [25] —, On families of Hopf algebras without the dual Chevalley property, Rev. Un. Mat. Argentina 59 (2) (2018), 443—469.
- [26] A.G. Iglesias and C. Vay, Finite-dimensional pointed or copointed Hopf algebras over affine racks, J. Algebra 397 (2014), 379—406.
- [27] I. Kaplansky, *Bialgebras*, Lecture Notes in Mathematics. Department of Mathematics, University of Chicago, Chicago, Ill., 1975.
- [28] Y. Kashina, Classification of semisimple Hopf algebras of dimension 16, J. Algebra 232 (2) (2000), 617—663.
- [29] C. Kassel, Quantum Groups, Graduate Texts in Mathematics, Vol. 155, Springer-Verlag, New York/Berlin, 1995.
- [30] S. Montgomery, Hopf Algebras and their Actions on Rings, CMBS Reg. Conf. Ser. in Math. 82, Amer. Math. Soc., 1993.
- [31] D.E. Radford, The structure of Hopf algebras with a projection, J. Algebra 92 (2) (1985), 322—347.
- [32] —, Hopf Algebras, Knots and Everything 49, World Scientific, 2012.

- [33] Y.X. Shi, Finite-dimensional Hopf algebras over the Kac-Paljutkin algebra H₈, Rev. Un. Mat. Argentina 60 (1) (2019), 265-298.
- [34] R.C. Xiong, Some classification results on finite-dimensional Hopf algebras, Ph. D. Dissertation of East China Normal University (pp. 347), May, 2019.
- [35] M. Sweedler, Hopf Algebras, Benjamin, New York, 1969.