

A Polynomial Kernel for Funnel Arc Deletion Set

Marcelo Garlet Milani
Technische Universität Berlin

November 14, 2019

Abstract

In DIRECTED FEEDBACK ARC SET (DFAS) we search for a set of at most k arcs which intersect every cycle in the input digraph. It is a well-known open problem in parameterized complexity to decide if DFAS admits a kernel of polynomial size. We consider \mathcal{C} -ARC DELETION SET (\mathcal{C} -ADS), a variant of DFAS where we want to remove at most k arcs from the input digraph in order to turn it into a digraph of a class \mathcal{C} . In this work, we choose \mathcal{C} to be the class of *funnels*. FUNNEL-ADS is NP-hard even if the input is a DAG, but is fixed-parameter tractable with respect to k . So far no polynomial kernels for this problem were known. Our main result is a kernel for FUNNEL-ADS with $\mathcal{O}(k^6)$ many vertices and $\mathcal{O}(k^7)$ many arcs, computable in linear time.

1 Introduction

In graph editing problems, we are given a (directed or undirected) graph G and a number k , and we search for a set of at most k vertices, edges or arcs whose removal or addition produces a graph with a desired property. There are several variants of these problems, and in this paper we consider the problem of removing arcs from a digraph in order to obtain a digraph in a given class \mathcal{C} . When \mathcal{C} is the class of all directed acyclic graphs (DAGs), the problem is called DIRECTED FEEDBACK ARC SET (DFAS). If we remove vertices instead of arcs, the problem is called DIRECTED FEEDBACK VERTEX SET (DFVS).

There are simple reductions between DFAS and DFVS. We can reduce DFAS to DFVS by taking the line digraph of the input. Removing a vertex from the reduced instance corresponds to removing an arc from the input instance and vice versa. For a reduction in the other direction, we split each vertex v into two vertices, say, v_o and v_i , connect them with an arc (v_i, v_o) and shift all outgoing arcs of v to v_o and all incoming arcs to v_i . In the context of *parameterized complexity*, such reductions are called *parameterized* as the parameter k is preserved. Hence, parameterized results are often stated for DFVS.

In a breakthrough paper it was proven that there is an algorithm for DFVS with running time $f(k) \cdot n^{\mathcal{O}(1)}$ for some function f [4], showing that the problem is *fixed-parameter tractable* (FPT) with respect to k . After obtaining an FPT result, it is natural to ask if the problem also admits a polynomial *kernel*, that is, if there is a polynomial-time algorithm which reduces the input instance to an

instance of size at most $\mathcal{O}(k^c)$ for some constant c . Such an algorithm is called a *kernelization* algorithm.

The existence of a polynomial kernel for DFVS is a fundamental open question in the field of parameterized complexity. One approach towards solving this question is to consider different parametrizations or restrictions of the input digraph. By considering smaller parameters or more general digraph classes, one can hope to eventually close the gap between the restricted cases and the general case of DFVS.

On tournaments, DFVS admits a polynomial kernel [1]; this was extended to generalizations of tournaments as well [3]. When parameterized by solution size k and the size ℓ of a treewidth η -modulator, DFVS admits a kernel of size $(k \cdot \ell)^{\mathcal{O}(\eta^2)}$ [10].


One can also restrict the output instead, that is, we can consider \mathcal{C} -VERTEX DELETION SET (\mathcal{C} -VDS) or \mathcal{C} -ARC DELETION SET (\mathcal{C} -ADS), where, for a fixed digraph class \mathcal{C} , we search for a set of at most k vertices (arcs) whose removal turns the input into a digraph in \mathcal{C} . Unlike DFVS and DFAS, \mathcal{C} -VDS and \mathcal{C} -ADS can belong to different complexity classes depending on \mathcal{C} : While OUT-FOREST-ADS can be solved in polynomial time, OUT-FOREST-VDS is NP-hard [12]. Further, note that even if $\mathcal{C}' \subseteq \mathcal{C}$, a polynomial kernel for \mathcal{C} -ADS does not immediately imply a polynomial kernel for \mathcal{C}' -ADS, and the implication also does not work in the other direction. Indeed, while the problem is trivial when \mathcal{C} is the class of all independent sets or the class of all digraphs, it is NP-hard if \mathcal{C} is the class of DAGs, which contains all independent sets and is a subclass of all digraphs. In a sense, the complexity landscapes of \mathcal{C} -ADS and \mathcal{C} -VDS are much more fine-grained than the landscape of DFVS, and may allow for smaller steps towards more general results.

OUT-FOREST-ADS and PUMPKIN-ADS can be solved in polynomial time [12], while OUT-FOREST-VDS and PUMPKIN-VDS are NP-hard and admit polynomial kernels [2; 12] of size $\mathcal{O}(k^2)$ and $\mathcal{O}(k^3)$, respectively [2]. \mathcal{F}_η -VDS admits a polynomial kernel for constant η , where \mathcal{F}_η is the class of all digraphs with (undirected) treewidth at most η [10].

In this work we consider FUNNEL-ADS and provide a polynomial kernel with $\mathcal{O}(k^6)$ vertices and $\mathcal{O}(k^7)$ arcs. A digraph is a funnel if it is a DAG and every source to sink path has an arc which is not in any other source to sink path. FUNNEL-ADS is NP-hard even if the input is DAG, but it is FPT with respect to solution size [11]. Out-forests and pumpkins are also funnels, but there are also dense funnels like complete bipartite digraphs (where all arcs go from the first partition to the second but not back).

Our results rely on characterizations for funnels based on forbidden subgraphs and on a “labeling” of the vertices [11]. We believe the techniques used here can be generalized to other digraph classes which are also similarly characterized, and hope they provide further insight about the classes \mathcal{C} for which \mathcal{C} -ADS admits a polynomial kernel.

Figure 1: The forbidden subgraph for funnels D_1 .



2 Preliminaries

A function $f : A \rightarrow B$ is a set of tuples $(a, f(a)) \in A \times B$ where for every $a \in A$ there is at most one $b \in B$ with $(a, b) \in f$ (that is, $f(a) = b$). We write $\text{Dom}(f)$ for the set of values $a \in A$ for which f is defined. Hence, \emptyset is the undefined function, and $f' \supseteq f$ if $f'(x) = f(x)$ for every $x \in \text{Dom}(f)$. All our functions are *partial*, that is, $\text{Dom}(f)$ is not necessarily A .

A *parameterized language* L is *fixed-parameter tractable* with respect to the parameter k if there is some algorithm with running time $f(k) \cdot n^{\mathcal{O}(1)}$ deciding whether $(x, k) \in L$, where f is some computable function, $n = |x|$ and k is the parameter (refer to [5; 6] for an introduction to parameterized complexity). We say that L *admits a problem kernel* if there is a polynomial-time algorithm which transforms an instance (x, k) into an instance (x', k') such that $(x, k) \in L$ if and only if $(x', k') \in L$, $k' \leq k$ and $|x'| \leq f(k)$ for some computable function f . If f is a polynomial, we say that L *admits a polynomial kernel* with respect to k .

When describing a kernelization algorithm, it is common to define *reduction rules*. These rules have a *condition* and an *effect*, and we say that a reduction rule is not *applicable* if the condition is not true. The effect of the reduction rule produces a new instance (x', k') of the problem, and a rule is said to be *safe* if $(x', k') \in L$ if and only if the original instance is in L . We refer the reader to [8; 9] for surveys on kernelization and to [7] for a book on the topic.

We only consider directed graphs (digraphs) without loops or parallel arcs in this paper. Let D be a digraph. The set of arcs of D is given by $A(D)$, and its set of vertices is $V(D)$. The set of outneighbors (inneighbors) in D of a vertex $v \in V(D)$ is given by $\text{out}_D(v)$ ($\text{in}_D(v)$); the outdegree (indegree) of v is $\text{outdeg}_D(v) = |\text{out}_D(v)|$ ($\text{indeg}_D(v) = |\text{in}_D(v)|$). If the digraph D is clear from context, we omit it from the index. For a set $U \subseteq V(D)$ we write $\text{out}(U)$ for the set $\{\text{out}(u) \mid u \in U\}$ (and analogously for $\text{in}(U)$). A vertex v is a *source* if $\text{indeg}(v) = 0$, and it is a *sink* if $\text{outdeg}(v) = 0$. We write $H \subseteq D$ if H a *subgraph* of D ; the subgraph of D *induced* by U is given by $D[U]$. We write $D - X$ for the operation of deleting a set of vertices or arcs X from D . Similarly, we add a set of arcs or vertices to D with $D + X$.

A *directed acyclic graph* (DAG) is a digraph which does not contain any directed cycle. A digraph D is a *funnel* if D is a DAG and for every path P from a source to a sink of D of length at least one there is some arc $a \in A(P)$ such that for any different path Q from a (possibly different) source to a sink we have $a \notin A(P) \cap A(Q)$. We repeat below several known characterizations for funnels, as they are particularly useful for our results.

Theorem 2.1 ([11]). *Let D be a DAG. The following statements are equivalent.*

a. D is a *funnel*.

- b. $V(D)$ can be partitioned into two sets¹ F and M such that: (1) F induces an out-forest; (2) M induces an in-forest; and (3) $(M \times F) \cap A(D) = \emptyset$.
- c. No digraph in $\mathcal{F} = \{D_i\}_{i=0}^{\infty}$ is contained in D as a (not necessarily induced) subgraph, where (see Figure 1)
- $V(D_k) = \{u_1, u_2, v_0, w_1, w_2\} \cup \{v_i\}_{i=0}^k$, and
 - $A(D_k) = \{(u_1, v_0), (u_2, v_0), (v_k, w_1), (v_k, w_2)\} \cup \{(v_i, v_{i+1})\}_{i=1}^{k-1}$.
- d. D does not contain D_0 as a butterfly minor.

The digraphs in \mathcal{F} are called *forbidden subgraphs for funnels*. For a digraph D we define a *labeling* as a function $\ell : V(D) \rightarrow \{F, M\}$. We say that ℓ is a *funnel labeling* for D if $\text{Dom}(\ell) = V(D)$, the set $F = \{v \in V(D) \mid \ell(v) = F\}$ induces an out-forest in D , the set $M = \{v \in V(D) \mid \ell(v) = M\}$ induces an in-forest in D and $(M \times F) \cap A(D) = \emptyset$. Due to [Theorem 2.1\(b\)](#), a digraph D is a funnel if and only if there exists a funnel labeling for D .

In the *feedback arc set* problem, we are given a digraph D and a $k \in \mathbb{N}$ as an input, and we search for a set $S \subseteq A(D)$ such that $D - S$ is a DAG and $|S| \leq k$. We consider a variant of this problem where we want $D - S$ to be a funnel instead, which is formally defined below.

FUNNEL ARC DELETION SET (FADS)

Input A digraph D and a number $k \in \mathbb{N}$.

Question Is there a set $S \subseteq A(D)$ with $|S| \leq k$ such that $D - S$ is a funnel?

To better make use of [Theorem 2.1\(b\)](#), we consider a more general problem in which some vertices might already be labeled with F or M, and the funnel we obtain in the end must respect this labeling. Formally, the problem is defined as follows.

FUNNEL ARC DELETION LABELING (FADL)

Input A digraph D , a number $k \in \mathbb{N}$ and a labeling $\ell : V(D) \rightarrow \{F, M\}$.

Question Are there a set $S \subseteq A(D)$ and a labeling $\hat{\ell} \supseteq \ell$ such that $\hat{\ell}$ is a funnel labeling for $D - S$ and $|S| \leq k$?

We say that (D, ℓ, k) is the *input instance* and $(S, \hat{\ell})$ is a solution for the input instance. This more general version of the problem allows us to decide which label a vertex will take and encode this in the instance itself. While technically not necessary, using FADL instead of FADS simplifies the kernelization algorithm and also the proofs. Due to space constraints, proofs marked with (\star) are deferred to the appendix.

3 Basic reduction rules

We construct our kernelization algorithm by defining a series of reduction rules and then showing that, if no reduction rule is applicable, the input size is bounded in a polynomial of k . Our strategy is to partition the vertex set into labeled and unlabeled vertices, then bound the number of unlabeled vertices ([Section 3.1](#))

¹ F and M stand for *fork* and *merge*, as vertices in F “fork” paths and those in M “merge” paths.

and use this to bound the number of labeled vertices (Section 3.2) as well. In this section we define some reduction rules which are useful both in Section 3.1 as well as in Section 3.2. For brevity, we assume that a reduction rule is no longer applicable to the input instance after it has been defined.

Let (D, ℓ, k) be the input instance. From Theorem 2.1 we can see that a funnel has no vertex v with $\text{indeg}(v) > 1$ and $\text{outdeg}(v) > 1$. Further, $\text{indeg}(v) \leq 1$ if $\ell(v) = \text{F}$, and $\text{outdeg}(v) \leq 1$ if $\ell(v) = \text{M}$. Hence, by simply counting the number of vertices disrespecting each case, we can obtain a lower bound for the number of arcs that need to be removed from D in order to obtain a funnel. As removing one arc changes the degree of two vertices, we obtain a bound of at most $2k$ such vertices. The safety of the following reduction rule follows easily from Theorem 2.1.

Reduction Rule 3.1 (Lower Bound). *Let $V_I \subseteq V(D)$ be the set of vertices with indegree greater than one, let V_O be the set of vertices with outdegree greater than one and let $V_X = V_O \cap V_I$. Output a trivial “no” instance if*

$$\sum_{u \in V_O, \ell(u) = \text{M}} (\text{outdeg}(u) - 1) + \sum_{u \in V_I, \ell(u) = \text{F}} (\text{indeg}(u) - 1) + \sum_{u \in V_X, u \notin \text{Dom}(\ell)} (\min\{\text{indeg}(u), \text{outdeg}(u)\} - 1) > 2k.$$

The following reduction rule is based on [11], with some modifications since the original reduction rule is applied as an intermediate step in an FPT algorithm and is not safe for kernelization. For certain vertices it is possible to optimally decide which label they should receive in an optimal solution. For example, vertices with outdegree greater than $k + 1$ can always be labeled with F, as otherwise we would need to remove at least $k + 1$ of its outgoing arcs, which is not possible.

Reduction Rule 3.2 (Set Label). *Let $v \in V(D)$ be an unlabeled vertex.*

Set $\ell(v) := \text{F}$ if at least one of the following is true: (1) $\text{indeg}(v) = 0$; (2) v has a single inneighbor u and $\ell(u) = \text{F}$; (3) there are at least $\text{indeg}(v) + 1$ vertices $u \in \text{out}(v)$ with $\ell(u) = \text{M}$ or $\ell(u) = \text{F} \wedge \text{indeg}(u) = 1$; or (4) $\text{outdeg}(v) > k + 1$.


Set $\ell(v) := \text{M}$ if at least one of the following is true: (1) $\text{outdeg}(v) = 0$; (2) v has a single outneighbor u and $\ell(u) = \text{M}$; (3) there are at least $\text{outdeg}(v) + 1$ vertices $u \in \text{in}(v)$ with $\ell(u) = \text{F}$ or $\ell(u) = \text{M} \wedge \text{outdeg}(u) = 1$; or (4) $\text{indeg}(v) > k + 1$.

Proof. We consider the case where we set $\ell(v) := \text{F}$, as the other case is symmetric. Let ℓ_r be the labeling obtained by the reduction rule. Let $(S, \hat{\ell})$ be a solution for the original instance. We set $\hat{\ell}_r := \hat{\ell}$ and $\hat{\ell}_r(v) := \text{F}$. If $\hat{\ell}(v) = \text{F}$, then clearly $(S, \hat{\ell}_r)$ is a solution for the reduced instance. So assume $\hat{\ell}(v) = \text{M}$.

If $\text{indeg}(v) = 0$, or $\text{indeg}(v) = 1$ and there is some $u \in \text{in}(v)$ with $\ell(u) = \text{F}$, then $\hat{\ell}_r$ is clearly also a funnel labeling for $D - S$.

If there are at least $\text{indeg}(v) + 1$ vertices $u \in \text{out}(v)$ with $\ell(u) = \text{M}$ or $\ell(u) = \text{F} \wedge \text{indeg}(u) = 1$, then we construct an S_r from S as follows. We add all incoming arcs of v to S_r and remove from S_r all outgoing arcs (v, u) where $\ell(u) = \text{M}$ or

Figure 2: A digraph which is not a funnel. Removing the arcs (v, u) and (u, w) results in a funnel.



$\ell(u) = F \wedge \text{indeg}(u) = 1$. Since $\hat{\ell}(v) = M$, at least $\text{outdeg}(v) \geq \text{indeg}(v)$ many outgoing arcs of v are in S . Hence, we remove at least $\text{indeg}(v)$ arcs from S and add at most $\text{indeg}(v)$. Thus, $|S_r| \leq |S|$.

The digraph $D - S_r$ does not contain cycles, as all incoming arcs of v were removed, so any cycle in $D - S_r$ is also in $D - S$, which is a funnel. To see that ℓ_r is a funnel labeling of $D - S_r$, first note that we can always keep arcs (v, u) in $D - S_r$ where $\ell(u) = M$. We can also keep arcs (v, u) in $D - S_r$ where $\ell(u) = F$ and $\text{indeg}(u) = 1$. As v has no incoming arcs in $D - S_r$, it lies in an out-forest. Hence, ℓ_r is a funnel labeling of $D - S_r$. \square

Replacing an arc in a funnel by a directed path cannot create any cycles nor any forbidden subgraph for funnels. The next reduction rule reverses this operation: We can contract certain paths where all vertices have in- and outdegree one by a single arc. However, we cannot replace any such path: In the example in Figure 2, if we remove v and add the arc (u, w) , then the size of an optimal solution set decreases by one. Some cases where contracting an arc is safe are identified below.

Reduction Rule 3.3 (Dissolve Vertex). *Let u, v, w be a path such that the following is true: (1) $v, u \in \text{Dom}(\ell)$ implies $\ell(v) = \ell(u)$; and (2) $v, w \in \text{Dom}(\ell)$ implies $\ell(v) = \ell(w)$.*

If $\text{indeg}(v) = \text{outdeg}(v) = 1$ and $\text{indeg}(w) = 1 \vee \text{outdeg}(u) = 1$, delete the vertex v and add the arc (u, w) .

Proof. Let D' be the reduced digraph. It is easy to see that we can obtain a solution for the reduced instance from a solution for the original instance: If we remove (u, v) or (v, w) from D , we can instead remove (u, w) from D' . As this is equivalent to removing v from D , the result is also a funnel and we can keep the same labeling (up to v , which is not in D'). If neither (u, v) nor (v, w) were removed, we simply keep the same arc-deletion set and labeling.

Now let $(S_r, \hat{\ell}_r)$ be a solution for the reduced instance. We start by setting $\hat{\ell} := \hat{\ell}_r$. If $(u, w) \notin S_r$, we set $\hat{\ell}(v) := \hat{\ell}_r(u)$. It is easy to see that $\hat{\ell}$ is a funnel labeling for $D - S_r$.

If $(u, w) \in S_r$, we distinguish two cases. If $\text{outdeg}_D(u) = 1$, we set $\hat{\ell}(v) := \hat{\ell}(u)$ and $S := (S_r \setminus \{(u, w)\}) \cup \{(v, w)\}$. Regardless of whether $\hat{\ell}(u) = M$ or $\hat{\ell}(u) = F$, we do not need to remove (u, v) . Since the neighborhood of w did not change and any cycle in $D - S$ is also a cycle in $D' - S_r$, it follows that $\hat{\ell}$ is a funnel labeling for $D - S$.

If $\text{outdeg}_D(u) > 1$ and $\text{indeg}_D(w) = 1$, we set $\hat{\ell}(v) := \hat{\ell}(w)$ and $S = (S_r \setminus \{(u, w)\}) \cup \{(u, v)\}$. As before, we may keep the arc (w, u) in $D - S$, and $\hat{\ell}$ is

a funnel labeling for $D - S$. Since the case $\text{outdeg}_D(u) > 1$ and $\text{indeg}_D(w) > 1$ does not occur, this concludes the proof. \square

3.1 Bounding the number of unlabeled vertices

From [Lower Bound \(Theorem 3.1\)](#) we know there are few vertices with both in- and outdegree greater than one. In this section we bound the number of unlabeled vertices by considering the remaining unlabeled vertices, that is, vertices v with $\text{indeg}(v) \leq 1$ or $\text{outdeg}(v) \leq 1$. Our strategy is to group such vertices into subgraphs of D with specific properties which we define later, and then develop reduction rules to both bound the maximum number of such subgraphs and also their size in any “yes” instance of FADL.

Even if the previous reduction rules are not applicable, there can still exist some “large” subgraph $H \subseteq D$ for which there is a “small” set $S \subseteq A(D)$ such that the weakly-connected component of H is a funnel in $D - S$. Our goal is to bound the size of such subgraphs H .

We first define a specific type of subgraph of D which behaves like a funnel in the sense that the degrees of the vertices match [Theorem 2.1\(b\)](#). We call such subgraphs *local funnels* and formally define them below.

Definition 3.4. An induced subgraph $H \subseteq D$ is a *local funnel* in D if H is a funnel, H has only one source and its vertex set can be partitioned into $F \uplus M = V(H)$ such that $\text{indeg}_D(v) \leq 1$ for all $v \in F$; $\text{outdeg}_D(v) \leq 1$ for all $v \in M$; and $(M \times F) \cap A(H) = \emptyset$.


Unlike *local funnels*, we might still have to remove many arcs from an *induced* funnel in D , as it can have, for example, several vertices v with $\text{indeg}_D(v) > 1$ and $\text{outdeg}_D(v) > 1$. Our goal is to bound the size of each unlabeled local funnel and the number of unlabeled local funnels in D . We start by “pushing” as many vertices as we can to the neighborhood of the roots of the in- and outforests of a local funnel. Consider for example a path as in [Figure 3](#). Intuitively, a cycle containing v and x must also contain u . To destroy this cycle, we can remove the unique incoming arc of u , as this will potentially destroy further cycles that contain u but not v . Hence, replacing the arc (v, x) with (u, x) in this case does not change the solution.

By moving vertices in an outtree towards its root s , we increase the outdegree of s . If the outdegree of s increases beyond $k + 1$, we can apply [Set Label \(Theorem 3.2\)](#) to s , giving it a label. By further applying [Set Label \(Theorem 3.2\)](#) to the neighbors of s which are in its outtree, we can label the entire tree. As we are only considering unlabeled local funnels in this section, we can use the idea above to limit the branching of any in- or outtree of a unlabeled local funnel.

We provide here a somewhat more general reduction rule which can also be applied if some vertices are labeled. Later, this reduction rule will again be useful to bound the number of labeled vertices. However, we need to carefully consider the possible labels of the vertices, as in some cases the rule would not be safe.

Reduction Rule 3.5 (Shift Neighbors). *Let u, v, w be a path.*

Figure 3: Example application of **Shift Neighbors** (Theorem 3.5).



- If $\text{indeg}(u) = \text{indeg}(v) = \text{indeg}(w) = 1$, $(u, M) \notin \ell$, $(v, M) \notin \ell$ and there is an $x \in \text{out}(v) \setminus \text{out}(u)$ with $w \neq x$, then remove the arc (v, x) and add the arc (u, x) .
- If $\text{outdeg}(u) = \text{outdeg}(v) = \text{outdeg}(w) = 1$, $(v, F) \notin \ell$, $(w, F) \notin \ell$ and there is an $x \in \text{in}(v) \setminus \text{in}(w)$ with $u \neq x$, then remove the arc (x, v) and add the arc (x, w) .

Before proving that **Shift Neighbors** (Theorem 3.5) is safe, we need two simple observations about certain cases where we can safely exchange two arcs or add an arc.

Observation 1. Let H be a funnel with funnel labeling ℓ and let $x, u, v \in V(H)$ such that $(v, x) \in A(H)$, $(u, x) \notin A(H)$ and at least one of the following is true: (1) $\ell(u) = F$; or (2) $\ell(u) = M = \ell(v)$ and $\text{outdeg}_H(u) = 0$. Let $H' = H - (v, x) + (u, x)$. Then ℓ is also a funnel labeling for H' if H' is a DAG.

Proof. Assume H' is a DAG.

Case 1: $\ell(u) = F$. If $\ell(x) = M$, then $H + (u, x)$ is still a funnel and so is $H - (v, x) + (u, x)$. If $\ell(x) = F$, then H' is the result of switching the unique inneighbour of x in the out-forest induced by vertices labeled with F . This is clearly an out-forest, and hence ℓ is a funnel labeling for H' .

Case 2: $\ell(u) = M = \ell(v)$ and $\text{outdeg}_H(u) = 0$. As $(v, x) \in A(H)$, we have $\ell(x) = M$. Since $\text{outdeg}_H(u) = 0$, ℓ is a funnel labeling for $H + (u, x)$ and, hence, also for H' . \square

Observation 2. Let H be a DAG and $x, u, v \in V(H)$ such that $\{u\} = \text{in}(v)$. Then $H + (u, x)$ contains a cycle if and only if $H + (v, x)$ contains a cycle.

Proof. Assume $H + (v, x)$ contains a cycle C . As $\{u\} = \text{in}(v)$, it follows that $(u, v) \in A(C)$. Hence, replacing (u, v) and (v, x) with (u, x) in C produces a cycle in $H + (u, x)$.

If $H + (u, x)$ contains a cycle C , then we can replace (u, x) by the path from u to x in $H + (v, x)$ going through v . This constructs a cycle in $H + (v, x)$, as desired. \square

Proof of safety of Shift Neighbors (Theorem 3.5). Consider the case where $1 = \text{indeg}(u) = \text{indeg}(v) = \text{indeg}(w)$, $(u, M) \notin \ell$, $(v, M) \notin \ell$ and there is an $x \in \text{out}(v) \setminus \text{out}(u)$ with $w \neq x$. The other case follows analogously. Let (D', ℓ, k) be the reduced instance and $(S_r, \hat{\ell}_r)$ be a solution for it. We construct a solution $(S, \hat{\ell})$ for the input instance (D, ℓ, k) .

First observe that, if $(u, x) \in S_r$, we can replace it with (v, x) in S , which means that $D' - S_r$ and $D - S$ are isomorphic. By setting $\hat{\ell} = \hat{\ell}_r$, we obtain the desired solution. If $(u, x) \notin S_r$, we consider the following cases.

Case 1: $\hat{\ell}_r(v) = \text{F}$. We set $\hat{\ell} := \hat{\ell}_r$ and $S := S_r$. Let $D^* = D - S$. Clearly, $D^* = D' - S_r - (u, x) + (v, x)$. As u is the only inneighbor of v , from [Observation 2](#) we know D^* is a DAG. From [Observation 1](#), it follows that $\hat{\ell} = \hat{\ell}_r$ is a funnel labeling for D^* .

Case 2: $\hat{\ell}_r(v) = \text{M} = \hat{\ell}_r(u)$. If $D' - S_r + (u, v)$ is a DAG, then we can assume that $(u, v) \notin S_r$, implying $(u, x) \in S_r$ (which was already considered).

If $D' - S_r + (u, v)$ is not a DAG, then it contains a cycle with v and u , implying $(u, v) \in S_r$. In particular, $\text{indeg}_{D' - S_r}(v) = 0$. We set $\hat{\ell} := \hat{\ell}_r$ and $\hat{\ell}(v) := \text{F}$. Clearly, $\hat{\ell}$ is a funnel labeling for $D' - S_r$. From [Observation 1](#) we have that $\hat{\ell}$ is a funnel labeling for $D - S_r$ as well.

Case 3: $\hat{\ell}_r(v) = \text{M}$ and $\hat{\ell}_r(u) = \text{F}$. We set $\hat{\ell} := \hat{\ell}_r$, $\hat{\ell}(v) := \text{F}$ and $S := S_r$. As $\{u\} = \text{in}_D(v)$ and $\hat{\ell}_r(u) = \text{F}$, $\hat{\ell}$ is a funnel labeling for $D' - S_r$. Let $D^* = D - S$.

From [Observation 2](#) we know $D^* = D' - S_r - (u, x) + (v, x)$ is a DAG since $D' - S_r$ is a DAG. Hence, from [Observation 1](#) it follows that $(S_r, \hat{\ell})$ is a solution for the input instance. In all cases a solution for the reduced instance implies a solution for the original instance.

Now assume there is a solution $(S, \hat{\ell})$ for the original instance. We show that there is solution $(S_r, \hat{\ell}_r)$ for the reduced instance. As in the previous direction, if $(v, x) \in S$, we can replace it with (u, x) and obtain the desired solution. So assume $(v, x) \notin S$.

If $(u, v) \in S$, let $S_1 = S \cup \{(y, u)\}$, where $\{y\} = \text{in}(u)$. Clearly, $\hat{\ell}$ is a funnel labeling for $D - S_1$. We set $\hat{\ell}_r := \hat{\ell}$ and $\hat{\ell}_r(u) := \text{F}$. As $\text{indeg}_{D - S_1}(u) = 0$, $\hat{\ell}_r$ is also a funnel labeling for $D - S_1$. From [Observation 1](#) we know that $\hat{\ell}_r$ is a funnel labeling for $D_1 = D' - S_1$. Since $\text{indeg}_{D_1}(v) = 0 = \text{indeg}_{D_1}(u)$ and $\hat{\ell}_r(u) = \text{F}$, we have that $\hat{\ell}_r$ is a funnel labeling for $D_1 + (u, v)$. Hence, $(S \setminus \{(u, v)\}, \hat{\ell}_r)$ is a solution for the reduced instance.

In the following we consider the remaining cases where $\{(u, v), (v, x)\} \cap S = \emptyset$. Note that the case $\hat{\ell}(u) = \text{M}$ and $\hat{\ell}(v) = \text{F}$ does not happen under this assumption.

Case 1: $\hat{\ell}(v) = \text{F} = \hat{\ell}(u)$. Let $\hat{\ell}_r = \hat{\ell}$ and $S_r = S$. Clearly, $D' - S_r = D - S - (v, x) + (u, x)$. From [Observation 2](#) there is no cycle in $D - S + (u, x)$ and, hence, $D' - S_r$ is a DAG. Thus, from [Observation 1](#) it follows that $\hat{\ell}_r$ is a funnel labeling for $D' - S_r$.

Case 2: $\hat{\ell}(v) = \text{M} = \hat{\ell}(u)$. Since $(v, x) \notin S$, we have $(v, w) \in S$ and $\hat{\ell}(x) = \text{M}$. Further, we know that $D' - S$ is a DAG due to [Observation 2](#). Let $S_1 = S \cup \{(u, v)\}$. Clearly, $\hat{\ell}$ is a funnel labeling for $D - S_1$, and $D' - S_1$ is also a DAG. From [Observation 1](#) follows that $\hat{\ell}$ is a funnel labeling for $D' - S_1$.

We set $\hat{\ell}_r := \hat{\ell}$ and $\hat{\ell}_r(v) = \text{F}$. Since $\text{indeg}_{D' - S_1}(w) = 0 = \text{indeg}_{D' - S_1}(v)$, we have that $\hat{\ell}_r$ is a funnel labeling for $D' - S_1 + (v, w)$, regardless of the label of w . By setting $S_r = (S \setminus \{(v, w)\}) \cup \{(u, v)\}$, we get that $\hat{\ell}_r$ is a funnel labeling for $D' - S_r$ and $|S_r| \leq |S|$.

Case 3: $\hat{\ell}(v) = \text{M}$ and $\hat{\ell}(u) = \text{F}$. Let $S_r = S$ and $\hat{\ell}_r = \hat{\ell}$. Since $(u, v) \notin S_r$, from [Observation 2](#) we know that $D - S_r - (v, x) + (u, x)$ is a DAG. From

Observation 1 we have that $\hat{\ell}_r$ is a funnel labeling for $D' - S_r$.

In all cases we found a solution $(S_r, \hat{\ell}_r)$ for the reduced instance, concluding the proof. \square

If **Shift Neighbors** (Theorem 3.5) is not applicable, then many vertices in a long path P in a local funnel must share a common out- or inneighbor w . However, from **Set Label** (Theorem 3.2) we know that w receives a label if it has too many neighbors. The next and final reduction rule needed for bounding the number of unlabeled vertices exploits this property and allows us to label some vertex u in P if its predecessor v in P is adjacent to a labeled vertex w .

Reduction Rule 3.6 (Labeled Neighbor). *Let v, u be a path of unlabeled vertices. Set $\ell(u) := F$ if $\text{indeg}(u) = \text{indeg}(v) = 1$ and $\exists w \in \text{out}(v) : \ell(w) = M$. Set $\ell(v) := M$ if $\text{outdeg}(u) = \text{outdeg}(v) = 1$ and $\exists w \in \text{in}(u) : \ell(w) = F$.*

Proof. Assume, without loss of generality, that the first case of the rule was applied. The proof for the second case follows analogously (note that it is not possible for both cases to be applied simultaneously). Let (D, ℓ_r, k) be the reduced instance. First note that $\ell_r \supseteq \ell$, which means that a solution for the reduced instance is already a solution for the original instance. Hence, it suffices to show that a solution $(S, \hat{\ell})$ for the original instance implies a solution $(S_r, \hat{\ell}_r)$ for the reduced instance.

If $\hat{\ell}(u) = F$, we can set $\hat{\ell}_r = \hat{\ell}$ and $S_r = S$ and we are done. So assume that $\hat{\ell}(u) = M$.

Case 1: $(v, u) \in S$. We set $S_r := S$, $\hat{\ell}_r := \hat{\ell}$ and $\hat{\ell}_r(u) = F$. As $\text{indeg}_{D-\hat{\ell}}(u) = 0$, it follows that $\hat{\ell}_r$ is also a funnel labeling for $D - S$.

Case 2: $(v, u) \notin S$ and $\hat{\ell}(v) = F$. We set $S_r := S$, $\hat{\ell}_r := \hat{\ell}$ and $\hat{\ell}_r(u) = F$. As $\hat{\ell}_r(v) = F = \hat{\ell}_r(u)$, we may keep the arc (v, u) and $\hat{\ell}_r$ is a funnel labeling for $D - S_r$.

Case 3: $(v, u) \notin S$ and $\hat{\ell}(v) = M$. Then $(v, w) \in S$. We set $\hat{\ell}_r := \hat{\ell}$, $\hat{\ell}_r(u) := F$, $\hat{\ell}_r(v) = F$, $S_r := (S \setminus \{(v, w)\}) \cup \{(y, v)\}$, where y is the unique inneighbor of v .

The digraph $D - S_r$ is a DAG: if it has a cycle, the cycle would have to use the arc (v, w) , yet $\text{indeg}_{D-S_r}(v) = 0$, a contradiction. We now argue that $\hat{\ell}_r$ is a funnel labeling for $D - S_r$. Since $\text{indeg}_{D-S_r}(v) = 0$, $\text{indeg}_{D-S_r}(u) = 1$ and $\hat{\ell}_r(u) = F$, the vertex v is the unique inneighbor of u in the out-forest of the funnel $D - S_r$. Finally, as $\hat{\ell}_r(w) = M$, the arc (v, w) is allowed in the funnel. Hence, $\hat{\ell}_r$ is a funnel labeling for $D - S_r$. In all cases we find a solution $(\hat{\ell}_r, S_r)$ for the reduced instance, concluding the proof. \square

Lemma 1. *Let s be some source (sink) of some unlabeled local funnel H . Let P_1, P_2, \dots, P_a be a sequence of paths in H starting (ending) at s such that $\text{indeg}(u) \leq 1$ ($\text{outdeg}(u) \leq 1$) for each u in each P_i , and $V(P_j) \not\subseteq V(P_i)$ for all $1 \leq i, j \leq a$ where $i \neq j$. Let E be the set of endpoints of all P_i . Then all of the following hold.*

1. $\text{outdeg}(u) > 1$ ($\text{indeg}(u) > 1$) for any inner vertex u of any P_i .
2. $\text{out}(\bigcup_{i=1}^a V(P_i) \setminus E) \subseteq \text{out}(s)$ ($\text{in}(\bigcup_{i=1}^a V(P_i) \setminus E) \subseteq \text{in}(s)$),

3. $V(P_i) \cap V(P_j) = \{s\}$ for each $1 \leq i, j \leq a$ where $i \neq j$, and

4. $a \leq k + 1$ and $|V(P_i)| \leq k + 2$ for each $1 \leq i \leq a$.

Proof. We consider the case where s is a source of H . The other case follows analogously.

Let u be some inner vertex of some P_i and w the unique outneighbor of u in P_i . By assumption on P_i , we have $\text{indeg}_D(w) = 1$. As [Dissolve Vertex \(Theorem 3.3\)](#) is not applicable, it follows that $\text{outdeg}(u) > 1$ (proving 1). In particular, u has some outneighbor x not in P_i .

Let v be the inneighbor of u in P_i . Since $\text{indeg}_D(v) = \text{indeg}_D(u) = \text{indeg}_D(w) = 1$ and [Shift Neighbors \(Theorem 3.5\)](#) is not applicable, it follows that $x \in \text{out}_D(v)$. By repeating this argument to the predecessors of u in P_i , we prove 2 (and also that $a \leq k + 1$, as $\text{outdeg}_D(s) \leq k + 1$).

Assume there are two paths P_i and P_j intersecting at more than one vertex. Let u be the last vertex of the intersection. Then u has two outneighbors w_i and w_j lying on P_i and P_j , respectively, and $w_i \neq w_j$. But due to 2, we have $w_i, w_j \in \text{out}_D(s)$, implying $\text{indeg}_D(w_i) > 1$ and $\text{indeg}_D(w_j) > 1$, a contradiction to our assumptions on P_i and P_j (proving 3).

Let v_1, v_2, \dots, v_m be the sequence of vertices of a path P_i . From 1 we know that there is some $w \in \text{out}_D(v_{m-1})$ outside of P_i . We also have $w \in \text{out}_D(v_j)$ for all $1 \leq j \leq m - 1$, implying $\text{indeg}_D(w) \geq m - 1$. If $m - 1 > k + 1$, then $\ell(w) = M$, as [Set Label \(Theorem 3.2\)](#) is not applicable. However, as $\text{indeg}_D(v_{m-1}) = 1 = \text{indeg}_D(v_{m-2})$, $w \in \text{out}_D(v_{m-2})$ and [Labeled Neighbor \(Theorem 3.6\)](#) is not applicable, we have $\ell(v_{m-1}) = F$, a contradiction to the assumption that H is unlabeled. Hence, $m - 1 \leq k + 1$, implying $|V(P_i)| \leq k + 2$ (proving 4). \square

Lemma 2. *Let H be a unlabeled local funnel in D . Then $|V(H)| \in \mathcal{O}(k^3)$.*

Proof. Let s be the source of H . Consider a partitioning of the vertices of H into an out-tree (since H has only one source) and an in-forest where the out-tree is maximal. Let P_1, P_2, \dots, P_a be a sequence of paths such that the out-tree is the union of all P_i . From [Lemma 1](#) we know that $a \leq k$ and that $V(P_i) \cap V(P_j) = \{s\}$ for all $i \neq j$. Let v_i be the endpoint of P_i which is not s and let $X = \bigcup_{i=1}^a \text{out}(v_i)$. As [Set Label \(Theorem 3.2\)](#) is not applicable, we have $|X| \leq a \cdot k \leq k^2$. Further, as $|V(P_i)| \leq k + 1$, the out-tree of H has at most $k(k + 1)$ many vertices.

Let Y be the set of sinks of H lying on its in-forest. Since H has only one source s , for every sink $t \in Y$ there is a path Q from s to t . Let Q_1, Q_2, \dots, Q_b be the set of all paths from s to each sink in Y , and let R_i be the subpath of Q_i contained in the in-forest of H .

Let Q_i be one of such paths, and let u be the first vertex of R_i (which is not in any P_j). Note that this implies $\text{indeg}(u) > 1$, otherwise the out-tree would not be maximal. Due to [Lemma 1](#) we have that no other R_ℓ contains u and if $u \notin x$, then $u \in \text{out}(s)$. Since [Set Label \(Theorem 3.2\)](#) is not applicable and each distinct R_ℓ implies the existence of a distinct outneighbor of s , there are at most k paths R_ℓ not ending in a vertex in X . By definition, all inneighbors of any vertex of X lie in some P_i . This implies that there are at most k paths Q_ℓ not containing any vertex of X .

If Q_i contains a vertex of X , then R_i ends on a vertex $u \in X$. Due to Lemma 1, no other R_ℓ contains u . As $|X| \leq k^2$, we have that there are at most k^2 paths R_ℓ which contain some vertex of X . Adding both cases, we obtain that there are at most $k^2 + k$ paths Q_ℓ .

Due to Lemma 1, the subpath R_i of Q_i has at most $k + 1$ vertices. Since the in-forest of H is the union over all R_i , we have that this in-forest has at most $(k + 1)(k^2 + k) = k^3 + 2k^2 + k$ many vertices. Thus, $|V(H)| \leq k(k + 1) + (k + 1)(k^2 + k) \in \mathcal{O}(k^3)$, concluding the proof. \square

We conclude by bounding the number of maximal vertex-disjoint unlabeled local funnels in D . Since we can always partition unlabeled vertices with in- or outdegree at most one into local funnels, by bounding the number of local funnels in such a partitioning, together with the bound on the size of each local funnel, we obtain a bound for the number of unlabeled vertices with in- and outdegree at most one.

Let $\mathcal{H} = \{H_1, H_2, \dots, H_a\}$ be a set of maximal vertex-disjoint unlabeled local funnels in D (in this context, maximal means that $H_i \cup H_j$ is not a local funnel for any two distinct $H_i, H_j \in \mathcal{H}$). Let s_i be the unique source of H_i for each i . We now show that, if there is a solution removing at most k arcs, then $|\mathcal{H}|$ is “small”. By contraposition this means that, if $|\mathcal{H}|$ is “large”, then we have a “no” instance and can stop the kernelization process.

We start with the simple observation that cycles cannot intersect only at a local funnel.

Observation 3. *Let C_i and C_j be two distinct cycles in D such that $V(C_i) \cap V(C_j) \subseteq H_\ell$ for some $H_\ell \in \mathcal{H}$. Then $V(C_i) \cap V(C_j) = \emptyset$.*

Proof. Assume towards a contradiction that $V(C_i) \cap V(C_j) \neq \emptyset$. Let $v \in V(C_i) \cap V(C_j)$ such that the predecessor of v in C_i is different from the predecessor of v in C_j . Then $\text{indeg}(v) > 1$. As H_ℓ is a local funnel, v can only reach one sink t of H_ℓ , implying that t is in both C_i and C_j . The unique neighbor of t is however not in H_ℓ , but it has to be in both C_i and C_j , a contradiction. \square

We partition the set of maximal unlabeled local funnels \mathcal{H} into three sets (1) $\mathcal{F} = \{H_i \in \mathcal{H} \mid \text{there is some } v \in V(H_i) \text{ with } \text{outdeg}_D(v) > 1\}$; (2) $\mathcal{M} = \{H_i \in \mathcal{H} \mid \text{indeg}_D(s_i) > 1\}$; and (3) $\mathcal{X} = \{H_i \in \mathcal{H} \mid \text{indeg}_D(s_i) = 1 \text{ and } \forall v \in V(H_i) : \text{outdeg}_D(v) = 1\}$.

Lemma 3. *If there is a solution $(S, \hat{\ell})$ for (D, ℓ, k) , then $|\mathcal{X}| \leq 2k^2$.*

Proof. Let $H_i \in \mathcal{X}$ and u be the unique inneighbor of s_i . Note that $\text{outdeg}_D(s_i) = 1$. As Dissolve Vertex (Theorem 3.3) is not applicable, we have that $\text{outdeg}_D(u) > 1$ and $\text{indeg}_D(w) > 1$, where w is the unique outneighbor of s_i .

Case 1: $u \in \text{Dom}(\ell)$. Then $\ell(u) = M$ since Set Label (Theorem 3.2) is not applicable. As $\text{outdeg}(u) > 1$, each $H_j \in \mathcal{X}$ with $s_j \in \text{out}_D(u)$ requires one more arc of u to be in S .

Case 2: $u \notin \text{Dom}(\ell)$. If $\text{indeg}_D(u) = 1$, then there is some $v_i \in V(H_i)$ such that $(v_i, u) \in A(D)$, otherwise H_i would not be maximal. Hence, there is a cycle C_i containing u, s_i and v_i . If there is any other $H_j \in \mathcal{X}$ with $s_j \in \text{out}_D(u)$ and

with some $v_j \in V(H_j)$ such that (v_j, u) , then the cycle C_j containing u, s_j and v_j is arc-disjoint to the cycle C_i due to [Observation 3](#). Thus, S must contain at least one arc of each such C_j , implying there are at most k local funnels H_j that fall into this case.

If $\text{indeg}_D(u) > 1$, one arc of u is in S as $\text{outdeg}_D(u) > 1$. Further, $\text{outdeg}_D(u) \leq k$. This means that there are at most k local funnels $H_j \in \mathcal{X}$ with $s_j \in \text{out}_D(u)$. As there can be at most $2k$ such vertices u , we have that there are at most $2k^2$ local funnels $H_j \in \mathcal{X}$ which fall into this case. In the worst case, we have $|\mathcal{X}| \leq \max\{k+1, 2k^2\} \leq 2k^2$. \square

Lemma 4. *If there is a solution $(S, \hat{\ell})$ for (D, ℓ, k) , then $|\mathcal{F}| \leq 2k^2 + 3k$.*

Proof. Let $H_i \in \mathcal{F}$ and let u be the unique inneighbor of s_i in D . Assume u is in some local funnel $H_j \in \mathcal{H}$ and let $D_i = D[V(H_i) \cup V(H_j)]$.

Case 1: D_i is a DAG. Then there are $w_j \in V(H_j)$ and $w_i \in V(H_i)$ such that $\text{indeg}_D(w_j) > 1$, $\text{outdeg}_D(w_i) > 1$ and there is a path P from w_j to w_i . If this were not the case, H_i and H_j would not be maximal, as D_i would be an unlabeled local funnel containing H_i and H_j . Let $G_i \subseteq D$ be a subgraph containing P , two incoming neighbors of w_j and two outgoing neighbors of w_i . Clearly, S contains some arc of G_i . Since H_j and H_i are local funnels, w_j can only reach one sink of H_j , namely u , and w_i can be reached by only one source of H_i , namely s_i . This means in particular that P is the only path from w_j to w_i . Hence, if there is any other $H_\ell \in \mathcal{F}$ that falls into this case, then the corresponding G_ℓ constructed is arc-disjoint to G_i . As there can be at most k arc-disjoint forbidden subgraphs for funnels in D , there are at most k local funnels in \mathcal{F} that fall into this case.

Case 2: D_i is not a DAG. Then there is some cycle C_i containing some $w_i \in V(H_i)$ and some $w_j \in V(H_j)$. Clearly, S contains some arc of C_i . Assume there is some other $H_\ell \in \mathcal{F}$ such that $\text{in}_D(s_\ell) \cap V(H_j) \neq \emptyset$ and $D_\ell = D[V(H_j) \cup V(H_\ell)]$ is not a DAG. Let C_ℓ be a cycle in D_ℓ . From [Observation 3](#) we know C_i and C_ℓ are arc disjoint. As we need one arc in S for each such cycle, we get that there are at most k local funnels falling into this case.

Now assume u is not in any local funnel in \mathcal{H} . We have two cases.

Case 1: $u \in \text{Dom}(\ell)$. Then $\ell(u) = \text{M}$, as [Set Label \(Theorem 3.2\)](#) is not applicable to s_i . Since there is some $v_i \in V(H_i)$ with $\text{outdeg}(v_i) > 1$ and s_i can reach v_i , we have that u can also reach v_i and so $(u, s_i) \in S$ or some arc of H_i is in S . Hence, there are at most k local funnels $H_j \in \mathcal{F}$ with $s_j \in \text{out}(u)$.

Case 2: $u \notin \text{Dom}(\ell)$. As u is not in a local funnel, we have $\text{indeg}(u) > 1$ and $\text{outdeg}(u) > 1$. Since [Set Label \(Theorem 3.2\)](#) is not applicable, $\text{outdeg}(u) \leq k$. Hence, there can be at most k local funnels $H_j \in \mathcal{F}$ with $u \in \text{in}(s_j)$. Because [Lower Bound \(Theorem 3.1\)](#) is not applicable, we know there are at most $2k$ vertices u' with $\text{indeg}(u') > 1$ and $\text{outdeg}(u') > 1$. Thus, there can be at most $2k^2$ local funnels $H_j \in \mathcal{F}$ that fall into this case.

By adding the bounds obtained in each case, we get $\mathcal{F} \leq k + k + k + 2k^2 \in \mathcal{O}(k^2)$. \square

Lemma 5. *If there is a solution $(S, \hat{\ell})$ for (D, ℓ, k) , then $|\mathcal{M}| \leq k^2 + 2k$.*

Proof. Let $H_i \in \mathcal{M}$.

Case 1: $\forall u \in \text{in}(s_i) : u \in \text{Dom}(\ell)$. As **Set Label (Theorem 3.2)** is not applicable, there is some $u \in \text{in}(s_i)$ with $\ell(u) = M$ and $\text{outdeg}(u) > 1$. Hence, S contains some outgoing arc of u . Any additional $H_j \in \mathcal{M}$ that falls into this case increases the outdegree of some u' with $\ell(u') = M$ and $\text{outdeg}(u') > 1$. Thus, if there are more than k local funnels $H_j \in \mathcal{M}$ that fall into this case, then $|S| > k$.

Case 2: There is some $u \in \text{in}(s_i)$ and some $H_j \in \mathcal{H}$ such that $u \in V(H_j)$. As $\text{indeg}(s_i) > 1$ and H_i is maximal, we have that $D_i = D[V(H_j) \cup V(H_i)]$ is not a DAG. Let C_i be the cycle in D_i . If there is some other $H_\ell \in \mathcal{M}$ that falls into this case, we know from **Observation 3** that the corresponding cycle C_ℓ and C_i are arc disjoint. As S must contain one arc of each C_ℓ , if there are more than k local funnels H_ℓ falling into this case, then $|S| > k$.

Case 3: There is some $u \in \text{in}(s_i)$ such that u is not in any local funnel and $u \notin \text{Dom} \ell$. Then $\text{indeg}(u) > 1$ and $\text{outdeg}(u) > 1$. As **Lower Bound (Theorem 3.1)** is not applicable, there can be at most $2k$ such vertices u in D . Further, $\text{outdeg}(u) \leq k$ as **Set Label (Theorem 3.2)** is not applicable. Hence, there can be at most $2k^2$ local funnels $H_j \in \mathcal{M}$ that fall into this case.

By adding all cases together we obtain $|\mathcal{M}| \leq k + k + 2k^2 \in \mathcal{O}(k^2)$, as desired. \square

From **Lemmas 3 to 5**, we easily obtain a bound for the number of vertices in unlabeled local funnels. Together with the fact that **Lower Bound (Theorem 3.1)** is not applicable, we obtain a bound for the number of unlabeled vertices in D .

Lemma 6. *Let D be a reduced digraph. Then there are $\mathcal{O}(k^5)$ vertices $v \in V(D)$ with $v \notin \text{Dom}(\ell)$ and $\text{indeg}(v) = 1 \vee \text{outdeg}(v) = 1$.*

Proof. Let \mathcal{H} be a maximal set of maximal vertex-disjoint local funnels in D . Clearly, every vertex $v \in V(D)$ with $v \notin \text{Dom}(\ell)$ and $\text{indeg}(v) = 1 \vee \text{outdeg}(v) = 1$ is in some local funnel. From **Lemmas 3 to 5** we know that $|\mathcal{H}| \leq |F| + |M| + |X| \in \mathcal{O}(k^2)$. Due to **Lemma 2**, each local funnel has at most $k^3 + 3k^2 + 2k \in \mathcal{O}(k^3)$ many vertices. Hence, there are at most $(5k^2 + 5k)(k^3 + 3k^2 + 2k) \in \mathcal{O}(k^5)$ vertices v lying in some unlabeled local funnel. \square

3.2 Bounding the number of labeled vertices

In **Section 3.1** we exploited the property that unlabeled vertices have bounded degree, and that we can label them if their neighborhood has some special structure captured by the reduction rules. For the labeled vertices, however, we can apply neither of those strategies. Instead, we first exploit the fact that we know the label of a vertex and use this to decide if an arc is never in an optimal solution or if it is always in an optimal solution.

Arcs from M to F vertices clearly need to be removed. We show that we can also ignore arcs from F to M vertices, that is, we can remove them without changing k .

Reduction Rule 3.7 (Remove Arcs). *Let $(v, u) \in A(D)$. If $\ell(v) = F$ and $\ell(u) = M$, remove (v, u) . If $\ell(v) = M$ and $\ell(u) = F$, remove (v, u) and decrease k by 1.*

Proof. Clearly, a solution for (D, ℓ, k) is also a solution for the reduced instance (D', ℓ_r, k') . So let $(\hat{\ell}_r, S_r)$ be a solution for the reduced instance.

If $\ell(v) = M$ and $\ell(u) = F$, then (v, u) is in any solution. Hence, $(\hat{\ell}_r, S_r \cup \{(v, u)\})$ is a solution for the input instance and $|S_r \cup \{(v, u)\}| \leq k$.

If $\ell(v) = F$ and $\ell(u) = M$, then we claim $\hat{\ell}_r$ is a funnel labeling for $D - S_r$. If $D - S_r$ is a DAG, then the claim trivially holds.

Now assume towards a contradiction that $D - S_r$ is not a DAG. Then there is a cycle C using the arc (v, u) . This implies that there is a path P from u to v in $D - S_r$. In particular, this path also exists in $D' - S_r$ since it does not use the arc (v, u) . However, as $\ell(u) = M$ and $\ell(v) = F$, it follows that there is some arc (v', u') in P with $\hat{\ell}_r(v') = M$ and $\hat{\ell}_r(u') = F$. But then $\hat{\ell}_r$ is not a funnel labeling for $D' - S_r$, a contradiction. Hence, a solution for the reduced instance implies a solution for the input instance, proving the rule is safe. \square

We now identify certain vertices that can be removed safely. Clearly, sources and sinks cannot be in any cycle in D . By carefully considering the neighborhood of a source or sink v , we can also prove that v is not “relevant” for any forbidden subgraph for funnels in D .

Reduction Rule 3.8 (Sources and Sinks). *Let $v \in V(D)$ be a labeled vertex where $\text{out}(v) \cup \text{in}(v) \subseteq \text{Dom}(\ell)$. Remove v if one of the following holds.*

1. $\text{indeg}(v) = 0$ and no $u \in \text{out}(v)$ exists with $\ell(u) = F$ and $\text{indeg}(u) > 1$, or
2. $\text{outdeg}(v) = 0$ and no $u \in \text{in}(v)$ exists with $\ell(u) = M$ and $\text{outdeg}(u) > 1$.

Proof. Let (D', ℓ_r, k) be the reduced instance. Clearly, if $(S, \hat{\ell})$ is a solution for (D, ℓ, k) , then, after restricting $(S, \hat{\ell})$ to the vertices in D' , we also have a solution for (D', ℓ_r, k) .

Now let $(S_r, \hat{\ell}_r)$ be a solution for the reduced instance. We consider the case where $\text{indeg}_D(v) = 0$ and there is no $u \in \text{out}(v)$ such that $\ell(u) = F$ and $\text{indeg}(u) > 1$. The other case follows analogously. We claim that the labeling $\hat{\ell} \supseteq \hat{\ell}_r$ with $\hat{\ell}(v) = \ell(v)$ is a funnel labeling for $D - S_r$. Since $\text{indeg}_{D-S_r}(v) = 0$, there can be no cycle containing v . There can be no vertex $u \in V(D)$ with $\hat{\ell}_r(u) = M$ and $\text{outdeg}_{D-S_r}(u) > 1$ as this would imply $\text{outdeg}_{D'-S_r}(u) > 1$ because $\text{indeg}_{D-S_r}(v) = 0$, a contradiction. There can also be no vertex $u \in V(D)$ with $\hat{\ell}_r(u) = F$ and $\text{indeg}_{D-\hat{\ell}_r}(u) > 1$ as all $u \in \text{out}_{D-S_r}(v)$ with $\ell(u) = F$ have $\text{indeg}_{D-S_r}(u) = 1$, and $\text{out}_D(v) \subseteq \text{Dom}(\ell)$.

Because **Remove Arcs** (Theorem 3.7) is not applicable, if $\ell(v) = M$, there is no $u \in \text{out}_{D-S_r}(v)$ with $\ell(u) = F$. Hence, there is no arc $(w, x) \in A(D - S_r)$ with $\hat{\ell}_r(w) = M$ and $\hat{\ell}_r(x) = F$. From the labeling characterization from Theorem 2.1 we have that $(S_r, \hat{\ell}_r)$ is solution for (D, ℓ, k) and the reduction rule is safe. \square

Reduction Rule 3.9 (Remove Cycles). *If there is a weakly-connected component C of D which is a cycle, remove C and decrease k by one.*

Proof. Any solution must remove some arc a of C . As $C - a$ is a path, it is also a funnel and no further arcs need to be removed since all arcs from M to F vertices were removed by **Remove Arcs** (Theorem 3.7). Hence, the reduction rule is safe. \square

Having exhaustively applied [Theorems 3.7 to 3.9](#), we can bound the number of labeled vertices in D . Since [Lower Bound \(Theorem 3.1\)](#) is not applicable, we already have a bound for the number of vertices v with $\ell(v) = F \wedge \text{indeg}(v) > 1$ or $\ell(v) = M \wedge \text{outdeg}(v) > 1$. Hence, we only need to consider vertices in the set $L = \{v \in \text{Dom}(\ell) \mid \ell(v) = F \wedge \text{indeg}(v) \leq 1 \text{ or } \ell(v) = M \wedge \text{outdeg}(v) \leq 1\}$.

To bound $|L|$, we exploit the bound on the number of unlabeled vertices from [Lemma 6](#) and also the fact that such vertices have small degree as [Set Label \(Theorem 3.2\)](#) is not applicable. We first partition L into two subsets $L_1 = \{v \in L \mid \text{in}(v) \cup \text{out}(v) \not\subseteq \text{Dom}(\ell)\}$ and $L_2 = L \setminus L_1$.

Lemma 7. $|L_1| \in \mathcal{O}(k^6)$.

Proof. Let U be the set of unlabeled vertices. Clearly $L_1 \subseteq \text{in}(U) \cup \text{out}(U)$. As [Set Label \(Theorem 3.2\)](#) is not applicable, we have $\text{indeg}(v) \leq k$ and $\text{outdeg}(v) \leq k$ for every $v \in U$. From [Lemma 6](#) we know $|U| \in \mathcal{O}(k^5)$. Hence, $|L_1| \leq |\text{in}(U) \cup \text{out}(U)| \in \mathcal{O}(k^6)$. \square

Lemma 8. $|L_2| \in \mathcal{O}(k)$.

Proof. Let $V_F = \{v \mid \ell(v) = F\}$ and $L_F = V_F \cap L_2$. The case for the vertices labeled with M follows analogously.

Since [Remove Arcs \(Theorem 3.7\)](#) is not applicable, we have $\ell(u) = F$ for all $u \in \text{out}(L_F) \cup \text{in}(L_F)$. Let $R_1 = \{u \in V_F \mid \text{indeg}(u) > 1\}$, $R_2 = \{u \in L_F \mid \text{indeg}(u) \leq 1, \text{out}(u) \cap R_1 \neq \emptyset\}$ and $R_3 = \{u \in L_F \mid \text{indeg}(u) \leq 1, \text{out}(u) \cap R_1 = \emptyset\}$. Note that $L_2 = R_2 \cup R_3$ and $R_1 \cap L_2 = \emptyset$.

A solution set $S \subseteq A(D)$ must contain $\text{indeg}(v) - 1$ many incoming arcs of v for every $v \in R_1$. As each $u \in R_2$ has some $v \in R_1$ as outneighbor, it follows that $|R_2| \leq 2k$.

Let $v \in R_3$. We claim that v can reach some vertex of R_2 . Since [Sources and Sinks \(Theorem 3.8\)](#) is not applicable and $\text{out}(v) \cap R_1 = \emptyset$, we have $\text{indeg}(v) = 1$ and $\text{outdeg}(v) \geq 1$. This means that, if we successively follow the outneighbors of v , we reach a vertex of R_2 or find a cycle C using only vertices of R_3 . Assume there is such a cycle C . As $\text{indeg}(u) \leq 1$ for all $u \in R_3$, we have that v is in C . As [Theorem 3.9](#) is not applicable, there is some $w \in V(C)$ with $\text{outdeg}(w) > 1$. Let $x \in \text{out}(w) \setminus V(C)$. By following a maximal path starting with the arc (w, x) , we necessarily reach some vertex of R_2 , as otherwise there would be some vertex $y \in R_3$ with $\text{indeg}(y) > 1$. Hence, every vertex of R_3 can reach some $u \in R_2$.

We greedily construct vertex-disjoint paths P_1, P_2, \dots, P_a ending in R_2 whose inner vertices lie in R_3 . For a vertex $v \in R_3$ take an arbitrary $u \in R_2$ such that v can reach u . Consider a path P from v to u . If none of its vertices lie in any already constructed P_i , we just take the path P into our set of paths. Otherwise, assume that P intersects some P_i at w and let w be the first such vertex in P . Since the indegree of any vertex in $R_3 \cup R_2$ is at most one, it follows that w is the starting point of P_i . Hence, we can obtain a path P_j by taking the path from v to w in P and then concatenating P_i . As w is the first vertex of P intersecting any other path, it follows that P_j only intersects P_i . By replacing P_i with P_j , we obtain a path that also contains v . We repeat this process until we covered all $v \in R_3$.

Since $|R_2| \leq 2k$, it follows that $a \leq 2k$. We now prove that $|V(P_i)| \leq 4$ for any P_i in our set of vertex-disjoint paths. Note that $\text{indeg}(u) \leq 1$ for any vertex $u \in V(P_i)$.

Since **Dissolve Vertex** (Theorem 3.3) is not applicable, any inner vertex u of P_i has $\text{outdeg}(u) > 1$. Let w be the successor of u in P_i . As **Shift Neighbors** (Theorem 3.5) is not applicable, it follows that $\text{indeg}(w) > 1$ or $v \in \text{out}(u)$ where v is the unique inneighbor of u . If $\text{indeg}(w) > 1$, then $u \in R_2$ and is the endpoint of P_i , a contradiction to the assumption that u is an inner vertex of P_i . Otherwise, we know that $u \notin \text{out}(w)$ as $\text{indeg}(u) = 1$. If u is the only inner vertex of P_i , then $|V(P_i)| \leq 3$. Otherwise, its successor x in P_i is an inner vertex of P_i (since v is the starting point of P_i). Further, there is no arc (x, u) as $\text{indeg}(u) = 1$. Hence, we can apply the same argumentation to x and conclude that it has some outneighbor w with $\text{indeg}(w) > 1$, implying $w \in R_2$ and $|V(P_i)| \leq 4$.

Since $|V(P_i)| \leq 4$ and $a \leq 2k$, we have that $|L_2| \leq 6k$. Because $L_2 = R_2 \cup R_3$, we have that $|L_2| \leq 8k \in \mathcal{O}(k)$, as desired. \square

Lemma 9. *Let (D, ℓ, k) be a FADL instance where Theorems 3.1, 3.2, 3.3, 3.5, 3.6, 3.7, 3.8 and 3.9 are not applicable. Then $|V(D)| \in \mathcal{O}(k^6)$ and $|A(D)| \in \mathcal{O}(k^6)$.*

Proof. As **Lower Bound** (Theorem 3.1) is not applicable, there are at most $2k$ vertices v with $\text{indeg}(v) > 1$ and $\text{outdeg}(v) > 1$, and also at most $2k$ many vertices v with $\ell(v) = \text{F} \wedge \text{indeg}(v) > 1$ or $\ell(v) = \text{M} \wedge \text{outdeg}(v) > 1$. From Lemma 6 we know there are $\mathcal{O}(k^5)$ many unlabeled vertices $v \in V(D)$ with $\text{indeg}(v) \leq 1$ or $\text{outdeg}(v) \leq 1$. Finally, due to Lemmas 7 and 8 there are $\mathcal{O}(k^6)$ vertices v with $\ell(v) = \text{F} \wedge \text{indeg}(v) \leq 1$ or $\ell(v) = \text{M} \wedge \text{outdeg}(v) = 1$. As any vertex in D falls into one of these groups, we have $|V(D)| \in \mathcal{O}(k^6)$.

As **Remove Arcs** (Theorem 3.7) is not applicable, there is no arc (v, u) where $v, u \in \text{Dom}(\ell)$ and $\ell(v) \neq \ell(u)$. Since there are $\mathcal{O}(k^5)$ many unlabeled vertices and every unlabeled vertex has in- and outdegree at most $k + 1$, there are $\mathcal{O}(k^6)$ arcs (v, u) where $v \notin \text{Dom}(\ell)$ or $u \notin \text{Dom}(\ell)$.

Now let (v, u) be some arc where $v, u \in \text{Dom}(\ell)$. Note that $\ell(v) = \ell(u)$.

Case 1: $v, u \in L$. Then $\text{outdeg}(v) = 1$ (if $\ell(v) = \text{M}$) or $\text{indeg}(u) = 1$ (if $\ell(u) = \text{F}$). Thus, there can be at most $|L| \in \mathcal{O}(k^6)$ many arcs (v, u) where $v, u \in L$.

Case 2: $v, u \notin L$. As **Lower Bound** (Theorem 3.1) is not applicable, there can be at most $2k$ such vertices. Thus, there are at most $4k^2$ arcs between labeled vertices not in L .

Case 3: Exactly one of v, u is in L .

Case 3.1: $v \notin L \wedge \ell(v) = \text{F}$ or $u \notin L \wedge \ell(u) = \text{M}$. Then $\text{indeg}(u) = 1$ or $\text{outdeg}(v) = 1$. Hence, there can be at most $|L| \in \mathcal{O}(k^6)$ such arcs.

Case 3.2: $v \notin L \wedge \ell(v) = \text{M}$ or $u \notin L \wedge \ell(u) = \text{F}$. If $v \notin L$, then at least half of its outgoing arcs need to be in a solution set. Similarly, if $u \notin L$, at least half of its incoming arcs need to be in a solution set. Hence, there can be at most $2k$ many arcs falling into this case. By adding all cases together, we obtain that $|A(D)| \in \mathcal{O}(k^6)$, concluding the proof. \square

4 Computing the Kernel

In [Sections 3.1](#) and [3.2](#) we defined the reduction rules for the kernelization process and showed that, if none of the reduction rules are applicable to a digraph D , then the size of D is polynomially bounded on k . To conclude the proof that FADS admits a polynomial problem kernel, we show that it is possible to apply all reduction rules in linear time and also reduce the FADL instance back into a FADS instance.

Theorem 4.1. *FADS admits a kernel with $\mathcal{O}(k^6)$ vertices and $\mathcal{O}(k^7)$ arcs which can be computed in $\mathcal{O}(n + m)$ time, where $n = |V(D)|$, $m = |A(D)|$ and D is the input digraph.*

Proof. We start by reducing the FADS instance into a FADL instance by adding an empty labeling ℓ . We apply the reduction rules exhaustively in the order they are defined. To apply [Lower Bound \(Theorem 3.1\)](#) and [Set Label \(Theorem 3.2\)](#) we only need to check the labels and degrees of a vertex and its neighbors. Whenever the label of a vertex changes, we need to recheck if we can apply the reduction rules to its neighbors. Hence, we need to visit each arc constantly many times, as the label of a vertex can change at most once.

For [Dissolve Vertex \(Theorem 3.3\)](#), [Shift Neighbors \(Theorem 3.5\)](#) and [Labeled Neighbor \(Theorem 3.6\)](#) we need to search for a path P in D . However, all vertices of P have in-degree one or all have out-degree one. Thus, it suffices to find one of the endpoints of P and then follow its unique outneighbor or its unique inneighbor (depending on which case of the rule is being applied). Once P is obtained, we only need to check the labels and degrees of the vertices.

[Remove Arcs \(Theorem 3.7\)](#) is applied by iterating through all arcs. [Sources and Sinks \(Theorem 3.8\)](#) is applied by breadth-first search from all labeled sources and sinks of D . [Remove Cycles \(Theorem 3.9\)](#) is applied by breadth-first search in each weakly connected component of D .

Since we only need to recheck the conditions of a rule if something in the neighborhood of a vertex changed, we can exhaustively apply all reduction rules in linear time.

From [Lemma 9](#) we know $|V(D)| \in \mathcal{O}(k^6)$ and $|A(D)| \in \mathcal{O}(k^6)$. We now reduce the FADL instance back into a FADS instance (D', k) in order to obtain a kernel for the original problem. We first set $D' := D$ and add $k + 2$ vertices f_1, f_2, \dots, f_{k+2} and $k + 2$ vertices m_1, m_2, \dots, m_{k+2} to D' . Let $v \in \text{Dom}(\ell)$. If $\ell(v) = \text{F}$, we add the arc (v, f_i) for each $1 \leq i \leq k + 2$. If $\ell(v) = \text{M}$, we add the arc (m_i, v) for each $1 \leq i \leq k + 2$.

Trivially, a solution for the FADL instance is also a solution for the FADS instance. It is also easy to see that, if there is some arc set $S_r \subseteq A(D')$ and some funnel labeling $\hat{\ell}_r$ for $D' - S_r$ such that $\ell(v) \neq \hat{\ell}_r(v)$ for some $v \in \text{Dom}(\ell)$, then $|S_r| > k$. Hence, a solution for (D', k) implies a solution for (D, ℓ, k) .

We added $2k + 4$ vertices and $\mathcal{O}(k^7)$ many arcs to D' , and so $|V(D')| \in \mathcal{O}(k^6)$ and $|A(D')| \in \mathcal{O}(k^7)$, thus concluding the proof. \square

5 Conclusion

The kernelization algorithm provided in this paper heavily relies on the characterizations of [Theorem 2.1](#) for funnels. Both the characterization by forbidden subgraphs as well as the labeling characterization allowed us to derive reduction rules based only on “local” substructures as the degree or neighborhood of a vertex. In a sense, this “locality” property saved us from computing any set of vertex-disjoint local funnels, despite the fact that the results and reduction rules from [Section 3.1](#) heavily rely on local funnels.

The polynomial kernels for OUT-FOREST-VDS and PUMPKIN-VDS due to [\[12\]](#) also rely on “localized” forbidden substructures. We consider that generalizing these results to certain digraph classes characterized by forbidden substructures to be a very interesting direction for future research.

References

- [1] Faisal N Abu-Khzam. A kernelization algorithm for d -hitting set. *Journal of Computer and System Sciences*, 76(7):524–531, 2010.
- [2] Akanksha Agrawal, Saket Saurabh, Roohani Sharma, and Meirav Zehavi. Kernels for deletion to classes of acyclic digraphs. *Journal of Computer and System Sciences*, 92:9–21, 2018.
- [3] Jørgen Bang-Jensen, Alessandro Maddaloni, and Saket Saurabh. Algorithms and kernels for feedback set problems in generalizations of tournaments. *Algorithmica*, 76(2):320–343, 2016.
- [4] Jianer Chen, Yang Liu, Songjian Lu, Barry O’sullivan, and Igor Razgon. A fixed-parameter algorithm for the directed feedback vertex set problem. *Journal of the ACM (JACM)*, 55(5):21, 2008.
- [5] Marek Cygan, Fedor V Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized algorithms*, volume 4. Springer, 2015.
- [6] Rodney G Downey and Michael R Fellows. *Fundamentals of parameterized complexity*, volume 4. Springer, 2013.
- [7] Fedor V Fomin, Daniel Lokshtanov, Saket Saurabh, and Meirav Zehavi. *Kernelization: theory of parameterized preprocessing*. Cambridge University Press, 2019.
- [8] Stefan Kratsch. Recent developments in kernelization: A survey. *Bulletin of EATCS*, 2(113), 2014.
- [9] Daniel Lokshtanov, Neeldhara Misra, and Saket Saurabh. Kernelization–preprocessing with a guarantee. In *The Multivariate Algorithmic Revolution and Beyond*, pages 129–161. Springer, 2012.

- [10] Daniel Lokshtanov, MS Ramanujan, Saket Saurabh, Roohani Sharma, and Meirav Zehavi. Wannabe bounded treewidth graphs admit a polynomial kernel for dfvs. In *Workshop on Algorithms and Data Structures*, pages 523–537. Springer, 2019.
- [11] Marcelo Garlet Millani, Hendrik Molter, Rolf Niedermeier, and Manuel Sorge. Efficient algorithms for measuring the funnel-likeness of DAGs. In Jon Lee, Giovanni Rinaldi, and A. Ridha Mahjoub, editors, *Combinatorial Optimization*, pages 183–195, Cham, 2018. Springer International Publishing. ISBN 978-3-319-96151-4.
- [12] Matthias Mnich and Erik Jan van Leeuwen. Polynomial kernels for deletion to classes of acyclic digraphs. *Discrete Optimization*, 25:48–76, 2017.