

LOW REGULARITY WELL-POSEDNESS FOR THE YANG-MILLS SYSTEM IN FOURIER-LEBESGUE SPACES

HARTMUT PECHER

FAKULTÄT FÜR MATHEMATIK UND NATURWISSENSCHAFTEN
BERGISCHE UNIVERSITÄT WUPPERTAL
GAUSSSTR. 20
42119 WUPPERTAL
GERMANY
E-MAIL PECHER@MATH.UNI-WUPPERTAL.DE

ABSTRACT. The Cauchy problem for the Yang-Mills system in three space dimensions with data in Fourier-Lebesgue spaces $\dot{H}^{s,r}$, $1 < r \leq 2$, is shown to be locally well-posed, where we have to assume only almost optimal minimal regularity for the data with respect to scaling as $r \rightarrow 1$. This is true despite of the fact that no null condition is known for one of the critical quadratic nonlinearities, which prevented by now the corresponding result in the classical case $r = 2$ with data in standard Sobolev spaces.

1. INTRODUCTION

Let \mathcal{G} be the Lie group $SO(n, \mathbb{R})$ (the group of orthogonal matrices of determinant 1) or $SU(n, \mathbb{C})$ (the group of unitary matrices of determinant 1) and g its Lie algebra $so(n, \mathbb{R})$ (the algebra of trace-free skew symmetric matrices) or $su(n, \mathbb{C})$ (the algebra of trace-free skew hermitian matrices) with Lie bracket $[X, Y] = XY - YX$ (the matrix commutator). For given $A_\alpha : \mathbb{R}^{1+n} \rightarrow g$ we define the curvature $F = F[A]$ by

$$F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha + [A_\alpha, A_\beta], \quad (1)$$

where $\alpha, \beta \in \{0, 1, \dots, n\}$ and $D_\alpha = \partial_\alpha + [A_\alpha, \cdot]$.

Then the Yang-Mills system is given by

$$D^\alpha F_{\alpha\beta} = 0 \quad (2)$$

in Minkowski space $\mathbb{R}^{1+n} = \mathbb{R}_t \times \mathbb{R}_x^n$, where $n \geq 3$, with metric $diag(-1, 1, \dots, 1)$. Greek indices run over $\{0, 1, \dots, n\}$, Latin indices over $\{1, \dots, n\}$, and the usual summation convention is used. We use the notation $\partial_\mu = \frac{\partial}{\partial x_\mu}$, where we write $(x^0, x^1, \dots, x^n) = (t, x^1, \dots, x^n)$ and also $\partial_0 = \partial_t$.

Setting $\beta = 0$ in (2) we obtain the Gauss-law constraint

$$\partial^j F_{j0} + [A^j, F_{j0}] = 0.$$

The total energy for Yang-Mills at time t is given by

$$\mathcal{E}(t) = \sum_{0 \leq \alpha, \beta \leq n} \int_{\mathbb{R}^n} |F_{\alpha\beta}(t, x)|^2 dx,$$

and is conserved for a smooth solution decaying sufficiently fast at spatial infinity.

The Yang-Mills system is invariant with respect to the scaling

$$A_\lambda(t, x) = \lambda A(\lambda t, \lambda x) \quad , \quad F_\lambda(t, x) = \lambda^2 F(\lambda t, \lambda x).$$

2010 Mathematics Subject Classification: 35Q40, 35L70

Key words and phrases: Yang-Mills, local well-posedness, Lorenz gauge

This implies

$$\begin{aligned} \|A_\lambda(0, \cdot)\|_{\widehat{H}^{s,r}} &= \lambda^{1+s-\frac{n}{r}} \|a_\lambda\|_{\widehat{H}^{s,r}}, \\ \|F_\lambda(0, \cdot)\|_{\widehat{H}^{l,r}} &= \lambda^{2+l-\frac{n}{r}} \|f_\lambda\|_{\widehat{H}^{l,r}}. \end{aligned}$$

Here $\|u\|_{\widehat{H}^{s,r}} := \|\langle \xi \rangle^s \widehat{u}(\xi)\|_{L^{r'}}$, where r and r' are dual exponents, and \widehat{H} denotes the homogeneous space. Therefore the scaling critical exponent is $s = \frac{n}{r} - 1$ for A and $l = \frac{n}{r} - 2$ for F .

The system is gauge invariant. Given a sufficiently smooth function $U : \mathbb{R}^{1+n} \rightarrow \mathcal{G}$ we define the gauge transformation T by $TA_0 = A'_0$, $T(A_1, \dots, A_n) = (A'_1, \dots, A'_n)$, where

$$A_\alpha \mapsto A'_\alpha = UA_\alpha U^{-1} - (\partial_\alpha U)U^{-1}.$$

It is well-known that if (A_0, \dots, A_n) satisfies (1),(2) so does (A'_0, \dots, A'_n) .

Hence we may impose a gauge condition. We exclusively study the case $n = 3$ and Lorenz gauge $\partial^\alpha A_\alpha = 0$. Other convenient gauges are the Coulomb gauge $\partial^j A_j = 0$ and the temporal gauge $A_0 = 0$. Our aim is to obtain local well-posedness for data with minimal regularity. Up to now there exists no result for data arbitrarily close to the critical scaling regularity.

The classical case $r = 2$ with data in standard Sobolev spaces was considered by Klainerman and Machedon [KM], who made the decisive detection that the nonlinearity satisfies a so-called null condition, which enabled them to prove global well-posedness in temporal and in Coulomb gauge in energy space. The corresponding result in Lorenz gauge, where the Yang-Mills equations can be formulated as a system of nonlinear wave equations, was shown by Selberg and Tesfahun [ST], who discovered that also in this case some of the nonlinearities have a null structure. Tesfahun [T] improved this result to data without finite energy, namely for $(A(0), (\partial_t A)(0)) \in H^s \times H^{s-1}$ and $(F(0), (\partial_t F)(0)) \in H^l \times H^{l-1}$ with $s = \frac{6}{7} + \epsilon$ and $l = -\frac{1}{14} + \epsilon$ for any $\epsilon > 0$ by discovering an additional partial null structure. A further improvement was achieved by the author [P], namely to $(s, l) = (\frac{5}{7} + \epsilon, -\frac{1}{7} + \epsilon)$ by modifying the solution spaces appropriately.

As the critical case with respect to scaling is $(s, l) = (\frac{1}{2}, -\frac{1}{2})$, there is however still a gap, a phenomenon, which is also present in other gauges. The present paper closes this gap in the sense that as $r \rightarrow 1$ we almost reach the critical case $(s, l) = (2, 1)$.

Local well-posedness in energy space was also given by Oh [O] using a new gauge, namely the Yang-Mills heat flow. He was also able to show that this solution can be globally extended [O1]. The Cauchy problem was also treated in higher space dimensions by several authors ([KS],[KT],[KrT],[KrSt],[P1]).

In the present paper we treat the local well-posedness problem for the Yang-Mills system in Lorenz gauge and space dimension $n = 3$ in the case of data $(A(0), (\partial_t A)(0)) \in \widehat{H}^{s,r} \times \widehat{H}^{s-1,r}$ and $(F(0), (\partial_t F)(0)) \in \widehat{H}^{l,r} \times \widehat{H}^{l-1,r}$ in Fourier-Lebesgue spaces for $r \neq 2$, which coincide with the classical Sobolev spaces H^s for $r = 2$. The assumption is that $s = \frac{16}{7r} - \frac{2}{7} + \delta$ and $l = \frac{15}{7r} - \frac{8}{7} + \delta$, where any $\delta > 0$ is admissible. Thus we obtain $s \rightarrow 2 + \delta$ and $l \rightarrow 1 + \delta$ as $r \rightarrow 1$, which is almost optimal with respect to scaling.

Such an approach was used by several authors already, starting with Vargas-Vega [VV] for 1D Schrödinger equations. Grünrock showed LWP for the modified KdV equation [G], a result which was improved by Grünrock and Vega [GV]. Grünrock treated derivative nonlinear wave equations in 3+1 dimensions [G1] and obtained an almost optimal result as $r \rightarrow 1$ with respect to scaling. Systems of nonlinear wave equations in the 2+1 dimensional case for nonlinearities which

fulfill a null condition were considered by Grigoryan-Nahmod [GN]. The latter two results are based on estimates by Foschi and Klainerman [FK].

In Chapter 2 we start by giving the formulation of the Yang-Mills equations as a system of semilinear wave equations and formulate the main theorem (Theorem 2.1 and Cor. 2.1). In chapter 3 we recall some basic facts about our solution spaces and a general local well-posedness theorem for the Cauchy problem for systems of nonlinear wave equations with data in Fourier-Lebesgue spaces, which allows to reduce it to estimates for the nonlinearities. In chapter 4 we give the final formulation of the system in terms of null forms as far as possible. The bi-, tri- and quadrilinear estimates sufficient for the local well-posedness result are formulated, where we rely on Tesfahun's paper [T]. In chapter 5 we prove bilinear estimates for the null forms and for general bilinear terms in generalized Bourgain-Klainerman-Machedon spaces $H_{s,b}^r$ (and $X_{s,b,\pm}^r$) based on elementary but non-trivial estimates by Foschi and Klainerman [FK], Grünrock [G] and Grigoryan-Nahmod [GN]. In the final chapter 6 we prove the multilinear estimates formulated in chapter 4 by reduction to the bilinear estimates of chapter 5.

2. MAIN RESULTS

Expanding (2) in terms of the gauge potentials $\{A_\alpha\}$, we obtain:

$$\square A_\beta = \partial_\beta \partial^\alpha A_\alpha - [\partial^\alpha A_\alpha, A_\beta] - [A^\alpha, \partial^\alpha A_\beta] - [A^\alpha, F_{\alpha\beta}]. \quad (3)$$

If we now impose the Lorenz gauge condition, the system (3) reduces to the nonlinear wave equation

$$\square A_\beta = -[A^\alpha, \partial_\alpha A_\beta] - [A^\alpha, F_{\alpha\beta}]. \quad (4)$$

In addition, regardless of the choice of gauge, F satisfies the wave equation

$$\begin{aligned} \square F_{\beta\gamma} &= -[A^\alpha, \partial_\alpha F_{\beta\gamma}] - \partial^\alpha [A_\alpha, F_{\beta\gamma}] - [A^\alpha, [A_\alpha, F_{\beta\gamma}]] \\ &\quad - 2[F_{\beta\alpha}, F_{\gamma\alpha}], \end{aligned} \quad (5)$$

where we refer to [ST], chapter 3.2.

Expanding the second and fourth terms in (5), and also imposing the Lorenz gauge, yields

$$\begin{aligned} \square F_{\beta\gamma} &= -2[A^\alpha, \partial_\alpha F_{\beta\gamma}] + 2[\partial_\gamma A^\alpha, \partial_\alpha A_\beta] - 2[\partial_\beta A^\alpha, \partial_\alpha A_\gamma] \\ &\quad + 2[\partial^\alpha A_\beta, \partial_\alpha A_\gamma] + 2[\partial_\beta A^\alpha, \partial_\gamma A_\alpha] - [A^\alpha, [A_\alpha, F_{\beta\gamma}]] \\ &\quad + 2[F_{\alpha\beta}, [A^\alpha, A_\gamma]] - 2[F_{\alpha\gamma}, [A^\alpha, A_\beta]] - 2[[A^\alpha, A_\beta], [A_\alpha, A_\gamma]]. \end{aligned} \quad (6)$$

Note on the other hand by expanding the last term in the right hand side of (4), we obtain

$$\square A_\beta = -2[A^\alpha, \partial_\alpha A_\beta] + [A^\alpha, \partial_\beta A_\alpha] - [A^\alpha, [A_\alpha, A_\beta]]. \quad (7)$$

We want to solve the system (6)-(7) simultaneously for A and F . So to pose the Cauchy problem for this system, we consider initial data for (A, F) at $t = 0$:

$$A(0) = a, \quad \partial_t A(0) = \dot{a}, \quad F(0) = f, \quad \partial_t F(0) = \dot{f}. \quad (8)$$

In fact, the initial data for F can be determined from (a, \dot{a}) as follows:

$$\begin{cases} f_{ij} = \partial_i a_j - \partial_j a_i + [a_i, a_j], \\ f_{0i} = \dot{a}_i - \partial_i a_0 + [a_0, a_i], \\ \dot{f}_{ij} = \partial_i \dot{a}_j - \partial_j \dot{a}_i + [\dot{a}_i, a_j] + [a_i, \dot{a}_j], \\ \dot{f}_{0i} = \partial^j f_{ji} + [a^\alpha, f_{\alpha i}] \end{cases} \quad (9)$$

where the first three expressions come from (1) whereas the last one comes from (2) with $\beta = i$.

Note that the Lorenz gauge condition $\partial^\alpha A_\alpha = 0$ and (2) with $\beta = 0$ impose the constraints

$$\dot{a}_0 = \partial^i a_i, \quad \partial^i f_{i0} + [a^i, f_{i0}] = 0. \quad (10)$$

Now we formulate our main theorem.

Theorem 2.1. *Let $1 < r \leq 2$, $\delta > 0$. Assume that s and l satisfy the following conditions:*

$$s = \frac{16}{7r} - \frac{2}{7} + \delta, \quad l = \frac{15}{7r} - \frac{8}{7} + \delta.$$

Given initial data $(a, \dot{a}) \in \widehat{H}^{s,r} \times \widehat{H}^{s-1,r}$, $(f, \dot{f}) \in \widehat{H}^{l,r} \times \widehat{H}^{l-1,r}$, there exists a time $T > 0$, $T = T(\|a\|_{\widehat{H}^{s,r}}, \|\dot{a}\|_{\widehat{H}^{s-1,r}}, \|f\|_{\widehat{H}^{l,r}}, \|\dot{f}\|_{\widehat{H}^{l-1,r}})$, such that the Cauchy problem (6),(7),(8) has a unique solution $A_\mu \in X_{s,b,+}^r[0, T] + X_{s,b,-}^r[0, T]$, $F \in X_{l,a,+}^r[0, T] + X_{l,a,-}^r[0, T]$ (these spaces are defined in Def. 2.1). Here $a = \frac{1}{r}+$ and $b = \frac{1}{2} + \frac{1}{2r}+$. This solution has the regularity

$$A_\mu \in C^0([0, T], \widehat{H}^{s,r}) \cap C^1([0, T], \widehat{H}^{s-1,r}), \quad F \in C^0([0, T], \widehat{H}^{l,r}) \cap C^1([0, T], \widehat{H}^{l-1,r}).$$

The solution depends continuously on the data and persistence of higher regularity holds.

Corollary 2.1. *Let s, r fulfill the assumptions of Theorem 2.1. Moreover assume that the initial data fulfill (9) and (10). Given any $(a, \dot{a}) \in \widehat{H}^{s,r} \times \widehat{H}^{s-1,r}$, there exists a time $T = T(\|a\|_{\widehat{H}^{s,r}}, \|\dot{a}\|_{\widehat{H}^{s-1,r}}, \|f\|_{\widehat{H}^{l,r}}, \|\dot{f}\|_{\widehat{H}^{l-1,r}})$, such that the solution (A, F) of Theorem 2.1 satisfies the Yang-Mills system (1),(2) with Cauchy data (a, \dot{a}) and the Lorenz gauge condition $\partial^\alpha A_\alpha = 0$.*

Proof of the Corollary. If $(a, \dot{a}) \in \widehat{H}^{s,r} \times \widehat{H}^{s-1,r}$, then (f, \dot{f}) , defined by (9), fulfill $(f, \dot{f}) \in \widehat{H}^{l,r} \times \widehat{H}^{l-1,r}$, as one easily checks. Thus we may apply Theorem 2.1. The solution (A, F) does not necessarily fulfill the Lorenz gauge condition and (1), i.e. $F = F[A]$. If however the conditions (9) and (10) are assumed then these properties are satisfied and (A, F) is a solution of the Yang-Mills system (1),(2) with Cauchy data (a, \dot{a}) . This was shown in [ST], Remark 2. \square

Let us fix some notation. We denote the Fourier transform with respect to space and time by $\widehat{\cdot}$. $\square = \partial_t^2 - \Delta$ is the d'Alembert operator, $a_\pm := a \pm \epsilon$ for a sufficiently small $\epsilon > 0$, and $\langle \cdot \rangle := (1 + |\cdot|^2)^{\frac{1}{2}}$.

Let Λ^α be the multiplier with symbol $\langle \xi \rangle^\alpha$. Similarly let D^+ , and D^- be the multipliers with symbols $|\xi|^\alpha$ and $|\tau| - |\xi|^\alpha$, respectively.

Definition 2.1. *Let $1 \leq r \leq 2$, $s, b \in \mathbb{R}$. The wave-Sobolev spaces $H_{s,b}^r$ are the completion of the Schwarz space $\mathcal{S}(\mathbb{R}^{1+3})$ with norm*

$$\|u\|_{H_{s,b}^r} = \|\langle \xi \rangle^s \langle |\tau| - |\xi| \rangle^b \widehat{u}(\tau, \xi)\|_{L_{\tau\xi}^{r'}},$$

where r' is the dual exponent to r . We also define $H_{s,b}^r[0, T]$ as the space of the restrictions of functions in $H_{s,b}^r$ to $[0, T] \times \mathbb{R}^3$. Similarly we define $X_{s,b,\pm}^r$ with norm

$$\|\phi\|_{X_{s,b,\pm}^r} := \|\langle \xi \rangle^s \langle \tau \pm |\xi| \rangle^b \widehat{\phi}(\tau, \xi)\|_{L_{\tau\xi}^{r'}}$$

and $X_{s,b,\pm}^r[0, T]$. $\dot{H}_{s,b}^r$ and $\dot{X}_{s,b,\pm}^r$ are the corresponding homogeneous spaces, where $\langle \xi \rangle$ is replaced by $|\xi|$.

3. PRELIMINARIES

We start by collecting some fundamental properties of the solution spaces. We rely on [G]. The spaces $X_{s,b,\pm}^r$ with norm

$$\|\phi\|_{X_{s,b,\pm}^r} := \|\langle \xi \rangle^s \langle \tau \pm |\xi| \rangle^b \tilde{\phi}(\tau, \xi)\|_{L_{\tau\xi}^{r'}}$$

for $1 < r < \infty$ are Banach spaces with \mathcal{S} as a dense subspace. The dual space is $X_{-s,-b,\pm}^{r'}$, where $\frac{1}{r} + \frac{1}{r'} = 1$. The complex interpolation space is given by

$$(X_{s_0,b_0,\pm}^{r_0}, X_{s_1,b_1,\pm}^{r_1})_{[\theta]} = X_{s,b,\pm}^r,$$

where $s = (1 - \theta)s_0 + \theta s_1$, $\frac{1}{r} = \frac{1-\theta}{r_0} + \frac{\theta}{r_1}$, $b = (1 - \theta)b_0 + \theta b_1$. Similar properties has the space $H_{s,b}^r$.

If $u = u_+ + u_-$, where $u_{\pm} \in X_{s,b,\pm}^r[0, T]$, then $u \in C^0([0, T], \hat{H}^{s,r})$, if $b > \frac{1}{r}$.

The "transfer principle" in the following proposition, which is well-known in the case $r = 2$, also holds for general $1 < r < \infty$ (cf. [GN], Prop. A.2 or [G], Lemma 1). We denote $\|u\|_{\hat{L}_t^p(\hat{L}_x^q)} := \|\tilde{u}\|_{L_{\tau\xi}^{p'}(L_{\xi}^q)}$.

Proposition 3.1. *Let $1 \leq p, q \leq \infty$. Assume that T is a bilinear operator which fulfills*

$$\|T(e^{\pm 1itD} f_1, e^{\pm 2itD} f_2)\|_{\hat{L}_t^p(\hat{L}_x^q)} \lesssim \|f_1\|_{\hat{H}^{s_1,r}} \|f_2\|_{\hat{H}^{s_2,r}}$$

for all combinations of signs \pm_1, \pm_2 , then for $b > \frac{1}{r}$ the following estimate holds:

$$\|T(u_1, u_2)\|_{\hat{L}_t^p(\hat{L}_x^q)} \lesssim \|u_1\|_{H_{s_1,b}^r} \|u_2\|_{H_{s_2,b}^r}.$$

The general local well-posedness theorem is the following (cf. [G], Thm. 1).

Theorem 3.1. *Let $N(u, v)$ and $M(u, v)$ be multilinear functions. Assume that for given $s, l \in \mathbb{R}$, $1 < r < \infty$ there exist $b, a > \frac{1}{r}$ such that the estimates*

$$\|N(u, v)\|_{X_{s,b-1+,\pm}^r} \leq \omega_1(\|u\|_{X_{s,b}^r}, \|v\|_{X_{l,a}^r})$$

and

$$\|M(u, v)\|_{X_{l,a-1+,\pm}^r} \leq \omega_2(\|u\|_{X_{s,b}^r}, \|v\|_{X_{l,a}^r})$$

are valid with nondecreasing functions ω_j , where $\|u\|_{X_{s,b}^r} := \|u\|_{X_{s,b,-}^r} + \|u\|_{X_{s,b,+}^r}$. Then there exist $T = T(\|u_0\|_{\hat{H}^{s,r}}, \|v_0\|_{\hat{H}^{l,r}}) > 0$ and a unique solution $(u, v) \in X_{s,b,\pm}^r[0, T] \times X_{l,a,\pm}^r[0, T]$ of the Cauchy problem

$$\partial_t u \pm i\Lambda u = N(u, v) \quad , \quad \partial_t v \pm i\Lambda v = M(u, v)$$

$$u(0) = u_0 \in \hat{H}^{s,r} \quad , \quad v(0) = v_0 \in \hat{H}^{l,r}.$$

This solution is persistent and the mapping data upon solution $(u_0, v_0) \mapsto (u, v)$, $\hat{H}^{s,r} \times \hat{H}^{l,r} \rightarrow X_{s,b,\pm}^r[0, T_0] \times X_{l,a,\pm}^r[0, T_0]$ is locally Lipschitz continuous for any $T_0 < T$.

4. REFORMULATION OF THE PROBLEM AND NULL STRUCTURE

The reformulation of the Yang-Mills equations and the reduction of our main theorem to nonlinear estimates is completely taken over from Tesfahun [T] (cf. also the fundamental paper by Selberg and Tesfahun [ST]).

The standard null forms are given by

$$\begin{cases} Q_0(u, v) = \partial_\alpha u \partial^\alpha v = -\partial_t u \partial_t v + \partial_i u \partial^i v, \\ Q_{\alpha\beta}(u, v) = \partial_\alpha u \partial_\beta v - \partial_\beta u \partial_\alpha v. \end{cases} \quad (11)$$

For g -valued u, v , define a commutator version of null forms by

$$\begin{cases} Q_0[u, v] = [\partial_\alpha u, \partial^\alpha v] = Q_0(u, v) - Q_0(v, u), \\ Q_{\alpha\beta}[u, v] = [\partial_\alpha u, \partial_\beta v] - [\partial_\beta u, \partial_\alpha v] = Q_{\alpha\beta}(u, v) + Q_{\alpha\beta}(v, u). \end{cases} \quad (12)$$

Note the identity

$$[\partial_\alpha u, \partial_\beta u] = \frac{1}{2}([\partial_\alpha u, \partial_\beta u] - [\partial_\beta u, \partial_\alpha u]) = \frac{1}{2}Q_{\alpha\beta}[u, u]. \quad (13)$$

Define

$$\mathcal{Q}[u, v] = -\frac{1}{2}\varepsilon^{ijk}\varepsilon_{klm}Q_{ij}[R^l u^m, v] - Q_{0i}[R^i u_0, v], \quad (14)$$

where ε_{ijk} is the antisymmetric symbol with $\varepsilon_{123} = 1$ and $R_i = \Lambda^{-1}\partial_i$ are the Riesz transforms.

Now we refer to Tesfahun [T], who showed that the system (6),(7) in Lorenz gauge can be written in the following form

$$\begin{aligned} \square A_\beta &= \mathcal{M}_\beta(A, \partial_t A, F, \partial_t F), \\ \square F_{\beta\gamma} &= \mathcal{N}_{\beta\gamma}(A, \partial_t A, F, \partial_t F), \end{aligned} \quad (15)$$

where

$$\begin{aligned} \mathcal{M}_\beta(A, \partial_t A, F, \partial_t F) &= -2\mathcal{Q}[\Lambda^{-1}A, A_\beta] + \sum_{i=1}^4 \Gamma_\beta^i(A, \partial A, F, \partial F) - 2[\Lambda^{-2}A^\alpha, \partial_\alpha A_\beta] \\ &\quad - [A^\alpha, [A_\alpha, A_\beta]], \end{aligned}$$

$$\begin{aligned} \mathcal{N}_{ij}(A, \partial_t A, F, \partial_t F) &= -2\mathcal{Q}[\Lambda^{-1}A, F_{ij}] + 2\mathcal{Q}[\Lambda^{-1}\partial_j A, A_i] - 2\mathcal{Q}[\Lambda^{-1}\partial_i A, A_j] \\ &\quad + 2Q_0[A_i, A_j] + Q_{ij}[A^\alpha, A_\alpha] - 2[\Lambda^{-2}A^\alpha, \partial_\alpha F_{ij}] \\ &\quad + 2[\Lambda^{-2}\partial_j A^\alpha, \partial_\alpha A_i] - 2[\Lambda^{-2}\partial_i A^\alpha, \partial_\alpha A_j] \\ &\quad - [A^\alpha, [A_\alpha, F_{ij}]] + 2[F_{\alpha i}, [A^\alpha, A_j]] - 2[F_{\alpha j}, [A^\alpha, A_i]] \\ &\quad - 2[[A^\alpha, A_i], [A_\alpha, A_j]], \end{aligned}$$

$$\begin{aligned} \mathcal{N}_{0i}(A, \partial_t A, F, \partial_t F) &= -2\mathcal{Q}[\Lambda^{-1}A, F_{0i}] + 2\mathcal{Q}[\Lambda^{-1}\partial_i A, A_0] - 2Q_{0j}[A^j, A_i] \\ &\quad + 2Q_0[A_0, A_i] + Q_{0i}[A^\alpha, A_\alpha] - 2[\Lambda^{-2}A^\alpha, \partial_\alpha F_{0i}] \\ &\quad + 2[\Lambda^{-2}\partial_i A^\alpha, \partial_\alpha A_0] - [A^\alpha, [A_\alpha, F_{0i}]] + 2[F_{\alpha 0}, [A^\alpha, A_i]] \\ &\quad - 2[F_{\alpha i}, [A^\alpha, A_0]] - 2[[A^\alpha, A_0], [A_\alpha, A_i]] \end{aligned}$$

where

$$\begin{aligned} \Gamma_\beta^1(A, \partial A, F, \partial F) &= -[A_0, \partial_\beta A_0] + [\Lambda^{-1}R_j(\partial_t A_0), \Lambda^{-1}R^j\partial_t(\partial_\beta A_0)], \\ \Gamma_\beta^2(A, \partial A, F, \partial F) &= -\frac{1}{2}\varepsilon^{ijk}\varepsilon_{klm}\left\{Q_{ij}[\Lambda^{-1}R^n\mathbf{A}_n, \Lambda^{-1}R^l\partial_\beta\mathbf{A}^m] \right. \\ &\quad \left. + Q_{ij}[\Lambda^{-1}R^n\partial_\beta\mathbf{A}_n, \Lambda^{-1}R^l\mathbf{A}^m]\right\}, \\ \Gamma_\beta^3(A, \partial A, F, \partial F) &= [\Lambda^{-2}\nabla \times \mathbf{F}, \Lambda^{-2}\nabla \times \partial_\beta\mathbf{F}] \\ &\quad - [\Lambda^{-2}\nabla \times \mathbf{F}, \Lambda^{-2}\partial_\beta\nabla \times (\mathbf{A} \times \mathbf{A})] \\ &\quad - [\Lambda^{-2}\nabla \times (\mathbf{A} \times \mathbf{A}), \Lambda^{-2}\nabla \times \partial_\beta\mathbf{F}] \\ &\quad + [\Lambda^{-2}\nabla \times (\mathbf{A} \times \mathbf{A}), \Lambda^{-2}\partial_\beta\nabla \times (\mathbf{A} \times \mathbf{A})], \\ \Gamma_\beta^4(A, \partial A, F, \partial F) &= [\mathbf{A}^{\text{cf}} + \mathbf{A}^{\text{df}}, \Lambda^{-2}\partial_\beta\mathbf{A}] + [\Lambda^{-2}\mathbf{A}, \partial_\beta\mathbf{A}]. \end{aligned} \quad (16)$$

Here $\mathbf{F} = (F_1, F_2, F_3)$, where $F_i = \sum_{j < k, j, k \neq i} F_{jk}$, $(\nabla \times \mathbf{A})_i = \varepsilon_{ijk}\partial^j A^k$ and $(\mathbf{A} \times \mathbf{B})_k = \varepsilon_{ijk}A^i B^j$.

Here especially the splitting of the spatial part $\mathbf{A} = (A_1, A_2, A_3)$ of the potential into divergence-free and curl-free parts and a smoother part is used

$$\mathbf{A} = \mathbf{A}^{\text{df}} + \mathbf{A}^{\text{cf}} + \Lambda^{-2}\mathbf{A}, \quad (17)$$

where

$$\begin{aligned}\mathbf{A}^{\text{df}} &= \Lambda^{-2} \nabla \times \nabla \times \mathbf{A}, \\ \mathbf{A}^{\text{cf}} &= -\Lambda^{-2} \nabla (\nabla \cdot \mathbf{A}).\end{aligned}$$

In a standard way we rewrite the system (15) as a first order (in t) system. Defining $A_{\pm} = \frac{1}{2}(A \pm (i\Lambda)^{-1} \partial_t A)$, $F_{\pm} = \frac{1}{2}(F \pm (i\Lambda)^{-1} \partial_t F)$, so that $A = A_+ + A_-$, $\partial_t A = i\Lambda(A_+ - A_-)$, $F = F_+ + F_-$, $\partial_t F = i\Lambda(F_+ - F_-)$ the system transforms to

$$(i\partial_t \pm \Lambda) A_{\pm}^{\beta} = -A^{\beta} \mp (2\Lambda)^{-1} \mathcal{M}_{\beta}(A, \partial_t A, F, \partial_t F), \quad (18)$$

$$(i\partial_t \pm \Lambda) F_{\pm}^{\beta\gamma} = -F^{\beta\gamma} \mp (2\Lambda)^{-1} \mathcal{N}_{\beta\gamma}(A, \partial_t A, F, \partial_t F). \quad (19)$$

The initial data transform to

$$A_{\pm}(0) = \frac{1}{2}(a \pm (i\Lambda)^{-1} \dot{a}) \in \widehat{H}^{s,r}, \quad F_{\mp}(0) = \frac{1}{2}(f \pm (i\Lambda)^{-1} \dot{f}) \in \widehat{H}^{l,r}.$$

Now, looking at the terms in \mathcal{M}_{β} and $\mathcal{N}_{\beta\gamma}$ and noting the fact that the Riesz transforms R_i are bounded in the spaces involved, the estimates in Theorem 3.1 reduce to proving:

1. the estimates for the null forms Q_{ij} , Q_0 and $Q \in \{Q_{0i}, Q_{ij}\}$:

$$\|Q[\Lambda^{-1}A, A]\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (20)$$

$$\|Q_{ij}[\Lambda^{-1}A, \Lambda^{-1}\partial A]\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (21)$$

$$\|Q[\Lambda^{-1}A, F]\|_{H_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s-1, b}^r} \|F\|_{X_{l, a}^r}, \quad (22)$$

$$\|Q[A, A]\|_{H_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (23)$$

$$\|Q_0[A, A]\|_{X_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (24)$$

the following estimate for Γ^1 and other bilinear terms

$$\|\Gamma^1(A, \partial A)\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (25)$$

$$\|\Pi(A, \Lambda^{-2}\partial A)\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (26)$$

$$\|\Pi(\Lambda^{-2}A, \partial A)\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (27)$$

$$\|\Pi(\Lambda^{-1}F, \Lambda^{-1}\partial F)\|_{H_{s-1, b-1+}^r} \lesssim \|F\|_{X_{l, a}^r} \|F\|_{X_{l, a}^r}, \quad (28)$$

$$\|\Pi(\Lambda^{-2}A, \partial F)\|_{H_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|F\|_{X_{l, a}^r}, \quad (29)$$

$$\|\Pi(\Lambda^{-1}A, \partial A)\|_{H_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (30)$$

and

2. the following trilinear and quadrilinear estimates:

$$\|\Pi(\Lambda^{-1}F, \Lambda^{-1}\partial(AA))\|_{H_{s-1, b-1+}^r} \lesssim \|F\|_{X_{l, a}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (31)$$

$$\|\Pi(\Lambda^{-1}\partial F, \Lambda^{-1}(AA))\|_{H_{s-1, b-1+}^r} \lesssim \|F\|_{X_{l, a}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (32)$$

$$\|\Pi(\Lambda^{-1}(AA), \Lambda^{-1}\partial(AA))\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (33)$$

$$\|\Pi(A, A, A)\|_{H_{s-1, b-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}, \quad (34)$$

$$\|\Pi(A, A, F)\|_{H_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r} \|F\|_{X_{l, a}^r}, \quad (35)$$

$$\|\Pi(A, A, A, A)\|_{H_{l-1, a-1+}^r} \lesssim \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r} \|A\|_{X_{s,b}^r}. \quad (36)$$

$\Pi(\dots)$ denotes a multilinear operator in its arguments and $\|u\|_{X_{s,b}^r} := \|u\|_{X_{s,b,-}^r} + \|u\|_{X_{s,b,+}^r}$.

The matrix commutator null forms are linear combinations of the ordinary ones, in view of (12). Since the matrix structure plays no role in the estimates under consideration, we reduce (20)–(24) to estimates of the ordinary null forms for \mathbb{C} -valued functions u and v (as in (11)).

Next we consider the term Γ_β^1 . We may ignore its matrix form and treat

$$\Gamma_k^1(A_0, \partial_k A_0) = -A_0(\partial_k A_0) + \Lambda^{-1} R_j(\partial_t A_0) \Lambda^{-1} R^j \partial_t(\partial_k A_0)$$

for $k = 1, 2, 3$ and

$$\begin{aligned} \Gamma_0^1(A_0, \partial^i A_i) &= -A_0(\partial_0 A_0) + \Lambda^{-1} R_j(\partial_t A_0) \Lambda^{-1} R^j \partial_t(\partial_0 A_0) \\ &= -A_0(\partial^i A_i) + \Lambda^{-1} R_j(\partial_t A_0) \Lambda^{-1} R^j \partial_t(\partial^i A_i), \end{aligned}$$

where we used the Lorenz gauge $\partial_0 A_0 = \partial^i A_i$ in the last line in order to eliminate one time derivative. Thus we have to consider

$$\Gamma^1(u, v) = -uv + \Lambda^{-1} R_j(\partial_t u) \Lambda^{-1} R^j(\partial_t v),$$

where $u = A_0$ and $v = \partial^i A_i$ or $v = \partial_k A_0$.

Next we show that Γ^1 also has a null structure. The proof of the following lemma was essentially given by Tesfahun [T]. In fact the detection of this null structure was the main progress of his paper over Selberg-Tesfahun [ST].

Lemma 4.1. *Let $q_{\mu\nu}(u, v) := Q_{\mu\nu}(D^{-1}u, D^{-1}v)$, $q_0(u, v) := Q_0(D^{-1}u, D^{-1}v)$. The following estimate holds:*

$$\Gamma^1(u, v) \lesssim \sum_{i,j} q_{ij}(u, v) + q_0(u, v) + (\Lambda^{-2}u)v + u(\Lambda^{-2}v). \quad (37)$$

Here $u \preceq v$ means $|\widehat{u}| \lesssim |\widehat{v}|$.

Proof. $\Gamma^1(u, v)$ has the symbol

$$p(\xi, \tau, \eta, \lambda) = -1 + \frac{\langle \xi, \eta \rangle \tau \lambda}{\langle \xi \rangle^2 \langle \eta \rangle^2} = \left(-1 + \frac{\langle \xi, \eta \rangle \langle \xi, \eta \rangle}{\langle \xi \rangle^2 \langle \eta \rangle^2} \right) + \frac{(\tau \lambda - \langle \xi, \eta \rangle) \langle \xi, \eta \rangle}{\langle \xi \rangle^2 \langle \eta \rangle^2} = I + II$$

Now we estimate

$$\begin{aligned} |I| &= \left| \frac{|\xi|^2 |\eta|^2 \cos^2 \angle(\xi, \eta)}{\langle \xi \rangle^2 \langle \eta \rangle^2} - 1 \right| \\ &\leq \left| \frac{\langle \xi \rangle^2 \langle \eta \rangle^2 \cos^2 \angle(\xi, \eta)}{\langle \xi \rangle^2 \langle \eta \rangle^2} - 1 \right| + \left| \frac{|\xi|^2 |\eta|^2 - \langle \xi \rangle^2 \langle \eta \rangle^2}{\langle \xi \rangle^2 \langle \eta \rangle^2} \right| \\ &= \sin^2 \angle(\xi, \eta) + \left| \frac{|\xi|^2 |\eta|^2 - \langle \xi \rangle^2 \langle \eta \rangle^2}{\langle \xi \rangle^2 \langle \eta \rangle^2} \right|, \end{aligned}$$

where $\angle(\xi, \eta)$ denotes the angle between ξ and η . We have

$$\left| \frac{|\xi|^2 |\eta|^2 - \langle \xi \rangle^2 \langle \eta \rangle^2}{\langle \xi \rangle^2 \langle \eta \rangle^2} \right| = \frac{|\xi|^2 + |\eta|^2 + 1}{\langle \xi \rangle^2 \langle \eta \rangle^2} \leq \frac{1}{\langle \xi \rangle^2} + \frac{1}{\langle \eta \rangle^2}$$

and

$$\sin \angle(\xi, \eta) = \frac{|\xi \times \eta|}{|\xi| |\eta|}.$$

Thus the operator belonging to the symbol I is controlled by $\sum_{i,j} q_{ij}(u, v) + (\Lambda^{-2}u)v + u(\Lambda^{-2}v)$. Moreover

$$|II| \leq \frac{|\tau \lambda - \langle \xi, \eta \rangle|}{\langle \xi \rangle \langle \eta \rangle} \leq |q_0(\xi, \eta)|.$$

Thus we obtain (37). \square

5. BILINEAR ESTIMATES

The proof of the following bilinear estimates relies on estimates given by Foschi and Klainerman [FK]. We first treat the case $r > 1$, but close to 1.

Lemma 5.1. *Assume $0 \leq \alpha_1, \alpha_2$, $\max(\alpha_1, \alpha_2) \neq \frac{2}{r} - \frac{1}{2}$, $\alpha_1 + \alpha_2 \geq \frac{2}{r}$ and $b > \frac{1}{r}$. The following estimate applies*

$$\|q_{ij}(u, v)\|_{H_{0,0}^r} \lesssim \|u\|_{X_{\alpha_1, b, \pm_1}^r} \|v\|_{X_{\alpha_2, b, \pm_2}^r}.$$

Proof. The left hand side equals

$$\|\mathcal{F}(q_{ij}(u, v))\|_{L_{\tau\xi}^{r'}} = \left\| \int q_{ij}(\eta, \eta - \xi) \widehat{u}(\lambda, \eta) \widehat{v}(\tau - \lambda, \xi - \eta) d\lambda d\eta \right\|_{L_{\tau\xi}^{r'}}.$$

Let now $u(t, x) = e^{\pm_1 i D} u_0^{\pm_1}(x)$, $v(t, x) = e^{\pm_2 i D} v_0^{\pm_2}(x)$, so that

$$\widehat{u}(\tau, \xi) = c\delta(\tau \mp_1 |\xi|) \widehat{u_0^{\pm_1}}(\xi), \quad \widehat{v}(\tau, \xi) = c\delta(\tau \mp_2 |\xi|) \widehat{v_0^{\pm_2}}(\xi).$$

This implies

$$\begin{aligned} & \|\mathcal{F}(q_{ij}(u, v))\|_{L_{\tau\xi}^{r'}} \\ &= c^2 \left\| \int q_{ij}(\eta, \eta - \xi) \widehat{u_0^{\pm_1}}(\eta) \widehat{v_0^{\pm_2}}(\xi - \eta) \delta(\lambda \mp_1 |\eta|) \delta(\tau - \lambda \mp_2 |\xi - \eta|) d\lambda d\eta \right\|_{L_{\tau\xi}^{r'}} \\ &= c^2 \left\| \int q_{ij}(\eta, \eta - \xi) \widehat{u_0^{\pm_1}}(\eta) \widehat{v_0^{\pm_2}}(\xi - \eta) \delta(\tau \mp_1 |\eta| \mp_2 |\xi - \eta|) d\lambda d\eta \right\|_{L_{\tau\xi}^{r'}}. \end{aligned}$$

By symmetry we only have to consider the elliptic case $\pm_1 = \pm_2 = +$ and the hyperbolic case $\pm_1 = +, \pm_2 = -$.

Elliptic case. We obtain by [FK], Lemma 13.2:

$$|q_{ij}(\eta, \xi - \eta)| \leq \frac{|\eta \times (\xi - \eta)|}{|\eta| |\xi - \eta|} \lesssim \frac{|\xi|^{\frac{1}{2}} (|\eta| + |\xi - \eta| - |\xi|)^{\frac{1}{2}}}{|\eta|^{\frac{1}{2}} |\xi - \eta|^{\frac{1}{2}}}.$$

By Hölder's inequality we obtain

$$\begin{aligned} & \|\mathcal{F}(q_{ij}(u, v))\|_{L_{\tau\xi}^{r'}} \\ & \lesssim \left\| \int \frac{|\xi|^{\frac{1}{2}} |\tau| - |\xi|^{\frac{1}{2}}}{|\eta|^{\frac{1}{2}} |\xi - \eta|^{\frac{1}{2}}} \delta(\tau - |\eta| - |\xi - \eta|) |\widehat{u_0^+}(\eta)| |\widehat{v_0^+}(\xi - \eta)| d\eta \right\|_{L_{\tau\xi}^{r'}} \\ & \lesssim \sup_{\tau, \xi} I \|\widehat{D^{\alpha_1} u_0^+}\|_{L^{r'}} \|D^{\alpha_2} v_0^+\|_{L^{r'}}, \end{aligned}$$

where

$$I = |\xi|^{\frac{1}{2}} |\tau| - |\xi|^{\frac{1}{2}} \left(\int \delta(\tau - |\eta| - |\xi - \eta|) |\eta|^{-(\alpha_1 + \frac{1}{2})r} |\xi - \eta|^{-(\alpha_2 + \frac{1}{2})r} d\eta \right)^{\frac{1}{r}}.$$

We want to prove $\sup_{\tau, \xi} I \lesssim 1$. By [FK], Lemma 4.3 we obtain

$$\int \delta(\tau - |\eta| - |\xi - \eta|) |\eta|^{-(\alpha_1 + \frac{1}{2})r} |\xi - \eta|^{-(\alpha_2 + \frac{1}{2})r} d\eta \sim \tau^A |\tau| - |\xi|^B,$$

where $A = \max((\alpha_1 + \frac{1}{2})r, 2) - (\alpha_1 + \alpha_2 + 1)r$ and $B = 2 - \max((\alpha_1 + \frac{1}{2})r, 2)$, if we assume without loss of generality $\alpha_1 \geq \alpha_2$ and $\max(\alpha_1, \alpha_2) \neq \frac{2}{r} - \frac{1}{2}$.

We assume from now on $\alpha_1 + \alpha_2 = \frac{2}{r}$ so that $\alpha_1, \alpha_2 \leq \frac{2}{r}$.

If $\alpha_1, \alpha_2 < \frac{2}{r} - \frac{1}{2}$, so that $A = 2 - (\alpha_1 + \alpha_2 + 1)r$ and $B = 0$, we obtain

$$I \lesssim |\xi|^{\frac{1}{2}} |\tau| - |\xi|^{\frac{1}{2}} \tau^{\frac{2}{r} - (\alpha_1 + \alpha_2 + 1)} \lesssim \tau^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1$$

using $|\xi| \leq |\tau|$.

If $\alpha_1 > \frac{2}{r} - \frac{1}{2}$, we obtain $A = -(\alpha_2 + \frac{1}{2})r$ and $B = 2 - (\alpha_1 + \frac{1}{2})r$, and therefore

$$\begin{aligned} I &\lesssim |\xi|^{\frac{1}{2}}|\tau| - |\xi|^{\frac{1}{2}}\tau^{-(\alpha_2 + \frac{1}{2})}|\tau| - |\xi|^{\frac{2}{r} - (\alpha_1 + \frac{1}{2})} \\ &\lesssim |\tau|^{\alpha_2}|\tau| - |\xi|^{\frac{2}{r} - \alpha_1} \lesssim \tau^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1, \end{aligned}$$

because $\alpha_1 + \alpha_2 = \frac{2}{r}$ and $\alpha_1 \leq \frac{2}{r}$.

Hyperbolic case. We start with the following bound (cf. [FK], Lemma 13.2):

$$|q_{ij}(\eta, \xi - \eta)| \leq \frac{|\eta \times (\xi - \eta)|}{|\eta| |\xi - \eta|} \lesssim \frac{|\xi|^{\frac{1}{2}} (|\xi| - |\eta| - |\eta - \xi|)^{\frac{1}{2}}}{|\eta|^{\frac{1}{2}} |\xi - \eta|^{\frac{1}{2}}},$$

so that similarly as in the elliptic case we have to estimate

$$I = |\xi|^{\frac{1}{2}}|\tau| - |\xi|^{\frac{1}{2}} \left(\int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-(\alpha_1 + \frac{1}{2})r} |\xi - \eta|^{-(\alpha_2 + \frac{1}{2})r} d\eta \right)^{\frac{1}{r}}.$$

In the subcase $|\eta| + |\xi - \eta| \leq 2|\xi|$ we apply [FK], Prop. 4.5 and obtain

$$\int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-(\alpha_1 + \frac{1}{2})r} |\xi - \eta|^{-(\alpha_2 + \frac{1}{2})r} d\eta \sim |\xi|^A |\xi| - |\tau|^B.$$

Assuming without loss of generality $\alpha_1 \geq \alpha_2$ we obtain $A = \max((\alpha_1 + \frac{1}{2})r, 2) - (\alpha_1 + \alpha_2 + 1)r$ and $B = 2 - \max((\alpha_1 + \frac{1}{2})r, 2)$, provided $\max(\alpha_1, \alpha_2) \neq \frac{2}{r} - \frac{1}{2}$.

If $\alpha_1 < \frac{2}{r} - \frac{1}{2}$, this means $A = 2 - (\alpha_1 + \alpha_2 + 1)r$ and $B = 0$, so that by $|\tau| \leq |\xi|$ this implies

$$I \lesssim |\xi|^{\frac{1}{2}}|\tau| - |\xi|^{\frac{1}{2}}|\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2 + 1)} \lesssim |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1.$$

If $\alpha_1 > \frac{2}{r} - \frac{1}{2}$, we obtain $A = -(\alpha_2 + \frac{1}{2})r$ and $B = 2 - (\alpha_1 + \frac{1}{2})r$, and thus

$$I \lesssim |\xi|^{-\alpha_2}|\xi| - |\tau|^{\frac{2}{r} - \alpha_1} \lesssim |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1,$$

because $|\tau| \leq |\xi|$, $\alpha_1 \leq \frac{2}{r}$ and $\alpha_1 + \alpha_2 = \frac{2}{r}$.

In the subcase $|\eta| + |\xi - \eta| \geq 2|\xi|$ we obtain by [FK], Lemma 4.4:

$$\begin{aligned} &\int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-(\alpha_1 + \frac{1}{2})r} |\xi - \eta|^{-(\alpha_2 + \frac{1}{2})r} d\eta \\ &\sim \int_2^\infty (|\xi|x + \tau)^{-(\alpha_1 + \frac{1}{2})r+1} (|\xi|x - \tau)^{-(\alpha_2 + \frac{1}{2})r+1} dx \\ &\sim \int_2^\infty \left(x + \frac{\tau}{|\xi|}\right)^{-(\alpha_1 + \frac{1}{2})r+1} \left(x - \frac{\tau}{|\xi|}\right)^{-(\alpha_2 + \frac{1}{2})r+1} dx \cdot |\xi|^{-(\alpha_1 + \alpha_2 + 1)r+2}. \end{aligned}$$

We remark that in fact the lower limit of the integral can be chosen as 2 by inspection of the proof in [FK]. The integral converges, because $|\tau| \leq |\xi|$ and $(\alpha_1 + \alpha_2 + 1)r - 2 > 1$ assuming that $\alpha_1 + \alpha_2 = \frac{2}{r}$ and $r > 1$. This implies the bound

$$I \lesssim |\xi|^{\frac{1}{2}}|\tau| - |\xi|^{\frac{1}{2}}|\xi|^{-(\alpha_1 + \alpha_2 + 1) + \frac{2}{r}} \lesssim |\xi|^{-(\alpha_1 + \alpha_2) + \frac{2}{r}} = 1.$$

Summarizing we obtain

$$\|q_{ij}(u, v)\|_{X_{0,0}^r} \lesssim \|D^{\alpha_1} u_0^{\pm 1}\|_{L^{r'}} \|D^{\alpha_2} v_0^{\pm 2}\|_{L^{r'}}.$$

By the transfer principle Prop. 3.1 we obtain the claimed result first in the case $\alpha_1 + \alpha_2 = \frac{2}{r}$, but then trivially also in the case $\alpha_1 + \alpha_2 \geq \frac{2}{r}$, because we consider inhomogeneous spaces. The assumption $\alpha_1, \alpha_2 \leq \frac{2}{r}$ is therefore also redundant. \square

In a similar manner we can also estimate the nullform $q_{0j}(u, v)$.

Lemma 5.2. *Assume $0 \leq \alpha_1, \alpha_2$, $\max(\alpha_1, \alpha_2) \neq \frac{2}{r} - \frac{1}{2}$, $\max(\alpha_1 + \frac{1}{2}, \alpha_2) \neq \frac{2}{r}$, $\max(\alpha_2 + \frac{1}{2}, \alpha_1) \neq \frac{2}{r}$, $\alpha_1 + \alpha_2 \geq \frac{2}{r}$ and $b > \frac{1}{r}$. The following estimate applies*

$$\|q_{0j}(u, v)\|_{H_{0,0}^r} \lesssim \|u\|_{X_{\alpha_1, b, \pm 1}^r} \|v\|_{X_{\alpha_2, b, \pm 2}^r}.$$

Proof. Arguing as in the proof of Lemma 5.1 we use in the elliptic case the estimate (cf. [FK], Lemma 13.2):

$$|q_{0j}(\eta, \xi - \eta)| \lesssim \frac{(|\eta| + |\xi - \eta| - |\xi|)^{\frac{1}{2}}}{\min(|\eta|^{\frac{1}{2}}, |\xi - \eta|^{\frac{1}{2}})}.$$

Assuming w.l.o.g. $|\eta| \leq |\xi - \eta|$ we have to estimate

$$\begin{aligned} I &= \|\tau\| - |\xi|^{\frac{1}{2}} \left(\int \delta(\tau - |\eta| - |\xi - \eta|) |\eta|^{-(\alpha_1 + \frac{1}{2})r} |\xi - \eta|^{-\alpha_2 r} d\eta \right)^{\frac{1}{r}} \\ &\sim \|\tau\| - |\xi|^{\frac{1}{2}} |\tau|^{\frac{A}{r}} \|\tau\| - |\xi|^{\frac{B}{r}}, \end{aligned}$$

where $A = \max((\alpha_1 + \frac{1}{2})r, \alpha_2 r, 2) - (\alpha_1 + \alpha_2 + \frac{1}{2})r$ and $B = 2 - \max((\alpha_1 + \frac{1}{2})r, \alpha_2 r, 2)$, provided $\max(\alpha_1 + \frac{1}{2}, \alpha_2) \neq \frac{2}{r}$.

Assume now that $\alpha_1 + \alpha_2 = \frac{2}{r}$ and $\alpha_1, \alpha_2 < \frac{2}{r}$.

If $\alpha_1 < \frac{2}{r} - \frac{1}{2}$ and $\alpha_2 < \frac{2}{r}$, we obtain $A = 2 - (\alpha_1 + \alpha_2 + \frac{1}{2})r$ and $B = 0$, thus using $|\tau| \geq |\xi|$:

$$I \sim \|\tau\| - |\xi|^{\frac{1}{2}} |\tau|^{\frac{2}{r} - (\alpha_1 + \alpha_2 + \frac{1}{2})r} \lesssim |\tau|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1.$$

If $\alpha_1 > \frac{2}{r} - \frac{1}{2}$ we obtain $A = -\alpha_2 r$ and $B = 2 - (\alpha_1 + \frac{1}{2})r$, thus:

$$I \sim \|\tau\| - |\xi|^{\frac{1}{2}} |\tau|^{\alpha_2} \|\tau\| - |\xi|^{\frac{2}{r} - (\alpha_1 + \frac{1}{2})r} \lesssim |\tau|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1.$$

In the hyperbolic case we obtain by [FK], Lemma 13.2:

$$|q_{0j}(\eta, \xi - \eta)| \lesssim |\xi|^{\frac{1}{2}} \frac{(|\xi| - |\eta| - |\eta - \xi|)^{\frac{1}{2}}}{|\eta|^{\frac{1}{2}} |\xi - \eta|^{\frac{1}{2}}}$$

and argue exactly as in the proof of Lemma 5.1. The proof is completed as before. \square

We also need the same result for $q_0(u, v)$.

Lemma 5.3. *Assume $0 \leq \alpha_1, \alpha_2$, $\max(\alpha_1, \alpha_2) \neq \frac{2}{r} - 1$, $\max(\alpha_1 + 1, \alpha_2) \neq \frac{2}{r}$, $\max(\alpha_2 + 1, \alpha_1) \neq \frac{2}{r}$, $\alpha_1 + \alpha_2 \geq \frac{2}{r}$ and $b > \frac{1}{r}$. The following estimate applies*

$$\|q_0(u, v)\|_{H_{0,0}^r} \lesssim \|u\|_{X_{\alpha_1, b, \pm 1}^r} \|v\|_{X_{\alpha_2, b, \pm 2}^r}.$$

Proof. Arguing as before we use in the elliptic case the estimate (cf. [FK], Lemma 13.2):

$$|q_0(\eta, \xi - \eta)| \lesssim \frac{(|\eta| + |\xi - \eta| - |\xi|)}{\min(|\eta|, |\xi - \eta|)}.$$

Assuming w.l.o.g. $|\eta| \leq |\xi - \eta|$ we have to estimate

$$\begin{aligned} I &= \|\tau\| - |\xi| \left(\int \delta(\tau - |\eta| - |\xi - \eta|) |\eta|^{-(\alpha_1 + 1)r} |\xi - \eta|^{-\alpha_2 r} d\eta \right)^{\frac{1}{r}} \\ &\sim \|\tau\| - |\xi| |\tau|^{\frac{A}{r}} \|\tau\| - |\xi|^{\frac{B}{r}}, \end{aligned}$$

where $A = \max((\alpha_1 + 1)r, \alpha_2 r, 2) - (\alpha_1 + \alpha_2 + 1)r$ and $B = 2 - \max((\alpha_1 + 1)r, \alpha_2 r, 2)$, provided $\max(\alpha_1 + 1, \alpha_2) \neq \frac{2}{r}$. Assume from now on $\alpha_1 + \alpha_2 = \frac{2}{r}$ and $\alpha_1, \alpha_2 < \frac{2}{r}$.

If $\alpha_1 < \frac{2}{r} - 1$ and $\alpha_2 < \frac{2}{r}$, we obtain:

$$I \sim \|\tau\| - |\xi| |\tau|^{\frac{2}{r} - (\alpha_1 + \alpha_2 + 1)r} \lesssim |\tau|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1.$$

If $\alpha_1 > \frac{2}{r} - 1$ we obtain

$$I \sim \|\tau\| - |\xi| \|\tau\|^{-\alpha_2} \|\tau\| - |\xi|^{\frac{2}{r} - (\alpha_1 + 1)r} \lesssim |\tau|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1,$$

using $\alpha_1 + \alpha_2 = \frac{2}{r}$ and $\alpha_1 \leq \frac{2}{r}$.

In the hyperbolic case we obtain by [FK], Lemma 13.2:

$$|q_{0j}(\eta, \xi - \eta)| \lesssim |\xi| \frac{|\xi| - \|\eta\| - |\eta - \xi|}{|\eta| |\xi - \eta|}.$$

In the subcase $|\eta| + |\xi - \eta| \leq 2|\xi|$ we apply [FK], Prop. 4.5 and obtain

$$\int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-(\alpha_1+1)r} |\xi - \eta|^{-(\alpha_2+1)r} d\eta \sim |\xi|^A \|\xi\| - |\tau| |^B.$$

Assuming without loss of generality $\alpha_1 \geq \alpha_2$ we obtain $A = \max((\alpha_1 + 1)r, 2) - (\alpha_1 + \alpha_2 + 2)r$ and $B = 2 - \max((\alpha_1 + 1)r, 2)$, provided $\max(\alpha_1, \alpha_2) \neq \frac{2}{r} - 1$.

If $\alpha_1 < \frac{2}{r} - 1$, this implies

$$I \lesssim |\xi| \|\tau\| - |\xi| \|\xi\|^{\frac{2}{r} - (\alpha_1 + \alpha_2 + 2)} \lesssim |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1.$$

If $\alpha_1 > \frac{2}{r} - 1$, we obtain

$$I \lesssim |\xi| \|\tau\| - |\xi| \|\xi\|^{-(\alpha_2+1)} \|\xi\| - |\tau| \|\xi\|^{\frac{2}{r} - (\alpha_1+1)} \lesssim |\xi|^{-\alpha_2} \|\xi\| - |\tau| \|\xi\|^{\frac{2}{r} - \alpha_1} \lesssim |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2)} = 1,$$

because $|\tau| \leq |\xi|$, $\alpha_1 \leq \frac{2}{r}$ and $\alpha_1 + \alpha_2 = \frac{2}{r}$.

In the subcase $|\eta| + |\xi - \eta| \geq 2|\xi|$ we obtain by [FK], Lemma 4.4:

$$\begin{aligned} & \int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-(\alpha_1+1)r} |\xi - \eta|^{-(\alpha_2+1)r} d\eta \\ & \sim \int_2^\infty (x + \frac{\tau}{|\xi|})^{-(\alpha_1+1)r+1} (x - \frac{\tau}{|\xi|})^{-(\alpha_2+1)r+1} dx \cdot |\xi|^{-(\alpha_1 + \alpha_2 + 2)r + 2}. \end{aligned}$$

The integral converges, because $|\tau| \leq |\xi|$ and $(\alpha_1 + \alpha_2 + 2)r - 2 > 1$, because $\alpha_1 + \alpha_2 = \frac{2}{r}$ and $r > 1$. This implies the bound

$$I \lesssim |\xi| \|\tau\| - |\xi| \|\xi\|^{-(\alpha_1 + \alpha_2 + 2) + \frac{2}{r}} \lesssim |\xi|^{-(\alpha_1 + \alpha_2) + \frac{2}{r}} = 1.$$

The proof is completed as in the proof of Lemma 5.1, where the assumption $\alpha_1, \alpha_2 < \frac{2}{r}$ is redundant. \square

Lemma 5.4. *Assume $0 \leq \alpha_0 \leq \alpha_1, \alpha_2$, $\alpha_1, \alpha_2 \neq \frac{2}{r}$, $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r}$, $\alpha_1 + \alpha_2 > \frac{3}{r}$, $b > \frac{1}{r}$. Then the following estimate applies:*

$$\|uv\|_{\dot{H}_{\alpha_0, 0}^r} \lesssim \|u\|_{\dot{X}_{\alpha_1, b, \pm 1}^r} \|v\|_{\dot{X}_{\alpha_2, b, \pm 2}^r}.$$

Proof. By similar calculations as before we have to estimate in the elliptic case:

$$\begin{aligned} I &= |\xi|^{\alpha_0} \left(\int \delta(\tau - |\eta| - |\xi - \eta|) |\eta|^{-\alpha_1 r} |\xi - \eta|^{-\alpha_2 r} d\eta \right)^{\frac{1}{r}} \\ &\sim |\xi|^{\alpha_0} \tau^{\frac{A}{r}} \|\tau\| - |\xi| \|\xi\|^{\frac{B}{r}} \lesssim \tau^{\alpha_0 + \frac{2}{r} - (\alpha_1 + \alpha_2)} = 1, \end{aligned}$$

if we assume from now on $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r}$ and $\alpha_1, \alpha_2 < \frac{2}{r}$, because by [FK], Lemma 4.3 we obtain $A = \max(\alpha_1 r, \alpha_2 r, 2) - (\alpha_1 + \alpha_2)r = 2 - (\alpha_1 + \alpha_2)r$ and $B = 2 - \max(\alpha_1 r, \alpha_2 r, 2) = 0$.

In the hyperbolic case we obtain in the subcase $|\eta| + |\xi - \eta| \leq 2|\xi|$

$$\begin{aligned} I &= \left(\int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-\alpha_1 r} |\xi - \eta|^{-\alpha_2 r} d\eta \right)^{\frac{1}{r}} |\xi|^{\alpha_0} \\ &\sim |\xi|^{\frac{A}{r}} \|\xi\| - |\tau| \|\xi\|^{\frac{B}{r}} |\xi|^{\alpha_0} \lesssim |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2) + \alpha_0} = 1, \end{aligned}$$

where we applied [FK], Prop. 4.5 with $A = \max(\alpha_1 r, 2) - (\alpha_1 + \alpha_2)r = 2 - (\alpha_1 + \alpha_2)r$ and $B = 2 - \max(\alpha_1 r, 2) = 0$ provided $\alpha_1 < \frac{2}{r}$ and $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r}$ (or similarly

with α_1 and α_2 interchanged). In the subcase $|\eta| + |\xi - \eta| \geq 2|\xi|$ we obtain by [FK], Lemma 4.4 the estimate

$$\begin{aligned} I &\sim \left(\int_2^\infty (|\xi|x + \tau)^{-\alpha_1 r + 1} (|\xi|x - \tau)^{-\alpha_2 r + 1} dx \right)^{\frac{1}{r}} |\xi|^{\alpha_0} \\ &\lesssim \left(\int_2^\infty \left(x + \frac{\tau}{|\xi|}\right)^{-\alpha_1 r + 1} \left(x - \frac{\tau}{|\xi|}\right)^{-\alpha_2 r + 1} dx \right)^{\frac{1}{r}} \cdot |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2) + \alpha_0} \lesssim 1. \end{aligned}$$

Because $|\tau| \leq |\xi|$ the integral converges provided $\alpha_1 + \alpha_2 > \frac{3}{r}$. Moreover we used again $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r}$.

By the transfer principle the claimed estimate results as before. \square

Lemma 5.5. *If $\alpha_1 + \alpha_2 > \frac{3}{r}$ and $b_1 + b_2 > \frac{1}{r}$ the following estimate applies:*

$$\|uv\|_{H_{0,0}^r} \lesssim \|u\|_{H_{\alpha_1, b_1}^r} \|v\|_{H_{\alpha_2, b_2}^r}.$$

Proof. By Young's and Hölder's inequalities we obtain

$$\begin{aligned} \|uv\|_{X_{0,0}^r} &= \|\widehat{uv}\|_{L_{\tau\xi}^{r'}} \lesssim \|\widehat{u}\|_{L_\xi^{p_1} L_\tau^{q_1}} \|\widehat{v}\|_{L_\xi^{p_2} L_\tau^{q_2}} \\ &\lesssim \|\langle \xi \rangle^{-\alpha_1} \langle |\tau| - |\xi| \rangle^{-b_1}\|_{L_\xi^{s_1} L_\tau^{r_1}} \|\langle \xi \rangle^{\alpha_1} \langle |\tau| - |\xi| \rangle^{b_1} \widehat{u}\|_{L_{\xi\tau}^{r'}} \\ &\quad \cdot \|\langle \xi \rangle^{-\alpha_2} \langle |\tau| - |\xi| \rangle^{-b_2}\|_{L_\xi^{s_2} L_\tau^{r_2}} \|\langle \xi \rangle^{\alpha_2} \langle |\tau| - |\xi| \rangle^{b_2} \widehat{v}\|_{L_{\xi\tau}^{r'}} \\ &\lesssim \|u\|_{H_{\alpha_1, b_1}^r} \|v\|_{H_{\alpha_2, b_2}^r}. \end{aligned}$$

Here $1 + \frac{1}{r'} = \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{q_1} + \frac{1}{q_2}$ and $\frac{1}{q_j} = \frac{1}{r_j} + \frac{1}{r'}$, $\frac{1}{p_j} = \frac{1}{s_j} + \frac{1}{r'}$. This implies $1 + \frac{1}{r'} = \frac{1}{s_1} + \frac{1}{s_2} + \frac{2}{r'} = \frac{1}{r_1} + \frac{1}{r_2} + \frac{2}{r'}$. For the last estimate we need $r_j b_j > 1$ and $s_j \alpha_j > 3$. This can be fulfilled, if $1 + \frac{1}{r'} < \frac{\alpha_1 + \alpha_2}{3} + \frac{2}{r'} \Leftrightarrow \alpha_1 + \alpha_2 > \frac{3}{r}$ and $1 + \frac{1}{r'} < b_1 + b_2 + \frac{2}{r'} \Leftrightarrow b_1 + b_2 > \frac{1}{r}$. \square

An immediate consequence of Lemma 5.4 and Lemma 5.5 is the following corollary.

Corollary 5.1. *Let $0 \leq \alpha_0 \leq \alpha_1, \alpha_2$, $\alpha_1, \alpha_2 \neq \frac{2}{r}$, $\alpha_1 + \alpha_2 - \alpha_0 \geq \frac{2}{r}$, $\alpha_1 + \alpha_2 > \frac{3}{r}$ and $b > \frac{1}{r}$. Then the following estimate applies:*

$$\|uv\|_{H_{\alpha_0, 0}^r} \lesssim \|u\|_{X_{\alpha_1, b, \pm 1}^r} \|v\|_{X_{\alpha_2, b, \pm 2}^r}.$$

Lemma 5.6. *Let $0 \leq \alpha_0 \leq \alpha_1, \alpha_2$, $\alpha_1, \alpha_2 \neq \frac{2}{r}$, $\alpha_1 + \alpha_2 - \alpha_0 \geq \frac{2}{r} + b$, $\alpha_1 + \alpha_2 > \frac{3}{r}$. Then the following estimate applies:*

$$\|uv\|_{H_{\alpha_0, b}^r} \lesssim \|u\|_{X_{\alpha_1, b}^r} \|v\|_{X_{\alpha_2, b}^r}.$$

Proof. We apply the "hyperbolic Leibniz rule" (cf. [AFS], p. 41):

$$|\tau| - |\xi| \lesssim |\rho| - |\eta| + |\tau - \rho| - |\xi - \eta| + b_\pm(\xi, \eta), \quad (38)$$

where

$$b_+(\xi, \eta) = |\eta| + |\xi - \eta| - |\xi|, \quad b_-(\xi, \eta) = |\xi| - |\eta| - |\xi - \eta|.$$

Let us first consider the term $b_\pm(\xi, \eta)$ in (38). Decomposing as before $uv = u_+ v_+ + u_+ v_- + u_- v_+ + u_- v_-$, where $u_\pm(t) = e^{\pm itD} f$, $v_\pm(t) = e^{\pm itD} g$, we use

$$\widehat{u}_\pm(\tau, \xi) = c\delta(\tau \mp |\xi|)\widehat{f}(\xi), \quad \widehat{v}_\pm(\tau, \xi) = c\delta(\tau \mp |\xi|)\widehat{g}(\xi)$$

and have to estimate

$$\begin{aligned} & \left\| \int |\xi|^{\alpha_0} b_{\pm}^b(\xi, \eta) \delta(\tau - |\eta| \mp |\xi - \eta|) \widehat{f}(\xi) \widehat{g}(\xi - \eta) d\eta \right\|_{L_{\tau\xi}^{r'}} \\ &= \left\| \int |\xi|^{\alpha_0} |\tau| - |\xi|^b \delta(\tau - |\eta| \mp |\xi - \eta|) \widehat{f}(\xi) \widehat{g}(\xi - \eta) d\eta \right\|_{L_{\tau\xi}^{r'}} \\ &\lesssim \sup_{\tau, \xi} I \|\widehat{D^{\alpha_1} f}\|_{L^{r'}} \|\widehat{D^{\alpha_2} g}\|_{L^{r'}}. \end{aligned}$$

Here we used Hölder's inequality, where

$$I = \|\tau| - |\xi|^b |\xi|^{\alpha_0} \left(\int \delta(\tau - |\eta| \mp |\xi - \eta|) |\eta|^{-\alpha_1 r} |\xi - \eta|^{-\alpha_2 r} d\eta \right)^{\frac{1}{r}}.$$

We assume for the time being $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r} + b$ and $\max(\alpha_1, \alpha_2) \leq \frac{2}{r} + b$. In order to obtain $I \lesssim 1$ we first consider the elliptic case $\pm_1 = \pm_2 = +$ and use [FK], Prop. 4.3. This gives

$$I \sim \|\tau| - |\xi|^b |\xi|^{\alpha_0} \tau^{\frac{A}{r}} \|\tau| - |\xi|^{\frac{B}{r}}$$

with $A = \max(\alpha_1 r, \alpha_2 r, 2) - (\alpha_1 + \alpha_2)r$ and $B = 2 - \max(\alpha_1 r, \alpha_2 r, 2)$ for $\max(\alpha_1, \alpha_2) \neq \frac{2}{r}$. Assume w.l.o.g. $\alpha_1 \geq \alpha_2$. If $\alpha_1 < \frac{2}{r}$ we obtain $A = 2 - (\alpha_1 + \alpha_2)r$ and $B = 0$, so that

$$I \sim \|\tau| - |\xi|^b |\xi|^{\alpha_0} \tau^{\frac{2}{r} - (\alpha_1 + \alpha_2)r} \lesssim \tau^{b + \alpha_0 - (\alpha_1 + \alpha_2)r + \frac{2}{r}} = 1$$

using $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r} + b$ and $|\xi| \leq \tau$. If $\alpha_1 > \frac{2}{r}$ we obtain $A = -\alpha_2 r$ and $b = 2 - \alpha_1 r$, which implies

$$I \sim \|\tau| - |\xi|^b |\xi|^{\alpha_0} \tau^{-\alpha_2 r} \|\tau| - |\xi|^{\frac{2}{r} - \alpha_1 r} \lesssim \tau^{b + \alpha_0 - (\alpha_1 + \alpha_2)r + \frac{2}{r}} = 1$$

as before, if $\alpha_1 \leq \frac{2}{r} + b$.

Next we consider the hyperbolic case $\pm_1 = +, \pm_2 = -$.

First we assume $|\eta| + |\xi - \eta| \leq 2|\xi|$ and use [FK], Prop. 4.5 which gives

$$\int \delta(\tau - |\eta| + |\xi - \eta|) |\eta|^{-\alpha_1 r} |\xi - \eta|^{\alpha_2 r} d\eta \sim |\xi|^A \|\xi| - |\tau|\|^B,$$

where $A = \max(\alpha_2 r, 2) - (\alpha_1 + \alpha_2)r$ and $B = 2 - \max(\alpha_2 r, 2)$, if $\alpha_2 \neq \frac{2}{r}$. If $\alpha_2 < \frac{2}{r}$ we obtain $A = 2 - (\alpha_1 + \alpha_2)r$ and $B = 0$, so that by $|\tau| \leq |\xi|$:

$$I \lesssim |\xi|^{\alpha_0} \|\tau| - |\xi|^b |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2)r} \lesssim |\xi|^{b + \frac{2}{r} + \alpha_0 - (\alpha_1 + \alpha_2)r} = 1,$$

if $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r} + b$. If $\alpha_2 > \frac{2}{r}$ we obtain $A = -\alpha_1 r$ and $B = 2 - \alpha_2 r$ so that

$$I \lesssim |\xi|^{\alpha_0} \|\tau| - |\xi|^b |\xi|^{-\alpha_1 r} \|\xi| - |\tau|\|^{\frac{2}{r} - \alpha_2 r} \lesssim |\xi|^{\alpha_0 + b - \alpha_1 r + \frac{2}{r} - \alpha_2 r} = 1,$$

if $\alpha_2 \leq \frac{2}{r} + b$. Similarly we estimate for interchanged roles of α_1 and α_2 .

Next we assume $|\eta| + |\xi - \eta| \geq 2|\xi|$, use [FK], Lemma 4.4 and obtain

$$\begin{aligned} & \int \delta(\tau - |\eta| - |\xi - \eta|) |\eta|^{-\alpha_1 r} |\xi - \eta|^{-\alpha_2 r} d\eta \\ &\sim \int_2^\infty (|\xi|x + \tau)^{-\alpha_1 r + 1} (|\xi|x - \tau)^{-\alpha_2 r + 1} dx \\ &\sim \int_2^\infty \left(x + \frac{\tau}{|\xi|}\right)^{-\alpha_1 r + 1} \left(x - \frac{\tau}{|\xi|}\right)^{-\alpha_2 r + 1} dx \cdot |\xi|^{2 - (\alpha_1 + \alpha_2)r}. \end{aligned}$$

This integral converges, because $\tau \leq |\xi|$ and by our assumption $\alpha_1 + \alpha_2 > \frac{3}{r}$. This implies

$$I \lesssim |\xi|^{\alpha_0} \|\tau| - |\xi|^b |\xi|^{\frac{2}{r} - (\alpha_1 + \alpha_2)r} \lesssim |\xi|^{\alpha_0 + b + \frac{2}{r} - (\alpha_1 + \alpha_2)r} = 1,$$

if $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r} + b$.

By the transfer principle we obtain

$$\|D^{\alpha_0} B_{\pm}^b(u, v)\|_{X_{0,0}^r} \lesssim \|u\|_{X_{\alpha_1, b}^r} \|v\|_{X_{\alpha_2, b}^r}$$

provided our assumptions are fulfilled, where the upper bounds for α_1 and α_2 are redundant, and the condition $\alpha_1 + \alpha_2 - \alpha_0 = \frac{2}{r} + b$ may be replaced by $\alpha_1 + \alpha_2 - \alpha_0 \geq \frac{2}{r} + b$, because we consider inhomogeneous spaces. Here B_{\pm}^b denotes the operator with Fourier symbol b_{\pm} .

Consider now the term $\|\rho\| - \|\eta\|$ (or similarly $\|\tau - \rho\| - \|\xi - \eta\|$) in (38). We have to prove

$$\|D^{\alpha_0}((D_-^b u)v)\|_{X_{0,0}^r} \lesssim \|u\|_{X_{\alpha_1,b}^r} \|v\|_{X_{\alpha_2,b}^r},$$

which is implied by

$$\|D^{\alpha_0}(uv)\|_{X_{0,0}^r} \lesssim \|u\|_{X_{\alpha_1,0}^r} \|v\|_{X_{\alpha_2,b}^r}.$$

After application of the fractional Leibniz rule this results from Lemma 5.5, because $\alpha_1 + \alpha_2 - \alpha_0 \geq \frac{2}{r} + b > \frac{3}{r}$, which completes the proof. \square

6. PROOF OF (20) - (36) AND THEOREM 2.1:

Proof of (20) - (36). We prove the estimates first for the case $r = 1+$ by the results of chapter 5. We assume $2 \geq s \geq 1 + \frac{1}{r}$, $l = 1$ and $b = a = \frac{1}{r} +$. Later we interpolate with the case $r = 2$ given by Tesfahun [T].

Proof of (20) and (21): This reduces to

$$\|q(u, v)\|_{H_{s-1,0}^r} \lesssim \|u\|_{X_{s,b}^r} \|u\|_{X_{s-1,b}^r}.$$

By the fractional Leibniz rule this results from Lemma 5.1 or Lemma 5.2 for $s > \frac{2}{r}$.

Proof of (22): This reduces to

$$\|q(u, v)\|_{H_{l-1,0}^r} \lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{l-1,b}^r}.$$

If $l = 1$ and $s > \frac{2}{r}$ this results from Lemma 5.1 or Lemma 5.2.

Proof of (23) and (24): We have to show

$$\|q(u, v)\|_{H_{0,0}^r} \lesssim \|u\|_{X_{s-1,b}^r} \|u\|_{X_{s-1,b}^r}.$$

This is implied by Lemma 5.1 or Lemma 5.2, if $s \geq 1 + \frac{1}{r}$, which is fulfilled.

Proof of (25): By Lemma 4.1 we have to prove

$$\|q_{ij}(u, \Lambda v)\|_{X_{s-1,0}^r} \lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{s,b}^r}$$

which by the fractional Leibniz rule results from Lemma 5.1. Moreover we need

$$\|q_0(u, \Lambda v)\|_{X_{s-1,0}^r} \lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{s,b}^r}$$

which is given by Lemma 5.3. Finally

$$\|(\Lambda^{-2}u)v\|_{X_{s-1,0}^r} + \|u(\Lambda^{-2}v)\|_{X_{s-1,0}^r} \lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{s-1,b}^r}$$

by Lemma 5.5 for $s + 2 > \frac{3}{r}$, which is fulfilled.

Proof of (26) and (27): The estimates result from Lemma 5.5, because $s + 2 > \frac{3}{r}$.

Proof of (28): We reduce to

$$\|uv\|_{H_{s-1,0}^r} \lesssim \|u\|_{X_{l+1,b}^r} \|v\|_{X_{l,b}^r}.$$

This is implied by Cor. 5.1 with parameters $\alpha_0 = s - 1$, $\alpha_1 = l + 1 = 2$, $\alpha_2 = l = 1$, so that $\alpha_1 + \alpha_2 = 3 > \frac{3}{r}$ and $\alpha_1 + \alpha_2 - \alpha_0 = 4 - s \geq \frac{2}{r}$ for $s \leq 2$.

Proof of (29): The estimate reduces to

$$\|uv\|_{H_{l-1,0}^r} \lesssim \|u\|_{X_{s+2,b}^r} \|v\|_{X_{l-1,b}^r},$$

which results from Lemma 5.5, because $s + 2 > \frac{3}{r}$.

Proof of (30): We reduce to

$$\|uv\|_{H_{l-1,0}^r} \lesssim \|u\|_{X_{s+1,b}^r} \|v\|_{X_{s-1,b}^r},$$

which results from Lemma 5.5, because $2s > \frac{3}{r}$.

Proof of (31): The estimate

$$\|uvw\|_{H_{s-1,0}^r} \lesssim \|u\|_{X_{l+1,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r}$$

reduces by the fractional Leibniz rule to the estimates

$$\|uvw\|_{X_{0,0}^r} \lesssim \|u\|_{X_{l-s+2,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r}$$

and

$$\|uvw\|_{X_{0,0}^r} \lesssim \|u\|_{X_{l+1,b}^r} \|v\|_{X_{1,b}^r} \|w\|_{X_{s,b}^r}.$$

Now we obtain by Lemma 5.5 :

$$\begin{aligned} \|uvw\|_{X_{0,0}^r} &\lesssim \|u\|_{X_{l-s+2,b}^r} \|vw\|_{X_{s-l-2+\frac{3}{r},0}^r} \\ &\lesssim \|u\|_{X_{l-s+2,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r}. \end{aligned}$$

The last estimate results from Cor. 5.1, if $2s - (s - l - 2 + \frac{3}{r}) > \frac{2}{r} \Leftrightarrow s > \frac{5}{r} - l - 2 = \frac{5}{r} - 3$ and $2s > \frac{3}{r}$, which is fulfilled for $s > \frac{2}{r}$. Moreover in exactly the same way we obtain

$$\begin{aligned} \|uvw\|_{X_{0,0}^r} &\lesssim \|u\|_{X_{l+1,b}^r} \|vw\|_{X_{\frac{3}{r}-l-1+0}^r} \\ &\lesssim \|u\|_{X_{l+1,b}^r} \|v\|_{X_{1,b}^r} \|w\|_{X_{s,b}^r}. \end{aligned}$$

Proof of (32): We apply Lemma 5.5 and Cor. 5.1 which implies

$$\|u\Lambda^{-1}(vw)\|_{H_{s-1,0}^r} \lesssim \|u\|_{X_{l,b}^r} \|\Lambda^{-1}(vw)\|_{X_{\frac{3}{r}+s-1-l+0}^r} \lesssim \|u\|_{X_{l,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r}$$

for $2s - \frac{3}{r} - s + 1 + l + 1 > \frac{2}{r} \Leftrightarrow s > \frac{5}{r} - 3$, which is fulfilled for $s > \frac{2}{r}$.

Proof of (34): Lemma 5.5 and Cor. 5.1 imply

$$\|uvw\|_{H_{s-1,0}^r} \lesssim \|u\|_{X_{s,b}^r} \|vw\|_{X_{\frac{3}{r}-1+0}^r} \lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r},$$

because $2s - \frac{3}{r} + 1 > \frac{2}{r}$ for $s > \frac{2}{r}$, and $2s > \frac{3}{r}$.

Proof of (35): Similarly we obtain

$$\|uvw\|_{H_{l-1,0}^r} \lesssim \|w\|_{X_{l,b}^r} \|uv\|_{X_{\frac{3}{r}-1+0}^r} \lesssim \|w\|_{X_{l,b}^r} \|u\|_{X_{s,b}^r} \|v\|_{X_{s,b}^r},$$

because $2s - \frac{3}{r} + 1 > \frac{2}{r} \Leftrightarrow 2s > \frac{5}{r} - 1$, which is fulfilled for $s > \frac{2}{r}$, and $2s > \frac{3}{r}$.

Proof of (33): We obtain

$$\begin{aligned} \|\Lambda^{-1}(uv)wz\|_{H_{s-1,0}^r} &\lesssim \|\Lambda^{-1}(uv)\|_{H_{\frac{3}{r}+s-1-2s+\frac{2}{r}+b}^r} \|wz\|_{H_{2s-\frac{2}{r},0}^r} \\ &\lesssim \|uv\|_{H_{\frac{5}{r}-2-s+b}^r} \|wz\|_{H_{2s-\frac{2}{r},0}^r} \\ &\lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r} \|z\|_{X_{s,b}^r}. \end{aligned}$$

We applied Lemma 5.5 first, then Lemma 5.6, which requires $2s - \frac{5}{r} + 2 + s > \frac{2}{r} + b \Leftrightarrow 3s > \frac{8}{r} - 2$ for $b = \frac{1}{r} +$ and $2s > \frac{3}{r}$. This is fulfilled for $s > \frac{2}{r}$. Finally Cor. 5.1 is used.

Proof of (36): We estimate as follows:

$$\begin{aligned} \|uvwz\|_{H_{0,0}^r} &\lesssim \|uv\|_{H_{2s-\frac{3}{r}-b}^r} \|wz\|_{H_{\frac{6}{r}-2s+0}^r} \\ &\lesssim \|u\|_{X_{s,b}^r} \|v\|_{X_{s,b}^r} \|w\|_{X_{s,b}^r} \|z\|_{X_{s,b}^r}. \end{aligned}$$

We applied Lemma 5.5 for the first step, using $\frac{6}{r} - 2s + 2s - \frac{3}{r} + > \frac{3}{r}$, and for the second step Lemma 5.6, using $2s - (2s - \frac{3}{r}) + > \frac{3}{r}$, and Cor. 5.1, using $2s - \frac{6}{r} + 2s > \frac{2}{r} \Leftrightarrow s > \frac{2}{r}$. \square

Proof of Theorem 2.1. Now we recall the results of Tesfahun [T], who proved the same estimates in the case $r = 2$ with $s = \frac{6}{7} + \epsilon$, $l = -\frac{1}{14} + \epsilon$, $a = \frac{1}{2} +$, $b = \frac{3}{4} +$ for $\epsilon > 0$. Thus we may apply bilinear interpolation between the case $r = 1 +$ and $r = 2$.

Let $\delta > 0$ be given and $s = \frac{16}{7r} - \frac{2}{7} + \delta$, $l = \frac{15}{7r} - \frac{8}{7} + \delta$. Then for $r > 1$ sufficiently close to 1 we have $\delta \geq \frac{15}{7} - \frac{15}{7r} + \omega$ with $\omega \geq 0$. In the case $\omega = 0$ we have $s = \frac{13}{7} + \frac{1}{7r}$ and $l = 1$. Because in this case $s \geq 1 + \frac{1}{r}$, we have proven that the estimates (20) - (36) are true. By the fractional Leibniz rule they remain true for $\omega > 0$, thus for the given δ and $r > 1$ close enough to 1. Bilinear interpolation between this case and $r = 2$ implies the estimates for the whole range $1 < r \leq 2$. This gives the estimates (20) - (36) for the following choice of parameters for the whole range $1 < r \leq 2$:

$$s = \frac{16}{7r} - \frac{2}{7} + \delta, \quad l = \frac{15}{7r} - \frac{8}{7} + \delta, \quad a = \frac{1}{r} +, \quad b = \frac{1}{2} + \frac{1}{2r} + .$$

Finally, an application of Theorem 3.1 to the Cauchy problem (6),(7),(8) completes the proof of Theorem 2.1. \square

REFERENCES

- [AFS] P. d’Ancona, D. Foschi and S. Selberg: *Atlas of products for wave-Sobolev spaces on \mathbb{R}^{1+3}* . Transactions AMS 364 (2012), 31-63
- [FK] D. Foschi and S. Klainerman: *Bilinear space-time estimates for homogeneous wave equations*. Ann. Sc. ENS. 4. serie, 33 (2000), 211-274
- [GN] V. Grigoryan and A. Nahmod: *Almost critical well-posedness for nonlinear wave equation with $Q_{\mu\nu}$ null forms in 2D*. Math. Res. Letters 21 (2014), 313-332
- [G] A. Grünrock: *An improved local well-posedness result for the modified KdV equation*. Int. Math. Res. Not. (2004), no.61, 3287-3308
- [G1] A. Grünrock: *On the wave equation with quadratic nonlinearities in three space dimensions*. Hyperbolic Diff. Equ. 8 (2011), 1-8
- [GV] A. Grünrock and L. Vega: *Local well-posedness for the modified KdV equation in almost critical \dot{H}_s^r -spaces*. Trans. Amer. Mat. Soc. 361 (2009), 5681-5694
- [KM] S. Klainerman and M. Machedon: *Finite energy solutions of the Yang-Mills equations in \mathbb{R}^{3+1}* . Ann. Math. 142 (1995), 39-119
- [KS] S. Klainerman and S. Selberg: *Bilinear estimates and applications to nonlinear wave equations*. Communications in Contemporary Mathematics. 4 (2002) 223-295
- [KT] S. Klainerman and D. Tataru: *On the optimal local regularity for the Yang-Mills equations in \mathbb{R}^{4+1}* . Journal of the AMS 12 (1999), 93-116
- [KrSt] J. Krieger and J. Sterbenz: *Global regularity for the Yang-Mills equations on high dimensional Minkowski space*. Mem. AMS 223 (2013), No. 1047
- [KrT] J. Krieger and D. Tataru: *Global well-posedness for the Yang-Mills equations in 4+1 dimensions. Small energy*. Annals of Math. 185 (2017), 831-893
- [O] S. Oh: *Gauge choice for the Yang-Mills equations using the Yang-Mills heat flow and local well-posedness in H^1* . J. Hyperbolic Differ. Equ. 11 (2014), 1-108.
- [O1] S. Oh: *Finite energy global well-posedness of the Yang-Mills equations on \mathbb{R}^{1+3} : an approach using the Yang-Mills heat flow*. Duke Math. J. 164 (2015), 1669-1732.
- [P] H. Pecher: *Infinite energy solutions for the (3+1)-dimensional Yang-Mills equation in Lorenz gauge*. Comm. Pure Appl. Analysis 18 (2019), 663-688
- [P1] H. Pecher: *Low regularity local well-posedness for the higher-dimensional Yang-Mills equation in Lorenz gauge*. Adv. Diff. Equ. 24 (2019), 283-320
- [ST] S. Selberg and A. Tesfahun: *Null structure and local well-posedness in the energy class for the Yang-Mills equations in Lorenz gauge*. Journal of the European Mathematical Society 18 (2016), 1729-1752.
- [T] A. Tesfahun: *Local well-posedness of Yang-Mills equations in Lorenz gauge below the energy norm*. Nonlin. Diff. Equ. Appl. 22 (2015), 849-875
- [VV] A. Vargas and L. Vega: *Global wellposedness for 1D non-linear Schrödinger equation for data with an infinite L^2 -norm*. J. Math. Pures Appl. 80 (2001), 1029-1044