

Thermodynamic Implementations of Quantum Processes

Philippe Faist,^{1,2,3} Mario Berta,^{4,5} and Fernando G. S. L. Brandao^{1,5}

¹*Institute for Quantum Information and Matter, Caltech, Pasadena, CA 91125, U.S.A.*

²*Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland*

³*Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, 14195 Berlin, Germany*

⁴*Department of Computing, Imperial College London, London SW7 2AZ, United Kingdom*

⁵*AWS Center for Quantum Computing, Pasadena, CA 91125, U.S.A.*

July 21, 2021

Abstract

Recent understanding of the thermodynamics of small-scale systems have enabled the characterization of the thermodynamic requirements of implementing quantum processes for fixed input states. Here, we extend these results to construct optimal universal implementations of a given process, that is, implementations that are accurate for any possible input state even after many independent and identically distributed (i.i.d.) repetitions of the process. We find that the optimal work cost rate of such an implementation is given by the thermodynamic capacity of the process, which is a single-letter and additive quantity defined as the maximal difference in relative entropy to the thermal state between the input and the output of the channel. As related results we find a new single-shot implementation of time-covariant processes and conditional erasure with nontrivial Hamiltonians, a new proof of the asymptotic equipartition property of the coherent relative entropy, and an optimal implementation of any i.i.d. process with thermal operations for a fixed i.i.d. input state. Beyond being a thermodynamic analogue of the reverse Shannon theorem for quantum channels, our results introduce a new notion of quantum typicality and present a thermodynamic application of convex-split methods.

Contents

1	Introduction	2
2	Preliminaries	4
2.1	Quantum states, quantum processes, and distance measures	4
2.2	Entropy measures	5
2.3	Schur-Weyl duality	6
2.4	Estimating entropy	8
2.5	Estimating energy	8
2.6	Post-selection technique	9
3	Resource theory of thermodynamics	10
3.1	Gibbs-preserving maps	10
3.2	Thermal operations	12

4	Thermodynamic capacity	13
4.1	Definition	13
4.2	Properties	14
4.3	Optimality	16
5	Construction #1: Trivial Hamiltonians	16
5.1	Statement and proof sketch	16
5.2	Challenges for extension to non-trivial Hamiltonians	18
6	Construction #2: Gibbs-preserving maps	19
6.1	Statement and proof sketch	19
6.2	Construction via universal conditional and relative typicality	20
6.3	Universal conditional and relative typical smoothing operator	22
7	Construction #3: Thermal operations	26
7.1	Statement and proof sketch	26
7.2	Universal conditional erasure	27
7.3	Construction via universal conditional erasure	32
8	Related Results	36
8.1	Alternative proof of the asymptotic equipartition property of the coherent relative entropy	36
8.2	Optimal implementation of any i.i.d. channel with thermal operations on an i.i.d. input state	40
8.3	Single-shot erasure protocol for fixed input state and for noninteracting system and memory	43
8.4	Single-shot protocol for implementing a time-covariant processes using thermal operations	45
9	Discussion	46
A	Missing proofs	47
B	Technical lemmas	50
C	Dilation of energy-conserving operators to unitaries	51
D	Robust counterexample against extensions of Construction #1	53
E	Universal conditional typical projector for trivial Hamiltonians	54
F	Universal conditional erasure for n copies and trivial Hamiltonians	56

1 Introduction

In the information-theoretic approach to thermodynamics, a careful analysis of the resources required to perform thermodynamic tasks has allowed to consistently and systematically describe the

thermodynamic behaviour of quantum systems at the nano-scale [1]. In particular, thermodynamics can be phrased as a resource theory [2–4]. In a resource theory, one specifies which operations can be carried out at no cost—the *free operations*—and then one studies how much of external resources (e.g., thermodynamic work) one needs to provide to carry out operations that are not free. Two established resource theories for quantum thermodynamics are *thermal operations* [2, 3] and *Gibbs-preserving maps* [5, 6]. In the former, the free operations consist of energy-conserving interactions of the system with a heat bath, while in the latter, the free operations are any quantum operation that preserves the thermal state. It is reasonable to assume that thermal operations can be realized in an idealized setting, making them a good choice of framework for constructing explicit protocols, whereas Gibbs-preserving maps encompass a broader class of operations, allowing us to derive stronger fundamental limits.

The resource theory approach to thermodynamics has revealed close connections with measures of information known from quantum information theory [7, 8]. Namely, single-shot thermodynamic and information-theoretic tasks are both quantified by relevant entropy measures [9–11]. Consequently, tools from quantum Shannon theory can be used to characterize tasks in thermodynamics, for instance to derive second-order asymptotics of the work cost of state transformations [12]. Recently, focus was shifted to understand the resource costs of quantum processes, rather than state transformations [13–16]. The information measure associated with quantum processes is the quantum capacity, along with its many variants [17]. A natural question arises: What is the thermodynamic analogue of the quantum capacity?

Here, we ask how much work is required to implement a given quantum process, with the requirement that the implementation is accurate for any possible input state. In the single-instance regime, we find that the answer is a variation of the results obtained in Ref. [16]. However, in the regime where we consider many independent and identically distributed (i.i.d.) copies of the process, important differences arise due to typicality. We find that the optimal work cost of such an implementation in the i.i.d. regime is given by the thermodynamic capacity, defined as the maximal difference between the input and output free energy of the process over all possible input states. The fact that no implementation can perform better than the thermodynamic capacity follows fairly straightforwardly from the results of Ref. [16]. The technically challenging part of the present paper is to show that there exist protocols that achieve this limit.

We provide three different constructions of such protocols, each valid in different settings. In the first construction, we make the simplifying assumption that Hamiltonian of the system is trivial as in Ref. [13]. We then show that simple properties of one-shot entropy measures, coupled with the post-selection technique [18], provide an existence proof of the required implementation. The implementation is given in terms of thermal operations. In our second construction, we develop novel quantum typicality tools which we use along with the post-selection technique to explicitly construct an implementation in terms of Gibbs-preserving maps for any i.i.d. process and for any system Hamiltonian. In our third construction, we assume that the i.i.d. process is time-covariant, i.e., commutes with the time evolution. We then use recent results on the convex-split lemma and position-based decoding [19] to construct an implementation of a time-covariant i.i.d. process with thermal operations.

Our results imply that the thermodynamic resource theory of channels becomes reversible in the i.i.d. limit [20]. Namely, invoking the results in Ref. [21], we see that the work rate that is required to implement a given i.i.d. process is the same as what can be extracted if the i.i.d. process is provided to us as a black box. This provides a thermodynamic analogue of the reverse Shannon theorem from quantum information theory. This theorem states that the quantum mutual information of the channel uniquely characterizes the resources required to simulate the channel with noiseless channel uses and shared entanglement, as well as to distil a noiseless channel from many uses of the channel and shared entanglement [22, 23]. Indeed, our proof techniques are inspired by Refs. [22, 24–26].

Finally, we provide some additional results that can be of independent interest: A significantly simpler proof of the asymptotic equipartition property of the quantity studied in Ref. [16], a new protocol for Landauer erasure with side information for systems with a general non-interacting Hamiltonian and in a time-covariant state, and a protocol for implementing any general i.i.d. process for a fixed i.i.d. input state, using thermal operations and a small source of coherence.

The remainder of this paper is structured as follows. Section 2 gives the necessary preliminaries and fixes some notation. Section 3 introduces two resource theories for thermodynamics, thermal operations and Gibbs-preserving maps. In Section 4 we introduce the thermodynamic capacity and present some elementary properties. In Section 5, we provide our first construction for a trivial Hamiltonian. In Section 6 we provide our second construction, which is valid in the general setting and provides an implementation in terms of Gibbs-preserving maps. Section 7 provides our third construction, valid for time-covariant i.i.d. processes, and built with thermal operations. In Section 8, we provide three related results that use techniques developed in the above constructions. Our conclusions are presented in Section 9. Various more technical proof details are deferred to Appendices A–F.

2 Preliminaries

2.1 Quantum states, quantum processes, and distance measures

Each quantum system considered lives in a finite-dimensional Hilbert space. A quantum state is a positive semi-definite operator ρ satisfying $\text{tr}[\rho] = 1$. A sub-normalized quantum state is a positive semi-definite operator ρ satisfying $\text{tr}[\rho] \leq 1$. To each system S is associated a standard basis, usually denoted by $\{|k\rangle_S\}$. For any two systems A, A' , we denote by $A \simeq A'$ the fact that they are isometric. In that case, we consider a representation in which the isometry maps the standard basis onto the standard basis, i.e., $\text{id}_{A \rightarrow A'}(|k\rangle\langle k|_A) = |k\rangle\langle k|_{A'}$ for all k , where $\text{id}_{A \rightarrow A'}$ denotes the identity process. For any two systems $A \simeq A'$, we define the non-normalized maximally entangled reference ket $|\Phi\rangle_{A:A'} = \sum_k |k\rangle_A \otimes |k\rangle_{A'}$. Matrix inequalities are with respect to the positive semi-definite cone: $A \leq B$ signifies that $B - A$ is positive semi-definite. A completely positive map $\mathcal{E}_{X \rightarrow X'}$ is a linear mapping that maps Hermitian operators on X to Hermitian operators on X' and that satisfies $\mathcal{E}_{X \rightarrow X'}(\Phi_{X:R_X}) \geq 0$, where $R_X \simeq X$. The adjoint $\mathcal{E}_{X \leftarrow X'}^\dagger$ of a completely positive map $\mathcal{E}_{X \rightarrow X'}$ is the unique completely positive map $X' \rightarrow X$ that satisfies $\text{tr}[\mathcal{E}(Y)Z] = \text{tr}[Y\mathcal{E}^\dagger(Z)]$ for all operators Y, Z .

A completely positive map $\mathcal{E}_{X \rightarrow X'}$ is trace-preserving if $\mathcal{E}^\dagger(\mathbb{1}_{X'}) = \mathbb{1}_X$ and trace non-increasing if $\mathcal{E}^\dagger(\mathbb{1}_{X'}) \leq \mathbb{1}_X$.

Proximity of quantum states can be measured by the fidelity $F(\rho, \sigma) = \|\sqrt{\rho}\sqrt{\sigma}\|_1$, where the one-norm of an operator is defined as $\|A\|_1 = \text{tr}[\sqrt{A^\dagger A}]$. The fidelity is extended to sub-normalized states ρ, σ as the generalized fidelity, $\bar{F}(\rho, \sigma) = \|\sqrt{\rho}\sqrt{\sigma}\|_1 + \sqrt{(1 - \text{tr}[\rho])(1 - \text{tr}[\sigma])}$, noting that $F(\cdot, \cdot) = \bar{F}(\cdot, \cdot)$ whenever at least one of the states is normalized. An associated metric can be defined for any sub-normalized states as $P(\rho, \sigma) = \sqrt{1 - \bar{F}^2(\rho, \sigma)}$, called the purified distance [10, 11, 27], or root infidelity, and is closely related to the Bures distance and the quantum angle [28]. The proximity of two sub-normalized quantum states ρ, σ may also be measured in the trace distance $D(\rho, \sigma) = \frac{1}{2}\|\rho - \sigma\|_1$. We note that the one-norm of a Hermitian operator A can be expressed as

$$\|A\|_1 = \max_{\|Z\|_\infty \leq 1} \text{tr}[ZA] = \min_{\substack{\Delta_\pm \geq 0 \\ A = \Delta_+ - \Delta_-}} \text{tr}[\Delta_+] + \text{tr}[\Delta_-] , \quad (2.1)$$

where the first optimization ranges over Hermitian Z operators and where the second over positive semi-definite operators Δ_\pm . For any two states ρ, σ (one can even be sub-normalized), the purified distance and the trace distance are related via

$$D(\rho, \sigma) \leq P(\rho, \sigma) \leq \sqrt{2D(\rho, \sigma)} . \quad (2.2)$$

Similarly, we may define a distance measure for channels: For two completely positive, trace non-increasing maps $\mathcal{T}_{X \rightarrow X'}$ and $\mathcal{T}'_{X \rightarrow X'}$, the diamond norm distance is defined as

$$\frac{1}{2} \|\mathcal{T}_{X \rightarrow X'} - \mathcal{T}'_{X \rightarrow X'}\|_\diamond = \max_{\sigma_{XR}} D(\mathcal{T}_{X \rightarrow X'}(\sigma_{XR}), \mathcal{T}'_{X \rightarrow X'}(\sigma_{XR})) , \quad (2.3)$$

where the optimization ranges over all bipartite quantum states over X and a reference system $R \simeq X$. The optimization may be restricted to pure states without loss of generality.

2.2 Entropy measures

The von Neumann entropy of a quantum state ρ is $H(\rho) = -\text{tr}[\rho \ln \rho]$. In this work, all entropies are defined in units of nats, using the natural logarithm $\ln(\cdot)$, instead of units of (qu)bits. A number of nats is equal to $\ln(2)$ times the corresponding number of qubits. The conditional von Neumann entropy of a bipartite state ρ_{AB} is given by

$$H(A|B)_\rho = H(AB)_\rho - H(B)_\rho = H(\rho_{AB}) - H(\rho_B) . \quad (2.4)$$

The quantum relative entropy is defined as

$$D(\rho \| \sigma) = \text{tr}[\rho (\ln \rho - \ln \sigma)] , \quad (2.5)$$

where ρ is a quantum state and where σ is any positive semi-definite operator whose support contains the support of ρ .

One of our proofs relies on the hypothesis testing relative entropy [29–33] in its form as presented in [34]. For any sub-normalized quantum state ρ and for any positive semi-definite operator σ whose support contains the support of ρ , we define it via the following equivalent optimizations, which are semi-definite programs [35] in terms of the primal variable $Q \geq 0$ and the dual variables $\mu, X \geq 0$:

$$\begin{aligned} \exp\{-D_{\text{H}}^{\eta}(\rho \parallel \sigma)\} &= \text{minimize : } \eta^{-1} \text{tr}[Q\sigma] \\ &\text{subject to : } Q \leq \mathbb{1} \\ &\text{tr}[Q\rho] \geq \eta \end{aligned} \quad (2.6)$$

$$\begin{aligned} &= \text{maximize : } \mu - \eta^{-1} \text{tr}[X] \\ &\text{subject to : } \mu\rho \leq \sigma + X . \end{aligned} \quad (2.7)$$

We also define for convenience the closely related quantity

$$D_{\text{h}}^{\eta}(\rho \parallel \sigma) = D_{\text{H}}^{\eta}(\rho \parallel \sigma) - \ln(\eta) , \quad (2.8)$$

which is sometimes also referred to as the hypothesis testing relative entropy.

2.3 Schur-Weyl duality

Consider a Hilbert space \mathcal{H}_A and $n \in \mathbb{N}$. The group $\text{GL}(d_A) \times \text{S}_n$ acts naturally on $\mathcal{H}_A^{\otimes n}$, where $X \in \text{GL}(d_A)$ acts as $X^{\otimes n}$ and where the permutation group permutes the tensor factors. We follow closely the notation of Refs. [24, 25]. Schur-Weyl tells us that the full Hilbert space decomposes as

$$\mathcal{H}_A \simeq \bigoplus_{\lambda} \mathcal{V}_{\lambda} = \bigoplus_{\lambda} \mathcal{Q}_{\lambda} \otimes \mathcal{P}_{\lambda} , \quad (2.9)$$

where $\lambda \in \text{Young}(n, d)$ are Young diagrams with n boxes and (at most) d rows, and where \mathcal{Q}_{λ} , \mathcal{P}_{λ} are irreducible representations of $\text{GL}(d_A)$ and S_n , respectively. The number of Young diagrams in the decomposition above is at most $\text{poly}(n)$, if d_A is kept constant. We write $\text{poly}(n) = O(\text{poly}(n))$ in big O notation for terms whose absolute value is upper bounded by some polynomial n^c for $c \in \mathbb{N}$ in the asymptotic limit $n \rightarrow \infty$.

We denote by $\Pi_{A^n}^{\lambda}$ the projector in $\mathcal{H}_A^{\otimes n}$ onto the term labelled by λ in the decomposition above. We denote by $q_{\lambda}(X)$ a representing matrix of $X \in \text{GL}(d_A)$ in the irreducible representation labelled by λ ; the operator $q_{\lambda}(X)$ lives in \mathcal{Q}_{λ} . We furthermore introduce the following notation, for any $Y \in \mathcal{Q}_{\lambda} \otimes \mathcal{P}_{\lambda}$,

$$[Y]_{\lambda} = \mathbb{1}_{(\mathcal{Q}_{\lambda} \otimes \mathcal{P}_{\lambda}) \rightarrow A^n} Y \mathbb{1}_{(\mathcal{Q}_{\lambda} \otimes \mathcal{P}_{\lambda}) \leftarrow A^n}^{\dagger} , \quad (2.10)$$

which represents the canonical embedding of an operator Y on $\mathcal{Q}_{\lambda} \otimes \mathcal{P}_{\lambda}$ into the space $\mathcal{H}_A^{\otimes n}$, i.e., mapping Y onto the corresponding block in (2.9). In particular,

$$\Pi_{A^n}^{\lambda} [Y]_{\lambda} \Pi_{A^n}^{\lambda} = [Y]_{\lambda} . \quad (2.11)$$

Any operator X_{A^n} acting on the n copies which commutes with all the permutations admits a decomposition of the form

$$X_{A^n} = \sum_{\lambda} [X_{\lambda} \otimes \mathbb{1}_{\mathcal{P}_{\lambda}}]_{\lambda} \quad (2.12)$$

for some set of operators $X_{\lambda} \in \mathcal{Q}_{\lambda}$. In particular, $[X_{A^n}, \Pi_{A^n}^{\lambda}] = 0$. We can make this more precise for i.i.d. states. For any $X \in \text{GL}(d_A)$, we have that

$$[\Pi_{A^n}^{\lambda}, X^{\otimes n}] = 0 \quad (2.13)$$

$$X^{\otimes n} = \sum_{\lambda} [q_{\lambda}(X) \otimes \mathbb{1}_{\mathcal{P}_{\lambda}}]_{\lambda} . \quad (2.14)$$

For a given $\lambda \in \text{Young}(n, d)$, it is often useful to consider the corresponding normalized probability distribution $\lambda/n = (\lambda_i/n)_i$. The entropy of this distribution is given by

$$\bar{H}(\lambda) = H(\lambda/n) = - \sum_i \frac{\lambda_i}{n} \ln \frac{\lambda_i}{n} , \quad (2.15)$$

where λ_i is the number of boxes in the i -th row of the diagram.

If we have n copies of a bipartite system $\mathcal{H}_A \otimes \mathcal{H}_B$, then we may Schur-Weyl decompose $\mathcal{H}_A^{\otimes n}$, $\mathcal{H}_B^{\otimes n}$ and $(\mathcal{H}_A \otimes \mathcal{H}_B)^{\otimes n}$ under the respective actions of $\text{GL}(d_A) \times S_n$, $\text{GL}(d_B) \times S_n$ and $\text{GL}(d_A d_B) \times S_n$. A useful property we will need here is that the projectors onto the respective Schur-Weyl blocks commute between these decompositions.

Lemma 2.1. *Consider two spaces $\mathcal{H}_A, \mathcal{H}_B$ and let $\Pi_{A^n B^n}^{\lambda}$ and $\Pi_{A^n}^{\lambda'}$ be the projectors onto Schur-Weyl blocks of $\mathcal{H}_{AB}^{\otimes n}$ and $\mathcal{H}_A^{\otimes n}$, respectively, with $\lambda \in \text{Young}(d_A d_B, n)$ and $\lambda' \in \text{Young}(d_A, n)$. Then, we have*

$$[\Pi_{A^n B^n}^{\lambda}, \Pi_{A^n}^{\lambda'} \otimes \mathbb{1}_{B^n}] = 0 . \quad (2.16)$$

Proof. $\Pi_{A^n}^{\lambda'} \otimes \mathbb{1}_{B^n}$ is invariant under the action of S_n permuting the copies of $A \otimes B$, and so it admits a decomposition of the form (2.12) and commutes with $\Pi_{A^n B^n}^{\lambda}$. \blacksquare

The following is another lemma about how much overlap Schur-Weyl blocks have on a bipartite system versus on one of the two systems. This lemma forms the basis of our universal typical subspace.

Lemma 2.2. *Consider $n \in \mathbb{N}$ copies of a bipartite system $\mathcal{H}_A \otimes \mathcal{H}_B$. Then, for any $\lambda \in \text{Young}(d_A d_B, n)$ and $\lambda' \in \text{Young}(d_B, n)$, we have*

$$\Pi_{B^n}^{\lambda'} \text{tr}_{A^n} [\Pi_{A^n B^n}^{\lambda}] \Pi_{B^n}^{\lambda'} \leq \text{poly}(n) e^{n(\bar{H}(\lambda) - \bar{H}(\lambda'))} \Pi_{B^n}^{\lambda'} \quad (2.17)$$

noting that $[\mathbb{1}_{A^n} \otimes \Pi_{B^n}^{\lambda'}, \Pi_{A^n B^n}^{\lambda}] = 0$.

(Proof on page 47.)

The proof is provided in [Appendix A](#).

2.4 Estimating entropy

Measuring the Young diagram λ — that is, performing the projective measurement with operators $\{\Pi_{A^n}^\lambda\}_\lambda$ — yields a good estimation of the spectrum of a state ρ_A when given $\rho_A^{\otimes n}$ [25]. An estimate for the entropy of ρ is thus obtained by calculating the entropy $H(\lambda/n)$ corresponding to the probability distribution λ/n .

Proposition 2.1 (Spectrum and entropy estimation [22, 24, 25]). *Consider $n \in \mathbb{N}$ copies of a system \mathcal{H}_A . Then, the family of projectors $\{\Pi_{A^n}^\lambda\}_\lambda$ given by Schur-Weyl duality forms a POVM obeying the following property: For any $\delta > 0$, there exists an $\eta > 0$ such that for any state ρ_A , we have*

$$\text{tr} \left[\left(\sum_{\lambda: \bar{H}(\lambda) \in [H(\rho) \pm \delta]} \Pi_{A^n}^\lambda \right) \rho_A^{\otimes n} \right] \geq 1 - \text{poly}(n) \exp(-n\eta). \quad (2.18)$$

(Proof on [page 47](#).)

The proof is provided in [Appendix A](#).

2.5 Estimating energy

Proposition 2.2. *Consider any observable H_A on \mathcal{H}_A and write $\Gamma_A = e^{-H_A}$. Then, the set of projectors $\{R_{A^n}^k\}$ onto the eigenspaces of $\Gamma_A^{\otimes n}$ forms a POVM satisfying the following properties:*

- (i) *There are at most $\text{poly}(n)$ POVM elements, with the label k running over a set $k \in \mathcal{K}_n(H_A) \subset \mathbb{R}$;*
- (ii) *We have $[R_{A^n}^k, \Gamma_A^{\otimes n}] = 0$ and $e^{-nk} R_{A^n}^k = R_{A^n}^k \Gamma_A^{\otimes n}$;*
- (iii) *For any $\delta > 0$ and for any state ρ_A ,*

$$\text{tr} \left[R_{A^n}^{\approx \delta \text{tr}[\rho_A H_A]} \rho_A^{\otimes n} \right] \geq 1 - 2e^{-n\eta} \quad (2.19)$$

with $\eta = \delta^2 / (2\|H_A\|_\infty^2)$ and where we defined for any $h \in \mathbb{R}$ that

$$R_{A^n}^{\approx \delta h} = \sum_{k \in \mathcal{K}_n(H_A) : |k-h| \leq \delta} R_{A^n}^k; \quad (2.20)$$

- (iv) *For any $h \in \mathbb{R}$, we have*

$$e^{-n(k+\delta)} R_{A^n}^{\approx \delta h} \leq R_{A^n}^{\approx \delta h} \Gamma_A^{\otimes n} \leq e^{-n(k-\delta)} R_{A^n}^{\approx \delta h}. \quad (2.21)$$

(Proof on [page 48](#).)

The proof is provided in [Appendix A](#).

2.6 Post-selection technique

The post-selection technique is useful for bounding the diamond norm of a candidate smoothed channel to a target ideal i.i.d. channel.

Theorem 2.1 (Post-selection technique [18]). *Let X, X' be quantum systems, $\mathcal{E}_{X \rightarrow X'}$ be a completely positive, trace-preserving map, and $\mathcal{T}_{X^n \rightarrow X'^n}$ be a completely positive, trace non-increasing map. Furthermore, let $\bar{R} \simeq X$,*

$$\zeta_{X^n} = \text{tr}_{\bar{R}^n} \left[\int d\phi_{X\bar{R}} |\phi\rangle\langle\phi|_{X\bar{R}}^{\otimes n} \right] = \int d\sigma_X \sigma_X^{\otimes n}, \quad (2.22)$$

where $d\phi_{X\bar{R}}$ denotes the Haar-induced measure on the pure states on $X \otimes \bar{R}$, and $d\sigma_X$ its induced measure on X after partial trace, and let $|\zeta\rangle_{X^n R}$ be a purification of ζ_{X^n} . Then, we have

$$\frac{1}{2} \|\mathcal{T} - \mathcal{E}^{\otimes n}\|_{\diamond} \leq \text{poly}(n) D(\mathcal{T}(\zeta_{X^n R}), \mathcal{E}^{\otimes n}(\zeta_{X^n R})). \quad (2.23)$$

Moreover, for all $n \in \mathbb{N}$ there exists a set $\{|\phi_i\rangle_{X\bar{R}}\}$ of at most $\text{poly}(n)$ states, and a probability distribution $\{p_i\}$, providing a purification of ζ_{X^n} as

$$|\zeta\rangle_{X^n \bar{R}^n R'} = \sum_i \sqrt{p_i} |\phi_i\rangle_{X\bar{R}}^{\otimes n} \otimes |i\rangle_{R'} \quad (2.24)$$

with a register R' of size $\text{poly}(n)$.

The first part of the theorem is [18, Eq. (4)] and the second part is, e.g., found as [23, Cor. D.6]. The following proposition shows that a given channel is close to an i.i.d. channel, if it behaves as expected on all i.i.d. states with exponentially good accuracy.

Proposition 2.3. *For three systems X, X', E , let $V_{X \rightarrow X'E}$ be an isometry and $W_{X^n \rightarrow X'^n E^n}$ be an isometry which commutes with the permutations of the n systems. Furthermore, assume that there exists $\eta > 0$ independent of n such that for all pure states $|\sigma\rangle\langle\sigma|_{X R_X}$ with a reference system $R_X \simeq X$, we have*

$$\text{Re} \left\{ \langle \sigma |_{X R_X}^{\otimes n} (V_{X \rightarrow X'E}^\dagger)^{\otimes n} W_{X^n \rightarrow X'^n E^n} | \sigma \rangle_{X R_X}^{\otimes n} \right\} \geq 1 - \text{poly}(n) \exp(-n\eta). \quad (2.25)$$

For $\mathcal{E}_{X \rightarrow X'}(\cdot) = \text{tr}_E [V_{X \rightarrow X'E}(\cdot) V^\dagger]$ and $\mathcal{T}_{X^n \rightarrow X'^n}(\cdot) = \text{tr}_{E^n} [W_{X^n \rightarrow X'^n E^n}(\cdot) W^\dagger]$ we then have

$$\frac{1}{2} \|\mathcal{T}_{X^n \rightarrow X'^n} - \mathcal{E}_{X \rightarrow X'}^{\otimes n}\|_{\diamond} \leq \text{poly}(n) \exp(-n\eta/2). \quad (2.26)$$

(Proof on page 48.)

The proof is provided in [Appendix A](#).

3 Resource theory of thermodynamics

3.1 Gibbs-preserving maps

We consider the framework of Ref. [16], where for each system S considered a positive semi-definite operator $\Gamma_S \geq 0$ is associated. A trace non-increasing, completely positive map $\Phi_{A \rightarrow B}$ is allowed for free if it satisfies $\Phi_{A \rightarrow B}(\Gamma_A) \leq \Gamma_B$. In the case of a system S with Hamiltonian H_S , and in the presence of a single heat bath at inverse temperature β , the relevant thermodynamic framework is given by setting $\Gamma_S = e^{-\beta H_S}$. In the remainder of this paper, when using the present framework, it is convenient to work with the Γ operators on an abstract level. The results then also apply to situations where several different thermodynamic baths are considered, or in more general settings where a specific operator needs to be conserved by the spontaneous evolution of the system [16].

The resources required to enable non-free operations are counted using an explicit system that provides these resources, such as an *information battery*. An information battery is a large register W whose associated operator Γ_W is simply $\Gamma_W = \mathbb{1}_W$ (i.e., $H_W = 0$). The information battery is required to be in a state of the special form $\tau_W^m = P_W^m / \text{tr}[P_W^m]$ where P_W^m is a projector of rank e^m . That is, τ_W^m has uniform eigenvalues over a given rank e^m . We denote the *charge* or *resource value* of a battery state τ_W^m by $w(\tau_W^m) = \ln(d) - m$, where d is the dimension of the information battery. The value $w(\tau)$ measures the amount of purity present in the state τ , which is the basic resource required to implement maps that are not already Gibbs-preserving maps. We choose to measure $w(\tau)$ in units of number of pure nats, equal to $\ln(2)$ times a number of pure qubits. A Gibbs-preserving map that acts jointly on a system and an information battery, and which maps the input battery state τ to the output battery state τ' , is deemed to *consume an amount of work* $w = w(\tau) - w(\tau')$.

The resources can be counted in terms of thermodynamic work in units of energy if we are given a heat bath at inverse temperature T . Recall that a pure qubit can be converted to $kT \ln(2)$ work using a Szilárd engine, where k is Boltzmann's constant [36]. By counting purity in nats instead of qubits, we get rid of the $\ln(2)$ factor: A number λ of pure nats can be converted into λkT thermodynamic work using a Szilárd-type engine. We count work exclusively in equivalent of pure nats, for simplicity, as opposed to units of energy. The two are directly related by a factor $\beta^{-1} = kT$. Furthermore, this eliminates the factor β from otherwise essentially information-theoretic expressions, and our theorems thus directly apply to cases where $\Gamma_X, \Gamma_{X'}$ are any abstract positive semi-definite operators which are not necessarily defined via a Hamiltonian.

Let $\Phi_{XW \rightarrow X'W}$ be a Gibbs-preserving map acting on an information battery W , and let $\tau_W^m, \tau_W^{m'}$ be two information battery states. An implementation running the operation $\Phi_{XW \rightarrow X'W}$ with the given input and output battery states is tasked to (a) make available the input battery state, (b) apply the operation $\Phi_{XW \rightarrow X'W}$, and (c) check that the output battery state is appropriate (e.g., for possible future re-use). For the verification in Point (c) it is sufficient to measure the two-outcome POVM $\{P_W^{m'}, \mathbb{1} - P_W^{m'}\}$; as long as the first outcome is observed, it is always possible to bring the state to $\tau_W^{m'}$ by applying a completely thermalizing operation on the support of $P_W^{m'}$ (here, this is a completely randomizing or completely symmetrizing operation). In the constructions presented in the present paper, we allow this verification measurement to fail with a small fixed probability $\varepsilon > 0$.

A convenient mathematical object to characterize what the operation does on the system is the following. The *effective work process* $\mathcal{T}_{X \rightarrow X'F}$ associated with $\Phi_{XW \rightarrow X'W}$ and $(\tau_W^m, \tau_W^{m'})$ is the trace non-increasing map defined as

$$\mathcal{T}_{X \rightarrow X'}(\cdot) = \text{tr}_W \left[P_W^{m'} \Phi_{XW \rightarrow X'W}((\cdot) \otimes \tau_W^m) \right]. \quad (3.1)$$

The question of implementing a process \mathcal{E} becomes the issue of finding a Gibbs-preserving map along with battery states such that the associated effective work process is close to \mathcal{E} . Specifically, if $\|\mathcal{T}_{X \rightarrow X'} - \mathcal{E}_{X \rightarrow X'}\|_\diamond \leq \varepsilon$, then we can assert that the failure probability in Point (c) above is bounded by ε for all possible inputs on X ; the operation therefore implements $\mathcal{E}_{X \rightarrow X'}$ accurately with high success probability.

A useful characterization of which processes can be implemented using an information battery is given by the following proposition.

Proposition 3.1 ([16, Proposition I]). *Let $\Gamma_X, \Gamma_{X'} \geq 0$, $\mathcal{T}_{X \rightarrow X'}$ be a completely positive, trace non-increasing map, and $w \in \mathbb{R}$. Then, the following are equivalent:*

- (i) *We have $\mathcal{T}_{X \rightarrow X'}(\Gamma_X) \leq e^w \Gamma_{X'}$;*
- (ii) *For all $\delta > 0$ there exists an information battery W and two battery states τ_W, τ'_W such that $w(\tau_W) - w(\tau'_W) \leq w + \delta$, and there exists a Gibbs-preserving map $\Phi_{XW \rightarrow X'W}$ with $\mathcal{T}_{X \rightarrow X'}$ the effective work process associated with $\Phi_{XW \rightarrow X'W}$ and (τ_W, τ'_W) .*

Therefore, to show that one can implement $\mathcal{E}_{X \rightarrow X'}$ with Gibbs-preserving maps while expending work w , it suffices to exhibit a map $\mathcal{T}_{X \rightarrow X'}$ that is ε -close to $\mathcal{E}_{X \rightarrow X'}$ in diamond distance and that satisfies $\mathcal{T}_{X \rightarrow X'}(\Gamma_X) \leq e^w \Gamma_{X'}$. From the proof in [16] we know in Point (ii) above that $W, \tau_W \equiv \tau_W^m$ and $\tau'_W \equiv \tau_W^{m'}$ can be chosen freely as long as $m' - m = w(\tau_W) - w(\tau'_W) \geq w$ and that the corresponding Gibbs-preserving map is given by

$$\Phi_{XW \rightarrow X'W}(\cdot) = \mathcal{T}_{X \rightarrow X'}[\text{tr}_W(P_W^m(\cdot))] \otimes \tau_W^{m'}. \quad (3.2)$$

In Ref. [16], the resource cost w of implementing a process $\mathcal{E}_{X \rightarrow X'}$ (any completely positive, trace-preserving map) up to an accuracy $\varepsilon \geq 0$ in terms of proximity of the process matrix given a fixed input state σ_X , counted in pure nats, was shown to be given by the *coherent relative entropy*

$$w = -\hat{D}_{X \rightarrow X'}^\varepsilon(\mathcal{E}_{X \rightarrow X'}(\sigma_{XR_X}) \parallel \Gamma_X, \Gamma_{X'}) = \ln \min_{\substack{\mathcal{T}(\Gamma_X) \leq \alpha \Gamma_{X'} \\ P(\mathcal{T}(\sigma_{XR_X}), \mathcal{E}(\sigma_{XR_X})) \leq \varepsilon}} \alpha, \quad (3.3)$$

where σ_{XR_X} is the purification of σ_X on a system $R_X \simeq X$ given by $|\sigma\rangle_{XR} = \sigma_X^{1/2} |\Phi\rangle_{X:R_X}$, and where the optimization ranges over completely positive, trace non-increasing maps $\mathcal{T}_{X \rightarrow X'}$. The coherent relative entropy enjoys a collection of properties in relation to the conditional min- and max-entropy, and to the min- and max-relative entropy. It satisfies the following asymptotic equipartition property:

For a completely positive, trace-preserving map $\mathcal{E}_{X \rightarrow X'}$ and quantum state σ_X we have for $0 < \varepsilon < 1$ that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \hat{D}_{X^n \rightarrow X'^n}^\varepsilon \left(\mathcal{E}_{X \rightarrow X'}^{\otimes n}(\sigma_{XX}^{\otimes n}) \parallel \Gamma_X^{\otimes n}, \Gamma_{X'}^{\otimes n} \right) = D(\sigma_X \parallel \Gamma_X) - D(\mathcal{E}(\sigma_X) \parallel \Gamma_{X'}) . \quad (3.4)$$

3.2 Thermal operations

The framework of Gibbs-sub-preserving maps is technically convenient, but it is unclear whether any Gibbs-sub-preserving operation can be implemented at no work cost using other frameworks. This includes for example thermal operations that might be considered more operational

Here, we consider the alternative framework of *thermal operations* [2, 3, 8]. Each system S of interest has an associated Hamiltonian H_S and is not interacting with the other systems. For a given fixed inverse temperature β , we allow the following operations to be carried out for free:

- (i) Apply any unitary operation that commutes with the total Hamiltonian;
- (ii) Bring in any ancillary system in its Gibbs state at inverse temperature β ; and
- (iii) Discard any system.

The most general transformation a system S can undergo under this set of rules is a *thermal operation*. A thermal operation is any process that can be implemented using an additional system B with any Hamiltonian H_B and with any unitary U_{SB} satisfying $[U_{SB}, H_S + H_B] = 0$, resulting in the completely positive, trace-preserving map

$$\Phi_S(\cdot) = \text{tr}_B [U_{SB} ((\cdot) \otimes \gamma_B) U_{SB}^\dagger] , \quad (3.5)$$

where $\gamma_B = e^{-\beta H_B} / \text{tr}[e^{-\beta H_B}]$ is the Gibbs state of the bath system B . Observe that any concatenation of thermal operations is again a thermal operation.

Clearly, any thermal operation Φ_S leaves the thermal state $\gamma_S = e^{-\beta H_S} / \text{tr}[e^{-\beta H_S}]$ on S invariant. Hence, any lower bound on the work cost of an implementation derived in the framework of Gibbs-preserving maps also applies to thermal operations. We use the same definitions of work and the effective work process for thermal operations as we defined for Gibbs-preserving maps earlier: an information battery is used to account for work, and the effective work process associated with a thermal operation $\Phi_{XW \rightarrow XW}$ with respect to battery states $(\tau_W^m, \tau_W^{m'})$ is also defined by (3.1).

When considering only states that commute with the Hamiltonian, a powerful tool to characterize possible state transformations is the notion of thermomajorization [8]. In the fully quantum regime, there is in contrast no known simple mathematical characterization of the work required to implement a quantum process with thermal operations. In fact, because thermal operations are time-covariant, it is impossible to implement processes that are not time-covariant, even if the latter might admit an implementation with a Gibbs-preserving map [6].

We will later use a primitive that transforms a thermal state into a pure energy eigenstate. The next statement follows directly from [8, Eq. (8) and Suppl. Note 4].

Proposition 3.2. *Let $\gamma_X = e^{-\beta H_X} / \text{tr}[e^{-\beta H_X}]$ be the thermal state on a system X with Hamiltonian H_X , and let $|E\rangle_X$ be a pure energy eigenstate of H_X . There exists a thermal operation Φ_{XW} on an information battery with battery states (τ_W, τ'_W) such that $\Phi_{XW}(\gamma_X \otimes \tau_W) = |E\rangle_X \langle E|_X \otimes \tau'_W$ and such that $w(\tau_W) - w(\tau'_W)$ can be chosen arbitrarily close to $\beta E + \ln \text{tr}[e^{-\beta H_X}]$.*

4 Thermodynamic capacity

4.1 Definition

Let X, X' be quantum systems, $\mathcal{E}_{X \rightarrow X'}$ be a quantum process, and $\varepsilon > 0$. We seek a free thermodynamic operation (either a thermal operation or a Gibbs preserving map) $\Phi_{X^n W \rightarrow X'^n W}$ that acts on $X^{\otimes n}$ and a battery W , with output on $X'^{\otimes n}$ and W , as well as information battery states $\tau_W^{(i)}$ and $\tau_W^{(f)}$, such that:

- (i) The effective work process $\mathcal{T}_{X^n \rightarrow X'^n}$ of $\Phi_{X^n W \rightarrow X'^n W}$ with respect to $(\tau_W^{(i)}, \tau_W^{(f)})$ is ε -close in diamond distance to $\mathcal{E}_{X \rightarrow X'}^{\otimes n}$;
- (ii) We seek to minimize the work consumption per copy w given by

$$w = \frac{1}{n} \left[w(\tau_W^{(i)}) - w(\tau_W^{(f)}) \right]. \quad (4.1)$$

Our main result is a collection of three independent constructions of such implementations in different regimes, using either Gibbs-preserving maps or thermal operations. In each case, the amount of work consumed per copy is given by a quantity which we call the *thermodynamic capacity* of the process, and which turns out to be the minimal work cost an implementation satisfying the above conditions can achieve. The thermodynamic capacity of a completely positive, trace-preserving map $\mathcal{E}_{X \rightarrow X'}$ relative to operators $\Gamma_X, \Gamma_{X'} > 0$ is defined as

$$T(\mathcal{E}) = \sup_{\sigma_X} \left\{ D(\mathcal{E}_{X \rightarrow X'}(\sigma_X) \parallel \Gamma_{X'}) - D(\sigma_X \parallel \Gamma_X) \right\}. \quad (4.2)$$

In a fully thermodynamic context where $\Gamma_X = e^{-\beta H_X}$ and $\Gamma_{X'} = e^{-\beta H_{X'}}$, one can choose to express the thermodynamic capacity in units of energy rather than in nats, in which case a pre-factor β^{-1} may be included in the definition above such that the thermodynamic capacity is a difference of free energies

$$T(\mathcal{E}) = \sup_{\sigma} \left\{ F_{H'}(\mathcal{E}(\sigma)) - F_H(\sigma) \right\} \quad \text{with} \quad F_H(\rho) = \beta^{-1} D(\rho \parallel e^{-\beta H}). \quad (4.3)$$

Construction for trivial Hamiltonians First, in Section 5 we consider the special case where $\Gamma_X = \mathbb{1}_X$ and $\Gamma_{X'} = \mathbb{1}_{X'}$ corresponding to trivial Hamiltonians and show that simple considerations based on properties of known entropy measures guarantee the existence of a universal implementation of $\mathcal{E}^{\otimes n}$ with either thermal operations or Gibbs-preserving maps.

Construction using Gibbs-preserving maps Second, in Section 6 we consider the case of general $\Gamma_X, \Gamma_{X'}$ and we construct a universal implementation of $\mathcal{E}_{X \rightarrow X'}^{\otimes n}$ with Gibbs-preserving maps, based on new typicality considerations.

Construction using thermal operations Third, for arbitrary Hamiltonians we construct in Section 7 a universal implementation of $\mathcal{E}_{X \rightarrow X'}^{\otimes n}$ with thermal operations, assuming that \mathcal{E} is time-covariant, i.e., that it commutes with the time evolution operation.

4.2 Properties

The thermodynamic capacity is a convex optimization program. Namely, the objective function of the optimization in (4.2) can be written as

$$\begin{aligned} D(\mathcal{E}_{X \rightarrow X'}(\sigma_X) \parallel \Gamma_{X'}) - D(\sigma_X \parallel \Gamma_X) \\ &= -H(\mathcal{E}_{X \rightarrow X'}(\sigma_X)) + H(\sigma_X) - \text{tr}[\mathcal{E}_{X \rightarrow X'}(\sigma_X) \ln \Gamma_{X'}] + \text{tr}[\sigma_X \ln \Gamma_X] \\ &= H(E \mid X')_\rho - \text{tr}[\mathcal{E}_{X \rightarrow X'}(\sigma_X) \ln \Gamma_{X'}] + \text{tr}[\sigma_X \ln \Gamma_X], \end{aligned} \quad (4.4)$$

where we defined the state $\rho_{EX'} = V_{X \rightarrow X'E} \sigma_X V^\dagger$ using a Stinespring dilation $V_{X \rightarrow X'E}$ of $\mathcal{E}_{X \rightarrow X'}$ into an environment system E , satisfying $\mathcal{E}_{X \rightarrow X'}(\cdot) = \text{tr}_E[V(\cdot)V^\dagger]$. The conditional entropy is concave in the quantum state as $H(E \mid X')_\rho = -D(\rho_{EX'} \parallel \mathbb{1}_E \otimes \rho_{X'})$ and the quantum relative entropy is jointly convex. The other terms in (4.4) are linear. Hence, the optimization (4.2) is a convex optimization that can be carried out efficiently for small system sizes [37]. Indeed, we have successfully computed the thermodynamic capacity of simple example quantum channels acting on few qubits with Python code, using the QuTip framework [38, 39] and the CVXOPT optimization software [40] (see also [41] for a direct algorithm).

The thermodynamic capacity is additive [21]. As a consequence of this property, it is not necessary to include a stabilization over a reference system in the definition of the thermodynamic capacity. That is, had we optimized over bipartite states σ_{XR} with a reference system R for any Γ_R , on which the process acts as the identity process, we would be effectively computing $T(\mathcal{E} \otimes \text{id}_R)$. However, additivity implies that $T(\mathcal{E} \otimes \text{id}_R) = T(\mathcal{E})$.

Proposition 4.1 (Additivity of thermodynamic capacity [21]). *For $\Gamma_X, \Gamma_{X'}, \Gamma_Z, \Gamma_{Z'} > 0$ and quantum channels $\mathcal{E}_{X \rightarrow X'}, \mathcal{F}_{Z \rightarrow Z'}$ we have*

$$T(\mathcal{E} \otimes \mathcal{F}) = T(\mathcal{E}) + T(\mathcal{F}). \quad (4.5)$$

For completeness we provide an independent proof of additivity, to ensure validity in the general setting of abstract Γ operators.

Proof. Let σ_X, τ_Z be states achieving the thermodynamic capacity of $T(\mathcal{E})$ and $T(\mathcal{F})$, respectively. Then, $\sigma_X \otimes \tau_Z$ is a candidate for $T(\mathcal{E} \otimes \mathcal{F})$, yielding

$$\begin{aligned} T(\mathcal{E} \otimes \mathcal{F}) &\geq D(\mathcal{E}(\sigma) \otimes \mathcal{F}(\tau) \parallel \Gamma_{X'} \otimes \Gamma_{Z'}) - D(\sigma \otimes \tau \parallel \Gamma_X \otimes \Gamma_Z) \\ &= D(\mathcal{E}(\sigma) \parallel \Gamma_{X'}) - D(\sigma \parallel \Gamma_X) + D(\mathcal{F}(\tau) \parallel \Gamma_{Z'}) - D(\tau \parallel \Gamma_Z) \\ &= T(\mathcal{E}) + T(\mathcal{F}). \end{aligned} \quad (4.6)$$

Now, let ζ_{XZ} achieve the optimum for $T(\mathcal{E} \otimes \mathcal{F})$. Let $V_{X \rightarrow E_1 X'}, W_{Z \rightarrow E_2 Z'}$ be Stinespring isometries of \mathcal{E} and \mathcal{F} respectively, such that $\mathcal{E}(\cdot) = \text{tr}_{E_1}[V(\cdot)V^\dagger]$ and $\mathcal{F}(\cdot) = \text{tr}_{E_2}[W(\cdot)W^\dagger]$. Let $\rho_{E_1 E_2 X' Z'} = (V \otimes W) \zeta (V \otimes W)^\dagger$. Then, we have

$$\begin{aligned} T(\mathcal{E} \otimes \mathcal{F}) &= D((\mathcal{E} \otimes \mathcal{F})(\zeta) \parallel \Gamma_{X'} \otimes \Gamma_{Z'}) - D(\zeta_{XZ} \parallel \Gamma_X \otimes \Gamma_Z) \\ &= H(E_1 E_2 | X' Z')_\rho - \text{tr}[\rho_{X' Z'} \ln(\Gamma_{X'} \otimes \Gamma_{Z'})] + \text{tr}[\zeta_{XZ} \ln(\Gamma_X \otimes \Gamma_Z)] , \\ &= H(E_1 E_2 | X' Z')_\rho - \text{tr}[\rho_{X'} \ln(\Gamma_{X'})] - \text{tr}[\rho_{Z'} \ln(\Gamma_{Z'})] \\ &\quad + \text{tr}[\zeta_X \ln(\Gamma_X)] + \text{tr}[\zeta_Z \ln(\Gamma_Z)] \end{aligned} \quad (4.7)$$

since $\ln(A \otimes B) = \ln(A) \otimes \mathbb{1} + \mathbb{1} \otimes \ln(B)$. Invoking the chain rule of the von Neumann entropy, and then strong sub-additivity of the entropy, we see that $H(E_1 E_2 | X' Z')_\rho = H(E_1 | X' Z')_\rho + H(E_2 | E_1 X' Z')_\rho \leq H(E_1 | X')_\rho + H(E_2 | Z')_\rho$. Hence, we have

$$\begin{aligned} (4.7) &\leq H(E_1 | X')_\rho - \text{tr}[\rho_{X'} \ln(\Gamma_{X'})] + \text{tr}[\zeta_X \ln(\Gamma_X)] \\ &\quad + H(E_2 | Z')_\rho - \text{tr}[\rho_{Z'} \ln(\Gamma_{Z'})] + \text{tr}[\zeta_Z \ln(\Gamma_Z)] \\ &\leq T(\mathcal{E}) + T(\mathcal{F}), \end{aligned} \quad (4.8)$$

where the last inequality holds because the reduced states ζ_X, ζ_Z are optimization candidates for $T(\mathcal{E})$ and $T(\mathcal{F})$, respectively. \blacksquare

A special case worth mentioning is when $\Gamma_X = \mathbb{1}_X, \Gamma_{X'} = \mathbb{1}_{X'}$, which corresponds to the situation where the Hamiltonians of X and X' are trivial. For any quantum channel $\mathcal{E}_{X \rightarrow X'}$, let $V_{X \rightarrow X' E}$ be a Stinespring dilation isometry with $\mathcal{E}_{X \rightarrow X'}(\cdot) = \text{tr}_E[V(\cdot)V^\dagger]$. Then, we have

$$T(\mathcal{E}) = \sup_{\sigma} \{H(\sigma_X) - H(\mathcal{E}(\sigma_X))\} = \sup_{\sigma} H(E | X')_{V \sigma V^\dagger}. \quad (4.9)$$

That is, the thermodynamic capacity characterizes by how much the channel is capable of reducing the entropy of its input, or equivalently, how much entropy the channel is capable of dumping into the environment when conditioned on the output. We note that the quantity $-T(\mathcal{E})$ has previously been studied in the information theory literature as the entropy gain of quantum channels [42–49]. Our work can be seen as giving a precise operational interpretation to this quantity.

4.3 Optimality

Here, we show that any universal implementation that obeys our stated conditions in Section 4.1 must necessarily consume an amount of work that is lower bounded by the thermodynamic capacity. That is, any universal implementation that consumes an amount of work equal to the thermodynamic capacity is optimal. This lower bound is simple to prove, because a universal implementation of a process must necessarily be a good implementation for any individual i.i.d. input state, a situation where the optimal work cost is known [16]. Furthermore, any scheme that satisfies the requirements of Section 4 at work cost w per copy counted with standard battery states of Ref. [16], has an effective process $\mathcal{T}_{X^n \rightarrow X'^n}$ on the systems that obeys $\mathcal{T}(\Gamma_X^{\otimes n}) \leq e^{nw} \Gamma_{X'}^{\otimes n}$. This is because any thermal operation is in particular a Gibbs-preserving map, and the work cost is characterized by Proposition 3.1. The following shows that for any such implementation, the work consumed w per copy cannot be less than the thermodynamic capacity of the process.

Proposition 4.2. *Let $\varepsilon > 0$, $\Gamma_X, \Gamma_{X'} > 0$, $\mathcal{E}_{X \rightarrow X'}$ a completely positive, trace-preserving map, and $\mathcal{T}_{X^n \rightarrow X'^n}$ a completely positive, trace non-increasing map such that we have $\|\mathcal{T} - \mathcal{E}^{\otimes n}\|_\diamond / 2 \leq \varepsilon$. For $w \in \mathbb{R}$ such that $\mathcal{T}_{X^n \rightarrow X'^n}(\Gamma_X^{\otimes n}) \leq e^{nw} \Gamma_{X'}^{\otimes n}$, we have in the limit $n \rightarrow \infty$ that $w \geq T(\mathcal{E})$.*

Proof. Let \mathcal{T} with $\frac{1}{2}\|\mathcal{E} - \mathcal{T}\|_\diamond \leq \varepsilon$, σ_X be a quantum state, and $|\sigma\rangle_{XR_X} = \sigma_X^{1/2} |\Phi\rangle_{X:R_X}$. Then, by definition of the diamond norm it must hold that $D(\mathcal{E}(\sigma_{XR_X}), \mathcal{T}(\sigma_{XR_X})) \leq \varepsilon$, which implies that $P(\mathcal{E}(\sigma_{XR_X}), \mathcal{T}(\sigma_{XR_X})) \leq \sqrt{2\varepsilon}$. We have that \mathcal{T} is a valid optimization candidate for the definition of the coherent relative entropy and thus

$$-\hat{D}_{X^n \rightarrow X'^n}^{\sqrt{2\varepsilon}}(\mathcal{E}_{X \rightarrow X'}^{\otimes n}(\sigma_{XR_X}^{\otimes n}) \| \Gamma_X^{\otimes n}, \Gamma_{X'}^{\otimes n}) \leq nw. \quad (4.10)$$

For $n \rightarrow \infty$, we can employ the asymptotic equipartition of the coherent relative entropy (3.4) to see that

$$D(\mathcal{E}(\sigma_X) \| \Gamma_{X'}) - D(\sigma_X \| \Gamma_X) \leq w. \quad (4.11)$$

Since this inequality holds for all σ_X , we deduce that $T(\mathcal{E}) \leq w$. ■

5 Construction #1: Trivial Hamiltonians

5.1 Statement and proof sketch

Instead of constructing explicitly an implementation that satisfies the requirements of Section 4, one might hope that the implementation could be given implicitly as the solution of a semi-definite program representing an entropy measure. This proof idea was indeed exploited in other contexts in Refs. [23, 50]. Here, we define the one-shot entropy-like quantity

$$W_{X \rightarrow X'}^\varepsilon(\mathcal{E}_{X \rightarrow X'} \| \Gamma_X, \Gamma_{X'}) = \min_{\substack{\mathcal{T}(\Gamma_X) \leq e^w \Gamma_{X'} \\ \frac{1}{2}\|\mathcal{T} - \mathcal{E}\|_\diamond \leq \varepsilon}} w, \quad (5.1)$$

where $\mathcal{T}_{X \rightarrow X'}$ ranges over all trace non-increasing, completely positive maps. The proof strategy would then be to relate this entropy measure to the coherent relative entropy, and to exploit known properties of the latter in the i.i.d. regime to provide an upper bound to the expression

$$\frac{1}{n} W_{X^n \rightarrow X'^n}^\varepsilon(\mathcal{E}_{X^n \rightarrow X'^n}^{\otimes n} \| \Gamma_X^{\otimes n}, \Gamma_{X'}^{\otimes n}) . \quad (5.2)$$

Should this upper bound behave like $T(\mathcal{E})$ to leading order, then the \mathcal{T} equal to the optimal solution to (5.1) defines an implementation in terms of Gibbs-preserving maps thanks to Proposition 3.1. It turns out that this proof strategy works well in the special case of trivial Hamiltonians, but fails in the general case.

The core technical statement that underlies our Construction #1 is summarized in the following theorem.

Theorem 5.1. *Let $\mathcal{E}_{X \rightarrow X'}$ be a completely positive, trace-preserving map, and $\varepsilon > 0$. Then we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{X^n \rightarrow X'^n}^\varepsilon(\mathcal{E}_{X^n \rightarrow X'^n}^{\otimes n} \| \mathbb{1}_{X^n}, \mathbb{1}_{X'^n}) = T(\mathcal{E}) , \quad (5.3)$$

where $T(\mathcal{E}) = \max_{\sigma_X} \{H(\sigma_X) - H(\mathcal{E}(\sigma_X))\}$.

This implementation is constructed by taking the implicit optimal solution $\mathcal{T}_{X^n \rightarrow X'^n}$ in the semi-definite program (5.1) for $\frac{1}{n} W_{X^n \rightarrow X'^n}^\varepsilon(\mathcal{E}_{X^n \rightarrow X'^n}^{\otimes n} \| \mathbb{1}_{X^n}, \mathbb{1}_{X'^n})$, and using Proposition 3.1 to construct an associated Gibbs-preserving map acting on battery states via (3.2). In summary, for any $\delta' > 0$, for n large enough and choosing any m, m' such that $m - m' \leq nT(\mathcal{E}) + \delta'$, the full implementation map in terms of $\mathcal{T}_{X^n \rightarrow X'^n}$ becomes

$$\Phi_{X^n W \rightarrow X'^n W}(\cdot) = \mathcal{T}_{X^n \rightarrow X'^n}(\text{tr}_W[P_W^m(\cdot)]) \otimes \tau_W^{m'} . \quad (5.4)$$

We emphasise that Theorem 5.1 exactly covers the entropy gain of quantum channels as studied in [42–49].

Proof of Theorem 5.1. By using the post-selection technique (Theorem 2.1) and recalling that the fixed-input state case is given by the coherent relative entropy, we find

$$W_{X^n \rightarrow X'^n}^\varepsilon(\mathcal{E}_{X^n \rightarrow X'^n}^{\otimes n} \| \mathbb{1}_{X^n}, \mathbb{1}_{X'^n}) \leq -\hat{D}_{X^n \rightarrow X'^n}^{\varepsilon/\text{poly}(n)}(\mathcal{E}_{X^n \rightarrow X'^n}^{\otimes n}(\zeta_{X^n R_X^n}) \| \mathbb{1}_{X^n}, \mathbb{1}_{X'^n}) . \quad (5.5)$$

In the case of trivial Hamiltonians, the coherent relative entropy reduces to the smooth max-entropy (cf. [16, Props. 28 and 26] and also Ref. [51]). More precisely, we have

$$\hat{D}_{X \rightarrow X'}^\varepsilon(\rho_{X' R_X} \| \mathbb{1}_X, \mathbb{1}_{X'}) \geq -H_{\max}^{c\varepsilon^\alpha}(E | X')_\rho + g(\varepsilon) , \quad (5.6)$$

where $|\rho\rangle_{X' R_X E}$ is a pure state, where $c > 0$, $0 < \alpha < 1$, $g(\varepsilon)$ are universal and do not depend on the

state or the dimensions of the systems, and the smooth max-entropy is defined as

$$H_{\max}^{\varepsilon}(E | X')_{\rho} = \min_{P(\hat{\rho}, \rho) \leq \varepsilon} H_{\max}(E | X')_{\hat{\rho}} ; \quad (5.7)$$

$$H_{\max}(E | X')_{\rho} = \max_{0 \leq \omega_{X'} \leq \mathbb{1}} \ln \|\hat{\rho}_{EX'}^{1/2} \omega_{X'}^{1/2}\|_1^2 . \quad (5.8)$$

Thus, we have

$$(5.5) \leq H_{\max}^{\varepsilon/\text{poly}(n)}(E^n | X'^n)_{\rho} + g(\varepsilon) , \quad (5.9)$$

where $\rho_{X'^n E^n} = V_{X \rightarrow X'E}^{\otimes n} \zeta_{X^n}(V^{\dagger})^{\otimes n} = \int d\sigma (V \sigma V^{\dagger})^{\otimes n}$ and $V_{X \rightarrow X'E}$ is a Stinespring dilation isometry of $\mathcal{E}_{X \rightarrow X'}$ as $\mathcal{E}_{X \rightarrow X'}(\cdot) = \text{tr}_E [V_{X \rightarrow X'E}(\cdot) V^{\dagger}]$. At this point we invoke two facts. First, note that the de Finetti state can be written as a mixture of only $\text{poly}(n)$ i.i.d. states, instead of a continuous average (Theorem 2.1): There exists a set $\{\sigma_i\}$ of at most $\text{poly}(n)$ states and a distribution $\{p_i\}$ such that $\zeta_{X^n} = \sum_i p_i \sigma_i^{\otimes n}$. Second, we invoke the property that the conditional max-entropy is quasi-convex up to a penalty term, namely, that the conditional max-entropy of $\sum_i p_i \rho_i$ is less than or equal to the maximum over the set of max-entropies corresponding to each ρ_i , plus a term proportional to the logarithm of the number of terms in the sum [52, Lemma 11]. Hence, with $\rho_i = V \sigma_i V^{\dagger}$, we get

$$(5.9) \leq \max_i H_{\max}^{\varepsilon/\text{poly}(n)}(E^n | X'^n)_{\rho_i^{\otimes n}} + \ln(\text{poly}(n)) + g(\varepsilon) . \quad (5.10)$$

Now, we are in business because the max-entropy is evaluated on an i.i.d. state, and we know that it asymptotically goes to the von Neumann entropy in this regime [53]. Also, $\lim_{n \rightarrow \infty} (1/n) \{ \ln(\text{poly}(n)) + g(\varepsilon) \} = 0$ and hence

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} W_{X^n \rightarrow X'^n}^{\varepsilon}(\mathcal{E}_{X \rightarrow X'}^{\otimes n} \parallel \mathbb{1}_{X^n}, \mathbb{1}_{X'^n}) &\leq \max_i H(E | X')_{\rho_i} \\ &= \max_i \{ H(\sigma_i) - H(\mathcal{E}(\sigma_i)) \} \\ &\leq \max_{\sigma} \{ H(\sigma) - H(\mathcal{E}(\sigma)) \} \\ &= T(\mathcal{E}) \end{aligned} \quad (5.11)$$

noting that $H(E | X') = H(EX') - H(X') = H(X) - H(X')$. ■

5.2 Challenges for extension to non-trivial Hamiltonians

Naturally, one might ask whether it is possible to extend this proof to the case of non-trivial Γ operators. Interestingly, this is not possible, at least not in a naive way. The problem is that we need a quasi-convexity property of the form

$$\begin{aligned} -\hat{D}_{X \rightarrow X'}^{\varepsilon}(\mathcal{E}_{X \rightarrow X'}(\sigma_{XR_X}) \parallel \Gamma_X, \Gamma_{X'}) \\ \stackrel{?}{\leq} \max_i (-\hat{D}_{X \rightarrow X'}^{\varepsilon}(\mathcal{E}_{X \rightarrow X'}(\sigma_{XR_X}^i) \parallel \Gamma_X, \Gamma_{X'})) + (\text{penalty}) , \end{aligned} \quad (5.12)$$

where $\sigma_X = \sum p_i \sigma_X^i$ and $|\sigma\rangle_{XR} = \sigma_X^{1/2} |\Phi\rangle_{X:R_X}$, $|\sigma^i\rangle_{XR} = (\sigma_X^i)^{1/2} |\Phi\rangle_{X:R_X}$, and where the (penalty) term scales in a favourable way in n , say of order $\ln(\text{poly}(M))$ where M is the number of terms in the convex decomposition as for the max-entropy. In fact, Eq. (5.12) is false, as can be shown using an explicit counterexample on a two-level system which we present below. As this example is based on physical reasons, the coherent relative entropy is not even approximately quasi-convex. We note that a priori we cannot rule out a quasi-convexity property that might have a penalty term that depends on properties of the Γ operators, yet such a term is likely to scale unfavourably with n .

Our example is as follows. Consider a two-level system with a Hamiltonian H with energy levels $|0\rangle, |1\rangle$ at corresponding energies $E_0 = 0$ and $E_1 > 0$. The corresponding Γ operator is $\Gamma = g_0|0\rangle\langle 0| + g_1|1\rangle\langle 1|$ with $g_0 = 1$, $g_1 = e^{-\beta E_1}$. Consider the process consisting in erasing the input and creating the output state $|+\rangle$, where we define $|\pm\rangle = [|0\rangle \pm |1\rangle]/\sqrt{2}$. That is, we consider the process $\mathcal{E}(\cdot) = \text{tr}[\cdot] |+\rangle\langle +|$. Suppose the input state is maximally mixed, $\sigma = \mathbb{1}/2$, such that $\rho_{X'R_X} = |+\rangle\langle +|_{X'} \otimes \mathbb{1}_{R_X}/2$. If $E_0 = 0$ and $E_1 \rightarrow \infty$, then this process requires a lot of work; intuitively, with probability $1/2$ we start in the ground state $|0\rangle$ and need to prepare the output state $|+\rangle$ which has high energy.

For $\varepsilon = 0$, we can see this because the input state is full rank, hence $\mathcal{T} = \mathcal{E}$; then $\mathcal{E}(\Gamma) = \text{tr}[\Gamma] |+\rangle\langle +|$ and the smallest α such that $\mathcal{E}(\Gamma) \leq \alpha \Gamma$ is given by

$$\begin{aligned} \alpha / \text{tr}[\Gamma] &= \|\Gamma^{-1/2} |+\rangle\langle +| \Gamma^{-1/2}\|_\infty = \langle + | \Gamma^{-1} | + \rangle = (g_0^{-1} + g_1^{-1})/2 \\ &= (1 + e^{\beta E_1})/2 \geq e^{\beta E_1}/2. \end{aligned} \quad (5.13)$$

Noting that $\text{tr}[\Gamma] \geq 1$, we have $\alpha \geq e^{\beta E_1}/2$, and hence the energy cost of the transformation $\mathbb{1}/2 \rightarrow |+\rangle$ is

$$\text{energy cost} = -\beta^{-1} \hat{D}_{X \rightarrow X'}(\mathcal{E}_{X \rightarrow X'}(\sigma_{X R_X}) \|\Gamma, \Gamma) = \beta^{-1} \ln \alpha \geq E_1 - \beta^{-1} \ln(2). \quad (5.14)$$

Clearly, this work cost can become arbitrarily large if $E_1 \rightarrow \infty$. On the other hand, we can perform the transformation $|+\rangle \rightarrow |+\rangle$ obviously at no work cost; similarly, $|-\rangle \rightarrow |+\rangle$ can be carried out by letting the system time-evolve under its own Hamiltonian for exactly the time interval required to pick up a relative phase (-1) between the $|0\rangle$ and $|1\rangle$ states. This also costs no work because it is a unitary operation that commutes with the Hamiltonian. We thus have our counter-example to the quasi-convexity of the coherent relative entropy. The transformation $\mathbb{1}/2 \rightarrow |+\rangle$ is very hard, but the individual transformations $|\pm\rangle \rightarrow |+\rangle$ are trivial, noting that $\mathbb{1}/2 = (1/2)|+\rangle\langle +| + (1/2)|-\rangle\langle -|$.

We show in [Appendix D](#) how to make the above claim robust against an accuracy tolerance $\varepsilon \geq 0$.

6 Construction #2: Gibbs-preserving maps

6.1 Statement and proof sketch

Here, we present a general construction of a universal implementation of an i.i.d. process using Gibbs-preserving maps according to the requirements of [Section 4.1](#). The idea is to explicitly

construct an implementation using a novel notion of quantum typicality. We introduce notions of quantum typicality that apply to quantum processes and universally capture regions of the Hilbert space where the conditional entropy (respectively the relative entropy difference) has a given value. This generalizes existing notions of typical projectors to a quantum typical operator that applies to bipartite states, is relative to a Γ operator, and universal.

The main result behind the construction in this section is the following theorem.

Theorem 6.1. *Let $\Gamma_X, \Gamma_{X'} > 0$, $\mathcal{E}_{X \rightarrow X'}$ be a completely positive, trace-preserving map, and $\varepsilon > 0$. Then, for $\delta > 0$ and $n \in \mathbb{N}$ large enough there exists a completely positive map $\mathcal{T}_{X^n \rightarrow X'^n}$ such that:*

- (i) $\mathcal{T}_{X^n \rightarrow X'^n}$ is trace non-increasing;
- (ii) $\|\mathcal{T}_{X^n \rightarrow X'^n} - \mathcal{E}_{X \rightarrow X'}^{\otimes n}\|_{\diamond} \leq \varepsilon$;
- (iii) $\mathcal{T}_{X^n \rightarrow X'^n}(\Gamma_X^{\otimes n}) \leq e^{n[T(\mathcal{E}) + 4\delta + n^{-1} \ln(\text{poly}(n))]} \Gamma_{X'}^{\otimes n}$.

Note that we have $n^{-1} \ln(\text{poly}(n)) \rightarrow 0$ as $n \rightarrow \infty$, and that we can take $\delta \rightarrow 0$ after taking $n \rightarrow \infty$. Thanks to [Proposition 3.1](#), the mapping $\mathcal{T}_{X^n \rightarrow X'^n}$ defines an implementation of the i.i.d. process $\mathcal{E}_{X \rightarrow X'}^{\otimes n}$ in terms of Gibbs-preserving maps and a battery, whose work cost rate is given to leading order by the thermodynamic capacity $T(\mathcal{E})$ after taking $\delta \rightarrow 0$.

As for Construction #1, the full Gibbs-preserving map implementing the required process is assembled in two steps, first constructing the map $\mathcal{T}_{X^n \rightarrow X'^n}$ in [Theorem 6.1](#) and then using [Proposition 3.1](#) to obtain the full Gibbs-preserving map. Let $V_{X \rightarrow X'E}$ be a Stinespring dilation isometry of $\mathcal{E}_{X \rightarrow X'}$. For $\delta > 0$, we introduce a universal conditional and relative typical smoothing operator $M_{E^n X'^n}^{x, \delta}$ (see later [Definition 6.1](#) and [Proposition 6.1](#)) with $x = -nT(\mathcal{E})$ and relative to $\Gamma_{X'E} \equiv V \Gamma_X V^\dagger$ and $\Gamma_{X'}$. The map $\mathcal{T}_{X^n \rightarrow X'^n}$ is then constructed as

$$\mathcal{T}_{X^n \rightarrow X'^n}(\cdot) = \text{tr}_{E^n} \left[M_{E^n X'^n}^{x, \delta} V_{X \rightarrow X'E}^{\otimes n} (\cdot) V_{X \leftarrow X'E}^{\dagger \otimes n} M_{E^n X'^n}^{x, \delta \dagger} \right]. \quad (6.1)$$

Finally, we employ [Proposition 3.1](#) to construct an associated Gibbs-preserving map acting on battery states via [\(3.2\)](#). For any $\delta' > 0$, for n large enough and choosing any m, m' such that $m - m' \leq nT(\mathcal{E}) + 4\delta + n^{-1} \ln \text{poly}(n) + \delta'$, the full implementation map in terms of $\mathcal{T}_{X^n \rightarrow X'^n}$ becomes

$$\Phi_{X^n W \rightarrow X'^n W}(\cdot) = \mathcal{T}_{X^n \rightarrow X'^n}(\text{tr}_W[P_W^m(\cdot)]) \otimes \tau_W^{m'}. \quad (6.2)$$

6.2 Construction via universal conditional and relative typicality

The main ingredient of our proof is a notion of a universal conditional and relative typical smoothing operator that enables us to discard events that are very unlikely to appear in the process while accounting for how much they contribute to the overall work cost. This operator is inspired by similar constructions in Refs. [\[54, 55\]](#). However, in addition to being “relative” as in [\[54\]](#) our smoothing operator is also simultaneously “conditional” and “universal”.

Definition 6.1. Let $\Gamma_{AB}, \Gamma'_B \geq 0$ and $x \in \mathbb{R}$. A *universal conditional and relative typical smoothing operator* $M_{A^n B^n}^{x, \delta}$ with parameter $\delta > 0$ is an operator on $A^n B^n$ that satisfies the following conditions:

- (i) $(M_{A^n B^n}^{x, \delta})^\dagger M_{A^n B^n}^{x, \delta} \leq \mathbb{1}$;
- (ii) There exists $\xi > 0$ independent of n with the following property: For any pure state $|\rho\rangle_{ABR}$ with ρ_{AB} (respectively ρ_B) in the support of Γ_{AB} (respectively Γ'_B) and such that $D(\rho_{AB} \parallel \Gamma_{AB}) - D(\rho_B \parallel \Gamma'_B) \geq x$, it holds that

$$\text{Re} \left\{ \langle \rho |_{ABR}^{\otimes n} M_{A^n B^n}^{x, \delta} | \rho \rangle_{ABR}^{\otimes n} \right\} \geq 1 - \text{poly}(n) \exp(-n\xi) ; \quad (6.3)$$

- (iii) $\text{tr}_{A^n} \left[M_{A^n B^n}^{x, \delta} \Gamma_{AB}^{\otimes n} (M_{A^n B^n}^{x, \delta})^\dagger \right] \leq \text{poly}(n) e^{-n(x-4\delta)} \Gamma_B'^{\otimes n}$.

Note that the smoothing operator is defined as a general operator of norm bounded by one, as opposed to the usual definition of typical subspaces or typical projectors. The main reason is that it is not known to us in general if such an object can be chosen to be a projector. By using the real part in Point (ii) above, we ensure that a process that applies the operator $M_{A^n B^n}^{x, \delta}$ preserves coherences when it is applied to a superposition of several states $\{|\rho\rangle_{ABR}^{\otimes n}\}$. This property would not have been ensured if instead, we had merely asserted that $M_{A^n B^n}^{x, \delta} |\rho\rangle_{ABR}^{\otimes n}$ and $|\rho\rangle_{ABR}^{\otimes n}$ have high absolute value overlap or are close in fidelity. If $M_{A^n B^n}^{x, \delta}$ is a projector then the expression reduces to $\text{tr}(M_{A^n B^n}^{x, \delta} \rho)$ as one usually considers for projectors on typical subspaces.

The core technical statement of Construction #2 is to show the existence of a universal conditional and relative smoothing operator, which is as follows.

Proposition 6.1. Let $\Gamma_{AB}, \Gamma'_B \geq 0$, $x \in \mathbb{R}$, as well as $n \in \mathbb{N}$ and $\delta > 0$. There exists a universal conditional and relative typical smoothing operator $M_{A^n B^n}^{x, \delta}$ that is furthermore permutation-invariant. Moreover, if $[\Gamma_{AB}, \mathbb{1}_A \otimes \Gamma'_B] = 0$, then $M_{A^n B^n}^{x, \delta}$ can be chosen to be a projector satisfying $[M_{A^n B^n}^{x, \delta}, \Gamma_B'^{\otimes n}] = 0$ and $[M_{A^n B^n}^{x, \delta}, \Gamma_{AB}^{\otimes n}] = 0$.

In the following, we present the proof of [Theorem 6.1](#) based on the existence of such the smoothing operator from [Proposition 6.1](#). The more technical proof of [Proposition 6.1](#) is then given in [Section 6.3](#).

Proof of Theorem 6.1. Let $V_{X \rightarrow X'E}$ be a Stinespring dilation of $\mathcal{E}_{X \rightarrow X'}$ into an environment system $E \simeq X \otimes X'$. For $n \in \mathbb{N}$ we need to find a suitable candidate implementation $\mathcal{T}_{X^n \rightarrow X'^n}$. Let

$$x = -\max_{\sigma_X} \left\{ D(\mathcal{E}(\sigma_X) \parallel \Gamma_{X'}) - D(\sigma_X \parallel \Gamma_X) \right\} = -T(\mathcal{E}) . \quad (6.4)$$

For any $\delta > 0$ let $M_{E^n X'^n}^{x, \delta}$ be the operator constructed by [Proposition 6.1](#), with the system E playing the role of the system A , with $V_{X \rightarrow X'E} \Gamma_X V_{X \leftarrow X'E}^\dagger$ as Γ_{AB} and with $\Gamma_{X'}$ as Γ'_B . Now, define

$$\mathcal{T}_{X^n \rightarrow X'^n}(\cdot) = \text{tr}_{E^n} \left[M_{E^n X'^n}^{x, \delta} V_{X \rightarrow X'E}^{\otimes n} (\cdot) (V_{X \leftarrow X'E}^\dagger)^{\otimes n} (M_{E^n X'^n}^{x, \delta})^\dagger \right] \quad (6.5)$$

noting that $\mathcal{T}_{X^n \rightarrow X^m}$ is trace non-increasing by construction thanks to Property (i) of Definition 6.1.

Let $|\sigma\rangle_{XR_X}$ be any pure state, and define $|\rho\rangle_{X'ER_X} = V_{X \rightarrow X'E} |\sigma\rangle_{XR_X}$. By construction, $D(\rho_{EX'} \| (V_{X \rightarrow X'E} \Gamma_X V^\dagger)) - D(\rho_{X'} \| \Gamma_{X'}) = D(\sigma_X \| \Gamma_X) - D(\mathcal{E}(\sigma_X) \| \Gamma_{X'}) \geq x$. Then Property (ii) of Proposition 6.1 tells us that there exists a $\xi > 0$ independent of both ρ and n such that

$$\operatorname{Re} \left\{ \langle \rho |_{X'ER_X}^{\otimes n} M_{E^n X^m}^{x, \delta} | \rho \rangle_{X'ER_X}^{\otimes n} \right\} \geq 1 - \operatorname{poly}(n) \exp(-n\xi). \quad (6.6)$$

The conditions of Proposition 2.3 are fulfilled, with $W_{X^n \rightarrow X^m E^n} = M_{A^n B^n}^{x, \delta} V_{X \rightarrow X'E}^{\otimes n}$, thanks furthermore to the fact that $M_{E^n X^m}^{x, \delta}$ is permutation-invariant as guaranteed by Proposition 6.1. Hence, we have

$$\frac{1}{2} \left\| \mathcal{T}_{X^n \rightarrow X^m} - \mathcal{E}_{X \rightarrow X'}^{\otimes n} \right\|_\diamond \leq \operatorname{poly}(n) \exp(-n\xi/2). \quad (6.7)$$

For $n \in \mathbb{N}$ large enough this becomes smaller than any fixed $\varepsilon > 0$. Furthermore, by Property (iii) of Definition 6.1, we have that

$$\begin{aligned} \mathcal{T}_{X^n \rightarrow X^m}(\Gamma_X^{\otimes n}) &= \operatorname{tr}_{E^n} [M_{E^n X^m}^{x, \delta} (V_{X \rightarrow X'E} \Gamma_X V_{X \leftarrow X'E}^\dagger)^{\otimes n} (M_{E^n X^m}^{x, \delta})^\dagger] \\ &\leq \operatorname{poly}(n) e^{-n(x-4\delta)} \Gamma_{X'}^{\otimes n} \end{aligned} \quad (6.8)$$

as required. \blacksquare

6.3 Universal conditional and relative typical smoothing operator

We now turn to the proof of Proposition 6.1, giving an explicit construction of a universal conditional and relative typical smoothing operator. As the proof of Proposition 6.1 is quite lengthy, it can be instructive to consider a simpler version of our typical smoothing operator which applies in the case where the Hamiltonians are trivial. We carry out this analysis in Appendix E.

Proof of Proposition 6.1. First, we claim that we can assume $\Gamma_{AB} > 0$ and $\Gamma'_B > 0$ without loss of generality. Indeed, if either operator is not positive definite, then we can first construct the operator $\tilde{M}_{A^n B^n}^{x, \delta}$ associated with modified operators $\tilde{\Gamma}_{AB} > 0$ and $\tilde{\Gamma}'_B > 0$ where all the zero eigenvalues of Γ_{AB} and Γ'_B are replaced by some arbitrary fixed strictly positive constant (e.g., one); we can then set $M_{A^n B^n}^{x, \delta} = P_{B^n}^{\Gamma'} \tilde{M}_{A^n B^n}^{x, \delta} P_{A^n B^n}^{\Gamma}$, where $P_{A^n B^n}^{\Gamma}$ (respectively $P_{B^n}^{\Gamma'}$) is the projector onto the support of $\Gamma_{AB}^{\otimes n}$ (respectively Γ'_B). The operator $M_{A^n B^n}^{x, \delta}$ constructed in this way satisfies all of the required properties. For the remainder of this proof we thus assume that $\Gamma_{AB} > 0$ and $\Gamma'_B > 0$.

Let $\{R_{A^n B^n}^k\}$ be the POVM constructed by Proposition 2.2 for $H_{AB} = -\ln(\Gamma_{AB})$. Similarly, let $\{S_{B^n}^\ell\}$ be the corresponding POVM constructed in Proposition 2.2 for $H'_B = -\ln(\Gamma'_B)$. Also, as before, we denote by $\Pi_{A^n B^n}^\lambda$ and by $\Pi_{B^n}^\mu$ the projectors on the Schur-Weyl blocks labelled by the Young diagrams $\lambda \in \operatorname{Young}(d_A d_B, n)$ and $\mu \in \operatorname{Young}(d_B, n)$. Let

$$M_{A^n B^n}^{x, \delta} = \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k. \quad (6.9)$$

Note that $[S_{B^n}^\ell, \Pi_{B^n}^\mu] = 0$ because $S_{B^n}^\ell$ is permutation-invariant, and $[\mathbb{1}_{A^n} \otimes S_{B^n}^\ell, \Pi_{A^n B^n}^\lambda] = 0$ because $\mathbb{1}_{A^n} \otimes S_{B^n}^\ell$ is permutation-invariant. Recall also that $[\mathbb{1}_{A^n} \otimes \Pi_{B^n}^\mu, \Pi_{A^n B^n}^\lambda] = 0$ for the same reason. The operator $M_{A^n B^n}^{x, \delta}$ is permutation-invariant by construction. Then, we have

$$\begin{aligned}
M_{A^n B^n}^{x, \delta} M_{A^n B^n}^{x, \delta} &= \sum_{\substack{k, \ell, \lambda, \mu, \\ k', \ell', \lambda', \mu' : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta \\ k' - \bar{H}(\lambda') - \ell' + \bar{H}(\mu') \geq x - 4\delta}} R_{A^n B^n}^k \Pi_{A^n B^n}^\lambda \Pi_{B^n}^\mu S_{B^n}^\ell S_{B^n}^{\ell'} \Pi_{B^n}^{\mu'} \Pi_{A^n B^n}^{\lambda'} R_{A^n B^n}^{k'} \\
&= \sum_{\substack{k, k', \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta \\ k' - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} R_{A^n B^n}^k (\Pi_{A^n B^n}^\lambda \Pi_{B^n}^\mu S_{B^n}^\ell) R_{A^n B^n}^{k'} \\
&= \sum_{k, k'} R_{A^n B^n}^k \left(\sum_{\substack{\ell, \lambda, \mu \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta \\ k' - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} \Pi_{A^n B^n}^\lambda \Pi_{B^n}^\mu S_{B^n}^\ell \right) R_{A^n B^n}^{k'} \\
&\leq \sum_{k, k'} R_{A^n B^n}^k R_{A^n B^n}^{k'} \\
&= \sum_k R_{A^n B^n}^k = \mathbb{1}_{A^n B^n} \tag{6.10}
\end{aligned}$$

recalling that the operators $(\Pi_{A^n B^n}^\lambda, \Pi_{B^n}^\mu, S_{B^n}^\ell)$ form a commuting set of projectors, and where in the third line the inner sum is taken to be the zero operator if no triplet (ℓ, λ, μ) satisfies the given constraints. This shows Property (i).

Now, consider any state $|\rho\rangle_{ABR}$, where R is any reference system, and assume that $D(\rho_{AB} \parallel \Gamma_{AB}) - D(\rho_B \parallel \Gamma'_B) \geq x$. Rewrite this condition as

$$x \leq -H(\rho_{AB}) - \text{tr}[\rho_{AB} \ln \Gamma_{AB}] + H(\rho_B) + \text{tr}[\rho_B \ln \Gamma'_B] . \tag{6.11}$$

We write

$$\begin{aligned}
\langle \rho |_{ABR}^{\otimes n} M_{A^n B^n}^{x, \delta} | \rho \rangle_{ABR}^{\otimes n} &= \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} \langle \rho |_{ABR}^{\otimes n} (S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k) | \rho \rangle_{ABR}^{\otimes n} \\
&= \blacksquare_1 + \blacksquare_2 , \tag{6.12}
\end{aligned}$$

where we define

$$\blacksquare_1 = \sum_{\substack{k, \ell, \lambda, \mu : \\ k \geq -\text{tr}[\rho_{AB} \ln \Gamma_{AB}] - \delta \\ \bar{H}(\lambda) \leq H(\rho_{AB}) + \delta \\ \ell \leq -\text{tr}[\rho_B \ln \Gamma'_B] + \delta \\ \bar{H}(\mu) \geq H(\rho_B) - \delta}} \langle \rho |_{ABR}^{\otimes n} (S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k) | \rho \rangle_{ABR}^{\otimes n} ; \tag{6.13a}$$

$$\blacksquare_2 = \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta \text{ AND} \\ [k < -\text{tr}[\rho_{AB} \ln \Gamma_{AB}] - \delta \text{ OR} \\ \bar{H}(\lambda) > H(\rho_{AB}) + \delta \text{ OR} \\ \ell > -\text{tr}[\rho_B \ln \Gamma'_B] + \delta \text{ OR} \\ \bar{H}(\mu) < H(\rho_B) - \delta]}} \langle \rho|_{ABR}^{\otimes n} (S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k) | \rho \rangle_{ABR}^{\otimes n} , \quad (6.13b)$$

further noting that the conditions in the sum defining \blacksquare_1 indeed imply that $k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq -\text{tr}[\rho_{AB} \ln \Gamma_{AB}] - H(\rho_{AB}) + \text{tr}[\rho_B \ln \Gamma'_B] + H(\rho_B) - 4\delta \geq x - 4\delta$. We first consider \blacksquare_1 . Define the projectors

$$X_1 = \sum_{k \geq -\text{tr}[\rho_{AB} \ln \Gamma_{AB}] - \delta} R_{A^n B^n}^k ; \quad X_1^\perp = \mathbb{1} - X_1 ; \quad (6.14a)$$

$$X_2 = \sum_{\bar{H}(\lambda) \leq H(\rho_{AB}) + \delta} \Pi_{A^n B^n}^\lambda ; \quad X_2^\perp = \mathbb{1} - X_2 ; \quad (6.14b)$$

$$X_3 = \sum_{\bar{H}(\mu) \geq H(\rho_B) - \delta} \Pi_{B^n}^\mu ; \quad X_3^\perp = \mathbb{1} - X_3 ; \quad (6.14c)$$

$$X_4 = \sum_{\ell \leq -\text{tr}[\rho_B \ln \Gamma'_B] + \delta} S_{B^n}^\ell ; \quad X_4^\perp = \mathbb{1} - X_4 , \quad (6.14d)$$

and observe that

$$\text{Re}\{ \blacksquare_1 \} = \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} (X_4 X_3 X_2 X_1) | \rho \rangle_{ABR}^{\otimes n} \} . \quad (6.15)$$

Thanks to [Proposition 2.2](#), we have $\|X_1^\perp | \rho \rangle_{ABR}^{\otimes n}\| \leq 2 \exp(-n\eta/2)$, recalling that $\|P|\psi\rangle\| = \sqrt{\text{tr}[P\psi]}$, and hence

$$\begin{aligned} & \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 X_2 X_1 | \rho \rangle_{ABR}^{\otimes n} \} \\ &= \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 X_2 | \rho \rangle_{ABR}^{\otimes n} \} - \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 X_2 X_1^\perp | \rho \rangle_{ABR}^{\otimes n} \} \\ &\geq \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 X_2 | \rho \rangle_{ABR}^{\otimes n} \} - 2 \exp(-n\eta/2) \end{aligned} \quad (6.16)$$

using Cauchy-Schwarz to assert that $\text{Re}(\langle \chi | \psi \rangle) \leq |\langle \chi | \psi \rangle| \leq \|\chi\| \|\psi\|$. Similarly, using [Proposition 2.1](#), we have $\|X_2^\perp | \rho \rangle_{ABR}^{\otimes n}\| \leq \text{poly}(n) \exp(-n\eta/2)$. Also, we have $\|X_3^\perp | \rho \rangle_{ABR}^{\otimes n}\| \leq \text{poly}(n) \exp(-n\eta/2)$, and $\|X_4^\perp | \rho \rangle_{ABR}^{\otimes n}\| \leq 2 \exp(-n\eta/2)$, yielding

$$\text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 X_2 | \rho \rangle_{ABR}^{\otimes n} \} \geq \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 | \rho \rangle_{ABR}^{\otimes n} \} - \text{poly}(n) \exp(-n\eta/2) ; \quad (6.17)$$

$$\text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 X_3 | \rho \rangle_{ABR}^{\otimes n} \} \geq \text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 | \rho \rangle_{ABR}^{\otimes n} \} - \text{poly}(n) \exp(-n\eta/2) ; \quad (6.18)$$

$$\text{Re}\{ \langle \rho|_{ABR}^{\otimes n} X_4 | \rho \rangle_{ABR}^{\otimes n} \} \geq 1 - 2 \exp(-n\eta/2) . \quad (6.19)$$

We take all these η 's to be the same, by choosing if necessary the minimum of the four possibly different η s. Hence, we have

$$\text{Re}\{ \blacksquare_1 \} \geq 1 - \text{poly}(n) \exp(-n\eta/2) . \quad (6.20)$$

Now we consider the term \blacksquare_2 . We know that

$$\|R_{A^n B^n}^k |\rho\rangle_{ABR}^{\otimes n}\| \leq \exp(-n\eta/2) \quad \text{if } k < -\text{tr}[\rho_{AB} \ln \Gamma_{AB}] - \delta; \quad (6.21a)$$

$$\|\Pi_{A^n B^n}^\lambda |\rho\rangle_{ABR}^{\otimes n}\| \leq \text{poly}(n) \exp(-n\eta/2) \quad \text{if } \bar{H}(\lambda) > H(\rho_{AB}) + \delta; \quad (6.21b)$$

$$\|S_{B^n}^\ell |\rho\rangle_{ABR}^{\otimes n}\| \leq \exp(-n\eta/2) \quad \text{if } \ell > -\text{tr}[\rho_B \ln \Gamma_B'] + \delta; \quad (6.21c)$$

$$\|\Pi_{B^n}^\mu |\rho\rangle_{ABR}^{\otimes n}\| \leq \text{poly}(n) \exp(-n\eta/2) \quad \text{if } \bar{H}(\mu) < H(\rho_B) - \delta \quad (6.21d)$$

recalling that $\|P|\psi\rangle\| = \sqrt{\text{tr}[P\psi]}$. So, for each term in the sum (6.13b), we have

$$\begin{aligned} \left| \langle \rho|_{ABR}^{\otimes n} (S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k) |\rho\rangle_{ABR}^{\otimes n} \right| &= \left| \langle \rho|_{ABR}^{\otimes n} S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda (R_{A^n B^n}^k |\rho\rangle_{ABR}^{\otimes n}) \right| \\ &\leq \|R_{A^n B^n}^k |\rho\rangle_{ABR}^{\otimes n}\| \cdot \|(S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda) |\rho\rangle_{ABR}^{\otimes n}\| \\ &\leq \text{poly}(n) \exp(-n\eta/2) \end{aligned} \quad (6.22)$$

using the Cauchy-Schwarz inequality and because at least one of the four conditions is violated, causing at least one of the two the norms to decay exponentially (noting also that $S_{B^n}^\ell, \Pi_{B^n}^\mu, \Pi_{A^n B^n}^\lambda$ all commute). Because there are only at most $\text{poly}(n)$ terms, we have

$$\begin{aligned} |\blacksquare_2| &\leq \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta \text{ AND} \\ [k < -\text{tr}[\sigma_X \ln \Gamma_X] - \delta \text{ OR} \\ \bar{H}(\lambda) > H(\sigma_X) + \delta \text{ OR} \\ \ell > -\text{tr}[\rho_{X'} \ln \Gamma_{X'}] + \delta \text{ OR} \\ \bar{H}(\mu) < H(\mathcal{E}(\sigma_X)) - \delta]} \left| \langle \rho|_{ABR}^{\otimes n} (S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k) |\rho\rangle_{ABR}^{\otimes n} \right| \\ &\leq \text{poly}(n) \exp(-n\eta/2). \end{aligned} \quad (6.23)$$

Hence, we have

$$\begin{aligned} \text{Re} \left\{ \langle \rho|_{ABR}^{\otimes n} M_{A^n B^n}^{x, \delta} |\rho\rangle_{ABR}^{\otimes n} \right\} &= \text{Re} \{ \blacksquare_1 \} + \text{Re} \{ \blacksquare_2 \} \\ &\geq \text{Re} \{ \blacksquare_1 \} - |\blacksquare_2| \\ &\geq 1 - \text{poly}(n) \exp(-n\eta/2) \end{aligned} \quad (6.24)$$

proving Property (ii) for $\xi = \eta/2$. Note that ξ does not depend on the state $|\sigma\rangle_{XR}$. Now, we prove Property (iii). Using Lemma B.1 and dropping some subsystem indices for readability, we have

$$\begin{aligned} \text{tr}_{A^n} [M_{A^n B^n}^{x, \delta} \Gamma_{AB}^{\otimes n} (M_{A^n B^n}^{x, \delta})^\dagger] \\ \leq \text{poly}(n) \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} \text{tr}_{A^n} [S_{B^n}^\ell \Pi_{B^n}^\mu \Pi_{A^n B^n}^\lambda R_{A^n B^n}^k \Gamma_{AB}^{\otimes n} R_{A^n B^n}^k \Pi_{A^n B^n}^\lambda \Pi_{B^n}^\mu S_{B^n}^\ell]. \end{aligned} \quad (6.25)$$

Recall that, using Proposition 2.2 and Lemma 2.2,

$$R_{A^n B^n}^k \Gamma_{AB}^{\otimes n} \leq e^{-nk} R_{A^n B^n}^k \leq e^{-nk} \mathbb{1}_{A^n B^n}; \quad (6.26)$$

$$\Pi_{B^n}^\mu \text{tr}_{A^n} [\Pi_{A^n B^n}^\lambda] \Pi_{B^n}^\mu \leq \text{poly}(n) \exp(n(\bar{H}(\lambda) - \bar{H}(\mu))) \mathbb{1}_{B^n} ; \quad (6.27)$$

$$S_{B^n}^\ell \leq e^{n\ell} S_{B^n}^\ell \Gamma_B^{\prime \otimes n} \leq e^{n\ell} \Gamma_B^{\prime \otimes n} \quad (6.28)$$

further recalling that $[R_{A^n B^n}^k, \Gamma_{AB}^{\otimes n}] = 0$ and $[S_{B^n}^\ell, \Gamma_B^{\prime \otimes n}] = 0$. Combining these together yields

$$\begin{aligned} (6.25) &\leq \text{poly}(n) \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} e^{-nk} S^\ell \Pi^\mu \text{tr}_{A^n} [\Pi_{A^n B^n}^\lambda] \Pi^\mu S^\ell \\ &\leq \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} \text{poly}(n) e^{-nk + n(\bar{H}(\lambda) - \bar{H}(\mu))} S_{B^n}^\ell \\ &\leq \sum_{\substack{k, \ell, \lambda, \mu : \\ k - \bar{H}(\lambda) - \ell + \bar{H}(\mu) \geq x - 4\delta}} \text{poly}(n) e^{-n(k - \bar{H}(\lambda) + \bar{H}(\mu) - \ell)} \Gamma_B^{\prime \otimes n} \\ &\leq \text{poly}(n) e^{-n(x - 4\delta)} \Gamma_B^{\prime \otimes n} . \end{aligned} \quad (6.29)$$

Finally, suppose that $[\Gamma_{AB}, \Gamma_B'] = 0$, meaning that we can choose a simultaneous eigenbasis for Γ_{AB} and Γ_B' . Then the operator $M_{A^n B^n}^{x, \delta}$ is a projector, as can be seen in (6.9) since in that case $\{S_{B^n}^\ell\}, \{\Pi_{B^n}^\mu\}, \{\Pi_{A^n B^n}^\lambda\}, \{R_{A^n B^n}^k\}$ are all complete sets of projectors all elements of which commute pairwise between different sets. Furthermore, $\Gamma_{B'}^{\otimes n}$ and $\Gamma_{AB}^{\otimes n}$ both commute with all of these projectors and therefore also with $M_{A^n B^n}^{x, \delta}$. ■

7 Construction #3: Thermal operations

7.1 Statement and proof sketch

We now present a construction of a universal thermodynamic implementation of a time-covariant i.i.d. process, using the framework of thermal operations instead of Gibbs-preserving maps.

Theorem 7.1. *Let X be a quantum system, H_X a Hermitian operator, $\beta \geq 0$, $\mathcal{E}_{X \rightarrow X}$ a completely positive, trace-preserving map satisfying*

$$\mathcal{E}_{X \rightarrow X}(e^{-iH_X t}(\cdot)e^{iH_X t}) = e^{-iH_X t} \mathcal{E}_{X \rightarrow X}(\cdot) e^{iH_X t} \quad \text{for all } t \in \mathbb{R}. \quad (7.1)$$

Let $\varepsilon > 0$. Let $\delta > 0$ be small enough and $n \in \mathbb{N}$ be large enough. Then, there exists an information battery W , a thermal operation $\Phi_{X^n W}$, and battery states $\tau_W^{(i)}$ and $\tau_W^{(f)}$ such that:

(i) *The effective work process $\mathcal{T}_{X^n \rightarrow X^n}$ associated with $\Phi_{X^n W}$ and $(\tau_W^{(i)}, \tau_W^{(f)})$ satisfies*

$$\frac{1}{2} \|\mathcal{T}_{X^n \rightarrow X^n} - \mathcal{E}_{X \rightarrow X}^{\otimes n}\|_\diamond \leq \varepsilon ; \quad (7.2)$$

(ii) *The work cost per copy satisfies*

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \left[w \left(\tau_W^{(i)} \right) - w \left(\tau_W^{(f)} \right) \right] = T(\mathcal{E}) . \quad (7.3)$$

(Proof on page 34.)

The main idea in the present construction is to first carry out a Stinespring dilation unitary explicitly using suitable ancillas as the environment system, and then to apply a conditional erasure process that resets the ancillas to a standard state while using the output of the process as side information. The idea of implementing a process in this fashion was also employed in Ref. [13].

Our core technical contribution for Construction #3 is to show how to build a thermodynamic protocol for universal conditional erasure, using the idea of position-based decoding [19, 56–62]. The assembly of the full thermal operation is slightly more involved than Constructions #1 and #2, because we cannot use Proposition 3.1. The construction will be illustrated in Figure 2, using a conditional erasure primitive whose construction is illustrated in Figure 1.

7.2 Universal conditional erasure

Conditional erasure is a task that is of independent interest because it generalizes Landauer’s erasure principle to situations where a quantum memory is available. A protocol for thermodynamic conditional erasure of a system using a memory as quantum side information was given in ref. [63] for trivial Hamiltonians. Here, we study the problem of finding a universal protocol for conditional erasure, whose accuracy is guaranteed for any input state on n copies of a system, and where the system and memory Hamiltonians can be arbitrary.

Definition 7.1 (Universal conditional erasure). Consider two systems S, M . Let σ_S be a fixed state, let $\mathcal{S}_{SM} = \{\rho_{SM}\}$ be an arbitrary set of states on $S \otimes M$, and let $\delta' \geq 0$. A *universal conditional δ' -erasure process* of S using M as side information is a completely positive, trace non-increasing map $\mathcal{T}_{SM \rightarrow SM}$ such that for all $\rho_{SM} \in \mathcal{S}_{SM}$, and writing $|\rho\rangle_{SMR}$ a purification of ρ_{SM} , we have

$$F \left(\mathcal{T}_{SM \rightarrow SM}(\rho_{SMR}), \sigma_S \otimes \rho_{MR} \right) \geq 1 - \delta' . \quad (7.4)$$

We provide a thermodynamic protocol for universal conditional erasure.

Proposition 7.1. Let S, M be systems with Hamiltonians H_S, H_M and let γ_S refer to the thermal state on S . Let \mathcal{S}_{SM} be an arbitrary set of states on $S \otimes M$. Let $m \geq 0$ such that e^m is integer. Let P_{SM} be a Hermitian operator satisfying $0 \leq P_{SM} \leq \mathbb{1}$ and $[P_{SM}, H_S + H_M] = 0$, and assume that there exists $\kappa, \kappa' \geq 0$ such that for all $\rho_{SM} \in \mathcal{S}_{SM}$ we have

$$\text{tr} [P_{SM} \rho_{SM}] \geq 1 - \kappa ; \quad (7.5a)$$

$$\text{tr} [P_{SM} (\gamma_S \otimes \rho_M)] \leq \frac{\kappa'}{e^m} . \quad (7.5b)$$

Then, there exists a thermal operation $\mathcal{R}_{SMJ \rightarrow SMJ}$ acting on the systems SM and an information battery J , such that the effective work process $\mathcal{T}_{SM \rightarrow SM}$ of $\mathcal{R}_{SMJ \rightarrow SMJ}$ with respect to the battery

states $(\tau_J^m, |0\rangle_J)$ is a universal conditional $(2\kappa + 4\kappa')$ -erasure process with $\sigma_S = \gamma_S$ for the set of states \mathcal{S}'_{SM} , where \mathcal{S}'_{SM} is the convex hull of \mathcal{S}_{SM} . (Proof on page 31.)

The proof of [Proposition 7.1](#) is developed in the rest of this section. We start by reformulating the ideas of the convex-split lemma, the position-based decoding, and the catalytic decoupling schemes [19, 56–62] to form a protocol for universal conditional erasure. The underlying ideas of the following proposition are the same as, e.g., in Ref. [19]. Yet, our technical statement differs in some aspects and that is why we provide a proof for completeness. The setting is depicted in [Figure 1](#).

Lemma 7.1 (Conditional erasure unitary using position-based decoding). *Consider two systems S, M and fix $m \geq 0$ such that e^m is integer. Let J be a large register of dimension at least $2e^m$, and choose a fixed basis $\{|j\rangle_J\}$. Now, let γ_S be any state, \mathcal{S}_{SM} an arbitrary set of quantum states on $S \otimes M$, P_{SM} a Hermitian operator satisfying $0 \leq P_{SM} \leq \mathbb{1}$, and assume that there exists $\kappa, \kappa' \geq 0$ such that for all $\rho_{SM} \in \mathcal{S}_{SM}$ the conditions (7.5) hold. Furthermore, let $A = A_1 \otimes \dots \otimes A_{e^m}$ be a collection of ancilla systems with each $A_j \simeq S$, and let $A' = A'_1 \otimes \dots \otimes A'_{e^m}$ be a copy of the full collection of ancilla systems. We write a purification of γ_{A_j} on A'_j as $|\gamma\rangle_{A_j A'_j} = \gamma_{A_j}^{1/2} |\Phi\rangle_{A_j A'_j}$. Let \mathcal{S}'_{SM} be the convex hull of \mathcal{S}_{SM} . Then, there exists a unitary operator $W_{SMAJ \rightarrow SMAJ}^{(m)}$ satisfying the following property: For any reference system R , for any pure tripartite state $|\rho\rangle_{SMR}$ with $\rho_{SM} \in \mathcal{S}'_{SM}$, and for any $|j\rangle_J$ with $1 \leq j \leq e^m$, we have*

$$\text{Re} \left\{ \left(\langle \hat{\tau}^j(\rho_{SMR}) |_{RMSAA'} \otimes \langle 0 |_J \right) W_{SMAJ}^{(m)} (|\rho\rangle_{RMS} \otimes |\gamma\rangle_{AA'}^{\otimes e^m} \otimes |j\rangle_J) \right\} \geq 1 - (2\kappa + 4\kappa'), \quad (7.6)$$

where we have defined

$$|\hat{\tau}^j(\rho_{SMR})\rangle_{RMSAA'} = |\rho\rangle_{A_j MR} \otimes |\gamma\rangle_{SA'_j} \otimes [|\gamma\rangle^{\otimes (e^m - 1)}]_{AA' \setminus A_j A'_j} \quad (7.7)$$

and by the notation $AA' \setminus A_j A'_j$ we refer to all AA' systems except $A_j A'_j$. Moreover, for any observables H_S, H_M such that $[P_{SM}, H_S + H_M] = 0$, the unitary $W_{SMAJ}^{(m)}$ may be chosen such that $[H_S + H_M + \sum H_{A_j}, W_{SMAJ}^{(m)}] = 0$, where $H_{A_j} = H_S$.

Intuitively, we absorb the initial randomness present in the register J , e.g., given to us by the environment in a mixed state, and return it in a pure state; J can therefore be identified as an information battery. Similarly, A can be identified as a heat bath.

Proof. First observe that we can assume \mathcal{S}_{SM} to be a convex set, because any convex combination of states in \mathcal{S}_{SM} also satisfies the conditions (7.5). For the rest of the proof we assume without loss of generality that $\mathcal{S}_{SM} = \mathcal{S}'_{SM}$.

The operator W is defined in two steps. The first operation simply consists on conditionally swapping S with A_j , depending on the value stored in J . Then, we infer again from MA which j we swapped S with, in order to coherently reset the register J back to the zero state (approximately). We define the first unitary operation as $W^{(1)}$, acting on systems SAJ

$$W_{SAJ}^{(1)} = \sum_j \mathbb{F}_{SA_j} \otimes |j\rangle\langle j|_J, \quad (7.8)$$




Figure 1: Construction of the thermal operation for universal conditional erasure using position-based decoding [19], illustrating the construction in the proof of Proposition 7.1 and Lemma 7.1. We define a map \mathcal{R}_{SMJ} that acts on a system S to reset, a quantum memory M and a register J , which is promised to be initialized in the uniformly mixed state $e^{-m} \mathbb{1}_{e^m}$ of rank e^m for a fixed and known value of m . A state ρ_{SM} of the system and the memory is purified by a reference system R (not pictured). The map \mathcal{R}_{SMJ} outputs the system S in a state close to the thermal state γ_S and the register J in a state close to the pure state $|0\rangle_J$, all while ensuring that ρ_{MR} remains unchanged (up to small errors), for all states ρ_{SM} in a given class of states \mathcal{S}_{SM} . The routine is provided a POVM effect P_{SM} whose task is to distinguish ρ_{SM} from $\gamma_S \otimes \rho_M$ in a hypothesis test for all $\rho_{SM} \in \mathcal{S}_{SM}$. As long as m is not too large (as determined by how well P_{SM} can perform this distinguishing), the procedure completes successfully. To implement \mathcal{R}_{SMJ} (shaded region) we involve e^m ancillas $A = A_1 \dots A_{e^m}$ with $A_j \simeq S$, each initialized in the thermal state $\gamma_{A_j} = \gamma_S$. Then S and A_j are coherently swapped (\mathbb{F}_{SA_j}) conditioned on the value stored in J . If m is not too large, a POVM $\{\Omega_{MA}^j\}$ can infer the value j stored in J , up to a small error; the POVM is constructed from P_{SM} . We then coherently reset the J register to zero by conditioning on this outcome (up to a small error). The full procedure is a thermal operation where the ancillas are the heat bath and J is an information battery such that m work has been extracted in units of pure nats (see main text).

where \mathbb{F}_{SA_j} denotes the swap operator between the two designated systems. Observe that $W^{(1)}$ maps ρ onto $\hat{\tau}^j$ according to

$$\begin{aligned} W_{SQJ}^{(1)} & \left(|\rho\rangle_{RMS} \otimes |\gamma\rangle_{A.A'}^{\otimes e^m} \otimes |j\rangle_J \right) \\ &= |\rho\rangle_{RMA_j} \otimes |\gamma\rangle_{SA'_j} \otimes \left[|\gamma\rangle^{\otimes (e^m-1)} \right]_{AA' \setminus A_j A'_j} \otimes |j\rangle_J \\ &= |\hat{\tau}^j\rangle_{SRMAA'} \otimes |j\rangle_J . \end{aligned} \quad (7.9)$$

The second step is more tricky. We need to infer from the systems MA alone which j was stored in J . Fortunately the answer is provided in the form of position-based decoding [19], using a pretty good measurement. Define

$$\Lambda_{MA}^j = P_{MA_j} \otimes \mathbb{1}_{A \setminus A_j} \quad (7.10)$$

such that $\{\Lambda_{MA}^j\}$ is a set of positive operators. We can form a POVM $\{\Omega_{MA}^j\}_j \cup \{\Omega_{MA}^\perp\}$ by normalizing the Λ^j 's as follows:

$$\Omega_{MA}^j = \Lambda_{MA}^{-1/2} \Lambda_{MA}^j \Lambda_{MA}^{-1/2} ; \quad \Lambda_{MA} = \sum_j \Lambda_{MA}^j ; \quad \Omega_{MA}^\perp = \mathbb{1} - \sum_j \Omega_{MA}^j . \quad (7.11)$$

We would now like to lower bound $\text{tr}[\Omega_{MA}^j \hat{\tau}_{MA}^j]$. Following the proof of [19, Theorem 2], we first invoke the Hayashi-Nagaoka inequality [64], which states that for any operators $0 \leq A \leq \mathbb{1}$, $B \geq 0$, we have

$$\mathbb{1} - (A + B)^{-1/2} A (A + B)^{-1/2} \leq 2(\mathbb{1} - A) + 4B . \quad (7.12)$$

Applying this inequality with $A = \Lambda_{MA}^j$ and $B = \sum_{j' \neq j} \Lambda_{MA}^{j'}$ we obtain

$$\begin{aligned} \text{tr} \left[(\mathbb{1} - \Omega^j) \hat{\tau}_{MA}^j \right] &\leq 2 \text{tr} \left[(\mathbb{1} - \Lambda_{MA}^j) \hat{\tau}_{MA}^j \right] + 4 \sum_{j' \neq j} \text{tr} \left[\Lambda_{MA}^{j'} \hat{\tau}_{MA}^j \right] \\ &\leq 2 \text{tr}[(\mathbb{1} - P_{SM}) \rho_{SM}] + 4m \text{tr}[P_{SM}(\gamma_S \otimes \rho_M)] \\ &\leq 2\kappa + 4\kappa' . \end{aligned} \quad (7.13)$$

Now, let $\text{SHIFT}_J(x) = \sum_j |j+x\rangle \langle j|_J$ denote the SHIFT operation on the J register, modulo e^m ; note that $(\text{SHIFT}_J(x))^\dagger = \text{SHIFT}_J(-x)$. We define

$$W_{MAJ}^{(2)} = \left(\sum_j \Omega_{MA}^j \otimes \text{SHIFT}_J(-j) \right) ; \quad W'_{SMAJ} = W_{MAJ}^{(2)} W_{SAJ}^{(1)} \quad (7.14)$$

and we see that $W'^\dagger W' \leq \mathbb{1}$ thanks to [Proposition B.1](#). Then, we have

$$\begin{aligned}
W'_{SMAJ} \left(|\rho\rangle_{RMS} \otimes |\phi\rangle_{A.A'}^{\otimes e^m} \otimes |j\rangle_J \right) \\
&= \left(\sum_{j'} \Omega_{MA}^{j'} \otimes \text{SHIFT}_J(-j') \right) (|\hat{\tau}^j\rangle_{SRMAA'} \otimes |j\rangle_J) \\
&= \sum_{j'} \left(\Omega_{MA}^{j'} |\hat{\tau}^j\rangle_{RMSAA'} \right) \otimes |j-j'\rangle.
\end{aligned} \tag{7.15}$$

Thanks to [Proposition C.1](#), the operator W'_{SMAJ} can be completed to a full unitary W_{SMAJ} by using an extra qubit in the J register, and such that $\langle 0|_J W_{SMAJ} |j\rangle_J = \langle 0|_J W'_{SMAJ} |j\rangle_J$ for all $j = 1, \dots, e^m$ (with the convention that $|j\rangle_J$ for $j \leq e^m$ forces the extra qubit to be in the zero state). So, recalling [\(7.13\)](#),

$$\begin{aligned}
&(\langle \hat{\tau}^j |_{RMSAA'} \otimes \langle 0 |_J) W_{SMAJ} \left(|\rho\rangle_{RMS} \otimes |\phi\rangle_{A.A'}^{\otimes e^m} \otimes |j\rangle_J \right) \\
&= (\langle \hat{\tau}^j |_{RMSAA'} \otimes \langle 0 |_J) W'_{SMAJ} \left(|\rho\rangle_{RMS} \otimes |\phi\rangle_{A.A'}^{\otimes e^m} \otimes |j\rangle_J \right) \\
&= \langle \hat{\tau}^j | \Omega_{MA}^j | \hat{\tau}^j \rangle_{RMSAA'} \\
&\geq 1 - (2\kappa + 4\kappa').
\end{aligned} \tag{7.16}$$

To prove the last part of the claim, let H_S, H_M be observables such that $[P_{SM}, H_S + H_M] = 0$ and $[H_S, \gamma_S] = 0$. Let $H_{A_j} = H_S$ and we write $H_A = \sum_j H_{A_j}$. For all j , we have

$$[H_S + H_M + H_A, \Lambda_{MA}^j] = [H_S + \sum_{j' \neq j} H_{A_{j'}}, \Lambda_{MA}^j] + [H_M + H_{A_j}, P_{MA_j}] = 0. \tag{7.17}$$

This implies that $[H_S + H_M + H_A, \Lambda_{MA}] = 0$, and in turn $[H_S + H_M + H_A, \Lambda_{MA}^{-1/2}] = 0$, and thus also $[H_S + H_M + H_A, \Omega^j] = 0$. Hence, we have

$$[H_S + H_M + H_A, W_{MAJ}^{(2)}] = 0. \tag{7.18}$$

Clearly, $[H_S + H_M + H_A, W_{SAJ}^{(1)}] = 0$, and hence $[H_S + H_M + H_A, W'_{SMAJ}] = 0$. Using [Proposition C.2](#) instead of [Proposition C.1](#), we may further enforce $[H_S + H_M + H_A, W_{SMAJ}] = 0$, as required. ■

We now give the proof of [Proposition 7.1](#).

Proof of Proposition 7.1. Let $W_{SMAJ}^{(m)}$ be the energy-conserving unitary as in [Lemma 7.1](#) and define the thermal operation

$$\mathcal{R}_{SMJ}(\cdot) = \text{tr}_A [W_{SMAJ}^{(m)} ((\cdot) \otimes \gamma_A) W_{SMAJ}^{(m)\dagger}]. \tag{7.19}$$

Identifying J as an information battery, the associated effective work process of \mathcal{R}_{SMJ} with respect to $(\tau_J^m, |0\rangle_J)$ is

$$\mathcal{T}_{SM \rightarrow SM}(\cdot) = \text{tr}_A [\langle 0 |_J W_{SMAJ}^{(m)} ((\cdot) \otimes \gamma_A \otimes \tau_J^m) W_{SMAJ}^{(m)\dagger} |0\rangle_J]. \tag{7.20}$$

Let $\rho_{SM} \in \mathcal{S}'_{SM}$ and let $|\rho\rangle_{SMR}$ be a purification of ρ_{SM} . We have that the state vector

$$e^{-m/2} \sum_{j=1}^{e^m} \langle 0|_J W_{SMAJ}^m (|\rho\rangle_{SMR} \otimes |\gamma\rangle_{AA'}^{\otimes e^m} \otimes |j\rangle_J) \otimes |j\rangle_{R_J} \quad (7.21)$$

is a purification of $\mathcal{T}_{SM \rightarrow SM}(\rho_{SMR})$, where R_J is an additional register. Similarly, the state vector

$$e^{-m/2} \sum_{j=1}^{e^m} |\hat{\tau}^j(\rho_{SMR})\rangle_{RMSAA'} \otimes |j\rangle_{R_J} \quad (7.22)$$

is a purification of $\gamma_S \otimes \rho_{MR}$. Then, with Uhlmann's theorem we find

$$\begin{aligned} & F(\mathcal{T}_{SM \rightarrow SM}(\rho_{SMR}), \gamma_S \otimes \rho_{MR}) \\ & \geq e^{-m} \sum_{j=1}^{e^m} \text{Re} \left\{ \left(\langle \hat{\tau}^j(\rho_{SMR}) |_{RMSAA'} \otimes \langle 0 |_J \right) W_{SMAJ}^{(m)} (|\rho\rangle_{RMS} \otimes |\gamma\rangle_{AA'}^{\otimes e^m} \otimes |j\rangle_J) \right\} \\ & \geq 1 - (2\kappa + 4\kappa') , \end{aligned} \quad (7.23)$$

making use of (7.6). ■

7.3 Construction via universal conditional erasure

This section is devoted to the proof of [Theorem 7.1](#). The strategy is to exploit the fact that time-covariant processes admit a Stinespring dilation with an energy-conserving unitary using an environment system with a separate Hamiltonian. This property enables us to map the problem of implementing such a process directly to a conditional erasure problem with a system and memory that are non-interacting.

The following lemma formalizes the property of time-covariant processes we make use of. Various proofs of this lemma can be found in [65], [66, Appendix B] and [67, Theorem 25].

Lemma 7.2 (Stinespring dilation of covariant processes [65–67]). *Let X be a quantum system with Hamiltonian H_X , and $\mathcal{E}_{X \rightarrow X}$ be a completely positive, trace-preserving map that is covariant with respect to time evolution. That is, for all t we have*

$$\mathcal{E}_{X \rightarrow X}(e^{-iH_X t}(\cdot)e^{iH_X t}) = e^{-iH_X t} \mathcal{E}_{X \rightarrow X}(\cdot) e^{iH_X t} . \quad (7.24)$$

Then, there exists a system E with Hamiltonian H_E including an eigenstate $|0\rangle_E$ of zero energy, as well as a unitary $V_{EX \rightarrow EX}$ such that

$$\mathcal{E}_{X \rightarrow X}(\cdot) = \text{tr}_E [V(|0\rangle\langle 0|_E \otimes (\cdot)) V^\dagger] \quad (7.25)$$

as well as $V(H_X + H_E)V^\dagger = H_X + H_E$. (Proof on [page 49](#).)

We provide an additional proof in [Appendix A](#). The main idea behind the construction in the following proof of [Theorem 7.1](#) is depicted in [Figure 2](#).




Figure 2: The conditional erasure procedure in Figure 1 can be used to construct an i.i.d. implementation of a given time-covariant process (Theorem 7.1). First we apply an energy-conserving Stinespring dilation of the process on all input copies, using a zero-initialized ancilla as environment system E for each copy. We then invoke the conditional erasure procedure $\mathcal{R}_{E^n X^n J}$ to reset E^n to the thermal state $\gamma_E^{\otimes n}$ using X^n as a memory, while extracting work using an information battery J . Here, the projector that can distinguish $\rho_{E X^n}^{\otimes n}$ from $\mathbb{1}_{E^n} \otimes \rho_{X^n}$ is the universal conditional typical projector given by Proposition E.1. The fact that $\mathcal{R}_{E^n X^n J}$ preserves the correlations $[\mathcal{E}(\sigma_{XR})]^{\otimes n}$ between the memory (output systems X^n) and the reference R^n ensures that the process is implemented accurately. The amount of work extracted by $\mathcal{R}_{E^n X^n J}$ is $m \sim n[\beta F_E + T(\mathcal{E})]$ but $\sim n\beta F_E$ work has to be paid to prepare the initially pure E^n ancillas, where $\beta F_E = -\ln \text{tr}(e^{-\beta H_E})$. The overall work extracted is $\sim T(\mathcal{E})$ per copy.

Proof of Theorem 7.1. Thanks to Lemma 7.2, there exists an environment system E with Hamiltonian H_E , as well as an energy-conserving unitary V_{XE} and a state $|0\rangle_E$ of zero energy such that (7.25) holds. Let $F_E = -\beta^{-1} \ln(Z_E)$ with $Z_E = \text{tr}[e^{-\beta H_E}]$. We define

$$x = \min_{\sigma} \left\{ D(\sigma \| e^{-\beta H_X}) - D(\mathcal{E}(\sigma) \| e^{-\beta H_X}) \right\} = -T(\mathcal{E}). \quad (7.26)$$

Writing $\rho_{XE} = V_{XE}(|0\rangle\langle 0|_E \otimes \sigma_X) V_{XE}^\dagger$, we have that $x = \min_{\sigma_X} \{ -H(\sigma_X) + \beta \text{tr}[\sigma_X H_X] + H(\rho_X) - \beta \text{tr}[\rho_X H_X] \}$. By $\text{tr}[\sigma_X H_X] = \text{tr}[(|0\rangle\langle 0|_E \otimes \sigma_X)(H_X + H_E)] = \text{tr}[\rho_{XE}(H_X + H_E)]$, we see that

$$x = \min_{\sigma_X} \{ -H(\rho_{XE}) + H(\rho_X) + \beta \text{tr}[\rho_E H_E] \}. \quad (7.27)$$

Observe that for any such ρ_{XE} , we have

$$\begin{aligned} -H(E|X)_\rho + \beta \text{tr}[\rho_E H_E] &\geq -H(E)_\rho + \beta \text{tr}[\rho_E H_E] + \ln(Z) - \ln(Z) \\ &= D(\rho_E \| \gamma_E) + \beta F_E \geq \beta F_E \end{aligned} \quad (7.28)$$

using the sub-additivity of the von Neumann entropy and the fact that relative entropy is positive for normalized states. Hence, we have $x \geq \beta F_E$.

Let

$$\mathcal{S}_{E^n X^n} = \{ \rho_{EX}^{\otimes n} : \rho_{EX} = V_{XE}(|0\rangle\langle 0|_E \otimes \sigma_X) V_{XE}^\dagger \text{ for some } \sigma_X \}, \quad (7.29)$$

noting that for all $\rho_{EX}^{\otimes n} \in \mathcal{S}_{E^n X^n}$, we have $D(\rho_{EX} \| e^{-\beta(H_X + H_E)}) - D(\rho_X \| e^{-\beta H_X}) = D(\sigma \| e^{-\beta H_X}) - D(\mathcal{E}(\sigma) \| e^{-\beta H_X}) \geq x$. Let $P_{E^n X^n}^{x, \delta}$ be the universal typical and relative conditional operator furnished by Proposition 6.1, where $\Gamma_X = e^{-\beta H_X}$ and $\Gamma_{XE} = e^{-\beta(H_X + H_E)} = \Gamma_X \otimes \Gamma_E$ with $\Gamma_E = e^{-\beta H_E}$. Since Γ_{XE} commutes with $\mathbb{1}_E \otimes \Gamma_X$, Proposition 6.1 guarantees that $P_{E^n X^n}^{x, \delta}$ is a projector which furthermore commutes with $\Gamma_{XE}^{\otimes n}$ and $\Gamma_X^{\otimes n}$. We proceed to show that $P_{E^n X^n}^{x, \delta}$ can perform a hypothesis test between $\rho_{EX}^{\otimes n}$ and $\gamma_E^{\otimes n} \otimes \rho_X^{\otimes n}$. Recalling Definition 6.1 we have

$$\text{tr}[P_{E^n X^n}^{x, \delta} \rho_{EX}^{\otimes n}] \geq 1 - \kappa, \quad (7.30)$$

with $\kappa = \text{poly}(n) e^{-n\eta}$ for some $\eta > 0$ independent of ρ and n . By construction we have $\mathbb{1}_X \otimes \Gamma_E = \Gamma_X^{-1/2} \Gamma_{XE} \Gamma_X^{-1/2}$, and so thanks to Point (iii) of Definition 6.1 we can compute

$$\begin{aligned} \text{tr}_{E^n} [P_{E^n X^n}^{x, \delta} \Gamma_E^{\otimes n}] &= (\Gamma_X^{-1/2})^{\otimes n} \text{tr}_{E^n} [P_{E^n X^n}^{x, \delta} \Gamma_{XE}^{\otimes n}] (\Gamma_X^{-1/2})^{\otimes n} \\ &\leq \text{poly}(n) \exp(-n(x - 4\delta)) \mathbb{1}_{X^n}, \end{aligned} \quad (7.31)$$

where we furthermore used the fact that $P_{E^n X^n}^{x, \delta}$ commutes with $\Gamma_{XE}^{\otimes n}$ and with $\Gamma_X^{\otimes n}$. We therefore see using $\gamma_E = \Gamma_E / \text{tr}[\Gamma_E]$ that

$$\begin{aligned} \text{tr}[P_{E^n X^n}^{x, \delta} \rho_X^{\otimes n} \otimes \gamma_E^{\otimes n}] &\leq \frac{1}{\text{tr}[\Gamma_E^{\otimes n}]} \text{poly}(n) \exp(-n(x - 4\delta)) \text{tr}[\rho_X^{\otimes n}] \\ &= \text{poly}(n) \exp(-n(x - \beta F_E - 4\delta)). \end{aligned} \quad (7.32)$$

Let

$$e^m = \lfloor \exp\{n(x - \beta F_E - 4\delta - \eta)\} \rfloor, \quad (7.33)$$

such that $\text{tr}[P_{E^n X^n}^{x, \delta} \rho_X^{\otimes n} \otimes \gamma_E^{\otimes n}] \leq e^{-m} \kappa'$ by choosing $\kappa' = \text{poly}(n) e^{-n\eta}$.

Now let J be a register of dimension at least $2e^m$ and let $\mathcal{R}_{E^n X^n J}$ be the thermal operation furnished by [Proposition 7.1](#) for $S = E^n$, $M = X^n$, $\mathcal{S}_{E^n X^n}$, $P_{E^n X^n}^{x, \delta}$, m , κ , and κ' as defined above. Here, we have assumed that $x > \beta F_E$, and that furthermore δ, η are small enough such that $4\delta + \eta < (x - \beta F_E)$; if instead $x = \beta F_E$ then we can set $e^m = 1$ and $\mathcal{R}_{E^n X^n J}(\cdot) = \text{tr}_{E^n}(\cdot) \otimes \gamma_E^{\otimes n}$ (which is a thermal operation) in the following.

We proceed to show that the effective work process $\mathcal{T}_{E^n X^n \rightarrow E^n X^n}^{\mathcal{R}}$ of $\mathcal{R}_{E^n X^n J}$ with respect to $(\tau_J^m, |0\rangle_J)$ is close to the partial trace map $\mathcal{T}_{E^n X^n \rightarrow E^n X^n}^{(0)}(\cdot) = \text{tr}_{E^n}(\cdot) \otimes \gamma_E^{\otimes n}$ in diamond distance. We invoke the post-selection technique ([Theorem 2.1](#)) to show this. Let $\zeta_{E^n X^n}$ be the de Finetti state which via (2.24) can be written as the convex combination of a finite number of i.i.d. states

$$\zeta_{E^n X^n} = \sum p_i \phi_i^{\otimes n}. \quad (7.34)$$

Hence $\zeta_{E^n X^n}$ lies in the convex hull of $\mathcal{S}_{E^n X^n}$, and from [Proposition 7.1](#) and [Definition 7.1](#) we see that for a purification $|\zeta\rangle_{E^n X^n R}$ of $\zeta_{E^n X^n}$ we have

$$F(\mathcal{T}_{E^n X^n \rightarrow E^n X^n}^{\mathcal{R}}(\zeta_{E^n X^n R}), \gamma_E^{\otimes n} \otimes \text{tr}_{E^n}(\zeta_{E^n X^n R})) \geq 1 - (2\kappa + 4\kappa'). \quad (7.35)$$

Using $D(\rho, \sigma) \leq \sqrt{1 - F(\rho, \sigma)}$ along with [Theorem 2.1](#) we find

$$\frac{1}{2} \|\mathcal{T}_{E^n X^n \rightarrow E^n X^n}^{\mathcal{R}} - \mathcal{T}_{E^n X^n \rightarrow E^n X^n}^{(0)}\|_{\diamond} \leq \sqrt{2\kappa + 4\kappa'} = \text{poly}(n) e^{-n\eta/2}. \quad (7.36)$$

We can start piecing together the full process. Our overall protocol needs to (a) bring in a heat bath E^n , i.e., ancillas initialized in their thermal state, (b) prepare the states $|0\rangle_E^{\otimes n}$ on the ancillas using an auxiliary information battery (denoted by W' below), (c) apply the energy-conserving unitary $V_{XE}^{\otimes n}$, (d) apply $\mathcal{R}_{E^n X^n J}$ using an information battery J initialized in the state τ_J^m , and (e) discard the ancillas.

As explained in [Section 3](#), there exists a thermal operation $\tilde{\Phi}_{E^n W'}$ on the ancillas and an information battery W' along with battery states $(\tau_{W'}^{(1)}, \tau_{W'}^{(2)})$ such that $\tilde{\Phi}_{E^n W'}(\gamma_E^{\otimes n} \otimes \tau_{W'}^{(1)}) = |0\rangle\langle 0|_E^{\otimes n} \otimes \tau_{W'}^{(2)}$ and with $w(\tau_{W'}^{(1)}) - w(\tau_{W'}^{(2)})$ arbitrarily close to $-\beta n F_E$. Now let $W = J \otimes W'$, $\tau_W^{(i)} = \tau_{W'}^{(i)} \otimes \tau_J^m$, $\tau_W^{(f)} = \tau_{W'}^{(2)} \otimes |0\rangle\langle 0|_J$, and define

$$\Phi_{X^n W}(\cdot) = \text{tr}_{E^n} \left[\mathcal{R}_{E^n X^n J} \left(V_{XE}^{\otimes n} \tilde{\Phi}_{E^n W'}((\cdot) \otimes \gamma_E^{\otimes n}) (V_{XE}^{\otimes n})^\dagger \right) \right]. \quad (7.37)$$

The map $\Phi_{X^n W}$ is a thermal operation because it is a concatenation of thermal operations. The overall heat bath is formed of the systems E^n , the ancillas A^n used in the implementation of $\mathcal{R}_{E^n X^n J}$, as well as the implicit heat bath used in the implementation of $\tilde{\Phi}_{E^n W'}$. The system $W = J \otimes W'$ is

the information battery. We can verify that the associated effective work process with respect to $(\tau_W^{(i)}, \tau_W^{(f)})$ is

$$\begin{aligned}
\mathcal{T}_{X^n}(\cdot) &= \langle 0 |_J \text{tr}_{E^n} \left[\mathcal{R}_{E^n X^n J} \left(V_{XE}^{\otimes n} \text{tr}_{W'} \left[P_{W'}^{(2)} \tilde{\Phi}_{E^n W'}((\cdot) \otimes \tau_{W'}^{(1)} \otimes \tau_J^m \otimes \gamma_E^{\otimes n}) \right] (V_{XE}^{\otimes n})^\dagger \right) \right] |0\rangle_J \\
&= \text{tr}_{E^n} \left[\langle 0 |_J \mathcal{R}_{E^n X^n J} \left([V_{XE}^{\otimes n} ((\cdot) \otimes |0\rangle\langle 0|_E^{\otimes n}) (V_{XE}^{\otimes n})^\dagger] \otimes \tau_J^m \right) |0\rangle_J \right] \\
&= \text{tr}_{E^n} \left[\mathcal{T}_{E^n X^n}^{\mathcal{R}} \left(V_{XE}^{\otimes n} ((\cdot) \otimes |0\rangle\langle 0|_E^{\otimes n}) (V_{XE}^{\otimes n})^\dagger \right) \right] \\
&= \text{tr}_{E^n} \left[V_{XE}^{\otimes n} ((\cdot) \otimes |0\rangle\langle 0|_E^{\otimes n}) (V_{XE}^{\otimes n})^\dagger \right] + \Delta_{X^n}(\cdot) \\
&= \mathcal{E}_{X \rightarrow X}^{\otimes n}(\cdot) + \Delta_{X^n}(\cdot), \tag{7.38}
\end{aligned}$$

where $\Delta_{X^n}(\cdot) = \text{tr}_{E^n}(\mathcal{T}_{X^n E^n}^{\mathcal{R}}(\cdot) - \mathcal{T}_{X^n E^n}^{(0)}(\cdot))$ satisfies $(1/2)\|\Delta_{X^n}\|_\diamond \leq \text{poly}(n)e^{-n\eta/2}$. Therefore for any fixed ε and for n large enough we have $(1/2)\|\mathcal{T}_{X^n} - \mathcal{E}_{X \rightarrow X}^{\otimes n}\|_\diamond \leq \varepsilon$.

The associated work cost per copy satisfies

$$\begin{aligned}
\lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} [w(\tau_W^{(i)}) - w(\tau_W^{(f)})] &= \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} [w(\tau_{W'}^{(1)}) - w(\tau_{W'}^{(2)}) - m] \\
&= \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} [-n\beta F_E - n(x - \beta F_E - 4\delta + \eta) + v] \\
&= T(\mathcal{E}), \tag{7.39}
\end{aligned}$$

recalling (7.33), where $0 \leq v \leq 2$ accounts for the rounding error in (7.33) and a possible arbitrarily small difference between $-n\beta F_E$ and $w(\tau_{W'}^{(1)}) - w(\tau_{W'}^{(2)})$, and recalling that $\eta \rightarrow 0$ as $\delta \rightarrow 0$. ■

8 Related Results

8.1 Alternative proof of the asymptotic equipartition property of the coherent relative entropy

Here we have a new proof of the asymptotic equipartition property (AEP) for the coherent relative entropy, with an explicit expression of a smoothing process which does the job. This isn't directly useful for our universality result, but it uses similar ideas and gives some intuition about the AEP of the coherent relative entropy. Furthermore, we get an explicit process that is near-optimal in the definition of the coherent relative entropy for an i.i.d. process matrix.

Recall the definition of the relative typical subspace [54, 55]:

Proposition 8.1 (Relative typical projector [54, 55]). *Let $\rho, \tau \geq 0$ be operators on a finite dimensional Hilbert space \mathcal{H} with $\text{tr}(\rho) = 1$, and let $\delta > 0$. There exists a constant $\eta > 0$, and for all n there*

exists a projector $\Pi_{\rho|\tau}^{n,\delta}$ such that the following conditions hold:

$$[\Pi_{\rho|\tau}^{n,\delta}, \tau^{\otimes n}] = 0 ; \quad (8.1a)$$

$$e^{-n(M(\rho \parallel \tau) + \delta)} \Pi_{\rho|\tau}^{n,\delta} \leq \Pi_{\rho|\tau}^{n,\delta} \tau^{\otimes n} \Pi_{\rho|\tau}^{n,\delta} \leq e^{-n(M(\rho \parallel \tau) - \delta)} \Pi_{\rho|\tau}^{n,\delta} ; \quad (8.1b)$$

$$\text{tr}(\Pi_{\rho|\tau}^{n,\delta} \rho^{\otimes n}) \geq 1 - 2e^{-n\eta} , \quad (8.1c)$$

where we have defined

$$M(\rho \parallel \tau) := -\text{tr}(\rho \ln \tau) . \quad (8.2)$$

The usual (weakly) typical projector for a state ρ is obtained by choosing $\tau = \rho$:

$$\Pi_{\rho}^{n,\delta} = \Pi_{\rho|\rho}^{n,\delta} . \quad (8.3)$$

The construction of the relative typical projector, as well as the proof of properties (8.1a) and (8.1b) are presented in Refs. [54, 55]. Here we show property (8.1c).

Proof of Proposition 8.1. The construction of Refs. [54, 55] satisfies properties (8.1a) and (8.1b); it remains to prove (8.1c). Consider the quantity $\text{tr}((\mathbb{1} - \Pi_{\rho|\tau}^{n,\delta}) \rho^{\otimes n})$, and note that it corresponds to the probability that a sequence of measurements of copies ρ of the observable $-\ln(\tau)$ ensemble averages to a quantity that is δ -far from $M(\rho \parallel \tau) = -\text{tr}(\rho \ln \tau)$. Let Z_j for $j = 1, \dots, n$ be random variables where Z_j is the outcome of the measurement of $-\ln(\tau)$ on the j -th system of $\rho^{\otimes n}$. Then using Hoeffding's inequality we find

$$\text{tr}((\mathbb{1} - \Pi_{\rho|\tau}^{n,\delta}) \rho^{\otimes n}) = \Pr \left[\left| \frac{1}{n} \sum Z_j - (-\text{tr}(\rho \ln \tau)) \right| > \delta \right] \leq 2 \exp(-n\eta) , \quad (8.4)$$

for some $\eta \geq \delta^2 / \|\ln \tau\|_{\infty}$, noting that the difference between the maximum and minimum eigenvalues of $-\ln \tau$ is upper bounded by $2\|\ln \tau\|_{\infty}$. \blacksquare

Now we may present the new proof of the asymptotic equipartition property of the coherent relative entropy.

Proposition 8.2. Let $\Gamma_X, \Gamma_{X'} \geq 0$, let $R_X \simeq X$ and let $|\sigma\rangle_{X:R_X}$ be any state. Write $\rho_{X'R_X} = \mathcal{E}(\sigma_{X:R_X}) = \rho_{R_X}^{1/2} E_{X'R_X} \rho_{R_X}^{1/2}$ writing $E_{X'R_X} = \mathcal{E}(\Phi_{X:R_X})$. Let $\Gamma_X, \Gamma_{X'} \geq 0$. For any $\delta > 0$, and for any n , let

$$S_{X^m} = \Pi_{\rho_{X'}|\Gamma_{X'}^{-1}}^{n,\delta} ; \quad Q_{X^m} = \Pi_{\rho_{X'}}^{n,\delta} ; \quad P_{X^n} = \Pi_{\sigma_X|\Gamma_{R_X}}^{n,\delta} , \quad R_{X^n} = \Pi_{\sigma_X}^{n,\delta} . \quad (8.5)$$

Then the completely positive map

$$\mathcal{T}_{X^n \rightarrow X^m}(\cdot) = S_{X^m} Q_{X^m} \mathcal{E}_{X \rightarrow X'}^{\otimes n} (R_{X^n} P_{X^n}(\cdot) P_{X^n} R_{X^n}) Q_{X^m} S_{X^m} , \quad (8.6)$$

is trace-nonincreasing and satisfies

$$\mathcal{T}_{X^n \rightarrow X'^n}(\Gamma_X^{\otimes n}) \leq e^{-n[D(\sigma_X \parallel \Gamma_X) - D(\rho_{X'} \parallel \Gamma_{X'}) - 4\delta]} \Gamma_{X'}^{\otimes n}; \quad (8.7a)$$

$$P(\mathcal{T}_{X^n \rightarrow X'^n}(\sigma_{R_X}^{\otimes n}), \rho_{X'R_X}^{\otimes n}) \leq 4e^{-n\eta'}, \quad (8.7b)$$

for some $\eta' > 0$. This implies that for any $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \hat{D}_{X^n \rightarrow X'^n}^\varepsilon(\rho_{X'R_X}^{\otimes n} \parallel \Gamma_X^{\otimes n}, \Gamma_{X'}^{\otimes n}) \geq D(\sigma_X \parallel \Gamma_X) - D(\rho_{X'} \parallel \Gamma_{X'}). \quad (8.8)$$

Proof of Proposition 8.2. It will prove convenient to work in the purified space, so let $E \simeq X' \otimes R_X$, and let $|E\rangle_{X'R_X E}$ be a purification of $E_{X'R_X}$. Let $V_{X \rightarrow X'E}$ be the corresponding isometry which satisfies $|E\rangle_{X'R_X E} = V_{X \rightarrow X'E} |\Phi\rangle_{X:R_X}$; this isometry is just a Stinespring dilation of \mathcal{E} . Now define

$$|T\rangle_{X^m R_X^n E^n} = S_{X^m} Q_{X^m} P_{R_X^n} R_{R_X^n} |E\rangle_{X'R_X E}^{\otimes n}, \quad (8.9)$$

where $P_{R_X^n} = t_{X^n \rightarrow R_X^n}(P_{X^n})$ and $R_{R_X^n} = t_{X^n \rightarrow R_X^n}(R_{X^n})$. We begin by showing (8.7a). Writing $\Gamma_{R_X} = t_{X \rightarrow R_X}(\Gamma_X)$, we have

$$\begin{aligned} & (\Gamma_{X'}^{-1/2})^{\otimes n} \text{tr}_{R_X^n} [T_{X^m R_X^n} \Gamma_{R_X}^{\otimes n}] (\Gamma_{X'}^{-1/2})^{\otimes n} \\ &= (\Gamma_{X'}^{-1/2})^{\otimes n} S_{X^m} Q_{X^m} \text{tr}_{R_X^n} \left[R_{R_X^n} E_{X'R_X}^{\otimes n} R_{R_X^n} (P_{R_X^n} \Gamma_{R_X}^{\otimes n} P_{R_X^n}) \right] Q_{X^m} S_{X^m} (\Gamma_{X'}^{-1/2})^{\otimes n} \\ &\leq e^{-n(M(\sigma_X \parallel \Gamma_X) - \delta)} (\Gamma_{X'}^{-1/2})^{\otimes n} S_{X^m} Q_{X^m} \text{tr}_{R_X^n} \left[R_{R_X^n} E_{X'R_X}^{\otimes n} R_{R_X^n} \right] Q_{X^m} S_{X^m} (\Gamma_{X'}^{-1/2})^{\otimes n}, \end{aligned} \quad (8.10)$$

recalling that $P_{R_X^n} \Gamma_{R_X}^{\otimes n} P_{R_X^n} \leq e^{-n(M(\sigma_X \parallel \Gamma_X) - \delta)} \mathbb{1}_{R_X^n}$. Now define $R_{X^m E^n}$ as the dual projector of $R_{R_X^n}$ with respect to $|E\rangle_{X'R_X E}^{\otimes n}$: Indeed, we have $|E\rangle_{X'R_X E} = V_{X \rightarrow X'E} |\Phi\rangle_{X:R_X}$; we may thus define $R_{X^m E^n} = (V_{X \rightarrow X'E})^{\otimes n} R_{X^n} (V^\dagger)^{\otimes n}$ in such a way that $R_{X^m E^n} |E\rangle_{X'R_X E}^{\otimes n} = V_{X \rightarrow X'E}^{\otimes n} (R_{X^n} \otimes \mathbb{1}_{R_X^n}) |\Phi\rangle_{X:R_X} = R_{R_X^n} |E\rangle_{X'R_X E}^{\otimes n}$. Then compute

$$\begin{aligned} & Q_{X^m} \text{tr}_{R_X^n} \left[R_{R_X^n} E_{X'R_X}^{\otimes n} R_{R_X^n} \right] Q_{X^m} \\ &= Q_{X^m} \text{tr}_{R_X^n E^n} \left[R_{X^m E^n} E_{X'R_X E}^{\otimes n} R_{X^m E^n} \right] Q_{X^m} \\ &= Q_{X^m} \text{tr}_{E^n} \left[R_{X^m E^n} E_{X'E}^{\otimes n} R_{X^m E^n} \right] Q_{X^m} \\ &\leq Q_{X^m} \text{tr}_{E^n} [R_{X^m E^n}] Q_{X^m} \\ &\leq e^{n(S(\rho_{X'E}) + \delta)} Q_{X^m} \text{tr}_{E^n} [R_{X^m E^n} \rho_{X'E}^{\otimes n} R_{X^m E^n}] Q_{X^m} \\ &\leq e^{n(S(\rho_{X'E}) + \delta)} Q_{X^m} \text{tr}_{E^n} [\rho_{X'E}^{\otimes n}] Q_{X^m} \\ &= e^{n(S(\rho_{X'E}) + \delta)} Q_{X^m} \rho_{X'}^{\otimes n} Q_{X^m} \\ &\leq e^{n(S(\rho_{X'E}) - S(\rho_{X'}) + 2\delta)} Q_{X^m} \\ &\leq e^{n(S(\sigma_X) - S(\rho_{X'}) + 2\delta)} \mathbb{1}_{X^m}. \end{aligned} \quad (8.11)$$

where we have used the fact that $E_{X'E} \leq \mathbb{1}_{X'E}$ (since $E_{R_X} \leq \mathbb{1}_{R_X}$), the usual properties of the typical projectors, the fact that $[R_{X'^n E^n}, \rho_{X'E}^{\otimes n}] = 0$, as well as the fact that $S(\rho_{X'E}) = S(\rho_{R_X}) = S(\sigma_X)$ because $\rho_{X'R_X E}$ is a pure state. We may then return to

$$\begin{aligned} (8.10) &\leq e^{-n(M(\sigma_X \parallel \Gamma_X) - S(\rho_{X'E}) + S(\rho_{X'}) - 3\delta)} S_{X'^n} (\Gamma_{X'}^{-1})^{\otimes n} S_{X'^n} \\ &\leq e^{-n(M(\sigma_X \parallel \Gamma_X) + M(\rho_{X'} \parallel \Gamma_{X'}^{-1}) - S(\rho_{X'E}) + S(\rho_{X'}) - 4\delta)} \mathbb{1}, \end{aligned} \quad (8.12)$$

recalling that $S_{X'^n}$ and $(\Gamma_{X'}^{-1})^{\otimes n}$ commute. A simple calculation then yields

$$\begin{aligned} &M(\sigma_X \parallel \Gamma_X) + M(\rho_{X'} \parallel \Gamma_{X'}^{-1}) - S(\sigma_X) + S(\rho_{X'}) \\ &= -\text{tr}(\sigma_X \ln \Gamma_X) + \text{tr}(\rho_{X'} \ln \Gamma_{X'}) + \text{tr}(\sigma_X \ln \sigma_X) - \text{tr}(\rho_{X'} \ln \rho_{X'}) \\ &= D(\sigma_X \parallel \Gamma_X) - D(\rho_{X'} \parallel \Gamma_{X'}) , \end{aligned} \quad (8.13)$$

which proves (8.7a). Now we go for (8.7b). Let $\eta_R, \eta_P, \eta_Q, \eta_S > 0$ be the corresponding parameters provided by Proposition 8.1 for $R_{X'^n}$, $P_{X'^n}$, $Q_{X'^n}$, and $S_{X'^n}$, and let $\eta = \min(\eta_R, \eta_P, \eta_Q, \eta_S)$. Then we may compute

$$\begin{aligned} &\text{Re} \left\{ \langle \rho |_{X'E R_X}^{\otimes n} (\sigma_{R_X}^{1/2})^{\otimes n} | T \rangle_{X'^n R_X^n E^n} \right\} \\ &= \text{Re} \left\{ \langle \rho |_{X'E R_X}^{\otimes n} S_{X'^n} Q_{X'^n} R_{X'^n E^n} P_{X'^n E^n} | \rho \rangle_{X'R_X E}^{\otimes n} \right\}, \end{aligned} \quad (8.14)$$

where analogously to $R_{X'^n E^n}$ we define $P_{X'^n E^n} = V^{\otimes n} P_{X'^n} (V^\dagger)^{\otimes n}$, and noting that $(\sigma_{R_X}^{1/2})^{\otimes n} | E \rangle_{X'R_X E}^{\otimes n} = | \rho \rangle_{X'R_X E}^{\otimes n}$. Compute

$$\begin{aligned} \left\| (\mathbb{1} - P_{X'^n E^n}) | \rho \rangle_{X'R_X E}^{\otimes n} \right\|^2 &= \langle \rho |_{X'R_X E}^{\otimes n} (\mathbb{1} - P_{X'^n E^n}) | \rho \rangle_{X'R_X E}^{\otimes n} \\ &= \text{tr}(\sigma_X (\mathbb{1} - P_{X'^n})) \leq 2 \exp(-n\eta); \end{aligned} \quad (8.15a)$$

$$\left\| (\mathbb{1} - R_{X'^n E^n}) | \rho \rangle_{X'R_X E}^{\otimes n} \right\|^2 = \text{tr}(\sigma_X (\mathbb{1} - R_{X'^n})) \leq 2 \exp(-n\eta); \quad (8.15b)$$

$$\left\| (\mathbb{1} - Q_{X'^n}) | \rho \rangle_{X'R_X E}^{\otimes n} \right\|^2 = \text{tr}(\rho_{X'} (\mathbb{1} - Q_{X'^n})) \leq 2 \exp(-n\eta); \quad (8.15c)$$

$$\left\| (\mathbb{1} - S_{X'^n}) | \rho \rangle_{X'R_X E}^{\otimes n} \right\|^2 = \text{tr}(\rho_{X'} (\mathbb{1} - S_{X'^n})) \leq 2 \exp(-n\eta); \quad (8.15d)$$

exploiting each time property (8.1c) of the corresponding relative typical projector. Now we use the fact that for any states $|\psi\rangle, |\psi'\rangle$, and for any $0 \leq X \leq \mathbb{1}$, we have $\text{Re}\{\langle \psi' | X | \psi \rangle\} = \text{Re}\{\langle \psi' | \psi \rangle\} - \text{Re}\{\langle \psi' | (\mathbb{1} - X) | \psi \rangle\} \geq \text{Re}\{\langle \psi' | \psi \rangle\} - \|\psi'\| \|\langle \mathbb{1} - X | \psi \rangle\|$, where the last inequality holds by Cauchy-Schwarz. Then

$$\begin{aligned} (8.14) &\geq \text{Re} \left\{ \langle \rho |_{X'E R_X}^{\otimes n} S_{X'^n} Q_{X'^n} R_{X'^n E^n} | \rho \rangle_{X'R_X E}^{\otimes n} \right\} - \sqrt{2} \exp(-n\eta/2) \\ &\geq \text{Re} \left\{ \langle \rho |_{X'E R_X}^{\otimes n} S_{X'^n} Q_{X'^n} | \rho \rangle_{X'R_X E}^{\otimes n} \right\} - 2\sqrt{2} \exp(-n\eta/2) \\ &\geq \dots \\ &\geq 1 - 4\sqrt{2} \exp(-n\eta/2). \end{aligned} \quad (8.16)$$

Since $|\rho\rangle_{X'R_X E}^{\otimes n}$ is a purification of $\rho_{X'R_X}^{\otimes n}$, and $(\sigma_{R_X}^{1/2})^{\otimes n}|T\rangle_{X'^n R_X^n E^n}$ is a purification of $\mathcal{T}_{X^n \rightarrow X'^n}(\sigma_{X R_X}^{\otimes n})$, we have

$$\begin{aligned} F(\rho_{X'R_X}^{\otimes n}, \mathcal{T}_{X^n \rightarrow X'^n}(\sigma_{X R_X}^{\otimes n})) &\geq |\langle \rho|_{X'E R_X}^{\otimes n} (\sigma_{R_X}^{1/2})^{\otimes n} |T\rangle_{X'^n R_X^n E^n}| \\ &\geq \text{Re}\{\langle \rho|_{X'E R_X}^{\otimes n} (\sigma_{R_X}^{1/2})^{\otimes n} |T\rangle_{X'^n R_X^n E^n}\} \\ &\geq 1 - 4\sqrt{2}\exp(-n\eta/2). \end{aligned} \quad (8.17)$$

Hence

$$P(\rho_{X'R_X}^{\otimes n}, \mathcal{T}_{X^n \rightarrow X'^n}(\sigma_{X R_X}^{\otimes n})) \leq \sqrt{1 - (1 - 4\sqrt{2}\exp(-n\eta/2))^2} \leq \sqrt{8\sqrt{2}\exp(-n\eta/4)}, \quad (8.18)$$

using the fact that $\sqrt{1 - (1 - x)^2} \leq \sqrt{2x}$, thus proving (8.7b) noting that $(8\sqrt{2})^{1/2} \leq 4$. \blacksquare

8.2 Optimal implementation of any i.i.d. channel with thermal operations on an i.i.d. input state

Here, we show that for any channel $\mathcal{E}_{X \rightarrow X'}$, mapping a system X with Hamiltonian H_X to a system X' with Hamiltonian $H_{X'}$, and for any fixed input state σ_X , then there exists an optimal implementation with thermal operations which uses a small amount of coherence, and an amount of work per copy which is asymptotically equal to

$$W_{\text{T.O.}} = \beta^{-1} D(\mathcal{E}(\sigma) \| e^{-\beta H_{X'}}) - \beta^{-1} D(\sigma \| e^{-\beta H_X}). \quad (8.19)$$

The coherence is counted using a half-infinite energy ladder with spacing x , i.e., with Hamiltonian $H_C = \sum_{k=0}^{d_C} kx|k\rangle\langle k|_C$, as considered in Ref. [68]. The state is initialized in the state $|\eta^L\rangle := L^{-1/2} \sum_{k=0}^{L-1} |\ell_0 + k\rangle$, where ℓ_0 is a base energy offset. Such a system may be consumed entirely by the process, i.e., at the end of the process we may return it in any state we like. We assume that, in some reasonable setting, such a state on such a system can be prepared using an amount of work of the same order as the dimension of the system d_C . In the following, d_C will be sublinear in the number of copies n , so asymptotically for $n \rightarrow \infty$, the coherence source will require negligible resources to create when counted per copy. Actually, we will use two such sources C_1, C_2 : We think of each of these as single-use disposable systems, and we need such systems at two stages in our protocol.

Proposition 8.3. *Let X, X' be systems, let $H_X, H_{X'}$ be the corresponding Hamiltonians, and let $\beta \geq 0$. Let $\mathcal{E}_{X \rightarrow X'}$ be a completely positive, trace-preserving map, and let σ_X be any input state. Let $R \simeq X$ and let $|\sigma\rangle_{XR} = \sigma_X^{1/2} |\Phi\rangle_{X:R}$ be a purification of σ_X . Then for any $0 < \theta < 1/3$, for any n and for any $\delta > 0$ there exists a thermal operation which acts on $X \rightarrow X'$ and an information battery W whose process matrix is ε -close to $\mathcal{E}_{X \rightarrow X'}^{\otimes n}(\sigma_{XR}^{\otimes n})$ and which uses two coherence sources C_1, C_2 and an*

amount of work $W_{\text{T.O.}}^n$, with

$$\frac{1}{n} W_{\text{T.O.}}^n \leq F(\rho_{X'}, H_{X'}) - F(\sigma_X, H_X) + O(n^{-1/2}) + O(n^{-1} \log(1/\theta)) ; \quad (8.20a)$$

$$\varepsilon \leq 3\theta + O(\exp(-cn\delta^2)) ; \quad (8.20b)$$

$$d_{C_i}/n \leq O(\delta/\theta^2) , \quad (8.20c)$$

with $F(\sigma, H) = D(\sigma \| e^{-\beta H})$ and for some $c > 0$ depending only on $H_X, H_{X'}$.

The following corollary is obtained straightforwardly from the above proposition by choosing $\delta = n^{-1/2+\xi}$, $\theta = n^{-\xi/2}$ for any choice of $0 < \xi < 1/4$.

Corollary 8.1. Any i.i.d. channel $\mathcal{E}_{X \rightarrow X'}^{\otimes n}$ between n copies of systems X, X' with Hamiltonians $H_X, H_{X'}$ can be implemented on a fixed i.i.d. input state $\sigma_X^{\otimes n}$ using thermal operations at a work cost rate per copy which is asymptotically equal to

$$W_{\text{T.O.}, \text{asympt.}} = F(\mathcal{E}(\sigma_X), H_{X'}) - F(\sigma_X, H_X) , \quad (8.21)$$

and using a vanishing amount of coherence per copy.

We need a technical lemma that tells us that whenever we have a Hamiltonian whose eigenvalues are all close, we may replace that Hamiltonian by an exactly flat Hamiltonian at a cost of investing an amount of coherence of the order of the energy spread we would like to flatten out.

Lemma 8.1 (Flattening Hamiltonians using [68]). *Let A be a system with a Hamiltonian H_A with m eigenvalues all lying in a range $[h_-, h_+]$. Let $B \simeq A$ be a system with a completely degenerate Hamiltonian $H_B = [(h_- + h_+)/2] \mathbb{1}_B$. We assume that the spacings of the eigenvalues of H_A as well as the value $(h_- + h_+)/2$ are multiple of some unit x . Fix $\theta > 0$. Consider a coherence source C of dimension d_C , with Hamiltonian $H_C = \sum_{k=0}^{d_C} kx|k\rangle\langle k|_C$. Assume that the coherence source starts in the state $|\eta\rangle_C = L^{-1/2} \sum_{k=0}^{L-1} |\ell_0 + k\rangle$ such that $\ell_0 \geq (h_- + h_+)/x$ and $L \geq \theta^{-2}(h_+ - h_-)/x$ and $d_C \geq L + \ell_0 + (h_- + h_+)/x$. Then there exists a partial isometry $U_{AC \rightarrow BC}$ which commutes exactly with the total Hamiltonians $(U_{AC \rightarrow BC}(H_A + H_C) = (H_B + H_C)U_{AC \rightarrow BC})$, such that for any ρ_{AR} on any reference system R , we have that $\rho_{AR} \otimes |\eta\rangle\langle\eta|_C$ is in the support of $U_{AC \rightarrow BC} \otimes \mathbb{1}_R$ and*

$$P\left(\text{tr}_C(U(\rho_{AR} \otimes \eta)U^\dagger), \rho_{BR}\right) \leq \theta , \quad (8.22)$$

where $\rho_{BR} = \text{id}_{A \rightarrow B}(\rho_{AR})$ denotes the same state as the initial state, but on the systems BR .

The reverse operation $B \rightarrow A$ may also be carried out with the consumption of a similar coherence source, at the same accuracy.

Proof of Lemma 8.1. First, we can reduce the problem to a system dimension m : Embed the system into a bipartite system with a ancilla with trivial Hamiltonian storing the degeneracy index, and a second system with nondegenerate Hamiltonian storing the actual energy. Then the problem reduces to change the Hamiltonian of the second system. So we may assume without loss of generality that the Hamiltonian H_A is nondegenerate with $m = d_A$.

Consider the protocol of Åberg [68]. We would like to apply a result in the spirit of Åberg's Supplementary Proposition 2, but we need a tighter bound. Denoting the energy levels of A by $H_A = \sum x z_j |j\rangle\langle j|$ for integers z_j , we apply the global energy-conserving unitary on A and C given by

$$U_{AC \rightarrow BC} = \sum_j |j\rangle_B \langle j|_A \otimes \Delta^{z' - z_j}, \quad (8.23)$$

where $z' = (h_- + h_+)/2x$ represents the fixed energy of the output. Then, starting in the state $|\psi\rangle_{AA'} = \sum \psi_{jj'} |j\rangle_{AA'}$ using a reference system $A' \simeq A$ and the initial state $|\eta\rangle_C$ on C , we have

$$U_{AC \rightarrow BC}(|\psi\rangle_{AA'} |\eta\rangle_C) = \sum_{jj'} \psi_{jj'} |j\rangle_{BA'} \otimes (\Delta^{z' - z_j} |\eta\rangle_C). \quad (8.24)$$

We may calculate the overlap with the initial state,

$$\text{Re}\{\langle \psi|_{BA'} \langle \eta|_C U_{AC \rightarrow BC} |\psi\rangle_{AA'} |\eta\rangle_C\} = \sum_{jj'} |\psi_{jj'}|^2 \text{Re}\{\langle \eta|_C \Delta^{z' - z_j} |\eta\rangle_C\} \geq 1 - \frac{h_+ - h_-}{xL}, \quad (8.25)$$

where we used the fact that $\text{tr}(\Delta^{-a} \eta_C) = \max(0, 1 - a/L)$ and $|z' - z_j| \leq (h_+ - h_-)/x$. Hence, using the fact that the partial trace can only increase the fidelity,

$$F^2(\Phi(|\psi\rangle\langle\psi|_{AA'}), |\psi\rangle\langle\psi|_{BA'}) \geq 1 - \frac{h_+ - h_-}{xL}, \quad (8.26)$$

where $\Phi(\cdot) = \text{tr}_C\{U_{AC \rightarrow BC}[(\cdot) \otimes |\eta\rangle\langle\eta|_C] U^\dagger\}$. Note that $|\psi\rangle_{AA'}$ is arbitrary at this point. For any state ρ_{AR} , using the joint concavity of the fidelity function, further noting that we may consider without loss of generality reference systems of the form $A' \simeq A$, we have

$$F^2(\Phi_{A \rightarrow B}(\rho_{AR}), \rho_{BR}) \geq \min_{|\psi\rangle_{AA'}} F^2(\Phi_{A \rightarrow B}(|\psi\rangle\langle\psi|_{AA'}), |\psi\rangle\langle\psi|_{BA'}) \geq 1 - \frac{h_+ - h_-}{xL}, \quad (8.27)$$

and thus, since $L \geq \theta^{-2}(h_+ - h_-)/x$,

$$P(\Phi_{A \rightarrow B}(\rho_{AR}), \rho_{BR}) \leq \sqrt{\frac{h_+ - h_-}{xL}} \leq \theta. \quad (8.28)$$

The same argument can be applied to the operation $B \rightarrow A$. ■

Proof of Proposition 8.3. Let $\{R_{X^n}^k\}$ be the POVM elements from Proposition 2.2 for the input energy (for the Hamiltonian H_X) over the n systems, and let $\{L_{X^n}^\ell\}$ be the corresponding output measurement.

We exhibit a protocol as a sequence of gentle measurements and thermal operations. For any $\delta > 0$, we measure the projector $R_{X^n}^{\approx_{\delta \text{tr}}(H_X \sigma_X)}$ on the n inputs. This measurement fails with probability $\leq 2 \exp(-\eta_1 n)$ for $\eta_1 = 2\delta^2/(\Delta H_X)^2$, where ΔH_X is the difference between the maximal and minimal eigenvalues of H_X .

Assume that the input and output Hamiltonians H_X and $H_{X'}$ have eigenvalues that are multiples

of a spacing x (x may be very small). Now the state lies in a subspace of energies in the interval $[n(\text{tr}(H_X \sigma_X) \pm \delta)]$. We invoke [Lemma 8.1](#) to change this n -system Hamiltonian to one which is entirely flat, $H'_{X^n} := h \mathbb{1}_{X^n}$ with $h = n \text{tr}(H_X \sigma_X)$. Given the target approximation parameter $\theta > 0$, the cost of this operation is the consumption of a coherence source C_1 of size $d_{C_1} = (\theta^{-2} + 2)(2n\delta/x) \leq O(n\delta/\theta^2)$, because $m = 2n\delta/x$.

Then we invoke the achievability result of Ref. [13], that one can implement any channel over trivial Hamiltonians using thermal operations, and an amount of work given approximately by the conditional entropy of the environment conditioned on the output. We use this step to implement the channel

$$S_{X^m}^{\approx \delta \text{tr}(H_{X'} \rho_{X'})} \mathcal{E}_{X^n \rightarrow X^m}^{\otimes n} (R_{X^n}^{\approx \delta \text{tr}(H_X \sigma_X)}(\cdot) R_{X^n}^{\approx \delta \text{tr}(H_X \sigma_X)}) S_{X^m}^{\approx \delta \text{tr}(H_{X'} \rho_{X'})} \quad (8.29)$$

up to an accuracy θ and investing an amount of work $nH(E|X')_\rho + O(\sqrt{n}) + \Delta(\theta)$ with $\Delta(\theta) = O(\log(1/\theta))$. We can now trivially shift the whole Hamiltonian $H'_{X^n} \rightarrow H'_{X^m} := h' \mathbb{1}_{X^m}$ with $h' = n \text{tr}(H_{X'} \rho_{X'})$, investing an amount of work $h' - h$.

Finally, we have the state $S_{X^m}^{\approx \delta \text{tr}(H_{X'} \rho_{X'})} \rho_{X'R}^{\otimes n} S_{X^m}^{\approx \delta \text{tr}(H_{X'} \rho_{X'})}$ but the Hamiltonian is still the trivial H'_{X^m} . Again we invoke [Lemma 8.1](#) to change to the final Hamiltonian, up to accuracy θ , by consuming another coherence source C_2 of size $d_{C_2} \leq O(n\delta/\theta^2)$.

Because the final state is $S_{X^m}^{\approx \delta \text{tr}(H_{X'} \rho_{X'})} \rho_{X'R}^{\otimes n} S_{X^m}^{\approx \delta \text{tr}(H_{X'} \rho_{X'})}$ instead of $\rho_{X'R}^{\otimes n}$, we again pay a “gentle measurement penalty” of $O(e^{-n\eta_2/2})$ where $\eta_2 \sim \delta^2$ (cf. [Lemma B.2](#)).

Finally, counting the total work, total failure probability and total use of coherence proves the claim, noting that $\text{tr}(H_{X'} \rho_{X'}) - \text{tr}(H_X \sigma_X) + H(E|X')_\rho = F(\rho_{X'}, H_{X'}) - F(\sigma_X, H_X)$. \blacksquare

8.3 Single-shot erasure protocol for fixed input state and for noninteracting system and memory

In the position-based decoding of Ref. [19], one uses the optimal distinguishing POVM for P_{SM} obtained from the hypothesis testing entropy, and we see that for a constant error we can choose $\ln(m)$ to be proportional to the hypothesis testing entropy. In fact, this gives us directly a new erasure protocol in the case of a fixed input state, and in the case where the system and memory are not interacting ($H_{SM} = H_S + H_M$):

Corollary 8.2. *Let S, M, R be quantum systems, with H_S and H_M the Hamiltonians on S and M . Let $|\rho\rangle_{SMR}$ be any pure state such that $[\rho_{SM}, H_S + H_M] = 0$. Let $\varepsilon > 0$. Suppose that $D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M) \geq \ln(2/\varepsilon)$. Then there exists an information battery J with battery states τ_J, τ'_J , and there exists a thermal operation $\mathcal{R}_{SMJ \rightarrow SMJ}$ acting on SMJ such that:*

- The effective work process $\mathcal{T}_{SM \rightarrow SM}$ associated with $\mathcal{R}_{SMJ \rightarrow SMJ}$ and (τ_J, τ'_J) satisfies

$$F(\mathcal{T}_{SM \rightarrow SM}(\rho_{SMR}), \gamma_S \otimes \rho_{MR}) \geq 1 - 6\varepsilon; \quad (8.30)$$

- The associated work cost (in units of pure nats) satisfies

$$w(\tau_J) - w(\tau'_J) \leq -D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M) + \ln(2/\varepsilon) . \quad (8.31)$$

If the register S is to be returned to a pure state instead of a thermal state, then this can be done separately as a final step, and there is a fixed cost associated to this. For instance, for trivial Hamiltonians, we have that

$$\begin{aligned} \text{total work extracted in nats,} \\ \text{also return } S \text{ to pure state} \end{aligned} \geq D_h^{1-\varepsilon} \left(\rho_{SM} \parallel \left\| \frac{\mathbb{1}_S}{d_S} \otimes \rho_M \right\| \right) - \ln(d_S) - \ln(2/\varepsilon) \\ = D_h^{1-\varepsilon}(\rho_{SM} \parallel \mathbb{1}_S \otimes \rho_M) - \ln(2/\varepsilon) , \quad (8.32)$$

and recalling that $D_h^{1-\varepsilon}(\rho_{SM} \parallel \mathbb{1}_S \otimes \rho_M) \approx -H_{\max}^\varepsilon(S|M)_\rho$ [10, 34], we recover the expression in Ref. [63] up to approximation terms.

Proof of Corollary 8.2. Let P_{SM} be the optimal positive semidefinite operator satisfying $0 \leq P_{SM} \leq \mathbb{1}_{SM}$ given by $D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)$, i.e., such that

$$\text{tr}(P_{SM} \rho_{SM}) \geq 1 - \varepsilon ; \quad (8.33)$$

$$\text{tr}(P_{SM} (\gamma_S \otimes \rho_M)) = \exp\{-D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)\} . \quad (8.34)$$

We can always take $[P_{SM}, H_S + H_M] = 0$ by dephasing P_{SM} in energy blocks, if necessary, since ρ_{SM} is time-covariant. Let

$$m = \ln \left[\varepsilon \exp\{D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)\} \right] . \quad (8.35)$$

Then $e^m \leq \varepsilon \exp\{D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)\}$ and

$$\text{tr}(P_{SM} (\gamma_S \otimes \rho_M)) \leq \frac{\varepsilon}{e^m} . \quad (8.36)$$

We may set $\kappa = \kappa' = \varepsilon$, and we are in the setting of Proposition 7.1 for the single-element set $\mathcal{S}_{SM} = \{\rho_{SM}\}$. Proposition 7.1 ensures that there exists a thermal operation $\mathcal{R}_{SMJ \rightarrow SMJ}$ and battery states τ_J, τ'_J that satisfies the condition (8.30). Furthermore according to Proposition 7.1, the associated work cost is

$$\begin{aligned} w(\tau_J) - w(\tau'_J) &= -m \leq -\ln \left(\varepsilon \exp\{D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)\} - 1 \right) \\ &= -D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M) - \ln(\varepsilon) - \ln \left(1 - \frac{1}{\varepsilon} e^{-D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)} \right) \\ &\leq -D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M) - \ln(\varepsilon) - \ln(1/2) , \end{aligned} \quad (8.37)$$

using in the last inequality our assumption that $\varepsilon \exp\{D_h^{1-\varepsilon}(\rho_{SM} \parallel \gamma_S \otimes \rho_M)\} \geq 2$. We have proven (8.31), thereby completing the proof of Corollary 8.2. ■

8.4 Single-shot protocol for implementing a time-covariant processes using thermal operations

We can translate our protocol for conditional erasure into a protocol for implementing more general processes. (A similar approach was taken in Ref. [13] in the case of trivial Hamiltonians). We obtain the following result.

Corollary 8.3 (Single-shot protocol for a time-covariant process and input state). *Let X be a quantum system with a Hamiltonian H_X , and let $\beta \geq 0$. Let $\mathcal{E}_{X \rightarrow X}$ be a completely positive, trace-preserving map satisfying*

$$\mathcal{E}_{X \rightarrow X}(e^{-iH_X t}(\cdot)e^{iH_X t}) = e^{-iH_X t} \mathcal{E}_{X \rightarrow X}(\cdot) e^{iH_X t} \quad \text{for all } t \in \mathbb{R}. \quad (8.38)$$

Let σ_X be any quantum state that satisfies $[\sigma_X, H_X] = 0$ and let $\varepsilon > 0$. There exists an information battery W , a thermal operation Φ_{XW} and battery states τ_W, τ'_W such that:

- *The effective work process $\mathcal{T}_{X \rightarrow X}$ associated with Φ_{XW} and (τ_W, τ'_W) satisfies*

$$F(\mathcal{T}_{X \rightarrow X}(|\sigma\rangle\langle\sigma|_{XR}), \mathcal{E}_{X \rightarrow X}(|\sigma\rangle\langle\sigma|_{XR})) \geq 1 - 6\varepsilon, \quad (8.39)$$

where $|\sigma\rangle_{XR}$ is a purification of σ_X ;

- *The work cost (in units of pure nats) satisfies*

$$w(\tau_W) - w(\tau'_W) \leq -D_h^{1-\varepsilon}(\rho_{EX} \parallel \gamma_E \otimes \rho_X) + \ln(\text{tr} e^{-\beta H_X}) + \ln(2/\varepsilon), \quad (8.40)$$

where E is an additional system with Hamiltonian H_E and where $\rho_{EX} = V(|0\rangle\langle 0|_E \otimes \sigma_X)V^\dagger$ is the output of a time-covariant Stinespring unitary $V_{XE \rightarrow XE}$ of $\mathcal{E}_{X \rightarrow X}$ with $|0\rangle_E$ an eigenstate of H_E with eigenvalue 0; i.e., the operator $V_{X \rightarrow E}$ satisfies $\mathcal{E}_{X \rightarrow X}(\cdot) = \text{tr}_E[V(|0\rangle\langle 0|_E \otimes (\cdot))V^\dagger]$. We note that $\rho_X = \mathcal{E}_{X \rightarrow X}(\sigma_X)$.

Observe that for any time-covariant process $\mathcal{E}_{X \rightarrow X}$, the existence of $V_{XE \rightarrow XE}$ with the desired properties is guaranteed by Lemma 7.2.

Proof. We start with the system X and a reference system R in the pure state $|\sigma_{XR}\rangle$. The protocol is assembled using the following steps:

- Bring in an ancilla E which can be used as the Stinespring environment system of $\mathcal{E}_{X \rightarrow X}$. The system E is initialized in its thermal state γ_E . This is a free operation;
- Reset the ancilla E to the state $|0\rangle_E$, using Proposition 3.2, at a work cost arbitrarily close to $\ln \text{tr}[e^{-\beta H_X}]$;
- Apply the energy-conserving unitary $V_{XE \rightarrow XE}$ at no work cost. The resulting state on XE is ρ_{XE} ;

- Invoke [Corollary 8.2](#) to transform the state ρ_{XE} into a state ρ'_{XE} that is close to $\gamma_E \otimes \rho_X$, with

$$F(\rho'_{XE}, \gamma_E \otimes \rho_X) \geq 1 - 6\epsilon, \quad (8.41)$$

and at a work cost w satisfying $w \leq -D_h^{1-\epsilon}(\rho_{EM} \| \gamma_E \otimes \rho_M) + \ln(2/\epsilon)$;

- Discard the system E , at no work cost.

The condition (8.39) is satisfied by construction. Furthermore, collecting the contributions to the work cost yields (8.40). \blacksquare

9 Discussion

Our results fits in the line of research extending results in thermodynamics from state-to-state transformations to quantum processes. Implementations of quantum processes are difficult to construct because they need to reproduce the correct correlations between the output and the reference system, and not only produce the correct output state. Here, we have seen that it is nevertheless possible to implement any quantum process at an optimal work cost: Any implementation that would use less work would violate the second law of thermodynamics on a macroscopic scale. As a special case this also provides an operational interpretation of the minimal entropy gain of a channel [42–49].

Our three constructions of optimal implementations of processes are valid in different settings, and it remains unclear if they can be unified in a single protocol that presents the advantages of all three constructions. Namely, is it possible to use a physically well-justified framework, e.g. thermal operations, to universally implement any i.i.d. process? We expect this to be possible only if an arbitrary amount of coherence is allowed, in analogy with the entanglement embezzling state required in the reverse Shannon theorem [22, 23].

Finally, the notion of quantum typicality that we have introduced in [Definition 6.1](#) and [Proposition 6.1](#) might be interesting in its own right. We anticipate that similar considerations might provide pathways to smooth other information-theoretic quantities [61, 69, 70] and to study the joint typicality conjecture [26, 71–74].

Acknowledgements. The authors thank Álvaro Alhambra, David Ding, Patrick Hayden, Rahul Jain, David Jennings, Martí Perarnau-Llobet, Mark Wilde, and Andreas Winter for discussions. PhF acknowledges support from the Swiss National Science Foundation (SNSF) through the Early PostDoc.Mobility Fellowship No. P2EZP2_165239 hosted by the Institute for Quantum Information and Matter (IQIM) at Caltech, from the IQIM which is a National Science Foundation (NSF) Physics Frontiers Center (NSF Grant PHY-1733907), from the Department of Energy Award DE-SC0018407, from the Swiss National Science Foundation (SNSF) through the NCCR QSIT and through Project No. 200020_16584, and from the Deutsche Forschungsgemeinschaft (DFG) Research Unit FOR 2724. FB is supported by the NSF. This work was completed prior to MB and FB joining the AWS Center for Quantum Computing.

Appendix

A Missing proofs

Proof of Lemma 2.2. A useful expression for $\Pi_{A^n B^n}^\lambda$ may be obtained following [25, Section V]

$$\begin{aligned}\Pi_{A^n B^n}^\lambda &= \frac{\dim(\mathcal{Q}_\lambda)}{s_\lambda(\text{diag}(\lambda/n))} \int dU_{AB} \Pi_{A^n B^n}^\lambda \left(U_{AB} \text{diag}(\lambda/n)_{AB} U_{AB}^\dagger \right)^{\otimes n} \Pi_{A^n B^n}^\lambda \\ &\leq \text{poly}(n) e^{n\bar{H}(\lambda)} \int dU_{AB} \left(U_{AB} \text{diag}(\lambda/n)_{AB} U_{AB}^\dagger \right)^{\otimes n},\end{aligned}\quad (\text{A.1})$$

recalling that $\Pi_{A^n B^n}^\lambda$ commutes with any i.i.d. state, with $s_\lambda(X) = \text{tr}[q_\lambda(X)]$ and using bounds on $\dim(\mathcal{Q}_\lambda)$ and $s_\lambda(\text{diag}(\lambda/n))$ derived in Ref. [25]. Here, dU_{AB} denotes the Haar measure over all unitaries acting on \mathcal{H}_{AB} , normalized such that $\int dU_{AB} = 1$. We then have

$$\text{tr}_{A^n} \left[\Pi_{A^n B^n}^\lambda \right] \leq \text{poly}(n) e^{n\bar{H}(\lambda)} \int dU_{AB} \text{tr}_{A^n} \left[\left(U_{AB} \text{diag}(\lambda/n)_{AB} U_{AB}^\dagger \right)^{\otimes n} \right]. \quad (\text{A.2})$$

Observe that for any state ω_B , we have

$$\begin{aligned}\| \Pi_{B^n}^{\lambda'} \omega_B^{\otimes n} \Pi_{B^n}^{\lambda'} \|_\infty &= \| [q_{\lambda'}(\omega_B) \otimes \mathbf{1}_{\mathcal{H}_{\lambda'}}]_{\lambda'} \|_\infty = \| q_{\lambda'}(\omega_B) \|_\infty \leq \text{tr}[q_{\lambda'}(\omega_B)] \\ &\leq \text{poly}(n) e^{-n\bar{H}(\lambda')} \quad (\text{A.3})\end{aligned}$$

as derived e.g. in [25, Eq. (9)], and thus for any state ω_B ,

$$\Pi_{B^n}^{\lambda'} \omega_B^{\otimes n} \Pi_{B^n}^{\lambda'} \leq \text{poly}(n) e^{-n\bar{H}(\lambda')} \Pi_{B^n}^{\lambda'}. \quad (\text{A.4})$$

Hence, we get

$$\begin{aligned}\Pi_{B^n}^{\lambda'} \text{tr}_{A^n} \left[\Pi_{A^n B^n}^\lambda \right] \Pi_{B^n}^{\lambda'} &\leq \text{poly}(n) e^{n\bar{H}(\lambda)} \int dU_{AB} \Pi_{B^n}^{\lambda'} \left(\text{tr}_A \left[U_{AB} \text{diag}(\lambda/n)_{AB} U_{AB}^\dagger \right] \right)^{\otimes n} \Pi_{B^n}^{\lambda'} \\ &\leq \text{poly}(n) e^{n\bar{H}(\lambda)} \int dU_{AB} \text{poly}(n) e^{-n\bar{H}(\lambda')} \Pi_{B^n}^{\lambda'} \\ &= \text{poly}(n) e^{n(\bar{H}(\lambda) - \bar{H}(\lambda'))} \Pi_{B^n}^{\lambda'},\end{aligned}\quad (\text{A.5})$$

as required. ■

Proof of Proposition 2.1. The Fannes-Audenaert continuity bound [75, 76] of the entropy states that for any $\delta' > 0$ there exists $\xi(\delta') > 0$ such that for any quantum states ρ, σ with $D(\rho, \sigma) \leq \delta'$ we have

$$|H(\rho) - H(\sigma)| \leq \xi(\delta'), \quad (\text{A.6})$$

and furthermore $\xi(\delta')$ is monotonically strictly decreasing and $\xi(\delta') \rightarrow 0$ if $\delta' \rightarrow 0$. Now, let $\delta > 0$, let ξ^{-1} be the inverse function of ξ , and let $\delta' = \xi^{-1}(\delta)$. Consider the set of Young diagrams $\Lambda_{\delta'} = \{\lambda \in \text{Young}(d_A, n) : D(\text{diag}(\lambda/n), \rho) \leq \delta'\}$. For all $\lambda \in \Lambda_{\delta'}$, we have that $|H(\rho) - \bar{H}(\lambda)| \leq \delta$ thanks to the Fannes-Audenaert inequality. Then, we have

$$\text{tr} \left[\left(\sum_{\lambda: \bar{H}(\lambda) \in [H(\rho) \pm \delta]} \Pi_{A^n}^\lambda \right) \rho_A^{\otimes n} \right] \geq \text{tr} \left[\left(\sum_{\lambda \in \Lambda_{\delta'}} \Pi_{A^n}^\lambda \right) \rho_A^{\otimes n} \right] \quad (\text{A.7})$$

because all terms in the sum in the right hand side are included in the sum on the left hand side. We may now invoke [24, Eq. (6.23)] to see that

$$(\text{A.7}) \geq 1 - \text{poly}(n) \exp\{-n\eta\}, \quad (\text{A.8})$$

where $\eta = \delta'^2/2$. ■

Proof of Proposition 2.2. The fact that there are only $\text{poly}(n)$ elements follows because there are only so many types. Property (ii) holds by definition. Property (iv) holds because $e^{-n(k \pm \delta)}$ is the minimum / maximum eigenvalue of $\Gamma_A^{\otimes n}$ in the subspace spanned by $R_{A^n}^{\approx \delta^h}$. Finally, we need to show Property (iii): This follows from a large deviation analysis. More precisely, let Z_j for $j = 1, \dots, n$ be random variables where Z_j represents the measurement outcome of H_A on the j -th system of the i.i.d. state $\rho_A^{\otimes n}$. By Hoeffding's inequality, we have that

$$\Pr \left[\left| (1/n) \sum Z_j - \text{tr}[\rho_A H_A] \right| > \delta \right] \leq 2 \exp \left(-\frac{2n\delta^2}{\Delta H_A^2} \right) \leq 2 \exp \left(-\frac{n\delta^2}{2\|H_A\|_\infty^2} \right), \quad (\text{A.9})$$

where ΔH_A is the difference between the maximum and minimum eigenvalue of H_A , and $\Delta H_A \leq 2\|H_A\|_\infty$. Thus, the event consisting of the outcomes k satisfying $|k - \text{tr}[\rho_A H_A]| \leq \delta$ happens with probability at least $1 - 2e^{-n\eta}$, proving (2.19). ■

Proof of Proposition 2.3. We use the post-selection technique (Theorem 2.1) to bound the diamond norm distance between $\mathcal{T}_{X^n \rightarrow X^n}$ and $\mathcal{E}_{X \rightarrow X'}^{\otimes n}$. Let $|\zeta\rangle_{X^n \bar{R}^n R'}$ be the purification of the de Finetti state given by (2.24). Calculate

$$\begin{aligned} & \text{Re} \left\{ \langle \zeta |_{X^n \bar{R}^n R'} (V_{X \rightarrow EX'}^{\otimes n})^\dagger W_{X^n \rightarrow E^n X^n} | \zeta \rangle_{X^n \bar{R}^n R'} \right\} \\ &= \sum p_i \text{Re} \left\{ \langle \phi_i |_{X \bar{R}}^{\otimes n} (V_{X \rightarrow EX'}^{\otimes n})^\dagger W_{X^n \rightarrow E^n X^n} | \phi_i \rangle_{X \bar{R}}^{\otimes n} \right\} \\ &\geq 1 - \text{poly}(n) \exp(-n\eta) \end{aligned} \quad (\text{A.10})$$

which implies, recalling that $F(|\psi\rangle, |\phi\rangle) = |\langle \psi | \phi \rangle| \geq \text{Re}\{\langle \psi | \phi \rangle\}$ and that $(1-x)^2 \geq 1-2x$,

$$F^2(V_{X \rightarrow EX'}^{\otimes n} | \zeta \rangle_{X^n \bar{R}^n R'}, W_{X^n \rightarrow E^n X^n} | \zeta \rangle_{X^n \bar{R}^n R'}) \geq 1 - \text{poly}(n) \exp(-n\eta) \quad (\text{A.11})$$

and hence

$$P(V_{X \rightarrow EX'}^{\otimes n} | \zeta \rangle_{X^n \bar{R}^n R'}, W_{X^n \rightarrow E^n X^n} | \zeta \rangle_{X^n \bar{R}^n R'}) \leq \text{poly}(n) \exp(-n\eta/2). \quad (\text{A.12})$$

Recalling the relations between the trace distance and the purified distance, and noting that these distance measures cannot increase under the partial trace, we obtain

$$\begin{aligned} D(\mathcal{T}(\zeta_{X^n \bar{R}^n R'}), \mathcal{E}^{\otimes n}(\zeta_{X^n \bar{R}^n R'})) &\leq P(\mathcal{T}(\zeta_{X^n \bar{R}^n R'}), \mathcal{E}^{\otimes n}(\zeta_{X^n \bar{R}^n R'})) \\ &\leq P(W_{X^n \rightarrow E^n X^n} |\zeta\rangle_{X^n \bar{R}^n R'}, V_{X \rightarrow EX'}^{\otimes n} |\zeta\rangle_{X^n \bar{R}^n R'}) \leq \text{poly}(n) \exp(-n\eta/2). \end{aligned} \quad (\text{A.13})$$

The post-selection technique then asserts that

$$\frac{1}{2} \|\mathcal{T} - \mathcal{E}^{\otimes n}\|_{\diamond} \leq \text{poly}(n) \exp(-n\eta/2) \quad (\text{A.14})$$

as claimed. \blacksquare

Proof of Lemma 7.2. Let $V'_{X \rightarrow XE}$ be any Stinespring dilation isometry of $\mathcal{E}_{X \rightarrow X}$, such that $\mathcal{E}_{X \rightarrow X}(\cdot) = \text{tr}_E[V'_{X \rightarrow XE}(\cdot)V'^{\dagger}]$. For the input state $|\Phi\rangle_{X:R_X}$, consider the output state $|\varphi\rangle_{XER_X}$ corresponding to first time-evolving by some time t , and then applying V'

$$|\varphi\rangle_{XER_X} = V' e^{-iH_X t} |\Phi\rangle_{X:R_X} = e^{-iV' H_X V'^{\dagger} t} V' |\Phi\rangle_{X:R_X}. \quad (\text{A.15})$$

Now, let us define $|\varphi'\rangle_{XER_X} = e^{-iH_X t} V' |\Phi\rangle_{X:R_X}$. By the covariance property of $\mathcal{E}_{X \rightarrow X}$ both $|\varphi\rangle$ and $|\varphi'\rangle$ have the same reduced state on XR_X . Hence, they are related by some unitary $W_E^{(t)}$ on the system E which in general depends on t

$$|\varphi\rangle_{XER_X} = W_E^{(t)} |\varphi'\rangle_{XER_X}. \quad (\text{A.16})$$

We have

$$\text{tr}_X[V' e^{-iH_X t} \Phi_{X:R_X} e^{iH_X t} V'^{\dagger}] = W_E^{(t)} \text{tr}_X[V' \Phi_{X:R_X} V'^{\dagger}] W_E^{(t)\dagger} \quad (\text{A.17})$$

so $W_E^{(t)}$ must define a representation of time evolution, at least on the support of the operator $\text{tr}_X[V' \Phi_{X:R_X} V'^{\dagger}]$. Hence, we may write $W_E^{(t)} = e^{-iH_E t}$ for some Hamiltonian H_E , and from (A.16), we have for all t

$$V'_{X \rightarrow XE} e^{-iH_X t} = e^{-i(H_X + H_E)t} V'_{X \rightarrow XE}. \quad (\text{A.18})$$

Expanding for infinitesimal t we obtain

$$V'_{X \rightarrow XE} H_X = (H_X + H_E) V'_{X \rightarrow XE}. \quad (\text{A.19})$$

Let $|0\rangle_E$ be an eigenvector of H_E corresponding to the eigenvalue zero; if H_E does not contain an eigenvector with eigenvalue equal to zero, we may trivially add a dimension to the system E to accommodate this vector. Then, the operator $V'_{X \rightarrow XE} \langle 0|_E$ maps each state of a subset of energy levels of XE to a corresponding energy level of same energy on XE ; it may thus be completed to a fully energy-preserving unitary $V_{XE \rightarrow XE}$. More precisely, let $|j\rangle_X$ be a complete set of eigenvectors of H_X with energies h_j . Then $|\psi'_j\rangle = V'_{X \rightarrow XE} |j\rangle_X$ is an eigenvector of $H_X + H_E$ of energy h_j thanks

to (A.19). We have two orthonormal sets $\{|0\rangle_E \otimes |j\rangle_X\}$ and $\{|\psi'_j\rangle_X\}$ in which the j -th vector of each set has the same energy; we can thus complete these sets into two bases $\{|\chi_i\rangle_{XE}\}$, $\{|\chi'_i\rangle_{XE}\}$ of eigenvectors of $H_X + H_E$, where the i -th element of either basis has exactly the same energy. This defines a unitary $V_{XE \rightarrow XE} = \sum_i |\chi'_i\rangle_{XE} \langle \chi_i|_{XE}$ that is an extension of $V'_{X \rightarrow XE} \langle 0|_E$, and that satisfies all the conditions of the claim. \blacksquare

B Technical lemmas

Lemma B.1 (Pinching-like operator inequality). *Let $\{E^i\}_{i=1}^M$ be a collection of M operators and $T \geq 0$. Then, we have*

$$\left(\sum E^i\right) T \left(\sum E^{j\dagger}\right) \leq M \sum E^i T E^{i\dagger}. \quad (\text{B.1})$$

Proof. Call our system S and consider an additional register C of dimension $|C| = M$, and let $|\chi\rangle_C = M^{-1/2} \sum_{k=1}^M |k\rangle_C$. Then, we have

$$\begin{aligned} \left(\sum E_S^i\right) T_S \left(\sum E_S^{j\dagger}\right) &= \text{tr}_C \left[\left(\sum E_S^i \otimes |i\rangle_C\right) T_S \left(\sum E_S^{j\dagger} \otimes \langle j|_C\right) (\mathbb{1}_S \otimes (M |\chi\rangle_C \langle \chi|_C)) \right] \\ &\leq M \text{tr}_C \left[\left(\sum E_S^i \otimes |i\rangle_C\right) T_S \left(\sum E_S^{j\dagger} \otimes \langle j|_C\right) (\mathbb{1}_S \otimes \mathbb{1}_C) \right] \\ &= M \sum E_S^i T_S E_S^{i\dagger}, \end{aligned} \quad (\text{B.2})$$

using $|\chi\rangle_C \langle \chi|_C \leq \mathbb{1}_C$. \blacksquare

Lemma B.2 (Gentle measurement). *Let ρ be a sub-normalized quantum state and $0 \leq Q \leq \mathbb{1}$. For $\text{tr}[Q\rho] \geq 1 - \delta$ we then have*

$$P(\rho, Q^{1/2} \rho Q^{1/2}) \leq \sqrt{2\delta}. \quad (\text{B.3})$$

This is a cruder statement than that of, e.g., [77, Lemma 7], allowing for a more straightforward proof.

Proof. We have

$$\begin{aligned} \bar{F}(\rho, Q^{1/2} \rho Q^{1/2}) &\geq F(\rho, Q^{1/2} \rho Q^{1/2}) = \text{tr} \left[\sqrt{\rho^{1/2} (Q^{1/2} \rho Q^{1/2}) \rho^{1/2}} \right] \\ &= \text{tr} \left[Q^{1/2} \rho \right] \geq \text{tr}[Q\rho] \geq 1 - \delta. \end{aligned} \quad (\text{B.4})$$

Then, we get $P(\rho, Q^{1/2} \rho Q^{1/2}) \leq \sqrt{1 - (1 - \delta)^2} \leq \sqrt{2\delta}$. \blacksquare

Proposition B.1 (Controlled-unitary using a POVM). *Let $\{Q^j\}$ be a set of positive semi-definite operators on a system X satisfying $\sum Q^j \leq \mathbb{1}$, $\{U^j\}$ be a collection of unitaries on a system Y , and*

$$W_{XY} = \sum_j Q_X^j \otimes U_Y^j. \quad (\text{B.5})$$

Then, we have $W^\dagger W \leq \mathbb{1}$.

Proof. Using an additional register K , define

$$V_{X \rightarrow XK} = \sum_j \sqrt{Q^j} \otimes |j\rangle_K. \quad (\text{B.6})$$

Then, we have $V^\dagger V = \sum Q^j \leq \mathbb{1}$. Clearly, $VV^\dagger \leq \mathbb{1}_{XK}$ because VV^\dagger and $V^\dagger V$ have the same non-zero eigenvalues. Now, let

$$W = V^\dagger \left(\sum \mathbb{1}_X \otimes U_Y^j \otimes |j\rangle\langle j|_K \right) V. \quad (\text{B.7})$$

Because the middle term in parentheses is unitary, we manifestly have $W^\dagger W \leq \mathbb{1}$. ■

C Dilation of energy-conserving operators to unitaries

This appendix collects a few technical lemmas on constructing an energy-conserving unitary that extends a given operator of norm less than one.

Proposition C.1. *Let W_X be an operator on a system X , such that $W^\dagger W \leq \mathbb{1}$. Then, there exists a unitary operator U_{XQ} acting on X and a qubit Q such that for any $|\psi\rangle_X$,*

$$\langle 0|_Q U_{XQ} (|\psi\rangle_X \otimes |0\rangle_Q) = W_X |\psi\rangle_X. \quad (\text{C.1})$$

That is, any operator W with $\|W\|_\infty \leq 1$ can be dilated to a unitary, with a post-selection on the output.

Proof. Setting $V_{X \rightarrow XQ} = W \otimes |0\rangle_Q + \sqrt{\mathbb{1} - W^\dagger W} \otimes |1\rangle_Q$, we see that $V^\dagger V = W^\dagger W + \mathbb{1} - W^\dagger W = \mathbb{1}_X$, and hence $V_{X \rightarrow XQ}$ is an isometry. We can complete this isometry to a unitary U_{XQ} that acts as V on the support of $\mathbb{1}_X \otimes |0\rangle\langle 0|_Q$ and that maps the support of $\mathbb{1}_X \otimes |1\rangle\langle 1|_Q$ onto the complementary space to the image of V . It then follows that for any $|\psi\rangle_X$, we have $U_{XQ} (|\psi\rangle_X \otimes |0\rangle_Q) = V_{X \rightarrow XQ} |\psi\rangle_X = (W_X |\psi\rangle_X) \otimes |0\rangle_Q + (\dots) \otimes |1\rangle_Q$, and the claim follows. ■

Proposition C.2. *Let X be a quantum system with Hamiltonian H_X and W_X be an operator with $W^\dagger W \leq \mathbb{1}$ as well as $[W_X, H_X] = 0$. Then, there exists a unitary operator U_{XQ} acting on X and a*

qubit Q with $H_Q = 0$, that satisfies $[U_{XQ}, H_X] = 0$ such that

$$\langle 0|_Q U_{XQ} |0\rangle_Q = W_X. \quad (\text{C.2})$$

That is, any energy-preserving operator W with $\|W\|_\infty \leq 1$ can be dilated to an energy-preserving unitary on an ancilla with a post-selection on the output.

Proof. First we calculate $[W^\dagger W, H_X] = W^\dagger [W, H_X] + [W^\dagger, H_X] W = 0 - [W, H_X]^\dagger W = 0$. This implies that $[\sqrt{\mathbb{1} - W^\dagger W}, H_X] = 0$, as $W^\dagger W$ and $\sqrt{\mathbb{1} - W^\dagger W}$ have the same eigenspaces. We define

$$V_{X \rightarrow XQ} = W \otimes |0\rangle_Q + \sqrt{\mathbb{1} - W^\dagger W} \otimes |1\rangle_Q. \quad (\text{C.3})$$

The operator $V_{X \rightarrow XQ}$ is an isometry, because $V^\dagger V = W^\dagger W + \mathbb{1} - W^\dagger W = \mathbb{1}_X$. Furthermore, we have

$$V_{X \rightarrow XQ} H_X = (W_X H_X) \otimes |0\rangle + (\sqrt{\mathbb{1} - W^\dagger W} H_X) \otimes |1\rangle \quad (\text{C.4})$$

$$= (H_X W_X) \otimes |0\rangle + (H_X \sqrt{\mathbb{1} - W^\dagger W}) \otimes |1\rangle = H_X V_{X \rightarrow XQ} \quad (\text{C.5})$$

and thus we find $[V_{X \rightarrow XQ}, H_X] = 0$. Let $\{|j\rangle_X\}$ be an eigenbasis of H_X , and let $|\psi'_j\rangle_{XQ} = V_{X \rightarrow XQ} |j\rangle_X$, noting that both $|j\rangle_X$ and $|\psi'_j\rangle_{XQ}$ have the same energy. The two collections of vectors $\{|j\rangle_X \otimes |0\rangle_Q\}$ and $\{|\psi'_j\rangle_{XQ}\}$ can thus be completed into two bases $\{|\chi_i\rangle_{XQ}\}$ and $\{|\chi'_i\rangle_{XQ}\}$ of eigenvectors of $H_X + H_Q$ where the i -th element of both bases have the same energy. Define finally $U_{XQ} = \sum_i |\chi'_i\rangle\langle\chi_i|_{XQ}$, noting that by construction $U_{XQ} |0\rangle_Q = V_{X \rightarrow XQ}$ and $[U_{XQ}, H_X] = 0$. ■

Proposition C.3. *Let X be a quantum system with Hamiltonian H_X and, W_X an operator satisfying $W^\dagger W \leq \mathbb{1}$ such that $[W_X, H_X] = 0$. Then, for any $\varepsilon > 0$, there exists a unitary operator U_X satisfying $[U_X, H_X] = 0$ such that for any states $|\psi\rangle_X, |\psi'\rangle_X$ satisfying $\text{Re}\{\langle\psi'|W|\psi\rangle\} \geq 1 - \varepsilon$, we have $\text{Re}\{\langle\psi'|U|\psi\rangle\} \geq 1 - 6\varepsilon^{1/4}$.*

Proof of Proposition C.3. Let $F = W^\dagger W \leq \mathbb{1}$ noting that $F^\dagger = F$. For some ν with $0 < \nu < 1$ to be determined later, let P be the projector onto the eigenspaces of F corresponding to eigenvalues greater or equal to ν . Define $V = WF^{-1/2}P$. The operator V is a partial isometry, meaning that its singular values are all equal to one or to zero, because $V^\dagger V = PF^{-1/2}W^\dagger WF^{-1/2}P = PF^{-1/2}FF^{-1/2}P = P$, since P lies within the support of F . Observe that $[F, H] = [W^\dagger W, H] = W^\dagger [W, H] - [W, H]^\dagger W = 0$ and hence $[F^{-1/2}, H] = 0$ and $[P, H] = 0$. Hence, we have $[V, H] = [WF^{-1/2}P, H] = 0$. So we may complete the partial isometry V into a full unitary U that also commutes with H by acting as the identity on the remaining elements of the eigenbasis of H in which V is diagonal. We may thus write $U = V + X = WF^{-1/2}P + X$ for some operator X satisfying $XP = 0$ and $[X, H] = 0$.

Now, let $|\psi\rangle, |\psi'\rangle$ such that $\text{Re}\{\langle\psi'|W|\psi\rangle\} \geq 1 - \varepsilon$, and write

$$\begin{aligned} \text{Re}\{\langle\psi'|U|\psi\rangle\} &= \text{Re}\{\langle\psi'|(U - W)|\psi\rangle\} + \text{Re}\{\langle\psi'|W|\psi\rangle\} \\ &\geq \text{Re}\{\langle\psi'|(U - W)|\psi\rangle\} + 1 - \varepsilon. \end{aligned} \quad (\text{C.6})$$

We have $\langle\psi|P|\psi\rangle \geq \langle\psi|PFP|\psi\rangle = \langle\psi|F|\psi\rangle - \langle\psi|(\mathbb{1} - P)F|\psi\rangle \geq \langle\psi|F|\psi\rangle - \nu$, recalling that $\mathbb{1} - P$ projects onto the eigenspaces of F whose eigenvalues are less than ν . Then,

$\langle \psi | F | \psi \rangle = \langle \psi | W^\dagger W | \psi \rangle \geq \langle \psi | W^\dagger | \psi' \rangle \langle \psi' | W | \psi \rangle = |\langle \psi' | W | \psi \rangle|^2 \geq (1 - \varepsilon)^2 \geq 1 - 2\varepsilon$, and hence $\|(\mathbb{1} - P)|\psi\rangle\|^2 = \langle \psi | (\mathbb{1} - P) | \psi \rangle \leq 1 - (\langle \psi | F | \psi \rangle - \nu) \leq 1 - (1 - 2\varepsilon) + \nu = 2\varepsilon + \nu$. Hence, we get

$$\begin{aligned} \operatorname{Re}\{\langle \psi' | (U - W) | \psi \rangle\} &= \operatorname{Re}\{\langle \psi' | (U - W)P | \psi \rangle\} + \operatorname{Re}\{\langle \psi' | (U - W)(\mathbb{1} - P) | \psi \rangle\} \\ &\geq \operatorname{Re}\{\langle \psi' | (U - W)P | \psi \rangle\} - 2\sqrt{2\varepsilon + \nu} \end{aligned} \quad (\text{C.7})$$

since by Cauchy-Schwarz $|\langle \psi' | (U - W)(\mathbb{1} - P) | \psi \rangle| \leq \|(U - W)^\dagger |\psi'\rangle\| \|(\mathbb{1} - P)|\psi\rangle\|$, where $\|(U - W)^\dagger |\psi'\rangle\| \leq 2$. In order to continue, let $|\chi\rangle = W^\dagger |\psi'\rangle - |\psi\rangle$, and calculate $\langle \chi | \chi \rangle = \langle \psi' | WW^\dagger | \psi' \rangle + \langle \psi | \psi \rangle - 2\operatorname{Re}\{\langle \psi' | W | \psi \rangle\} \leq 2 - 2(1 - \varepsilon) = 2\varepsilon$, and hence we deduce that $\| |\chi\rangle \| = \|W^\dagger |\psi'\rangle - |\psi\rangle\| \leq \sqrt{2\varepsilon}$. Then, with $\langle \psi' | W = \langle \psi | + \langle \chi |$ we have

$$\begin{aligned} \operatorname{Re}\{\langle \psi' | (U - W)P | \psi \rangle\} &= \operatorname{Re}\{\langle \psi' | W(F^{-1/2} - \mathbb{1})P | \psi \rangle\} \\ &= \operatorname{Re}\{\langle \chi | (F^{-1/2} - \mathbb{1})P | \psi \rangle\} + \operatorname{Re}\{\langle \psi | (F^{-1/2} - \mathbb{1})P | \psi \rangle\} \\ &\geq \operatorname{Re}\{\langle \chi | (F^{-1/2} - \mathbb{1})P | \psi \rangle\} \end{aligned} \quad (\text{C.8})$$

since $\langle \psi | (F^{-1/2} - \mathbb{1})P | \psi \rangle = \langle \psi | P(F^{-1/2} - \mathbb{1})P | \psi \rangle \geq 0$ as P commutes with $F^{-1/2}$ and since $F = W^\dagger W \leq \mathbb{1}$ implies that $F^{-1/2} \geq \mathbb{1}$. To bound the remaining term we first write $|\langle \chi | (F^{-1/2} - \mathbb{1})P | \psi \rangle| \leq \| |\chi\rangle \| \| (F^{-1/2} - \mathbb{1})P | \psi \rangle \| \leq \sqrt{2\varepsilon/\nu}$; the last inequality follows since P projects onto the eigenspaces of F with eigenvalues larger than or equal to ν , thus $F^{-1/2}P \leq \nu^{-1/2}P$ and hence $\| (F^{-1/2} - \mathbb{1})P | \psi \rangle \| \leq \nu^{-1/2} - 1 \leq \nu^{-1/2}$. Hence, we have

$$(\text{C.8}) \geq -\sqrt{\frac{2\varepsilon}{\nu}} \quad (\text{C.9})$$

Following the inequalities from (C.6), invoking (C.7) and with the above, we finally obtain

$$\operatorname{Re}\{\langle \psi' | U | \psi \rangle\} \geq 1 - \varepsilon - 2\sqrt{2\varepsilon + \nu} - \sqrt{\frac{2\varepsilon}{\nu}}. \quad (\text{C.10})$$

Choosing $\nu = 2\varepsilon^{1/2}$, we obtain, using $\varepsilon \leq \sqrt{\varepsilon}$,

$$1 - \operatorname{Re}\{\langle \psi' | U | \psi \rangle\} \leq \varepsilon + 2\sqrt{2\varepsilon + 2\varepsilon^{1/2}} + \sqrt{\frac{2\varepsilon}{2\varepsilon^{1/2}}} \leq (1 + 4 + 1)\varepsilon^{1/4} = 6\varepsilon^{1/4} \quad (\text{C.11})$$

as claimed. \blacksquare

D Robust counterexample against extensions of Construction #1

In this appendix we show that the counterexample of Section 5.2 is robust to small errors on the process. The process is $\mathcal{E}_{X \rightarrow X'}(\cdot) = \operatorname{tr}[\cdot] |+\rangle\langle +|$, where $|+\rangle = [|0\rangle + |1\rangle]/\sqrt{2}$ with $|0\rangle, |1\rangle$ energy eigenstates of respective energies $E_0 = 0, E_1 > 0$; we write $H_X = \sum_{j=0,1} E_j |j\rangle\langle j|$ and $\Gamma_X = e^{-\beta H_X}$.

The initial state on X and a reference system $R_X \simeq X$ is the maximally entangled state $|\sigma\rangle_{XR_X} = [|00\rangle + |11\rangle]/\sqrt{2} = |\Phi\rangle_{X:R_X}/\sqrt{2}$.

We seek a map $\mathcal{T}_{X \rightarrow X'}$ such that

$$P(\mathcal{T}_{X \rightarrow X'}(\sigma_{XR_X}), \mathcal{E}_{X \rightarrow X'}(\sigma_{XR_X})) \leq \varepsilon \quad \text{and} \quad \mathcal{T}_{X \rightarrow X'}(\Gamma_X) \leq \alpha \Gamma_{X'}, \quad (\text{D.1})$$

for a α that is independent of E_0, E_1 . Here we have $X \simeq X'$ and $\Gamma_X = \Gamma_{X'}$.

Let $\rho_{X'R_X} = \mathcal{E}_{X \rightarrow X'}(\sigma_{XR_X})$. From (D.1) we find $\frac{1}{2} \|\mathcal{T}_{X \rightarrow X'}(\sigma_{XR_X}) - \rho_{X'R_X}\|_1 \leq \varepsilon$, which in turn implies that $(1/4) \|\mathcal{T}_{X \rightarrow X'}(\Phi_{X:R_X}) - |+\rangle\langle+|_{X'} \otimes \mathbb{1}_{R_X}\|_1 \leq \varepsilon$, and hence that $\mathcal{T}_{X \rightarrow X'}(\cdot) = \text{tr}[\cdot] |+\rangle\langle+|_{X'} + \Delta(\cdot)$ for some Hermiticity preserving map $\Delta(\cdot)$ satisfying $\frac{1}{2} \|\Delta(\Phi_{X:R_X})\|_1 \leq 2\varepsilon$.

Let $\Delta_{\pm} \geq 0$ be the positive and negative parts of $\Delta(\Gamma) = \Delta_+ - \Delta_-$, noting that $\text{tr}(\Delta_-) \leq \text{tr}(\Delta_-) + \text{tr}(\Delta_+) = \|\Delta(\Gamma)\|_1 = \|\text{tr}_{R_X}(\Gamma_{R_X}^{1/2} \Delta(\Phi_{X:R_X}) \Gamma_{R_X}^{1/2})\|_1$, defining Γ_{R_X} as the transpose of Γ_X onto the system R_X , and continuing the computation we obtain $\text{tr}(\Delta_-) \leq \|\Gamma_{R_X}^{1/2} \Delta(\Phi_{X:R_X}) \Gamma_{R_X}^{1/2}\|_1 \leq \|\Gamma_{R_X}\|_{\infty} \|\Delta(\Phi_{X:R_X})\|_1 \leq 4\varepsilon$, using the fact that $\|\Gamma_X\|_{\infty} = \max_j \{e^{-\beta E_j}\} = 1$.

The condition $\mathcal{T}_{X \rightarrow X'}(\Gamma) \leq \alpha \Gamma$ implies that $\alpha \Gamma \geq \text{tr}[\Gamma] |+\rangle\langle+| + \Delta(\Gamma) \geq |+\rangle\langle+| - \Delta_-$. Hence, we have that $\alpha^{-1} |+\rangle\langle+| \leq \Gamma + \Delta_-/\alpha$. Hence, for any $0 < \eta \leq 1$ to be fixed later, $\mu = \alpha^{-1}$ is feasible for the dual problem (2.7) defining the hypothesis testing entropy $D_H^{\eta}(|+\rangle\langle+| \|\Gamma)$, and $e^{-D_H^{\eta}(|+\rangle\langle+| \|\Gamma)} \geq \alpha^{-1} - \text{tr}[\Delta_-/\alpha]/\eta \geq \alpha^{-1}(1 - 4\varepsilon/\eta)$. Thus, we have $\ln(\alpha) \geq D_H^{\eta}(|+\rangle\langle+| \|\Gamma) + \ln(1 - 4\varepsilon/\eta)$. Choosing $\eta = 8\varepsilon$ yields $\ln(1 - 4\varepsilon/\eta) = -\ln(2)$.

On the other hand, by definition we have $e^{-D_H^{\eta}(|+\rangle\langle+| \|\Gamma)} \leq \text{tr}[Q\Gamma]/\eta$ for any $0 \leq Q \leq \mathbb{1}$ satisfying $\text{tr}[Q|+\rangle\langle+|] \geq \eta$; with $Q = 2\eta|1\rangle\langle 1|$ we obtain $e^{-D_H^{\eta}(|+\rangle\langle+| \|\Gamma)} \leq 2e^{-\beta E_1}$ and thus $D_H^{\eta}(|+\rangle\langle+| \|\Gamma) \geq \beta E_1 - \ln(2)$.

Then, $\ln(\alpha) \geq -\ln(2) + \beta E_1 - \ln(2) = -2\ln(2) + \beta E_1$. Now let α be the optimal candidate in the coherent relative entropy $\hat{D}_{X \rightarrow X'}^{\varepsilon}(\rho_{X'R_X} \|\Gamma, \Gamma) = -\ln(\alpha)$. We finally see that the transformation $\mathbb{1}/2 \rightarrow |+\rangle$ may require arbitrarily much energy if $E_1 \rightarrow \infty$, even for a small $\varepsilon > 0$, since

$$\text{energy cost} = -\beta^{-1} \hat{D}_{X \rightarrow X'}^{\varepsilon}(\rho_{X'R_X} \|\Gamma, \Gamma) = \beta^{-1} \ln(\alpha) \geq E_1 - 2\beta^{-1} \ln(2). \quad (\text{D.2})$$

E Universal conditional typical projector for trivial Hamiltonians

In the case of trivial Hamiltonians, Definition 6.1 can be simplified. We call the corresponding object a universal conditional typical projector

Definition E.1. Consider two systems with Hilbert spaces $\mathcal{H}_A, \mathcal{H}_B$ and let $s \in \mathbb{R}$. We define a *universal conditional typical projector* $P_{A^n B^n}^{s, \delta}$ with parameter $\delta > 0$ as a projector acting on $(\mathcal{H}_A \otimes \mathcal{H}_B)^{\otimes n}$ such that:

- (i) There exists $\eta > 0$ independent of n such that for any quantum state ρ_{AB} with $H(A|B)_{\rho} \leq s$, we have

$$\text{tr}[P_{A^n B^n}^{s, \delta} \rho_{AB}^{\otimes n}] \geq 1 - \text{poly}(n) \exp(-n\eta); \quad (\text{E.1})$$

$$(ii) \quad \text{tr}_{A^n} [P_{A^n B^n}^{s, \delta}] \leq \text{poly}(n) e^{n(s+2\delta)} \mathbb{1}_{B^n}.$$

Observe that we choose to define the object in [Definition E.1](#) as a projector whereas we only require the object in [Definition 6.1](#) to be an operator of norm at most 1. The reason is that while we can prove that a projector satisfying the conditions of [Definition E.1](#) exists, we are currently not able to guarantee the existence of a projector satisfying the criteria of [Definition 6.1](#).

Proposition E.1. *Consider two systems A, B and let $s \in \mathbb{R}$. For any $\delta > 0$ and $n \in \mathbb{N}$ there exists a universal conditional typical projector $P_{A^n B^n}^{s, \delta}$ that is permutation-invariant.*

The proof of [Proposition E.1](#) is developed in the rest of this appendix. To understand why the projector of [Definition E.1](#) is conditional—as well as for a simple illustration of its use—consider the smooth Rényi-zero conditional max-entropy, also known as the smooth alternative max-entropy [11]. It is defined for a bipartite state ρ_{AB} as

$$\hat{H}_{\max}^\varepsilon(A|B)_\rho = \min_{\hat{\rho} \approx_\varepsilon \rho} \ln \left\| \text{tr}_A [\Pi_{AB}^{\hat{\rho}_{AB}}] \right\|_\infty, \quad (\text{E.2})$$

where $\Pi_{AB}^{\hat{\rho}_{AB}}$ is the projector onto the support of $\hat{\rho}_{AB}$, and where the optimization ranges over sub-normalized states $\hat{\rho}_{AB}$ which are ε -close to ρ_{AB} in purified distance. We may understand the i.i.d. behaviour of this quantity as follows. For $\delta > 0$ and $n \in \mathbb{N}$ let $P_{A^n B^n}^{s, \delta}$ be a universal conditional typical projector with $s = H(A|B)_\rho$. We define $\hat{\rho}_{A^n B^n} = P_{A^n B^n}^{s, \delta} \rho_{AB}^{\otimes n} P_{A^n B^n}^{s, \delta}$. Then, we have $\hat{\rho}_{A^n B^n} \approx_\varepsilon \rho_{AB}^{\otimes n}$ for $n \in \mathbb{N}$ large enough, thanks to Property (i) and the gentle measurement lemma ([Lemma B.2](#)). On the other hand, using Property (ii) we have

$$\frac{1}{n} \hat{H}_{\max}^\varepsilon(A^n | B^n)_{\rho^{\otimes n}} \leq \frac{1}{n} \ln \left\| \text{tr}_{A^n} [P_{A^n B^n}^{s, \delta}] \right\|_\infty \leq H(A|B)_\rho + 2\delta + \frac{1}{n} \ln(\text{poly}(n)) \quad (\text{E.3})$$

such that taking the limits $n \rightarrow \infty$ and $\delta \rightarrow 0$, we get that the smooth Rényi-zero conditional entropy is asymptotically upper bounded by the von Neumann conditional entropy in the i.i.d. regime.

We proceed to construct a universal conditional typical projector based on ideas from Schur-Weyl duality. The construction presented here is similar to, and inspired by, techniques put forward in earlier work [22, 24–26, 54, 55].

Proof of Proposition E.1. Let

$$P_{A^n B^n}^{s, \delta} = \sum_{\substack{\lambda, \lambda' : \\ \bar{H}(\lambda) - \bar{H}(\lambda') \leq s + 2\delta}} (\mathbb{1}_{A^n} \otimes \Pi_{B^n}^{\lambda'}) \Pi_{A^n B^n}^\lambda, \quad (\text{E.4})$$

where the respective projectors $\Pi_{B^n}^{\lambda'}$, $\Pi_{A^n B^n}^\lambda$ refer to Schur-Weyl decompositions of $\mathcal{H}_B^{\otimes n}$ and of $(\mathcal{H}_A \otimes \mathcal{H}_B)^{\otimes n}$, respectively, $\lambda \in \text{Young}(d_A d_B, n)$ and $\lambda' \in \text{Young}(d_B, n)$. Observe that $P_{A^n B^n}^{s, \delta}$ is a projector: Each term in the sum is a projector as a product of two commuting projectors ([Lemma 2.1](#)), and each term of the sum acts on a different subspace of $(\mathcal{H}_A \otimes \mathcal{H}_B)^{\otimes n}$. The projector $P_{A^n B^n}^{s, \delta}$ corresponds to the measurement of the two commuting POVMs $\{\Pi_{A^n B^n}^\lambda\}$ and $\{\Pi_{B^n}^{\lambda'}\}$, and testing whether or not the event $\bar{H}(\lambda) - \bar{H}(\lambda') \leq s + 2\delta$ is satisfied. Also by construction $P_{A^n B^n}^{s, \delta}$ is permutation-invariant.

For any ρ_{AB} with $H(A|B)_\rho \leq s$, the probability that the measurement of $P_{A^n B^n}^{s, \delta}$ fails on $\rho_{AB}^{\otimes n}$ can be upper bounded as follows. The passing event $\bar{H}(\lambda) - \bar{H}(\lambda') \leq s + 2\delta$ is implied in particular by the two events (a) $\bar{H}(\lambda) \leq H(AB)_\rho + \delta$ and (b) $\bar{H}(\lambda') \geq H(B)_\rho - \delta$ happening simultaneously, recalling that $H(AB)_\rho - H(B)_\rho = H(A|B)_\rho \leq s$. The probability of event (a) failing is

$$\Pr[\bar{H}(\lambda) > H(AB)_\rho + \delta] \leq \text{poly}(n) \exp(-n\eta) \quad (\text{E.5})$$

as given by [Proposition 2.1](#), and similarly for event (b)

$$\Pr[\bar{H}(\lambda') < H(B)_\rho - \delta] \leq \text{poly}(n) \exp(-n\eta). \quad (\text{E.6})$$

We can use the same η in both cases by picking the lesser of the two values given by [Proposition 2.1](#), if necessary. Note furthermore that $\eta > 0$ does not depend on ρ . Hence with this η , for any ρ_{AB} we have

$$\text{tr}[P_{A^n B^n}^{s, \delta} \rho_{AB}^{\otimes n}] \geq 1 - \text{poly}(n) \exp(-n\eta) \quad (\text{E.7})$$

as required.

For the second property, we use [Lemma 2.2](#) to write

$$\begin{aligned} \text{tr}_{A^n} [P_{A^n B^n}^{s, \delta}] &= \sum_{\lambda, \lambda' : \bar{H}(\lambda) - \bar{H}(\lambda') \leq s + 2\delta} \Pi_{B^n}^{\lambda'} \text{tr}_{A^n} [\Pi_{A^n B^n}^\lambda] \Pi_{B^n}^{\lambda'} \\ &\leq \sum_{\lambda, \lambda' : \bar{H}(\lambda) - \bar{H}(\lambda') \leq s + 2\delta} \text{poly}(n) e^{n(\bar{H}(\lambda) - \bar{H}(\lambda'))} \mathbb{1}_{B^n} \\ &\leq \text{poly}(n) e^{n(s + 2\delta)} \mathbb{1}_{B^n} \end{aligned} \quad (\text{E.8})$$

recalling that there are only $\text{poly}(n)$ many possible Young diagrams and hence at most so many terms in the sum. \blacksquare

F Universal conditional erasure for n copies and trivial Hamiltonians

Corollary F.1 (Thermodynamic protocol for universal conditional erasure for n copies). *Let S, M be systems, let σ_S be the maximally mixed state on S . Let $s < \ln(d_S)$, where d_S is the dimension of S , and let $\delta > 0$ small enough. Let $n \in \mathbb{N}$ be large enough. Let J be a large enough information battery and let any $m \leq n(\ln(d_S) - s - 3\delta)$ such that e^m is integer.*

Then, there exists $\eta' > 0$ and a thermal operation $\mathcal{R}_{S^n M^n J \rightarrow S^n M^n J}$ acting on the systems $S^n M^n J$, such that the effective work process $\mathcal{T}_{S^n M^n \rightarrow S^n M^n}$ of $\mathcal{R}_{S^n M^n J \rightarrow S^n M^n J}$ with respect to the battery states $(\tau_J^m, |0\rangle_J)$ is a universal conditional $(\text{poly}(n) e^{-n\eta'})$ -erasure process resetting S^n to the state $\sigma_S^{\otimes n}$ with respect to the set of states $\mathcal{S}'_{S^n M^n}$, where $\mathcal{S}'_{S^n M^n}$ is the convex hull of $\mathcal{S}_{S^n M^n} = \{\rho_{SM}^{\otimes n} : H(S|M)_\rho \leq s\}$.

The case where $s = \ln(d_S)$ is uninteresting as we cannot hope to extract any work. In such cases one can simply set $m = 0$ and take $\mathcal{R}_{S^n M^n J}$ to be the thermal operation that completely thermalizes S^n .

Proof. This is in fact a relatively straightforward application of [Proposition 7.1](#) over n copies of SM . Let $P_{S^n M^n}^{s, \delta}$ be given by [Proposition E.1](#). We seek κ, κ' that satisfy (7.5). We can choose $\kappa = \text{poly}(n) \exp\{-n\eta(\delta)\}$ thanks to [Definition E.1](#). Furthermore for any $\rho_{SM}^{\otimes n} \in \mathcal{S}_{S^n M^n}$ we have

$$\begin{aligned} \text{tr} \left[P_{S^n M^n}^{s, \delta} \left(\frac{\mathbb{1}_S}{d_S} \otimes \rho_M \right)^{\otimes n} \right] &\leq \text{poly}(n) e^{n(s+2\delta)} d_S^{-n} \text{tr}[\rho_M^{\otimes n}] = \text{poly}(n) e^{-n(\ln(d_S) - s - 2\delta)} \\ &\leq \frac{\text{poly}(n) e^{-n\delta}}{e^m} \end{aligned} \quad (\text{F.1})$$

and thus we may take $\kappa' = \text{poly}(n) e^{-n\delta}$. Finally, η' is given as $\eta' = \min\{\delta, \eta(\delta)\}$. \blacksquare

References

- [1] J. Goold, M. Huber, A. Riera, L. d. Rio, and P. Skrzypczyk, The role of quantum information in thermodynamics—a topical review, [Journal of Physics A: Mathematical and Theoretical](#) **49**, 143001 (2016)
- [2] F. G. S. L. Brandão, M. Horodecki, J. Oppenheim, J. M. Renes, and R. W. Spekkens, Resource theory of quantum states out of thermal equilibrium, [Physical Review Letters](#) **111**, 250404 (2013)
- [3] F. Brandão, M. Horodecki, N. Ng, J. Oppenheim, and S. Wehner, The second laws of quantum thermodynamics, [Proceedings of the National Academy of Sciences](#) **112**, 3275–3279 (2015)
- [4] E. Chitambar and G. Gour, Quantum resource theories, [Reviews of Modern Physics](#) **91**, 025001 (2019)
- [5] D. Janzing, P. Wocjan, R. Zeier, R. Geiss, and T. Beth, Thermodynamic cost of reliability and low temperatures: Tightening Landauer’s principle and the second law, [International Journal of Theoretical Physics](#) **39**, 2717–2753 (2000)
- [6] P. Faist, J. Oppenheim, and R. Renner, Gibbs-preserving maps outperform thermal operations in the quantum regime, [New Journal of Physics](#) **17**, 043003 (2015)
- [7] J. Åberg, Truly work-like work extraction via a single-shot analysis, [Nature Communications](#) **4**, 1925 (2013)
- [8] M. Horodecki and J. Oppenheim, Fundamental limitations for quantum and nanoscale thermodynamics, [Nature Communications](#) **4**, 2059 (2013)
- [9] R. Renner, *Security of Quantum Key Distribution*, Ph.D. thesis, ETH Zürich (2005), [arXiv:quant-ph/0512258](#)
- [10] M. Tomamichel, *A Framework for Non-Asymptotic Quantum Information Theory*, [Quantum physics; mathematical physics; mathematical physics](#), ETH Zurich (2012), [arXiv:1203.2142](#)

- [11] M. Tomamichel, *Quantum Information Processing with Finite Resources*, Springer-Briefs in Mathematical Physics, Vol. 5 (Springer International Publishing, Cham, 2016) [arXiv:1504.00233](#)
- [12] C. T. Chubb, M. Tomamichel, and K. Korzekwa, Beyond the thermodynamic limit: finite-size corrections to state interconversion rates, *Quantum* **2**, 108 (2018)
- [13] P. Faist, F. Dupuis, J. Oppenheim, and R. Renner, The minimal work cost of information processing, *Nature Communications* **6**, 7669 (2015)
- [14] C. Cîrstoiu and D. Jennings, Global and local gauge symmetries beyond lagrangian formulations (2017), [arXiv:1707.09826](#)
- [15] K. Ben Dana, M. García Díaz, M. Mejatty, and A. Winter, Resource theory of coherence: Beyond states, *Physical Review A* **95**, 062327 (2017)
- [16] P. Faist and R. Renner, Fundamental work cost of quantum processes, *Physical Review X* **8**, 021011 (2018)
- [17] G. Smith, Quantum channel capacities, in *IEEE Information Theory Workshop* (2010) pp. 1–5
- [18] M. Christandl, R. König, and R. Renner, Postselection technique for quantum channels with applications to quantum cryptography, *Physical Review Letters* **102**, 20504 (2009)
- [19] A. Anshu, R. Jain, and N. A. Warsi, Building blocks for communication over noisy quantum networks, *IEEE Transactions on Information Theory* **65**, 1287–1306 (2019)
- [20] P. Faist, M. Berta, and F. Brandão, Thermodynamic capacity of quantum processes, *Physical Review Letters* **122**, 200601 (2019)
- [21] M. Navascués and L. P. García-Pintos, Nonthermal quantum channels as a thermodynamical resource, *Physical Review Letters* **115**, 010405 (2015)
- [22] C. H. Bennett, I. Devetak, A. W. Harrow, P. W. Shor, and A. Winter, The quantum reverse Shannon theorem and resource tradeoffs for simulating quantum channels, *IEEE Transactions on Information Theory* **60**, 2926–2959 (2014)
- [23] M. Berta, M. Christandl, and R. Renner, The quantum reverse Shannon theorem based on one-shot information theory, *Communications in Mathematical Physics* **306**, 579–615 (2011)
- [24] A. W. Harrow, *Applications of coherent classical communication and the Schur transform to quantum information theory*, Ph.D. thesis, Massachusetts Institute of Technology (2005), [arXiv:quant-ph/0512255](#)
- [25] J. Haah, A. W. Harrow, Z. Ji, X. Wu, and N. Yu, Sample-optimal tomography of quantum states, *IEEE Transactions on Information Theory* **63**, 5628–5641 (2017)
- [26] J. Nötzl, A solution to two party typicality using representation theory of the symmetric group (2012), [arXiv:1209.5094](#)
- [27] M. Tomamichel, R. Colbeck, and R. Renner, Duality between smooth min- and max-entropies, *IEEE Transactions on Information Theory* **56**, 4674–4681 (2010)
- [28] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000)
- [29] L. Wang and R. Renner, One-shot classical-quantum capacity and hypothesis testing, *Physical Review Letters* **108**, 200501 (2012)

- [30] M. Tomamichel and M. Hayashi, A hierarchy of information quantities for finite block length analysis of quantum tasks, *IEEE Transactions on Information Theory* **59**, 7693–7710 (2013)
- [31] W. Matthews and S. Wehner, Finite blocklength converse bounds for quantum channels, *IEEE Transactions on Information Theory* **60**, 7317–7329 (2014)
- [32] F. Buscemi and N. Datta, The quantum capacity of channels with arbitrarily correlated noise, *IEEE Transactions on Information Theory* **56**, 1447–1460 (2010)
- [33] F. G. S. L. Brandão and N. Datta, One-shot rates for entanglement manipulation under non-entangling maps, *IEEE Transactions on Information Theory* **57**, 1754–1760 (2011)
- [34] F. Dupuis, L. Kraemer, P. Faist, J. M. Renes, and R. Renner, Generalized entropies, in *XVIIIth International Congress on Mathematical Physics* (World Scientific, Singapore, 2013) pp. 134–153, [arXiv:1211.3141](#)
- [35] J. Watrous, Semidefinite programs for completely bounded norms, *Theory of Computing* **5**, 217–238 (2009)
- [36] L. Szilard, Über die entropieverminderung in einem thermodynamischen system bei eingriffen intelligenter wesen, *Zeitschrift für Physik* **53**, 840–856 (1929)
- [37] S. P. Boyd and L. Vandenberghe, *Convex Optimization* (Cambridge University Press, Cambridge, UK, 2004)
- [38] A. Pitchford, C. Granade, P. D. Nation, and R. J. Johansson, *QuTip 4.1.0* (2016)
- [39] J. Johansson, P. Nation, and F. Nori, QuTip 2: A Python framework for the dynamics of open quantum systems, *Computer Physics Communications* **184**, 1234–1240 (2013)
- [40] M. S. Andersen, J. Dahl, and L. Vandenberghe, *CVXOPT 1.1.9* (2016)
- [41] N. Ramakrishnan, R. Iten, V. B. Scholz, and M. Berta, Computing quantum channel capacities, *IEEE Transactions on Information Theory* **67**, 946–960 (2021)
- [42] R. Alicki, Isotropic quantum spin channels and additivity questions (2004), [arXiv:quant-ph/0402080](#)
- [43] I. Devetak, M. Junge, C. King, and M. B. Ruskai, Multiplicativity of completely bounded p-norms implies a new additivity result, *Communications in Mathematical Physics* **266**, 37–63 (2006)
- [44] A. S. Holevo, The entropy gain of quantum channels, in *Proceedings of the 2011 IEEE International Symposium on Information Theory*, Vol. 82 (IEEE, 2011) pp. 289–292
- [45] A. S. Holevo, The entropy gain of infinite-dimensional quantum evolutions, *Doklady Mathematics* **82**, 730–731 (2010)
- [46] A. S. Holevo, Entropy gain and the Choi-Jamiolkowski correspondence for infinite-dimensional quantum evolutions, *Theoretical and Mathematical Physics* **166**, 123–138 (2011)
- [47] A. S. Holevo, *Quantum Systems, Channels, Information* (De Gruyter, Berlin, Boston, 2012)
- [48] F. Buscemi, S. Das, and M. M. Wilde, Approximate reversibility in the context of entropy gain, information gain, and complete positivity, *Physical Review A* **93**, 062314 (2016)
- [49] G. Gour and M. M. Wilde, Entropy of a quantum channel, *Physical Review Research* **3**, 023096 (2021)
- [50] M. Berta, J. M. Renes, and M. M. Wilde, Identifying the information gain of a quantum

- measurement, *IEEE Transactions on Information Theory* **60**, 7987–8006 (2014)
- [51] P. Faist, *Quantum Coarse-Graining: An Information-Theoretic Approach to Thermodynamics*, Ph.D. thesis, ETH Zürich (2016), [arXiv:1607.03104](#)
 - [52] C. Morgan and A. Winter, “Pretty strong” converse for the quantum capacity of degradable channels, *IEEE Transactions on Information Theory* **60**, 317–333 (2014)
 - [53] M. Tomamichel, R. Colbeck, and R. Renner, A fully quantum asymptotic equipartition property, *IEEE Transactions on Information Theory* **55**, 5840–5847 (2009)
 - [54] I. Bjelakovic and R. Siegmund-Schultze, Quantum Stein’s lemma revisited, inequalities for quantum entropies, and a concavity theorem of Lieb (2003), [arXiv:quant-ph/0307170](#)
 - [55] M. Berta, M. Lemm, and M. M. Wilde, Monotonicity of quantum relative entropy and recoverability, *Quantum Information and Computation* **15**, 1333–1354 (2015)
 - [56] A. Anshu, V. K. Devabathini, and R. Jain, Quantum communication using coherent rejection sampling, *Physical Review Letters* **119**, 120506 (2017)
 - [57] A. Anshu, R. Jain, and N. A. Warsi, A one-shot achievability result for quantum state redistribution, *IEEE Transactions on Information Theory* **64**, 1425–1435 (2018)
 - [58] A. Anshu, R. Jain, and N. A. Warsi, A generalized quantum Slepian–Wolf, *IEEE Transactions on Information Theory* **64**, 1436–1453 (2018)
 - [59] A. Anshu, R. Jain, and N. A. Warsi, Convex-split and hypothesis testing approach to one-shot quantum measurement compression and randomness extraction, *IEEE Transactions on Information Theory* **65**, 5905–5924 (2019)
 - [60] C. Majenz, M. Berta, F. Dupuis, R. Renner, and M. Christandl, Catalytic decoupling of quantum information, *Physical Review Letters* **118**, 080503 (2017)
 - [61] A. Anshu, M. Berta, R. Jain, and M. Tomamichel, Partially smoothed information measures, *IEEE Transactions on Information Theory* **66**, 5022–5036 (2020)
 - [62] M. Berta and C. Majenz, Disentanglement cost of quantum states, *Physical Review Letters* **121**, 190503 (2018)
 - [63] L. del Rio, J. Åberg, R. Renner, O. Dahlsten, and V. Vedral, The thermodynamic meaning of negative entropy, *Nature* **474**, 61–63 (2011)
 - [64] M. Hayashi and H. Nagaoka, General formulas for capacity of classical-quantum channels, *IEEE Transactions on Information Theory* **49**, 1753–1768 (2003)
 - [65] H. Scutaru, Some remarks on covariant completely positive linear maps on C^* -algebras, *Reports on Mathematical Physics* **16**, 79–87 (1979)
 - [66] M. Keyl and R. F. Werner, Optimal cloning of pure states, testing single clones, *Journal of Mathematical Physics* **40**, 3283–3299 (1999)
 - [67] I. Marvian Mashhad, *Symmetry, Asymmetry and Quantum Information*, Ph.D. thesis, University of Waterloo (2012)
 - [68] J. Åberg, Catalytic coherence, *Physical Review Letters* **113**, 150402 (2014)
 - [69] K. Fang, X. Wang, M. Tomamichel, and M. Berta, Quantum channel simulation and the channel’s smooth max-information, *IEEE Transactions on Information Theory* **66**, 2129 – 2140 (2020)

- [70] G. Gour and A. Winter, How to quantify a dynamical resource?, *Physical Review Letters* **123**, 150401 (2019)
- [71] N. Dutil, *Multiparty quantum protocols for assisted entanglement distillation*, Ph.D. thesis, McGill University, Montréal (2011), [arXiv:1105.4657](#)
- [72] L. Drescher and O. Fawzi, On simultaneous min-entropy smoothing, in *2013 IEEE International Symposium on Information Theory* (IEEE, 2013) pp. 161–165, [arXiv:1312.7642](#)
- [73] P. Sen, A one-shot quantum joint typicality lemma (2018), [arXiv:1806.07278](#)
- [74] A. Anshu, M. Berta, R. Jain, and M. Tomamichel, A minimax approach to one-shot entropy inequalities, *Journal of Mathematical Physics* **60**, [10.1063/1.5126723](#) (2019)
- [75] M. Fannes, A continuity property of the entropy density for spin lattice systems, *Communications in Mathematical Physics* **31**, 291–294 (1973)
- [76] K. M. R. Audenaert, A sharp continuity estimate for the von Neumann entropy, *Journal of Physics A: Mathematical and Theoretical* **40**, 8127–8136 (2007)
- [77] M. Berta, M. Christandl, R. Colbeck, J. M. Renes, and R. Renner, The uncertainty principle in the presence of quantum memory, *Nature Physics* **6**, 659–662 (2010)