

Approximation by Exponential Type Neural Network Operators

Bajpeyi Shivam * · A. Sathish Kumar

Received: date / Accepted: date

Abstract In the present article, we introduce and study the behaviour of the new family of exponential type neural network operators activated by the sigmoidal functions. We establish the point-wise and uniform approximation theorems for these NN (Neural Network) operators in $\mathcal{C}[a, b]$. Further, the quantitative estimates of order of approximation for the proposed NN operators in $C^{(N)}[a, b]$ are established in terms of the modulus of continuity. We also analyze the behaviour of the family of exponential type quasi-interpolation operators in $\mathcal{C}(\mathbb{R}^+)$. Finally, we discuss the multivariate extension of these NN operators and some examples of the sigmoidal functions.

Keywords Exponential sampling type operators · Neural Network operators · Order of convergence · Sigmoidal functions · Mellin transform

Mathematics Subject Classification (2010) 41A35 · 92B20 · 94A20 · 41A25

1 Introduction

The exponential sampling methods are powerful tool to solve problems arising in the areas of optical physics and engineering, precisely in the phenomena like Fraunhofer diffraction, light scattering etc. [21, 23, 38, 45]. It all started when a group of optical physicists and engineers Bartero, Pike [21] and Gori [38] proposed a representation formula known as *exponential sampling formula* ([13]), for the class of Mellin band-limited function having exponentially spaced

S. Bajpeyi *(corresponding author)
Department of Mathematics, Visvesvaraya National Institute of Technology, Nagpur-440010
E-mail: shivambajpai1010@gmail.com

A. Sathish Kumar
Department of Mathematics, Visvesvaraya National Institute of Technology, Nagpur-440010
E-mail: mathsathish9@gmail.com

sample points. This exponential sampling formula is also considered as the Mellin-version of the well known *Shannon sampling theorem* ([16]). Butzer and Jansche ([19]) pioneered the mathematical study of the exponential sampling formula. They established a rigorous proof of this formula mathematically using the theory of Mellin transform and Mellin approximation, which was first studied separately by Mamedov [43], and then developed by Butzer and Jansche [17,18,19,20]. We mention some of the work related to the theory of Mellin transform and approximation in [9,11,12].

Bardaro et al.[13] introduced the generalized version of the exponential sampling series, acting on functions which are not necessarily Mellin band-limited, by replacing the lin_c function in the exponential sampling formula, by more general kernel function. The generalized exponential sampling series is defined by

$$(S_w^\chi f)(x) = \sum_{k=-\infty}^{+\infty} \chi(e^{-k} x^w) f(e^{\frac{k}{w}}), \quad \forall x \in \mathbb{R}^+, w > 0 \quad (1)$$

where $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be any function for which the series is absolutely convergent. Subsequently, Bardaro et.al. extended the study of convergence of the above series in Mellin-Lebesgue spaces in [14]. Recently, Balsamo and Mantellini [8] considered the linear combination of these operators and analyzed its approximation results.

Let $C(\mathbb{R}^+)$ denotes the space of all uniformly continuous functions defined on \mathbb{R}^+ , where \mathbb{R}^+ denotes the set of all positive real numbers. We call a function $f \in C(\mathbb{R}^+)$ log-uniformly continuous on \mathbb{R}^+ , if for any given $\epsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(y)| < \epsilon$ whenever $|\log x - \log y| \leq \delta$, for any $x, y \in \mathbb{R}^+$. We denote the space of all log-uniformly continuous functions defined on \mathbb{R}^+ by $\mathcal{C}(\mathbb{R}^+)$. It is to mention that a log-uniformly continuous function need not uniformly continuous. We consider $M(\mathbb{R}^+)$ as the class of all Lebesgue measurable functions on \mathbb{R}^+ and $L^\infty(\mathbb{R}^+)$ as the space of all bounded functions on \mathbb{R}^+ .

Let $L^p(\mathbb{R}^+)$, $1 \leq p < +\infty$ denotes the space of all the Lebesgue measurable functions defined on \mathbb{R}^+ , equipped with the usual norm $\|f\|_p$. For $c \in \mathbb{R}$, we define the space

$$X_c = \{f : \mathbb{R}^+ \rightarrow \mathbb{C} : f(\cdot)(\cdot)^{c-1} \in L^1(\mathbb{R}^+)\}$$

equipped with the norm

$$\|f\|_{X_c} = \|f(\cdot)(\cdot)^{c-1}\|_1 = \int_0^{+\infty} |f(u)|u^{c-1} du.$$

The Mellin transform of a function $f \in X_c$ is defined by

$$\hat{M}[f](s) := \int_0^{+\infty} u^{s-1} f(u) du, \quad (s = c + it, t \in \mathbb{R}).$$

We call a function $f \in X_c \cap C(\mathbb{R}^+)$, $c \in \mathbb{R}$, Mellin band-limited in the interval $[-\eta, \eta]$, if $\hat{M}[f](c + iw) = 0$ for all $|w| > \eta$, $\eta \in \mathbb{R}^+$.

For a differentiable function $f : \mathbb{R}^+ \rightarrow \mathbb{C}$, the Mellin differential operator θ_c , is defined as

$$\theta_c f(x) := x f'(x) + c f(x), \quad x \in \mathbb{R}^+ \text{ and } c \in \mathbb{R}.$$

We set $\theta_0 f(x) := \theta f(x)$. The Mellin differential operator of order $r \in \mathbb{N}$ is defined by the following relation,

$$\theta_c^1 := \theta_c, \quad \theta_c^{(r)} = \theta_c(\theta_c^{(r-1)}).$$

2 Auxiliary results

In this section, we give some preliminary definitions and related results which will be helpful to study the approximation properties of the family of operators $(E_n^{\chi_\sigma} f)$. We begin with the definition of *sigmoidal function*. A function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is sigmoidal function if and only if $\lim_{x \rightarrow -\infty} \sigma(x) = 0$ and $\lim_{x \rightarrow \infty} \sigma(x) = 1$. In the present context, we assume $\sigma(x)$ to be non-decreasing function with $\sigma(2) > \sigma(0)$ satisfying the following conditions,

- (1) $\sigma \in C^{(2)}(\mathbb{R})$ and concave on \mathbb{R}^+ .
- (2) $\sigma(x) = \mathcal{O}(|x|^{-1-\nu})$, as $x \rightarrow -\infty$, for some $\nu > 0$.
- (3) The function $(\sigma(x) - \frac{1}{2})$ is an odd function.

Now, we define the density function activated by the sigmoidal function σ by the following linear combination,

$$\chi_\sigma(x) := \frac{1}{2} [\sigma(\log x + 1) - \sigma(\log x - 1)], \quad x \in \mathbb{R}^+.$$

The algebraic moment of order ν for the density function χ_σ is defined as,

$$m_\nu(\chi_\sigma, u) := \sum_{k=-\infty}^{+\infty} \chi_\sigma(e^{-k}u) (k - \log(u))^\nu, \quad \forall u \in \mathbb{R}^+.$$

Similarly, the absolute moment of order ν can be defined as,

$$M_\nu(\chi_\sigma, u) := \sum_{k=-\infty}^{+\infty} |\chi_\sigma(e^{-k}u)| |k - \log(u)|^\nu, \quad \forall u \in \mathbb{R}^+.$$

We define $M_\nu(\chi_\sigma) := \sup_{u \in \mathbb{R}^+} M_\nu(\chi_\sigma, u)$.

Now, we derive some basic results related to the density function χ_σ .

Lemma 1 *The condition $\sum_{k=-\infty}^{+\infty} \chi_\sigma(e^{-k}u) = 1$, holds for every $u \in \mathbb{R}^+$ and $k \in \mathbb{Z}$.*

Proof For any fixed $x \in \mathbb{R}^+$ and $n \in \mathbb{N}$, we have

$$\sum_{k=-n}^n (\sigma(\log x + 1 - k) - \sigma(\log x - k)) = \sigma(\log x + 1 + n) - \sigma(\log x - n).$$

Similarly,

$$\sum_{k=-n}^n (\sigma(\log x - k) - \sigma(\log x - 1 - k)) = \sigma(\log x + n) - \sigma(\log x - n - 1).$$

Using the definition of χ_σ , we obtain

$$\begin{aligned} \sum_{k=-\infty}^{+\infty} \chi_\sigma(e^{-k}u) &= \frac{1}{2} [\sigma(\log(ue^{-k}) + 1) - \sigma(\log(ue^{-k}) - 1)] \\ &= \frac{1}{2} [\sigma(\log u - k + 1) - \sigma(\log u - k - 1)] \\ &= \frac{1}{2} [(\sigma(\log u - k + 1) - \sigma(\log u - k)) + (\sigma(\log u - k) - \sigma(\log u - k - 1))] \\ &= \frac{1}{2} [\sigma(\log u + n + 1) - \sigma(\log u - n) + \sigma(\log u + n) - \sigma(\log u - n - 1)] \end{aligned}$$

As $n \rightarrow +\infty$, we obtain the desired result.

Lemma 2 Let $x \in [a, b] (\subset \mathbb{R}^+)$ and $n \in \mathbb{N}$, $k \in \mathbb{Z}$. Then, we have

$$\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k}x^n) \geq \chi_\sigma(e).$$

Proof Since $\sum_{k=-\infty}^{+\infty} \chi_\sigma(e^{-k}x^n) = 1$, we have $\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k}x^n) \leq 1$. Now, we have

$$\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k}x^n) = \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(|e^{-k}x^n|) \geq \chi_\sigma(|e^{-\hat{k}}x^n|), \quad \forall x \in \mathbb{R}^+.$$

Now by choosing $\hat{k} \in [\lceil na \rceil, \lfloor nb \rfloor] \cap \mathbb{Z}$ such that $2 < |e^{-\hat{k}}x^n| < 3$, we get

$$\chi_\sigma(|e^{-\hat{k}}x^n|) \geq \chi_\sigma(e) > 0, \quad \text{as } \sigma(2) > \sigma(0).$$

This establishes the result.

Remark 1: Using the condition (1), it is easy to see that $\chi_\sigma(x)$ is non-increasing for $x > 1$.

Lemma 3 For every $\eta > 0$ and $u \in \mathbb{R}^+$, the following condition holds

$$\lim_{n \rightarrow \infty} \sum_{|k - \log u| > n\eta} \chi_\sigma(e^{-k}u) = 0.$$

Proof Let $\{\log u\}$ denotes the fractional part of $\log u$ for every $u \in \mathbb{R}^+$, i.e., $\{\log u\} = \log u - [\log u]$, where, $[\cdot]$ denotes the *greatest integer function*. Define $\hat{k} := k - [\log u]$. This gives $|k - \log u| = |\hat{k} - \{\log u\}|$. Now, for any fixed $\eta > 0$, we have

$$\begin{aligned} \sum_{|k - \log u| > n\eta} \chi_\sigma(e^{-k}u) &= \sum_{|\hat{k} - \{\log u\}| > n\eta} \chi_\sigma(e^{-k}u) \\ &< \sum_{|\hat{k}| > n\eta+1} \chi_\sigma(e^{-k}u) \\ &= \left(\sum_{\hat{k} > n\eta+1} \chi_\sigma(e^{-k}u) + \sum_{\hat{k} < -(n\eta+1)} \chi_\sigma(e^{-k}u) \right) \\ &:= I_1 + I_2. \end{aligned}$$

Using the condition (2), we say that there exists $K > 0$ such that

$$\sigma(x) \leq K(|x|^{-1-\nu}), \quad \forall x < -L, \quad \text{for some } L \in \mathbb{R}^+ \text{ and } \nu > 0.$$

Since $\chi_\sigma(x) \leq \sigma(1 + \log x)$, we have

$$I_1 \leq \sum_{\hat{k} > n\eta+1} \sigma(1 + \log u - k) \leq \sum_{\hat{k} > n\eta+1} \sigma(1 + \{\log u\} - \hat{k}) < K \sum_{\hat{k} > n\eta+1} |2 - \hat{k}|^{-1-\nu}.$$

Moreover, from condition (3) we obtain

$$\chi_\sigma(x) \leq 1 - \sigma(\log x - 1) = \sigma(1 - \log x), \quad \text{which gives}$$

$$I_2 \leq \sum_{\hat{k} < -(n\eta+1)} \sigma(1 - \log u + k) \leq \sum_{\hat{k} < -(n\eta+1)} \sigma(1 - \{\log u\} + \hat{k}) \leq K \sum_{\hat{k} < -(n\eta+1)} |1 + \hat{k}|^{-1-\nu}.$$

As $\lim n \rightarrow +\infty$, we get the desired result.

3 Neural Network Operators

The feed-forward neural network operators (FNNs) with one hidden layer can be defined as

$$N_n(x) = \sum_{j=0}^n c_j \sigma(\langle a_j, x \rangle + b_j), \quad x \in \mathbb{R}^s, \quad s \in \mathbb{N},$$

where for $0 \leq j \leq n$, $b_j \in \mathbb{R}$ are thresholds, $a_j \in \mathbb{R}^s$ are the connection weights, $c_j \in \mathbb{R}$ are the coefficients, $\langle a_j, x \rangle$ is the inner product of a_j and x , and σ is the activation function of the network. The functions of one or several variables have been approximated by these NN (neural network) operators extensively from the last three decades. Cybenko [35] and Funahashi [37] have shown that any continuous function can be approximated on a compact set with uniform topology by the FNN-operators, using any continuous sigmoidal activation

function. Further, Hornik et al. in [40], proved that any measurable function can be approached with such a network. Cheney [32] and Lenze [42] used the idea of convolution kernel from the sigmoidal function to approximate the functions which is based on some results related to the theory of ridge functions.

The theory of NN operators has been introduced in order to study a constructive approximation process by neural networks. Anastassiou [1] pioneered the study of neural network approximation of continuous functions, where he extended the results proved by Cardalignet and Euvrard in [22]. Subsequently, Anastassiou analyzed the rate of approximation of the univariate and multivariate NN-operators activated by various sigmoidal functions in-terms of the modulus of smoothness (see[1,2,3]). Costarelli and Spigler extended the results proved by Anastassiou in ([24,26]). After this, many researchers analyzed the FNN-operators and established the various density results using different approaches, see ([28,29,34,36,39,41,44], etc.). Further, Costarelli and Spigler ([27]) studied the approximation of discontinuous functions by means of Kantorovich version of these NN-operators. Recently, the quantitative estimates of order of approximation was studied for kantorovich type NN operators in [30].

Let $f : [a, b](\subset \mathbb{R}^+) \rightarrow \mathbb{R}$ be any bounded function and $n \in \mathbb{N}$ such that $\lceil na \rceil \leq \lfloor nb \rfloor$. The exponential type neural network operators activated by the sigmoidal function σ , are defined as

$$E_n^{\chi_\sigma}(f, x) = \frac{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} f(e^{\frac{k}{n}}) \chi_\sigma(e^{-k} x^n)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n)}, \quad x \in \mathbb{R}^+.$$

Now, we derive the point-wise and uniform convergence results for the above family of operators ($E_n^{\chi_\sigma} f$).

3.1 Approximation Results

Theorem 1 *Let $f : [a, b](\subset \mathbb{R}^+) \rightarrow \mathbb{R}$ be any bounded function. Then, $(E_n^{\chi_\sigma}(f, x))$ converges to $f(x)$ at every point of continuity of f . Moreover, if $f \in \mathcal{C}[a, b]$, then*

$$\lim_{n \rightarrow \infty} \|E_n^{\chi_\sigma}(f, \cdot) - f\|_\infty = 0.$$

Proof We see that,

$$\begin{aligned}
|E_n^{\chi_\sigma}(f, x) - f(x)| &= \frac{\left| \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) (f(e^{\frac{k}{n}}) - f(x)) \right|}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |\chi_\sigma(e^{-k} x^n)|} \\
&\leq \frac{1}{|\chi_\sigma(e)|} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| \\
&\leq \frac{1}{|\chi_\sigma(e)|} \sum_{|\frac{k}{n} - \log(x)| < \delta} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| + \\
&\quad \frac{1}{|\chi_\sigma(e)|} \sum_{|\frac{k}{n} - \log(x)| \geq \delta} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| := I_1 + I_2.
\end{aligned}$$

First we evaluate I_1 . Since $f \in \mathcal{C}[a, b]$, we have $|I_1| < \frac{\epsilon}{|\chi_\sigma(e)|}$. Now for I_2 , using Lemma 3, we have

$$|I_2| \leq \frac{2\|f\|_\infty}{|\chi_\sigma(e)|} \epsilon.$$

On combining the estimates $I_1 - I_2$, we establish the desired result.

Now, we obtain the order of convergence of $(E_n^{\chi_\sigma} f)$ for the functions belonging to the class of *log-holderian functions* of order λ , for $0 < \lambda \leq 1$ (see [13]), which is defined as follows

$$L_\lambda(f) = \{f : \mathbb{R}^+ \rightarrow \mathbb{R} : |f(x) - f(y)| < H |\log x - \log y|^\lambda; x, y, H \in \mathbb{R}^+\}.$$

Theorem 2 *Let $f \in L_\lambda(f)$ and $M_\lambda(\chi_\sigma) < +\infty$, for some $\lambda > 0$. Then, we have*

$$\lim_{n \rightarrow \infty} \|E_n^{\chi_\sigma}(f, \cdot) - f\|_\infty = \mathcal{O}(n^{-\lambda}).$$

Proof Since $f \in L_\lambda(f)$, then $|f(e^{\frac{k}{n}}) - f(x)| \leq H |\frac{k}{n} - \log(x)|^\lambda$, whenever $|\frac{k}{n} - \log(x)| < \delta$, $\delta > 0$. Now, we have

$$\begin{aligned}
|E_n^{\chi_\sigma}(f, x) - f(x)| &\leq \frac{1}{|\chi_\sigma(e)|} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| \\
&\leq \frac{1}{|\chi_\sigma(e)|} \sum_{|\frac{k}{n} - \log(x)| < \delta} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| + \\
&\quad \frac{1}{|\chi_\sigma(e)|} \sum_{|\frac{k}{n} - \log(x)| \geq \delta} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| \\
&:= I_1 + I_2.
\end{aligned}$$

It easily follows that, $|I_1| < \frac{Hn^{-\lambda}}{|\chi_\sigma(1)|} M_\lambda(\chi_\sigma)$. Next, we estimate I_2 .

$$\begin{aligned} |I_2| &\leq \frac{2\|f\|_\infty}{|\chi_\sigma(e)|} \sum_{|\frac{k}{n} - \log(x)| \geq \delta} |\chi_\sigma(e^{-k}x^n)| \\ &\leq \frac{1}{|\chi_\sigma(e)|} \frac{2\|f\|_\infty}{n^\lambda \delta^\lambda} M_\lambda(\chi_\sigma). \end{aligned}$$

As $M_\lambda(\chi_\sigma) < +\infty$, it completes the proof.

3.2 Quantitative Estimates

In this section, we obtain a quantitative estimate of order of convergence for the family of operators $(E_n^{\chi_\sigma} f)$ for a particular sigmoidal function. There are several examples of the sigmoidal function σ satisfying all the assumptions of the presented theory (see [3, 4, 5, 6, 24, 25, 31, 33, 35]). We begin with an example of such sigmoidal function known as *hyperbolic tangent* sigmoidal function [4], is defined as

$$\sigma_h(x) := \frac{1}{2}(\tanh x + 1), \quad x \in \mathbb{R}.$$

The corresponding density function activated by the above sigmoidal function is as follows,

$$\chi_\sigma(x) = \frac{1}{2} \left[\frac{x^2(e^4 - 1)}{x^2(1 + e^4 + e^2x^2) + e^2} \right], \quad \forall x \in \mathbb{R}^+.$$

Now, let $0 < \nu < 1$ and $n \in \mathbb{N}$. Indeed,

$$\begin{aligned} \sum_{|\frac{k}{n} - \log(x)| \geq n^{1-\nu}} \chi_\sigma(e^{-k}x^n) &= \sum_{|\frac{k}{n} - \log(x)| \geq n^{1-\nu}} \chi_\sigma(|e^{-k}x^n|) \\ &< \frac{(e^4 - 1)}{2e^2} \int_{n^{1-\nu}}^{+\infty} \frac{1}{x^2} dx \\ &= \frac{(e^4 - 1)}{2e^2} \frac{1}{n^{1-\nu}} \\ &= \frac{3.6268}{n^{1-\nu}}. \end{aligned}$$

Using the above estimate, we establish a quantitative estimate of order of convergence in $C([a, b])$ using the notion of *logarithmic modulus of continuity*.

The logarithmic modulus of continuity is defined as

$$\omega(f, \delta) := \sup\{|f(x) - f(y)|; \text{whenever } |\log(x) - \log(y)| \leq \delta, \delta \in \mathbb{R}^+\}.$$

The properties of logarithmic modulus of continuity can be seen in [10].

Theorem 3 Let $[a, b] \subset \mathbb{R}^+$ and $f \in C([a, b])$. Then, for $0 < \nu < 1$, we have the following estimate,

$$|E_n^{\chi_\sigma}(f, x) - f(x)| \leq (4.14925) \left(\omega \left(f, \frac{1}{n^\nu} \right) + 7.2536 \|f\|_\infty n^{(\nu-1)} \right),$$

where, $0 < \nu < 1$, $n \in \mathbb{N}$.

Proof We have

$$\begin{aligned} |E_n^{\chi_\sigma}(f, x) - f(x)| &= \frac{\left| \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) (f(e^{\frac{k}{n}}) - f(x)) \right|}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |\chi_\sigma(e^{-k} x^n)|} \\ &\leq \frac{1}{|\chi_\sigma(e)|} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |\chi_\sigma(e^{-k} x^n)| \left| f(e^{\frac{k}{n}}) - f(x) \right| \quad (2) \end{aligned}$$

Now, we can write

$$\begin{aligned} \left| \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} (f(e^{\frac{k}{n}}) - f(x)) \chi_\sigma(e^{-k} x^n) \right| &\leq \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |f(e^{\frac{k}{n}}) - f(x)| |\chi_\sigma(e^{-k} x^n)| \\ &= \sum_{|\frac{k}{n} - \log(x)| < \frac{1}{n^\nu}} |\chi_\sigma(e^{-k} x^n)| |f(e^{\frac{k}{n}}) - f(x)| + \\ &\quad \sum_{|\frac{k}{n} - \log(x)| \geq \frac{1}{n^\nu}} |\chi_\sigma(e^{-k} x^n)| |f(e^{\frac{k}{n}}) - f(x)| \\ &\leq \sum_{|\frac{k}{n} - \log(x)| < \frac{1}{n^\nu}} |\chi_\sigma(e^{-k} x^n)| \omega \left(f, \frac{1}{n^\nu} \right) + 2 \|f\|_\infty \sum_{|\frac{k}{n} - \log(x)| \geq \frac{1}{n^\nu}} |\chi_\sigma(e^{-k} x^n)| \\ &\leq \omega \left(f, \frac{1}{n^\nu} \right) + 2 \|f\|_\infty \left(\frac{3.6268}{n^{1-\nu}} \right). \end{aligned}$$

Using this estimate in (2), we establish the result.

In order to derive the higher order of approximation, we use the Taylor's formula in terms of Mellin's derivatives as the Mellin's analysis is the suitable frame to put the rigorous theory of the exponential sampling (see [43, 13]).

Theorem 4 Let $[a, b] \subset \mathbb{R}^+$ and $f \in C^{(2)}([a, b])$. Then, for $0 < \nu < 1$, the following estimate holds

$$|E_n^{\chi_\sigma}(f, x) - f(x)| \leq (4.14925) \left\{ \sum_{i=1}^2 \frac{\|\theta^{(i)} f\|_\infty}{i!} \left(n^{-i\nu} + |b-a|^i \frac{3.6268}{n^{1-\nu}} \right) + \frac{\omega(\theta^{(2)} f, n^{-\nu})}{2n^{2\nu}} + \frac{(3.6268)}{n^{1-\nu}} \|\theta^{(2)} f\|_\infty (b-a)^2 \right\}.$$

Proof Let $f \in C^{(2)}[a, b]$. Then, by the Taylor's formula in terms of Mellin's derivatives with integral form of remainder, we have

$$f(e^u) = f(x) + (\theta f)(x)(u - \log x) + \int_{\log x}^u (\theta^{(2)} f)(e^\xi)(u - \xi) d\xi, \quad \text{where, } \xi \in (\log x, u).$$

Now, we can write

$$\begin{aligned} & \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} f(e^{\frac{k}{n}}) \chi_\sigma(e^{-k} x^n) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) \right) \\ &= (\theta f)(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) \left(\frac{k}{n} - \log x \right) + \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) \int_{\log x}^{\frac{k}{n}} (\theta^{(2)} f)(e^\xi) \left(\frac{k}{n} - \xi \right) d\xi \\ &= \sum_{i=1}^2 \frac{(\theta^{(i)} f)(x)}{i!} \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) \right) \left(\frac{k}{n} - \log x \right)^i + \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) \int_{\log x}^{\frac{k}{n}} ((\theta^{(2)} f)(e^\xi) - (\theta^{(2)} f)(x)) \left(\frac{k}{n} - \xi \right) d\xi. \\ &= \sum_{i=1}^2 \frac{(\theta^{(i)} f)(x)}{i!} \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \chi_\sigma(e^{-k} x^n) \right) \left(\frac{k}{n} - \log x \right)^i + I \text{ (say)}. \end{aligned}$$

Now, I can be written as

$$I = \left(\sum_{\left| \frac{k}{n} - \log(x) \right| < \frac{1}{n^\nu}} + \sum_{\left| \frac{k}{n} - \log(x) \right| \geq \frac{1}{n^\nu}} \right) \chi_\sigma(e^{-k} x^n) \int_{\log x}^{\frac{k}{n}} ((\theta^{(2)} f)(e^\xi) - (\theta^{(2)} f)(x)) \left(\frac{k}{n} - \xi \right) d\xi.$$

In the view of modulus of continuity and the estimate of $\sum_{\left| \frac{k}{n} - \log(x) \right| \geq \frac{1}{n^\nu}} |\chi_\sigma(e^{-k} x^n)|$,

we obtain

$$|I| \leq \left(\frac{1}{2n^{2\nu}} \omega(\theta^{(2)} f, n^{-\nu}) + \frac{(3.6268)}{n^{1-\nu}} \|\theta^{(2)} f\|_\infty (b-a)^2 \right).$$

Further, we estimate $\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} |\chi_\sigma(e^{-k} x^n)| \left| \frac{k}{n} - \log x \right|^i$

$$\begin{aligned} &= \sum_{\left| \frac{k}{n} - \log x \right| \leq n^{-\nu}} |\chi_\sigma(e^{-k} x^n)| \left| \frac{k}{n} - \log x \right|^i + \sum_{\left| \frac{k}{n} - \log x \right| \geq n^{-\nu}} |\chi_\sigma(e^{-k} x^n)| \left| \frac{k}{n} - \log x \right|^i \\ &\leq n^{-i\nu} \sum_{\left| \frac{k}{n} - \log x \right| \leq n^{-\nu}} |\chi_\sigma(e^{-k} x^n)| + |b-a|^i \sum_{\left| \frac{k}{n} - \log x \right| \geq n^{-\nu}} |\chi_\sigma(e^{-k} x^n)| \\ &\leq n^{-i\nu} + |b-a|^i \frac{3.6268}{n^{1-\nu}}. \end{aligned}$$

On combining all the above estimates, we get the desired result.

Remark 2: Let $f \in C^{(N)}([a, b])$, $N \in \mathbb{N}$. Then, the higher order Taylor's formula in terms of Mellin's derivatives with integral form of remainder is given by

$$f(e^u) = f(x) + (\theta f)(x)(u - \log x) + \frac{(\theta^{(2)} f)(x)}{2!} (u - \log(x))^2 + \dots + \int_{\log x}^u \frac{(\theta^{(n)} f)(e^\xi)}{(n-1)!} (u - \xi)^{n-1} d\xi$$

where, $\xi \in (\log x, u)$. The proof of this Mellin's Taylor formula can be easily obtained by using the fundamental theorem of integral calculus.

4 Exponential Type Quasi-Interpolation Operators

In the present section, we define the notion of exponential analogue of quasi-interpolation operators ([4, 7, 24, 33]). Let σ be the sigmoidal function satisfying all the conditions (1) – (3) and $\chi_\sigma(\cdot)$ be the corresponding density function generated by the sigmoidal function σ . Then, for every $x \in \mathbb{R}^+$, $k \in \mathbb{Z}$ and $f : \mathbb{R}^+ \rightarrow \mathbb{R}$, the exponential type quasi-interpolation operators are defined as,

$$Q_n(f, x) := \sum_{k=-\infty}^{\infty} f\left(e^{\frac{k}{n}}\right) \chi_\sigma(e^{-k} x^n), \quad \forall n \in \mathbb{N}.$$

The above operators are well-defined for the class of all real valued bounded functions defined on \mathbb{R}^+ .

First, we derive the point-wise and uniform convergence theorem for these operators in $\mathcal{C}(\mathbb{R}^+)$.

Theorem 5 *Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be any bounded function. Then, $(Q_n^{\chi_\sigma}(f, x))$ converges to $f(x)$ at every point of continuity of f . Moreover, if $f \in \mathcal{C}(\mathbb{R}^+)$, then*

$$\lim_{n \rightarrow \infty} \|Q_n^{\chi_\sigma}(f, \cdot) - f\|_\infty = 0.$$

Proof The proof follows in the similar manner as discussed in Theorem 1, with the application of Lemma 1.

Moreover, we obtain a similar type of quantitative estimate of order of approximation for the family of operators $(Q_n^{\chi_\sigma} f)$, provided f is bounded on \mathbb{R}^+ .

Theorem 6 *Let $f \in \mathcal{C}(\mathbb{R}^+)$ be any bounded function. Then, the following estimate holds*

$$|Q_n^{\chi_\sigma}(f, x) - f(x)| \leq \left(\omega\left(f, \frac{1}{n^\nu}\right) + 7.2536 \|f\|_\infty n^{(\nu-1)} \right)$$

where, $0 < \nu < 1$, $n \in \mathbb{N}$.

Proof The proof can be obtained by following the proof of Theorem 3.

5 Multivariate Exponential Type Neural Network Operators

In this section, we extend the concept of exponential type neural network operators in the multivariate setting. Let $\mathbf{x} \in (\mathbb{R}^+)^N$, $N \in \mathbb{N}$. Then, the multivariate density function activated by the sigmoidal function σ satisfying the assumptions (1) – (3), is defined as

$$\bar{\chi}_\sigma^N(\mathbf{x}) := \prod_{i=1}^N \chi_\sigma(x_i), \quad x_i \in \mathbb{R}^+$$

where, each $\chi_\sigma(x_i)$ is the univariate density function defined on \mathbb{R}^+ . As the multivariate density function $\bar{\chi}_\sigma^N(\mathbf{x})$ is the finite product of univariate density functions $\chi_\sigma(x_i)$, it can easily be shown that $\bar{\chi}_\sigma^N(\mathbf{x})$ also satisfies the multivariate extensions of Lemma 1, 2 and 3 (see [5, 6]). Now, the multivariate exponential type neural network operators can be defined as follows,

$$E_n^{\bar{\chi}_\sigma^N}(f, \mathbf{x}) = \frac{\sum_{k_1=\lceil na \rceil}^{\lfloor nb \rfloor} \dots \sum_{k_N=\lceil na \rceil}^{\lfloor nb \rfloor} f(e^{\frac{\hat{k}}{n}}) \hat{\chi}_\sigma^N(e^{-\hat{k}} \mathbf{x}^n)}{\sum_{k_1=\lceil na \rceil}^{\lfloor nb \rfloor} \dots \sum_{k_N=\lceil na \rceil}^{\lfloor nb \rfloor} \hat{\chi}_\sigma^N(e^{-\hat{k}} \mathbf{x}^n)}, \quad n \in \mathbb{N}$$

where, $\hat{k} = (k_1, k_2, \dots, k_N) \in \mathbb{Z}^N$. One can study the point-wise and uniform approximation theorems for the multivariate extension of exponential type neural network operators activated by sigmoidal functions, by following the same proof technique as discussed earlier.

6 Examples

In this section, we present few examples of the sigmoidal function $\sigma(x)$ which satisfy the assumptions of the proposed theory. We begin with an example of sigmoidal function which is known as *logistic function* ([7]) and defined as,

$$\sigma_l(x) = \frac{1}{1 + e^{-x}}, \quad x \in \mathbb{R}.$$

It is easy that the logistic function satisfies all the conditions (1) – (3) of section 1 (see [24]).

Another example of such smooth sigmoidal function is *hyperbolic tangent* sigmoidal function ([4]), which is defined as,

$$\sigma_h(x) := \frac{1}{2}(\tanh x + 1), \quad x \in \mathbb{R}.$$

Moreover, we can construct the example of sigmoidal function satisfying all the assumptions (1) – (3) with the help of well-known *B-spline functions* ([15,24]) of order n , by defining as follows,

$$\sigma_{M_n}(x) := \int_{-\infty}^x M_n(t)dt, \quad x \in \mathbb{R}.$$

It is clear that $\sigma_{M_n}(x)$ is non-decreasing on \mathbb{R}^+ and $0 \leq \sigma_{M_n}(x) \leq 1$ for every $n \in \mathbb{N}$ and $x \in \mathbb{R}^+$. Indeed,

$$\sigma_{M_1}(x) = \begin{cases} 0, & \text{if, } x < -\frac{1}{2} \\ x + \frac{1}{2} & \text{if, } -\frac{1}{2} \leq x \leq \frac{1}{2} \\ 1, & \text{if, } x > \frac{1}{2} \end{cases}$$

This function is also known as *ramp-function* (see [4,5]) which is an example of a discontinuous sigmoidal function.

Similarly, we have

$$\sigma_{M_2}(x) = \begin{cases} \frac{(1+x)^2}{2} & \text{if, } -1 < x < 0 \\ \left(1 - \frac{(x-1)^2}{2}\right) & \text{if, } 0 \leq x < 1. \end{cases}$$

Conclusion: In this paper, we introduced and studied the behaviour of the new family of Exponential type neural network operators activated by the sigmoidal functions. We established the point-wise and uniform approximation theorems for these NN (Neural Network) operators in $\mathcal{C}[a, b]$. We also studied the order of convergence for these operators in the class of *log-holderian* functions. Further, the quantitative estimates of order of approximation for the proposed NN operators in $C^N[a, b]$ established in terms of logarithmic modulus of continuity. We also analyzed the behaviour of the family of exponential type quasi-interpolation operators in $\mathcal{C}(\mathbb{R}^+)$ and the quantitative estimate of order of convergence in $C(\mathbb{R}^+)$. Finally, we propose the multivariate extension of exponential type NN operators and few examples of the sigmoidal functions to which the present theory can be applied.

References

1. Anastassiou, G. A.: Rate of convergence of some neural network operators to the unit-univariate case. *J. Math. Anal. Appl.* **212**(1), 237-262(1997).
2. Anastassiou, G. A.: *Quantitative Approximations*. New York, Chapman and Hall/CRC (2001).
3. Anastassiou, G. A.: *Intelligent systems: approximation by artificial neural networks*. Intelligent Systems Reference Library. **19**. Springer-Verlag, Berlin (2011).
4. Anastassiou, G. A.: Univariate hyperbolic tangent neural network approximation. *Math. Comput. Modelling.* **53**, 1111-1132(2011).

5. Anastassiou, G.A.: Multivariate hyperbolic tangent neural network approximation. *Comput. Math. Appl.* **61**(4), 809-821(2011).
6. Anastassiou, G.A.: Multivariate sigmoidal neural network approximation, *Neural Networks.* **24**, 378-386(2011).
7. Anastassiou, G. A.: Univariate sigmoidal neural network approximation. *J. Comput. Anal. Appl.* **14**(4), 659-690(2012).
8. Balsamo, S., Mantellini, I.: On Linear Combinations of General Exponential Sampling Series. *Results Math.* **74**(4), 180(2019).<https://doi.org/10.1007/s00025-019-1104-x>.
9. Bardaro, C., Butzer, P.L., Mantellini, I.: The exponential sampling theorem of signal analysis and the reproducing kernel formula in the Mellin transform setting. *Sampl. Theory Signal Image Process.* **13**(1), 35-66 (2014).
10. Bardaro, C., Mantellini, I.: On Mellin convolution operators: A direct approach to the asymptotic formulae. *Integral Transforms Spec. Funct.* **25**(3), 182-195 (2014).
11. Bardaro, C., Butzer, P.L., Mantellini, I.: The Mellin-Parseval formula and its interconnections with the exponential sampling theorem of optical physics. *Integral Transforms Spec. Funct.* **27** (1), 17-29(2016).
12. Bardaro, C., Butzer, P.L., Mantellini, I., Schmeisser, G.: On the Paley-Wiener theorem in the Mellin transform setting. *J. Approx. Theory.* **207** , 60-75(2016).
13. Bardaro, C., Faina, L., Mantellini, I.: A generalization of the exponential sampling series and its approximation properties. *Math. Slovaca.* **67**(6), 1481-1496 (2017).
14. Bardaro, C., Mantellini, I., Schmeisser, G.: Exponential sampling series: convergence in Mellin-Lebesgue spaces. *Results Math.* **74**(3), (2019).<https://doi.org/10.1007/s00025-019-1044-5>.
15. Butzer, P. L., Nessel, R. J. : Fourier analysis and approximation. Volume 1: One-dimensional theory. *Pure and Applied Mathematics.* **40**. Academic Press, New York-London(1971).
16. Butzer P.L., Stens, R.L.: Linear prediction by samples from the past. In: *Advanced Topics in Shannon Sampling and Interpolation Theory* (Ed. R. J. Marks II) Springer Texts Electrical Eng., Springer, New York(1993).
17. Butzer, P. L., Jansche, S.: A direct approach to the Mellin transform. *J. Fourier Anal. Appl.* **3**(4), 325-376(1997).
18. Butzer, P.L., Jansche, S.: The finite Mellin transform, Mellin-Fourier series, and the Mellin-Poisson summation formula. *Proceedings of the Third International Conference on Functional Analysis and Approximation Theory, Vol. I* (Acquafredda di Maratea, 1996). *Rend. Circ. Mat. Palermo* (2) Suppl. No. 52, Vol. I , 55-81(1998).
19. Butzer, P.L., Jansche, S.: The exponential sampling theorem of signal analysis, Dedicated to Prof. C. Vinti (Italian) (Perugia, 1996), *Atti. Sem. Mat. Fis. Univ. Modena, Suppl.* **46**, 99-122(1998).
20. Butzer, P.L., Stens, R.L.: A self contained approach to Mellin transform analysis for square integrable functions; applications. *Integral Transform. Spec. Funct.* **8**(3-4), 175-198(1999).
21. Bertero, M., Pike, E.R.: Exponential-sampling method for Laplace and other dilationally invariant transforms. II. Examples in photon correlation spectroscopy and Fraunhofer diffraction. *Inverse Problems.* **7**(1), 21-41(1991).
22. Cardaliaguet, P., Euvrard, G.: Approximation of a function and its derivative with a neural network. *Neural Networks,* **5**(2), 207220(1992).
23. Casasent, D.: Optical signal processing. In *Optical Data Processing.* Springer, Berlin, Heidelberg, 241-282(1978).
24. Costarelli D., Spigler, R.: Approximation results for neural network operators activated by sigmoidal functions, *Neural Networks.* **44**, 101-106(2013).
25. Costarelli D., Spigler, R.: Constructive approximation by superposition of sigmoidal functions, *Anal. Theory Appl.* **29**(2), 169-196(2013).
26. Costarelli, D., Spigler, R.: Multivariate neural network operators with sigmoidal activation functions. *Neural Networks.* **48**, 72-77(2013).
27. Costarelli, D., Spigler, R.: Convergence of a family of neural network operators of the Kantorovich type. *J. Approx. Theory.* **185**, 80-90(2014).
28. Costarelli, D.: Neural network operators: constructive interpolation of multivariate functions. *Neural Networks.* **67**, 28-36(2015).

29. Costarelli, D., Spigler, R., Vinti, G. : A survey on approximation by means of neural network operators. *Journal of NeuroTechnology*. **1**(1),29-52(2016)
30. Costarelli, D., Vinti, G.: Quantitative estimates involving K -functionals for neural network-type operators. *Appl. Anal.* **98**(15), 2639-2647(2019).
31. Cheang, Gerald H. L.: Approximation with neural networks activated by ramp sigmoids. *J. Approx. Theory*. **162**(8), 1450-1465(2010).
32. Cheney, E. W., Light, W. A., Xu, Y.: On kernels and approximation orders. *Approximation theory (Memphis, TN, 1991)*, 227-242, *Lecture Notes in Pure and Appl. Math.* **138**, Dekker, New York, 1992.
33. Chen, Z., Cao, F.: The construction and approximation of a class of neural networks operators with ramp functions. *J. Comput. Anal. Appl.* **14**(1), 101-112(2012).
34. Chen, T., Chen, H.: Universal approximation capability of EBF neural networks with arbitrary activation functions. *Circuits Systems Signal Process.* **15**(5), 671-683(1996).
35. Cybenko, G.: Approximation by superpositions of sigmoidal function. *Math. of Control Signals and System.* **2**, 303-314 (1989).
36. Chui, Charles K., Li, Xin.: Approximation by ridge functions and neural networks with one hidden layer. *J. Approx. Theory*. **70** (2), 131-141(1992).
37. Funahashi, K. I.: On the approximate realization of continuous mappings by neural networks. *Neural Networks*, **2**, 183-192 (1989).
38. Gori, F.: Sampling in optics. *Advanced topics in Shannon sampling and interpolation theory*. Springer Texts Electrical Engrg., Springer, New York, 37-83(1993).
39. Hahm, N., Hong, B. I.: An approximation by neural networks with a fixed weight. *Comput. Math. Appl.* **47**(12), 1897-1903(2004).
40. Hornik, K., Stinchcombe, M., White, H.: Multilayer feedforward networks are universal approximators. *Neural Networks*. **2**, 359-366 (1989).
41. Leshno, M., Lin, V.Y., Pinks, A., Schocken, S.: Multilayer feedforward networks with a nonpolynomial activation function can approximate any function. *Neural Networks*. **6**, 861-867 (1993).
42. Lenze, B.: Constructive multivariate approximation with sigmoidal functions and applications to neural networks. *Numerical methods in approximation theory*, Vol. 9 (Oberwolfach, 1991), 155-175, *Internat. Ser. Numer. Math.* **105**, Birkhuser, Basel, 1992.
43. Mamedov, R. G.: *The Mellin transform and approximation theory*,(in Russian), 'Im', Baku, 1991.273 pp.
44. Mhaskar, H. N.; Micchelli, Charles A. Approximation by superposition of sigmoidal and radial basis functions. *Adv. in Appl. Math.* **13**, 350-373(1992) .
45. Ostrowsky, N., Sornette, D., Parke, P., Pike, E.R.: Exponential sampling method for light scattering polydispersity analysis, *Opt. Acta.* **28** , 1059-1070(1994).