

QUANTITATIVE REDUCTION THEORY AND UNLIKELY INTERSECTIONS

CHRISTOPHER DAW AND MARTIN ORR

ABSTRACT. We prove quantitative versions of Borel and Harish-Chandra’s theorems on reduction theory for arithmetic groups. Firstly, we obtain polynomial bounds on the lengths of reduced integral vectors in any rational representation of a reductive group. Secondly, we obtain polynomial bounds in the construction of fundamental sets for arithmetic subgroups of reductive groups, as the latter vary in a $\mathbf{G}(\mathbb{R})$ -conjugacy class of subgroups of a fixed reductive group \mathbf{G} .

Our results allow us to apply the Pila–Zannier strategy to the Zilber–Pink conjecture for the moduli space of principally polarised abelian surfaces. Building on our previous paper [DO], we prove this conjecture under a Galois orbits hypothesis. Finally, we establish the Galois orbits hypothesis for points corresponding to abelian surfaces with quaternionic multiplication, under certain geometric conditions.

1. INTRODUCTION

Reduction theory is concerned with finding small representatives for each orbit in actions of arithmetic groups, for example through constructing fundamental sets. In the 18th and 19th centuries, the focus was on the action of $\mathrm{SL}_n(\mathbb{Z})$ on quadratic forms, motivated at first by questions about which integers can be represented by a given quadratic form and subsequently by applications to algebraic number theory. (For a historical account of the reduction theory of quadratic forms, see [SO85].)

In the 20th century, Siegel saw that the reduction theory of positive definite quadratic forms was equivalent to the construction of a fundamental set for $\mathrm{SL}_n(\mathbb{Z})$ in $\mathrm{SL}_n(\mathbb{R})$. Borel and Harish-Chandra generalised this to arithmetic lattices in arbitrary semisimple Lie groups, which has had wide-ranging applications including the theory of automorphic forms and locally symmetric spaces [AMRT75], the arithmetic of algebraic groups [PR94] and finiteness theorems for abelian varieties [Mil86].

The first goal of this paper is to prove quantitative bounds for the group elements used in Borel and Harish-Chandra’s construction of fundamental sets. These bounds are polynomial in terms of suitable input parameters, although they are

2010 *Mathematics Subject Classification.* 11F06, 11G18.

Key words and phrases. Reduction theory, arithmetic groups, Zilber–Pink conjecture, unlikely intersections.

not fully effective. They complement the second author’s polynomial bounds for the Siegel property [Orr18] and generalise Li and Margulis’s polynomial bounds for the reduction theory of quadratic forms [LM16]. It should be noted that while Borel and Harish-Chandra’s reduction theory is algorithmic in nature, as made explicit by Grunewald and Segal [GS80], their arguments do not give bounds on the running time or output size of these algorithms.

The second goal of the paper is to apply these quantitative bounds in reduction theory to the Zilber–Pink conjecture on unlikely intersections. We will prove the Zilber–Pink conjecture for \mathcal{A}_2 , the moduli space of principally polarised abelian surfaces, subject to a large Galois orbits conjecture which we show holds if the curve under consideration satisfies a multiplicative reduction hypothesis at the boundary of the moduli space. The new case of the Zilber–Pink conjecture which we consider in this paper is that of intersections with (quaternionic) Shimura curves in \mathcal{A}_2 ; this combines with the authors’ previous work on intersections with “ $E \times \text{CM}$ curves” [DO] and Pila and Tsimerman’s proof of the André–Oort conjecture for \mathcal{A}_2 [PT13] to cover all cases of the Zilber–Pink conjecture for \mathcal{A}_2 .

The results on quantitative reduction theory in this paper will be an important tool for proving the Zilber–Pink conjecture for other Shimura varieties, which will be the subject of future work by the authors. We will also apply these results to obtain a uniform version of the second author’s bounds for polarisations and isogenies of abelian varieties [Orr17]. By combining the results of this paper with techniques from homogeneous dynamics, as in [LM16], we will obtain bounds for the heights of generators of arithmetic groups.

1.A. Reduction theory for quadratic forms. The group $\text{SL}_n(\mathbb{Z})$ acts on the set of integral quadratic forms in n variables, by acting on the variables. The classical reduction theory of quadratic forms defines a set of *reduced quadratic forms* with the following properties.

Properties 1.1.

- (i) *Each $\text{SL}_n(\mathbb{Z})$ -orbit of non-degenerate integral quadratic forms contains at least one reduced form.*
- (ii) *Each $\text{SL}_n(\mathbb{Z})$ -orbit of non-degenerate integral quadratic forms contains only finitely many reduced forms.*

A variety of definitions of reduced quadratic forms are used, possessing varying properties in addition to Properties 1.1. The most important definitions are due to Lagrange and Gauss for binary quadratic forms, and to Hermite, Minkowski and Siegel for quadratic forms in any number of variables. In the definitions of Lagrange, Gauss, Minkowski and Siegel, the sets of reduced quadratic forms can be defined by finite collections of polynomial inequalities in the coefficients of the forms. In other words, if we consider quadratic forms with real coefficients (not just integral ones) then the sets of reduced forms are semialgebraic.

When we generalise to quadratic forms with real coefficients, positive (or negative) definite forms are better behaved than indefinite forms. Properties 1.1(i) and (ii) both remain true for definite real quadratic forms, and indeed they satisfy a much stronger version of (ii) called the Siegel property which guarantees that there is a uniform bound on the number of reduced quadratic forms in each $\mathrm{SL}_n(\mathbb{Z})$ -orbit (for fixed n). In the nicest case of all, definite binary quadratic forms using Gauss's definition of reduced, each $\mathrm{SL}_2(\mathbb{Z})$ -orbit contains *exactly one* reduced form.

For indefinite quadratic forms, Property 1.1(i) holds for forms with real coefficients. However there is no topologically reasonable definition of reduced quadratic forms which satisfies Property 1.1(ii) for indefinite real quadratic forms. Furthermore, an $\mathrm{SL}_n(\mathbb{Z})$ -orbit of integral indefinite quadratic forms can contain arbitrarily many reduced forms.

Quantitative reduction theory for quadratic forms. One way of proving Property 1.1(ii) is via the following stronger statement.

Lemma 1.2. *For each integer $\Delta \neq 0$, there are only finitely many reduced integral quadratic forms in n variables of discriminant Δ .*

The following quantitative version of Lemma 1.2 is classical for anisotropic quadratic forms [Cas78, p. 287, Cor. 1], using the definitions of reduction due to Hermite and Minkowski (note that anisotropic forms are always definite when $n \geq 5$). Indeed, for binary anisotropic forms, it goes back to Lagrange. For Siegel reduced forms (definite or indefinite), it can be proved by adapting the proof of Lemma 1.2 found on [Cas78, pp. 322–324] but we are unsure whether this was classically known.

Proposition 1.3. *For each positive integer n , there exists a positive real number $C_1(n)$ such that, for every integer $\Delta \neq 0$, all reduced integral n -ary quadratic forms of discriminant Δ have coefficients whose absolute values are at most $C_1(n)|\Delta|$.*

Lemma 1.2 is equivalent to the conjunction of Property 1.1(ii) with the following statement, which does not mention reduced forms, and which was historically one of the most important consequences of reduction theory.

Lemma 1.4. *For each integer $\Delta \neq 0$, there are only finitely many $\mathrm{SL}_n(\mathbb{Z})$ -orbits of integral quadratic forms in n variables of discriminant Δ .*

Li and Margulis have proved a quantitative version of Lemma 1.4. The bound is stronger than in Proposition 1.3, but it applies only to at least one form in each $\mathrm{SL}_n(\mathbb{Z})$ -orbit, rather than to all reduced forms.

Proposition 1.5. [LM16, Theorem 3] *For each integer $n \geq 3$, there exists a constant $C_2(n)$ such that every $\mathrm{SL}_n(\mathbb{Z})$ -orbit of indefinite quadratic forms in n variables of discriminant $\Delta \neq 0$ contains a form whose coefficients have absolute value at most $C_2(n)|\Delta|^{1/n}$.*

1.B. **Siegel sets.** Siegel shifted the emphasis in reduction theory from quadratic forms to arithmetic groups. In the simplest case, there is a direct link between the reduction theory of positive definite quadratic forms and fundamental sets for $\mathrm{SL}_n(\mathbb{Z})$ in $\mathrm{SL}_n(\mathbb{R})$. Let v_0 denote the standard diagonal positive definite quadratic form in n variables:

$$v_0(x_1, \dots, x_n) = x_1^2 + \dots + x_n^2.$$

If \mathcal{F} is a fundamental set of *positive definite real* quadratic forms in n variables (that is, a set which satisfies Properties 1.1(i) and (ii) for all orbits of positive definite real forms), then

$$\mathfrak{S} = \{g \in \mathrm{SL}_n(\mathbb{R}) : gv_0 \in \mathcal{F}\}$$

is a fundamental set in $\mathrm{SL}_n(\mathbb{R})$ for the action of $\mathrm{SL}_n(\mathbb{Z})$ by multiplication on the left, that is, every right $\mathrm{SL}_n(\mathbb{Z})$ -coset intersects \mathfrak{S} in at least one, and at most finitely many elements. Conversely, if $\mathfrak{S} \subset \mathrm{SL}_n(\mathbb{R})$ is a fundamental set for $\mathrm{SL}_n(\mathbb{Z})$ which is invariant under right multiplication by $K = \mathrm{SO}_n(\mathbb{R}) = \mathrm{Stab}_{\mathrm{SL}_n(\mathbb{R})}(v_0)$, then $\mathbb{R}_{>0}\mathfrak{S}v_0$ is a fundamental set of positive definite real quadratic forms.

We could apply this construction to the set of Minkowski reduced positive definite forms in order to obtain a fundamental set in $\mathrm{SL}_n(\mathbb{R})$. However, Siegel instead defined a different fundamental set in $\mathrm{SL}_n(\mathbb{R})$ and used the reverse construction to obtain a fundamental set of positive definite quadratic forms [Sie40]. In fact, Siegel defined a family of sets $\mathfrak{S} = \mathfrak{S}_{t,u} \subset \mathrm{SL}_n(\mathbb{R})$, depending on two parameters $t, u \in \mathbb{R}_{>0}$. We call the set $\mathfrak{S}_0 = \mathfrak{S}_{2/\sqrt{3}, 1/2}$ the **standard Siegel set** in $\mathrm{SL}_n(\mathbb{R})$ – this is the smallest of Siegel’s sets to form a fundamental set for $\mathrm{SL}_n(\mathbb{Z})$. We say that a positive definite quadratic form is **Siegel reduced** if it lies in $\mathbb{R}_{>0}\mathfrak{S}_0v_0$. The set of Siegel reduced positive definite forms is slightly larger than the set of Minkowski reduced forms.

We also define Siegel reduced indefinite quadratic forms: a form of signature (p, q) is **Siegel reduced** if it lies in $\mathbb{R}_{>0}\mathfrak{S}_0v_0^{(p,q)}$, where

$$v_0^{(p,q)}(x_1, \dots, x_{p+q}) = x_1^2 + \dots + x_p^2 - x_{p+1}^2 - \dots - x_{p+q}^2. \tag{1}$$

Note that the set of Siegel reduced indefinite quadratic forms satisfies Properties 1.1(i) and (ii), but it is not a fundamental set because it does not satisfy the generalisation of Properties 1.1(ii) to real forms. The Siegel reduced indefinite quadratic forms are the same as those obtained from Hermite’s recipe for reduced indefinite quadratic forms, providing we use Siegel’s definition for reduced positive definite forms.

Borel and Harish-Chandra generalised the notion of Siegel set from $\mathrm{SL}_n(\mathbb{R})$ to all reductive Lie groups [BHC62, sec. 4.1] (which they called **Siegel domains**). However Borel and Harish-Chandra’s Siegel domains do not give rise to fundamental sets for general arithmetic subgroups – in general, one can only say that there is a fundamental set *contained in* a finite union of translates of Siegel domains

[BHC62, Thm. 6.5, Lemma 7.5] (we will discuss this construction of fundamental sets further below).

Borel subsequently gave a new definition of Siegel sets for reductive \mathbb{Q} -algebraic groups [Bor69, 12.3], taking into account the \mathbb{Q} -algebraic group structure and not just the Lie group structure. For any reductive \mathbb{Q} -algebraic group \mathbf{G} and any arithmetic subgroup $\Gamma \subset \mathbf{G}(\mathbb{Q})$, there is a finite union of $\mathbf{G}(\mathbb{Q})$ -translates of a Siegel set which forms a fundamental set for Γ in $\mathbf{G}(\mathbb{R})$ [Bor69, Thm. 13.1]. In this paper we shall use a minor modification of Borel’s definition of Siegel sets, described in section 2.B.

1.C. Reduction theory for representations of reductive groups. The following result is a key step in Borel and Harish-Chandra’s construction of fundamental sets for arithmetic groups.

Proposition 1.6. [BHC62, Lemma 5.4] *Let \mathbf{G} be a reductive \mathbb{Q} -algebraic group whose \mathbb{Q} -rank is equal to its \mathbb{R} -rank. Let $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$ be a Siegel set. Let $\rho: \mathbf{G} \rightarrow \mathbf{GL}(V)$ be a representation of \mathbf{G} defined over \mathbb{Q} . Let $\Lambda \subset V$ be a \mathbb{Z} -lattice. Let $v \in V_{\mathbb{R}}$ be such that:*

- (i) *the orbit $\rho(\mathbf{G}(\mathbb{R}))v$ is closed in $V_{\mathbb{R}}$;*
- (ii) *the stabiliser $\text{Stab}_{\mathbf{G}(\mathbb{R}),\rho}(v)$ is self-adjoint.*

Then $\rho(\mathfrak{S})v \cap \Lambda$ is finite.

The restriction on the \mathbb{Q} -rank of \mathbf{G} in Proposition 1.6 can be removed with only minor alterations to the proof, provided we use the definition of Siegel sets from [Bor69, 12.3] (or the definition in section 2.B of this paper) instead of the definition of Siegel domains from [BHC62, 4.1] (the condition on the \mathbb{Q} -rank is precisely the condition under which the definitions of Siegel sets in [BHC62] and [Bor69] are essentially equivalent).

As noted in [BHC62, Example 5.5], Proposition 1.6 implies Lemma 1.2, by applying it to the representation of $\mathbf{SL}_n(\mathbb{R})$ on the vector space of real quadratic forms in n variables, with $v = \lambda v_0^{(p,q)}$ where $\lambda \in \mathbb{R}_{>0}$ and $v_0^{(p,q)}$ is defined by equation (1). Then the orbit $\mathbf{SL}_n(\mathbb{R})v$ is the set of all quadratic forms of signature (p, q) and discriminant $(-1)^q \lambda^n$, and $\mathfrak{S}v \cap L$ is the set of Siegel reduced integral quadratic forms of given signature and discriminant.

Quantitative reduction theory for representations. This paper’s first main theorem is a quantitative version of Proposition 1.6, bounding the length of vectors in the finite set $\rho(\mathfrak{S})v \cap L$ as we vary v (while fixing the group \mathbf{G} and representation ρ). We are not able to prove such a bound for all $v \in V_{\mathbb{R}}$: we must restrict to a set of v for which $\mathbf{G}(\mathbb{R})$ acts “in the same way” on all of the permitted vectors v . This is achieved through the condition that v must lie in the $\text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$ -orbit of a fixed vector.

Theorem 1.7. *Let \mathbf{G} be a reductive \mathbb{Q} -algebraic group and let $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$ be a Siegel set. Let $\rho: \mathbf{G} \rightarrow \mathbf{GL}(V)$ be a representation of \mathbf{G} defined over \mathbb{Q} . Let $\Lambda \subset V$ be a \mathbb{Z} -lattice. Let $v_0 \in V_{\mathbb{R}}$ be such that:*

- (i) $\rho(\mathbf{G}(\mathbb{R}))v_0$ is closed in $V_{\mathbb{R}}$;
- (ii) the stabiliser $\text{Stab}_{\mathbf{G}(\mathbb{R}),\rho}(v_0)$ is self-adjoint.

Then there exist constants C_3, C_4 such that, for every $v \in \text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})v_0$ and every $w \in \rho(\mathfrak{S})v \cap \Lambda$, we have $|w| \leq C_3|v|^{C_4}$.

For example, taking $v_0 = v_0^{(p,q)}$ and noting that any scalar $\lambda \in \mathbb{R}$ is in $\text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$, we deduce that there are constants C_5 and C_6 (depending on n) such that

$$|w| < C_5|\lambda|^{C_6} \text{ for all } \lambda \in \mathbb{R} \text{ and } w \in \mathfrak{S}\lambda v_0 \cap L.$$

Thus Theorem 1.7 implies a weakened version of Proposition 1.3 – Proposition 1.3 is stronger because it gives a bound which is linear in the discriminant, while the constants in Theorem 1.7, even the exponent, are ineffective (see Remark 3.3).

In this example, the closed orbits in $V_{\mathbb{R}}$ are those which consist of non-degenerate quadratic forms. These orbits are parameterised by their signature and discriminant, so every closed orbit intersects $\text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})v_0^{(p,q)}$ for some signature (p, q) . Thus, in this case, Theorem 1.7 is sufficient to give a polynomial bound for integral elements of a reduced set in every closed orbit. In general however, there is no finite subset of $V_{\mathbb{R}}$ whose $\text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$ -orbit intersects every closed $\mathbf{G}(\mathbb{R})$ -orbit, and then Theorem 1.7 does not allow us to compare all closed orbits.

In this example, the representation ρ is absolutely irreducible so its only endomorphisms are scalars. In general, there may be more endomorphisms of ρ , and it will be important for our applications that we allow v to be any element of $\text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})v_0$, not just a scalar multiple of v_0 .

1.D. Fundamental sets for arithmetic groups. The central result of Borel and Harish-Chandra's reduction theory was the construction of fundamental sets for $\Gamma_{\mathbf{H}} \backslash \mathbf{H}(\mathbb{R})$, where \mathbf{H} is a reductive \mathbb{Q} -algebraic group and $\Gamma_{\mathbf{H}} \subset \mathbf{H}(\mathbb{Q})$ is an arithmetic subgroup. These fundamental sets are constructed by embedding \mathbf{H} into some \mathbf{GL}_n , where we already know how to obtain fundamental sets using standard Siegel sets. (Note that Borel and Harish-Chandra wrote \mathbf{G} where we write \mathbf{H} in this theorem; we use \mathbf{H} because in the next few paragraphs we will introduce a different reductive group which we will call \mathbf{G} .)

Theorem 1.8. [BHC62, Thm. 6.5] *Let \mathbf{H} be a reductive \mathbb{Q} -algebraic subgroup of $\mathbf{GL}_{n,\mathbb{Q}}$ and let $\Gamma_{\mathbf{H}} = \mathbf{GL}_n(\mathbb{Z}) \cap \mathbf{H}(\mathbb{R})$. Let $\mathfrak{S}_{\mathbf{GL}}$ be a standard Siegel set in $\mathbf{GL}_n(\mathbb{R})$ which forms a fundamental set for $\mathbf{GL}_n(\mathbb{Z})$. Let $u \in \mathbf{GL}_n(\mathbb{R})$ be such that $u^{-1}\mathbf{H}(\mathbb{R})u$ is self-adjoint.*

Then there exists a finite set $B \subset \mathbf{GL}_n(\mathbb{Z})$ such that

$$B\mathfrak{S}_{\mathbf{GL}}u^{-1} \cap \mathbf{H}(\mathbb{R})$$

is a fundamental set for $\Gamma_{\mathbf{H}}$ in $\mathbf{H}(\mathbb{R})$.

The ambient group \mathbf{GL}_n in Theorem 1.8 can be replaced by any reductive \mathbb{Q} -algebraic group \mathbf{G} such that $\mathbf{H} \subset \mathbf{G}$ with only minor alterations to the proof. However in order to apply Theorem 1.8 we need to know that some Siegel set (or the union of finitely many translates of a Siegel set) in this ambient group \mathbf{G} forms a fundamental set for an arithmetic subgroup of \mathbf{G} . For \mathbf{GL}_n , this is the content of Siegel’s reduction theory. For a general reductive group \mathbf{G} , this was not known at the time of [BHC62]. It is proved in [Bor69, §§ 12–15] but the proof makes use of Theorem 1.8! This argument is not circular because the proof that we can construct a fundamental set in $\mathbf{G}(\mathbb{R})$ from a Siegel set only requires Theorem 1.8 for \mathbf{H} replaced by \mathbf{G} , considered as a subgroup of some larger \mathbf{GL}_n . (Borel also outlines a second proof that Siegel sets give rise to fundamental sets in [Bor69, § 16], which can be made independent of Theorem 1.8.)

Quantitative fundamental sets for arithmetic groups. Our second main theorem is a quantitative version of Theorem 1.8, where we allow \mathbf{H} to vary over the \mathbb{Q} -algebraic members of a $\mathbf{G}(\mathbb{R})$ -conjugacy class of subgroups of some fixed reductive group \mathbf{G} . This theorem is “quantitative” in the sense that it controls the size of the elements of the finite set B . Ideally we would like to bound the height of elements of B but we have not yet achieved this (it may be possible by combining the methods of this paper with tools of homogeneous dynamics as in [LM16]). Instead we control the size of elements of B in terms of how they act on a vector v_u (whose stabiliser is \mathbf{H}) in a suitable representation of \mathbf{G} (with a similar flavour to Theorem 1.7). This turns out to be sufficient for our applications to unlikely intersections.

We only consider subgroups $\mathbf{H} \subset \mathbf{G}$ which are defined over \mathbb{Q} because this is necessary for $\Gamma \cap \mathbf{H}(\mathbb{R})$ to be a lattice in $\mathbf{H}(\mathbb{R})$. However it is very important that we allow \mathbf{H} to vary over a $\mathbf{G}(\mathbb{R})$ -conjugacy class, not just a $\mathbf{G}(\mathbb{Q})$ -conjugacy class, because this allows the \mathbb{Q} -algebraic subgroups in the conjugacy class to come from more than one isomorphism class over \mathbb{Q} ; indeed some of the groups \mathbf{H} in the conjugacy class may be \mathbb{Q} -anisotropic, meaning that the fundamental set in $\mathbf{H}(\mathbb{R})$ is compact, while others may be non- \mathbb{Q} -anisotropic, so that their fundamental set is not compact (for example the unit groups of quaternion algebras and $\mathbf{SL}_{2,\mathbb{Q}}$ can be found in the same $\mathbf{SL}_4(\mathbb{R})$ -conjugacy class of subgroups of \mathbf{SL}_4).

We should also remark that, while this theorem deals with a single $\mathbf{G}(\mathbb{R})$ -conjugacy class of subgroups of \mathbf{G} , all semisimple subgroups of \mathbf{G} belong to only finitely many such conjugacy classes [BDR, Cor. 0.2]. Note that this is not true for reductive subgroups, as may be seen by considering the torus \mathbb{G}_m^2 , which contains infinitely many non-conjugate subgroups isomorphic to \mathbb{G}_m – see [BDR, Remark 1.2].

Theorem 1.9. *Let \mathbf{G} be a reductive \mathbb{Q} -algebraic group. Let $\Gamma \subset \mathbf{G}(\mathbb{Q})$ be an arithmetic subgroup. Let $\mathcal{S} \subset \mathbf{G}(\mathbb{R})$ be a Siegel set such that $C\mathcal{S}$ is a fundamental set for Γ in $\mathbf{G}(\mathbb{R})$, for some finite set $C \subset \mathbf{G}(\mathbb{Q})$.*

Let $\rho: \mathbf{G} \rightarrow \mathbf{GL}(\Lambda_{\mathbb{Q}})$ be a \mathbb{Q} -algebraic representation of \mathbf{G} , where Λ is a finitely generated free \mathbb{Z} -module. Let $\mathbf{H}_0 \subset \mathbf{G}$ be a self-adjoint reductive \mathbb{Q} -algebraic subgroup and let $v_0 \in \Lambda$ be a vector such that:

- (i) $\text{Stab}_{\mathbf{G}, \rho}(v_0) = \mathbf{H}_0$;
- (ii) the orbit $\rho(\mathbf{G}(\mathbb{R}))v_0$ is closed in $\Lambda_{\mathbb{R}}$;
- (iii) for each $u \in \mathbf{G}(\mathbb{R})$, if the group $u\mathbf{H}_{0, \mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , then there exists $v_u \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ such that $\rho(u)v_u \in \Lambda$.

Then there exist positive constants C_7 and C_8 (depending only on \mathbf{G} , \mathbf{H}_0 , ρ and v_0) with the following property: for every $u \in \mathbf{G}(\mathbb{R})$, if $\mathbf{H}_u = u\mathbf{H}_{0, \mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , then there exists a fundamental set for $\Gamma \cap \mathbf{H}_u(\mathbb{R})$ in $\mathbf{H}_u(\mathbb{R})$ of the form

$$B_u C \mathfrak{S} u^{-1} \cap \mathbf{H}_u(\mathbb{R}),$$

where $B_u \subset \Gamma$ is a finite set such that

$$|\rho(b^{-1}u)v_u| \leq C_7 |v_u|^{C_8}$$

for every $b \in B_u$.

For example, consider the case where $\mathbf{G} = \mathbf{GL}_n$ and $\mathbf{H}_0 \subset \mathbf{GL}_n$ is the orthogonal group associated with the quadratic form $v_0^{p,q}$. The representation $\rho: \mathbf{G} \rightarrow \mathbf{GL}(\Lambda_{\mathbb{Q}})$, where Λ is the lattice of integral quadratic forms of signature (p, q) , satisfies the conditions of Theorem 1.9. In particular, (iii) holds because if \mathbf{H}_u is defined over \mathbb{Q} then it is the orthogonal group of some integral quadratic form v of the form $\rho(u)\lambda v_0$, where $\lambda \in \mathbb{R}^{\times}$ is a scalar such that $\text{disc}(v) = \pm \lambda^n$. As noted in [BHC62, 6.7], $\Sigma_u = \mathbf{H}_u(\mathbb{R}) / (u\mathbf{O}_n(\mathbb{R})u^{-1} \cap \mathbf{H}_u(\mathbb{R}))$ is the space of Hermite majorants of v and we get a fundamental set for $\Gamma \cap \mathbf{H}_u(\mathbb{R})$ in Σ_u which is the image in Σ_u of $B_u C \mathfrak{S} u^{-1} \cap \mathbf{H}_u(\mathbb{R})$. Now Theorem 1.9 allows us to control B_u in the sense that for each $b \in B_u$, the coefficients of the quadratic form $\rho(b^{-1})v$ are polynomially bounded in terms of $\text{disc}(v)$. A related result can be found in [LM16, §9.5], which bounds the entries of the matrices $b \in B_u$ (stronger than bounding $\rho(b^{-1})v$ directly), although its bound involves the entries of v as well as the discriminant.

As with Theorem 1.7, the constants in Theorem 1.9 are ineffective.

1.E. Unlikely intersections. Our interest in quantitative reduction theory came from so-called problems of unlikely intersections, specifically in the context of Shimura varieties. As an application of Theorem 1.9, we obtain some new results in these problems.

Let \mathcal{A}_2 denote the moduli space of principally polarised abelian surfaces over \mathbb{C} , which is a Shimura variety of dimension 3. Let $V \subset \mathcal{A}_2$ be an irreducible algebraic curve. The Zilber–Pink conjecture predicts that if V is Hodge generic in \mathcal{A}_2 (that is, V is not contained in a proper special subvariety of \mathcal{A}_2) then it should have only finitely many points of intersection with the special subvarieties of \mathcal{A}_2 having dimension ≤ 1 .

The fact that V contains only finitely many special points (special subvarieties of dimension 0) – a special case of the André–Oort conjecture – was proved by Pila and Tsimerman [PT13]. Furthermore, any point P on a special curve Z is either a special point or Z is the unique special curve containing P (in which case, we refer to P as a Hodge-generic point of Z). Therefore, the Zilber–Pink conjecture in this setting is implied by the statement that V contains only finitely many points that are Hodge-generic in some special curve Z .

The special curves in \mathcal{A}_2 are of three types:

- (1) curves parametrising abelian surfaces with quaternionic multiplication (we refer to these as quaternionic curves);
- (2) curves parametrising abelian surfaces isogenous to the square of an elliptic curve (“ E^2 curves”);
- (3) curves parametrising abelian surfaces isogenous to the product of two elliptic curves, at least one of which has complex multiplication (“ $E \times \text{CM}$ curves”).

In a previous article [DO], we considered the $E \times \text{CM}$ curves. We showed that V contains only finitely many points that are Hodge-generic in some $E \times \text{CM}$ curve provided V satisfies a so-called Galois orbits hypothesis. Furthermore, we showed that V satisfies this Galois orbits hypothesis if it is defined over $\overline{\mathbb{Q}}$ and its Zariski closure in the Baily–Borel compactification of \mathcal{A}_2 intersects the 0-dimensional stratum of the boundary.

In this article, we will consider the quaternionic curves and E^2 curves. As in [DO], the general strategy follows the proof of [DR18, Theorem 14.2], which was an application of the Pila–Zannier method to the Zilber–Pink conjecture for general Shimura varieties. However, we will have to make several modifications, and the end of the proof is closer to [Orr, Proposition 3.5]. Our main theorems on unlikely intersections are as follows. These theorems refer to Conjectures 5.2 and 5.3, which are Galois orbits hypotheses and are stated in section 5.

Theorem 1.10. *Let $V \subset \mathcal{A}_2$ denote an irreducible Hodge generic algebraic curve and let Σ_{Quat} denote the set of points of \mathcal{A}_2 which are Hodge generic in some quaternionic curve. Assume that V satisfies Conjecture 5.2.*

Then $V \cap \Sigma_{\text{Quat}}$ is finite.

Theorem 1.11. *Let $V \subset \mathcal{A}_2$ denote an irreducible Hodge generic algebraic curve and let Σ_{E^2} denote the set of points of \mathcal{A}_2 which are Hodge generic in some E^2 curve. Assume that V satisfies Conjecture 5.3.*

Then $V \cap \Sigma_{E^2}$ is finite.

Theorem 1.12. *Let V and Σ_{Quat} be as in Theorem 1.10.*

Suppose that V is defined over $\overline{\mathbb{Q}}$ and that the Zariski closure of V in the Baily–Borel compactification of \mathcal{A}_2 intersects the 0-dimensional stratum of the boundary of the compactification.

Then Conjecture 5.2 holds for V , and so $V \cap \Sigma_{\text{Quat}}$ is finite.

The proof of Theorem 1.12 uses a height bound for abelian varieties with large endomorphism algebras due to André [And89, Ch. X, Thm. 1.3], following a similar method to the proof of the analogous result for $E \times \text{CM}$ curves as in [DO]. Note that we have not been able to prove Conjecture 5.3 for curves which meet the 0-dimensional stratum of the Baily–Borel compactification, because André’s height bound does not apply to abelian surfaces isogenous to the square of an elliptic curve.

1.F. Outline of the paper. In section 2, we define Cartan involutions (verifying the equivalence between several definitions), Siegel sets and some notation which will be used throughout the paper. In section 3, we prove our main theorems on quantitative reduction theory, namely Theorems 1.7 and 1.9, which apply to all reductive \mathbb{Q} -algebraic groups and all representations.

In order to apply Theorem 1.9 to a specific situation, we need a suitable representation of the ambient reductive group. This is the topic of section 4 – we construct a representation of \mathbf{GSp}_4 which has the properties required to apply Theorem 1.9 to the subgroups associated with quaternionic curves and E^2 curves.

Finally section 5 contains the proofs of our theorems on unlikely intersections – Theorems 1.10 to 1.12, making use of the results from sections 3 and 4 to prove the parameter height bounds needed for the Pila–Zannier method.

Acknowledgements. The first author would like to thank the EPSRC for its support via a New Investigator Award (EP/S029613/1). He would also like to thank the Universities of Reading and Oxford. The second author would like to thank the EPSRC for its support via a New Investigator Award (EP/T010134/1). He would also like to thank the University of Warwick.

2. PRELIMINARIES

2.A. Cartan involutions. The theory of Cartan involutions is well-known for connected semisimple groups. However for reductive real algebraic groups, several definitions of Cartan involutions are used in the literature. The commonly used definitions are equivalent to each other, but the equivalences are not obvious and it is difficult to find proofs for all of the equivalences. Furthermore, many papers state without proof basic properties such as the existence and uniqueness of the Cartan involution associated with a given maximal compact subgroup. For convenience, we therefore give several definitions and prove that they are equivalent.

The following seems to us to be the most elegant definition.

Definition. Let \mathbf{G} be a reductive \mathbb{R} -algebraic group. A **Cartan involution** of \mathbf{G} is an involution $\theta: \mathbf{G} \rightarrow \mathbf{G}$ in the category of \mathbb{R} -algebraic groups such that the set of fixed points of θ in $\mathbf{G}(\mathbb{R})$ is a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$.

The fundamental example is the standard Cartan involution $x \mapsto (x^t)^{-1}$ on \mathbf{GL}_n , whose real fixed point set is $\mathbf{O}_n(\mathbb{R})$.

The fact that we require θ to be a morphism in the category of \mathbb{R} -algebraic groups, not just of real Lie groups, is important (unlike in the case of semisimple groups, where this makes no difference). Indeed, if we allow morphisms in the category of real reductive Lie groups, then it is not true in general that there is a unique involution for each maximal compact subgroup of $\mathbf{G}(\mathbb{R})$, as discussed in [Bor91, sec. 24.6].

In the following lemma, (ii) is the definition of Cartan involutions used in [Bor69, 11.17]. The importance of the \mathbb{R} -algebraic group structure is hidden in this definition; it lies in the fact that the maximal \mathbb{R} -split subtorus \mathbf{Z}_d of $Z(\mathbf{G})$ is well-defined only for an \mathbb{R} -algebraic group. (iii) is the definition used in [BHC62] and [RS90], while (iv) is the definition from [Del79, p. 255], commonly used in the study of Shimura varieties.

Lemma 2.1. *Let \mathbf{G} be a reductive \mathbb{R} -algebraic group and let $\theta: \mathbf{G}(\mathbb{R}) \rightarrow \mathbf{G}(\mathbb{R})$ be an involution in the category of real Lie groups. Let \mathbf{Z}_d denote the maximal \mathbb{R} -split torus in the centre of \mathbf{G} . The following are equivalent:*

- (i) θ is a Cartan involution as defined above;
- (ii) the set of fixed points of θ in $\mathbf{G}(\mathbb{R})$ is a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ and $\theta(z) = z^{-1}$ for all $z \in \mathbf{Z}_d(\mathbb{R})$;
- (iii) there exists a faithful representation $\rho: \mathbf{G} \rightarrow \mathbf{GL}_{n,\mathbb{R}}$ in the category of \mathbb{R} -algebraic groups such that

$$\rho(\theta(g)) = (\rho(g)^t)^{-1}$$

for all $g \in \mathbf{G}(\mathbb{R})$;

- (iv) θ is a morphism in the category of \mathbb{R} -algebraic groups and the real form $G^\theta = \{g \in \mathbf{G}(\mathbb{C}) : \theta(\bar{g}) = g\}$ is compact and intersects every connected component of $\mathbf{G}(\mathbb{C})$, where $\bar{}$ denotes complex conjugation.

Furthermore, for each maximal compact subgroup $K \subset \mathbf{G}(\mathbb{R})$, there exists a unique Cartan involution of \mathbf{G} whose set of real fixed points is K .

Proof. We begin by proving the existence and uniqueness of the Cartan involution having a given maximal compact subgroup as its real fixed points. In fact, we will prove existence for definition (iii) and uniqueness for definition (ii), and subsequently use these properties in proving the equivalence between the definitions.

Existence for (iii). Let $K \subset \mathbf{G}(\mathbb{R})$ be a maximal compact subgroup. Choose any faithful representation $\rho': \mathbf{G} \rightarrow \mathbf{GL}(V)$ where V is a real vector space. By [Mos55, Thm. 7.1], there exists a positive definite quadratic form on V with respect to which $\rho'(\mathbf{G})$ is self-adjoint, that is, if we choose an orthonormal basis for V with respect to this quadratic form and thereby identify $\mathbf{GL}(V)$ with $\mathbf{GL}_{n,\mathbb{R}}$, then $\rho'(\mathbf{G})$ is stable under the standard Cartan involution of \mathbf{GL}_n . Pulling back by ρ' , the standard Cartan involution induces an involution θ' of \mathbf{G} in the category of \mathbb{R} -algebraic groups. By [BHC62, Prop. 1.10], the set of fixed points of θ' in $\mathbf{G}(\mathbb{R})$ is a maximal compact subgroup $K' \subset \mathbf{G}(\mathbb{R})$. By [Mos55, Thm 3.1], there exists

$u \in \mathbf{G}(\mathbb{R})$ such that $K' = uKu^{-1}$. Letting $\rho = \rho' \circ \text{Inn}(u): \mathbf{G} \rightarrow \mathbf{GL}_n$, we obtain the desired property: the involution of \mathbf{G} obtained by pulling back the standard involution by ρ has set of real fixed points equal to K .

Uniqueness for (ii). Let $\theta: \mathbf{G}(\mathbb{R}) \rightarrow \mathbf{G}(\mathbb{R})$ be an involution in the category of real Lie groups satisfying (ii), and such that its set of real fixed points is K . Now $\mathbf{G}^{\text{der}}(\mathbb{R})^+$ is a connected semisimple Lie group, so by the classical theory of Cartan involutions of connected semisimple Lie groups, the restriction of θ to $\mathbf{G}^{\text{der}}(\mathbb{R})^+$ is uniquely determined by the fact that its set of real fixed points is $K \cap \mathbf{G}^{\text{der}}(\mathbb{R})^+$.

Meanwhile $Z_K = K \cap Z(\mathbf{G})^0(\mathbb{R})$ is the unique maximal compact subgroup of the torus $Z(\mathbf{G})^0(\mathbb{R})$. By [Bor69, §10.8], we have an almost direct product $Z(\mathbf{G})^0(\mathbb{R}) = Z_K \cdot \mathbf{Z}_d(\mathbb{R})$. The restriction of θ to $\mathbf{Z}_d(\mathbb{R})$ is determined by (ii), so the restriction of θ to $Z(\mathbf{G})^0(\mathbb{R})$ is uniquely determined by (ii) and the hypothesis that its set of real fixed points is Z_K .

Since $Z(\mathbf{G})(\mathbb{R})^+ \subset Z(\mathbf{G})^0(\mathbb{R})$ and $\mathbf{G}^{\text{der}}(\mathbb{R})^+$ generate $\mathbf{G}(\mathbb{R})^+$, we conclude that the restriction of θ to $\mathbf{G}(\mathbb{R})^+$ is uniquely determined. Finally by [Mos55, Lemma 3.1], $\mathbf{G}(\mathbb{R}) = K\mathbf{G}(\mathbb{R})^+$ so in fact θ is uniquely determined on all of $\mathbf{G}(\mathbb{R})$.

(iii) \Rightarrow (iv). The formula (iii) shows that θ is a morphism of \mathbb{R} -algebraic groups. Extending scalars to \mathbb{C} , we get

$$G^\theta = \{g \in \mathbf{G}(\mathbb{C}) : \theta(\bar{g}) = g\} = \{g \in \mathbf{G}(\mathbb{C}) : (\bar{g}^t)^{-1} = g\} = \mathbf{G}(\mathbb{C}) \cap U(n).$$

Since $U(n)$ is compact, this proves that G^θ is compact.

Furthermore, by [BHC62, Prop. 1.10], the set K of real fixed points of G^θ is a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$. Hence by [Mos55, Lemma 3.1], K intersects every connected component of $\mathbf{G}(\mathbb{R})$. Since $K \subset G^\theta$, we conclude that G^θ intersects every connected component of $\mathbf{G}(\mathbb{R})$.

(iv) \Rightarrow (i). $\text{Lie}(G^\theta)$ is the set of fixed points of the conjugate-linear involution of $\text{Lie}(\mathbf{G}(\mathbb{C}))$ defined by $X \mapsto d\theta(\bar{X})$. Consequently it satisfies

$$\text{Lie}(\mathbf{G}(\mathbb{C})) = \text{Lie}(G^\theta) \oplus i \text{Lie}(G^\theta).$$

Furthermore, G^θ is stable under complex conjugation and, by (iv), G^θ intersects every connected component of $\mathbf{G}(\mathbb{C})$. Thus G^θ is an invariant compact form of \mathbf{G} as defined in [Mos55, p. 45].

Therefore by [Mos55, Lemma 2.2], $\mathbf{G}(\mathbb{R}) \cap G^\theta$ is a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$. Since $\mathbf{G}(\mathbb{R}) \cap G^\theta = \{g \in \mathbf{G}(\mathbb{R}) : \theta(g) = g\}$, this proves (i).

(i) \Rightarrow (ii). Since θ is a morphism of algebraic groups, θ maps \mathbf{Z}_d to itself. Hence it induces $\theta_*: X_*(\mathbf{Z}_d) \rightarrow X_*(\mathbf{Z}_d)$ (where $X_*(\mathbf{Z}_d)$ denotes the space of cocharacters of \mathbf{Z}_d defined over \mathbb{R}). Since θ_* is an involution, its action on $X_*(\mathbf{Z}_d) \otimes_{\mathbb{Z}} \mathbb{Q}$ is diagonalisable with eigenvalues ± 1 . If there is any cocharacter $\alpha \in X_*(\mathbf{Z}_d)$ on which θ_* acts with eigenvalue $+1$, then θ fixes the image of α . But $\alpha(\mathbb{G}_m(\mathbb{R})) \cong \mathbb{R}^\times$ is not compact, so this contradicts (i). We conclude that θ_* acts on $X_*(\mathbf{Z}_d)$ as multiplication by -1 .

Since \mathbf{Z}_d is a split torus, $\theta_*: X_*(\mathbf{Z}_d) \rightarrow X_*(\mathbf{Z}_d)$ determines the restriction of θ to \mathbf{Z}_d . Thus $\theta|_{\mathbf{Z}_d}$ is the morphism $z \mapsto z^{-1}$.

(ii) \Rightarrow (iii). Let K be the set of fixed points of θ . We have shown that there exists an involution θ' of \mathbf{G} in the category of \mathbb{R} -algebraic groups satisfying (iii) and whose set of real fixed points is K . Since (iii) \Rightarrow (iv) \Rightarrow (i) \Rightarrow (ii), θ' satisfies (ii). We have shown that there is at most one involution of \mathbf{G} in the category of real Lie groups which satisfies (ii) and whose set of fixed points is K , so $\theta = \theta'$. \square

Given a reductive \mathbb{R} -algebraic group \mathbf{G} and a Cartan involution θ of \mathbf{G} , we say that an algebraic subgroup $\mathbf{H} \subset \mathbf{G}$ is **self-adjoint** (with respect to θ) if $\theta(\mathbf{H}) = \mathbf{H}$.

In several of our theorem statements (including Theorems 1.7 and 1.9), we are given a reductive \mathbb{Q} -algebraic group \mathbf{G} and a Siegel set $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$. It will be seen in section 2.B that the definition of a Siegel set involves the choice of a maximal compact subgroup $K \subset \mathbf{G}(\mathbb{R})$. In such a situation, we say that a subgroup of \mathbf{G} is self-adjoint if it is self-adjoint with respect to the Cartan involution whose fixed point set is the K used in the construction of the Siegel set.

2.B. Siegel sets. We use the definition of Siegel sets from [Orr18, sec. 2.2], which is a minor modification of definitions used in [Bor69, Def. 12.3] and [AMRT75, Ch. II, sec. 4.1]. For an explanation of the differences between these definitions, see [Orr18, sec. 2.3].

Let \mathbf{G} be a reductive \mathbb{Q} -algebraic group. In order to define a Siegel set in $\mathbf{G}(\mathbb{R})$, we begin by making choices of the following subgroups of \mathbf{G} :

- (1) \mathbf{P} a minimal parabolic \mathbb{Q} -subgroup of \mathbf{G} ;
- (2) K a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$.

As a consequence of the lemma in [AMRT75, chapter II, section 3.7], there is a unique \mathbb{R} -torus $\mathbf{S} \subset \mathbf{P}$ satisfying the following conditions:

- (i) \mathbf{S} is $\mathbf{P}(\mathbb{R})$ -conjugate to a maximal \mathbb{Q} -split torus in \mathbf{P} .
- (ii) \mathbf{S} is self-adjoint with respect to the Cartan involution associated with K .

These conditions could equivalently be stated as:

- (i) \mathbf{S} is a lift of the unique maximal \mathbb{Q} -split torus in $\mathbf{P}/R_u(\mathbf{P})$.
- (ii) $\text{Lie } \mathbf{S}(\mathbb{R})$ is orthogonal to $\text{Lie } K$ with respect to the Killing form of \mathbf{G} .

Define the following further pieces of notation:

- (1) \mathbf{U} is the unipotent radical of \mathbf{P} .
- (2) \mathbf{M} is the preimage in $Z_{\mathbf{G}}(\mathbf{S})$ of the maximal \mathbb{Q} -anisotropic subgroup of \mathbf{P}/\mathbf{U} . (Note that by [BT65, Corollaire 4.16], $Z_{\mathbf{G}}(\mathbf{S})$ is a Levi subgroup of \mathbf{P} and hence maps isomorphically onto \mathbf{P}/\mathbf{U} .)
- (3) Δ is the set of simple roots of \mathbf{G} with respect to \mathbf{S} , using the ordering induced by \mathbf{P} . (The roots of \mathbf{G} with respect to \mathbf{S} form a root system because \mathbf{S} is conjugate to a maximal \mathbb{Q} -split torus in \mathbf{G} .)
- (4) $A_t = \{\alpha \in \mathbf{S}(\mathbb{R})^+ : \chi(\alpha) \geq t \text{ for all } \chi \in \Delta\}$ for any real number $t > 0$.

A **Siegel set** in $\mathbf{G}(\mathbb{R})$ (with respect to $(\mathbf{P}, \mathbf{S}, K)$) is a set of the form

$$\mathfrak{S} = \Omega A_t K$$

where

- (1) Ω is a compact subset of $\mathbf{U}(\mathbb{R})\mathbf{M}(\mathbb{R})^+$; and
- (2) t is a positive real number.

We say that a set $\Omega \subset \mathbf{G}(\mathbb{R})$ is a **fundamental set** for Γ if the following conditions are satisfied:

- (F1) $\Gamma\Omega = \mathbf{G}(\mathbb{R})$; and
- (F2) for every $\theta \in \mathbf{G}(\mathbb{R})$, the set $\{\gamma \in \Gamma : \gamma\Omega \cap \theta\Omega \neq \emptyset\}$ is finite (the **Siegel property**).

The following two theorems show that, if we make suitable choices of Siegel set $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$ and finite set $C \subset \mathbf{G}(\mathbb{Q})$, then $C\mathfrak{S}$ is a fundamental set for Γ in $\mathbf{G}(\mathbb{R})$.

Theorem 2.2. [Bor69, Théorème 13.1] *Let Γ be an arithmetic subgroup of $\mathbf{G}(\mathbb{Q})$. For any minimal parabolic \mathbb{Q} -subgroup $\mathbf{P} \subset \mathbf{G}$ and maximal compact subgroup $K \subset \mathbf{G}(\mathbb{R})$, there exist a Siegel set $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$ with respect to $(\mathbf{P}, \mathbf{S}, K)$ and a finite set $C \subset \mathbf{G}(\mathbb{Q})$ such that*

$$\mathbf{G}(\mathbb{R}) = \Gamma C\mathfrak{S}.$$

Theorem 2.3. [Bor69, Théorème 15.4] *Let Γ be an arithmetic subgroup of $\mathbf{G}(\mathbb{Q})$. Let $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$ be a Siegel set. For any finite set $C \subset \mathbf{G}(\mathbb{Q})$ and any element $\theta \in \mathbf{G}(\mathbb{Q})$, the set*

$$\{\gamma \in \Gamma : \gamma C\mathfrak{S} \cap \theta C\mathfrak{S} \neq \emptyset\}$$

is finite.

A quantitative version of Theorem 2.3 can be found at [Orr18, Thm. 1.1].

2.C. Notation. If Λ is a \mathbb{Z} -module, we write $\Lambda_{\mathbb{Q}}$ for $\Lambda \otimes_{\mathbb{Z}} \mathbb{Q}$ and $\Lambda_{\mathbb{R}}$ for $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$.

We write $|\cdot|$ for a norm on an \mathbb{R} -vector space V . Unless otherwise specified, it does not matter which norm we choose, except that the values of constants will depend on the norm. Whenever the statement of a theorem involves a norm $|\cdot|$, we implicitly assume that such a norm has been chosen, and the constants in the theorem implicitly depend on this choice.

Given an \mathbb{R} -vector space with a norm $|\cdot|$, we write $\|\cdot\|$ for the associated operator norm on $\text{End}_{\mathbb{R}}(V)$. In other words, for $f \in \text{End}_{\mathbb{R}}(V)$,

$$\|f\| = \sup\{|f(v)| : v \in V, |v| = 1\}.$$

Given a group G , a representation $\rho: G \rightarrow \text{GL}(V)$ and $v \in V$, we write $\text{Stab}_{G,\rho}(v)$ for the stabiliser of v in G with respect to ρ , that is,

$$\text{Stab}_{G,\rho}(v) = \{g \in G : \rho(g)v = v\}.$$

3. QUANTITATIVE REDUCTION THEORY

In this section, we will prove Theorems 1.7 and 1.9. The proof follows the same strategy as that of [BHC62, Lemma 5.3 and Thm. 6.5]. We replace the purely topological proof of [BHC62, Prop. 5.2] by an orbit growth bound of Eberlein [Ebe14] using Riemannian geometry in $\mathbf{GL}_n(\mathbb{R})^+$. We also prove a lemma bounding the norm of $\tau \in \text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$ in terms of the length of $v = \tau(v_0)$ – this is a calculation in a semisimple \mathbb{R} -algebra. For the rest, the proof closely follows the method of Borel and Harish-Chandra, keeping track of quantitative information and the action of $\text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$ throughout and with some small adaptations to generalise to reductive groups whose \mathbb{R} -rank is greater than their \mathbb{Q} -rank.

3.A. Bound for orbits of real reductive groups. We begin by proving the following bound for orbits in representations of real reductive groups, not yet considering any arithmetic subgroup.

Proposition 3.1. *Let \mathbf{G} be a reductive \mathbb{R} -algebraic group and let $\rho: \mathbf{G} \rightarrow \mathbf{GL}(V)$ be a \mathbb{Q} -algebraic representation. Let $v_0 \in V$ be a non-zero vector whose orbit $\rho(\mathbf{G}(\mathbb{R}))v_0$ is closed. Then there exist constants C_9 and C_{10} (depending on \mathbf{G} , ρ and v_0) such that, for every $w \in \rho(\mathbf{G}(\mathbb{R}))v_0$, there exists $g \in \mathbf{G}(\mathbb{R})$ satisfying $w = \rho(g)v_0$ and*

$$\max(\|\rho(g)\|, \|\rho(g^{-1})\|) \leq C_9|w|^{C_{10}}.$$

Proposition 3.1 provides a quantitative version of [BHC62, Prop. 5.2], which asserts that if $w \in \rho(\mathbf{G}(\mathbb{R}))v_0 \cap Q$ for some compact subset $Q \subset V$, then in fact $w \in \rho(\Omega)v_0$ for some compact subset $\Omega \subset \mathbf{G}(\mathbb{R})$. Here we show that the operator norm of elements of $\rho(\Omega)$ is polynomially bounded with respect to the length of vectors in Q .

We define a Riemannian metric on $\mathbf{GL}_n(\mathbb{R})^+$ as follows. The positive definite bilinear form $(A, B) \mapsto \text{tr}(AB^t)$ on $M_n(\mathbb{R}) = T_I\mathbf{GL}_n(\mathbb{R})$ induces a right-invariant Riemannian metric on the Lie group $\mathbf{GL}_n(\mathbb{R})^+$. (Here and henceforth we denote by I the identity matrix.) Let d_R denote the distance function on $\mathbf{GL}_n(\mathbb{R})^+$ induced by this Riemannian metric.

Eberlein’s theorem relates $|\rho(g)|v_0$ to the Riemannian distance between g and the stabiliser of v_0 . We will combine this with the following lemma bounding $\|\rho(g)\|$ in terms of the Riemannian distance.

Lemma 3.2. *There exists a constant $C_{11}(n)$ such that every $g \in \mathbf{GL}_n(\mathbb{R})^+$ satisfies*

$$\|g\| \leq C_{11}(n) \exp(d_R(g, I)).$$

Proof. Let $|g|_F$ denote the Frobenius norm of g , that is,

$$|g|_F = \sqrt{\text{tr}(gg^t)}.$$

Using the Cartan decomposition, we can write $g = k \exp(X)$ for some $k \in \mathbf{SO}_n(\mathbb{R})$ and some symmetric matrix $X \in M_n(\mathbb{R})$. Let λ_{\max} denote the largest

eigenvalue of X (note that X is diagonalisable and all its eigenvalues are real because it is symmetric).

By [Ebe14, Prop. 4.8], we have

$$n^{-1/2} \exp(-c) \exp(|X|_F - \lambda_{\max}) \leq \frac{\exp(d_R(g, I))}{|g|_F} \tag{2}$$

for some constant c which depends only on n . Since X is symmetric, we have

$$|X|_2 = \sqrt{\operatorname{tr}(XX^t)} = \sqrt{\operatorname{tr}(X^2)} = \sqrt{\sum_{i=1}^n \lambda_i^2} \geq \lambda_{\max}$$

where $\lambda_1, \dots, \lambda_n$ denote the eigenvalues of X . Hence $\exp(|X|_F - \lambda_{\max}) \geq 1$, so (2) implies that

$$|g|_F \leq C_{12}(n) \exp(d_R(g, I))$$

where $C_{12}(n) = n^{-1/2} \exp(-c)$.

Since $|\cdot|_F$ and $\|\cdot\|$ are norms on the finite-dimensional vector space $M_n(\mathbb{R})$, they are equivalent, so this proves the lemma. \square

Proof of Proposition 3.1. Let $G = \mathbf{G}(\mathbb{R})^+$. Fix a finite list of representatives a_1, \dots, a_r for the connected components of $\mathbf{G}(\mathbb{R})$. Then, given $w \in \rho(\mathbf{G}(\mathbb{R}))v_0$, we can write $w = a_i w'$ for some $i \leq r$ and some $w' \in \rho(G)v_0$. Hence it suffices to prove the proposition for $w \in \rho(G)v_0$.

By [Mos55] we can choose an inner product on V with respect to which $\rho(G)$ is self-adjoint. Since all norms on a finite-dimensional vector space are equivalent, it suffices to prove the proposition under the assumption that the norm on V is induced by such an inner product. In particular, the stabiliser of the norm in G is a maximal compact subgroup K and if \mathfrak{p} denotes the -1 eigenspace of the associated Cartan involution θ on $\operatorname{Lie}(G)$, then for every $X \in \mathfrak{p}$, $d\rho(X)$ is self-adjoint. Thus the conditions of [RS90, sec. 3] are satisfied.

Hence by a theorem of Richardson and Slodowy [RS90, Thm. 4.4], $\rho(G)v_0$ is closed if and only if it contains a minimal vector, that is, a vector whose length is minimal among all elements of the orbit. Replacing v_0 by another vector in its orbit changes the g such that $w = \rho(g)v_0$ by a fixed element of G , so we may assume that v_0 itself is a minimal vector.

Let $H_0 = \operatorname{Stab}_{\rho(G)}(v_0) \subset \mathbf{GL}_n(\mathbb{R})$. Thanks to [RS90, Thm. 4.3], H_0 is self-adjoint with respect to our chosen norm on V .

If $\rho(G)v_0$ is bounded, then it is compact so by [BHC62, Prop. 5.2] there exists a compact set $\Omega \subset G$ such that $\rho(G)v_0 = \rho(\Omega)v_0$. The elements $g \in \Omega$ satisfy a uniform bound for $\max(\|\rho(g)\|, \|\rho(g^{-1})\|)$, proving the proposition in this case.

From now on, the orbit $\rho(G)v_0$ is unbounded. Then [Ebe14, sec. 2.5] defines an associated value $\lambda^-(v_0) \in \mathbb{R}$. By [Ebe14, Thm. 3.1](1), since v_0 is minimal,

$\lambda^-(v_0) > 0$. By [Ebe14, Thm. 3.1](2), we also have

$$\lambda^-(v_0) \leq \liminf_{d_R(\rho(g), H_0) \rightarrow \infty} \frac{\log|\rho(g)v_0|}{d_R(\rho(g), H_0)}.$$

Hence there exists a constant $C_{13} > 0$ (depending only on G , ρ and v_0) such that

$$\frac{\log|\rho(g)v_0|}{d_R(\rho(g), H_0)} > \frac{1}{2}\lambda^-(v_0)$$

for all $g \in G$ satisfying $d_R(\rho(g), H_0) > C_{13}$.

On the other hand, if $d_R(\rho(g), H_0) \leq C_{13}$, then because v is minimal we have

$$\frac{\log|\rho(g)v_0|}{d_R(\rho(g), H_0)} \geq \frac{\log|v_0|}{d_R(\rho(g), H_0)} \geq \frac{\log|v_0|}{C_{13}}$$

which is a positive constant.

Combining the above two inequalities, we deduce that there is a positive constant C_{14} such that the following inequality holds for all $g \in G$:

$$\frac{\log|\rho(g)v_0|}{d_R(\rho(g), H_0)} \geq C_{14}$$

or in other words,

$$|\rho(g)v_0|^{1/C_{14}} \geq d_R(\rho(g), H_0). \quad (3)$$

Given $w \in \rho(G)v_0$, write $w = \rho(g')v_0$ where $g' \in G$. Since H_0 is closed, we can choose $h \in H_0$ such that $d_R(\rho(g'), H_0) = d_R(\rho(g'), h)$.

Since $H_0 \subset \rho(G)$, we can choose $g \in G$ such that $\rho(g) = \rho(g')h^{-1}$. Since $h \in H_0$, we have $\rho(g)v_0 = w$. Since d_R is right invariant, we have

$$d_R(\rho(g'), h) = d_R(\rho(g), e) = d_R(e, \rho(g^{-1})).$$

Thus (3) (applied to g') becomes

$$|w|^{1/C_{14}} \geq d_R(\rho(g), e) = d_R(\rho(g^{-1}), e).$$

Applying Lemma 3.2 to both $\rho(g)$ and $\rho(g^{-1})$ completes the proof of the proposition. \square

Remark 3.3. The proof of Proposition 3.1 is ineffective for two reasons.

- (1) It depends on the value of $\lambda^-(v_0)$. We do not know a general method of calculating this value, although it seems to be feasible to calculate it in particular cases.
- (2) The value C_{13} depends on the speed of convergence of the limit in [Ebe14, Thm. 3.1], which is ineffective.

3.B. Quantitative reduction theory for representations. We now prove Theorem 1.7. The proof follows that of [BHC62, Lemma 5.3], keeping track of quantitative information and some minor generalisations. For the sake of clarity, we have broken it down into a series of lemmas, each proved by a short calculation.

We use the notation for Siegel sets from section 2.B. We also adopt some notation from the proof of [BHC62, Lemma 5.3] (bearing in mind that we have reversed the order of multiplication in our Iwasawa decomposition relative to [BHC62]). By the Iwasawa and Langlands decompositions, the multiplication map

$$\mathbf{U}(\mathbb{R}) \times \mathbf{S}(\mathbb{R})^+ \times \mathbf{M}(\mathbb{R})^+ K \rightarrow \mathbf{G}(\mathbb{R})$$

is bijective. Given $x \in \mathbf{G}(\mathbb{R})$, we write it as $x = n_x a_x k_x$ according to this decomposition. Let

$$y_x = a_x^{-1} x, \quad z_x = a_x^{-2} x.$$

For each character $\chi \in X^*(\mathbf{S})$, let V_χ denote the corresponding eigenspace in V . Since \mathbf{S} is \mathbb{Q} -split, the eigenspaces V_χ are defined over \mathbb{Q} and $V = \bigoplus_\chi V_\chi$. Let $\pi_\chi: V \rightarrow V_\chi$ denote the projection maps. Since all norms on the finite-dimensional vector space $V_\mathbb{R}$ are equivalent, we may assume without loss of generality that the norm is chosen so that the spaces V_χ are orthogonal to each other.

In the lemmas which follow, τ denotes an element of $\text{End}_{\rho(\mathbf{G})}(V_\mathbb{R})$. The notation C_n denotes constants which depend only on \mathbf{G} , \mathfrak{S} , ρ , Λ and v_0 , not τ , x , v or w .

Lemma 3.4. *There exists a constant $C_{15} > 0$ such that for all $v' \in V_\mathbb{R}$ and all $\chi \in X^*(\mathbf{S})$, if $\tau(v') \in \Lambda$, then either $\pi_\chi(v') = 0$ or $|\pi_\chi(v')| \geq C_{15}/\|\tau\|$.*

Proof. Since the eigenspace V_χ is defined over \mathbb{Q} , the projection $\pi_\chi(\Lambda)$ is a lattice in V_χ . Therefore there is a constant $C_{15} > 0$ such that, if $\tau(v') \in \Lambda$, then

$$\pi_\chi(\tau(v')) = 0 \text{ or } |\pi_\chi(\tau(v'))| \geq C_{15}.$$

Since $V_\chi \neq 0$ for only finitely many characters χ , it is possible to choose a single constant $C_{15} > 0$ which works for every χ .

Since τ commutes with $\rho(\mathbf{G}(\mathbb{R}))$, it preserves the eigenspaces V_χ and hence commutes with π_χ . We conclude by noting that

$$\|\tau\| |\pi_\chi(v')| \leq |\tau(\pi_\chi(v'))| = |\pi_\chi(\tau(v'))|. \quad \square$$

Lemma 3.5. *There exists a constant C_{16} such that, for all $x \in \mathfrak{S}$, we have*

$$|\rho(y_x)v_0| \leq C_{16}.$$

Proof. From the definition of a Siegel set, $\{n_x : x \in \mathfrak{S}\}$ is relatively compact. Therefore by [Bor69, Lemme 12.2], $\{a_x^{-1}n_x a_x : x \in \mathfrak{S}\}$ is relatively compact. Furthermore $\{k_x : x \in \mathfrak{S}\}$ is also relatively compact. Since

$$y_x = a_x^{-1} x = a_x^{-1} n_x a_x k_x$$

we conclude that $\{y_x : x \in \mathfrak{S}\}$ is relatively compact. □

Lemma 3.6. *There exists a constant C_{17} such that, for all $x \in \mathfrak{S}$, if $\tau(\rho(x)v_0) \in \Lambda$, then*

$$|\rho(z_x)v_0| \leq C_{17}\|\tau\|.$$

Proof. Let $\chi \in X^*(\mathbf{S})$. From the definitions of y_x and z_x , we can calculate

$$\pi_\chi(\rho(y_x)v_0) = \chi(a_x)^{-1}\pi_\chi(\rho(x)v_0), \quad \pi_\chi(\rho(z_x)v_0) = \chi(a_x)^{-2}\pi_\chi(\rho(x)v_0).$$

Therefore either $\pi_\chi(\rho(x)v_0) = 0$, in which case $\pi_\chi(\rho(z_x)v_0) = 0$, or else by Lemma 3.4 (applied to $v' = \rho(x)v_0$) and Lemma 3.5, we can calculate

$$|\pi_\chi(\rho(z_x)v_0)| = \frac{|\pi_\chi(\rho(y_x)v_0)|^2}{|\pi_\chi(\rho(x)v_0)|} \leq \frac{C_{16}^2}{C_{15}/\|\tau\|} = C_{18}\|\tau\|.$$

Since V is the orthogonal direct sum of the V_χ , the lemma follows by squaring and summing over χ . \square

Lemma 3.7. *There exist constants C_{19} and C_{20} such that, for every $x \in \mathfrak{S}$, if $\tau(\rho(x)v_0) \in \Lambda$, then there exists $g \in \mathbf{G}(\mathbb{R})$ satisfying*

$$\rho(g)v_0 = \rho(a_x^{-1}k_x)v_0 \text{ and } \max(\|\rho(g)\|, \|\rho(g)^{-1}\|) \leq C_{19}\|\tau\|^{C_{20}}.$$

Proof. By Proposition 3.1 and Lemma 3.6, there exists $g' \in \mathbf{G}(\mathbb{R})$ such that $\rho(z_x)v_0 = \rho(g')v_0$ and

$$\max(\|\rho(g')\|, \|\rho(g')^{-1}\|) \leq C_9|\rho(z_x)v_0|^{C_{10}} \leq C_9C_{17}\|\tau\|^{C_{10}}. \quad (4)$$

Let

$$g = a_x^{-2}n_x^{-1}a_x^{-2}g'.$$

Then

$$\begin{aligned} \rho(g)v_0 &= \rho(a_x^{-2}n_x^{-1}a_x^{-2})\rho(g')v_0 = \rho(a_x^{-2}n_x^{-1}a_x^{-2})\rho(z_x)v_0 \\ &= \rho(a_x^{-2}n_x^{-1}x)v_0 = \rho(a_x^{-1}k_x)v_0. \end{aligned}$$

Meanwhile by [Bor69, Lemma 12.2], $\{a_x^{-2}n_xa_x^2 : x \in \mathfrak{S}\}$ is relatively compact so (4) implies the required bound on $\max(\|\rho(g)\|, \|\rho(g)^{-1}\|)$. \square

Let θ denote the Cartan involution of \mathbf{G} whose set of real fixed points is K .

Lemma 3.8. *There exists a compact set $\Delta \subset \mathbf{G}(\mathbb{R})$ such that, for all $x \in \mathfrak{S}$, we have $\theta(a_x^{-1}k_x) \in \Delta x$.*

Proof. By definition $k_x \in \mathbf{M}(\mathbb{R})^+K$, so we can write $k_x = m_x\ell_x$ where $m_x \in \mathbf{M}(\mathbb{R})^+$ and $\ell_x \in K$. This is not a unique decomposition, but the definition of Siegel set guarantees that we can choose m_x in a fixed compact subset of $\mathbf{M}(\mathbb{R})^+$. (Recall that, by definition, \mathbf{M} commutes with \mathbf{S} .)

By definition, θ acts trivially on K and stabilises $\mathbf{S}(\mathbb{R})$. Since \mathbf{S} is an \mathbb{R} -split torus, the latter implies that $\theta(a) = a^{-1}$ for all $a \in \mathbf{S}(\mathbb{R})$. Hence

$$\theta(a_x^{-1}k_x) = \theta(m_xa_x^{-1}\ell_x) = \theta(m_x)a_x\ell_x = \theta(m_x)m_x^{-1}n_xx.$$

Since m_x and n_x lie in compact sets independent of x , this proves the lemma. \square

We are now ready to prove a version of Theorem 1.7 in which the bound is expressed in terms of the operator norm of $\tau \in \text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$, instead of the length of $v = \tau(v_0)$.

Proposition 3.9. *Let \mathbf{G} be a reductive \mathbb{Q} -algebraic group and let $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})$ be a Siegel set. Let $\rho: \mathbf{G} \rightarrow \mathbf{GL}(V)$ be a representation of \mathbf{G} defined over \mathbb{Q} . Let $\Lambda \subset V$ be a \mathbb{Z} -lattice. Let $v_0 \in V_{\mathbb{R}}$ be such that:*

- (i) $\rho(\mathbf{G}(\mathbb{R}))v_0$ is closed in $V_{\mathbb{R}}$;
- (ii) the stabiliser $\text{Stab}_{\mathbf{G}(\mathbb{R}),\rho}(v_0)$ is self-adjoint.

Then there exist constants C_{21}, C_{22} such that, for every $\tau \in \text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$ and every $w \in \rho(\mathfrak{S})\tau(v_0) \cap \Lambda$, we have $|w| \leq C_{21}\|\tau\|^{C_{22}}$.

Proof. Write $w = \rho(x)\tau(v_0) = \tau(\rho(x)v_0)$, with $x \in \mathfrak{S}$. Then we get g as in Lemma 3.7.

By [BHC62, Prop. 13.5], there is a Cartan involution θ' of $\mathbf{GL}(V_{\mathbb{R}})$ such that $\theta' \circ \rho = \rho \circ \theta$. With respect to a suitable basis of $V_{\mathbb{R}}$, θ' is given by $g \mapsto (g^{-1})^t$. The norms $\|X\|$ and $\|X^t\|$ on the finite dimensional vector space $\text{End}(V_{\mathbb{R}})$ are equivalent, so there exists a constant C_{23} such that $\|\rho(\theta(g))\| \leq C_{23}\|\rho(g^{-1})\|$.

We have $g = a_x^{-1}k_x h$ where $h \in H_0 = \text{Stab}_{\mathbf{G}(\mathbb{R}),\rho}(v_0)$. Hence by Lemma 3.8, we get

$$\theta(g) = \theta(a_x^{-1}k_x)\theta(h) \in \Delta x\theta(h)$$

where Δ is a fixed finite set. Hence we get

$$\|\rho(x\theta(h))\| \leq C_{24}\|\rho(\theta(g))\| \leq C_{25}\|\rho(g^{-1})\| \leq C_{26}\|\tau\|^{C_{27}}.$$

By hypothesis, H_0 is self-adjoint so $\theta(h) \in H_0$. Hence $\rho(x\theta(h))v_0 = \rho(x)v_0$ so $w = \tau(\rho(x\theta(h))v_0)$ and

$$|w| \leq \|\tau\| \|\rho(x\theta(h))\| |v_0|$$

which is polynomially bounded with respect to $\|\tau\|$, as required. □

To conclude, we show that it is possible to choose τ such that $\|\tau\|$ is bounded in terms of $|v|$. Theorem 1.7 follows by combining Proposition 3.9 with Lemma 3.10, applied to $E = \text{End}_{\rho(\mathbf{G})}(V_{\mathbb{R}})$.

Lemma 3.10. *Let V be a real vector space and let E be a semisimple \mathbb{R} -subalgebra of $\text{End}_{\mathbb{R}}(V)$. Let $v_0 \in V$.*

Then there exists a constant C_{28} such that, for every $v \in Ev_0$, there exists $e \in E$ satisfying $v = ev_0$ and $\|e\| \leq C_{28}|v|$.

Note that E^\times is the group of \mathbb{R} -points of a reductive \mathbb{R} -algebraic group, but even if we restrict to $v \in E^\times v_0$, this lemma does not follow from Proposition 3.1 because the orbit $E^\times v_0$ is not closed.

Proof of Lemma 3.10. Write E as a product of simple \mathbb{R} -algebras $\prod_{i=1}^m E_i$. There is a corresponding decomposition $V = \bigoplus_{i=1}^m V_i$, where the action of E_i on V factors through V_i . If $v_0 = \sum_{i=1}^m v_i \in V$ and $e = (e_1, \dots, e_m) \in E$, then $ev_0 = \sum_{i=1}^m e_i v_i$

and $\|e\| = \max\{\|e_i\| : 1 \leq i \leq m\}$. Hence it suffices to prove the lemma for each pair (E_i, V_i) .

In other words, we may assume that E is a simple \mathbb{R} -algebra. Thus $E = M_n(D)$ for some positive integer n , where D is a division algebra \mathbb{R} , \mathbb{C} or \mathbb{H} . There is a unique simple E -module, namely D^n . We can identify V (as a left E -module) with $(D^n)^r$ for some positive integer r .

Since all norms on a finite-dimensional real vector space are equivalent, we may assume that the norm on V is induced by a norm on D .

Via this identification, write

$$v_0 = (x_1, \dots, x_r), \quad v = (y_1, \dots, y_r)$$

where $x_1, \dots, x_r, y_1, \dots, y_r \in D^n$.

Reordering x_1, \dots, x_r (and the corresponding y_1, \dots, y_r), we may assume that $\{x_1, \dots, x_s\}$ forms a maximal right D -linearly independent subset of $\{x_1, \dots, x_r\}$ for a suitable positive integer $s \leq r$. Note that $s \leq n$. Then there exist $a_{ij} \in D$ ($1 \leq i \leq s < j \leq r$) such that

$$x_j = \sum_{i=1}^s x_i a_{ij} \text{ for } s+1 \leq j \leq r. \tag{5}$$

By hypothesis, there exists $e' \in E$ such that $v = e'v_0$ or in other words $y_i = e'x_i$ for all i . Consequently

$$y_j = e'x_j = \sum_{i=1}^s e'x_i a_{ij} = \sum_{i=1}^s y_i a_{ij} \text{ for } s+1 \leq j \leq r. \tag{6}$$

Since the set $\{x_1, \dots, x_s\}$ is right D -linearly independent, it can be extended to form a right D -basis of D^n . Let $h \in M_n(D)$ denote the matrix formed using such a basis as its columns. Then h is invertible and $hb_i = x_i$ for $1 \leq i \leq s$, where $\{b_1, \dots, b_n\}$ denotes the standard D -basis of D^n . Note that the choices made in constructing h can be made depending only on v_0 , not on v .

Let $f \in M_n(D)$ be the matrix which has y_1, \dots, y_s as its first s columns, and 0 for the remaining columns. Then

$$fh^{-1}x_i = fb_i = y_i \text{ for } 1 \leq i \leq s.$$

Using (5) and (6), we deduce that also

$$fh^{-1}x_j = y_j \text{ for } s+1 \leq j \leq r.$$

In other words, $e = fh^{-1} \in E$ satisfies $ev_0 = v$.

By construction, $\|f\| \leq |y_1| + \dots + |y_s| \leq s|v|$. Since h is independent of v , we conclude that $\|e\|$ is bounded by a constant multiple of $|v|$. □

3.C. Quantitative fundamental sets for arithmetic groups. The proof of Theorem 1.9 follows the proof of [BHC62, Thm. 6.5]. All we have to do is use the quantitative information from Theorem 1.7 in place of the finiteness statement [BHC62, Lemma 5.4]. There are also some minor additional technical steps due to the need to keep track of the finite set $C \subset \mathbf{G}(\mathbb{Q})$ such that $C\mathfrak{S}$ is a fundamental set in the ambient group $\mathbf{G}(\mathbb{R})$ – this was not needed in [BHC62, Thm. 6.5] because there $\mathbf{G} = \mathbf{GL}_n$ and so $C = \{1\}$.

Proof of Theorem 1.9. Suppose we are given $u \in \mathbf{G}(\mathbb{R})$ such that $\mathbf{H}_u = u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ is defined over \mathbb{Q} . By hypothesis (iii) of the theorem, there exists $v_u \in \text{End}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ such that $\rho(u)v_u \in \Lambda$. Let $v_u = \tau(v_0)$ where $\tau \in \text{End}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})$, and let $v = \rho(u)v_u$.

Thanks to [BHC62, Cor. 6.3], we may enlarge the lattice $\Lambda \subset \Lambda_{\mathbb{Q}}$ so that it is $\rho(\Gamma)$ -stable. For each $c \in C$, $c^{-1}\Lambda$ is a lattice in $\Lambda_{\mathbb{Q}}$. Hence we can choose a lattice $\Lambda' \subset \Lambda_{\mathbb{Q}}$ such that $c^{-1}\Lambda \subset \Lambda'$ for all $c \in C$.

By Theorem 1.7, every $w \in \rho(\mathfrak{S})v_u \cap \Lambda'$ has length polynomially bounded with respect to $|v_u|$. In particular, for each $c \in C$, the set

$$\rho(\mathfrak{S})v_u \cap \rho(c^{-1}\Gamma)v \subset \rho(\mathfrak{S})v_u \cap \Lambda'$$

is finite so we can choose a finite set $\{b_{c,1}, \dots, b_{c,m_c}\} \subset \Gamma$ such that

$$\rho(\mathfrak{S})v_u \cap \rho(c^{-1}\Gamma)v = \{\rho(c^{-1}b_{c,1}^{-1})v, \dots, \rho(c^{-1}b_{c,m_c}^{-1})v\}.$$

Let $B_u = \bigcup_{c \in C} \{b_{c,1}, \dots, b_{c,m_c}\}$, which is a finite subset of Γ .

By Theorem 1.7, we have

$$|\rho(c^{-1}b_{c,i}^{-1})v| \leq C_{29}|v_u|^{C_{30}}$$

for all $c \in C$ and $i \leq m_c$. Since c comes from a fixed finite set, we deduce that

$$|\rho(b_{c,i}^{-1})v| \leq C_{31}|v_u|^{C_{32}}.$$

This is the length bound on $\rho(b^{-1}u)v_u$ for $b \in B_u$ which is required by the statement of the lemma.

Let $\Gamma_u = \Gamma \cap \mathbf{H}_u(\mathbb{R})$ and $\mathcal{F}_{\mathbf{H}_u} = B_u C \mathfrak{S} u^{-1} \cap \mathbf{H}_u(\mathbb{R})$. It remains to show that $\mathcal{F}_{\mathbf{H}_u}$ is a fundamental set for Γ_u in $\mathbf{H}_u(\mathbb{R})$.

Let $h \in \mathbf{H}_u(\mathbb{R}) \subset \mathbf{G}(\mathbb{R})$. By hypothesis, $C\mathfrak{S}$ is a fundamental set for Γ in $\mathbf{G}(\mathbb{R})$ so we can write

$$hu = bcs$$

where $b \in \Gamma$, $c \in C$ and $s \in \mathfrak{S}$. Since $h \in \mathbf{H}_u(\mathbb{R}) = \text{Stab}_{\mathbf{G}(\mathbb{R}), \rho}(\rho(u)v_0)$, we obtain

$$\rho(bcs)v_0 = \rho(hu)v_0 = \rho(u)v_0.$$

Applying τ we get

$$\rho(bcs)v_u = v$$

or in other words

$$\rho(s)v_u = \rho(c^{-1}b^{-1})v \in \rho(\mathfrak{S})v_u \cap \rho(c^{-1}\Gamma)v.$$

Hence there exists $b_{c,i} \in B_u$ such that

$$\rho(c^{-1}b_{c,i}^{-1})v = \rho(s)v_u = \rho(c^{-1}b^{-1})v.$$

In particular, $bb_{c,i}^{-1} \in \text{Stab}_{\mathbf{G}(\mathbb{R}),\rho}(v)$, and we also have $bb_{c,i}^{-1} \in \Gamma$. Since $\tau \in \text{Aut}_{\rho}(\mathbf{G})(\Lambda_{\mathbb{R}})$, we have $\text{Stab}_{\mathbf{G}(\mathbb{R}),\rho}(v) = \mathbf{H}_u(\mathbb{R})$. Thus $bb_{c,i}^{-1} \in \Gamma_u$ and

$$h = bb_{c,i}^{-1}.b_{c,i}csu^{-1} \in \Gamma_u \mathcal{F}_{\mathbf{H}_u}.$$

Thus the Γ_u -translates of $\mathcal{F}_{\mathbf{H}_u}$ cover $\mathbf{H}_u(\mathbb{R})$. The fact that there are only finitely many $\gamma \in \Gamma_u$ for which $\gamma\mathcal{F}_{\mathbf{H}_u} \cap \mathcal{F}_{\mathbf{H}_u} \neq \emptyset$ follows from the Siegel property for \mathfrak{S} (and indeed this implies that $\mathcal{F}_{\mathbf{H}_u}$ also satisfies the Siegel property). Thus $\mathcal{F}_{\mathbf{H}_u}$ is a fundamental set for Γ_u in $\mathbf{H}_u(\mathbb{R})$. \square

4. QUANTITATIVE REDUCTION THEORY FOR QUATERNION ALGEBRAS

In order to apply Theorem 1.9, it is necessary to choose a representation ρ and a vector v_0 having the properties described in the theorem. In this section, we will explain how to construct a suitable representation for our application to unlikely intersections with E^2 and quaternionic curves. This illustrates a method for constructing representations which will be useful for applying Theorem 1.9 to other problems of unlikely intersections in the future, while avoiding many technical complications which occur in more general situations.

Borel and Harish-Chandra's reduction theory considered only a fixed reductive subgroup $\mathbf{H}_0 \subset \mathbf{G}$ (and not its conjugates $u\mathbf{H}_0u^{-1}$), so they needed only properties (i) and (ii) of Theorem 1.9. [BHC62, Thm. 3.8] constructs a representation satisfying properties (i) and (ii) but not (iii). Another construction of representations satisfying (i) is given by [Del82, Prop. 3.1] (based on [Che58, Exp. 10, Prop. 5]); Deligne's construction can be easily modified to satisfy (iii).

However, it is not enough to know just that there exist vectors v_u as in property (iii). Theorem 1.9 gives bounds in terms of $|v_u|$ so in order to apply these, we need to control the length $|v_u|$ in terms of some more intrinsic quantity attached to the subgroup $u\mathbf{H}_0u^{-1}$. For example, in our application the subgroups $u\mathbf{H}_0u^{-1}$ will be associated with quaternion algebras and we will bound $|v_u|$ in terms of the discriminants of these algebras.

4.A. The set-up: quaternionic subgroups of \mathbf{GSp}_4 . We consider the following situation: $\mathbf{G} = \mathbf{GSp}_4$, by which we mean the group of linear transformations of \mathbb{Q}^4 which multiply by a scalar the symplectic form $\psi: \mathbb{Q}^4 \times \mathbb{Q}^4 \rightarrow \mathbb{Q}$ represented by the matrix

$$\begin{pmatrix} J & 0 \\ 0 & J \end{pmatrix}, \text{ where } J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The subgroup \mathbf{H}_0 is equal to \mathbf{GL}_2 , embedded block diagonally in \mathbf{GSp}_4 :

$$\mathbf{H}_0 = \left\{ \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix} \in \mathbf{GSp}_4 : A \in \mathbf{GL}_2 \right\}. \quad (7)$$

If a $\mathbf{G}(\mathbb{R})$ -conjugate $u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , then its \mathbb{Q} -points form the multiplicative group of an indefinite quaternion algebra over \mathbb{Q} .

The inclusion $\mathbf{H}_0 \rightarrow \mathbf{GSp}_4$ given by (7) induces a morphism of Shimura varieties $\mathcal{A}_1 \rightarrow \mathcal{A}_2$ (where \mathcal{A}_1 is the moduli space of elliptic curves and \mathcal{A}_2 is the moduli space of principally polarised abelian varieties). In terms of moduli, this morphism of Shimura varieties sends an elliptic curve E with its unique principal polarisation λ to the principally polarised abelian surface $(E \times E, \lambda \times \lambda)$. The special subvarieties of \mathcal{A}_2 corresponding to $\mathbf{G}(\mathbb{R})$ -conjugates $u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ defined over \mathbb{Q} are images of Shimura curves parameterising principally polarised abelian surfaces with quaternionic multiplication.

The precise statement we shall prove is as follows.

Proposition 4.1. *Let $\mathbf{G} = \mathbf{GSp}_{4,\mathbb{Q}}$ and let $\mathbf{H}_0 = \mathbf{GL}_{2,\mathbb{Q}}$, embedded in \mathbf{G} as in (7). Let $\Gamma = \mathbf{Sp}_4(\mathbb{Z})$. Let $L = \mathbb{Z}^4$ and let \mathbf{G} act on $L_{\mathbb{Q}}$ in the natural way.*

There exist a \mathbb{Q} -algebraic representation $\rho: \mathbf{G} \rightarrow \mathbf{GL}(\Lambda_{\mathbb{Q}})$, where Λ is a finitely generated free \mathbb{Z} -module stabilised by Γ , a vector $v_0 \in \Lambda$ and constants $C_{33}, C_{34}, C_{35}, C_{36}$ such that:

- (i) $\text{Stab}_{\mathbf{G},\rho}(v_0) = \mathbf{H}_0$;
- (ii) the orbit $\rho(\mathbf{G}(\mathbb{R}))v_0$ is closed in $\Lambda_{\mathbb{R}}$;
- (iii) for each $u \in \mathbf{G}(\mathbb{R})$, if the group $\mathbf{H}_u = u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , then
 - (a) there exists $v_u \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ such that $\rho(u)v_u \in \Lambda$ and

$$|v_u| \leq C_{33}|\text{disc}(R_u)|^{C_{34}};$$

- (b) there exists $\gamma \in \Gamma$ and $h \in \mathbf{H}_0(\mathbb{R})$ such that

$$\|\gamma uh\| \leq C_{35}|\text{disc}(R_u)|^{C_{36}},$$

where R_u denotes the ring $\text{End}_{\mathbf{H}_u}(L) \subset M_4(\mathbb{Z})$.

The condition $v_u \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ in Proposition 4.1(iii)(a) ensures that the element $\rho(u)v_u \in \Lambda$ satisfies $\text{Stab}_{\mathbf{G},\rho}(\rho(u)v_u) = \mathbf{H}_u$. Proposition 4.1(iii)(a) is the bound we need to apply Theorem 1.9. Proposition 4.1(iii)(b) is not required for our application to unlikely intersections, but may be useful in its own right – we can replace u by γuh if we replace \mathbf{H}_u by $\gamma\mathbf{H}_u\gamma^{-1}$, a subgroup of \mathbf{G} which gives rise to the same special subvariety of \mathcal{A}_2 as \mathbf{H}_u .

The proof of Proposition 4.1 will proceed in three steps: first we construct ρ and v_0 satisfying property (i), then we show that the representation we have constructed possesses property (ii) and finally we prove (iii). The proofs of properties (ii) and (iii) are independent of each other, while (iii)(b) is a by-product of the proof of (iii)(a).

4.B. Construction of representation of \mathbf{GSp}_4 . We construct the representation of \mathbf{GSp}_4 and the vector v_0 which we shall use, and define notation which we shall use throughout the rest of the section.

Let $W = M_4(\mathbb{Q})$, considered as a \mathbb{Q} -vector space. Define two representations $\sigma_L, \sigma_R: \mathbf{G} = \mathbf{GSp}_4 \rightarrow \mathbf{GL}(W)$ by multiplication on the left and on the right:

$$\sigma_L(g)w = gw, \quad \sigma_R(g)w = wg^{-1}.$$

(The inverse in the formula for σ_R is so that σ_R is a left representation of \mathbf{G} .)

Let $E_0 = M_2(\mathbb{Q})$ and define $\iota_0: E_0 \rightarrow M_4(\mathbb{Q})$ by

$$\iota_0(A) = \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}.$$

Thus $\mathbf{H}_0 = \iota_0(\mathbf{GL}_2)$. Let $Z = \iota_0(E_0)$, a 4-dimensional \mathbb{Q} -linear subspace of W . Observe that

$$\text{Stab}_{\mathbf{G}, \sigma_L}(Z) = \iota_0(E_0) \cap \mathbf{G} = \mathbf{H}_0.$$

Similarly, $\text{Stab}_{\mathbf{G}, \sigma_R}(Z) = \mathbf{H}_0$ but we shall not need this latter fact.

Let $V = \bigwedge^4 W$ and let $\rho_L, \rho_R: \mathbf{G} \rightarrow \mathbf{GL}(V)$ be the representations

$$\rho_L = \bigwedge^4 \sigma_L \otimes \det^{-1}, \quad \rho_R = \bigwedge^4 \sigma_R \otimes \det.$$

Then $\bigwedge^4 Z$ is a 1-dimensional \mathbb{Q} -linear subspace of V , with

$$\text{Stab}_{\mathbf{G}, \rho_L}(\bigwedge^4 Z) = \text{Stab}_{\mathbf{G}, \sigma_L}(Z) = \mathbf{H}_0.$$

The action of $\mathbf{GL}_2(\mathbb{Q})$ on Z via $\sigma_L \circ \iota_0$ is the restriction of the left regular representation of $M_2(\mathbb{Q})$. Hence the action of $\mathbf{GL}_2(\mathbb{Q})$ on $\bigwedge^4 Z$ via $\bigwedge^4 \sigma_L \circ \iota_0$ is multiplication by $(\det_{\mathbf{GL}_2})^2 = \det_{\mathbf{GL}_4} \circ \iota_0$. Therefore the action of \mathbf{H}_0 on $\bigwedge^4 Z$ via ρ_L is trivial, so each non-zero vector in $\bigwedge^4 Z$ has stabiliser equal to \mathbf{H}_0 .

Let $\Lambda = \bigwedge^4 M_4(\mathbb{Z}) \subset V$. For later use, we choose a specific element $v_0 \in \bigwedge^4 Z \cap \Lambda$. Let e_1, e_2, e_3, e_4 denote the following \mathbb{Z} -basis for $M_2(\mathbb{Z})$:

$$e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad e_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad e_4 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (8)$$

Then $\iota(e_1), \iota(e_2), \iota(e_3), \iota(e_4)$ form a \mathbb{Z} -basis for $Z \cap M_4(\mathbb{Z})$, so

$$v_0 = \iota_0(e_1) \wedge \iota_0(e_2) \wedge \iota_0(e_3) \wedge \iota_0(e_4)$$

is a generator of the rank-1 \mathbb{Z} -module $\bigwedge^4 Z \cap \Lambda$. Then ρ_L and v_0 satisfy Proposition 4.1(i).

Given $u \in \mathbf{G}(\mathbb{R})$, we can easily find a vector $v_u \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ such that $\rho(u)v_u \in \Lambda$ (that is, the first part of Proposition 4.1(iii)(a)). The vector

$$\rho_L(u)\rho_R(u)v_0 = \iota_0(e_1)u^{-1} \wedge \iota_0(e_2)u^{-1} \wedge \iota_0(e_3)u^{-1} \wedge \iota_0(e_4)u^{-1} \in V_{\mathbb{R}}$$

generates the line $\bigwedge^4 uZ_{\mathbb{R}}u^{-1} \subset V_{\mathbb{R}}$. If the subgroup $u\mathbf{H}_{0,\mathbb{R}}u^{-1} \subset \mathbf{G}_{\mathbb{R}}$ is defined over \mathbb{Q} , then so is the linear subspace $uZ_{\mathbb{R}}u^{-1} \subset W_{\mathbb{R}}$. Consequently $\bigwedge^4 uZ_{\mathbb{R}}u^{-1} \cap \Lambda$ is non-empty, so there exists $d_u \in \mathbb{R}^{\times}$ such that

$$d_u \rho_L(u)\rho_R(u)v_0 \in \Lambda.$$

Now $d_u \rho_R(u) \in \text{Aut}_{\rho_L}(V_{\mathbb{R}})$ so $d_u \rho_R(u)v_0$ has the required property. This algebraic construction does not control the size of d_u , and hence does not control $|v_u|$. Later, in section 4.E, we will choose a slightly different v_u (making use of γ as in Proposition 4.1(iii)(b)) allowing us to bound $|v_u|$.

To place this representation in a more general context, we compare it with [Che58, Exp. 10, Prop. 5]. Let \mathbf{G} be an arbitrary affine \mathbb{Q} -algebraic group and $\mathbf{H}_0 \subset \mathbf{G}$ an algebraic subgroup. Chevalley considers the ring of regular functions $\mathbb{Q}[\mathbf{G}]$, on which \mathbf{G} acts by right translations. The stabiliser of the ideal $I(\mathbf{H}_0)$ is equal to \mathbf{H}_0 . Choose a finite-dimensional subrepresentation $W \subset \mathbb{Q}[\mathbf{G}]$ which contains a generating set for $I(\mathbf{H}_0)$. Then \mathbf{H}_0 is also the stabiliser of $Z = W \cap I(\mathbf{H}_0)$. Let $d = \dim_{\mathbb{Q}}(Z)$. Then $\wedge^d W$ is a representation of \mathbf{G} in which the line $\wedge^d Z$ is defined over \mathbb{Q} and has stabiliser equal to \mathbf{H}_0 . If \mathbf{H}_0 is semisimple, it has no non-trivial characters so each non-zero vector in $\wedge^d Z$ also has stabiliser equal to \mathbf{H}_0 . [Del82, Prop. 3.1] describes how this construction can be modified to obtain a vector (not just a line) with stabiliser equal to \mathbf{H}_0 whenever \mathbf{H}_0 is reductive.

If we choose W to be stable under left as well as right translations (denoting the representations by ρ_L and ρ_R respectively), then the same argument as in the special case above shows that the line $\mathbb{R}^\times \rho_L(u) \rho_R(u)v_0$ is defined over \mathbb{Q} whenever $u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , and so this line contains non-zero rational vectors.

Comparing this general construction with our special case of $\mathbf{G} = \mathbf{GSp}_4$, $\mathbf{H}_0 = \iota_0(\mathbf{GL}_2)$, we note that in the special case $I(\mathbf{H}_0)$ is generated by linear functions on M_4 . Thus following Chevalley’s method, we could choose W to be the linear dual of $M_4(\mathbb{Q})$. In fact, we chose W to be $M_4(\mathbb{Q})$ itself, and Z to be the linear subspace of $M_4(\mathbb{Q})$ which is annihilated by $I(\mathbf{H}_0) \cap M_4(\mathbb{Q})^\vee$. The choice of $M_4(\mathbb{Q})$ instead of its dual is a matter of convenience.

The representations constructed by Chevalley’s method do not necessarily have a closed orbit $\rho_L(\mathbf{G}(\mathbb{R}))v_0$, although this can often be achieved by carefully choosing $W \subset \mathbb{Q}[\mathbf{G}]$ and perhaps making some minor modifications using linear algebra constructions. On the other hand, finding a suitable v_u with bounded length requires much more detailed arithmetic information about the groups \mathbf{H}_u , and we carry this out only for our special case.

4.C. Closed orbit. We now show that Proposition 4.1(ii) holds, that is, the orbit $\rho_L(\mathbf{G}(\mathbb{R}))v_0$ is closed in $V_{\mathbb{R}}$. Note that by [BHC62, Prop. 2.3], it suffices to prove that $\rho_L(\mathbf{G}(\mathbb{C}))v_0$ is closed in $V_{\mathbb{C}}$.

We use the following definitions. If V is a vector space over \mathbb{C} , we say that a subset of V is **homogeneous** if it is non-empty and stable under multiplication by scalars. In other words, a subset of V is homogeneous if and only if it is the cone over some subset of $\mathbb{P}(V)$. For a non-negative integer t , a set-theoretic function between vector spaces $f: V' \rightarrow V$ is **homogeneous of degree d** if

$$f(\lambda v) = \lambda^d f(v) \text{ for all } \lambda \in \mathbb{C}, v \in V'.$$

Homogeneous sets and homogeneous maps are useful because of the following lemma, which is equivalent to the fact that a morphism of algebraic varieties between projective spaces maps Zariski closed sets to Zariski closed sets.

Lemma 4.2. *Let V, V' be vector spaces over \mathbb{C} (or any algebraically closed field), let $X \subset V'$ be a homogeneous Zariski closed subset and let $f: V' \rightarrow V$ be a morphism of algebraic varieties which is homogeneous of positive degree. Then $f(X)$ is a homogeneous Zariski closed subset of V .*

We begin the proof of Proposition 4.1(ii) by showing that the smallest homogeneous set containing the orbit $\rho_L(\mathbf{G}(\mathbb{C}))v_0$ is the image of a homogeneous Zariski closed set under a homogeneous morphism of varieties, and so is Zariski closed.

Lemma 4.3. $\mathbb{C}\rho_L(\mathbf{G}(\mathbb{C}))v_0$ is a Zariski closed subset of $V_{\mathbb{C}}$.

Proof. Let $U = \mathbb{C}^4$. Define two linear maps $\beta_1, \beta_2: U^2 \rightarrow W_{\mathbb{C}} = M_4(\mathbb{C})$ by

$$\begin{aligned}\beta_1(u_1, u_2) &= \begin{pmatrix} u_1 & 0 & u_2 & 0 \end{pmatrix}, \\ \beta_2(u_1, u_2) &= \begin{pmatrix} 0 & u_1 & 0 & u_2 \end{pmatrix}.\end{aligned}$$

This notation means that $\beta(u_1, u_2)$ is the 4×4 matrix with columns $u_1, 0, u_2, 0$, and similarly for β_2 .

If z_1, z_2, z_3, z_4 denote the standard basis of U , then

$$\iota_0(e_1) = \beta_1(z_1, z_3), \iota_0(e_2) = \beta_2(z_1, z_3), \iota_0(e_3) = \beta_1(z_2, z_4), \iota_0(e_4) = \beta_2(z_2, z_4). \quad (9)$$

The maps β_1 and β_2 commute with the action of \mathbf{G} by left multiplication in the sense that

$$g\beta_i(u_1, u_2) = \beta_i(gu_1, gu_2)$$

for all $u_1, u_2 \in U$ and $g \in \mathbf{G}(\mathbb{C})$. Consequently

$$\begin{aligned}\mathbb{C}\rho_L(\mathbf{G}(\mathbb{C}))v_0 &= \{\lambda \det(g)^{-1} g\iota_0(e_1) \wedge g\iota_0(e_2) \wedge g\iota_0(e_3) \wedge g\iota_0(e_4) : \\ &\quad \lambda \in \mathbb{C}, g \in \mathbf{GSp}_4(\mathbb{C})\} \\ &= \{\lambda \det(g)^{-1} \beta_1(gz_1, gz_3) \wedge \beta_2(gz_1, gz_3) \wedge \beta_1(gz_2, gz_4) \wedge \beta_2(gz_2, gz_4) : \\ &\quad \lambda \in \mathbb{C}, g \in \mathbf{GSp}_4(\mathbb{C})\} \\ &= \{\beta_1(u_1, u_3) \wedge \beta_2(u_1, u_3) \wedge \beta_1(u_2, u_4) \wedge \beta_2(u_2, u_4) : \\ &\quad u_1, u_2, u_3, u_4 \text{ form a scalar multiple of a symplectic basis for } (U, \psi)\}.\end{aligned}$$

Thus if we let $V' = U^4$ and

$$X = \{(u_1, u_2, u_3, u_4) \in V' : \text{scalar multiple of a symplectic basis for } (U, \psi)\},$$

and define $f: V' \rightarrow V_{\mathbb{C}}$ by

$$f(u_1, u_2, u_3, u_4) = \beta_1(u_1, u_3) \wedge \beta_2(u_1, u_3) \wedge \beta_1(u_2, u_4) \wedge \beta_2(u_2, u_4).$$

then $\mathbb{C}\rho_L(\mathbf{G}(\mathbb{C}))v_0 = f(X)$.

Now X is a homogeneous Zariski closed subset of V' because it is defined by the equations

$$\begin{aligned} \psi(u_1, u_3) = \psi(u_1, u_4) = \psi(u_2, u_3) = \psi(u_2, u_4) = 0, \\ \psi(u_1, u_2) = \psi(u_3, u_4) \end{aligned}$$

and f is a homogeneous morphism of algebraic varieties. Consequently $f(X)$ is Zariski closed. \square

We complete the proof that $\rho_L(\mathbf{G}(\mathbb{C}))v_0$ is closed in $V_{\mathbb{C}}$ by showing that it is a hyperplane section of $\mathbb{C}\rho_L(\mathbf{G}(\mathbb{C}))v_0$.

Lemma 4.4. *There exists a linear map $s: V_{\mathbb{C}} \rightarrow \mathbb{C}$ such that*

$$\rho_L(\mathbf{G}(\mathbb{C}))v_0 = \mathbb{C}\rho_L(\mathbf{G}(\mathbb{C}))v_0 \cap s^{-1}(1).$$

Proof. We continue to use the notation U, β_1, β_2 from the proof of Lemma 4.3.

Let $\delta: W_{\mathbb{C}} \rightarrow U$ be the linear map which sends the matrix with columns $(C_1 \ C_2 \ C_3 \ C_4)$ to the sum $C_1 + C_4$. This map is \mathbf{G} -equivariant and the compositions $\delta \circ \beta_1, \delta \circ \beta_2: U^2 \rightarrow U$ are the projections onto the two copies of U .

Taking the fourth exterior power, δ induces a linear map $s: V_{\mathbb{C}} = \Lambda^4 W_{\mathbb{C}} \rightarrow \Lambda^4 U \cong \mathbb{C}$. By (9) and the descriptions of $\delta \circ \beta_1, \delta \circ \beta_2$, we have

$$s(v_0) = \delta\beta_1(z_1, z_3) \wedge \delta\beta_2(z_1, z_3) \wedge \delta\beta_1(z_2, z_4) \wedge \delta\beta_2(z_2, z_4) = z_1 \wedge z_2 \wedge z_3 \wedge z_4.$$

Since z_1, z_2, z_3, z_4 form a basis for U , $s(v_0) \neq 0$. Hence we can choose the isomorphism $\Lambda^4 U \cong \mathbb{C}$ so that $s(v_0) = 1$.

The linear map δ is \mathbf{G} -equivariant with respect to left multiplication. Consequently s is \mathbf{G} -equivariant with respect to $\Lambda^4 \sigma_L$ on $V_{\mathbb{C}}$ and multiplication by the determinant on $\Lambda^4 U$. Twisting by \det^{-1} , we deduce that s is \mathbf{G} -equivariant with respect to ρ_L on $V_{\mathbb{C}}$ and the trivial action on $\Lambda^4 U$. Thus

$$s(\rho_L(g)v_0) = s(v_0) = 1 \text{ for all } g \in \mathbf{G}(\mathbb{C}).$$

Conversely, if $v \in \mathbb{C}\rho_L(\mathbf{G}(\mathbb{C}))v_0 \cap s^{-1}(1)$, then we can write $v = \lambda\rho_L(g)v_0$ for some $\lambda \in \mathbb{C}$ and $g \in \mathbf{G}(\mathbb{C})$. Since s is a linear map, we get $s(v) = \lambda$ so the fact that $s(v) = 1$ forces $v \in \rho_L(\mathbf{G}(\mathbb{C}))v_0$. \square

4.D. Discriminants and orders. Before the proof of Proposition 4.1(iii), we prove some results on discriminants, involutions and orders in semisimple algebras.

Let D be a semisimple \mathbb{Q} -algebra. We define a symmetric \mathbb{Q} -bilinear form $\phi: D \times D \rightarrow \mathbb{Q}$ by

$$\phi(x, y) = \text{Tr}_{D/\mathbb{Q}}(xy),$$

where $\text{Tr}_{D/\mathbb{Q}}$ is the (non-reduced) trace of the regular representation of D . This form is non-degenerate by [KMRT98, Ch. I, Prop. (1.8)] (note that [KMRT98] uses reduced trace, but $\text{Tr}_{D/\mathbb{Q}}$ is a non-zero multiple of reduced trace on each simple factor so this is fine).

Let R be an order in D . We define the **discriminant** of R , denoted $\text{disc}(R)$, to be the discriminant of the bilinear form ϕ on R . In other words, $\text{disc}(R)$ is the determinant of the matrix $(\phi(e_i e_j))_{i,j}$ where $\{e_1, \dots, e_n\}$ is a \mathbb{Z} -basis for R (this determinant is independent of the choice of basis).

For any involution \dagger of D , we define another \mathbb{Q} -bilinear form $\phi^\dagger: D \times D \rightarrow \mathbb{Q}$ by

$$\phi^\dagger(x, y) = \text{Tr}_{D/\mathbb{Q}}(xy^\dagger). \tag{10}$$

This form is symmetric by [KMRT98, Ch. I, Cor. (2.2) and Cor. (2.16)].

Lemma 4.5. *Let R be an order in a semisimple \mathbb{Q} -algebra D and let \dagger be any involution of D . Then $\text{disc}(R, \phi^\dagger) = \pm \text{disc}(R)$.*

Proof. This is based on the proof of [GR14, Prop. 2.9].

Let $\{d_1, \dots, d_n\}$ be a \mathbb{Z} -basis for R . Let $A \in \mathbf{GL}_n(\mathbb{Q})$ be the matrix such that $d_j^\dagger = \sum_{i=1}^n A_{ij} d_i$. Since \dagger is \mathbb{Q} -linear and an involution, we also have $d_j = \sum_{i=1}^n A_{ij} d_i^\dagger$ and so $A^2 = \mathbb{1}$. In particular $\det(A) = \pm 1$.

Now $\phi^\dagger(e_i, e_j) = \sum_{k=1}^n A_{kj} \phi(e_i, e_k)$ so

$$\text{disc}(R, \phi^\dagger) = \det(A) \text{disc}(R) = \pm \text{disc}(R). \quad \square$$

The following lemma is restricted to quaternion algebras because its proof makes use of the fact that the reduced norm is a quadratic form on a quaternion algebra.

Lemma 4.6. *There exists an absolute constant C_{37} with the following property.*

Let R be an order in a quaternion algebra D over \mathbb{Q} . Let L be a left R -module such that $L_{\mathbb{Q}}$ is isomorphic to the left regular representation of D .

Then there exists a left R -ideal $I \subset R$ such that I is isomorphic to L as a left R -module and

$$[R : I] \leq C_{37} |\text{disc}(R)|^{3/2}.$$

Proof. This is a generalisation of Minkowski's bound for ideal classes in a number field to quaternion algebras, and the proof is similar.

Choose an isomorphism of D -modules $\eta_1: D \rightarrow L_{\mathbb{Q}}$ and let $I_1 = \eta_1^{-1}(L)$.

Since D is a quaternion algebra, it possesses a canonical involution $*$ defined by $d^* = \text{Tr}_{D/\mathbb{Q}}(d) - d$. The canonical involution has the property that

$$\phi^*(d, d) = \text{Tr}_{D/\mathbb{Q}}(dd^*) = 4 \text{Nrd}_{D/\mathbb{Q}}(d) = \pm 4 \text{Nm}_{D/\mathbb{Q}}(d)^{1/2}. \tag{11}$$

By [Bla48, Lemma 1], there is an element $s \in I_1$ satisfying

$$0 < |\phi^*(s, s)| \leq C_{38} |\text{disc}(I_1, \phi^*)|^{1/4}$$

(the exponent is $1/4$ because I_1 has rank 4 as a \mathbb{Z} -module). Hence by (11), there is a constant C_{39} such that

$$0 < |\text{Nm}_{D/\mathbb{Q}}(s)| \leq C_{39} |\text{disc}(I_1, \phi^*)|^{1/2}. \tag{12}$$

Since $\text{Nm}_{D/\mathbb{Q}}(s) \neq 0$, s is invertible in D . Let $I_2 = I_1 s^{-1} \subset E$. Then I_2 is a left R -module isomorphic to L . Since $1 = s s^{-1} \in I_2$, we have $R \subset I_2$.

Using (12), we can calculate

$$|\operatorname{disc}(I_2, \phi^*)| = |\operatorname{Nm}_{D/\mathbb{Q}}(s^{-1})|^2 |\operatorname{disc}(I_1, \phi^*)| \geq C_{39}^{-2}.$$

Consequently, using Lemma 4.5,

$$[I_2 : R]^2 = \frac{|\operatorname{disc}(R, \phi^*)|}{|\operatorname{disc}(I_2, \phi^*)|} \leq C_{39}^2 |\operatorname{disc}(R)|.$$

Finally let $I = [I_2 : R]I_2$. This is contained in R and is a left R -submodule of D , so it is a left R -ideal. It satisfies

$$[R : I] = \frac{[I_2 : I]}{[I_2 : R]} = \frac{[I_2 : R]^4}{[I_2 : R]} \leq C_{39}^3 |\operatorname{disc}(R)|^{3/2}. \quad \square$$

Lemma 4.7. *There exists an absolute constant C_{40} with the following property.*

Let R be an order in a quaternion algebra D over \mathbb{Q} . Let L be a left D -module such that $L_{\mathbb{Q}}$ is isomorphic to the left regular representation of D . Let $S = \operatorname{End}(L)^D$. Then

$$|\operatorname{disc}(S)| \leq C_{40} |\operatorname{disc}(R)|^4.$$

Proof. By Lemma 4.6, L is isomorphic to a left R -ideal $I \subset R$ which satisfies

$$[R : I] \leq C_{37} |\operatorname{disc}(R)|^{3/2}.$$

Then $S = \operatorname{End}(L)^D = \operatorname{End}(I)^D \subset \operatorname{End}(D)^D$, where the latter denotes endomorphisms of D as a left D -module.

We can define a multiplication-reversing function $\mu: D \rightarrow \operatorname{End}(D)^D$ by

$$\mu(d)x = xd \text{ for all } d, x \in D.$$

This is a \mathbb{Q} -algebra isomorphism $D^{\operatorname{op}} \rightarrow \operatorname{End}(D)^D$ [Sch85, Ch. 8, Lemma 1.10].

If $r \in R$ and $x \in I$, then we have $\mu(r)x = xr \in R$ since $I \subset R$ and R is closed under multiplication. Hence

$$[R : I]\mu(r)x \in I.$$

Thus $[R : I]\mu(r) \in \operatorname{End}(I)^D$ for all $r \in R$.

Let ϕ_S denote the trace form on $\operatorname{End}(D)^D = S_{\mathbb{Q}}$. Since μ is an algebra isomorphism, it pulls back ϕ_S to the trace form on D^{op} , which is equal to the trace form on D . Hence $\operatorname{disc}(\mu(R), \phi_S) = \operatorname{disc}(R)$.

Since $[R : I]\mu(R) \subset S$, we conclude that

$$|\operatorname{disc}(S)| \leq |\operatorname{disc}([R : I]\mu(R), \phi_S)| = [R : I]^2 |\operatorname{disc}(R)| \leq C_{37}^2 |\operatorname{disc}(R)|^4. \quad \square$$

4.E. Choice of v_u, γ and h . Throughout this section, C_n will denote absolute constants (in particular, independent of u).

We will now prove Proposition 4.1(iii). Thus, we are given $u \in \mathbf{G}(\mathbb{R}) = \mathbf{GSp}_4(\mathbb{R})$ such that the algebraic group $\mathbf{H}_u = u\mathbf{H}_{0,\mathbb{R}}u^{-1} \subset \mathbf{G}_{\mathbb{R}}$ is defined over \mathbb{Q} . Multiplying u by a scalar does not change \mathbf{H}_u , so we may assume that u multiplies the symplectic form ψ by ± 1 ; consequently $\det(u) = 1$.

Since \mathbf{H}_u is defined over \mathbb{Q} , the \mathbb{R} -vector space $u\iota_0(E_{0,\mathbb{R}})u^{-1}$ is also defined over \mathbb{Q} . Hence the \mathbb{Q} -algebra

$$E = M_4(\mathbb{Q}) \cap u\iota_0(E_{0,\mathbb{R}})u^{-1}$$

satisfies $E_{\mathbb{R}} = u\iota_0(E_{0,\mathbb{R}})u^{-1}$.

Let D_0 and D denote the centralisers of $\iota_0(E_0)$ and E respectively in $M_4(\mathbb{Q})$. Then $D_{\mathbb{R}} = uD_{0,\mathbb{R}}u^{-1}$, so there is an isomorphism of \mathbb{R} -algebras $\alpha: D_{\mathbb{R}} \rightarrow D_{0,\mathbb{R}}$ defined by

$$\alpha(d) = u^{-1}du.$$

Note that

$$D_0 = \left\{ \begin{pmatrix} aI & bI \\ cI & dI \end{pmatrix} \in M_4(\mathbb{Q}) : a, b, c, d \in \mathbb{Q} \right\}.$$

Hence the ‘‘transpose’’ involution of $M_4(\mathbb{Q})$ restricts to an involution of D_0 , which we denote by t . This involution is positive. Let

$$\dagger = \alpha^{-1} \circ t \circ \alpha: D_{\mathbb{R}} \rightarrow D_{\mathbb{R}}.$$

Let $L = \mathbb{Z}^4$. Let $R = \text{End}(L)^E = \text{Stab}_D(L)$, which is an order in D .

Lemma 4.8. *The quadratic form ϕ^\dagger takes integer values on R .*

Proof. Thanks to our choice of symplectic form ψ on \mathbb{Q}^4 , we have $\psi(d_0x, y) = \psi(x, d_0^t y)$ for all $x, y \in \mathbb{R}^4$ and $d_0 \in D_{0,\mathbb{R}}$. Using this, the fact that u multiplies ψ by ± 1 , and the definition of \dagger , we can calculate

$$\begin{aligned} \psi(dx, y) &= \pm \psi(u\alpha(d)u^{-1}x, y) = \pm \psi(\alpha(d)u^{-1}x, u^{-1}y) \\ &= \pm \psi(u^{-1}x, \alpha(d)^t u^{-1}y) = \pm \psi(u^{-1}x, \alpha(d^\dagger)u^{-1}y) \\ &= \pm \psi(x, u\alpha(d^\dagger)u^{-1}y) = \pm \psi(x, d^\dagger y). \end{aligned}$$

Since ψ is a perfect pairing on L , this implies that \dagger maps R into R . It follows that $\phi^\dagger(x, y) = \text{Tr}_{D/\mathbb{Q}}(xy^\dagger) \in \mathbb{Z}$ for all $x, y \in R$. \square

By Lemma 4.6, there is a left R -ideal $I \subset R$ which is isomorphic to L as a left R -module, such that

$$[R : I] \leq C_{37} |\text{disc}(R)|^{3/2}.$$

Choose a left R -module isomorphism $\eta: I \rightarrow L$.

Fix an isomorphism $\eta_0: D_0 \rightarrow \mathbb{Q}^4$ of left D_0 -modules (independent of u).

Lemma 4.9. *There exists $h \in \mathbf{H}_0(\mathbb{R})$ such that $\eta\alpha^{-1}\eta_0^{-1} = uh$ in $\text{Aut}(\mathbb{R}^4)$.*

Proof. We can view $D_{\mathbb{R}}$ as a left $D_{0,\mathbb{R}}$ -module with the action given by

$$d.x = \alpha^{-1}(d)x. \tag{13}$$

Now $\alpha^{-1}\eta_0^{-1}: \mathbb{R}^4 \rightarrow D_{\mathbb{R}}$ is an isomorphism of left $D_{0,\mathbb{R}}$ -modules with respect to the natural action on \mathbb{R}^4 and the action (13) on $D_{\mathbb{R}}$.

Since $\eta: D_{\mathbb{R}} \rightarrow \mathbb{R}^4$ is an isomorphism of $D_{\mathbb{R}}$ -modules, it is also an isomorphism of $D_{0,\mathbb{R}}$ -modules with respect to the action (13) on $D_{\mathbb{R}}$ and the natural action conjugated by u on \mathbb{R}^4 . (We use here the fact that α^{-1} is conjugation by u .)

Finally $u^{-1}: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ is an isomorphism of $D_{0,\mathbb{R}}$ -modules with respect to the natural action conjugated by u on the domain and the natural action on the target.

Composing these, we deduce that $u^{-1}\eta\alpha^{-1}\eta_0^{-1}$ is an automorphism of \mathbb{R}^4 with its natural action of $D_{0,\mathbb{R}}$. In other words, $u^{-1}\eta\alpha^{-1}\eta_0^{-1}$ lies in the centraliser of $D_{0,\mathbb{R}}$ in $M_4(\mathbb{R})$. By the double centraliser theorem, this centraliser is equal to $\iota_0(E_{0,\mathbb{R}})$ and so its group of invertible elements is equal to $\iota_0(\mathbf{GL}_2(\mathbb{R})) = \mathbf{H}_0(\mathbb{R})$. \square

Lemma 4.10. *The absolute value of $\det(h)$ is uniformly bounded.*

Proof. Let ϕ_0^\dagger denote the bilinear form $\phi_0^\dagger(x, y) = \text{Tr}_{D_0/\mathbb{Q}}(xy^t)$ on D_0 , and let ϕ_L denote the bilinear form on \mathbb{Q}^4 given by $\phi_L(x, y) = \phi_0^\dagger(\eta^{-1}(x), \eta^{-1}(y))$.

Since α is an isomorphism of \mathbb{R} -algebras, it preserves traces, so

$$\phi^\dagger(x, y) = \phi_0^\dagger(\alpha(x), \alpha(y)) = \phi_L(\eta_0\alpha(x), \eta_0\alpha(y)). \quad (14)$$

Consequently

$$\text{disc}(\phi^\dagger, I) = \text{disc}(\phi^\dagger, \eta^{-1}(L)) = \text{disc}(\phi_L, \eta_0\alpha\eta^{-1}(L)) = \det(\eta_0\alpha\eta^{-1})^2 \text{disc}(\phi_L, L).$$

Thanks to Lemma 4.9 and noting that $\det(u) = 1$, this can be rewritten as

$$\det(h)^2 = \det(uh)^2 = \text{disc}(\phi_L, L) / \text{disc}(\phi^\dagger, I).$$

By Lemma 4.8, ϕ^\dagger takes integer values on R and hence on I . So $\text{disc}(\phi^\dagger, I)$ is a positive integer. So $\det(h)^2 \leq \text{disc}(\phi_L, L)$, which is a constant. \square

Lemma 4.11. *There exists a \mathbb{Z} -basis $\{d'_1, d'_2, d'_3, d'_4\}$ for I such that the coordinates of the vectors $\eta_0\alpha(d'_1), \eta_0\alpha(d'_2), \eta_0\alpha(d'_3), \eta_0\alpha(d'_4) \in \mathbb{R}^4$ are polynomially bounded in terms of $|\text{disc}(R)|$.*

Proof. By Lemma 4.8, ϕ^\dagger takes integer values on I . Furthermore \dagger is a positive involution (because t is a positive involution on D_0) so ϕ^\dagger is a positive definite quadratic form. Hence by [Wey40, Thm. 5] there exists a \mathbb{Z} -basis $\{d'_1, d'_2, d'_3, d'_4\}$ for I satisfying

$$\phi^\dagger(d'_i, d'_i) \leq C_{41} \text{disc}(I, \phi^\dagger)$$

for $i = 1, 2, 3, 4$. Hence by (14), the values $\phi_L(\eta_0\alpha(d'_i), \eta_0\alpha(d'_i))$ are bounded by a constant multiple of $\text{disc}(I, \phi^\dagger)$. Since ϕ_L is a fixed positive definite quadratic form on \mathbb{R}^4 , this implies that the coordinates of the vectors $\eta_0\alpha(d'_i)$ are polynomially bounded in terms of $\text{disc}(I, \phi^\dagger)$.

Finally, by Lemmas 4.5 and 4.6, we have

$$\text{disc}(I, \phi^\dagger) = [R : I]^2 |\text{disc}(R)| \leq C_{42} |\text{disc}(R)|^4. \quad \square$$

Let $\{\ell_1, \ell_2, \ell_3, \ell_4\}$ denote the standard basis for L . Since $\{\eta(d'_1), \eta(d'_2), \eta(d'_3), \eta(d'_4)\}$ is a \mathbb{Z} -basis for L , there is a matrix $\gamma' \in \mathbf{GL}_4(\mathbb{Z})$ such that $\ell_i = \gamma'\eta(d'_i)$ for each i .

Lemma 4.12. *The entries of the matrices $\gamma'uh, (\gamma'uh)^{-1} \in \mathbf{GL}_4(\mathbb{R})$ are polynomially bounded in terms of $|\text{disc}(R)|$.*

Proof. Let $A = \gamma'uh = \gamma'\eta\alpha^{-1}\eta_0^{-1} \in \mathbf{GL}_4(\mathbb{R})$. We have

$$\ell_i = \gamma'\eta(d'_i) = A\eta_0\alpha(d'_i).$$

By Lemma 4.11, the coordinates of the vectors $A^{-1}\ell_i = \eta_0\alpha(d'_i)$ are polynomially bounded, or in other words, the entries of the matrix A^{-1} are polynomially bounded in terms of $|\text{disc}(R)|$.

Meanwhile, $|\det(\gamma')| = \det(u) = 1$ so $|\det(A)| = |\det(h)|$. By Lemma 4.10 we deduce that $|\det(A^{-1})|$ is bounded below by a positive constant. Hence by Cramer's rule, the entries of the matrix A are also polynomially bounded in terms of $|\text{disc}(R)|$. \square

The following lemma establishes Proposition 4.1(iii)(b). Note that it is not required for the proof of Proposition 4.1(iii)(a) (the subsequent arguments proving Proposition 4.1(iii)(a) do not use the fact that $\gamma \in \mathbf{Sp}_4(\mathbb{Z})$, so they would still work with γ' instead of γ).

Lemma 4.13. *There exists $\gamma \in \Gamma = \mathbf{Sp}_4(\mathbb{Z})$ such that the entries of γuh and $(\gamma uh)^{-1}$ are polynomially bounded in terms of $|\text{disc}(R)|$.*

Proof. We have $uh \in \mathbf{H}_0(\mathbb{R}) \subset \mathbf{GSp}_4(\mathbb{R})$. Consequently

$$\psi(\gamma'^{-1}\ell_i, \gamma'^{-1}\ell_j) = \pm \det(uh)^{1/2} \psi((uh)^{-1}\gamma'^{-1}\ell_i, (uh)^{-1}\gamma'^{-1}\ell_j).$$

Using Lemmas 4.10 and 4.12, we conclude that the values $\psi(\gamma'^{-1}\ell_i, \gamma'^{-1}\ell_j)$ are polynomially bounded in terms of $|\text{disc}(R)|$.

Hence by [Orr15, Lemma 4.3], there exists a symplectic \mathbb{Z} -basis $\{s_1, s_2, s_3, s_4\}$ for (L, ψ) whose coordinates with respect to $\{\gamma'^{-1}\ell_1, \gamma'^{-1}\ell_2, \gamma'^{-1}\ell_3, \gamma'^{-1}\ell_4\}$ are polynomially bounded in terms of $|\text{disc}(R)|$. Applying γ' , we deduce that the coordinates of $\gamma's_1, \gamma's_2, \gamma's_3, \gamma's_4$ with respect to the standard basis are polynomially bounded.

Let $\gamma \in \mathbf{GL}_4(\mathbb{Z})$ be the matrix such that $\ell_i = \gamma s_i$ for each i . Since $\{s_1, s_2, s_3, s_4\}$ is a symplectic basis, we have $\gamma \in \Gamma$. We have just shown that the coordinates of $\gamma's_i = \gamma'\gamma^{-1}\ell_i$ are polynomially bounded. In other words, the entries of the matrix $\gamma'\gamma^{-1}$ are polynomially bounded in terms of $|\text{disc}(R)|$.

Multiplying $(\gamma'uh)^{-1}$ by $\gamma'\gamma^{-1}$ and applying Lemma 4.12, we deduce that the entries of $(\gamma uh)^{-1}$ are polynomially bounded in terms of $|\text{disc}(R)|$. Thanks to Lemma 4.10, $|\det((\gamma uh)^{-1})|$ is bounded below by a positive constant, so it follows that the entries of $(\gamma uh)^{-1}$ are also polynomially bounded in terms of $|\text{disc}(R)|$. \square

Let $S = E \cap M_4(\mathbb{Z}) = \text{Stab}_E(L) = \text{End}(L)^D$ and $S_0 = \iota_0(E_0) \cap M_4(\mathbb{Z})$. Set $d_u = (\text{disc}(S)/\text{disc}(S_0))^{1/2}$ and

$$v_u = d_u \rho_R(\gamma u) v_0 \in V_{\mathbb{R}}.$$

We shall use this v_u to prove Proposition 4.1(iii)(a). Note first that $d_u \rho_R(\gamma u) \in \text{Aut}_{\rho_L}(\Lambda_{\mathbb{R}})$.

Lemma 4.14. $\rho_L(u)v_u \in \Lambda$.

Proof. Since $\rho_R(\gamma)$ stabilises Λ , in order to show that $\rho_L(u)v_u \in \Lambda$, it suffices to show that

$$d_u \rho_R(u) \rho_L(u) v_0 \in \Lambda.$$

Since v_0 is a generator for the rank-1 \mathbb{Z} -module $\bigwedge^4 Z \cap \Lambda = \bigwedge^4 S_0$, it follows that $\rho_R(u) \rho_L(u) v_0$ is a generator for $\bigwedge^4 u S_0 u^{-1} \subset \bigwedge^4 \iota(E_{\mathbb{R}})$.

Consider a matrix $B \in \mathbf{GL}(E_{\mathbb{R}})$ such that $B(u S_0 u^{-1}) = S$. Conjugation by u maps the trace form ϕ_{E_0} on $\iota_0(E_{0,\mathbb{R}})$ to the trace form ϕ_E on $E_{\mathbb{R}}$, so we have

$$\text{disc}(u S_0 u^{-1}, \phi_E) = \text{disc}(S_0, \phi_{E_0}) = \text{disc}(S_0).$$

Consequently $\det(B)^2 = \text{disc}(S) / \text{disc}(S_0)$. In other words, $\det(B) = \pm d_u$.

It follows that

$$\bigwedge^4 S = \det(B) \bigwedge^4 u S_0 u^{-1} = \pm d_u \bigwedge^4 u S_0 u^{-1}$$

and so $d_u \rho_R(u) \rho_L(u) v_0$ is a generator for $\bigwedge^4 S \subset \Lambda$. \square

Lemma 4.15. $|v_u| \leq C_{43} |\text{disc}(R)|^{C_{44}}$.

Proof. The action of \mathbf{H}_0 on the line $\bigwedge^4 Z$ via ρ_R is trivial, for the same reasons as the action via ρ_L is trivial. Therefore

$$\begin{aligned} v_u &= d_u \rho_R(\gamma u) v_0 \\ &= d_u \rho_R(\gamma u h) v_0 \\ &= d_u \iota_0(e_1) (\gamma u h)^{-1} \wedge \cdots \wedge \iota_0(e_4) (\gamma u h)^{-1}. \end{aligned}$$

By Lemma 4.7, d_u is polynomially bounded in terms of $|\text{disc}(R)|$. By Lemma 4.13, the entries of $(\gamma u h)^{-1}$ are polynomially bounded in terms of $|\text{disc}(R)|$. We conclude that the coordinates, and hence the length, of v_u are polynomially bounded in terms of $|\text{disc}(R)|$. \square

Lemmas 4.14 and 4.15 complete the proof of Proposition 4.1(iii)(a).

5. UNLIKELY INTERSECTIONS IN \mathcal{A}_2

In this section, we prove Theorems 1.10, 1.11, and 1.12. We first need some preliminary material.

5.A. Realising \mathcal{A}_2 as a Shimura variety. Recall that \mathcal{A}_2 denotes the moduli space of principally polarised abelian surfaces. To realise \mathcal{A}_2 as a Shimura variety, we let (\mathbf{G}, X) denote the Shimura datum for which $\mathbf{G} = \mathbf{GSp}_4$ and X is isomorphic to the disjoint union of the Siegel upper and lower half-spaces of genus 2. We let $K = \mathbf{G}(\hat{\mathbb{Z}})$, where $\hat{\mathbb{Z}} = \prod_p \mathbb{Z}_p$, the product ranging over all finite primes p . Then K is a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$, where \mathbb{A}_f denotes the finite rational adèles, and \mathcal{A}_2 is equal to the Shimura variety whose complex points are given by

$$\text{Sh}_K(\mathbf{G}, X) = \mathbf{G}(\mathbb{Q}) \backslash X \times \mathbf{G}(\mathbb{A}_f) / K.$$

As is easily seen, this is isomorphic to the quotient $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathcal{H}_2$, where \mathcal{H}_2 denotes the Siegel upper half-space of genus 2.

5.B. Quaternionic curves and E^2 curves. Quaternionic curves and E^2 curves are the images in \mathcal{A}_2 of Shimura varieties of PEL type. We recall the construction of Shimura varieties of PEL type in the case of quaternion algebras over \mathbb{Q} , following [Mil05, sec. 8].

Let B denote a quaternion algebra over \mathbb{Q} such that $B \otimes_{\mathbb{Q}} \mathbb{R}$ is isomorphic to $\mathbf{M}_2(\mathbb{R})$ and let \dagger be a positive involution of B . (Positive involutions exist for any such B , as explained in [Mum74, pp. 195–196].) As explained in [Mum74, p. 196], we can choose the isomorphism $B \otimes_{\mathbb{Q}} \mathbb{R} \rightarrow \mathbf{M}_2(\mathbb{R})$ in such a way that \dagger corresponds to transpose of matrices, so $(B \otimes_{\mathbb{Q}} \mathbb{C}, \dagger)$ has type C in the sense of [Mil05, Prop. 8.3].

Choose $\alpha \in B$ such that $\alpha = -\alpha^\dagger$. Then we can define a symplectic form on B by the formula

$$\psi_\alpha(x, y) = \mathrm{Tr}_{B/\mathbb{Q}}(x\alpha y^\dagger).$$

If $\alpha \in B^\times$, then ψ_α is non-degenerate and (B, ψ_α) (with B acting via the left regular representation) is a symplectic (B, \dagger) -module as defined in [Mil05, sec. 8].

Let \mathbf{H}_B denote the centraliser of B (acting on itself via the left regular representation) in $\mathbf{GL}(B)$, seen as a \mathbb{Q} -algebraic group. Since every symplectic form $B \times B \rightarrow \mathbb{Q}$ which satisfies $\psi(bx, y) = \psi(x, b^\dagger y)$ has the form ψ_α for some α such that $\alpha = -\alpha^\dagger$, and (for a given positive involution \dagger) such α form a one-dimensional \mathbb{Q} -linear subspace of B , every element of \mathbf{H}_B preserves ψ_α up to multiplication by a scalar. Therefore, $\mathbf{H}_B = \mathbf{GAut}_B(B, \psi_\alpha)$. In other words, \mathbf{H}_B is equal to the group denoted G in [Mil05, sec. 8]. (Note: we have $\mathbf{H}_B(\mathbb{Q}) \cong (B^{\mathrm{op}})^\times \cong B^\times$ where the second isomorphism uses the fact that B is a quaternion algebra.)

By [Mil05, Prop. 8.14], there is a unique Shimura datum (\mathbf{H}_B, X_B) such that each $h \in X_B$ defines a complex structure on $B \otimes_{\mathbb{Q}} \mathbb{R}$ for which the symmetric form $\psi(x, h(i)y)$ is positive or negative definite. As a Hermitian symmetric domain, X_B is isomorphic to the union of the upper and lower half-planes in \mathbb{C} .

Choosing a symplectic \mathbb{Q} -basis for (B, ψ_α) , the tautological action of \mathbf{H}_B on B gives rise to an injective group homomorphism $\mathbf{H}_B \rightarrow \mathbf{G}$. Thanks to the properties of X_B given to us by [Mil05, Prop. 8.14], this induces an embedding of Shimura data $(\mathbf{H}_B, X_B) \rightarrow (\mathbf{G}, X)$. Letting $K_B = \mathbf{H}_B(\mathbb{A}_f) \cap K$, we obtain a morphism

$$\mathrm{Sh}_{K_B}(\mathbf{H}_B, X_B) \rightarrow \mathrm{Sh}_K(\mathbf{G}, X)$$

of algebraic varieties. The irreducible components of the images of such morphisms are, by definition, special curves in \mathcal{A}_2 . If B is isomorphic (over \mathbb{Q}) to $\mathbf{M}_2(\mathbb{Q})$, we obtain E^2 curves, and otherwise we obtain quaternionic curves. Any such curve parametrises abelian surfaces with multiplication by an order in B .

The Shimura data (\mathbf{G}, X) and (\mathbf{H}_B, X_B) all have reflex field \mathbb{Q} . Therefore, $\mathrm{Sh}_{K_B}(\mathbf{H}_B, X_B)$, $\mathrm{Sh}_K(\mathbf{G}, X)$ and $\mathrm{Sh}_{K_B}(\mathbf{H}_B, X_B) \rightarrow \mathrm{Sh}_K(\mathbf{G}, X)$ are all defined over \mathbb{Q} (although $\mathrm{Sh}_{K_B}(\mathbf{H}_B, X_B)$ often has geometrically irreducible components which

are not defined over \mathbb{Q}). In particular, the action of $\text{Aut}(\mathbb{C}/\mathbb{Q})$ on $\mathcal{A}_2(\mathbb{C})$ induces actions on the set of quaternionic curves and on the set of E^2 curves in \mathcal{A}_2 . From the theory of complex multiplication of abelian varieties, we know that $\text{Aut}(\mathbb{C}/\mathbb{Q})$ acts on the set of special points in \mathcal{A}_2 . Consequently, $\text{Aut}(\mathbb{C}/\mathbb{Q})$ acts on

$$\Sigma_{\text{Quat}} = \bigcup_{Z \in \mathcal{S}} Z \setminus Z^{\text{sp}},$$

where \mathcal{S} denotes the set of quaternionic curves in \mathcal{A}_2 and Z^{sp} denotes the union of the special points contained in Z . Similarly, $\text{Aut}(\mathbb{C}/\mathbb{Q})$ acts on Σ_{E^2} .

Another way to obtain these families of special subvarieties is as follows. Let $B_0 = \text{M}_2(\mathbb{Q})$. Let B_0 act on \mathbb{Q}^4 via the left regular representation, with respect to the basis given by (8) (which is a symplectic basis with respect to the form ψ_α where $\alpha = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$). Let $\mathbf{H}_0 \subset \mathbf{G}$ be the centraliser of this action of B_0 in \mathbf{G} . Then \mathbf{H}_0 is equal to the image of \mathbf{GL}_2 embedded block diagonally, as in (7). Let

$$X_0 = \left\{ \begin{pmatrix} \tau & 0 \\ 0 & \tau \end{pmatrix} \in \mathcal{H}_2 : \text{Im}(\tau) > 0 \right\},$$

$$X_0^\pm = \left\{ \begin{pmatrix} \tau & 0 \\ 0 & \tau \end{pmatrix} \in \mathcal{H}_2 : \text{Im}(\tau) \neq 0 \right\}.$$

Then (\mathbf{H}_0, X_0^\pm) is the unique Shimura subdatum of (\mathbf{G}, X) with underlying group \mathbf{H}_0 , and X_0 is the only connected component of X_0 contained in \mathcal{H}_2 .

For any point $x_0 \in X_0$, we have $X_0 = \mathbf{H}_0^{\text{der}}(\mathbb{R})x_0$ (recall that $\mathbf{H}_0^{\text{der}}(\mathbb{R}) = \mathbf{SL}_2(\mathbb{R})$ is connected) and its image in \mathcal{A}_2 is an E^2 curve. For any $g \in \mathbf{G}(\mathbb{R})$ such that $\mathbf{H} = g\mathbf{H}_{0,\mathbb{R}}g^{-1}$ is defined over \mathbb{Q} the image of gX_0 in \mathcal{A}_2 is a special curve, and \mathbf{H} is isomorphic (as a \mathbb{Q} -group) to \mathbf{H}_B for some quaternion algebra B as above. If \mathbf{H} is isomorphic to $\mathbf{GL}_{2,\mathbb{Q}}$, then we obtain an E^2 curve, and, otherwise, we obtain a quaternionic curve.

Lemma 5.1. *Every quaternionic or E^2 curve in \mathcal{A}_2 is the image of gX_0 for some $g \in \mathbf{G}(\mathbb{R})$ such that $g\mathbf{H}_{0,\mathbb{R}}g^{-1}$ is defined over \mathbb{Q} .*

Proof. Let Z be a quaternionic or E^2 curve in \mathcal{A}_2 . Let B be the generic endomorphism algebra of the abelian surfaces parametrised by Z , and let \dagger be the Rosati involution of B . Choose an analytic irreducible component Y of the preimage of Z in X .

The inclusion $\mathbf{G} \rightarrow \mathbf{GL}_4$ induces a variation \mathcal{V} of \mathbb{Q} -Hodge structures on X with trivial underlying local system $X \times \mathbb{Q}^4$. The restriction $\mathcal{V}|_Y$ has endomorphism algebra B and its generic Mumford–Tate group $\mathbf{H} \subset \mathbf{G}$ is the centraliser of B in \mathbf{G} . Thus \mathbf{H} is the image of one of the homomorphisms $\mathbf{H}_B \rightarrow \mathbf{G}$ defined above.

The choice of basis (8) induces an isomorphism of \mathbb{Q} -vector spaces $\mathbb{Q}^4 \rightarrow B_0$. Choose an isomorphism of \mathbb{R} -algebras with involutions $(B_0 \otimes_{\mathbb{Q}} \mathbb{R}, t) \rightarrow (B \otimes_{\mathbb{Q}} \mathbb{R}, \dagger)$. The action of B on $\mathcal{V}|_Y$ gives rise to a B -module structure on \mathbb{Q}^4 , and this is isomorphic to the left regular representation of B on itself, i.e. we get

an isomorphism of B -modules $B \rightarrow \mathbb{Q}^4$. Composing these isomorphisms (after extending scalars to \mathbb{R}), we get an isomorphism of \mathbb{R} -vector spaces

$$\mathbb{R}^4 \rightarrow B_0 \otimes_{\mathbb{Q}} \mathbb{R} \rightarrow B \otimes_{\mathbb{Q}} \mathbb{R} \rightarrow \mathbb{R}^4$$

or in other words an element $g \in \mathbf{GL}_4(\mathbb{R})$. Via the isomorphism $\mathbb{R}^4 \rightarrow B_0 \otimes_{\mathbb{Q}} \mathbb{R}$, the standard symplectic form on \mathbb{R}^4 satisfies $\psi(bx, y) = \psi(x, b^t y)$ for all $b \in B_0$, and the isomorphism $B \otimes_{\mathbb{Q}} \mathbb{R} \rightarrow \mathbb{R}^4$ behaves similarly with respect to (B, \dagger) . Since the spaces of symplectic forms satisfying these conditions are one-dimensional, the composed isomorphism maps the standard symplectic form to a multiple of itself; in other words, $g \in \mathbf{G}(\mathbb{R})$.

Comparing the actions of $B \otimes_{\mathbb{Q}} \mathbb{R}$ and $B_0 \otimes_{\mathbb{Q}} \mathbb{R}$, we see that $\mathbf{H}_{\mathbb{R}} = g\mathbf{H}_{0,\mathbb{R}}g^{-1}$. It follows that $g^{-1}Y$ is a connected component of a Shimura subdatum of (\mathbf{G}, X) with underlying group \mathbf{H}_0 . The only such Shimura subdatum is $(\mathbf{H}_0, X_0^{\pm})$, so $g^{-1}Y$ is a connected component of X_0^{\pm} .

If $g^{-1}Y \neq X_0$, replace g by $g \operatorname{diag}(1, -1, 1, -1)$. Since $\operatorname{diag}(1, -1, 1, -1) \in \mathbf{H}_0(\mathbb{R})$ and it swaps the two connected components of X_0^{\pm} , after this replacement we will still have $\mathbf{H}_{\mathbb{R}} = g\mathbf{H}_{0,\mathbb{R}}g^{-1}$ but now $g^{-1}Y = X_0$.

Thus we get $Y = gX_0$ and Z is the image of Y in \mathcal{A}_2 . □

5.C. Complexity. As in [DR18], we will need to define a notion of complexity. That is, to each E^2 or quaternionic curve Z in \mathcal{A}_2 , we attach a natural number $\Delta(Z)$, which we refer to as the complexity of Z . The complexity should be defined in terms of arithmetic information associated with Z and possess the fundamental property that only finitely many E^2 or quaternionic curves in \mathcal{A}_2 have complexity $\leq b$ for any $b \in \mathbb{R}$. We proceed as follows.

Let $g \in \mathbf{G}(\mathbb{R})$ be such that $\mathbf{H} = g\mathbf{H}_{0,\mathbb{R}}g^{-1}$ is defined over \mathbb{Q} . Then the image Z of gX_0 in \mathcal{A}_2 is an E^2 or quaternionic curve and, by Lemma 5.1 every E^2 or quaternionic curve is obtained this way. We define the complexity $\Delta(Z)$ of Z to be $|\operatorname{disc}(R)|$, where R denotes the ring $\operatorname{End}_{\mathbf{H}}(\mathbb{Z}^4)$ of \mathbb{Z} -linear endomorphisms of $\mathbb{Z}^4 \subset \mathbb{Q}^4$ commuting with $\mathbf{H}(\mathbb{Q}) \subset \mathbf{G}(\mathbb{Q}) \subset \mathbf{M}_4(\mathbb{Q})$. Note that this ring R is the generic endomorphism ring of the abelian surfaces parameterised by Z . Indeed, for every non-special point of Z , the associated abelian surface (over \mathbb{C}) has endomorphism ring isomorphic to R . We will show that $\Delta(Z)$ is a suitable complexity with Corollary 5.5 below.

We are now in a position to make the conjectures alluded to above. Note that these conjectures are identical except that Conjecture 5.2 refers to Σ_{Quat} and Conjecture 5.3 refers to Σ_{E^2} .

Conjecture 5.2. *Let V and Σ_{Quat} be as in Theorem 1.10. Let L be a finitely generated subfield of \mathbb{C} over which V is defined.*

There exist positive constants C_{45} and C_{46} such that, for any point $s \in V \cap \Sigma_{\text{Quat}}$, if we let Z denote the (unique) special curve containing s , then

$$\# \operatorname{Aut}(\mathbb{C}/L) \cdot s \geq C_{45} \Delta(Z)^{C_{46}}.$$

Conjecture 5.3. *Let V and Σ_{E^2} be as in Theorem 1.11. Let L be a finitely generated subfield of \mathbb{C} over which V is defined.*

There exist positive constants C_{47} and C_{48} such that, for any point $s \in V \cap \Sigma_{E^2}$, if we let Z denote the (unique) special curve containing s , then

$$\# \text{Aut}(\mathbb{C}/L) \cdot s \geq C_{47} \Delta(Z)^{C_{48}}.$$

5.D. The fixed data. We write Γ for the congruence subgroup $\mathbf{Sp}_4(\mathbb{Z}) \subset \mathbf{G}(\mathbb{Q})$. We let $\pi : \mathcal{H}_2 \rightarrow \mathcal{A}_2$ denote the (transcendental) uniformisation map. We choose a Siegel set $\mathfrak{S} \subset \mathbf{G}(\mathbb{R})^+$ (associated with the standard Siegel triple) such that, for some finite set $C \subset \mathbf{G}(\mathbb{Q})$, $\mathcal{F}_{\mathbf{G}} = C\mathfrak{S}$ is a fundamental set for Γ in $\mathbf{G}(\mathbb{R})^+$. We write \mathcal{F} for $\mathcal{F}_{\mathbf{G}}x_0$, where $x_0 \in \mathcal{H}_2$ is the point associated with $\mathcal{F}_{\mathbf{G}}$.

By Proposition 4.1, we can fix a finitely generated, free \mathbb{Z} -module Λ , a representation $\rho : \mathbf{G} \rightarrow \mathbf{GL}(\Lambda_{\mathbb{Q}})$ such that Λ is stabilised by $\rho(\Gamma)$, an element $v_0 \in \Lambda$, and positive constants C_{49} and C_{50} such that

- (i) $\text{Stab}_{\mathbf{G},\rho}(v_0) = \mathbf{H}_0$;
- (ii) the orbit $\rho(\mathbf{G}(\mathbb{R}))v_0$ is closed in $\Lambda_{\mathbb{R}}$;
- (iii) for each $u \in \mathbf{G}(\mathbb{R})$, if the group $\mathbf{H}_u = u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , then there exists $v_u \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ such that $\rho(u)v_u \in \Lambda$ and

$$|v_u| \leq C_{49} |\text{disc}(R_u)|^{C_{50}},$$

where R_u denotes the ring $\text{End}_{\mathbf{H}_u}(L) \subset M_4(\mathbb{Z})$.

By Theorem 1.9, we can then fix positive constants C_{51} and C_{52} with the following property: for every $u \in \mathbf{G}(\mathbb{R})$, if $\mathbf{H}_u = u\mathbf{H}_{0,\mathbb{R}}u^{-1}$ is defined over \mathbb{Q} , then there exists a fundamental set for $\Gamma \cap \mathbf{H}_u(\mathbb{R})$ in $\mathbf{H}_u(\mathbb{R})$ of the form

$$B_u C \mathfrak{S} u^{-1} \cap \mathbf{H}_u(\mathbb{R}),$$

where $B_u \subset \Gamma$ is a finite set such that

$$|\rho(b^{-1}u)v_u| \leq C_{51} |v_u|^{C_{52}}$$

for every $b \in B_u$.

Choosing a basis, we obtain $\Lambda = \mathbb{Z}^d$ and we may refer to the height $H(v)$ of any $v \in \Lambda$ (defined as the maximum of the absolute values of the coordinates). For any $v \in \Lambda_{\mathbb{R}}$, we write $\mathbf{G}(v) = \text{Stab}_{\mathbf{G},\rho}(v)$.

Proposition 5.4. *Let $P \in \Sigma_{\text{quat}} \cup \Sigma_{E^2}$. Then there exists $z \in \pi^{-1}(P) \cap \mathcal{F}$ and $v \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})\rho(\mathbf{G}(\mathbb{R}))v_0 \cap \Lambda$ such that $z(\mathbb{S}) \subset \mathbf{G}(v)$ and*

$$H(v) \leq C_{51} C_{49}^{C_{50}} |\text{disc}(R)|^{C_{50}C_{52}},$$

where R denotes the ring $\text{End}_{\mathbf{G}(v)}(L) \subset M_4(\mathbb{Z})$.

Proof. Let $z \in \pi^{-1}(P) \cap \mathcal{F}$ and let Y denote the smallest pre-special subvariety of \mathcal{H}_2 containing z . Then $\pi(Y)$ is an E^2 or quaternionic curve and so

$$Y = g\mathbf{H}_0^{\text{der}}(\mathbb{R})x_0 = g\mathbf{H}_0^{\text{der}}(\mathbb{R})g^{-1} \cdot gx_0 = \mathbf{H}^{\text{der}}(\mathbb{R}) \cdot gx_0,$$

where $g \in \mathbf{G}(\mathbb{R})$, \mathbf{H} is a \mathbb{Q} -subgroup of \mathbf{G} isomorphic to \mathbf{H}_B for some quaternion algebra B , as above, $\mathbf{H}_{\mathbb{R}} = g\mathbf{H}_{0,\mathbb{R}}g^{-1}$ and $z(\mathbb{S}) \subset \text{MT}(Y) = \mathbf{H}(\mathbb{R})$. By Proposition 4.1, we obtain $v \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})v_0$ such that $\rho(g)v \in \Lambda$ and

$$|v| \leq C_{49}|\text{disc}(R)|^{C_{50}},$$

where $R = \text{End}_{\mathbf{H}}(L)$. Note that

$$\mathbf{G}(\rho(g)v) = g\mathbf{G}(v)g^{-1} = g\mathbf{G}(v_0)g^{-1} = \mathbf{H}_{\mathbb{R}}.$$

By Theorem 1.9 (with $u = g$), we obtain a finite set $B_g \subset \Gamma$ such that

$$\mathcal{F}_{\mathbf{H}} = B_g C\mathfrak{S}g^{-1} \cap \mathbf{H}(\mathbb{R})$$

is a fundamental set for $\Gamma_{\mathbf{H}} = \Gamma \cap \mathbf{H}(\mathbb{R})$ in $\mathbf{H}(\mathbb{R})$, and

$$|\rho(b^{-1}g)v| \leq C_{51}|v|^{C_{52}}$$

for all $b \in B_g$. In particular, since $z \in \mathbf{H}(\mathbb{R})gx_0$, we can write

$$z = \gamma bsx_0$$

for some $\gamma \in \Gamma_{\mathbf{H}}$, $b \in B_g$, and $s \in C\mathfrak{S}$. Hence,

$$z' := b^{-1}\gamma^{-1}z \in \mathcal{F} \cap \pi^{-1}(P).$$

Furthermore, we have

$$z'(\mathbb{S}) \subset \mathbf{G}(\rho(b^{-1}\gamma^{-1}g)v) = \mathbf{G}(\rho(b^{-1}g)v) = b^{-1}\mathbf{H}_{\mathbb{R}}b,$$

where we use the fact that $\gamma \in \mathbf{H}(\mathbb{R}) = \mathbf{G}(\rho(g)v)(\mathbb{R})$. Finally, from the above, we have

$$|\rho(b^{-1}g)v| \leq C_{51}|v|^{C_{52}} \leq C_{51}(C_{49}|\text{disc}(R)|^{C_{50}})^{C_{52}} = C_{51}C_{49}^{C_{50}}|\text{disc}(R)|^{C_{50}C_{52}}$$

and

$$R = \text{End}_{\mathbf{H}}(L) = \text{End}_{b^{-1}\mathbf{H}b}(b^{-1}L) = \text{End}_{b^{-1}\mathbf{H}b}(L) = \text{End}_{\mathbf{G}(\rho(b^{-1}g)v)}(L).$$

Therefore, since $\rho(b^{-1}g)v \in \Lambda$, we conclude that z' and $\rho(b^{-1}g)v$ satisfy the conditions of the proposition. \square

Corollary 5.5. *Let $b \in \mathbb{R}$. The set of E^2 or quaternionic curves Z satisfying $\Delta(Z) \leq b$ is finite.*

Proof. Let Z be an E^2 or quaternionic curve satisfying $\Delta(Z) \leq b$ and let $P \in Z$ be a Hodge generic point on Z . Therefore, $P \in \Sigma_{\text{Quat}} \cup \Sigma_{E^2}$ and, by Proposition 5.4, we obtain $z \in \pi^{-1}(P) \cap \mathcal{F}$ and $v \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})\rho(\mathbf{G}(\mathbb{R}))v_0 \cap \Lambda$ such that $z(\mathbb{S}) \subset \mathbf{G}(v)$ and

$$\mathbf{H}(v) \leq C_{51}C_{49}^{C_{50}}|\text{disc}(R)|^{C_{50}C_{52}},$$

where R denotes the ring $\text{End}_{\mathbf{G}(v)}(L) \subset \text{M}_4(\mathbb{Z})$. Let Y denote the smallest pre-special subvariety of \mathcal{H}_2 containing z . Since $\pi(Y) = Z$ is an E^2 or quaternionic curve, we have

$$Y = g\mathbf{H}_0^{\text{der}}(\mathbb{R})x_0 = g\mathbf{H}_0^{\text{der}}(\mathbb{R})g^{-1} \cdot gx_0 = \mathbf{H}^{\text{der}}(\mathbb{R}) \cdot gx_0,$$

where $g \in \mathbf{G}(\mathbb{R})$, \mathbf{H} is a \mathbb{Q} -subgroup of \mathbf{G} isomorphic to \mathbf{H}_B for some quaternion algebra B , as above, $\mathbf{H}_{\mathbb{R}} = g\mathbf{H}_{0,\mathbb{R}}g^{-1}$ and $z(\mathbb{S}) \subset \text{MT}(Y) = \mathbf{H}(\mathbb{R})$. Therefore, $\mathbf{H}(\mathbb{R}) \subset \mathbf{G}(v)$ and since $\mathbf{G}(v)$ is also a conjugate of $\mathbf{H}_0(\mathbb{R})$, we conclude that $\mathbf{G}(v) = \mathbf{H}(\mathbb{R})$. The fact that $\Delta(Z) \leq b$ thus implies that $\mathbf{H}(v)$ is bounded, which implies that v belongs to a finite set and, therefore, so does \mathbf{H} . As in the proof of Lemma 5.1, there is only one Shimura subdatum of (\mathbf{G}, X) associated with each \mathbf{H} and so the result follows. \square

5.E. Proof of Theorem 1.10. Let $\mathcal{V} = \pi^{-1}(V) \cap \mathcal{F}$ – a set definable in the o-minimal structure $\mathbb{R}_{\text{an,exp}}$ (see [KUY16] for more details). Let L be a finitely generated field of definition for V . Let $P \in V \cap \Sigma_{\text{Quat}}$. Varying over $\sigma \in \text{Aut}(\mathbb{C}/L)$, we obtain points $\sigma(P) \in V \cap \Sigma_{\text{Quat}}$ and, for each, we let $z_{\sigma} \in \mathcal{F} \cap \pi^{-1}(\sigma(P))$ and we let $v_{\sigma} \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})\rho(\mathbf{G}(\mathbb{R}))v_0 \cap \Lambda$ be the elements afforded to us by Proposition 5.4. That is, $z_{\sigma}(\mathbb{S}) \subset \mathbf{G}(v_{\sigma})$ and

$$\mathbf{H}(v_{\sigma}) \leq C_{51}C_{49}^{C_{50}}|\text{disc}(R_{\sigma})|^{C_{50}C_{52}},$$

where R_{σ} denotes the ring $\text{End}_{\mathbf{G}(v_{\sigma})}(L) \subset \text{M}_4(\mathbb{Z})$. As above, $|\text{disc}(R_{\sigma})| = \Delta(\sigma(Z))$. Note that we also have $z_{\sigma} \in \mathcal{V}$.

We obtain a set Θ of tuples $(v_{\sigma}, z_{\sigma}) \in \Lambda \times \mathcal{V}$ belonging to the definable set

$$D = \{(v, z) \in \Lambda_{\mathbb{R}} \times \mathcal{V} : v \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})\rho(\mathbf{G}(\mathbb{R}))v_0, z(\mathbb{S}) \subset \mathbf{G}(v)\}.$$

Let $\pi_1 : D \rightarrow \Lambda_{\mathbb{R}}$ and let $\pi_2 : D \rightarrow \mathcal{V}$ denote the projection maps. By Conjecture 5.2, we have

$$\begin{aligned} \#\pi_2(\Theta) &= \#\text{Aut}(\mathbb{C}/L) \cdot P = \#\text{Aut}(\mathbb{C}/L) \cdot \sigma(P) \\ &\geq C_{45}|\text{disc}(R_{\sigma})|^{C_{46}} \geq C_{53}H(v_{\sigma})^{C_{46}/C_{50}C_{52}}. \end{aligned}$$

Applying [DR18, Theorem 9.1] (a variant of [HP16, Corollary 7.2]) with $\varepsilon < C_{46}/C_{50}C_{52}$, we conclude that either

- (1) $\#\pi_2(\Theta)$ is bounded, hence $\Delta(Z)$ is bounded and the theorem holds,
- (2) or there exists a continuous definable function

$$\beta : [0, 1] \rightarrow D$$

such that $\beta_1 = \pi_1 \circ \beta$ is semi-algebraic, $\beta_2 = \pi_2 \circ \beta$ is non-constant, $\beta(0) \in \Theta$, and $\beta|_{(0,1)}$ is real analytic. Therefore, it suffices to rule out the latter possibility.

To that end, suppose that we have such a function. By definable choice, there exists a semi-algebraic function

$$\tilde{\beta}_1 : [0, 1] \rightarrow \text{Aut}_{\rho}(\Lambda_{\mathbb{R}})\rho(\mathbf{G}(\mathbb{R}))$$

such that $\tilde{\beta}_1(t) \cdot v_0 = \beta_1(t)$ for all $t \in [0, 1]$. We let $v_t = \beta_1(t)$, $g_t = \tilde{\beta}_1(t)$, and $z_t = \beta_2(t)$. Since $z_t(\mathbb{S}) \subset \mathbf{G}(v_t)$ and $g_t \in \text{Aut}_{\rho(\mathbf{G})}(\Lambda_{\mathbb{R}})\rho(\mathbf{G}(\mathbb{R}))$, we have

$$(g_t^{-1}z_t)(\mathbb{S}) \subset g_t^{-1}\mathbf{G}(v_t)g_t = \mathbf{G}(g_t^{-1}v_t) = \mathbf{G}(v_0) = \mathbf{H}_0(\mathbb{R}).$$

We conclude that $g_t^{-1}z_t$ lies on the unique pre-special subvariety of \mathcal{H}_2 associated with $\mathbf{H}_0^{\text{der}}(\mathbb{R})$, namely, X_0 .

On the other hand, there exists $0 < t_1 \leq 1$ such that $\beta_2([0, t_1])$ is contained in a single irreducible analytic component V_1 of $\pi^{-1}(V)$. By [UY18, Theorem 1.3] (the inverse Ax–Lindemann conjecture), \mathcal{H}_2 is the smallest algebraic subset of \mathcal{H}_2 containing V_1 .

Let $B \subset \Lambda_{\mathbb{C}}$ denote the Zariski closure of $\beta_1([0, t_1]) \subset \Lambda_{\mathbb{R}}$. By definable choice, there exists a complex algebraic set $\tilde{B} \subset \text{Aut}_{\rho}(\Lambda_{\mathbb{C}})\rho(\mathbf{G}(\mathbb{C}))$ of dimension at most 1 whose image under the algebraic map $g \mapsto g \cdot v_0$ is B . Using the superscript $^{\vee}$ to denote the compact dual of a hermitian symmetric domain, we obtain a complex algebraic set $\tilde{B} \times X_0^{\vee}$ of dimension at most 2. Note that $\tilde{B} \cdot \mathcal{H}_2^{\vee} = \mathcal{H}_2^{\vee}$. Hence, $\tilde{B} \cdot X_0^{\vee} \subset \mathcal{H}_2^{\vee}$ is algebraic of dimension at most 2.

On the other hand, V_1 is an irreducible complex analytic curve having an uncountable intersection with $\tilde{B} \cdot X_0^{\vee}$ (in particular, it includes $\beta_2([0, t_1])$) because $g_t^{-1}z_t \in X_0$ and $z_t \in V_1$ for all $t \in [0, t_1]$. Therefore, V_1 is contained in $\tilde{B} \cdot X_0^{\vee}$, hence, so is \mathcal{H}_2^{\vee} . However, $\dim \mathcal{H}_2^{\vee} = 3$, and we arrive at a contradiction.

5.F. Proof of Theorem 1.11. The proof proceeds exactly as in the previous section, working with Σ_{E^2} instead of Σ_{Quat} and Conjecture 5.3 instead of Conjecture 5.2.

5.G. Proof of Theorem 1.12. We claim that it suffices to prove the following Theorem. In the below, if V is an algebraic curve over a number field and $\mathfrak{A} \rightarrow V$ is an abelian scheme of relative dimension 2, we say that $s \in V(\overline{\mathbb{Q}})$ is a quaternionic point if the endomorphism algebra of the fiber \mathfrak{A}_s is a quaternion algebra over \mathbb{Q} not isomorphic to $M_2(\mathbb{Q})$.

Theorem 5.6. *Let V be an irreducible algebraic curve and let $\mathfrak{A} \rightarrow V$ be a principally polarised non-isotrivial abelian scheme of relative dimension 2 such that $\text{End}(\mathfrak{A}_{\bar{\eta}}) = \mathbb{Z}$, where $\bar{\eta}$ denotes a geometric generic point of V .*

Suppose that V and \mathfrak{A} are defined over a number field L and that there exist a curve V' , a semiabelian scheme $\mathfrak{A}' \rightarrow V'$ and an open immersion $\iota: V \rightarrow V'$, all defined over $\overline{\mathbb{Q}}$, such that $\mathfrak{A} \cong \iota^\mathfrak{A}'$ and there is a point $s_0 \in V'(\overline{\mathbb{Q}}) \setminus V(\overline{\mathbb{Q}})$ for which the fibre \mathfrak{A}'_{s_0} is a torus.*

Then there exist positive constants C_{54} and C_{55} such that, for any quaternionic point $s \in V$,

$$\# \text{Aut}(\mathbb{C}/L) \cdot s \geq C_{54} |\text{disc}(\text{End}(\mathfrak{A}_s))|^{C_{55}}.$$

To see that Theorem 5.6 implies Theorem 1.12, consider V as in Theorem 1.12. Then V is defined over a number field L and, furthermore, we can construct a curve \tilde{V}' , a finite surjective morphism $q: \tilde{V} \rightarrow V$, $s_0 \in \tilde{V}'$, and a semiabelian scheme $\mathfrak{A}' \rightarrow \tilde{V}'$ as in [DO, Proposition 10.2]. We can find a finite extension \tilde{L}/L such that \tilde{V}' , $q: \tilde{V} \rightarrow V$, s_0 and \mathfrak{A}' are all defined over \tilde{L} . The abelian scheme

$\mathfrak{A}'_{|\tilde{V}} \rightarrow \tilde{V}$ and the point $s_0 \in \tilde{V}'(\overline{\mathbb{Q}})$ satisfy the conditions of Theorem 5.6 and so, for any quaternionic point $\tilde{s} \in \tilde{V}$,

$$\# \text{Aut}(\mathbb{C}/\tilde{L}) \cdot \tilde{s} \geq C_{54} |\text{disc}(\text{End}(\mathfrak{A}'_{\tilde{s}}))|^{C_{55}}.$$

If $s \in V \cap \Sigma_{\text{Quat}}$, then we can find a quaternionic point $\tilde{s} \in \tilde{V}$ such that $q(\tilde{s}) = s$, and since q is finite,

$$\# \text{Aut}(\mathbb{C}/L) \cdot s \geq C_{56} \# \text{Aut}(\mathbb{C}/\tilde{L}) \cdot \tilde{s}.$$

Let Z denote the unique special curve in \mathcal{A}_2 containing s . Since s is a Hodge generic point of Z , the endomorphism ring of the associated abelian surface A_s is isomorphic to the generic endomorphism ring of Z and so $\Delta(Z) = |\text{disc}(\text{End}(A_s))|$. Since also $A_s = \mathfrak{A}'_{\tilde{s}}$, we can combine the above inequalities to obtain

$$\# \text{Aut}(\mathbb{C}/L) \cdot s \geq C_{54} C_{56} \Delta(Z)^{C_{55}},$$

that is, Conjecture 5.2.

Therefore, it remains to prove Theorem 5.6.

Proof of Theorem 5.6. After a finite extension, we may assume that $V', \mathfrak{A}' \rightarrow V', \iota: V \rightarrow V'$, and s_0 are all defined over L . Since $\text{End}(\mathfrak{A}_{\overline{\eta}}) = \mathbb{Z}$ and $\dim(\mathfrak{A}_{\overline{\eta}}) = 2$, the Mumford–Tate group of $\mathfrak{A}_{\overline{\eta}}$ is $\mathbf{GSp}_{4, \mathbb{Q}}$ (see [DO, Section 2.F]). Thus $\mathfrak{A} \rightarrow V$ satisfies the conditions of [DO, Theorem 9.1], as modified in [DO, Remark 9.6].

Let $s \in V$ be a quaternionic point. The image of s under the map $V \rightarrow \mathcal{A}_2$ induced by $\mathfrak{A} \rightarrow V$ is in the intersection between the image of V and a quaternionic curve. We deduce that $s \in V(\overline{\mathbb{Q}})$.

Now $\text{End}(\mathfrak{A}_s) \otimes \mathbb{Q}$ is a non-split quaternion algebra, so cannot inject into $M_2(\mathbb{Q})$. Hence, A_s is exceptional in the sense of [DO, Section 9]. Therefore, by [DO, Theorem 9.1], $h(s)$ is polynomially bounded in terms of $[L(s) : L]$, where h denotes a Weil height on V' . Let h_F denote the stable Faltings height. As proved in [Fal83, p. 356],

$$|h_F(\mathfrak{A}_s) - h(s)| = O(\log h(s)).$$

We conclude that $h_F(\mathfrak{A}_s)$ is polynomially bounded in terms of $[L(s) : L]$.

In order to deduce a bound for $\text{disc}(\text{End}(A_s))$, we use the following theorem of Masser and Wüstholz.

Theorem. [MW94, p. 641] *Given positive integers n, d and δ , there are constants $C_{57} = C_{57}(n, d, \delta)$ and $C_{58} = C_{58}(n)$, with the following property. Let A be an abelian variety of dimension n defined over a number field k of degree d , equipped with a polarisation of degree δ . Let \dagger be the Rosati involution of $\text{End}(A)$ associated with this polarisation and let ϕ^\dagger be the bilinear form on $\text{End}(A)$ defined by (10). Then $\text{disc}(\text{End}_k(A), \phi^\dagger)$ is at most $C_{57} \max(1, h_F(A))^{C_{58}}$.*

As remarked in [MW94] immediately following the statement of this theorem, one can replace $\text{End}_k(A)$ by $\text{End}_{\mathbb{C}}(A)$ (the endomorphism ring which appears in the statement of Theorem 5.6) because one can find a finite extension K/k of degree

bounded only in terms of n such that $\text{End}_K(A) = \text{End}_{\mathbb{C}}(A)$. Furthermore, as stated near the bottom of [MW94, p. 650], the constant $C_{57}(n, d, \delta)$ is polynomial in d and δ , for a polynomial which depends only on n . Using also Lemma 4.5 to see that $|\text{disc}(\text{End}(A), \phi^\dagger)| = |\text{disc}(\text{End}(A))|$, we conclude that there are constants C_{59}, C_{60} depending only on n such that, for all A as in the theorem, we have

$$|\text{disc}(\text{End}(A))| \leq C_{59} \max(\delta, d, h_F(A))^{C_{60}}. \tag{15}$$

In our case, we have always $n = 2$ and $\delta = 1$. So applying (15) together with the fact that $h_F(A_s)$ is polynomially bounded in terms of $[L(s) : L]$ completes the proof of Theorem 5.6. \square

REFERENCES

[AMRT75] A. Ash, D. Mumford, M. Rapoport, and Y. Tai, *Smooth compactification of locally symmetric varieties*, Math. Sci. Press, Brookline, Mass., 1975, Lie Groups: History, Frontiers and Applications, Vol. IV.

[And89] Y. André, *G-functions and geometry*, Aspects of Mathematics, E13, Friedr. Vieweg & Sohn, Braunschweig, 1989.

[BDR] M. Borovoi, C. Daw, and J. Ren, *Conjugation of semisimple subgroups over real number fields of bounded degree*, to appear in Proc. Amer. Math. Soc., available at arXiv:1802.05894.

[BHC62] A. Borel and Harish-Chandra, *Arithmetic subgroups of algebraic groups*, Ann. of Math. (2) **75** (1962), 485–535.

[Bla48] H. Blaney, *Indefinite quadratic forms in n variables*, J. London Math. Soc. **23** (1948), 153–160.

[Bor69] A. Borel, *Introduction aux groupes arithmétiques*, Publications de l’Institut de Mathématique de l’Université de Strasbourg, XV. Actualités Scientifiques et Industrielles, No. 1341, Hermann, Paris, 1969.

[Bor91] ———, *Linear algebraic groups*, second ed., Graduate Texts in Mathematics, vol. 126, Springer-Verlag, New York, 1991.

[BT65] A. Borel and J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. (1965), no. 27, 55–150.

[Cas78] J. W. S. Cassels, *Rational quadratic forms*, London Mathematical Society Monographs, vol. 13, Academic Press, Inc., London-New York, 1978.

[Che58] C. Chevalley, *Classification des groupes de Lie algébriques*, vol. 1, Secrétariat mathématique, Paris, 1958, Séminaire C. Chevalley 1956–1958.

[Del79] P. Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, Automorphic forms, representations and L -functions (Part 2), Proc. Sympos. Pure Math., XXXIII, AMS, Providence, R.I., 1979, pp. 247–289.

[Del82] ———, *Hodge cycles on abelian varieties*, Hodge cycles, motives, and Shimura varieties, Lecture Notes in Mathematics, vol. 900, Springer-Verlag, 1982, Notes by J. S. Milne, pp. 9–100.

[DO] C. Daw and M. Orr, *Unlikely intersections with $E \times CM$ curves in \mathcal{A}_2* , preprint, available at arXiv:1902.10483.

[DR18] C. Daw and J. Ren, *Applications of the hyperbolic Ax–Schanuel conjecture*, Compos. Math. **154** (2018), 1843–1888.

- [Ebe14] Patrick Eberlein, *Growth estimates for orbits of self-adjoint groups*, *Geom. Dedicata* **170** (2014), 87–117.
- [Fal83] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, *Invent. Math.* **73** (1983), no. 3, 349–366.
- [GR14] É. Gaudron and G. Rémond, *Polarisations et isogénies*, *Duke Math. J.* **163** (2014), no. 11, 2057–2108.
- [GS80] F. Grunewald and D. Segal, *Some general algorithms. I. Arithmetic groups*, *Ann. of Math. (2)* **112** (1980), no. 3, 531–583.
- [HP16] P. Habegger and J. Pila, *O-minimality and certain atypical intersections*, *Ann. Sci. Éc. Norm. Supér. (4)* **49** (2016), no. 4, 813–858.
- [KMRT98] M.-A. Knus, A. Merkurjev, M. Rost, and J.-P. Tignol, *The book of involutions*, American Mathematical Society Colloquium Publications, vol. 44, American Mathematical Society, Providence, RI, 1998.
- [KUY16] B. Klingler, E. Ullmo, and A. Yafaev, *The hyperbolic Ax-Lindemann-Weierstrass conjecture*, *Publications mathématiques de l’IHÉS* **123** (2016), no. 1, 333–360.
- [LM16] H. Li and G. A. Margulis, *Effective estimates on integral quadratic forms: Masser’s conjecture, generators of orthogonal groups, and bounds in reduction theory*, *Geom. Funct. Anal.* **26** (2016), no. 3, 874–908.
- [Mil86] J. S. Milne, *Abelian varieties*, Arithmetic geometry (Storrs, Conn., 1984), Springer, New York, 1986, pp. 103–150.
- [Mil05] ———, *Introduction to Shimura varieties*, Harmonic analysis, the trace formula, and Shimura varieties, *Clay Math. Proc.*, vol. 4, Amer. Math. Soc., Providence, RI, 2005, pp. 265–378.
- [Mos55] G. D. Mostow, *Self-adjoint groups*, *Ann. of Math. (2)* **62** (1955), 44–55.
- [Mum74] D. Mumford, *Abelian varieties*, second ed., Oxford University Press, 1974.
- [MW94] D. W. Masser and G. Wüstholz, *Endomorphism estimates for abelian varieties*, *Math. Z.* **215** (1994), no. 4, 641–653. MR 1269495
- [Orr] M. Orr, *Unlikely intersections with Hecke translates of a special subvariety*, to appear in *J. Eur. Math. Soc.*, available at arXiv:1710.04092v2.
- [Orr15] ———, *Families of abelian varieties with many isogenous fibres*, *J. Reine Angew. Math.* **705** (2015), 211–231.
- [Orr17] ———, *On compatibility between isogenies and polarizations of abelian varieties*, *Int. J. Number Theory* **13** (2017), no. 3, 673–704.
- [Orr18] ———, *Height bounds and the Siegel property*, *Algebra & Number Theory* **12** (2018), no. 2, 455–478.
- [PR94] V. Platonov and A. Rapinchuk, *Algebraic groups and number theory*, Pure and Applied Mathematics, vol. 139, Academic Press, Inc., Boston, MA, 1994, Translated from the 1991 Russian original by R. Rowen.
- [PT13] J. Pila and J. Tsimerman, *The André–Oort conjecture for the moduli space of abelian surfaces*, *Compos. Math.* **149** (2013), no. 2, 204–216.
- [RS90] R. W. Richardson and P. J. Slodowy, *Minimum vectors for real reductive algebraic groups*, *J. London Math. Soc. (2)* **42** (1990), no. 3, 409–429.
- [Sch85] W. Scharlau, *Quadratic and Hermitian forms*, Grundlehren der Mathematischen Wissenschaften, vol. 270, Springer-Verlag, Berlin, 1985.
- [Sie40] C. L. Siegel, *Einheiten quadratischer Formen*, *Abh. Math. Sem. Hansischen Univ.* **13** (1940), 209–239.
- [SO85] W. Scharlau and H. Opolka, *From Fermat to Minkowski: Lectures on the theory of numbers and its historical development*, Undergraduate Texts in Mathematics, Springer-Verlag, 1985, Translated from the German by W. K. Bühler and G. Cornell.

- [UY18] Emmanuel Ullmo and Andrei Yafaev, *Algebraic flows on Shimura varieties*, *Manuscripta Math.* **155** (2018), no. 3-4, 355–367. MR 3763410
- [Wey40] H. Weyl, *Theory of reduction for arithmetical equivalence*, *Trans. Amer. Math. Soc.* **48** (1940), 126–164.

DAW: DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF READING, WHITE-KNIGHTS, PO BOX 217, READING, BERKSHIRE RG6 6AH, UNITED KINGDOM
E-mail address: `chris.daw@reading.ac.uk`

ORR: MATHEMATICS INSTITUTE, ZEEMAN BUILDING, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UNITED KINGDOM
E-mail address: `martin.orr@warwick.ac.uk`