

**ON THE RELATIVIZED ALON SECOND EIGENVALUE  
CONJECTURE III: ASYMPTOTIC EXPANSIONS FOR  
TANGLE-FREE HASHIMOTO TRACES**

JOEL FRIEDMAN AND DAVID KOHLER

ABSTRACT. This is the third in a series of articles devoted to showing that a typical covering map of large degree to a fixed, regular graph has its new adjacency eigenvalues within the bound conjectured by Alon for random regular graphs.

In this paper we consider random graphs that are random covering graphs of large degree  $n$  of a fixed base graph. We prove the existence of asymptotic expansion in  $1/n$  for the expected value of the number of strictly non-backtracking closed walks of length  $k$  times the indicator function that the graph is free of certain *tangles*; moreover, we prove that the coefficients of these expansions are “nice functions” of  $k$ , namely approximately equal to a sum of polynomials in  $k$  times exponential functions of  $k$ .

Our results use the methods of Friedman used to resolve Alon’s original conjecture, combined with the results of Article II in this series of articles. One simplification in this article over the previous methods of Friedman is that the “regularized traces” used in this article, which we call *certified traces*, are far easier to define and work with than the previously utilized *selective traces*.

CONTENTS

1. Introduction	2
2. Review of the Main Definitions	4
2.1. Basic Notation and Conventions	4
2.2. Graphs, Our Basic Models, Walks	5
2.3. Asymptotic Expansions	7
2.4. Tangles	8
2.5. $B$ -Graphs, Ordered Graphs, and Strongly Algebraic Models	8
2.6. Homotopy Type	10
2.7. $B$ -graphs and Wordings	11
2.8. Algebraic Models	11
2.9. SNBC Counting Functions	12
3. Main Theorems in This Article	13
4. Theorems From Article II	14
5. Certified Traces and Theorem 3.1	15
5.1. Motivation for Modified Traces	15
5.2. Definition of Certified Traces	15
5.3. Statement of the Expansion Theorems for Certified Traces	16

---

*Date:* Thursday 14<sup>th</sup> November, 2019.

2010 *Mathematics Subject Classification.* Primary 68R10.

Research supported in part by an NSERC grant.

Research supported in part by an NSERC grant.

5.4.	Generalizations of Tangle Free Sets	17
5.5.	Main Theorem for Indicator Functions	17
5.6.	Main Theorems for Certified Traces with Indicator Functions	18
5.7.	Remarks on Theorems 5.8 and 5.9 and Isomorphism Classes of $B$ -Graphs	18
6.	Proof of Theorem 5.2	19
6.1.	VLG Comparisons	19
6.2.	The Finiteness of Minimal Elements in an Upper Subset of $\mathbb{N}^n$	19
6.3.	The Zeroth Order Coefficient	20
6.4.	Conclusion of The Proof of Theorem 5.2	21
7.	Finiteness of Minimal Tangles	23
8.	Indicator Function Approximation	26
8.1.	The Main Results	26
8.2.	Proof of Lemma 8.4	27
8.3.	Proof of Lemma 8.5	28
8.4.	The Injection Count and Resulting Partial Order	29
8.5.	The Möbius Function	29
8.6.	The Truncated Indicator Function and the Proof of Theorem 8.6	30
9.	Proof of Theorem 5.9	31
10.	Proof of Theorem 5.8	33
11.	Conclusion of the Proofs of Theorems 3.1 and 3.2	34
	References	36

## 1. INTRODUCTION

This is the third article in a series of six devoted to proving two main results of a generalization of Alon’s second eigenvalue conjecture.

Alon’s original conjecture ([Alo86], Conjecture 5.1) is that for fixed integer  $d$ , and real  $\epsilon > 0$ , the second adjacency matrix eigenvalue of “most”  $d$ -regular graphs on  $n$  vertices is at most  $2\sqrt{d-1} + \epsilon$ . Graphs with so small a second eigenvalue provably “expand” sets of vertices of certain sizes when passing to their set of neighbours (e.g., Section 1 of [Alo86] and also [AM85, Tan84, Dod84]); see [HLW06] for many other applications of expanders and the explicit construction of expanders [LPS88, Mar88, Mor94]. We do not know the motivation for Alon’s conjecture, although Section 4 of [Alo86] states the result of Alon and Boppana [Nil91] that  $2\sqrt{d-1} + o_n(1)$  is a lower bound for all  $d$ -regular graphs on  $n$  vertices. Alon’s conjecture with weaker upper bounds appears in [BS87, FKS89, Fri91], and was finally proven in [Fri08]. Both the counting argument of Kahn and Szemerédi [FKS89], and the trace methods of Broder [BS87] and Friedman [Fri91, Fri08] give the same lower bound on the absolute value of all the negative eigenvalues.

In this series of articles we study a generalization of Alon’s conjecture, where one fixes a “base” graph  $B$ , and studies random covering maps of degree  $n$  to  $B$ . The *new eigenvalues* [Fri93a, Fri03] of the adjacency matrix of the covering graph are those not arising from (and therefore orthogonal to) the pullback of eigenfunctions on  $B$ ; [Fri03] Section 5 conjectures that the largest absolute of the new eigenvalues of “most” such covering maps is  $\rho(B) + o_n(1)$  where  $\rho(B)$  is the spectral radius of the adjacency matrix of the universal cover of  $B$  (and  $\rho(B) =$

$2\sqrt{d-1}$  if  $B$  is  $d$ -regular). This conjecture generalizes Alon’s conjecture, which is the special case where  $B$  consists of one vertex. This conjecture was proven with a weaker upper bound in [Fri03], by adapting the trace methods of Broder-Shamir [BS87] to this more general situation; counting methods improved these results in [LSV11, ABG10] for regular base graphs, as did trace methods [LP10, Pud15]. The conjecture was proven when  $B$  is a regular graph in [FK14], and with a much simpler type of trace method, but involving more sophisticated probabilistic methods, by Bordenavé [Bor15]. Recently Bordenavé and Collins [BC19] have proven spectral bounds for a large class of sums of random permutation matrices, and their result resolves the entire relativized Alon conjecture, where the base graph need not be regular. The point of this series of articles is to “factor” the proof in [FK14] into independent parts; this factorization includes a number of significant simplifications to the methods in [Fri91, Fri08].

The above generalization of Alon’s conjecture is also a *relativization* of this conjecture (now a theorem) in the sense that one associates with Grothendieck, in that this generalization is a theorem regarding covering morphisms in categories where the special case regarding morphisms to a final element of the category (i.e. graphs with one vertex) specialize to the original conjecture (which is viewed as a conjecture regarding objects in these categories).

The study of relative expanders of fixed degree two [BL06] have lead to a number of exciting recent results [MSS15a, MSS15b], proving the existence of relatively Ramanujan bipartite of all degrees and all even number of vertices; see also [HPS18].

Let us briefly describe the main technical result of this article, in rough terms.

In this article we fix a graph  $B$  (not necessarily regular) and consider a family  $\{\mathcal{C}_n(B)\}$  of random covering map of degree  $n$  to  $B$  (defined for at least an infinite set of integers,  $n$ ) that is *algebraic* in a sense we will define. The main and most difficult result in this paper concerns the function

$$f(k, n) = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}(G) \text{Trace}(H_G^k)],$$

where  $H_B$  is the Hashimoto matrix (also called the non-backtracking matrix) of  $G$ , and  $\mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}$  is the indicator function of a graph being free of certain *tangles*. We prove that such a function has an asymptotic expansion

$$c_0(k) + c_1(k)/n + \dots + c_{r-1}(k)/n^{r-1} + O_k(1/n^r),$$

for  $k \ll n^{1/2}$ , where  $c_0(k)$  is explicitly computed to within a small error term, and  $c_1, \dots, c_{r-1}$  are certain well-behaved functions of  $k$ . In Article IV we study such a result purely in terms of probability theory to prove that the results in this article can be used to control the eigenvalues of  $H_G$ , which in Article V allows us to prove the relativized Alon conjecture when  $B$  is  $d$ -regular, and in Article VI is used to determine probability that new adjacency eigenvalues exceeding  $2\sqrt{d-1} + \epsilon$  occur in  $\mathcal{C}_n(B)$  to within a multiplicative constant (with depending on  $\epsilon$  but not on  $n$ ) provided that  $B$  is regular and *Ramanujan*.

This article uses the methods of [Fri08] but makes some simplifying modifications, the most significant of which is the introduction of *certified traces* that replace the *selective traces* of [Fri08]. It is not only simpler to define certified traces, but their properties (and proofs using certified traces) can be “factor” into more independent parts.

The theorems in this paper requires the results of Article II. We assume that the reader is familiar with definitions in Article I. For ease of reading, we have

summarized all definitions that we need in Section 2; however, to understand the motivation and the subtleties of these definitions we refer the reader to Article I.

The rest of this article is organized as follows. After reviewing the definitions needed in Section 2, we state the main results in this article in Section 3. We review the theorems needed from Article II in Section 4. In Section 5 we explain our strategy to prove our main theorems, and state two more general theorems from which our main theorems easily follow. In Section 6 we define  $(\geq \nu, < r)$ -tangles and certified traces, and discuss how they are related to traces of the Hashimoto matrix of a graph. In Section 6 we prove an asymptotic expansion theorem regarding certified traces; in the process we prove that each certified trace has a finite number of “certificates,” which is why it is easy to use them in proofs (much easier than the *selective traces* in [Fri08]). In Section 7 we prove that there are a finite number of minimal  $(\geq \nu, < r)$ -tangles for any fixed  $\nu, r$  with  $\nu > 1$ ; this was done in [Fri08], but we correct a minor error there. In Section 8 we define a series of approximations to the indicator function of a graph having  $(\geq \nu, < r)$ -tangles, and give their basic properties. In Sections 9 and Section 10 we prove the generalizations of the main theorems in this paper stated in Section 5, and in Section 11 we prove our main theorems using these generalizations.

## 2. REVIEW OF THE MAIN DEFINITIONS

We refer the reader to Article I for the definitions used in this article, the motivation of such definitions, and an appendix there that lists all the definitions and notation. In this section we briefly review these definitions and notation.

**2.1. Basic Notation and Conventions.** We use  $\mathbb{R}, \mathbb{C}, \mathbb{Z}, \mathbb{N}$  to denote, respectively, the real numbers, the complex numbers, the integers, and positive integers or natural numbers; we use  $\mathbb{Z}_{\geq 0}$  ( $\mathbb{R}_{>0}$ , etc.) to denote the set of non-negative integers (of positive real numbers, etc.). We denote  $\{1, \dots, n\}$  by  $[n]$ .

If  $A$  is a set, we use  $\mathbb{N}^A$  to denote the set of maps  $A \rightarrow \mathbb{N}$ ; we will refer to its elements as *vectors*, denoted in bold face letters, e.g.,  $\mathbf{k} \in \mathbb{N}^A$  or  $\mathbf{k}: A \rightarrow \mathbb{N}$ ; we denote its *component* in the regular face equivalents, i.e., for  $a \in A$ , we use  $k(a) \in \mathbb{N}$  to denote the  $a$ -component of  $\mathbf{k}$ . As usual,  $\mathbb{N}^n$  denotes  $\mathbb{N}^{[n]} = \mathbb{N}^{\{1, \dots, n\}}$ . We use similar conventions for  $\mathbb{N}$  replaced by  $\mathbb{R}, \mathbb{C}$ , etc.

If  $A$  is a set, then  $\#A$  denotes the cardinality of  $A$ . We often denote a set with all capital letters, and its cardinality in lower case letters; for example, when we define  $\text{SNBC}(G, k)$ , we will write  $\text{snbc}(G, k)$  for  $\#\text{SNBC}(G, k)$ .

If  $A' \subset A$  are sets, then  $\mathbb{I}_{A'}: A \rightarrow \{0, 1\}$  (with  $A$  understood) denotes the characteristic function of  $A'$ , i.e.,  $\mathbb{I}_{A'}(a)$  is 1 if  $a \in A'$  and otherwise is 0; we also write  $\mathbb{I}_{A'}$  (with  $A$  understood) to mean  $\mathbb{I}_{A' \cap A}$  when  $A'$  is not necessarily a subset of  $A$ .

All probability spaces are finite; hence a probability space is a pair  $\mathcal{P} = (\Omega, P)$  where  $\Omega$  is a finite set and  $P: \Omega \rightarrow \mathbb{R}_{>0}$  with  $\sum_{\omega \in \Omega} P(\omega) = 1$ ; hence an *event* is any subset of  $\Omega$ . We emphasize that  $\omega \in \Omega$  implies that  $P(\omega) > 0$  with strict inequality; we refer to the elements of  $\Omega$  as the atoms of the probability space. We use  $\mathcal{P}$  and  $\Omega$  interchangeably when  $P$  is understood and confusion is unlikely.

A *complex-valued random variable* on  $\mathcal{P}$  or  $\Omega$  is a function  $f: \Omega \rightarrow \mathbb{C}$ , and similarly for real-, integer-, and natural-valued random variable; we denote its  $\mathcal{P}$ -expected value by

$$\mathbb{E}_{\omega \in \Omega}[f(\omega)] = \sum_{\omega \in \Omega} f(\omega)P(\omega).$$

If  $\Omega' \subset \Omega$  we denote the probability of  $\Omega'$  by

$$\text{Prob}_{\mathcal{P}}[\Omega'] = \sum_{\omega \in \Omega'} P(\omega') = \mathbb{E}_{\omega \in \Omega}[\mathbb{I}_{\Omega'}(\omega)].$$

At times we write  $\text{Prob}_{\mathcal{P}}[\Omega']$  where  $\Omega'$  is not a subset of  $\Omega$ , by which we mean  $\text{Prob}_{\mathcal{P}}[\Omega' \cap \Omega]$ .

**2.2. Graphs, Our Basic Models, Walks.** A *directed graph*, or simply a *digraph*, is a tuple  $G = (V_G, E_G^{\text{dir}}, h_G, t_G)$  consisting of sets  $V_G$  and  $E_G^{\text{dir}}$  (of *vertices* and *directed edges*) and maps  $h_G, t_G$  (*heads* and *tails*)  $E_G^{\text{dir}} \rightarrow V_G$ . Therefore our digraphs can have multiple edges and self-loops (i.e.,  $e \in E_G^{\text{dir}}$  with  $h_G(e) = t_G(e)$ ). A *graph* is a tuple  $G = (V_G, E_G^{\text{dir}}, h_G, t_G, \iota_G)$  where  $(V_G, E_G^{\text{dir}}, h_G, t_G)$  is a digraph and  $\iota_G: E_G^{\text{dir}} \rightarrow E_G^{\text{dir}}$  is an involution with  $t_G \iota_G = h_G$ ; the *edge set* of  $G$ , denoted  $E_G$ , is the set of orbits of  $\iota_G$ , which (notation aside) can be identified with  $E_G^{\text{dir}} / \iota_G$ , the set of equivalence classes of  $E_G^{\text{dir}}$  modulo  $\iota_G$ ; if  $\{e\} \in E_G$  is a singleton, then necessarily  $e$  is a self-loop with  $\iota_G e = e$ , and we call  $e$  a *half-loop*; other elements of  $E_G$  are sets  $\{e, \iota_G e\}$  of size two, i.e., with  $e \neq \iota_G e$ , and for such  $e$  we say that  $e$  (or, at times,  $\{e, \iota_G e\}$ ) is a *whole-loop* if  $h_G e = t_G e$  (otherwise  $e$  has distinct endpoints).

Hence these definitions allow our graphs to have multiple edges and two types of self-loops—whole-loops and half-loops—as in [Fri93b, Fri08]. The *indegree* and *outdegree* of a vertex in a digraph is the number of edges whose tail, respectively whose head, is the vertex; the *degree* of a vertex in a graph is its indegree (which equals its outdegree) in the underlying digraph; therefore a whole-loop about a vertex contributes 2 to its degree, whereas a half-loop contributes 1.

An *orientation* of a graph,  $G$ , is a choice  $E_G^{\text{or}} \subset E_G^{\text{dir}}$  of  $\iota_G$  representatives; i.e.,  $E_G^{\text{or}}$  contains every half-loop,  $e$ , and one element of each two-element set  $\{e, \iota_G e\}$ .

A *morphism*  $\pi: G \rightarrow H$  of directed graphs is a pair  $\pi = (\pi_V, \pi_E)$  where  $\pi_V: V_G \rightarrow V_H$  and  $\pi_E: E_G^{\text{dir}} \rightarrow E_H^{\text{dir}}$  are maps that intertwine the heads maps and the tails maps of  $G, H$  in the evident fashion; such a morphism is *covering* (respectively, *étale*, elsewhere called an *immersion*) if for each  $v \in V_G$ ,  $\pi_E$  maps those directed edges whose head is  $v$  bijectively (respectively, injectively) to those whose head is  $\pi_V(v)$ , and the same with tail replacing head. If  $G, H$  are graphs, then a morphism  $\pi: G \rightarrow H$  is a morphism of underlying directed graphs where  $\pi_E \iota_G = \iota_H \pi_E$ ;  $\pi$  is called *covering* or *étale* if it is so as a morphism of underlying directed graphs. We use the words *morphism* and *map* interchangeably.

A walk in a graph or digraph,  $G$ , is an alternating sequence  $w = (v_0, e_1, \dots, e_k, v_k)$  of vertices and directed edges with  $t_G e_i = v_{i-1}$  and  $h_G e_i = v_i$  for  $i \in [k]$ ;  $w$  is *closed* if  $v_k = v_0$ ; if  $G$  is a graph,  $w$  is *non-backtracking*, or simply *NB*, if  $\iota_G e_i \neq e_{i+1}$  for  $i \in [k-1]$ , and *strictly non-backtracking closed*, or simply *SNBC*, if it is closed, non-backtracking, and  $\iota_G e_k \neq e_1$ . The *visited subgraph* of a walk,  $w$ , in a graph  $G$ , denoted  $\text{VisSub}_G(w)$  or simply  $\text{VisSub}(w)$ , is the smallest subgraph of  $G$  containing all the vertices and directed edges of  $w$ ;  $\text{VisSub}_G(w)$  generally depends on  $G$ , i.e.,  $\text{VisSub}_G(w)$  cannot be inferred from the sequence  $v_0, e_1, \dots, e_k, v_k$  alone without knowing  $\iota_G$ .

The adjacency matrix,  $A_G$ , of a graph or digraph,  $G$ , is defined as usual (its  $(v_1, v_2)$ -entry is the number of directed edges from  $v_1$  to  $v_2$ ); if  $G$  is a graph on  $n$  vertices, then  $A_G$  is symmetric and we order its eigenvalues (counted with multiplicities) and denote them

$$\lambda_1(G) \geq \cdots \geq \lambda_n(G).$$

If  $G$  is a graph, its Hashimoto matrix (also called the non-backtracking matrix),  $H_G$ , is the adjacency matrix of the *oriented line graph* of  $G$ ,  $\text{Line}(G)$ , whose vertices are  $E_G^{\text{dir}}$  and whose directed edges are the subset of  $E_G^{\text{dir}} \times E_G^{\text{dir}}$  consisting of pairs  $(e_1, e_2)$  such that  $e_1, e_2$  form the directed edges of a non-backtracking walk (of length two) in  $G$  (the tail of  $(e_1, e_2)$  is  $e_1$ , and its head  $e_2$ ); therefore  $H_G$  is the square matrix indexed on  $E_G^{\text{dir}}$ , whose  $(e_1, e_2)$  entry is 1 or 0 according to, respectively, whether or not  $e_1, e_2$  form a non-backtracking walk (i.e.,  $h_G e_1 = t_G e_2$  and  $\iota_G e_1 \neq e_2$ ). We use  $\mu_1(G)$  to denote the Perron-Frobenius eigenvalue of  $H_G$ , and use  $\mu_i(G)$  with  $1 < i \leq \#E_G^{\text{dir}}$  to denote the other eigenvalues of  $H_G$  (which are generally complex-valued) in any order.

If  $B, G$  are both digraphs, we say that  $G$  is a *coordinatized graph over  $B$  of degree  $n$*  if

$$(1) \quad V_G = V_B \times [n], \quad E_G^{\text{dir}} = E_B^{\text{dir}} \times [n], \quad t_G(e, i) = (t_B e, i), \quad h_G(e, i) = (h_B e, \sigma(e)i)$$

for some map  $\sigma: E_B^{\text{dir}} \rightarrow \mathcal{S}_n$ , where  $\mathcal{S}_n$  is the group of permutations on  $[n]$ ; we call  $\sigma$  (which is uniquely determined by (1)) *the permutation assignment associated to  $G$* . [Any such  $G$  comes with a map  $G \rightarrow B$  given by “projection to the first component of the pair,” and this map is a covering map of degree  $n$ .] If  $B, G$  are graphs, we say that a graph  $G$  is a *coordinatized graph over  $B$  of degree  $n$*  if (1) holds and also

$$(2) \quad \iota_G(e, i) = (\iota_B e, \sigma(e)i),$$

which implies that

$$(3) \quad (e, i) = \iota_G \iota_G(e, i) = (e, \sigma(\iota_B e)\sigma(e)i) \quad \forall e \in E_B^{\text{dir}}, i \in [n],$$

and hence  $\sigma(\iota_B e) = \sigma(e)^{-1}$ ; we use  $\text{Coord}_n(B)$  to denote the set of all coordinatized covers of a graph,  $B$ , of degree  $n$ .

The *order* of a graph,  $G$ , is  $\text{ord}(G) \stackrel{\text{def}}{=} (\#E_G) - (\#V_G)$ . Note that a half-loop and a whole-loop each contribute 1 to  $\#E_G$  and to the order of  $G$ . The *Euler characteristic* of a graph,  $G$ , is  $\chi(G) \stackrel{\text{def}}{=} (\#V_G) - (\#E_G^{\text{dir}})/2$ . Hence  $\text{ord}(G) \geq -\chi(G)$ , with equality iff  $G$  has no half-loops.

If  $w$  is a walk in any  $G \in \text{Coord}_n(B)$ , then one easily sees that  $\text{VisSub}_G(w)$  can be inferred from  $B$  and  $w$  alone.

If  $B$  is a graph without half-loops, then the *permutation model over  $B$*  refers to the probability spaces  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  where the atoms of  $\mathcal{C}_n(B)$  are coordinatized coverings of degree  $n$  over  $B$  chosen with the uniform distribution. More generally, a *model* over a graph,  $B$ , is a collection of probability spaces,  $\{\mathcal{C}_n(B)\}_{n \in N}$ , defined for  $n \in N$  where  $N \subset \mathbb{N}$  is an infinite subset, and where the atoms of each  $\mathcal{C}_n(B)$  are elements of  $\text{Coord}_n(B)$ . There are a number of models related to the permutation model, which are generalizations of the models of [Fri08], that we call *our basic models* and are defined in Article I; let us give a rough description.

All of *our basic models* are *edge independent*, meaning that for any orientation  $E_B^{\text{or}} \subset E_B^{\text{dir}}$ , the values of the permutation assignment,  $\sigma$ , on  $E_B^{\text{or}}$  are independent

of one another (of course,  $\sigma(\iota_G e) = (\sigma(e))^{-1}$ , so  $\sigma$  is determined by its values on any orientation  $E_B^{\text{or}}$ ); for edge independent models, it suffices to specify the ( $\mathcal{S}_n$ -valued) random variable  $\sigma(e)$  for each  $e \in E_B^{\text{or}}$  or  $E_B^{\text{dir}}$ . The permutation model can be alternatively described as the edge independent model that assigns a uniformly chosen permutation to each  $e \in E_B^{\text{dir}}$  (which requires  $B$  to have no half-loops); the *full cycle* (or simply *cyclic*) model is the same, except that if  $e$  is a whole-loop then  $\sigma(e)$  is chosen uniformly among all permutations whose cyclic structure consists of a single  $n$ -cycle. If  $B$  has half-loops, then we restrict  $\mathcal{C}_n(B)$  either to  $n$  even or  $n$  odd and for each half-loop  $e \in E_B^{\text{dir}}$  we choose  $\sigma(e)$  as follows: if  $n$  is even we choose  $\sigma(e)$  uniformly among all perfect matchings, i.e., involutions (maps equal to their inverse) with no fixed points; if  $n$  is odd then we choose  $\sigma(e)$  uniformly among all *nearly perfect matchings*, meaning involutions with one fixed point. We combine terms when  $B$  has half-loops: for example, the term *full cycle-involution* (or simply *cyclic-involution*) *model of odd degree over  $B$*  refers to the model where the degree,  $n$ , is odd, where  $\sigma(e)$  follows the full cycle rule when  $e$  is not a half-loop, and where  $\sigma(e)$  is a near perfect matching when  $e$  is a half-loop; similarly for the *full cycle-involution* (or simply *cyclic-involution*) *model of even degree* and the *permutation-involution model of even degree* or *of odd degree*.

If  $B$  is a graph, then a model,  $\{\mathcal{C}_n(B)\}_{n \in N}$ , over  $B$  may well have  $N \neq \mathbb{N}$  (e.g., our basic models above when  $B$  has half-loops); in this case many formulas involving the variable  $n$  are only defined for  $n \in N$ . For brevity, we often do not explicitly write  $n \in N$  in such formulas; for example we usually write

$$\lim_{n \rightarrow \infty} \quad \text{to abbreviate} \quad \lim_{n \in N, n \rightarrow \infty} .$$

Also we often write simply  $\mathcal{C}_n(B)$  or  $\{\mathcal{C}_n(B)\}$  for  $\{\mathcal{C}_n(B)\}_{n \in N}$  if confusion is unlikely to occur.

A graph is *pruned* if all its vertices are of degree at least two (this differs from the more standard definition of *pruned* meaning that there are no leaves). If  $w$  is any SNBC walk in a graph,  $G$ , then we easily see that  $\text{VisSub}_G(w)$  is necessarily pruned: i.e., any of its vertices must be incident upon a whole-loop or two distinct edges [note that a walk of length  $k = 1$  about a half-loop,  $(v_0, e_1, v_1)$ , by definition, is not SNBC since  $\iota_G e_k = e_1$ ]. It easily follows that  $\text{VisSub}_G(w)$  is contained in the graph obtained from  $G$  by repeatedly “pruning any leaves” (i.e., discarding any vertex of degree one and its incident edge) from  $G$ . Since our trace methods only concern (Hashimoto matrices and) SNBC walks, it suffices to work with models  $\mathcal{C}_n(B)$  where  $B$  is pruned. It is not hard to see that if  $B$  is pruned and connected, then  $\text{ord}(B) = 0$  iff  $B$  is a cycle, and  $\mu_1(B) > 1$  iff  $\chi(B) < 0$ ; this is formally proven in Article III (Lemma 6.4). Our theorems are not usually interesting unless  $\mu_1(B) > \mu_1^{1/2}(B)$ , so we tend to restrict our main theorems to the case  $\mu_1(B) > 1$  or, equivalently,  $\chi(B) < 0$ ; some of our techniques work without these restrictions.

**2.3. Asymptotic Expansions.** A function  $f: \mathbb{N} \rightarrow \mathbb{C}$  is a *polyexponential* if it is a sum of functions  $p(k)\mu^k$ , where  $p$  is a polynomial and  $\mu \in \mathbb{C}$ , with the convention that for  $\mu = 0$  we understand  $p(k)\mu^k$  to mean any function that vanishes for sufficiently large  $k$ <sup>1</sup>; we refer to the  $\mu$  needed to express  $f$  as the *exponents* or *bases*

<sup>1</sup> This convention is used because then for any fixed matrix,  $M$ , any entry of  $M^k$ , as a function of  $k$ , is a polyexponential function of  $k$ ; more specifically, the  $\mu = 0$  convention is due to the fact that a Jordan block of eigenvalue 0 is nilpotent.

of  $f$ . A function  $f: \mathbb{N} \rightarrow \mathbb{C}$  is of growth  $\rho$  for a  $\rho \in \mathbb{R}$  if  $|f(k)| = o(1)(\rho + \epsilon)^k$  for any  $\epsilon > 0$ . A function  $f: \mathbb{N} \rightarrow \mathbb{C}$  is  $(B, \nu)$ -bounded if it is the sum of a function of growth  $\nu$  plus a polyexponential function whose bases are bounded by  $\mu_1(B)$  (the Perron-Frobenius eigenvalue of  $H_B$ ); the *larger bases* of  $f$  (with respect to  $\nu$ ) are those bases of the polyexponential function that are larger in absolute value than  $\nu$ . Moreover, such an  $f$  is called  $(B, \nu)$ -Ramanujan if its larger bases are all eigenvalues of  $H_B$ .

We say that a function  $f = f(k, n)$  taking some subset of  $\mathbb{N}^2$  to  $\mathbb{C}$  has a  $(B, \nu)$ -bounded expansion of order  $r$  if for some constant  $C$  we have

$$(4) \quad f(k, n) = c_0(k) + \cdots + c_{r-1}(k) + O(1)c_r(k)/n^r,$$

whenever  $f(k, n)$  is defined and  $1 \leq k \leq n^{1/2}/C$ , where for  $0 \leq i \leq r-1$ , the  $c_i(k)$  are  $(B, \nu)$ -bounded and  $c_r(k)$  is of growth  $\mu_1(B)$ . Furthermore, such an expansion is called  $(B, \nu)$ -Ramanujan if for  $0 \leq i \leq r-1$ , the  $c_i(k)$  are  $(B, \nu)$ -Ramanujan.

Typically our functions  $f(k, n)$  as in (4) are defined for all  $k \in \mathbb{N}$  and  $n \in N$  for an infinite set  $N \subset \mathbb{N}$  representing the possible degrees of our random covering maps in the model  $\{\mathcal{C}_n(B)\}_{n \in N}$  at hand.

**2.4. Tangles.** A  $(\geq \nu)$ -tangle is any connected graph,  $\psi$ , with  $\mu_1(\psi) \geq \nu$ , where  $\mu_1(\psi)$  denotes the Perron-Frobenius eigenvalue of  $H_B$ ; a  $(\geq \nu, < r)$ -tangle is any  $(\geq \nu)$ -tangle of order less than  $r$ ; similarly for  $(> \nu)$ -tangles, i.e.,  $\psi$  satisfying the weak inequality  $\mu_1(\psi) > \nu$ , and for  $(> \nu, r)$ -tangles. We use  $\text{TangleFree}(\geq \nu, < r)$  to denote those graphs that don't contain a subgraph that is  $(\geq \nu, < r)$ -tangle, and  $\text{HasTangles}(\geq \nu, < r)$  for those that do; we never use  $(> \nu)$ -tangles in defining  $\text{TangleFree}$  and  $\text{HasTangles}$ , for the technical reason (see Article III or Lemma 9.2 of [Fri08]) that for  $\nu > 1$  and any  $r \in \mathbb{N}$  that there are only finitely many  $(\geq \nu, < r)$ -tangles, up to isomorphism, that are minimal with respect to inclusion<sup>2</sup>.

**2.5. B-Graphs, Ordered Graphs, and Strongly Algebraic Models.** An *ordered graph*,  $G^\leq$ , is a graph,  $G$ , endowed with an *ordering*, meaning an orientation (i.e.,  $\iota_G$ -orbit representatives),  $E_G^{\text{or}} \subset E_G^{\text{dir}}$ , and total orderings of  $V_G$  and  $E_G$ ; a walk,  $w = (v_0, \dots, e_k, v_k)$  in a graph endows  $\text{VisSub}(w)$  with a *first-encountered* ordering: namely,  $v \leq v'$  if the first occurrence of  $v$  comes before that of  $v'$  in the sequence  $v_0, v_1, \dots, v_k$ , similarly for  $e \leq e'$ , and we orient each edge in the order in which it is first traversed (some edges may be traversed in only one direction). We use  $\text{VisSub}^\leq(w)$  to refer to  $\text{VisSub}(w)$  with this ordering.

A *morphism*  $G^\leq \rightarrow H^\leq$  of ordered graphs is a morphism  $G \rightarrow H$  that respects the ordering in the evident fashion. We are mostly interested in *isomorphisms* of ordered graphs; we easily see that any isomorphism  $G^\leq \rightarrow G^\leq$  must be the identity morphism; it follows that if  $G^\leq$  and  $H^\leq$  are isomorphic, then there is a unique isomorphism  $G^\leq \rightarrow H^\leq$ .

If  $B$  is a graph, then a  $B$ -graph,  $G_{/B}$ , is a graph  $G$  endowed with a map  $G \rightarrow B$  (its  $B$ -graph structure). A *morphism*  $G_{/B} \rightarrow H_{/B}$  of  $B$ -graphs is a morphism  $G \rightarrow H$  that respects the  $B$ -structures in the evident sense. An *ordered B-graph*,  $G_{/B}^\leq$ , is a graph

<sup>2</sup> By contrast, there are infinitely many minimal  $(> \nu, < r)$ -tangles for some values of  $\nu > 1$  and  $r$ : indeed, consider any connected pruned graph  $\psi$ , and set  $r = \text{ord}(\psi) + 2$ ,  $\nu = \mu_1(\psi)$ . Then if we fix two vertices in  $\psi$  and let  $\psi_s$  be the graph that is  $\psi$  with an additional edge of length  $s$  between these two vertices, then  $\psi_s$  is an  $(> \nu, < r)$ -tangle. However, if  $\psi'$  is  $\psi$  with any single edge deleted, and  $\psi'_s$  is  $\psi_s$  with this edge deleted, then one can show that  $\mu_1(\psi'_s) < \nu$  for  $s$  sufficiently large. It follows that for  $s$  sufficiently large,  $\psi_s$  are minimal  $(> \nu, < r)$ -tangles.



endowed with both an ordering and a  $B$ -graph structure; a morphism of ordered  $B$ -graphs is a morphism of the underlying graphs that respects both the ordering and  $B$ -graph structures. If  $w$  is a walk in a  $B$ -graph,  $G_{/B}$ , we use  $\text{VisSub}_{/B}(w)$  to denote  $\text{VisSub}(w)$  with the  $B$ -graph structure it inherits from  $G$  in the evident sense; we use  $\text{VisSub}_{/B}^{\leq}(w)$  to denote  $\text{VisSub}_{/B}(w)$  with its first-encountered ordering.

At times we drop the superscript  $\leq$  and the subscript  $/B$ ; for example, we write  $G \in \text{Coord}_n(B)$  instead of  $G_{/B} \in \mathcal{C}_n(B)$  (despite the fact that we constantly utilize the  $B$ -graph structure on elements of  $\text{Coord}_n(B)$ ).

A  $B$ -graph  $G_{/B}$  is *covering* or *étale* if its structure map  $G \rightarrow B$  is.

If  $\pi: S \rightarrow B$  is a  $B$ -graph, we use  $\mathbf{a} = \mathbf{a}_{S_{/B}}$  to denote the vector  $E_B^{\text{dir}} \rightarrow \mathbb{Z}_{\geq 0}$  given by  $a_{S_{/B}}(e) = \#\pi^{-1}(e)$ ; since  $a_{S_{/B}}(\iota_B e) = a_{S_{/B}}(e)$  for all  $e \in E_B^{\text{dir}}$ , we sometimes view  $\mathbf{a}$  as a function  $E_B \rightarrow \mathbb{Z}_{\geq 0}$ , i.e., as the function taking  $\{e, \iota_B e\}$  to  $a_{S_{/B}}(e) = a_{S_{/B}}(\iota_B e)$ . We similarly define  $\mathbf{b}_{S_{/B}}: V_B \rightarrow \mathbb{Z}_{\geq 0}$  by setting  $b_{S_{/B}}(v) = \#\pi^{-1}(v)$ . If  $w$  is a walk in a  $B$ -graph, we set  $\mathbf{a}_w$  to be  $\mathbf{a}_{S_{/B}}$  where  $S_{/B} = \text{VisSub}_{/B}(w)$ , and similarly for  $\mathbf{b}_w$ . We refer to  $\mathbf{a}, \mathbf{b}$  (in either context) as  *$B$ -fibre counting functions*.

If  $S_{/B}^{\leq}$  is an ordered  $B$ -graph and  $G_{/B}$  is a  $B$ -graph, we use  $[S_{/B}^{\leq}] \cap G_{/B}$  to denote the set of ordered graphs  $G'_{/B}^{\leq}$  such that  $G'_{/B} \subset G_{/B}$  and  $G'_{/B}^{\leq} \simeq S_{/B}^{\leq}$  (as ordered  $B$ -graphs); this set is naturally identified with the set of injective morphisms  $S_{/B} \rightarrow G_{/B}$ , and the cardinality of these sets is independent of the ordering on  $S_{/B}^{\leq}$ .

A  $B$ -graph,  $S_{/B}$ , or an ordered  $B$ -graph,  $S_{/B}^{\leq}$ , *occurs in a model*  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  if for all sufficiently large  $n \in \mathbb{N}$ ,  $S_{/B}$  is isomorphic to a  $B$ -subgraph of some element of  $\mathcal{C}_n(B)$ ; similarly a graph,  $S$ , *occurs in*  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  if it can be endowed with a  $B$ -graph structure,  $S_{/B}$ , that occurs in  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$ .

A model  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  of coverings of  $B$  is *strongly algebraic* if

- (1) for each  $r \in \mathbb{N}$  there is a function,  $g = g(k)$ , of growth  $\mu_1(B)$  such that if  $k \leq n/4$  we have

$$(5) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{snbc}_{\geq r}(G, k)] \leq g(k)/n^r$$

where  $\text{snbc}_{\geq r}(G, k)$  is the number of SNBC walks of length  $k$  in  $G$  whose visited subgraph is of order at least  $r$ ;

- (2) for any  $r$  there exists a function  $g$  of growth 1 and real  $C > 0$  such that the following holds: for any ordered  $B$ -graph,  $S_{/B}^{\leq}$ , that is pruned and of order less than  $r$ ,

- (a) if  $S_{/B}$  occurs in  $\mathcal{C}_n(B)$ , then for  $1 \leq \#E_S^{\text{dir}} \leq n^{1/2}/C$ ,

$$(6) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#([S_{/B}^{\leq}] \cap G) \right] = c_0 + \cdots + c_{r-1}/n^{r-1} + O(1)g(\#E_S)/n^r$$

where the  $O(1)$  term is bounded in absolute value by  $C$  (and therefore independent of  $n$  and  $S_{/B}$ ), and where  $c_i = c_i(S_{/B}) \in \mathbb{R}$  such that  $c_i$  is 0 if  $i < \text{ord}(S)$  and  $c_i > 0$  for  $i = \text{ord}(S)$ ; and

- (b) if  $S_{/B}$  does not occur in  $\mathcal{C}_n(B)$ , then for any  $n$  with  $\#E_S^{\text{dir}} \leq n^{1/2}/C$ ,

$$(7) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#([S_{/B}^{\leq}] \cap G) \right] = 0$$

(or, equivalently, no graph in  $\mathcal{C}_n(B)$  has a  $B$ -subgraph isomorphic to  $S_{/B}^{\leq}$ );

- (3)  $c_0 = c_0(S_{/B})$  equals 1 if  $S$  is a cycle (i.e.,  $\text{ord}(S) = 0$  and  $S$  is connected) that occurs in  $\mathcal{C}_n(B)$ ;
- (4)  $S_{/B}$  occurs in  $\mathcal{C}_n(B)$  iff  $S_{/B}$  is an étale  $B$ -graph and  $S$  has no half-loops; and

- (5) there exist polynomials  $p_i = p_i(\mathbf{a}, \mathbf{b})$  such that  $p_0 = 1$  (i.e., identically 1), and for every étale  $B$ -graph,  $S_{/B}^{\leq}$ , we have that

$$(8) \quad c_{\text{ord}(S)+i}(S_{/B}) = p_i(\mathbf{a}_{S_{/B}}, \mathbf{b}_{S_{/B}}) .$$

Notice that condition (3), regarding  $S$  that are cycles, is implied by conditions (4) and (5); we leave in condition (3) since this makes the definition of *algebraic* (below) simpler. Notice that (6) and (8) are the main reasons that we work with ordered  $B$ -graphs: indeed, the coefficients depend only on the  $B$ -fibre counting function  $\mathbf{a}, \mathbf{b}$ , which depend on the structure of  $S_{/B}^{\leq}$  as a  $B$ -graph; this is not true if we don't work with ordered graphs: i.e., (6) fails to hold if we replace  $[S_{/B}^{\leq}]$  with  $[S_{/B}]$  (when  $S_{/B}$  has nontrivial automorphisms), where  $[S_{/B}] \cap G$  refers to the number of  $B$ -subgraphs of  $G$  isomorphic to  $S_{/B}$ ; the reason is that

$$\#[S_{/B}^{\leq}] \cap G_{/B} = (\#\text{Aut}(S_{/B}))(\#[S_{/B}] \cap G_{/B})$$

where  $\text{Aut}(S_{/B})$  is the group of automorphisms of  $S_{/B}$ , and it is  $[S_{/B}^{\leq}] \cap G_{/B}$  rather than  $[S_{/B}] \cap G_{/B}$  that turns out to have the “better” properties; see Section 6 of Article I for examples. Ordered graphs are convenient to use for a number of other reasons.

**2.6. Homotopy Type.** The homotopy type of a walk and of an ordered subgraph are defined by *suppressing* its “uninteresting” vertices of degree two; examples are given in Section 6 of Article I. Here is how we make this precise.

A *bead* in a graph is a vertex of degree two that is not incident upon a self-loop. Let  $S$  be a graph and  $V' \subset V_S$  be a *proper bead subset* of  $V_S$ , meaning that  $V'$  consists only of beads of  $V$ , and that no connected component of  $S$  has all its vertices in  $V'$  (this can only happen for connected components of  $S$  that are cycles); we define the *bead suppression*  $S/V'$  to be the following graph: (1) its vertex set  $V_{S/V'}$  is  $V'' = V_S \setminus V'$ , (2) its directed edges,  $E_{S/V'}^{\text{dir}}$ , consist of the  *$V'$ -beaded paths*, i.e., non-backtracking walks in  $S$  between elements of  $V''$  whose intermediate vertices lie in  $V'$ , (3)  $t_{S/V'}$  and  $h_{S/V'}$  give the first and last vertex of the beaded path, and (4)  $\iota_{S/V'}$  takes a beaded path to its reverse walk (i.e., takes  $(v_0, e_1, \dots, v_k)$  to  $(v_k, \iota_S e_k, \dots, \iota_S e_1, v_0)$ ). One can recover  $S$  from the suppression  $S/V'$  for pedantic reasons, since we have defined its directed edges to be beaded paths of  $S$ . If  $S^{\leq} = \text{VisSub}^{\leq}(w)$  where  $w$  is a non-backtracking walk, then the ordering of  $S$  can be inferred by the naturally corresponding order on  $S/V'$ , and we use  $S^{\leq}/V'$  to denote  $S/V'$  with this ordering.

Let  $w$  be a non-backtracking walk in a graph, and  $S^{\leq} = \text{VisSub}^{\leq}(w)$  its visited subgraph; the *reduction* of  $w$  is the ordered graph,  $R^{\leq}$ , denoted  $S^{\leq}/V'$ , whose underlying graph is  $S/V'$  where  $V'$  is the set of beads of  $S$  except the first and last vertices of  $w$  (if one or both are beads), and whose ordering is naturally arises from that on  $S^{\leq}$ ; the *edge lengths* of  $w$  is the function  $E_{S/V'} \rightarrow \mathbb{N}$  taking an edge of  $S/V'$  to the length of the beaded path it represents in  $S$ ; we say that  $w$  is of *homotopy type*  $T^{\leq}$  for any ordered graph  $T^{\leq}$  that is isomorphic to  $S^{\leq}/V'$ ; in this case the lengths of  $S^{\leq}/V'$  naturally give lengths  $E_T \rightarrow \mathbb{N}$  by the unique isomorphism from  $T^{\leq}$  to  $S^{\leq}/V'$ . If  $S^{\leq}$  is the visited subgraph of a non-backtracking walk, we define the reduction, homotopy type, and edge-lengths of  $S^{\leq}$  to be that of the walk, since these notions depend only on  $S^{\leq}$  and not the particular walk.

If  $T$  is a graph and  $\mathbf{k}: E_T \rightarrow \mathbb{N}$  a function, then we use  $\text{VLG}(T, \mathbf{k})$  (for *variable-length graph*) to denote any graph obtained from  $T$  by gluing in a path of length

$k(e)$  for each  $e \in E_T$ . If  $S^\leq$  is of homotopy type  $T^\leq$  and  $\mathbf{k}: E_T \rightarrow \mathbb{N}$  its edge lengths, then  $\text{VLG}(T, \mathbf{k})$  is isomorphic to  $S$  (as a graph). Hence the construction of variable-length graphs is a sort of inverse to bead suppression.

If  $T^\leq$  is an ordering on  $T$  that arises as the first encountered ordering of a non-backtracking walk on  $T$  (whose visited subgraph is all of  $T$ ), then this ordering gives rise to a natural ordering on  $\text{VLG}(T, \mathbf{k})$  that we denote  $\text{VLG}^\leq(T^\leq, \mathbf{k})$ . Again, this ordering on the variable-length graph is a sort of inverse to bead suppression on ordered graphs.

**2.7.  $B$ -graphs and Wordings.** If  $w_B = (v_0, e_1, \dots, e_k, v_k)$  with  $k \geq 1$  is a walk in a graph  $B$ , then we can identify  $w_B$  with the string  $e_1, e_2, \dots, e_k$  over the alphabet  $E_B^{\text{dir}}$ . For technical reasons, the definitions below of a  $B$ -wording and the *induced wording*, are given as strings over  $E_B^{\text{dir}}$  rather than the full alternating string of vertices and directed edges. The reason is that doing this gives the correct notion of the *eigenvalues* of an algebraic model (defined below).

Let  $w$  be a non-backtracking walk in a  $B$ -graph, whose reduction is  $S^\leq/V'$ , and let  $S_B^\leq = \text{VisSub}_{/B}^\leq$ . Then the *wording induced by  $w$*  on  $S^\leq/V'$  is the map  $W$  from  $E_{S/V'}^{\text{dir}}$  to strings in  $E_B^{\text{dir}}$  of positive length, taking a directed edge  $e \in E_{S/V'}^{\text{dir}}$  to the string of  $E_B^{\text{dir}}$  edges in the non-backtracking walk in  $B$  that lies under the walk in  $S$  that it represents. Abstractly, we say that a  $B$ -wording of a graph  $T$  is a map  $W$  from  $E_T^{\text{dir}}$  to words over the alphabet  $E_B^{\text{dir}}$  that represent (the directed edges of) non-backtracking walks in  $B$  such that (1)  $W(\iota_T e)$  is the reverse word (corresponding to the reverse walk) in  $B$  of  $W(e)$ , (2) if  $e \in E_T^{\text{dir}}$  is a half-loop, then  $W(e)$  is of length one whose single letter is a half-loop, and (3) the tail of the first directed edge in  $W(e)$  (corresponding to the first vertex in the associated walk in  $B$ ) depends only on  $t_{Te}$ ; the *edge-lengths* of  $W$  is the function  $E_T \rightarrow \mathbb{N}$  taking  $e$  to the length of  $W(e)$ . [Hence the wording induced by  $w$  above is, indeed, a  $B$ -wording.]

Given a graph,  $T$ , and a  $B$ -wording  $W$ , there is a  $B$ -graph, unique up to isomorphism, whose underlying graph is  $\text{VLG}(T, \mathbf{k})$  where  $\mathbf{k}$  is the edge-lengths of  $W$ , and where the  $B$ -graph structure maps the non-backtracking walk in  $\text{VLG}(T, \mathbf{k})$  corresponding to an  $e \in E_T^{\text{dir}}$  to the non-backtracking walk in  $B$  given by  $W(e)$ . We denote any such  $B$ -graph by  $\text{VLG}(T, W)$ ; again this is a sort of inverse to starting with a non-backtracking walk and producing the wording it induces on its visited subgraph.

Notice that if  $S_B^\leq = \text{VLG}(T^\leq, W)$  for a  $B$ -wording,  $W$ , then the  $B$ -fibre counting functions  $\mathbf{a}_{S_B}$  and  $\mathbf{b}_{S_B}$  can be inferred from  $W$ , and we may therefore write  $\mathbf{a}_W$  and  $\mathbf{b}_W$ .

**2.8. Algebraic Models.** By a  $B$ -type we mean a pair  $T^{\text{type}} = (T, \mathcal{R})$  consisting of a graph,  $T$ , and a map from  $E_T^{\text{dir}}$  to the set of regular languages over the alphabet  $E_B^{\text{dir}}$  (in the sense of regular language theory) such that (1) all words in  $\mathcal{R}(e)$  are positive length strings corresponding to non-backtracking walks in  $B$ , (2) if for  $e \in E_T^{\text{dir}}$  we have  $w = e_1 \dots e_k \in \mathcal{R}(e)$ , then  $w^R \stackrel{\text{def}}{=} \iota_B e_k \dots \iota_B e_1$  lies in  $\mathcal{R}(\iota_T e)$ , and (3) if  $W: E_T^{\text{dir}} \rightarrow (E_B^{\text{dir}})^*$  (where  $(E_B^{\text{dir}})^*$  is the set of strings over  $E_B^{\text{dir}}$ ) satisfies  $W(e) \in \mathcal{R}(e)$  and  $W(\iota_T e) = W(e)^R$  for all  $e \in E_T^{\text{dir}}$ , then  $W$  is a  $B$ -wording. A  $B$ -wording  $W$  of  $T$  is of *type  $T^{\text{type}}$*  if  $W(e) \in \mathcal{R}(e)$  for each  $e \in E_T^{\text{dir}}$ .

Let  $\mathcal{C}_n(B)$  be a model that satisfies (1)–(3) of the definition of strongly algebraic. If  $\mathcal{T}$  a subset of  $B$ -graphs, we say that the model is *algebraic restricted to  $\mathcal{T}$*  if

either all  $S_{/B} \in \mathcal{T}$  occur in  $\mathcal{C}_n(B)$  or they all do not, and if so there are polynomials  $p_0, p_1, \dots$  such that  $c_i(S_{/B}) = p_i(S_{/B})$  for any  $S_{/B} \in \mathcal{T}$ . We say that  $\mathcal{C}_n(B)$  is *algebraic* if

- (1) setting  $h(k)$  to be the number of  $B$ -graph isomorphism classes of étale  $B$ -graphs  $S_{/B}$  such that  $S$  is a cycle of length  $k$  and  $S$  does not occur in  $\mathcal{C}_n(B)$ , we have that  $h$  is a function of growth  $(d-1)^{1/2}$ ; and
- (2) for any pruned, ordered graph,  $T^{\leq}$ , there is a finite number of  $B$ -types,  $T_j^{\text{type}} = (T^{\leq}, \mathcal{R}_j)$ ,  $j = 1, \dots, s$ , such that (1) any  $B$ -wording,  $W$ , of  $T$  belongs to exactly one  $\mathcal{R}_j$ , and (2)  $\mathcal{C}_n(B)$  is algebraic when restricted to  $T_j^{\text{type}}$ .

[In Article I we show that if instead each  $B$ -wording belong to *at least one*  $B$ -type  $T_j^{\text{type}}$ , then one can choose a another set of  $B$ -types that satisfy (2) and where each  $B$ -wording belongs to a *unique*  $B$ -type; however, the uniqueness is ultimately needed in our proofs, so we use uniqueness in our definition of algebraic.]

We remark that one can say that a walk,  $w$ , in a  $B$ -graph, or an ordered  $B$ -graphs,  $S_{/B}^{\leq}$ , is of *homotopy type*  $T^{\leq}$ , but when  $T$  has non-trivial automorphism one *cannot* say that is of  $B$ -type  $(T, \mathcal{R})$  unless—for example—one orders  $T$  and speaks of an *ordered  $B$ -type*,  $(T^{\leq}, \mathcal{R})$ . [This will be of concern only in Article II.]

We define the *eigenvalues* of a regular language,  $R$ , to be the minimal set  $\mu_1, \dots, \mu_m$  such that for any  $k \geq 1$ , the number of words of length  $k$  in the language is given as

$$\sum_{i=1}^m p_i(k) \mu_i^k$$

for some polynomials  $p_i = p_i(k)$ , with the convention that if  $\mu_i = 0$  then  $p_i(k) \mu_i^k$  refers to any function that vanishes for  $k$  sufficiently large (the reason for this is that a Jordan block of eigenvalue 0 is a nilpotent matrix). Similarly, we define the eigenvalues of a  $B$ -type  $T^{\text{type}} = (T, \mathcal{R})$  as the union of all the eigenvalues of the  $\mathcal{R}(e)$ . Similarly a *set of eigenvalues* of a graph,  $T$  (respectively, an algebraic model,  $\mathcal{C}_n(B)$ ) is any set containing the eigenvalues containing the eigenvalues of some choice of  $B$ -types used in the definition of algebraic for  $T$ -wordings (respectively, for  $T$ -wordings for all  $T$ ).

[In Article V we prove that all of our basic models are algebraic; some of our basic models, such as the permutation-involution model and the cyclic models, are not strongly algebraic.]

We remark that a homotopy type,  $T^{\leq}$ , of a non-backtracking walk, can only have beads as its first or last vertices; however, in the definition of algebraic we require a condition on *all pruned graphs*,  $T$ , which includes  $T$  that may have many beads and may not be connected; this is needed when we define homotopy types of pairs in Article II.

**2.9. SNBC Counting Functions.** If  $T^{\leq}$  is an ordered graph and  $\mathbf{k}: E_T \rightarrow \mathbb{N}$ , we use  $\text{SNBC}(T^{\leq}, \mathbf{k}; G, k)$  to denote the set of SNBC walks in  $G$  of length  $k$  and of homotopy type  $T^{\leq}$  and edge lengths  $\mathbf{k}$ . We similarly define

$$\text{SNBC}(T^{\leq}, \geq \boldsymbol{\xi}; G, k) \stackrel{\text{def}}{=} \bigcup_{\mathbf{k} \geq \boldsymbol{\xi}} \text{SNBC}(T^{\leq}, \mathbf{k}; G, k)$$

where  $\mathbf{k} \geq \boldsymbol{\xi}$  means that  $k(e) \geq \xi(e)$  for all  $e \in E_T$ . We denote the cardinality of these sets by replacing SNBC with snbc; we call  $\text{snbc}(T^{\leq}, \geq \boldsymbol{\xi}; G, k)$  the set of

$\xi$ -certified traces of homotopy type  $T^\leq$  of length  $k$  in  $G$ ; in Article III we will refer to certain  $\xi$  as *certificates*.

### 3. MAIN THEOREMS IN THIS ARTICLE

The main theorems in this article were stated in Article I. For ease of reading we restate them here.

Recall that if  $A' \subset A$  are sets, then  $\mathbb{I}_{A'}$  denotes the indicator function of  $A'$ .

We also recall that a model,  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$ , over a graph  $B$  may well have  $N \neq \mathbb{N}$  (e.g., our basic models above when  $B$  has half-loops); in this case many formulas involving the variable  $n$  are only defined for  $n \in N$ . For brevity, we often do not explicitly write  $n \in N$  in such formulas; for example we usually write

$$\lim_{n \rightarrow \infty} \quad \text{to abbreviate} \quad \lim_{n \in N, n \rightarrow \infty};$$

as another example,  $(B, \nu)$ -bounded expansions for a function  $f(k, n)$  only hold where  $f$  is defined, which is typically for all  $k \in \mathbb{N}$  but only  $n \in N$ .

**Theorem 3.1.** *Let  $B$  be a connected graph with  $\mu_1(B) > 1$ , and let  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  be an algebraic model over  $B$ . Let  $r > 0$  be an integer and  $\nu \geq \mu_1^{1/2}(B)$  be a real number. Then*

$$(9) \quad f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}(G) \text{Trace}(H_G^k)]$$

has a  $(B, \nu)$ -bounded expansion to order  $r$ ,

$$f(k, n) = c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

where

$$(10) \quad c_0(k) = \sum_{k'|k} \text{Trace}(H_B^{k'})$$

where the sum is over all positive integers,  $k'$ , dividing  $k$ ; hence

$$c_0(k) = \text{Trace}(H_B^k) + O(k)\mu_1^{k/2}(B);$$

furthermore, the larger bases of each  $c_i(k)$  (with respect to  $\mu_1^{1/2}(B)$ ) is some subset of the eigenvalues of the model. Finally, for any  $r' \in \mathbb{N}$  the function

$$(11) \quad \tilde{f}(n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{TangleFree}(\geq \nu, < r')}(G)] = \text{Prob}_{\mathcal{C}_n(B)}[\text{TangleFree}(\geq \nu, < r')]$$

has an asymptotic expansion in  $1/n$  to any order  $r$ ,

$$\tilde{c}_0 + \cdots + \tilde{c}_{r-1}/n^{r-1} + O(1)/n^r;$$

where  $\tilde{c}_0 = 1$ ; furthermore, if  $j_0$  is the smallest order of a  $(\geq \nu)$ -tangle occurring in  $\mathcal{C}_n(B)$ , then  $c_j = 0$  for  $1 \leq j < j_0$  and  $c_j > 0$  for  $j = j_0$  (provided that  $r \geq j_0 + 1$  so that  $\tilde{c}_{j_0}$  is uniquely defined).

Notice that a model may have—at least in principle—an infinite number of eigenvalues, which means that for each  $r, \nu$ , the number of bases of the  $c_i(k)$  may be unbounded as  $i \rightarrow \infty$ ; however there are a few remarks to consider:

- (1) taking  $\nu = \mu_1^{1/2}(B)$ , for each  $r$ , the  $c_i(k)$  with  $i < r$  have a finite number of bases;

(2) since for any fixed  $k$  we have

$$c_i(k) = \lim_{n \rightarrow \infty} n^i (f(k, n) - c_0(k) - \cdots - n^{1-i} c_{i-1}(k)),$$

the  $c_i(k)$  is uniquely defined and independent of  $r > i$ ; hence a fixed  $c_i(k)$  has a finite number of larger (than  $\mu_1^{1/2}(B)$ ) bases;

(3) in all our basic models, the eigenvalues of the model can be taken to be the  $\mu_j(B)$  and possibly the eigenvalue 1; hence all larger bases of any  $c_i(k)$  lie in this finite set of eigenvalues.

The other result that will be needed in later articles, namely Article V, is proven similarly to our proof of an expansion for the function in (11).

**Theorem 3.2.** *Let  $\mathcal{C}_n(B)$  be an algebraic model over a graph,  $B$ , and let  $S_{/B}$  be a connected, pruned graph of positive order that occurs in this model (recall that this means that for some  $n$  and some  $G \in \mathcal{C}_n(B)$ ,  $G_{/B}$  has a subgraph isomorphic to  $S_{/B}$ ). Then for some constant,  $C'$ , and  $n$  sufficiently large (and  $n \in \mathbb{N}$ ),*

$$\text{Prob}_{G \in \mathcal{C}_n(B)} [S_{/B} \cap G \neq \emptyset] \geq C' n^{-\text{ord}(S_{/B})}.$$

#### 4. THEOREMS FROM ARTICLE II

For ease of reading, let us recall the main theorems of Article II, which we will use here.

**Theorem 4.1.** *Let  $\mathcal{C}_n(B)$  be an algebraic model over a graph  $B$ . Let  $T^\leq$  be an ordered graph, let  $\xi: E_T \rightarrow \mathbb{N}$  be a function, and let*

$$\nu = \max\left(\mu_1^{1/2}(B), \mu_1(\text{VLG}(T, \xi))\right).$$

Let  $\psi_{/B}^\leq$  be any pruned, ordered  $B$ -graph. Then for any  $r \geq 1$  we have

$$(12) \quad f(k, n) = \mathbb{E}_{G \in \mathcal{C}_n(B)} [(\#\psi_{/B}^\leq \cap G) \text{snbc}(T^\leq; \geq \xi, G, k)]$$

has a  $(B, \nu)$ -bounded expansion

$$c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

to order  $r$ ; the bases of the coefficients in the expansion are some subset of the eigenvalues of the model, and  $c_i(k) = 0$  for  $i$  less than the order of all  $B$ -graphs that contain both a walk of homotopy type  $T^\leq$  and a subgraph isomorphic to  $\psi_{/B}$ .

We note that the conclusions of this theorem also hold for the function

$$(13) \quad f(k, n) = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\text{snbc}(T^\leq; \geq \xi, G, k)]$$

(for the first part of this paper, dealing with expansions for the expected certified traces, we use only need this particular  $f(k, n)$ ). The reason is, as we now explain, in that the special case where  $\psi_{/B}^\leq = \emptyset_{/B}$  is the empty graph, (12) reduces to (13); the reader who dislikes the empty graph is free to simply view the above theorem as also applying to (13) (this special case is stated both in Articles I and II), and with  $c_i(k) = 0$  if  $i < \text{ord}(T)$ . The empty graph refers to the graph whose vertex and directed edge sets are both the empty set,  $\emptyset$ ; since there is a unique map from  $\emptyset$  to any set, there are unique heads, tails, and edge involution, and a unique  $B$ -structure and ordering for this graph. Hence  $[\emptyset_{/B}^\leq]$  consists of this single graph, and the  $B$ -graph  $\emptyset_{/B}$  is a subgraph of any  $B$ -graph; for this reason  $[\emptyset_{/B}^\leq] \cap G_{/B}$  equals  $\emptyset_{/B}^\leq$  for any  $B$ -graph,  $G$ , and hence  $\#[\emptyset_{/B}^\leq] \cap G_{/B} = 1$  for all  $G_{/B}$ ; hence (12) reduces

to (13) in this case. Moreover, any  $B$ -graph contains the empty  $B$ -graph, and so the condition on  $i$  to have  $c_i(k) = 0$  amounts to  $i$  being less than the order of any  $B$ -graph of homotopy type  $T^\leq$ , which implies  $i < \text{ord}(T)$  (and is equivalent to  $i < \text{ord}(T)$  assuming at least one ordered  $B$ -graph of homotopy type  $T^\leq$  occurs in  $\{\mathcal{C}_n(B)\}$ ).

## 5. CERTIFIED TRACES AND THEOREM 3.1

In this section we describe our approach to proving the main part of Theorem 3.1, which is the existence of asymptotic expansions for (9) and facts about the coefficients  $c_i = c_i(k)$ ; the existence of asymptotic expansions for (11) follows easily from the facts we develop for (9).

**5.1. Motivation for Modified Traces.** If  $B$  is a bouquet of  $d/2$  whole-loops (so  $d$  is even), and  $\mathcal{C}_n(B)$  is the permutation model, then [Fri08] proves that

$$f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{snbc}(G, k)] = \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{Trace}(H_G^k)]$$

fails to have a  $(B, \nu)$ -Ramanujan expansion to all orders; the reason is mainly due to the existence of  $(\geq \nu, < r)$ -tangles that occur as subgraphs of graphs in  $\mathcal{C}_n(B)$ , where  $\nu > \sqrt{d-1}$  and  $r$  is of order  $d^{1/2}$ ; see the proof of Theorem 2.12 of [Fri08]. Our remedy, as in [Fri08], will be to replace  $\text{snbc}(G, k)$  by a “modification” or “regularization” of this count, by counting elements of  $\text{SNBC}(G, k)$  that satisfy a restrictive condition. In [Fri08], these modified versions of  $\text{snbc}(G, k)$  were called *selective traces*; in this series of articles we use the simpler *certified traces*.

## 5.2. Definition of Certified Traces.

**Definition 5.1.** Let  $\nu > 1$  be a real number,  $r, k \in \mathbb{Z}_{\geq 0}$ , and  $G$  be a graph. We define the set of  $(< \nu, < r)$  (*strictly certified walks*) (respectively,  $(\leq \nu, < r)$  (*weakly certified*)), denoted  $\text{CERT}_{<\nu, <r}(G, k)$  (respectively  $\text{CERT}_{\leq \nu, <r}(G, k)$ ) to be the set of  $\text{SNBC}$  walks in  $G$  of length  $k$  whose visited subgraph  $S$  satisfies  $\mu_1(S) < \nu$  (respectively,  $\mu_1(S) \leq \nu$ ) and  $\text{ord}(S) < r$ . We define the  $(< \nu, < r)$  (*weakly certified trace*) (respectively,  $(\nu, < r)$  (*strictly certified trace*)) of  $G$  of length  $k$ , denoted  $\text{cert}_{<\nu, <r}(G, k)$  (respectively  $\text{cert}_{\leq \nu, <r}(G, k)$ ) to be the cardinality of  $\text{CERT}_{<\nu, <r}(G, k)$  (respectively,  $\text{CERT}_{\leq \nu, <r}(G, k)$ ).

The fundamental fact about certified traces is that

$$(14) \quad G \in \text{TangleFree}(\geq \nu, < r) \Rightarrow \text{cert}_{<\nu', <r}(G, k) = \text{cert}_{\leq \nu', <r}(G, k) = \text{snbc}_{<r}(G, k)$$

for any  $r$  and  $\nu' \geq \nu$ . For this reason, the certified traces are sort of “regularized”  $\text{SNBC}$  count, that agrees with  $\text{snbc}_{<r}(G, k)$  for  $G$  without  $\nu$ -tangles of small order, but remains well controlled for  $G$  with such tangles. The *selective traces* of [Fri08] are another collection of “regularized traces,” but are more cumbersome to define and utilize. In this article we work with strongly-certified traces  $\text{cert}_{<\nu, <r}(G, k)$ , although one could equally well work with weakly-certified traces  $\text{cert}_{\leq \nu, <r}(G, k)$ . All that our trace methods require is that we apply (14) with  $r \rightarrow \infty$  and  $(d-1)^{1/2} < \nu \leq \nu' \leq (d-1)^{1/2} + \epsilon$  with  $\epsilon \rightarrow 0$ , and that we work with either  $\text{cert}_{<\nu', <r}(G, k)$  or  $\text{cert}_{\leq \nu', <r}(G, k)$  there; we prefer to take  $\nu' = \nu$  for simplicity. By contrast, we must work with  $\text{TangleFree}(\geq \nu, < r)$ , rather than the analogously defined  $\text{TangleFree}(> \nu, < r)$ , since we need the number of (isomorphism classes of) *minimal*  $(\geq \nu, < r)$ -tangles to be finite (see the remarks concerning Lemma 5.7 in Section 11).

**5.3. Statement of the Expansion Theorems for Certified Traces.** The main theorem in this paper is proven using the following two results.

**Theorem 5.2.** *Let  $B$  be a connected graph with  $\chi(B) < 0$ , and let  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  be an algebraic model over  $B$ . Let  $r' \in \mathbb{N}$  and  $\nu \geq \mu_1^{1/2}(B)$  be a real number. Then*

$$(15) \quad f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\text{cert}_{<\nu, <r'}(G, k)]$$

has a  $(B, \nu)$ -bounded asymptotic expansion to any order  $r \in \mathbb{N}$ ,

$$f(k, n) = c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

where for some function,  $h = h(k)$ , of growth  $(d-1)^{1/2}$  we have

$$(16) \quad c_0(k) = \sum_{k'|k} \text{Trace}(H_B^{k'}) - h(k)$$

(the sum being over all positive integers,  $k'$ , dividing  $k$ ), and where the larger bases of each  $c_i(k)$  (with respect to  $\mu_1^{1/2}(B)$ ) is some subset of any set of eigenvalues of the model. Also, the function  $h(k)$  in (16) is precisely the function described in condition (1) of the definition of algebraic model. The same theorem holds if the (strictly-)certified trace in (15) is replaced with the weakly-certified trace  $\text{cert}_{\leq\nu, <r}(G, k)$ .

Let  $\text{HasTangles}(\geq \nu, < r)$  denote the set of graphs,  $G$ , that contain a  $(\geq \nu, < r)$ -tangle (as a subgraph); then  $\text{HasTangles}(\geq \nu, < r)$  is the complement of  $\text{TangleFree}(\geq \nu, < r)$ , and so

$$\mathbb{I}_{\text{HasTangles}(\geq\nu, <r)}(G) = 1 - \mathbb{I}_{\text{TangleFree}(\geq\nu, <r)}(G).$$

**Theorem 5.3.** *Let  $B$  be a connected graph with  $\chi(B) < 0$ , and let  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  be an algebraic model over  $B$ . Let  $r, r', r'' > 0$  be integers and  $\nu > 1$  be a real number. Then*

$$(17) \quad f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{HasTangles}(\geq\nu, <r'')} (G) \text{cert}_{<\nu, <r'}(G, k)]$$

has a  $(B, \nu)$ -bounded asymptotic expansion to order  $r$ ,

$$f(k, n) = c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

such that the bases of the  $c_i(k)$  are the eigenvalues of the model; moreover,  $c_i$  vanishes if  $i$  is less than the smallest order of a  $(\geq \nu, < r'')$ -tangle that occurs in  $\mathcal{C}_n(B)$  provided that  $i < r$  (i.e., occurs with positive probability in  $\mathcal{C}_n(B)$  for some  $n$ , and hence for every  $n$  sufficiently large). In particular,  $c_0(k) = 0$  since  $\nu > 1$ . The same theorem holds if the strictly certified trace in (15) is replaced with the weakly-certified trace  $\text{cert}_{\leq\nu, <r'}(G, k)$ .

Subtracting the above two results yields an expansion theorem to order  $r$  for

$$f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{TangleFree}(\geq\nu, <r)}(G) \text{cert}_{<\nu, <r}(G, k)]$$

with  $c_0(k)$  given as in (16) for  $\nu > 1$ ; in view of (14), this function is the same as

$$f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{TangleFree}(\geq\nu, <r)}(G) \text{snbc}_{<r}(G, k)];$$

by (5) we may replace  $\text{snbc}_{<r}(G, k)$  by  $\text{snbc}(G, k)$  at an additive cost bounded by  $Ck^{2r} \text{Trace}(H_B^k)/n^r$ . This proves the expansion theorem in Theorem 3.1 for (9). The expansion theorem (11) easily follows from the methods we use to prove Theorem 5.3.



**5.4. Generalizations of Tangle Free Sets.** Theorems 5.2 and 5.3 involve indicator function for the set  $\text{TangleFree}(\geq \nu, < r)$ . In fact, we will prove more general results where  $\text{TangleFree}(\geq \nu, < r)$  is replaced with any set of graphs,  $\mathcal{T}$ , subject to certain restrictions, which we now describe.

**Definition 5.4.** We say that a graph is *pruned* if each of its vertices is of degree at least two, and *positive* if it is pruned and moreover each of its connected components is of positive order. We say that a  $B$ -graph (or ordered graph, etc.) is positive if its underlying graph is positive.

**Definition 5.5.** Let  $\mathcal{T}$  be a class of graphs (respectively, of  $B$ -graphs, of ordered graphs, etc.). We say that a graph ( $B$ -graph, etc.)  $G$  *meets*  $\mathcal{T}$  if  $G$  has a non-empty subgraph ( $B$ -subgraph, etc.) that is isomorphic to an element of  $\mathcal{T}$ , and otherwise we say  $G$  *avoids*  $\mathcal{T}$ ; we use  $\text{Meets}(\mathcal{T})$  and  $\text{Avoids}(\mathcal{T})$  respectively to be the class of graphs (or  $B$ -graphs, etc.) that meet and avoid  $\mathcal{T}$ .

Although  $\text{TangleFree}(\geq \nu, < r)$  describes a class of graphs, for various reasons we will want to work with the class of  $B$ -graphs whose underlying graph lies in  $\text{TangleFree}(\geq \nu, < r)$ ; for this reason we make the above definition for class of graphs and  $B$ -graphs. The above notion of meeting and avoiding also makes sense for ordered graphs and ordered  $B$ -graphs (and in other settings), but we will only be interested in graphs and  $B$ -graphs.

**Definition 5.6.** We say that a class of graphs ( $B$ -graphs, etc.)  $\mathcal{T}$  is *finitely generated* if there is a finite set  $\mathcal{T}'$  for which  $\text{Meets}(\mathcal{T}) = \text{Meets}(\mathcal{T}')$ , and *finitely positively generated* if there exists such a  $\mathcal{T}'$  such that each of its elements is positive.

It is easy to see that if  $\mathcal{T}' \subset \mathcal{T}$ , then  $\text{Meets}(\mathcal{T}) = \text{Meets}(\mathcal{T}')$  iff  $\mathcal{T}'$  contains at least one graph in each isomorphism class of graphs that are minimal with respect to inclusion (of graphs, of  $B$ -graphs, etc.) in  $\mathcal{T}$ .

Our interest in finitely positively generated classes is due to the following proposition.

**Lemma 5.7.** *For any real  $\nu > 1$  and  $r \in \mathbb{Z}$ ,  $\text{TangleFree}(\geq \nu, < r)$  is finitely positively generated.*

We will prove this in Section 11, using the ideas of Lemma 9.2 of [Fri08] and its proof there.

### 5.5. Main Theorem for Indicator Functions.

**Theorem 5.8.** *Let  $B$  be a graph,  $\{\mathcal{C}_n(B)\}_{n \in \mathbb{N}}$  an algebraic model of  $B$ , and let  $\mathcal{T}$  be a finitely positively generated class of graphs or of  $B$ -graphs; let  $j$  be the smallest order of a graph in  $\mathcal{T}$  that occurs in  $\mathcal{C}_n(B)$  (if  $j$  doesn't exist we take  $j = +\infty$ ). Then the function*

$$f(n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\mathcal{T})}(G)] = \text{Prob}_{G \in \mathcal{C}_n(B)} [\text{Meets}(\mathcal{T})]$$

has an asymptotic expansion in  $1/n$  to any order  $r$

$$c_0 + c_1 n^{-1} + \cdots + c_{r-1} n^{-r+1} + O(n^{-r})$$

with  $c_i = 0$  if  $i < j$  and, if  $j \neq +\infty$ , then  $c_j > 0$ .

We remark that the case  $j = +\infty$  in the above theorem is not particularly interesting, since then  $f(n) = 0$  for all  $n$ . Also, similar to a remark after Theorem 3.1, the above theorem implies that  $c_i$  are given inductively as

$$c_i = \lim_{n \rightarrow \infty} (f(n) - c_0 + c_1 n^{-1} + \cdots + c_{i-1} n^{-i+1}) n^i,$$

so that fact that  $f(n)$  is a probability of some event implies that  $f(n) \in [0, 1]$ , and hence  $c_i \in \mathbb{R}$  for all  $i$ , and the first nonzero  $c_i$  must be positive.

### 5.6. Main Theorems for Certified Traces with Indicator Functions.

**Theorem 5.9.** *Let  $\{\mathcal{C}_n\}_{n \in \mathbb{N}}$  be an algebraic model of random covering maps over a graph  $B$ , with  $\chi(B) < 0$ . Let  $r' \in \mathbb{N}$  and  $\nu \geq \mu_1^{1/2}(B)$ , and let  $\mathcal{T}$  be a finitely positively generated class of graphs (or of  $B$ -graphs). Then*

$$f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\mathcal{T})}(G) \text{cert}_{<\nu, <r'}(G; k)]$$

has a  $(B, \nu)$ -bounded asymptotic expansion to any order  $r$ ,

$$f(k, n) = c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

where the bases of the  $c_i$  are a subset of any set of eigenvalues of the model; moreover  $c_i(k)$  vanishes for all  $i$  less than the minimum order of a graph (or of a  $B$ -graph) that contains both some element of  $\mathcal{T}$  and a  $(< \nu, < r')$ -certified walk, provided that  $i < r$ .

**5.7. Remarks on Theorems 5.8 and 5.9 and Isomorphism Classes of  $B$ -Graphs.** In Sections 6–9 we prove Theorems 5.8 and 5.9, where  $\mathcal{T}$  is a finitely positively generated class of  $B$ -graphs (rather than graphs). There is no harm in passing to  $B$ -graphs, since our models are algebraic, and all our proof techniques work with  $B$ -graphs. The case where  $\mathcal{T}$  is a set of graphs is equivalent to the case where  $\mathcal{T}$  is replaced with the set of all  $B$ -graphs whose underlying graph lies in  $\mathcal{T}$ : if  $\mathcal{T}$  is a finitely generated set of graphs, then the set of all possible  $B$ -graph structures on the finite set of generators is finite.

We warn the reader of another change in Sections 6–9: we work sets,  $\Psi$ , of *isomorphism classes* of  $B$ -graphs, rather than  $B$ -graphs. It is simpler to state Theorems 5.8 and 5.9 with  $\Psi$  being a set of  $B$ -graphs, which is why we have done so. However, to prove these theorems we will work with formulas—including those for *Möbius functions* and *indicator function approximations*—that are simpler to define using *isomorphism classes* of  $B$ -graphs. Moreover, some notions discussed already, such as being *finitely generated*, can be stated in terms of a finite number of *isomorphism classes* of graphs (or of  $B$ -graphs). In [Fri08], the symbol  $\Psi$  with various subscripts refers either to isomorphism classes of graphs, or to a set of representatives in each isomorphism classes; in this article, we find it conceptually simpler to give the proofs of the above theorems using  $\Psi$  to refer to a set of *isomorphism classes* of  $B$ -graphs.

The small cost of working with isomorphism classes of  $B$ -graphs is that one has to get used to slightly different terminology. So in the next few sections we replace a class of graphs or  $B$ -graphs,  $\mathcal{T}$ , with a finite set of *isomorphism classes* of  $B$ -graphs

$$\Psi = \{[\psi_B^1], \dots, [\psi_B^m]\}.$$

One has to get used to speaking of  $B$ -graphs,  $\psi_B$ , lying in *an element* of  $\Psi$  (or of  $\Psi^+$  or  $\Psi_{<r}^+$ , defined in Section 8), meaning  $\psi_B \in [\psi_B^i]$  for some  $i$  (rather than  $\psi_B \in \mathcal{T}$ ). Hence we make the following definition.

**Definition 5.10.** Let  $B$  be a graph and  $\Psi$  be a set of isomorphism classes of  $B$ -graphs. We use  $\text{Meets}(\Psi)$  to denote the class of graphs,  $G$ , such that some nonempty subgraph of  $G$  is contained in an element  $\Psi$ .

## 6. PROOF OF THEOREM 5.2

In this section we prove Theorem 5.2. The first two subsections each prove an easy preliminary lemma.

### 6.1. VLG Comparisons.

**Lemma 6.1.** *Let  $T$  be a graph, and  $\mathbf{k}, \mathbf{k}'$  be two maps  $E_T \rightarrow \mathbb{N}$  with  $\mathbf{k} \leq \mathbf{k}'$  (i.e.,  $k(e) \leq k'(e)$  for all  $e \in E_T$ ). Then*

$$(18) \quad \mu_1(\text{VLG}(T, \mathbf{k})) \geq \mu_1(\text{VLG}(T, \mathbf{k}')) .$$

Its proof is a standard consequence of ‘‘Shannon’s algorithm,’’ and *majorization* as described just above Theorem 3.5 of [Fri08]. In the terminology there, each entry of the matrix  $Z_G(z)$ , where  $G$  is the oriented line graph of  $\text{VLG}(T, \mathbf{k})$ , majorizes each of  $Z_H(z)$  where  $H$  is the oriented line graph of  $\text{VLG}(T, \mathbf{k}')$ ; hence each entry of  $M_G(z)$  majorizes that of  $M_H(z)$ ; hence equation (12) and Theorem 3.5 of [Fri08] imply (18).

[One can also prove (18) without Shannon’s algorithm: note that every SNBC walk in  $G = \text{VLG}(T, \mathbf{k})$  can be cyclically shifted by at most  $\#E_G$  places to an SNBC walk in  $G$  beginning at some vertex of  $T$  (viewing  $V_T$  as a subset of  $V_G$ ); it follows that  $\mu_1(\text{VLG}(T, \mathbf{k}))$  is the limit as  $m \rightarrow \infty$  of  $W_m^{1/m}$ , where  $W_m$  is the number of SNBC walks beginning and ending at a vertex of  $T$  of length at most  $m$ . But if  $W'_m$  is the same quantity for  $G' = \text{VLG}(T, \mathbf{k}')$ , then we have  $W_m \geq W'_m$  in view of the one-to-one correspondence of such SNBC walks in  $G'$  with those in  $G$  (and with those in  $T$ ), for which the length of the walk in  $G$  is at most the length of that in  $G'$  since  $\mathbf{k} \leq \mathbf{k}'$ .]

**6.2. The Finiteness of Minimal Elements in an Upper Subset of  $\mathbb{N}^n$ .** The basis of our analysis of certified traces is the following finiteness lemma, which we give after some definitions.

**Definition 6.2.** For an integer  $n \geq 1$ , endow  $\mathbb{N}^n$  with the partial order  $\mathbf{k} \leq \mathbf{k}'$  to mean that  $k(i) \leq k'(i)$  for all  $i = 1, \dots, n$ . By an *upper set* in  $\mathbb{N}^n$  we mean a subset,  $U$ , such that if  $\mathbf{u} \in U$  and  $\mathbf{u} \leq \mathbf{u}'$ , then  $\mathbf{u}' \in U$ .

**Lemma 6.3.** *Any upper set of  $\mathbb{N}^n$  has a finite number of minimal elements.*

*Proof.* Let  $U \subset \mathbb{N}^n$  be an upper set. Let  $x_1, \dots, x_n$  be  $n$  indeterminates, and let  $I \subset \mathbb{C}[x_1, \dots, x_n]$  be the set of polynomials that are linear combinations of monomials

$$\mathbf{x}^{\mathbf{u}} = x_1^{u_1} \dots x_n^{u_n}$$

with  $\mathbf{u} \in U$ . Since  $U$  is an upper set,  $I$  is an ideal of the ring  $\mathbb{C}[x_1, \dots, x_n]$ ; by Hilbert’s Basis Theorem,  $I$  is finitely generated by polynomials,  $p_1, \dots, p_m \in I$ . For any  $i = 1, \dots, m$  and  $q_i \in \mathbb{C}[x_1, \dots, x_n]$ , each monomial  $\mathbf{x}^{\mathbf{w}}$  appearing in  $p_i q_i$  arises as the product of some monomial in  $p_i$  and some monomial in  $q_i$ . It follows that any monomial  $\mathbf{x}^{\mathbf{w}}$  that appears in a sum

$$p_1 q_1 + \dots + p_m q_m$$

has a corresponding monomial  $\mathbf{x}^{\mathbf{u}}$  that appears in one of  $p_1, \dots, p_m$ , with  $\mathbf{u} \leq \mathbf{w}$ . But for any  $\mathbf{w} \in U$ , since  $\mathbf{x}^{\mathbf{w}} \in I$ , we have

$$\mathbf{x}^{\mathbf{w}} = p_1 q_1 + \dots + p_m q_m,$$

for some  $q_1, \dots, q_m \in \mathbb{C}[x_1, \dots, x_n]$ , and hence there is a monomial  $\mathbf{x}^{\mathbf{u}}$  appearing in  $p_1, \dots, p_m$  for which  $\mathbf{u} \leq \mathbf{w}$ . Hence the finite set of  $\mathbf{u}$  such that  $\mathbf{x}^{\mathbf{u}}$  appears in one of  $p_1, \dots, p_m$ , is a set of certificates for  $U$ .  $\square$

One can alternatively prove the above lemma directly: clearly it holds for  $n = 1$ ; one can then prove the more general lemma that if  $\mathcal{P}_1, \mathcal{P}_2$  are posets where any upper set has a finite number of minimal elements, then the same is true of  $\mathcal{P}_1 \times \mathcal{P}_2$ .

**6.3. The Zeroth Order Coefficient.** In this section we make the following observations about algebraic models. First we need a simple lemma.

**Lemma 6.4.** *Let  $S$  be a connected and pruned graph. Then either  $S$  is a cycle, or  $S$  is of positive order. Also,  $\mu_1(S) > 1$  iff  $\chi(S) < 0$ .*

We will specifically need the first statement of the lemma in this section. The second statement in the above lemma is used in a number of places in this series of articles, to interchangeably use the conditions  $\mu_1(S) > 1$  and  $\chi(S) < 0$  for pruned, connected graphs. Since both statements are based on a similar principle, we prove both of them here.

*Proof.* To prove the first claim, note that

$$\text{ord}(S) = (1/2) \sum_{v \in V_S} (\text{deg}'_S(v) - 2),$$

where  $\text{deg}'$  is the usual degree of a vertex except that each half-loop about a vertex,  $v$ , contributes 2 to its degree (instead of 1 used for the usual degree and Euler characteristic). Since  $S$  is pruned, we have  $\text{deg}'_S(v) \geq 2$  for each  $v \in V_S$ , and hence  $\text{ord}(S) = 0$  iff for  $\text{deg}'_S(v) = 2$  for all  $v \in V_S$ . It follows that each vertex of  $V_S$  is either (1) incident upon two edges that are not self-loops, or (2) incident upon exactly one self-loop. In case any vertex is incident upon a self-loop, then the graph has one vertex and must be incident upon a whole-loop (for otherwise the vertex would be of degree one); hence  $S$  is a cycle of length 1. Otherwise all vertices are of case (1), and therefore  $S$  is a cycle.

To prove the second claim, we similarly note that

$$\chi(S) = (1/2) \sum_{v \in V_S} (\text{deg}_S(v) - 2).$$

So if  $S$  is pruned and  $\chi(S) \leq 0$ , then  $\text{deg}_S(v) = 2$  for all  $v$ ; it follows that  $\chi(S) \leq 0$  implies that  $S$  is either (1) a cycle, (2) a path where each endpoint is incident upon an additional half-loop, (3) a single vertex incident upon a single whole-loop, or (4) a single vertex incident upon two half-loops. In all these cases we easily check that  $\mu_1(S) \leq 1$ , since we easily see that there are at most two SNBC walks of a given length about any vertex of  $S$ .

If  $\chi(S) < 0$ , then some vertex of  $S$  has degree 3, say  $v$ . Let us show that  $\mu_1(S) > 1$ .

First we claim that for any  $e \in E_S^{\text{dir}}$  with  $t_{se} = v$ , there is a non-backtracking walk,  $w$ , about  $v$  beginning with  $e$ : to see this, we keep walking in a non-backtracking fashion, which we can do since each vertex is of degree two, until

we reach a repeated vertex; then return to  $v$ . Similarly, for any  $e$  with  $h_S e = v$ , there is a non-backtracking walk about  $v$  ending in  $e$ . So consider all pairs  $(e, e')$  such that  $t_S e = h_S e' = v$  and that there exists a non-backtracking walk beginning in  $e$  and ending in  $e'$ ; for each such pair, choose such a non-backtracking walk,  $w_{e, e'}$ ; let  $m$  be an upper bound on the lengths of all these walks. We now claim that for any  $k$ , the number of SNBC walks about  $v$  of length at most  $km$  is at least  $2^{k-2}3$ . To see this, consider which walks of the form

$$w_{e_1, e_2} w_{e_3, e_4} \cdots w_{e_{2k-1}, e_{2k}}$$

are SNBC: we may choose  $e_1$  to be any of at least 3 edges leaving  $v$ ; choosing some  $e_2$  such that  $w_{e_1, e_2}$  exists, we choose  $e_3$  to be any of at least 2 edges leaving  $v$  not equal to  $\iota e_2$ ; for  $i = 2, \dots, k-2$  we similarly choose  $e_{2i+1}$  to be an edge leaving  $v$  not equal to  $\iota e_{2i}$ , of which there are at least 2 choices; finally we choose  $e_{2k-1}$  to be an edge leaving  $v$  such that  $w_{e_{2k-1}, e_{2k}}$  exists and  $e_{2k-1} \neq \iota e_{2k-2}$  and  $e_{2k} \neq \iota e_1$ , of which there must be at least one choice. Hence the trace of  $H_S^{km}$  must be at least  $2^{k-2}3$ , and hence

$$2^{k-2}3 \leq \text{Trace}(H_S^{km}) \leq (\#E_S^{\text{dir}}) \mu_1^{km}(S);$$

taking  $k \rightarrow \infty$  we have  $\mu_1(S) \geq 2^{1/m} > 1$ .  $\square$

We easily see that no vertex of an SNBC walk can be of degree zero or one, and hence we conclude the following corollary.

**Corollary 6.5.** *Let  $T^\leq$  be the homotopy type of an SNBC walk. Then either  $T^\leq$  is the homotopy type of a cycle (i.e.,  $T$  is the bouquet of a single whole-loop), or else  $\text{ord}(T) \geq 1$ .*

**Corollary 6.6.** *Let  $B$  be a graph and  $\mathcal{C}_n(B)_{n \in \mathbb{N}}$  an algebraic model over  $B$ . Then there is a constant,  $C$ , and a function  $g$  of growth  $\mu_1(B)$  such that for  $1 \leq k \leq C/n^{1/2}$  we have*

$$\begin{aligned} \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{snbc}_0(G, k)] &= c_0(k) + g(k)O(1)/n, \\ \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{snbc}_{\geq 1}(G, k)] &= g(k)O(1)/n, \end{aligned}$$

where  $c_0(k)$  is given in (10).

*Proof.* The second equation follows from (5) (in the definition of strongly algebraic and algebraic). If  $S_{\bar{B}}^\leq$  is a  $B$ -graph than is the visited subgraph of an SNBC walk,  $w$ , of order 0, then  $S_{\bar{B}}^\leq$  is necessarily of the homotopy type of a cycle, and therefore the length  $k'$ , of  $S$ , must divide  $k$ ; furthermore, the directed edges over  $B$  that lie over  $w$  in the first encountered ordering yield an SNBC walk in  $B$  of length  $k'$ . Conversely, every SNBC walk in  $B$  of length  $k'$  gives rise to an ordered graph  $S_{\bar{B}}^\leq$ , unique up to an isomorphism of  $B$ -graphs. Since each such  $S_{\bar{B}}^\leq$  has  $c_0(S_{\bar{B}}) = 1$ , the first equation of the corollary follows, since  $\text{snbc}(B, k') = \text{Trace}(H_B^{k'})$ .  $\square$

#### 6.4. Conclusion of The Proof of Theorem 5.2.

*Proof of Theorem 5.2.* The elements of  $\text{CERT}_{<\nu, <r'}(G; k)$  are walks of a finite number of homotopy types,  $T^\leq$ . Hence it suffices to prove that for any fixed  $T^\leq$

$$f(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^\leq; G, k)) \right]$$

has a  $(B, \nu)$ -asymptotic expansion to any order  $r$ , and that its zero-th coefficient,  $c_0(k)$  is given by

- (1) the formula (10) when  $T^\leq$  is the homotopy type of a cycle (i.e.,  $T$  is the bouquet of a single whole-loop), and
- (2)  $c_0(k) = 0$  otherwise.

So fix an ordered graph,  $T^\leq$ , and consider the set

$$U = \{\mathbf{k}: E_T \rightarrow \mathbb{N} \mid \text{VLG}(T^\leq, \mathbf{k}) < \nu\}$$

which is clearly an upper set by Lemma 6.1; according to Lemma 6.3, this upper set has a finite number of minimal elements

$$\xi^1, \dots, \xi^s.$$

[Intuitively we think of each  $\xi_j$  as a *certificate* for belonging in  $U$ , in that the condition  $\mathbf{k} \geq \xi_j$  certifies (or guarantees) that  $\mathbf{k} \in U$ . The usefulness of the certified trace is due, in part, to the fact that the condition  $\text{VLG}(T^\leq, \mathbf{k}) < \nu$  is equivalent to being certified so by one of finitely many certificates. This is why we use the name *certified trace*.]

For each  $M \subset [s]$ , we have

$$\bigcap_{m \in M} \{\mathbf{k} \mid \mathbf{k} \geq \xi^m\} = \{\mathbf{k} \mid \mathbf{k} \geq \xi^M\}$$

where

$$\xi^M \stackrel{\text{def}}{=} \max_{m \in M}(\xi^m)$$

is the component-wise maximum. By inclusion/exclusion we have

$$(19) \quad \#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^\leq; G, k)) = \sum_{M \subset [s], M \neq \emptyset} (-1)^{1+(\#M)} \text{snbc}(T^\leq, \geq \xi^M; G, k).$$

By Theorem 4.1, we have that for any  $r > 0$  and any  $M \subset [s]$  with  $M \neq \emptyset$ ,

$$f_M(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{snbc}(T^\leq, \geq \xi^M; G, k)]$$

has a  $(B, \nu_M)$ -bounded expansion to order  $r$  with

$$\nu_M = \max\left(\mu_1^{1/2}(B), \mu_1(\text{VLG}(T, \xi^M))\right).$$

Since  $U$  is an upper set, we have  $\xi^M \in U$  for all  $M \neq \emptyset$ , and hence  $\nu_M \leq \nu$ . It follows that each  $f_M(k, n)$  has an expansion that satisfies the conditions in the statement of the theorem. Since  $\mathcal{C}_n(B)$  is a finite probability space, we may take expected values in (19) to conclude that

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^\leq; G, k)) \right] = \sum_{M \subset [s], M \neq \emptyset} (-1)^{1+(\#M)} f_M(k, n);$$

since the RHS of this equation is a finite sum of functions with  $(B, \nu)$ -expansions to any order  $r$ , so is each function

$$f(k, n) = \mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^\leq; G, k)) \right].$$

Summing over all the types,  $T^\leq$ , of walks of order less than  $r'$ , we conclude the same for

$$f(k, n) = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\text{cert}_{<\nu, <r'}(G, k)],$$

which proves the theorem for the expected value of the strictly-certified trace in (15).

It remains to compute  $c_0(k)$ . For  $T$  of order one or greater,

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^{\leq}; G, k)) \right] \leq \mathbb{E}_{G \in \mathcal{C}_n(B)} [\text{snbc}_{\geq 1}(G, k)]$$

which is bounded by  $g(k)O(1)/n$  by Corollary 6.6. On the other hand, any graph that is the homotopy type of a cycle has  $\mu_1 = 1 < \nu$ , and therefore if  $T^{\leq}$  is the homotopy type of a cycle, we have

$$\#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^{\leq}; G, k)) = \text{snbc}_0(G, k)$$

for  $k \geq 1$ . Hence for this  $T$  we have

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} \left[ \#(\text{CERT}_{<\nu, <r'}(G, k) \cap \text{SNBC}(T^{\leq}; G, k)) \right] = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\text{snbc}_0(G, k)]$$

whose zeroth order coefficient,  $c_0(k)$ , is given by Corollary 6.6 to be as in (10).

This concludes the proof for strongly-certified traces. The proof for weakly-certified traces is the same, with  $U$  replaced by

$$U' = \{\mathbf{k}: E_T \rightarrow \mathbb{N} \mid \text{VLG}(T^{\leq}, \mathbf{k}) \leq \nu\}.$$

This may change the set of certificates, i.e., of minimal elements,  $\xi^1, \dots, \xi^s$ , but everything else in the proof remains the same.  $\square$

## 7. FINITENESS OF MINIMAL TANGLES

We show that there are, up to isomorphism, only a finite number of minimal ( $\geq \nu, < r$ )-tangles for any real  $\nu > 0$  and integer  $r > 0$ . This fact is not generally true of ( $> \nu, < r$ )-tangles. This fact is essentially Lemma 9.2 of [Fri08], stated in the terms we use in this article.

**Definition 7.1.** Let  $\nu > 0$  be a real number and  $r > 0$  an integer. We say that a graph,  $\psi$ , is a *minimal* ( $\geq \nu, < r$ )-tangle if  $\psi$  is a ( $\geq \nu, < r$ )-tangle, i.e.,  $\mu_1(\psi) \geq \nu$  and  $\text{ord}(\psi) < r$ , but all of proper subgraphs of  $\psi$  are not ( $\geq \nu, < r$ )-tangles.

Let us recall Lemma 9.2 of [Fri08] and its simple proof based on the following “continuity lemma” regarding VLG’s.

**Lemma 7.2.** *Let  $T$  be a fixed graph, and  $E', E''$  partition of  $E_T$  into two sets. Let  $\mathbf{k}^1, \mathbf{k}^2, \dots$  a sequence of elements of  $\mathbb{N}^{E_T}$  such that*

- (1)  $\mathbf{k}^i(e')$  is independent of  $i$  for  $e' \in E'$ , and
- (2)  $\mathbf{k}^i(e'') \rightarrow \infty$  as  $i \rightarrow \infty$  for each  $e'' \in E''$ .

*Let  $T'$  be the graph obtained from  $T$  by discarding  $E''$  from  $E_T$ , and let  $\mathbf{k}'$  be the restriction of  $\mathbf{k}^i$  to  $E'$ . Say that there is a  $\nu > 1$  such that for all  $i$  we have*

$$(20) \quad \mu_1(\text{VLG}(T, \mathbf{k}^i)) \geq \nu.$$

*Then*

$$\lim_{i \rightarrow \infty} \mu_1(\text{VLG}(T, \mathbf{k}^i)) = \mu_1(\text{VLG}(T', \mathbf{k}')).$$

We use *Shannon’s algorithm*, as in [Fri08], Theorem 3.6 and its proof; we correct a minor error there: it is necessary that (20) hold with  $\nu > 1$ , since if  $T$  has one vertex, one whole-loop, and  $E' = \emptyset$ , then  $\mu_1(\text{VLG}(T, \mathbf{k}^i)) = 1$  for all  $i$ , but  $\mu_1(\text{VLG}(T', \mathbf{k}')) = 0$ .

*Proof.* For any  $\mathbf{k} \in \mathbb{N}^{E_T}$ , let  $M_{\mathbf{k}}(z)$  be the square matrix indexed on  $E_T^{\text{dir}}$  whose  $(e, e')$  entry is 0 if the corresponding of  $H_T$  is zero (i.e.,  $e, e'$  are not the directed edges of a non-backtracking walk of length 2), and otherwise this entry is  $z^{k(e)}$ . Let us prove that  $\mu_1(\text{VLG}(T, \mathbf{k}))$  is the reciprocal of the smallest positive root  $z = z_{\mathbf{k}}$  of the polynomial equation

$$(21) \quad \det(I - M(z)) = 0, \quad M(z) = M_{\mathbf{k}}(z).$$

To see this, let  $G = \text{VLG}(T, \mathbf{k})$ , and let  $G' = \text{Line}(G)$  be the oriented line graph of  $G$ ; hence  $V_{G'} = E_G^{\text{dir}}$ , and for  $e, e' \in V_{G'} = E_G^{\text{dir}}$  there is one or zero edges from  $e$  to  $e'$  according to whether or not  $e, e'$  are the directed edges of a non-backtracking walk of length two. For each directed edge of  $T$ ,  $e_T \in E_T^{\text{dir}}$ , let  $\tilde{e}_T \in V_{G'}$  denote the first directed edge in the (beaded) directed walk path in  $G'$  corresponding to  $e_T$ ; let  $\tilde{E}$  be the union of all the  $\tilde{e}_T$  with  $e_T \in E_T^{\text{dir}}$ . Then we easily see that  $G'$  is the same graph as the variable-length graph on its subset of vertices,  $\tilde{E}$ , where two vertices,  $\tilde{e}_T, \tilde{e}'_T$  have either one or zero edges from  $\tilde{e}_T$  to  $\tilde{e}'_T$  iff they form a non-backtracking walk of length two in  $E_T^{\text{dir}}$ , and, if so, the length of the edge is  $k(e_T)$ . Hence  $\mu_1(G)$ , which equals the Perron-Frobenius eigenvalue of  $G'$ , is given by (21) in view of *Shannon's algorithm* (e.g., Theorem 3.5 of [Fri08], but see the much earlier references in Section 3.2 of [Fri08]).

Next say that an element of  $V_{\text{Line}(T)} = E_T^{\text{dir}}$  belongs to  $E'$  if its  $\nu_T$  orbit, i.e., its corresponding edge, lies in  $E'$ , and similarly for  $E''$ . Then we may partition  $E_T^{\text{dir}}$  into two sets: those belonging to  $E'$  and those belonging to  $E''$ , and this gives a block representation of  $M(z)$  as

$$M_{\mathbf{k}^i}(z) = \begin{bmatrix} M_1(z) & M_{3,i}(z) \\ M_2(z) & M_{4,i}(z) \end{bmatrix},$$

where  $M_1(z), M_2(z)$  are blocks that are independent of  $i$ , since the edge-lengths  $\mathbf{k}^i$  are constant on directed edges belonging to  $E'$ , and each non-zero entry of  $M_{3,i}(z), M_{4,i}(z)$  is a power of  $z$  than tends to infinity as  $i \rightarrow \infty$ . In view of the last paragraph, we have that  $\mu_1(\text{VLG}(T, \mathbf{k}^i))$  is the reciprocal of the smallest positive root  $z_i$  to

$$\det(I - M_{\mathbf{k}^i}(z)) = 0.$$

On the other hand, the above paragraph also shows that  $\mu_1(\text{VLG}(T', \mathbf{k}'))$  is the reciprocal of the smallest positive root,  $z$ , of

$$(22) \quad \det(I' - M_1(z)) = 0$$

where  $I'$  is the identity matrix indexed on directed edges belonging to  $E'$ ; by convention, we allow  $z = +\infty$  if (22) has no positive roots, in which case  $\mu_1(\text{VLG}(T', \mathbf{k}')) = 0$ . It remains to prove that

$$(23) \quad \lim_{i \rightarrow \infty} z_i = z_0.$$

Let us first show that the above limit exists. Since  $z_i < 1/\nu$ , the sequence  $\{z_i\}$  is bounded above; let  $z_\infty$  be its least upper bound; clearly  $z_\infty \leq 1/\nu < 1$ . By definition, there exist  $i_1, i_2, \dots$  such that  $z_{i_n} \rightarrow z_\infty$  as  $n \rightarrow \infty$ . Let  $i = i_a$  for a fixed  $a \in \mathbb{N}$ . We have  $\mathbf{k}^i \leq \mathbf{k}^j$  for  $j$  sufficiently large, and hence then Lemma 6.1 implies that for such  $j$ ,  $1/z_i > 1/z_j$ , i.e.,  $z_i < z_j$ . Hence

$$\liminf_{n \rightarrow \infty} z_j \geq z_i = z_{i_a}.$$



Taking  $a \rightarrow \infty$  shows that

$$\liminf_{j \rightarrow \infty} z_j \geq z_\infty = \limsup_{n \rightarrow \infty} z_n.$$

Hence the limit in (23) exists. To prove the lemma it suffices to show that

$$z_\infty \leq z_0 \quad \text{and} \quad z_\infty \geq z_0.$$

Since  $\text{VLG}(T', \mathbf{k}')$  is a subgraph of each  $\text{VLG}(T, \mathbf{k}^i)$ , we have

$$1/z_0 = \mu_1(\text{VLG}(T', \mathbf{k}')) \leq \mu_1(\text{VLG}(T, \mathbf{k}^i)) = 1/z_i,$$

and hence, taking  $i \rightarrow \infty$ ,  $1/z_0 \leq 1/z_\infty$ . Hence  $z_\infty \leq z_0$ . Now let us show  $z_0 \leq z_\infty$ .

Since  $z_i \leq z_\infty \leq 1/\nu < 1$  for each  $i$ , we have that  $M_{3,i}(z), M_{4,i}(z)$  tend to zero as  $i \rightarrow \infty$ , and hence

$$M_{\mathbf{k}^i}(z_i) = \begin{bmatrix} M_1(z_i) & M_{3,i}(z_i) \\ M_2(z_i) & M_{4,i}(z_i) \end{bmatrix} \rightarrow \begin{bmatrix} M_1(z_\infty) & 0 \\ M_2(z_\infty) & 0 \end{bmatrix}$$

as  $i \rightarrow \infty$ ; then (21) (and the continuity of the determinant) implies that

$$\det \left( \begin{bmatrix} I' & 0 \\ 0 & I'' \end{bmatrix} - \begin{bmatrix} M_1(z_\infty) & 0 \\ M_2(z_\infty) & 0 \end{bmatrix} \right) = 0,$$

where  $I''$  is the block identity matrix corresponding to directed edges belonging to  $E''$ . Hence

$$\det \left( \begin{bmatrix} I' - M_1(z_\infty) & 0 \\ -M_2(z_\infty) & I'' \end{bmatrix} \right) = 0$$

and therefore

$$\det(I' - M_1(z_\infty)) = \det \left( \begin{bmatrix} I' - M_1(z_\infty) & 0 \\ -M_2(z_\infty) & I'' \end{bmatrix} \right) = 0.$$

Hence  $z_\infty$  is a positive root of (22). Therefore  $z_0 \leq z_\infty$ .  $\square$

**Theorem 7.3.** *Let  $\nu > 1$  be a real number and  $r \geq 0$  an integer. The number of (isomorphism classes of) minimal ( $\geq \nu, < r$ )-tangles is finite.*

*Proof.* Assume, to the contrary, that the theorem is false for some  $r$ ; since there are only finitely many homotopy types of order less than  $r$ , then must exist an ordered graph,  $T^\leq$ , and an infinite sequence of distinct minimal  $\nu$ -tangles  $\psi^1, \psi^2, \dots$  of homotopy type  $T^\leq$  such that  $\mu_1(\psi^i) \geq \nu$ . Then we have  $\psi^i = \text{VLG}(T; \mathbf{k}^i)$  for a sequence of distinct vectors  $\mathbf{k}^i \in \mathbb{N}^{E_T}$ . Let  $E_T = \{e_1, \dots, e_m\}$ ; by passing to a subsequence of the  $\mathbf{k}^i$ , we may assume that either (1)  $k^i(e_1) \rightarrow \infty$  as  $i \rightarrow \infty$ , or that (2) the  $k^i(e_1)$  are bounded, and hence—by passing to a further subsequence—that  $k^i(e_1)$  is independent of  $i$ ; we then repeat this process to show that for  $j = 2, \dots, m$ , by passing to a subsequence we may assume that  $k^i(e_j) \rightarrow \infty$  as  $i \rightarrow \infty$  or  $k^i(e_j)$  is independent of  $i$ . At this point the  $\mathbf{k}^i$  satisfy the hypotheses of Lemma 7.2. It follows that  $\text{VLG}(T', \mathbf{k}')$  is a proper subgraph of  $\text{VLG}(T, \mathbf{k}^i) \simeq \psi^i$  for all  $i$ , and  $\mu_1(\text{VLG}(T', \mathbf{k}')) \geq \nu$ , contradicting the minimality of the  $\psi^i$ .  $\square$

We remark that the above lemma would be false for ( $> \nu, < r$ )-tangles, defined the same but with strict inequality between  $\mu_1$  and  $\nu$ , as was explained in a footnote in Subsection 2.4.

## 8. INDICATOR FUNCTION APPROXIMATION

In this section we develop some foundations regarding approximations  $I_r(\Psi, G)$  we shall use for the indicator function

$$\mathbb{I}_{\text{Meets}(\Psi)}(G_{/B})$$

where  $\Psi$  is a collection of  $B$ -graphs that satisfy slighter milder assumptions than those of Theorems 5.8 and 5.9. The results in this section are adaptations of results of Section 9 of [Fri08]. In the first subsection we will state all the definitions and results we will use in Sections 9 and 10; the remaining subsections are devoted to their proofs.

## 8.1. The Main Results.

**Definition 8.1.** Let  $B$  be a graph, and let

$$\Psi = \{[\psi_{/B}^1], \dots, [\psi_{/B}^m]\}$$

be a finite set of isomorphism classes of  $B$ -graphs. By *the set derived  $B$ -graphs of  $\Psi$* , denoted  $\Psi^+$ , we mean the isomorphism classes  $[\psi_{/B}]$  of  $B$ -graphs such that  $\psi$  can be written as the union of  $B$ -subgraphs each isomorphic to some  $\psi_{/B}^i$ . We use  $\Psi_{<r}^+$  to denote classes  $[\psi_{/B}]$  of  $\Psi^+$  with  $\text{ord}(\psi) < r$ . If  $G_{/B}$  is any  $B$ -graph, the  $\Psi$ -*image in  $G$* , denoted,  $\Psi^+ \cap G_{/B}$ , is the union of all  $B$ -subgraphs of  $G_{/B}$  that lie in some class,  $[\psi_{/B}^i]$ , of  $\Psi$  (clearly  $\Psi^+ \cap G_{/B}$  is largest subgraph of  $G_{/B}$  that lies in a class of  $\Psi^+$ ); we use  $\text{ord}_{\Psi}(G)$  to denote  $\text{ord}(\Psi^+ \cap G_{/B})$ .

**Definition 8.2.** Let  $B$  be a graph and  $\psi_{/B}, G_{/B}$  be two  $B$ -graphs. By an *injective morphism*  $\psi_{/B} \rightarrow G_{/B}$  we mean a morphism that is injective as a map of vertex sets and of edge sets. We use  $N(\psi_{/B}, G_{/B})$  to denote the number of injective maps  $\psi_{/B} \rightarrow G_{/B}$ .

The proposition below is worth stating, but easy to prove.

**Proposition 8.3.** *For any graph  $B$  and any two  $B$ -graphs  $\psi_{/B}, G_{/B}$  we have that for any ordering  $\psi_{/B}^{\leq}$  on  $\psi_{/B}$ ,*

$$N(\psi_{/B}, G_{/B}) = \#([\psi_{/B}^{\leq}] \cap G_{/B}).$$

*Proof.* Fix an ordering  $\psi_{/B}^{\leq}$ ; each injective morphism  $u: \psi_{/B} \rightarrow G_{/B}$  determines an element  $S_{/B}^{\leq} = u(\psi_{/B}^{\leq})$  of  $[\psi_{/B}^{\leq}] \cap G_{/B}$ ; furthermore, for  $S_{/B}^{\leq} \in [\psi_{/B}^{\leq}] \cap G_{/B}$  there is a unique isomorphism  $\psi_{/B}^{\leq} \rightarrow S_{/B}^{\leq}$  giving rise to an injection  $\psi_{/B} \rightarrow G_{/B}$ . We easily check that this correspondence between injective maps  $u$  and elements of  $[\psi_{/B}^{\leq}] \cap G_{/B}$  are inverses of each other.  $\square$

We now state three results that will be proven in the subsections that follow this one.

**Lemma 8.4.** *Let  $B$  be a graph, and let*

$$\Psi = \{[\psi_{/B}^1], \dots, [\psi_{/B}^m]\}$$

*be a finite set of isomorphism classes of  $B$ -graphs. If each  $\psi^i$  is positive, then for each  $r \in \mathbb{N}$ ,  $\Psi_{<r}^+$  is finite.*

**Lemma 8.5.** *Let  $B$  be a graph, and let*

$$\Psi = \{[\psi_B^1], \dots, [\psi_B^m]\}$$

*be a finite set of isomorphism classes of  $B$ -graphs such  $\Psi_{<r}^+$  is finite for all  $r \in \mathbb{N}$ ; let  $s$  be the largest number of edges among the graphs  $\psi^1, \dots, \psi^m$ .*

(1) *If  $[\psi_B] \in \Psi^+$  and  $G_B$  is any  $B$ -graph, then any injective morphism  $\psi_B \rightarrow G_B$  factors as an injective map  $\psi_B \rightarrow \Psi^+ \cap G_B$  followed by the inclusion of  $\Psi^+ \cap G_B$  in  $G_B$ .*

(2) *If  $r \in \mathbb{N}$  and  $G_B$  is a  $B$ -graph with  $\text{ord}_\Psi(G) \geq r$ , and  $[\psi_B] \in \Psi_{<r}^+$ , then any injective map  $\psi_B \rightarrow G_B$  factors as two injective maps  $\psi_B \rightarrow \psi'_B$  and  $\psi'_B \rightarrow G_B$  where  $\psi'_B$  lies in an element of  $\Psi_{<r+s}^+ \setminus \Psi_{<r}^+$ , i.e.,  $[\psi'_B]$  is an element of  $\Psi^+$  whose order is between  $r$  and  $r + s - 1$ . In particular*

$$(24) \quad N(\psi_B, G_B) \leq \sum_{\psi'_B \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} N(\psi_B, \psi'_B) N(\psi'_B, B_B).$$

**Theorem 8.6.** *Let  $B$  be a graph, and let*

$$\Psi = \{[\psi_B^1], \dots, [\psi_B^m]\}$$

*be a finite set of isomorphism classes of  $B$ -graphs such  $\Psi_{<s}^+$  is finite for all  $s \in \mathbb{N}$  and each  $\psi^i$  is pruned. Then for each  $[\psi_B] \in \Psi^+$  there is a rational number  $\mu[\psi_B]$  such that for any  $r \in \mathbb{N}$  the function*

$$I_r(\Psi, G_B) = \sum_{[\psi_B] \in \Psi_{<r}^+} \#([\psi_B] \cap G) \mu[\psi_B],$$

*satisfies*

(1)

$$(25) \quad I_r(\Psi, G_B) = \mathbb{I}_{\text{Meets}(\Psi)}(G_B)$$

*whenever  $\text{ord}_\Psi(G_B) < r$ , and*

(2) *there is a constant  $C \in \mathbb{R}$  and  $s \in \mathbb{N}$  such that if  $\text{ord}_\Psi(G_B) \geq r$ , then*

$$(26) \quad |I_r(\Psi, G_B)| \leq C \sum_{[\psi_B] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} \#([\psi_B] \cap G).$$

**8.2. Proof of Lemma 8.4.** Lemma 8.4 follows from the proof Lemma 9.2 of [Fri08], which proves the same in the context of graphs as opposed to  $B$ -graphs. For convenience we provide a complete proof here. Our proof will use the following graph theoretic lemma.

**Lemma 8.7.** *Let  $G_1 \subset G$  be pruned graphs, i.e., no vertex is isolated or is incident upon a single edge that is not a self-loop. Then if  $G_1 \neq G$ ,  $\text{ord}(G_1) < \text{ord}(G)$ .*

This is Lemma 4.10 of [Fri08]; for ease of reading we provide a proof here.

*Proof.* Assume that  $G_1 \neq G$ . If  $V_{G_1} = V_G$ , then there is some edge in  $G$  that is not in  $G_1$ , and hence  $\text{ord}(G_1) < \text{ord}(G)$ .

Otherwise  $V_{G_1} \neq V_G$ . Since  $G$  is connected, there is at least one vertex  $v_1 \in V_G \setminus V_{G_1}$  of distance one to  $V_{G_1}$ ; let  $e_1$  be an edge connecting  $v_1$  to a vertex in  $V_{G_1}$ . Continuing in this fashion, we get edges  $e_2, \dots, e_t$  and vertices  $v_2, \dots, v_t$  such that the graph  $G'$ , obtained as the union of  $G_1$  and  $e_1, \dots, e_t$  and  $v_1, \dots, v_t$  has  $t$  more

vertices than  $V_{G_1}$ ,  $t$  more edges (that are not self-loops), and  $V_{G'} = V_G$ . Note that  $\text{ord}(G_1) = \text{ord}(G')$ , since  $G'$  contains  $t$  more vertices and  $t$  more edges than  $G_1$ .

Since  $v_t$  is a leaf in  $G'$ , and it cannot be a leaf in  $G$ , it follows that  $G$  has an edge that is not in  $G'$ ; hence  $\text{ord}(G) > \text{ord}(G')$ . But  $\text{ord}(G') = \text{ord}(G_1)$ .  $\square$

*Proof of Lemma 8.4.* Let  $\psi$  be an element of  $\Psi_{<r}^+$ . Then there is a sequence of strictly increasing graphs

$$\psi_0 \subset \psi_1 \subset \cdots \subset \psi_t = \psi$$

where  $\psi_0$  is isomorphic to some  $\psi_i$ , and for each  $j \in [t]$ ,  $\psi_j$  is the union of  $\psi_{j-1}$  and a graph  $\tilde{\psi}_j$  that is isomorphic to one of the  $\psi_i$ . It follows that each  $\psi_j$  is pruned; the lemma about increasing order of pruned graphs shows that  $\text{ord}(\psi_t) > t + \text{ord}(\psi_0) \geq t + 1$ . Hence  $t \leq r - 2$ , and hence  $\psi$  is the union of  $r - 1$  graphs, each of which lies in a class in  $\Psi$ . Hence

$$\#E_\psi \leq (r - 1) \max_{i \in [t]} (\#E_{\psi_i})$$

which is bounded. Since  $\psi$  is positive,

$$\#V_\psi \leq \#E_\psi - \text{ord}(\psi) \leq \#E_\psi$$

so  $\#V_\psi$  is bounded. Hence there are only finitely many possible isomorphism classes of graphs,  $\psi$ , as graphs, and hence only finitely many possible  $B$ -graph classes in  $\Psi_{<r}^+$ .  $\square$

### 8.3. Proof of Lemma 8.5.

*Proof of Lemma 8.5.* The first claim is easy: if  $[\psi_{/B}] \in \Psi^+$ , then  $\psi_{/B}$  is the union of injective morphisms  $\psi_{/B}^i \rightarrow \psi_{/B}$ ; hence the image of any injective map  $u: \psi_{/B} \rightarrow G_{/B}$  is the union of the images of the compositions  $\psi_{/B}^i \rightarrow \psi_{/B}$  with  $u$ , which are injective morphisms. Since  $\Psi^+ \cap G$  contains all of these images, it contains  $u(\Psi_{/B})$ . Hence  $u$  factors through  $\Psi^+ \cap G$ .

For the second claim, let  $G_{/B}^0$  be the image of the injective morphism  $\psi_{/B} \rightarrow G_{/B}$ . Since  $\Psi^+ \cap G_{/B}$  is the union of images of injective maps from the  $\psi_{/B}^i$  to  $G_{/B}$ , there must be a strict inclusion of graphs

$$G_{/B}^0 \subset G_{/B}^1 \subset \cdots \subset G_{/B}^t = \Psi^+ \cap G_{/B}$$

where each  $G_{/B}^i$  is the union of  $G_{/B}^{i-1}$  and a  $B$ -subgraph of  $G_{/B}$  that lies in some element of  $\Psi$ . Let  $i$  be the smallest value such that  $\text{ord}(G^i) \geq r$ ; then  $i \geq 1$  (since  $\text{ord}(\psi) < r$  by assumption and  $G_{/B}^0$  and  $\psi_{/B}$  are isomorphic, so  $\text{ord}(G^0) < r$ ). Since  $G^i$  has at most  $s$  more edges than  $G^{i-1}$ , we have

$$\text{ord}(G^i) \leq \text{ord}(G^{i-1}) + s \leq r - 1 + s.$$

Hence the injection  $\psi_{/B} \rightarrow G$  factors through  $\psi'_{/B} \stackrel{\text{def}}{=} G_{/B}^i$ , and

$$\psi'_{/B} \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+$$

as desired; (24) follows because the number of injective morphisms  $\psi_{/B} \rightarrow G_{/B}$  that factor through  $\psi'_{/B}$  is at most the number of injective morphisms  $\psi_{/B} \rightarrow \psi'_{/B}$  times those  $\psi'_{/B} \rightarrow G_{/B}$ .  $\square$

**8.4. The Injection Count and Resulting Partial Order.** In this section we describe the partial order which we will later use to define the rational numbers  $\mu[\psi_{/B}]$  that are fundamental to our construction of “approximate indicator functions” (see (29)).

**Definition 8.8.** Let  $B$  be a graph. Recall that  $[S_{/B}]$  denotes the class of  $B$ -graphs that are isomorphic to  $S_{/B}$ ; write  $[S_{/B}] \leq_B [T_{/B}]$  if  $N(S_{/B}, T_{/B}) > 0$  (Definition 8.2), i.e., if there exists an injection  $S_{/B} \rightarrow T_{/B}$ ; we sometimes refer to  $\leq_B$  as  $\leq$  when confusion is unlikely to occur.

**Lemma 8.9.** *Let  $B$  be a graph. Then the relation  $\leq_B$  in Definition 8.8 is a partial order (of isomorphism classes of  $B$ -graphs).*

*Proof.* The relation  $\leq = \leq_B$  is clearly reflexive and transitive, so we need only show that it is anti-symmetric: so assume that  $[S_{/B}] \leq [T_{/B}]$  and  $[T_{/B}] \leq [S_{/B}]$ ; let us prove that  $[S_{/B}] = [T_{/B}]$ . Since  $[S_{/B}] \leq [T_{/B}]$ , there is a morphism  $\nu: S_{/B} \rightarrow T_{/B}$  that is injective; hence  $\#V_S \leq \#V_T$  and  $\#E_S^{\text{dir}} \leq \#E_T^{\text{dir}}$ ; then  $[T_{/B}] \leq [S_{/B}]$  provides the reverse inequalities, and hence  $\#V_S = \#V_T$  and  $\#E_S^{\text{dir}} = \#E_T^{\text{dir}}$ . It follows that  $\nu$  is bijective on the vertex sets and directed edge sets, and we easily check that the inverse map on these sets yields an isomorphism of  $B$ -graphs. Hence  $S_{/B}$  and  $T_{/B}$  are isomorphic, and hence  $[S_{/B}] = [T_{/B}]$ .  $\square$

**8.5. The Möbius Function.** Now let  $B$  be a graph, and let  $\mathcal{O}$  be any subset of the set of isomorphism classes of  $B$ -graphs; then the partial order  $\leq = \leq_B$  above restricts to give a partial order on  $\mathcal{O}$ . We now define a *Möbius function* for the partially ordered set,  $\mathcal{O}$ , in the usual way: we define a function  $\mu = \mu_{\mathcal{O}}: \mathcal{O} \rightarrow \mathbb{R}$  with the property that

$$(27) \quad \sum_{\substack{[S_{/B}] \in \mathcal{O} \\ [S_{/B}] \leq [T_{/B}]}} N([S_{/B}], [T_{/B}]) \mu_{\mathcal{O}}([S_{/B}]) = 1$$

for each  $[T_{/B}]$ , by defining

$$\mu([T_{/B}]) = 1/N([T_{/B}], [T_{/B}]) = 1/(\#\text{Aut}(T_{/B}))$$

when  $[T_{/B}]$  is a minimal element of  $\mathcal{O}$ , and then—by structural induction on  $T$ , or regular induction on  $(\#V_T) + (\#E_T^{\text{dir}})$ —we set

$$(28) \quad \mu_{\mathcal{O}}([T_{/B}]) = \frac{1}{N([T_{/B}], [T_{/B}])} \left( 1 - \sum_{\substack{[S_{/B}] \in \mathcal{O} \\ [S_{/B}] < [T_{/B}]}} N([S_{/B}], [T_{/B}]) \mu_{\mathcal{O}}([S_{/B}]) \right).$$

Notice that  $N(S_{/B}, S_{/B})$  can be strictly greater than one (e.g., which can happen if  $S$  is a cycle, or a “barbell graph” or “theta graph”; see Section 6 of Article I). For this reason, the  $\mu_{\mathcal{O}}([S_{/B}])$  are not necessarily integers.

**Definition 8.10.** Let  $B$  be a graph, and  $\mathcal{O}$  a subset of the set of isomorphism classes of  $B$ -graphs. By the *Möbius function on  $\mathcal{O}$*  we mean the unique function  $\mu([S_{/B}]) = \mu_{\mathcal{O}}([S_{/B}])$  satisfying (27), defined inductively by (28).

We remark that [Fri08] works with  $B$ -graphs (and graphs); here we work with  $B$ -graphs simply because our definition of algebraic model makes this convenient.

### 8.6. The Truncated Indicator Function and the Proof of Theorem 8.6.

**Definition 8.11.** Let  $B$  be a graph, and  $\Psi$  a set of isomorphism classes of  $B$ -graphs. Then  $\Psi^+$  is a set of isomorphism classes of  $B$ -graphs, and gives rise to a Möbius function  $\mu_{\Psi^+}$ . For each  $r$  we define the *order  $r$  truncated  $\Psi$ -indicator function* to be the function defined on  $B$ -graphs,  $G_{/B}$ , given by

$$(29) \quad I_r(\Psi, G_{/B}) \stackrel{\text{def}}{=} \sum_{[S_{/B}] \in \Psi_{<r}^+} N(S_{/B}, G_{/B}) \mu_{\Psi^+}([S_{/B}]).$$

This sum is finite for any given  $G_{/B}$  since  $N(S_{/B}, G_{/B}) = 0$  if  $S$  has more vertices than  $G$ ; in cases of interest to us we will require  $\Psi_{<r}^+$  to be finite for all  $r$ , so that the above sum involves finitely many  $[S_{/B}]$ .

*Proof of Theorem 8.6.* Let  $\mu[S_{/B}] = \mu_{\Psi^+}[S_{/B}]$  be the Möbius function for the partially ordered set  $\Psi^+$ . Let us prove the various claims in Theorem 8.6 regarding  $I_r(\Psi, G_{/B})$  for any  $B$ -graph  $G_{/B}$ .

Set  $\psi_{/B} = \Psi^+ \cap G_{/B}$ , and consider the three cases where  $\text{ord}_{\Psi}(G) = \text{ord}(\psi)$  is 0, between 1 and  $r-1$ , and at least  $r$ . In all cases it will be useful to note that by Lemma 8.5, for all  $[S_{/B}] \in \Psi^+$  we have that

$$N(S_{/B}, G_{/B}) = N(S_{/B}, \Psi^+ \cap G_{/B}) = N(S_{/B}, \psi_{/B}),$$

and hence  $I_r(\Psi, G_{/B}) = I_r(\Psi, \psi_{/B})$ .

First, consider the case where  $\text{ord}(\psi) = 0$ . Then  $\Psi^+ \cap G_{/B} = \emptyset_{/B}$ , the empty graph, and  $N(S_{/B}, G_{/B}) = N(S_{/B}, \psi_{/B}) = 0$  for all  $S_{/B} \in \Psi^+$ , and hence

$$I_r(\Psi, G_{/B}) = 0 = \mathbb{I}_{\text{Meets}(\Psi)}(G).$$

This proves (25) in this case.

Second, consider the case where  $1 \leq \text{ord}(\psi) < r$ , and hence  $\mathbb{I}_{\text{Meets}(\Psi)}(G) = 1$ . By Lemma 8.7, if  $[\psi_{/B}] \in \Psi^+$  and  $\text{ord}(\psi) < r$ , then for  $S_{/B} \in \Psi^+$  with  $\text{ord}(S) \geq r$  we have

$$N(S_{/B}, \psi_{/B}) = 0,$$

and hence

$$\begin{aligned} I_r(\Psi, G_{/B}) &= I_r(\Psi, \psi_{/B}) = \sum_{[S_{/B}] \in \Psi_{<r}^+} N(S_{/B}, \psi_{/B}) \mu[S_{/B}] \\ &= \sum_{[S_{/B}] \in \Psi^+} N(S_{/B}, \psi_{/B}) \mu[S_{/B}] = 1 = \mathbb{I}_{\text{Meets}(\Psi)}(G_{/B}) \end{aligned}$$

by (27). This proves (25) in this case.

Third and lastly, consider the case that  $\text{ord}(\psi) \geq r$ . Then (24) of Lemma 8.5 implies that for any  $[S_{/B}] \in \Psi^+$  with  $\text{ord}(S) < r$  we have

$$N(S_{/B}, G_{/B}) \leq \sum_{\psi'_{/B} \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} N(S_{/B}, \psi'_{/B}) N(\psi'_{/B}, B_{/B}).$$

This implies (26) for

$$C = \max_{\psi'_{/B} \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} \sum_{[S_{/B}] \in \Psi_{<r}^+} |\mu[S_{/B}]| N(S_{/B}, \psi'_{/B}),$$

which is a finite sum since  $\Psi_{<r}^+$  and  $\Psi_{<r+s}^+$  are finite sets by assumption.  $\square$

## 9. PROOF OF THEOREM 5.9

In this section we prove Theorem 5.9. We build up the proof with a sequence of lemmas; we start the following lemma based on Theorem 4.1.

**Lemma 9.1.** *Let  $B$  be a graph, and  $\Psi$  a finite family of  $B$ -graphs such that  $\Psi_{<s}^+$  is finite for all  $s \in \mathbb{N}$ . Then for any fixed  $\nu > 0$  and  $r', r'' \in \mathbb{N}$  we have that*

$$f(k, n) = \mathbb{E}_{G \in \mathcal{C}}[I_{r'}(\Psi, G_{/B}) \text{cert}_{<\nu, <r''}(G, k)]$$

has a  $(B, \nu)$ -Ramanujan expansion to any order  $r \in \mathbb{N}$

$$c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

where the bases of the  $c_i = c_i(k)$  are a subset of any set of eigenvalues of the model, and where  $c_i(k) = 0$  provided that  $i < r$  and  $i$  is less than the order of any  $B$ -graph occurring in  $\mathcal{C}_n(B)$  that meets  $\Psi$ , i.e., that lies in an element of  $\Psi$ .

As usual, the same lemma holds for the weakly-certified trace  $\text{cert}_{\leq\nu, <r''}(G, k)$ , but we shall not need this result.

*Proof.* Since  $\Psi_{<r'}^+$  is finite, (29) implies that  $I_{r'}(\Psi, G_{/B})$  is a finite linear combination of functions  $G_{/B} \mapsto N(\psi_{/B}, G_{/B})$ ; hence to prove this lemma, it suffices to prove such expansions exist for functions of the form

$$f(k, n) = \mathbb{E}_{G \in \mathcal{C}}[N(\psi_{/B}, G_{/B}) \text{cert}_{<\nu, <r''}(G, k)]$$

with  $[\psi_{/B}] \in \Psi_{<r'}^+$ . By subdividing the certified walks by their homotopy type and applying (19), it suffices to prove such expansions exist for functions of the form

$$\begin{aligned} f(k, n) &= \mathbb{E}_{G \in \mathcal{C}}[N(\psi_{/B}, G_{/B}) \text{snbc}(T^{\leq}, \geq \xi; G, k)] \\ &= \mathbb{E}_{G \in \mathcal{C}}[\#([\psi_{/B}] \cap G_{/B}) \text{snbc}(T^{\leq}, \geq \xi; G, k)] \end{aligned}$$

for a fixed  $[\psi_{/B}] \in \Psi_{<r'}^+$ , a fixed ordered graph,  $T^{\leq}$ , and a fixed  $\xi$  with

$$(30) \quad \mu_1(\text{VLG}(T, \xi)) < \nu.$$

But this follows from Theorem 4.1.  $\square$

**Lemma 9.2.** *Let  $B$  be a graph and  $\Psi$  a collection of finite isomorphism classes of  $B$ -graphs such that  $\Psi_{<r}^+$  is finite for all  $r$ . Let  $\mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)}(G_{/B})$  denote the indicator function of those  $B$ -graphs  $G_{/B}$  with  $\text{ord}_{\Psi}(G) \geq r$ . Then for any algebraic model  $\mathcal{C}_n(B)$  and  $r \in \mathbb{N}$  there is a  $C$  such that*

$$\begin{aligned} \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)}(G_{/B}) \text{cert}_{<\nu, <r}(G, k)] &\leq Cn^{-r+1} \mu_1(B)^k \\ \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)}(G_{/B}) I(G, \Psi, r) \text{cert}_{<\nu, <r}(G, k)] &\leq Cn^{-r+1} \mu_1(B)^k \\ \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)}(G_{/B}) I(G, \Psi, r)] &\leq Cn^{-r} \end{aligned}$$

for all  $k \in \mathbb{N}$ .

*Proof.* For any graph  $G$  we have

$$0 \leq \text{cert}_{<\nu, <r}(G, k) \leq \text{snbc}(G, k) \leq \text{Trace}(H_G^k) \leq n \text{Trace}(H_B^k) \leq n \mu_1^k(B) (\#E_B^{\text{dir}}).$$

Hence it suffices to show that both

$$(31) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{ord}_{\Psi}(\geq r)}(G_{/B})] \quad \text{i.e.,} \quad \text{Prob}_{G \in \mathcal{C}_n(B)}[\text{ord}_{\Psi}(G_{/B}) \geq r],$$

and

$$(32) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{ord}_{\Psi}(\cdot) \geq r}(G_{/B}) I_r(\Psi, G_{/B})]$$

are bounded by  $O(n^{-r})$ .

By Lemma 8.5, if  $\text{ord}_\Psi(G_{/B}) \geq r$  then  $N(\psi_{/B}, G_{/B}) \geq 1$  for some  $\psi_{/B} \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+$ . Hence

$$\begin{aligned} \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{ord}_\Psi(\cdot) \geq r}(G_{/B})] &\leq \sum_{[\psi_{/B}] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} \text{Prob}_{G \in \mathcal{C}_n(B)} [N(\psi_{/B}, G_{/B}) > 0] \\ &\leq \sum_{[\psi_{/B}] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} \mathbb{E}_{G \in \mathcal{C}_n(B)} [N(\psi_{/B}, G_{/B})]. \end{aligned}$$

Since the above sum is over finitely many classes  $\psi_{/B}$ , and since for each  $\psi_{/B}$  we have that

$$(33) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} [N(\psi_{/B}, G_{/B})] = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\#([\psi_{/B}^\leq] \cap G_{/B})] = O(n^{-\text{ord}(\psi)}) = O(n^{-r})$$

since  $\mathcal{C}_n(B)$  is algebraic, we get the desired bound on (31).

To get the desired bound on (32), since  $I_r(\Psi, G_{/B})$  is a linear combination of the functions  $N(\psi'_{/B}, G_{/B})$  with  $\psi'_{/B} \in \Psi_{<r}^+$ , it suffices to prove such a bound for each function

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{ord}_\Psi^{-1}(\geq r)}(G_{/B}) N(\psi'_{/B}, G_{/B})].$$

But if  $\text{ord}_\Psi(G) \geq r$ , then we have

$$N(\psi'_{/B}, G_{/B}) \leq C \sum_{[\psi_{/B}] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} \mathbb{E}_{G \in \mathcal{C}_n(B)} [N(\psi_{/B}, G_{/B})]$$

according to (24), where  $C$  is the maximum value of  $N(\psi'_{/B}, \psi_{/B})$  over  $[\psi_{/B}] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+$ ; hence it suffices to bound

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} [N(\psi_{/B}, G_{/B})]$$

for each  $\psi_{/B}$  with  $[\psi_{/B}] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+$ , which again is implied by (33).  $\square$

We will see that the following lemma almost immediately implies Theorem 5.9.

**Lemma 9.3.** *Let  $B$  be a graph, and  $\Psi$  a finite family of  $B$ -graphs such that  $\Psi_{<s}^+$  is finite for all  $s \in \mathbb{N}$ . Then for any fixed  $\nu > 0$  and  $r'' \in \mathbb{N}$  we have that*

$$(34) \quad f(k, n) = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\Psi)}(G_{/B}) \text{cert}_{<\nu, <r''}(G, k)]$$

has a  $(B, \nu)$ -Ramanujan expansion to any order  $r \in \mathbb{N}$

$$c_0(k) + \cdots + c_{r-1}(k)/n^{r-1} + O(1)c_r(k)/n^r,$$

where the bases of the  $c_i = c_i(k)$  are the exponents of the model, and where  $c_i(k) = 0$  provided that  $i$  is less than the order of any  $B$ -graph occurring in  $\mathcal{C}_n(B)$  that meets  $\Psi$ , i.e., that lies in an element of  $\Psi$ .

*Proof.* Let  $r' = r + 1$ . We have

$$(35) \quad 1 = \mathbb{I}_{\text{ord}_\Psi^{-1}(\geq r')}(G_{/B}) + \mathbb{I}_{\text{ord}_\Psi^{-1}(<r')}(G_{/B})$$

where  $\text{ord}_\Psi^{-1}(\geq r')$  and  $\text{ord}_\Psi^{-1}(<r')$  denote the set of  $B$ -graphs,  $G_{/B}$ , for which  $\text{ord}_\Psi(G_{/B})$  is, respectively,  $\geq r'$  and  $< r'$ . Using the facts that

- (1)  $\text{ord}_\Psi(G_{/B}) \geq r'$  implies that  $G \in \text{Meets}(\Psi)$ , and
- (2)  $\text{ord}_\Psi(G_{/B}) < r'$  implies that  $I_{r'}(\Psi, G_{/B}) = \mathbb{I}_{\text{Meets}(\Psi)}(G_{/B})$ ,



we have

$$\begin{aligned}
(36) \quad \mathbb{I}_{\text{Meets}(\Psi)}(G_{/B}) &= \mathbb{I}_{\text{Meets}(\Psi)}(G) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r')} (G_{/B}) + \mathbb{I}_{\text{Meets}(\Psi)}(G) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r')} (G_{/B}) \\
&= I_{r'}(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r')} (G_{/B}) + \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r')} (G_{/B}) \\
(37) \quad &= I_{r'}(\Psi, G_{/B}) - I_{r'}(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r')} (G_{/B}) + \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r')} (G_{/B})
\end{aligned}$$

Now we multiply the left-hand-side of (36) and (37) by  $\text{cert}_{<\nu, <r''}(G, k)$  and taking expected values: the left-hand-side becomes (34), and the individual summands of (37) become a sum of

$$(38) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} [I_{r'}(\Psi, G_{/B}) \text{cert}_{<\nu, <r''}(G, k)]$$

plus terms bounded by  $Cn^{-r'+1} \mu_1(B)^k = Cn^{-r} \mu_1(B)$  by Lemma 9.2. By Lemma 9.1, (38) has an  $(B, \nu)$ -bounded order  $r$  expansion, and hence so does (34).  $\square$

*Proof of Theorem 5.9.* If  $\mathcal{T}$  is a set of  $B$ -graphs, then let  $\Psi$  be the isomorphism classes of a finite set of positive generators of  $\mathcal{T}$ ; if  $\mathcal{T}$  is a set of graphs, let  $\Psi$  consist of all possible  $B$ -graph structures on elements of a finite set of positive generators of  $\mathcal{T}$ .

Then for all  $G = G_{/B} \in \mathcal{C}_n(B)$ ,  $G$  meets  $\mathcal{T}$  iff  $G_{/B}$  meets  $\Psi$ . According to Lemma 8.4,  $\Psi_{<r}^+$  is finite for all  $r$ . Now we apply Lemma 9.1.  $\square$

## 10. PROOF OF THEOREM 5.8

In this section we easily prove Theorem 5.8 based on the methods we have already developed.

*Proof of Theorem 5.8.* If  $\mathcal{T}$  is a set of  $B$ -graphs, then let  $\Psi$  be the isomorphism classes of a finite set of positive generators of  $\mathcal{T}$ ; if  $\mathcal{T}$  is a set of graphs, let  $\Psi$  consist of all possible  $B$ -graph structures on elements of a finite set of positive generators of  $\mathcal{T}$ . Hence  $\text{Meets}(\mathcal{T}) = \text{Meets}(\Psi)$ .

Fix an  $r \in \mathbb{N}$ . Let  $s$  be the largest number of edges in a graph in a class of  $\Psi$ . According to Lemma 8.4,  $\Psi_{r+s}^+$  is finite.

With notation as in (35), we have

$$\begin{aligned}
\mathbb{I}_{\text{Meets}(\Psi)}(G) &= \mathbb{I}_{\text{Meets}(\Psi)}(G) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} (G_{/B}) + \mathbb{I}_{\text{Meets}(\Psi)}(G) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)} (G_{/B}) \\
&= \mathbb{I}_{\text{Meets}(\Psi)}(G) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} (G_{/B}) + \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)} (G_{/B})
\end{aligned}$$

since  $\text{ord}_{\Psi}(G_{/B}) \geq r$  implies that  $G_{/B}$  meets  $\Psi$ . With  $I_r(\Psi, G_{/B})$  as in (29), we have

$$\mathbb{I}_{\text{Meets}(\Psi)}(G) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} (G_{/B}) = I_r(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} (G_{/B}),$$

since  $I_r(\Psi, G_{/B}) = \mathbb{I}_{\text{Meets}(\Psi)}(G)$  provided that  $\text{ord}_{\Psi}(G_{/B}) < r$ . Combining the above two displayed equations we have

$$\mathbb{I}_{\text{Meets}(\Psi)}(G) = I_r(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} (G_{/B}) + \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)} (G_{/B})$$

Taking expected values yields

$$(39) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\Psi)}(G)] = \mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} (G_{/B})] + O(n^{-r})$$

in view of Lemma 9.2. In view of the fact that  $\Psi_{<r+s}^+ \setminus \Psi_{<r}^+$  is finite, taking expected values in (26) yields

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)}(G_{/B})] \leq \sum_{[\psi_{/B}] \in \Psi_{<r+s}^+ \setminus \Psi_{<r}^+} O(n^{-\text{ord}(\psi)}) = O(n^{-r}).$$

Adding this to (39) yields

$$\begin{aligned} & \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\Psi)}(G)] + O(n^{-r}) \\ &= \mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)}(G_{/B})] + \mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B}) \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)}(G_{/B})] + O(n^{-r}), \end{aligned}$$

and, since  $\mathbb{I}_{\text{ord}_{\Psi}^{-1}(<r)} + \mathbb{I}_{\text{ord}_{\Psi}^{-1}(\geq r)} = 1$ ,

$$(40) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\Psi)}(G)] = \mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B})] + O(n^{-r}).$$

Next consider

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B})].$$

For each  $S_B \in \Psi_{<r}^+$ , we have

$$(41) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} [N(S_B, G_{/B})] = \mathbb{E}_{G \in \mathcal{C}_n(B)} [\# [S_B] \cap G_{/B}] = c_0 + c_1/n + \dots + c_{r-1}/n^{r-1} + O(1/n^r),$$

with  $c_i = 0$  for  $i < \text{ord}(S)$  since  $\mathcal{C}_n(B)$  is algebraic; furthermore  $c_i > 0$  if  $i = \text{ord}(S)$  and  $S_B$  occurs in  $\mathcal{C}_n(B)$ . Since  $\Psi_{<r}^+$  is finite and  $I_r(\Psi, G_{/B})$  is, by definition, a finite linear combination of functions of the form (41), we have

$$(42) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)} [I_r(\Psi, G_{/B})] = c_0 + c_1/n + \dots + c_{r-1}/n^{r-1} + O(1/n^r),$$

where if  $j$  is the minimum order of an element of  $\Psi_{<r}^+$  occurring in  $\mathcal{C}_n(B)$ , then with  $c_i = 0$  for  $i < j$  (assuming that  $r > i$  so that  $c_i$  is uniquely defined).

Now let  $j$  be the minimum order of an element of  $\Psi^+$  occurring in  $\mathcal{C}_n(B)$ . By Lemma 8.7, if  $[S_B] \in \Psi^+$  occurs in  $\mathcal{C}_n(B)$  and  $\text{ord}(S) = j$ , then  $[S_B]$  is a minimal element of the partially ordered set  $\Psi^+$ . [This implies that  $[S_B] \in \Psi$ , but this is inconsequential here.] It follows that

$$\mu[S_B] = 1/(\#\text{Aut}(S_B)) > 0.$$

It follows that  $c_j$  in (42) is the sum of  $\mu[S_B]$  over all such  $S_B$ , and hence  $c_j > 0$  (again, assuming  $r > j$  so that  $c_j$  is uniquely determined).

Combining (42) with (40) yields

$$\mathbb{E}_{G \in \mathcal{C}_n(B)} [\mathbb{I}_{\text{Meets}(\Psi)}(G)] = c_0 + c_1/n + \dots + c_{r-1}/n^{r-1} + O(1/n^r),$$

for the same  $c_i$  as in (42), as desired.  $\square$

## 11. CONCLUSION OF THE PROOFS OF THEOREMS 3.1 AND 3.2

In this section we complete the proof of Theorem 3.1. We will need the following straightforward fact.

**Lemma 11.1.** *Let  $\nu > 1$  and  $r \in \mathbb{N}$ , and let  $\psi$  be a minimal ( $\geq \nu, < r$ )-tangle (Definition 7.1, i.e.,  $\psi$  is a ( $\geq \nu, < r$ )-tangle but no proper subgraph of  $\psi$  is). Then  $\text{ord}(\psi) \geq 1$ .*

*Proof.* By definition a tangle is necessarily connected, and hence  $\psi$  is connected. Let us show that  $\psi$  is pruned: if  $v \in V_\psi$ ,  $v$  cannot be of degree one, incident upon an edge,  $e$ , joining  $v$  to some other vertex,  $w$ , for then we could “prune”  $\psi$ , discarding  $v$  and  $e$ ; this pruning preserves the  $\mu_1$  and the order, and does not affect the connectedness, which would contradict the minimality of  $\psi$ . Furthermore,  $v$  cannot be of degree one incident upon a half-loop, or of degree zero, since in either case  $V_\psi = \{v\}$  (since  $\psi$  is connected), and then  $\mu_1(\psi)$ , in either case, is 0. Hence each vertex of  $\psi$  is of degree at least two, and so  $\psi$  is pruned.

Since  $\mu_1$  of a cycle of any length equals 1, and  $\nu > 1$ ,  $\psi$  cannot be a cycle. Hence Lemma 6.4 shows that  $\text{ord}(\psi) \geq 1$ .  $\square$

*Proof of Theorem 3.1.* Let  $\mathcal{T} = \text{HasTangles}(\geq \nu, < r')$ . Each element of  $\mathcal{T}$  contains an  $(\geq \nu, < r')$ -tangle, and by Theorem 7.3 there are finitely many such tangles, up to isomorphism, that are minimal with respect to inclusion. According to Lemma 11.1, each minimal tangle has order at least one, and by definition any tangle is connected. Hence  $\mathcal{T}$  is finitely positively generated.

Hence we can apply Theorems 5.8 and 5.9 to  $\mathcal{T} = \text{HasTangles}(\geq \nu, < r)$ . Theorem 5.8 implies that

$$\text{Prob}_{\mathcal{C}_n(B)}[\text{HasTangles}(\geq \nu, < r')]$$

has an expansion to any order  $r$

$$c_0 + c_1/n + \cdots + c_{r-1}/n^{r-1} + O(1/n^r)$$

where  $c_i = 0$  for  $i < r$  and  $i < i_0$  where  $i_0$  is the smallest order of a  $(\geq \nu)$ -tangle. Since  $\nu > 1$ , Lemma 6.4 implies that  $i_0 \geq 1$ . Taking  $r \geq i_0$  (which we are free to do), since  $c_0 = 0$  and

$$(43) \quad \mathbb{I}_{\text{TangleFree}(\geq \nu, < r')} = 1 - \mathbb{I}_{\text{HasTangles}(\geq \nu, < r')},$$

it follows that

$$\text{Prob}_{\mathcal{C}_n(B)}[\text{TangleFree}(\geq \nu, < r')] = 1 - c_1/n - \cdots - c_{r-1}/n^{r-1} + O(1/n^r)$$

which, along with  $c_i = 0$  for  $i < i_0$ , establishes the part of Theorem 3.1 regarding the asymptotic expansions (and their coefficients) for (11).

Similarly we use (43), and subtract the result in Theorem 5.9 from that of Theorem 5.2 to obtain that for any  $r', r''$

$$\mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{TangleFree}(\geq \nu, < r')}(G) \text{cert}_{< \nu, r''}(G)]$$

has an expansion to any order  $r$ , whose coefficients  $c_i(k)$  have  $c_0(k)$  as in (10), since the  $1/n^i$ -coefficients in Theorem 5.9 vanish for  $i < r$  and  $i$  less than the smallest order of a  $(\geq \nu)$ -tangle, which is at least 1. Now we take  $r = r' = r'' \geq 1$ ; according to (14) we have

$$\mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}(G) \text{cert}_{< \nu, < r}(G, k) = \mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}(G) \text{snbc}_{< r}(G, k),$$

whereupon we have that

$$(44) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}(G) \text{snbc}_{< r}(G, k)]$$

has an expansion to order  $r$  with  $c_0(k)$  as in (10). Finally we note that

$$(45) \quad \mathbb{E}_{G \in \mathcal{C}_n(B)}[\mathbb{I}_{\text{TangleFree}(\geq \nu, < r)}(G) \text{snbc}_{\geq r}(G, k)] \leq \mathbb{E}_{G \in \mathcal{C}_n(B)}[\text{snbc}_{\geq r}(G, k)] \leq g(k)O(1)/n^r$$

where  $g$  is a function of growth  $\mu_1(B)$ . Adding (44) and (45), and using  $\text{snbc}(G, k) = \text{Trace}(H_G^k)$  establishes the claim in Theorem 3.1 for asymptotic expansions of (9).  $\square$

*Proof of Theorem 3.2.* We apply Theorem 5.8 with  $\mathcal{T} = [S_B]$ , which is positively generated since  $S$  is positive. Since  $S_B$  occurs in  $\mathcal{C}_n(B)$  by assumption, and  $[S_B]$  is the unique generator of  $\mathcal{T}$ , Theorem 5.8 implies that

$$\text{Prob}_{G \in \mathcal{C}_n(B)} \left[ [S_B] \cap G \neq \emptyset \right] = c_0 + c_1/n + \cdots + c_{r-1}/n^{r-1} + O(1/n^r)$$

for any  $r$ , where for  $i < r$  we have  $c_i = 0$  if  $i < \text{ord}(S)$  and  $c_i \neq 0$  if  $i = \text{ord}(S)$ . Taking any  $r > \text{ord}(S)$  yields

$$\text{Prob}_{G \in \mathcal{C}_n(B)} \left[ [S_B] \cap G \neq \emptyset \right] = c_i/n^i + O(1/n^{i+1})$$

for  $i = \text{ord}(S)$  with  $c_i > 0$ , which is bounded below by  $C'/n^i$  for any  $C' < c_i$  and  $n$  sufficiently large.  $\square$

#### REFERENCES

- [ABG10] Louigi Addario-Berry and Simon Griffiths, *The spectrum of random lifts*, December 2010, available as <http://arxiv.org/abs/1012.4097>, 35 pages.
- [Alo86] N. Alon, *Eigenvalues and expanders*, *Combinatorica* **6** (1986), no. 2, 83–96, Theory of computing (Singer Island, Fla., 1984). MR 88e:05077
- [AM85] N. Alon and V. D. Milman,  $\lambda_1$ , *isoperimetric inequalities for graphs, and superconcentrators*, *J. Combin. Theory Ser. B* **38** (1985), no. 1, 73–88. MR 782626 (87b:05092)
- [BC19] Charles Bordenave and Benoît Collins, *Eigenvalues of random lifts and polynomials of random permutation matrices*, *Ann. of Math. (2)* **190** (2019), no. 3, 811–875. MR 4024563
- [BL06] Yonatan Bilu and Nathan Linial, *Lifts, discrepancy and nearly optimal spectral gap*, *Combinatorica* **26** (2006), no. 5, 495–519. MR 2279667 (2008a:05160)
- [Bor15] Charles Bordenave, *A new proof of friedman’s second eigenvalue theorem and its extension to random lifts*, 2015.
- [BS87] Andrei Broder and Eli Shamir, *On the second eigenvalue of random regular graphs*, *Proceedings 28th Annual Symposium on Foundations of Computer Science*, 1987, pp. 286–294.
- [Dod84] Jozef Dodziuk, *Difference equations, isoperimetric inequality and transience of certain random walks*, *Trans. Amer. Math. Soc.* **284** (1984), no. 2, 787–794. MR 85m:58185
- [FK14] Joel Friedman and David-Emmanuel Kohler, *The relativized second eigenvalue conjecture of alon*, Available at <http://arxiv.org/abs/1403.3462>.
- [FKS89] J. Friedman, J. Kahn, and E. Szemerédi, *On the second eigenvalue of random regular graphs*, 21st Annual ACM Symposium on Theory of Computing, 1989, pp. 587–598.
- [Fri91] Joel Friedman, *On the second eigenvalue and random walks in random  $d$ -regular graphs*, *Combinatorica* **11** (1991), no. 4, 331–362. MR 93i:05115
- [Fri93a] ———, *Relative expansion and an extremal degree two cover of the boolean cube*, preprint, unpublished, available at [http://www.math.ubc.ca/~jfk/pubs/web\\_stuff/cover.html](http://www.math.ubc.ca/~jfk/pubs/web_stuff/cover.html).
- [Fri93b] ———, *Some geometric aspects of graphs and their eigenfunctions*, *Duke Math. J.* **69** (1993), no. 3, 487–525. MR 94b:05134
- [Fri03] ———, *Relative expanders or weakly relatively Ramanujan graphs*, *Duke Math. J.* **118** (2003), no. 1, 19–35. MR 1978881
- [Fri08] ———, *A proof of Alon’s second eigenvalue conjecture and related problems*, *Mem. Amer. Math. Soc.* **195** (2008), no. 910, viii+100. MR 2437174
- [HLW06] Shlomo Hoory, Nathan Linial, and Avi Wigderson, *Expander graphs and their applications*, *Bulletin of the American Mathematical Society* **43** (2006), no. 4, 439–561.
- [HPS18] Chris Hall, Doron Puder, and William F. Sawin, *Ramanujan coverings of graphs*, *Adv. Math.* **323** (2018), 367–410. MR 3725881

- [LP10] Nati Linial and Doron Puder, *Word maps and spectra of random graph lifts*, Random Structures Algorithms **37** (2010), no. 1, 100–135. MR 2674623
- [LPS88] A. Lubotzky, R. Phillips, and P. Sarnak, *Ramanujan graphs*, Combinatorica **8** (1988), no. 3, 261–277. MR 89m:05099
- [LSV11] Eyal Lubetzky, Benny Sudakov, and Van Vu, *Spectra of lifted Ramanujan graphs*, Adv. Math. **227** (2011), no. 4, 1612–1645. MR 2799807 (2012f:05181)
- [Mar88] G. A. Margulis, *Explicit group-theoretic constructions of combinatorial schemes and their applications in the construction of expanders and concentrators*, Problemy Peredachi Informatsii **24** (1988), no. 1, 51–60. MR 89f:68054
- [Mor94] Moshe Morgenstern, *Existence and explicit constructions of  $q + 1$  regular Ramanujan graphs for every prime power  $q$* , J. Combin. Theory Ser. B **62** (1994), no. 1, 44–62. MR 95h:05089
- [MSS15a] Adam W. Marcus, Daniel A. Spielman, and Nikhil Srivastava, *Interlacing families I: Bipartite Ramanujan graphs of all degrees*, Ann. of Math. (2) **182** (2015), no. 1, 307–325. MR 3374962
- [MSS15b] ———, *Interlacing families IV: Bipartite Ramanujan graphs of all sizes*, 2015 IEEE 56th Annual Symposium on Foundations of Computer Science—FOCS 2015, IEEE Computer Soc., Los Alamitos, CA, 2015, pp. 1358–1377. MR 3473375
- [Nil91] A. Nilli, *On the second eigenvalue of a graph*, Discrete Math. **91** (1991), no. 2, 207–210. MR 92j:05124
- [Pud15] Doron Puder, *Expansion of random graphs: new proofs, new results*, Invent. Math. **201** (2015), no. 3, 845–908. MR 3385636
- [Tan84] R. Michael Tanner, *Explicit concentrators from generalized  $N$ -gons*, SIAM J. Algebraic Discrete Methods **5** (1984), no. 3, 287–293. MR 752035 (85k:68080)

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z4, CANADA

*E-mail address:* `jf@cs.ubc.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA

*Current address:* 422 Richards St, Suite 170, Vancouver BC V6B 2Z4, CANADA

*E-mail address:* `David.kohler@3.epfl.ch`