

A RELATIVIZED ALON SECOND EIGENVALUE CONJECTURE FOR REGULAR BASE GRAPHS IV: AN IMPROVED SIDESTEPPING THEOREM

JOEL FRIEDMAN AND DAVID KOHLER

ABSTRACT. This is the fourth in a series of articles devoted to showing that a typical covering map of large degree to a fixed, regular graph has its new adjacency eigenvalues within the bound conjectured by Alon for random regular graphs.

In this paper we prove a *Sidestepping Theorem* that is more general and easier to use than earlier theorems of this kind. Such theorems concerns a family probability spaces $\{\mathcal{M}_n\}$ of $n \times n$ matrices, where n varies over some infinite set, N , of natural numbers. Many trace methods use simple “Markov bounds” to bound the expected spectral radius of elements of \mathcal{M}_n : this consists of choosing one value, $k = k(n)$, for each $n \in N$, and proving expected spectral radius bounds based on the expected value of the trace of the $k = k(n)$ -power of elements of \mathcal{M}_n . *Sidestepping* refers to bypassing such simple Markov bounds, obtaining improved results using a number of values of k for each fixed $n \in N$.

In more detail, if the $M \in \mathcal{M}_n$ expected value of $\text{Trace}(M^k)$ has an asymptotic expansion in powers of $1/n$, whose coefficients are “well behaved” functions of k , then one can get improved bounds on the spectral radius of elements of \mathcal{M}_n that hold with high probability. Such asymptotic expansions are shown to exist in the third article in this series for the families of matrices that interest us; in the fifth and sixth article in this series we will apply the Sidestepping Theorem in this article to prove the main results in this series of articles.

This article is independent of all other articles in this series; it can be viewed as a theorem purely in probability theory, concerning random matrices or, equivalently, the n random variables that are the eigenvalues of the elements of \mathcal{M}_n .

CONTENTS

1. Introduction	2
1.1. Motivation and Rough Idea Behind the Sidestepping Theorem	3
2. Main Results	4
2.1. Preliminary Notation	4
2.2. \mathbb{E}_{in} and \mathbb{E}_{out}	5
2.3. Polyexponentials and Approximate Polyexponentials	5
2.4. (Λ_0, Λ_1) -Bounded Matrix Models	6
2.5. The Sidestepping Theorem	7
2.6. The Exceptional Eigenvalue Bound	7

Date: Thursday 14th November, 2019.

2010 *Mathematics Subject Classification.* Primary 68R10.

Research supported in part by an NSERC grant.

Research supported in part by an NSERC grant.

2.7. The Sidestepping Lemma	8
3. The Shift Operator and The Proof of Lemma 2.5	9
3.1. The Shift Operator	9
3.2. Real Eigenvalue Bounds	10
3.3. Intermediate Lemmas	12
3.4. The Proof of Lemma 3.3	13
3.5. Proof of Lemma 3.4	14
3.6. Proof of Lemma 2.5	14
4. Proof of The Sidestepping Lemma (Lemma 2.6)	15
4.1. A Variant of Lemma 3.3	15
4.2. A Variant of Lemma 3.4	16
4.3. A Shift Computation	19
4.4. Proof of Lemma 2.6	20
5. Proof of the Sidestepping Theorem	24
References	24

1. INTRODUCTION

This is the fourth article in a series of six articles devoted to developing trace methods to prove a relativization of the Alon Second Eigenvalue Conjecture for covering maps of a fixed base graph that is regular; we also get a sharper theorem when the base graph is *Ramanujan*. This article is independent of all the other articles and most of the terminology they use; the results here are purely theorems in probability theory.

The main goal of this article is to prove a strengthening of the *Sidestepping Lemma* of [Fri08], i.e., Lemma 11.1 there, that allows us to infer—with very high probability—bounds on the eigenvalues of random $n \times n$ matrices, provided that the expected values of their k -power traces satisfy certain “asymptotic expansions.” Such asymptotic expansions are proven in Articles I—III, the first three articles in this series, for random matrices that express the *new Hashimoto* matrix (i.e., *new non-backtracking* matrix) of random matrices of the random covering maps to a fixed base graph. Applying the main result in this article to those of Articles III (i.e., the third article in this series) proves fundamental results about the location of the eigenvalues of such random matrices when the base graph is regular. This application forms the basis of Articles V and VI of this series, where the two main results of this series of articles are proven.

After proving the *Sidestepping Lemma* in this article, akin to that of [Fri08], we will give a consequence of this lemma which we call the *Sidestepping Theorem*; this theorem is simpler to state, and suffices for our applications in Articles V and VI. Hence we state the Sidestepping Theorem as the main result in this paper, and it is this theorem that we will quote in Article V.

The rest of this article is organized as follows. In Section 2 we precisely state the Sidestepping Theorem, as well as the two main lemmas that are used to prove it; we call these lemmas the Exceptional Eigenvalue Bound and the Sidestepping Lemma, and these lemmas have direct analogs in [Fri08]. In Section 3 we introduce a number of tools—most of which are connected to the *shift operator*—and prove the Exceptional Eigenvalue Bound. In Section 4 we prove the Sidestepping Lemma,

which uses the Exceptional Eigenvalue Bound and develops additional properties of the shift operator. In Section 5 we easily prove the Sidestepping Theorem as a consequence of the Sidestepping Lemma.

For the rest of the section we motivate the Sidestepping Theorem in this article and make some rough remarks on its proof.

1.1. Motivation and Rough Idea Behind the Sidestepping Theorem. The most basic trace methods use a simple Markov-type bound as follows: if M is an $n \times n$ matrix with real or complex entries and with real eigenvalues, then if we use $\rho(M)$ to denote the spectral radius of M (i.e., the largest absolute value among M 's eigenvalues), then for any even integer $k > 0$ we have

$$\rho(M)^k \leq \text{Trace}(M^k);$$

it follows that if \mathcal{M} is a probability space of such matrices, then for any $\alpha > 0$

$$(1) \quad \text{Prob}_{M \in \mathcal{M}}[\rho(M) \geq \alpha] \leq \alpha^{-k} \mathbb{E}_{M \in \mathcal{M}}[\text{Trace}(M^k)].$$

This is a simple Markov-type bound, used in [BS87, Fri91, Fri03, LP10, Pud15] to bound the *new adjacency spectral radius* of random covers of a base graph. Unfortunately a direct Markov method like this cannot yield the optimal eigenvalue bound conjectured to hold with high probability by Alon (although [Pud15] obtains results that are quite close). The problem is that certain *tangles* cause

$$\mathbb{E}_{M \in \mathcal{M}}[\text{Trace}(M^k)]$$

to be too large for (1) to be useful. We use the term “sidestepping” for results that avoid this direct Markov-type approach.

In more detail, consider the situation where \mathcal{M}_n is a probability space of such $n \times n$ matrices, defined for an infinite number of positive integers, n , such that for any some integer $r > 1$ one has an asymptotic expansion

$$(2) \quad \mathbb{E}_{M \in \mathcal{M}_n}[\text{Trace}(M^k)] = f_0(k) + f_1(k)/n + \cdots + f_{r-1}(k)/n^{r-1} + O(1)f_r(k)/n^r$$

valid for all $k \leq n^{1/2}$. To fix ideas, say that there are real $\Lambda_1 > \Lambda_0 > 0$ such that for $i \leq r-1$ we have $f_i(k) = O(\Lambda_0^k)$ and that $f_r(k) = O(\Lambda_1^k)$. Then choosing k to balance the bounds on $f_0(k)$ and $f_r(k)/n^r$ (so k will be proportional to $\log n$), the Markov-type bound (1) gives a high probability bound for $\rho(M)$ of at most

$$\Lambda_0^{(r-1)/r} \Lambda_1^{1/r},$$

which is a function tending to Λ_0 as $r \rightarrow \infty$.

We remark that later in our main theorems, the upper bounds $O(\Lambda_0^k)$ and $O(\Lambda_1^k)$ (that appear in this section) will be replaced by slightly weaker upper bounds; we write the bounds $O(\Lambda_0^k)$ and $O(\Lambda_1^k)$ for simplicity, to illustrate the main ideas.

The Sidestepping Theorem deals with the following situation: say that for any r we have an expansion (2), but that for some $i < r$ we have that $f_i(k)$ can be written as $O(\Lambda_0^k)$ plus a finite number of terms of the form $Ck^a \lambda^k$ for an integer a , $C \in \mathbb{C}$, and λ is real with $\Lambda_0 < |\lambda| \leq \Lambda_1$; we call any sum of such terms *polyexponentials in k* . Then we can still infer a high probability bound of $\Lambda_0 + g(r)$ where $g(r) \rightarrow 0$ as $r \rightarrow \infty$, under appropriate conditions that will be made precise in Section 2.

Our proof uses the following ideas. Let S denote the “shift operator in k ,” i.e., S acts on function of k via $(Sf)(k) \stackrel{\text{def}}{=} f(k+1)$. Then the function $f(k) \stackrel{\text{def}}{=} Ck^a \lambda^k$ as above is annihilated by the operator $(S - \lambda)^{a+1}$. It follows that there is some polynomial $P = P(S)$ that annihilates all the polyexponential terms in the $f_i(k)$

for $i = 0, \dots, r-1$ in (2). Any fixed such polynomial $P = P(S)$ takes a function in k bounded by $O(\rho^k)$ to function bounded by $O(k^d \rho^k)$ (where $d = \deg(P)$). So now we apply $P(S)$ to both sides of (2) and try to infer information on the location of the eigenvalues of $M \in \mathcal{M}_n$. Furthermore, it is easy to see that for any polynomial, P ,

$$P(S)(\lambda^k) = P(\lambda)\lambda^k$$

(where λ^k on the left-hand-side is shorthand for the function $k \mapsto \lambda^k$); this allows us to control the effect of $P(S)$ on the expected value of $\text{Trace}(M^k)$ provided that λ is real and $\Lambda_0 < |\lambda| \leq \Lambda_1$.

Of course, in the above one is using the fact that (2) holds for many values of k for a given n , and hence one can apply polynomials in S to (2); furthermore, the left-hand-side becomes the expected value of the sum of $P(S)\lambda_i(M)^k$ over the eigenvalues, $\lambda_i(M)$, of M , rather than merely $\lambda_i(M)^k$. By contrast, the Markov-type bound only uses a single value of k . [In Markov-type methods and our sidestepping methods, all the values of k that we use are proportional to $\log n$.]

Our Sidestepping Theorem is stated somewhat more generally, because the non-backtracking matrix or Hashimoto matrix of a d -regular graph can also have complex eigenvalues, but only those of absolute value $(d-1)^{1/2}$. In our applications, $\Lambda_1 = d-1 + \epsilon$ and $\Lambda_0 = (d-1)^{1/2} + \epsilon$, where d is fixed and $\epsilon > 0$ is small, and $f_i(k)$ begins to contain polyexponential terms $Ck^a \lambda^k$ with $\lambda > \Lambda_0$ for i at least of order roughly $d^{1/2}$.

We remark that our Sidestepping Theorem concerns only the random variables $\lambda_i(M)$, so that one could formulate this theorem without any reference to matrices, only referring to the random variables $\lambda_1(M), \dots, \lambda_n(M)$ of $M \in \mathcal{M}_n$. However, we prefer to state the Sidestepping Theorem in (its equivalent form) involving eigenvalues of matrices, because this is closer to our intuition and is the context in which we apply the theorem.

2. MAIN RESULTS

In this section we state the main results in the paper, namely Lemmas 2.5 and 2.6, and the result that we need for Article V, namely Theorem 2.4. For convenience, we repeat some of the notation and definitions in Article I.

2.1. Preliminary Notation. In this subsection we introduce some notation that is completely standard, except for possibly (3) below.

We use $\mathbb{R}, \mathbb{C}, \mathbb{Z}, \mathbb{N}$ to denote, respectively, the the real numbers, the complex numbers, the integers, and positive integers or natural numbers; we use $\mathbb{Z}_{\geq 0}$ to denote the set of non-negative integers. If $n \in \mathbb{N}$ we use $[n]$ to denote $\{1, 2, \dots, n\}$.

If $\rho \in \mathbb{R}$ is non-negative and $z \in \mathbb{C}$, we use $B_\rho(z)$ to denote the *closed* ball of radius ρ about z , i.e., the set $\{x \in \mathbb{C} \mid |x - z| \leq \rho\}$. If $a, b \in \mathbb{R}$, we use $[a, b]$ to denote the closed interval $\{x \mid a \leq x \leq b\} \subset \mathbb{R}$; we often use the inclusion $\mathbb{R} \subset \mathbb{C}$ (as the complex numbers with vanishing imaginary part), to view subsets of \mathbb{R} as subsets of \mathbb{C} .

If $L \subset \mathbb{C}$ and $\rho \geq 0$ is real, we use the (fairly standard) notation

$$(3) \quad B_\rho(L) = \bigcup_{\ell \in L} B_\rho(\ell) .$$

All probability spaces are finite; hence a probability space is a pair $\mathcal{P} = (\Omega, P)$ where Ω is a finite set and $P: \Omega \rightarrow \mathbb{R}_{>0}$ with $\sum_{\omega \in \Omega} P(\omega) = 1$; hence an *event*

is any subset of Ω . We emphasize that $\omega \in \Omega$ implies that $P(\omega) > 0$ with strict inequality (i.e., no atoms have zero probability). We use \mathcal{P} and Ω interchangeably when P is understood and confusion is unlikely.

A *complex-valued random variable* on \mathcal{P} or Ω is a function $f: \Omega \rightarrow \mathbb{C}$, and similarly for real-, integer-, and natural-valued random variable; we denote its \mathcal{P} -expected value by

$$\mathbb{E}_{\omega \in \Omega}[f(\omega)] = \sum_{\omega \in \Omega} f(\omega)P(\omega).$$

If $\Omega' \subset \Omega$ we denote the probability of Ω' by

$$\text{Prob}_{\mathcal{P}}[\Omega'] = \sum_{\omega \in \Omega'} P(\omega') = \mathbb{E}_{\omega \in \Omega}[\mathbb{I}_{\Omega'}(\omega)],$$

where $\mathbb{I}_{\Omega'}$ denotes the indicator function of Ω' . At times we write $\text{Prob}_{\mathcal{P}}[\Omega']$ and $\mathbb{I}_{\Omega'}$ where Ω' is not a subset of Ω , by which we mean $\text{Prob}_{\mathcal{P}}[\Omega' \cap \Omega]$ and $\mathbb{I}_{\Omega' \cap \Omega}$.

A *matrix-valued random variable* (with entries in \mathbb{R} or \mathbb{C}) is similarly defined.

2.2. $\mathbb{E}\text{in}$ and $\mathbb{E}\text{out}$. If \mathcal{M} is a probability space of square matrices, and $R \subset \mathbb{C}$, it will be very useful to define

$$(4) \quad \mathbb{E}\text{in}_{\mathcal{M}}[R] \quad \text{and} \quad \mathbb{E}\text{out}_{\mathcal{M}}[R],$$

respectively, as the expected number of eigenvalues of $M \in \mathcal{M}$ (counted with multiplicity) that lie, respectively, in R and not in R . Of course, these two expected values are non-negative; furthermore, if \mathcal{M} consists entirely of $n \times n$ matrices, then

$$\mathbb{E}\text{in}_{\mathcal{M}}[R] + \mathbb{E}\text{out}_{\mathcal{M}}[R] = n.$$

2.3. Polyexponentials and Approximate Polyexponentials.

Definition 2.1. We say that a function $Q: \mathbb{N} \rightarrow \mathbb{C}$ is a *polyexponential function* if it is a linear combination of functions of the form

$$f(k) = k(k-1)\dots(k-m+1)\ell^{k-m}$$

for some $m \in \mathbb{Z}_{\geq 0}$ and $\ell \in \mathbb{C}$, or, equivalently, if we have

$$(5) \quad Q(k) = \sum_{\ell \in L} p_{\ell}(k)\ell^k$$

where $p_{\ell} = p_{\ell}(k)$ are polynomials and $L \subset \mathbb{C}$ is a finite set, and where the expression $p_{\ell}(k)\ell^k$ with $\ell = 0$ is taken to mean any function that vanishes for k sufficiently large. It is easy to see that the representation (5) is unique if we insist that L is minimal, i.e., $\ell \in L$ appears only if it is needed, i.e., if $p_{\ell}(k)\ell^k$ is a nonzero function; we refer to the minimal set L as the *set of bases of Q* , and if $\ell \in L$ we refer to the function $p_{\ell}(k)\ell^k$ as the *ℓ -part of Q* .

For example, if $n \in \mathbb{N}$, $i, j \in [n]$, and M is any $n \times n$ matrix over the complex numbers, then $f(k) = (M^k)_{i,j}$ (the i, j -th entry of M^k) is a polyexponential function of k . The fact that a Jordan block with eigenvalue 0 is nilpotent justifies our convention that for $\ell = 0$, $p_{\ell}(k)\ell^k$ refers to any function of finite support.

Definition 2.2. Let $\rho \in \mathbb{R}$. We say that a function $f: \mathbb{N} \rightarrow \mathbb{C}$ is *of growth ρ* if for every $\epsilon > 0$ we have that $|f(k)| < (\rho + \epsilon)^k$ for k sufficiently large. By an *approximate polyexponential with error growth ρ* we mean a function $Q: \mathbb{N} \rightarrow \mathbb{C}$ that can be written as $Q(k) = f(k) + g(k)$, where g is of growth ρ and f is a polyexponential; given such a Q and ρ , it is easy to see that f and g are uniquely

determined if we insist that all bases of f are greater than ρ in absolute value (if $|\ell| \leq \rho$, we may move a term $p_\ell(k)\ell^k$ of f to g); we refer to this unique f as the *polyexponential part* of Q (with respect to ρ), and to the bases of f as the *larger bases* of Q (with respect to ρ); if ℓ is one of these larger bases, then the ℓ -part of Q is the ℓ -part of f , i.e., the term $p_\ell(k)\ell^k$ in the sum

$$f(k) = \sum_{\ell \in L} p_\ell(k)\ell^k.$$

2.4. (Λ_0, Λ_1) -Bounded Matrix Models. We now define the type of families of random matrices that are of interest to us in this article.

Definition 2.3. Let $\Lambda_0 < \Lambda_1$ be positive real numbers. By a (Λ_0, Λ_1) -bounded matrix model we mean a collection of finite probability spaces $\{\mathcal{M}_n\}_{n \in N}$ where $N \subset \mathbb{N}$ is an infinite subset, and where the atoms of \mathcal{M}_n are $n \times n$ real-valued matrices whose eigenvalues lie in the set

$$B_{\Lambda_0}(0) \cup [-\Lambda_1, \Lambda_1]$$

in \mathbb{C} . Let $r \geq 0$ be an integer and $K: \mathbb{N} \rightarrow \mathbb{N}$ be a function such that $K(n)/\log n \rightarrow \infty$ as $n \rightarrow \infty$. We say that this model has an *order r expansion* with *range $K(n)$* (with Λ_0, Λ_1 understood) if as $n \rightarrow \infty$ we have that

$$(6) \quad \mathbb{E}_{M \in \mathcal{M}_n}[\text{Trace}(M^k)] = c_0(k) + c_1(k)/n + \cdots + c_{r-1}(k)/n^{r-1} + O(c_r(k))/n^r$$

for all $n \in N$ and integers k with $1 \leq k \leq K(n)$, where (1) $c_r = c_r(k)$ is of growth Λ_1 , (2) the constant in the $O(c_r(k))$ is independent of k and n , and (3) for $0 \leq i < r$, $c_i = c_i(k)$ is an approximate polyexponential with Λ_0 error term and whose larger bases (i.e., larger than Λ_0 in absolute value) lie in $[-\Lambda_1, \Lambda_1]$; at times we speak of an *order r expansion* without explicitly specifying K . When the model has such an expansion, then we use the notation L_r to refer to the union of all larger bases of $c_i(k)$ (with respect to Λ_0) over all i between 0 and $r-1$, and call L_r the *larger bases (of the order r expansion)*.

[We remark that since Λ_0 is positive, it is easy to see that the expansion (6) also holds for $k = 0$ (by possibly redefining $c_r(0)$ so that it is positive); we prefer to insist that $k \geq 1$ so that in Articles I–III we don't need to worry about certain technicalities when we define *algebraic models* there (such as the empty graph and non-backtracking walks of length 0).]

The main theorems in this paper concern a single, fixed matrix model (with fixed Λ_0, Λ_1 as above), and we will reserve the symbols \mathcal{M}_n and L_r (and Λ_0, Λ_1) to have the above meaning. We also note that (6) implies that for fixed $k \in \mathbb{N}$,

$$(7) \quad c_i(k) = \lim_{n \in N, n \rightarrow \infty} \left(\mathbb{E}_{M \in \mathcal{M}_n}[\text{Trace}(M^k)] - (c_0(k) + \cdots + c_{i-1}(k)/n^{i-1}) \right) n^i$$

for all $i \leq r-1$; we conclude that the $c_i(k)$ are uniquely determined, and that $c_i(k)$ is independent of r for any $r > i$ for which (6) holds. We also see that if (6) holds for some value of r , then it also holds for smaller values of r . It follows that if (6) holds for some r , then L_i is defined for each $i < r$ (as the set of larger bases of the functions $c_0(k), \dots, c_{i-1}(k)$). Furthermore L_i is empty iff $c_0(k), \dots, c_{i-1}(k)$ are all functions of growth Λ_0 .

We remark that in our applications (see Article V), N , the set of n for which \mathcal{M}_n is defined, will generally be a proper subset of \mathbb{N} ; indeed, n will be multiples

of the number of directed edges of a fixed *base graph*, B , since \mathcal{M}_n are related to the Hashimoto matrix, H_G , of random covers, G , of B .

For brevity we often write $n \rightarrow \infty$ or $\lim_{n \rightarrow \infty}$ understanding that n is confined to the set N for which \mathcal{M}_n is defined. Similarly, when we write an equation involving n , especially one involving \mathcal{M}_n , we understand n to be confined to N .

2.5. The Sidestepping Theorem. For Article V we need only the following theorem. This theorem will be proven in two main steps, represented in Lemmas 2.5 and 2.6 below; the theorem follows almost immediately from Lemma 2.6.

Theorem 2.4. *Let $\{\mathcal{M}_n\}_{n \in N}$ be a (Λ_0, Λ_1) -bounded matrix model, for some real $\Lambda_0 < \Lambda_1$, that for all $r \in \mathbb{N}$ has an order r expansion; let $p_i(k)$ denote the polyexponential part of $c_i(k)$ (with respect to Λ_0) in (6) (which is independent of $r \geq i + 1$ by (7)). If $p_i(k) = 0$ for all $i \in \mathbb{Z}_{\geq 0}$, then for all $\epsilon > 0$ and $j \in \mathbb{Z}_{\geq 0}$*

$$(8) \quad \mathbb{E}\text{out}_{\mathcal{M}_n}[B_{\Lambda_0+\epsilon}(0)] = O(n^{-j}).$$

Otherwise let j be the smallest integer for which $p_j(k) \neq 0$. Then for all $\epsilon > 0$, and for all $\theta > 0$ sufficiently small we have

$$(9) \quad \mathbb{E}\text{out}_{\mathcal{M}_n}[B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1})] = o(n^{-j});$$

moreover, if $L = L_{j+1}$ is the (necessarily nonempty) set of bases of p_j , then for each $\ell \in L$ there is a real $C_\ell > 0$ such that

$$(10) \quad p_j(k) = \sum_{\ell \in L} \ell^k C_\ell,$$

and for all $\ell \in L$ for sufficiently small $\theta > 0$,

$$(11) \quad \mathbb{E}\text{in}_{\mathcal{M}_n}[B_{n^{-\theta}}(\ell)] = n^{-j} C_\ell + o(n^{-j}).$$

In the above theorem, $L = L_{j+1}$ as in Definition 2.3, and this definition implies that

$$L \subset [-\Lambda_1, \Lambda_1] \setminus [-\Lambda_0, \Lambda_0].$$

We remark that in our applications $p_i(k) \neq 0$ for some i ; the theorem then states that the smallest j for which $p_j(k) \neq 0$ corresponds to the power of $1/n$ such that eigenvalues begin to appear with probability order n^{-j} near the bases of $p_j(k)$.

2.6. The Exceptional Eigenvalue Bound. The first step in proving the Sidestepping Theorem is what we call the *Exceptional Eigenvalue Bound*; this was called the *First Exception Bound* in Section 2.6 of [FK14] (see Subsection 2.6.3 there). It is interesting in its own right.

Lemma 2.5. *Let $\epsilon, \alpha, \Lambda_0, \Lambda_1 > 0$ be positive real numbers with $\Lambda_0 < \Lambda_1$. Then there is a positive integer $r_0 = r_0(\Lambda_0, \Lambda_1, \epsilon, \alpha)$ and a real $\theta_0 = \theta_0(\Lambda_0, \Lambda_1, \epsilon, \alpha) > 0$ with the following property: let $\{\mathcal{M}_n\}_{n \in N}$ be a (Λ_0, Λ_1) -bounded matrix model that has an order r expansion (6) for some $r \geq r_0$, and let L_r be the set of larger bases of $c_0(k), \dots, c_{r-1}(k)$. Then for any $\theta \leq \theta_0$,*

$$(12) \quad \mathbb{E}\text{out}_{\mathcal{M}_n}[B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L_r)] \leq n^{-\alpha}$$

for n sufficiently large.

In other words, an eigenvalue of $M \in \mathcal{M}_n$ that lies outside of $B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L_r)$ is “exceptional” in the sense that the expected number of such eigenvalues decays larger than any fixed power of n (for $\epsilon, \theta > 0$ sufficiently small), for models with expansions of sufficiently large order r .

Of course, if (12) holds for θ, r , then it holds for any larger r and smaller $\theta > 0$ since $B_{n^{-\theta}}(L_r)$ is non-decreasing as r increases and θ decreases.

We remark that to prove the above lemma we will take $r_0 = r_0(\Lambda_0, \Lambda_1, \epsilon, \alpha)$ to be any integer such that

$$(13) \quad r_0 > \alpha + (\alpha + 1) \frac{\log(\Lambda_1) - \log(\Lambda_0 + \epsilon)}{\log(\Lambda_0 + \epsilon) - \log(\Lambda_0)}$$

(see (37) and (38) in the next Section). Hence for $\Lambda_0, \Lambda_1, \alpha$ fixed and ϵ small, which is the case in our applications, we have that r_0 is proportional to $1/\epsilon$, in view of the denominator in the fraction in (13). Hence for small ϵ we need an asymptotic expansion of order proportional to $1/\epsilon$ to be able to apply the above lemma.

2.7. The Sidestepping Lemma. We now state the main lemma of this article, which is a version of the Sidestepping Lemma of [Fri08] that is more general and easier to apply in our situation. It easily implies Theorem 2.4, which is easier to state and sufficient for us in Articles V and VI.

Lemma 2.6. *Let $\Lambda_0 < \Lambda_1$ and ϵ be positive real numbers with $\Lambda_0 + \epsilon \leq \Lambda_1$, and let $j \geq 0$ be an integer. Then there are $r_1 = r_1(\Lambda_0, \Lambda_1, j, \epsilon)$ and $\theta_1 = \theta_1(\Lambda_0, \Lambda_1, j, \epsilon)$ with the following properties: let $\{\mathcal{M}_n\}_{n \in \mathbb{N}}$ be a (Λ_0, Λ_1) -bounded matrix model that has an order r expansion (6) for some $r \geq r_1(\Lambda_0, \Lambda_1, j, \epsilon)$. Recall that for each $i < r$, L_i denotes the larger bases of $c_0(k), \dots, c_{i-1}(k)$ (with respect to Λ_0). If $L_j = \emptyset$, then*

$$(14) \quad \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0+\epsilon}(0)] = O(n^{-j}).$$

In more detail, for any positive $\theta \leq \theta_1$ we have

$$(15) \quad \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1})] = o(n^{-j}).$$

Furthermore, for each $\ell \in L_{j+1}$ with $|\ell| \geq \Lambda_0 + 2\epsilon/3$ we have

$$(16) \quad \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] = C_\ell n^{-j} + o(n^{-j}),$$

for some real $C_\ell \geq 0$. Furthermore, if $q_{\ell,j} = q_{\ell,j}(k)$ are the polynomials in k such that

$$p_j(k) = \sum_{\ell \in L_{j+1}} q_{\ell,j}(k) \ell^k$$

is the polyexponential part of $c_j(k)$ (with respect to Λ_0), then if $|\ell| \geq \Lambda_0 + 2\epsilon/3$ we have

$$(17) \quad q_{\ell,j}(k) = C_\ell$$

is a constant function in k .

It other words, if $c_0(k), \dots, c_{j-1}(k)$ are of growth Λ_0 , then we know the “expected location” of the eigenvalues of $M \in \mathcal{M}_n$ “to order n^{-j} ” in terms of the polyexponential part, $p_j(k)$, of $c_j(k)$ with respect to Λ_0 (this requires that ϵ above is taken small enough so that all the larger bases (with respect to Λ_0) of c_j are larger than $\Lambda_0 + 2\epsilon/3$); furthermore, this $p_j(k)$ is a linear combination of purely exponential functions of k , and the coefficients in this linear combination reflects the

excepted number (times n^{-j}) of eigenvalues near each base of $p_j(k)$. Of course, it is possible that $p_j(k)$ is identically zero, i.e., also L_{j+1} is empty; in such a situation the hypotheses of the Sidestepping Lemma also hold when j is replaced with $j+1$, and we get more precise results provided that \mathcal{M}_n has expansions of sufficiently large order.

We remark that in the above lemma, if $\theta \leq 0$, then the sets $B_{n-\theta}(\ell)$ may contain elements of L_{j+1} other than ℓ ; for this reason we generally need $\theta > 0$ in order for (16) to hold.

We remark that in our applications, our models \mathcal{M}_n can be assumed to have expansions of arbitrarily high order, and so the particular value of $r_1(\Lambda_0, \Lambda_1, j, \epsilon)$ above is unimportant.

3. THE SHIFT OPERATOR AND THE PROOF OF LEMMA 2.5

In this section we develop some simple properties of the shift operator. Then we separate $\text{Trace}(M^k)$ into the contribution given by its real and non-real eigenvalues, and develop some fundamental facts about the $M \in \mathcal{M}_n$ -expected contribution due to the real eigenvalues. This allows us to prove Lemma 2.5, in a way that we will easily adapt in Section 4 to prove Lemma 2.6. The main idea in both proofs, just as in [Fri08, FK14], is to apply polynomials of the *shift in k* operator to both sides of (6). These polynomials of the shift operator will be used to annihilate the polyexponential parts of the coefficients of the asymptotic expansion; one simply needs to keep track of how these shift-operator polynomials will affect the expected value of $\text{Trace}(M^k)$ and the other terms in (6).

3.1. The Shift Operator.

Definition 3.1. By the *shift operator in k* , denoted $S = S_k$, we mean the operator on functions, $f = f(k)$, defined on non-negative integers, k , taking real or complex values, that takes f to Sf given by

$$(Sf)(k) = f(k+1).$$

For any integer $i \geq 0$ we let S^i denote the i -fold application of S , so that

$$(S^i f)(k) = f(k+i).$$

If $Q(z) = q_0 + q_1 z + \cdots + q_t z^t$ is any polynomial, with real or complex coefficients, we define

$$Q(S) = q_0 S^0 + q_1 S^1 + \cdots + q_t S^t.$$

If $f = f(k, n)$ depends on two variables, k and n , we apply S and S^i to $f(k, n)$ understanding that the shift is applied to the first variable, with the second variable, n , fixed.

Throughout this paper, $S = S_k$, will always be used to shift in k ; hence we never write the subscript k .

We remark that we often write formulas such as

$$(18) \quad S(\mu^k) = \mu \mu^k$$

with $k \in \mathbb{N}$ being a specific value; it would be more correct to write

$$(19) \quad (S(\mu^k)) \Big|_{k=k_0} = \mu \mu^{k_0},$$

for a specific $k_0 \in \mathbb{N}$, because the μ^k in $S(\mu^k)$ refers to a function of k . However, we generally use (18) for a particular value of k rather than the more tedious (19), unless confusion is likely.

There are a number of easy and useful results on the polynomials of shift operator, which we leave to the reader:

- (1) if $Q(z) = Q_1(z)Q_2(z)$ are polynomials, then $Q(S) = Q_1(S)Q_2(S) = Q_2(S)Q_1(S)$;
(2) if $Q(z) = q_0 + q_1z + \cdots + q_tz^t$, then
- $$(20) \quad (Q(S))(\mu^k) = Q(\mu)\mu^k$$
- (where μ^k refers to this function of k);
(3) if $p = p(k)$ is any polynomial, then
- $$(21) \quad (S - \mu)^D(p(k)\mu^k) = 0 \quad \text{if } D > \deg(p).$$

We will be interested in one particular type of polynomial.

Definition 3.2. Let $D \in \mathbb{Z}$ and $L \subset \mathbb{C}$ be a finite set. We define the *annihilator of L of degree D* to be the polynomial

$$\text{Ann}_{D,L}(z) \stackrel{\text{def}}{=} \prod_{\ell \in L} (z - \ell)^D.$$

We easily verify the following claims:

- (1) If $p(k)$ is any polyexponential function whose bases lie in a finite set $L \subset \mathbb{C}$, then $\text{Ann}_{D,L}(S)p = 0$ for $D \in \mathbb{N}$ sufficiently large (in view of (21)).
(2) If $f(k)$ is a function of growth ρ , then Sf is also of growth ρ , since for any fixed $\epsilon > 0$, for k sufficiently large we have
- $$(22) \quad |f(k+1)| < (\rho + \epsilon/2)^{k+1} \leq (\rho + \epsilon/2)^k(\rho + \epsilon/2) \leq (\rho + \epsilon)^k$$
- for k sufficiently large; hence for any fixed $i \in \mathbb{N}$, $S^i f$ is also of growth ρ ; hence for any fixed polynomial, Q , $Q(S)f$ is also of growth ρ .
(3) If $L \subset \mathbb{C}$ is a finite set that is closed under conjugation, and $\mu \in \mathbb{R}$, then $\text{Ann}_{1,L}(\mu)$ is real-valued, since

$$\overline{\text{Ann}_{1,L}(\mu)} = \prod_{\ell \in L} \overline{\mu - \ell} = \prod_{\ell \in L} (\mu - \bar{\ell}) = \prod_{\ell \in L} (\mu - \ell) = \text{Ann}_{1,L}(\mu);$$

if, in addition, $D \in \mathbb{N}$ is even then we have

$$\text{Ann}_{D,L}(\mu) = (\text{Ann}_{1,L}(\mu))^D \geq 0;$$

in particular, under these assumptions

$$(23) \quad \text{Ann}_{D,L}(\mu^k) = \text{Ann}_{D,L}(\mu)\mu^k \geq 0$$

when evaluated at any even integer k ; this positivity will be crucial to our proof of Lemma 2.5.

3.2. Real Eigenvalue Bounds. For each $M \in \mathcal{M}_n$, let

$$(24) \quad \text{RealTrace}(M, k), \quad \text{NonRealTrace}(M, k)$$

respectively denote the sum of the k -powers of the real and, respectively, non-real eigenvalues of M . Our proof of (12) is based on the following two conceptual steps which we state as lemmas; both lemmas will be proven in later subsections.

Lemma 3.3. *Let $\{\mathcal{M}_n\}_{n \in \mathbb{N}}$ be a (Λ_0, Λ_1) -bounded matrix model that has an order r expansion with range $K(n)$ ((6), Definition 2.3). Let $L \subset \mathbb{C}$ be any finite set including all the bases of $c_0(k), \dots, c_{r-1}(k)$ in an expansion (6). For any D such that $\text{Ann}_{D,L}$ annihilates the polyexponential parts of $c_i(k)$ with respect to Λ_0 (therefore any sufficiently large D), we have that for all $n \in \mathbb{N}$ and integers k with $0 \leq k \leq K(n) - D(\#L)$.*

$$(25) \quad \left| \text{Ann}_{D,L}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] \right) \right| \leq f_0(k)n + f_1(k)n^{-r},$$

where f_0, f_1 (depending on D, L, r) are functions of respective growths Λ_0, Λ_1 .

The proof of this lemma will be given in Subsection 3.4.

We remark that although [BS87, Fri91, Fri03, LP10] work with adjacency matrices, one could get similar estimates (for regular graphs) by working with Hashimoto matrices; if we work with Hashimoto matrices, then these older works would essentially use the above lemma in the (simple) special case where $D = 0$, where the c_0, \dots, c_{r-1} have vanishing polyexponential part. Hence one can view the above lemma as a generalization of a lemma used by older trace methods, where the older methods use the special case $D = 0$.

It is easy to see roughly why the above lemma is true: we write

$$\text{RealTrace}(M, k) + \text{NonRealTrace}(M, k) = \text{Trace}(M^k)$$

whose expected value has a power series expansion, namely the right-hand-side of (6). The non-real trace term is bounded by n (the maximum possible number of eigenvalues non-real eigenvalues) times Λ_0^n . We apply $\text{Ann}_{D,L}(S)$ where D is large enough to annihilate the polyexponential parts of the c_i ; this leaves (1) $\text{Ann}_{D,L}(S)$ applied to the $c_i(k)/n^i$, which is $1/n^i$ times a function of growth Λ_0 , plus (2) $\text{Ann}_{D,L}(S)$ applied to the expected non-real trace, which is bounded by n times another function of growth Λ_0 , plus (3) $\text{Ann}_{D,L}(S)$ applied to $c_r(k)/n^r$ which is $1/n^r$ times a function of growth Λ_1 . To prove the above lemma rigorously one has to verify that $\text{Ann}_{D,L}(S)$ acts in this way on the various terms.

Lemma 3.4. *Let $\{\mathcal{M}_n\}_{n \in \mathbb{N}}$ be a (Λ_0, Λ_1) -bounded matrix model. Let $L \subset \mathbb{C}$ be any finite set closed under conjugation, and let both D and k be positive even integers. Then for any $\theta, \epsilon > 0$, for all $n \in \mathbb{N}$ and $k \in \mathbb{Z}$,*

$$(26) \quad n^{-\theta D(\#L)} (\Lambda_0 + \epsilon)^k \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L)] \leq \text{Ann}_{D,L}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] \right)$$

(and both sides of the inequality are purely real).

The proof of this lemma will be given in Subsection 3.5. Note that this lemma does not require that $\{\mathcal{M}_n\}$ have any asymptotic expansion; its proof uses only the location of the eigenvalues of the matrices in \mathcal{M}_n .

Once we have proven the above two lemmas, we will combine them to conclude that

$$n^{-\theta D(\#L)} (\Lambda_0 + \epsilon)^k \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L)] \leq f_0(k)n + f_1(k)n^{-r},$$

with the notation and conditions of the above lemmas; then, with D, L fixed, we will judiciously choose a fixed r and variable $k = k(n)$ to prove the Exceptional Eigenvalue Bound (Lemma 2.5).

3.3. Intermediate Lemmas. In this subsection we state some lemmas that will be used both to prove Lemma 3.3 and a variant thereof needed to complete the proof of the Sidestepping Lemma. They are conceptually helpful and very easy to prove.

Lemma 3.5. *Let $c = c(k): \mathbb{N} \rightarrow \mathbb{C}$ be an approximate polyexponential function with error growth $\Lambda_0 \in \mathbb{R}$, whose larger bases (with respect to Λ_0) lie in a finite set $L \subset \mathbb{C}$. Then for sufficiently large D we have $\text{Ann}_{D,L}(S)c(k)$ is of growth Λ_0 .*

Proof. We have $c(k) = f(k) + g(k)$ where f is a polyexponential function whose bases lie in L , and g is of growth Λ_0 . If D is sufficiently large then $\text{Ann}_{D,L}(f) = 0$ (by the first claim after Definition 3.2). Therefore for such D we have

$$\text{Ann}_{D,L}(S)c(k) = \text{Ann}_{D,L}(S)g(k),$$

and the right-hand-side is a function of growth Λ_0 , according to the discussion around (22). \square

Notice that if $c(k) = 1$ for all k , then c is of growth 1; we easily check that $(S+1)^D c$ is the function $2^D c(k) = 2^D$, which is not uniformly bounded in D . So it is important to understand that in all the lemmas in this section, D, L are arbitrary but fixed, and the growth bounds are not uniform in D ; for similar reasons the bounds are not uniform in L : taking c as above and $L = \{-1, -2, \dots, -m\}$, we have $\text{Ann}_{1,L} c = (m+1)!$.

Lemma 3.6. *Let $r \geq 0$ be an integer, $K: \mathbb{N} \rightarrow \mathbb{N}$ a function, $N \subset \mathbb{Z}$ a subset, and $c = c(k): \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ a function of growth $\Lambda_1 \in \mathbb{R}$. Let $f = f(k, n)$ be any function $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{C}$ such that $|f(k, n)| \leq c(k)n^{-r}$ provided that $k, n \in \mathbb{N}$ satisfy $n \in N$ and $1 \leq k \leq K(n)$. Then for any finite set $L \subset \mathbb{C}$ and any integer $D \geq 0$, there is a function $g = g(k)$ of growth Λ_1 such that*

$$\left| (\text{Ann}_{D,L}(S)f)(k, n) \right| \leq g(k)n^{-r}$$

provided that $n \in N$ and $1 \leq k \leq K(n) - D(\#L)$.

Proof. We have

$$\text{Ann}_{D,L}(z) = a_0 + a_1 z + \dots + a_t z^t$$

for some $a_i \in \mathbb{C}$ and $t = D(\#L)$. For a fixed $n \in N$, the triangle inequality shows that

$$\left| (\text{Ann}_{D,L}(S)f)(k, n) \right| \leq (|a_0| |f(k, n)| + |a_1| |f(k+1, n)| + \dots + |a_t| |f(k+t, n)|);$$

provided that $1 \geq k$ and $k+t \leq K(n)$, the right-hand-side above is bounded above by

$$\leq (|a_0| |c(k)| + |a_1| |c(k)| + \dots + |a_t| |c(k+t)|) / n^{-r}.$$

But for each fixed $i \in \mathbb{N}$, $c(k+i)$ is also a function of growth Λ_1 (by the discussion around (22)). \square

Lemma 3.7. *Let $\{\mathcal{M}_n\}_{n \in \mathbb{N}}$ be a (Λ_0, Λ_1) -bounded matrix model with range $K(n)$. Then for any finite set $L \subset \mathbb{C}$ and integer $D \geq 0$ we have that for all $n \in \mathbb{Z}$ and k with $1 \leq k \leq K(n) - D(\#L)$,*

$$(27) \quad \left| \text{Ann}_{D,L}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{NonRealTrace}(M, k)] \right) \right| \leq f_0(k)n,$$

where f_0 is a function of growth Λ_0 .

Proof. For each $M \in \mathcal{M}_n$ we have

$$\text{NonRealTrace}(M, k) = \sum_{\mu \in \text{NonReal}(M)} \mu^k$$

where $\text{NonReal}(M)$ are the eigenvalues of M that are not real. Hence

$$\text{Ann}_{D,L}(S)\text{NonRealTrace}(M, k) = \sum_{\mu \in \text{NonReal}(M)} \text{Ann}_{D,L}(\mu)\mu^k,$$

and hence

$$(28) \quad \left| \text{Ann}_{D,L}(S)\text{NonRealTrace}(M, k) \right| \leq Mn\Lambda_0^k,$$

where

$$M = \max_{|z| \leq \Lambda_0} |\text{Ann}_{D,L}(z)|.$$

But the triangle inequality implies that for any random variable $g: \mathcal{M}_n \rightarrow \mathbb{C}$ we have

$$|\mathbb{E}_{M \in \mathcal{M}_n} g(M)| \leq \mathbb{E}_{M \in \mathcal{M}_n} |g(M)|;$$

taking $g = \text{Ann}_{D,L}(S)\text{NonRealTrace}(M, k)$ and exchanging the order of $\text{Ann}_{D,L}(S)$ and $\mathbb{E}_{M \in \mathcal{M}_n}$ (which clearly commute), (28) implies that

$$\left| \text{Ann}_{D,L}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{NonRealTrace}(M, k)] \right) \right| \leq Mn\Lambda_0^k.$$

□

3.4. The Proof of Lemma 3.3.

Proof of Lemma 3.3. Consider (6) for any fixed r . For any $0 \leq i \leq r-1$ we have

$$\text{Ann}_{D,L}(S)(c_i(k)/n^i) = \text{Ann}_{D,L}(S)(c_i(k))/n^i,$$

so Lemmas 3.5 implies that for any fixed, sufficiently large $D \in \mathbb{N}$ there is a function $f_{0,i}(k)$ of growth Λ_0 for which

$$\left| \text{Ann}_{D,L}(S)(c_i(k)/n^i) \right| = \left| \text{Ann}_{D,L}(S)(c_i(k)) \right| / n^i \leq f_{0,i}(k) / n^i \leq f_{0,i}(k)$$

since $i \geq 0$. Adding the above equation applied to $i = 0, \dots, r-1$ and applying the triangle inequality shows that

$$\left| \text{Ann}_{D,L}(S)(c_0(k) + \dots + c_{r-1}(k)/n^{r-1}) \right| \leq F_0(k), \quad \text{where } F_0(k) = f_{0,0}(k) + \dots + f_{0,r-1}(k),$$

is a function of growth Λ_0 . The term $O(c_r(k))n^{-r}$ in (6) refers to a function of k, n that is bounded by n^{-r} times a function of growth Λ_1 ; Lemma 3.6 implies the same bound when $\text{Ann}_{D,L}(S)$ is applied; hence, if $\text{RHS} = \text{RHS}(k, n)$ denotes the right-hand-side of (6),

$$(29) \quad \left| \text{Ann}_{D,L}(S)\text{RHS}(k, n) \right| \leq F_0(k) + n^{-r} F_1(k),$$

where for $i = 0, 1$, F_i is of growth Λ_i , and provided that $1 \leq k \leq K(n) - D(\#L)$ (since applying $\text{Ann}_{D,L}(S)$ requires (6) to hold for $k, k+1, \dots, k+D(\#L)$). Set

$$(30) \quad g_1(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)], \quad g_2(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{M \in \mathcal{M}_n} [\text{NonRealTrace}(M, k)].$$

Lemma 3.7 implies that

$$\left| \text{Ann}_{D,L}(S)g_2(k, n) \right| \leq n\tilde{f}(k)$$

where $\tilde{f}(k)$ is a function of growth Λ_0 . Since $g_1(k, n) + g_2(k, n)$ equals the left-hand-side of (6), we have

$$(31) \quad \text{Ann}_{D,L}(S)g_1(k, n) = -\text{Ann}_{D,L}(S)g_2(k, n) + \text{Ann}_{D,L}(S)(\text{RHS}(k, n)).$$

Using (29) and (27) we establish (25) with $f_0(k) = \tilde{f}(k) + F_0(k)$ and $f_1(k) = F_1(k)$. \square

3.5. Proof of Lemma 3.4.

Proof of Lemma 3.4. Since D is even and L is closed under complex conjugation, then $\text{Ann}_{D,L}$ maps real values to non-negative values (see (23) and the discussion above it), and hence

$$\text{Ann}_{D,L}(\mu) \geq 0,$$

and therefore

$$\text{Ann}_{D,L}(S)\mu^k = \text{Ann}_{D,L}(\mu)\mu^k \geq 0$$

if evaluated on an even $k \in \mathbb{N}$ and $\mu \in \mathbb{R}$. It follows that if k is even, n is fixed, and $X_n \subset \mathbb{R}$ is a measurable set, then

$$(32) \quad \left(\inf_{x \in X_n} (\text{Ann}_{D,L}(x)x^k) \right) \mathbb{E}_{M \in \mathcal{M}_n} [\#(\text{Spec}(M) \cap X_n)] \leq \text{Ann}_{D,L}(S) \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)]$$

(and both the above numbers are non-negative reals).

Now take

$$X_n = [-\Lambda_1, \Lambda_1] \setminus (B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L))$$

and consider any $x \in X_n$. Then $|x - \ell| \geq n^{-\theta}$ for each $\ell \in L$, and hence

$$\text{Ann}_{D,L}(x) \geq n^{-\theta D(\#L)};$$

in addition $|x| \geq \Lambda_0 + \epsilon$ since $x \in X_n$, and therefore

$$(\text{Ann}_{D,L}(x)x^k) \geq n^{-\theta D(\#L)}(\Lambda_0 + \epsilon)^k.$$

Taking the minimum over all $x \in X_n$ and combining this with (32) yields (26). \square

3.6. Proof of Lemma 2.5.

Proof of Lemma 2.5. By definition L_r , the set of larger bases (with respect to Λ_0) of the polyexponential parts of $c_0(k), \dots, c_{r-1}(k)$ in an expansion (6), consists entirely of real numbers; hence L_r is closed under conjugation. In view of Lemmas 3.3 and 3.4 we have that for sufficiently large even $D \in \mathbb{N}$, for any $\theta > 0$ and all positive, even $k \leq K(n) - D(\#L)$,

$$(33) \quad n^{-\theta D(\#L)}(\Lambda_0 + \epsilon)^k \mathbb{E}\text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] \leq f_0(k)n + f_1(k)n^{-r},$$

where f_i are of growth Λ_i , for a function $K(n) \gg \log n$. Since the f_i are of growth Λ_i , for every $\delta > 0$ we have

$$n^{-\theta D(\#L)}(\Lambda_0 + \epsilon)^k \mathbb{E}\text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] \leq (\Lambda_0 + \delta)^k n + (\Lambda_1 + \delta)^k n^{-r}$$

for n large; hence

$$\mathbb{E}\text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] \leq n^{\theta D(\#L)}(\Lambda_0 + \epsilon)^{-k} ((\Lambda_0 + \delta)^k n + (\Lambda_1 + \delta)^k n^{-r}).$$

Hence, if for some even k we can show that

$$(34) \quad n^{\theta D(\#L)}(\Lambda_0 + \epsilon)^{-k} \max((\Lambda_0 + \delta)^k n, (\Lambda_1 + \delta)^k n^{-r}) \leq n^{-\alpha - \delta'}$$

for some $\delta' > 0$, then it would follow that

$$\mathbb{E}\text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] \leq n^{-\alpha}$$

for large n , which is just (12).

Fix a $\kappa \in \mathbb{R}$ —to be specified later—and let k be the nearest even integer to $\kappa \log n$ (which is typical in trace methods for regular graphs), where \log is with respect to any fixed base (common to all logarithms used). Taking logarithms in (34) and dividing by $\log n$, it suffices to choose $\delta, \kappa, \theta, r$ so that

$$\theta D(\#L) - \kappa \log(\Lambda_0 + \epsilon) + \max(1 + \kappa \log(\Lambda_0 + \delta), -r + \kappa \log(\Lambda_1 + \delta)) < -\alpha - \delta'.$$

By continuity of the above expression in δ', δ, θ , for a fixed κ, r it would be possible to find $\delta', \delta, \theta > 0$ satisfying the above provided that the above inequality holds (strictly) for $\delta' = \delta = \theta = 0$, i.e., that

$$(35) \quad -\kappa \log(\Lambda_0 + \epsilon) + 1 + \kappa \log \Lambda_0 < -\alpha,$$

$$(36) \quad -\kappa \log(\Lambda_0 + \epsilon) - r + \kappa \log \Lambda_1 < -\alpha.$$

To choose such κ, r , it suffices to first choose κ to satisfy (35), which is possible since $\log(\Lambda_0 + \epsilon) > \log \Lambda_0$ since $\epsilon > 0$; then we choose r to satisfy (36). It follows that there exist κ, r and $\delta', \delta, \theta > 0$ that depend only on $\Lambda_0, \Lambda_1, \epsilon, \alpha$ that satisfy (34), and therefore (12) holds for n sufficiently large.

If we set r_0, θ_0 to be those particular values of r, θ (which depend only on $\Lambda_0, \Lambda_1, \epsilon, \alpha$), then for any other $r \geq r_0$ and $\theta \leq \theta_0$ we also have (12), since in this case we have

$$B_{n-\theta_0}(L_{r_0}) \subset B_{n-\theta}(L_r)$$

for all $n \in \mathbb{N}$. □

We remark that (35) holds iff

$$(37) \quad \kappa > (\alpha + 1) / \log \left(\frac{\Lambda_0 + \epsilon}{\Lambda_0} \right),$$

$$(38) \quad r > \alpha + \kappa \log \left(\frac{\Lambda_1}{\Lambda_0 + \epsilon} \right).$$

Hence if Λ_0, Λ_1 are fixed, then κ must be at least proportional to α/ϵ as $\alpha \rightarrow \infty$ and $\epsilon \rightarrow 0$, and hence the same is true of r ; hence we need arbitrarily high order expansions for the model \mathcal{M}_n as $\alpha \rightarrow \infty$ and $\epsilon \rightarrow 0$.

4. PROOF OF THE SIDESTEPPING LEMMA (LEMMA 2.6)

In this section we give a variant of Lemma 3.3 that allows us to isolate the \mathcal{M}_n expected value of eigenvalues near points outside of $B_{\Lambda_0+\epsilon}$; we then combine this result with that of (12) (in the Exceptional Eigenvalue Bound, i.e., Lemma 2.5) to prove Lemma 2.6.

4.1. A Variant of Lemma 3.3.

Lemma 4.1. *Let $\{\mathcal{M}_n\}_{n \in \mathbb{N}}$ be a (Λ_0, Λ_1) -bounded matrix model that has an order r expansion (6) with range $K(n)$ (as in Definition 2.3). Let $L \subset \mathbb{C}$ be any finite set including all the bases of $c_0(k), \dots, c_{r-1}(k)$ in an expansion (6). For some $\ell \in L$, let $\tilde{L} = L \setminus \{\ell\}$, and for $i = 0, \dots, r-1$ let $p_i = p_i(k)$ be the polynomial such that $p_i(k)\ell^k$ is the ℓ -part of $c_i(k)$. Then for sufficiently large $\tilde{D} \in \mathbb{N}$, for all k with*

$1 \leq k \leq K(n) - \tilde{D}(\#\tilde{L})$ we have
(39)

$$\left| \text{Ann}_{\tilde{D}, \tilde{L}}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] - \ell^k \sum_{i=0}^{r-1} p_i(k)/n_i \right) \right| \leq f_0(k)n + f_1(k)n^{-r},$$

where f_0, f_1 (depending on \tilde{D}, \tilde{L}, r and the c_i , but not on n, k) are functions of respective growths Λ_0, Λ_1 .

Proof. Setting

$$(40) \quad \tilde{g}_1(k, n) \stackrel{\text{def}}{=} \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] - \sum_{i=0}^{r-1} p_i(k)\ell^k/n^{-i}$$

and $g_2(k, n)$ as in (30), we have

$$(41) \quad \tilde{g}_1(k, n) + g_2(k, n) = \sum_{i=0}^{r-1} \frac{\tilde{c}_i(k)}{n^i} + \frac{O(c_r(k))}{n^r},$$

where $\tilde{c}_i(k) = c_i(k) - p_i(k)\ell^k$; therefore all the larger bases (with respect to Λ_0) of the $\tilde{c}_i(k)$ lie in \tilde{L} . Now to both sides of (41) we apply $\text{Ann}_{\tilde{D}, \tilde{L}}$, subtract $g_2(k, n)$, take absolute values, and apply the triangle inequality to get

$$|\text{Ann}_{\tilde{D}, \tilde{L}} \tilde{g}_1(k, n)| \leq |\text{Ann}_{\tilde{D}, \tilde{L}} g_2(k, n)| + |\text{Ann}_{\tilde{D}, \tilde{L}} c_r(k)/n^r| + \sum_{i=0}^{r-1} |\text{Ann}_{\tilde{D}, \tilde{L}} \tilde{c}_i(k)/n^i|$$

Applying Lemmas 3.5–3.7 (as they were in the proof of Lemma 3.3), we conclude that

$$|\text{Ann}_{\tilde{D}, \tilde{L}} g_2(k, n)| \leq n f_0(k), \quad |\text{Ann}_{\tilde{D}, \tilde{L}} c_r(k)/n^r| \leq n^{-r} f_1(k),$$

where $f_i(k)$ are functions of growth Λ_i ($i = 1, 2$), and

$$|\text{Ann}_{\tilde{D}, \tilde{L}} \tilde{c}_i(k)/n^i| \leq f_{0,i}(k)/n^i$$

where $f_{0,i}(k)$ is a function of growth Λ_0 for $i = 0, \dots, r-1$. Letting \tilde{f}_0 be f_0 plus the sum of the $f_{0,i}$ we have

$$|\text{Ann}_{\tilde{D}, \tilde{L}} \tilde{g}_1(k, n)| \leq \tilde{f}_0(k)n + f_1(k)n^{-r},$$

where \tilde{f}_0, f_1 are of respective growths Λ_0, Λ_1 . This bound and (40) imply (39). \square

4.2. A Variant of Lemma 3.4. Next we prove a variant of Lemma 3.4 that allows us to relate the term

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S) \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)]$$

(which implicitly forms part of the left-hand-side of (39)) to the quantity $\mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n-\theta}(\ell)]$, with notation as in Lemma 3.3. Let us state this result formally, in a way that will be useful to us.

Lemma 4.2. *Let $r \in \mathbb{N}$, and let $\{\mathcal{M}_n\}_{n \in \mathbb{N}}$ be a (Λ_0, Λ_1) -bounded matrix model that has an order r expansion of range $K(n)$ ((6), Definition 2.3). Let $\tilde{D} \geq 0$ be an integer and $\epsilon, \theta > 0$ real numbers; let $\ell \in \mathbb{R}$ with $|\ell| > \Lambda_0$ and $\tilde{L} \subset \mathbb{C} \setminus \{\ell\}$ a finite*

set such that $L = \ell \cup \tilde{L}$ lies in $B_{\Lambda_1}(0)$. Then there is a constant, C , independent of k and n , such that for large $n \in N$ and all $1 \leq k \leq K(n) - \tilde{D}(\#L)$

(42)

$$\left| \left(1 + O(n^{-\theta})\right) \text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \ell^k \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] - \text{Ann}_{\tilde{D}, \tilde{L}}(S) \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] \right|$$

(43)

$$\leq C(\Lambda_0 + \epsilon)^k n + C\Lambda_1^k \left(n^{1-\theta\tilde{D}} + \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L)] \right)$$

provided that $K(n) \ll n^\theta$ (i.e., $K(n)/n^\theta \rightarrow 0$ as $n \rightarrow \infty$).

Note that since we only need $K(n) \gg \log n$ in Definition 2.3 (and in our main lemmas and Theorem 2.4), we can always replace $K(n)$ by a smaller function (e.g., $\max(K(n), (\log n)^2)$) that is $\gg \log n$ and $\ll n^\theta$ (for all $\theta > 0$); hence the assumption that $K(n) \ll n^\theta$ is harmless to our main results. We remark that under further assumptions on \tilde{L} and θ , such as those in Lemma 3.3, we can make the $\mathbb{E} \text{out}$ term in (43) smaller than $n^{-\alpha}$ for any α ; however, the above lemma has no such assumptions.

Proof. For each square matrix M , each $\mathcal{R} \subset \mathbb{C}$, and $k \in \mathbb{N}$, let

$$(44) \quad \text{RealTrace}(M, k; \mathcal{R}) \stackrel{\text{def}}{=} \sum_{\lambda \in \text{Spec}(M) \cap \mathcal{R} \cap \mathbb{R}} \lambda^k.$$

For each n let

$$\begin{aligned} S_0 &= S_0(n) = B_{n^{-\theta}}(\ell), \\ S_1 &= S_1(n) = B_{n^{-\theta}}(\tilde{L}), \\ S_2 &= S_2(n) = B_{\Lambda_0+\epsilon}(0), \\ S_3 &= S_3(n) = \mathbb{R} \setminus (B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L)), \end{aligned}$$

and

$$g_0(k, n) = \text{RealTrace}(M, k; S_0(n)) = \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)].$$

We may write

$$\begin{aligned} \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] - \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k; S_0(n))] \\ = \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k; \mathbb{R} \setminus S_0(n))]. \end{aligned}$$

Noting that \mathbb{R} is contained in the union of the S_i , applying $\text{Ann}_{\tilde{D}, \tilde{L}}$ to both sides we have

$$(45) \quad \left| \text{Ann}_{\tilde{D}, \tilde{L}}(S) \left(g_0(k, n) - \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] \right) \right| \leq \sum_{i=1}^3 E_i M_i,$$

where for $i = 1, 2, 3$

$$E_i = \mathbb{E} \text{in}_{\mathcal{M}_n} [S_i(n)], \quad M_i = \max_{x \in S_i(n) \cap B_{\Lambda_1}(0)} |\text{Ann}_{\tilde{D}, \tilde{L}}(S)x^k|.$$

Let us estimate $E_i M_i$ for $i = 1, 2, 3$.

Since \tilde{D}, \tilde{L} are fixed, we have

$$M_2 \leq C \left(\max_{x \in S_2(n)} |x| \right)^k = C(\Lambda_0 + \epsilon)^k,$$

and so the trivial estimate $E_2 \leq n$ implies that

$$(46) \quad E_2 M_2 \leq Cn(\Lambda_0 + \epsilon)^k.$$

If $x \in B_{n^{-\theta}(\tilde{\ell})}$ for some $\tilde{\ell} \in \tilde{L}$, then

$$|(x - \tilde{\ell})^{\tilde{D}}| \leq n^{-\theta\tilde{D}},$$

and hence

$$|\text{Ann}_{\tilde{D}, \tilde{L}}(x)| = |x - \tilde{\ell}|^{\tilde{D}} \prod_{\ell' \in \tilde{L} \setminus \{\tilde{\ell}\}} |x - \ell'|^{\tilde{D}} \leq n^{-\theta\tilde{D}}(\Lambda_1 + |\ell'|)^{\tilde{D}}.$$

In view of (20) we have

$$|\text{Ann}_{\tilde{D}, \tilde{L}}(S)x^k| = |\text{Ann}_{\tilde{D}, \tilde{L}}(x)||x|^k \leq n^{-\theta\tilde{D}}C\Lambda_1^k.$$

Combining this with the trivial $E_1 \leq n$ yields

$$(47) \quad E_1 M_1 \leq Cn^{1-\theta\tilde{D}}\Lambda_1^k.$$

Finally

$$(48) \quad E_3 = \mathbb{E}\text{out}_{\mathcal{M}_n}[B_{\Lambda_0+\epsilon}(0) \cup B_{n^{-\theta}}(L)]$$

and

$$(49) \quad M_3 \leq \max_{|x| \leq \Lambda_1} (|\text{Ann}_{\tilde{D}, \tilde{L}}(x)||x|^k) \leq C\Lambda_1^k.$$

Combining (46), (47), (48), (49), with (45) we have

$$(50) \quad \left| \text{Ann}_{\tilde{D}, \tilde{L}}(S) \left(g_0(k, n) - \mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] \right) \right|$$

is bounded by (43).

Since (50) is bounded above by (43), and since (50) clearly equals

$$\left| \text{Ann}_{\tilde{D}, \tilde{L}}(S)g_0(k, n) - \text{Ann}_{\tilde{D}, \tilde{L}}(S)\mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] \right|,$$

to finish the proof of Lemma 4.2 it suffices to show that

$$(51) \quad \text{Ann}_{\tilde{D}, \tilde{L}}(S)g_0(k, n) = \left(1 + O(n^{-\theta})\right) \text{Ann}_{\tilde{D}, \tilde{L}}(\ell)\ell^k \mathbb{E}\text{in}_{\mathcal{M}_n}[B_{n^{-\theta}}(\ell)]$$

for fixed $\tilde{D}, \tilde{L}, \ell$, and large n, k with $k \ll n^\theta$. Let us prove this.

Since $\ell \in \mathbb{R}$, the set $B_{n^{-\theta}}(\ell) \cap \mathbb{R}$ is just the closed interval $I_n = [\ell - n^{-\theta}, \ell + n^{-\theta}]$; so setting

$$h(x) \stackrel{\text{def}}{=} \text{Ann}_{\tilde{D}, \tilde{L}}(x)x^k,$$

we have that for any $\lambda \in I_n$ there is a $\xi \in I_n$ for which

$$h(\lambda) - h(\ell) = (\lambda - \ell)h'(\xi).$$

The product rule shows that

$$|h'(x)| \leq C|\xi|^k \leq C(|\ell| + n^{-\theta})^k \leq C'|\ell|^k$$

provided that $k \ll n^\theta$. Hence

$$(52) \quad |h(\lambda) - h(\ell)| \leq |\lambda - \ell|C'|\ell|^k \leq C'n^{-\theta}|\ell|^k.$$

Since $\ell \notin \tilde{L}$, the quantity

$$A = \text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \neq 0$$

(and is, of course, independent of k, n); since $h(\ell) = A\ell^k$ (by (20)), (52) implies

$$h(\lambda) = h(\ell) + O(n^{-\theta})|\ell^k| = A\ell^k + O(n^{-\theta})|\ell^k| = A\ell^k \left(1 + O(n^{-\theta})\right),$$

which holds for all $\lambda \in I_n = B_{n^{-\theta}}(\ell) \cap \mathbb{R}$. It follows that for any $M \in \mathcal{M}_n$ we have

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S) \text{RealTrace} \left(M, k; B_{n^{-\theta}}(\ell) \right) = (\# \text{Spec}(M) \cap B_{n^{-\theta}}(\ell)) A\ell^k \left(1 + O(n^{-\theta})\right).$$

Taking expected values yields

$$\begin{aligned} \text{Ann}_{\tilde{D}, \tilde{L}}(S) g_0(k, n) &= \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] A\ell^k \left(1 + O(n^{-\theta})\right) \\ &= \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] \text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \ell^k \left(1 + O(n^{-\theta})\right) \end{aligned}$$

which proves (51); by the discussion in the paragraph containing (51), this completes the proof. \square

4.3. A Shift Computation. Here is a simple lemma regarding the shift operation which we will need regarding the left-hand-side of (39).

Lemma 4.3. *Let $p = p(k)$ be a non-zero polynomial with complex coefficients, let $\ell \in \mathbb{C}$, and let $\tilde{L} \subset \mathbb{C} \setminus \{\ell\}$ be a finite set. Then for each $\tilde{D} \in \mathbb{N}$ there is a polynomial $\tilde{p}(k)$ of the same degree as $p(k)$ such that*

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S) (p(k)\ell^k) = \tilde{p}(k)\ell^k;$$

more precisely, if $p(k) = c_0 + c_1k + \dots + c_tk^t$ with $c_t \neq 0$, then we have

$$\tilde{p}(k) = \tilde{c}_0 + \tilde{c}_1k + \dots + \tilde{c}_tk^t$$

where $\tilde{c}_t \neq 0$ and is given by

$$\tilde{c}_t = c_t \text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \neq 0.$$

Proof. For any $\tilde{\ell} \neq \ell$, we see that

$$(S - \tilde{\ell})(k^t \ell^k) = (k+1)^t \ell^{k+1} - k^t \tilde{\ell} \ell^k = \ell^k (k^t (\ell - \tilde{\ell}) + \dots)$$

where the \dots is a polynomial in k of degree less than t . It follows that

$$(S - \tilde{\ell})(p(k)\ell^k) = \ell^k (c_t k^t (\ell - \tilde{\ell}) + \dots)$$

where the \dots is a polynomial in k of degree less than t ; this proves the claim for $\#L = 1$ (i.e., $\tilde{L} = \{\tilde{\ell}\}$) and $\tilde{D} = 1$. Repeatedly applying $S - \tilde{\ell}$ as above proves the case $\tilde{L} = \{\tilde{\ell}\}$ and all $\tilde{D} \in \mathbb{N}$. For general \tilde{D} and \tilde{L} , we write $\tilde{L} = \{\ell_1, \dots, \ell_m\}$, and write

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S) (p(k)\ell^k) = (S - \ell_1)^{\tilde{D}} \dots (S - \ell_m)^{\tilde{D}} (p(k)\ell^k).$$

We then first apply $(S - \ell_m)^{\tilde{D}}$ to $p(k)\ell^k$, then apply $(S - \ell_{m-1})^{\tilde{D}}$ to the result, etc., and the lemma follows (more formally, the lemma follows by induction on $m = \#\tilde{L}$). \square

4.4. Proof of Lemma 2.6.

Proof of Lemma 2.6. Let $\tilde{\epsilon} = \epsilon/3$, and let κ_0 be given by

$$\kappa_0 = \frac{j+2}{\log\left(\frac{\Lambda_0 + 2\tilde{\epsilon}}{\Lambda_0 + \tilde{\epsilon}}\right)}$$

where the logarithm, and all logarithms below, are taken relative to a fixed base [this choice of base does not affect r_1, θ_1 below and merely scales κ_0]. Clearly κ_0 is positive and can be equivalently be described by the equation

$$(53) \quad \kappa_0 \log(\Lambda_0 + 2\tilde{\epsilon}) - j - 2 = \kappa_0 \log(\Lambda_0 + \tilde{\epsilon}).$$

Let κ be any real number with

$$\kappa \geq \kappa_0;$$

our proof will require us to do the present computation twice, with two different values of κ ; this is why we don't choose one specific value of κ . Then let

$$(54) \quad \tilde{\alpha} = j + 1 + \kappa \log(\Lambda_1) - \kappa \log(\Lambda_0 + 2\tilde{\epsilon})$$

$$(55) \quad \tilde{r} = j + 1 + \lceil \kappa \log(\Lambda_1 + \tilde{\epsilon}) - \kappa \log(\Lambda_0 + 2\tilde{\epsilon}) \rceil$$

where $\lceil \cdot \rceil$ denotes the ceiling function (the least upper bound that is an integer). Since $\Lambda_1 \geq 3\tilde{\epsilon} + \Lambda_0$, we have $\tilde{\alpha}, \tilde{r} > 0$.

We then set

$$(56) \quad r_1 = \max(\tilde{r}, r_0(\Lambda_0, \Lambda_1, \tilde{\epsilon}, \tilde{\alpha}))$$

$$(57) \quad \theta_1 = \theta_0(\Lambda_0, \Lambda_1, \tilde{\epsilon}, \tilde{\alpha}),$$

with r_0, θ_0 being the functions in Lemma 2.5. For now we have that r_1, θ_1 depend on κ as well as $\Lambda_0, \Lambda_1, j, \epsilon$ (since $\tilde{r}, \tilde{\alpha}$ depend on κ); we will ultimately take $\kappa = \kappa_0$ to produce the r_1, θ_1 in Lemma 2.6.

Next let $\theta \in \mathbb{R}$ and $r \in \mathbb{N}$ satisfy

$$0 < \theta \leq \theta_1(\Lambda_0, \Lambda_1, j, \epsilon, \kappa), \quad r \geq r_1(\Lambda_0, \Lambda_1, j, \epsilon, \kappa).$$

Let us prove (14)–(17).

Since

$$r \geq r_1 \geq r_0(\Lambda_0, \Lambda_1, \tilde{\epsilon}, \tilde{\alpha}), \quad \theta \leq \theta_1 \leq \theta_0(\Lambda_1, \Lambda_0, \tilde{\epsilon}, \tilde{\alpha}),$$

Lemma 2.5 implies that

$$(58) \quad \mathbb{E}\text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \tilde{\epsilon}}(0) \cup B_{n^{-\theta_1}}(L_{r_1})] \leq n^{-\tilde{\alpha}} \leq n^{-j-1}$$

for n sufficiently large; since $\theta \leq \theta_1$, $r \geq r_1$, and $\tilde{\epsilon} < \epsilon$, we have

$$B_{\Lambda_0 + \tilde{\epsilon}}(0) \cup B_{n^{-\theta_1}}(L_{r_1}) \subset B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)$$

and hence

$$(59) \quad \mathbb{E}\text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] \leq n^{-j-1}$$

for n sufficiently large.

Note that $r_1 \geq \tilde{r}$ by (56), and $\tilde{r} \geq j + 1$ by (55), and hence

$$(60) \quad r \geq r_1 \geq j + 1, \quad \text{and} \quad L_{j+1} \subset L_r.$$

Now let $\ell \in L_r$ with $|\ell| \geq \Lambda_0 + 2\tilde{\epsilon}$, and set

$$\tilde{L} = L_r \setminus \{\ell\}.$$

Let $\tilde{D} \in \mathbb{N}$ be sufficiently large so that Lemma 4.1 holds and that

$$(61) \quad \kappa \log(\Lambda_0 + 2\tilde{\epsilon}) - j - 1 \geq \kappa \log(\Lambda_1) + 1 - \theta\tilde{D}.$$

Lemma 4.1 implies that for some function $K(n)$ with $K(n) \gg \log n$ we have that for all $k \in \mathbb{N}$ with $1 \leq k \leq K(n) - \tilde{D}(\#\tilde{L})$ we have

$$(62) \quad \left| \text{Ann}_{\tilde{D}, \tilde{L}}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] - \sum_{i=0}^{r-1} \ell^k q_{\ell, i}(k) n^{-i} \right) \right| \leq f_0(k)n + f_1(k)n^{-r},$$

where f_0, f_1 are functions of respective growths Λ_0, Λ_1 .

Now for n such that \mathcal{M}_n is defined, let $k(n)$ be any function with $k(n) - \kappa \log n = O(1)$. Let us prove that for $k = k(n)$, as $n \rightarrow \infty$ (over n where \mathcal{M}_n is defined) we have

$$(63) \quad |f_0(k(n))n| = o(|\ell|^{k(n)} n^{-j}) \quad \text{and}$$

$$(64) \quad |f_1(k(n))n^{-r}| = o(|\ell|^{k(n)} n^{-j}).$$

Since f_0 is of growth Λ_0 , we have

$$(65) \quad f_0(k(n))n = O(1)(\Lambda_0 + \tilde{\epsilon})^{k(n)} n \leq O(1)(\Lambda_0 + \tilde{\epsilon})^{\kappa \log n + O(1)} n.$$

Since $\kappa \geq \kappa_0$, we have that (53) also holds with κ replacing κ_0 , and hence multiplying the equation by $\log n$ we have

$$(66) \quad (\Lambda_0 + \tilde{\epsilon})^{\kappa \log n} n \leq (\Lambda_0 + 2\tilde{\epsilon})^{\kappa \log n} n^{-j-2} \leq |\ell|^{\kappa \log n} n^{-j-2} \leq |\ell|^{k(n) + O(1)} n^{-j-2};$$

combining this with (65) yields

$$f_0(k(n))n = |\ell|^{k(n)} n^{-j} O(1/n)$$

which establishes (63). The proof of (64) is similar: in view of (55) and $r \geq \tilde{r}$ we have

$$\begin{aligned} n^{-r} &\leq n^{-\tilde{r}} \leq n^{-j-1} n^{-\kappa \log(\Lambda_1 + \tilde{\epsilon})} n^{\kappa \log(\Lambda_0 + 2\tilde{\epsilon})} \\ &= n^{-j-1} O(1)(\Lambda_1 + \tilde{\epsilon})^{-k(n)} (\Lambda_0 + 2\tilde{\epsilon})^{k(n)} \end{aligned}$$

where the last equality holds since $k(n) = \kappa \log n + O(1)$; since $f_1(k) \leq (\Lambda_1 + \tilde{\epsilon})^{k(n)}$ for large n , we have

$$f_1(k(n))n^{-r} = O(1/n) n^{-j} n^{\kappa \log(\Lambda_0 + 2\tilde{\epsilon})} = |\ell|^{k(n)} n^{-j} O(1/n)$$

which proves (64).

Combining (62)–(64) we have

$$(67) \quad \left| \text{Ann}_{\tilde{D}, \tilde{L}}(S) \left(\mathbb{E}_{M \in \mathcal{M}_n} [\text{RealTrace}(M, k)] - \sum_{i=0}^{r-1} \ell^k q_{\ell, i}(k) n^{-i} \right) \right| \leq o(|\ell|^{k(n)} n^{-j})$$

provided that $1 \leq k(n) \leq K(n) - \tilde{D}(\#\tilde{L})$. Since $k(n) = \kappa \log n + O(1)$, we have that $1 \leq k(n) \leq K(n) - \tilde{D}(\#\tilde{L})$ for sufficiently large n . With this in mind, we now finish the proof of Lemma 2.6 by estimating the left-hand-side of (67) up to terms of size $o(|\ell|^{k(n)} n^{-j})$.

Applying Lemma 4.3 we have

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S) \ell^{k(n)} \sum_{i=0}^{r-1} q_{\ell, i}(k(n)) / n^i = \ell^{k(n)} \sum_{i=0}^{r-1} Q_i(k(n)) / n^i$$

where $Q_i = Q_i(k)$ is the (unique) polynomial such that

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S)(\ell^k q_{\ell, i}(k)) = \ell^k Q_i(k);$$

hence $Q_i(k)$ is of the same degree as $q_{\ell, i}(k)$ and the leading coefficient of $Q_i(k)$ is $\text{Ann}_{\tilde{D}, \tilde{L}}(\ell)$ times that of $q_{\ell, i}(k)$. Since $L_j = \emptyset$ we have $q_{\ell, i}(k) = 0$ for all $i < j$; it follows that

$$\text{Ann}_{\tilde{D}, \tilde{L}}(S)\ell^{k(n)} \sum_{i=0}^{r-1} q_{\ell, i}(k(n))/n^i = \ell^{k(n)} Q_j(k(n))n^{-j} + |\ell|^{k(n)} n^{-j} O\left(\left(k(n)\right)^m/n\right)$$

where m is the degree of $q_{\ell, j+1}(k)$. Since $k(n) = O(\log n)$ we have

$$(68) \quad \text{Ann}_{\tilde{D}, \tilde{L}}(S)\ell^{k(n)} \sum_{i=0}^{r-1} q_{\ell, i}(k(n))/n^i = \ell^{k(n)} Q_j(k(n))n^{-j} + |\ell|^{k(n)} n^{-j} o(1)$$

Next consider

$$(69) \quad \text{Ann}_{\tilde{D}, \tilde{L}}(S)\mathbb{E}_{M \in \mathcal{M}_n}[\text{RealTrace}(M, k(n))].$$

Applying Lemma 4.2 with $\tilde{\epsilon}$ replacing ϵ , we have (69) equals

$$(70) \quad \left(1 + O(n^{-\theta})\right) \text{Ann}_{\tilde{D}, \tilde{L}}(\ell)\ell^{k(n)} \mathbb{E}_{\text{in}_{\mathcal{M}_n}[B_{n^{-\theta}}(\ell)]} + O(1)\text{Term}(n)$$

where

$$\text{Term}(n) = \text{Term}_1(n) + \text{Term}_2(n)$$

where

$$\text{Term}_1(n) = (\Lambda_0 + \tilde{\epsilon})^{k(n)} n, \quad \text{Term}_2(n) = \Lambda_1^{k(n)} \left(n^{1-\theta\tilde{D}} + \mathbb{E}_{\text{out}_{\mathcal{M}_n}[B_{\Lambda_0 + \tilde{\epsilon}}(0) \cup B_{n^{-\theta}}(\tilde{L})]} \right);$$

let us show that $\text{Term}_1(n), \text{Term}_2(n)$ are both $= o(|\ell|^{k(n)} n^{-j})$: we have already shown

$$(\Lambda_0 + \tilde{\epsilon})^{k(n)} n = o(|\ell|^{k(n)} n^{-j})$$

in (66), so $\text{Term}_1(n) = o(|\ell|^{k(n)} n^{-j})$. Also

$$\begin{aligned} \mathbb{E}_{\text{out}_{\mathcal{M}_n}[B_{\Lambda_0 + \tilde{\epsilon}}(0) \cup B_{n^{-\theta}}(L_r)]} &\leq \mathbb{E}_{\text{out}_{\mathcal{M}_n}[B_{\Lambda_0 + \tilde{\epsilon}}(0) \cup B_{n^{-\theta}}(L_{r_1})]} \leq n^{-\tilde{\alpha}} \\ &= n^{-j-1-\kappa \log(\Lambda_1) + \kappa \log(\Lambda_0 + 2\tilde{\epsilon})} \end{aligned}$$

by (58). Furthermore (61) implies that

$$1 - \theta\tilde{D} \leq \kappa \log(\Lambda_0 + 2\tilde{\epsilon}) - j - 1 - \kappa \log(\Lambda_1)$$

and therefore

$$n^{1-\theta\tilde{D}} \leq n^{-j-1-\kappa \log(\Lambda_1) + \kappa \log(\Lambda_0 + 2\tilde{\epsilon})}.$$

It follows that

$$\text{Term}_2(n) \leq \Lambda_1^{k(n)} n^{-j-1-\kappa \log(\Lambda_1) + \kappa \log(\Lambda_0 + 2\tilde{\epsilon})}$$

which—using $k(n) = \kappa \log n + O(1)$ (as before)—is bounded by

$$O(1/n)n^{-j}(\Lambda_0 + 2\tilde{\epsilon})^{k(n)} = o(|\ell|^{k(n)} n^{-j}).$$

Hence $\text{Term}(n) = o(|\ell|^{k(n)} n^{-j})$; combining this with (68), (69), (70), and (67) we have that

$$\left(1 + O(n^{-\theta})\right) \text{Ann}_{\tilde{D}, \tilde{L}}(\ell)\ell^k \mathbb{E}_{\text{in}_{\mathcal{M}_n}[B_{n^{-\theta}}(\ell)]} - \text{Ann}_{\tilde{D}, \tilde{L}}(S)(\ell^k q_{\ell, j}(k)n^{-j}) = o(|\ell|^{k(n)} n^{-j}).$$

Dividing by $\ell^k n^{-j}$ and applying Lemma 4.3 we have that

$$\text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \left(1 + O(n^{-\theta})\right) \mathbb{E}_{\text{in}_{\mathcal{M}_n}[B_{n^{-\theta}}(\ell)]} n^j - Q_j(k(n)) = o(1).$$

Dividing by the function that is $1 + O(n^{-\theta})$ above, we have

$$\text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] n^j = Q_j(k(n))(1 + o(1)) = Q_j(\kappa \log n)(1 + o(1))$$

where the last equality uses the fact that $k(n) = \kappa \log n + O(1)$.

Now we claim that $Q_j(k)$ must be a constant, rather than a polynomial of degree one or greater. Indeed, let $\kappa' \geq \kappa_0$ with $\kappa' \neq \kappa$. Repeating all the above with κ' replacing κ we have

$$\text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] n^j = Q_j(\kappa \log n)(1 + o(1)) = Q_j(\kappa' \log n)(1 + o(1))$$

as $n \rightarrow \infty$; the equality

$$Q_j(\kappa \log n)(1 + o(1)) = Q_j(\kappa' \log n)(1 + o(1))$$

as $n \rightarrow \infty$, and $\kappa' \neq \kappa$, implies that the polynomial Q_j must be a constant.

Since $Q_j = Q_j(k)$ is a constant, the limit

$$\lim_{n \rightarrow \infty} \left(\text{Ann}_{\tilde{D}, \tilde{L}}(\ell) \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] n^j \right)$$

exists and equals the constant Q_j . But Lemma 4.3 implies that $q_{\ell, j} = q_{\ell, j}(k)$ is also a constant and

$$Q_j = \text{Ann}_{\tilde{D}, \tilde{L}}(\ell) q_{\ell, j}.$$

It follows that

$$(71) \quad \lim_{n \rightarrow \infty} \left(\mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(\ell)] n^j \right) = q_{\ell, j}.$$

This proves (16) and (17).

Next let

$$L'_r = \{\ell \in L_r \mid |\ell| \geq \Lambda_0 + 2\epsilon/3\}.$$

For sufficiently large n —and more precisely for $n^{-\theta} < \epsilon/3$ (here we use $\theta > 0$)—we have

$$B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r) = B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1}) \cup B_{n^{-\theta}}(L'_r \setminus L_{j+1}).$$

It follows that

$$\mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1})] \leq \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] + \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(L'_r \setminus L_{j+1})].$$

But for all $\ell \in L'_r \setminus L_{j+1}$ we have $q_{\ell, j} = 0$ since $\ell \notin L_{j+1}$. Applying (71) to all $\ell \in L'_r \setminus L_{j+1}$ and summing over such ℓ we have

$$\mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1})] \leq \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_r)] + o(n^{-j}).$$

Combining this with (59) yields (15).

Similarly we have that if

$$L'_{j+1} = \{\ell \in L_{j+1} \mid |\ell| \geq \Lambda_0 + 2\epsilon/3\},$$

then for sufficiently large n

$$\begin{aligned} \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0)] &\leq \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1})] + \mathbb{E} \text{in}_{\mathcal{M}_n} [B_{n^{-\theta}}(L_{j+1})] \\ &\leq \mathbb{E} \text{out}_{\mathcal{M}_n} [B_{\Lambda_0 + \epsilon}(0) \cup B_{n^{-\theta}}(L_{j+1})] + O(n^{-j}), \end{aligned}$$

which proves (14). \square

5. PROOF OF THE SIDESTEPPING THEOREM

Proof of Theorem 5. Using induction on i , the formula (7) shows that for any $i \in \mathbb{Z}_{\geq 0}$, the function $c_i = c_i(k)$ appearing in the expansions (6) are independent of r over all $r \geq i + 1$ (for $r \leq i - 1$, the function c_i does not exist in (6), and for $r = i$ the term $O(c_i(k))/n^i$ does not uniquely determine c_i).

If $p_j(k)$, the polyexponential part of $c_j(k)$ with respect to Λ_0 , vanishes for all j , then according to Lemma 2.6, (14),

$$\mathbb{E}\text{out}_{\mathcal{M}_n}[B_{\Lambda_0+\epsilon}(0)] = O(n^{-j})$$

for all j .

Otherwise let j be the smallest integer such that $p_j(k) \neq 0$. Then $p_j(k)$ has a finite number of bases, so there exists an $\epsilon_0 > 0$ such that (1) all of these bases have absolute value at least $\Lambda_0 + 2\epsilon_0/3$, and (2) $\Lambda_0 + \epsilon_0 \leq \Lambda_1$. According to Lemma 2.6, for any $\epsilon > 0$ with $\epsilon \leq \epsilon_0$, the existence of an expansion of order $r_1 = r_1(\Lambda_0, \Lambda_1, j, \epsilon)$ for \mathcal{M}_n (which exist for any order) implies (15) and that

$$p_j(k) = \sum_{\ell \in L_{j+1}} C_\ell \ell^k$$

where the $C_\ell > 0$ are constants given by (16) for $\theta > 0$ satisfying $\theta \leq \theta_1(\Lambda_0, \Lambda_1, j, \epsilon)$. This proves (9)–(11), provided that $\epsilon \leq \epsilon_0$ and $\theta \leq \theta_1$; of course, if $\epsilon \geq \epsilon_0$, then (9) for $\epsilon = \epsilon_0$ also implies this for any larger value of ϵ . \square

REFERENCES

- [BS87] Andrei Broder and Eli Shamir, *On the second eigenvalue of random regular graphs*, Proceedings 28th Annual Symposium on Foundations of Computer Science, 1987, pp. 286–294.
- [FK14] Joel Friedman and David-Emmanuel Kohler, *The relativized second eigenvalue conjecture of alon*, Available at <http://arxiv.org/abs/1403.3462>.
- [Fri91] Joel Friedman, *On the second eigenvalue and random walks in random d -regular graphs*, *Combinatorica* **11** (1991), no. 4, 331–362. MR 93i:05115
- [Fri03] ———, *Relative expanders or weakly relatively Ramanujan graphs*, *Duke Math. J.* **118** (2003), no. 1, 19–35. MR 1978881
- [Fri08] ———, *A proof of Alon’s second eigenvalue conjecture and related problems*, *Mem. Amer. Math. Soc.* **195** (2008), no. 910, viii+100. MR 2437174
- [LP10] Nati Linial and Doron Puder, *Word maps and spectra of random graph lifts*, *Random Structures Algorithms* **37** (2010), no. 1, 100–135. MR 2674623
- [Pud15] Doron Puder, *Expansion of random graphs: new proofs, new results*, *Invent. Math.* **201** (2015), no. 3, 845–908. MR 3385636

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z4, CANADA

E-mail address: `jf@cs.ubc.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA

Current address: 422 Richards St, Suite 170, Vancouver BC V6B 2Z4, CANADA

E-mail address: `David.kohler@a3.epfl.ch`