

BASE SIZES FOR FINITE GROUPS OF LIE TYPE IN CROSS-CHARACTERISTIC

MELISSA LEE

ABSTRACT. Let $G \leq \mathrm{GL}(V)$ be a group with a unique subnormal quasisimple subgroup $E(G)$ that acts absolutely irreducibly on V . A base for G acting on V is a set of vectors with trivial pointwise stabiliser in G . In this paper we determine the minimal base size of G when $E(G)/Z(E(G))$ is a finite simple group of Lie type in cross-characteristic. We show that G has a regular orbit on V , with specific exceptions, for which we find the base size.

1. INTRODUCTION

Suppose $G \leq \mathrm{Sym}(\Omega)$ is a permutation group. We say that G has a *regular orbit* on Ω if there exists $\omega \in \Omega$ with trivial stabiliser in G . The study of regular orbits arises in a number of contexts, particularly when Ω is a vector space V , and G acts linearly on V . For example, if V is a finite vector space, $G \leq \mathrm{GL}(V)$, and all of the orbits of G on $V \setminus \{0\}$ are regular, then the affine group GV is a Frobenius group with Frobenius complement G , and such G were classified by Zassenhaus [33]. Regular orbits are also used heavily in the proof of the well-known $k(GV)$ -conjecture [11], which states that the number of conjugacy classes $k(GV)$ of GV , with $|G|$ coprime to $|V|$, is at most $|V|$. One of the major cases was where G is an almost quasisimple group acting irreducibly on V . Here, proving the existence of a regular orbit of G on V was sufficient to prove the $k(GV)$ -conjecture. A classification of such pairs (G, V) was completed by Köhler and Pahlings [26], based on work of Goodwin [12, 13] and Liebeck [28].

We say that $B \subset V$ is a *base* for G if its pointwise stabiliser in G is trivial, and call the minimal size of a base for G the *base size* and denote it by $b(G)$. Obviously if G has a regular orbit on V , then $b(G) = 1$. Since each element of a group is characterised by its action on a base, we have $b(G) \geq \lceil \log |G| / \log |V| \rceil$. Notice that if G has a regular orbit on an irreducible G -module V , then $GV \leq \mathrm{AGL}(V)$ has $b(GV) = 2$. In recent years there has been significant progress towards the more general aim of classifying finite primitive groups H with $b(H) = 2$. This includes a partial classification for diagonal type groups [7], a complete classification for primitive actions of S_m and A_m [4, 24] and sporadic groups [6] and also progress for almost simple classical groups [5].

We say that a group G is *quasisimple* if it is perfect and $G/Z(G)$ is a non-abelian simple group. We further define a group G to be *almost quasisimple* if $G/Z(G)$ is an almost simple group, or equivalently if it has a unique quasisimple subnormal subgroup, which forms the layer $E(G)$ of G . A large open case in the classification of primitive groups H with $b(H) = 2$ is where $H = GV$ is of affine type, $(|G|, |V|) > 1$ and $G \leq \mathrm{GL}(V)$ is an almost quasisimple group, whose quasisimple layer $E(G)$ acts absolutely irreducibly on V . The classification of such pairs (G, V) where G has a regular orbit on V has been completed for sporadic and alternating groups [8, 9]. In the infinite case, Guralnick and Lawther [15] classified (G, V) where G is a simple algebraic group that has a regular orbit on the irreducible G -module V , as well as determining their generic stabilisers in these cases. Their methods rely heavily on detailed analyses of highest weight representations of these simple algebraic groups.

This paper is the first in a series of three, which complete the analysis of base sizes of (G, V) , where G has a unique subnormal quasisimple subgroup $E(G)$ which is a group of Lie type that acts absolutely irreducibly on V . The present paper deals with the case where V is an absolutely irreducible module for $E(G)$ in cross characteristic. The other two papers will address representations of the groups of Lie type in defining characteristic. Throughout, we will assume that $E(G)/Z(E(G)) \not\cong A_m$, since this has been covered in [9].

Our main result is as follows.

Theorem 1.1. *Let $V = V_d(r)$ be a vector space of dimension d over \mathbb{F}_q . Suppose $G \leq \mathrm{GL}(V)$ and that the layer $E(G)$ of G is quasisimple of Lie type in characteristic $p \nmid r$ acting absolutely irreducibly on V . Also suppose that $E(G)/Z(E(G)) \not\cong A_n$. Then one of the following holds.*

- i) $b(G) = \lceil \log |G| / \log |V| \rceil$,
- ii) $b(G) = \lceil \log |G| / \log |V| \rceil + 1$ and (G, V) is in Table 1.1,
- iii) $(G, V) = (2 \times \mathrm{PSp}_6(2), V_7(3)), (U_4(2).2, V_5(3)), (2 \circ U_4(2), V_5(3))$ or $(\mathrm{PSp}_4(3).2, V_6(2))$ and $b(G) = \lceil \log |G| / \log |V| \rceil + 2 = 4, 4, 4, 5$ respectively.

$b(G)$	G	d	r	$b(G)$	G	d	r	
2	$3 \times L_2(8)$	2	64	2	$c \times \mathrm{PSp}_4(3), c \in \{3, 9, 21, 63\}$	4	64	
	$7 \times L_2(8)$	4	8		$\mathrm{PSp}_4(3) \leq G < 2 \times \mathrm{PSp}_4(3).2$	5	9	
	$2 \times L_2(8)$	7	3		$\mathrm{PSp}_4(3) < G \leq 12 \circ \mathrm{PSp}_4(3)^\dagger$	5	13	
	$L_2(8).3$	7	3		$c \circ \mathrm{PSp}_4(3), c \in \{6, 18\}^\dagger$	5	19	
	$L_2(11)$	5	4,5		$c \times \mathrm{PSp}_4(3), c = 2k \mid 26$	5	27	
	$2.L_2(11)$	6	3		$c \times (\mathrm{PSp}_4(3).2), c \mid 26$	5	27	
	$L_2(11).c, c \in \{1, 2\}$	10	2		$\mathrm{PSp}_4(3) \leq G \leq 2 \circ \mathrm{PSp}_4(3).2^\dagger$	6	7	
	$3 \times L_2(13)$	6	4		$3 \times \mathrm{PSp}_4(3).2^\dagger$	6	7	
	$c \times L_2(13), c \in \{1, 2\}$	7	3		$\mathrm{PSp}_4(3).c, c \in \{1, 2\}$	6	8	
	$L_2(13).c, c \in \{1, 2\}$	14	2		$10 \times \mathrm{PSp}_4(3)^\dagger$	6	11	
	$L_2(31)$	15	2		$\mathrm{PSp}_4(3).2^\dagger$	6	11	
	$c \circ (2.L_3(2)), c \in \{4, 8, 12\}$	2	49		$\mathrm{PSp}_4(3).2 < G \leq \mathbb{F}_r^\times \times \mathrm{PSp}_4(3).2^\dagger$	6	11,13	
	$4 \circ (2.L_3(2).2)$	2	49		$\mathrm{PSp}_4(3).2 \leq G \leq 15 \times (\mathrm{PSp}_4(3).2)$	6	16	
	$L_3(2) \leq G < 2 \times L_3(2).2$	3	7		$\mathrm{PSp}_4(3) \leq G < 2 \times (\mathrm{PSp}_4(3).2)$	10	3	
	$c \times L_3(2), c \in \{2, 4\}$	3	9		$c \times \mathrm{PSp}_6(2), c \mid 8$	7	9	
	$2 \times L_3(2)^\dagger$	3	11		$\mathrm{PSp}_6(2) \leq G \leq \mathbb{F}_r^\times \times \mathrm{PSp}_6(2)^\dagger$	7	11,13	
	$L_3(2) \leq G \leq 2 \times L_3(2).2$	6	3		$\mathbb{F}_r^\times \times \mathrm{PSp}_6(2)^\dagger$	7	17,19	
	$L_3(2)$	8	2		$c \circ (2.\mathrm{PSp}_6(2)), c \mid (r-1)$	8	7,9	
	$2.L_3(4) \leq G \leq 8 \circ (2.L_3(4).2^2)$	6	9		$c \times \mathrm{PSp}_6(2), c \in \{1, 2\}$	14	3	
	$4_1.L_3(4).2_3$	8	5		$c \times \mathrm{PSp}_6(3), c \in \{2, 3, 6\}$	13	7	
	$L_4(3) \leq G \leq L_4(3).2_2$	26	2		$2.\mathrm{PSp}_6(3) \leq G \leq 6 \circ (2.\mathrm{PSp}_6(3).2)$	14	7	
	$c \times U_3(3), c = 2k \mid 80$	3	81		$c \circ (2.\mathrm{P}\Omega_8^+(2))^\dagger, c \mid 12$	8	13	
	$U_3(3) < G < 2 \times U_3(3).2^\dagger$	6	5		$2.\mathrm{P}\Omega_8^+(2) \leq G \leq \mathbb{F}_r^\times \circ (2.\mathrm{P}\Omega_8^+(2).2)^\dagger$	8	17, 19, 23	
	$6 \times U_3(3)$	6	7		$2.\mathrm{P}\Omega_8^+(2) \leq G \leq \mathbb{F}_r^\times \circ (2.\mathrm{P}\Omega_8^+(2).2)$	8	25, 27	
	$c \times (U_3(3).2), c \in \{2, 6\}$	6	7		$c \circ (2.^2B_2(8)), c \mid 4$	8	5	
	$U_3(3) \leq G \leq 4 \circ U_3(3)^\dagger$	7	5*		$6 \circ (2.G_2(4).c), c \in \{1, 2\}$	12	7	
	$4 \circ U_3(3).2^\dagger$	7	5					
	$U_3(3).c, c \in \{1, 2\}$	14	2		3	$2.L_3(4) \leq G \leq 2.L_3(4).2^2$	6	3
	$2 \times U_3(4)$	12	3			$c \times U_3(3), c \in \{2, 4, 8\}$	3	9
	$U_3(5).A, 1 \neq A \leq S_3$	20	2			$3_1.U_4(3)$	6	4
	$3_1.U_4(3)$	6	16			$2 \times U_5(2).2$	10	3
	$6_1.U_4(3) \leq G \leq 6_1.U_4(3).2_2^\dagger$	6	19					
$c \circ (6_1.U_4(3)), c \mid 24$	6	25		$\mathrm{PSp}_4(3)$	4	4		
$c \circ (6_1.U_4(3).2_2), c \mid (r-1)$	6	25,49		$\mathrm{PSp}_4(3)$	5	3		
$c \circ (6_1.U_4(3)), c \mid (r-1)^\dagger$	6	31		$\mathrm{PSp}_4(3).2 \leq G \leq 3 \times \mathrm{PSp}_4(3).2$	6	4		
$c \circ (6_1.U_4(3).2_2), c \mid (r-1)^\dagger$	6	31,37		$c \times \mathrm{PSp}_4(3), c \in \{1, 3\}$	6	4		
$U_4(3).D_8$	20	2		$\mathrm{PSp}_6(2)$	7	3		
$2 \times U_5(2).2i$	10	9		$2.\mathrm{PSp}_6(2)$	8	3		
$c \times U_5(2).2, 1 < c \mid 8$	10	9		$2.\mathrm{P}\Omega_8^+(2) \leq G \leq 4 \circ (2.\mathrm{P}\Omega_8^+(2).2)$	8	5		
$U_5(2) < G \leq 6 \circ U_5(2)^\dagger$	10	7						
$U_6(2)$	21	3		4	$2.\mathrm{PSp}_4(3).c, c \in \{1, 2\}$	4	3	
					$3 \times \mathrm{PSp}_4(3)$	4	4	
$\mathrm{PSp}_4(3)$	4	16			$2 \times \mathrm{PSp}_4(3).2$	5	3	
$c \times \mathrm{PSp}_4(3), c \in \{6, 12, 24\}$	4	25			$\mathrm{PSp}_4(3)$	6	2	
$2.\mathrm{PSp}_4(3) \leq G \leq 2.\mathrm{PSp}_4(3).2$	4	27			$2.\mathrm{P}\Omega_8^+(2) \leq G \leq 2.\mathrm{P}\Omega_8^+(2).2$	8	3	
$\mathbb{F}_r^\times \circ (2.\mathrm{PSp}_4(3))^\dagger$	4	31,37						

TABLE 1.1. Pairs (G, V) with G an almost quasisimple group of Lie type for which $b(G) = \lceil \log |G| / \log |V| \rceil + 1$.

Combining the cases in Table 1.1 with the cases where $b(G) = \lceil \log |G| / \log |V| \rceil > 1$, we obtain the following corollary.

$b(G)$	G	d	r	$b(G)$	G	d	r	
2	$c \times L_2(8), c \mid 7$	2	8	2	$2.\text{PSp}_4(3) \leq G \leq 8 \circ (2.\text{PSp}_4(3).2)$	4	9	
	$c \times L_2(8), c = 3k \mid 63$	2	64		$c \times \text{PSp}_4(3), c \mid 15$	4	16	
	$7 \times L_2(8)$	4	8		$c \circ (2.\text{PSp}_4(3)), 1 \neq c \mid 18$	4	19	
	$2 \times L_2(8)$	7	3		$c \times \text{PSp}_4(3), c \in \{6, 12, 24\}$	4	25	
	$L_2(8).c, c \in \{1, 3\}$	8	2		$2.\text{PSp}_4(3) \leq G \leq 26 \circ (2.\text{PSp}_4(3).2)$	4	27	
	$c \times (L_2(8).3), c \in \{1, 2\}$	7	3		$\mathbb{F}_r^\times \circ (2.\text{PSp}_4(3))^\dagger$	4	31,37	
	$c \times L_2(11), c \mid (r-1)$	5	3,4,5		$c \times \text{PSp}_4(3), c \in \{3, 9, 21, 63\}$	4	64	
	$2.L_2(11)$	6	3		$\text{PSp}_4(3) < G \leq \mathbb{F}_r^\times \times \text{PSp}_4(3)^\dagger$	5	7,13	
	$L_2(11).c, c \in \{1, 2\}$	10	2		$\text{PSp}_4(3) \leq G \leq 8 \times (\text{PSp}_4(3).2)$	5	9	
	$2.L_2(13)$	6	3		$c \circ \text{PSp}_4(3), c \in \{6, 18\}^\dagger$	5	19	
	$3 \times L_2(13)$	6	4		$\text{PSp}_4(3) \leq G \leq 26 \times (\text{PSp}_4(3).2)$	5	27	
	$c \times L_2(13), c \in \{1, 2\}$	7	3		$c \times (\text{PSp}_4(3)), c = 2k \mid 26$	5	27	
	$L_2(13).c, c \in \{1, 2\}$	14	2		$c \times (\text{PSp}_4(3).2), c \mid 26$	5	27	
	$L_2(17)$	8	2		$c \times \text{PSp}_4(3), c \in \{1, 3\}$	6	4	
	$L_2(23)$	11	2		$\text{PSp}_4(3) \leq G \leq 4 \times (\text{PSp}_4(3).2)$	6	5	
	$L_2(25) \leq G \leq L_2(25).2_2$	12	2		$\text{PSp}_4(3) \leq G \leq 6 \times \text{PSp}_4(3).2^\dagger$	6	7	
	$L_2(31)$	15	2		$\text{PSp}_4(3) \leq G \leq 7 \times (\text{PSp}_4(3).2)$	6	8	
	$c \circ (2.L_3(2)), c \mid 6$	2	7		$10 \times \text{PSp}_4(3)^\dagger$	6	11	
	$c \circ (2.L_3(2)), c \mid 48$ and $c \neq \{2, 3, 6\}$	2	49		$\text{PSp}_4(3).2 < G \leq \mathbb{F}_r^\times \times \text{PSp}_4(3).2^\dagger$	6	11,13	
	$c \circ (2.L_3(2).2), c \mid 48$ and $c \neq \{2, 3, 6\}$	2	49		$c \times (\text{PSp}_4(3).2), c \mid 15$	6	16	
	$c \times L_3(2), c \mid 3$	3	4		$\text{PSp}_4(3) \leq G \leq 2 \times (\text{PSp}_4(3).2)$	10	3	
	$L_3(2) \leq G \leq 6 \times L_3(2).2$	3	7		$\text{PSp}_4(3).c, c \in \{1, 2\}$	14	2	
	$7 \times L_3(2)$	3	8		$\text{PSp}_4(7)$	24	2	
	$c \times L_3(2), c \mid 8$	3	9		$c \times \text{PSp}_6(2), c \mid (r-1)$	7	5,9	
	$c \times L_3(2)^\dagger, c \in \{2, 10\}$	3	11		$\text{PSp}_6(2) \leq G \leq \mathbb{F}_r^\times \times \text{PSp}_6(2)^\dagger$	7	11,13	
	$L_3(2) \leq G \leq 2 \times L_3(2).2$	6	3		$\mathbb{F}_r^\times \times \text{PSp}_6(2)^\dagger$	7	17,19	
	$L_3(2).c, c \in \{1, 2\}$	8	2		$c \circ (2.\text{PSp}_6(2)), c \mid (r-1)$	8	5,7,9	
	$L_3(3) \leq G \leq L_3(3).2$	12	2		$c \times \text{PSp}_6(2), c \in \{1, 2\}$	14	3	
	$2.L_3(4) \leq G \leq (r-1) \circ (2.L_3(4).2^2)$	6	3,9		$c \times \text{PSp}_6(3), c \mid 3$	13	4	
	$6.L_3(4) \leq G \leq 6.L_3(4).2_2$	6	7		$c \times \text{PSp}_6(3), c \in \{2, 3, 6\}$	13	7	
	$4_1.L_3(4).2_3$	8	5		$2.\text{PSp}_6(3) \leq G \leq 6 \circ (2.\text{PSp}_6(3).2)$	14	7	
	$L_4(3) \leq G \leq L_4(3).2_2$	26	2					
	$U_3(3)$	3	9		$2.\text{P}\Omega_8^+(2) \leq G \leq \mathbb{F}_r^\times \circ (2.\text{P}\Omega_8^+(2).2)$	8	7,9	
	$c \times U_3(3), c = 2k \mid 80$	3	81		$2.\text{P}\Omega_8^+(2) \leq G \leq \mathbb{F}_r^\times \circ (2.\text{P}\Omega_8^+(2).2)^\dagger$	8	13,17	
	$U_3(3) < G < 3 \times U_3(3).2$	6	4		$2.\text{P}\Omega_8^+(2) \leq G \leq \mathbb{F}_r^\times \circ (2.\text{P}\Omega_8^+(2).2)^\dagger$	8	19,23	
	$U_3(3) < G < 4 \times U_3(3).2^\dagger$	6	5		$2.\text{P}\Omega_8^+(2) \leq G \leq \mathbb{F}_r^\times \circ (2.\text{P}\Omega_8^+(2).2)$	8	25,27	
	$6 \times U_3(3)$	6	7					
	$c \times (U_3(3).2), c \in \{2, 6\}$	6	7		$c \circ (2.^2B_2(8)), c \mid 4$	8	5	
	$U_3(3) \leq G \leq 2 \times U_3(3).2$	7	3		$2.G_2(4).c, c \in \{1, 2\}$	12	3	
	$c \times U_3(3).2i, c \mid 4^{1*}$	7	5		$2.G_2(4) \leq G \leq 4 \circ (2.G_2(4).2)$	12	5	
	$U_3(3) \leq G \leq 4 \circ U_3(3)^{1*}$	7	5		$6 \circ (2.G_2(4).c), c \in \{1, 2\}$	12	7	
	$U_3(3).c, c \in \{1, 2\}$	14	2					
	$2 \times U_3(4)$	12	3		3	$L_2(7)$	2	2
	$U_3(5).A, 1 \neq A \leq S_3$	20	2			$2.L_3(4) \leq G \leq 2.L_3(4).2^2$	6	3
	$6_1.U_4(3)$	6	7					
	$c \circ (3_1.U_4(3)), c \mid 15$	6	16			$c \times U_3(3), c \in \{2, 4, 8\}$	3	9
	$c \circ (3_1.U_4(3).2_2), c \mid 15$	6	16			$U_3(3).c, c \in \{1, 2\}$	6	2
$c \circ (6_1.U_4(3)), c \mid (r-1)^\dagger$	6	13,19		$3_1.U_4(3)$	6	4		
$c \circ (6_1.U_4(3)), 1 \neq c \mid (r-1)^\dagger$	6	31		$2 \times U_5(2).2$	10	3		
$c \circ (6_1.U_4(3).2_2), c \mid (r-1)^\dagger$	6	13,19						
$c \circ (6_1.U_4(3).2_2), c \mid (r-1)^\dagger$	6	31,37		$\text{PSp}_4(3)$	4	4		
$c \circ (6_1.U_4(3)), c \mid 24$	6	25		$\text{PSp}_4(3)$	5	3		
$c \circ (6_1.U_4(3).2_2), c \mid (r-1)$	6	25,49		$\text{PSp}_4(3).2 \leq G \leq 3 \times \text{PSp}_4(3).2$	6	4		
$U_4(3).D_8$	20	2		$\text{PSp}_6(2)$	7	3		
$U_5(2) \leq G < 2 \times U_5(2).2$	10	3		$2.\text{PSp}_6(2)$	8	3		
$U_5(2) \leq G \leq 4 \times U_5(2).2$	10	5						
$U_5(2) < G \leq 6 \times U_5(2)^\dagger$	10	7		$2.\text{P}\Omega_8^+(2) \leq G \leq 4 \circ (2.\text{P}\Omega_8^+(2).2)$	8	5		
$8 \times U_5(2)$	10	9						
$2 \times U_5(2).2i$	10	9						
$c \times U_5(2).2, 1 < c \mid 8$	10	9		4	$2.\text{PSp}_4(3).c, c \in \{1, 2\}$	4	3	
$U_6(2) \leq G \leq 2 \times (U_6(2).S_3)$	21	3		$3 \times \text{PSp}_4(3)$	4	4		
				$\text{PSp}_4(3) < G \leq 2 \times \text{PSp}_4(3).2$	5	3		
				$\text{PSp}_4(3)$	6	2		
				$2 \times \text{PSp}_6(2)$	7	3		
				$2.\text{P}\Omega_8^+(2) \leq G \leq 2.\text{P}\Omega_8^+(2).2$	8	3		
				5	$\text{PSp}_4(3).2$	6	2	

TABLE 1.2. Pairs (G, V) with G an almost quasisimple group of Lie type for which $b(G) > 1$.

Corollary 1.2. *Let $V = V_d(r)$ be a vector space. Suppose $G \leq \mathrm{GL}(V)$ and that the layer $E(G)$ of G is quasisimple of Lie type in characteristic $p \nmid r$ acting absolutely irreducibly on V . Also suppose that $E(G)/Z(E(G)) \not\cong A_n$. Then either G has a regular orbit on V , or (G, V) appear in Table 1.2.*

- Remark 1.3.**
- (1) The entries in Tables 1.1 and 1.2 marked with \dagger can be deduced from [26] and constructions of the relevant modules in GAP [10]. This deals entirely with the case where $(|G|, |V|) = 1$, and the remainder of this paper is devoted to the case where $(|G|, |V|) > 1$.
 - (2) In the cases $L_3(2) \cong L_2(7)$, $\mathrm{PSp}_4(3) \cong U_4(2)$, $L_2(8) \cong {}^2G_2(3)'$ and $U_3(3) \cong G_2(2)'$ where there is an exceptional isomorphism between two groups of different characteristics, only one of the groups appears in Tables 1.1, and 1.2, and we complete the analysis in all characteristics dividing the order of the group. As mentioned earlier, we do not consider groups of Lie type that are isomorphic to alternating groups.
 - (3) We require some additional information for the asterisked entries in Tables 1.1 and 1.2. From the character table in [25], $U_3(3)$ has three absolutely irreducible 7-dimensional modules. Only one of these may be extended to $U_3(3).2$. The first asterisked line in Table 1.2 refers to this module, while the asterisked line in Table 1.1, and the second such line in Table 1.2 refers to one of the two other modules.

The following result is an immediate consequence of Table 1.2.

Corollary 1.4. *Let $V = V_d(r)$ be a vector space. Suppose $G \leq \mathrm{GL}(V)$ and that the layer $E(G)$ of G is quasisimple of Lie type in characteristic $p \nmid r$ acting absolutely irreducibly on V . If either $d > 24$, or $|V| > 27^8$ then G has a regular orbit on V .*

The remainder of the paper is organised as follows. In Section 2, we give some preliminary results from the literature concerning properties of groups of Lie type, such as minimal degrees of irreducible representations, generation by conjugates and bounds on the number of certain prime order elements. In Section 3, we outline the techniques that we will use to prove Theorem 1.1, and present some key propositions which provide the mechanism for the vast majority of the proof. We then proceed with the proof of Theorem 1.1 in Sections 4–8 by analysing, in order, the linear, unitary, symplectic, orthogonal and exceptional groups.

2. PRELIMINARIES

From now on, we will let G be an almost quasisimple group, and $V = V_d(r)$ be an irreducible $\mathbb{F}_r G$ -module, that is also absolutely irreducible under the action of $E(G)$. By Remark 1.3, we will also assume that $(|G|, |V|) > 1$. For $g \in G$, let $\mathcal{E}_{\max}^V(g)$ (or just $\mathcal{E}_{\max}(g)$) denote the dimension of the largest eigenspace of g on $\overline{V} = V \otimes \overline{\mathbb{F}_r}$. For a prime r_0 dividing $|G|$, we will denote by g_{r_0} an element of order r_0 , and by $g_{r'_0}$ an element of prime order not r_0 . For a set of primes P , we will denote g_P an element whose order lies in P , and will use $g_{P'}$ to denote an element of prime order whose order does not lie in P .

2.1. Minimal degrees of cross-characteristic representations. The degrees of irreducible cross-characteristic representations of quasisimple groups of Lie type are not known in general. However, we are able to rely on some partial results. Hiss and Malle [21] give a complete list of irreducible representations of quasisimple groups of degree at most 250. This suffices for our examination of the exceptional groups and small classical groups. For large classical groups, we instead utilise results giving the minimal degree of a representation in cross-characteristic, and lower bounds on the degree of the next smallest representation. We outline some of these results below.

Proposition 2.1 ([31, Tables I, II]). *Suppose $H = H(q)$, $q = p^e$, is a quasisimple group of Lie type acting irreducibly on the $\mathbb{F}_r H$ -module V with $p \nmid r$. Then $\dim V \geq d_1(H)$, where $d_1(H)$ is given in Table 2.1.*

For linear groups, the following result will also be useful.

		Exceptions				Exceptions																																																																															
$H/Z(H)$	$d_1(H)$	(n, q)	$d_1(H)$	$H/Z(H)$	$d_1(H)$	q	$d_1(H)$																																																																														
$\mathrm{PSL}_2(q)$	$(q-1)/(2, q-1)$	(2, 4)	2	${}^3D_4(q)$	$q^5 - q^3 + q - 1$																																																																																
		(2, 9)	3					$\mathrm{PSL}_n(q), n \geq 3$	$\frac{q^n - q}{q-1} - 1$	(3, 2)	2	$E_6(q)$	$(q^5 + q)(q^6 + q^3 + 1) - 1$			(3, 4)	4	(4, 2)	6	(4, 3)	26	$\mathrm{PSU}_n(q)$	$\lfloor \frac{q^n - 1}{q+1} \rfloor$	(4, 2)	4	$F_4(q), q \text{ odd}$	$q^8 + q^4 - 2$			(4, 3)	6	$F_4(q), q \text{ even}$	$\frac{1}{2}(q^3 - 1)(q^8 - q^7)$	2	52	$\mathrm{PSp}_{2n}(q), 2 \nmid q$	$\frac{1}{2}(q^n - 1)$			$G_2(q), q \equiv 0(3)$	$q^4 + q^2$	3	14	$\mathrm{PSp}_{2n}(q), 2 \mid q$	$\frac{(q^n - 1)(q^n - q)}{2(q+1)}$	(2, 2)	4	$G_2(q), q \equiv 1(3)$	$q^3 - 1$	4	12	$\mathrm{P}\Omega_{2n}^+(q), q > 3$	$\frac{(q^n - 1)(q^{n-1} + q)}{q^2 - 1} - 2$			$G_2(q)', q \equiv 2(3)$	q^3			$\mathrm{P}\Omega_{2n}^+(q), q \leq 3$	$\frac{(q^n - 1)(q^{n-1} - 1)}{q^2 - 1} - 2$	(4, 2)	8	${}^2B_2(q)$	$(q-1)\sqrt{q/2}$	8	8	$\mathrm{P}\Omega_{2n}^-(q)$	$\frac{(q^n + 1)(q^{n-1} - q)}{q^2 - 1} - 1$			${}^2G_2(q)'$	$q(q-1)$			$\mathrm{P}\Omega_{2n+1}(q), q > 3$	$\frac{q^{2n} - 1}{q^2 - 1} - 2$			${}^2E_6(q)$	$(q^5 + q)(q^6 - q^3 + 1) - 1$	2	1938	$\mathrm{P}\Omega_{2n+1}(q), q = 3$	$\frac{(q^n - 1)(q^n - q)}{q^2 - 1}$
$\mathrm{PSL}_n(q), n \geq 3$	$\frac{q^n - q}{q-1} - 1$	(3, 2)	2	$E_6(q)$	$(q^5 + q)(q^6 + q^3 + 1) - 1$																																																																																
		(3, 4)	4																																																																																		
		(4, 2)	6																																																																																		
		(4, 3)	26																																																																																		
$\mathrm{PSU}_n(q)$	$\lfloor \frac{q^n - 1}{q+1} \rfloor$	(4, 2)	4	$F_4(q), q \text{ odd}$	$q^8 + q^4 - 2$																																																																																
		(4, 3)	6					$F_4(q), q \text{ even}$	$\frac{1}{2}(q^3 - 1)(q^8 - q^7)$	2	52																																																																										
$\mathrm{PSp}_{2n}(q), 2 \nmid q$	$\frac{1}{2}(q^n - 1)$			$G_2(q), q \equiv 0(3)$	$q^4 + q^2$	3	14																																																																														
$\mathrm{PSp}_{2n}(q), 2 \mid q$	$\frac{(q^n - 1)(q^n - q)}{2(q+1)}$	(2, 2)	4	$G_2(q), q \equiv 1(3)$	$q^3 - 1$	4	12																																																																														
$\mathrm{P}\Omega_{2n}^+(q), q > 3$	$\frac{(q^n - 1)(q^{n-1} + q)}{q^2 - 1} - 2$			$G_2(q)', q \equiv 2(3)$	q^3																																																																																
$\mathrm{P}\Omega_{2n}^+(q), q \leq 3$	$\frac{(q^n - 1)(q^{n-1} - 1)}{q^2 - 1} - 2$	(4, 2)	8	${}^2B_2(q)$	$(q-1)\sqrt{q/2}$	8	8																																																																														
$\mathrm{P}\Omega_{2n}^-(q)$	$\frac{(q^n + 1)(q^{n-1} - q)}{q^2 - 1} - 1$			${}^2G_2(q)'$	$q(q-1)$																																																																																
$\mathrm{P}\Omega_{2n+1}(q), q > 3$	$\frac{q^{2n} - 1}{q^2 - 1} - 2$			${}^2E_6(q)$	$(q^5 + q)(q^6 - q^3 + 1) - 1$	2	1938																																																																														
$\mathrm{P}\Omega_{2n+1}(q), q = 3$	$\frac{(q^n - 1)(q^n - q)}{q^2 - 1}$	(3, 3)	27	${}^2F_4(q)'$	$(q^5 - q^4)\sqrt{q/2}$	2	26																																																																														

TABLE 2.1. Lower bounds $d_1(G)$ on representation degrees in cross-characteristic r .

Proposition 2.2 ([31, Theorem 4.1]). *Suppose $G = \mathrm{SL}_n(q)$ with $n \geq 3$, and $(n, q) \neq (3, 2), (3, 4), (4, 2), (4, 3), (6, 2)$ or $(6, 3)$. If V is a non-trivial absolutely irreducible $\mathbb{F}_r G$ -module of dimension less than*

$$\begin{cases} (q-1)(q^2-1)/(3, q-1) & \text{if } n = 3 \\ (q-1)(q^3-1)/(2, q-1) & \text{if } n = 4 \\ (q^{n-1}-1) \left(\frac{q^{n-2}-q}{q-1} - 1 \right) & \text{if } n \geq 5 \end{cases}$$

then V has dimension $(q^n - q)/(q - 1)$ or $(q^n - 1)/(q - 1)$.

2.2. Properties of prime order elements in groups of Lie type. For G_0 a simple group, and non-identity $x \in \mathrm{Aut}(G_0)$, define $\alpha(x)$ to be the minimal number of G_0 -conjugates of x needed to generate the group $\langle G_0, x \rangle$, and for almost quasisimple G , define

$$\alpha(G) = \max \{ \alpha(x) \mid 1 \neq x \in G/Z(G) \}.$$

We now give some results which compute bounds on $\alpha(G)$ for G a group of Lie type.

Proposition 2.3 ([17, Lemma 3.1]). *Let H be an almost simple group with $\mathrm{soc}(H) \cong \mathrm{PSL}_2(q)$, and let $x \in H$ have prime order r . Then $\alpha(x) \leq 3$ unless:*

- i) x is an involutory field automorphism and $\alpha(x) \leq 4$, except $\alpha(x) = 5$ for $q = 9$, or*
- ii) $q = 5$, x is an involutory diagonal automorphism, and $\alpha(x) = 4$.*

Moreover, if the order of x is odd, then $\alpha(x) = 2$, unless $q = 9$, $r = 3$ and $\alpha(x) = 3$.

Proposition 2.4 ([17]). *Let G be an almost simple classical group over \mathbb{F}_q with socle $\mathrm{soc}(G) = G_0$ and natural module of dimension $n \geq 3$, taking $n \geq 7$ for G an orthogonal group. Then for $x \in G \setminus \{1\}$, one of the following holds:*

- i) $\alpha(x) \leq n$,*
- ii) $G_0 = \mathrm{PSp}_n(q)$, q even, and $\alpha(x) \leq n + 1$,*
- iii) $G_0 = L_3(q)$, x is an involutory graph-field automorphism and $\alpha(x) \leq 4$,*

- iv) $G_0 = L_4^\epsilon(q)$ with $q \geq 3$, and x is an involutory graph automorphism and $\alpha(x) \leq 6$,
- v) $G_0 = PSp_4(q)$ and $\alpha(x) \leq 5$, or
- vi) $G_0 = U_3(3)$, x is an inner involution with $\alpha(x) = 4$,
- vii) $G_0 = L_4^\epsilon(2)$.

Furthermore, if G_0 is a simple exceptional group of Lie type of untwisted Lie rank l , then $\alpha(G) \leq l+3$, except $\alpha(G) \leq 8$ for $G_0 = F_4(q)$, and $\alpha(G) \leq 3$ for $G_0 = {}^2G_2(q)$ or ${}^2B_2(q)$.

We also have these more precise bounds for symplectic and orthogonal groups.

Proposition 2.5 ([17, Theorem 4.3]). *Suppose $G_0 = \mathrm{PSp}_{2m}(q)$, and $x \in \mathrm{Aut}(G_0)$. Then either:*

- i) $\alpha(x) \leq m+3$,
- ii) x is a transvection and $\alpha(x) = 2m$ or $2m+1$ if q is odd or even respectively, or
- iii) $(m, q) = (2, 3)$ and x is an involution with $\alpha(x) = 6$.

Proposition 2.6 ([17, Theorem 4.4]). *Suppose $G_0 = \mathrm{P}\Omega_n^\epsilon(q)$ with $n = 2m$ where $m \geq 4$, or $n = 2m+1$ with $m \geq 3$ and q odd. Then if $x \in \mathrm{Aut}(G_0)$, either*

- i) $\alpha(x) \leq m+3$,
- ii) q is odd, x is a reflection and $\alpha(x) = n$, or
- iii) q is even and x is a transvection with $\alpha(x) = n$.

The remaining lemmas in this section count certain types of elements of prime order in groups of Lie type.

Proposition 2.7 ([27, Proposition 1.3]). *Let \bar{G} be a simple algebraic group over $\bar{\mathbb{F}}_p$ with associated root system Φ . Let σ be a Frobenius endomorphism of \bar{G} such that $G_0 = \bar{G}_\sigma'$ is a finite simple group of Lie type over \mathbb{F}_q . Assume that G_0 is not of type 2F_4 , 2G_2 or 2B_2 . Then*

- i) $i_2(\mathrm{Aut}(G_0)) < 2(q^{N_2} + q^{N_2-1})$, where $N_2 = \dim \bar{G} - \frac{1}{2}|\Phi|$, and
- ii) $i_3(\mathrm{Aut}(G_0)) < 2(q^{N_3} + q^{N_3-1})$, where $N_3 = \dim \bar{G} - \frac{1}{3}|\Phi|$.

For $N = \mathrm{Aut}(\mathrm{PSL}_n^\epsilon(q))$, we define $\gamma(N)$ denote the number of (involutory) graph automorphisms in N .

Lemma 2.8. i)

$$\gamma(\mathrm{Aut}(\mathrm{PSL}_n(q))) \leq \begin{cases} q^{(n^2+n)/2} & \text{if } n \text{ even, } q \text{ odd} \\ 2q^{(n^2-n)/2-1} & \text{if } n \text{ even, } q \text{ even} \\ q^{n^2/2-5/4} & \text{if } n \text{ odd} \end{cases}$$

and the number of graph-field automorphisms in $\mathrm{Aut}(\mathrm{PSL}_n(q))$ is less than $2q^{(n^2-1)/2}$.

ii)

$$\gamma(\mathrm{Aut}(\mathrm{PSU}_n(q))) \leq \begin{cases} 2q^{\frac{n^2-n}{2}+1}(q^n+1) & n \text{ even, } q \text{ odd} \\ 4q^{\frac{n(n+1)}{2}} & n \text{ even, } q \text{ even} \\ 2q^{\frac{n^2+n}{2}-1} & n \text{ odd} \end{cases}$$

Proof. We give the proof for part i); the others are similar. Let $H = \mathrm{PGL}_n(q)$. We adopt the notation of [3, §3.2.5]. When n is even and q is odd there are three distinct H -conjugacy classes of graph automorphisms [3, Table B.3] satisfying

$$|\gamma_1^H| = \frac{|\mathrm{PGL}_n(q)|}{|\mathrm{PGSp}_n(q)|} = q^{\frac{n^2-n}{4}-\frac{n}{2}} \prod_{i=1}^{n/2-1} (q^{2i+1} - 1) < q^{\frac{n^2-n}{2}-1},$$

$$|\gamma_2^H| = \frac{|\mathrm{PGL}_n(q)|}{|\mathrm{PGO}_n^+(q)|} = \frac{1}{2}q^{\frac{n^2}{4}}(q^{n/2}+1) \prod_{i=2}^{n/2} (q^{2i-1} - 1) < \frac{1}{2}q^{\frac{n^2}{2}-1}(q^{n/2}+1),$$

$$|(\gamma_2')^H| = \frac{|\mathrm{PGL}_n(q)|}{|\mathrm{PGO}_n^-(q)|} = \frac{1}{2}q^{\frac{n^2}{4}}(q^{n/2}-1) \prod_{i=2}^{n/2} (q^{2i-1} - 1) < \frac{1}{2}q^{\frac{n^2}{2}-1}(q^{n/2}-1).$$

We have $|\gamma_1^H| + |\gamma_2^H| + |(\gamma_2')^H| < q^{\frac{n^2-n}{2}-1} + \frac{1}{2}q^{\frac{n^2}{2}-1}(q^{n/2}+1) + \frac{1}{2}q^{\frac{n^2}{2}-1}(q^{n/2}-1) < q^{\frac{n^2+n}{2}}$ as required. If instead n is even and q is even, then there are two $\mathrm{PGL}_n(q)$ -conjugacy classes of graph automorphisms, namely γ_1^H (as above) and γ_3^H . The total number of graph automorphisms in this case is

$$|\gamma_1^H| + |\gamma_3^H| < q^{\frac{n^2-n}{2}-1} + \frac{|\mathrm{PGL}_n(q)|}{q^{n^2/4} \prod_{i=1}^{n/2-1} (q^{2i}-1)} = q^{\frac{n^2-n}{2}-1} + q^{\frac{n^2}{4}-\frac{n}{2}} \prod_{i=2}^{n/2} (q^{2i-1}-1) < 2q^{\frac{n^2-n}{2}-1}.$$

If n is odd, then there is only one $\mathrm{PGL}_n(q)$ -conjugacy class of graph automorphisms, satisfying

$$|\gamma_1^H| \leq \frac{|\mathrm{PGL}_n(q)|}{|\mathrm{Sp}(n-1, q)|} = q^{\frac{n^2-1}{4}} \prod_{i=2}^{(n+1)/2} (q^{2i-1}-1) < q^{\frac{n^2}{2}-\frac{5}{4}}.$$

Finally, if q is a square, there is one $\mathrm{PGL}_n(q)$ -conjugacy class of graph-field automorphisms and the size of this class is

$$\frac{|\mathrm{PGL}_n(q)|}{|\mathrm{PGU}(n, q^{1/2})|} < 2q^{\frac{n^2-1}{2}}$$

□

2.3. Weil representations of symplectic and unitary groups. We give some background on Weil representations of $\mathrm{Sp}_{2m}(q)$, q odd, and $\mathrm{SU}_m(q)$, beginning with the former. Suppose $q = p^e$ is odd, and let $r_0 \nmid q$ be a prime. Let $S = q^{1+2m}$ be a special group with exponent p and $Z(S) = S'$ of order q . For every non-trivial irreducible character χ of $Z(S)$, there is a unique irreducible q^m -dimensional representation of S over $\overline{\mathbb{F}}_{r_0}$ where $Z(S)$ acts via χ [16, §5]. Now, $\mathrm{Sp}_{2m}(q)$ acts as a group of automorphisms of S , so preserves the character of the q^m -dimensional representation. Let $H = S : \mathrm{Sp}_{2m}(q)$. We may construct an irreducible $\overline{\mathbb{F}}_{r_0}H$ -module V_0 which restricts to S as the q^m -dimensional representation.

The restriction of V_0 to $\mathrm{Sp}_{2m}(q)$ is called a *Weil module* for $\mathrm{Sp}_{2m}(q)$, with corresponding *Weil character*. The non-trivial irreducible composition factors of this Weil representation correspond to the minimal degree representations of $\mathrm{Sp}_{2m}(q)$, q odd, in characteristic r_0 , unless $(m, q) = (2, 2)$ (cf. Table 2.1). In our work, we will additionally use the terms *Weil representation* and *Weil module* to refer to one of these (absolutely) irreducible composition factors. If $r_0 \neq 2$, there are four inequivalent such Weil modules of $\mathrm{Sp}_{2m}(q)$ in characteristic r_0 – two of dimension $(q^m - 1)/2$, and two of dimension $(q^m + 1)/2$. If instead $r_0 = 2$, then there are two Weil modules up to equivalence, both of dimension $(q^m - 1)/2$ [30].

The Weil modules for a unitary group $\mathrm{SU}_m(q) \leq \mathrm{Sp}_{2m}(q)$ for q odd are the composition factors of the restriction of V_0 to $\mathrm{SU}_m(q)$. These Weil modules for $\mathrm{SU}_m(q)$ are of dimension $(q^n - (-1)^n)/(q+1)$ or $(q^n + q(-1)^n)/(q+1)$, with the number of modules of each dimension dependent on r_0 , n and q [22]. If $m \geq 4$ and $(m, q) \neq (4, 3)$, then these irreducible Weil representations are the minimal degree representations of $\mathrm{SU}_m(q)$, q odd, in characteristic r_0 .

For unitary groups, we may additionally define Weil modules for q even [32, §4]. In this case, there is an irreducible embedding of the symplectic type group $4 \circ 2^{1+2m}$ into $\mathrm{GL}_{2m}(\overline{\mathbb{F}}_{r_0})$, with normaliser $\overline{\mathbb{F}}_{r_0}^\times \circ 2^{1+2m} \cdot \mathrm{Sp}_{2m}(q)$. This is a non-split extension. However, $4 \circ 2^{1+2m} \cdot \mathrm{SU}_m(q)$ is a split extension, and the composition factors of the restriction of $V_0 = V_{2m}(\overline{\mathbb{F}}_{r_0})$ to $\mathrm{SU}_m(q)$ give a set of *Weil modules* for $\mathrm{SU}_m(q)$. Similarly to q odd, if $m \geq 4$ and $(m, q) \neq (4, 2)$, then these irreducible Weil representations are the minimal degree representations of $\mathrm{SU}_m(q)$ in characteristic r_0 .

As in the case of the symplectic groups, we will also sometimes use the term *Weil module* to refer to the q^m -dimensional restriction of V_0 to $\mathrm{SU}_m(q)$.

We can often infer the values of the Brauer character of a q^m dimensional Weil representation over $\overline{\mathbb{F}}_r$ from those of the ordinary (complex) character. This allows us to place upper bounds on the eigenspace dimensions of group elements on the irreducible composition factors. The following results give us the means to compute these ordinary character values.

Proposition 2.9 ([23, Theorem 4.8]). *Suppose q is an odd prime power. Then $\mathrm{Sp}_{2m}(q)$ has irreducible ordinary Weil characters χ_1, χ_2 of degrees $(q^m \pm 1)/2$. Moreover, $|\chi_1(g) + \chi_2(g)|^2 = |C_W(g)|$ for all $g \in \mathrm{Sp}_{2m}(q)$, where W is the natural module for $\mathrm{Sp}_{2m}(q)$ over \mathbb{F}_q .*

Proposition 2.10 ([18, Corollary 4.9.2]). *For $g \in \mathrm{SU}_n(q)$, the ordinary Weil character of degree q^n of $\mathrm{SU}_n(q)$ has value $(-1)^n(-q)^{\dim C_W(g)}$, where W is the natural module for $\mathrm{SU}_n(q)$.*

Proposition 2.11. *Suppose G is a group acting irreducibly on the $\overline{\mathbb{F}}_r G$ -module $V = V_d(\overline{\mathbb{F}}_r)$ with corresponding Brauer character ρ . Suppose $x \in G$ is of prime order $p \nmid r$. Let E_x denote the set of eigenvalues of x (including multiplicities), $\overline{\Omega}$ denote the set of primitive p th roots of unity in $\overline{\mathbb{F}}_r$, and $\Omega = \overline{\Omega} \cup \{1\}$. Let $c \in \mathbb{N}$. Then:*

- i) *If $\rho(x) = c$, then $E_x = (1^c, \Omega^{(d-c)/p})$.*
- ii) *If $\rho(x) = -c$, then $E_x = (\overline{\Omega}^c, \Omega^{(d-cp+c)/p}) = (\overline{\Omega}^{(d+c)/p}, 1^{(d-cp+c)/p})$.*
- iii) *If $|\rho(y)| = |\rho(z)|$ for y, z elements of the same projective prime order in G , then $E_y = \lambda E_z$ for some $\lambda \in \overline{\mathbb{F}}_r$. In particular, y and z have the same set of eigenspace dimensions.*

Proof. Notice that the sets E_x in parts i) and ii) sum to $\rho(x)$ in each case, so it remains to show that E_x is uniquely determined. The primitive p th roots of unity are linearly independent over \mathbb{Z} , which implies uniqueness. The argument for iii) is similar. \square

If g is of odd prime order not equal to r_0 , we can use [3, §3.3.1, 3.4.1] to limit the dimension of $C_W(g)$.

Proposition 2.12. *Suppose g is a semisimple element of odd prime order r_0 in $\mathrm{Sp}_n(q)$ or $\mathrm{SU}_n(q)$. Define $i = \min\{k \in \mathbb{N} \mid r_0 \mid q^k - 1\}$. Let $t = 1/2$ if $g \in \mathrm{SU}_n(q)$ has $i \equiv 2(4)$ and $i > 2$, let $t = 1$ if i is even otherwise, and let $t = 2$ if i is odd. Then unless $g \in \mathrm{SU}_n(q)$ and $i = 2$,*

$$\dim C_W(g) \in \left\{ n - tij \mid 1 \leq j \leq \left\lfloor \frac{n}{ti} \right\rfloor \right\}$$

Proof. Follows immediately from [3, Propositions 3.3.2, 3.4.3]. \square

For large symplectic and unitary groups it will also be useful to bound the degree of the smallest representation that is not a Weil representation.

Proposition 2.13 ([22]). *Suppose $G = \mathrm{SU}_n(q)$ with $n \geq 4$ and $(n, q) \neq (4, 2), (4, 3)$. Let \mathbb{F} be an algebraically closed field of characteristic $r \neq p$. Suppose that V is a non-trivial absolutely irreducible $\mathbb{F}G$ -module of degree less than*

$$\begin{cases} (q^2 + 1)(q^2 - q + 1)/(2, q - 1) - 1 & \text{if } n = 4, \\ (q^{n-2} - 1)(q - 1)(q^{n-2} - 1)/(q + 1) & \text{if } n \geq 5 \end{cases}$$

Then V is a Weil module of degree $(q^n - (-1)^n)/(q + 1)$ or $(q^n + q(-1)^n)/(q + 1)$.

Proposition 2.14 ([16]). *Suppose $G = \mathrm{Sp}_{2m}(q)$ with $n \geq 2$, q odd and $(m, q) \neq (2, 3)$. Let \mathbb{F} be an algebraically closed field of characteristic $r \neq p$. If V is a non-trivial absolutely irreducible $\mathbb{F}G$ -module for G of dimension less than $(q^m - 1)(q^m - q)/2(q + 1)$, then V is a Weil module of dimension $(q^m \pm 1)/2$.*

3. TECHNIQUES

Let G be an almost quasisimple group with $E(G)$ a quasisimple group of Lie type acting absolutely irreducibly on $V_d(r)$ in cross characteristic. Recall that we also assume $(r, |G|) > 1$, since the co-prime case was completed by [26]. Our method of proof for Theorem 1.1 relies heavily on the observation that if G has no regular orbit on V , then V is the union of the fixed points spaces $C_V(g)$ for $g \in G \setminus \{1\}$. Since conjugates in G have fixed point spaces of the same dimension, we have

$$|V| \leq \sum_{x \in \mathcal{X}} |x^G| |C_V(x)|, \quad (1)$$

where \mathcal{X} is a set of non-identity conjugacy class representatives of G .

We say that an element $g \in G$ is of *projective prime order* if the coset $gZ(G)$ has prime order in $G/Z(G)$.

The following proposition gives the main results used to prove Theorem 1.1. Let H_s denote the set of elements of a group H of projective prime order s , and $H_{s'}$ denote the set of elements of H with projective prime order coprime to s .

Proposition 3.1. *Let $G \leq \mathrm{GL}(V)$ be an almost quasisimple group, with layer $E(G)$ that acts absolutely irreducibly on the d -dimensional module $V = V_d(r)$ over \mathbb{F}_r . Set $H = G/Z(G)$ and let \mathcal{P} be a set of conjugacy class representatives of elements of projective prime order in G . Denote the order of $x \in H$ by $o(x)$, and let $\overline{V} = V \otimes \overline{\mathbb{F}}_r$. Then the following statements hold.*

i) *For $\nu \in \overline{\mathbb{F}}_r$, the ν -eigenspace $E_\nu(x)$ of $x \in G$ on \overline{V} satisfies*

$$\dim_{\overline{\mathbb{F}}_r}(E_\nu(x)) \leq \left\lfloor \dim_{\overline{\mathbb{F}}_r}(\overline{V}) \left(1 - \frac{1}{\alpha(x)}\right) \right\rfloor$$

Further, if G has no regular orbit on $V = V_d(r)$ with $r = s^e$ for s prime,

ii)

$$|V| \leq \sum_{x \in \mathcal{P}} \sum_{\kappa \in \overline{\mathbb{F}}_r} \frac{1}{o(x) - 1} |xZ(G)^H| |C_V(\kappa x)|$$

iii)

$$|V| \leq \sum_{x \in \mathcal{P} \cap G_{s'}} \frac{o(x)}{o(x) - 1} |xZ(G)^H| \max\{|C_V(\kappa x)| \mid \kappa \in \overline{\mathbb{F}}_r^\times\} + \sum_{x \in \mathcal{P} \cap G_s} \frac{1}{o(x) - 1} |xZ(G)^H| |C_V(x)|$$

iv)

$$|V| \leq 2 \sum_{x \in \mathcal{P} \cap G_{s'}} |xZ(G)^H| \max\{|C_V(\kappa x)| \mid \kappa \in \overline{\mathbb{F}}_r^\times\} + \sum_{x \in \mathcal{P} \cap G_s} \frac{1}{o(x) - 1} |xZ(G)^H| |C_V(x)|$$

v)

$$|V| = r^d \leq 2 \sum_{x \in \mathcal{P} \cap G_{s'}} |xZ(G)^H| |r|^{\lfloor (1-1/\alpha(x))d \rfloor} + \sum_{x \in \mathcal{P} \cap G_s} \frac{1}{o(x) - 1} |xZ(G)^H| |r|^{\lfloor (1-1/\alpha(x))d \rfloor} \quad (2)$$

and for fixed r , if this inequality fails for a given d , then it fails for any $d_1 \geq d$.

Proof. Part i) follows from [19, p. 454]. We will prove the other parts of the proposition simultaneously. Let $x \in G$ be of projective prime order. Then there exists a prime a such that $x^a = z \in Z(G)$. Now, we may choose $\hat{z} \in \overline{\mathbb{F}}_r$ so that $\hat{z}^a = z^{-1}$, and so $g = \hat{z}x \in \overline{\mathbb{F}}_r \cdot G$ has prime order a . Then g has at most a eigenspaces on $V \otimes \overline{\mathbb{F}}_r$, and therefore x has at most a eigenspaces on V . Let \mathcal{H} be a set of coset representatives of elements of prime order in H . Now assuming that G has no regular orbit on V ,

$$V = \bigcup_{x \in \mathcal{H}} \bigcup_{\kappa \in Z(G)} C_V(\kappa x) \quad (3)$$

From (3), noting that $C_V(\kappa x)$ is precisely the κ^{-1} eigenspace of x , and that $C_V(g) \subseteq C_V(g^i)$ for all $g \in G \setminus Z(G)$, we have

$$\begin{aligned} |V| &\leq \sum_{x \in \mathcal{H}} \sum_{\kappa \in \overline{\mathbb{F}}_r} \frac{1}{o(x) - 1} |C_V(\kappa x)| \\ &= \sum_{x \in \mathcal{P}} \sum_{\kappa \in \overline{\mathbb{F}}_r} \frac{1}{o(x) - 1} |x^H| |C_V(\kappa x)| \\ &\leq \sum_{x \in \mathcal{P} \cap G_{s'}} \frac{o(x)}{o(x) - 1} |x^H| \max\{|C_V(\kappa x)| \mid \kappa \in \overline{\mathbb{F}}_r^\times\} + \sum_{x \in \mathcal{P} \cap G_s} \frac{1}{o(x) - 1} |x^H| |C_V(x)|, \end{aligned}$$

since unipotent elements in V have only one eigenspace. We have thus proved ii) and iii). Further, iv) follows from iii) since $o(x) \geq 2$, and equation (2) in v) follows from iv) and i). Finally, we prove that if we fix r and (2) fails for a given d , then it fails for any $d_1 \geq d$. Write $d_1 = d + k$. Comparing (2) for d and d_1 , we see that the proof reduces to showing that for $x \in \mathcal{P}$ we have $\lfloor (1-1/\alpha(x))d \rfloor + k \geq \lfloor (1-1/\alpha(x))(d+k) \rfloor$. We observe that $\lfloor (1-1/\alpha(x))d \rfloor + k = \lfloor (1-1/\alpha(x))d + k \rfloor$ and the result follows. \square

The next two results present additional methods of bounding $C_V(g)$ for g of projective prime order in G .

Proposition 3.2. *Suppose that G is a finite group and V is an $\mathbb{F}_r G$ -module. Let V_1, \dots, V_k be the composition factors of V under the action of G . Then for an element $x \in G$,*

$$\dim C_V(x) \leq \sum_{i=1}^k \dim C_{V_i}(x).$$

Proof. The proof follows from an elementary induction argument on the number of composition factors of V . \square

Proposition 3.3 ([29, Lemma 3.7]). *Let V_1 and V_2 be vector spaces over \mathbb{F}_r of dimension d_1 and d_2 respectively. Then if $g = g_1 \otimes g_2 \in \mathrm{GL}(V_1) \otimes \mathrm{GL}(V_2)$ is an element of projective prime order that acts on $V = V_1 \otimes V_2$, then*

$$\mathcal{E}_{\max}^V(g) \leq \min \left\{ d_2 \mathcal{E}_{\max}^{\overline{V_1}}(g_1), d_1 \mathcal{E}_{\max}^{\overline{V_2}}(g_2) \right\}.$$

where $\overline{V_i} = V_i \otimes \overline{\mathbb{F}_r}$.

Proof. This follows from [29, Lemma 3.7] with the observation that any element of projective prime order is of the form λx for $\lambda \in \overline{\mathbb{F}_r}$ and x an element of prime order in $\mathrm{GL}(V_1) \otimes \mathrm{GL}(V_2)$. \square

Given a family of almost quasisimple groups of Lie type G , our proof of Theorem 1.1 proceeds as follows. We first determine an upper bound on $\alpha(G)$ and a lower bound on $d_1(G)$ for groups G in the family, and note that from Proposition 3.1v), if G has no regular orbit on $V = V_d(r)$, then

$$r^d - 2 \sum_{x \in \mathcal{P}} |x^H|_{r^{\lfloor (1-1/\alpha(x))d \rfloor}} \leq 0.$$

The left hand side is increasing with respect to $\dim V$, so by substituting in $d_1(G)$ and $\alpha(G)$, we may easily prove that most groups in the family satisfy Theorem 1.1, leaving us with a finite collection of groups G and irreducible modules $V = V_d(r)$ to consider. We consider these cases individually. We start by checking whether $|G| > |V|$, since this implies the non-existence of a regular orbit. If this is the case, we can usually determine the base size either computationally in GAP [10] or MAGMA [2], or by using the following lemma.

Lemma 3.4. *Suppose V is a d -dimensional vector space over \mathbb{F}_r , and let $G \leq \mathrm{GL}(V)$. If G has a base of size c on $V \otimes \mathbb{F}_{r^i}$, then G has a base of size at most ci on V .*

Proof. Let $\{v_1, \dots, v_c\}$ be a base for G acting on $V \otimes \mathbb{F}_{r^i}$. Write each v_j as $v_j = \sum_{k=1}^i \eta_k w_{j,k}$, where $w_{j,k} \in V$ and η_1, \dots, η_i is a basis for \mathbb{F}_{r^i} over \mathbb{F}_r . Then $\{w_{j,k} \mid 1 \leq j \leq c, 1 \leq k \leq i\}$ is a base for G , since any non-trivial $g \in G$ stabilising the set must also stabilise $\{v_1, \dots, v_c\}$. \square

If instead $|G| < |V|$, we aim to apply Proposition 3.1 to prove the existence of a regular orbit. In order to apply Proposition 3.1, we usually need to compute the sizes of conjugacy classes of elements of prime order in $G/Z(G)$, and the eigenspace dimensions of their preimages in G on V . We determine conjugacy class orders using GAP [10] or [3]. Eigenspace dimensions of projective prime order elements are determined (or at least bounded from above) using one of a number of methods. We may use Proposition 3.1i) along with results from Section 2.2, or refine upper bounds of $\alpha(x)$ for $x \in G$ in GAP. If the Brauer character of V is available in [25], we may use it to determine eigenspace dimensions for semisimple elements. We may also seek to restrict V to a subgroup of G and apply Proposition 3.2. If we are not able to use these methods to successfully complete a proof by contradiction with an application of Proposition 3.1, we resort to computational methods. We use the generators of some irreducible modules in the online ATLAS [1] and construct them in GAP, or compute generators explicitly in MAGMA [2]. Once we have constructed V , we then either determine eigenspace dimensions of projective prime order elements to enable us to apply Proposition 3.1, or determine computationally whether G has a regular orbit on V . We are grateful to Eamonn O'Brien and Jürgen Müller for their assistance with certain computationally intensive calculations for groups G with $E(G)/Z(E(G))$ equal to one of: $L_2(q)$ with $11 \leq q \leq 81$, $L_3(4)$, $L_4(3)$, $U_3(3)$, $U_3(7)$, $U_4(3)$, $U_5(2)$, $U_5(3)$, $U_6(2)$, $\mathrm{PSp}_4(9)$, $\mathrm{PSp}_6(2)$, $\mathrm{PSp}_6(3)$, $\mathrm{P}\Omega_8^+(2)$, $G_2(4)$ or ${}^3D_4(2)$.

4. PROOF OF THEOREM 1.1: LINEAR GROUPS

In this section, we set out to prove the following proposition.

Theorem 4.1. *Suppose G is an almost quasisimple group with $E(G)/Z(E(G)) \cong \mathrm{PSL}_n(q)$, with $(n, q) \neq (2, 4), (2, 5), (2, 9), (4, 2)$. Let $V = V_d(r)$, $(r, q) = 1$ be a module for G , with absolutely irreducible restriction to $E(G)$. Also suppose that $(r, |G|) > 1$. Then either:*

- i) $b(G) = \lceil \log |G| / \log |V| \rceil$, or*
- ii) $b(G) = \lceil \log |G| / \log |V| \rceil + 1$ and (G, V) appear in Table 1.1.*

The excluded values of (n, q) in Theorem 4.1 are due to exceptional isomorphisms with alternating groups, which have already been treated in [9].

Let G be an almost quasisimple group with $E(G)/Z(E(G)) = \mathrm{PSL}_n(q)$.

Proposition 4.2. *Theorem 4.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \mathrm{PSL}_2(q)$.*

Proof. As above we assume $q > 7$ and $q \neq 9$. In addition to cross-characteristic representations, we must also consider representations of $L_2(8)$ in characteristic 2, due to the exceptional isomorphism $L_2(8) \cong {}^2G_2(3)'$. Now [21, Table 2] provides a list of cross-characteristic representations of $E(G)$. The number of involutions in $G/Z(G)$ is at most $2(q^2 + q)$ by Proposition 2.7, and the number of field automorphisms of order 2 is at most $q^{1/2}(q^{1/2} + 1)(q + 1)$ by [3, Table B.3]. If G has no regular orbit on $V = V_d(r)$ then by Propositions 2.3 and 3.1v),

$$r^d \leq 4(q^2 + q)r^{\lfloor 2d/3 \rfloor} + 2q^{1/2}(q^{1/2} + 1)(q + 1)r^{\lfloor 3d/4 \rfloor} + 2|\mathrm{P}\Gamma\mathrm{L}(2, q)|r^{\lfloor d/2 \rfloor}$$

since the number of odd prime order elements in $G/Z(G)$ is bounded above by $|\mathrm{P}\Gamma\mathrm{L}(2, q)|$. By Proposition 2.1, $d_1(G) = (q - 1)/(2, q - 1)$ and when we substitute this in to the equation, we see that this gives a contradiction for all q except $q \leq 113$ for q odd and $q \leq 32$ for q even. If $83 \leq q \leq 113$, then q is prime, we only need to consider $r = 2$, and $d = (q \pm 1)/2$. Let $i_P(H)$ denote the number of prime order elements in a finite group H . If G has no regular orbit on V then

$$r^d \leq i_2(\mathrm{Aut}(L_2(q)))r^{\lfloor 2d/3 \rfloor} + 2i_P(\mathrm{Aut}(L_2(q)))r^{\lfloor d/2 \rfloor}, \quad (4)$$

where we calculate $i_2(G)$ and $2i_P(\mathrm{Aut}(L_2(q)))$ in GAP. This is a contradiction for q with $83 \leq q \leq 113$. We further use this equation (with the exponent $\lfloor 2d/3 \rfloor$ replaced with $\lfloor 3d/4 \rfloor$ if q is a square to account for involutory field automorphisms) to determine the values of r we need to consider for the remaining values of q . We further note that the characters corresponding to the representations of dimension $(q \pm 1)/2$ for q odd contain b_q irrationalities. The remaining cross-characteristic cases to consider are summarised in Table 4.1. We also apply (4) to show that if $E(G)/Z(E(G)) \cong L_2(8)$ and V is an absolutely irreducible $\mathbb{F}_{2^k}E(G)$ -module for some k , then G has a regular orbit on V unless possibly $d = 2$ and $r = 8, 84, 512$, or (d, r) is one of $(4, 8)$ or $(8, 2)$.

We obtain the set of absolutely irreducible modules to consider from [21] and also the relevant Brauer character tables. We do not include the modules that fail (4). Apart from some cases addressed in Table 4.2, which were determined using constructions in MAGMA, all of the remaining cases are included in Table 4.3. The first column of Table 4.3 gives the layer $E(G)$ of G . The second column gives a list of primes (apart from 2 and 3) which divide the order of $E(G)$, given in ascending order. The third column gives the dimension d and the field size r of the irreducible module $V = V_d(r)$ being considered. The remaining columns give the eigenspace dimensions of elements of projective prime order on V . The eigenspace dimensions for the semisimple elements (i.e., those elements whose order is coprime to r) are sourced from the corresponding Brauer character, found either in [25] or constructed in GAP. We write the eigenspace dimensions using index notation so that, for example, we denote a set of eigenspaces with dimensions 16, 12, 12 as $(16, 12^2)$. The eigenspace bounds for elements of prime order dividing r are derived from Lemma 2.3 or from an explicit construction of the module in GAP. We then apply Proposition 3.1ii) to each module in Table 4.3. For example, suppose $E(G) = L_2(81)$

Upper bound for r			
q	$d = (q - 1)/2$	$d = (q + 1)/2$	$d \geq q - 1$
11	17	17	4
13	19	8	4
17	9	7	3
19	9	5	3
23	5	3	2
25	11	3	3
27	4	—	2
29	4	3	2
31	4	3	2
37	3	3	—
41	3	3	—
43	3	—	—
47	2	—	—
49	4	3	2
81	2	—	—

Upper bound for r			
q	$d = q - 1$	$d = q$	$d = q + 1$
8	5	5	3
16	3	3	3

TABLE 4.1. Remaining cases for $L_2(q)$ in cross-characteristic.

and $V = V_{40}(2)$, and let $H = G/Z(G)$. If G has no regular orbit on V , then by Proposition 3.1ii),

$$\begin{aligned}
r^{40} &\leq (i_2(H) - |2C|)r^{20} + |2C|r^{30} + \frac{1}{2}i_3(H)(r^{16} + 2r^{12}) + \frac{1}{4}i_5(H)(5r^8) + \frac{1}{40}i_{41}(H)(40r) \\
&\leq (7299 - 3240)r^{20} + 3240r^{30} + \frac{1}{2}6560(r^{16} + 2r^{12}) + \frac{1}{4}13284(5r^8) + \frac{1}{40}129600(40r)
\end{aligned}$$

This is a contradiction for $r = 2$, so G has a regular orbit on V .

Applying Proposition 3.1ii) to each module in Table 4.3, we see that in all cases, G has a regular orbit on V . The asterisked cases in Table 4.2 require some extra explanation. If $E(G)/Z(E(G)) \cong L_2(13)$ and $V = V_6(4)$, then G has a regular orbit on V unless $G = \mathbb{F}_4^\times \circ L_2(13)$ where $b(G) = 2$. If instead $E(G)/Z(E(G)) \cong L_2(8)$ and $V = V_7(3)$ or $V_4(8)$, then $b(G) = 2$ unless $G = L_2(8)$, where $b(G) = 1$. Finally, if $V = V_2(64)$, so $G = c \times L_2(8)$ for some $c \mid 63$, then $\lceil \log |G| / \log |V| \rceil > 1$ if and only if $c \geq 9$, and moreover $b(G) = 2$ if $3 \mid c$ and $b(G) = 1$ otherwise.

$E(G)/Z(E(G))$	(d, r)	$\lceil \frac{\log G }{\log V } \rceil$	$b(G)$	$E(G)/Z(E(G))$	(d, r)	$\lceil \frac{\log G }{\log V } \rceil$	$b(G)$
$L_2(31)$	(15, 2)	1	2	$L_2(11)$	(10, 3)	1	1
$L_2(25)$	(13, 3)	1	1		(10, 2)	1	2
	(12, 2)	2	2		(6, 3)	1	2
$L_2(23)$	(11, 3)	1	1		(5, 5)	1	2
	(12, 3)	1	1		(5, 4)	1	2
	(11, 2)	2	2		(5, 3)	2	2
$L_2(17)$	(16, 2)	1	1		(6, 5)	1	1
	(8, 2)	2	2	$L_2(8)$	(8, 2)	2	2
$L_2(13)$	(7, 3)	1	2		(7, 3)	1	2*
	$(6, 3^k), k \geq 2$	1	1		(4, 8)	1	2*
	(6, 3)	2	2		(2, 512)	1	1
	(14, 2)	1	2		(2, 64)	1	2*
	(6, 16)	1	1		(2, 8)	2	2
	(6, 4)	1	2*				

TABLE 4.2. Base sizes for some absolutely irreducible modules for $E(G)/Z(E(G)) \cong L_2(q)$.

$E(G)$	(p_3, p_4, p_5)	(d, r)	Projective prime order				
			2	3	p_3	p_4	p_5
$L_2(81)$	$(5, 41, -)$	$(40, 2)$	30 (2C), 20 (o/w)	$(16, 12^2)$	(8^5)	(1^{40})	–
$L_2(49)$	$(5, 7, -)$	$(48, 2)$	36	(16^3)	$(10^4, 8)$	$(7^6, 6)$	–
		$(48, 2)$	36	(16^3)	$(10^3, 9^2)$	$(7^6, 6)$	–
		$(25, 3)$	$(16, 9)$	12	(5^5)	$(7, 6^6)$	–
		$(24, r), r = 2, 4$	12	(8^3)	$(5^4, 4)$	$(6, 3^6)$	–
$2.L_2(49)$		$(24, 3)$	(12^2)	12	$(5^4, 4)$	$(6, 3^6)$	–
$L_2(47)$	$(23, 47, -)$	$(23, 2)$	12	$(8^2, 7)$	(1^{23})	(1^{23})	–
$L_2(43)$	$(7, 11, 43)$	$(21, 3)$	$(11, 10)$	10	(3^7)	$(2^{10}, 1)$	(1^{21})
$L_2(41)$	$(5, 7, 41)$	$(21, 3)$	$(11, 10)$	10	$(5, 4^4)$	(3^7)	(1^{21})
		$(20, 2)$	10	7	4	3	1
$2.L_2(41)$		$(20, 3)$	(10^2)	10	(4^5)	$(3^6, 2)$	(1^{20})
$L_2(37)$	$(19, 37, -)$	$(19, 3)$	$(10, 9)$	9	(1^{19})	(1^{19})	–
$2.L_2(37)$		$(18, 3)$	(9^2)	9	(1^{18})	(1^{18})	–
$L_2(31)$	$(5, 31, -)$	$(32, 2)$	21	$(11^2, 10)$	$(7^2, 6^3)$	$(2, 1^{30})$	–
		$(15, 4)$	8	(5^3)	(3^5)	(1^{15})	–
$2.L_2(31)$		$(15, 3)$	$(8, 7)$	7	(3^5)	(1^{15})	–
$L_2(29)$	$(5, 7, 29)$	$(28, 2)$	18	$(10, 9^2)$	$(6^4, 4)$	(7^4)	(1^{28})
		$(15, 3)$	$(8, 7)$	7	(3^5)	$(3, 2^6)$	(1^{15})
		$(14, 4)$	7	$(5^2, 4)$	$(3^4, 2)$	(2^7)	(1^{14})
$2.L_2(29)$		$(14, 3)$	(7^2)	7	$(3^4, 2)$	(2^7)	(1^{14})
$L_2(27)$	$(7, 13, -)$	$(13, 4)$	7	$(6, 4, 3)$	$(2^6, 1)$	(1^{13})	–
$L_2(25)$	$(5, 13, -)$	$(12, r), r = 4, 8$	8 (2D) 6 (o/w)	(4^3)	$(4, 2^4)$	(1^{12})	–
		$(12, 3) (\chi_{12})$	(6^2)	4	(3^4)	(1^{12})	–
		$(12, 9) (\chi_{12})$	(6^2)	4	(3^4)	(1^{12})	–
$L_2(23)$	$(11, 23, -)$	$(22, 2)$	12	$(8, 7^2)$	(2^{11})	(1^{22})	–
		$(11, 4)$	6	$(4^2, 3)$	(1^{11})	(1^{11})	–
$L_2(19)$	$(5, 19, -)$	$(20, 2)$	11	$(8, 6^2)$	(4^5)	$(2, 1^{17})$	–
		$(19, 3)$	12	9	9	9	–
		$(9, 9)$	$(5, 4)$	4	$(2^4, 1)$	(1^9)	–
		$(9, 4)$	5	(3^3)	$(2^4, 1)$	(1^9)	–
		$(9, 5)$	$(5, 4)$	(3^3)	4	(1^9)	–
		$(10, 5)$	(5^2)	$(4, 3^2)$	5	(1^{10})	–
$2.L_2(19)$		$(10, 9)$	(5^2)	5	(2^5)	(1^{10})	–
	$L_2(17)$	$(17, -, -)$	$(18, 3)$	$(10, 8)$	6	$(2, 1^{16})$	–
			$(16, 3)$	$(9, 7)$	6	(1^{16})	–
			$(9, 9)$	$(5, 4)$	3	(1^9)	–
$(8, 4)$			4	$(3^2, 2)$	(1^8)	–	
$2.L_2(17)$		$(8, 9)$	(4^2)	3	(1^8)	–	
$L_2(16)$	$(5, 17, -)$	$(16, 3)$	$(10, 6)$	8	$(4, 3^4)$	(1^{16})	–
$L_2(13)$		$(13, 3)$	$(7, 6)$	5	$(2^6, 1)$	(1^{13})	–
$2.L_2(13)$	$(7, 13, -)$	$(14, 3)$	(7^2)	7	(2^7)	$(2, 1^{12})$	–
$L_2(11)$	$(5, 11, -)$	$(12, 4)$	8	(4^3)	$(3^2, 2^3)$	$(2, 1^{10})$	–
		$(5, 9)$	$(3, 3)$	2	$(2, 1^4)$	(1^5)	–
		$(5, 16)$	3	$(2^2, 1)$	(1^5)	(1^5)	–
		$(10, 2)$	6 (2A), 5 (2B)	$(4, 3^2)$	(2^5)	(1^{10})	–
		$(10, 3)$	(5^2)	5	(2^5)	(1^{10})	–
		$(6, 9)$	$(3, 3)$	2	(1^6)	(1^6)	–

TABLE 4.3. Eigenspaces of projectively prime order elements on absolutely irreducible $V = V_d(r)$ for $E(G)/Z(E(G)) = \text{PSL}_2(q)$.

□

Proposition 4.3. *Theorem 4.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \text{PSL}_3(q)$.*

Proof. Suppose $n = 3$ and $q \neq 2, 4$. Then by Proposition 2.4, $\alpha(x) \leq 3$ unless x is a graph-field involutory automorphism where $\alpha(x) \leq 4$. Therefore, if G has no regular orbit on an absolutely irreducible $V = V_d(r)$, then by Proposition 3.1v) and Lemma 2.8i),

$$r^d \leq 2|\text{P}\Gamma\text{L}_3(q)|r^{\lfloor 2d/3 \rfloor} + 4q^4r^{\lfloor 3d/4 \rfloor},$$

since the number of non-graph prime order elements in $G/Z(G)$ is bounded above by $|\text{P}\Gamma\text{L}_3(q)|$. If we substitute in $d_1(G)$ from Proposition 2.1 (as described in section 3, then we obtain a contradiction except for when $(q, r) = (7, 2)$, $q = 5$ and $r \leq 4$, and $q = 3$ with $r \leq 11$, so G has a regular orbit on all choices of V in all other cases.

We now examine each of the remaining cases, along with the cases with $q = 2, 4$ in detail.

Suppose that $E(G)/Z(E(G)) \cong \text{PSL}_3(7)$ and $r = 2$. From [21], the smallest non-trivial representation of G when $r = 2$ has degree 56. The number of prime order elements in $\text{Aut}(\text{PSL}_3(7))$ is 853481. Therefore, by Proposition 3.1v), if G has no regular orbit on V , then

$$2^d \leq 2 \times 853481 \times 2^{\lfloor 2d/3 \rfloor}$$

This gives a contradiction for all V except for $V = V_{56}(2)$. Therefore, G has a regular orbit on V except perhaps when $d = 56$. In the latter case, examining the corresponding Brauer character in [25], we determine that no element of G of odd prime order has eigenspace of dimension greater than 24, i.e., $\mathcal{E}_{\max}(g_{2'}) \leq 24$. We also compute that $i_2(\text{Aut}(\text{PSL}_3(7))) = 19551$. So if G has no regular orbit on $V = V_{56}(2)$, then by Proposition 3.1iv),

$$2^{56} \leq 2 \times 853481 \times 2^{24} + 19551 \times 2^{\lfloor 2 \times 56/3 \rfloor}$$

This is a contradiction, so G has a regular orbit on V .

Now let $E(G)/Z(E(G)) \cong \text{PSL}_3(5)$ and $r \leq 4$. Since G has no graph-field automorphisms, $\alpha(x) \leq 3$ for all projective prime order $x \in G$. We also compute that the number of prime order elements in $G/Z(G)$ is at most 154999. If G has no regular orbit on $V = V_d(r)$, then $r^d \leq 2 \times 154999r^{\lfloor 2d/3 \rfloor}$ which implies that $d \leq 54, 33, 27$ for $r = 2, 3$ and 4 respectively. From [21] we deduce that we only need to examine one representation of degree 30 for $r = 2$, while if $r = 3$, then there is one representation of degree 30 and another of degree 31 to consider. First suppose $r = 3$. The Brauer character for the 30-dimensional representation over $L_3(5).2$ contains $r_5 \notin \mathbb{F}_3$ so we only need to consider this representation for $L_3(5)$. There are at most 15500 elements of order 3 in $G/Z(G)$, and at most 3875 involutions. Examining the Brauer characters of the 30- and 31- dimensional representations, we find that $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 10$ and 11 respectively. In addition, $\mathcal{E}_{\max}(g_2) \leq 18$ and 19 respectively. Therefore, if G has no regular orbit on the 30-dimensional module, then by Proposition 3.1iv),

$$r^{30} \leq 2 \times 154999r^{10} + 2 \times 3875r^{18} + 15500r^{\lfloor 2d/3 \rfloor}$$

which gives a contradiction for $r = 3$. A similar inequality gives the result for $d = 31$.

Now suppose that $r = 2$ and $V = V_{30}(2)$ is an absolutely irreducible $\mathbb{F}_2E(G)$ -module. We determine in MAGMA that $\mathcal{E}_{\max}(g_2) \leq 18$. Examining the Brauer character in [25], we see that elements in classes 3A and 5A have eigenspaces of dimension at most 10, and no other element of prime order has an eigenspace of dimension greater than 6. Therefore, if G has no regular orbit on V , by Proposition 3.1iv),

$$r^{30} \leq 2 \times 154999r^6 + 2(|3A| + |5A|)r^{10} + 3875r^{18}$$

this is a contradiction for $r = 2$, so G has a regular orbit on V .

Suppose that $E(G)/Z(E(G)) \cong \text{PSL}_3(4)$. By [3, Table B.3], the number of graph-field involutory automorphisms in $L_3(4).D_{12}$ is $|\text{PGL}(3, 4)|/|\text{PGU}(3, 2)| = 280$, and we compute in GAP that the number of prime order elements is 20619. Therefore, if G has no regular orbit on the irreducible module V then by Proposition 3.1v),

$$r^d \leq 2 \times 20619r^{\lfloor 2d/3 \rfloor} + 2 \times 280r^{\lfloor 3d/4 \rfloor}$$

Therefore, either G has a regular orbit on V , or V appears in Table 4.4. We will investigate the remaining modules in decreasing order of dimension. We summarise our analysis across Tables 4.5 and 4.6. If $V = V_d(r)$ is an absolutely irreducible module for G , then in Table 4.5, the eigenspaces for

d	$E(G)$	Characteristic	Irrationalities	Notes
4	$4_2.L_3(4)$	3	i_1, r_7	$r = 9^k$
6	$6.L_3(4)$	$\neq 2, 3$	z_3	$r = 25^k$ or 7^k
6	$2.L_3(4)$	3		$r = 3^k$
8	$4_1.L_3(4)$	$\neq 2, 5$	i_1, b_5	$r = 9$
8	$4_1.L_3(4)$	5	i_1	$r = 5, 25$
10	$2.L_3(4)$	$\neq 2, 7$	b_7	$r = 9$
10	$2.L_3(4)$	7		$r = 7$
15	$L_3(4)$	3		$r = 3$
15	$3.L_3(4)$	$\neq 2, 3$	z_3	$r = 7$
19	$L_3(4)$	3,7		$r = 3$

TABLE 4.4. List of remaining cases for $E(G)/Z(E(G)) \cong \text{PSL}_3(4)$.

$E(G)$	(d, r)	2A	3A	5A	5B	7A	7B	2B	3B	3C	2C	2D
$L_3(4)$	(19, 3)	(11, 8)	7	$(4^4, 3)$	$(4^4, 3)$	$(3^6, 1)$	$(3^6, 1)$	(10, 9)	12	9	(12, 7)	(10, 9)
$3.L_3(4)$	(15, 7)	(8, 7)	(5^3)	(3^5)	(3^5)	10	10	(9, 6)	–	–	–	–
$L_3(4)$	(15, 3)	(8, 7)	5	(3^5)	(3^5)	$(3, 2^6)$	$(3, 2^6)$	(9, 6)	–	–	(9, 6)	(9, 6)
$2.L_3(4)$	(10, 7)	(6, 4)	$(4, 3^2)$	(2^5)	(2^5)	2	2	(5^2)	–	–	(7, 3)	(6, 4)
	(10, 9)	(6, 4)	4	(2^5)	(2^5)	$(2^3, 1^4)$	$(2^3, 1^4)$	–	–	–	(7, 3)	–
$4_1.L_3(4)$	(8, 9)	(4^2)	3	$(2^3, 1^2)$	$(2^3, 1^2)$	$(2, 1^6)$	$(2, 1^6)$	–	–	–	–	(5, 3)
	(8, 25)	(4^2)	$(3^2, 2)$	5	5	$(2, 1^6)$	$(2, 1^6)$	–	–	–	–	(5, 3)
$2.L_3(4)$	$(6, 3^k)$	(4, 2)	2	$(2, 1^4)$	$(2, 1^4)$	(1^6)	(1^6)	(3^2)	–	–	(3^2)	(3^2)
$6.L_3(4)$	$(6, 5^k)$	(4, 2)	(3^2)	2	2	(1^6)	(1^6)	(3^2)	–	–	–	–
$6.L_3(4)$	$(6, 7^k)$	(4, 2)	(3^2)	$(2, 1^4)$	$(2, 1^4)$	1	1	(3^2)	–	–	–	–
$4_2.L_3(4)$	$(4, 9^k)$	(2^2)	2	(1^4)	(1^4)	(1^4)	(1^4)	–	–	–	(3, 1)	–
# elements		315	2240	4032	4032	2880	2880	280	672	1920	360	1008

TABLE 4.5. Eigenspace dimensions of projectively prime order elements in G with $E(G)/Z(E(G)) \cong \text{PSL}_3(4)$.

elements of projective prime order coprime to r are derived from the character table, while those of projective prime order dividing r are derived from Proposition 3.1i), except if they are italicised, in which case the eigenspaces are computed from a construction in GAP. Using Proposition 3.1 ii), we see that if V lies in Table 4.4, either V also appears in Table 4.6, where the base size $b(G)$ is determined computationally from a construction in GAP, or G appears in Table 4.5 and G has a regular orbit on V by Proposition 3.1ii).

G	(d, r)	$\lceil \log G / \log V \rceil$	$b(G)$
$2 \times L_3(4).2^2$	(15, 3)	1	1
$4_1.L_3(4)$	(8, 5)	1	1
$4_1.L_3(4).2_3$	(8, 5)	1	2
$2.L_3(4) \leq G \leq (F_r^\times \circ 2).L_3(4).2^2$	(6, 3)	2	3
	(6, 9)	1	2
$6.L_3(4) \leq G \leq 6.L_3(4).2^2$	(6, 7)	2	2
$4_2.L_3(4) \leq G \leq (\mathbb{F}_9^\times \circ 4_2).L_3(4).2_2$	(4, 9)	2	2
$4_2.L_3(4) \leq G \leq 20 \times (\mathbb{F}_{81}^\times \circ 4_2).L_3(4).2_2$	(4, 81)	1	1

TABLE 4.6. Some base size results for groups G with $E(G)/Z(E(G)) \cong \text{PSL}_3(4)$.

Now let $E(G)/Z(E(G)) \cong \text{PSL}_3(3)$ and $r \in \{2, 4, 8\}$. We compute that the number of prime order elements in $\text{Aut}(\text{PSL}_3(3))$ is 2807, and from Proposition 2.1, $d_1(G/Z(G)) = 12$. Therefore, if G has no regular orbit on V then by Proposition 3.1v) $r^d \leq 2 \times 2807r^{\lfloor 2d/3 \rfloor}$. Substituting in the degrees of the irreducible representations of G from [25], we see that G has a regular orbit on all V except possibly when $d \leq 12, 18, 36$ for $r = 8, 4$ and 2 respectively. From [21] we deduce that the modules we need to

consider are of dimensions 12 and 26. If $d = 26$ and $r = 2$, then constructing the representation in GAP, we find that there is a regular orbit. We may also construct the 12-dimensional representation in GAP. Note that G has no regular orbit on the 12-dimensional representation when $r = 2$, since $|V| < |L_3(3)|$. When $r = 2$, we find that there is a base of size two and when $r = 4, 8$, there is a regular orbit.

Finally, let $E(G)/Z(E(G)) \cong \text{PSL}_3(2)$. Since $L_3(2) \cong L_2(7)$, we will consider representations in all characteristics 2,3,7. By Proposition 2.1, we have $d_1(G/Z(G)) \geq 2$, $\alpha(x) \leq 3$ for non-trivial $x \in G/Z(G)$ by Proposition 2.4, and we compute that there are 153 prime order elements in $\text{Aut}(\text{PSL}_3(2))$. If G has no regular orbit on $V = V_d(r)$ then by Proposition 3.1v),

$$r^d \leq 2 \times 153r^{\lfloor 2d/3 \rfloor}$$

Substituting in $d = 2$ to find an upper bound on r to consider for each representation, we see that G has a regular orbit on V unless possibly when $r \leq 306$.

We construct all of the modules V that are left to consider in GAP and determine their base sizes computationally. We summarise our findings in Table 4.7. The first column of Table 4.7 gives the layer $E(G)$ and the second column gives the dimension of the module V being considered. The third column gives information about the finite fields where V exists. Here

$$\nu = \begin{cases} 1 & \text{if } G/Z(G) \text{ is simple,} \\ 2 & \text{otherwise.} \end{cases}$$

The remaining columns give the base sizes of groups G with the specified $E(G)$ on $V = V_d(r)$. If the entry is asterisked, then G has a regular orbit on $V = V_d(r)$ if $G = L_3(2)$, and $b(G) = 2$ otherwise. The entry marked with a dagger represents that G has a regular orbit on $V = V_2(49)$ if $Z(G)$ has order 2 or 6, and $b(G) = 2$ otherwise. There are some cases included in the table where $b(G) = \lceil \log |G| / \log |V| \rceil + 1$, namely $G = L_3(2), L_3(2).2$ or $2 \circ L_3(2)$ and $V = V_3(7)$; $G = L_3(2)$ and $V = V_8(2)$; G quasisimple with $|Z(G)| = 4, 8, 12$, or $G/Z(G) = L_3(2).2$, $|Z(G)| = 4$, and $V = V_2(49)$. All of these are included in Table 1.1.

$E(G)$	d	Field	$b(G) = 1$	$b(G) = 2$	$b(G) = 3$	
$L_3(2)$	3	$r = 2^k$	$r = 8^*, r \geq 16$	$r = 4$	$r = 2$	
		$r = 9^k$	$r \geq 9^*$			
		$r = 7^k$	$r \geq 49$	$r = 7$		
	5	$r = 7^k$	$r \geq 7$			
	6	$r = 3^k$	$r \geq 9$	$r = 3$		
	7	$r = 3^k$	$r \geq 3$			
		$r = 7^k$	$r \geq 7$			
	8	$r = 2^k$	$r \geq 4$	$r = 2$		
	$2.L_3(2)$	2	$r = 7^{\nu k}$	$r = 49^\dagger$	$r = 7$	
			$r = 9^k$	$r \geq 9$		
4		$r = 7^{\nu k}$	$r \geq 7$			
6		$r = 9^{\nu k}$	$r \geq 9$			
		$r = 7^{\nu k}$	$r \geq 7$			

TABLE 4.7. Base sizes of groups G with $E(G)/Z(E(G)) \cong L_3(2)$ on irreducible V .

□

Proposition 4.4. *Theorem 4.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \text{PSL}_4(q)$.*

Proof. Suppose $n = 4$, so $q \geq 3$. By Proposition 2.4, $\alpha(x) \leq 4$ unless x is an involutory graph automorphism where $\alpha(x) \leq 6$, so if G has no regular orbit on $V = V_d(r)$ then by Proposition 3.1v) and Lemma 2.8i),

$$r^d \leq 2|\text{PTL}(4, q)|r^{\lfloor 3d/4 \rfloor} + 2q^{10}r^{\lfloor 5d/6 \rfloor}$$

Substituting in $d = d_1(G/Z(G)) = \frac{q^4 - q}{q - 1} - 1$ for $q > 3$ and $d = 26$ for $q = 3$, we see that G has a regular orbit on all irreducible V unless possibly when $q = 3$ and $r \leq 11$.

So suppose $(n, q) = (4, 3)$. We first determine which modules we must consider. We compute that the number of prime order elements in $\text{Aut}(L_4(3))$ is 2279303, and the number of involutions in $G/Z(G)$ is at most 27639. If G has no regular orbit on the d -dimensional module V , then by Proposition 3.1v) $r^d \leq 2 \times 2279303r^{\lfloor 3d/4 \rfloor} + 27639r^{\lfloor 5d/6 \rfloor}$ which implies that $d \leq 96, 48, 40, 30$ for $r = 2, 4, 5$ and 8 respectively. From [21], we deduce that the only modules left to consider are those in Table 4.8, and also $V = V_{26}(2)$ or $V_{38}(2)$, where a construction in MAGMA shows that there is no regular orbit of G on V if $d = 26$, but there is for $d = 38$. The bounds in Table 4.8 were obtained from the Brauer characters, constructing the modules in GAP and Proposition 3.1i). The asterisked entry has a bound of 31 for involutions in classes 2D and 2E, and a bound of 25 otherwise. Therefore, applying Proposition 3.1ii), we find that G has a regular orbit on V for all cases in Table 4.8.

$E(G)$	(d, r)	2A	2C	Other $o(x) = 2$	3A	3B	Other $o(x) = 3$	$o(x) = 5$	$o(x) = 13$
$2.L_4(3)$	$(40, 5)$	20	26	20	21	16	22	26	4
$L_4(3)$	$(38, 5)$	20	25	23	20	11	14	8	3
$L_4(3)$	$(26, 5)$	16	–	16	9	14	14	6	2
$L_4(3)$	$(26, 2^k), k \geq 2$	16	–	14	9	14	9	6	2
	$(38, 2^k), k \geq 2$	20	31	31*	20	11	14	8	3
#elements		2106	1080	24453	1040	3120	78000	303264	1866240

TABLE 4.8. Upper bounds on \mathcal{E}_{\max} for element of projective prime order in G where $E(G)/Z(E(G)) \cong \text{PSL}_4(3)$.

□

Proposition 4.5. *Theorem 4.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \text{PSL}_n(q)$ with $n \geq 5$.*

Proof. Suppose $n \geq 5$. We begin by applying Proposition 3.1v) with $\alpha(x) \leq n$ from Proposition 2.4 and $d_1(G) \geq \frac{q^n - q}{q - 1} - 1$ from Proposition 2.1. Therefore, if G has no regular orbit on V then

$$r^d \leq 2 \times |\text{P}\Gamma\text{L}_n(q).2|r^{\lfloor \frac{n-1}{n}d \rfloor}$$

We find that G has a regular orbit on V except possibly when (n, q, r) lies in Table 4.9.

n	q	r
8	2	3
7	2	≤ 5
6	2	≤ 11
5	3	2
5	2	≤ 17

TABLE 4.9. Remaining cases to consider when $n \geq 5$.

We consult [21] to construct a list of modules to check, also making use of Proposition 2.2 for $(n, q) = (8, 2)$. We summarise our analysis of these remaining cases in Table 4.10. The bold entries denote those upper bounds obtained from the corresponding Brauer characters, while the italicised entries are computed from an explicit construction in GAP. The remaining entries are obtained by applying Propositions 3.1i) and 2.4. Applying Proposition 3.1iii) to each module in Table 4.10 shows that G has a regular orbit on V in all cases. The only cases not addressed in Table 4.10, are where $G = \text{PSL}_5(2)$ or $\mathbb{F}_3^\times \circ \text{PSL}_5(2)$ and $V = V_{30}(3)$. These cases were investigated computationally using a construction in MAGMA and there is a regular orbit in both cases.

□

$E(G)/Z(E(G))$	V	$o(x) = 2$	$o(x) = 3$	$o(x) \mid r$	Remaining prime order elements
$\mathrm{PSL}_8(2)$	$V_{253}(3)$	221	221	221	61
$\mathrm{PSL}_7(2)$	$V_{125}(3)$	94	93	93	94
$\mathrm{PSL}_7(2)$	$V_{126}(5)$	94	94	108	94
$\mathrm{PSL}_6(2)$	$V_{61}(r), r = 3, 9$	46	29	29	13
$\mathrm{PSL}_6(2)$	$V_{61}(7)$	46	46	50	46
$\mathrm{PSL}_6(2)$	$V_{62}(5)$	46	46	51	46
$\mathrm{PSL}_5(3)$	$V_{120}(2)$	96	66	66	66
$\mathrm{PSL}_5(2)$	$V_{30}(5)$	22	14	6	6
$\mathrm{PSL}_5(2)$	$V_{30}(7)$	22	14	6	6
$\mathrm{PSL}_5(2)$	$V_{30}(9)$	22	14	14	6

TABLE 4.10. Upper bounds on \mathcal{E}_{\max} for elements of projective prime order elements in linear groups, $n \geq 5$.

5. PROOF OF THEOREM 1.1: UNITARY GROUPS

In this section we set out to prove the following result.

Theorem 5.1. *Suppose G is an almost quasisimple group with $E(G)/Z(E(G)) \cong \mathrm{PSU}_n(q)$, and with $n \geq 3$ and $(n, q) \neq (3, 2), (4, 2)$. Let $V = V_d(r)$, $(r, q) = 1$ be a module for G , with absolutely irreducible restriction to $E(G)$. Also suppose that $(r, |G|) > 1$. Then either:*

- i) $b(G) = \lceil \log |G| / \log |V| \rceil$, or*
- ii) $b(G) = \lceil \log |G| / \log |V| \rceil + 1$ and (G, V) lies in Table 1.1.*

We exclude the cases where $E(G)/Z(E(G)) \cong U_3(2)$ since it is soluble, and $U_4(2)$ since it is isomorphic to $\mathrm{PSP}_4(3)$ and is dealt with later in Proposition 6.17.

Proposition 5.2. *Theorem 5.1 holds for $n = 3$.*

Proof. We begin by determining the values of d, q, r we must consider. By Proposition 2.4, $\alpha(x) \leq 3$, unless $q = 3$ and x is an inner involution with $\alpha(x) = 4$. So by Proposition 3.1v), if G has no regular orbit on V , and $q \neq 3$, then

$$r^d \leq 2|\mathrm{PGU}_3(q)|r^{\lfloor 2d/3 \rfloor}.$$

Substituting in $d_1(G) = \lfloor (q^3 - 1)/(q + 1) \rfloor$, we see that this gives a contradiction for $q \geq 9$ and $r \geq 2$, so there is a regular orbit of G on all irreducible modules V when $q \geq 9$ by Proposition 3.1v). When we replace $|\mathrm{PGU}_3(q)|$ by the number of prime order elements in $\mathrm{PGU}_3(q)$, we also eliminate $q = 8, 9$, and the remaining cases to consider are given in Table 5.1, obtained by examining [21]. Note that for $q = 3$, we use $\alpha(x) \leq 4$, as in Proposition 2.4.

q	d	r	$\mathrm{char}(\mathbb{F}_r)$
7	42	2	2
5	20	≤ 4	2,3
	21	3	3
	28	≤ 3	2,3
4	12	≤ 13	3,5,13
3	3	$r \leq 729$	3
	6	≤ 64	2,3,7
	7	≤ 49	3,7
	14	≤ 8	2,7
	27	3	3
	32	2	2

TABLE 5.1. Remaining cases for $E(G)/Z(E(G)) \cong \mathrm{PSU}_3(q)$.

Suppose $E(G)/Z(E(G)) \cong \text{PSU}_3(7)$. From Table 5.1, we are left to consider one 42-dimensional representation V , and from the Brauer character in [25], $\mathcal{E}_{\max}(g_{2'}) \leq 14$ on the corresponding module. From a construction of V in GAP, we determine that involutions in classes 2A and 2B have eigenspaces of dimension 24 and 21 respectively. The number of prime order elements in $\text{Aut}(U_3(7))$ is 2098571. Therefore, if G has no regular orbit on V then by Proposition 3.1iv), $r^{42} \leq 2 \times 2098571r^{14} + |2A|r^{24} + |2B|r^{20}$, which gives a contradiction for $r = 2$.

Now let $E(G)/Z(E(G)) \cong \text{PSU}_3(5)$. The number of prime order elements in $\text{Aut}(U_3(5))$ is 71849, and we proceed by examining each of the relevant cases in Table 5.1. First suppose $r = 3$ and $d = 28$. There are at most 16550 elements of order 3 in $G/Z(G)$, and we infer from the Brauer character in [25] that $\mathcal{E}_{\max}(g_{3'}) \leq 16$ on both of the 28-dimensional modules. Therefore, if G has no regular orbit on V ,

$$r^{28} \leq (71849 - 16550)(r^{16} + r^{12}) + 16550r^{\lfloor 2 \times 16/3 \rfloor}$$

which is a contradiction for $r = 3$. If instead $V = V_{20}(3)$, then we compute that $\mathcal{E}_{\max}(g_2) \leq 12$ and $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 5$. The number of involutions in $\text{Aut}(U_3(5))$ is 3674. In addition, in GAP we compute that elements x in classes 3A or 3C have $\alpha(x) = 2$. Therefore, if G has no regular orbit on V ,

$$r^{20} \leq 2 \times 71849r^5 + 3674(r^{12} + r^8) + (|3A| + |3C|)r^{10} + |3B|r^{13},$$

which gives a contradiction for $r = 3$. Now suppose V is a 21-dimensional Weil module. Again, we find from the Brauer character in [25] that $\mathcal{E}_{\max}(g_2) \leq 13$ and $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 5$. Therefore, if G has no regular orbit on V ,

$$r^{21} \leq 2 \times 71849r^5 + 3674(r^{13} + r^8) + (|3A| + |3C|)r^{10} + |3B|r^{14}$$

This also gives a contradiction for $r = 3$. Now suppose $r = 2$ or 4. From Table 5.1, $d = 20$ or 28. Let $V = V_{20}(r)$. From the Brauer character we compute that $\mathcal{E}_{\max}(g_3) \leq 8$ and $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 5$. Therefore, if G has no regular orbit on V ,

$$r^{20} \leq 2 \times (71849 - 3674 - 16550)r^5 + 2 \times 16550r^8 + 3674r^{\lfloor 2 \times 20/3 \rfloor}$$

This is a contradiction for $r = 4$. When $r = 2$, we find in GAP that there is a single regular orbit under $U_3(5)$ and there is no regular orbit for any G with $U_3(5) < G \leq U_3(5).S_3$ and $b(G) = 2$ here by Lemma 3.4. Now suppose $V = V_{28}(2)$. In GAP we compute that $\mathcal{E}_{\max}(g_2) \leq 16$. Let $H = G/Z(G)$. We deduce bounds on eigenspace dimensions from the Brauer character and find that if G has no regular orbit on V ,

$$r^{28} \leq 2i_7(H)r^4 + 2i_5(H)r^8 + 2|3A|r^{10} + i_2(H)r^{16}$$

and this is false for $r = 2$, so G has a regular orbit on V .

Suppose now that $q = 4$. From Table 5.1, we see that we only need to consider 12-dimensional modules for $r \leq 13$. When $r = 3$, the representation is only realised on $U_3(4)$ and $2 \circ U_3(4)$ because of an i irrationality in the corresponding character. Constructing the representation in GAP shows there is a regular orbit under $U_3(4)$, however, there is no regular orbit under $2 \circ U_3(4)$ and $b(G) = 2$ here. For $r = 5, 9, 13$, we summarise information about bounds on eigenspace dimensions of projective prime order elements in Table 5.2. Applying Proposition 3.1iv), we see that G has a regular orbit on V in all cases.

Bounds on eigenspace dimensions					
r	2A	2B	$o(x) = 3$	$o(x) = 5$	$o(x) = 13$
5	8	6	4	5	1
9	8	6	4	4	1
13	8	6	4	4	1

TABLE 5.2. Upper bounds on \mathcal{E}_{\max} for projective prime order elements acting on $V_{12}(r)$.

Finally, suppose $q = 3$. Since $U_3(3) \cong G_2(2)'$, we will consider representations in characteristics 2,3,7. The number of prime order elements in $U_3(3).2$ is 2771. We determine in GAP that if x is

of order 7 or in class 3A, then $\alpha(x) = 2$. Otherwise, $\alpha(x) \leq 3$ unless x is an inner involution and $\alpha(x) \leq 4$. Therefore, if G has no regular orbit on an absolutely irreducible $\mathbb{F}_r E(G)$ -module V ,

$$\begin{aligned} r^d &\leq 2(i_7(G) + |3A|)r^{\lfloor d/2 \rfloor} + 2(|3B| + |2B|)r^{\lfloor 2d/3 \rfloor} + 2|2A|r^{\lfloor 3d/4 \rfloor} \\ &\leq 2(1728 + 672)r^{\lfloor d/2 \rfloor} + 2(56 + 252)r^{\lfloor 2d/3 \rfloor} + 2 \times 63r^{\lfloor 3d/4 \rfloor} \end{aligned}$$

Comparing this to the list in Table 5.1, we only now need to consider $d = 3, 6, 7, 14$ for $r = 9, 81, 729$, $r \leq 16$, $r = 3, 7, 9$ and $r = 2, 4$ respectively. We resolve all of the remaining cases by constructing the modules in GAP and computing the base size. If $V = V_{14}(r)$ for $r = 2, 4$, we find there is no regular orbit on V when $r = 2$ and $b(G) = 2$ here. If $r = 4$, then there is a regular orbit. If $V = V_7(r)$ for $r = 3, 7, 9$, then there is a regular orbit for $r = 7, 9$. If $r = 3$ then $|V| < |G|$ and $b(G) = 2$ by Lemma 3.4. Now let $V = V_6(r)$ for $r \leq 16$. Note that in characteristic 3, $r = 9$ due to an i irrationality in the corresponding character. When $r = 2$, $|V|^2 < |G|$ so $b(G) \geq 3$, and we compute that indeed $b(G) = 3$. If $r = 4$, we have $|V| < |G|$ and $b(G) = 2$. If instead $r = 8, 9$ or 16 , we compute that G has a regular orbit on V . When $r = 7$, we compute that $\langle \rho \rangle.U_3(3)$, $\rho \in \mathbb{F}_7^\times$ has a regular orbit on V , except when $\langle \rho \rangle \cong \mathbb{F}_7^\times$. In this case, $b(G) = 2$ by Lemma 3.4. Moreover, $U_3(3).2$ and $3 \times U_3(3).2$ have a regular orbit on V , while $2 \times U_3(3).2$ and $6 \times U_3(3).2$ do not. Again, these have $b(G) = 2$ by Lemma 3.4. Finally, let $V = V_3(r)$ for $r = 9, 81, 729$. If $r = 9$ then $|V| < |G|$ and $b(G) = 2$ if $G = U_3(3)$, and $b(G) = 3$ if $c \circ U_3(3)$ with $c \in \{2, 4, 8\}$. If instead $r = 81$, then G has a regular orbit on V if $G = U_3(3)$ or $5 \circ U_3(3)$, and $b(G) = 2$ if $G = c \circ U_3(3)$ for even c dividing 80. Finally, if $r = 729$, then G always has a regular orbit on V . \square

Proposition 5.3. *Theorem 5.1 holds for $n = 4$.*

Proof. Suppose that G is an almost quasisimple group with $\text{soc}(G/Z(G)) \cong \text{PSU}_4(q)$. Applying Proposition 3.1 v) with $d_1(G) \geq (q^4 - 1)/(q + 1)$ from Proposition 2.1 and $\alpha(G) \leq 6$ from Proposition 2.4, we deduce that G has a regular orbit on all V if $q \geq 7$. We must further consider when $q \leq 5$.

Suppose $q = 5$. The number of prime order elements in $\text{Aut}(U_4(5))$ is 1572224649, while the number of involutions is 2391775. By Proposition 2.4, $\alpha(x) \leq 4$ for $x \in G/Z(G)$, unless x is an involutory graph automorphism with $\alpha(x) \leq 6$. Therefore, if G has no regular orbit on V ,

$$r^d \leq 2 \times 1572224649r^{\lfloor \frac{3d}{4} \rfloor} + 2 \times 2391775r^{\lfloor \frac{5d}{6} \rfloor},$$

so by examining [21] we see that G has a regular orbit on V unless possibly $r = 2$ and $V = V_{104}(2)$. We compute in GAP that this representation is a composition factor of the reduction modulo 2 of the 105-dimensional irreducible representation of G over \mathbb{C} , so we can deduce the Brauer character of V from the ordinary character. From the Brauer character we find that $\mathcal{E}_{\max}(g_{2'}) \leq 63$. We also find that the number of graph automorphisms in $G/Z(G)$ is at most 3150. Therefore, if G has no regular orbit on V then by Proposition 3.1iv),

$$r^{104} \leq 2 \times 1572224649r^{63} + 2391775r^{\lfloor \frac{3d}{4} \rfloor} + 3150r^{\lfloor \frac{5d}{6} \rfloor}$$

This is a contradiction, so G has a regular orbit on V .

Now suppose $q = 4$. The number of prime order elements in $\text{PSU}_4(4).4$ is 156437999, and 335855 of these are involutions. If G has no regular orbit on an irreducible module V ,

$$r^d \leq 2 \times 156437999r^{\lfloor \frac{3d}{4} \rfloor} + 2 \times 335855r^{\lfloor \frac{5d}{6} \rfloor}$$

This gives a contradiction for all r, d except $r = 3$ and $d \leq 72$. By [21], this leaves a 52-dimensional module V to consider. This module can be constructed by reducing a 52-dimensional module for G over \mathbb{C} , so we can deduce the Brauer character corresponding to V . The number of elements of order 3 in G is 1214720, and examining the Brauer character we see that $\mathcal{E}_{\max}(g_{3'}) \leq 28$, except if the element x is in class 2A, where $\dim C_V(x) \leq 32$. Therefore, if G has no regular orbit on V ,

$$r^{52} \leq 2 \times 156437999r^{28} + 2 \times |2A|r^{32} + 1214720r^{\lfloor \frac{3 \times 52}{4} \rfloor}$$

This is a contradiction for $r = 3$, so G has a regular orbit on V .

Now suppose $q = 3$. We summarise the numbers of prime order elements in $H = \text{PSU}_4(3).D_8$ computed using GAP in Table 5.3. Let G be an almost quasisimple group with $\text{soc}(G/Z(G)) \cong \text{PSU}_4(3)$. Then $d \geq 6$ by Proposition 2.1, and the number of prime order elements in $G/Z(G)$ is at

r_0	$i_{r_0}(H)$
2 (graph auts)	20664
2 (not graph auts)	7911
3	47600
5	653184
7	933120
Total	1662479

TABLE 5.3. Elements of prime order in $\text{PSU}_4(3).D_8$.

most 1662479, with at most of 20664 of these being graph automorphisms. Therefore, if G has no regular orbit on V ,

$$r^d \leq 2(1662479 - 20664)r^{\lfloor \frac{3d}{4} \rfloor} + 2 \times 20664r^{\lfloor \frac{5d}{6} \rfloor}$$

This gives a contradiction except for the cases in Table 5.4, which also take irrationalities into account. We summarise our analysis of these remaining cases in Tables 5.5 and 5.6. The italicised entries in

d	$E(G)$	Characteristic	Irrationalities	$r \leq$
6	$3_1.U_4(3)$	2	z_3	2^{15}
6	$6_1.U_4(3)$	$\neq 2, 3$	z_3	41407
15	$3_1.U_4(3)$	$\neq 3$	z_3	32
20	$U_4(3)$	2		16
20	$2.U_4(3)$	$\neq 2, 3$		7
20	$4.U_4(3)$	$\neq 2, 3$	i_1	5
21	$U_4(3)$	$\neq 2, 3$		7
21	$3_1.U_4(3)$	$\neq 2, 3$	z_3	7
34	$U_4(3)$	2		4
35	$U_4(3)$	$\neq 2, 3$		5

TABLE 5.4. Remaining modules to consider for G with $\text{soc}(G/Z(G)) \cong \text{PSU}_4(3)$.

Table 5.6 were obtained by a construction in GAP, the underlined entries in MAGMA, the bold entries by restricting to a maximal subgroup and applying Proposition 3.2, and the remaining entries were obtained from the Brauer characters and Proposition 3.1i). An application of Proposition 3.1ii) implies that G has a regular orbit on V in all of the cases listed in Table 5.6, except for those listed in Table 5.5, where the value of $b(G)$ is computed explicitly in MAGMA [2].

□

Proposition 5.4. *Theorem 5.1 holds for $n \geq 5$.*

Proof. Suppose that $n \geq 5$. We begin by applying Proposition 3.1v) with $d_1(G) \geq \lfloor (q^n - 1)/(q + 1) \rfloor$ and $\alpha(G) \leq n$. It follows that G has a regular orbit on V except possibly if G is a group listed in Table 5.7.

Suppose $(n, q) = (11, 2)$. From Propositions 3.1v) and 2.13, we only need consider the 682-dimensional Weil module V for $r = 3$. In GAP we compute that $i_2(H) < 3^{42.02}$ and $i_3(H) < 3^{53.58}$ for $H = \text{Aut}(\text{PSU}_{11}(2))$. We will bound the eigenspace dimensions of elements of projective prime order at least 5 on the 2048-dimensional Weil module \overline{V} for G using Propositions 2.10 and 2.11. Since V is a composition factor of \overline{V} , this also gives upper bounds for eigenspace dimensions in V . Let ρ be the Brauer character corresponding to \overline{V} and let W be the natural module for $U_{11}(2)$. Now, let $x \in G/Z(G)$, and suppose $o(x) = 5$. By Proposition 2.12, $\dim C_W(x) \in \{3, 7\}$ and therefore by Proposition 2.10, $\rho(x) = 8$ or 128, which, in the notation of Proposition 2.11, corresponds to eigenvalues $(\Omega^{408}, 1^8)$ or $(\Omega^{384}, 1^{128})$. Therefore, $\dim C_V(x) \leq \dim C_{\overline{V}}(x) \leq 512$. We summarise the analogous calculations for other prime order elements x in Table 5.8.

G	(d, r)	$\lceil \frac{\log G }{\log V } \rceil$	$b(G)$
$U_4(3).D_8$	(20, 2)	1	2
$3_1.U_4(3).2_2$	(15, 4)	1	1
$c \circ (6_1.U_4(3)), c \mid 24$	(6, 25)	1	2
$c \circ (6_1.U_4(3).2_2), c \mid 24$	(6, 25)	1	2
$6_1.U_4(3).2_2$	(6, 7)	2	2
$c \circ (6_1.U_4(3).2_2), c \mid 48$	(6, 49)	1	2
$3_1.U_4(3)$	(6, 4)	2	3
$3_1.U_4(3).2_2$	(6, 4)	3	3
$3_1.U_4(3)$	(6, 16)	1	2
$15 \circ (3_1.U_4(3))$	(6, 16)	2	2
$c \circ (3_1.U_4(3).2_2), c \mid 15$	(6, 16)	2	2
$3_1.U_4(3).2_2$	(6, 64)	1	1
$3_1.U_4(3).2_2$	(6, 256)	1	1

TABLE 5.5. Base sizes of some absolutely irreducible modules for G with $\text{soc}(G/Z(G)) \cong U_4(3)$.

$E(G)$	(d, r)	2A	2B	2C	2D	2E	2F	3A	3B	3C/3D	5A	7A/7B
$U_4(3)$	(35, 5)	(19, 16)	(21, 14)	(20, 15)	(25, 10)	(18, 17)	–	(17, 9 ²)	(17, 9 ²)	(12 ² , 11)	7	(5 ⁷)
$U_4(3)$	(34, 2 ^k)	18	<u>20</u>	<u>19</u>	<u>20</u>	<u>19</u>	–	(16, 9 ²)	(16, 9 ²)	(12 ² , 10)	(7 ⁴ , 6)	(5 ⁶ , 4)
$3_1.U_4(3)$	(21, 7)	(13, 8)	–	–	(16, 5)	(12, 9)	–	(9, 6 ²)	(11, 5 ²)	(7 ³)	(5, 4 ⁴)	5
$U_4(3)$	(21, 7)	(13, 8)	(14, 7)	(11, 10)	(15, 6)	(11, 10)	(11, 10)	(9 ² , 3)	(9, 6 ²)	(9, 6 ²)	(5, 4 ⁴)	3
$U_4(3)$	(21, 5)	(13, 8)	(14, 7)	(11, 10)	(15, 6)	(11, 10)	(11, 10)	(9 ² , 3)	(9, 6 ²)	(9, 6 ²)	5	(3 ⁷)
$4.U_4(3)$	(20, 5)	(10 ²)	(13, 7)	(10 ²)	–	–	–	(9 ² , 2)	(8, 6 ²)	(8, 6 ²)	10	(3 ⁶ , 2)
$2.U_4(3)$	(20, 7)	(12, 8)	(14, 6)	(10 ²)	(10 ²)	(10 ²)	(10 ²)	(9 ² , 2)	(8, 6 ²)	(8, 6 ²)	(4 ⁵)	4
$2.U_4(3)$	(20, 5)	(12, 8)	(14, 6)	(10 ²)	(10 ²)	(10 ²)	(10 ²)	(9 ² , 2)	(8, 6 ²)	(8, 6 ²)	10	(3 ⁶ , 2)
$U_4(3)$	(20, 2 ^k)	<u>12</u>	<u>14</u>	<u>10</u>	<u>14</u>	<u>10</u>	<u>11</u>	(9 ² , 2)	(8, 6 ²)	(8, 6 ²)	(4 ⁵)	(3 ⁶ , 2)
$3_1.U_4(3)$	(15, 4 ^k)	<u>9</u>	–	–	<u>11</u>	<u>9</u>	–	(9, 3 ²)	(7, 4 ²)	(5 ³)	(3 ⁵)	(3, 2 ⁶)
$3_1.U_4(3)$	(15, 25)	(8, 7)	–	–	(10, 5)	(9, 6)	–	(9, 3 ²)	(7, 4 ²)	(5 ³)	3	(3, 2 ⁶)
$3_1.U_4(3)$	(15, 7)	(8, 7)	–	–	(10, 5)	(9, 6)	–	(9, 3 ²)	(7, 4 ²)	(5 ³)	(3 ⁵)	3
$6_1.U_4(3)$	(6, 25 ^k)	(4, 2)	–	–	(5, 1)	(3 ²)	–	(3 ²)	(4, 1 ²)	(2 ³)	2	(1 ⁶)
$6_1.U_4(3)$	(6, 7 ^k)	(4, 2)	–	–	(5, 1)	(3 ²)	–	(3 ²)	(4, 1 ²)	(2 ³)	(2, 1 ⁴)	1
$3_1.U_4(3)$	(6, 4 ^k)	4	–	–	5	3	–	(3 ²)	(4, 1 ²)	(2 ³)	(2, 1 ⁴)	(1 ⁶)
#elements		2835	540	4536	126	5670	4536	560	3360	3360/40320 (2 classes)	653184	466560 (each)

TABLE 5.6. Eigenspace dimensions of elements of projective prime order.

n	5	6	7	8	9	10	11
q	2, 3	2, 3	2, 3	2	2	2	2

TABLE 5.7. Remaining groups to consider when $n \geq 5$.

From Table 5.8, we therefore deduce that $\mathcal{E}_{\max}(g_{\{2,3\}^r}) \leq 512$ on \overline{V} and therefore V . Therefore, if G has no regular orbit on V ,

$$3^{682} \leq 2(3^{42.02} + 3^{53.58})3^{\lfloor 10 \times 682/11 \rfloor} + 2|U_{11}(2).2| \times 3^{512}$$

This gives a contradiction for $r = 3$, so G has a regular orbit on V .

Now let $(n, q) = (10, 2)$. By Propositions 3.1v) and 2.13, we must consider the 341- and 342-degree Weil representations for $r = 3, 5, 7$. We begin with $r = 7$. By Propositions 2.10, 2.11 and 2.12, $\mathcal{E}_{\max}(g_{\{2,3,7\}^r}) \leq 256$ on a Weil representation. Let $H = G/Z(G)$. If G has no regular orbit on the 342-dimensional Weil module V ,

$$r^{342} \leq 2(i_2(H) + i_3(H) + i_7(H))7^{\lfloor 9 \times 342/10 \rfloor} + 2|U_{10}(2).2|7^{256}$$

and this gives a contradiction for $r = 7$. An analogous inequality gives the result when $d = 341$. Now suppose $r = 5$. By Propositions 2.10, 2.11 and 2.12, $\mathcal{E}_{\max}(g_{\{2,3,5\}^r}) \leq 160$ on a Weil representation. Let $H = G/Z(G)$. If G has no regular orbit on the 342-dimensional Weil module V , then by Proposition

$o(x)$	$\dim C_W(x)$	$\rho(x)$	Eigsp.
7	5	32	$(\Omega^{288}, 1^{32})$
11	1	2	$(\Omega^{186}, 1^2)$
	6	-64	$(\overline{\Omega}^{64}, \Omega^{128})$
17	3	8	$(\Omega^{120}, 1^8)$
19	2	-4	$(\overline{\Omega}^4, \Omega^{104})$
31	1	2	$(\Omega^{66}, 1^2)$
43	4	-16	$(\overline{\Omega}^{16}, \Omega^{32})$
683	0	-1	$(\overline{\Omega}, \Omega^3)$

 TABLE 5.8. Eigenspace dimensions of prime order elements of $U_{11}(2)$ on $\overline{V} = V_{2048}(3)$.

3.1iv)

$$r^{342} \leq \frac{1}{4}i_5(H) \times 5^{\lfloor 9 \times 342/10 \rfloor} + 2(i_2(H) + i_3(H))5^{\lfloor 9 \times 342/10 \rfloor} + 2|U_{10}(2).2|5^{160}$$

and this gives a contradiction for $r = 5$. An analogous inequality gives the result when $d = 341$. Finally, when $r = 3$, we use Propositions 2.10, 2.11 and 2.12 to deduce that $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 256$ on any of the Weil modules. We observe that every class of elements of order 3 in G has a representative in the subgroup $H = \text{SU}_6(2) \times \text{SU}_4(2)$. The 2^{10} -dimensional irreducible Weil module for $4 \circ 2^{1+20}.\text{SU}_{10}(2)$ restricts to the subgroup $\text{SU}_6(2)$ as the sum of composition factors of dimensions $(21 + 22 + 21)$ and to $\text{SU}_4(2)$ as the sum of composition factors of dimensions $(5 + 6 + 5)$. The restriction to $H = \text{SU}_6(2) \times \text{SU}_4(2)$ then is a tensor product $(21 + 22 + 21) \otimes (5 + 6 + 5)$ (cf. [14, Lemma 9.3.1]). Thus $V_{341}(3) \downarrow H$ has composition factors $(21 \otimes 5)^3/21 \otimes 1/1 \otimes 5$, while $V_{341}(3) \downarrow H$ has composition factors $(21 \otimes 5)^3/21 \otimes 1/1 \otimes 5/1 \otimes 1$. So any element of order 3 has a fixed point space of dimension at most $3 \times (\lfloor \frac{5 \times 21}{6} \rfloor \times 5) + \lfloor \frac{5 \times 21}{6} \rfloor + 1 \times 5 = 273$ on $V_{341}(3)$, and 274 on $V_{342}(3)$. We have by [3, Table B.4] that $i_2(G/Z(G)) < 2^{45}$ and $i_3(G/Z(G)) < 3^{44.54}$. Therefore, if G has no regular orbit on V ,

$$r^d \leq 2|\text{PSU}_{10}(2).2|r^{256} + 2i_2(G)r^{\lfloor 9d/10 \rfloor} + i_3(G)r^{d-68}$$

This gives a contradiction for $r = 3$, so G has a regular orbit on all of the Weil modules V .

Let $(n, q) = (9, 2)$. We need to consider $r \leq 17$, and by [21], we only need consider the Weil representations of degree 170 and 171. The degree 171 representations for $3.U_9(2)$ contain the z_3 irrationality, so we only have $r = 7$ here. By Propositions 2.10, 2.11 and 2.12, $\mathcal{E}_{\max}(g_{\{2,3,5\}'}) \leq 80$ for semisimple $g_{\{2,3,5\}'}$, and by [14, Lemma 8.3.10] inner elements $g \in G$ of projective order 2,3 have $\mathcal{E}_{\max}(g) \leq 128$. We compute using GAP that $\alpha(g_5) \leq 5$, and $\alpha(x) \leq 8$ for x an outer element of order 3. We also calculate from [3, Table B.4] that the number of outer elements of order 3 in $G/Z(G)$ is at most $7^{19.29}$, and by Lemma 2.8ii), the number of graph automorphisms x in $G/Z(G)$ is at most 2^{45} .

We bound the fixed point space dimensions of unipotent elements by restricting the Weil representations to appropriate subgroups of the form $\text{SU}_m(2) \times \text{SU}_{9-m}(2)$. From this technique we find that $\dim C_V(g_r) \leq 138$ for $r \leq 9$ on $V = V_{170}(r)$, while if $r = 11, 17$, we find using GAP that $\alpha(g_r) \leq 5$, so $\dim C_V(g_r) \leq 136$ by Proposition 3.1i). We also deduce using this technique that $\dim C_V(g_7) \leq 139$ on $V = V_{171}(7)$. If $d = 171$, $r = 7$ and G has no regular orbit on V ,

$$r^{171} \leq 2|\text{PSU}_9(2).6|r^{139} + 2 \times 7^{19.29}r^{\lfloor 7 \times 171/8 \rfloor} + 2^{45}r^{\lfloor 8 \times 171/9 \rfloor}$$

which is false, implying the existence of a regular orbit. Let $H = G/Z(G)$. If $d = 170$ and G has no regular orbit on V , then

$$r^{170} \leq 2|\text{PSU}_9(2).6|r^{138} + 2 \times i_3(H)r^{\lfloor 7 \times 170/8 \rfloor} + 2 \times i_2(H)r^{\lfloor 8 \times 170/9 \rfloor}$$

which gives a contradiction for $r = 7, 9, 11, 17$. If instead $r = 5$, we use GAP to determine the character values of graph automorphisms $x \in G$ and infer that $\mathcal{E}_{\max}(x) \leq 88$. If G has no regular orbit on V ,

$$r^{170} \leq 2|U_9(2).6|r^{80} + i_5(H)r^{\lfloor 4 \times 170/5 \rfloor} + 2(i_2(H) + i_3(H))r^{128}$$

This gives a contradiction for $r = 5$. Finally, when $r = 3$, we restrict the representation to $K = 3(U_3(2) \times U_6(2)).3$ and find that the composition factors of the 170-dimensional representation are $(3 \otimes 21)^2/2 \otimes 21/2 \otimes 1$. Every class of elements of order 3 in G has a representative in $\text{SU}_6(2) : 3$. In GAP we find that all elements of order 3 in $\text{SU}_6(2) : 3$ have fixed point spaces of dimension at

most 11 on the 21-dimensional representation. Therefore, for $g \in G$ of order 3, we have $\dim C_V(g) \leq 2(11 \times 3) + 2(11 + 1) = 90$. Therefore, if G has no regular orbit on V ,

$$r^{170} \leq 2|U_9(2).6|r^{80} + 2i_5(H)r^{\lfloor 4 \times 170/5 \rfloor} + i_3(H)r^{90} + 2i_2(G)r^{128}$$

This gives a contradiction for $r = 3$, so G has a regular orbit on V .

Now suppose $(n, q) = (8, 2)$. The number of prime order elements in $\text{PFU}_8(2)$ is less than $2^{52.94}$. By Proposition 3.1v) and examining [21], we see that we only need consider the Weil representations of G of degrees 85 and 86 for $r \leq 43$. By Propositions 2.10, 2.11 and 2.12, we compute that semisimple elements of prime order at least 7 have $\mathcal{E}_{\max}(g_{\{2,3,5\}'}) \leq 40$. Now, constructing the character table of $U_8(2).2$ using MAGMA, we determine that $\mathcal{E}_{\max}(g_{\{2,3,5\}'}) \leq 54$, except that elements in class $2A$ have $\mathcal{E}_{\max}(g) \leq 64$. We summarise upper bounds on the number of prime order elements in Table 5.9. We now find upper bounds for fixed point spaces of elements of order r_0 in representations in

Order r_0	2 (inner)	2 (outer)	3	5	7	11	17	43
$\log_{r_0}(\#r_0\text{-elts}) <$	32.31	35.23	26.45	20.86	19.29	14.72	14.02	10.74

TABLE 5.9. Bounds on the number of elements of prime order in $\text{Aut}(U_8(2))$.

characteristic r_0 . These bounds are summarised in Table 5.10. Let $H = G/Z(G)$. If G has no regular

r_0	Technique	Bound on 85-dim reps ν_{85}	Bound on 86-dim reps ν_{86}
3	Restrict to $\text{SU}_4(2) \times \text{SU}_4(2)$	53	54
5	Compute in GAP $\alpha(x) = 2$	42	43
7	Compute in GAP $\alpha(x) \leq 3$	56	57
11	Restrict to $\text{SU}_6(2) \times \text{SU}_6(2)$	69	70
17	Compute in GAP $\alpha(x) \leq 3$	57	57
43	Compute in GAP $\alpha(x) \leq 8$	74	74

TABLE 5.10. Bounds on unipotent elements in absolutely irreducible modules.

orbit on one of the 86-dimensional modules $V = V_d(r)$ in characteristic r_0 ,

$$r^{86} \leq 2 \times 2^{52.94} r^{40} + 2 \times (i_2(H) + i_3(H) + i_5(H)) r^{54} + 2 \times |2A| r^{64} + i_{r_0}(H) r^{\nu_{86}}$$

This gives a contradiction for $r \geq 3$. A similar calculation gives the result for the 85-dimensional representation.

Now let $(n, q) = (7, 3)$. By Propositions 3.1v) and 2.13, we need only consider the 546-dimensional Weil module for $r = 2$. By Propositions 2.10, 2.11 and 2.12, $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 459$. Therefore, if G has no regular orbit on V , then

$$r^{546} \leq 2|U_7(3).2|r^{459} + 2(i_2(G) + i_3(G))r^{\lfloor 6 \times 546/7 \rfloor}$$

which gives a contradiction for $r = 3$, so G has a regular orbit on V .

Suppose now that $(n, q) = (7, 2)$. Let $H = G/Z(G)$. We begin by summarising some information about the prime order elements in $U_7(2).2$ in Table 5.11.

If G has no regular orbit on an irreducible V , then by Proposition 3.1v),

$$\frac{1}{2}r^d \leq (i_{43}(H) + i_{11}(H) + i_7(H) + i_5(H))r^{\lfloor d/2 \rfloor} + (i_2(H) + i_3(H))r^{\lfloor 6d/7 \rfloor}$$

From [21], we see that we only need to consider Weil representations of dimensions 42, 43 for $r \leq 43$ and $r = 7, 25$ respectively, due to a z_3 irrationality in the character of the 43-dimensional representation. First let $V = V_{43}(r)$. We compute using GAP that $\alpha(g_{\{5,7\}'}) = 2$. Constructing the character table of $U_7(2).2$ using MAGMA [2], we deduce that if G has no regular orbit on V ,

$$\frac{1}{2}r^{43} \leq i_{43}(H)r + i_{11}(H)r^4 + i_7(H)r^{\lfloor 43/2 \rfloor} + i_5(H)r^{\lfloor 43/2 \rfloor} + i_3(H)r^{26} + |2A|(r^{32} + r^{11}) + (|2B| + |2C|)r^{26} + |2D|r^{\lfloor 2 \times 43/3 \rfloor}$$

Prime r_0	$\log_{r_0} i_{r_0}(H) <$	$\alpha(x) \leq$
2	27.32	3 (2D), 7
3	20.16	2 (3J/3K), 7
5	15.52	2
7	14.87	2
11	11.88	2
43	8.27	2
Total	$2^{48.43}$	

TABLE 5.11. Bounds on the number of prime order elements in $U_7(2).2$.

This is a contradiction for both $r = 7, 25$. Now suppose V is the 42-dimensional Weil module for $U_7(2).2$. From the Brauer character corresponding to V , we see that if G has no regular orbit on V in characteristics 5,7 then

$$\frac{1}{2}r^{42} \leq i_{43}(H)r + i_{11}(H)r^4 + i_7(H)r^{21} + i_5(H)r^{21} + i_3(H)r^{26} + |2A|(r^{32} + r^{10}) + (|2B|)r^{26} + |2C|r^{24} + |2D|r^{21}$$

This is a contradiction for $r \geq 5$. Similar computations give the results for other characteristics greater than 7. If instead $V = V_{42}(3^k)$, then we bound $\mathcal{E}_{\max}(g_3)$ by restricting the representation to $U_5(2).2$ and applying Proposition 3.2. So if G has no regular orbit on V ,

$$\begin{aligned} r^{42} &\leq 2i_{43}(H)r + 2i_{11}(H)r^4 + 2i_7(H)r^6 + 2i_5(H)r^{21} \\ &\quad + (|3A| + |3B|)r^{22} + (|3C| + |3D| + |3E|)r^{26} + (|3F| + |3G|)r^{22} + (|3H| + |3I|)r^{18} \\ &\quad + (|3J| + |3K|)r^{21} + |2A|(r^{32} + r^{10}) + 2(|2B|)r^{26} + 2|2C|r^{24} + 2|2D|r^{23} \end{aligned}$$

This gives a contradiction for $r \geq 3$.

Now let $(n, q) = (6, 3)$. We have $d_1(G) \geq 182$ by Proposition 2.1, while $\alpha(G) \leq 6$ by Proposition 2.4. Therefore, applying Proposition 3.1v) and examining [21], we see that we only need to consider $r = 2$ and $V = V_{182}(2)$.

From the Brauer character (computed using MAGMA [2]), we deduce that $\mathcal{E}_{\max}(g_{2'}) \leq 81$. Using GAP, we determine that $\alpha(g_2) \leq 5$, except for elements in class 2A where $\alpha(x) \leq 6$ by Proposition 2.4. Therefore, if G has no regular orbit on V then

$$r^{182} \leq 2|U_6(3).2^2|r^{81} + |2A|r^{\lfloor 5 \times 182/6 \rfloor} + i_2(G/Z(G))r^{\lfloor 4 \times 182/5 \rfloor}$$

This is a contradiction, so G has a regular orbit on V .

Now suppose $(n, q) = (6, 2)$ and let $H = U_6(2).S_3$. We summarise some information about elements of prime order in H in Table 5.12. Therefore, if G has no regular orbit on an irreducible module V ,

Prime r_0	$i_{r_0}(H)$	$\alpha(g_{r_0}) \leq$
2	1529055	6 (2A/2D), 3 (O/W)
3	22061888	6 (3D), 3 (O/W)
5	306561024	2
7	1313832960	2
11	1672151040	2

TABLE 5.12. Elements of prime order in $U_6(2).S_3$.

$$\frac{1}{2}r^d \leq (i_{11}(H) + i_7(H) + i_5(H))r^{\lfloor d/2 \rfloor} + (i_3(H) + i_2(H))r^{\lfloor 2d/3 \rfloor} + (|2A| + |2D| + |3D|)r^{\lfloor 5d/6 \rfloor}$$

Consulting [21], we see that the only representations that remain are the Weil representations of dimension 21 and 22 for $r \leq 11$. First suppose V is a 21-dimensional Weil representation for $U_6(2).S_3$ in characteristic 3. When $r = 3$, we compute using MAGMA that there is no regular orbit of G on V . Now suppose $r = 9$. Then from constructing the representation in GAP, we find that $\dim C_V(g_3) \leq 11$

and all involutions have eigenspaces of dimension less than 14, except elements in class 2A, which have an eigenspace of dimension 16. Therefore, if G has no regular orbit on V ,

$$r^{21} \leq 2(i_{11}(H) + i_7(H) + i_5(H))r^{10} + i_3(H)r^{11} + 2 \times i_2(H)r^{14} + 693(r^{16} + r^5)$$

This gives a contradiction for $r = 9$, so G has a regular orbit on V and $b(G) = 2$ for $r = 3$ by Lemma 3.4. Now suppose that V is a 21-dimensional Weil representation for $3.U_6(2).3$ in characteristic $\neq 2, 3$. Since the character contains a z_3 irrationality, we only consider $r = 7$. Constructing the module in GAP, we find that $\dim C_V(g_7) = 3$. This along with observations from the Brauer character (constructed in MAGMA [2]) show that if G has no regular orbit on V ,

$$r^{21} \leq 2(i_{11}(H) + i_7(H) + i_5(H))r^5 + 2 \times i_3(H)r^{11} + 2 \times i_2(H)r^{12}$$

This gives a contradiction, so G has a regular orbit on V .

Now suppose that V is a 22-dimensional Weil representation for $U_6(2).S_3$. From [21] we observe that this representation does not exist in characteristic 3, so we only need consider V in characteristic 5, 7 or 11. Examining the Brauer characters and constructions in GAP, we see that if G has no regular orbit on V ,

$$\begin{aligned} r^{22} \leq & 2 \times i_{11}(G)r^2 + 2 \times i_7(G)r^4 + 2 \times i_5(G)r^6 + |3A|(r^{10} + 2r^6) + |3B|(2r^9 + r^4) + |3C|(r^{10} + 2r^6) \\ & + |3D|(2r^{11}) + |3E|(r^{12} + 2r^5) + |3F|(2r^8 + r^6) + |3G|(r^8 + 2r^7) + |2A|(r^{16} + r^6) + |2B|(r^{14} + r^8) \\ & + |2C|(r^{12} + r^{10}) + |2D|(r^{15} + r^7) + |2E|(2r^{11}) \end{aligned}$$

This gives a contradiction for $r = 5, 7, 11$.

Now let $(n, q) = (5, 3)$. By Proposition 3.1v) and 2.13, $r \leq 7$ and we must consider 60- and 61-dimensional Weil modules. We calculate that $i_2(U_5(3).2) = 5430159$, and the total number of prime order elements in $U_5(3).2$ is 54826442783. We begin with $r = 7$. Constructing the group in GAP, we deduce that $i_7(U_5(3).2) < 7^{10.52}$. From the character table of $U_5(3).2$, we find that $\mathcal{E}_{\max}(g_2) \leq 40$, and $\mathcal{E}_{\max}(g_{\{2,7\}'}) \leq 27$. Therefore, if G has no regular orbit on the 60-dimensional Weil module V , then

$$r^{60} \leq 2 \times 54826442783r^{27} + 2i_2(G)r^{40} + 7^{10.52}r^{\lfloor 4 \times 60/5 \rfloor}.$$

This gives a contradiction for $r = 7$, so G has a regular orbit on V . Analogous calculations show that G also has a regular orbit on the three 61-dimensional Weil modules in characteristic 7. Suppose now that $r = 5$, and let V be the 60-dimensional Weil module. Again, we find that $\mathcal{E}_{\max}(g_2) \leq 40$ from the character, and $\mathcal{E}_{\max}(g_{\{2,5\}'}) \leq 27$. We also compute that $i_5(U_5(3).2) < 5^{13.61}$, and these elements all have $\alpha(x) = 2$. Therefore, if G has no regular orbit on V ,

$$r^{60} \leq 2 \times 54826442783r^{27} + 2 \times 5430159r^{40} + 5^{13.61}r^{30}$$

This is a contradiction for $r = 5$, and analogous calculations give a contradiction for the 61-dimensional modules as well. Finally, suppose that V is a 60-dimensional Weil module in characteristic 2. Examining the character we see that $\mathcal{E}_{\max}(g_3) \leq 27$, and $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 21$. By restricting the representation to $4.U_4(3).4$, we infer that for classes 2A and 2B, $\dim C_V(g) \leq 39$. Moreover, we compute from a construction of the module in MAGMA that outer involutions in class 2C have fixed point spaces of dimension 30. Therefore, if G has no regular orbit on V ,

$$r^{60} \leq 2 \times 54826442783r^{21} + 2i_3(G)r^{27} + (|2A| + |2B|)r^{39} + |2C|r^{30}$$

This gives a contradiction for $r \geq 2$.

Now suppose $(n, q) = (5, 2)$. The number of prime order elements in $H = U_5(2).2$ is 3479519. We summarise some further information about prime order elements in $U_5(2).2$ in Table 5.13. Therefore,

Prime r_0	$i_{r_0}(H)$	$\alpha(x) \leq$
2 (inner)	3135	5
2 (outer)	19008	3
3	56672	2 (3F) 5 (O/W)
5	912384	2
11	2488320	2

TABLE 5.13. Elements of prime order in $H = U_5(2).2$.

applying Proposition 3.1i) and iv), it follows from [21] that we only need to further consider an irreducible 44-dimensional representation when $r = 3$, 11-dimensional Weil representations for $r = 25$ and 10-dimensional Weil representations for $r \leq 187$. If V is the 44-dimensional representation and $r = 3$, we infer from the Brauer character that $\mathcal{E}_{\max}(g_{3V}) \leq 28$. Therefore, if G has no regular orbit on V ,

$$r^{44} \leq 2 \times 3479519r^{28} + |3F|r^{\lfloor 44/2 \rfloor} + (56672 - 42240)r^{\lfloor 4 \times 44/5 \rfloor}$$

which is false, so G has a regular orbit on V . Now suppose $V = V_{11}(25)$. From the Brauer characters of the two conjugate modules of dimension 11, we see that if G has no regular orbit on V ,

$$r^{11} \leq 912384r^5 + 3135(r^8 + r^3) + 2 \times 56672r^6 + 2488320r$$

This is a contradiction for $r = 25$. Now suppose V is the 10-dimensional irreducible Weil module. We summarise information about eigenspace dimensions in $U_5(2).2$ obtained from the Brauer tables and constructions in GAP in Table 5.14. Now, if G has no regular orbit on V ,

Characteristic r_0	Class in $U_5(2).2$								
	2A	2B	3A/3B	3C/3D	3E	3F	5A	11A/11B	2C
3	(8,2)	(6,4)	5	5	6	4	(2 ⁵)	(1 ¹⁰)	(5 ²)
5	(8,2)	(6,4)	(5 ²)	(4, 3 ²)	(6, 2 ²)	(4 ² , 2)	2	(1 ¹⁰)	(5 ²)
11	(8,2)	(6,4)	(5 ²)	(4, 3 ²)	(6, 2 ²)	(4 ² , 2)	(2 ⁵)	1	(5 ²)
Size of class in $U_5(2).2$	165	2970	352	7040	7040	42240	912384	2488320	19008

TABLE 5.14. Eigenspace dimensions on $V = V_{10}(r)$.

$$r^{10} \leq 165(r^8 + r^2) + 2970(r^6 + r^4) + 352(2r^5) + 7040(r^4 + 3r^2 + r^6 + 2r^2) + 42240(2r^4 + r^2) + 912384(5r^2) + 2488320(10r) + 19008(2r^5).$$

This gives a contradiction for $r \geq 25$. If $r = 3$ then $|V| < |G|$ and we compute using GAP that $b(U_5(2).2) = 2$ for both isoclinism types. We also compute that $b(2 \times U_5(2)) = 2$ and $b(2 \times U_5(2).2) = 3$. If $r = 5$ then we have $|V| < |G|$ as well, and $b(G) = 2$ by Lemma 3.4. If $r = 9$ and $\rho \in \mathbb{F}_9^\times$, then we compute that $\langle \rho \rangle.U_5(2)$ has a regular orbit on V unless $\langle \rho \rangle \cong \mathbb{F}_9^\times$, in which case $b(G) = 2$ by Lemma 3.4. Moreover, both of the isoclinic variants of shape $U_5(2).2$ has a regular orbit on V , but $\langle \rho \rangle.U_5(2).2$ does not for non-identity ρ . Again, in these cases we have $b(G) = 2$ by Lemma 3.4. If instead $r = 11$, then we compute that $\langle \rho \rangle.U_5(2)$ and $\langle \rho \rangle.U_5(2).2$ both always have a regular orbit on V . \square

6. PROOF OF THEOREM 1.1: SYMPLECTIC GROUPS

The following is the main result of this section.

Theorem 6.1. *Suppose G is an almost quasisimple group with $E(G)/Z(E(G)) \cong \text{PSp}_n(q)$, $n \geq 4$, with $(n, q) \neq (4, 2)$. Let $V = V_d(r)$, $(r, q) = 1$ be a module for G , with absolutely irreducible restriction to $E(G)$. Also suppose that $(r, |G|) > 1$. Then one of the following holds.*

- i) $b(G) = \lceil \log |G| / \log |V| \rceil$,
- ii) $b(G) = \lceil \log |G| / \log |V| \rceil + 1$ and (G, V) lies in Table 1.1, or
- iii) (G, V) is one of $(2 \times \text{PSp}_6(2), V_7(3))$, $(\text{PSp}_4(3).2, V_5(3))$, $(2 \circ \text{PSp}_4(3), V_5(3))$, or $(\text{PSp}_4(3).2, V_6(2))$ and $b(G) = \lceil \log |G| / \log |V| \rceil + 2 = 4, 4, 4, 5$ respectively.

Note that $(n, q) \neq (4, 2)$, since $\text{PSp}_4(2) \cong S_6$, which is dealt with in [9]. Moreover, recall that $\text{PSp}_4(3) \cong U_4(2)$, so we also consider modules for this group in characteristic 3.

6.1. Odd characteristic.

Proposition 6.2. *Theorem 6.1 holds for $E(G)/Z(E(G)) \cong \text{PSp}_n(q)$, q odd.*

Suppose $(n, q) \neq (4, 3)$ (this case will be dealt with separately), and let $m = n/2$. From [3, Table B.7] we compute that the number of transvections in $G/Z(G)$ is at most $2q^{n-1}$. If G has no regular orbit on an irreducible module $V = V_d(r)$, then by Propositions 2.5 and 3.1v)

$$r^d \leq 2|\mathrm{P}\Gamma\mathrm{Sp}_n(q)|r^{\lfloor \frac{m+2}{m+3}d \rfloor} + 2(q^n - 1)r^{\lfloor \frac{n-1}{n}d \rfloor} \quad (5)$$

This gives a contradiction in all cases except those listed in Table 6.1.

n	14	12	10	8	6	4
q	3	3	3	3,5	≤ 9	≤ 19

TABLE 6.1. Remaining symplectic groups in odd characteristic.

Proposition 6.3. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_{14}(3)$.*

Proof. By [16] and (5), we only need consider Weil representations V of dimension 1093 for $r = 2$. Let \bar{V} be the 2187-dimensional Weil module (as described in Section 2.3) for G with corresponding character ρ . For an element $x \in G$ of projective prime order at least 5, we use Propositions 2.9, 2.12, and 2.11 to find the eigenspace dimensions of x on \bar{V} over \mathbb{F}_2 , and summarise our findings in Table 6.2. These eigenspace dimensions of x on V form upper bounds for the eigenspace dimensions of x on V , and therefore $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 459$ on V . From the proof of [17, Theorem 4.3], $\alpha(g_3) \leq 9$, unless g_3 is a transvection, where $\alpha(g_3) \leq 14$ by Proposition 2.4. From Proposition 2.5, the number of transvections in $G/Z(G)$ is at most $3^{14} - 1$, and for x an involution, $\alpha(x) \leq 10$ [17]. Therefore, if G has no regular orbit on V ,

$$r^d \leq 2|\mathrm{PSp}(14, 3)|r^{459} + i_2(G/Z(G))r^{\lfloor 9d/10 \rfloor} + 2(3^{14} - 1)r^{\lfloor 13d/14 \rfloor} + 2i_3(G/Z(G))r^{\lfloor 8d/9 \rfloor}$$

This gives a contradiction for $r = 2$ and $d = 1093$.

Proj, prime order	$\dim C_W(x)$	$ \rho(x) $	Eigenvalue mults
5	2	3	$(438^4, 435)$
	6	27	$(459, 432^4)$
	10	243	$(486^4, 243)$
7	2	3	$(315, 312^6)$
	8	81	$(324^6, 243)$
11	4	9	$(207, 198^{10})$
13	2	3	$(171, 168^{12})$
	8	81	$(243, 162^{12})$
41	6	27	$(54^{40}, 27)$
61	4	9	$(36^{60}, 27)$
73	2	3	$(30^{72}, 27)$
547	0	1	$(4, 3^{546})$
1093	0	1	$(3, 2^{1092})$

TABLE 6.2. Eigenspace dimensions of some prime order elements in $\mathrm{PSp}_{14}(3)$ on a 2187-dimensional Weil module.

□

Proposition 6.4. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_{12}(3)$.*

Proof. Here by (5) and Propositions 2.14 and 3.1v), we only need consider the Weil modules of dimension 364,365 for $r = 4, 5$ and $r = 5$ respectively. First suppose $r = 5$. We restrict the Weil representations to $H = \mathrm{Sp}_8(3) \times \mathrm{Sp}_4(3)$ and by [14, Lemma 9.3.13], we see that $V_{364} \downarrow H = 41 \otimes 4 + 40 \otimes 5$, while $V_{365} \downarrow H = 41 \otimes 5 + 40 \otimes 4$, where each integer denotes an irreducible module of that dimension. Now, H contains a Sylow 5-subgroup of G , so all classes of elements of order 5 in G have representatives in H . Therefore, by Propositions 2.5 and 3.3, any element of order 5 in G has a fixed point space of dimension at most $41 \times 3 + 40 \times 4 = 283$ on the 364-dimensional module,

and at most $41 \times 4 + 40 \times 3 = 284$ on the 365-dimensional module. Examining the Brauer character and using [14, Lemma 9.3.13], we see that $\mathcal{E}_{\max}(g_{\{5,73\}'}) \leq 273$ on either of the Weil modules. Since 73 is a primitive prime divisor of $3^{12} - 1$, $\alpha(x) \leq 3$ for any $x \in G$ of order 73 by the proof of [17, Theorem 4.3]. Therefore, if G has no regular orbit on V , then $r^d \leq 2|2.\text{PSp}_{12}(3).2|r^{284}$, and this is a contradiction for $r = 5$, $d = 364$ or 365 .

Now suppose $V = V_{364}(4)$. Since the two Weil modules of this dimension fuse on $\text{PSp}_{12}(3).2$, we assume that $G/Z(G) \cong \text{PSp}_{12}(3)$. We observe that all classes of involutions have representatives in $H = \text{Sp}_8(3) \times \text{Sp}_4(3)$. Now, $V_{364} \downarrow H$ has composition factors $(40 \otimes 4)^2$, $1 \otimes 4$ and $40 \otimes 1$ [14, Lemma 9.3.13]. By Propositions 3.2 and 3.3, every involution in G has a fixed point space of dimension at most $2 \times 40 \times 3 + 3 + 40 = 283$. Using the same argument as the characteristic 5 case, we deduce that $\mathcal{E}_{\max}(g_{2'}) \leq 273$. We also compute that the number of inner involutions of G is less than 2^{68} . Therefore, if G has no regular orbit on V ,

$$r^{364} \leq 2|\text{PSp}(12, 3).2|r^{273} + 2^{68}r^{283}$$

This is a contradiction for $r = 4$. □

Proposition 6.5. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_{10}(3)$.*

Proof. By (5) and Proposition 2.14, we only need to consider the Weil representations of G of dimensions 121 and 122 with $r \in \{4, 7, 13, 16, 25, 49\}$, due to z_3 irrationalities in the corresponding Brauer characters.

We summarise upper bounds for eigenspace dimensions of elements of projective prime order acting on $V = V_{121}(r)$ in Table 6.3. We obtain the bounds for elements of projective order $r_0 \mid r$ by restricting V to $\text{Sp}_8(3) \times \text{Sp}_2(3)$ and then applying Propositions 3.2 and 3.1i). When $r_0 \nmid r$, we achieve the bounds by applying Propositions 2.9 and 2.12 for $r_0 = 11, 41, 61$, and by [14, Lemma 9.3.12] otherwise. Applying Proposition 3.1iv) it follows that G has a regular orbit on V for all valid r . If instead $V = V_{122}(r)$, we compute upper bounds on \mathcal{E}_{\max} which are at most one greater than those listed in Table 6.3, so another application of Proposition 3.1iv) yields the result.

Prime r_0	$r_0 \nmid r$	$r_0 \mid r$
2	82 (2A), 70	94
3	81 (3A/3B) 68 (O/W)	–
5	27	67
7	18	76
11	23	–
13	14	94
41	8	–
61	4	–

TABLE 6.3. Bounds on eigenspace dimensions for projective prime order elements in G with $\text{soc}(G/Z(G)) \cong \text{PSp}_{10}(3)$ acting on $V_{121}(r)$.

□

Proposition 6.6. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_8(5)$.*

Proof. From Proposition 2.1, we deduce that the smallest representations of G are Weil modules of dimensions 362 and 363, and the next smallest module has dimension at least 32240 by Proposition 2.14. By (5), we see that we only need to consider the Weil representations for $r = 2, 3$; however the Weil representations are not realised over either field by [16, §5]. □

Proposition 6.7. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_8(3)$.*

Proof. The number of prime order elements in $H = \text{PSp}_8(3).2$ is 8658865029997007, and of these, 8022531299328000 are elements of order 41 with $\alpha(x) = 2$ (computed in GAP), and 6560 are transvections. Therefore, by Propositions 2.5 and 3.1 if G has no regular orbit on the irreducible module

$$V = V_d(r),$$

$$\frac{1}{2}r^d \leq i_{41}(H)r^{\lfloor d/2 \rfloor} + 6560r^{\lfloor 7d/8 \rfloor} + (8658865029997007 - i_{41}(H) - 6560)r^{\lfloor 6d/7 \rfloor}$$

Therefore, by Proposition 2.14, we only need to check the 40- and 41- dimensional Weil representations for $r \leq 329$. In characteristic 41, there is nothing to consider because the Weil characters contain $z_3 \notin \mathbb{F}_{41}$. Moreover, $z_3 \notin \mathbb{F}_5$, so we only need to consider $r = 25$ in characteristic 5. The computed upper bounds for \mathcal{E}_{\max} on $V = V_{40}(r)$ are summarised in Table 6.4. The values in the third column were computed from the associated Brauer character inferred from [13, Table XV], and the bounds in the final column were determined by refining $\alpha(g_{r_0})$ and applying Proposition 3.1i) for $r = 2, 7, 13$, and by restricting $V \downarrow \mathrm{Sp}_6(3) \times \mathrm{Sp}_2(3)$ for $r_0 = 5$.

Prime r_0	$i_{r_0}(\mathrm{P}\mathrm{Sp}(8, 3).2) <$	$r_0 \nmid r$	$r_0 \mid r$
2	$2^{31.91}$	28	24
3	$3^{24.11}$	27 (3A/3B), 23 (O/W)	–
5	$5^{18.99}$	9	22
7	$7^{16.92}$	6	20
13	$13^{13.13}$	4	20
41	$41^{9.87}$	2	–

TABLE 6.4. Upper bounds on \mathcal{E}_{\max} for elements of projective prime order acting on $V_{40}(r)$.

Therefore, applying Proposition 3.1ii), we see that G has a regular orbit on V in all cases where r is odd.

If V is a Weil module in characteristic 2, then $d = 40$ and $r = 4^k$. Examining the Brauer character, we see that if G has no regular orbit on V then

$$\begin{aligned} r^{40} \leq & 2^{31.91}r^{24} + |5A|(4r^9 + r^4) + |5B|(5r^8) + |7A|(6r^6 + r^4) + i_{13}(G)(r^4 + 12r^3) + i_{41}(G)(20r^2) \\ & + (|3A| + |3B|)(r^{27} + r^{13}) + |3C|(2r^{18} + r^4) + |3D|(r^{22} + 2r^9) + (|3E| + |3F|)(r^{18} + r^{13} + r^9) \\ & + |3G|(r^{16} + 2r^{12}) + |3H|(2r^{15} + r^{10}) + |3I|(r^{16} + 2r^{12}) + (|3J| + |3K|)(r^{15} + r^{13} + r^{12}) \end{aligned}$$

This is a contradiction for $r \geq 4$. Finally, if $d = 41$ and r is odd, then we compute upper bounds for \mathcal{E}_{\max} and find that they are at most one greater than those given in Table 6.4. An application of Proposition 3.1ii) gives the result. \square

Proposition 6.8. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_6(9)$.*

Proof. By (5) and Proposition 2.14, we only need to consider the Weil representations of G of dimensions 364 for $r = 2$. Now $\mathrm{PSp}_6(9) < \mathrm{PSp}_{12}(3)$ and we can consider this Weil module as the restriction of the 364-dimensional Weil module V_2 of $\mathrm{PSp}_{12}(3)$ to $\mathrm{PSp}_6(9)$. By the proof of Proposition 6.4, we see that for projective prime order $g \in G$, $\mathcal{E}_{\max}(g) \leq 283$. Therefore if G has no regular orbit on V , $r^{364} \leq 2|\mathrm{PSp}_6(9).2|r^{284}$ and this gives a contradiction for $r = 2$. \square

Proposition 6.9. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_6(7)$.*

Proof. By (5) and Proposition 2.14, we only need consider the two 171-dimensional Weil modules for $\mathrm{PSp}_6(7)$ with $r = 2$. Using Proposition 2.9, we deduce that $\mathcal{E}_{\max}(g_{\{2,3\}^\nu}) \leq 91$. In GAP we determine that $\alpha(g_{\{2,3\}}) \leq 4$, so $\dim C_V(g_{\{2,3\}}) \leq 128$ by Proposition 3.1i). We compute that $i_2(G/Z(G)) < 2^{33.72}$ and $i_3(G/Z(G)) < 2^{40}$. So if G has no regular orbit on V ,

$$r^{171} \leq 2|\mathrm{PSp}(6, 7).2|r^{91} + (2^{33.72} + 2^{40})(r^{128} + r^{43})$$

This is a contradiction for $r = 2$. \square

Proposition 6.10. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_6(5)$.*

Proof. By (5), we only need to consider Weil representations with $d = 62, 63$ for $r \in \{4, 9, 16\}$ due to a b_5 irrationality. From Propositions 2.9 and 2.12, we see that elements $g \in G$ of projective order 7, 13 or 31 have $\dim C_V(g) \leq 18, 10$ and 5 respectively. Furthermore, for $g \in G$ of order 5, $\dim C_V(g) \leq 30$ unless g is one of the 15624 elements that have a 5-dimensional fixed point space on the natural module for G , in which case $\dim C_V(g) \leq 50$. In GAP we compute that all elements of order 2, 3 have $\alpha(x) \leq 4, 3$ respectively. Therefore, if G has no regular orbit on a Weil module V in either characteristic 2 or 3,

$$\frac{1}{2}r^{62} \leq i_2(G)r^{[3 \times 63/4]} + i_3(G)r^{[2 \times 63/3]} + i_5(G)r^{30} + 15624r^{50} + i_7(G)r^{18} + i_{13}(G)r^{10} + i_{31}(G)r^5$$

and this gives a contradiction for $r \geq 4$. □

Proposition 6.11. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_6(3)$.*

Proof. We summarise information about elements of prime order in $H = \text{PSp}_6(3).2$ in Table 6.5. Therefore, if G has no regular orbit on the d -dimensional irreducible module V ,

Prime r_0	# elts x of order r_0 in $\text{PSp}_6(3).2$	$\alpha(x) \leq$
2	605151	4 (2A), 3 (2B), 5 (2C)
3	5307848	6 (3A/3B), 3 (O/W)
5	38211264	2
7	327525120	2
13	705438720	2
Total	1077088103	

TABLE 6.5. Elements of prime order in $\text{PSp}_6(3).2$.

$$\frac{1}{2}r^d \leq (i_5(H) + i_7(H) + i_{13}(H))r^{[d/2]} + i_3(H)r^{[2d/3]} + (|3A| + |3B|)r^{[5d/6]} + |2A|r^{[3d/4]} + |2B|^{[2d/3]} + |2C|r^{[4d/5]}$$

This gives a contradiction except for $d = 78$ and $r = 2$ or $d = 13, 14$ for $r \leq 93$. If $V = V_{78}(2)$, then from a GAP construction and the Brauer character we deduce that every element g of projective prime order has $\mathcal{E}_{\max}(g) \leq 46$ on V . Therefore if G has no regular orbit on V , $r^{78} \leq 2 \times 1077088103r^{46}$, which is a contradiction.

Now suppose $V = V_{14}(r)$. These representations contain a z_3 irrationality, so we only need to consider $r = 7, 13, 25, 49$. We bound the eigenspace dimensions of unipotent elements g of order r_0 by restricting V to a subgroup K . We summarise our findings in Table 6.6. Therefore, if G has no

Characteristic r_0	K	$\dim C_V(g) \leq$
5	$\text{PSp}_2(3) \times \text{PSp}_4(3)$	8
7	$\text{SL}_2(13)$	2
13	$\text{SL}_3(3).2$	2

TABLE 6.6. Upper bounds for $\dim C_V(g)$ for $V = V_{14}(r)$ obtained by restricting to $K < G$.

regular orbit on $V = V_{14}(13)$,

$$r^{14} \leq |2A|(r^{10} + r^4) + |2B|(2r^7) + (|3A| + |3B|)(r^9 + r^5) + |3C|(2r^6 + r^2) + |3D|(r^8 + 2r^3) + (|3E| + |3F|)(r^6 + r^5 + r^3) + |3G|(r^6 + 2r^4) + i_5(G)(4r^3 + r^2) + i_7(G)(7r^2) + i_{13}(G)r^2$$

This gives a contradiction for $r = 13$, and the calculations for $r = 25, 49$ are similar. If $r = 7$, then using a construction in MAGMA, we show that there is no regular orbit and $b(G) = 2$ by Proposition 3.4. Now suppose that V is a 13-dimensional Weil module. Again, there is a z_3 irrationality, so $r \in \{4, 7, 13, 16, 25, 49, 64\}$. Table 6.7 summarises some fixed point space bounds for unipotent elements. The values were obtained from constructions of the relevant modules in GAP.

Characteristic r_0	$\dim C_V(g) \leq$
2	9 (2A), 7 (2B)
5	3
7	2
13	1

TABLE 6.7. Bounds on $\dim C_V(g)$ for unipotent elements g acting on $V_{13}(r)$.

If G has no regular orbit on $V = V_{13}(13)$, then

$$r^{13} \leq 705438720r + |2A|(r^8 + r^5) + |2B|(r^7 + r^6) + |3A|(r^9 + r^4) + |3B|(r^9 + r^4) + |3C|(2r^6 + r) \\ + |3D|(r^7 + 2r^3) + (|3E| + |3F|)(r^6 + r^4 + r^3) + |3G|(r^5 + 2r^4) + |5A|(4r^3 + r) + |7A|(6r^2 + r)$$

This gives a contradiction for $r = 13$. Similar calculations give the results for $r \geq 16$. If $V = V_{13}(4)$, then $|V| < |G|$ and by Proposition 3.4, $b(G) = 2$. If $r = 7$, then we compute from constructions in MAGMA that $\text{PSP}_6(3)$ has a regular orbit on V , however $c \times \text{PSP}_6(3)$ does not for $c \in \{2, 3, 6\}$. In these cases, $b(G) = 2$ by the result for $r = 49$ and Proposition 3.4. \square

Proposition 6.12. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_4(q)$, $q \in \{13, 17, 19\}$.*

Proof. If $q = 19$, then by (5) and Proposition 2.14, we only need to consider Weil modules for $r = 2$, but these are not realised over \mathbb{F}_2 due to a b_{19} irrationality in the Brauer character. Now let $q = 17$. There are 62466181283 elements of prime order in $\text{PSp}_4(17).2$ and of these, 24304611 are involutions and 48492960768 are elements of order 29. In GAP we compute that $\alpha(g_{29}) = 2$. Therefore, if G has no regular orbit on some irreducible module V in cross-characteristic,

$$\frac{1}{2}r^d \leq (62466181283 - 48492960768)r^{\lfloor 3d/4 \rfloor} + 48492960768r^{\lfloor d/2 \rfloor} + 24304611r^{\lfloor 4d/5 \rfloor}.$$

This gives a contradiction for $d = d_1(E(G))$ by Proposition 2.1, and so for all V by Proposition 3.1v). Similarly, the number of prime order elements in $\text{PSp}_4(13).2$ is 6986011811, and of these, 4883931 are involutions and 3224422656 are elements of order 17. In GAP, we compute that $\alpha(g_{17}) = 2$. Therefore, if G has no regular orbit on an irreducible module V ,

$$\frac{1}{2}r^d \leq (6986011811 - 3224422656)r^{\lfloor 3d/4 \rfloor} + 3224422656r^{\lfloor d/2 \rfloor} + 4883931r^{\lfloor 4d/5 \rfloor}.$$

By Propositions 2.14 and 3.1v), we see that this is a contradiction for $d = d_1(E(G))$ by Proposition 2.1, and so for all V . \square

Proposition 6.13. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_4(11)$.*

Proof. The number of prime order elements in $H = \text{PSp}_4(11).2$ is 3646870787 and of these, 3162456000 are elements of order 61 which, from computations in GAP, have $\alpha(x) = 2$. We also have $i_2(H) = 1800843$. Therefore, if G has no regular orbit on $V = V_d(r)$, then

$$r^d \leq 2 \times i_{61}(H)r^{\lfloor d/2 \rfloor} + 2i_2(H)r^{\lfloor 4d/5 \rfloor} + (3645889235 - i_{61}(H) - i_2(H))(r^{\lfloor 3d/4 \rfloor} + r^{\lfloor d/4 \rfloor})$$

This gives a contradiction for all V , except for the 60- and 61- dimensional Weil representations for $r = 3$. There are 3247640 elements of order 3 in $\text{PSp}_4(11)$. From the Brauer character, we deduce that $\mathcal{E}_{\max}(g_{3'}) \leq 36$ on a Weil module. Therefore, if G has no regular orbit on either of the Weil modules V ,

$$r^{60} \leq 2 \times 3645889235r^{36} + 3247640r^{\lfloor 3 \times 61/4 \rfloor}$$

And this gives a contradiction for $r = 3$. \square

Proposition 6.14. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_4(9)$.*

Proof. The number of prime order elements in $H = \text{PSp}_4(9).2^2$ is 457397927 and of these 419904000 are elements of order 41 and have $\alpha(x) = 2$ from GAP computations. In addition, $i_2(H) = 610893$. Therefore, if G has no regular orbit on $V = V(d, r)$,

$$r^d \leq 2i_{41}(H)r^{\lfloor d/2 \rfloor} + 2i_2(H)r^{\lfloor 4d/5 \rfloor} + 2(457397927 - i_{41}(H) - i_2(H))r^{\lfloor 3d/4 \rfloor}$$

This gives a contradiction for all V except for the 40- and 41- dimensional Weil modules for $r \leq 5$. Since $\mathrm{Sp}_4(9).2_1 < \mathrm{Sp}_8(3)$, we can consider these Weil representations as the restrictions of those of the same dimension for $\mathrm{Sp}_8(3)$. If $r = 4, 5$, then we deduce that G has a regular orbit on V because the same is true for the Weil module of $\mathrm{Sp}_8(3)$. This leaves the 40-dimensional Weil module for $r = 2$. Of the 610983 involutions in $\mathrm{PSP}_4(9).2^2$, only 368631 preserve the Weil module (i.e., those in $\mathrm{PSP}_4(9).2_1$). Constructing the module in GAP, we find that elements g in classes 2A or 2B have $\dim C_V(g) = 20$ and 24 respectively. Moreover, the two classes of field automorphisms in $\mathrm{PSP}(4, 9).2_1$ also have $\dim C_V(g) = 20$ and 24. These observations, along with those from the Brauer character imply that if G has no regular orbit on V ,

$$r^{40} \leq (|2A| + |2C|)r^{20} + (|2B| + |2D|)r^{24} + |3A|(r^{22} + 2r^9) + |3B|(r^4 + 2r^{18}) + |3C|(r^{10} + 2r^{15}) \\ + |3D|(r^{16} + 2r^{12}) + 2i_5(G)r^9 + i_5(G)(40r)$$

This gives a contradiction for $r = 2$ and so G has a regular orbit on V . \square

Proposition 6.15. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_4(7)$.*

Proof. We summarise some information about elements of prime order in $H = \mathrm{PSP}_4(7).2$ in Table 6.8. Therefore, if G has no regular orbit on an irreducible module $V = V_d(r)$,

Prime r_0	$i_{r_0}(H)$	$\alpha(g_{r_0})$
2	122451	3 (2B/2D), 5 (2A/2C)
3	274400	3
5	5531904	2
7	5764800	2 (7E/7F), 4 (7A/7B), 3 (O/W)

TABLE 6.8. Information about elements of prime order in $\mathrm{PSp}_4(7).2$.

$$\frac{1}{2}r^d \leq (|2B| + |2D|)r^{\lfloor 2d/3 \rfloor} + (|2A| + |2C|)r^{\lfloor 4d/5 \rfloor} + i_3(H)r^{\lfloor 2d/3 \rfloor} + i_5(H)r^{\lfloor d/2 \rfloor} + (|7E| + |7F|)r^{\lfloor d/2 \rfloor} \\ + (|7A| + |7B|)r^{\lfloor 3d/4 \rfloor} + (|7C| + |7D|)r^{\lfloor 2d/3 \rfloor}$$

From [21] we see that we need only consider the 24-dimensional Weil modules for $r = 2, 4$. When $r = 4$, from the Brauer character and constructing the representation in GAP, we see that if G has no regular orbit on V ,

$$r^{24} \leq |3A|(r^9 + 2r^8) + |3B|(r^{11} + 2r^7) + |5A|(5r^5) + (|7A| + |7B|)(3r^7 + r^4) + |7C|(r + 6r^4) \\ + |7D|(r^7 + 6r^3) + (|7E| + |7F|)(4r^4 + 3r^3) + |2A|r^{15} + |2B|r^{12}$$

This gives a contradiction for $r = 4$. When $r = 2$, $|V| < |G|$ and by Proposition 3.4, $b(G) = 2$. \square

Proposition 6.16. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \mathrm{PSp}_4(5)$.*

Proof. We summarise information about elements of prime order in $H = \mathrm{PSp}_4(5).2$ in Table 6.9. Therefore, if G has no regular orbit on an irreducible module V ,

Prime r_0	$i_{r_0}(H)$	$\alpha(x)$
2	16875	5 (2A/2C), 3 (2B/2D)
3	26000	3
5	390624	4 (5A/5B), 3 (5C/5D), 2 (5E/5F)
13	1080000	2

TABLE 6.9. Elements of prime order in $H = \mathrm{PSp}_4(5).2$.

$$\frac{1}{2}r^d \leq (|2B| + |2D|)r^{\lfloor 2d/3 \rfloor} + (|2A| + |2C|)r^{\lfloor 4d/5 \rfloor} + i_3(H)r^{\lfloor 2d/3 \rfloor} + i_{13}(H)r^{\lfloor d/2 \rfloor} + (|5E| + |5F|)r^{\lfloor d/2 \rfloor} \\ + (|5A| + |5B|)r^{\lfloor 3d/4 \rfloor} + (|5C| + |5D|)r^{\lfloor 2d/3 \rfloor}$$

It follows from [21] that we only need to consider 12- and 13-dimensional Weil modules for $r = 9$ and $r = 4, 9, 16$ respectively, as well as $d = 40$ for $r = 2$. We begin by supposing $V = V_{40}(2)$. Examining the corresponding Brauer character, we see that $\mathcal{E}_{\max}(g_{2'}) \leq 16$, and constructing the representation in GAP, we find that elements in class 2C have a 26-dimensional fixed point space, and all other involutions g have $\dim C_V(g) \leq 24$. Therefore, if G has no regular orbit on V ,

$$r^d \leq 2(1080000 + 390624 + 26000)r^{16} + 16875r^{24} + 300r^{26}$$

This is a contradiction for $r = 2$, so G has a regular orbit on V . Now suppose that $V = V_{13}(9)$. Examining the Brauer character and constructing the representations in GAP, we find that if G has no regular orbit on one of the two 13-dimensional Weil representations,

$$\begin{aligned} r^{13} \leq & |2A|(r^9 + r^4) + |2B|(r^7 + r^6) + (|5A| + |5B|)(r^3 + 2r^5) + |5C|(4r^3 + r) + |5D|(r^5 + 4r^2) \\ & + (|5E| + |5F|)(3r^3 + 2r^2) + (|13A| + |13B|)(13r) + (|3A| + |3B|)r^5 \end{aligned}$$

This is a contradiction for $r = 9$.

Now suppose that V is one of the 12-dimensional Weil modules for $r = 4, 9, 16$. If $r = 9$ and G has no regular orbit on V , then from the Brauer character and constructions in GAP,

$$\begin{aligned} r^{12} \leq & |2A|(2r^6) + |2B|(2r^6) + (|5A| + |5B|)(r^2 + 2r^5) + |5C|(4r^3) + |5D|(r^4 + 4r^2) \\ & + (|5E| + |5F|)(2r^3 + 3r^2) + (|13A| + |13B|)(12r) + (|3A| + |3B|)r^7 \end{aligned}$$

This gives a contradiction for $r = 9$. If $r = 4$ or 16 then construct the representation in GAP and find a representative of a regular orbit. \square

Proposition 6.17. *Proposition 6.2 holds for $E(G)/Z(E(G)) \cong \text{PSp}_4(3)$.*

Proof. We summarise some information about elements of prime order in $H = \text{PSp}_4(3).2$ in Table 6.10. This information was obtained using GAP.

Prime r_0	$i_{r_0}(H)$	$\alpha(x)$
2	891	3 (2B/2D), 5 (2A), 6 (2D)
3	800	4 (3A), 3 (O/W)
5	5184	2

TABLE 6.10. Elements of prime order in $H = \text{PSp}_4(3).2$.

If G has no regular orbit on the irreducible module V ,

$$\frac{1}{2}r^d \leq 5184r^{\lfloor d/2 \rfloor} + 80r^{\lfloor 3d/4 \rfloor} + (240 + 480)r^{\lfloor 2d/3 \rfloor} + (270 + 540)r^{\lfloor 2d/3 \rfloor} + 45r^{\lfloor 4d/5 \rfloor} + 36r^{\lfloor 5d/6 \rfloor}$$

This gives a contradiction except for the modules set out in Table 6.11. In Table 6.11 we also set out our analysis of these cases. We give the eigenspace dimensions of elements of projective prime order, obtained by examining the Brauer characters of the modules and constructing the modules themselves in GAP.

Applying Proposition 3.1ii), or alternatively constructing the module with GAP, we see that G has a regular orbit on V unless (G, V) appear in Table 1.1, where the base size is computed explicitly. \square

6.2. Even characteristic.

Proposition 6.18. *Theorem 6.1 holds for $E(G)/Z(E(G)) \cong \text{PSp}_n(q)$, q even.*

Proof. By Proposition 2.1 we have $d_1(G) \geq (q^{n/2} - 1)(q^{n/2} - q)/2(q + 1)$ and also $\alpha(G) \leq n + 1$ by Proposition 2.4. Applying Proposition 3.1v), we see that G has a regular orbit on V except possibly when $q = 2$ and $4 \leq n \leq 10$, or $(n, q) = (4, 4)$. We consider each of these cases in turn.

First let $E(G)/Z(E(G)) \cong \text{PSp}_{10}(2)$. We include information about elements of prime order in $H = G/Z(G)$ in Table 6.12.

	Class in PSp(4, 3).2							
	2A	2B	3A/3B	3C	3D	5A	2C	2D
Size	45	270	80	240	480	5184	36	540
$V_4(4^k)$	3	2	(3, 1)	(2 ²)	(2, 1 ²)	(1 ⁴)	–	–
$V_6(2^k)$	4	4	(3 ²)	(4, 1 ²)	(2 ³)	(2, 1 ⁴)	5	3
$V_{14}(2^k)$	8	8	(8, 3 ²)	(6, 4 ²)	(5 ² , 4)	(3 ⁴ , 2)	10	8
$V_5(3^k)$	(4, 1)	(3, 2)	3	3	3	(1 ⁵)	(4, 1)	(3, 2)
$V_{10}(3^k)$	(6, 4)	(6, 4)	6	4	4	(2 ⁵)	(6, 4)	(6, 4)
$V_{14}(3^k)$	(10, 4)	(8, 6)	6	6	6	(3 ⁴ , 2)	(10, 4)	(8, 6)
$V_4(3^k)$	(2 ²)	(2 ²)	3	2	2	(1 ⁴)	(2 ²)	(2 ²)
$V_{16}(3^k)$	(8 ²)	(8 ²)	8	6	6	(4, 3 ⁴)	(8 ²)	(8 ²)
$V_4(25^k)$	(2 ²)	(2 ²)	(1, 3)	(2 ²)	(2, 1 ²)	1	–	–
$V_5(25^k)$	(4, 1)	(3, 2)	(2, 3)	(1, 2 ²)	(3, 1 ²)	1	–	–
$V_6(5^k)$	(4, 2)	(4, 2)	(3 ²)	(4, 1 ²)	(2 ³)	2	(5, 1)	(3 ²)
$V_{10}(25^k)$	(6, 4)	(6, 4)	(1, 6, 3)	(4, 3 ²)	(4, 3 ²)	2	–	–
$V_{15a}(5^k)$	(8, 7)	(8, 7)	(9, 3 ²)	(7, 4 ²)	(5 ³)	3	(10, 5)	(9, 6)
$V_{15b}(5^k)$	(11, 4)	(9, 6)	(3, 6 ²)	(5 ³)	(7, 4 ²)	3	(10, 5)	(8, 7)

TABLE 6.11. Eigenspace dimensions of projective prime order elements in G .

Prime r_0	$i_{r_0}(H)$	$\alpha(x) \leq$
2	1424671743	11 (transvection), 6 (O/W)
3	107929774592	6
5	13789672833024	3
7	4923662008320	2
11	751977470361600	2
17	486573657292800	2
31	2401476437606400	2

TABLE 6.12. Information about elements of prime order in $H = \text{PSp}_{10}(2)$.

Therefore, if G has no regular orbit on V , then by Proposition 3.1v),

$$\frac{1}{2}r^d \leq (i_7(H) + i_{11}(H) + i_{17}(H) + i_3 1(H))r^{\lfloor d/2 \rfloor} + i_5(H)r^{\lfloor 2d/3 \rfloor} + i_3(H)r^{\lfloor 5d/6 \rfloor} + i_2(H)r^{\lfloor 5d/6 \rfloor} + 1023r^{\lfloor 10d/11 \rfloor}$$

This gives a contradiction for all V , so G has a regular orbit in all cases.

Now suppose that $E(G)/Z(E(G)) \cong \text{PSp}_8(2)$. By Proposition 3.1v), we only need to further consider modules listed in Table 6.13, which was compiled with help from [21].

Dimension	Characteristic	$r \leq$
35	$\neq 2$	106
50	3	9
51	$\neq 2, 3$	17
85	$\neq 2, 3$	5
118	3, 5	3
135	$\neq 2, 17$	3

TABLE 6.13. Absolutely irreducible modules to consider for $E(G)/Z(E(G)) \cong \text{PSp}_8(2)$

We summarise our analysis of the relevant modules $V = V_d(r)$ with dimension greater than 35 in Table 6.14. Entries where $o(x) \nmid r$ were derived from the Brauer table if the entry is bold, and from Proposition 3.1i) otherwise. Entries where $o(x) \mid r$ were derived from Proposition 3.1i) if $d \geq 85$ and

from a construction in GAP otherwise. Applying Proposition 3.1iv), we see that G has a regular orbit on V in all cases.

	$o(x) = 2$	Transvections	$o(x) = 3$	$o(x) = 5$	$o(x) = 7$	$o(x) = 17$
# elements	1371135	255	16461440	171085824	1128038400	5573836800
$V_{135}(3)$	115	120	115	31	31	31
$V_{118}(3)$	101	104	101	30	30	30
$V_{85}(5)$	64	64	43	72	43	43
$V_{51}(r), r \in \{5, 7, 17\}$	36	36	21	21 – $10\epsilon_5$	21 – $12\epsilon_7$	21 – $18\epsilon_{17}$
$V_{50}(r) r = 3, 9$	36	36	22	22	22	22

TABLE 6.14. Upper bounds on $\mathcal{E}_{\max}(g)$ for prime order g acting on some absolutely irreducible modules of $\text{PSP}_8(2)$.

Finally, suppose that $V = (35, r)$ for $r \leq 106$. We will give the details here for this module in characteristic 3, the other cases are similar. Constructing the representation in GAP, we find that elements in classes $3A, 3B, 3C, 3D$ have fixed point spaces of dimensions 21, 15, 13 and 13 respectively. This, along with the Brauer character of the representation allows us to deduce that if G has no regular orbit on V , then

$$r^{35} \leq |2A|(r^{28} + r^7) + (|2B| + |2F|)(r^{19} + r^{16}) + |2C|(r^{23} + r^{12}) + (|2D| + |2E|)(r^{20} + r^{15}) + |7A|(7r^5) \\ + |5A|(r^{11} + 4r^6) + |5B|(5r^7) + (|17A| + |17B|)(r^3 + 16r^2) + |3A|r^{21} + |3B|r^{15} + (|3C| + |3D|)r^{13}$$

and this gives a contradiction for $r \geq 3$.

Next suppose that $E(G)/Z(E(G)) \cong \text{PSP}_6(2)$. We first compute that the number of prime order elements in $G/Z(G)$ is at most 277199. Therefore, by Propositions 2.5 and 3.1v), if G has no regular orbit on $V = V_d(r)$,

$$r^d \leq 2 \times 277199r^{\lfloor \frac{5d}{6} \rfloor} + 2 \times 63r^{\lfloor \frac{6d}{7} \rfloor}$$

After inspecting [21], we find that we only need to further consider the absolutely irreducible modules that lie in Table 6.16 for $d \geq 21$, and Table 6.17 for $d < 21$.

In both tables we also present information about eigenspaces of elements of projective prime order on these remaining modules. In Table 6.16, upper bounds on \mathcal{E}_{\max} for elements of projective prime order are given. The boldface entries were obtained from the corresponding Brauer characters, while the italicised entries were obtained from explicit constructions in GAP. The remainder were obtained from Proposition 3.1 i). In Table 6.17, we give explicit eigenspace dimensions computed from the corresponding Brauer characters, and for elements whose order divides r , we obtain the dimension of their fixed point spaces from an explicit construction in GAP. Applying Proposition 3.1iv) in each case, we see that G has a regular orbit on V except for the cases given in Table 6.15, where the base size is computed from an explicit construction of the module in either GAP or MAGMA.

	$V_{14}(3)$	$V_8(3)$	$V_8(9)$	$V_8(5)$	$V_8(7)$	$V_7(3)$	$V_7(3)$	$V_7(9)$	$V_7(27)$	$V_7(81)$	$V_7(5)$	$V_7(25)$	$V_7(7)$	$V_7(49)$
$\left\lceil \frac{\log G }{\log V } \right\rceil$	1	2	1	2	1	2	2	1	1	1	1	1	1	1
$b(G)$	2	3	2	2	2	3	4	2	1	1	2	1	2	1

TABLE 6.15. Some information about $b(G)$ for small absolutely irreducible modules for $E(G)/Z(E(G)) \cong \text{PSP}_6(2)$.

This concludes our analysis of almost quasisimple G with $E(G)/Z(E(G)) \cong \text{PSP}_6(2)$.

$E(G)$	Module V	$o(x) = 2$	Transvections	$o(x) = 3$	$o(x) = 5$	$o(x) = 7$
2.PSp ₆ (2)	$V_{56}(3)$	28	28	46	28	28
PSp ₆ (2)	$V_{49}(3)$	34	34	40	34	34
2.PSp ₆ (2)	$V_{48}(3)$	24	24	<i>20</i>	24	24
2.PSp ₆ (2)	$V_{48}(5)$	24	24	24	40	24
PSp ₆ (2)	$V_{35}(7)$	25	25	25	25	28
PSp ₆ (2)	$V_{35}(5)$	25	25	25	17	25
PSp ₆ (2)	$V_{35}(r), r = 3, 9$	20	20	<i>15</i>	20	20
PSp ₆ (2)	$V_{34}(r), r = 3, 9$	24	24	<i>14</i>	7	7
PSp ₆ (2)	$V_{27}(5)$	17	21	9	7	9
PSp ₆ (2)	$V_{27}(r), r = 3, 9$	17	21	<i>15</i>	9	9
PSp ₆ (2)	$V_{26}(7)$	20	20	14	14	13
PSp ₆ (2)	$V_{21}(7)$	13	16	11	11	<i>3</i>
PSp ₆ (2)	$V_{21}(5^k), k \in \{1, 2\}$	13	16	11	5	11
PSp ₆ (2)	$V_{21}(3^k)$	12	15	<i>11</i>	5	5
# elements		5103	63	16352	48384	207360

TABLE 6.16. Upper bounds on \mathcal{E}_{\max} of projectively prime order elements in PSp₆(2) for remaining modules with $d \geq 21$.

	2A	2B	2C	2D	3A	3B	3C	5A	7A
Size	63	315	945	3780	672	2240	13440	48384	207360
$V_{15}(5^k)$	(10,5)	(11,4)	(9,6)	(8,7)	(5 ³)	(6 ² , 3)	(7, 4 ²)	7	(3, 2 ⁶)
$V_{15}(7^k)$	(10,5)	(11,4)	(9,6)	(8,7)	(5 ³)	(6 ² , 3)	(7, 4 ²)	(3 ⁵)	7
$V_{14}(3^k)$	(10,4)	(10,4)	(8,6)	(8,6)	6	6	6	(3 ⁴ , 2)	(2 ⁷)
$V_8(3^k)$	(2 ⁴)	(4 ²)	(2 ⁴)	(4 ²)	4	4	4	(2 ⁴)	(2, 1 ⁶)
$V_8(5^k)$	(2 ⁴)	(4 ²)	(2 ⁴)	(4 ²)	(4 ²)	(3 ² , 2)	(4, 2 ²)	2	(2, 1 ⁶)
$V_8(7^k)$	(2 ⁴)	(4 ²)	(2 ⁴)	(4 ²)	(4 ²)	(3 ² , 2)	(4, 2 ²)	(2 ⁴)	2
$V_7(3^k)$	(6, 1)	(4, 3)	(5, 2)	(4, 3)	5	3	3	(3, 1 ⁴)	(1 ⁷)
$V_7(5^k)$	(6, 1)	(4, 3)	(5, 2)	(4, 3)	(5, 1 ²)	(3 ² , 1)	(3, 2 ²)	3	(1 ⁷)
$V_7(7^k)$	(6, 1)	(4, 3)	(5, 2)	(4, 3)	(5, 1 ²)	(3 ² , 1)	(3, 2 ²)	(3, 1 ⁴)	1

TABLE 6.17. Eigenspace dimensions of projectively prime order elements for $E(G)/Z(E(G)) \cong \text{PSp}_6(2)$.

Now let $E(G)/Z(E(G)) \cong \text{PSp}_4(4)$. The number of prime order elements in PSp₄(4).4 is 299199. Therefore, if G has no regular orbit on the irreducible module $V = V_d(r)$,

$$r^d \leq 2 \times 299199r^{\lfloor 4d/5 \rfloor}$$

and this gives a contradiction for all choices of d, r except for those listed in Table 6.18. In this table, we also provide a set of upper bounds for \mathcal{E}_{\max} for elements of prime order. As before, the bold entries are obtained from the corresponding Brauer characters, the italicised entries are obtained from a construction in GAP, and the remainder originate from Proposition 3.1i). Applying Proposition 3.1iv) or ii) in each case implies the existence of a regular orbit for all choices of G preserving each module V . The only exception is $V = V_{18}(3)$. Here we compute in GAP that PSp₄(4) has 14 regular orbits on V . If PSp₄(4).4 has no regular orbit on V , all of these elements of regular orbits of PSp₄(4) must lie in the fixed point spaces of elements in class 2D. Therefore, $14|\text{PSp}_4(4)| \leq |2D|(2 \times 3^9)$. This is false, so PSp₄(4).4 has a regular orbit on V .

	2A/2B	2C	2D	$o(x) = 3$	$o(x) = 5$	$o(x) = 17$
# elements	255 (each)	3825	1360	10880	52224	230400
$V_{50}(3)$	30	30	30	37	30	30
$V_{34}(3)$	22	22	22	25	10	10
$V_{33}(5)$	21	21	21	13	9	13
$V_{18}(3^k), k \leq 3$	12	10	9	6	6	6
$V_{18}(5^k), k = 1, 2$	12	10	9	6	6	6
$V_{18}(7)$	12	10	9	6	6	2

TABLE 6.18. Bounds on \mathcal{E}_{\max} for prime order elements in $\mathrm{PSP}_4(4)$.

□

7. PROOF OF THEOREM 1.1: ORTHOGONAL GROUPS

7.1. Odd dimensional orthogonal groups.

Theorem 7.1. *Suppose G is an almost quasisimple group with $E(G)/Z(E(G)) \cong \mathrm{P}\Omega_{2m+1}(q)$, with $m \geq 3$ and q odd. Let $V = V_d(r)$, $(r, q) = 1$ be a module for G , with absolutely irreducible restriction to $E(G)$. Also suppose that $(r, |G|) > 1$. Then G has a regular orbit on V .*

Proof. Let V be a d -dimensional irreducible $\mathbb{F}_r G$ -module. If $q > 3$, then by Proposition 3.1v) with $|G| < q^{2m^2+m+2}$, $\alpha(G) \leq 2m+1$ from Proposition 2.6 and $d_1(G) \geq (q^{2m}-1)/(q^2-1)-2$ by Proposition 2.1, we see that G always has a regular orbit on an absolutely irreducible V . Similarly Proposition 3.1v) also shows that if $q = 3$ and $m > 3$ with identical $|G|$ and $\alpha(G)$, and $d_1(G) \geq (q^m-1)(q^m-q)/(q^2-1)$, then G has a regular orbit on V .

Finally, suppose G has $\mathrm{soc}(G/Z(G)) = \mathrm{P}\Omega_7(3)$. By Proposition 2.6, if G has no regular orbit on V , then

$$r^d \leq 2 \times |\mathrm{P}\Omega(7, 3).2| r^{\lfloor \frac{5d}{6} \rfloor} + 2i_2(G/Z(G)) r^{\lfloor \frac{6d}{7} \rfloor} \quad (6)$$

Using [21] we deduce that G has a regular orbit on V except possibly in one of the cases in Table 7.1, or $E(G) = 3.\mathrm{P}\Omega_7(3)$ on $V_{27}(4^k)$, which we handle separately.

For conciseness, we also summarise the analysis of all cases except the action of $3.\mathrm{P}\Omega_7(3)$ on $V_{27}(4^k)$ in Table 7.1. In the table, the bold entries indicate eigenspace bounds computed using the corresponding Brauer character, the italicised entries were computed from a construction in GAP, and the remaining entries originate from determining $\alpha(g)$ for g in the collection, and subsequently applying Proposition 3.1i). Applying Proposition 3.1iv) with the values given in Table 7.1 shows that G has

	Projective class									
	2A	2B	2C	2D	2E	2F	$o(x) = 3$	$o(x) = 5$	$o(x) = 7$	$o(x) = 13$
# elements	351	22113	331695	378	44226	265356	5307848	38211264	327525120	705438729
$V_{104}(2)$	89	78	78	89	78	78	50	50	50	50
$V_{90}(r), r = 2, 4$	77	67	67	77	67	67	42	42	42	42
$V_{78}(r), r = 2, 4$	<i>56</i>	<i>56</i>	<i>56</i>	<i>56</i>	<i>56</i>	<i>56</i>	36	36	36	36
$V_{78}(5)$	56	56	56	56	56	56	36	65	36	36
$V_{27}(13^k)$	21	17	17	–	–	–	15	15	15	13
$V_{27}(7^k)$	21	17	17	–	–	–	15	15	13	15
$V_{27}(25^k)$	21	17	17	–	–	–	15	18	15	15

TABLE 7.1. Sizes and \mathcal{E}_{\max} upper bounds of elements of collections of projectively prime order elements in $\mathrm{P}\Omega_7(3).2$.

a regular orbit on V in all cases.

Finally, suppose that $V = V_{27}(4^k)$. When we construct V in GAP, we find that involutions in classes 2A, 2B and 2C have 21-, 17- and 5-dimensional fixed point spaces respectively. This, along

with information from the corresponding Brauer character imply that if G has no regular orbit on V , $r^{27} \leq |2A|r^{21} + |2B|r^{17} + |2C|r^{15} + i_3(G)(3r^9) + |3B|(r^{15} + 2r^6) + |5A|(r^7 + 4r^5) + |7A|(r^3 + 6r^4) + \frac{13}{12}i_{13}(G)r^3$ and this is a contradiction for $r \geq 4$, so G has a regular orbit on V . \square

7.2. Even dimensional orthogonal groups.

Theorem 7.2. *Suppose G is an almost quasisimple group with $E(G)/Z(E(G)) \cong \text{P}\Omega_{2m}^\epsilon(q)$ with $m \geq 4$. Let $V = V_d(r)$, $(r, q) = 1$ be a module for G , with absolutely irreducible restriction to $E(G)$. Also suppose that $(r, |G|) > 1$. Then one of the following holds.*

- i) $b(G) = \lceil \log |G| / \log |V| \rceil$, or
- ii) $E(G)/Z(E(G)) \cong \text{P}\Omega_8^+(2)$, $d = 8$, $r \in \{3, 5, 25, 27\}$. and $b(G) = \lceil \log |G| / \log |V| \rceil + 1$.

By Proposition 2.1, excluding the case $(\epsilon, m, q) = (+, 4, 2)$, we have $d_1(G) \geq \frac{(q^m-1)(q^{m-1}-q)}{q^2-1} - 1$.

Therefore, if G has no regular orbit on V then by Propositions 2.4 and 3.1,

$$r^{\frac{(q^m-1)(q^{m-1}-q)}{q^2-1} - 1} \leq 2|G/Z(G)|r^{\lfloor \frac{2m-1}{2m}d \rfloor}$$

This inequality gives a contradiction for all valid choices of (m, q, r, ϵ) except for $(5, 2, r_1, \pm)$, $(4, 3, r_2, \pm)$, and $(4, 2, r_3, -)$, where $r_1 \leq 7$, $r_2 = 2$ and $r_3 \leq 154$.

We will consider each pair (m, q) individually.

Proposition 7.3. *Theorem 7.2 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \text{P}\Omega_{10}^\epsilon(2)$.*

Proof. We first determine the range of d to consider. The number of prime order elements $i_P(\text{P}\Omega_{10}^\epsilon(2).2)$ in $\text{P}\Omega_{10}^\epsilon(2).2$ is 5642583264255 if $\epsilon = +$ and 4547966108159 if $\epsilon = -$. The number of reflections $i_R(\text{P}\Omega_{10}^\epsilon(2).2)$ is 496 and 528 for $\epsilon = +, -$ respectively. If G has no regular orbit on V then by Proposition 2.6,

$$r^d \leq 2i_P(\text{P}\Omega_{10}^\epsilon(2).2)r^{\lceil 7d/8 \rceil} + 2i_R(\text{P}\Omega_{10}^\epsilon(2).2)r^{\lceil 9d/10 \rceil}$$

So we only need consider $d \leq 216$ for $r = 3$, and G has a regular orbit on the remaining modules. From the Brauer character table, we find that we only need to consider two representations for each ϵ , one each of degrees $154 - \epsilon$ and $186 - (1 + \epsilon)/2$. Examining the Brauer characters, we determine that $\mathcal{E}_{\max}(g_{3'}) \leq 118 + (1 - \epsilon)$ and $136 - (1 + \epsilon)/2$ respectively. Moreover, by Proposition 2.6, all elements of order 3 have $\alpha(x) \leq 8$. Therefore, if G has no regular orbit on the 153-dimensional representation when $\epsilon = -$ then

$$r^{153} \leq 2i_P(G/Z(G))r^{116} + i_3(G/Z(G))r^{\lceil 7 \times 153/8 \rceil}$$

which gives a contradiction for $r = 3$. Similar calculations give the results for the other three modules. \square

Proposition 7.4. *Theorem 7.2 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \text{P}\Omega_8^\epsilon(3)$.*

Proof. Recall that we only need to consider $r = 2$. Using GAP, we compute that every element $x \in G/Z(G)$ of prime order at least 5 has $\alpha(x) = 2$. We also determine that the number $i_R(G/Z(G))$ of reflections in $G/Z(G)$ is at most 2214 and 6480 for $\epsilon = -, +$ respectively. Denote the number of prime order elements in $H = G/Z(G)$ by $i_P(H)$. If G has no regular orbit on V then

$$\frac{1}{2}r^d \leq (i_P(H) - i_2(H) - i_3(H))r^{\lfloor d/2 \rfloor} + (i_2(H) + i_3(H))r^{\lfloor 6d/7 \rfloor} + i_r(H)r^{\lfloor 7d/8 \rfloor}$$

This is a contradiction for all $d \geq 245$ (cf. Proposition 2.1) with $r = 2$. \square

Proposition 7.5. *Theorem 7.2 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong \text{P}\Omega_8^\epsilon(2)$.*

Proof. First suppose that $E(G)/Z(E(G)) \cong \text{P}\Omega_8^-(2)$. The number of prime order elements in $\text{P}\Omega_8^-(2).2$ is 57548943; of these, 112591 are involutions and of those, 136 are transvections. If G has no regular orbit on V then

$$r^d \leq 2 \times 57548943r^{\lfloor 6d/7 \rfloor} + 2 \times 136r^{\lfloor 7d/8 \rfloor}$$

Thus G has a regular orbit on V , or V lies in Table 7.2.

We also summarise our analysis of the remaining cases in Table 7.2. In all cases, the bounds on $\mathcal{E}_{\max}(x)$ are obtained from the Brauer character if $o(x) \nmid r$, and from constructing the corresponding module in GAP otherwise.

Projective prime order						
	$o(x) = 2$	Transvections	$o(x) = 3$	$o(x) = 5$	$o(x) = 7$	$o(x) = 17$
# elements	112591	136	1285200	1096704	9400320	46448640
$V_{51}(5)$	36	36	21	11	21	21
$V_{51}(7)$	36	36	21	11	9	21
$V_{50}(3^k), r \in \{1, 2\}$	36	36	22	10	10	10
$V_{34}(3^k), k \in \{1, 2, 3\}$	22	27	20	10	10	10
$V_{34}(5^k), k \in \{1, 2\}$	22	27	20	10	20	20
$V_{33}(7)$	21	26	19	19	5	19

TABLE 7.2. Maximum eigenspace dimensions on absolutely irreducible modules of $\mathrm{P}\Omega_8^-(2)$.

Using the information in Table 7.2 to apply Proposition 3.1iv), we find that G has a regular orbit on V in all cases.

Now suppose that $E(G)/Z(E(G)) \cong \mathrm{P}\Omega_8^+(2)$. The number of prime order elements in $\mathrm{P}\Omega_8^+(2)$ is 28815119 and so if G has no regular orbit on V ,

$$r^d \leq 2 \times 28815119r^{\lfloor 7d/8 \rfloor}.$$

Using [21], we give a complete list of the modules we need to consider in the Table 7.3. In Table 7.4 give information on the number of elements of various prime orders in $\mathrm{P}\Omega_8^+(2).S_3$.

d	$E(G)$	Characteristic	$r \leq$	Order	# elements
104	$2.\mathrm{P}\Omega_8^+(2)$	3,5	3	2	183375
83	$\mathrm{P}\Omega_8^+(2)$	5	5	3	2006720
56	$2.\mathrm{P}\Omega_8^+(2)$	$\neq 2$	13	5	1741824
50	$\mathrm{P}\Omega_8^+(2)$	$\neq 2, 3$	13	7	24883200
48	$\mathrm{P}\Omega_8^+(2)$	3	9		
35	$\mathrm{P}\Omega_8^+(2)$	$\neq 2$	31		
28	$\mathrm{P}\Omega_8^+(2)$	$\neq 2$	96		
8	$2.\mathrm{P}\Omega_8^+(2)$	$\neq 2$	86445357	Total	28815119

TABLE 7.3. Cases left to consider when $E(G)/Z(E(G)) \cong \mathrm{P}\Omega_8^+(2)$.

TABLE 7.4. Summary of prime order elements in $\mathrm{P}\Omega_8^+(2).S_3$.

As before, we summarise our analysis of the modules listed in Table 7.3 in a Table 7.5. The bold entries were obtained from the corresponding Brauer characters of the absolutely irreducible modules. The italicised entries were obtained from an explicit construction of the module in GAP. The entries that are both bold and italicised were obtained using the former method when $(o(x), r) = 1$, and the latter otherwise. The underlined entries were computed by restricting $V_{28} = V_{28}(3^k)$ to the maximal subgroup $H = U_3(3) : 2 \times S_3 \leq \mathrm{P}\Omega_8^+(2).S_3$ and applying Propositions 3.1i) and 3.2. The remaining entries were computed using Proposition 3.1i). Applying Proposition 3.1iv) to each of the modules in the table shows that G has a regular orbit on V for all valid choices of G .

Finally, suppose that V is the 8-dimensional irreducible representation of $2.\mathrm{P}\Omega_8^+(2)$. If $r \leq 9$, then $|V| < |G|$ and there is no regular orbit. If $r = 3$ then also $|V|^2 < |G|$, so $b(G) \geq 3$. If G has no regular orbit on V in characteristics 5 or 7 then

$$\begin{aligned} r^8 \leq & (|2A| + |2C| + |2D| + |2E|)(2r^4) + |2B|(r^6 + r^2) + |3A|(r^6 + 2r) + (|3B| + |3C|)(2r^4) \\ & + |3D|(r^2 + 2r^3) + |3E|(r^4 + 2r^2) + |5A|(r^4 + 4r) + (|5B| + |5C|)(4r^2) + |7A|(r^2 + 6r) \\ & + |2F|(r^7 + r) + |2G|(r^5 + r^3). \end{aligned}$$

Class	Size	$V_{104}(3)$	$V_{83}(5)$	$V_{56}(r)$	$V_{50}(r)$	$V_{48}(r)$	$V_{35}(r)$	$V_{28}(3^k)$	$V_{28}(5^k)$ or $V_{28}(7^k)$
2A	1575	77	72	35	35	34	20	16	16
2B	3780	77	72	35	35	34	23	16	16
2C	3780	77	72	35	35	34	20	16	16
2D	3780	77	72	35	35	34	20	16	16
2E	56700	77	72	35	35	34	20	16	16
3A	2240	89	71	$35 - 7\epsilon_3$	35	<i>26</i>	<i>21</i>	<i>16</i>	16
3B	2240	89	71	$35 - 7\epsilon_3$	35	<i>26</i>	<i>15</i>	<i>16</i>	16
3C	2240	89	71	$35 - 7\epsilon_3$	35	<i>26</i>	<i>15</i>	<i>16</i>	16
3D	89600	89	71	$35 - 7\epsilon_3$	35	<i>26</i>	<i>15</i>	<i>12</i>	10
3E	268800	89	71	$35 - 7\epsilon_3$	35	<i>26</i>	<i>15</i>	<i>12</i>	10
5A	580608	77	71	$35 - 7\epsilon_5$	25	10	<i>11</i>	8	8
5B	580608	77	71	$35 - 7\epsilon_5$	25	10	<i>11</i>	8	8
5C	580608	77	71	$35 - 7\epsilon_5$	25	10	<i>11</i>	8	8
7A	24883200	77	71	$35 - 7\epsilon_7$	25	10	<i>5</i>	7	<i>4</i>
2F	120	77	72	35	35	34	28	21	21
2G	37800	77	72	35	35	34	20	15	15
3F	14400	–	–	–	35	<i>26</i>	–	<u><i>14</i></u>	14
3G	806400	–	–	–	35	<i>26</i>	–	<u><i>14</i></u>	10

TABLE 7.5. Sizes and upper bounds on \mathcal{E}_{\max} for conjugacy classes of elements of prime order in $\text{P}\Omega_8^+(2).S_3$.

This gives a contradiction for $r > 125$. When $r = 5$, we construct the representation in MAGMA and find that $b(G) = 3$, which implies from Lemma 3.4 that $b(G) = 2$ when $r = 25$. We also compute from constructions of the modules in MAGMA that there is a regular orbit of G on $V = V_8(r)$ when $r = 49, 125$.

If instead V is in characteristic 3, and G has no regular orbit on V , then

$$r^8 \leq (|2A| + |2C| + |2D| + |2E|)(2r^4) + |2B|(r^6 + r^2) + \frac{1}{2}i_3(G)r^4 + |5A|(r^4 + 4r) + (|5B| + |5C|)(4r^2) + |7A|(r^2 + 6r) + |2F|(r^7 + r) + |2G|(r^5 + r^3).$$

This gives a contradiction for $r \geq 243$. We compute that $b(G) = 4, 2, 2, 1$ when $r = 3, 9, 27, 81$ using a combination of GAP, MAGMA and Lemma 3.4. \square

8. PROOF OF THEOREM 1.1: EXCEPTIONAL GROUPS

The main result in this section is as follows.

Theorem 8.1. *Suppose G is an almost quasisimple group with $E(G)/Z(E(G))$ an exceptional simple group of Lie type. Let $V = V_d(r)$, $(r, q) = 1$ be a module for G , with absolutely irreducible restriction to $E(G)$. Also suppose that $(r, |G|) > 1$. Then one of the following holds:*

- i) $b(G) = \lceil \log |G| / \log |V| \rceil$,*
- ii) $E(G) = 2 \cdot {}^2B_2(8)$, $(d, r) = (8, 5)$ and $b(G) = 2$,*
- iii) $G = 6 \circ (2.G_2(4))$ or $6 \circ (2.G_2(4).2)$, $(d, r) = (12, 7)$ and $b(G) = 2$.*

We now proceed with with proof of Theorem 8.1. If G has no regular orbit on $V = V_d(r)$, then by Proposition 3.1i) and v),

$$r^d \leq 2|G/Z(G)|r^{\lfloor (1-1/\alpha(G))d \rfloor}.$$

From Propositions 2.1 and 2.4 we deduce that either G has a regular orbit on V , or $E(G)/Z(E(G))$ is one of ${}^2B_2(8)$, ${}^2F_4(2)'$, ${}^3D_4(2)$, ${}^3D_4(3)$, $G_2(3)$, $G_2(4)$, $G_2(5)$ or $F_4(2)$.

We examine the remaining cases in a series of propositions.

Proposition 8.2. *Theorem 8.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong {}^2B_2(8)$.*

Proof. By [21], we see that we only need to consider representations with $(d, r) = (8, 5)$ or $(14, 5)$. Generators for G in this representation are given in [1] and we find that there is no regular orbit of G on V , but there is a base of size 2. Finally, for $(d, r) = (14, 5)$, using generators from [1], we find that $b(G) = 1$. \square

Proposition 8.3. *Theorem 8.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong {}^2F_4(2)'$.*

Proof. The number of prime order elements in ${}^2F_4(2)'.2$ is 3304079. By the proof of [17, Proposition 5.5], $\alpha(x) \leq 4$ for involutions and $\alpha(x) \leq 3$ for odd prime order elements. Moreover, $d \geq 26$ by [21]. Therefore, if G has no regular orbit on V , then by Proposition 3.1v),

$$\frac{d}{4} \leq \log_r(2 \times 3304079)$$

which gives a contradiction unless $d \leq 57$ for $r = 3$, $d \leq 39$ for $r = 5$ and $d \leq 28$ for $r = 9$. Consulting [21], we find that we have only 26- and 27-dimensional representations of ${}^2F_4(2)'$ to consider. First let $d = 27$. Examining the Brauer character tables for G , we find that $\mathcal{E}_{\max}(g_{r'}) \leq 16$. Therefore, if $r = 5$, and G has no regular orbit on V ,

$$r^{27} \leq 2 \times 3304079r^{16} + 359424r^{18}$$

which gives a contradiction, and a similar calculation gives the result for $r = 9$. When $r = 3$, the representation is not realised due to a i_1 irrationality in the corresponding Brauer character. We now consider the 26-dimensional representations. The corresponding Brauer characters contain i_2 irrationalities which do not exist for $r = 5$, so we only need consider $r = 3, 9$. Examining the Brauer character table in characteristic 3, we deduce that $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 6$ and $\mathcal{E}_{\max}(g_2) \leq 16$. We further compute in GAP that $\alpha(x) = 2$ for $x \in G$ of order 3. Therefore, since $i_2({}^2F_4(2)) = 13455$ and $i_3({}^2F_4(2)) = 166400$, if G has no regular orbit on V then

$$r^{26} \leq 2 \times 3304079r^6 + 166400r^{13} + 13455(r^{16} + r^{10})$$

which gives a contradiction for $r = 3, 9$. \square

Proposition 8.4. *Theorem 8.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong {}^3D_4(q)$, $q \in \{2, 3\}$.*

Proof. Suppose that $q = 3$. By Proposition 3.1v), we are left to examine the cases where $r = 2$, $d \leq 315$. According to [20, A.15], we only need consider the 218-dimensional representation of G . Using the construction of V in GAP, and computing that the number of prime order elements in ${}^3D_4(3) = 6101289775853$ is we deduce that if G has no regular orbit on V , then

$$r^{218} \leq (i_2(G) + i_3(G))r^{\lfloor \frac{6 \times 218}{7} \rfloor} + 6101289775853r^{32}$$

and this gives a contradiction for $r = 2$. Now suppose that $q = 2$. By [21] and Proposition 3.1v), we must consider $d = 25, 26, 52$ for $r \leq 102, 102$ and 9 respectively. Note that if $d = 25$ then $r = 3^k$. The numbers of prime order elements in ${}^3D_4(2).3$ are given in Table 8.1. We split the arguments

Prime r_0	2	3	7	13	Total
Number of elements	69615	2457728	4852224	48771072	56150639

TABLE 8.1. Prime order elements in ${}^3D_4(2).3$.

up by characteristic. In characteristic 13, we only need consider $V = V_{26}(13)$. From the Brauer character we observe that $\mathcal{E}_{\max}(g_{13'}) \leq 16$. Moreover, in GAP, we find that elements g of order 13 have $\dim C_V(g) = 2$. Therefore, if G has no regular orbit on V ,

$$r^{26} \leq 2 \times 56150639r^{16} + 48771072r^2.$$

This is a contradiction for $r = 13$. We now consider representations in characteristic 7. We need to consider $d = 26$ for $r = 7, 49$ and $d = 52$ for $r = 7$. We find that when $d = 26$, we have $\mathcal{E}_{\max}(g_7) \leq 8$ and $\mathcal{E}_{\max}(g_{7'}) \leq 16$. On the other hand, when $d = 52$, $\mathcal{E}_{\max}(g_7) \leq 44$ and $\mathcal{E}_{\max}(g_{7'}) \leq 36$. Therefore, if G has no regular orbit on one of these modules,

$$r^d \leq 2 \times 56150639r^{\mathcal{E}_{\max}(g_{7'})} + 4852224r^{\mathcal{E}_{\max}(g_7)}.$$

This gives a contradiction in both cases for $r \geq 7$. Now suppose V is an irreducible module in characteristic 3. Here we must consider $d = 52$ for $r = 3, 9$, and $d = 25$ for $r \leq 81$. The classes of elements of order 3 in ${}^3D_4(2).3$ and their sizes are given in Table 8.2.

Class	3A	3B	3C	3C'	3D	3D'
Size	139776	326144	17472	17472	978432	978432

TABLE 8.2. Orders of conjugacy classes of elements of order 3 in ${}^3D_4(2).3$.

Let V be the irreducible 52-dimensional representation. From the Brauer character we see that $\mathcal{E}_{\max}(g_2) \leq 36$, and $\mathcal{E}_{\max}(g_{\{2,3\}'}) \leq 10$. This, in addition to dimensions of fixed point spaces of elements of order 3 computed in GAP show that if G has no regular orbit on V ,

$$r^{52} \leq 2 \times 56150639r^{10} + 2 \times 69615r^{36} + 2 \times 978432r^{18} + (139776 + 2 \times 17472)r^{22} + 326144r^{18}.$$

This is a contradiction for $r \geq 3$, so G has a regular orbit on V . Finally, suppose that $d = 25$. From the Brauer character we observe that $\mathcal{E}_{\max}(g_2) \leq 16$, and that $\mathcal{E}_{\max}(g_{7,13}) \leq 7$. Constructing the representation in GAP, we observe that elements in classes 3A and 3B have 9-dimensional fixed point spaces. We see that elements in classes 3C and 3D appear in the maximal subgroup $H = S_3 \times L_2(8).3 < {}^3D_4(2).3$. Now $V \downarrow H$ has four trivial composition factors, and three composition factors of degree 7. Elements of order 3 have at most a 3-dimensional fixed point space on the 7-dimensional composition factor, so by Proposition 3.2, $\dim C_V(g) \leq 4 \times 1 + 3 \times 3 = 13$ for an element in class 3C or 3D. Since 3C' and 3D' are the squares of elements in classes 3C, 3D, they have the same eigenspaces, so we need not consider them in our sum.

Therefore, if G has no regular orbit on V ,

$$r^{25} \leq 2 \times 56150639r^7 + 2 \times 69615r^{10} + (17472 + 978432)r^{13} + (139776 + 326144)r^9$$

This gives a contradiction for $r \geq 9$. If $r = 3$, then we prove using a construction of the module in MAGMA that G has a regular orbit on V . \square

Proposition 8.5. *Theorem 8.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong G_2(q)$, with $3 \leq q \leq 5$.*

Proof. First suppose $E(G)/Z(E(G)) = G_2(5)$. The number of elements of prime order in $G_2(5)$ is 1242340499, and so by Proposition 3.1v) and [21], we only need to consider the irreducible 124-dimensional module for G over \mathbb{F}_2 . Examining the Brauer character, we find that $\mathcal{E}_{\max}(g_{2'}) \leq 48$. Therefore, along with our observation that there are 406875 involutions in $G_2(5)$, we find that if G has no regular orbit on this module then

$$2^{124} \leq |G_2(5)|2^{48} + 406875 \times 2^{\lfloor \frac{4}{5} \times 124 \rfloor}$$

which gives a contradiction. Therefore, G has a regular orbit on V .

We now consider G with $\text{soc}(G/Z(G)) = G_2(4)$. The number of elements of prime order in $G_2(4).2$ is 55534959. Using the same arguments as above along with [21], we see that we only need consider modules of dimension 12, 64, 78 if $r = 3$, and $d = 12$ for $5 \leq r \leq 484$. If V is the 78-dimensional irreducible module of $G_2(4).2$, $\mathcal{E}_{\max}(g_{3'}) \leq 46$. Therefore, since the number of 3-elements in $G_2(4).2$ is 1401920, if G has no regular orbit on V ,

$$3^{78} \leq 2 \times 55534959 \times 3^{46} + 1401920 \times 3^{\lfloor 4 \times 78/5 \rfloor}$$

which gives a contradiction. Similarly, if $d = 64$, then $\mathcal{E}_{\max}(g_{3'}) \leq 36$, and an analogous calculation shows that G has a regular orbit on V here. Finally, we consider the 12-dimensional representation of $2.G_2(4)$ for $r \leq 484$. This extends to a representation of $2.G_2(4).2$. If $r = 3, 5$ then $|G| > |V|$ and G has no regular orbit on V . When $r = 3$, constructing the representation in GAP we find that there is a base of size 2. For the remaining cases, we will summarise our analysis in Table 8.3. The entries in the third column were obtained from the corresponding Brauer tables, while the entries in the fourth were derived from constructions in GAP. Applying Proposition 3.1iv) with the entries in the table, we see that G has a regular orbit on V for $r \geq 9$ and a base of size two for $r = 5$. When $r = 7$, we determine using orbit enumeration techniques in MAGMA that $2.G_2(4)$ and $2.G_2(4).2$ have regular orbits on V , while $6 \circ (2.G_2(4))$ and $6 \circ (2.G_2(4).2)$ do not, and so $b(G) = 2$ by Proposition 3.4.

Proj. Class	Size	Eigenspace Dimensions	
		$o(x) \nmid r$	$o(x) \mid r$
2A	4095	(8,4)	–
2B	65520	(6 ²)	–
3A	4160	(6 ²)	6
3B	1397760	(4 ³)	4
5A	838656	(4, 2 ⁴)	4
5B	838656	(4, 2 ⁴)	4
5C	838656	(3 ⁴)	3
5D	838656	(3 ⁴)	3
7A	11980800	(2 ⁶)	2
13A	19353600	(1 ¹²)	1
13B	19353600	(1 ¹²)	1

TABLE 8.3. Eigenspace dimensions of projectively prime order elements in $2.G_2(4).2$

Finally, we investigate G with $\text{soc}(G/Z(G)) \cong G_2(3)$. The number of prime order elements in $G_2(3).2$ is 1329587. Using analogous calculations to those above and inspecting [21], we deduce that G has a regular orbit on V unless possibly if $r = 2$ and $d = 64, 78, 90$, $r = 2, 4, 7$ and $d = 27$, or $d = 14$ and $r \leq 128$. By the proof of [17, Proposition 5.6], $\alpha(x) \leq 3$ if x is an inner involution, $\alpha(x) \leq 5$ if x is of order 3, and $\alpha(x) \leq 4$ otherwise. Therefore, if G has no regular orbit on the 90-dimensional representations of $G_2(3)$, then

$$r^{90} \leq 2 \times 1326779r^{[3 \times 90/4]} + 2 \times 59696r^{[4 \times 90/5]} + 10179r^{[2 \times 90/3]}$$

which is a contradiction for $r = 2$. If V is the 78-dimensional \mathbb{F}_2G -module, then examining the Brauer character we find that $\mathcal{E}_{\max}(g_{2'}) \leq 30$. So if G has no regular orbit on V ,

$$r^{78} \leq 2 \times 1326779r^{30} + 10179r^{[2 \times 78/3]}$$

which again is a contradiction for $r = 2$. The 64-dimensional representations of $G_2(3)$ in characteristic 2 are not realised over \mathbb{F}_2 due to a b_{27} irrationality. Similarly, the 27-dimensional representation of $3.G_2(3)$ is not realised over \mathbb{F}_2 because of a z_3 irrationality. If $d = 27$ and $r = 4$ or 7 , then from the Brauer character we determine that $\mathcal{E}_{\max}(g_{\{2,p\}'}) \leq 9$, so if G has no regular orbit on V ,

$$r^{27} \leq 2 \times 1326779r^9 + 2i_2(G/Z(G))r^{[3 \times 27/4]} + i_7(G/Z(G))r^{[3 \times 27/4]}$$

which gives a contradiction for $r = 4, 7$. It therefore remains to consider the 14-dimensional representation of $G_2(3).2$ for $r \leq 128$. If $r = 2$, then $|V| < |G|$ and we compute in GAP that $b(G) = 2$. We also compute that if $r = 4$, then G has a regular orbit on V . Otherwise, if G has no regular orbit on

	$o(x) = 2$	$o(x) = 3$	$o(x) = 7$	$o(x) = 13$
Number of elements	10179	59696	606528	653184
Largest eigsp. in char. 2	8	8	2	2
Largest eigsp. in char. 7	8	8	2	2
Largest eigsp. in char. 13	8	8	2	2

TABLE 8.4. Information about projectively prime order elements acting on an absolutely irreducible 14-dimensional module for $G_2(3).2$.

V then by Proposition 3.iv) and the information in Table 8.4,

$$r^{14} \leq 2(10179r^8 + 606528r^2 + 653184r^2) + 59696(r^8 + 2r^3)$$

which is a contradiction for $r \geq 7$. □

Proposition 8.6. *Theorem 8.1 holds for almost quasisimple G with $E(G)/Z(E(G)) \cong F_4(2)$.*

Proof. The number of prime order elements in $F_4(2).2$ is 650870214978815. So if G has no regular orbit on V then

$$r^d \leq 2 \times 650870214978815r^{\lfloor 7d/8 \rfloor}$$

which gives a contradiction unless $d \leq 248$. Examining [21], we find that we only need to consider the 52-dimensional module of $2.F_4(2)$. We record the classes of elements of projective prime order in $2.F_4(2).2$ and their eigenspace dimensions in cases where they are semisimple in Table 8.5. For each characteristic dividing the order of $2.F_4(2).2$, we also compute a bound of the fixed point space dimensions on unipotent elements and these are also recorded in Table 8.5. Applying Proposition

Class in $F_4(2).2$	Eigensp. dims			
2A	(36,16)			
2B	(32,20)			
2C	(28,24)			
3A	(22, 15 ²)	3	22	Construction in GAP
3B	(22, 15 ²)	5	12	Construction in GAP
3C	(16, 18 ²)	7	10	Restrict V to $\mathrm{PSp}_8(2)$
5A	(12, 10 ⁴)	13	4	Restrict V to ${}^2F_4(2)$
7A	(10, 7 ⁶)	17	35	Restrict V to $\mathrm{PSp}_4(4)$
13A	(4 ¹³)			
17A	(4, 3 ¹⁶)			
2D	(26 ²)			

TABLE 8.5. Eigenspace dimensions of projectively prime order elements on an absolutely irreducible 52-dimensional module.

3.1iv), we deduce that G has a regular orbit on V in each case. \square

This completes the proof of Theorem 8.1, and hence Theorem 1.1.

ACKNOWLEDGEMENTS

This paper represents part of the PhD work of the author under the supervision of Professor Martin W. Liebeck, and the author would like to thank Professor Liebeck for his guidance and careful reading of this paper. The author acknowledges the support of an EPSRC International Doctoral Scholarship at Imperial College London. The author would also like to thank Professors Eamonn O'Brien and Jürgen Müller for aiding with a number of computationally intensive cases included in this paper, and Dr. Tim Burness for providing some helpful suggestions.

REFERENCES

- [1] Rachel Abbott, John Bray, Steve Linton, Simon Nickerson, Simon Norton, Richard Parker, Stephen Rogers, Ibrahim Suleiman, Jonathan Tripp, Peter Walsh, and Robert Wilson. ATLAS of Finite Group Representations – Version 3. <http://brauer.maths.qmul.ac.uk/Atlas/v3/>. Accessed: 29/7/2019.
- [2] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [3] Timothy C. Burness and Michael Giudici. *Classical groups, derangements and primes*, volume 25 of *Australian Mathematical Society Lecture Series*. Cambridge University Press, Cambridge, 2016.
- [4] Timothy C. Burness, Robert M. Guralnick, and Jan Saxl. On base sizes for symmetric groups. *Bull. Lond. Math. Soc.*, 43(2):386–391, 2011.
- [5] Timothy C. Burness, Robert M. Guralnick, and Jan Saxl. Base sizes for \mathcal{S} -actions of finite classical groups. *Israel J. Math.*, 199(2):711–756, 2014.
- [6] Timothy C. Burness, E. A. O'Brien, and Robert A. Wilson. Base sizes for sporadic simple groups. *Israel J. Math.*, 177:307–333, 2010.
- [7] Joanna B. Fawcett. The base size of a primitive diagonal group. *J. Algebra*, 375:302–321, 2013.
- [8] Joanna B. Fawcett, Jürgen Müller, E. A. O'Brien, and Robert A. Wilson. Regular orbits of sporadic simple groups. *J. Algebra*, 522:61–79, 2019.
- [9] Joanna B. Fawcett, E. A. O'Brien, and Jan Saxl. Regular orbits of symmetric and alternating groups. *J. Algebra*, 458:21–52, 2016.
- [10] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.9.3*, 2018.

- [11] David Gluck, Kay Magaard, Udo Riese, and Peter Schmid. The solution of the $k(GV)$ -problem. *J. Algebra*, 279(2):694–719, 2004.
- [12] Dominic P. M. Goodwin. Regular orbits of linear groups with an application to the $k(GV)$ -problem. I. *J. Algebra*, 227(2):395–432, 2000.
- [13] Dominic P. M. Goodwin. Regular orbits of linear groups with an application to the $k(GV)$ -problem.II. *J. Algebra*, 227(2):433–473, 2000.
- [14] Dominic P.M. Goodwin. *Regular Orbits of Linear Groups*. PhD thesis, Imperial College London, 1998.
- [15] R. M. Guralnick and R. Lawther. Generic stabilizers in actions of simple algebraic groups I: modules and the first Grassmanian varieties. *arXiv e-prints*, page arXiv:1904.13375, Apr 2019.
- [16] Robert M. Guralnick, Kay Magaard, Jan Saxl, and Pham Huu Tiep. Cross characteristic representations of symplectic and unitary groups. *J. Algebra*, 257(2):291–347, 2002.
- [17] Robert M. Guralnick and Jan Saxl. Generation of finite almost simple groups by conjugates. *J. Algebra*, 268(2):519–571, 2003.
- [18] Paul Gérardin. Weil representations associated to finite fields. *Journal of Algebra*, 46(1):54 – 101, 1977.
- [19] J. I. Hall, Martin W. Liebeck, and Gary M. Seitz. Generators for finite simple groups, with applications to linear groups. *Quart. J. Math. Oxford Ser. (2)*, 43(172):441–458, 1992.
- [20] Frank Himstedt. On the 2-decomposition numbers of Steinberg’s triality groups ${}^3D_4(q)$, q odd. *J. Algebra*, 309(2):569–593, 2007.
- [21] Gerhard Hiss and Gunter Malle. Low-dimensional representations of quasi-simple groups. *LMS J. Comput. Math.*, 4:22–63, 2001.
- [22] Gerhard Hiss and Gunter Malle. Low-dimensional representations of special unitary groups. *J. Algebra*, 236(2):745–767, 2001.
- [23] I. M. Isaacs. Characters of solvable and symplectic groups. *Amer. J. Math.*, 95:594–635, 1973.
- [24] J. P. James. Partition actions of symmetric groups and regular bipartite graphs. *Bull. London Math. Soc.*, 38(2):224–232, 2006.
- [25] Christoph Jansen, Klaus Lux, Richard Parker, and Robert Wilson. *An atlas of Brauer characters*, volume 11 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 1995. Appendix 2 by T. Breuer and S. Norton, Oxford Science Publications.
- [26] Christoph Köhler and Herbert Pahlings. Regular orbits and the $k(GV)$ -problem. In *Groups and computation, III (Columbus, OH, 1999)*, volume 8 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 209–228. de Gruyter, Berlin, 2001.
- [27] Ross Lawther, Martin W. Liebeck, and Gary M. Seitz. Fixed point ratios in actions of finite exceptional groups of Lie type. *Pacific J. Math.*, 205(2):393–464, 2002.
- [28] Martin W. Liebeck. Regular orbits of linear groups. *J. Algebra*, 184(3):1136–1142, 1996.
- [29] Martin W. Liebeck and Aner Shalev. Simple groups, permutation groups, and probability. *J. Amer. Math. Soc.*, 12(2):497–520, 1999.
- [30] Pham Huu Tiep. Low dimensional representations of finite quasisimple groups. *Groups, Combinatorics and Geometry, Durham*, pages 277–294, 2001.
- [31] Pham Huu Tiep. Low dimensional representations of finite quasisimple groups. In *Groups, combinatorics & geometry (Durham, 2001)*, pages 277–294. World Sci. Publ., River Edge, NJ, 2003.
- [32] Pham Huu Tiep and Alexander E. Zalesskii. Some characterizations of the Weil representations of the symplectic and unitary groups. *J. Algebra*, 192(1):130–165, 1997.
- [33] Hans Zassenhaus. Über endliche Fastkörper. *Abh. Math. Sem. Univ. Hamburg*, 11(1):187–220, 1935.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE, LONDON SW7 2BZ, UK

E-mail address: m.lee16@imperial.ac.uk