

Haar wavelets collocation on a class of Emden-Fowler equation via Newton's quasilinearization and Newton-Raphson techniques

Amit K. Verma^a, Narendra Kumar^b

^{a,b} *Department of Mathematics,
Indian Institute of Technology Patna,
Bihta, Patna 801103, (BR) India.*

November 19, 2019

Abstract

In this paper we have considered generalized Emden-Fowler equation,

$$y''(t) + \sigma t^\gamma y^\beta(t) = 0, \quad t \in]0, 1[$$

subject to the following boundary conditions

$$y(0) = 1, \quad y(1) = 0; \quad \& \quad y(0) = 1, \quad y'(1) = y(1),$$

where γ, β and σ are real numbers, $\gamma < -2$, $\beta > 1$. We proposed to solve the above singular nonlinear BVPs with the aid of Haar wavelet coupled with quasilinearization approach as well as Newton-Raphson approach. We have also considered the special case of Emden-Fowler equation ($\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$) which is popularly known as Thomas-Fermi equation. We have analysed different cases of generalised Emden-Fowler equation and compared our results with existing results in literature. We observe that small perturbations in initial guesses does not affect the the final solution significantly.

Keywords: Haar wavelet; Emden-Fowler equation; Thomas-Fermi equation; Quasilinearization; Newton-Raphson method

AMS Subject Classification: 34B15, 34B16, 65Z05

1 Introduction

The following singular boundary value problem,

$$y''(t) = t^{-\frac{1}{2}} y^{\frac{3}{2}}(t), \quad t \in]0, 1[,$$

with the following boundary conditions

$$y(0) = 1, \quad y(1) = 0,$$

is known as Thomas-Fermi equation ([3]). This equation was studied by Thomas [14] and Fermi [2]. One can impose three types of boundary conditions, which reflects three different states of atoms:

a) ionized atom

$$y(0) = 1, \quad y(a) = 0,$$

b) neutral atom

$$y(0) = 1, \quad ay'(a) = y(a),$$

c) isolated neutral atom

$$y(0) = 1, \quad \lim_{x \rightarrow \infty} y(x) = 0.$$

Thomas-Fermi equation may be written in a generalised form known as Emden-Fowler equation ([3, 5])

$$y''(t) + \sigma t^\gamma y^\beta(t) = 0, \quad t \in]0, 1[, \tag{1}$$

$$y(0) = 1; \quad y(1) = 0, \tag{2}$$

$$y(0) = 1; \quad y(1) = y'(1), \tag{3}$$

where γ, β and σ are real numbers, $\gamma < -2$, $\beta > 1$. Mehta et al. [9] studied it by using hypergeometric function and studied the Emden-Fowler equations. Wong [16] presented survey on Emden-Fowler equations. Rosenau [13] by simple change of variable generated a one parameter family of integrable Emden-Fowler equations. P M Lima [5, 6, 7] analyzed numerical methods based on finite differences/shooting methods for solving Emden-Fowler equation. Mohammadi et al. [10] used an iterative method to the singular and nonlinear fractional partial differential version of Emden-Fowler equations. Haar wavelets operational matrix of fractional integration is be applied to compute the solution with the Picard technique.

Pikulin [11] considered the Emden-Fowler equations on the half-line and on the interval and by assuming that the exponent in the coefficient of the nonlinear term is rational, he derived new parametric representations.

In this paper we consider (1) subject to the boundary conditions (2) and (3). We develop two approaches. In the first approach we use Newton's quasilinearization and arrive at an iteration scheme which is linear at each iteration. At each iteration we apply Haar wavelet collocation approach. After certain iterations the solutions does not vary, so we stop and the solution that we get at this stage is the solution of the Emden-Fowler equation. In the second approach we first use Haar wavelet collocation and arrive at the set of nonlinear equations. We then solve the system of nonlinear equation to get the solution of the Emden-Fowler equation. To the best of our knowledge the results do not exists in literature.

The paper is divided into the following sections. In section 1 we discuss the literature survey. In section 2 we discuss basics of Haar Wavelets. In section 3 we discuss about proposed methodologies. Section 4 is devoted to computations. Finally in section 5 we conclude the paper.

2 Haar Wavelet

[17, pp.7-10] Let us consider the interval $0 \leq x \leq 1$. Let us define $M = 2^J$, where J is maximum level of resolution. Let us divide $[0, 1]$ into $2M$ subintervals of equal length $\Delta x = \frac{1}{2M}$. The wavelet number i is calculated by $i = m + k + 1$, here $j = 0, 1, \dots, J$ and $k = 0, 1, \dots, m - 1$ (here $m = 2^j$). The Haar wavelet's mother wavelet function is defined as,

$$h_i(x) = \begin{cases} 1, & \alpha_1(i) \leq x < \alpha_2(i), \\ -1, & \alpha_2(i) \leq x < \alpha_3(i), \\ 0, & \text{else,} \end{cases} \quad (4)$$

where,

$$\alpha_1(i) = 2k\mu\Delta x, \quad \alpha_2(i) = (2k + 1)\mu\Delta x, \quad \alpha_3(i) = 2(k + 1)\mu\Delta x, \quad \mu = \frac{M}{m}, \quad (5)$$

if $i > 2$. For $i = 1$, it is defined as,

$$h_1(x) = \begin{cases} 1, & 0 \leq x \leq 1, \\ 0, & \text{else.} \end{cases} \quad (6)$$

For $i = 2$,

$$\alpha_1(2) = 0, \quad \alpha_2(2) = 0.5, \quad \alpha_3(2) = 1. \quad (7)$$

The width of the i^{th} wavelet is,

$$\alpha_3(i) - \alpha_1(i) = 2\mu\Delta x = 2^{-j}.$$

The integral $P_{v,i}(x)$ is defined as,

$$P_{v,i}(x) = \int_0^x \int_0^x \dots \int_0^x h_i(t) dt^v = \frac{1}{(v-1)!} \int_0^x (x-t)^{v-1} h_i(t) dt, \quad (8)$$

here v is the order of integration.

Using (4), we will calculate these integrals analytically and by doing it we obtain,

$$P_{v,i}(x) = \begin{cases} 0, & x < \alpha_1(i), \\ \frac{1}{v!} [x - \alpha_1(i)]^v, & \alpha_1(i) \leq x \leq \alpha_2(i), \\ \frac{1}{v!} [x - \alpha_1(i)]^v - 2[x - \alpha_2(i)]^v, & \alpha_2(i) \leq x \leq \alpha_3(i), \\ \frac{1}{v!} [x - \alpha_1(i)]^v - 2[x - \alpha_2(i)]^v + [x - \alpha_3(i)]^v, & x > \alpha_3(i), \end{cases} \quad (9)$$

for $i > 1$. For $i = 1$ we have $\alpha_1 = 0, \alpha_2 = \alpha_3 = 1$ and,

$$P_{v,1}(x) = \frac{x^v}{v!}. \quad (10)$$

For computation by using Haar wavelet, we make use of the method of collocation. Here, the grid points are defined by,

$$\tilde{x}_c = c\Delta x, \quad c = 0, 1, \dots, 2M, \quad (11)$$

and collocation points are defined by,

$$x_c = 0.5(\tilde{x}_{c-1} + \tilde{x}_c), \quad c = 1, \dots, 2M. \quad (12)$$

We substitute $x \rightarrow x_c$ in (4),(9),(10).

For computational point of view, we introduce the Haar matrices H, P_1, P_2, \dots, P_v . The order of these matrices are $2M \times 2M$. These matrices are defined by,

$$H(i, c) = h_i(x_c), P_v(i, c) = p_{v,i}(x_c) \quad v = 1, 2, \dots.$$

For $J = 1$, the matrices H, P_1 and P_2 are defined by,

$$H = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}, \quad P_1 = \frac{1}{8} \begin{bmatrix} 1 & 3 & 5 & 7 \\ 1 & 3 & 3 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}, \quad P_2 = \frac{1}{128} \begin{bmatrix} 1 & 9 & 25 & 49 \\ 1 & 9 & 23 & 31 \\ 1 & 7 & 8 & 8 \\ 0 & 0 & 1 & 7 \end{bmatrix}.$$

3 Haar Wavelet Collocation Method

In this section we develop two methods for solving generalized Emden-Fowler equation.

3.1 Quasilinearization Approach

By quasilinearization ([1, 15]) generalized Emden-Fowler equation is linearized then at each iteration Haar wavelet collocation method is used to compute the solution. After some iteration the solution converges to the solution of the Emden-Fowler equation. Consider the problem considered by Lima [5],

$$y''(t) + \sigma t^\gamma y^\beta(t) = 0, \quad t \in]0, 1[\quad (13)$$

$$y(0) = 1; \quad y(1) = 0, \quad (14)$$

where γ, β and σ are real numbers, $\gamma < -2, \beta > 1$. Now applying quasilinearization in above equation,

$$y''_{r+1}(t) = \sigma t^\gamma [-y_r^\beta(t) + (y_{r+1}(t) - y_r(t))(-\beta y_r^{\beta-1}(t))], \quad (15)$$

$$y_{r+1}(0) = 1; \quad y_r(1) = 0. \quad (16)$$

To apply Haar wavelet method [4], we assume that

$$y''_{r+1}(t) = \sum_{i=1}^{2M} a_i h_i(t), \quad (17)$$

where a_i are wavelet coefficients. Now integrate above equation from 0 to t twice we get,

$$y'_{r+1}(t) = \sum_{i=1}^{2M} a_i P_{1,i}(t) + y'_{r+1}(0), \quad (18)$$

$$y_{r+1}(t) = \sum_{i=1}^{2M} a_i P_{2,i}(t) + t y'_{r+1}(0) + y_{r+1}(0). \quad (19)$$

Now apply boundary conditions (16) in (19) and find $y'_{r+1}(0)$ then substitute (19),(17) in (15) and we get

$$\sum_{i=1}^{2M} a_i [h_i(t) + \sigma \beta t^\gamma y_r^{\beta-1}(t) (P_{2,i}(t) - t P_{2,i}(1))] = \sigma (\beta - 1) t^\gamma y_r^\beta(t) - \sigma \beta (1 - t) t^\gamma y_r^{\beta-1}(t), \quad (20)$$

Now, we can solve above equation by collocation method.

3.2 Newton Raphson Method

[17, pp.33] Consider a nonlinear ordinary differential equation,

$$\psi(x, y, y', y'') = 0, \quad x \in [0, 1], \quad (21)$$

where ψ is a nonlinear function. Now again applying Haar wavelet method [4], let us assume

$$y''(t) = \sum_{i=1}^{2M} a_i h_i(t), \quad (22)$$

and integrate above equation 2-times. Obtained results are discretized by collocation method then substitute these results in (21). By doing this we will get a system of non-linear equations,

$$\phi_c(a_1, a_2, \dots, a_{2M}) = 0, \quad c = 1, 2, \dots, 2M. \quad (23)$$

To solve above equation here we are using Newton's method and using this method wavelet coefficients (a_i) can be calculated.

3.3 Convergence

The convergence of the proposed methods is an immediate consequence of the results stated by Majak et al. [8].

Theorem 3.1. ([8]) Let us assume that $g(x) = \frac{d^2y}{dx^2} \in L^2(\mathbb{R})$ is a continuous function on $[0, 1]$ and its first derivative is bounded for all $x \in [0, 1]$, there exists ξ such that $\left| \frac{dg}{dt} \right| \leq \xi$, then both approaches discussed in this paper converge.

Proof. The proof is similar to the proof given in [15].

4 Numerical Illustration

4.1 Example 1

Consider (13), (14) with $\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$. Solving by quasilinearization method (subsection 3.1) substitute σ, γ and β and replace $t \rightarrow t_c \forall c = 1, 2, \dots, 2M$ in (20) where t_c are collocation points (section 2). Now we get (20) in terms of H, P_2, A and Y_r where $A = [a_1, a_2, \dots, a_{2M}]$ and $Y_r = [y_r(t_1), y_r(t_2), \dots, y_r(t_{2M})]$ are row vectors from which unknowns $[a_1, a_2, \dots, a_{2M}]$ are calculated.

Taking initial guess $[y_0(t_1), y_0(t_2), \dots, y_0(t_{2M})]$ and by Haar wavelet collocation method, we will get required solution. Taking initial guess $[0, 0, \dots, 0]$ solution is given in Table 1 and graph for $J = 3, J = 5, J = 7$ and $J = 8$ is given in figure 1. Taking different initial guess $[0.01, 0.01, \dots, 0.01]$ and $[0.1, 0.1, \dots, 0.1]$ solution still remain same.

Now by Newton's method (subsection 3.2), assume

$$y''(t) = \sum_{i=1}^{2M} a_i h_i(t), \quad (24)$$

where a_i are wavelet coefficients. Now integrate above equation from 0 to t twice we get,

$$y'(t) = \sum_{i=1}^{2M} a_i P_{1,i}(t) + y'(0), \quad (25)$$

$$y(t) = \sum_{i=1}^{2M} a_i P_{2,i}(t) + ty'(0) + y(0). \quad (26)$$

Now apply boundary conditions (16) in (26) and we get $y(t)$ as,

$$y(t) = 1 - t + \sum_{i=1}^{2M} a_i (P_{2,i}(t) - tP_{2,i}(1)). \quad (27)$$

Substituting (27),(24) in (21), we get system of non-linear equations. Thus we arrive at (23). To solve the non-linear equation we use Newton's method to calculate wavelet coefficients (a_i).

For initial guess $[1, 1, \dots, 1]$ computed solution is given in Table 2 and graph for $J = 3, J = 5, J = 7$ and $J = 8$ is given in figure 2. Taking varying initial guesses to $[1.01, 1.01, \dots, 1.01]$ and $[1.1, 1.1, \dots, 1.1]$ we observe that solution does not change significantly.

t	$J = 3, r = 3$	$J = 5, r = 3$	$J = 7, r = 3$	$J = 8, r = 3$	E-algo.[5, pp.262]
0.1	0.849094	0.849401	0.849464	0.84947	0.849474
0.2	0.726803	0.727162	0.727222	0.727228	0.727231
0.3	0.618867	0.619233	0.619286	0.619292	0.619294
0.4	0.520041	0.520361	0.520408	0.520412	0.520414
0.5	0.427227	0.427506	0.427544	0.427548	0.42755
0.6	0.338421	0.338651	0.338682	0.338685	0.338686
0.7	0.252197	0.252371	0.252395	0.252397	0.252398
0.8	0.16751	0.167631	0.167647	0.167648	0.167649
0.9	0.0836165	0.0836776	0.0836856	0.0836864	0.083686

Table 1: Solutions for $\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$ for $J = 3, J = 5, J = 7$ and $J = 8$ by Quasilinearization method




Figure 1: Graph of Quasilinearization method for $J = 3, J = 5, J = 7, J = 8$ and E-algo ([5]).

t	$J = 3, r = 3$	$J = 5, r = 3$	$J = 7, r = 3$	$J = 8, r = 3$	E-algo.[5, pp.262]
0.1	0.849094	0.849401	0.849464	0.84947	0.849474
0.2	0.726803	0.727162	0.727222	0.727228	0.727231
0.3	0.618867	0.619233	0.619286	0.619292	0.619294
0.4	0.520041	0.520361	0.520408	0.520412	0.520414
0.5	0.427227	0.427506	0.427544	0.427548	0.42755
0.6	0.338421	0.338651	0.338682	0.338685	0.338686
0.7	0.252197	0.252371	0.252395	0.252397	0.252398
0.8	0.16751	0.167631	0.167647	0.167648	0.167649
0.9	0.0836165	0.0836776	0.0836856	0.0836864	0.083686

Table 2: Solutions for $\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$ for $J = 3, J = 5, J = 7$ and $J = 8$ by Newton's method




Figure 2: Graph of Newton's method for $J = 3$, $J = 5$, $J = 7$, $J = 8$ and E-algo ([5]).

4.2 Example 2

Consider (13), (14) with $\sigma = -1, \gamma = -1$ and $\beta = 2$. Here we follow same procedure similar to example 1. For Quasilinearization method we take $[0, 0, \dots, 0]$ as initial guess. Computed solution is given in Table 3 and graph for $J = 3$, $J = 5$, $J = 7$ and $J = 8$ is given in figure 3. Taking different initial guesses $[0.01, 0.01, \dots, 0.01]$ and $[0.1, 0.1, \dots, 0.1]$ we observe that solution does not vary which proves that method is stable.

For Newton's method we take $[1, 1, \dots, 1]$ as initial guess. solution is given in Table 4 and graph for $J = 3$, $J = 5$, $J = 7$ and $J = 8$ is given in figure 4. Taking different initial guess $[1.01, 1.01, \dots, 1.01]$ and $[1.1, 1.1, \dots, 1.1]$ solution still remains same.

4.3 Example 3

Consider (13), (14) with $\sigma = -1, \gamma = \frac{-5}{4}$ and $\beta = \frac{9}{4}$. Here we follow same procedure similar to example 1. For Quasilinearization method we take $[0, 0, \dots, 0]$ as initial guess. Computed solution is displayed in Table 5 and graph for $J = 3$, $J = 5$, $J = 7$ and $J = 8$ is given in figure 5. Small perturbations in initial guesses $[0.01, 0.01, \dots, 0.01]$ and $[0.1, 0.1, \dots, 0.1]$ does not change the solution significantly.

For Newton's method we take $[1, 1, \dots, 1]$ as initial guess. Computed solution is given in Table 6 and graph for $J = 3$, $J = 5$, $J = 7$ and $J = 8$ is given in figure 6. Taking different initial guess $[1.01, 1.01, \dots, 1.01]$ and $[1.1, 1.1, \dots, 1.1]$ we observe that solution remains same.

t	$J = 3, r = 3$	$J = 5, r = 3$	$J = 7, r = 3$	$J = 8, r = 3$	E-algo.[5, pp.264]
0.1	0.781336	0.780201	0.78013	0.780126	0.780125
0.2	0.657973	0.657498	0.657471	0.657469	0.657468
0.3	0.558596	0.558365	0.55835	0.558349	0.558348
0.4	0.470276	0.470119	0.470109	0.470109	0.470108
0.5	0.387687	0.387587	0.387581	0.387581	0.38758
0.6	0.308217	0.30815	0.308146	0.308146	0.308145
0.7	0.23039	0.230345	0.230342	0.230342	0.230342
0.8	0.153353	0.153328	0.153327	0.153326	0.153326
0.9	0.0766365	0.0766246	0.0766239	0.0766238	0.076623

Table 3: Solutions for $\sigma = -1, \gamma = -1$ and $\beta = 2$ for $J = 3$, $J = 5$, $J = 7$ and $J = 8$ by Quasilinearization method




Figure 3: Graph of Quasilinearization method for $J = 3$, $J = 5$, $J = 7$, $J = 8$ and E-algo ([5]).

t	$J = 3, r = 3$	$J = 5, r = 3$	$J = 7, r = 3$	$J = 8, r = 3$	E-algo.[5, pp.264]
0.1	0.781336	0.780201	0.78013	0.780126	0.780125
0.2	0.657973	0.657498	0.657471	0.657469	0.657468
0.3	0.558596	0.558365	0.55835	0.558349	0.558348
0.4	0.470276	0.470119	0.470109	0.470109	0.470108
0.5	0.387687	0.387587	0.387581	0.387581	0.38758
0.6	0.308217	0.30815	0.308146	0.308146	0.308145
0.7	0.23039	0.230345	0.230342	0.230342	0.230342
0.8	0.153353	0.153328	0.153327	0.153326	0.153326
0.9	0.0766365	0.0766246	0.0766239	0.0766238	0.076623

Table 4: Solutions for $\sigma = -1, \gamma = -1$ and $\beta = 2$ for $J = 3$, $J = 5$, $J = 7$ and $J = 8$ by Newton's method




Figure 4: Graph of Newton's method for $J = 3$, $J = 5$, $J = 7$, $J = 8$ and E-algo ([5]).

t	$J = 3, r = 3$	$J = 5, r = 3$	$J = 7, r = 3$	$J = 8, r = 3$	E-algo.[5, pp.265]
0.1	0.716249	0.708156	0.7057	0.70517	0.704396
0.2	0.598674	0.593057	0.591181	0.590768	0.590163
0.3	0.508462	0.504055	0.502524	0.502185	0.501688
0.4	0.428945	0.425335	0.424071	0.42379	0.423379
0.5	0.354347	0.351425	0.350393	0.350164	0.349827
0.6	0.282173	0.27987	0.279053	0.278872	0.278605
0.7	0.211155	0.20944	0.208831	0.208696	0.208497
0.8	0.140637	0.1395	0.139095	0.139005	0.138872
0.9	0.0702995	0.0697316	0.0695291	0.0694841	0.069418

Table 5: Solutions for $\sigma = -1, \gamma = \frac{-5}{4}$ and $\beta = \frac{9}{4}$ for $J = 3, J = 5, J = 7$ and $J = 8$ by Quasilinearization method

t	$J = 3, r = 3$	$J = 5, r = 3$	$J = 7, r = 3$	$J = 8, r = 3$	E-algo.[5, pp.265]
0.1	0.716249	0.708156	0.7057	0.70517	0.704396
0.2	0.598674	0.593057	0.591181	0.590768	0.590163
0.3	0.508462	0.504055	0.502524	0.502185	0.501688
0.4	0.428945	0.425335	0.424071	0.42379	0.423379
0.5	0.354347	0.351425	0.350393	0.350164	0.349827
0.6	0.282173	0.27987	0.279053	0.278872	0.278605
0.7	0.211155	0.20944	0.208831	0.208696	0.208497
0.8	0.140637	0.1395	0.139095	0.139005	0.138872
0.9	0.0702995	0.0697316	0.0695291	0.0694841	0.069418

Table 6: Solutions for $\sigma = -1, \gamma = \frac{-5}{4}$ and $\beta = \frac{9}{4}$ for $J = 3, J = 5, J = 7$ and $J = 8$ by Newton's method




Figure 5: Graph of Quasilinearization method for $J = 3, J = 5, J = 7, J = 8$ and E-algo ([5]).




Figure 6: Graph of Newton's method for $J = 3$, $J = 5$, $J = 7$, $J = 8$ and E-algo ([5]).

4.4 Example 4

Consider (13) with $\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$ along with the boundary conditions,

$$y(0) = 1, \quad y'(1) = y(1). \quad (28)$$

This problem is considered in [12] to compute various atomic states.

Now, by applying Haar wavelet method ([4]), we arrive at the following nonlinear system:

$$y(t) = 1 - t + \sum_{i=1}^{2M} a_i (P_{2,i}(t) - tP_{2,i}(1)) + x \left(\sum_{i=1}^{2M} a_i h(1) \right)^{\frac{2}{3}}. \quad (29)$$

In this case we compute residual error. Residual error for the differential equation $y''(t) - t^{-1/2}y^{3/2} = 0$ is defined as $R(t) = \tilde{y}''(t) - t^{-1/2}\tilde{y}^{3/2}$, where \tilde{y} is approximate Haar wavelet solution. Maximum absolute residual error is defined as

$$R^\infty = \max_i |R(t_i)|.$$

To solve nonlinear system obtained by substituting (29) into considered Thomas Fermi equation (13) - (28) with $\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$, we apply Newton's method and compute the solutions. Taking $[1, 1, \dots, 1]$ as initial guess, we get solutions which is depicted in Table 7. Graph of computed solution and plots for residual errors for $J = 3$, $J = 5$ and $J = 7$ is given in figure 7 and 8, respectively. To check the continuous dependence on data and stability of the proposed method we take different initial guesses $[1.01, 1.01, \dots, 1.01]$ and $[1.1, 1.1, \dots, 1.1]$, and observe that final computed solution does not vary significantly.

As J increases R^∞ decreases and solution is further improved (see Fig. 8). Since final solution is independent of initial guesses and R^∞ decreases with increase in J , our method is a powerful technique which can be used to compute the solution of Emden-Fowler or Thomas Fermi type equations.

t	$J = 3, N = 3$	$J = 5, N = 3$	$J = 7, N = 3$
0.1	0.970377	0.975761	0.978282
0.2	0.974685	0.985461	0.990554
0.3	1.00098	1.01744	1.02527
0.4	1.04576	1.06831	1.07909
0.5	1.10753	1.13672	1.15072
0.6	1.18584	1.22232	1.23985
0.7	1.28087	1.32541	1.34684
0.8	1.39328	1.44677	1.47256
0.9	1.5241	1.58762	1.61829
R^∞	0.102233	0.0237877	0.00601627

Table 7: Solutions for $\sigma = -1, \gamma = \frac{-1}{2}$ and $\beta = \frac{3}{2}$ for $J = 3$, $J = 5$ and $J = 7$ by Newton's method.




Figure 7: Graph of solutions by Newton's method for $J = 3$, $J = 5$ and $J = 7$.




Figure 8: Graph of Residual error for Newton's method at $J = 3$, $J = 5$ and $J = 7$.

5 Conclusion

In this paper we have used Haar wavelet collocation method on a class of Emden-Fowler equation which originates in Physics. These equations are also referred as Thomas-Fermi equations. We have proposed two methods: in one approach we first linearise it by Newton's quasilinearization method and then solve it by Haar wavelet collocation method. In the second approach we use Haar wavelet collocation method and arrive at system of non-linear equation which we then solve by Newton-Raphson method. We observed that as value of J is increased beyond certain value computed solution does not vary significantly. We also observed that small perturbation in initial guess does not vary final computed solution at all. So therefore both methods are stable and robust. Higher the resolution we use, our computed solution is closer to the exact solution.

References

- [1] Eman S. Alaidarous, Malik Zaka Ullah, Fayyaz Ahmad, and A.S. Al-Fhaid. An efficient higher-order quasilinearization method for solving nonlinear bvps. *Journal of Applied Mathematics*, 2013:11 pages, 2013.
- [2] E. Fermi. Un metodo statistico per la determinazione di alcune proprieta dell'atome. *Rend. Accad. Naz. Lincei*, 6:602–607, 1927.

- [3] A.C. Lemos, M.J. Baines, and N.K. Nichols. Numerical methods for the emden-fowler equation, 1998.
- [4] Ulo Lepik. Haar wavelet method for solving higher order differential equations. *Int. J. Math. Comput*, 1:84–94, 2008.
- [5] Pedro M Lima. Numerical methods and asymptotic error expansions for the emden-fowler equations. *Journal of computational and applied mathematics*, 70:245–266, 1996.
- [6] P.M. Lima and M.P. Carpentier. Asymptotic expansions and numerical approximation of nonlinear degenerate boundary-value problems. *Applied Numerical Mathematics*, 30(1):93 – 111, 1999.
- [7] P.M. Lima and L. Morgado. Analytical-numerical investigation of a singular boundary value problem for a generalized emdenfowler equation. *Journal of Computational and Applied Mathematics*, 229(2):480 – 487, 2009. Special Issue: Analysis and Numerical Approximation of Singular Problems.
- [8] J. Majak, B. S. Shvartsman, M. Kirs, M. Pohlak, and H. Herranen. Convergence theorem for the Haar wavelet based discretization method. *Composite Structures*, 126:227–232, 2015.
- [9] B.N Mehta and R Aris. A note on a form of the emden-fowler equation. *Journal of Mathematical Analysis and Applications*, 36(3):611 – 621, 1971.
- [10] Amir Mohammadi, Nasser Aghazadeh, and Shahram Rezapour. Haar wavelet collocation method for solving singular and nonlinear fractional time-dependent emden–fowler equations with initial and boundary conditions. *Mathematical Sciences*, 13(3):255–265, Sep 2019.
- [11] S. V. Pikulin. The thomas–fermi problem and solutions of the emden–fowler equation. *Computational Mathematics and Mathematical Physics*, 59(8):1292–1313, Aug 2019.
- [12] R.C.Flagg, C.D.Luning, and W.L.Perry. Implementation of new iterative techniques for solutions of thomas-fermi and emden-fowler equations. *Journal of Computational Physics*, 38:396–405, 1980.
- [13] Philip Rosenau. A note on integration of the emden-fowler equation. *International Journal of Non-Linear Mechanics*, 19(4):303 – 308, 1984.
- [14] L. H. Thomas. The calculation of atomic fields. *Mathematical Proceedings of the Cambridge Philosophical Society*, 23(5):542548, 1927.
- [15] Amit K Verma and Diksha Tiwari. Higher resolution methods based on quasilinearization and haar wavelets on lane-emden equations. *International Journal of Wavelets Multiresolution and Information Processing*, 17:1950005, 2019.
- [16] James S. W. Wong. On the generalized emdenfowler equation. *SIAM Review*, 17(2):339–360, 1975.
- [17] Ulo Lepik and Helle Hein. *Haar Wavelets With Applications*. Springer, Cham, 2014.