

Exclusive quarkonium production or decay in soft gluon factorization

Rong Li^a Yu Feng^b Yan-Qing Ma^{c,d,e}

^a*School of Science, Xi'an Jiaotong University, Xi'an 710049, China*

^b*Department of Physics, College of Basic Medical Sciences, Army Medical University, Chongqing 400038, China*

^c*School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

^d*Center for High Energy Physics, Peking University, Beijing 100871, China*

^e*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

E-mail: rongli@xjtu.edu.cn, yfeng@ihep.ac.cn, yqma@pku.edu.cn

ABSTRACT: In this paper, we study the application of the recently proposed soft gluon factorization (SGF) to exclusive quarkonium production or decay. We find that in the nonrelativistic QCD factorization framework there are too many nonperturbative parameters. Thanks to the factorization of kinematical physics from dynamical physics, the SGF significantly reduces the number of nonperturbative parameters. Therefore, the SGF can improve our predictive power of exclusive quarkonium production or decay.

Contents

1	Introduction	1
2	Factorization formula for exclusive processes	2
2.1	From inclusive formula to exclusive formula	2
2.2	Definition of nonperturbative matrix elements	2
2.3	Factorization formula at the amplitude level	3
2.4	Perturbative matching	4
2.5	Factorization formula for quarkonium exclusive decay	5
3	Applications	6
3.1	Leading order calculation of $e^+e^- \rightarrow \eta_c + \gamma$	6
3.2	Leading order calculation of $\eta_c \rightarrow \gamma\gamma$	7
3.3	Numerical result	8
4	Comparison with NRQCD factorization	8
5	Summary and outlook	10

1 Introduction

The study of heavy quarkonium production or decay is useful to understand both perturbative and nonperturbative physics of QCD. In the past two decades, the widely used theory for quarkonium physics is the non-relativistic quantum chromodynamics (NRQCD) factorization [1], which factorizes processes to perturbatively calculable hard parts multiplied by nonperturbative long-distance matrix elements (LDMEs). Because LDMEs are simply numbers, the NRQCD factorization has strong predictive power.

However, recent studies reveal that NRQCD factorization encounters some difficulties in describing inclusive quarkonium production data. In ref. [2], it was argued that the difficulties of NRQCD may be caused by the bad convergence of relativistic expansion. To overcome this problem, a new factorization method called soft gluon factorization (SGF) was proposed [2]. In this method, a series of relativistic corrections can be resummed to all orders and thus a better convergence in relativistic expansion is expected.

Exclusive quarkonium processes have also been paid a lot of attentions to. For example, the leading order (LO) in α_s calculation of $J/\psi + \eta_c$ production at B factories in NRQCD [3, 4] conflicts with experimental data [5], which attracts a lot of studies of relativistic corrections [6, 7] and high order QCD corrections [8–10]. Later on, the factorization theorem for double charmonia production was proved rigorously [11–13]. The exclusive

production of $\eta_c + \gamma$ at B factories has also been studied extensively within the NRQCD factorization [14–20].

Due to the advantage of SGF, it is interesting to see what happens if one applies it to exclusive quarkonium processes, which is the aim of this paper. In the rest of the paper, we first establish the general method to apply SGF on exclusive quarkonium processes. We then take $\sigma_{e^+e^- \rightarrow \eta_c + \gamma}$ and $\Gamma(\eta_c \rightarrow \gamma + \gamma)$ as examples to do phenomenological study. We will find that the SGF can not only factorize perturbative physics from nonperturbative physics but also factorize kinematical physics from dynamical physics. Because of the later effect, the SGF has much less free parameters comparing with NRQCD factorization, and therefore it has much stronger predictive power.

2 Factorization formula for exclusive processes

2.1 From inclusive formula to exclusive formula

In SGF, the differential cross section for inclusive production of a quarkonium H is given by [2]

$$(2\pi)^3 2P^0 \frac{d\sigma_H}{d^3P} = \sum_n \int \frac{d^4P_c}{(2\pi)^4} \mathcal{H}_n(P_c) F_{n \rightarrow H}(P_c, P), \quad (2.1)$$

where $\mathcal{H}_n(P_c)$ are perturbatively calculable hard parts that, roughly speaking, produce an intermediate state n with total momentum P_c , and $F_{n \rightarrow H}(P_c, P)$ are called soft gluon distributions (SGDs) that describe the hadronization of the intermediate state to the quarkonium H with momentum P .

Exclusive process can be thought of as a special case of inclusive process, where there is no real emission during the hadronization process, described by $F_{n \rightarrow H}(P_c, P)$. As a result, all conserved quantities are the same between the intermediate state n and the quarkonium H , which include total momentum, color charge, and J^{PC} . The momentum conservation enables the decomposition

$$F_{n \rightarrow H}(P_c, P) \rightarrow (2\pi)^4 \delta^4(P_c - P) \langle \overline{\mathcal{O}}_n^{H\text{ex}} \rangle, \quad (2.2)$$

where the subscript “ex” means exclusive production. Then we get the factorization formula for exclusive production of polarization summed quarkonium

$$(2\pi)^3 2P^0 \frac{d\sigma_H}{d^3P} = \sum_n \mathcal{H}_n(P) \langle \overline{\mathcal{O}}_n^{H\text{ex}} \rangle. \quad (2.3)$$

2.2 Definition of nonperturbative matrix elements

As discussed in ref. [2], the intermediate state n should at least contain a $Q\overline{Q}$ pair. In addition to that, n can also contain dynamical soft partons, i.e., gluons or light (anti-)quarks. Contributions from intermediate states with soft partons are inevitably suppressed by powers of v , the typical velocity of heavy quark inside of the quarkonium, and usually there is no mechanism to enhance this kind of contributions [2]. Therefore, as the first

approximation, we only consider intermediate states that do not contain dynamical soft partons and denote quantum numbers of n in terms of spectroscopic notation $^{2S+1}L_{J,J_z}^{[1]}$, with superscript 1 denoting color singlet. Then, we have

$$\langle \overline{\mathcal{O}}_n^{H\text{ex}} \rangle = \sum_{J_z^H} \langle 0 | [\overline{\Psi} \mathcal{K}_n \Psi]^\dagger(0) | H \rangle \langle H | [\overline{\Psi} \mathcal{K}_n \Psi](0) | 0 \rangle, \quad (2.4)$$

where the quarkonium state has standard relativistic normalization and \mathcal{K}_n are defined as [2]

$$\mathcal{K}_n = \frac{\sqrt{M}}{M+2m} \frac{M+\not{P}}{2M} \Gamma_n \frac{M-\not{P}}{2M} \mathcal{C}^{[1]}, \quad (2.5)$$

where m is the mass of heavy quark, M is the mass of heavy quarkonium, and color projector $\mathcal{C}^{[1]} = 1/\sqrt{N_c}$. If n is S-wave, we have

$$\Gamma_n = \begin{cases} \gamma_5, & \text{if } S = 0, \\ \epsilon_{S_z}^\mu \gamma_\mu, & \text{if } S = 1, \end{cases} \quad (2.6)$$

where $\epsilon_{S_z}^\mu$ is polarization vector with $P \cdot \epsilon_{S_z} = 0$. For other cases, we refer the definition of Γ_n in ref. [2]. Following the discussion in ref. [2], we can obtain

$$\langle \overline{\mathcal{O}}_n^{H\text{ex}} \rangle = \frac{1}{2N_c} \langle \mathcal{O}_n^{H\text{ex}} \rangle [1 + O(v^2)], \quad (2.7)$$

where $\langle \mathcal{O}_n^{H\text{ex}} \rangle$ are standard NRQCD LDMEs for exclusive quarkonium production, which can be related to inclusive quarkonium production LDMEs via vacuum saturation approximation [1],

$$\langle \mathcal{O}_n^{H\text{ex}} \rangle = \langle \mathcal{O}_n^H \rangle [1 + O(v^4)]. \quad (2.8)$$

2.3 Factorization formula at the amplitude level

Based on the above definition, it is clear that the factorization formula eq. (2.3) can be expressed at the amplitude level as ¹

$$\mathcal{A}_H(P) = \sum_n \hat{\mathcal{A}}_n(P) \overline{R}_H^n, \quad (2.9)$$

where

$$\overline{R}_H^n = \langle H | [\overline{\Psi} \mathcal{K}_n \Psi](0) | 0 \rangle. \quad (2.10)$$

with

$$\langle \overline{\mathcal{O}}_n^{H\text{ex}} \rangle = \sum_{J_z^H} |\overline{R}_H^n|^2, \quad (2.11)$$

$$\sigma_H = \frac{1}{2s} \int \overline{\sum} |\mathcal{A}_H(P)|^2 d\text{PS}, \quad (2.12)$$

where “ $\overline{\sum}$ ” means averaging over initial states and summing over final states. Note that, based on the relation eq. (2.7), \overline{R}_H^n can be related to wave function at the origin of the heavy quarkonium [1] up to $O(v^2)$ corrections.

¹Considered the fact that different intermediate states can interfere with each others, the factorization formula at the amplitude level is more precise.

2.4 Perturbative matching

The factorization formula eq. (2.9) will be our start point to study exclusive quarkonium production in the following. To use this formula, we need to calculate $\hat{\mathcal{A}}_n(P)$ in perturbation theory. To this end, we relate n in eq. (2.9) by n' and then project H to a color-singlet state $n = Q\bar{Q} \left({}^{2S+1}L_J^{[1]} \right)$ in both sides of the equation, which results in

$$\mathcal{A}_n(P) = \sum_{n'} \hat{\mathcal{A}}_{n'}(P) \bar{R}_n^{n'}, \quad (2.13)$$

where the definition of projection will be explained below.

Following ref. [2], we define a complete set of on-shell color-singlet state $n = Q\bar{Q} \left({}^{2S+1}L_J^{[1]} \right)$ with momenta

$$p_Q = P/2 + q, \quad (2.14)$$

$$p_{\bar{Q}} = P/2 - q. \quad (2.15)$$

We project the pair to color-singlet state by $\mathcal{C}^{[1]}$, and we project the pair to state with total spin S and S_z by replacing spinors of $Q\bar{Q}$ pair by

$$\tilde{\Pi}_{SS_z} = \frac{(\not{p}_{\bar{Q}} - m) \frac{M - \not{P}}{2M} \tilde{\Gamma}_{SS_z}^s \frac{M + \not{P}}{2M} (\not{p}_Q + m)}{\sqrt{M}(M/2 + m)}, \quad (2.16)$$

with

$$\tilde{\Gamma}_{00}^s = -\gamma_5, \quad (2.17a)$$

$$\tilde{\Gamma}_{1S_z}^s = \epsilon_{S_z}^{*\mu} \gamma_\mu. \quad (2.17b)$$

On-shell conditions $p_Q^2 = p_{\bar{Q}}^2 = m^2$ results in

$$P \cdot q = 0, \quad (2.18)$$

$$q^2 = m^2 - M^2/4, \quad (2.19)$$

which constrain two degrees of freedom of q . The other two degrees of freedom of q , defined as spatial angles in the rest frame of the pair, can be removed by partial wave expansion.

Eventually, we get our definition

$$\mathcal{A}_n(P) = \sum_{L_z, S_z} \langle L, L_z; S, S_z | J, J_z \rangle \int d^2\Omega |\mathbf{q}|^{-L} \sqrt{\frac{(2L+1)!!}{4\pi(L!)}} Y_L^{*L_z} \text{Tr} \left[\mathcal{C}^{[1]} \tilde{\Pi}_{SS_z} \mathcal{A}_{Q+\bar{Q}}(P) \right], \quad (2.20)$$

where $\mathcal{A}_{Q+\bar{Q}}(P)$ is the $Q\bar{Q}$ production amplitude with spinors of $Q\bar{Q}$ removed. Similarly,

$$\bar{R}_n^{n'} = \sum_{L_z, S_z} \langle L, L_z; S, S_z | J, J_z \rangle \int d^2\Omega |\mathbf{q}|^{-L} \sqrt{\frac{(2L+1)!!}{4\pi(L!)}} Y_L^{*L_z} \text{Tr} \left[\mathcal{C}^{[1]} \tilde{\Pi}_{SS_z} \bar{R}_{Q+\bar{Q}}^{n'} \right]. \quad (2.21)$$

Following the derivation in ref. [2], we have

$$\overline{R}_n^{n'(0)} = \delta_{nn'}, \quad (2.22)$$

where the superscript “(0)” denotes leading order in α_s expansion.

By inserting perturbative expansions

$$\mathcal{A}_n = \mathcal{A}_n^{(0)} + \alpha_s \mathcal{A}_n^{(1)} + \alpha_s^2 \mathcal{A}_n^{(2)} + \dots, \quad (2.23a)$$

$$\hat{\mathcal{A}}_{n'} = \hat{\mathcal{A}}_{n'}^{(0)} + \alpha_s \hat{\mathcal{A}}_{n'}^{(1)} + \alpha_s^2 \hat{\mathcal{A}}_{n'}^{(2)} + \dots, \quad (2.23b)$$

$$\overline{R}_n^{n'} = \overline{R}_n^{n'(0)} + \alpha_s \overline{R}_n^{n'(1)} + \alpha_s^2 \overline{R}_n^{n'(2)} + \dots \quad (2.23c)$$

into eq. (2.13) and using the orthogonal relations eq. (2.22), we get the following relations

$$\hat{\mathcal{A}}_n^{(0)} = \mathcal{A}_n^{(0)}, \quad (2.24a)$$

$$\hat{\mathcal{A}}_n^{(1)} = \mathcal{A}_n^{(1)} - \sum_{n'} \mathcal{A}_{n'}^{(0)} \overline{R}_n^{n'(1)}, \quad (2.24b)$$

$$\hat{\mathcal{A}}_n^{(2)} = \mathcal{A}_n^{(2)} - \sum_{n'} \mathcal{A}_{n'}^{(1)} \overline{R}_n^{n'(1)} - \sum_{n'} \mathcal{A}_{n'}^{(0)} \overline{R}_n^{n'(2)}, \quad (2.24c)$$

and so on. Based on these relations, to get the perturbative expansion of $\hat{\mathcal{A}}_n$, we need to calculate \mathcal{A}_n and $\overline{R}_n^{n'}$ perturbatively.

2.5 Factorization formula for quarkonium exclusive decay

For quarkonium exclusive decay, we have similar factorization formula at the amplitude level

$$\mathcal{A}^H = \sum_n \hat{\mathcal{A}}^n \overline{R}_H^{n*}, \quad (2.25)$$

with decay width

$$\Gamma^H = \frac{1}{2M} \int \overline{\sum} |\mathcal{A}^H|^2 d\text{PS}. \quad (2.26)$$

Similar to quarkonium production, $\hat{\mathcal{A}}^n$ can be perturbatively calculated with

$$\hat{\mathcal{A}}^{n(0)} = \mathcal{A}^{n(0)}, \quad (2.27a)$$

$$\hat{\mathcal{A}}^{n(1)} = \mathcal{A}^{n(1)} - \sum_{n'} \mathcal{A}^{n'(0)} \overline{R}_n^{n'*(1)}, \quad (2.27b)$$

and so on. In these relations, \mathcal{A}^n is defined by

$$\mathcal{A}^n = \sum_{L_z, S_z} \langle L, L_z; S, S_z | J, J_z \rangle \int d^2\Omega |\mathbf{q}|^{-L} \sqrt{\frac{(2L+1)!!}{4\pi(L!)}} Y_L^{L_z} \text{Tr} \left[\mathcal{C}^{[1]} \Pi_{SS_z} \mathcal{A}^{Q+\overline{Q}} \right], \quad (2.28)$$

where $\mathcal{A}^{Q+\overline{Q}}$ is the $Q\overline{Q}$ decay amplitude with spinors of $Q\overline{Q}$ removed and

$$\Pi_{SS_z} = \frac{(\not{p}_Q + m) \frac{M+\not{P}}{2M} \Gamma_{SS_z}^s \frac{M-\not{P}}{2M} (\not{p}_{\overline{Q}} - m)}{\sqrt{M}(M/2 + m)}, \quad (2.29)$$

with

$$\Gamma_{00}^s = -\gamma_5, \quad (2.30a)$$

$$\Gamma_{1S_z}^s = \epsilon_{S_z}^\mu \gamma_\mu. \quad (2.30b)$$

3 Applications

3.1 Leading order calculation of $e^+e^- \rightarrow \eta_c + \gamma$

For η_c production, we have $J^{PC} = 0^{-+}$, which demands $n = 1S_0^{[1]}$. Thus the amplitude of $e^+e^- \rightarrow \eta_c + \gamma$ is given by

$$\mathcal{A}_{e^+e^- \rightarrow \eta_c + \gamma} = \hat{\mathcal{A}}_{e^+e^- \rightarrow c\bar{c}(1S_0^{[1]}) + \gamma} \bar{R}_{\eta_c}^{1S_0^{[1]}}. \quad (3.1)$$

At the lowest order in α_s , we have

$$\hat{\mathcal{A}}_{e^+e^- \rightarrow c\bar{c}(1S_0^{[1]}) + \gamma}^{(0)} = \mathcal{A}_{e^+e^- \rightarrow c\bar{c}(1S_0^{[1]}) + \gamma}^{(0)}, \quad (3.2)$$

with

$$\mathcal{A}_{e^+e^- \rightarrow c\bar{c}(1S_0^{[1]}) + \gamma}^{(0)} = \int \frac{d^2\Omega}{4\pi} \text{Tr} \left[\mathcal{C}^{[1]} \tilde{\Pi}_{00} \mathcal{A}_{e^+e^- \rightarrow c\bar{c} + \gamma}^{(0)} \right]. \quad (3.3)$$




Figure 1. Feynman diagrams for the process $e^+e^- \rightarrow \eta_c + \gamma$ at leading order in α_s .

Therefore, we need to calculate the two Feynman diagrams in figure. 1. Since the leptonic current is independent of spatial angles, we separate it from the complete amplitude and rewrite eq. (3.3) as

$$\mathcal{A}_{e^+e^- \rightarrow c\bar{c}(1S_0^{[1]}) + \gamma}^{(0)} = \frac{-i}{s} \mathcal{L}_\mu \epsilon_\nu^*(k) \mathcal{A}^{\mu\nu} \quad (3.4)$$

where $s = (k_{e^+} + k_{e^-})^2$ is the square of center of mass energy, $\epsilon_\nu^*(k)$ is the polarization vector of the final-state photon and the leptonic current \mathcal{L}_μ is defined by

$$\mathcal{L}_\mu = -ie \bar{v}(k_{e^+}) \gamma_\mu u(k_{e^-}). \quad (3.5)$$

Lorentz symmetry, C, P and T invariance imply that the hadronic current can be expressed as

$$\mathcal{A}^{\mu\nu} = \epsilon^{\mu\nu Pk} A, \quad (3.6)$$

where A is a lorentz invariant quantity. Then we multiply $\epsilon_{\mu\nu Pk}$ on both sides of eq. (3.6) and sum over Lorentz indexes, we obtain

$$A = \frac{\mathcal{A}^{\mu\nu}\epsilon_{\mu\nu Pk}}{2(P \cdot k)^2}, \quad (3.7)$$

where we have used $\epsilon^{\mu\nu Pk}\epsilon_{\mu\nu Pk} = 2(P \cdot k)^2$. Then the integration over spatial angles is very easy, which eventually gives

$$A = \frac{96\pi\alpha m e_c^2}{M\sqrt{3M\delta}(s - M^2)} \ln\left(\frac{1 + \sqrt{\delta}}{1 - \sqrt{\delta}}\right) \quad (3.8)$$

where $\delta = 1 - 4m^2/M^2$ and electronic charge of charm quark is denoted by e_c .

By substituting for leptonic current

$$\mathcal{L}_\mu \mathcal{L}_{\mu'}^* \rightarrow -\frac{4e^2 s}{3} g_{\mu\mu'} \quad (3.9)$$

and for the summation of photon polarization

$$\sum \epsilon^*(k)_\nu \epsilon(k)_{\nu'} \rightarrow -g_{\nu\nu'}, \quad (3.10)$$

we get the cross section

$$\begin{aligned} \sigma_{e^+e^- \rightarrow \eta_c + \gamma} &= \frac{1}{2s} \int |\overline{\mathcal{A}_{e^+e^- \rightarrow \eta_c + \gamma}}|^2 d\text{PS}_2 \\ &= \frac{1}{2s} \int \left(\frac{1}{s^2}\right) \left(\frac{1}{4}\right) \left(-\frac{4e^2 s}{3} g_{\mu\mu'}\right) (-g_{\nu\nu'}) \epsilon^{\mu\nu Pk} \epsilon^{\mu'\nu' Pk} |A|^2 \langle \overline{\mathcal{O}_{1S_0^{[1]}}^{\eta_{\text{cex}}}} \rangle d\text{PS}_2 \\ &= \frac{128\pi^2 \alpha^3 m^2 e_c^4 (s - M^2)}{s^3 M^3 \delta} \ln^2\left(\frac{1 + \sqrt{\delta}}{1 - \sqrt{\delta}}\right) \langle \overline{\mathcal{O}_{1S_0^{[1]}}^{\eta_{\text{cex}}}} \rangle. \end{aligned} \quad (3.11)$$

According to the proposal in SGF [2], one can expand m in the hard part around $M/2$ to simply the calculation, which is equivalent to expand eq. (3.11) in power series of δ , which gives

$$\sigma_{e^+e^- \rightarrow \eta_c + \gamma} = \frac{128\pi^2 \alpha^3 e_c^4 (s - M^2)}{s^3 M} \langle \overline{\mathcal{O}_{1S_0^{[1]}}^{\eta_{\text{cex}}}} \rangle \left(1 - \frac{1}{3}\delta - \frac{7}{45}\delta^2 - \frac{29}{315}\delta^3 + \dots\right), \quad (3.12)$$

where high order terms in this expansion can be thought of as relativistic corrections to the lowest order term.

3.2 Leading order calculation of $\eta_c \rightarrow \gamma\gamma$

We begin with the amplitude of $\eta_c \rightarrow \gamma\gamma$

$$\mathcal{A}^{\eta_c \rightarrow \gamma\gamma} = \hat{\mathcal{A}}^{c\bar{c}(1S_0^{[1]}) \rightarrow \gamma\gamma} \overline{R}_{\eta_c}^{1S_0^{[1]*}}, \quad (3.13)$$

with

$$\mathcal{A}^{c\bar{c}(1S_0^{[1]}) \rightarrow \gamma\gamma} = \int d\Omega \text{Tr} \left[\mathcal{C}^{[1]} \Pi_{00} \mathcal{A}^{e^+ \bar{c} \rightarrow \gamma\gamma} \right]. \quad (3.14)$$

The amplitude can be decomposed as

$$\mathcal{A}^{c\bar{c}(1S_0^{[1]})\rightarrow\gamma\gamma} = \epsilon_\mu^*(k_1)\epsilon_\nu^*(k_2)\mathcal{A}^{\mu\nu}, \quad (3.15)$$

where $k_i (i = 1, 2)$ are momenta of γ 's in the final state and the hadronic current $\mathcal{A}^{\mu\nu}$ can be calculated similarly to the production case.

The decay width of $\eta_c \rightarrow \gamma\gamma$ in SGF gives

$$\begin{aligned} \Gamma(\eta_c \rightarrow \gamma\gamma) &= \frac{1}{2M} \cdot \frac{1}{2!} \int |\overline{\mathcal{A}_{\eta_c \rightarrow \gamma\gamma}}|^2 d\text{PS}_2 \\ &= \frac{1}{2M} \cdot \frac{1}{2!} \cdot (-g_{\mu\mu'})(-g_{\nu\nu'})\epsilon^{\mu\nu Pk}\epsilon^{\mu'\nu'Pk}|A|^2 \frac{1}{8\pi} \langle \overline{\mathcal{O}}_{1S_0^{[1]}}^{\eta_{c\text{ex}}} \rangle \\ &= \frac{48\pi\alpha^2 m^2 e_c^4}{M^4 \delta} \ln^2\left(\frac{1+\sqrt{\delta}}{1-\sqrt{\delta}}\right) \langle \overline{\mathcal{O}}_{1S_0^{[1]}}^{\eta_{c\text{ex}}} \rangle. \end{aligned} \quad (3.16)$$

This expression is consistent with NRQCD calculation in ref. [21] and references therein.

3.3 Numerical result

Based on eqs. (3.11) and (3.16), we get

$$\sigma_{e^+e^- \rightarrow \eta_c + \gamma} = \frac{8\pi\alpha M(s - M^2)}{3s^3} \Gamma(\eta_c \rightarrow \gamma\gamma), \quad (3.17)$$

which relates the exclusive production cross section of η_c with the exclusive decay width. This relation holds at leading order in α_s and all orders in v . It is impressive that this relation has no dependence on heavy quark mass m .

In numerical calculation, we choose the fine structure constant $\alpha = 1/137$, the quarkonium mass $M = 2.98$ GeV and collision energy $\sqrt{s} = 10.58$ GeV for B factories. By using experiment result for decay width [22]

$$\Gamma(\eta_c \rightarrow \gamma\gamma) = 5.02 \pm 0.13 \pm 0.38 \text{ keV}, \quad (3.18)$$

we get the prediction for cross section of $e^+e^- \rightarrow \eta_c + \gamma$ in SGF:

$$\sigma_{e^+e^- \rightarrow \eta_c + \gamma} = 26.2 \pm 2.1 \text{ fb}. \quad (3.19)$$

If we further choose the heavy quark pole mass as $m = 1.4 \pm 0.2$ GeV, we obtain

$$2N_c \langle \overline{\mathcal{O}}_{1S_0^{[1]}}^{\eta_{c\text{ex}}} \rangle = 0.176 \pm 0.014_{-0.015}^{+0.021} = 0.176_{-0.021}^{+0.025} \text{ GeV}^3. \quad (3.20)$$

where the later uncertainty is due to the varying of heavy quark mass.

4 Comparison with NRQCD factorization

In the NRQCD factorization [1], the calculation of the cross section of $e^+e^- \rightarrow \eta_c + \gamma$ is very similar to that in the SGF method, but one needs to expand q^2 in the hard part and

then put it into the definition of LDMEs. By setting $M^2 = 4m^2 - 4q^2$, from eq. (3.11) we can obtain the cross section calculated in NRQCD:

$$\sigma_{e^+e^- \rightarrow \eta_c + \gamma} = \hat{\sigma}_{1S_0^{[1]}}^{v^0} \langle \mathcal{O}^{\eta_{cex}}(1S_0^{[1]}) \rangle + \hat{\sigma}_{1S_0^{[1]}}^{v^2} \langle \mathcal{P}^{\eta_{cex}}(1S_0^{[1]}) \rangle + \dots, \quad (4.1)$$

with short-distance coefficients

$$\hat{\sigma}_{1S_0^{[1]}}^{v^0} = \frac{32\pi^2 \alpha^3 e_c^4 (s - 4m^2)}{3ms^3}, \quad (4.2)$$

$$\hat{\sigma}_{1S_0^{[1]}}^{v^2} = -\frac{16\pi^2 \alpha^3 e_c^4 (5s + 4m^2)}{9ms^3}, \quad (4.3)$$

where $\hat{\sigma}_{1S_0^{[1]}}^{v^0}$ agrees with results in refs. [15, 23] and $\hat{\sigma}_{1S_0^{[1]}}^{v^2}$ agrees with results in refs. [15–19].

The terms in eq. (4.1) are organized by the power counting in v , e.g., the second term is v^2 suppressed comparing with the first term. As each term introduces one free parameter, in NRQCD factorization one needs a lot of nonperturbative parameters to provide a precise description of the cross section. In contrast, one needs only one parameter in SGF framework².

In table 1, we compare predictions based on SGF and that based on NRQCD factorization [16–18]. Because of different choice of parameters, results at lowest order in v are very different between these references. Especially, the $\langle \mathcal{O}_{1S_0^{[1]}}^{\eta_{cex}} \rangle$ is chosen as 0.437 GeV^3 in refs. [16, 17] and 0.694 GeV^3 in ref. [18]. The difference of the parameters can be well understood. For example, comparing our result in eq. (3.20) with $\langle \mathcal{O}_{1S_0^{[1]}}^{\eta_{cex}} \rangle = 0.437^{+0.111}_{-0.105} \text{ GeV}^3$ obtained in ref. [21], nearly half of the suppression is due to the change of experimental value $\Gamma(\eta_c \rightarrow \gamma\gamma)$ from the older one $7.2 \pm 0.7 \pm 2.0 \text{ keV}$ to the new one $5.02 \pm 0.13 \pm 0.38 \text{ keV}$. About another half of the suppression is due to the missing of α_s correction in our result. As we have not calculated α_s corrections, the value in eq. (3.20) can be only consistently used for leading order in α_s calculations, like that in eq. (3.12).

To improve theoretical precision in NRQCD factorization, in ref. [17] the authors resum the relativistic correction to all orders for $e^+e^- \rightarrow \eta_c + \gamma$. This resummation in principle needs infinity number of nonperturbative parameters, which are modeled by assuming a generalized Gremm-Kapustin relation [24] that relates all LDMEs at higher order in v to $\langle \mathcal{O}^{\eta_{cex}}(1S_0^{[1]}) \rangle$. The resummed result is also listed in table 1. One can find that the SGF prediction is smaller than all predictions using NRQCD.

It is interesting to understand why there are less parameters in SGF comparing with NRQCD. Let us begin with the Gremm-Kapustin relation [25],

$$\langle \mathcal{P}^{\eta_{cex}}(1S_0^{[1]}) \rangle = \frac{1}{2} m(M - 2m) \langle \mathcal{O}^{\eta_{cex}}(1S_0^{[1]}) \rangle (1 + O(v^2)), \quad (4.4)$$

where $O(v^2)$ terms are matrix elements of operators with dynamical gluon fields. The Gremm-Kapustin relation tells us that nonperturbative behavior (infrared divergences in perturbation theory) of $\langle \mathcal{P}^{\eta_{cex}}(1S_0^{[1]}) \rangle$ can be expressed as linear combination of that of

²Note that, both in NRQCD and SGF, we have ignored intermediate states that explicitly include dynamical soft partons.

Table 1. The cross section $\sigma_{e^+e^- \rightarrow \eta_c + \gamma}$ calculated in SGF and NRQCD.

	SGF	NRQCD [16]	NRQCD [18]	NRQCD [17]
v^0		83.3	117	83.2
v^2		73.5	92	
$v^{\infty*}$				68.9
v^∞	26.2			

$\langle \mathcal{O}^{\eta_{cex}}(1S_0^{[1]}) \rangle$ and other LDMEs with dynamical gluon fields. Therefore, for the purpose of a valid factorization³, the introduction of $\langle \mathcal{P}^{\eta_{cex}}(1S_0^{[1]}) \rangle$ into the factorization formula is unnecessary.

In NRQCD factorization, one introduces nonperturbative quantities like $\langle \mathcal{P}^{\eta_{cex}}(1S_0^{[1]}) \rangle$ into factorization formula so that each term in eq. (4.1) has a definite power counting in v . But the price to pay is that one has lots of nonperturbative parameters.

In SGF, we do not introduce nonperturbative quantities similar to $\langle \mathcal{P}^{\eta_{cex}}(1S_0^{[1]}) \rangle$ (see discussion in Appendix A of ref. [2]). Leading contributions of these quantities are purely kinematic and they have been taken into account by coefficients of $\langle \overline{\mathcal{O}}_{1S_0^{[1]}}^{\eta_{cex}} \rangle$. Other contributions can be taken into account by nonperturbative SGDs with dynamical gluon fields, which are neglected in this paper but can be systematically included. In this sense, besides factorizing perturbative physics from nonperturbative physics, the SGF also factorizes kinematical physics from dynamical physics. It is the later effect that enables SGF to have less free parameters and thus to have a stronger predictive power.

5 Summary and outlook

In summary, we set up a general framework to apply the newly proposed SGF to exclusive quarkonium production or decay. Comparing with the NRQCD factorization, our method resums a series of relativistic corrections to all orders, which can reduce theoretical uncertainties. More importantly, our method has much less number of nonperturbative parameters and therefore has a stronger predictive power. To make it possible, the SGF not only factorizes perturbative physics from nonperturbative physics but also factorizes kinematical physics from dynamical physics.

Taking $\sigma_{e^+e^- \rightarrow \eta_c + \gamma}$ and $\Gamma(\eta_c \rightarrow \gamma + \gamma)$ as examples, we show how to use our new method to do phenomenological study. We find that these two quantities can be related to each other at all orders in v , and the relation is independent of charm quark mass. Based on experimental inputs, we provide a prediction for $\sigma_{e^+e^- \rightarrow \eta_c + \gamma}$. We also compare our prediction with predictions from NRQCD.

³A factorization formula holds as far as it includes a complete set of nonperturbative quantities that can decompose nonperturbative physics of target quantities.

Acknowledgments

We thank An-Ping Chen for many useful discussions. The work is supported in part by the National Natural Science Foundation of China (Grants No. U1832160, No. 11875071, No. 11975029, and No. 11905292).

References

- [1] G. T. Bodwin, E. Braaten, and G. P. Lepage, *Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium*, *Phys. Rev.* **D51** (1995) 1125–1171 [[hep-ph/9407339](#)] [[InSPIRE](#)]. [Erratum: *Phys. Rev.*D55,5853(1997)].
- [2] Y.-Q. Ma and K.-T. Chao, *New factorization theory for heavy quarkonium production and decay*, *Phys. Rev.* **D100** (2019) 094007 [[arXiv:1703.08402](#)].
- [3] E. Braaten and J. Lee, *Exclusive double charmonium production from e^+e^- annihilation into a virtual photon*, *Phys.Rev.* **D67** (2003) 054007 [[hep-ph/0211085](#)] [[InSPIRE](#)].
- [4] K.-Y. Liu, Z.-G. He, and K.-T. Chao, *Problems of double charm production in e^+e^- annihilation at $\sqrt{s} = 10.6\text{-GeV}$* , *Phys.Lett.* **B557** (2003) 45–54 [[hep-ph/0211181](#)] [[InSPIRE](#)].
- [5] **Belle Collaboration**, K. Abe *et al.*, *Observation of double $c\bar{c}$ production in e^+e^- annihilation at \sqrt{s} approximately 10.6-GeV*, *Phys.Rev.Lett.* **89** (2002) 142001 [[hep-ex/0205104](#)] [[InSPIRE](#)].
- [6] Z.-G. He, Y. Fan, and K.-T. Chao, *Relativistic corrections to J/ψ exclusive and inclusive double charm production at B factories*, *Phys.Rev.* **D75** (2007) 074011 [[hep-ph/0702239](#)] [[InSPIRE](#)].
- [7] G. T. Bodwin, J. Lee, and C. Yu, *Resummation of Relativistic Corrections to $e^+e^- \rightarrow J/\psi + \eta_c$* , *Phys. Rev.* **D77** (2008) 094018 [[arXiv:0710.0995](#)].
- [8] Y.-J. Zhang, Y.-j. Gao, and K.-T. Chao, *Next-to-leading order QCD correction to $e^+e^- \rightarrow J/\psi + \eta_c$ at $\sqrt{s} = 10.6\text{-GeV}$* , *Phys.Rev.Lett.* **96** (2006) 092001 [[hep-ph/0506076](#)] [[InSPIRE](#)].
- [9] B. Gong and J.-X. Wang, *QCD corrections to $J/\psi + \eta_c$ production in e^+e^- annihilation at $\sqrt{S} = 10.6\text{-GeV}$* , *Phys.Rev.* **D77** (2008) 054028 [[arXiv:0712.4220](#)] [[InSPIRE](#)].
- [10] F. Feng, Y. Jia, and W.-L. Sang, *Next-to-next-to-leading-order QCD corrections to $e^+e^- \rightarrow J/\psi + \eta_c$ at B factories*, [[arXiv:1901.08447](#)].
- [11] G. T. Bodwin, X. Garcia i Tormo, and J. Lee, *Factorization theorems for exclusive heavy-quarkonium production*, *Phys. Rev. Lett.* **101** (2008) 102002 [[arXiv:0805.3876](#)].
- [12] G. T. Bodwin, X. Garcia i Tormo, and J. Lee, *Factorization of low-energy gluons in exclusive processes*, *Phys. Rev.* **D81** (2010) 114005 [[arXiv:0903.0569](#)].
- [13] G. T. Bodwin, X. Garcia i Tormo, and J. Lee, *Factorization in exclusive quarkonium production*, *Phys. Rev.* **D81** (2010) 114014 [[arXiv:1003.0061](#)].
- [14] H. S. Chung, J. Lee, and C. Yu, *Exclusive heavy quarkonium + gamma production from e^+e^- annihilation into a virtual photon*, *Phys. Rev.* **D78** (2008) 074022 [[arXiv:0808.1625](#)].
- [15] D. Li, Z.-G. He, and K.-T. Chao, *Search for $C=$ charmonium and bottomonium states in $e^+e^- \rightarrow \gamma + X$ at B factories*, *Phys. Rev.* **D80** (2009) 114014 [[arXiv:0910.4155](#)] [[InSPIRE](#)].

- [16] W.-L. Sang and Y.-Q. Chen, *Higher Order Corrections to the Cross Section of $e^+e^- \rightarrow$ Quarkonium + gamma*, *Phys. Rev.* **D81** (2010) 034028 [[arXiv:0910.4071](#)] [[InSPIRE](#)].
- [17] Y. Fan, J. Lee, and C. Yu, *Resummation of relativistic corrections to exclusive productions of charmonia in e^+e^- collisions*, *Phys. Rev.* **D87** (2013) 094032 [[arXiv:1211.4111](#)] [[InSPIRE](#)].
- [18] Y.-J. Li, G.-Z. Xu, K.-Y. Liu, and Y.-J. Zhang, *Search for $C = +$ charmonium and XYZ states in $e^+e^- \rightarrow \gamma + H$ at BESIII*, *JHEP* **01** (2014) 022 [[arXiv:1310.0374](#)] [[InSPIRE](#)].
- [19] G.-Z. Xu, Y.-J. Li, K.-Y. Liu, and Y.-J. Zhang, *$\alpha_s v^2$ corrections to η_c and χ_{cJ} production recoiled with a photon at e^+e^- colliders*, *JHEP* **10** (2014) 71 [[arXiv:1407.3783](#)] [[InSPIRE](#)].
- [20] L.-B. Chen, Y. Liang, and C.-F. Qiao, *NNLO QCD corrections to $\gamma + \eta_c(\eta_b)$ exclusive production in electron-positron collision*, *JHEP* **01** (2018) 091 [[arXiv:1710.07865](#)].
- [21] G. T. Bodwin, H. S. Chung, D. Kang, J. Lee, and C. Yu, *Improved determination of color-singlet nonrelativistic QCD matrix elements for S-wave charmonium*, *Phys. Rev.* **D77** (2008) 094017 [[arXiv:0710.0994](#)].
- [22] **Particle Data Group**, M. Tanabashi *et al.*, *Review of Particle Physics*, *Phys. Rev.* **D98** (2018) 030001.
- [23] V. V. Braguta, *Exclusive $C=+$ charmonium production in $e^+e^- \rightarrow H + \gamma$ at B-factories within light cone formalism*, *Phys. Rev.* **D82** (2010) 074009 [[arXiv:1006.5798](#)] [[InSPIRE](#)].
- [24] G. T. Bodwin, D. Kang, and J. Lee, *Reconciling the light-cone and NRQCD approaches to calculating $e^+e^- \rightarrow J/\psi + \eta_c$* , *Phys.Rev.* **D74** (2006) 114028 [[hep-ph/0603185](#)] [[InSPIRE](#)].
- [25] M. Gremm and A. Kapustin, *Annihilation of S wave quarkonia and the measurement of alpha-s*, *Phys. Lett.* **B407** (1997) 323–330 [[hep-ph/9701353](#)] [[InSPIRE](#)].