

# Evolution implications of neutron star magnetic fields: inferred from pulsars and cyclotron lines of HMXBs

Chang-Qing Ye<sup>1</sup> • De-Hua Wang<sup>1,\*</sup> •  
Cheng-Min Zhang<sup>2,3,4</sup> • Zhen-Qi Diao<sup>1</sup>

## Abstract

The evolution of neutron star (NS) magnetic field (B-field) has long been an important topic, which is still not yet settled down. Here, we analyze the NS B-fields inferred by the cyclotron resonance scattering features (CRSFs) for the high mass X-ray binaries (HMXBs) and by the magnetic dipole model for the spin-down pulsars. We find that the B-fields of both the 32 NS-HMXBs and 28 young pulsars with the supernova remnants follow the log-normal distributions, with the average values of  $3.4 \times 10^{12}$  G and  $4.1 \times 10^{12}$  G respectively, which are further verified to come from the same continuous distribution by the statistical tests. These results declaim that the two methods of measuring NS B-fields are reliable for the above two groups of samples. In addition, since the NS-HMXBs have experienced the spin-down phase as the normal pulsars without accretion and then the spin-up phase by accretion, their ages should be about million years (Myrs). Our statistical facts imply that the B-fields of NS-HMXBs have little decayed in their non-accretion spin-down phases of  $\sim$  Myrs, as well as in their accretion phases of  $\sim 0.1$  Myrs.

**Keywords** X-rays: binaries—stars: neutron—pulsars: general—magnetic field

---

Chang-Qing Ye

De-Hua Wang

Cheng-Min Zhang

Zhen-Qi Diao

<sup>1</sup>School of Physics and Electronic Science, Guizhou Normal University, Guiyang 550001, China. \*wangdh@gznu.edu.cn

<sup>2</sup>CAS Key Lab of FAST, National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China.


<sup>3</sup>School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China.

<sup>4</sup>Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, Beijing 100012, China.

## 1 Introduction

The magnetic fields (B-fields) of neutron stars (NSs) play the significant roles in many aspects of pulsar phenomena, e.g., the electromagnetic emissions from the radio to high energy wavebands (Harding & Kalapotharakos 2017; Manchester et al. 2017; Becker et al. 2018). Until now, more than 2700 radio pulsars (Lorimer et al. 2019) and over 300 accreting NSs in high/low mass X-ray binaries (HMXBs/LMXBs) (Liu et al. 2007; Walter et al. 2015) have been observed. It is generally accepted that the B-fields of NSs dramatically decay in the binary accretion phase, while those of the isolated NSs should not decay, based on which the low B-field distributions of the double NSs ( $B \sim 10^9 - 10^{10}$  G, see Yang et al. 2019) and millisecond pulsars (MSPs,  $B \sim 10^{7.5} - 10^9$  G, see Pan et al. 2018) are very well understood by the binary accretion (Bhattacharya & van den Heuvel 1991; Zhang 2016; van den Heuvel 2017). However, there are still debated issues on the evolutions of NS B-fields: In what extent the B-fields of the isolated NSs have no decay? If the B-fields decay in the accretion phase, can we estimate the extent of the decay for NSs in HMXBs as an inverse relation to the accretion mass proposed (Shibazaki et al. 1989; Kulkarni 1986; Zhang & Kojima 2006)? To answer the above questions, one could evaluate the evolution mechanism of NS B-fields, e.g., the Ohmic dissipation (Geppert & Urpin 1994), Hall-drift effect (Geppert & Rheinhardt 2002; Cumming et al. 2004), screening by diamagnetism of accreting matter (Cumming et al. 2001; Lovelace et al. 2005), and accretion flow to cause the B-field decay (van den Heuvel & Bizaraki 1995; Melatos & Phinney 2001; Konar & Choudhuri 2004; Payne & Melatos 2004; Zhang & Kojima 2006).

There exist the two methods for measuring the NS B-fields, the magnetic dipole model for the spin-down pulsars (Lyne & Graham-Smith 2012), and cyclotron resonance scattering features (CRSFs) for NS-HMXBs



**Fig. 1** Diagram of the NS B-field versus spin period. The data of NS-HMXBs and pulsars are taken from Table 1 and ATNF pulsar catalog (Manchester et al. 2005), respectively. The shadow area indicates the  $B$  range of  $\sim 10^{12} - 10^{13}$  G, in which the B-fields of NS-HMXBs are measured. The solid line and dashed-dot line represent the spin-up line and death line of pulsars as defined by Bhattacharya & van den Heuvel (1991).

(Mészáros 1992). To evaluate the B-field evolution of NSs, we should have the good measurements of both the B-fields and their ages, the young pulsars with the supernova remnants (SNRs) and old NSs in HMXBs. The age of SNR pulsar is less than 0.3 million years (Myrs) (Ferrand & Safi-Harb 2012), which may retain the B-field distribution similar to those as they are born, whereas the NSs in HMXBs have experienced the spin-down and accretion induced spin-up phase with the age of  $\sim$  Myrs. Thus, the comparison of B-fields of both groups could acquire the information of B-field evolutions of NSs.

This paper aims to probe the NS B-field evolution of NS-HMXBs and SNR pulsars by comparing their B-field distributions, inferred by CRSFs and by magnetic dipole model, respectively. The structure of the paper is organized as follows: In Section 2, we introduce the B-field data of NS-HMXBs and SNR pulsars. In Section 3, we analyze the B-field distributions of the two group of samples, where the statistical tests are performed to see their properties. Finally, we present the discussions and conclusions in Section 4.

## 2 Two methods of NS magnetic field measures

### 2.1 B-fields of NS-HMXBs by CRSFs

For the accretion powered X-ray pulsars in HMXBs, NSs emit X-rays (e.g.,  $\sim 1 - 100$  keV) by accreting

matters from their companions (Caballero et al. 2012; Walter et al. 2015). In the strong B-field regime of NS surface, the electron energy is quantized to the discrete Landau levels, which experiences the resonance scattering with photons and results in the absorption features in the X-ray energy spectrum (Trümper et al. 1978), i.e., the energy of CRSFs:

$$E_c = \hbar \frac{eB}{m_e c} = 11.6(\text{keV}) \times B_{12} \quad (1)$$

where  $B_{12} \equiv \frac{B}{10^{12} \text{ G}}$ , with the electron mass  $m_e$  and charge  $e$ , as well as the speed of light  $c$  and Planck constant  $\hbar$ . The CRSFs are generally considered to form around NS surface (Mészáros 1992; Schönherr et al. 2007; Becker et al. 2012), so the observed energy  $E_c$  are gravitationally red-shifted and corrected to

$$E_n = nE_c = (1+z)E_{n,\text{obs}} \quad (2)$$

where  $n$  is the quantum number of the energy level, corresponding to the fundamental and harmonics of CRSF given by  $n = 1, 2, 3, \dots$ , and the parameter  $z$  is the gravitational red-shift expressed as,

$$z = \frac{1}{\sqrt{1 - \frac{2GM}{Rc^2}}} - 1, \quad (3)$$

where  $G$  is the Newtonian gravitational constant. For the typical NS mass of  $M = 1.4 M_\odot$  and radius of  $R = 10$  km, the gravitational red-shift parameter is obtained as  $z = 0.3$  (Shapiro & Teukolsky 1983; Maitra et al. 2017).

We search for the NS-HMXBs which have been detected with CRSFs, and focus on the 32 sources with the confirmed detection. Table 1 lists the CRSF energies ( $E_c$ , fundamental and harmonic) and NS spin periods ( $P$ ) of these sources that are collected from the published literature (Makishima et al. 1999; Coburn et al. 2002; Caballero et al. 2012; Walter et al. 2015). Furthermore, we infer the B-fields of NS-HMXBs by equation (1)-(3) with the CRSFs in Table 1 by assuming they emit around the NS surface (Mészáros 1992; Becker et al. 2012). Figure 1 illustrates the NS B-field versus spin period distribution of the samples.

### 2.2 B-fields of pulsars by their periods and derivatives

For the spin-down pulsars, their losses of rotational energies is presumed by the magnetic dipole radiation, based on which their B-fields can be derived by their observed spin periods ( $P$ ) and derivatives  $\dot{P}$ , as given in the references in the case of perpendicular rota-

**Table 1** CRSF energy and inferred B-fields of NS-HMXBs

Source (32)	$E_c$ (keV)	B-field ( $10^{12}$ G)	$P$ (s)	REFS. <sup>a</sup>
Swift J1626.6-5156	10	1.12	15.35	[1, 1]
XMMU J05	10	1.12	61.23	[2, 33]
KS 1947+300	12.50	1.40	18.70	[3, 34]
4U 0115+634	14, 24, 36, 48, 62	1.57	3.61	[4, 35]
XTEJ1829-098	15	1.68	7.84	[5, 5]
IGR J17544-2619	17	1.91	–	[6, 36]
4U 1907+09	19, 39	2.12	437–440	[7, 37]
4U 1538-52	22, 47	2.47	528–530	[8, 38]
IGR J18179-1621	22	2.47	11.82	[9, 39]
IGRJ18027-2016	23	2.58	139.61	[10, 40]
2S 1553-542	23.50	2.63	9.27	[11, 11]
Vela X-1	25, 50	2.80	283	[12, 41]
V 0332+53	27, 51, 74	3.03	4.37	[13, 42]
SMC X-2	27	3.03	2.37	[14, 14]
Cep X-4	28,45	3.14	66.25	[15, 43]
X per	29	3.25	835	[16, 44]
IGR J16393-4643	29.3	3.28	904	[17, 17]
Cen X-3	30	3.36	4.82	[18, 45]
J 16493-4348	30	3.36	1093	[19, 46]
RX J0520.5-6932	31.50	3.53	8.03	[20, 20]
RX J0440.9-4431	32	3.59	202.50	[21, 47]
MXB 0656-072	33	3.70	160	[22, 48]
IGR J19294+1816	35.50	3.98	12.44	[23, 49]
XTE J11946+274	36	4.03	15.80	[24, 50]
4U 1626-67	37	4.15	7.67	[25, 51]
GX 301-2	37	4.15	675–700	[26, 52]
Her X-1	39, 73	4.37	1.24	[27, 53]
MAXI J1409-619	44, 73, 128	4.93	500	[28, 54]
1A 0535+262	45, 100	5.04	103	[29, 55]
GX 304-1	54	6.05	272	[30, 56]
1A 1118-615	55, 112?	6.16	405–407	[31, 57]
GRO J1008-57	76	8.52	93.60	[32, 58]

Note:  $E_c$ —CRSF energy of the fundamental and multiple harmonics;  $P$ —Spin period of NS; <sup>a</sup>—The first reference of each source is for  $E_c$  (for some sources it record the fundamental and multiple harmonics CRSFs observed simultaneously), and the second reference is for  $P$ .

REFS.: [1]. DeCesar et al. 2013; [2]. Manousakis et al. 2009; [3]. Fürst et al. 2014; [4]. Wheaton W A et al. 1979; Heindl et al. 1999; Ferrigno et al. 2011; [5]. Shtykovsky et al. 2019; [6]. Bhalerao et al. 2015; [7]. Rivers et al. 2010; Cusumano et al. 1998; [8]. Rodes-Roca et al. 2009; Clark et al. 1990; [9]. Li J et al. 2012; Tuerler et al. 2012; [10]. Lutovinov et al. 2017; [11]. Tsygankov et al. 2016; [12]. La Barbera et al. 2003; Kreykenbohm et al. 1999; Kendziorra et al. 1992; [13]. Nakajima et al. 2010; Makishima et al. 1990; Coburn et al. 2005; [14]. Jaisawal et al. 2016; [15]. Jaisawal et al. 2015; Mihara et al. 1991; [16]. Coburn et al. 2001; [17]. Bodaghee et al. 2016; [18]. Burderi et al. 2000; Nagase et al. 1992; [19]. D’A ‘i A et al. 2011; [20]. Tendulkar et al. 2014; [21]. Tsygankov et al. 2012; [22]. McBride et al. 2006; Heindl et al. 2003; [23]. Roy et al. 2017; [24]. Heindl et al. 2001; [25]. Orlandini et al. 1998; [26]. Suchy et al. 2012; Makishima et al. 1992; [27]. Enoto et al. 2008; Trüemper et al. 1978; [28]. Orlandini et al. 2012; [29]. Terada et al. 2006; [30]. Yamamoto et al. 2011; Mihara et al. 2010; [31]. Doroshenko et al. 2010; [32]. Yamamoto et al. 2014; [33]. Manousakis et al. 2009; [34]. Chakrabarty et al. 1995; Galloway et al. 2004; [35]. Rappaport et al. 1978 [36]. Clark et al. 2009 [37]. in’t Zand et al. 1998 [38]. Clark et al. 2000, 1994 [39]. Bozzo et al. 2012; [40]. Mason et al. 2011; [41]. Quintrell et al. 2003; [42]. Zhang et al. 2005; Qu et al. 2005; [43]. Mihara et al. 1991; [44]. White et al. 1976; Delgado-Martí et al. 2001; [45]. van der Meer et al. 2007; [46]. Pearlman et al. 2013; [47]. Liu et al. 2005; [48]. McBride et al. 2006; [49]. Rodriguez et al. 2009; Corbet et al. 2009; [50]. Smith et al. 1998; Wilson et al. 2003; [51]. Kii et al. 1986; [52]. Kaper et al. 2006; Koh et al. 1997; [53]. Trüemper et al. 1978; [54]. Yamamoto et al. 2010; [55]. Rosenberg et al. 1975; Finger et al. 1996; [56]. Huckle et al. 1977; Priedhorsky et al. 1983; [57]. Staubert et al. 2011; [58]. Stollberg et al. 1993; Kuehnel et al. 2012.

**Table 2** B-field statistics of 32 NS-HMXBs and 28 SNR pulsars

Category	Number	Range of B	$\langle B \rangle$	$\tilde{B}$	$\sigma_B$
( $\times 10^{12}$ G)					
NS-HMXBs	32	1.12 ~ 8.52	3.39	3.27	1.58
SNR pulsars	28	1.11 ~ 9.91	4.09	3.60	2.14

Note:  $\langle B \rangle$ —mean of  $B$ ;  $\tilde{B}$ —median of  $B$ ;  $\sigma_B$ —standard deviation of  $B$ .

tor (Ostriker & Gunn 1969; Manchester & Taylor 1977; Lyne & Graham-Smith 2012):

$$B = \left( \frac{3Ic^3P\dot{P}}{8\pi^2R^6} \right)^{1/2} \simeq 3.2 \times 10^{19} (P\dot{P})^{1/2} \text{ (G)}, \quad (4)$$

where  $I$  and  $R$  stand for the moment of inertia and radius of NS with the conventional values of  $I = 10^{45} \text{g}\cdot\text{cm}^2$  and  $R = 10 \text{ km}$ .

In order to investigate the B-field evolution of NSs, we compare the B-field distribution of the NS-HMXBs with that of the young pulsars with SNRs, whose ages are generally less than 0.3 Myr. Until now, there are 61 SNR pulsars have been recorded in ATNF pulsar catalog (Manchester et al. 2005), in which 28 sources share the B-fields in the ranges of  $\sim 10^{12} - 10^{13} \text{ G}$  (as compared to those of CRSF sources, see Table 1) that are analyzed in the paper. These samples are labeled in the diagram of B-field versus spin period, as plotted in Figure 1.

**Table 3** S-W Test Results of NS B-field Distributions

Category	Number	$S - W$ ( $p$ -value)	Reject $H_0$
NS-HMXB	32	0.56	No
SNR pulsar	28	0.97	No

Note:  $H_0$  is the null hypothesis that the data follows a log-normal distribution, with the confidence level parameter  $\alpha = 0.05$ .


**Table 4** K-S Test Results of NS B-field Distributions

Category	Number	$K - S$ ( $p$ -value)	Reject $H_0$
NS-HMXB	32	0.74	No
SNR pulsar	28		

Note:  $H_0$  is the null hypothesis that the two groups of data come from the same continuous distribution, with the confidence level parameter  $\alpha = 0.05$ .

### 3 Analysis on the B-field distributions

The NS B-field statistical quantities of the samples, including the range, mean ( $\langle B \rangle$ ), median ( $\tilde{B}$ ) and standard deviation ( $\sigma_B$ ), are obtained and summarized in Table 2. It can be seen that the NS-HMXBs share the similar  $\langle B \rangle$  and  $\tilde{B}$  values ( $\langle B \rangle \sim 3.39 \times 10^{12} \text{ G}$  and  $\tilde{B} \sim 3.27 \times 10^{12} \text{ G}$ ) to those of the SNR pulsars



**Fig. 2** The CDF curves of B-fields for the NS-HMXBs and SNR pulsars.

( $\langle B \rangle \sim 4.09 \times 10^{12} \text{ G}$  and  $\tilde{B} \sim 3.60 \times 10^{12} \text{ G}$ ). In order to investigate whether the two group samples of B-fields come from the same continuous distribution, we make the following two steps: (I). testing the two distributions to be the log-normal or not, and (II). testing the two group B-fields to come from the same continuous distribution or not if both distributions follow the log-normal.

(I). Table 3 lists the Shapiro-Wilk (S-W) test results (Corder & Foreman 2009) of the two group B-field samples, from which it can be noticed that the B-fields of both the NS-HMXBs and SNR pulsars are compatible with the log-normal distributions at the 95 percent confidence level. The expressions of the mean and standard deviation of the log-normal distributions are  $(28.75 \pm 0.47) \log_e(\text{G})$  for the NS-HMXBs, and  $(28.91 \pm 0.52) \log_e(\text{G})$  for the SNR pulsars, respectively.

(II). Furthermore, we take the Kolmogorov-Smirnov (K-S) test (Corder & Foreman 2009) to compare the B-field distributions of the NS-HMXBs and SNR pulsars, and show the results in Table 4, where the consistency of both distributions is obtained at the 95 percent confidence level. In order to more intuitively represent the two group B-fields come from the same log-normal distribution, we also show the corresponding B-field cumulative distribution function (CDF) curves of the two group samples in Figure 2, the appearance of which clearly strengthens the conclusion of the same distributions.

## 4 Discussions and Conclusions

In order to investigate whether the NS B-field decay or evolve in million years, we studied the two distinctive classes of NS samples: NS-HMXBs that are the accretion X-ray binary systems with the ages of  $\sim$  Myrs (Bhattacharya & van den Heuvel 1991), and SNR pulsars that are the pulsating objects with the ages of less than 0.3 Myrs (Ferrand & Safi-Harb 2012). The B-fields of the formers are measured directly by CRSFs and those of the latters are derived by the magnetic dipole model. The following facts and conclusions are noticed and obtained:

- The S-W test indicates that the B-fields of both the 32 NS-HMXBs and 28 SNR pulsars follow the log-normal distributions (see Table 3), and the two group B-fields are further verified to come from the same continuous distribution by the K-S test (see Table 4). According to the evolutionary theory of the X-ray binaries, the NSs in HMXBs may firstly experience the spin-down phase for  $\sim$  Myrs by the magnetic dipole emissions (Bhattacharya & van den Heuvel 1991; Lorimer 2008), and then they can switch to the accretion induced spin-up phase for  $\sim$  0.1 Myr (Bhattacharya & van den Heuvel 1991). Therefore, the statistical test facts imply that the B-fields of NSs in HMXBs have little decayed in their non-accretion spin-down phases of  $\sim$  Myrs, as well as in their accretion phases of  $\sim$  0.1 Myrs. Furthermore, it is true that the NS B-field decays in the binary accretion phase present a positive correlation with how much mass accreted, where the surface field decreases both by Ohmic decay and submergence by the inflowing matter (Bisnovatyi-Kogan et al. 1974; Shibazaki et al. 1989; Bhattacharya & van den Heuvel 1991; Geppert & Urpin 1994; Urpin & Geppert 1995; Zhang & Kojima 2006). In particular, it is suggested that for an accretion rate of  $\sim 10^{-10} M_{\odot} \text{ yr}^{-1}$ , which corresponds to an accreted mass of  $\sim 10^{-5} M_{\odot}$  during  $\sim$  0.1 Myr, the NS surface field may decay by  $\sim$  two orders of magnitude (Urpin & Geppert 1995; Urpin et al. 1998). Therefore, the consistent B-field distributions between the SNR pulsars and NS-HMXBs imply that the actually accreted mass of HMXBs may be less than  $\sim 10^{-5} M_{\odot}$ .
- It should be noticed that the B-field inferred from the CRSF measurement is the B-field around the NS surface, while the B-field obtained from the measurement of  $P$  and  $\dot{P}$  values is calculated by considering the energy loss at the NS light cylinder and then extrapolates the field strength back to the star's surface assuming the flat space-time  $1/r^3$  radial dependence of the field. Thus, for the HMXBs, equation (2) takes

into account gravitational red-shift, while general relativity effects are not taken into consideration for the B-field of isolated pulsars. The different B-field measures for the NS-HMXBs and SNR pulsars present the same distributions, and we declaim that the two methods, by the CRSF and magnetic dipole model, are successfully applied to obtain the B-fields around NS surfaces. In general, the CRSF indicates the local B-field around NS surface, while the B-field inferred by the magnetic dipole model represents the large scale value. The similarity of both B-field values might imply that both the measures and selected NS parameters should be in the reasonable regimes, at least valid in the statistical sense.

The B-field evolution mechanism of NS is critical for understanding the emissions and binary interactions of various pulsars, thus our conclusions, based on the B-field measures of different methods and statistical tests, would provide the robust clues or constraints for the time scale of B-field decay.

**Acknowledgements** This work is supported by the National Natural Science Foundation of China (Grant No. 11703003 and No. U1731238), the Science and Technology Foundation of Guizhou Province (Grant No. [2017]5726), NAOC-Y834081V01.

## References

- Becker, P. A., Klochkov, D., Schönherr, G., et al.: *Astron. Astrophys.*, 544, A123 (2012)
- Becker, W., Kramer, M., & Sesana, A.: *Space Sci. Rev.*, 214, 30 (2018)
- Bhalerao, V., Romano, P., Tomsick, J., et al.: *Mon. Not. R. Astron. Soc.*, 447, 2274 (2015)
- Bhattacharya, D. & van den Heuvel, E. P. J.: *Phys. Rep.*, 203, 1 (1991)
- Bisnovatyi-Kogan, G. S., Komberg, B. V.: *Soviet Astron.*, 18, 217, (1974),
- Bodaghee, A., Tomsick, J. A., Fornasini, F. M., et al.: *Astrophys. J.*, 2016, 823, 146 (2016)
- Bozzo, E., Ferrigno, C., Türler, M., et al.: *Astron. Astrophys.*, 545, A83 (2012)
- Burderi, L., Di Salvo, T., Robba, N. R., et al.: *Astrophys. J.*, 530, 429 (2000)
- Caballero, I., Wilms, J.: *Memorie della Societa Astronomica Italiana*, 83, 230 (2012)
- Chakrabarty, D., Koh, T., Bildsten, L., et al.: *Astrophys. J.*, 446, 826 (1995)
- Clark, G. W., Woo, J. W., Nagase, F., et al.: *Astrophys. J.*, 353, 274 (1990)
- Clark, G. W., Woo, J. W., Nagase, F., *Astrophys. J.*, 422, 336 (1994)
- Clark, G. W., *Astrophys. J.*, 542, L131 (2000)
- Clark, D. J., Hill, A. B., Bird, A. J., et al.: *Mon. Not. R. Astron. Soc.*, 399, L113 (2009)
- Coburn, W., Heindl, W. A., Gruber, D. E., et al.: *Astrophys. J.*, 552, 738 (2001)
- Coburn, W., Heindl, W. A., Rothschild, R. E., et al.: *Astrophys. J.*, 580, 394 (2002)
- Coburn, W., Kretschmar, P., Kreykenbohm, I., et al.: *The Astronomer's Telegram*. 381 (2005)
- Corbet, R. H. D., Krimm, H. A.: *The Astronomer's Telegram*, 2008 (2009)
- Corder, G. W., & Foreman, D. I: *Nonparametric Statistics for Non-Statisticians: A Step-by-Step Approach* (New York: Wiley) (2009)
- Cumming, A., Zweibel, E., & Bildsten, L.: *Astrophys. J.*, 557, 958 (2001)
- Cumming A., Arras P., Zweibel E.: *Astrophys. J.*, 609, 999 (2004)
- Cusumano, G., di Salvo, T., Burderi, L., et al.: *Astron. Astrophys.*, 338, L79 (1998)
- D'Ai, A., Cusumano, G., La Parola, V., et al.: *Astron. Astrophys.*, 532, A73 (2011)
- DeCesar, M. E., Boyd, P. T., Pottschmidt, K., et al.: *Astrophys. J.*, 762, 61 (2013)
- Delgado-Martí, H., Levine, A. M., Pfahl, E., et al, *Astrophys. J.*, 546, 455 (2001)
- Doroshenko, V., Suchy, S., Santangelo, A., et al.: *Astron. Astrophys.*, 515, 1 (2010)
- Enoto, T., Makishima, K., Terada, Y., et al.: *Publ. Astron. Soc. Jpn.*, 60, 57 (2008)
- Fürst, F., Pottschmidt, K., Wilms, J., et al.: *Astrophys. J.*, 784, L40 (2014)
- Ferrand, G., & Safi-Harb, S.: *Advances in Space Research*, 49, 1313 (2012)
- Ferrigno, C., Falanga, M., Bozzo, E., et al.: *Astron. Astrophys.*, 532, A76 (2011)
- Finger, M. H., Wilson, R. B., Harmon, B. A.: *Astrophys. J.*, 459, 288 (1996)
- Galloway, D. K., Morgan, E. H., Levine, A. M.: *Astrophys. J.*, 613, 1164 (2014)
- Geppert, U., & Rheinhardt, M.: *Astron. Astrophys.*, 392, 1015 (2002)
- Geppert, U., & Urpin, V.: *Mon. Not. R. Astron. Soc.*, 271, 490 (1994)
- Harding, A. K., & Kalapotharakos, C.: *Proceedings of the 7th International Fermi Symposium*, held 15-20 October 2017, in Garmisch-Partenkirchen, Germany (2017).
- Heindl, W. A., Coburn, W., Gruber, D. E., et al.: *Astrophys. J.*, 1999, 521, L49 (1999)
- Heindl W. A., Coburn, W., Gruber, D. E., et al.: *Astrophys. J.*, 563, L35 (2001)
- Heindl, W., Coburn, W., Kreykenbohm, I., et al.: *The Astronomer's Telegram*, 200 (2003)
- Huckle, H. E., Mason, K. O., White N E, et al.: *Mon. Not. R. Astron. Soc.*, 180, 21 (1977)
- in't Zand, J. J. M., Baykal, A., Strohmayer, T. E., *Astrophys. J.*, 496, 386 (1998)
- Jaisawal, G. K., Naik, S.: *Mon. Not. R. Astron. Soc.*, 2015, 453, L21 (2015)
- Jaisawal, G. K., Naik, S. *Mon. Not. R. Astron. Soc.*, 461, L97 (2016)
- Kaper, L., van der Meer, A., Najarro, F.: *Astron. Astrophys.*, 457, 595 (2006)
- Kendziorra, E., Mony, B., Kretschmar, P., et al.: *NASA Conference Publication*, 3137 (1992)
- Kii, T., Hayakawa, S., Nagase, F., et al.: *Publ. Astron. Soc. Jpn.*, 38, 751 (1986)
- Koh, D. T., Bildsten, L., Chakrabarty, D., et al.: *Astrophys. J.*, 479, 933 (1997)
- Konar, S., & Choudhuri, A. R.: *Mon. Not. R. Astron. Soc.*, 348, 661 (2004)
- Kreykenbohm, I., Kretschmar, P., Wilms, J., et al.: *Astron. Astrophys.*, 341, 141 (1999)
- Kuehnel, M., Mueller, S., Kreykenbohm, I., et al.: *The Astronomer's Telegram*. 4564 (2012)
- Kulkarni, S. R.: *Astrophys. J. Lett.*, 306, L85 (1986)
- La Barbera, A., Santangelo, A., Orlandini, M., et al.: *Astron. Astrophys.*, 400, 993 (2003)
- Li, J., Zhang, S., Torres, D. F., et al.: *Mon. Not. R. Astron. Soc.*, 426, L16 (2012)
- Liu, Q. Z., van Paradijs, J., van den Heuvel, E. P. J. *Astron. Astrophys.*, 442, 1135 (2005)
- Liu, Q. Z., van Paradijs, J., & van den Heuvel, E. P. J.: *Astron. Astrophys.*, 469, 807 (2007)
- Lorimer, D. R.: *Living Reviews in Relativity*, 11, 8 (2008)
- Lorimer, D. R., Pol, N., Rajwade, K., et al.: *Bull. Am. Astron. Soc.*, 51, 261 (2019)
- Lovelace, R. V. E., Romanova, M. M., & Bisnovatyi-Kogan, G. S.: *Astrophys. J.*, 625, 957 (2005)
- Lutovinov, A. A., Tsygankov, S. S., Postnov, K. A., et al.: *Mon. Not. R. Astron. Soc.*, 466, 593 (2017)
- Lyne, A. G., Francis Graham-Smith: *Pulsar Astronomy*, Cambridge; New York: Cambridge University Press (2012)

- Mészáros, P.: High-energy radiation from magnetized neutron stars. Chicago, Univ Chicago Press, 25 (1992)
- Maitra, C.: *Journal of Astrophysics and Astronomy*, 38, 50 (2017)
- Makishima, K., Mihara, T., Ishida, M., et al.: *Astrophys. J.*, 365, L59 (1990)
- Makishima, K., Mihara, T. *Frontiers Science Series*, 23 (1992)
- Makishima, K., Mihara, T., Nagase, F., et al.: *Astrophys. J.*, 525, 978 (1999)
- Manchester, R. N. & Taylor, J. H.: *Pulsars*, Freeman, San Francisco (1977)
- Manchester, R. N., Hobbs, G. B., Teoh, A., Hobbs, M.: *Astron. J.*, 129 (2005)
- Manchester, R. N.: *Journal of Astrophysics and Astronomy*, 38, 42 (2017)
- Manousakis, A., Walter, R., Audard, M., et al.: *American Institute of Physics Conference Series*, 1126, 325 (2009)
- Manousakis, A., Walter, R., Audard, M., et al.: *Astron. Astrophys.*, 498, 217 (2009)
- Mason, A. B., Norton, A. J., Clark, J. S., et al.: *Astron. Astrophys.*, 532, A124 (2011)
- McBride, V. A., Wilms, J., Coe, M. J., et al.: *Astron. Astrophys.*, 451, 267 (2006)
- Melatos, A., & Phinney, E. S.: *Proc. Astron. Soc. Aust.*, 18, 421 (2001)
- Mihara, T., Makishima, K., Kamijo, S., et al.: *Astrophys. J.*, 379, L61 (1991)
- Mihara, T., Yamamoto, T., Sugizaki, M., et al.: *The Astronomer's Telegram*, 2796 (2010)
- Nagase, F., Corbet, R. H. D., Day, C. S. R., et al.: *Astrophys. J.*, 396, 147 (1992)
- Nakajima, M., Mihara, T., Makishima, K.: *Astrophys. J.*, 710, 1755 (2010)
- Orlandini, M., Dal Fiume, D., Frontera, F., et al.: *Astrophys. J.*, 500, L163 (1998)
- Orlandini, M., Frontera, F., Masetti, N. et al.: *Astrophys. J.*, 748, 86 (2012)
- Ostriker, J. P. & Gunn, J. F.: *Astrophys. J.*, 157, 1395(1969)
- Pan, Y. Y., Zhang, C. M., Song, L. M., et al.: *Mon. Not. R. Astron. Soc.*, 480, 692 (2018)
- Payne, D. J. B., Melatos, A.: *Mon. Not. R. Astron. Soc.*, 351, 569 (2004)
- Pearlman, A. B., Corbet, R., Pottschmidt, K. In, *American Astronomical Society meeting abstracts*, 221, 142 (2013)
- Priedhorsky, W. C., Terrell, J.: *Astrophys. J.*, 273, 709 (1983)
- Qu, J. L., Zhang, S., Song, L. M., et al.: *Astrophys. J.*, 629, L33 (2005)
- Quaintrell, H., Norton, A. J., Ash, T. D. C., et al.: *Astron. Astrophys.*, 401, 313 (2003)
- Rappaport, S., Clark, G. W., Cominsky, L., et al, *Astrophys. J.*, 224, L1 (1978)
- Rivers, E., Markowitz, A., Pottschmidt, K., et al.: *Astrophys. J.*, 2010, 709, 179 (2010)
- Rodes-Roca, J. J., Torrejn, J. M., Kreykenbohm, I., et al.: *Astron. Astrophys.*, 508, 395 (2009)
- Rodriguez, J., Tuerler, M., Chaty, S., et al.: *The Astronomer's Telegram*, 1998 (2009)
- Rosenberg, F. D., Eyles, C. J., Skinner, G. K., et al.: *Nature*, 256, 628 (1975)
- Roy J, Choudhury, M., Agrawa, P. C., *Astrophys. J.*, 848, 124 (2017)
- Schönherr, G., Wilms, J., Kretschmar, P., et al.: *Astron. Astrophys.*, 472, 353 (2007)
- Shapiro, S. L. & Teukolsky, S. A.: *Black Holes, White Dwarfs and Neutron Stars*. Wiley, New York (1983)
- Shibazaki, N., Murakami, T., Shaham, J., & Nomoto, K.: *Nature*, 342, 656 (1989)
- Shtykovsky, A. E., Lutovinov, A. A., Tsygankov, S. S., & Molkov, S. V.: *Mon. Not. R. Astron. Soc.*, 482, L14 (2019)
- Smith, D. A., Takeshima, T.: *The Astronomer's Telegram*, 1998, 36 (1998)
- Staubert, R., Pottschmidt, K., Doroshenko, V., et al.: *Astron. Astrophys.*, 527, A7 (2011)
- Stollberg, M. T., Finger, M. H., Wilson, R. B., et al.: *IAU Circ.*, 5836, 1 (1993)
- Suchy, S., Fürst, F., Pottschmidt, K., et al.: *Astrophys. J.*, 745, 124 (2012)
- Tendulkar, S. P., Fürst, F., Pottschmidt, K., et al.: *Astrophys. J.*, 795, 154 (2014)
- Terada, Y., Mihara, T., Nakajima, M., et al, *Astrophys. J.*, 648, L139 (2006)
- Trümper, J., Pietsch, W., Reppin, C., et al.: *Astrophys. J.*, 219, 105 (1978)
- Trümper, J., Kahabka, P., Oegelman H, et al, *Astrophys. J.*, 300, L63 (1986)
- Tsygankov, S. S., Krivonos, R. A., Lutovinov, A. A. *Mon. Not. R. Astron. Soc.*, 2012, 421, 2407 (2012)
- Tsygankov, S. S., Lutovinov, A. A., Krivonos, R. A., et al.: *Mon. Not. R. Astron. Soc.*, 457, 258 (2016)
- Tuerler, M., Chenevez, J., Bozzo, E., et al.: *The Astronomer's Telegram*, 3947 (2012)
- Urpin, V., & Geppert, U.: *Mon. Not. R. Astron. Soc.*, 275, 1117 (1995)
- Urpin, V., Konenkov, D., & Geppert, U.: *Mon. Not. R. Astron. Soc.*, 299, 73 (1998)
- van den Heuvel E. P. J. & Bitzaraki, O.: *Astron. Astrophys.*, 297, 41 (1995)
- van den Heuvel E. P. J.: *Journal of Astrophysics and Astronomy*, 38, 45 (2017)
- van der Meer, A., Kaper, L., van Kerkwijk, M. H., et al.: *Astron. Astrophys.*, 473, 523 (2007)
- Walter, R., Lutovinov, A. A., Bozzo, E., et al.: *Astron. Astrophys. Rev.*, 23, 2 (2015)
- Wheaton, W. A., Doty, J. P., Primini, F. A., et al.: *Nature*, 282, 240 (1979)
- White, N. E., Mason, K. O., Sanford, P. W., et al.: *Mon. Not. R. Astron. Soc.*, 176, 201 (1976)
- Wilson, C. A., Finger, M. H., Coe, M. J., et al.: *Astrophys. J.*, 584, 996
- Yamamoto, T., Nakajima, M., Yamaoka, K., et al.: *The Astronomer's Telegram*, 3070, 1 (2010)
- Yamamoto, T., Sugizaki, M., Mihara, T., et al.: *Publ. Astron. Soc. Jpn.*, 63, 751 (2011)
- Yamamoto, T., Mihara, T., Sugizaki, M., et al.: *Publ. Astron. Soc. Jpn.*, 66, 59 (2014)
- Yang, Y. Y., Zhang, C. M., Li, D., et al.: *Publ. Astron. Soc. Pac.*, 131, 064201 (2019)
- Zhang, C. M. & Kojima, Y.: *Mon. Not. R. Astron. Soc.*, 366, 137 (2006)

Zhang, C. M., in Alsabti, A. W. & Murdin, P., eds, Handbook of Supernovae, Springer International Publishing, p.1375 (2016)

Zhang, S., Qu, J. L., Song, L. M., et al.: *Astrophys. J.*, 2005, 630, L65 (2005)