

The globalization problem of the Hamilton-DeDonder-Weyl equations on a local k -symplectic framework

Oğul Esen[†], Manuel de León[‡], Cristina Sardón^{*}, Marcin Zajac^{**}

Department of Mathematics[†],
Gebze Technical University, 41400 Gebze, Kocaeli, Turkey.
oesen@gtu.edu.tr

Instituto de Ciencias Matemáticas, Campus Cantoblanco[‡],
Consejo Superior de Investigaciones Científicas
and
Real Academia Española de las Ciencias.
C/ Nicolás Cabrera, 13–15, 28049, Madrid, Spain.
mdeleon@icmat.es

Instituto de Ciencias Matemáticas, Campus Cantoblanco^{*},
Consejo Superior de Investigaciones Científicas.
C/ Nicolás Cabrera, 13–15, 28049, Madrid, Spain.
cristinasardon@icmat.es

Department of Mathematical Methods in Physics^{**},
Faculty of Physics. University of Warsaw,
ul. Pasteura 5, 02-093 Warsaw, Poland.
marcin.zajac@fuw.edu.pl

Abstract

In this paper we aim at addressing the globalization problem of Hamilton-DeDonder-Weyl equations on a local k -symplectic framework and we introduce the notion of *locally conformal k -symplectic (l.c.k-s.) manifolds*. This formalism describes the dynamical properties of physical systems that locally behave like multi-Hamiltonian systems. Here, we describe the local Hamiltonian properties of such systems, but we also provide a global outlook by introducing the global Lee one-form approach. In particular, the dynamics will be depicted with the aid of the Hamilton–Jacobi equation, which is specifically proposed in a l.c.k-s manifold.

Contents

1	Introduction	2
1.1	Locally conformal symplectic manifolds	2
1.2	Geometric Hamilton-Jacobi theory	4
1.3	The goal and the contents	5

2	k-symplectic framework	5
2.1	k-tangent bundle	5
2.2	k-symplectic manifolds	7
2.3	Dynamics on k-symplectic manifolds	8
2.4	Hamilton–Jacobi theory	10
3	Locally conformal k-symplectic framework	13
3.1	Locally conformal k-symplectic manifolds	13
3.2	L.c.k-s. structures on the Whitney sum of cotangent bundles	15
3.3	Dynamics in l.c.k-s. manifolds	17
3.4	Hamilton-Jacobi theory	17
4	An example	20
5	Conclusion and future work	23
6	Acknowledgements	23

1 Introduction

In this work, we address the globalization problem of local Hamiltonian-De Donder-Weyl (HDW) dynamics on k -symplectic spaces, and we propose the associated Hamilton-Jacobi theory on k -locally conformal symplectic manifolds. In the classical symplectic framework, the glueing problem of local Hamiltonian dynamics was investigated by Vaismann [32] in terms of locally conformal symplectic (l.c.s.) manifolds, being the latter introduced by Lee in [17]. Our interest in this work is to generalize this approach to the case of k -symplectic manifolds. In that direction, we shall propose a new geometric structure, which we will call a locally conformal k -symplectic (l.c.k-s.) structure, to provide a globalization of the local HDW equations. Further, we shall propose a Hamilton-Jacobi formalism for the dynamics in such manifold. Let us now recall some introductory concepts to follow along the present work and to state our goal more explicitly.

1.1 Locally conformal symplectic manifolds

Consider an even dimensional manifold M equipped with a non-degenerate two-form ω . The manifold M is called a locally conformal symplectic (l.c.s.) if M has an open cover $\{U_\alpha\}$ and there exists a family of smooth functions σ_α on each chart such that $d(e^{-\sigma_\alpha}\omega|_\alpha)$ vanishes identically [6, 32]. Here, $\omega|_\alpha$ denotes the restriction of the two-form ω to the chart U_α . In this picture, the local two-form defined by $\omega_\alpha := e^{-\sigma_\alpha}\omega|_\alpha$ turns out to be symplectic and the pair $(U_\alpha, \omega_\alpha)$ becomes a symplectic manifold.

Referring to this symplectic framework, for a local Hamiltonian function h_α , the local geometric Hamilton equation is

$$\iota_{X_\alpha}\omega_\alpha = dh_\alpha. \tag{1}$$

Notice that the Hamiltonian vector fields X_α are still the same even if we multiply the local Hamilton equation (1) by scalars $\lambda_{\beta\alpha}$, in overlapping local charts U_α and U_β . These scalars can be written as $\lambda_{\beta\alpha} = e^{\sigma_\alpha - \sigma_\beta}$ for local functions σ_α 's. This leads to the determination of a line bundle L over the manifold M . Accordingly, we can glue up the local Hamiltonian functions according to the following transition relation

$$h_\beta = e^{\sigma_\alpha - \sigma_\beta} h_\alpha. \quad (2)$$

This family defines a line bundle valued function \tilde{h} on M . Further, we define local functions $h|_\alpha := e^{\sigma_\alpha} h_\alpha$ which can be considered as a local picture of a real valued global Hamiltonian function h . Let us now discuss how we can formulate geometrically the dynamics generated by h , not necessarily relying on coordinates. For this, we address an alternative definition of l.c.s. manifolds.

Lee form. An equivalent definition of a l.c.s. manifold is possible with the aid of a compatible one form by glueing up the exterior derivatives $d\sigma_\alpha$ of the local functions σ_α to a well defined closed one-form θ , called the Lee one-form [17]. This observation leads us to an alternative definition of l.c.s. manifolds in terms of global differential forms. An even dimensional manifold M endowed with a non-degenerate two-form ω is called a l.c.s. manifold if there exists a globally defined one-form θ such that

$$d\omega = \theta \wedge \omega, \quad d\theta = 0. \quad (3)$$

According to this definition, we denote a l.c.s. manifold by a triple (M, ω, θ) where ω is an almost symplectic two-form and θ is a Lee form satisfying (3). This realization of locally conformal symplectic manifolds implies that a l.c.s. manifold is symplectic if and only if the Lee form θ vanishes identically.

Lichnerowicz-de Rham differential. Let θ be a closed one-form on a manifold M . The Lichnerowicz-de Rham differential d_θ is defined on the space of differential forms $\Lambda(M)$

$$d_\theta : \Lambda^k(M) \rightarrow \Lambda^{k+1}(M) : \beta \mapsto d\beta - \theta \wedge \beta, \quad (4)$$

where d denotes the exterior (de Rham) derivative [14, 15]. A triple (M, ω, θ) involving a non-degenerate two-form ω and a closed one-form θ is a l.c.s. manifold if and only if $d_\theta\omega = 0$. Recall that if we employ the local equation of dynamics in (1), and we use the definitions of ω_α and h_α , then, glueing up the local one-forms $d\sigma_\alpha$ to the global Lee one-form θ , we arrive at

$$\iota_{X_h}\omega = d_\theta h. \quad (5)$$

Here, the Hamiltonian vector field X_h is the global realization of the local vector fields X_α in (1).

L.c.s. structures on cotangent bundles. Consider the canonical symplectic manifold T^*Q equipped with the canonical symplectic two-form $\Omega_Q = -d\Theta_Q$ where Θ_Q is the canonical one-form on T^*Q [1, 2, 23]. Let ϑ be a closed form on Q and pull it back to T^*Q by means of the cotangent bundle projection π_Q . This gives us a closed semi-basic one-form $\theta = \pi_Q^*(\vartheta)$. By means of the Lichnerowicz-deRham differential, we define a two-form

$$\Omega_\theta = -d_\theta(\Theta_Q) = -d\Theta_Q + \theta \wedge \Theta_Q = \Omega_Q + \theta \wedge \Theta_Q \quad (6)$$

on the cotangent bundle T^*Q . Notice that the triple

$$T_\theta^*Q := (T^*Q, \Omega_\theta, \theta), \quad \Omega_\theta = -d_\theta\Theta_Q \quad (7)$$

determines a l.c.s. manifold with the Lee-form θ , see [11, 27]. In short, we denote this l.c.s. manifold by T_θ^*Q . Note that all l.c.s. manifolds are locally T_θ^*Q for some Q and for a closed one-form ϑ .

1.2 Geometric Hamilton-Jacobi theory

Consider a Hamiltonian system (T^*Q, Ω_Q, h) generated by a Hamiltonian function h . In the Hamilton-Jacobi theory one aims at finding a function W , the so-called generating function, satisfying the Hamilton-Jacobi equation (HJE)

$$h\left(q^i, \frac{\partial W}{\partial q^i}\right) = \epsilon, \quad (8)$$

where ϵ is a constant [2]. The geometrical picture is the following [10]. Let X_h be the Hamiltonian vector field for the Hamiltonian function h then, consider the following projected vector field

$$X_h^{dW} = T\pi_Q \circ X_h \circ dW, \quad (9)$$

on Q . Here, $T\pi_Q$ is the tangent mapping of the cotangent bundle projection π_Q . The following theorem asserts that an integral curve of X_h^{dW} can be lifted to an integral curve of X_h if and only if W is a solution of (8):

Theorem 1. *For a closed one-form dW on Q the following conditions are equivalent:*

1. *The vector fields X_h and X_h^{dW} are dW -related.*
2. *$d(h \circ dW) = 0$.*

A Hamilton-Jacobi theory in l.c.s. manifolds. Recently, the authors of this same paper have proposed a Hamilton–Jacobi theory on a l.c.s. manifold [12]. Let us summarize here the results we obtained. Consider a Hamiltonian vector field X_h defined through the equation (5). Consider now a section γ of the cotangent bundle and define a vector field on Q as

$$X_h^\gamma = T\pi \circ X_h \circ \gamma. \quad (10)$$

One can define the vector field X_h^γ through the following commutative diagram

$$\begin{array}{ccc} T_\theta^*Q & \xrightarrow{X_h} & TT_\theta^*Q \\ \downarrow \pi_Q & & \downarrow T\pi_Q \\ Q & \xrightarrow{X_h^\gamma} & TQ \end{array}$$

Now, the Hamilton-Jacobi theorem for l.c.s. cotangent bundles reads as follows.

Theorem 2. *Consider a one-form γ whose image is a Lagrangian submanifold of the locally conformal symplectic manifold T_θ^*Q with respect to the almost symplectic two-form Ω_θ , that is $d_\theta\gamma = 0$. Then, the following conditions are equivalent:*

1. *The vector fields X_h and X_h^γ are γ -related,*
2. *$d_\theta(h \circ \gamma) = 0$.*

1.3 The goal and the contents

The aim of this work is two-fold. The first aim, rooting in Vaisman's discussion in [32], is to address the globalization problem of local HDW equations on local k -symplectic spaces. Such discussion will lead us to introduce the notion of what we call henceforth locally conformal k -symplectic (l.c.k-s.) manifolds. Our second goal is to present a geometric Hamilton-Jacobi formalism on l.c.k-s. manifolds. This work can be considered as a continuation or complementary to [12].

This paper contains three main sections. In the first section, we recall the notion of k -symplectic manifolds, HDW dynamics, and the Hamilton-Jacobi theorem on k -symplectic manifolds presented in [18]. We present a coordinate free proof of this Hamilton-Jacobi theorem. The next section 3 concerns the introduction of our novel concept, here presented as locally conformal k -symplectic manifolds. We formulate Hamiltonian dynamics on l.c.k-s. manifolds along with a Hamilton-Jacobi theorem, which constitutes our second main result in this paper. In the last section 4, we present a non-trivial example to illustrate the previous theoretical discussion. All manifolds are considered to be real, paracompact, connected and C^∞ . All maps are C^∞ . Sum over crossed repeated indices is understood.

2 k -symplectic framework

2.1 k -tangent bundle

Let M be an n -dimensional manifold. The tangent bundle (TM, τ_M, M) is a $2n$ -dimensional manifold. Here, τ_M is the tangent bundle projection. A section of the tangent bundle is a vector field X on M . The space of smooth vector fields $\mathfrak{X}(M)$ on M has a module structure on the space of smooth functions on M .

The Whitney sum of k copies of tangent bundles

$$T^k M := TM \oplus_M \cdots \oplus_M TM, \quad (11)$$

is a $nk + n$ dimensional manifold, and the triple $(T^k M, \boldsymbol{\tau}_M, M)$ is a fiber bundle [24]. Here, $\boldsymbol{\tau}_M$ is the projection from $T^k M$ to its base manifold M . Note that we reserve the bold symbol $\boldsymbol{\tau}_M$ in order to differentiate this projection from the tangent bundle projection τ_M . We refer to $T^k M$ as the k -tangent bundle of M . There exists a family of projections τ_κ (here, κ runs from 1 to k) from the k -tangent bundle $T^k M$ to the tangent bundle TM which is assumed to be the κ^{th} -copy TM in (11). A section of k -tangent bundle is a k -vector field \mathbf{X} on M . We denote the space of k -vector fields on M by $\mathfrak{X}^k(M)$. A k -vector field \mathbf{X} is equivalent to a family of k vector fields on M , accordingly, we denote a k -vector field by

$$\mathbf{X} := [X_\kappa] = (X_1, \dots, X_k). \quad (12)$$

Referring to the projections τ_κ we can determine the relation

$$\tau_\kappa \circ \mathbf{X} = X_\kappa, \quad \forall \kappa = 1, \dots, k. \quad (13)$$

According to this, we present the following commutative diagram.

$$\begin{array}{ccc}
T^k M & \xrightarrow{\tau_\kappa} & TM \\
\downarrow \mathbf{X} & \searrow \tau_M & \swarrow \tau_M \\
& & M \\
& & \uparrow X_\kappa
\end{array}
\tag{14}$$

Integrability of k -vector fields. We denote the space of k -vector fields on M by $\mathfrak{X}^k(M)$. As in the classical picture, the integrability of k -vector fields can also be discussed equivalently. For this, let U be an open neighborhood containing the origin $\mathbf{0}$ in a k -dimensional Euclidean space \mathbb{R}^k endowed with coordinates $\mathbf{t} = (t^\kappa)$. An integral section of a k -vector field \mathbf{X} passing through a point x is a differential map ϕ from U to the manifold M satisfying that

$$\phi(\mathbf{0}) = x, \quad \phi_* \left(\frac{\partial}{\partial t^\kappa} \right) = X_\kappa
\tag{15}$$

for all κ from 1 to k . Here, the notation ϕ_* refers to the push-forward operation. A k -vector field is integrable if there is an integral section passing through every point of M .

k -tangent lift. Consider a differentiable mapping φ from a manifold M to another manifold N . Then k -tangent lift of φ is a mapping from the k -tangent bundle $T^k M$ to the k -tangent bundle $T^k N$, and it is defined to be

$$T^k \varphi[X_\kappa(z)] = T^k \varphi(X_1(z), \dots, X_k(z)) = (T\varphi(X_1(z)), \dots, T\varphi(X_k(z)))
\tag{16}$$

where $T\varphi$ is the tangent lift of φ . Here, $X_\kappa(z)$ is an element of $T_z M$, and $[X_\kappa(z)]$ is an element of $T^k M$. Notice that the k -tangent lift of φ satisfies the following commutation rule

$$\tau_N \circ T^k \varphi = \varphi \circ \tau_M
\tag{17}$$

where τ is the k -tangent bundle projection. Let us provide another commutative diagram summarizing this:

$$\begin{array}{ccccc}
T^k M & \xrightarrow{T^k \varphi} & T^k N & & \\
\downarrow \tau_\kappa & & \downarrow \tau_\kappa & & \\
TM & \xrightarrow{T\varphi} & TN & & \\
\downarrow \tau_M & & \downarrow \tau_N & & \\
M & \xrightarrow{\varphi} & N & & \\
\downarrow \mathbf{X} & & \downarrow (\varphi^k)_* \mathbf{X} & & \\
& & M & & N \\
& & \uparrow X_\kappa & & \uparrow T\varphi \circ X_\kappa
\end{array}
\tag{18}$$

Note that, the commutation of this diagram additionally enables us to define the k -th push forward, denoted by $(\varphi^k)_* \mathbf{X}$, of a k -vector field \mathbf{X} by means of the differentiable mapping φ . Accordingly, the relationship between two k -vector fields is defined to be

$$(\varphi^k)_* \mathbf{X} \circ \varphi = T^k \varphi \circ \mathbf{X}.
\tag{19}$$

2.2 k-symplectic manifolds

Consider a n dimensional manifold M with k number of closed two-forms $[\Omega^\kappa]$, and nk -dimensional integrable distribution V . A triple $(M, [\Omega^\kappa], V)$ is called a k -symplectic manifold if the following two conditions are satisfied

$$(i) \Omega^\kappa|_{V \times V} = 0, \quad \forall \kappa = 1, \dots, k, \quad (ii) \bigcap_{\kappa=1}^k \ker \Omega^\kappa = \{0\}. \quad (20)$$

See [3, 4, 19, 20, 21, 22] for fundamentals on k -symplectic manifolds. The canonical model for a k -symplectic manifold is the k -cotangent bundle of a manifold. Let us depict this in detail.

Lagrangian submanifolds. Consider a submanifold N of a k -symplectic manifold $(M, [\Omega^\kappa], V)$. Then k -symplectic dual of the tangent space $T_z N$ is defined as follows

$$(T_z N)^\perp = \left\{ v \in T_z M : \Omega^1(v, w) = \dots = \Omega^k(v, w) = 0, \quad \forall w \in T_z N \right\}. \quad (21)$$

A submanifold N is called isotropic if $T_z N \subset (T_z N)^\perp$ for all z in N [24, 25]. N is called coisotropic if $(T_z N)^\perp \subset T_z N$ whereas it is called as Lagrangian if $(T_z N)^\perp = T_z N$, for all z . It is important to observe that the submanifold of M integrating the distribution V is a Lagrangian submanifold of M . We shall characterize the Lagrangian submanifolds more concretely in the generic example of k -symplectic manifolds defined in the upcoming paragraph.

k -cotangent bundle. The k -cotangent bundle of a manifold Q is defined to be the Whitney sum of k copies of the cotangent bundle T^*Q as

$$T_k^*Q := T^*Q \oplus_Q \dots \oplus_Q T^*Q \quad (22)$$

equipped with the canonical projection π_Q from T_k^*Q to the base Q . Additionally, there exists a family of projections π^κ from T_k^*Q to the κ^{th} -copy T^*Q in the decomposition (22). We define a family of closed two-forms on T_k^*Q by pulling the canonical symplectic two-forms on each T^*Q back to T_k^*Q by means of π^κ . We denote the set of such two-forms by $[\Omega^\kappa]$. Consider the kernel V of the tangent lift $T\pi_Q$ of the projection. It is an integrable distribution. Notice that the set

$$(T_k^*Q, [\Omega^\kappa], V) \quad (23)$$

determines a k -symplectic manifold by satisfying both conditions presented in (20).

In Darboux' coordinates (q^i, p_i^κ) on T_k^*Q , the closed two-forms and the distribution V are computed to be

$$\Omega^\kappa = dq^i \wedge dp_i^\kappa, \quad V = \left\langle \frac{\partial}{\partial p_i^1}, \dots, \frac{\partial}{\partial p_i^k} \right\rangle. \quad (24)$$

It is important to note that this local realization is generic for all k -symplectic manifolds. That is, any k -symplectic manifold, at every point of the manifold, admits a local coordinate system (q^i, p_i^κ) such that the closed two-forms and the distribution have the local expressions in (24), see [3, 22].

Sections of k -cotangent bundle. A section of a k -cotangent bundle is a differentiable mapping γ from the base manifold Q to the total space T_k^*Q satisfying that $\pi_Q \circ \gamma = \text{id}$. A decomposition of γ is

$$\gamma = [\gamma^\kappa] = (\gamma^1, \dots, \gamma^k) \quad (25)$$

where each of γ^κ is a differential one-form on Q . Notice that, the relationships between γ and the one-forms $[\gamma^\kappa]$ are given by

$$\pi^\kappa \circ \gamma = \gamma^\kappa, \quad \forall \kappa = 1, \dots, k. \quad (26)$$

The following commutative diagram summarizes the discussions

$$\begin{array}{ccc} T_k^*Q & \xrightarrow{\pi^\kappa} & T^*Q \\ & \searrow \pi_Q & \swarrow \pi_Q \\ & Q & \end{array} \quad (27)$$

A section γ of π_M is called a closed section if all of the constitutive differential one-forms are closed. We perform the following direct calculation

$$\gamma^* \Omega^\kappa = \gamma^* (\pi^\kappa)^* \Omega_Q = (\pi^\kappa \circ \gamma)^* \Omega_Q = (\gamma^\kappa)^* \Omega_Q = -d\gamma^\kappa \quad (28)$$

where we have employed (26) in the third identity. From (28) we can see that $\gamma^* \Omega^\kappa$ vanishes identically if and only if each γ^κ is closed. This implies that the image space of a section γ of the k -cotangent bundle is a Lagrangian submanifold if and only if it is closed.

2.3 Dynamics on k -symplectic manifolds

Consider the canonical k -symplectic manifold $(T_k^*Q, [\Omega^\kappa], V)$ given in (22). Let us first introduce some notations. Since T_k^*Q is a manifold, one can define its tangent bundle and cotangent bundle as usual TT_k^*Q and $T^*T_k^*Q$. A vector field on T_k^*Q takes values in the tangent bundle TT_k^*Q whereas a one-form on T_k^*Q takes values in the cotangent bundle $T^*T_k^*Q$. Accordingly, we denote the space of vector fields by $\mathfrak{X}(T_k^*Q)$, and the space of one-forms by $\Lambda^1(T_k^*Q)$. Further, we can define the k -tangent bundle $T^k T_k^*Q$ of the k -cotangent bundle T_k^*Q . Sections of this fibration are k -vector fields on T_k^*Q . Recalling (12), we define the following musical mapping

$$b : \mathfrak{X}^k(T_k^*Q) \longrightarrow \Lambda^1(T_k^*Q) : \mathbf{X} = (X_1, \dots, X_k) \mapsto \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa. \quad (29)$$

If $k = 1$, one arrives at the classical symplectic framework where the musical mapping becomes a symplectic isomorphism. But, for $k > 1$, the kernel of the musical mapping is far from being trivial, so it fails to be an isomorphism.

Dynamics on the k -cotangent bundle. Start with a k -cotangent bundle $(T_k^*Q, [\Omega^\kappa], V)$. Consider a smooth Hamiltonian function H defined on T_k^*Q . We define the Hamilton-DeDonder-Weyl (HDW) equation as

$$b(\mathbf{X}) = dH, \quad (30)$$

where \flat is the operator defined in (29). Notice that, in this framework, HDW equation determines a k -vector field \mathbf{X} on T_k^*Q . We call \mathbf{X} a HDW k -vector field. Let us write the HDW equation more explicitly by endowing the decomposition in (12), i.e.,

$$\sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa = dH. \quad (31)$$

It is important to note that it is not possible to determine a unique HDW k -vector field satisfying the HDW equation (30). Let us clarify this in local coordinates.

Dynamics in local coordinates. Recall the existence of Darboux' coordinates (q^i, p_i^κ) on T_k^*Q , and the local realizations of the closed two-forms Ω^κ in (24). In this chart, each vector field X_κ can be written as

$$X_\kappa = (X_\kappa)^i \frac{\partial}{\partial q^i} + (X_\kappa)_i^\lambda \frac{\partial}{\partial p_i^\lambda}, \quad (32)$$

where $(X_\kappa)^i$ and $(X_\kappa)_i^\lambda$ are real valued function on T_k^*Q . Then, we obtain that the HDW equation (31) is equivalent to the system of equations

$$\frac{\partial H}{\partial q^i} = - \sum_{\kappa=1}^k (X_\kappa)_i^\kappa, \quad \frac{\partial H}{\partial p_i^\kappa} = (X_\kappa)^i. \quad (33)$$

A k -vector field \mathbf{X} on T_k^*Q lies in the kernel of the musical mapping \flat in (29) if the components of the constitutive vector fields satisfy

$$(X_\kappa)^i = 0, \quad \sum_{\kappa=1}^k (X_\kappa)_i^\kappa = 0. \quad (34)$$

The existence of solutions of (33) is guaranteed. These depend on $n(k^2 - 1)$ arbitrary functions although they are not necessarily integrable, so the number of integrability conditions imply that the number of arbitrary functions will be less than $n(k^2 - 1)$.

Solutions of HDW equations. A solution to the HDW equations (33) is a mapping ϕ from an open neighbourhood U of the Euclidean space \mathbb{R}^k to the base manifold T_k^*Q . In Darboux' coordinates (q^i, p_i^κ) , we write such a solution as $\phi(\mathbf{t}) = (\phi^i(\mathbf{t}), \phi_i^\kappa(\mathbf{t}))$. In accordance with this, a solution must satisfy

$$\frac{\partial H}{\partial q^i} = - \frac{\partial \phi_i^\kappa}{\partial t^\kappa}, \quad \frac{\partial H}{\partial p_i^\kappa} = \frac{\partial \phi^i}{\partial t^\kappa}, \quad (35)$$

where we have assumed that $\mathbf{t} = (t^\kappa)$ is in \mathbb{R}^k . A solution ϕ of the HDW-equations (35) is said to be an admissible solution if $\text{Im } \phi$ is a closed embedded submanifold of T_k^*Q . An admissible solution of the HDW-equations is an integral section of an integrable k -vector field \mathbf{X} [28]. We say that the set $(T_k^*Q, [\Omega^\kappa], V, H)$ is an admissible k -symplectic Hamiltonian system if all solutions of the HDW-equation are admissible.

2.4 Hamilton–Jacobi theory

Let us briefly recall the Hamilton–Jacobi theory in the k -symplectic setting. One can find more detailed discussions on the issue e.g. in [18, 24]. Consider an admissible k -symplectic Hamiltonian system $(T_k^*Q, [\Omega^\kappa], V, H)$. In this framework, the Hamilton–Jacobi problem is the following partial differential equation

$$H\left(q^i, \frac{\partial W^1}{\partial q^i}, \dots, \frac{\partial W^k}{\partial q^i}\right) = \epsilon, \quad (36)$$

with k number of real valued functions W^κ on Q . Here, ϵ is a real constant. Let us exhibit this geometrically.

Assume that a section \mathbf{X} is a HDW k -vector field on T_k^*Q satisfying (30). Using a closed section γ of the k -cotangent bundle T_k^*Q , we are defining a section \mathbf{X}^γ of the k -tangent bundle T^kQ as follows

$$\mathbf{X}^\gamma := T^k\pi_Q \circ \mathbf{X} \circ \gamma : Q \longrightarrow T^kQ. \quad (37)$$

Here, $T^k\pi_Q$ is the k -th tangent mapping of the projection π_Q , defined according to (16). Let us consider the decompositions of the k -vector fields \mathbf{X} (defined on T_k^*Q) and \mathbf{X}^γ (defined on Q) given by

$$\mathbf{X} = [X_\kappa] = (X_1, \dots, X_k), \quad \mathbf{X}^\gamma = [X_\kappa^\gamma] = (X_1^\gamma, \dots, X_k^\gamma), \quad (38)$$

respectively. Notice that, in these decompositions, each X_κ is a classical vector field on T_k^*Q whereas each X_κ^γ is a classical vector field on Q . We have that

$$\mathbf{X} \circ \gamma = [X_\kappa \circ \gamma] = (X_1 \circ \gamma, \dots, X_k \circ \gamma) \quad (39)$$

By referring to the definition of $T^k\pi_Q$ and in terms of the decompositions, the definition in (37) turns out to be

$$X_\kappa^\gamma = T\pi_Q \circ X_\kappa \circ \gamma : Q \longrightarrow TQ \quad (40)$$

for any κ running from 1 to k . We present a commutative diagram summarizing the discussions.

$$\begin{array}{ccc} T_k^*Q & \xrightarrow{\mathbf{X}=(X_1, \dots, X_k)} & T^kT_k^*Q \\ \downarrow \pi_Q & & \downarrow T^k\pi_Q=(T\pi_Q, \dots, T\pi_Q) \\ Q & \xrightarrow{\mathbf{X}^\gamma=(X_1^\gamma, \dots, X_k^\gamma)} & T^kQ \end{array} \quad (41)$$

$T^k\gamma=(T^k\gamma^1, \dots, T^k\gamma^k)$

where $T^k\gamma$ is the k -tangent lift of γ .

As we previously indicated, each component X_κ of \mathbf{X} is a vector field on T_k^*Q . Consider Darboux' coordinates (q^i, p_i^κ) on T_k^*Q , assume that each X_κ has a local realization as defined in (32). In this frame, each component X_κ^γ of the section \mathbf{X}^γ defined in (40) can locally be computed as

$$X_\kappa^\gamma = (X_\kappa \circ \gamma)^i \frac{\partial}{\partial q^i}. \quad (42)$$

The space $V\pi_Q$ of vertical vectors on the total space of the k -cotangent bundle T_k^*Q are vectors in TT_k^*Q lying in the kernel of the mapping $T\pi_Q$. We have the following decomposition [10]

$$T_{\gamma(q)}T_k^*Q = T_{\gamma(q)}(\text{Im } \gamma) \oplus V_{\gamma(q)}\pi_Q. \quad (43)$$

We generalize this decomposition to the k -tangent bundles as follows. Define the space of k -vertical vectors $V^k\pi_Q$ as a subbundle of the k -tangent bundle $T^kT_k^*Q$

$$V^k\pi_Q = \left\{ \mathbf{V} \in T^kT_k^*Q : T^k\pi_Q(\mathbf{V}) = 0 \right\}. \quad (44)$$

Notice that we can write the space of k -vertical vectors $V^k\pi_Q$ as the Whitney sum of the space of vertical vectors $V\pi_Q$ that is

$$V^k\pi_Q = V\pi_Q \oplus_{T_k^*Q} \cdots \oplus_{T_k^*Q} V\pi_Q. \quad (45)$$

Referring to this identification, we decompose the k -tangent vectors on the image space of γ as follows

$$T_{\gamma(q)}^k T_k^*Q = T_{\gamma(q)}^k(\text{Im } \gamma) \oplus V_{\gamma(q)}^k\pi_Q. \quad (46)$$

Let us first concentrate the following lemma which is useful in the proof of some forthcoming theorems.

Lemma 3. *Consider the canonically k -symplectic k -cotangent bundle $(T_k^*Q, [\Omega^\kappa], V)$. Let $\mathbf{X} = [X_\kappa]$ be a k -vector field on T_k^*Q and $\mathbf{Y} = [Y_\kappa]$ be a k -vector field on Q . Assume also that $\mathbf{X} - T^k\gamma(\mathbf{Y})$ lies in the kernel of the mapping \flat in (29). Then the following equality holds*

$$\gamma^* \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa = \sum_{\kappa=1}^k \iota_{Y_\kappa} (\gamma^* \Omega^\kappa), \quad (47)$$

where γ is a section of π_Q .

Proof. Consider an arbitrary vector $v \in TT_k^*Q$ and compute the expression

$$\gamma^* \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa(v) = \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa(T\gamma(v)) = \sum_{\kappa=1}^k \Omega^\kappa(X_\kappa, T\gamma(v)) \quad (48)$$

and the following one

$$\sum_{\kappa=1}^k \iota_{Y_\kappa} \gamma^* \Omega^\kappa(v) = \sum_{\kappa=1}^k \gamma^* \Omega^\kappa(Y_\kappa, v) = \sum_{\kappa=1}^k \Omega^\kappa(T\gamma(Y_\kappa), T\gamma(v)). \quad (49)$$

The expressions (48) and (49) coincide if and only if

$$\sum_{\kappa=1}^k \Omega^\kappa(T\gamma(Y_\kappa) - X_\kappa, T\gamma(v)) = \flat(T^k\gamma(\mathbf{Y}) - \mathbf{X})(T\gamma(v)) = 0, \quad (50)$$

for all vectors in form $T\gamma(v)$. On the other hand $\mathbf{X} - T^k\gamma(\mathbf{Y})$ is a vertical vector field over the image space of γ with respect to the projection π_Q which reads that $\flat(\mathbf{X} - T^k\gamma(\mathbf{Y}))$ vanishes for all vertical tangent vectors with respect to π_Q . Since the decomposition (46), we deduce that (47) holds if $\flat(\mathbf{X} - T^k\gamma(\mathbf{Y})) = 0$. \square

Let us recall here a geometric version of the Hamilton-Jacobi formalism for k -symplectic framework, see [18, 24]. In the present work, we exhibit the proof in a coordinate free way.

Theorem 4. Let \mathbf{X} be an integrable solution of the HDW equation (31) and γ be a closed section of π_Q , then the following statements are equivalent:

1. If $\sigma : U \subset \mathbb{R}^k \mapsto Q$ is an integral section of \mathbf{X}^γ then $\gamma \circ \sigma$ is a solution \mathbf{X} .
2. $\mathbf{X} \circ \gamma - T^k\gamma(\mathbf{X}^\gamma) \in \ker \flat$.
3. $d(H \circ \gamma) = 0$,

where $T^k\gamma$ is the k -tangent lift of γ , whereas the operator \flat is the one in (29).

Proof. To prove this theorem we first show that (1) if and only if (3) and (2) if and only if (3).

(1) \implies (3) : Let σ be an integral curve of \mathbf{X}^γ . The condition (1) gives that

$$\sigma_*(\mathbf{t}) \left(\frac{\partial}{\partial t^\kappa} \right) = X_a^\gamma(\sigma(\mathbf{t})) \quad \Rightarrow \quad (\gamma \circ \sigma)_*(\mathbf{t}) \left(\frac{\partial}{\partial t^\kappa} \right) = X_\kappa(\gamma \circ \sigma(\mathbf{t})) \quad (51)$$

Applying the push forward γ_* to the former equation we see that X_a^γ and X_κ are γ -related over the solution hypersurfaces. So, we have

$$d(H \circ \gamma) = \gamma^* dH = \gamma^* \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa = \sum_{\kappa=1}^k i_{X_\kappa^\gamma} (\gamma^\kappa)^* \Omega_Q = 0. \quad (52)$$

(3) \implies (1) : Let us assume that $d(H \circ \gamma) = 0$ and $\sigma : U \subset \mathbb{R}^k \mapsto Q$ is an integral section of \mathbf{X}^γ . We have to show that $\gamma \circ \sigma$ is a solution of \mathbf{X} (a solution of Hamilton-De Donder-Weyl equation). Notice that $\gamma \circ \sigma$ is an integral section of the k -symplectic vector field \mathbf{X} defined on \mathbb{T}_k^*Q . From Theorem (4), \mathbf{X} and \mathbf{X}^γ are γ -related up to the kernel of the \flat map. We have to show that

$$\flat(\mathbf{X}) = dH$$

on the image of σ . From (3) we have

$$\begin{aligned} 0 &= d(H \circ \gamma)(\sigma(t)) = \gamma^* dH(\sigma(t)) = \sum_{\kappa=1}^k \iota_{X_\kappa^\gamma(\sigma(t))} (\gamma^\kappa)^* \Omega^\kappa(\sigma(t)) \\ &= \gamma^* \left(\sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa(\sigma(t)) \right) = \gamma^* \flat(\mathbf{X})(\sigma(t)), \end{aligned} \quad (53)$$

where the third equality comes from (28). Hence $\gamma \circ \sigma$ is a solution of \mathbf{X} .

(2) \implies (3) : Assume that (2) holds, that is $\mathbf{X} - T^k\gamma(\mathbf{X}^\gamma)$ belongs to $\ker \flat$. Then we have

$$d(H \circ \gamma) = \gamma^* dH = \gamma^* \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa = \sum_{\kappa=1}^k \iota_{X_\kappa^\gamma} \gamma^* \Omega_Q = 0 \quad (54)$$

where we have used Lemma (3). However, it is (28) which provides the last equality.

(3) \implies (2) : Assume that (3) holds. Define a k -vector field $\mathbf{D} = \mathbf{X} - T^k\gamma(\mathbf{X}^\gamma)$ on T_k^*Q . If the vector field \mathbf{D} belongs to $\ker \flat$, we have finished the proof. For this, first see that \mathbf{D} is a vertical vector field with respect to the projection π_Q , that is

$$\begin{aligned}
T^k\pi_Q \circ \mathbf{D} &= T^k\pi_Q(\mathbf{X} - T^k\gamma(\mathbf{X}^\gamma)) \\
&= T^k\pi_Q \circ \mathbf{X} - T^k\pi_Q \circ T^k\gamma(\mathbf{X}^\gamma) \\
&= T^k\pi_Q \circ \mathbf{X} - T^k\pi_Q \circ T^k\gamma \circ T^k\pi_Q \circ \mathbf{X} \\
&= T^k\pi_Q \circ \mathbf{X} - T^k\pi_Q \circ \mathbf{X} = 0,
\end{aligned} \tag{55}$$

where we have used that $T^k\gamma \circ T^k\pi_Q$ is equal to the k -tangent lift of the identity mapping $\pi_Q \circ \gamma$. Now, see that \mathbf{D} vanishes identically on the image space of γ for the vector fields on T_k^*Q in the form γ_*Y where Y is any vector field on Q .

$$\begin{aligned}
\flat(\mathbf{D})(\gamma_*Y) &= \sum_{\kappa=1}^k i_{D_\alpha} \Omega^\kappa(\gamma_*Y) = \sum_{\kappa=1}^k \Omega^\kappa(X_\kappa - \gamma_*X_\kappa^\gamma, \gamma_*Y) \\
&= \sum_{\kappa=1}^k \Omega^\kappa(X_\kappa, \gamma_*Y) - \sum_{\kappa=1}^k \Omega^\kappa(\gamma_*X_\kappa^\gamma, \gamma_*Y) \\
&= \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa(\gamma_*Y) - \sum_{\kappa=1}^k \gamma^*(\Omega^\kappa)(X_\kappa^\gamma, Y) \\
&= \gamma^* \sum_{\kappa=1}^k (\iota_{X_\kappa} \Omega^\kappa)(Y) - \sum_{\kappa=1}^k \gamma^*(\Omega^\kappa)(X_\kappa^\gamma, Y) \\
&= (\gamma^*dH)(Y) - \sum_{\kappa=1}^k \gamma^*(\Omega^\kappa)(X_\kappa^\gamma, Y) = 0
\end{aligned}$$

where we have employed condition (2) in the first term and the closedness of γ in the second term given in the last line. It follows that \mathbf{D} is in the kernel of \flat that implies condition (1). \square

3 Locally conformal k -symplectic framework

3.1 Locally conformal k -symplectic manifolds

Motivation. In [32], one can review a first discussion on the necessity of locally conformal symplectic manifolds to study Hamiltonian dynamics. In this section, we provide a similar discussion justifying the necessity of what we call locally conformal k -symplectic manifolds while studying HDW dynamics on the k -symplectic framework. We start working on a manifold M with an open covering $\{U_\alpha\}$. Assume also that on each open chart U_α , there exist k number of closed two-forms $[\Omega_\alpha^\kappa] = (\Omega_\alpha^1, \dots, \Omega_\alpha^k)$ satisfying the conditions in (20) for an integrable distribution V_α . In other words, we impose that the family $(U_\alpha, [\Omega_\alpha^\kappa], V_\alpha)$ is a k -symplectic manifold. For a local Hamiltonian function H_α on this local k -symplectic space, the local HDW equations can be written as

$$\sum_{\kappa=1}^k \iota_{(X_\alpha)_\kappa} \Omega_\alpha^\kappa = dH_\alpha. \tag{56}$$

The solutions $(X_\alpha)_\kappa$ of this covariant equation are vector fields on U_α , and the collection of these vector fields $(X_\alpha)_\kappa$ determines a k -vector field on U_α taking values in the k -tangent bundle $T^k U_\alpha$. We denote this local k -vector field by $\mathbf{X}_\alpha := [(X_\alpha)_\kappa]$.

We wish to glue the local k -vector fields \mathbf{X}_α determined in (56) up to a global k -vector field \mathbf{X} on M in order to arrive at global dynamics on M . In this case, as it is well-known, the allowed coordinate transformations are the ones preserving the family of two-forms $[\Omega_\alpha^\kappa]$. Notice that, multiplying the equality (56) by a scalar $\lambda_{\beta\alpha}$, we can preserve the structure on overlapping charts U_α and U_β . That is, the local k -vector fields coincide

$$(X_\kappa)_\alpha = (X_\kappa)_\beta, \quad (57)$$

if the closed two-forms $[\Omega_\alpha^\kappa]$ and the Hamiltonian function H_α obey conformal transformations

$$\Omega_\beta^\kappa = \lambda_{\beta\alpha} \Omega_\alpha^\kappa, \quad \forall \kappa = 1, \dots, k, \quad H_\beta = \lambda_{\beta\alpha} H_\alpha, \quad (58)$$

respectively. This can also be recognized by applying these conformal relations directly to the local equations of motion in (35). It is immediate to realize that the scalars must satisfy the cocycle condition

$$\lambda_{\delta\beta} \lambda_{\beta\alpha} = \lambda_{\delta\alpha}. \quad (59)$$

This leads us to define a real line bundle $L \mapsto M$ over the base manifold M . This way, one can glue up the local two-forms Ω_α^κ to the line bundle valued two-form $\tilde{\Omega}^\kappa$ on M . Further, we argue that the family $[\Omega_\alpha^\kappa]$ of closed one-forms on U_α determines a family of line bundle valued two-forms $[\tilde{\Omega}^\kappa]$ on M .

The cocycle condition (59) manifests the existence of a family of local functions σ_α on each U_α satisfying the relation

$$\lambda_{\beta\alpha} = e^{\sigma_\alpha} / e^{\sigma_\beta} = e^{-(\sigma_\beta - \sigma_\alpha)} \quad (60)$$

in overlapping charts. The substitution of this realization into the first equation in (58) motivates us to define local two-forms

$$\Omega^\kappa|_\alpha := e^{\sigma_\alpha} \Omega_\alpha^\kappa. \quad (61)$$

Since, these local forms give $\Omega^\kappa|_\alpha = \Omega^\kappa|_\beta$ on the intersections $U_\alpha \cap U_\beta$, they can be glued to a real valued two-form Ω^κ on M . So, the local family $[\Omega^\kappa|_\alpha]$ determines a global real valued two-form family $[\Omega^\kappa]$. To sum up, we say that there are two global two-form families, namely $[\Omega^\kappa]$ (real valued) and $[\tilde{\Omega}^\kappa]$ (line bundle valued), on M with local realizations $[\Omega^\kappa|_\alpha]$ and $[\Omega_\alpha^\kappa]$, respectively. These local two-forms are related as given in (61).

In the light of the present discussion, we can argue that if one only imposes the invariance of the HDW equations when we are glueing up the local k -symplectic structures, the result is not necessarily a global k -symplectic structure. Instead, we arrive at what we call locally conformal k -symplectic structures. Here we present the formal definition.

The formal definition. Consider a $n + kn$ dimensional manifold M with k number of two-forms Ω^κ , and nk -dimensional integrable distribution V . M is called a locally conformal k -symplectic (l.c.k-s.) manifold if M has an open cover $\{U_\alpha\}$ and there exists a family of smooth functions σ_α on each chart U_α providing that

$$(i) \quad d(e^{-\sigma_\alpha} \Omega^\kappa|_\alpha) = 0, \quad (ii) \quad \bigcap_{\kappa=1}^k \ker \Omega^\kappa = \{0\}, \quad (iii) \quad \Omega^\kappa|_{V \times V} = 0. \quad (62)$$

Here, $\Omega^\kappa|_\alpha$ denotes the restriction of the two-form Ω^κ to the chart U_α . So, the local two-forms defined by $\Omega_\alpha^\kappa := e^{-\sigma_\alpha} \Omega^\kappa|_\alpha$ are closed two-forms on U_α . Note that the first condition in (62) is local and the other two are global. By glueing up the local condition, we arrive at the following definition of the l.c.k-s. manifolds in terms of a Lee form.

Let M be a manifold of dimension $n + kn$, then $(M, [\Omega^\kappa], \theta, V)$ is called a l.c.k-s. manifold if θ is a closed one-form on M , V is an integrable nk -dimensional distribution on M , and each Ω^κ is a 2-form on M satisfying

$$(i) \quad d\Omega^\kappa = \theta \wedge \Omega^\kappa, \quad (ii) \quad \bigcap_{\kappa=1}^k \ker \Omega^\kappa = \{0\}, \quad (iii) \quad \Omega^\kappa|_{V \times V} = 0. \quad (63)$$

The first condition in (63) is obtained by glueing up the local one-forms, i.e., on each chart, $\theta|_\alpha = d\sigma_\alpha$. Here, we call the family $([\Omega^\kappa], \theta)$ of differential forms a l.c.k-s. structure. Let us discuss the first condition (63) in the realm of the Lichnerowicz-de Rham differential presented in (4). We state that a family $(M, [\Omega^\kappa], \theta, V)$ satisfying the second and the third conditions in (63) is a l.c.k-s. manifold if and only if the two-forms $[\Omega^\kappa]$ are lying in the kernel of the Lichnerowicz-de Rham differential d_θ for the closed one-form θ on M . A l.c.k-s. manifold is said to be globally conformal k -symplectic manifold if the Lee form is an exact one-form.

Exact l.c.k-s. structures. We say that a l.c.k-s. structure $([\Omega^\kappa], \theta)$ is exact if $\Omega^\kappa = d_\theta \Upsilon^\kappa$ for a family of differential one-forms $[\Upsilon^\kappa] = (\Upsilon^1, \dots, \Upsilon^k)$ on M . We call such a set $[\Upsilon^\kappa]$ as Liouville one-form family for the l.c.k-s. structure. We denote an exact l.c.k-s. manifold by $(M, [d_\theta \Upsilon^\kappa], \theta)$ generated by the family $[\Upsilon^\kappa]$. In this case, we define a set of vector fields Z_{Υ^κ} by

$$\iota_{Z_{\Upsilon^\kappa}} \Omega^\kappa = \Upsilon^\kappa, \quad \Omega^\kappa = d_\theta \Upsilon^\kappa. \quad (64)$$

Notice that this family determines a k -vector field $\mathbf{Z} = [Z_{\Upsilon^\kappa}]$ on M , which we call Liouville k -vector field. Applying $\iota_{Z_{\Upsilon^\kappa}}$ to the both hand side of the first equation in (64) we see that $\iota_{Z_{\Upsilon^\kappa}} \Upsilon^\kappa = 0$ where there is no sum on κ .

Lagrangian submanifolds of l.c.k-s. manifolds. We define Lagrangian submanifolds of l.c.k-s. manifolds similar to the Lagrangian submanifolds of k -symplectic manifolds. Let N be a submanifold of a l.c.k-s. manifold $(M, [\Omega^\kappa], V)$. Then l.c.k-s. dual of the tangent space $T_z N$ is defined as the same with (21) whereas in this time the family $[\Omega^\kappa]$ does not consist of closed two-forms but the ones satisfying (62). A submanifold N is called isotropic if $T_z N \subset (T_z N)^\perp$ for all z . N is called coisotropic if $(T_z N)^\perp \subset T_z N$ whereas it is called Lagrangian if $(T_z N)^\perp = T_z N$, for all z . Notice that, the definition of Lagrangian submanifold only related with the second and the third conditions in (62). These are the same with the ones in (20). So that one may expect that Lagrangian submanifolds of l.c.k-s. manifolds and k -symplectic manifolds be in an harmony. For example, as in the case of k -symplectic geometry, the submanifold of M integrating the distribution V is also a Lagrangian submanifold of l.c.k-s. manifold.

3.2 L.c.k-s. structures on the Whitney sum of cotangent bundles

Let Q be a manifold and ϑ be a closed form on it. Pull ϑ back to the cotangent bundle T^*Q by means of the cotangent bundle projection π_Q in order to define a semi-basic and closed one-form

θ . Recall the l.c.s. manifold T_θ^*Q determined in (7). Define the Whitney sum of k number of such bundles

$$T_{k,\theta}^*Q := T_\theta^*Q \oplus_Q T_\theta^*Q \oplus_Q \cdots \oplus_Q T_\theta^*Q. \quad (65)$$

This product is topologically the same with the one in (22), and the subscript θ is placed just to remind the Lee form.

There exist k number of projections π^κ from $T_{k,\theta}^*Q$ to the κ -th copy of T_θ^*Q in the sum. These projections satisfy the commutative diagram in (27). Using them, we introduce k number of differential two-forms

$$\Omega_\theta^\kappa := (\pi^\kappa)^*\Omega_\theta \quad (66)$$

on $T_{k,\theta}^*Q$. Here, Ω_θ is the almost symplectic two-form defined in (6). Another way to define these two-forms is as follows. Consider the closed one-form ϑ defining the Lee-form $\theta = \pi_Q^*\vartheta$ on T_θ^*Q . It can be pulled back to $T_{k,\theta}^*Q$ by means of the k -th order cotangent bundle projection π_Q . In this case, we arrive at a semi-basic one-form section on $T_{k,\theta}^*Q$. Referring to the commutative diagram (27), this one-form coincides with the pull-back of the Lee-form θ on T_θ^*Q by the projection π^κ , that is

$$\pi_Q^*(\vartheta) = (\pi^\kappa)^*(\pi_Q)^*(\vartheta) = (\pi^\kappa)^*\theta. \quad (67)$$

Let us make abuse of notation we denote $\pi_Q^*(\vartheta)$ by θ as well. Notice that, exterior derivatives of the two-forms (Ω_θ^κ) defined in (66) are closed up to the Lee-form θ , that is

$$d\Omega_\theta^\kappa = d(\pi^\kappa)^*\Omega_\theta = (\pi^\kappa)^*d\Omega_\theta = (\pi^\kappa)^*(\theta \wedge \Omega_\theta) = (\pi^\kappa)^*\theta \wedge (\pi^\kappa)^*\Omega_\theta = \theta \wedge \Omega_\theta^\kappa.$$

On the base manifold Q , consider a local coordinate system (q^i) . In this frame, the closed one-form is written as $\vartheta = \vartheta_i dq^i$. Accordingly, on the Darboux' coordinates (q^i, p_i^κ) on $T_{k,\theta}^*Q$, we have the following local realizations of the almost symplectic two-forms

$$\Omega_\theta^\kappa = dp_i^\kappa \wedge dq^i + \vartheta_i(q) p_j^\kappa dq^i \wedge dq^j. \quad (68)$$

One can see that the kernel of Ω_θ^κ is locally generated by the $k-1$ local vector fields $\langle \partial/\partial p_i^\lambda \rangle$ such that $\lambda \neq \kappa$. Therefore, the intersection of the kernels of almost symplectic two-forms is trivial, and the second condition in (63) is satisfied. Considering once more the vertical distribution V exhibited in (24) we have that the third condition in (63) is fulfilled. As a result we arrive at that the family $(T_{k,\theta}^*Q, [\Omega_\theta^\kappa], \theta, V)$ is a l.c.k-s. manifold.

d_ϑ closed Section of l.c.k-s. manifold $T_{k,\theta}^*Q$. Let us determine horizontal Lagrangian submanifolds of l.c.k-s. manifold $T_{k,\theta}^*Q$. For this we consider a section $\gamma = [\gamma^\kappa]$ of the bundle $T_{k,\theta}^*Q$. We perform the following calculation

$$\begin{aligned} (\gamma^\kappa)^*\Omega_\theta^\kappa &= (\gamma^\kappa)^*(-d\Theta_Q^\kappa + \theta \wedge \Theta_Q^\kappa) = -d(\gamma^\kappa)^*(\Theta_Q^\kappa) + (\gamma^\kappa)^*(\theta \wedge \Theta_Q^\kappa) \\ &= -d\gamma^\kappa + (\gamma^\kappa)^*\theta \wedge (\gamma^\kappa)^*\Theta_Q^\kappa = -d\gamma^\kappa + \vartheta \wedge \gamma^\kappa = -d_\vartheta\gamma^\kappa \end{aligned} \quad (69)$$

where we used the identities $(\gamma^\kappa)^*\Theta_Q^\kappa = (\gamma^\kappa)$ and $(\gamma^\kappa)^*\theta = \vartheta$. This gives that the image space of $\gamma = [\gamma^\kappa]$ is an isotropic submanifold of T_θ^*Q if and only if $d_\vartheta\gamma^\kappa = 0$ for all κ from 1 to k . We have discussed that the Lagrangian submanifolds of k -symplectic and l.c.k-s. manifolds are in

harmony. For another example of this, see that each two-form in $[\Omega_\theta^\kappa]$ is the sum of the canonical two-form and the wedge product of two semi-basic one-forms. This is given in (6). So that, each Ω_θ^κ vanishes on vector fields those parallel to the projection π_Q . That is, fibers of the projection are isotropic submanifolds of l.c.k-s. manifold $T_{k,\theta}^*Q$. Another way to see this directly refer to the local realization in (68). Since the tangent space to image of γ at each point is complement those tangent to the fibers, we argue that both the image of γ and the fibers of the projection are Lagrangian submanifolds. See [25] for a similar discussions done in k -symplectic geometry. Since d_θ^2 is identically zero, the image space of the one-form $d_\theta W$ is a Lagrangian submanifold of T_θ^*Q for a function W defined on Q . So that, for a family of functions (W^1, \dots, W^k) the set

$$\text{im}(d_\theta W^1) \oplus_Q \cdots \oplus_Q \text{im}(d_\theta W^k) \subset T^*Q \oplus_Q \cdots \oplus_Q T^*Q \quad (70)$$

is a Lagrangian submanifold of $T_{k,\theta}^*Q$.

3.3 Dynamics in l.c.k-s. manifolds

Recall the discussion done in the beginning of Subsection 3.1 regarding the globalization of the local HDW equations (56). We have exhibited that the two-forms Ω^κ in the global picture take the local realization $\Omega^\kappa|_\alpha = e^{\sigma_\alpha} \Omega_\alpha^\kappa$ on a local chart U_α . On the other hand, the collection of local Hamiltonian functions H_α generating the local HDW equations (56) determine a line bundle valued Hamiltonian (a "twisted Hamiltonian") function \tilde{H} . On the other hand, the local functions $H|_\alpha$, defined by

$$H|_\alpha = e^{\sigma_\alpha} H_\alpha, \quad (71)$$

give a real valued Hamiltonian function H on the manifold M . That is we have two global functions on M , the line bundle valued Hamiltonian function \tilde{H} and real valued Hamiltonian function H . On a chart U_α , these two functions reduce to h_α and $h|_\alpha$, respectively, and they satisfy the relation (71).

We are now ready to glue up the local HDW equations (56). Consider a l.c.k-s. manifold $(M, [\Omega^\kappa], \theta, V)$, and a Hamiltonian function H on M . DWH equation in this framework is

$$\flat(\mathbf{X}) = d_\theta H, \quad (72)$$

where d_θ is the Lichnerowicz-deRham differential defined on M . Here, the mapping \flat is defined as in (29). In other words, we have

$$\sum_{\kappa=1}^k \iota_{X_\kappa} \Omega^\kappa = d_\theta H. \quad (73)$$

3.4 Hamilton-Jacobi theory

We will formulate now a generalization of the Hamilton-Jacobi theory for k -symplectic manifolds to the l.c.k-s. case. For this, consider the locally conformal k -symplectic manifold $T_{k,\theta}^*Q$ presented in (65). Consider a section γ of the fibration π_Q . This section can be written as a combination of k number of differential one-forms $\gamma = (\gamma^1, \dots, \gamma^k)$ as defined in (26). We wish that each γ^κ is closed with respect to the almost symplectic two-form Ω_θ^κ defined in (6). In the light of the calculation done in (69), this is equivalent to the closure of the sections with respect to the Lichnerowicz-deRham

differential d_ϑ where ϑ is the closed one-form generating the Lee form $\theta = \pi_Q(\vartheta)$. We denote a k -vector field on $T_{k,\theta}^*Q$ by \mathbf{Z} . As in (37), using γ , we can construct a k -vector field \mathbf{Z}^γ on Q by

$$\mathbf{Z}^\gamma = T^k \pi_Q \circ \mathbf{Z} \circ \gamma,$$

where the mapping $T^k \pi_Q$ is k -tangent lift of the projection and it is defined as in (18). Notice that, the k -vector fields \mathbf{Z} and \mathbf{Z}^γ fulfill the commutativity diagram (41).

Theorem 5. *Let \mathbf{Z} be an integrable solution of HDW equation (73) and γ is a closed section. Then the followings statements are equivalent:*

1. If $\sigma : U \subset \mathbb{R}^k \mapsto Q$ is an integral section of \mathbf{Z}^γ then $\gamma \circ \sigma$ is a solution of \mathbf{Z} .
2. $\mathbf{Z} \circ \gamma - T^k \gamma(\mathbf{Z}^\gamma) \in \ker \flat$.
3. $d_\vartheta(H \circ \gamma) = 0$.

Proof. While proving this theorem we first show that (1) and (3) are in if and only if relationship then we show that (2) and (3) are in if and only if relationship.

(1) \implies (3) : Let σ be an integral curve of \mathbf{Z}^γ . The condition (1) gives that

$$\sigma_*(\mathbf{t}) \left(\frac{\partial}{\partial t^\kappa} \right) = X_\kappa^\gamma(\sigma(\mathbf{t})) \quad \Rightarrow \quad (\gamma \circ \sigma)_*(\mathbf{t}) \left(\frac{\partial}{\partial t^\kappa} \right) = X_\kappa(\gamma \circ \sigma(\mathbf{t})) \quad (74)$$

Applying push forward γ_* to the former equation we see that Z_a^γ and Z_κ are γ -related over the solution hypersurfaces. So that we have

$$d_\vartheta(H \circ \gamma) = \gamma^* d_\theta H = \gamma^* \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega_\theta^\kappa = \sum_{\kappa=1}^k i_{X_\kappa^\gamma} (\gamma^\kappa)^* \Omega_Q = 0. \quad (75)$$

(3) \implies (1) : Let us assume that $d_\vartheta(H \circ \gamma) = 0$ and $\sigma : U \subset \mathbb{R}^k \mapsto Q$ is an integral section of \mathbf{Z}^γ . We have to show that $\gamma \circ \sigma$ is a solution of \mathbf{Z} (a solution of Hamilton-De Donder-Weyl equation). Let us notice that $\gamma \circ \sigma$ is an integral section of the k -symplectic vector field \mathbf{Z} defined on $T_{k,\theta}^*Q$. Referring to Theorem (5), \mathbf{Z} and \mathbf{Z}^γ are γ -related up to kernel of \flat . We have to show that

$$\flat(\mathbf{Z}) = d_\vartheta H$$

on the image of σ . From (3) we have

$$\begin{aligned} 0 &= d_\vartheta(H \circ \gamma)(\sigma(t)) = \gamma^* d_\theta H(\sigma(t)) = \sum_{\kappa=1}^k \iota_{X_\kappa^\gamma(\sigma(t))} (\gamma^*)^* \Omega_\theta^\kappa(\sigma(t)) \\ &= \gamma^* \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega_\theta^\kappa(\sigma(t)) = \gamma^* \flat(\mathbf{Z})(\sigma(t)), \end{aligned} \quad (76)$$

where the third equality comes from (28). Hence $\gamma \circ \sigma$ is a solution of \mathbf{Z} .

(2) \implies (3) : Notice that the second condition (3) can be rewritten as $\gamma^*(d_\theta H) = 0$ according to the following calculation

$$\begin{aligned} d_\vartheta(H \circ \gamma) &= d(H \circ \gamma) - (H \circ \gamma)\vartheta = d(\gamma^* H) - \gamma^* H \vartheta \\ &= \gamma^* dH - \gamma^* H \gamma^* \theta = \gamma^*(dH - H\theta) = \gamma^*(d_\theta H), \end{aligned} \quad (77)$$

where we used the identity

$$\gamma^*\theta = \gamma^*\pi_Q^*\vartheta = (\pi_Q \circ \gamma)^*\vartheta = \vartheta \quad (78)$$

in the second line of the calculation (77). Assume now that the first condition holds, then Lemma (3) leads us to compute

$$\begin{aligned} \gamma^*(d_\theta H) &= \gamma^* \sum_{\kappa=1}^k \iota_{Z_\kappa} \Omega_\theta^\kappa = \sum_{\kappa=1}^k \iota_{Z_\kappa} \gamma^* \Omega_\theta^\kappa \\ &= \sum_{\kappa=1}^k \iota_{Z_\kappa} \gamma^* (\pi^\kappa)^* \Omega_\theta = \sum_{\kappa=1}^k \iota_{Z_\kappa} (\pi^\kappa \circ \gamma)^* \Omega_\theta \\ &= \sum_{\kappa=1}^k \iota_{Z_\kappa} (\gamma^\kappa)^* \Omega_\theta = 0, \end{aligned} \quad (79)$$

where the last equality comes from the calculation in (69) stating that $\gamma^* \Omega_\theta^\kappa = 0$ equivalent to $d_\theta \gamma = 0$.

(3) \implies (2) : To prove that, same as in k -symplectic case, we define a k -vector field $\mathbf{D} = \mathbf{Z} - T^k \gamma(\mathbf{Z}^\gamma)$ on $T_{k,\theta}^* Q$. Our goal is to show that \mathbf{D} belongs to $\ker \flat$ when it is restricted on the image space of γ . For this end, see that \mathbf{D} is a vertical k -vector field over the image space of γ with respect to the projection π_Q that is

$$T^k \pi_Q \circ \mathbf{D} \circ \gamma = 0.$$

This is due to the calculation (55). Now, consider a vector field Y on Q . By pushing Y forward by γ , we arrive at a vector field $\gamma_* Y$ on $T_{k,\theta}^* Q$. We compute the following

$$\begin{aligned} \flat(\mathbf{D})(\gamma_* Y) &= \sum_{\kappa=1}^k \iota_{D_\alpha} \Omega_\theta^\kappa(\gamma_* Y) = \sum_{\kappa=1}^k \Omega_\theta^\kappa(X_\kappa - \gamma_* X_\kappa^\gamma, \gamma_* Y) \\ &= \sum_{\kappa=1}^k \Omega_\theta^\kappa(X_\kappa, \gamma_* Y) - \sum_{\kappa=1}^k \Omega_\theta^\kappa(\gamma_* X_\kappa^\gamma, \gamma_* Y) \\ &= \sum_{\kappa=1}^k \iota_{X_\kappa} \Omega_\theta^\kappa(\gamma_* Y) - \sum_{\kappa=1}^k \gamma^*(\Omega_\theta^\kappa)(X_\kappa^\gamma, Y) \\ &= \gamma^* \left(\sum_{\kappa=1}^k \iota_{X_\kappa} \Omega_\theta^\kappa \right) (Y) - \sum_{\kappa=1}^k \gamma^*(\Omega_\theta^\kappa)(X_\kappa^\gamma, Y) \\ &= (\gamma^* d_\theta H)(Y) - \sum_{\kappa=1}^k \gamma^*(\Omega_\theta^\kappa)(X_\kappa^\gamma, Y) = 0 \end{aligned} \quad (80)$$

where we have employed the second condition for the first term at the last line, whereas we employed the closure of γ for the second term at the last line. Observing the decomposition of the k -vector spaces given by (46), we arrive at that \mathbf{D} is in the kernel of \flat this gives the condition (2). \square

4 An example

Let $Q = \mathbf{R}^2 - \{\mathbf{0}\}$ be the two dimensional punctured Euclidean space. Here, $\mathbf{0}$ denotes the origin. On this manifold Q , consider the following one-form

$$\vartheta = 2 \frac{xdy - ydx}{x^2 + y^2} \quad (81)$$

which is closed but not exact. Consider the k -cotangent bundle T_k^*Q over the base manifold Q equipped with the canonical projection π_Q and the local coordinates $(x, y, p_x^\kappa, p_y^\kappa)$ where κ runs from 1 to k . We pull the one-form ϑ in (81) back to T_k^*Q with the projection π_Q . This results in a one-form $\theta = (\pi_Q)^*\vartheta$ on the k -cotangent bundle T_k^*Q . The semi-basic character of θ manifests that its local realization is the same as the one-form in (81). We define a k number of one-forms and a k number of closed two-forms on T_k^*Q by pulling back the canonical one forms and the symplectic two-forms on each component of T_k^*Q , so that we have

$$\theta^\kappa = (\pi^\kappa)^*(\theta_Q) = p_x^\kappa dx + dp_y^\kappa dy, \quad \Omega^\kappa = (\pi^\kappa)^*(\Omega_Q) = dx \wedge dp_x^\kappa + dy \wedge dp_y^\kappa. \quad (82)$$

It is immediate to observe that minus the exterior derivative of θ^κ is precisely Ω^κ for all κ running from 1 to k . By means of the differential forms exhibited in (82), we are defining the following family of two-forms

$$\Omega_\theta^\kappa = \Omega^\kappa + \theta \wedge \theta^\kappa = dx \wedge dp_x^\kappa + dy \wedge dp_y^\kappa - 2 \frac{yp_y^\kappa + xp_x^\kappa}{x^2 + y^2} dx \wedge dy, \quad (83)$$

where θ is the one-form defined as $(\pi_Q)^*\vartheta$. After a direct calculation it is easy to observe that the set $[\Omega_\theta^\kappa]$ satisfies the first and the second condition in (63). To see that this family determines a l.c.k-s. structure we define the integrable distribution

$$V = \left\langle \frac{\partial}{\partial p_x^\kappa}, \frac{\partial}{\partial p_y^\kappa} \right\rangle. \quad (84)$$

Referring to this, observation easily validates the third condition in (63).

Now we introduce the following quadratic Hamiltonian function

$$H = \sum_{\kappa=1}^k \frac{1}{2} ((p_x^\kappa)^2 + (p_y^\kappa)^2) \quad (85)$$

on T_k^*Q and recall the HDW equation (73) in the k.l.c.s. framework defined in terms of the Lichnerowicz-deRham differential d_θ . We compute the Lichnerowicz-deRham differential of the Hamiltonian function as follows

$$d_\theta H = dH - H\theta = \sum_{\kappa=1}^k p_x^\kappa dp_x^\kappa + p_y^\kappa dp_y^\kappa - \sum_{\kappa=1}^k \frac{1}{2} ((p_x^\kappa)^2 + (p_y^\kappa)^2) \frac{xdy - ydx}{x^2 + y^2}. \quad (86)$$

For a HDW k -vector field $\mathbf{X} = (X_1, \dots, X_k)$, the HDW equation (73) determines the constitutive vector fields

$$X_\kappa = A_\kappa \frac{\partial}{\partial x} + B_\kappa \frac{\partial}{\partial y} + (C_\kappa)^\lambda \frac{\partial}{\partial p_x^\lambda} + (D_\kappa)^\lambda \frac{\partial}{\partial p_y^\lambda}, \quad (87)$$

with the following explicit computations

$$\begin{aligned} A_\kappa &= p_x^\kappa, & B_\kappa &= p_y^\kappa, & \sum_{\kappa=1}^k (C_\kappa)^\kappa &= \sum_{\kappa=1}^k \frac{y(p_y^\kappa)^2 - y(p_x^\kappa)^2 + 2xp_x^\kappa p_y^\kappa}{x^2 + y^2}, \\ \sum_{\kappa=1}^k (D_\kappa)^\kappa &= \sum_{\kappa=1}^k \frac{x(p_y^\kappa)^2 - x(p_x^\kappa)^2 - 2yp_x^\kappa p_y^\kappa}{x^2 + y^2}. \end{aligned} \quad (88)$$

Let us now apply the global HJ theorem (5) to the present example. Start with a family of one-forms

$$\gamma^\kappa = \beta^\kappa(x, y)dx + \rho^\kappa(x, y)dy \quad (89)$$

satisfying $d_\vartheta \gamma^\kappa = 0$ for each $\kappa = 1, \dots, k$ in order to guarantee that the image space of γ is a Lagrangian submanifold of the l.c.k.s. manifold $T_{k, \theta}^*Q$ equipped with the two-forms $[\Omega_\theta^\kappa]$ in (83). That is, we have

$$\frac{\partial \rho^\kappa}{\partial x} - \frac{\partial \beta^\kappa}{\partial y} + \frac{2x\beta^\kappa - 2y\rho^\kappa}{x^2 + y^2} = 0, \quad \forall \kappa = 1, \dots, k. \quad (90)$$

Referring to the third condition in the HJ theorem (5), we write the HJ equation as

$$d_\vartheta(H \circ \gamma) = \frac{1}{2}d\left(\sum_{\kappa=1}^k (\beta^\kappa)^2 + (\rho^\kappa)^2\right) - \left(\sum_{\kappa=1}^k (\beta^\kappa)^2 + (\rho^\kappa)^2\right) \left(\frac{xdy - ydx}{x^2 + y^2}\right) = 0. \quad (91)$$

Explicitly, we compute the following system of equations

$$\begin{aligned} \sum_{\kappa=1}^k \left(\beta^\kappa \frac{\partial \beta^\kappa}{\partial x} + \rho^\kappa \frac{\partial \rho^\kappa}{\partial x} + \left((\beta^\kappa)^2 + (\rho^\kappa)^2 \right) \frac{y}{x^2 + y^2} \right) &= 0, \\ \sum_{\kappa=1}^k \left(\beta^\kappa \frac{\partial \beta^\kappa}{\partial y} + \rho^\kappa \frac{\partial \rho^\kappa}{\partial y} - \left((\beta^\kappa)^2 + (\rho^\kappa)^2 \right) \frac{x}{x^2 + y^2} \right) &= 0. \end{aligned} \quad (92)$$

We write this system in a local coordinate chart. For it, we choose the polar coordinates $x = r \cos \phi$, $y = r \sin \phi$ on an open chart in Q , and the induced coordinates $(r, \phi, p_r^\kappa, p_\phi^\kappa)$ on the open chart U_α . In these coordinates, the Lee form θ turns out to be an exact one-form $d\sigma_\alpha$ for $\sigma_\alpha = 2\phi$ whereas the two-form in (83) reduces to

$$\Omega_\theta^\kappa|_\alpha = dr \wedge dp_r^\kappa + d\phi \wedge dp_\phi^\kappa - 2p_r^\kappa dr \wedge d\phi. \quad (93)$$

Notice that this two-form is not closed but the following one, which is defined according to the formula in (61),

$$\Omega_\alpha^\kappa = e^{-\sigma_\alpha} \Omega_\theta^\kappa|_\alpha = e^{-2\phi} \Omega_\theta^\kappa|_\alpha = e^{-2\phi} \left(dr \wedge dp_r^\kappa + d\phi \wedge dp_\phi^\kappa - 2p_r^\kappa dr \wedge d\phi \right) \quad (94)$$

is closed. One can check that the set $(U_\alpha, [\Omega_\alpha^\kappa], V_\alpha)$ determines a k -symplectic manifold. Here, V_α is the restriction of the integrable distribution V presented in (84). In the local chart, V_α is spanned by tangent vectors

$$V = \left\langle \left(\cos \phi \frac{\partial}{\partial p_r^\kappa} - r \sin \phi \frac{\partial}{\partial p_\phi^\kappa} \right), \left(\sin \phi \frac{\partial}{\partial p_r^\kappa} + r \cos \phi \frac{\partial}{\partial p_\phi^\kappa} \right) \right\rangle. \quad (95)$$

The Hamiltonian function (85) can be written as

$$H|_\alpha = \frac{1}{2} \sum_{\kappa=1}^k \left((p_r^\kappa)^2 + \frac{1}{r^2} (p_\phi^\kappa)^2 \right) \quad (96)$$

and then, according to (71), define the local function

$$H_\alpha = e^{-\sigma_\alpha} H|_\alpha = \frac{1}{2} e^{-2\phi} \sum_{\kappa=1}^k \left((p_r^\kappa)^2 + \frac{1}{r^2} (p_\phi^\kappa)^2 \right) \quad (97)$$

on the local chart U_α . We find the Hamiltonian dynamics generated by the local HDW equation

$$\sum_{\kappa=1}^k \iota_{(X_\kappa)_\alpha} \Omega_\alpha^\kappa = dH_\alpha. \quad (98)$$

For a vector field

$$X_\kappa = A_\kappa \partial_r + B_\kappa \partial_\phi + (C_\kappa)^\lambda \partial_{p_r^\lambda} + (D_\kappa)^\lambda \partial_{p_\phi^\lambda}$$

we obtain the coefficients

$$A_\kappa = p_r^\kappa, \quad B_\kappa = \frac{1}{r^2} p_\phi^\kappa, \quad \sum_{\kappa=1}^k (C_\kappa)^\kappa = \sum_{\kappa=1}^k \frac{1}{r^2} p_\phi^\kappa (2p_r^\kappa + \frac{1}{r} p_\phi^\kappa), \quad \sum_{\kappa=1}^k (D_\kappa)^\kappa = \sum_{\kappa=1}^k - (p_r^\kappa)^2 + \frac{1}{r^2} (p_\phi^\kappa)^2. \quad (99)$$

Now, we are in the realm of the Hamilton-Jacobi theorem (4) valid for k -symplectic framework. So, we consider a closed section $\gamma_\alpha = (\gamma_\alpha^1, \dots, \gamma_\alpha^k)$ which in local coordinates becomes

$$\gamma_\alpha^\kappa = \xi^\kappa(r, \phi) dr + \eta^\kappa(r, \phi) d\phi.$$

The closure of γ implies that for each κ

$$0 = (\gamma_\alpha^\kappa)^* \Omega_\alpha^\kappa = e^{-2\phi} \left(\frac{\partial \xi^\kappa}{\partial \phi} - \frac{\partial \eta^\kappa}{\partial r} - 2\xi^\kappa \right) dr \wedge d\phi. \quad (100)$$

Now, we calculate $d(H_\alpha \circ \gamma_\alpha)$ and expect that it vanishes identically. Then, the HJ equations would be

$$\begin{aligned} \sum_{\kappa=1}^k \left(\xi^\kappa \frac{\partial \xi^\kappa}{\partial \phi} + \eta^\kappa \frac{1}{r^2} \frac{\partial \eta^\kappa}{\partial \phi} - \xi^2 - \frac{1}{r^2} (\eta^\kappa)^2 \right) &= 0 \\ \sum_{\kappa=1}^k \left(\xi^\kappa \frac{\partial \xi^\kappa}{\partial r} + \eta^\kappa \frac{1}{r^2} \frac{\partial \eta^\kappa}{\partial r} - \frac{1}{r^3} (\eta^\kappa)^2 \right) &= 0. \end{aligned} \quad (101)$$

One can sum these two equations, and simplify the expression to

$$\frac{1}{2} d \left((\xi^\kappa)^2 + \frac{1}{r^2} (\eta^\kappa)^2 \right) = (\xi^\kappa)^2 + \frac{1}{r^2} (\eta^\kappa)^2 \quad (102)$$

where “ d ” denotes total derivative with respect to (r, ϕ) .

5 Conclusion and future work

In this work, we have addressed the glueing problem of HDW equations defined on local k -symplectic structures. Accordingly, we have introduced the notion of locally conformal k -symplectic (l.c.k-s.) manifolds in Subsection 3.1.

In Subsection 3.2, we exhibited the Whitney sum of a k number of cotangent bundles equipped with locally conformal symplectic structures and a HDW equation for this framework has been studied in Subsection 3.3. Further, we have depicted the HJ formalism for the HDW theory. To this end, we have first presented a coordinate free proof of the HJ theorem on k -symplectic spaces in Subsection 2.4, then we have taken this discussion to l.c.k-s. manifolds in Subsection 3.4 and we have provided an example. We plan on pursuing investigations on locally conformal k -symplectic (l.c.k-s.) manifolds in the following headlines:

- The reduction of locally conformal symplectic (l.c.s.) manifolds under Lie groups has been studied by many authors see, for example, [16, 31, 30], whereas reduction on k -symplectic manifolds has been given, for example, in [7, 26]. In similar fashion, it could be interesting to study the reduction of the HDW equation (73) in the l.c.k-s. framework under Lie group symmetry.
- The Hamiltonian realization of field theories is also available in the multisymplectic framework [8, 9, 13, 29]. We find interesting to discuss the glueing problem of local Hamiltonian dynamics in this framework.

6 Acknowledgements

This work has been partially supported by MINECO grants MTM2016-76-072-P and the ICMAT Severo Ochoa Project SEV-2011-0087 and SEV-2015-0554.

References

- [1] Abraham R and Marsden J E 1978 *Foundations of mechanics*. Second edition. Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, Mass.
- [2] Arnold V I 1989 *Mathematical methods of classical mechanics* **60** of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition.
- [3] Awane A 1992 k -symplectic structures, *J. Math. Phys.* **33** (12), 4046–4052.
- [4] Awane A 1994 G -spaces k -symplectic homogènes, *J. Geom. Phys.* **13** 139–157.
- [5] Balseiro P 2017 The Jacobiator of nonholonomic systems and the geometry of magentauced nonholonomic brackets, *Arch. Rat. Mech. Anal.* **214** 453–501.
- [6] Bazzoni G 2017 *Locally conformal symplectic and Kahler geometry* arXiv:1711.02440 (2017).
- [7] Blaga A M 2009 Remarks on Poisson reduction on k -symplectic manifolds, *J. Geom. Symm. Phys.* **13**, 1–7.

- [8] Cantrijn F, Ibort L A and de León M 1996 Hamiltonian structures on multisymplectic manifolds, *Rend. Sem. Mat. Univ. Politec. Torino* **54**(3), 225–236.
- [9] Cantrijn F, Ibort L A and de León M 1999 On the geometry of multisymplectic manifolds, *Journal of the Australian Mathematical Society* **66**(3), 303–330.
- [10] Carinena J F, Gracia X, Marmo G, Martínez E, Muñoz-Lecanda M C and Roman-Roy N 2006 Geometric Hamilton–Jacobi theory, *Int. J. Geom. Methods Mod. Phys.* **3**(07), 1417–1458.
- [11] Chantraine B and Murphy E 2016 *Conformal symplectic geometry of cotangent bundles*, arXiv:1606.00861.
- [12] Esen O, de León M, Sardon C and Zajac M 2019 *Hamiltonian mechanics on locally conformal symplectic manifolds*, arXiv:1910.02016.
- [13] Gotay M J 1991 A multisymplectic framework for classical field theory and the calculus of variations: I. Covariant Hamiltonian formalism, *In Mechanics, analysis and geometry: 200 years after Lagrange* (203-235). Elsevier.
- [14] Guedira F and Lichnerowicz A 1984 Géométrie des algèbres de Lie locales de Kirillov *J. Math. Pures Appl.* **63** 4, 407–484.
- [15] Haller S and Rybicki T 1999 On the Group of Diffeomorphisms Preserving a Locally Conformal Symplectic Structure, *Ann. Global Anal. Geom.* **17**, 475–502.
- [16] Haller S and Rybicki T 2001 Reduction for locally conformal symplectic manifolds, *J. Geom. Phys.* **37**(3), 262–271.
- [17] Lee H W 1943 A kind of even dimensional differential geometry and its applications to exterior calculus, *Amer. J. Math.* **65**, 433–438.
- [18] de León M, de Diego D M, Marrero J C, Salgado M and Vilariño S 2010 Hamilton-Jacobi Theory in k-symplectic Field Theories, *Int. J. Geom. Methods Mod. Phys.* **7**(8), 1491–1507.
- [19] de León M, Méndez I and Salgado M 1988 Regular p-almost cotangent structures, *J. Korean Math. Soc.* **25**(2), 273–287.
- [20] de León M, Méndez I and Salgado M 1988 p-almost tangent structures, *Rend. Circ. Mat. Palermo* **37**(2), 282-294.
- [21] de León M, Méndez I and Salgado M 1991 Integrable p-almost tangent manifolds and tangent bundles of p1-velocities, *Acta Math. Hungar.* **58**(1-2), 45-54.
- [22] de León M, Méndez I and Salgado M 1993 p-almost cotangent structures, *Boll. Un. Mat. Ital. A* **7**(1), 97-107
- [23] de León M and Rodrigues P R 1989 *Methods of differential geometry in analytical mechanics*. North-Holland Mathematics Studies **158**. North-Holland Publishing Co., Amsterdam.
- [24] de León M, Salgado M and Vilariño S 2015 *Methods of Differential Geometry in Classical Field Theories k-Symplectic and k-Cosymplectic Approaches* World Scientific, Singapore.

- [25] de León M and Vilariño S 2013 Lagrangian submanifolds in k -symplectic setting, *Monatsh Math* **170**, 381–404.
- [26] Munteanu F, Rey A M and Salgado M 2004 The Günthers formalism in classical field theory: momentum map and reduction. *J. Math. Phys.* **45**(5), 1730–1751.
- [27] Otíman A and Stanciu M 2017 Darboux-Weinstein theorem for locally conformal symplectic manifolds. *J. Geom. Phys.* **111**, 1–5.
- [28] Román-Roy N, Rey A M, Salgado M, Vilariño S 2007 On the k -symplectic, k -cosymplectic, and multisymplectic formalism of classical field theories, *J. Geom. Mech.* **3**, 113–137.
- [29] Román-Roy N 2009 Multisymplectic Lagrangian and Hamiltonian formalisms of classical field theories, *SIGMA: Symmetry, Integrability and Geometry: Methods and Applications* **5**(0), 100 (25 pages).
- [30] Stanciu M 2019 Locally conformal symplectic reduction, *Ann.Global Anal. Geom.* **56** (2), 245–275.
- [31] Stanciu M 2019 *Locally conformal symplectic reduction of the cotangent bundle*, arXiv:1905.02798.
- [32] Vaisman I 1985 Locally conformal symplectic manifolds *Internat. J. Math. Math. Sci.* **8**, 521–536.