

REDUCIBILITY OF PARAMETER IDEALS IN LOW POWERS OF THE MAXIMAL IDEAL

KATHARINE SHULTIS AND PEDER THOMPSON

To Roger and Sylvia Wiegand in celebration of their combined 150th birthday.

ABSTRACT. A commutative noetherian local ring (R, \mathfrak{m}) is Gorenstein if and only if every parameter ideal of R is irreducible. Although irreducible parameter ideals may exist in non-Gorenstein rings, Marley, Rogers, and Sakurai show there exists an integer ℓ (depending on R) such that R is Gorenstein if and only if there exists an irreducible parameter ideal contained in \mathfrak{m}^ℓ . We give upper bounds for ℓ that depend primarily on the existence of certain systems of parameters in low powers of the maximal ideal.

1. INTRODUCTION

Let (R, \mathfrak{m}, k) be a commutative noetherian local ring of dimension $\dim R = d$. It is known that R is Gorenstein if and only if every parameter ideal of R is irreducible, but we cannot characterize Gorenstein rings by the existence of an irreducible parameter ideal. For example, the non-Gorenstein ring $\mathbb{Q}[[x, y]]/(x^2, xy)$ has an irreducible parameter ideal (y) , although (y^j) is reducible for $j \geq 2$. Marley, Rogers, and Sakurai show [11], however, that the existence of a parameter ideal in a sufficiently high power of the maximal ideal *does* characterize Gorenstein rings:

Theorem 1.1 (Marley, Rogers, and Sakurai). *There exists an integer ℓ , depending on R , such that R is Gorenstein if and only if some parameter ideal contained in \mathfrak{m}^ℓ is irreducible.*

The integer ℓ in Theorem 1.1, considered previously by Goto and Sakurai [9, Lemma 3.12] may be taken to be the least integer i such that the canonical map

$$(1.2) \quad \text{Ext}_R^d(R/\mathfrak{m}^i, R) \longrightarrow \varinjlim_j \text{Ext}_R^d(R/\mathfrak{m}^j, R) \cong H_{\mathfrak{m}}^d(R)$$

becomes surjective after applying the socle functor $\text{Hom}_R(R/\mathfrak{m}, -)$. The existence of such an integer is guaranteed as the socle module $\text{Hom}_R(R/\mathfrak{m}, H_{\mathfrak{m}}^d(R))$ is finitely generated, but determining how large ℓ must be seems to be somewhat subtle. Indeed, we show in Example 2.8 that for each integer $a \geq 1$, there exists a ring which requires $\ell > a$. To understand how deep in the maximal ideal one must go before detecting whether R is Gorenstein in terms of reducibility of parameter ideals, we consider the problem, posed to the authors by Marley, of finding an upper bound for the integer ℓ in Theorem 1.1.

Date: June 8, 2020.

2020 Mathematics Subject Classification. 13C05, 13D45, 13H10.

Key words and phrases. System of parameters, reducible parameter ideal, Gorenstein ring.

This material is based upon work supported by the National Science Foundation under Grant #1321794, as part of the Mathematical Research Communities 2015 program in Snowbird, Utah.

For rings of dimension one, we take a direct approach to determine surjectivity of the maps in (1.2) after applying $\mathrm{Hom}_R(R/\mathfrak{m}, -)$. We show (see Theorem 2.4):

Theorem 1.3. *Assume $\dim R = 1$ and k is infinite. If n is the least integer such that $\mathfrak{m}^n = (x)\mathfrak{m}^{n-1}$ for some parameter x and $\mathfrak{m}^{n-1} \cap \Gamma_{\mathfrak{m}}(R) = 0$, then for $i \geq n$ the canonical map in (1.2) becomes surjective after applying $\mathrm{Hom}_R(R/\mathfrak{m}, -)$.*

Thus, in this setting, R is Gorenstein if and only if some parameter ideal contained in \mathfrak{m}^n is irreducible; see Corollary 2.7. This consequence of Theorem 1.3 can also be deduced from work of Rogers [13] and Marley, Rogers, and Sakurai [11]. The assumption that k is infinite is only needed to ensure the existence of a parameter x such that $\mathfrak{m}^n = (x)\mathfrak{m}^{n-1}$.

For a ring R of dimension d (not necessarily equal to 1) and a system of parameters x_1, \dots, x_d , we instead consider—in place of (1.2)—the least integer i such that the canonical map

$$(1.4) \quad R/(x_1^i, \dots, x_d^i) \longrightarrow \varinjlim_j R/(x_1^j, \dots, x_d^j) \cong H_{\mathfrak{m}}^d(R)$$

becomes surjective after applying $\mathrm{Hom}_R(R/\mathfrak{m}, -)$. We focus on the case where x_1, \dots, x_d is a p_s -standard system of parameters (see Definition 3.5); this is a variant of the p -standard systems of parameters considered by Cuong [3]. These systems of parameters (both p_s -standard and p -standard) are chosen in a way as to annihilate certain local cohomology modules. We show in Proposition 3.7 that if x_1, \dots, x_d is a p_s -standard system of parameters for some $s \geq 2$, then for $i \geq s$ the canonical map in (1.4) becomes surjective after applying $\mathrm{Hom}_R(R/\mathfrak{m}, -)$.

As a consequence of this surjectivity, we obtain a characterization of the Gorenstein property of R in terms of irreducibility of parameter ideals: the integer ℓ from Theorem 1.1 can be taken to be the integer n in the next result. In particular, we prove (as a special case of Theorem 3.9):

Theorem 1.5. *Assume R has a dualizing complex. If n is an integer such that $\mathfrak{m}^n \subseteq (x_1^2, \dots, x_d^2)$ for a p_2 -standard system of parameters x_1, \dots, x_d , then R is Gorenstein if and only if some parameter ideal contained in \mathfrak{m}^n is irreducible.*

The assumption that R has a dualizing complex is sufficient for the existence of a p_2 -standard system of parameters; see the discussion before Definition 3.5. Moreover, work of Cuong and Cuong [4, 5] implies that Theorem 1.5 holds if one replaces the assumption that R has a dualizing complex with the assumption that R is a quotient of a Cohen-Macaulay local ring; see Remark 3.6.

* * *

Throughout this paper, let (R, \mathfrak{m}, k) be a commutative noetherian local ring. Let $\dim R = d$ be the Krull dimension of R . We briefly recall a few facts and notation.

For an R -module M , submodule $N \subseteq M$, and ideal $\mathfrak{a} \subseteq R$, we consider the submodule $(N :_M \mathfrak{a}) = \{y \in M \mid \mathfrak{a}y \subseteq N\}$ of M . If $\mathfrak{a} = (x)$, just write $(N :_M x)$. The *socle* of M is $\mathrm{Soc} M = (0 :_M \mathfrak{m}) \cong \mathrm{Hom}_R(R/\mathfrak{m}, M)$. The *annihilator* of M is $\mathrm{ann}_R M = (0 :_R M)$.

For an ideal \mathfrak{a} of R , denote by $\Gamma_{\mathfrak{a}}(-)$ the \mathfrak{a} -torsion functor; its right derived functors yield the usual local cohomology functors, denoted $H_{\mathfrak{a}}^i(-)$ for $i \geq 0$; for additional background on local cohomology, refer to [1, 2, 10].

A *system of parameters of R* is a sequence of elements x_1, \dots, x_d in R such that $\mathfrak{m}^i \subseteq (x_1, \dots, x_d)$ for some integer i . More generally, if M is an R -module with

$\dim M = t$, then a sequence x_1, \dots, x_t in R is a *system of parameters* of M if $M/(x_1, \dots, x_t)M$ has finite length. In either case, an element of a system of parameters is called a *parameter* and an ideal generated by a system of parameters is a *parameter ideal*. For additional facts about systems of parameters, refer to [7, 12].

2. A BOUND IN DIMENSION ONE

Assume in this section that the ring (R, \mathfrak{m}, k) has an infinite residue field k and $\dim(R) = 1$. Moreover, we fix the next two invariants throughout this section; the first is finite because k is infinite¹ [2, Corollary 4.6.10], the second is finite because $\Gamma_{\mathfrak{m}}(R)$ has finite length [1, Theorem 7.1.3]:

$$(2.1) \quad \begin{aligned} c &= \inf\{i \mid \text{there exists a parameter } x \text{ of } R \text{ such that } \mathfrak{m}^{i+1} = (x)\mathfrak{m}^i\}; \\ g &= \inf\{i \mid \mathfrak{m}^i \cap \Gamma_{\mathfrak{m}}(R) = 0\}. \end{aligned}$$

These invariants have been considered elsewhere; c is the reduction number of \mathfrak{m} , and the bound we consider below, $\max\{c, g\} + 1$, is used by Rogers [13, Theorem 2.3]. We begin with two elementary lemmas involving these invariants:

Lemma 2.2. *Let $x \in R$ be a parameter and $y \in \mathfrak{m}^g$. If $x^i y = 0$ for some $i \geq 1$, then $y = 0$.*

Proof. As (x) is a parameter ideal, there exists an integer j such that $\mathfrak{m}^j \subseteq (x)$ hence $\mathfrak{m}^{ij} \subseteq (x^i)$ for $i \geq 1$. If $x^i y = 0$, then $\mathfrak{m}^{ij} y = 0$. It thus follows that $y \in \mathfrak{m}^g \cap (0 :_R \mathfrak{m}^{ij}) \subseteq \mathfrak{m}^g \cap \Gamma_{\mathfrak{m}}(R) = 0$, hence $y = 0$. \square

Lemma 2.3. *Let $x \in R$ be a parameter with $\mathfrak{m}^{c+1} = (x)\mathfrak{m}^c$ and set $n = \max\{c, g\}$. If $\{y_1, \dots, y_e\}$ is a minimal generating set of \mathfrak{m}^n , then $\{x^i y_1, \dots, x^i y_e\}$ is a minimal generating set of \mathfrak{m}^{n+i} for each $i \geq 1$.*

Proof. Let $\{y_1, \dots, y_e\}$ be a minimal generating set of \mathfrak{m}^n . As $n \geq c$, the equality $(x)\mathfrak{m}^n = \mathfrak{m}^{n+1}$ implies $(x^i)\mathfrak{m}^n = \mathfrak{m}^{n+i}$ by induction. Thus $(x^i y_1, \dots, x^i y_e) = \mathfrak{m}^{n+i}$. If there exists $r_q \in R$ such that $x^i y_j = \sum_{q \neq j} r_q x^i y_q$ for some j , then we have $x^i(y_j - \sum_{q \neq j} r_q y_q) = 0$. Since $y_j - \sum_{q \neq j} r_q y_q \in \mathfrak{m}^g$, we have $y_j - \sum_{q \neq j} r_q y_q = 0$ by Lemma 2.2; this contradicts the fact that y_1, \dots, y_e is a minimal generating set, hence we must have $\{x^i y_1, \dots, x^i y_e\}$ is a minimal generating set for \mathfrak{m}^{n+i} . \square

Theorem 2.4. *Assume k is infinite and $\dim(R) = 1$. For $i \geq \max\{c, g\} + 1$, the canonical map*

$$\varphi_i : \text{Ext}_R^1(R/\mathfrak{m}^i, R) \longrightarrow \varinjlim_j \text{Ext}_R^1(R/\mathfrak{m}^j, R) \cong H_{\mathfrak{m}}^1(R)$$

becomes surjective after applying $\text{Soc}(-) = \text{Hom}_R(R/\mathfrak{m}, -)$.

Proof. Let x be a parameter such that $(x)\mathfrak{m}^c = \mathfrak{m}^{c+1}$ and set $n = \max\{c, g\}$. Let $\{u_1, \dots, u_e\}$ be a minimal generating set for \mathfrak{m}^n . By Lemma 2.3, we know $\{x^i u_1, \dots, x^i u_e\}$ is a minimal generating set for \mathfrak{m}^{n+i} for $i \geq 0$. We will consider $\text{Soc} \text{Ext}_R^1(R/\mathfrak{m}^{n+i}, R)$ by examining a projective resolution of R/\mathfrak{m}^{n+i} .

We first show that, for $i \geq 0$, one may choose free resolutions of R/\mathfrak{m}^{n+i} which agree starting in degree 1. Set $\vec{u} = [u_1 \cdots u_e] : R^e \rightarrow R$. For $i \geq 0$, the containment $\ker(\vec{u}) \subseteq \ker(x^i \vec{u})$ is clear, and hence the equality $\ker(\vec{u}) = \ker(x^i \vec{u})$ holds because

¹For the purposes of this paper, the assumption of k being an infinite field may be replaced with the assumption that a parameter x exists so that $\mathfrak{m}^{i+1} = (x)\mathfrak{m}^i$ for some integer i .

if $\vec{r} \in \ker(x^i \vec{u})$, then $\vec{u} \vec{r} = 0$ by Lemma 2.2. Thus, for $i \geq 0$, there is a matrix $A : R^f \rightarrow R^e$ and a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} R^f & \xrightarrow{A} & R^e & \xrightarrow{x^{i+1} \vec{u}} & R & \longrightarrow & R/\mathfrak{m}^{n+i+1} & \longrightarrow & 0 \\ \downarrow & & \downarrow x & & \downarrow = & & \downarrow & & \\ R^f & \xrightarrow{A} & R^e & \xrightarrow{x^i \vec{u}} & R & \longrightarrow & R/\mathfrak{m}^{n+i} & \longrightarrow & 0 \end{array}$$

Applying $\text{Hom}_R(-, R)$ to this diagram yields a commutative diagram:

$$(2.5) \quad \begin{array}{ccccc} R & \xrightarrow{x^{i+1} \vec{u}^T} & R^e & \xrightarrow{A^T} & R^f \\ \uparrow = & & \uparrow x & & \uparrow \\ R & \xrightarrow{x^i \vec{u}^T} & R^e & \xrightarrow{A^T} & R^f \end{array}$$

Taking cohomology, we obtain that $\text{Ext}_R^1(R/\mathfrak{m}^{n+i}, R) = \ker(A^T)/\text{im}(x^i \vec{u}^T)$ for $i \geq 0$. Set $K = \ker(A^T) \subseteq R^e$, $I_i = \text{im}(x^i \vec{u}^T) \subseteq K$, and identify

$$\text{Soc Ext}_R^1(R/\mathfrak{m}^{n+i}, R) \cong \text{Soc}(K/I_i)$$

for $i \geq 0$. Moreover, for $i \geq 0$, the map $\text{Ext}_R^1(R/\mathfrak{m}^{n+i}, R) \rightarrow \text{Ext}_R^1(R/\mathfrak{m}^{n+i+1}, R)$ is induced by multiplication by x , see (2.5), as well as is the induced map after applying $\text{Soc}(-)$. Indeed, for $j \geq 1$, the map $x^j : \text{Soc}(K/I_i) \longrightarrow \text{Soc}(K/I_{i+j})$ is defined by $\vec{z} + I_i \longmapsto x^j \vec{z} + I_{i+j}$.

In order to show that $\text{Soc } \varphi_i$ is surjective for $i \geq n+1$, where φ_i is as in the statement, it will be enough to show that $\text{Soc } \varphi_{n+1}$ is surjective (this follows from the definition of direct systems). Note that for $\vec{v} + I_i \in \text{Soc } K/I_i$, the function $\text{Soc } \varphi_i$ is induced by φ_i and hence $(\text{Soc } \varphi_i)(\vec{v} + I_i) = \varphi_i(\vec{v} + I_i)$. Hereafter, we use the latter notation.

Let $\sigma \in \varinjlim_j \text{Soc } K/I_j$. As $\varinjlim_j \text{Soc } K/I_j (\cong \text{Soc } H_{\mathfrak{m}}^d(R))$ is finitely generated, $\text{Soc } \varphi_i$ is surjective for $i \gg 0$, thus $\sigma = \varphi_{n+p}(\vec{v} + I_p)$ for some $\vec{v} + I_p \in \text{Soc } K/I_p$ for some $p \geq 1$. If $p = 1$, then $\sigma \in \text{im } \text{Soc } \varphi_{n+1}$ as desired, so assume $p > 1$. We proceed by descending induction: that is, we aim to show there is an element $\vec{w} + I_{p-1} \in \text{Soc } K/I_{p-1}$ such that

$$x^i(\vec{v} + I_p) = x^{i+1}(\vec{w} + I_{p-1})$$

for some $i \geq 1$, and hence $\varphi_{n+p-1}(\vec{w} + I_{p-1}) = \varphi_{n+p}(\vec{v} + I_p)$.

We consider the element $x^g(\vec{v} + I_p) = x^g \vec{v} + I_{p+g}$, recalling that g is the least integer such that $\mathfrak{m}^g \cap \Gamma_{\mathfrak{m}}(R) = 0$. As $x^g \vec{v} + I_{p+g}$ is a socle element, we have:

$$\begin{aligned} x(x^g \vec{v} + I_{p+g}) &= 0 + I_{p+g} \\ \implies x^{g+1} \vec{v} &\in I_{p+g} \\ \implies x^{g+1} \vec{v} &= ax^{p+g} \vec{u}^T \text{ for some } a \in R, \text{ recalling } I_{p+g} = \text{im}(x^{p+g} \vec{u}^T), \\ \implies x(x^g \vec{v} - ax^{p+g-1} \vec{u}^T) &= 0 \\ \implies x^g \vec{v} &= ax^{p+g-1} \vec{u}^T, \text{ by Lemma 2.2.} \end{aligned}$$

Since $p \geq 2$, we may set $\vec{w} = ax^{p-2} \vec{u}^T$, and notice that $x^g \vec{v} = x^{g+1} \vec{w}$.

We claim $\vec{w} + I_{p-1} \in \text{Soc}(K/I_{p-1})$. First, $\vec{v} \in K$ implies that $A^T \vec{v} = 0$, hence $0 = x^g A^T \vec{v} = x^{g+1} A^T \vec{w}$. As the entries of \vec{w} are contained in \mathfrak{m}^g , so are the entries

of $A^T \bar{w}$. Lemma 2.2 yields $A^T \bar{w} = 0$, hence $\bar{w} \in K$. Next, for any $z \in \mathfrak{m}$, we have

$$zx^{g+1} \bar{w} = zx^g \bar{v} = bx^{p+g} \bar{u}^T, \text{ for some } b \in R,$$

since $x^g \bar{v} + I_{p+g}$ is a socle element in K/I_{p+g} . Thus $x^{g+1}(z\bar{w} - bx^{p-1} \bar{u}^T) = 0$. The entries of $z\bar{w} - bx^{p-1} \bar{u}^T$ are all in \mathfrak{m}^g , so Lemma 2.2 implies that $z\bar{w} = bx^{p-1} \bar{u}^T$. Therefore $\bar{w} + I_{p-1} \in \text{Soc}(K/I_{p-1})$, hence

$$\begin{aligned} \varphi_{n+p}(\bar{v} + I_p) &= \varphi_{n+p+g}(x^g \bar{v} + I_{g+p}) \\ &= \varphi_{n+p+g}(x^{g+1} \bar{w} + I_{g+p}) \\ &= \varphi_{n+p-1}(\bar{w} + I_{p-1}). \end{aligned}$$

By descending induction, there exists $\bar{w}' + I_1 \in \text{Soc } K/I_1$ such that

$$\varphi_{n+p}(\bar{v} + I_p) = \varphi_{n+1}(\bar{w}' + I_1).$$

The desired map $\text{Soc } \varphi_i$ is therefore surjective for $i \geq n+1 = \max\{c, g\} + 1$. \square

The next example shows that the bound in Theorem 2.4 is not sharp.

Example 2.6. Let k be an infinite field and $a \geq 1$ a fixed integer. Consider the local ring $Q = k[[x, y]]/(x^{a+1}, xy^a)$ with maximal ideal $\mathfrak{m} = (x, y)$. A direct computation shows that the invariants in (2.1) for the ring Q satisfy $c \leq a$, since $(y)\mathfrak{m}^a = \mathfrak{m}^{a+1}$, and $g = 2a$. Thus $\max\{c, g\} + 1 = 2a + 1$.

For $i \geq 2a$, one has $\mathfrak{m}^i = (y^i)$, and the natural surjection $Q/\mathfrak{m}^{i+1} \rightarrow Q/\mathfrak{m}^i$ induces a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & Q & \xrightarrow{x} & Q & \xrightarrow{y^{i+1}} & Q & \longrightarrow & Q/\mathfrak{m}^{i+1} & \longrightarrow & 0 \\ & & \downarrow y & & \downarrow y & & \downarrow = & & \downarrow & & \\ \cdots & \longrightarrow & Q & \xrightarrow{x} & Q & \xrightarrow{y^i} & Q & \longrightarrow & Q/\mathfrak{m}^i & \longrightarrow & 0. \end{array}$$

Applying $\text{Hom}_Q(-, Q)$ to this diagram, observe that $\text{Ext}_Q^1(Q/\mathfrak{m}^i, Q) = (x^a, y^a)/(y^i)$ for each $i \geq 2a$. Moreover, the induced maps $\text{Ext}_Q^1(Q/\mathfrak{m}^i, Q) \rightarrow \text{Ext}_Q^1(Q/\mathfrak{m}^{i+1}, Q)$ are multiplication by y .

Theorem 2.4 shows the canonical map $\text{Soc } \varphi_i : \text{Soc } \text{Ext}_Q^1(Q/\mathfrak{m}^i, Q) \rightarrow \text{Soc } H_{\mathfrak{m}}^1(Q)$ is surjective for $i \geq 2a + 1$; we claim that in fact $\text{Soc } \varphi_{2a}$ is surjective as well. First note that for $i \geq 2a$, one has $\text{Soc } \text{Ext}_Q^1(Q/\mathfrak{m}^i, Q) \cong (x^a y^{a-1}, y^{i-1})/(y^i)$. Thus it will suffice to consider the direct system

$$(x^a y^{a-1}, y^{2a-1})/(y^{2a}) \xrightarrow{y} (x^a y^{a-1}, y^{2a})/(y^{2a+1}) \xrightarrow{y} \cdots$$

In this direct system, the element $x^a y^{a-1} \in (x^a y^{a-1}, y^{2a})/(y^{2a+1})$ is sent to 0, hence surjectivity of $\text{Soc } \varphi_{2a+1}$ implies $\text{Soc } H_{\mathfrak{m}}^1(Q)$ is generated by $\text{Soc } \varphi_{2a+1}(y^{2a})$. Moreover, $y^{2a} \in (x^a y^{a-1}, y^{2a})/(y^{2a+1})$ is the image of $y^{2a-1} \in (x^a y^{a-1}, y^{2a-1})/(y^{2a})$, and so $\text{Soc } H_{\mathfrak{m}}^1(Q)$ is also generated by $\text{Soc } \varphi_{2a}(y^{2a-1})$, thus $\text{Soc } \varphi_{2a}$ is surjective.

Recall that an ideal \mathfrak{q} of R is reducible if $\mathfrak{q} = \mathfrak{b} \cap \mathfrak{c}$ for two ideals \mathfrak{b} and \mathfrak{c} of R strictly containing \mathfrak{q} ; if such a decomposition is not possible, then \mathfrak{q} is irreducible. The following consequence of Theorem 2.4 allows us to characterize Gorenstein rings in terms of the existence of irreducible parameter ideals in \mathfrak{m}^n for $n = \max\{c, g\} + 1$; it can also be obtained using Rogers' [13, Theorem 2.3] in place of Theorem 2.4.

Corollary 2.7. *Assume k is infinite and $\dim(R) = 1$. Set $n = \max\{c, g\} + 1$. The ring R is Gorenstein if and only if some parameter ideal in \mathfrak{m}^n is irreducible.*

Proof. This follows from Theorem 2.4 and [11, Theorem 2.7]. \square

The least integer ℓ required to determine whether R is Gorenstein in terms of the existence of an irreducible parameter ideal in \mathfrak{m}^ℓ depends on R and is thus at most $\max\{c, g\} + 1$ in the case of a dimension 1 local ring with an infinite residue field. We next show that given an integer a , there exists a ring with $a < \ell \leq 2a$.

Example 2.8. Let us return to the ring $Q = k[[x, y]]/(x^{a+1}, xy^a)$ and the setting of Example 2.6. The ring Q has dimension 1 and depth 0, hence is non-Gorenstein. As noted above, $\max\{c, g\} + 1 = 2a + 1$, so Corollary 2.7 implies that every parameter ideal in \mathfrak{m}^{2a+1} is reducible, hence $\ell \leq 2a + 1$. In fact, the computation in Example 2.6—which relies on Theorem 2.4—shows that $\ell \leq 2a$ and that every parameter ideal in \mathfrak{m}^{2a} is reducible.

On the other hand, the parameter ideal (y^a) is irreducible, thus $\ell > a$. To see this, it is enough to show that any ideal of Q properly containing (y^a) also contains the nonzero element $x^a y^{a-1}$. Let $\mathfrak{b} \subseteq Q$ be an ideal that properly contains (y^a) and fix $\beta \in \mathfrak{b} \setminus (y^a)$. Write $\beta = \sum_{s,t \geq 0} a_{s,t} x^s y^t$, with $a_{s,t} \in k$. Because $\beta \notin (y^a)$, the set $\Lambda = \{(s, t) \mid a_{s,t} \neq 0, s \leq a, \text{ and } t \leq a - 1\}$ is nonempty. Choose $(s_0, t_0) \in \Lambda$ with $s_0 + t_0 \leq s + t$ for all $(s, t) \in \Lambda$. Noting the element $x^{a-s_0} y^{a-1-t_0} \beta$ belongs to $\mathfrak{m}^{2a-1} = (x^a y^{a-1}, y^{2a-1})$, it follows that the ideal \mathfrak{b} contains the element $a_{s_0, t_0}^{-1} x^{a-s_0} y^{a-1-t_0} \beta = x^a y^{a-1} + \varepsilon$ with $\varepsilon \in (y^{2a-1}) \subseteq (y^a)$. Thus $x^a y^{a-1} \in \mathfrak{b}$.

3. A BOUND IN HIGHER DIMENSIONS

For rings of higher dimension, the problem of determining surjectivity of the socle of the map in (1.2) becomes more subtle, with obstructions similar to those noted by Fouli and Huneke [8, Discussion 4.5]. In particular, it is not clear to us whether the same type of “lifting” technique employed in Theorem 2.4 can be used to show surjectivity of the socle of the map in (1.2) if $\dim R > 1$. Our solution here is to instead consider surjectivity of the socle of the map in (1.4) for p_s -standard systems of parameters (defined below).

For this section, (R, \mathfrak{m}, k) is a commutative noetherian local ring with $\dim R = d$.

Remark 3.1. Let M be a finitely generated R -module with $\dim M = t > 0$ and let x be a parameter of M . The exact sequence $0 \rightarrow M/(0 :_M x) \xrightarrow{x} M \rightarrow M/xM \rightarrow 0$ induces a canonical connecting homomorphism $H_{\mathfrak{m}}^{t-1}(M/xM) \rightarrow H_{\mathfrak{m}}^t(M/(0 :_M x))$. Moreover, the containment $(x) + \text{ann}_R M \subseteq \text{ann}_R(0 :_M x)$ implies $\dim(0 :_M x) < t$, hence the exact sequence $0 \rightarrow (0 :_M x) \rightarrow M \rightarrow M/(0 :_M x) \rightarrow 0$ along with Grothendieck’s Vanishing Theorem (for example, [1, 6.1.2]) yields an isomorphism

$$(3.2) \quad H_{\mathfrak{m}}^t(M) \xrightarrow{\cong} H_{\mathfrak{m}}^t(M/(0 :_M x)).$$

Composing the connecting homomorphism $H_{\mathfrak{m}}^{t-1}(M/xM) \rightarrow H_{\mathfrak{m}}^t(M/(0 :_M x))$ from above with the inverse of the isomorphism in (3.2) yields a homomorphism

$$\delta_x^M : H_{\mathfrak{m}}^{t-1}(M/xM) \longrightarrow H_{\mathfrak{m}}^t(M).$$

The next two lemmas are the primary tools for proving one of our main results below, Proposition 3.7. In light of the isomorphism in (3.2), this first lemma essentially follows from a result of Cuong and Quy [6, Proposition 2.1], but we spell

out some of the details in order to keep track of the map inducing the surjection, which we will need later.

Lemma 3.3. *Let M be a finitely generated R -module with $\dim M = t > 0$. Let x be a parameter of M . If $x \in \text{ann}_R H_{\mathfrak{m}}^{t-1}(M)$, then for $s \geq 2$ the map $\delta_{x^s}^M$ defined in Remark 3.1 induces a split surjection*

$$\text{Soc } \delta_{x^s}^M : \text{Soc } H_{\mathfrak{m}}^{t-1}(M/x^s M) \longrightarrow \text{Soc } H_{\mathfrak{m}}^t(M).$$

Proof. The inclusion $(0 :_M x) \subseteq (0 :_M x^s)$ induces the left vertical map in the next commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \frac{M}{(0 :_M x)} & \xrightarrow{x} & M & \longrightarrow & M/xM & \longrightarrow & 0 \\ & & \downarrow & & \downarrow x^{s-1} & & \downarrow x^{s-1} & & \\ 0 & \longrightarrow & \frac{M}{(0 :_M x^s)} & \xrightarrow{x^s} & M & \longrightarrow & M/x^s M & \longrightarrow & 0 \end{array}$$

From (3.2), using that both x and x^s are parameters of M , there are isomorphisms $H_{\mathfrak{m}}^t(M/(0 :_M x^s)) \cong H_{\mathfrak{m}}^t(M) \cong H_{\mathfrak{m}}^t(M/(0 :_M x))$. We thus obtain a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} H_{\mathfrak{m}}^{t-1}(M) & \xrightarrow{x} & H_{\mathfrak{m}}^{t-1}(M) & \longrightarrow & H_{\mathfrak{m}}^{t-1}(M/xM) & \xrightarrow{\delta'} & H_{\mathfrak{m}}^t(M) & \xrightarrow{x} & H_{\mathfrak{m}}^t(M) \\ \downarrow & & \downarrow x^{s-1} & & \downarrow x^{s-1} & & \downarrow = & & \downarrow x^{s-1} \\ H_{\mathfrak{m}}^{t-1}(M) & \xrightarrow{x^s} & H_{\mathfrak{m}}^{t-1}(M) & \longrightarrow & H_{\mathfrak{m}}^{t-1}(M/x^s M) & \xrightarrow{\delta} & H_{\mathfrak{m}}^t(M) & \xrightarrow{x^s} & H_{\mathfrak{m}}^t(M) \end{array}$$

where $\delta = \delta_{x^s}^M$ and $\delta' = \delta_x^M$ are the maps defined in Remark 3.1. By assumption, $xH_{\mathfrak{m}}^{t-1}(M) = 0$, hence this yields the next commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H_{\mathfrak{m}}^{t-1}(M) & \longrightarrow & H_{\mathfrak{m}}^{t-1}(M/xM) & \longrightarrow & (0 :_{H_{\mathfrak{m}}^t(M)} x) & \longrightarrow & 0 \\ & & \downarrow 0 & & \downarrow x^{s-1} & & \downarrow \subseteq \downarrow \iota & & \\ 0 & \longrightarrow & H_{\mathfrak{m}}^{t-1}(M) & \longrightarrow & H_{\mathfrak{m}}^{t-1}(M/x^s M) & \xrightarrow{\delta} & (0 :_{H_{\mathfrak{m}}^t(M)} x^s) & \longrightarrow & 0 \end{array}$$

Following the argument in [6, proof of Proposition 2.1], note that the middle vertical map induces $\varepsilon : (0 :_{H_{\mathfrak{m}}^t(M)} x) \rightarrow H_{\mathfrak{m}}^{t-1}(M/x^s M)$ such that $\delta\varepsilon = \iota$. Since ι is the inclusion, and $\text{Soc}(0 :_{H_{\mathfrak{m}}^t(M)} x) = \text{Soc}(0 :_{H_{\mathfrak{m}}^t(M)} x^s) = \text{Soc } H_{\mathfrak{m}}^t(M)$, we see that $\text{Soc } \iota = 1_{\text{Soc}(H_{\mathfrak{m}}^t(M))}$ and thus $(\text{Soc } \delta)(\text{Soc } \varepsilon) = 1_{\text{Soc}(H_{\mathfrak{m}}^t(M))}$. It follows that $\text{Soc } \delta = \text{Soc } \delta_{x^s}^M$ is a split surjection. \square

Given a system of parameters x_1, \dots, x_d of R , the canonical map in (1.4) to the direct limit $R/(x_1^i, \dots, x_d^i) \longrightarrow \varinjlim_j R/(x_1^j, \dots, x_d^j)$ is defined by the direct system

$$R/(x_1, \dots, x_d) \xrightarrow{x_1 \cdots x_d} R/(x_1^2, \dots, x_d^2) \xrightarrow{x_1 \cdots x_d} \cdots$$

Moreover, there is a unique isomorphism, see [1, Theorem 5.2.9]:

$$\varinjlim_j R/(x_1^j, \dots, x_d^j) \xrightarrow{\cong} H_{\mathfrak{m}}^d(R).$$

Lemma 3.4. *Let x_1, \dots, x_d be a system of parameters of R . The canonical map $R/(x_1, \dots, x_d) \rightarrow \varinjlim_j R/(x_1^j, \dots, x_d^j) \cong H_m^d(R)$ agrees with the composition*

$$H_m^0(R/(x_1, \dots, x_d)) \rightarrow H_m^1(R/(x_2, \dots, x_d)) \rightarrow \cdots \rightarrow H_m^{d-1}(R/(x_d)) \rightarrow H_m^d(R)$$

of homomorphisms defined in Remark 3.1.

Proof. First note that $R/(x_1, \dots, x_d) = H_m^0(R/(x_1, \dots, x_d))$. Further, the canonical map $R/(x_1, \dots, x_d) \rightarrow \varinjlim_j R/(x_1^j, \dots, x_d^j)$ can be decomposed as the composition of the following canonical maps:

$$\frac{R}{(x_1, \dots, x_d)} \rightarrow \varinjlim_j \frac{R}{(x_1^j, x_2, \dots, x_d)} \rightarrow \cdots \rightarrow \varinjlim_j \frac{R}{(x_1^j, \dots, x_{d-1}^j, x_d)} \rightarrow \varinjlim_j \frac{R}{(x_1^j, \dots, x_d^j)}.$$

Fix $0 < t \leq d$. It is therefore sufficient to show that the next two maps agree up to isomorphism; indeed, by [1, Theorem 5.2.9] there is a unique isomorphism between the domains, and another between the codomains:

$$\alpha_t : \varinjlim_i \frac{R}{(x_1^i, \dots, x_{t-1}^i, x_t, \dots, x_d)} \longrightarrow \varinjlim_j \varinjlim_i \frac{R}{(x_1^i, \dots, x_{t-1}^i, x_t^j, x_{t+1}, \dots, x_d)}, \text{ and}$$

$$\delta_t : H_m^{t-1}\left(\frac{R}{(x_t, \dots, x_d)}\right) \longrightarrow H_m^t\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right),$$

where α_t is the canonical map to the direct limit and $\delta_t = \delta_{x_t}^{R/(x_{t+1}, \dots, x_d)}$ is the map defined in Remark 3.1.

For $0 \leq u \leq d-1$, there is by [1, Theorem 5.2.9] a natural equivalence of functors $\varinjlim_j \left(\frac{R}{(x_1^j, \dots, x_u^j)} \otimes_R -\right) \xrightarrow{\cong} H_{(x_1, \dots, x_u)}^u(-)$. This provides the isomorphisms and commutativity in the following diagram:

$$\begin{array}{ccc} \varinjlim_i \frac{R}{(x_1^i, \dots, x_{t-1}^i, x_t, \dots, x_d)} & \xrightarrow{\cong} & H_m^{t-1}\left(\frac{R}{(x_t, \dots, x_d)}\right) \\ \downarrow \alpha_t & & \downarrow \beta_t \\ \varinjlim_j \varinjlim_i \frac{R}{(x_1^i, \dots, x_{t-1}^i, x_t^j, x_{t+1}, \dots, x_d)} & \xrightarrow{\cong} & \varinjlim_j H_m^{t-1}\left(\frac{R}{(x_t^j, x_{t+1}, \dots, x_d)}\right), \end{array}$$

where β_t is the canonical map to the direct limit and the modules on the right have also utilized the fact that $H_{(x_1, \dots, x_{t-1})}^{t-1}\left(\frac{R}{(x_t^j, x_{t+1}, \dots, x_d)}\right) \cong H_m^{t-1}\left(\frac{R}{(x_t^j, x_{t+1}, \dots, x_d)}\right)$ for each $j \geq 1$; see [1, Exercise 2.1.9]. It remains only to show that β_t is isomorphic to δ_t . To see this, consider the short exact sequence of direct systems:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \frac{R/(x_{t+1}, \dots, x_d)}{(0:x_t)} & \xrightarrow{x_t} & R/(x_{t+1}, \dots, x_d) & \longrightarrow & R/(x_t, \dots, x_d) \longrightarrow 0 \\ & & \downarrow & & \downarrow x_t & & \downarrow x_t \\ 0 & \longrightarrow & \frac{R/(x_{t+1}, \dots, x_d)}{(0:x_t^2)} & \xrightarrow{x_t^2} & R/(x_{t+1}, \dots, x_d) & \longrightarrow & R/(x_t^2, x_{t+1}, \dots, x_d) \longrightarrow 0 \\ & & \downarrow & & \downarrow x_t & & \downarrow x_t \\ & & \vdots & & \vdots & & \vdots \end{array}$$

Each row of this diagram yields a long exact sequence in local cohomology; along with the homomorphisms defined in Remark 3.1, this gives the following commutative diagram with exact rows:

$$\begin{array}{ccccccc}
 H_{\mathfrak{m}}^{t-1}\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right) & \longrightarrow & H_{\mathfrak{m}}^{t-1}\left(\frac{R}{(x_t, \dots, x_d)}\right) & \xrightarrow{\delta_t} & H_{\mathfrak{m}}^t\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right) & \longrightarrow & H_{\mathfrak{m}}^t\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right) \\
 \downarrow x_t & & \downarrow x_t & & \downarrow = & & \downarrow x_t \\
 H_{\mathfrak{m}}^{t-1}\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right) & \longrightarrow & H_{\mathfrak{m}}^{t-1}\left(\frac{R}{(x_t^2, x_{t+1}, \dots, x_d)}\right) & \longrightarrow & H_{\mathfrak{m}}^t\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right) & \longrightarrow & H_{\mathfrak{m}}^t\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right) \\
 \downarrow x_t & & \downarrow x_t & & \downarrow = & & \downarrow x_t \\
 \vdots & & \vdots & & \vdots & & \vdots
 \end{array}$$

The map from the second term in the first row, $H_{\mathfrak{m}}^{t-1}\left(\frac{R}{(x_t, \dots, x_d)}\right)$, to the direct limit of the second column is β_t . The direct limits of the left and right columns are the localizations $(H_{\mathfrak{m}}^{t-1}\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right))_{x_t}$ and $(H_{\mathfrak{m}}^t\left(\frac{R}{(x_{t+1}, \dots, x_d)}\right))_{x_t}$, respectively. These are zero since they are both \mathfrak{m} -torsion [1, 2.1.3] and multiplication by $x_t \in \mathfrak{m}$ is invertible on either module. Hence the direct limit of the middle maps is an isomorphism, showing that β_t and δ_t are isomorphic. \square

In order to prove surjectivity of the socle of the map in (1.4), we introduce the following special types of systems of parameters. Let M be a finitely generated R -module with $\dim M = t > 0$. As is standard, denote the annihilator of $H_{\mathfrak{m}}^i(M)$ by $\mathfrak{a}_i(M) = \text{ann}_R H_{\mathfrak{m}}^i(M)$, and put $\mathfrak{a}(M) = \mathfrak{a}_0(M) \cdots \mathfrak{a}_{t-1}(M)$. A system of parameters x_1, \dots, x_t of M is called a *p-standard system of parameters* if $x_t \in \mathfrak{a}(M)$ and $x_i \in \mathfrak{a}(M/(x_{i+1}, \dots, x_t)M)$ for $i = 1, \dots, t-1$. Such systems were defined at this level of generality by Cuong [3], who noted that a result of Schenzel [14, Korollar 2.2.4] implies that every finitely generated R -module has a *p-standard system of parameters* provided R has a dualizing complex. Indeed, if R has a dualizing complex, then $\dim R/\mathfrak{a}(M) < t$ by [14, Korollar 2.2.4] and so prime avoidance provides an element $x_t \in \mathfrak{a}(M)$ that is a parameter of M . Inductively, this shows the existence of *p-standard systems of parameters*, as well as the existence of the following variant², provided R has a dualizing complex.

Definition 3.5. Let M be a finitely generated R -module with $\dim M = t$. For an integer $s \geq 1$, a system of parameters x_1, \dots, x_t of M is called a *p_s-standard system of parameters* if $x_t \in \mathfrak{a}(M)$ and $x_i \in \mathfrak{a}(M/(x_{i+1}^s, \dots, x_t^s)M)$ for $i = 1, \dots, t-1$.

Evidently, *p₁-standard* and *p-standard* systems of parameters are the same, but the relationship between *p_s-standard* and *p-standard* systems of parameters is less straightforward for $s \geq 2$.

Remark 3.6. If x_1, \dots, x_t is a *p_s-standard system of parameters* of M for an integer $s \geq 1$, then x_1^s, \dots, x_t^s is a *p-standard system of parameters* of M ; this follows straight from the definitions. Thus the existence of a *p_s-standard system of parameters* of M for some $s \geq 1$ implies the existence of a *p-standard system of parameters* of M .

²In fact, one may find by [14, Korollar 2.2.4] an element in $\mathfrak{a}_{t-1}(M)$ that is a parameter of M , provided R has a dualizing complex. Hence for our purposes, one may instead find a system of parameters x_1, \dots, x_t of M satisfying $x_i \in \mathfrak{a}_{i-1}(M/(x_{i+1}^s, \dots, x_t^s)M)$ for $i = 1, \dots, t$. The notion considered in Definition 3.5 is chosen to be reminiscent of *p-standard systems*.

Conversely, it follows from work of Cuong and Cuong [4] that the existence of a p -standard system of parameters of M implies the existence of a p_s -standard system of parameters of M for every $s \geq 1$: indeed, if x_1, \dots, x_t is a p -standard system of parameters of M and $s \geq 1$, then [4, Corollary 3.9] implies that $x_1^t, \dots, x_i^t, x_{i+1}^{st}, \dots, x_t^{st}$ is also p -standard for each $i = 1, \dots, t$ hence x_1^t, \dots, x_t^t is a p_s -standard system of parameters of M . Moreover, Cuong and Cuong prove in [5, Theorem 5.2] that the existence of a p -standard system of parameters is equivalent to the ring R being a quotient of a local Cohen-Macaulay ring.

In this paper, we consistently express our results in terms of p_s -standard systems of parameters so that the exposition is relatively self-contained, but combining with the aforementioned results one can reformulate our results in terms of p -standard systems of parameters and observe that Theorem 3.9 below applies to any ring R that is a quotient of a local Cohen-Macaulay ring.

We now come to proving surjectivity of the socle of the map in (1.4) for p_s -standard systems of parameters if $s \geq 2$. The main distinction between the next result and [11, Proposition 2.5] or [9, Lemma 3.12] is that here we have some control for the point at which the induced maps on socles are surjective.

Proposition 3.7. *Let x_1, \dots, x_d be a system of parameters of R . If x_1, \dots, x_d is a p_s -standard system of parameters for some $s \geq 2$, then the canonical map $R/(x_1^s, \dots, x_d^s) \rightarrow \varinjlim_j R/(x_1^j, \dots, x_d^j) \cong H_m^d(R)$ induces a split surjection*

$$\text{Soc } R/(x_1^s, \dots, x_d^s) \longrightarrow \text{Soc } H_m^d(R).$$

Hence the canonical map $\text{Soc } R/(x_1^i, \dots, x_d^i) \rightarrow \text{Soc } H_m^d(R)$ is surjective for $i \geq s$.

Proof. Suppose x_1, \dots, x_d is a p_s -standard system of parameters of R for some $s \geq 2$. By definition, we have $x_t \in \mathfrak{a}(R/(x_{t+1}^s, \dots, x_d^s)) \subseteq \text{ann}_R H_m^{t-1}(R/(x_{t+1}^s, \dots, x_d^s))$ for each $t = 1, \dots, d$, and so Lemma 3.3 yields that the induced map

$$\text{Soc } H_m^{t-1}(R/(x_t^s, x_{t+1}^s, \dots, x_d^s)) \longrightarrow \text{Soc } H_m^t(R/(x_{t+1}^s, \dots, x_d^s))$$

is a split surjection for each $t = 1, \dots, d$. Thus the composition

$$H_m^0(R/(x_1^s, \dots, x_d^s)) \rightarrow H_m^1(R/(x_2^s, \dots, x_d^s)) \rightarrow \dots \rightarrow H_m^{d-1}(R/(x_d^s)) \rightarrow H_m^d(R),$$

which agrees with the canonical map $R/(x_1^s, \dots, x_d^s) \rightarrow \varinjlim_j R/(x_1^j, \dots, x_d^j) \cong H_m^d(R)$ by Lemma 3.4, induces the desired split surjection on socles.

The final remark follows from the definition of the direct system. \square

Remark 3.8. Let y_1, \dots, y_d and x_1, \dots, x_d be systems of parameters of R such that $(y_1, \dots, y_d) \subseteq (x_1, \dots, x_d)$. Suppose $A = (a_{ij})$ and $B = (b_{ij})$ are matrices such that $y_i = \sum_{j=1}^d a_{ij} x_j = \sum_{j=1}^d b_{ij} x_j$. One has $\det A \cdot (x_1, \dots, x_d) \subseteq (y_1, \dots, y_d)$, thus multiplication by $\det A$ induces a well-defined map $R/(x_1, \dots, x_d) \xrightarrow{\det A} R/(y_1, \dots, y_d)$; see [8, p. 2681]. Moreover, the proof of [8, Corollary 2.5] shows that multiplication by either $\det A$ or $\det B$ determines the same map from $R/(x_1, \dots, x_d)$ to $R/(y_1, \dots, y_d)$.

After a reduction in order to apply Proposition 3.7 in place of [11, Proposition 2.5], the proof of the next result is similar to that of [11, Theorem 2.7]. Let $s \geq 1$ be an integer and recall from above that a p_s -standard system of parameters exists if R has a dualizing complex, for example if R is complete; see also Remark 3.6.

Theorem 3.9. *Suppose there exists a p_s -standard system of parameters x_1, \dots, x_d of R for some $s \geq 2$. The ring R is Gorenstein if and only if some parameter ideal contained in (x_1^s, \dots, x_d^s) is irreducible.*

Proof. Suppose x_1, \dots, x_d is a p_s -standard system of parameters of R for some $s \geq 2$. All parameter ideals in a Gorenstein ring are irreducible [12, Theorem 18.1], so it is sufficient to prove the converse.

Assume y_1, \dots, y_d is a system of parameters such that (y_1, \dots, y_d) is irreducible and contained in (x_1^s, \dots, x_d^s) . We first claim that, for the direct system

$$R/(y_1, \dots, y_d) \xrightarrow{y_1 \cdots y_d} R/(y_1^2, \dots, y_d^2) \xrightarrow{y_1 \cdots y_d} \cdots,$$

the canonical map $R/(y_1, \dots, y_d) \rightarrow \varinjlim_j R/(y_1^j, \dots, y_d^j) \cong H_m^d(R)$ is surjective when restricted to socles. As x_1^j, \dots, x_d^j and y_1^j, \dots, y_d^j are systems of parameters for all $j \geq 1$, there exist families of positive integers $\{1 = t_1 < t_2 < \cdots\}$, $\{u_1 < u_2 < \cdots\}$, and $\{v_1 < v_2 < \cdots\}$ such that for each $i \geq 1$ we have containments:

$$(x_1^{i_s}, \dots, x_d^{i_s}) \supseteq (y_1^{t_i}, \dots, y_d^{t_i}) \supseteq (x_1^{u_i}, \dots, x_d^{u_i}) \supseteq (y_1^{v_i}, \dots, y_d^{v_i}).$$

By Remark 3.8, we obtain maps (coming from determinants of matrices) making the following commutative diagram of direct systems:

$$\begin{array}{ccccccc} R/(x_1^s, \dots, x_d^s) & \xrightarrow{\sigma_s} & R/(y_1, \dots, y_d) & \xrightarrow{\tau_s} & R/(x_1^{u_1}, \dots, x_d^{u_1}) & \xrightarrow{\rho_s} & R/(y_1^{v_1}, \dots, y_d^{v_1}) \\ \downarrow x_1^s \cdots x_d^s & & \downarrow y_1^{t_2-1} \cdots y_d^{t_2-1} & & \downarrow x_1^{u_2-u_1} \cdots x_d^{u_2-u_1} & & \downarrow y_1^{v_2-v_1} \cdots y_d^{v_2-v_1} \\ R/(x_1^{2s}, \dots, x_d^{2s}) & \xrightarrow{\sigma_{2s}} & R/(y_1^{t_2}, \dots, y_d^{t_2}) & \xrightarrow{\tau_{2s}} & R/(x_1^{u_2}, \dots, x_d^{u_2}) & \xrightarrow{\rho_{2s}} & R/(y_1^{v_2}, \dots, y_d^{v_2}) \\ \downarrow x_1^s \cdots x_d^s & & \downarrow y_1^{t_3-t_2} \cdots y_d^{t_3-t_2} & & \downarrow x_1^{u_3-u_2} \cdots x_d^{u_3-u_2} & & \downarrow y_1^{v_3-v_2} \cdots y_d^{v_3-v_2} \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

Moreover, Remark 3.8 also yields that the compositions of horizontal maps are the familiar ones: $\tau_{is}\sigma_{is} = x_1^{u_i-i_s} \cdots x_d^{u_i-i_s}$ and $\rho_{is}\tau_{is} = y_1^{v_i-t_i} \cdots y_d^{v_i-t_i}$, for $i \geq 1$. The direct limits of all four columns are isomorphic to $H_m^d(R)$, and it thus follows from the universal property of direct limits that the induced maps on these direct limits are isomorphisms. Hence we obtain a commutative diagram of canonical maps:

$$\begin{array}{ccc} \text{Soc } R/(x_1^s, \dots, x_d^s) & \xrightarrow{\text{Soc } \sigma_s} & \text{Soc } R/(y_1, \dots, y_d) \\ \downarrow & & \downarrow \\ \text{Soc } H_m^d(R) & \xrightarrow{\cong} & \text{Soc } H_m^d(R) \end{array}$$

The left vertical map is a surjection by Proposition 3.7, hence it follows that the right vertical map is a surjection as well.

Let $\phi : R/(y_1, \dots, y_d) \rightarrow \varinjlim_j R/(y_1^j, \dots, y_d^j) \cong H_m^d(R)$ be the canonical map and proceed as in the proof of [11, Theorem 2.7]: Recall that the limit closure of y_1, \dots, y_d is defined as $\{y_1, \dots, y_d\}_R^{\text{lim}} = \bigcup_{n \geq 0} ((y_1^{n+1}, \dots, y_d^{n+1}) :_R y_1^n \cdots y_d^n)$. By [11, Remark 2.2], $\ker(\phi) = \{y_1, \dots, y_d\}_R^{\text{lim}}/(y_1, \dots, y_d)$. Applying $\text{Soc}(-)$ to the canonical maps, we obtain the next exact sequence, where surjectivity of the map on the right was shown above:

$$0 \longrightarrow \text{Soc } \ker(\phi) \longrightarrow \text{Soc } R/(y_1, \dots, y_d) \longrightarrow \text{Soc } H_m^d(R) \longrightarrow 0.$$

Irreducibility of (y_1, \dots, y_d) yields $\dim_{R/\mathfrak{m}} \text{Soc } R/(y_1, \dots, y_d) = 1$. As $H_{\mathfrak{m}}^d(R) \neq 0$ we obtain $\text{Socker}(\phi) = 0$ implying per [11, Proposition 2.3] that y_1, \dots, y_d is regular. Thus R is Cohen-Macaulay with $\dim_{R/\mathfrak{m}} \text{Soc } H_{\mathfrak{m}}^d(R) = 1$, hence Gorenstein. \square

As an immediate consequence of Theorem 3.9, we obtain Theorem 1.5 from the introduction.

Remark 3.10. If R has finite local cohomologies, in which case there is an integer $n_0 > 0$ such that $\mathfrak{m}^{n_0} H_{\mathfrak{m}}^i(R) = 0$ for $i < d$ (this is not assumed in the results above), a similar bound can be obtained by using [6, Corollary 4.3] in conjunction with [11, Theorem 2.7]; in particular, it follows from these that R is Gorenstein if and only if some parameter ideal contained in \mathfrak{m}^{2n_0} is irreducible. In particular, if R is Buchsbaum so that we have $\mathfrak{m} H_{\mathfrak{m}}^i(R) = 0$ for $i < d$, then R is Gorenstein if and only if some parameter ideal contained in \mathfrak{m}^2 is irreducible.

ACKNOWLEDGMENTS

The authors are grateful to Thomas Marley for comments on an early version of this paper and to the anonymous referee for helpful suggestions.

REFERENCES

- [1] M. P. Brodmann and R. Y. Sharp. *Local cohomology*, volume 136 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2013. An algebraic introduction with geometric applications.
- [2] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*. Cambridge University Press, 1998.
- [3] Nguyen Tu Cuong. p -standard systems of parameters and p -standard ideals in local rings. *Acta Math. Vietnam.*, 20(1):145–161, 1995.
- [4] Nguyen Tu Cuong and Doan Trung Cuong. dd-sequences and partial Euler-Poincaré characteristics of Koszul complex. *J. Algebra Appl.*, 6(2):207–231, 2007.
- [5] Nguyen Tu Cuong and Doan Trung Cuong. Local cohomology annihilators and Macaulayfication. *Acta Math. Vietnam.*, 42(1):37–60, 2017.
- [6] Nguyen Tu Cuong and Pham Hung Quy. A splitting theorem for local cohomology and its applications. *J. Algebra*, 331:512–522, 2011.
- [7] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [8] Louiza Fouli and Craig Huneke. What is a system of parameters? *Proc. Amer. Math. Soc.*, 139(8):2681–2696, 2011.
- [9] Shiro Goto and Hideto Sakurai. The equality $I^2 = QI$ in Buchsbaum rings. *Rend. Sem. Mat. Univ. Padova*, 110:25–56, 2003.
- [10] Srikanth B. Iyengar, Graham J. Leuschke, Anton Leykin, Claudia Miller, Ezra Miller, Anurag K. Singh, and Uli Walther. *Twenty-four hours of local cohomology*, volume 87 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2007.
- [11] Thomas Marley, Mark W. Rogers, and Hideto Sakurai. Gorenstein rings and irreducible parameter ideals. *Proc. Amer. Math. Soc.*, 136(1):49–53, 2008.
- [12] Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- [13] Mark W. Rogers. The index of reducibility of parameter ideals in low dimension. *J. Algebra*, 278(2):571–584, 2004.
- [14] Peter Schenzel. *Dualisierende Komplexe in der lokalen Algebra und Buchsbaum-Ringe*, volume 907 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1982. With an English summary.

(K. Shultis) DEPARTMENT OF MATHEMATICS, GONZAGA UNIVERSITY, SPOKANE, WA 99258,
USA

E-mail address: `shultis@gonzaga.edu`

(P. Thompson) INSTITUTT FOR MATEMATISKE FAG, NORWEGIAN UNIVERSITY OF SCIENCE AND
TECHNOLOGY, N-7491 TRONDHEIM, NORWAY

E-mail address: `peder.thompson@ntnu.no`