

# Demographics of disks around young very low-mass stars and brown dwarfs in Lupus

E. Sanchis<sup>1,2</sup>, L. Testi<sup>1,3,4</sup>, A. Natta<sup>4,5</sup>, C. F. Manara<sup>1</sup>, B. Ercolano<sup>2,3</sup>, T. Preibisch<sup>2</sup>, T. Henning<sup>6</sup>, S. Facchini<sup>1</sup>, A. Miotello<sup>1</sup>, I. de Gregorio-Monsalvo<sup>7,8</sup>, C. Lopez<sup>8</sup>, K. Mužić<sup>9</sup>, I. Pascucci<sup>10,11</sup>, A. Santamaría-Miranda<sup>7,12</sup>, A. Scholz<sup>13</sup>, M. Tazzari<sup>14</sup>, S. van Terwisga<sup>15</sup>, and J. P. Williams<sup>16</sup>

<sup>1</sup> European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei München, Germany

<sup>2</sup> Universitäts-Sternwarte, Ludwig-Maximilians-Universität München, Scheinerstrasse 1, D-81679 München, Germany

<sup>3</sup> Excellence Cluster Origins, Boltzmannstrasse 2, D-85748 Garching bei München, Germany

<sup>4</sup> INAF/Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy

<sup>5</sup> School of Cosmic Physics, Dublin Institute for Advanced Studies, 31 Fitzwilliams Place, Dublin 2, Ireland

<sup>6</sup> Max Planck Institute for Astronomy, Königstuhl 17, D-69117, Heidelberg, Germany

<sup>7</sup> Núcleo Milenio Formación Planetaria - NPF, Universidad de Valparaíso, Av. Gran Bretaña 1111, Valparaíso, Chile

<sup>8</sup> Atacama Large Millimeter/Submillimeter Array, Joint ALMA Observatory, Alonso de Córdova 3107, Vitacura 763-0355, Santiago, Chile

<sup>9</sup> CENTRA, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande, 1749-016 Lisboa, Portugal

<sup>10</sup> Lunar and Planetary Laboratory, The University of Arizona, Tucson, AZ 85721, USA

<sup>11</sup> Earths in Other Solar Systems Team, NASA Nexus for Exoplanet System Science, USA

<sup>12</sup> European Southern Observatory, 3107, Alonso de Córdova, Santiago de Chile

<sup>13</sup> SUPA, School of Physics & Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, UK

<sup>14</sup> Institute of Astronomy, University of Cambridge, Madingley Road, CB3 0HA Cambridge, UK

<sup>15</sup> Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

<sup>16</sup> Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA

Received ... / Accepted ...

## ABSTRACT

We present new 890  $\mu\text{m}$  continuum ALMA observations of 5 brown dwarfs (BDs) with infrared excess in Lupus I and III – which, in combination with 4 BDs previously observed, allowed us to study the mm properties of the full known BD disk population of one star-forming region. Emission is detected in 5 out of the 9 BD disks. Dust disk mass, brightness profiles and characteristic sizes of the BD population are inferred from continuum flux and modeling of the observations. Only one source is marginally resolved, allowing for the determination of its disk characteristic size. We conduct a demographic comparison between the properties of disks around BDs and stars in Lupus. Due to the small sample size, we cannot confirm or disprove if the disk mass over stellar mass ratio drops for BDs, as suggested for Ophiuchus. Nevertheless, we find that all detected BD disks have an estimated dust mass between 0.2 and 3.2  $M_{\oplus}$ ; these results suggest that the measured solid masses in BD disks can not explain the observed exoplanet population, analogous to earlier findings on disks around more massive stars. Combined with the low estimated accretion rates, and assuming that the mm-continuum emission is a reliable proxy for the total disk mass, we derive ratios of  $\dot{M}_{\text{acc}}/M_{\text{disk}}$  significantly lower than in disks around more massive stars. If confirmed with more accurate measurements of disk gas masses, this result could imply a qualitatively different relationship between disk masses and inward gas transport in BD disks.

**Key words.** Planets and satellites: formation – (Stars:) brown dwarfs – Stars: pre-main sequence

## 1. Introduction

Sub/mm wavelengths observations are particularly useful to study dust properties in protoplanetary disks, since the dust thermal emission of the outer disk, where the bulk of the dust mass resides, can be probed at these wavelengths (Testi et al. 2014; Andrews 2015). Demographic studies based on (sub)-mm-wavelengths surveys of the Class II population from nearby star-forming regions (Ansdell et al. 2016; Barenfeld et al. 2016; Pascucci et al. 2016; Cox et al. 2017; Cieza et al. 2018; Cazzoletti et al. 2019), have found positive correlations between various disk properties: disk mass with stellar mass ( $M_{\text{disk}}-M_{\star}$ , Andrews et al. 2013; Ansdell et al. 2016; Pascucci et al. 2016), disk size

with luminosity (Andrews et al. 2010; Tazzari et al. 2017; Tripathi et al. 2017; Andrews et al. 2018), and mass accretion rate onto the central star with the disk mass ( $\dot{M}_{\text{acc}}-M_{\text{disk}}$ , Manara et al. 2016b; Mulders et al. 2017; Rosotti et al. 2017).

These relations are poorly constrained in the brown dwarf (BD) and very low-mass (VLM) stars regime, since these surveys focused primarily on disks around more massive stars. Therefore, observations at (sub-)mm wavelengths targeting BDs and VLM stars are necessary in order to extend these demographic studies, and to investigate their formation mechanisms and ongoing physical processes in their disks.

General interest on BDs and VLM stars has increased substantially thanks to the recent exoplanet discoveries around VLM objects. The most thrilling cases are Trappist-1 (Gillon et al.

Send offprint requests to: E. Sanchis, e-mail: esanchis@eso.org

2017), a  $\sim 0.085 M_{\odot}$  VLM star that hosts 7 rocky planets in a packed orbital configuration, and Proxima B (Anglada-Escudé et al. 2016), an Earth-like planet found in our closest neighbor ( $M_{\star} = 0.12 M_{\odot}$ ), at a distance of only  $\sim 1.3$  pc from us. These and other discoveries (e.g., 2M1207b and 2M J044144b, Chauvin et al. 2004; Todorov et al. 2010) suggest that planets orbiting BDs and VLMs may be a common outcome of their formation.

The study of the early stages of BDs and VLM stars is crucial to understand the viability of planet formation around these objects and to determine the properties of the potential planetary systems that may form. In Klein et al. (2003), millimeter emission of dust from disks around BDs was detected for the first time. Like stars, BDs are often found surrounded by a protoplanetary disk in their early stages (Comeron et al. 1998; Natta & Testi 2001; Scholz 2008), where planet formation is expected to take place. The disk fraction for stellar and BD populations are found to be similar (Luhman 2012). Disk accretion (Jayawardhana et al. 2003; Scholz & Eislöffel 2004; Muzerolle et al. 2003, 2005) and outflows (Natta et al. 2004; Whelan et al. 2005) also occur in the early stages of BDs, analogous to those around more massive stars.

In this work we conducted a systematic survey of BD disks in the Lupus star-forming region, observing the full known sample of BD disks from a single region with the Atacama Large Millimeter/submillimeter Array (ALMA) in the same band for the first time. Previous ALMA observations of BD disks studied incomplete samples of the known BD population of other regions (Testi et al. 2016; van der Plas et al. 2016; Ward-Duong et al. 2018).

Dust disk masses, dust emission distribution profiles and dust disk characteristic sizes are determined from these observations. The last two properties are inferred from interferometric modeling of the dust disk emission. The characteristic size of the dusty disks is crucial to constrain the ongoing disk evolution processes (e.g., radial drift, grain growth). However, its determination is not straightforward. Firstly, the disk emission needs to be sufficiently resolved. For disks around BDs and VLM stars, this is only possible using state-of-art facilities, like ALMA, that allow for the high resolution and sensitivity required at these wavelengths. Besides, a general size definition is needed for a reliable comparison between observations and theoretical models. In this work we use the radius enclosing 68% of the object's emission distribution; this definition is representative of the physical size of the object (Tripathi et al. 2017), and independent of the model used to fit the observations. Another important disk property that can be derived from (sub-)mm observations is the disk mass. For the formation of rocky planets, the dust mass in disks should be larger than the mass of the resulting planets. However, comparing the mass derived from disk emission with the results from exoplanetary surveys, there is an apparent lack of material to produce the known planetary systems (Greaves & Rice 2010; Williams 2012; Najita & Kenyon 2014; Mulders et al. 2015; Pasucci et al. 2016; Testi et al. 2016; Manara et al. 2018).

The inferred disk properties of the young BD population are compared to the properties of disks around stars in the same region, with the aim of testing whether the known relations for stars hold for disks around BDs. In addition to the BD observations, other seven T Tauri star (TTS) disks are characterized and modeled in this work for the first time.

The work is organized as follows: the target selection is described in Section 2. A summary of the observations and the data processing can be found in Section 3. Section 4 provides a description of the modeling employed for the disk properties determination, together with the modeling results. The demographic

comparison of the inferred properties between BD and stellar disks, and the planet formation implications from the measured dust masses of the BD disks are discussed in Section 5, and the main conclusions of this work are included in Section 6.

## 2. Sample selection

The list of selected targets of the Lupus BD disks survey (Cycle 5; PI: L. Testi, Project ID: 2017.1.01243.S) encompasses all the known BDs in the Lupus region I-IV that were not observed previously with ALMA Band 7. Our population of BD disks in the Lupus star-forming region consists of all the known objects from the region census (Merín et al. 2008; Mužić et al. 2014, 2015) that show excess emission in at least two mid-infrared bands (Spitzer IRAC/MIPS), that have been spectroscopically classified as spectral type (SpT) M6 or later, and with estimated mass  $\leq 0.09 M_{\odot}$  (down to  $\sim 0.02 M_{\odot}$ ). Eleven sources in Lupus satisfied these selection criteria, seven were the targets for the new observations; the remaining four had been already observed in the Lupus disks survey (Cycle 2; PI: J. Williams, Project ID: 2013.1.00220.S). Based on radial velocity analysis and X-shooter spectra, two sources in the sample, IRAS 15567-4141 and SSTc2d J160034.4-422540, have been recently excluded from being Lupus members, and are likely background giants (Frasca et al. 2017; Alcalá et al. 2017); in agreement with the poorly constrained parallaxes from Gaia DR2 (Gaia Collaboration et al. 2018). Therefore, these will not be discussed further in this paper; only five new targets are discussed. All the studied BDs are isolated systems, except SONYC-Lup3-7, which might form a very wide ( $\sim 7''$ ) binary system with SONYC-Lup3-6, but this last object has no confirmed membership to the region (Mužić et al. 2014).

In Table 1, we list all the sources analyzed in this work: the known BD population (5 objects from the new observations and 4 from previous observations), together with the 7 disks around stars observed in the Lupus completion survey (Cycle 5; PI: S.E. van Terwisga, Project ID: 2016.1.01239.S). The last two objects in the table are those that were observed but later excluded from the Lupus census (Frasca et al. 2017; Alcalá et al. 2017). The names, sky position and main stellar properties of the central stars are included in the Table 1. The stellar properties shown in the table (SpT, effective temperature  $T_{\text{eff}}$ , extinction in V-band  $A_V$ , stellar luminosity  $L_{\star}$  and  $M_{\star}$ ) were reported by Alcalá et al. (2014, 2017) and Mužić et al. (2014). The methodology for the stellar luminosity derivation between these studies differs, nevertheless the agreement between the two methods is very good, as shown in Manara et al. (2016a).  $L_{\star}$  have been adjusted accounting for updated distances by Gaia DR2 (distance estimated as the inverse of the parallax, Gaia Collaboration et al. 2018). The stellar mass is derived from the pre-main sequence (MS) evolutionary models of Baraffe et al. (2015), estimated from the position in the Hertzsprung–Russell (HR) diagram. For objects with estimated mass  $> 1.4 M_{\odot}$  and objects laying above the 1 Myr isochrone, the tracks from Siess et al. (2000) are used instead. Stellar mass uncertainties are computed with a Monte Carlo approach (as described in Alcalá et al. 2017), which takes into account the associated uncertainties of the stellar properties  $L_{\star}$  and  $T_{\text{eff}}$  used to infer the mass.

The BD disk population is compared to the sample of young stellar objects (YSOs) in Lupus that have a protoplanetary disk and estimated stellar mass  $> 0.09 M_{\odot}$ . Thanks to the inclusion of the 7 stellar disks from the Lupus completion survey, we have the largest sample of stellar disks in the Lupus clouds (I-IV) observed with ALMA in Band 7. The stellar disk population

**Table 1.** Protoplanetary disks from Lupus modeled in this study.

Object	$\alpha$ (J2000)	$\delta$ (J2000)	Lupus cloud	Distance [pc]	SpT	$T_{\text{eff}}$ [K]	$A_V$ [mag]	$L_\star$ [ $L_\odot$ ]	$M_\star$ [ $M_\odot$ ]	Notes
<i>BDs from this survey:</i>										
J154518.5-342125	15:45:18.53	-34:21:24.8	I	$152 \pm 4$	M6.5	2935	0.0	0.04	$0.09 \pm 0.02$	1
SONYC-Lup3-7	16:08:59.53	-38:56:27.6	III	$150 \pm 6$	M8.5	2600	0.0	0.01	$0.02 \pm 0.01$	1, 4
Lup706	16:08:37.30	-39:23:10.8	III	158.5	M7.5	2795	0.0	0.002	$0.05 \pm 0.01$	1, 3, 5
AKC2006-18	15:41:40.82	-33:45:19.0	I	$149 \pm 8$	M6.5	2935	0.0	0.01	$0.07 \pm 0.02$	1
SONYC-Lup3-10	16:09:13.43	-38:58:04.9	III	158.5	M8.8	2650	1.5	0.003	$0.03 \pm 0.01$	2, 3
<i>BDs from Lupus disks survey (Ansdell et al. 2016):</i>										
Lup818s	16:09:56.29	-38:59:51.7	III	$157 \pm 3$	M6	2990	0.0	0.02	$0.09 \pm 0.02$	1
J161019.8-383607	16:10:19.84	-38:36:06.8	III	$159 \pm 3$	M6.5	2935	0.0	0.04	$0.09 \pm 0.02$	1
J160855.3-384848	16:08:55.29	-38:48:48.1	III	$158 \pm 3$	M6.5	2935	0.0	0.05	$0.09 \pm 0.02$	1
Lup607	16:08:28.10	-39:13:10.0	III	$175 \pm 6$	M6.5	2935	0.0	0.05	$0.10 \pm 0.02$	1
<i>Disks from Lupus completion survey:</i>										
Sz102	16:08:29.71	-39:03:11.0	III	158.5	K2	4900	0.7	0.01	-	1, 3, 5, 6
V1094 Sco	16:08:36.18	-39:23:02.5	III	$154 \pm 1$	K6	4205	1.7	1.15	$0.86 \pm 0.18$	1
GQ Lup	15:49:12.10	-35:39:05.1	I	$152 \pm 1$	K6	4205	0.7	1.48	$0.85 \pm 0.17$	1
Sz76	15:49:30.74	-35:49:51.4	I	$160 \pm 1$	M4	3270	0.2	0.18	$0.23 \pm 0.04$	1
Sz77	15:51:46.96	-35:56:44.1	I	$155 \pm 1$	K7	4060	0.0	0.59	$0.75 \pm 0.15$	1
RXJ1556.1-3655	15:56:02.10	-36:55:28.3	II	$158 \pm 1$	M1	3705	1.0	0.26	$0.5 \pm 0.14$	1
EX Lup	16:03:05.49	-40:18:25.4	III	$158 \pm 1$	M0	3850	1.1	0.76	$0.56 \pm 0.13$	1
<i>Observed objects rejected from being members of Lupus:</i>										
IRAS 15567-4141	16:00:07.42	-41:49:48.4	-	-	-	-	-	-	-	7
J160034.4-422540	16:00:34.40	-42:25:38.6	-	-	-	-	-	-	-	7

**Notes.**


- (1) Stellar properties from Alcalá et al. (2014, 2017) and adjusted to the new Gaia DR2 parallaxes.
- (2) Stellar properties from Mužić et al. (2014), and adjusted to the Gaia DR2 parallaxes.
- (3) Gaia parallax unknown, mean distance of Lupus region considered.
- (4) Two sets of stellar properties (Alcalá et al. 2017; Mužić et al. 2014).
- (5) Sub-luminous object (see Alcalá et al. 2014, 2017).
- (6) No estimation of the stellar mass since it falls below the zero-age main sequence in the HR diagram (see Alcalá et al. 2017).
- (7) Categorized as background sources (Frasca et al. 2017; Alcalá et al. 2017).

is assembled from different census of the region from 2MASS, Spitzer, and Herschel surveys (Hughes et al. 1994; Comerón 2008; Merín et al. 2008; Mortier et al. 2011; Dunham et al. 2015; Bustamante et al. 2015). 82 of these disks were observed with ALMA in Band 7 in the original Lupus disks survey (ALMA Cycle 2; PI: J. Williams), IM Lup and Sz 91 were observed separately (Cycle 2; PI: I. Cleeves, Project ID: 2013.1.00226.S and Cycle 2; PI: H. Canovas, Project ID: 2013.1.00663.S), and the seven remaining belong to the Lupus completion survey (Cycle 5; PI: S.E. van Terwisga).

Therefore, the BD and stellar disks samples of our demographic study consist of 9 and 91 sources respectively. The HR-diagram for the studied disk population is shown in Figure 1, using the stellar properties from previous studies as described before. The BD disks are marked in red, and the stellar disk population in blue. Sub-luminous sources, those with luminosities lower than those expected for YSOs of age  $\sim 3$  Myr (likely due to gray obscuration Alcalá et al. 2014, 2017), are illustrated with square symbols. From the re-adjusted  $L_\star$  using the more accurate Gaia DR2 parallaxes, 2 objects (J16081497-3857145 and J16085373-3914367) are now added to the list of sub-luminous objects of the region.

### 3. Observations

ALMA observed our targets on 2018 April 1 and April 2 with 44, and 42 12 m antennas respectively. The baselines ranged between 15.1 and 704.1 m for the array configuration of the first day, and between 15.1 and 629.2 m for the configuration of the second day. Four spectral windows were set for the continuum



**Fig. 1.** Hertzsprung–Russell diagram for the studied BD and stellar populations from Lupus. The stellar luminosity is taken from the literature and re-adjusted to the new distance associated to the parallaxes from Gaia DR2 (Gaia Collaboration et al. 2018). The BD population is shown in red, and the stellar population is indicated in blue. The pre-MS tracks of Baraffe et al. (2015) are overlaid in the figure. Objects with luminosities that would correspond to older ages than expected are considered sub-luminous and marked as squares. A number of points lay on top of each other (e.g., 4 BDs near the 1 Myr and the  $0.1 M_\odot$  lines.)

observation of the targets, centered at 334.432, 336.321, 345.889 and 347.821 GHz and bandwidths of 2, 1.875, 1.875 and 0.938 GHz respectively (total receivers bandwidth of  $\sim 6.688$  GHz).

The calibrators for the observations were J1517-2422 for flux and passband, and J1610-3958 for the complex gain calibration, the same in both executions. The flux density scale accuracy is expected to be of 10% for observations of the Lupus BD disks survey. 12 scans of 60-62 seconds duration each were performed for every target, for a total integration time of more than 12 minutes per source.

The CASA 5.3.0 software has been used for the interferometric visibilities calibration and imaging. The continuum maps are produced using the channels free from spectral line emission, with Briggs weighting of the visibilities ( $-1.0$  and  $+0.5$  robustness for resolved and unresolved objects respectively). None of the sources are bright enough to perform self-calibration. The full width at half maximum (FWHM) of the synthesized beam is  $0.27'' \times 0.24''$  for robust parameter =  $-1$ , and  $0.36'' \times 0.33''$  for robustness =  $+0.5$ , with average position angle (PA) of  $28^\circ$ . The continuum maps of the 5 BD disks observed are shown in Figure 2. The sensitivity for the BD disks survey is improved by a factor of  $\sim 3$  with respect to the previous Lupus disks surveys, this allowed us to detect fainter emission. Three BD disks are detected, J154518.5-342125, SONYC-Lup3-7 and Lup706, with respective signal-to-noise (S/N) of 42, 8 and 7. Emission is not detected from the two other disks (AKC2006-18 and SONYC-Lup3-10) nor from the two background objects (IRAS 15567-4141 and J160034.4-422540).

The main results of the observations are reported in Table 2. This table includes the total disk flux, peak intensity and the rms of the image. These values from the observations are obtained using identical methodology to the results presented in Ansdell et al. (2016) for the original Lupus disks survey. The continuum flux is inferred from the *uvmodelfit* task in CASA: emission is fitted with an elliptical gaussian in case that resulting FWHM along the major axis from the fit is at least five times its uncertainty, otherwise the emission is fitted as a point source. For sources with resolved structure, the flux is obtained from a curve of growth method with increasing circular apertures, centered at the peak emission of the object. The rms is computed from a  $4''\text{-}9''$  radius annulus centered on the detected emission, or on the expected source position if no disk emission is detected. The flux upper-limits of the non-detected BD disks are displayed in the  $F_{\text{cont}}$  column of the table. In this work, upper-limits of non-detected BD and stellar disks are computed as three times the rms level above the measured flux within the beam size are (centered at the expected source position), this corresponds to a 99.87% confidence level. This differs slightly from the upper-limits in Ansdell et al. (2016), considered to be  $3\sigma$ .

We have also derived the flux densities and image rms for the sources observed with ALMA in Band 7 in the Lupus completion survey (see also van Terwisga et al. 2018, 2019). For these sources we have followed the same procedure as for the BDs disks. As reported by van Terwisga et al. (2018), the flux calibrator used for these disks in Band 7 was highly variable, thus the absolute flux densities are very uncertain. To alleviate this problem, van Terwisga et al. (2018) compared the flux of one of these sources (GQ Lup) to a previous observation with high S/N and reliable flux calibration (MacGregor et al. 2017), obtaining a fluxes ratio of  $1.3 \pm 0.009$  between both observations. Since GQ Lup and the rest of the disks of the Lupus completion survey were observed at the same day, we applied that factor to the measured fluxes of all these sources. The results are included in Table 2. For the observations results from the Lupus disks survey (ALMA Project ID: 2013.1.00220.S), we refer to Ansdell et al. (2016).

**Table 2.** Inferred fluxes for all the studied disks.

Object	$F_{\text{cont}}$ [mJy]	$I_{\text{peak}}$ [mJy/beam]	rms [mJy/beam]
<i>BDs from this survey:</i>			
J154518.5-342125	5.65	5.18	0.12
SONYC-Lup3-7	0.52	0.62	0.07
Lup706	0.79	0.71	0.10
AKC2006-18	< 0.23	-	0.08
SONYC-Lup3-10	< 0.24	-	0.07
<i>BDs from Lupus disks survey:</i>			
Lup818s	7.44	7.44	0.24
J161019.8-383607	< 1.24	-	0.23
J160855.3-384848	1.81	1.81	0.26
Lup607	< 0.95	-	0.24
<i>Disks from Lupus disks completion survey:</i>			
Sz102	14.69	7.81	0.34
V1094 Sco	553.17	23.43	0.37
GQ Lup	78.00	42.36	0.34
Sz76	12.00	4.88	0.35
Sz77	4.88	3.88	0.20
RXJ1556.1-3655	57.52	16.70	0.35
EX Lup	41.78	9.80	0.35

**Notes.** From top to bottom: the 9 protoplanetary disks around BDs (5 from the new Lupus BDs disks survey) and 7 additional disks from the Lupus disks completion survey. The columns are the total disk flux ( $F_{\text{cont}}$ ), peak intensity ( $I_{\text{peak}}$ ) of each object and the rms of their continuum maps.


#### 4. Modeling

Previous work characterizing interferometric observations of protoplanetary disks modeled the continuum emission with either physical or empirical models. A physical model commonly used is the two-layer approximation model (Ricci et al. 2014; Testi et al. 2016; Tazzari et al. 2017). Although this model can successfully describe the spectral energy distribution (SED) of TTS disks (Chiang & Goldreich 1997; Dullemond et al. 2001), the model assumes a simplified physical structure of the disk. In order to provide an observational characterization of the emission, we prefer to fit empirical analytical functions to the emission profile, to allow a more straightforward comparison of the disk properties.

We model the extended disks around stars with the Nuker profile, used in Andrews et al. (2018) to characterize the Lupus disks observed with ALMA in Band 7 from Ansdell et al. (2016). By using the Nuker profile we ensure the homogeneity on the characterization of the Lupus disks sample, which is a key aspect of the demographic study discussed in the next section. We follow the modeling described in Tripathi et al. (2017), using the Lauer et al. (1995) formulation:

$$I_\nu(\rho) = I_0 \cdot \left( \frac{\rho}{\rho_t} \right)^{-\gamma} \cdot \left[ 1 + \left( \frac{\rho}{\rho_t} \right)^\alpha \right]^{(\gamma-\beta)/\alpha}, \quad (1)$$

where  $\rho_t$  is the transition radius that sets the boundary between the inner and outer regimes of the radial profile,  $\gamma$  and  $\beta$  are the inner and outer disk slopes and  $\alpha$  is a factor that determines the smoothness of the transition between both regimes. The disk is assumed to be azimuthally symmetric. The total number of parameters used to model the extended disks with the Nuker profile are 9:  $\rho_t$ ,  $\gamma$ ,  $\beta$ ,  $\alpha$ , the total disk flux density  $F_{\text{tot}}$ , and four additional geometrical parameters connected to the observation: inclination of the disk in the sky towards the observer ( $i$ ,  $0^\circ$  face-on



**Fig. 2.** Dust continuum images at  $890\,\mu\text{m}$  of the Lupus BDs disks survey from ALMA Band 7 observations. The beam size FWHM is  $0.27'' \times 0.24''$  for the J154518.5-342125 map (robust parameter of  $-1$ ), and  $0.36'' \times 0.33''$  for the rest of the maps (robustness =  $+0.5$ ). The average beam position angle is  $PA = 28^\circ$ . The contours are drawn at increasing (or decreasing)  $3\sigma$  intervals as solid (dashed) lines.

disk,  $90^\circ$  edge-on), the position angle in the sky plane ( $PA$ , defined east of north), and right ascension and declination off-sets to the phase center of the observations ( $\Delta RA$  and  $\Delta Dec$ ).

A simple parametrized gaussian function has been used to fit the BD disks, to reduce the number of free parameters, since the emission maps of these objects are extremely compact; only one BD disk is marginally resolved. This function can be used to model moderate-resolution observations to characterize the disk brightness profile and its size, as shown in Appendix A (see Figure A.1) where the Gaussian profile together with other models were used to fit the disk emission around RXJ1556.1-3655; the resulting profiles are alike and the values of the chosen size definition are indistinguishable. The expression of the gaussian function used to model the BDs surface brightness profile is:

$$I_\nu(\rho) = I_0 \cdot \exp \left[ -0.5 \cdot \left( \frac{\rho}{\sigma} \right)^2 \right], \quad (2)$$

where  $\rho$  is the projected radius in the sky in arcsec,  $I_0$  is a normalization factor, and  $\sigma$  is the standard deviation of the gaussian profile. The disk is assumed to be azimuthally symmetric. The free parameters to model the BD disks are 6 in total, two from the gaussian model ( $I_0$  and  $\sigma$ ), together with the observational parameters analogous to the Nuker model ( $i$ ,  $PA$ ,  $\Delta RA$  and  $\Delta Dec$ ).

To perform the fits, the `Galarion` (Tazzari et al. 2018) and `emcee` (Foreman-Mackey et al. 2013) packages have been used. For a detailed description of the methodology we refer to Tazzari et al. (2016, 2017). To run the affine invariant Markov chain Monte Carlo (MCMC) Goodman & Weare (2010), 200 walkers have been used, in order to investigate the parameter space for each disk ( $\gtrsim 20$ -30 walkers for each parameter); the computation ran for 20000 steps per walker, this guaranteed convergence in all fitted disks.

#### 4.1. Size definition

An appropriate definition of the disk size is necessary for a proper characterization of disks from observations, and for its comparison to theoretical models. An approach commonly used is to extract the radius from the models used to fit the emission. The transition radius  $\rho_t$  from the Nuker profile definition (Equation 1) provides misleading information of the disk size, since systems with very different architectures and extents might have similar values of  $\rho_t$ . When fitting to a power-law with an exponential cut-off, an analogous problem arises if the cut-off radius  $\rho_c$  from Guilloteau et al. (2011) parametrization is used. The outer radius  $R_{out}$  was used in Ricci et al. (2014) and in Testi et al. (2016) to fit BDs disks observations, defined as the outermost radius of their modeled surface density.

All these definitions may be useful in specific studies, but are not suitable for a general characterization of disk sizes from their emission. A more convenient size definition is the radius enclosing a certain fraction of the total disk emission. This definition with a fraction of 68% of the total disk emission has been used in many recent works (e.g., Tripathi et al. 2017; Andrews et al. 2018; Facchini et al. 2019; Long et al. 2019; Manara et al. 2019), in Tazzari et al. (2017) the 95% of the total disk emission was used. To avoid confusion with the different terminology used in the literature, we simply refer to them as 68% ( $R_{68\%}$ ) and 95% ( $R_{95\%}$ ) flux radii. In the nomenclature along this work,  $R$  refers to the radius in the system reference frame (typically in au), while  $\rho$  stands for the projected radius in the sky plane in arcsec.

We tested both  $R_{68\%}$  and  $R_{95\%}$  to determine the quality of each radius as the characteristic size for the disk emission (details in Appendix A). For this test we fitted the same disk to various models and inferred  $R_{68\%}$  and  $R_{95\%}$  for each model. This test shows that the dispersion on  $R_{68\%}$  is much smaller than for  $R_{95\%}$ . Thus, we consider  $R_{68\%}$  as the most reliable size definition

for our sample. Nevertheless, in the modeling results of the disks (Table 3), we include both  $R_{68\%}$  and  $R_{95\%}$  for completeness.

Additionally, we fitted several disks that were previously modeled in Andrews et al. (2018), in order to test the proper functioning of our modeling tool. This additional test is also included in Appendix A; the resulting  $\rho_{68\%}$  from this work and from Andrews et al. (2018) are in very good agreement.

#### 4.2. Dust disk masses

To ease the comparison with the existing surveys of disks around stars in the Lupus clouds (Ansdell et al. 2016), we provide an estimate of the disk dust masses using the simplifying assumption of optically thin emission and using an average temperature of 20 K. We note that these assumptions may lead to underestimating the disk mass in case the emission is optically thick or the average temperature is lower than the assumed value.


For the dust mass determination, assumptions on the dust temperature and opacity are needed. A dependence of the dust temperature with the stellar luminosity was first proposed from theoretical grounds (e.g., Yorke et al. 1993; Sonnhalter et al. 1995). More recently, from radiative transfer modeling of mm observations, Andrews et al. (2013) proposed a single-valued mean temperature for each disk that could be used to estimate the disk mass for objects with  $L_* \in [0.1, 100] L_\odot$ . Soon after, van der Plas et al. (2016) suggested a more flattened relation for VLM objects. In Ballering & Eisner (2019), the correlation of disk temperature with the stellar luminosity was derived using simplistic radiative transfer models from SED fitting of Taurus disks. Daemgen et al. (2016) and Tazzari et al. (2017) showed that, depending on assumptions on disk size and vertical structure, similar  $T_{\text{dust}}$  even with very different luminosities are compatible with the data. From the different studies, it is unclear whether there is a simple and general relation between dust temperature and stellar properties. Using additional relations of the temperature with other stellar properties as the proposed by Andrews et al. (2013); van der Plas et al. (2016) might introduce spurious results in our analysis, or even erase possible relations between different disk properties.

We therefore compute the dust mass of each BD and stellar disk assuming a constant dust opacity of  $\kappa_{890\mu\text{m}} = 2 \text{ cm}^2 \text{ g}^{-1}$ , following previous ALMA Band 7 observations for VLMs and BDs (Ricci et al. 2014; Testi et al. 2016), and an averaged dust temperature of  $T_{\text{dust}} = 20 \text{ K}$ , as in Pascucci et al. (2016); Ansdell et al. (2016, 2018). This value is the median temperature for protoplanetary disks in the Taurus region (Andrews & Williams 2005). Nevertheless, in Appendix B we show the main demographic results of this work using the dust temperature and stellar luminosity relations from Andrews et al. (2013); van der Plas et al. (2016).

#### 4.3. Modeling results

The results of the modeling and the derived properties of dust mass and disk size are presented in this section. The results for J154518.5-342125 are shown in Figures 3, 4, 5, 7, while the results for the remaining disks which fits converged are in Appendix C.

In Figure 3 we show the real and imaginary part of the observed and modeled visibilities as a function of baseline. The visibilities were first centered using  $\Delta\text{RA}$  and  $\Delta\text{Dec}$  from the model with lowest  $\chi^2$ , then de-projected taking  $i$  and  $PA$  (for a detailed description, see Tazzari et al. 2017, 2018).




**Fig. 3.** Observed and model visibilities of J154518.5-342125, plotted as real and imaginary part as a function of the baseline (in  $k\lambda$ ). The data from the observations are plotted as black data points with error bars, the model with the lowest  $\chi^2$  is shown as solid red curve, and a random set of models from the parameter space investigation are drawn as gray curves. Figure made with the `uvplot` Python package Tazzari (2017).


The posterior probability distribution functions (PDFs) of the free parameters (of the gaussian model, or from the Nuker profile), are shown in the top sub-panels of Figure 4. Vertical dashed lines represent the median values of each histogram, used as the best fitted parameter value, and the 16th and 84th percentiles, used to infer the lower and upper values of the uncertainties. From the figure,  $I_0$ ,  $\sigma$  and the sky plane off-sets are well determined, while the inclination and position angle are both degenerated, and their values are only poorly constrained. The rest of the sub-panels in the figure show the 2D histograms for the pairs of parameters, which indicate possible correlations between the different parameters. In a few disks of the Lupus completion survey, the smoothness parameter ( $\alpha$ ), and/or the inner and outer slopes ( $\beta$ ,  $\gamma$ ) of the Nuker profile can not be appropriately constrained from moderate resolution observations (see corner plots of the fits in Appendix C). This limitation does not affect the characterization of their disk sizes.

The modeled emission distribution of the disk is shown in the top panel of Figure 5. On the bottom panel, we show the normalized cumulative flux  $f_{\text{cumul}}$ , derived from Equation A.1 for the respective models of the top panel. In both plots, the inferred values of  $R_{68\%}$  and  $R_{95\%}$  radii are included as vertical dashed and dotted lines. These radii are computed for each model of the MCMC; the final values of  $R_{68\%}$  and  $R_{95\%}$  are the median of their respective PDFs, with upper and lower errors as the median  $\pm 1\sigma$  (example of a  $R_{68\%}$  PDF in Figure 6).


In Figure 7, we show the reconstructed continuum emission of the source in the sky plane: in the left panel the observed emission, in the central panel the reconstructed emission corresponding to the model with lowest  $\chi^2$ , and in the right panel the residuals. Residuals are at noise level on all fitted disks, indicating that the model represents the observation faithfully.



**Fig. 4.** 1D and 2D histograms of the free parameters used to model J154518.5-342125 ALMA observations, resulting from the MCMC analysis. The marginalized PDFs of the parameters are plotted in the top panels, including the 16th, 50th and 84th percentiles as vertical dashed lines. The  $I_0$  and  $\sigma$  parameters are connected to the gaussian model used, as defined in Equation 2;  $i$ ,  $PA$ ,  $\Delta RA$  and  $\Delta Dec$  are geometrical parameters linked to the observations.



**Fig. 5.** Radial brightness profile (top panel) and the associated cumulative flux (bottom) for J154518.5-342125 disk, resulting from the gaussian model used to fit the observed visibilities. The lowest  $\chi^2$  model and a random subset of models are drawn in both panels as a red and thin gray curves respectively.



**Fig. 6.** PDF of the  $R_{68\%}$  radius for the disk around J154518.5-342125.  $R_{68\%}$  is computed for each model of the parameter space investigation, the value of the radius is taken as the median of the PDF, while upper/lower error are the median  $\pm 1\sigma$  of the distribution. The three values are represented as vertical dashed lines.

The results of the free parameters from the fits and the derived disk properties can be found in Table 3. The values shown are the median of their respective PDF, lower (upper) uncertainties are obtained from the range between median and 16th (84th) percentiles of their posterior distribution. The missing values are for the cases in which the fit did not find a clear convergence. A note on the source Sz 102, from our fit results, gray obscuration due to its inclination ( $\sim 60^\circ$ ) would not explain its sub-luminous nature. Strong episodic accretion as suggested by Baraffe & Chabrier (2010) could explain its luminosity: this effect reduces the radius of the star, increasing its temperature and resulting in different pre-MS path and a relatively lower luminosity. Another viable explanation would be a misalignment of the inner and outer disk.

#### 4.3.1. Disk size results

The radii enclosing 68% and 95% of the total flux ( $R_{68\%}$  and  $R_{95\%}$ ) are specified in Table 3. The values are inferred from their respective PDF (example in Figure 6), as derived from the model parameters results. The radii of the detected BDs disks are unfortunately poorly determined, due to the compactness of the sources, combined with a low S/N of their continuum emission at this waveband. Only for J154518.5-342125 we can properly quantify its size, since its continuum emission is detected with enough S/N and is marginally resolved. We denote as marginally resolved disk the case where the emission is of similar spatial scale as the beam size in the image plane, but where the observed visibilities can be fitted by a gaussian function with well constrained  $\sigma$ . For all the other BD sources we provide upper limits of their sizes as 95% confidence level, inferred from the PDF of  $R_{68\%}$  and  $R_{95\%}$ . On the other hand, we have determined the emission distribution size for six out of seven disks of the Lupus completion survey; for the remaining one (Sz77) we provide an upper limit.

Previous ALMA observations of BD disks in other regions showed that most of the objects were too compact to be resolved (van der Plas et al. 2016; Testi et al. 2016). Using different methodology to define and derive disk radii, Ricci et al. (2014) and Testi et al. (2016) showed that some BD disk radii ( $R$ ) in Taurus may extend beyond  $R \gtrsim 80$  au, while in  $\rho$ -Oph BD disks seem to all have  $R \lesssim 25$  au.


Nevertheless, we should bear in mind that the few BD disks with well determined sizes are among the brightest and most massive of the BD population of their respective regions (Lu-

**Table 3.** Results from the modeling of the studied disks, together with inferred sizes ( $R_{68\%}$  and  $R_{95\%}$ , in au), and total disk dust mass ( $M_{\text{dust}}$ ).

Object	$\log_{10}(I_0)$ [Jy/sr]	$\sigma$ [arcsec]	$i$ [deg]	$PA$ [deg]	$\Delta RA$ [arcsec]	$\Delta Dec$ [arcsec]	$R_{68\%}$ [au]	$R_{95\%}$ [au]	$M_{\text{dust}}$ [ $M_\oplus$ ]
<i>BDS from this survey:</i>									
<i>BDS from Lupus disks survey:</i>									
J154518.5-342125	$10.56^{+0.16}_{-0.11}$	$0.035^{+0.005}_{-0.005}$	$25.8^{+18.7}_{-17.0}$	$90.2^{+54.7}_{-59.8}$	$-0.270^{+0.002}_{-0.002}$	$-0.352^{+0.002}_{-0.002}$	$8.2^{+1.0}_{-1.2}$	$13.2^{+1.7}_{-2.0}$	$2.3 \pm 0.2$
SONYC-Lup3-7	$9.84^{+0.55}_{-0.49}$	$0.03^{+0.02}_{-0.01}$	-	-	$-0.07^{+0.02}_{-0.02}$	$-0.24^{+0.02}_{-0.02}$	$<17.5$	$<28.3$	$0.21 \pm 0.02$
Lup706	$8.70^{+0.17}_{-0.15}$	$0.14^{+0.04}_{-0.03}$	-	$88.3^{+33.9}_{-38.5}$	$-0.12^{+0.02}_{-0.02}$	$-0.46^{+0.02}_{-0.02}$	$<56.4$	$<91.4$	$0.35 \pm 0.04$
AKC2006-18	-	-	-	-	-	-	-	-	$<0.06$
SONYC-Lup3-10	-	-	-	-	-	-	-	-	$<0.08$
<i>BDS from Lupus disks survey:</i>									
Lup818s	$10.99^{+0.41}_{-0.36}$	$0.03^{+0.01}_{-0.01}$	-	-	$-0.06^{+0.01}_{-0.01}$	$-0.207^{+0.005}_{-0.005}$	$<12.1$	$<19.7$	$3.2 \pm 0.3$
J161019.8-383607	-	-	-	-	-	-	-	-	$<0.45$
J160855.3-384848	$10.15^{+0.59}_{-0.47}$	$0.04^{+0.03}_{-0.02}$	-	-	$-0.16^{+0.02}_{-0.02}$	$-0.43^{+0.02}_{-0.02}$	$<21.7$	$<35.2$	$0.79 \pm 0.08$
Lup607	-	-	-	-	-	-	-	-	$<0.38$
Object	$\rho_i$ [arcsec]	$\gamma$ [-]	$\beta$ [-]	$\log_{10}\alpha$ [-]	$F_{\text{tot}}$ [mJy]	$i$ [deg]	$PA$ [deg]	$\Delta RA$ [arcsec]	$\Delta Dec$ [arcsec]
<i>Disk from Lupus disks completion survey:</i>									
Sz102	$0.55^{+0.45}_{-0.26}$	$-1.2^{+1.2}_{-1.2}$	$15.2^{+3.3}_{-4.0}$	$-0.1^{+0.2}_{-0.2}$	$27.6^{+2.4}_{-2.9}$	$57.6^{+2.9}_{-2.9}$	$14.0^{+3.0}_{-2.9}$	$-0.134^{+0.001}_{-0.002}$	$20.1^{+1.1}_{-1.0}$
V1094 Sco	$0.18^{+0.01}_{-0.01}$	$0.38^{+0.03}_{-0.03}$	$1.25^{+0.03}_{-0.01}$	$1.7^{+0.2}_{-0.3}$	$1038^{+5}_{-5}$	$56.2^{+0.3}_{-0.3}$	$110.9^{+0.2}_{-0.2}$	$-0.172^{+0.001}_{-0.001}$	$201.5^{+0.4}_{-0.4}$
GQ Lup	$0.21^{+0.04}_{-0.04}$	$0.2^{+0.3}_{-0.3}$	$0.5^{+0.2}_{-0.1}$	$158.7^{+2.5}_{-2.5}$	$60.6^{+0.5}_{-0.5}$	$-11.6^{+0.5}_{-0.5}$	$0.0301^{+0.0003}_{-0.0002}$	$0.1284^{+0.0004}_{-0.0003}$	$29.2^{+0.8}_{-0.8}$
Sz76	$0.45^{+0.12}_{-0.05}$	$1.38^{+0.04}_{-0.05}$	$1.26^{+0.5}_{-0.5}$	$1.72^{+2.4}_{-2.4}$	$38.9^{+7.6}_{-7.6}$	$113^{+11}_{-11}$	$0.001^{+0.003}_{-0.003}$	$0.003^{+0.003}_{-0.003}$	$40.9^{+0.2}_{-0.2}$
Sz77	-	-	-	$1.26^{+0.5}_{-0.5}$	$1.72^{+1.9}_{-1.9}$	$11.1^{+14}_{-14}$	$0.001^{+0.003}_{-0.003}$	$0.003^{+0.003}_{-0.003}$	$75^{+17}_{-8}$
RXJ1556.1-36555	$0.30^{+0.03}_{-0.04}$	$0.4^{+0.1}_{-0.1}$	$15.9^{+2.8}_{-3.8}$	$0.6^{+0.1}_{-0.1}$	$85.5^{+1.3}_{-1.3}$	$49.4^{+0.7}_{-0.7}$	$61.1^{+10.3}_{-10.3}$	$0.008^{+0.004}_{-0.004}$	$<55.5$
EX Lup	$0.23^{+0.01}_{-0.01}$	$0.38^{+0.05}_{-0.06}$	$14.4^{+3.9}_{-3.5}$	$1.4^{+0.4}_{-0.3}$	$50.0^{+0.9}_{-0.9}$	$30.8^{+1.5}_{-1.6}$	$69.2^{+2.9}_{-2.9}$	$-0.061^{+0.001}_{-0.001}$	$24.3 \pm 2.4$

**Notes.**

The first 9 objects are the full list of known BDs in Lupus, fitted to a gaussian model. The six free parameters are: normalization factor of the emission profile  $\log I_0$ , standard deviation of the gaussian profile  $\sigma$ , inclination  $i$ , position angle  $PA$ , and right ascension and declination offsets to the phase center of the observations  $\Delta RA$  and  $\Delta Dec$ . The last 7 disks were fitted to a Nuker profile. The nine free parameters of these fits are: transition radius  $\rho_i$ , inner and outer slopes  $\gamma$  and  $\beta$ , smoothness parameter  $\alpha$ , total disk flux density  $F_{\text{tot}}$ , and the geometrical parameters of the observation  $i$ ,  $PA$ ,  $\Delta RA$  and  $\Delta Dec$ .



**Fig. 7.** Observed (left), model (center) and respective residuals (right panel) for the continuum emission of J154518.5-342125 observed with ALMA. The modeled emission map is reconstructed from the synthetic visibilities with the lowest  $\chi^2$  from the interferometric modeling. The contours are drawn at increasing (or decreasing)  $3\sigma$  intervals as solid (dashed) lines.

pus, Taurus and Ophiuchus); likely they are not representative of the BD population.

#### 4.4. Total dust mass results

The total disk dust mass (last column in Table 3) is computed from the assumptions detailed in Section 4.2. As all detections have good S/N (Table 2), the main uncertainty when comparing samples observed at different times is the flux calibrator uncertainty ( $\sim 10\%$ , see Section 3). Dust mass upper limits for non-detected disks around BDs and stars are obtained from the respective continuum flux upper limits as described in Section 3.

The total dust mass for the detected BD disks range between 0.2 and  $3.2 M_{\oplus}$ . This means that our sources are within the lightest protoplanetary disks known up to date. In particular, SONYC-Lup3-7 is the BD disk with the lowest dust mass estimate, independent of the prescription used for the dust disk mass determination. Comparing our dust mass results of BD disks in Lupus to the results of BD disks in other regions, our results are found to be similar. In Testi et al. (2016), a sample of 17 BD disks in the  $\rho$  Ophiuchus region were observed and their dust mass estimates are within  $0.5$  and  $6.3 M_{\oplus}$ , with the same assumptions of dust temperature and opacity as for our results. The dust masses of Taurus disks around BD and VLMs (Ward-Duong et al. 2018) range between  $\sim 0.25$  and  $\sim 16.7 M_{\oplus}$ , using the same temperature and opacity values as in this work.

## 5. Discussion

### 5.1. Comparison to disks around T Tauri stars

We have performed a demographic comparison between the BD and stellar disk populations of Lupus. For this analysis we use the derived disk properties to test whether the known relations for disks around stars stand for BD disks. The observational dataset of both populations has been obtained from the same facility (ALMA, Band 7), and the derivation of the properties has been conducted with homogeneous methodology for the entire disk population. In this manner we eradicate systematic errors due to the mixing of diverse datasets handled with different methods.

In addition, we have updated the relations between disk properties in Lupus with the largest sample of disks in the region

observed with ALMA in Band 7, thanks to the incorporation of the seven stellar disks from the Lupus completion survey to the stellar population. In Table 4 we summarize all the correlations discussed along this section.

#### 5.1.1. Correlation between $M_{\star}$ and $M_{\text{dust}}$


As a previous step, we show in Figure 8 the relation of the respective observables of  $M_{\star}$  and  $M_{\text{dust}}$ , that is, the stellar luminosities  $L_{\star}$  and the fluxes at  $890 \mu\text{m}$  wavelength (scaled to a distance of 158.5 pc). From the figure, there is a continuity of the correlation for any range of  $L_{\star}$ , and it holds for the BD population. The linear regression shown in the figure is for the entire population (stars and BDs), obtained following the Bayesian method described in Kelly (2007)<sup>1</sup>. Uncertainties and upper limits are taken into account, while sub-luminous objects are excluded for the fit.

Testi et al. (2016) found potential evidence of BD disks being less massive than stellar disks, based on the analysis of an incomplete sample of BD disks in Ophiuchus. Our Lupus sample allow us to check whether similar results hold in this star forming region. In Figure 8, there is no obvious trend for BDs to have very significantly smaller  $890 \mu\text{m}$  fluxes than stars with similar luminosities. To make a quantitative statement, we follow a similar procedure as in Testi et al. (2016) based on a statistical comparison of the two populations, and analyze whether the distribution of the  $M_{\text{dust}}/M_{\star}$  ratios in the sample of BD disks is consistent with being drawn from the same distribution as for the stars.


Figure 9 shows the cumulative distributions and the histograms of the values of the  $M_{\text{dust}}/M_{\star}$  ratios for the Lupus samples. Dust mass of each object is derived following Section 4.2,  $M_{\star}$  as described in Section 2. The histogram shows that the values of the BD ratios are similar to the stellar population ratios, unlike the Ophiuchus sample in Testi et al. (2016). We have performed the Anderson-Darling test<sup>2</sup> to study the null hypothesis that the two samples are drawn from the same underlying population, obtaining a probability of 6% that the BD and stellar disk populations are drawn from the same distribution. Although it is

<sup>1</sup> implemented with the linmix Python package, <https://linmix.readthedocs.io/en/latest/index.html>


<sup>2</sup> using scipy.stats Python module, <https://docs.scipy.org/doc/scipy/reference/stats.html>



**Fig. 8.**  $890\mu\text{m}$  fluxes versus stellar luminosity for the Lupus population. Stellar population is shown in blue; the BD disk population is plotted in red. Detected sources from ALMA observations are represented as circles; upper limits of non-detections are shown as triangles; sub-luminous objects as inferred from X-Shooter spectra are plotted as squares. The linear regression shown has been obtained from the entire population, excluding sub-luminous sources.



**Fig. 10.** Disk dust masses versus central object mass for the BD (red) and stellar (blue) populations in Lupus. Detected sources from ALMA observations are represented as circles; upper limits of non-detections are shown as triangles. Dust mass uncertainties and upper limits as described in Section 4.4. Uncertainties of  $M_*$  are  $1\sigma$ . Linear regression shown for the entire disk population (stars and BDs).



**Fig. 9.** Statistical comparison between BD and TTS disk populations in the Lupus star forming region. The histograms and cumulative distributions of the dust mass-stellar mass ratio is shown for both populations. The results summing up both populations are also included (shown as gray). A bin-size of 0.2 has been used for the histogram of the populations.

a low percentage, it is below  $2\sigma$  significance. Moreover, the likelihood increases to  $\sim 80\text{-}90\%$  if we use the dependence of  $T_{\text{dust}}$  with  $L_*$  from Andrews et al. (2013); van der Plas et al. (2016). Thus, the data are consistent with the null hypothesis to be correct. Our analysis of the Lupus sample do not show a statistically different fraction of dust mass around BDs as compared to stars. We caution that our sample of VLM stars and BDs in Lupus is very limited and that the results of Testi et al. (2016) was based on highly incomplete and inhomogeneous samples. Further studies with larger and/or unbiased samples are needed to conclusively assess this point.


The result of the previous analysis is also confirmed by inspecting the dependence of  $M_{\text{dust}}$  on  $M_*$  (Figure 10). In Appendix B, this dependence is shown for the other  $T_{\text{dust}}$  prescriptions. The linear regression result is consistent to Ansdell et al. (2016); Pascucci et al. (2016) when using the same assumptions of dust opacity and temperature. The slope ( $\alpha$ ) and in-

tercept ( $\beta$ ) for the stellar population are  $\alpha = 1.73 \pm 0.25$  and  $\beta = -3.88 \pm 0.14$  respectively (inferred using `linmix` package, and including upper limits of ALMA non-detections). As consequence of incorporating the BD population into the fit, there is a substantial reduction of the uncertainty of  $\alpha$  and  $\beta$ , thanks to the extension of the mass range over one order of magnitude: the slope and intercept become  $1.69 \pm 0.19$  and  $-3.89 \pm 0.13$ . If we compute a linear regression only taking into account the BD sample, we obtain a slope and intercept that is in agreement with the stellar fit, although the uncertainties in this case are large due to the short range in both axes of the BD population.

### 5.1.2. Disk size-luminosity relation

The existence of a correlation between the disk luminosity and its size was first shown from pre-ALMA observations (Andrews et al. 2010; Piétu et al. 2014). For the Lupus disk population, this dependence was confirmed in Tazzari et al. (2017) and Andrews et al. (2018). In Figure 11, we show the updated relation for the Lupus disk population, including the seven new measurements from this paper (one BD disk and six disks around stars). The linear regression shown in the figure is obtained for the stellar disk population, excluding the upper-limits of the disks with poorly constrained sizes.

The BD disk with well constrained  $R_{68\%}$  (J154518.5-342125) is in very good agreement with the relation for stars. The result suggests that this BD disk is as a scaled down equivalent of the very extended disks around stars that show substructure. Nevertheless, since its central object mass is near the BD/VLM boundary, this result might not be representative of the full BD population. Higher angular resolution observations of the BD population are needed in order obtain reliable estimates of their sizes. The estimated size upper limits of the other BD disks provide little constraints on the relation. The compactness of the BD disks might be indicative that BD disks follow the size-luminosity relation of stars, as suggested by Helder et al. (2017) from SED fitting of disk observations, and also from the results in  $\rho$ -Ophiuchus Testi et al. (2016). If these objects follow



**Fig. 11.** ALMA 890  $\mu\text{m}$  fluxes, scaled to a common distance, versus observed size ( $R_{68\%}$ , see text), for stars (blue) and BDs (red) in the Lupus star forming region. Upper limits of disk sizes that could not be determined accurately from our disk modeling methodology are shown as triangles, they represent the 95% confidence level of the disk size.  $R_{68\%}$  error bars account for  $1\sigma$  from the mean value, flux error bars are associated to the 10% flux calibration uncertainty.


the same relation as stars, their  $R_{68\%}$  would range between 1 and 10 au.

There is now evidence of optically thick emission in the inner ( $\lesssim 50$  au) regions of disks around stars (Huang et al. 2018; Zhu et al. 2019). Likewise, BD disks in Lupus might be optically thick, as suggested by their compact continuum emission. In Figure 11, we show two optically thick (optical depth  $\tau >> 1$ ) fiducial models, the first one (green) assuming a constant  $T_{\text{dust}}$  of 20 K, and a second model (purple) with radial dependence of  $T_{\text{dust}}[\text{K}] \approx 30 \cdot \left(\frac{L_*}{L_\odot}\right)^{0.25} \cdot \left(\frac{R}{10}\right)^{-0.5}$  (Andrews et al. 2013). The emission of these models are described by  $I_\nu(R) = \mathcal{F} B_\nu(T_{\text{dust}})$ , where  $\mathcal{F}$  is a filling factor that describes the fraction of the disk emission distribution that is optically thin:  $\mathcal{F} = 1$  if the disk emission is optically thick,  $0 < \mathcal{F} < 1$  for a partially optically thick disk (analogous to Tripathi et al. 2017; Andrews et al. 2018). The optically thick curves in the figure are built considering a set of disks with increasing outer size. Objects laying on the line are compatible with being fully optically thick. Additionally, in optically thick disks, the inferred  $R_{68\%}$  trace the location of large grains rather than the physical outer radius of the disk (Rosotti et al. 2019).

The only BD with determined dust disk size (J154518.5-342125) lays below both fiducial models. Its dust emission can be understood as optically thin with a fraction of the disk emission distribution being optically thick. If its disk emission is partially optically thin, a portion of dust is not observed, thus the inferred dust mass is underestimated. The upper-limits of the remaining BD disks are far below the optically thick models, although their exact positions in the  $R_{68\%}$ - $F_{890\mu\text{m}}$  are unknown.

### 5.1.3. Correlation between $\dot{M}_{\text{acc}}$ and $M_{\text{dust}}$

A linear correlation between mass accretion rate onto the central object ( $\dot{M}_{\text{acc}}$ ) and the disk mass is expected if disks evolve viscously (e.g., Dullemond et al. 2006; Natta et al. 2007; Lodato et al. 2017). Observational evidence for this correlation was first



**Fig. 12.** Relation between the total disk mass (obtained assuming  $\kappa_{890\mu\text{m}} = 2 \text{ cm}^2 \text{ g}^{-1}$ , and  $T_{\text{dust}} = 20 \text{ K}$ ) and the mass accretion rate onto the central object.

reported by Manara et al. (2016b) for the Lupus disks, and Mulders et al. (2017) in the Chamaeleon I region.

The  $\dot{M}_{\text{acc}}\text{-}M_{\text{dust}}$  relation for Lupus disks is shown in Figure 12. The x-axis of the figure is an estimate of the total disk mass, based on our derivation of the disk dust mass and assuming a gas-to-dust ratio of 100. The  $\dot{M}_{\text{acc}}$  and its uncertainty is taken from the X-Shooter observations presented by Alcalá et al. (2014, 2017). The accretion rate values have been recomputed with the new accretion luminosities that correspond to the parallaxes from Gaia DR2. One BD disk (SONYC-Lup3-10) was not characterized from X-Shooter observations. Although the  $H_\alpha$  emission line is known (Mužić et al. 2014), we have excluded this BD from the analysis in order to have a fully homogeneous sample for our statistical study.

The linear regression for the stellar population in Figure 12 has been obtained excluding BDs (due to the different BD disks behavior compared to stellar disks, demonstration below), non-detections from ALMA, upper limits of  $\dot{M}_{\text{acc}}$  and sub-luminous sources. The resulting slope is  $\alpha = 0.69 \pm 0.14$ , while the intercept is  $\beta = -7.26 \pm 0.36$ . When using the same assumptions of dust temperatures and opacities, the linear regression is consistent with the results presented in Manara et al. (2016b). BDs have systematically lower accretion rates than stars for the same disk mass. This is also seen in the relation between the more direct observed properties (in Appendix D), and is independent to the considered prescription of the dust temperature (results using other prescriptions in Appendix B).


We have inspected the  $M_{\text{disk}}/\dot{M}_{\text{acc}}$  ratio for the Lupus disk population to confirm or deny this trend. This ratio can be understood as the accretion depletion timescale (or disk age, as in Jones et al. 2012, see also Rosotti et al. 2017), and provides an estimate of the survival time of the disk, assuming that accretion onto the central object remains constant and that accretion is the dominant mechanism for the depletion of the disk. In Figure 12, we plotted different lines indicating accretion depletion timescales of 0.1, 1 and 10 Myr.

Analogous to the Subsection 5.1.1, we have conducted a statistical analysis of the two populations in order to confirm the behavior of the BD disks. We compare the BD population with the sub-sample of stars with disk masses within the range of the BD disk masses (in other words, all disks with  $\log_{10}(M_{\text{disk}}) < -3$  in Figure 12). This is done in order to remove the more mas-

**Table 4.** Linear regression results of all the investigated disk properties correlations.

X-axis	Y-axis	$T_{\text{dust}}$ prescription	$\alpha$	$\beta$	Dispersion
$L_{\star} [L_{\odot}]$	$F_{890\mu\text{m}} [\text{mJy}]$	-	$1.27 \pm 0.13$	$2.02 \pm 0.13$	$0.60 \pm 0.07$
$F_{890\mu\text{m}} [\text{mJy}]$	$L_{\text{acc}} [L_{\odot}]$	-	$0.81 \pm 0.15$	$-3.28 \pm 0.22$	$0.68 \pm 0.08$
$M_{\star} [M_{\odot}]$	$M_{\text{dust}} [M_{\odot}]$	Constant (20 K)	$1.69 \pm 0.19$	$-3.89 \pm 0.13$	$0.60 \pm 0.05$
		Andrews et al. (2013)	$0.95 \pm 0.18$	$-4.08 \pm 0.12$	$0.57 \pm 0.05$
		van der Plas et al. (2016)	$1.25 \pm 0.18$	$-3.98 \pm 0.12$	$0.58 \pm 0.05$
$R_{68\%} [\text{au}]$	$F_{890\mu\text{m}} [\text{mJy}]$	-	$1.31 \pm 0.17$	$-0.41 \pm 0.27$	$0.41 \pm 0.05$
$100 \cdot M_{\text{dust}} [M_{\odot}]$	$M_{\text{acc}} [M_{\odot}/\text{yr}]$	Constant (20 K)	$0.69 \pm 0.14$	$-7.26 \pm 0.36$	$0.56 \pm 0.08$
		Andrews et al. (2013)	$0.63 \pm 0.17$	$-7.49 \pm 0.43$	$0.64 \pm 0.09$
		van der Plas et al. (2016)	$0.67 \pm 0.16$	$-7.38 \pm 0.42$	$0.62 \pm 0.09$

**Notes.** The values of  $\alpha$  and  $\beta$  correspond to the slope and intercept of the linear fit, following the linear relation  $\log(Y) = \beta + \alpha \cdot \log(X)$ . The dispersion is the standard deviation of the regression in dex.  $F_{890\mu\text{m}}$  has been scaled to the average distance of the region (158.5 pc).



**Fig. 13.** Histograms (bottom) and cumulative distributions (top) of the accretion depletion timescale are shown for both populations (BDs as red, stars as blue). The results summing up both populations are also included (shown as gray).

sive disks of the stellar sample, which may accentuate the difference between populations. Sub-luminous objects and those with upper-limits of the  $\dot{M}_{\text{acc}}$  are excluded from the studied samples. The median value of the central object mass is  $0.08 M_{\star}$  for the BD sample and  $0.19 M_{\star}$  for the stellar sub-sample. The histograms of the accretion depletion timescale and the respective cumulative distributions are shown in Figure 13. The Anderson-Darling test confirms that the BD disk population has a significantly larger accretion depletion timescale respect to the stellar population, with only a 0.6% probability that the BD and stellar disks populations are drawn from the same original distribution. The median value of this timescale for the BD sample is 9.5 Myr, while for the stellar sub-sample considered is 1.4 Myr. The results hold when using other prescriptions of the disk dust mass for this test, with even lower probabilities ( $\sim 0.05\%$ ).

The result of the accretion depletion timescale is obtained using the total disk mass, which is estimated assuming that the emission of the dust is optically thin, and that dust mass traces the total disk mass. If (sub-)mm emission of BD disks is optically thick, the disk masses are underestimated, and consequently the accretion depletion timescale would be larger than the estimated values. Thus, the difference on the accretion depletion timescale would be even more pronounced if BD disks are optically thick at these wavelengths.

A larger accretion depletion timescale may reflect a difference in the accretion process of BDs with respect to stars. If viscous evolution models are invoked to explain accretion onto

the central star, a weaker accretion would imply a lower  $\alpha$  parameter (Shakura & Sunyaev 1973) in BD disks. Since the turbulence in viscous disks depends on  $\alpha$ , a lower  $\dot{M}_{\text{acc}}$  implies a less turbulent disk. Thus, in viscous disks, our result suggests that the  $\alpha$  parameter of disks around BDs is lower than around stars (in contrast with the results of Mulders & Dominik 2012), and consequently BD disks being less turbulent. A bi-modal behavior of accretion has been suggested observationally in Alcalá et al. (2017); Manara et al. (2017), and predicted from theoretical modeling in Vorobyov & Basu (2009); in those studies the two suggested modes were between VLMs ( $M_{\star} < 0.2 M_{\odot}$ ) and more massive stars. When performing a statistical comparison in our sample between the VLM population ( $0.1 M_{\odot} < M_{\star} < 0.2 M_{\odot}$ ) and more massive stars, we obtain a likelihood in the Anderson-Darling test of  $\sim 9\%$  (averaged over the three different prescriptions of  $M_{\text{dust}}$  used in this work). Thus, VLM stars might show this behavior as well, but less pronounced, and with much lower statistical significance.

A lower viscosity in disks around BDs/VLM stars compared to disks around more massive stars could be explained with a globally lower ionization rate (e.g., see Mohanty et al. 2005). As a consequence of the general correlation between X-ray and bolometric luminosities, young BDs/VLM stars have slightly lower X-ray luminosities than more massive young low-mass stars (see, e.g., Gregory et al. 2016); this might cause slightly lower ionization rates in the disks of the BDs/VLM objects. Another tentative explanation of lower ionization rates might be if some BD/VLM disks are flatter than disks around more massive stars (as suggested from SED models and observations by Ercolelano et al. 2009; Pascucci et al. 2003; Apai et al. 2004; Allers et al. 2006), which would decrease the irradiation cross-section. However, there is evidence of disks around BDs being flared (e.g., Natta & Testi 2001; Natta et al. 2002; Furlan et al. 2011), thus this last option would need more detailed and extended investigation.

## 5.2. Planet formation around BDs

The exoplanetary systems recently discovered around BDs and VLMs can be used to study the ability of BDs to form planets. Since the planets hosted by Trappist-1 and Proxima Centauri are most likely of rocky composition, the total planetary mass of these systems can be compared to the estimates of the disk dust mass of the BD disk population in Lupus. The summed up mass of the seven known planets (Gillon et al. 2017) in Trappist is  $4 M_{\oplus}$  (Wang et al. 2017). Proxima B, the planet hosted by our closest neighbor (Anglada-Escudé et al. 2016), has a minimum

mass of  $1.3 M_{\oplus}$ , and (Bixel & Apai 2017) predicted a mass of  $1.63^{+1.66}_{-0.72} M_{\oplus}$  with 95% confidence level. The remaining known exoplanets around BDs have estimated masses of at least several times Earth's mass. A considerable fraction of them have been detected via microlensing (e.g., Jung et al. 2018), which typically much higher masses. Thus the picture for these planets is analogous to the Trappist-1 planets and Proxima B.

From theoretical predictions of planet formation around BDs via core accretion (Payne & Lodato 2007), disk masses (gas and dust) on the order of a few Jupiter masses are required in order to form Earth-like planets around BDs. Not only none of the BD disks in Lupus (this paper) and  $\rho$ -Oph (Testi et al. 2016) have enough mass available at their current stage to form a planetary system, but even the available mass in solids is smaller than the total planetary mass in Trappist-1. The efficiency to convert the available mass into the final planetary rocky cores might be boosted due to internal recycling of the disk material, but it is unlikely to reach an efficiency close to unity (Manara et al. 2018, and references therein). On the other hand, the tentative result from Section 5.1.3 of lower viscosity and ionization rates on BD disks might contribute to the presence of an extended dead zone in the disk, which would boost up the planet formation process.

A plausible explanation to alleviate this divergence is that the determination of dust mass from continuum emission flux might be underestimated, as pointed out in Ballering & Eisner (2019). This might be the case if the emission at this wavelength is optically thick, then the inferred  $M_{\text{dust}}$  provides only a lower limit of the disk dust mass. In Fig. 11, disks laying on the  $\tau \gg 1$  fiducial models are consistent with their emission being fully optically thick. The only Lupus BD disk with well-determined size (J154518.5-342125) is below these models. This suggests that its emission is optically thin with small regions of the disk being optically thick. The inferred dust mass of this disk is underestimating by an unknown fraction. This can help explaining the mass difference with exoplanetary systems. Nevertheless, it seems unlikely that only this indeterminacy can account for the large solid mass difference in disks with partially optically thin emission.

A likely possibility is that planets might have already formed at this stage of the disk evolution (Greaves & Rice 2010; Najita & Kenyon 2014; Manara et al. 2018; Dodds et al. 2015). If that is indeed the case, the formation of planetary rocky cores would have occurred within the first Myr (considering the estimated ages of Lupus and  $\rho$ -Ophiuchus). While direct detection of planets embedded in protoplanetary disks is extremely difficult (see Sanchis et al. 2019; Johns-Krull et al. 2016), the presence of circumplanetary disks sets, if confirmed, a strong indirect evidence of young planets in these disks (Keppler et al. 2018; Isella et al. 2019; Pérez et al. 2019). Analysis of the gas kinematics can also be used as an indirect method to study embedded planets (Teague et al. 2018; Pinte et al. 2018). Other indirect indications, such as the existence of gaps, spirals, assymmetries, and dust processing are observed frequently, and suggest that planets might already have formed (e.g., ALMA Partnership et al. 2015; Zhang et al. 2018; Lodato et al. 2019; Pinilla et al. 2018).

## 6. Conclusions

In this work we presented new Band 7 ALMA observations of five protoplanetary disks around BDs in Lupus. Combined with previous observations, we present and analyze the (sub-)mm disk properties of the known population of BDs/VLM objects with infrared excess.

- We infer total dust disk masses and characteristic sizes of the disk population, measuring the continuum fluxes and modeling the visibilities.
- We update the relations of  $M_{\star}\text{-}M_{\text{dust}}$ , size-luminosity and  $M_{\text{dust}}\text{-}\dot{M}_{\text{acc}}$  extending them down to the substellar regime.
- BD disks in Lupus follow the relation for stars between stellar mass and dust disk mass. There is no statistical difference with the stellar disk population on the disk mass fraction, however we note the apparent lack of massive BD disks.
- BD disks are extremely compact, size determination was only possible on one BD disk, while for the other detected disks we present upper limits on the size.
- The accretion depletion timescale (inferred assuming that dust mm continuum emission is a reliable proxy of the total disk mass) of the BD population is significantly longer than for stars (9.5 Myr vs. 1.4 Myr), which in viscously evolving disks may imply a lower  $\alpha$  value, possibly linked to a globally lower ionization rate.
- Estimated disk dust masses are very low, suggesting that either these systems are unable to form planets, or, more likely, rocky planetary cores are formed within the first Myr. Optically thick emission in BD disks can alleviate this mass discrepancy.

**Acknowledgements.** This paper makes use of the following ALMA data: ADS/JAO.ALMA#2018.1.00544.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan) and KASI (Republic of Korea), in co-operation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. This work was partly supported by the Italian Ministero dell'Istruzione, Università e Ricerca through the grant Progetti Premiali 2012 – iALMA (CUP C52I13000140001), by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Ref no. FOR 2634/1 TE 1024/1-1, and by the DFG cluster of excellence Origins ([www.origins-cluster.de](http://www.origins-cluster.de)), and by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No823823 (RISE DUSTBUSTERS project). T.H. acknowledges support from the European Research Council under the Horizon 2020 Framework Program via the ERC Advanced Grant Origins 83 24 28. KM acknowledges funding by the Science and Technology Foundation of Portugal (FCT), grants No. IF/00194/2015 and PTDC/FISAST/28731/2017. CM, SF, AM acknowledge an ESO Fellowship. M.T. has been supported by the UK Science and Technology research Council (STFC).

## References

- Alcalá, J. M., Manara, C. F., Natta, A., et al. 2017, *A&A*, 600, A20  
 Alcalá, J. M., Natta, A., Manara, C. F., et al. 2014, *A&A*, 561, A2  
 Allers, K. N., Kessler-Silacci, J. E., Cieza, L. A., & Jaffe, D. T. 2006, *ApJ*, 644, 364  
 ALMA Partnership, Brogan, C. L., Pérez, L. M., et al. 2015, *ApJ*, 808, L3  
 Andrews, S. M. 2015, *PASP*, 127, 961  
 Andrews, S. M., Rosenfeld, K. A., Kraus, A. L., & Wilner, D. J. 2013, *ApJ*, 771, 129  
 Andrews, S. M., Terrell, M., Tripathi, A., et al. 2018, *ApJ*, 865, 157  
 Andrews, S. M. & Williams, J. P. 2005, *ApJ*, 631, 1134  
 Andrews, S. M., Wilner, D., Hughes, M., Qi, C., & Dullemond, C. P. 2010, in *Bulletin of the American Astronomical Society*, Vol. 42, American Astronomical Society Meeting Abstracts #215, 527  
 Anglada-Escudé, G., Amado, P. J., Barnes, J., et al. 2016, *Nature*, 536, 437  
 Ansdell, M., Williams, J. P., Trapman, L., et al. 2018, *ApJ*, 859, 21  
 Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, *ApJ*, 828, 46  
 Apai, D., Pascucci, I., Sterzik, M. F., et al. 2004, *A&A*, 426, L53  
 Ballering, N. P. & Eisner, J. A. 2019, *AJ*, 157, 144  
 Baraffe, I. & Chabrier, G. 2010, *A&A*, 521, A44  
 Baraffe, I., Homeier, D., Allard, F., & Chabrier, G. 2015, *A&A*, 577, A42  
 Barenfeld, S. A., Carpenter, J. M., Ricci, L., & Isella, A. 2016, *ApJ*, 827, 142  
 Bixel, A. & Apai, D. 2017, *ApJ*, 836, L31  
 Bustamante, I., Merín, B., Ribas, Á., et al. 2015, *A&A*, 578, A23  
 Cazzoletti, P., Manara, C. F., Baobab Liu, H., et al. 2019, *A&A*, 626, A11  
 Chauvin, G., Lagrange, A. M., Dumas, C., et al. 2004, *A&A*, 425, L29  
 Chiang, E. I. & Goldreich, P. 1997, *ApJ*, 490, 368

- Cieza, L. A., Ruiz-Rodríguez, D., Perez, S., et al. 2018, MNRAS, 474, 4347  
 Comerón, F. 2008, The Lupus Clouds, ed. B. Reipurth, 295  
 Comerón, F., Rieke, G. H., Claes, P., Torra, J., & Laureijs, R. J. 1998, A&A, 335, 522  
 Cox, E. G., Harris, R. J., Looney, L. W., et al. 2017, ApJ, 851, 83  
 Daemgen, S., Natta, A., Scholz, A., et al. 2016, A&A, 594, A83  
 Dodds, P., Greaves, J. S., Scholz, A., et al. 2015, MNRAS, 447, 722  
 Dullemond, C. P., Dominik, C., & Natta, A. 2001, ApJ, 560, 957  
 Dullemond, C. P., Natta, A., & Testi, L. 2006, The Astrophysical Journal, 645, L69  
 Dunham, M. M., Allen, L. E., Evans, II, N. J., et al. 2015, ApJS, 220, 11  
 Ercolano, B., Clarke, C. J., & Robitaille, T. P. 2009, MNRAS, 394, L141  
 Facchini, S., van Dishoeck, E. F., Manara, C. F., et al. 2019, A&A, 626, L2  
 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306  
 Frasca, A., Biazzo, K., Alcalá, J. M., et al. 2017, A&A, 602, A33  
 Furlan, E., Luhman, K. L., Espaillat, C., et al. 2011, ApJS, 195, 3  
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1  
 Gillon, M., Triaud, A. H. M. J., Demory, B.-O., et al. 2017, Nature, 542, 456  
 Goodman, J. & Weare, J. 2010, Communications in Applied Mathematics and Computational Science, Vol. 5, No. 1, p. 65-80, 2010, 5, 65  
 Greaves, J. S. & Rice, W. K. M. 2010, MNRAS, 407, 1981  
 Gregory, S. G., Adams, F. C., & Davies, C. L. 2016, MNRAS, 457, 3836  
 Guilloteau, S., Dutrey, A., Piétu, V., & Boehler, Y. 2011, A&A, 529, A105  
 Hendler, N. P., Mulders, G. D., Pascucci, I., et al. 2017, ApJ, 841, 116  
 Huang, J., Andrews, S. M., Dullemond, C. P., et al. 2018, ApJ, 869, L42  
 Hughes, J., Hartigan, P., Krautter, J., & Kelemen, J. 1994, AJ, 108, 1071  
 Isella, A., Benisty, M., Teague, R., et al. 2019, arXiv e-prints, arXiv:1906.06308  
 Jayawardhana, R., Mohanty, S., & Basri, G. 2003, ApJ, 592, 282  
 Johns-Krull, C. M., McLane, J. N., Prato, L., et al. 2016, ApJ, 826, 206  
 Jones, M. G., Pringle, J. E., & Alexander, R. D. 2012, MNRAS, 419, 925  
 Jung, Y. K., Udalski, A., Gould, A., et al. 2018, AJ, 155, 219  
 Kelly, B. C. 2007, ApJ, 665, 1489  
 Keppler, M., Benisty, M., Müller, A., et al. 2018, A&A, 617, A44  
 Klein, R., Apai, D., Pascucci, I., Henning, T., & Waters, L. B. F. M. 2003, ApJ, 593, L57  
 Lauer, T. R., Ajhar, E. A., Byun, Y.-I., et al. 1995, AJ, 110, 2622  
 Lodato, G., Dipierro, G., Ragusa, E., et al. 2019, Monthly Notices of the Royal Astronomical Society, 486, 453  
 Lodato, G., Scardoni, C. E., Manara, C. F., & Testi, L. 2017, Monthly Notices of the Royal Astronomical Society, 472, 4700  
 Long, F., Herczeg, G. J., Harsono, D., et al. 2019, ApJ, 882, 49  
 Luhman, K. L. 2012, ARA&A, 50, 65  
 MacGregor, M. A., Wilner, D. J., Czekala, I., et al. 2017, ApJ, 835, 17  
 Manara, C. F., Fedele, D., Herczeg, G. J., & Teixeira, P. S. 2016a, A&A, 585, A136  
 Manara, C. F., Morbidelli, A., & Guillot, T. 2018, A&A, 618, L3  
 Manara, C. F., Rosotti, G., Testi, L., et al. 2016b, A&A, 591, L3  
 Manara, C. F., Tazzari, M., Long, F., et al. 2019, A&A, 628, A95  
 Manara, C. F., Testi, L., Herczeg, G. J., et al. 2017, A&A, 604, A127  
 Merín, B., Jørgensen, J., Spezzi, L., et al. 2008, ApJS, 177, 551  
 Mohanty, S., Jayawardhana, R., & Basri, G. 2005, ApJ, 626, 498  
 Mortier, A., Oliveira, I., & van Dishoeck, E. F. 2011, MNRAS, 418, 1194  
 Mulders, G. D. & Dominik, C. 2012, A&A, 539, A9  
 Mulders, G. D., Pascucci, I., & Apai, D. 2015, ApJ, 814, 130  
 Mulders, G. D., Pascucci, I., Manara, C. F., et al. 2017, ApJ, 847, 31  
 Mužić, K., Scholz, A., Geers, V. C., & Jayawardhana, R. 2015, ApJ, 810, 159  
 Mužić, K., Scholz, A., Geers, V. C., Jayawardhana, R., & López Martí, B. 2014, ApJ, 785, 159  
 Muzerolle, J., Hillenbrand, L., Calvet, N., Briceño, C., & Hartmann, L. 2003, ApJ, 592, 266  
 Muzerolle, J., Luhman, K. L., Briceño, C., Hartmann, L., & Calvet, N. 2005, ApJ, 625, 906  
 Najita, J. R. & Kenyon, S. J. 2014, MNRAS, 445, 3315  
 Natta, A. & Testi, L. 2001, A&A, 376, L22  
 Natta, A., Testi, L., Calvet, N., et al. 2007, in Protostars and Planets V, ed. B. Reipurth, D. Jewitt, & K. Keil, 767  
 Natta, A., Testi, L., Comerón, F., et al. 2002, A&A, 393, 597  
 Natta, A., Testi, L., Muzerolle, J., et al. 2004, A&A, 424, 603  
 Pascucci, I., Apai, D., Henning, T., & Dullemond, C. P. 2003, ApJ, 590, L111  
 Pascucci, I., Testi, L., Herczeg, G. J., et al. 2016, ApJ, 831, 125  
 Payne, M. J. & Lodato, G. 2007, MNRAS, 381, 1597  
 Pérez, S., Casassus, S., Hales, A., et al. 2019, arXiv e-prints, arXiv:1906.06305  
 Piétu, V., Guilloteau, S., Di Folco, E., Dutrey, A., & Boehler, Y. 2014, A&A, 564, A95  
 Pinilla, P., Natta, A., Manara, C. F., et al. 2018, A&A, 615, A95  
 Pinte, C., Price, D. J., Ménard, F., et al. 2018, ApJ, 860, L13  
 Ricci, L., Testi, L., Natta, A., et al. 2014, ApJ, 791, 20  
 Rosotti, G. P., Clarke, C. J., Manara, C. F., & Facchini, S. 2017, MNRAS, 468, 1631  
 Rosotti, G. P., Tazzari, M., Booth, R. A., et al. 2019, MNRAS, 486, 4829  
 Sanchis, E., Picogna, G., Ercolano, B., Testi, L., & Rosotti, G. 2019, MNRAS, submitted (not yet accepted)  
 Scholz, A. 2008, Reviews in Modern Astronomy, 20, 357  
 Scholz, A. & Eisloffel, J. 2004, A&A, 419, 249  
 Shakura, N. I. & Sunyaev, R. A. 1973, A&A, 500, 33  
 Siess, L., Dufour, E., & Forestini, M. 2000, A&A, 358, 593  
 Sonnhalter, C., Preibisch, T., & Yorke, H. W. 1995, A&A, 299, 545  
 Tazzari, M. 2017, mtazzari/uvplot: v0.1.1  
 Tazzari, M., Beaujean, F., & Testi, L. 2018, MNRAS, 476, 4527  
 Tazzari, M., Testi, L., Ercolano, B., et al. 2016, A&A, 588, A53  
 Tazzari, M., Testi, L., Natta, A., et al. 2017, A&A, 606, A88  
 Teague, R., Bae, J., Bergin, E. A., Birnstiel, T., & Foreman-Mackey, D. 2018, ApJ, 860, L12  
 Testi, L., Birnstiel, T., Ricci, L., et al. 2014, in Protostars and Planets VI, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 339  
 Testi, L., Natta, A., Scholz, A., et al. 2016, A&A, 593, A111  
 Todorov, K., Luhman, K. L., & McLeod, K. K. 2010, ApJ, 714, L84  
 Tripathi, A., Andrews, S. M., Birnstiel, T., & Wilner, D. J. 2017, ApJ, 845, 44  
 van der Plas, G., Ménard, F., Ward-Duong, K., et al. 2016, ApJ, 819, 102  
 van Terwisga, S. E., van Dishoeck, E. F., Ansdell, M., et al. 2018, A&A, 616, A88  
 van Terwisga, S. E., van Dishoeck, E. F., Cazzoletti, P., et al. 2019, A&A, 623, A150  
 Vorobyov, E. I. & Basu, S. 2009, ApJ, 703, 922  
 Wang, S., Wu, D.-H., Barclay, T., & Laughlin, G. P. 2017, arXiv e-prints, arXiv:1704.04290  
 Ward-Duong, K., Patience, J., Bulger, J., et al. 2018, AJ, 155, 54  
 Whelan, E. T., Ray, T. P., Bacciotti, F., et al. 2005, Nature, 435, 652  
 Williams, J. P. 2012, Meteoritics and Planetary Science, 47, 1915  
 Yorke, H. W., Bodenheimer, P., & Laughlin, G. 1993, ApJ, 411, 274  
 Zhang, S., Zhu, Z., Huang, J., et al. 2018, The Astrophysical Journal, 869, L47  
 Zhu, Z., Zhang, S., Jiang, Y.-F., et al. 2019, ApJ, 877, L18

## Appendix A: Tests comparing $R_{68\%}$ and $R_{95\%}$

We computed the radii enclosing 68% ( $R_{68\%}$ ) and 95% ( $R_{95\%}$ ) for different models fitting a disk with high S/N and well resolved continuum emission, in order to test which radius is better as the characteristic size of the disk. Additionally, we demonstrate that the gaussian function can be used to describe the interferometric data of moderate angular resolution observations. For the calculation of these radii we build the cumulative flux as a function of radius:

$$f_{\text{cumul}}(R) = 2\pi \cdot \int_0^R I_\nu(R') \cdot R' \cdot dR', \quad (\text{A.1})$$

which gives us the flux contained within the radius  $R$ . Therefore,  $R_{68\%}$  and  $R_{95\%}$  is obtained from  $f_{\text{cumul}}(R_{68\%}) = 0.68 \cdot F_{\text{tot}}$  and  $f_{\text{cumul}}(R_{95\%}) = 0.95 \cdot F_{\text{tot}}$ . The total disk flux density  $F_{\text{tot}}$  is one of the free parameters of our fit and is obtained from its PDF. We build the PDFs of the derived radii: the median value of each distribution is used as the inferred value for each radius.

The resulting radii using three empirical models (gaussian, function, Nuker function, and broken power-law) and one physical model (two-layer approximation) were used to fit RXJ1556.1-3655 disk, the results are shown in Figure A.1. The values of  $R_{68\%}$  for the various models range between [29.3, 30.1] au, while the values of  $R_{95\%}$  range between [40.5, 47.4] au.


These results show that the difference of  $R_{68\%}$  between different models is negligible (thus the value of  $R_{68\%}$  is independent of the model used to fit our data) and for the  $R_{95\%}$  the values may differ and indeed be considerable. This is in accordance with Tripathi et al. (2017), and it is a consequence of the low sensitivity of larger scales, thus the noise makes the fitting of the visibilities uncertain. Therefore, the  $R_{68\%}$  is favored over the  $R_{95\%}$  as the most reliable size definition, since the differences between models are negligible.

On the other hand, this test also shows that the gaussian function is a reliable model to describe emission of moderate angular resolution observations, since the  $R_{68\%}$  obtained from the gaussian model is in perfect agreement with the value inferred using other empirical models.


We also tested the quality of our disk size results, by performing additional fits of disks around stars in Lupus that were already characterized in Andrews et al. (2018). The comparison between the inferred radii with the radii presented in Andrews et al. (2018) is shown in Figure A.2; the disk size values are indeed in very good agreement. Therefore the combination of the disk size results of the new disks modeled in this work and the disks modeled in Andrews et al. (2018) can be done adequately for the demographic analysis in Section 5.1.2.

## Appendix B: Disk properties relations using other dust temperature prescriptions

The demographic analysis comparing BD and stellar disks for the relations between disk properties was investigated for various prescriptions of the dust temperature of the disk. The results shown along Section 5 in the text are obtained assuming a constant  $T_{\text{dust}}$  of 20 K. Here we show the disk properties relations for other dust temperature dependence with stellar luminosity. The prescription from Andrews et al. (2013) was designed for disks around central objects of  $L_\star \in [0.1, 100] L_\odot$ ; the one from van der Plas et al. (2016) is more suitable for very low-mass objects (VLMs and BDs).



**Fig. A.1.** Normalized cumulative fluxes of Various fits used to model the interferometric visibilities of RXJ1556.1-3655 disk, used for the size determination. The results are shown for the following models: a gaussian distribution as in Equation 2 (illustrated in red), the Nuker profile from Equation 1 (blue), a broken power-law (green), and for the "Two-layer" approximation physical model to describe the disk (orange). The radii enclosing 68% and 95% of the total flux for each model are shown as dashed and dotted lines respectively.




**Fig. A.2.** Comparison of the characteristic size between this work and Andrews et al. (2018). For this comparison we followed the modeling described in Section 4 and fitted the interferometric data of a random sub-set of Lupus disks that were previously analyzed in Andrews et al. (2018). The dashed line represents the 1:1 ratio of the radii values.

### Appendix B.1: Dust temperature from Andrews et al. (2013)


### Appendix B.2: Dust temperature from van der Plas et al. (2016)

## Appendix C: Results from fits


The results of the interferometric modeling for the disks that could be characterized in radius are included in this section. The plots shown for each fitted disk are: (top left) the corner figure composed of the 1D and 2D histograms of the parameter investigation, (top right) the model and observed visibilities (real and imaginary part as a function of  $K\lambda$ ), (center right) the modeled brightness profile with its respective cumulative distri-




**Fig. B.1.** Relation between the stellar mass and the dust disk mass for the BD and stellar populations in Lupus, using the dust temperature dependence with stellar luminosity from Andrews et al. (2013).



**Fig. B.2.** Relation between the inferred disk mass of the source and the mass accretion rate onto the central object, using the dust temperature dependence with stellar luminosity from Andrews et al. (2013), and assuming a gas-to-dust ratio of 100.



**Fig. B.3.** Relation between the stellar mass and the dust disk masses for the BD and stellar populations in Lupus, using the dust temperature dependence with stellar luminosity from van der Plas et al. (2016).



**Fig. B.4.** Relation between the inferred disk mass of the source and the mass accretion rate onto the central object, using the dust temperature dependence with stellar luminosity from van der Plas et al. (2016), and assuming a gas-to-dust ratio of 100.

bution, (bottom) and the observed, modeled and residuals reconstruction in the imaginary plane from the interferometric analysis. Detailed description of the panels can be found in the captions of Figures 3, 4, 5 and 7.

#### Appendix C.1: Results for Sz 102

#### Appendix C.2: Results for V 1094 Sco

#### Appendix C.3: Results for GQ Lup

#### Appendix C.4: Results for Sz 76


#### Appendix C.5: Results for RXJ 1556.1-3655

#### Appendix C.6: Results for EX Lup


### Appendix D: Accretion luminosity vs. scaled continuum flux

We inspected the relation between accretion luminosity and continuum flux (scaled to 158.5 pc, the average distance to Lupus region) for the BD and stellar disks population. The demographic analysis is analogous to the different relations studied in Section 5. The accretion luminosity used is inferred from X-Shooter observations (Alcalá et al. 2014, 2017) and corrected with the parallaxes from Gaia Collaboration et al. (2018); continuum flux at 890  $\mu$ m is obtained from ALMA observations in Band 7 (AnsdeLL et al. 2016, , also this work). The relation is shown in Figure D.1. Red datapoints represent the BD population, while blue datapoints indicate the stellar disks. The linear regression of the stellar population is shown in the figure, it has been obtained excluding non-detections and sub-luminous sources.


The stellar relation seems to describe poorly the behavior of the BD disk population. In order to verify this result, we build the histograms of the distance to the linear regression of the two populations (Figure D.2), and performed the Anderson-Darling test comparing the Bd and stellar populations. This test gives a 0.02% of probability that the BD and the stellar disk populations are drawn from the same distribution. Therefore, the difference of accretion between BD and stars is statistically significant based only on observed quantities.




**Fig. C.1.** Modeling results of the Sz 102 disk from ALMA observations. A Nuker profile is fitted to the continuum emission distribution of the disk in the  $uv$ -plane. The panels show: the real and imaginary part of the observed and modeled visibilities, the 1D and 2D histograms of the Nuker model free parameters, the brightness emission and cumulative distributions, the  $R_{68\%}$  and  $R_{95\%}$  radii, and the fit results in the image plane.




**Fig. C.2.** Modeling results of the V 1094 Sco disk from ALMA observations. A Nuker profile is fitted to the continuum emission distribution of the disk in the *uv*-plane. The panels show: the real and imaginary part of the observed and modeled visibilities, the 1D and 2D histograms of the Nuker model free parameters, the brightness emission and cumulative distributions, the  $R_{68\%}$  and  $R_{95\%}$  radii, and the fit results in the image plane. The very extended emission of V1094 Sco was studied in detail in van Terwisga et al. (2018), and fitted with a more detailed function. For a comprehensive characterization of this disk we refer to that work. Nevertheless, our fit using the Nuker profile allows us to infer a characteristic radius consistent with the rest of the Lupus disk population.




**Fig. C.3.** Modeling results of the GQ Lup disk from ALMA observations. A Nuker profile is fitted to the continuum emission distribution of the disk in the  $uv$ -plane. The panels show: the real and imaginary part of the observed and modeled visibilities, the 1D and 2D histograms of the Nuker model free parameters, the brightness emission and cumulative distributions, the  $R_{68\%}$  and  $R_{95\%}$  radii, and the fit results in the image plane.




**Fig. C.4.** Modeling results of the Sz 76 disk from ALMA observations. A Nuker profile is fitted to the continuum emission distribution of the disk in the  $uv$ -plane. The panels show: the real and imaginary part of the observed and modeled visibilities, the 1D and 2D histograms of the Nuker model free parameters, the brightness emission and cumulative distributions, the  $R_{68\%}$  and  $R_{95\%}$  radii, and the fit results in the image plane.




**Fig. C.5.** Modeling results of the RXJ 1556.1-3655 disk from ALMA observations. A Nuker profile is fitted to the continuum emission distribution of the disk in the  $uv$ -plane. The panels show: the real and imaginary part of the observed and modeled visibilities, the 1D and 2D histograms of the Nuker model free parameters, the brightness emission and cumulative distributions, the  $R_{68\%}$  and  $R_{95\%}$  radii, and the fit results in the image plane.



**Fig. C.6.** Modeling results of the EX Lup disk from ALMA observations. A Nuker profile is fitted to the continuum emission distribution of the disk in the  $uv$ -plane. The panels show: the real and imaginary part of the observed and modeled visibilities, the 1D and 2D histograms of the Nuker model free parameters, the brightness emission and cumulative distributions, the  $R_{68\%}$  and  $R_{95\%}$  radii, and the fit results in the image plane.



**Fig. D.1.** Relation between the accretion luminosity and the continuum flux of the disk (scaled to the average distance of Lupus). These are the observables used to infer the mass accretion rate and the disk mass as discussed in Section 5.



**Fig. D.2.** Histogram of the ratio between accretion luminosity and continuum flux of the disk (scaled to the average distance of Lupus).