

Thermoelectric transport in a three channel “charge” Kondo circuit

T. K. T. Nguyen^{1,2} and M. N. Kiselev³

¹*Institute of Physics, Vietnam Academy of Science and Technology, 10 Dao Tan, Hanoi, Vietnam*

²*Institute of Theoretical and Applied Research (ITAR), Duy Tan University, Hanoi 100 000, Vietnam*

³*The Abdus Salam International Centre for Theoretical Physics, Strada Costiera 11, I-34151, Trieste, Italy*

(Dated: November 26, 2019)

We investigate theoretically the thermoelectric transport through a circuit implementation of the three-channel “charge” Kondo model quantum simulator [Z. Iftikhar *et al.*, *Science* **360**, 1315 (2018)]. The universal temperature scaling law of the Seebeck coefficient is computed perturbatively approaching the non-Fermi liquid strong coupling fixed point using abelian bosonization technique. The predicted $T^{1/3} \log T$ scaling behaviour of the thermoelectric power sheds a light on the properties of Z_3 emerging parafermions and gives an access to exploring pre-fractionalized zero modes in the quantum transport experiments. We discuss a generalization of approach for investigating a multi-channel Kondo problem with emergent $Z_N \rightarrow Z_M$ crossovers between “weak” non-Fermi liquid regimes corresponding to different low-temperature fixed points.

Quantum thermoelectricity is one of the most rapidly developing directions of the quantum technology [1, 2]. Modern progress in fabrication of nano-devices operating at ultra low (milli-Kelvin range) temperatures opens an access to a broad variety of the charge, spin and heat transport phenomena governed entirely by the quantum effects [3, 4]. In particular, quantization effects in behaviour of quantum simulators (see. e.g. [5–12]) at the regimes affected by quantum criticality are challenging for both experimental and theoretical communities.

Among a large variety of available quantum devices, Quantum Dots (QD) play important and significant role. On the one hand, the QD devices [3, 4] are highly controllable and fine-tunable setups operating at the regimes adjustable by external electric and magnetic fields at both weak and strong out of equilibrium conditions. On the other hand, the QD devices as the quantum impurity simulators provide an important playground for understanding the influence of strong electron-electron interactions, interference effects and resonance scattering on the quantum transport.

One of the cornerstone effects showing both the resonance scattering and strong interactions as two sides of the same coin is the Kondo effect [13, 14]. While conventional Kondo phenomenon is attributed to a spin degree of freedom of the quantum impurity [15–17], the unconventional “charge” Kondo effect is dealing with an iso-spin implementation of the charge quantization [18–23]. Kondo model [14–16] is one of known realizations of the “minimal models” archetypal for description of both Fermi liquid (FL) and non-Fermi liquid (NFL) regimes associated with the collective many-body phenomena.

The FL paradigm is one of the most important achievements of twentieth-century condensed matter physics [24]. It provides a tool to account for the effects of interaction in the equilibrium and out-of-equilibrium correlation functions [25]. While FLs are well defined objects characterized by some universal properties of corresponding quantum field theory encoded in scaling behaviour of the correlation functions [25] or, equivalently, certain constrains in the phenomenological de-

scription, NFLs represent rather “*terra incognita*” unless some strong fingerprints of the quantum behaviour inconsistent with the FL paradigm directly follow from known classes of the models. Fortunately, the multi-channel Kondo (MCK) model gives an access to collective behaviour completely different from the FL theory predictions [26–28]. The beauty and “simplicity” of the Kondo model makes it attractive for both experimental implementation of the strongly correlated physics and theoretical benchmarking of the many-body approaches beyond conventional mean-field or perturbation theory techniques. The price one has to pay for using minimal model is in immense complications in experimental fabrication of the MCK devices [30] and necessity to use advanced and cumbersome theoretical tools for the description of the strong coupling regimes [31–33].

Recently, the breakthrough experiments [11, 12] convincingly demonstrated the paramount importance of the MCK physics for the quantum charge transport through the nano-device. The few-channel Kondo physics is shown to be extended beyond existing realization of a two-channel Kondo (2CK) effect [29, 30] to a three-channel Kondo (3CK) phenomenon. While NFL regime of 2CK [34–36] is explained by an emergent Z_2 symmetry attributed to Majorana fermions [31, 37], the 3CK Kondo physics is known to be associated with Z_3 parafermion states [38–43].

In this *Letter* we address a fundamental question on how the NFL physics of 3CK model influences the quantum thermoelectric transport through the quantum simulators reported in [11, 12]. In particular, we investigate theoretically a scaling behaviour of thermoelectric coefficients and analyse crossovers between the NFL regimes associated with different low temperature strong coupling fixed points of 3CK.

Model – In a nano-device (see Fig. 1) designed to be used for thermoelectric measurements [44, 45], the drain consists of a large metallic QD electrically connected to two-dimensional electron gas (2DEG) electrodes through three quantum point contacts (QPCs) as proposed in Refs. [11, 12]. The 2DEG is in the in-

teger quantum Hall (IQH) regime at the filling factor $\nu = 2$. The QPCs are fine-tuned to satisfy condition that only the outer spin polarized chiral edge current is partially transmitted across the QPCs. The drain is at the reference temperature T . The source is separated from the QD by a tunnel barrier with low transparency $|t| \ll 1$ as described by a tunnel Hamiltonian $H_{tun} = \sum_k (tc_k^\dagger d + \text{h.c.})$ with c and d denote the electrons in the left lead and in the dot. The temperature of the source can be controlled by the “floating island” technique [5]. A micron-sized metal island [5] is electrically connected by several channels at opposite voltages (to have a zero dc voltage) in the left electrode upstream to the tunnel contact to the Kondo island [46]. Electrons in the “floating island” are heated up with Joule heat. The resulting temperature is measured by the noise-based thermometry [5, 46–48]. The temperature difference ΔT across the tunnel barrier is assumed to be small compared to the reference temperature T to guarantee the linear response regime for the device at the weak link [49]. The central metallic island (QD) is in a regime of weak (mesoscopic) Coulomb blockade [21, 50] characterized by the charging energy E_C . The gate voltage V_g is used to tune charge degeneracy $N(V_g)$ to the regimes of Coulomb peaks (N is half-integer) and Coulomb valleys (N is integer). The Kondo physics is observed through the measurements of the QPCs differential conductances G_α at zero bias voltages $V_\alpha \rightarrow 0$ through the measurement of I_α/V_α [12] (see Fig. 1). The MCK regime is fine-tuned by setting transmission coefficients across QPCs to be equal. Applying a thermo-voltage ΔV_{th} to implement a zero-current condition for the electric current between the source (orange lead) and drain (QD and three blue leads) allows to access the thermoelectric coefficient G_T through the measurements of $I_\alpha/\Delta T$ and Seebeck coefficient aka thermopower (TP) $S = G_T/G|_{I=0} = -\Delta V_{th}/\Delta T$ [44].

The mapping of IQH setup to a MCK problem is explained in details in Refs. [51]. We assign the iso-spin \uparrow to the electrons in each QPC and the iso-spin \downarrow to the electrons in the QD. The “charge” iso-spin flips when the electrons move in- and out- of the QD. Backscattering transfers “moving in-” the QD electrons to “moving out-” from the QD electrons and vice versa. The number of QPCs is equivalent to the number of orbital channels in the conventional $S=1/2$ Kondo problem.

It is convenient to describe the interacting electrons in the QD and QPCs in the bosonized representation [19–23, 52]. We start with the Euclidean action $S = S_0 + S_C + S'$ describing QD and three QPCs. The action S_0 [33] stands for the free part representing three copies of free one-dimensional electrons in QPC $_\alpha$:

$$S_0 = \frac{v_F}{2\pi} \sum_{\alpha=1}^3 \int_0^\beta dt \int_{-\infty}^{\infty} dx \left[\frac{(\partial_t \phi_\alpha(x, t))^2}{v_F^2} + (\partial_x \phi_\alpha(x, t))^2 \right],$$

Here $\phi_\alpha(x, t)$ denotes bosonic field describing the transport through QPC $_\alpha$ (see also [51]) and v_F is a Fermi velocity, $\beta=1/T$ (we adopt the units $\hbar=c=k_B=1$).




FIG. 1. (Color online) Schematic of three channel “charge” Kondo (3CK) setup. Central metallic island, aka Quantum Dot (QD) is connected to four electrodes formed by two dimensional electron gas (2DEG). The state in QD is characterized by the iso-spin $\sigma=\downarrow$. The states in the electrodes are characterized by the iso-spin $\sigma=\uparrow$. The left (orange) electrode is heated to the temperature $T+\Delta T$ and connected to the rest of the setup through a tunnel contact (red dashed lines). The reference temperature of the QD and three blue electrodes is T . The yellow plunger gate is used to control a mesoscopic Coulomb blockade in the QD. The setup is fine tuned by external magnetic field to the Integer Quantum Hall (IQH) regime $\nu=2$. The current propagates along spin-polarized edge channels (red lines with arrows). Only one relevant (outer) chiral edge channel is shown. The transparencies of the quantum point contacts (QPCs) QPC $_1$ -QPC $_3$ are controlled by the surface split gates (not shown). The identical thermo-voltages are applied across the tunnel contacts to nullify the net electric current through the device. Insert shows schematically a renormalization group (RG) flow for 3-channel Kondo effect. The unstable strong coupling fixed point at $G_1=G_2=G_3 \approx 0.69e^2/h$ corresponds to the 3CK non-Fermi-liquid regime.

The effects of the weak mesoscopic Coulomb blockade in the QD are described by the Hamiltonian $H_C = E_C[\hat{n} - N(V_g)]^2$. In the spirits of Andreev-Matveev theory [52], the operator \hat{n} in the Hamiltonian H_C accounts for the electrons entering the dot through the left weak tunnel barrier and three QPCs ($\hat{n}=\hat{n}_L+\hat{n}_{QPC}$). The number of electrons entering QD from the QPCs is related to the bosonic fields ϕ_α as $\hat{n}_{QPC} \rightarrow \sum_{\alpha=1}^3 \phi_\alpha(0, t)/\pi$ [52], while the operator \hat{n}_L counting the number of electrons tunneling from the left electrode can be replaced by the function $n_\tau(t) = \theta(t)\theta(\tau - t)$ [52]. Here $\theta(t)$ is the unit step function (Heaviside function.) The Coulomb blockade action S_C in bosonized representation [19–23, 52] is given by:

$$S_C(\tau) = \int_0^\beta dt E_C \left[n_\tau(t) + \frac{1}{\pi} \sum_{\alpha=1}^3 \phi_\alpha(0, t) - N(V_g) \right]^2.$$

Finally, the action S'

$$S' = -\frac{D}{\pi}|r| \sum_{\alpha=1}^3 \int_0^\beta dt \cos[2\phi_\alpha(0,t)]$$

characterizes the backscattering at QPCs with r_α is reflection amplitude for the QPC α and D is the bandwidth (ultraviolet cutoff). We consider the symmetric situation, where $|r_1| = |r_2| = |r_3| \equiv |r|$.

Three flavour electrons – we introduce three linear combinations of the fields ϕ_α to represent charge, spin and flavour modes (see, e.g. [53]):

$$\begin{aligned} \phi_c(x,t) &= \frac{1}{\sqrt{3}}(\phi_1(x,t) + \phi_2(x,t) + \phi_3(x,t)), \\ \phi_s(x,t) &= \frac{1}{\sqrt{2}}(\phi_1(x,t) - \phi_3(x,t)), \\ \phi_f(x,t) &= \frac{1}{\sqrt{6}}(\phi_1(x,t) - 2\phi_2(x,t) + \phi_3(x,t)), \end{aligned} \quad (1)$$

and the same for the dual boson fields $\frac{1}{\pi}\partial_x\theta_\alpha = \Pi_\alpha = -\frac{1}{v_F}\partial_t\phi_\alpha$ satisfying equal-time commutation relations: $[\phi_\alpha(x), \Pi_{\alpha'}(x')] = i\delta(x-x')\delta_{\alpha\alpha'}$ [31–33]. Here α, α' denote charge, spin, and flavour. The spin and flavour modes are related to two diagonal Gell-Mann matrices of SU(3) group [4, 54].

The action in the charge, spin and flavour modes is given by

$$S_0 = \frac{v_F}{2\pi} \int_0^\beta dt \int_{-\infty}^\infty dx \sum_{\alpha=c,s,f} \left[\frac{(\partial_t \phi_\alpha(x,t))^2}{v_F^2} + (\partial_x \phi_\alpha(x,t))^2 \right], \quad (2)$$

$$S_C(\tau) = \int_0^\beta dt E_C \left[n_\tau(t) + \frac{\sqrt{3}}{\pi} \phi_c(0,t) - N(V_g) \right]^2, \quad (3)$$

$$\begin{aligned} S' &= -\frac{D}{\pi}|r| \int_0^\beta dt \left\{ \cos \left[\frac{2}{\sqrt{3}} \phi_c(0,t) - \frac{2\sqrt{2}}{\sqrt{3}} \phi_f(0,t) \right] \right. \\ &\quad \left. + 2 \cos \left[\frac{2}{\sqrt{3}} \phi_c(0,t) + \frac{\sqrt{2}}{\sqrt{3}} \phi_f(0,t) \right] \cos \left[\sqrt{2} \phi_s(0,t) \right] \right\}. \end{aligned} \quad (4)$$

Action S_0 is particle-hole (PH) symmetric. PH transformation in the action S_C corresponds to change N to $-N$ (electrons are replaced by holes). As a result, the transport coefficients G and G_T transform under PH transformation as follows: $G(N) = G(-N)$ and $G_T(-N) = -G_T(N)$. Besides, the thermoelectric transport requires breaking of the particle-hole symmetry described by the backscattering action S' .

Furthermore, due to Coulomb blockade effect all transport coefficients are periodic in $N(V_g)$ and the action is invariant with respect to the shift $N \rightarrow N + 1$. To show it we notice that the electron travels from/to QD to/from one of the QPCs. In the setup (see Fig. 1), there are three possible ways to do it: i) electron enters QD from the QPC1: $\phi_1 \rightarrow \phi_1 + \pi$, $\phi_2 \rightarrow \phi_2$, $\phi_3 \rightarrow \phi_3$. As a result $\phi_c \rightarrow \phi_c + \pi/\sqrt{3}$, $\phi_s \rightarrow \phi_s + \pi/\sqrt{2}$, $\phi_f \rightarrow \phi_f + \pi/\sqrt{6}$; ii) electron enters QD from the QPC2: $\phi_1 \rightarrow \phi_1$, $\phi_2 \rightarrow \phi_2 + \pi$, $\phi_3 \rightarrow \phi_3$ then $\phi_c \rightarrow \phi_c + \pi/\sqrt{3}$,

$\phi_s \rightarrow \phi_s$, $\phi_f \rightarrow \phi_f - 2\pi/\sqrt{6}$; and iii) electron enters QD from the QPC3: $\phi_1 \rightarrow \phi_1$, $\phi_2 \rightarrow \phi_2$, $\phi_3 \rightarrow \phi_3 + \pi$ then $\phi_c \rightarrow \phi_c + \pi/\sqrt{3}$, $\phi_s \rightarrow \phi_s - \pi/\sqrt{2}$, $\phi_f \rightarrow \phi_f + \pi/\sqrt{6}$. These discrete transformations keep the backscattering action S' invariant and increase charge of the QD by one. We rely upon these transformations in the perturbative calculations (see details in the Supplemental Materials [55]).

Perturbative calculations – The transport coefficients G and G_T are expressed in terms of the correlation function $K(\tau)$ [52]:

$$\begin{aligned} K(\tau) &= Z(\tau)/Z(0), \\ Z(\tau) &= \int \exp[-S_0 - S_C(\tau) - S'] \prod_\alpha \mathcal{D}\phi_\alpha(x,t). \end{aligned} \quad (5)$$

This correlation function is characterized by the following symmetries associated with PH and shift transformation: $K(\beta - \tau, N) = K(\tau, 1 - N)$ and $K(\beta - \tau, N) = K(\tau, -N)$.

The electric conductance G [20] is given by

$$G = \frac{G_L \pi T}{2} \int_{-\infty}^\infty \frac{1}{\cosh^2(\pi T t)} K\left(\frac{1}{2T} + it\right) dt. \quad (6)$$

Here $G_L \ll e^2/h$ denotes the tunnel conductance of the left barrier calculated ignoring influence of the dot. The thermoelectric coefficient G_T takes the form [52]

$$G_T = -\frac{i\pi^2 G_L T}{2e} \int_{-\infty}^\infty \frac{\sinh(\pi T t)}{\cosh^3(\pi T t)} K\left(\frac{1}{2T} + it\right) dt. \quad (7)$$

The correlator $K(\tau)$ acquires a simple form in the absence of the backscattering. The action $S_0 + S_C$ is Gaussian and the functional integrals are explicitly evaluated resulting in [56] (see details of calculations in the Supplemental Materials [55])

$$K(\tau)|_{r=0} = K^{(0)}(\tau) = \left[\frac{\pi^2 T}{3\gamma E_C} \frac{1}{|\sin(\pi T \tau)|} \right]^{\frac{2}{3}}. \quad (8)$$

The backscattering $r \neq 0$ explicitly breaks the PH symmetry. However, the mechanism of the PH symmetry breaking is different for the FL ($M = 1$) and MCK-NFL, ($M \geq 2$) states. Namely, for the FL case, there exists only one gapped mode associated with the charge. Therefore, the PH symmetry breaking occurs already in the first order of the perturbation theory [52]. If, however, there are $M - 1$ gapless modes describing spin and flavours for the MCK-NFL, the first order perturbative correction vanishes and PH symmetry breaking occurs in the second order. The non-vanishing contribution to the G_T and S is associated with the fluctuations of $M - 1$ gapless modes. We proceed with the perturbative calculations at the second order $K^{(2)}(\tau) = K_C(\tau) (\langle S'^2 \rangle_\tau - \langle S'^2 \rangle_0) / 2$. The validity of the perturbation theory at $|r|^2 \ll 1$ for 2CK [52] is justified by the condition for the temperature regime $T^* \ll T \ll E_C$ where $T^* = |r|^2 E_C$ [52]. We refer to this regime as the *weak NFL regime*.

Scaling of transport coefficients – The main contribution to the electric conductance does not depend on $|r|$. Its temperature scaling is fully determined by the form of $K^{(0)}(\tau)$ given by Eq.(8) (see [57, 62]):

$$G \sim G_L [T/E_C]^{\frac{2}{3}}. \quad (9)$$

The perturbative contribution to the thermoelectric coefficient G_T proportional to $|r|^2$ consists of two terms [55] computed with log-accuracy. The first term is associated with contribution of electrons entering the dot from QPC1 and QPC3:

$$G_T^{(1)} \sim \frac{G_L}{e} |r|^2 \sin(2\pi N) [1 + a \cos(2\pi N)] \left[\frac{T}{E_C} \right] \ln \left[\frac{E_C}{T} \right], \quad (10)$$

and the second term accounts for the electrons entering the dot from the QPC2:

$$G_T^{(2)} \sim \frac{G_L}{e} |r|^2 \sin(2\pi N) \cos(2\pi N) \left[\frac{T}{E_C} \right] \ln \left[\frac{E_C}{T} \right]. \quad (11)$$

While the temperature scaling in both terms is the same, the difference in the gate voltage dependence is associated with the parametrization of the spin and flavor modes (1) where the symmetry between (QCP1, QCP3) and QCP2 is explicitly broken [63]. Substituting Eqs. (10, 11) and asymptotic equation for G Eq.(9) into the definition of the TP $S = G_T/G$ we obtain [64]:

$$S \sim \frac{1}{e} |r|^2 \sin(2\pi N) [1 + 2a \cos(2\pi N)] \left[\frac{T}{E_C} \right]^{\frac{1}{3}} \ln \left[\frac{E_C}{T} \right], \quad (12)$$

with $a \sim 1$ [55]. The perturbative 3CK results for G_T (10, 11) and TP (12) do not diverge at the limit $T \rightarrow 0$ in contrast to 2CK predictions [52]. Besides, the temperature scaling of TP $S^{3CK} \propto T^{1/3} \log T$ is compatible with corresponding *non-perturbative* scaling of the TP maximums $S_{\max}^{2CK} \propto T^{1/2} \log T$ for 2CK. In both cases S vanishes when $T \rightarrow 0$. We therefore expect that the scaling (12) will survive at the limit $T \rightarrow 0$ and acquire only marginal modifications in the argument of log [52]. Eqs. (10-12) represent the central result of this *Letter*.

Channel symmetry breaking – We comment on possible ways to crossover $3CK \rightarrow 2CK$ and $3CK \rightarrow 1CK$ in the “charge” Kondo circuits. These crossovers have been experimentally reported in [11, 12] and numerically reproduced in [65–67] by using Numerical Renormalization Group (NRG) technique. The simplest way to describe continuous crossover $3CK \rightarrow 2CK$ is to imbalance e.g. the reflection amplitudes in QPC1 and QPC3. Having $a_{13} \equiv ||r_1| - |r_3||$ as a relevant perturbation to the symmetric state characterized by $s_{13} \equiv (|r_1| + |r_3|)/2 \approx |r|$ provides a condition for a crossover $a_{13} \sim s_{13}$ similar to theory of channel symmetry breaking of $2CK \rightarrow 1CK$ discussed in [49]. In addition, the condition $a_{13}s_{13} \ll |r_2|^2 \approx |r|^2$ is required. However, one needs to go beyond the perturbation theory for the quantitative description of the crossover. The mechanism of $3CK \rightarrow 1CK$ is more delicate. First of all, the experiment [12] shows the non-monotonous behaviour

of conductance evolution confirmed by non-monotonous NRG flow in numerical calculations [65–67]. Second, the crossover regime has to be fine tuned by the condition $|a_{13}s_{13} - |r_2|^2| \ll |r_2|^2$. Discussion of both crossovers goes beyond the scope of this paper and will be published elsewhere [68].

Discussion and open questions – Describing the quantum thermoelectricity in the NFL regime of the MCK model at the strong coupling limit $T \ll T^*$ is one of the main open questions. In particular, it is important to understand if there exists a re-(para)fermionization procedure for Z_3 fixed point similar to Emery-Kivelson (EK) approach [37] developed for $U(1) \rightarrow Z_2$ symmetry reduction [69]. The EK re-fermionization being a cornerstone for understanding of the emergence of the NFL state of 2CK is known to allow straightforward re-writing of the strong coupling Hamiltonian in terms of Z_2 Majorana (para)fermions. However, even if such a procedure does exist for Z_3 low temperature fixed point [70], the strong coupling fixed point Hamiltonian will not be quadratic in terms of the Z_3 parafermions [71]. Therefore, the non-perturbative treatment of the 3CK problem at its strong coupling will require some additional assumptions or approximations. Yet another challenging question is related to the generalization of the approach developed in this *Letter* for the description of the $M > 3$ MCK effect at the strong coupling. We expect that even- and odd- M -channel models behave significantly differently: while the ground state of the even- $M = 2k$ channel models can be represented in terms of the Majorana fermions [72], Z_{2k+1} parafermions are needed for the description of the odd- $M = 2k + 1$ -channel Kondo physics. Besides, switching between Z_{2k+1} and Z_{2k} low temperature fixed points opens an interesting possibility for investigation the crossovers between the states with different parafermion fractionalized zero modes. The same goal can be achieved by using the quantum simulators containing a tunnel contact between two different NFL states [51].

Conclusions – In summary, we derived the scaling relations for the TP of the 3CK “charge” Kondo model approaching the NFL strong coupling fixed point. We analysed perturbatively the temperature behaviour of the thermoelectric coefficient and TP in the high-temperature *weak NFL* regime. We predict that while the electric conductance vanishes as a power law with an exponent inversely proportional to the number of channels, the thermoelectric coefficient G_T scales as $T \log T$ when the temperature goes down. As a result, the scaling dependence of the TP of 3CK model shows a strong deviation from corresponding behaviour of both 2CK and 1CK opening a possibility of experimental observation of Z_3 pre-fractionalization in the quantum transport experiments. We propose to use experimental technique [12] providing the circuit implementation of quantum simulators of the MCK model for investigation of the parafermion contribution to the quantum thermoelectricity controlled by switching the quantum regimes between different low temperature fixed points.

Acknowledgements – We are grateful to Leonid Glazman, Yuval Gefen, Joel Moore and Alexander Nersesyan for illuminating discussions. We especially grateful to Frederic Pierre for the detailed explanation of the experiments [11, 12], suggestions [47] for the experimental realization of the thermoelectric measurements with quantum simulators [46] and discussion of the dynamical Coulomb blockade effects [57, 58]. We acknowledge the warm hospitality of the Center for Theoretical Physics of Complex Systems (PCS) of the Institute for Basic Science (IBS) and the International Cen-

tre of Physics (ICP) of the Institute of Physics (IOP - VAST). T.K.T.N. acknowledges support through the ICTP Associate Program and ICTP Asian Network on Condensed Matter and Complex Systems. This research in Hanoi is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 103.01-2016.34. The work of MK was performed in part at Aspen Center for Physics, which is supported by National Science Foundation grant PHY-1607611 and partially supported by a grant from the Simons Foundation.

-
- [1] G. Benenti, G. Casati, K. Saito, and R. S. Whitney, *Phys. Rep.* **694**, 1 (2017).
- [2] V. Zlatic and R. Monnier, *Modern Theory of Thermoelectricity* (Oxford University Press, Oxford, 2014).
- [3] Y. M. Blanter and Y. V. Nazarov, *Quantum Transport: Introduction to Nanoscience* (Cambridge University Press, Cambridge, 2009).
- [4] K. Kikoin, M. N. Kiselev, and Y. Avishai, *Dynamical Symmetry for Nanostructures. Implicit Symmetry in Single-Electron Transport Through Real and Artificial Molecules* (Springer, New York, 2012).
- [5] S. Jezouin, F.D. Parmentier, A. Anthore, U. Gennser, A. Cavanna, Y. Jin, F. Pierre, *Science* **342**, 601 (2013).
- [6] S. Jezouin, Z. Iftikhar, A. Anthore, F.D. Parmentier, U. Gennser, A. Cavanna, A. Ouerghi, I.P. Levkivskiy, E. Idrisov, E.V. Sukhorukov, L.I. Glazman, F. Pierre, *Nature* **536**, 58 (2016).
- [7] E. Sivre, A. Anthore, F.D. Parmentier, A. Cavanna, U. Gennser, A. Ouerghi, Y. Jin, F. Pierre, *Nature Physics* **14**, 145 (2018).
- [8] M. Banerjee, M. Heiblum, V. Umansky, D.E. Feldman, Y. Oreg, A. Stern, *Nature* **562**, 205 (2018).
- [9] M. Banerjee, M. Heiblum, A. Rosenblatt, Y. Oreg, D.E. Feldman, A. Stern, V. Umansky, *Nature* **545**, 75 (2017).
- [10] H. Inoue, A. Grivnin, N. Ofek, I. Neder, M. Heiblum, V. Umansky, D. Mahalu, *Phys. Rev. Lett.* **112**, 166801 (2014).
- [11] Z. Iftikhar, S. Jezouin, A. Anthore, U. Gennser, F. D. Parmentier, A. Cavanna, and F. Pierre, *Nature* **526**, 233 (2015).
- [12] Z. Iftikhar, A. Anthore, A. K. Mitchell, F. D. Parmentier, U. Gennser, A. Ouerghi, A. Cavanna, C. Mora, P. Simon, and F. Pierre, *Science* **360**, 1315 (2018).
- [13] J. Kondo, *Prog. Theor. Phys.* **32**, 37 (1964).
- [14] A. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, Cambridge, England, 1993).
- [15] A. M. Tsvelik and P. B. Wiegmann, *Advances in Physics*, **32**, 453 (1983).
- [16] N. Andrei, K. Furuya, and J. H. Lowenstein, *Rev. Mod. Phys.* **55**, 331 (1983).
- [17] L. Kouwenhoven and L.I. Glazman, *Physics World* **14**, 33 (2001).
- [18] K. Flensberg, *Phys. Rev. B* **48**, 11156 (1993).
- [19] K.A. Matveev, *Phys. Rev. B* **51**, 1743 (1995).
- [20] A. Furusaki and K.A. Matveev, *Phys. Rev. B* **52**, 16676 (1995).
- [21] I. L. Aleiner and L. I. Glazman *Phys. Rev. B* **57**, 9608 (1998).
- [22] K. Le Hur, *Phys. Rev. B* **64**, 161302(R) (2001).
- [23] K. Le Hur and G. Seelig, *Phys. Rev. B* **65**, 165338 (2002).
- [24] L. D. Landau, *Sov. Phys. JETP* **3**, 920 (1957); **5**, 101 (1957).
- [25] P. Coleman, *Introduction to Many-Body Physics* (Cambridge University Press, Cambridge, 2015).
- [26] Ph. Nozières and A. Blandin, *J. Phys.* **41**, 193 (1980).
- [27] D. Cox and A. Zawadowski, *Advances in Physics* **47**, 599 (1998).
- [28] I. Affleck and A. W. W. Ludwig, *Phys. Rev. B* **48**, 7297 (1993).
- [29] Y. Oreg, D. Goldhaber-Gordon, *Phys. Rev. Lett.* **90**, 136602 (2003).
- [30] R.M. Potok, I.G. Rau, H. Shtrikman, Y. Oreg, and D. Goldhaber-Gordon, *Nature* **446**, 167 (2007).
- [31] A.O. Gogolin, A.A. Nersesyan, and A.M. Tsvelik, *Bosonization and strongly correlated systems* (Cambridge University Press, 1998).
- [32] T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, Oxford, 2004).
- [33] D. Sènèchal, *An Introduction to Bosonization. In: Sènèchal D., Tremblay A.M., Bourbonnais C. (eds) Theoretical Methods for Strongly Correlated Electrons. CRM Series in Mathematical Physics.* (Springer, New York, 2004).
- [34] N. Andrei and C. Destri, *Phys. Rev. Lett.* **52**, 364 (1984).
- [35] M. Fabrizio, A. O. Gogolin, and Ph. Nozières, *Phys. Rev. Lett.* **74**, 4503 (1995).
- [36] M. Fabrizio, A. O. Gogolin, and Ph. Nozières, *Phys. Rev. B* **51**, 16088 (1995).
- [37] V.J. Emery and S. Kivelson, *Phys. Rev. B* **46**, 10812 (1992).
- [38] A.B. Zamolodchikov and V.A. Fateev, *Sov. Phys. JETP* **62**, 215 (1985).
- [39] H. Yi, C. L. Kane, *Phys. Rev. B* **57**, R5579 (1998).
- [40] H. Yi, *Phys. Rev. B* **65**, 195101 (2002).
- [41] I. Affleck, M. Oshikawa, H. Saleur, *Nucl. Phys. B* **594**, 535 (2001).
- [42] C. Nayak, S.H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *Rev. Mod. Phys.* **80**, 1083 (2008).
- [43] J. Alicea and P. Fendley, *Annu. Rev. Condens. Matter Phys.* **7**, 119 (2016).
- [44] R. Scheibner, H. Buhmann, D. Reuter, M. N. Kiselev, and L. W. Molenkamp, *Phys. Rev. Lett.* **95**, 176602 (2005).

- [45] R. Scheibner, E. G. Novik, T. Borzenko, M. König, D. Reuter, A.D. Wieck, H. Buhmann and L. W. Molenkamp, *Phys. Rev. B* **75**, 041301 (2007).
- [46] Frederic Pierre (private communication).
- [47] Alternatively, the heating of the left electrode (Fig. 1) can be achieved by injection of some energy with the voltage biased QPC located upstream the tunnel contact [46, 48]. The electrons relax after some adjustable propagation distance to a “hot” Fermi distribution characterized by the temperature $T + \Delta T$ measured by the noise thermometry [5].
- [48] H. le Sueur, C. Altimiras, U. Gennser, A. Cavanna, D. Mailly, and F. Pierre, *Phys. Rev. Lett.* **105**, 056803 (2010).
- [49] T.K.T. Nguyen, M.N. Kiselev, and V.E. Kravtsov, *Phys. Rev. B* **82**, 113306 (2010).
- [50] S. Amasha, I. G. Rau, M. Grobis, R. M. Potok, H. Shtrikman, and D. Goldhaber-Gordon, *Phys. Rev. Lett.* **107**, 216804 (2011).
- [51] T. K. T. Nguyen and M. N. Kiselev, *Phys. Rev. B* **97**, 085403 (2018).
- [52] A. V. Andreev and K. A. Matveev, *Phys. Rev. Lett.* **86**, 280 (2001); K. A. Matveev and A. V. Andreev, *Phys. Rev. B* **66**, 045301 (2002).
- [53] M. Fabrizio and A. O. Gogolin, *Phys. Rev. B* **50**, 17732(R) (1994).
- [54] In general, for M species of the electrons, the normal modes are charge, spin and $M-2$ flavour modes. Thus, $M-1$ modes describing spin and all flavours are connected to the Cartan (abelian) subgroup of corresponding $SU(M)$ group [4].
- [55] Supplemental materials are available at (web link will be provided here).
- [56] Importantly, for general MCK, we define the charge mode $\phi_c(x, t) = (\phi_1(x, t) + \dots + \phi_M(x, t))/\sqrt{M}$. Only the charge mode enters the charging energy action. Therefore, the correlator $K_C(\tau) = (\pi^2 T / M \gamma E_C |\sin(\pi T \tau)|)^{2/M}$.
- [57] The temperature dependence of the electric conductance $G \propto (T/E_C)^{2/3}$ given by (9) exactly corresponds to the dynamical Coulomb blockade scaling [46]. This dependence was experimentally observed recently [58]. The scaling (9) is consistent with the orthogonality catastrophe [59–61] explanation given in Ref. [20].
- [58] A. Anthore, Z. Iftikhar, E. Boulat, F.D. Parmentier, A. Cavanna, A. Ouerghi, U. Gennser, and F. Pierre, *Phys. Rev. X* **8**, 031075 (2018).
- [59] P. W. Anderson, *Phys. Rev. Lett.* **18**, 1049 (1967).
- [60] P. Nozières and C. T. de Dominicis, *Phys. Rev.* **178**, 1097 (1969).
- [61] G. D. Mahan, *Many Particle Physics* (Plenum, New York, 1991).
- [62] This scaling is also valid in the strong coupling regime $T \ll T^*$ since Eq. (9) corresponds to the *zero-th order* term of the perturbative expansion in $|r|^2$.
- [63] The sum of two terms in G_T is obviously invariant under rotation of the Cartan basis [54] resulting in renumeration of the QPCs.
- [64] The temperature scaling of G_T for $M \geq 2$ channel Kondo model is, in general, described by $G_T \propto T^{1-2/M+2/M} \log T \sim T \log T$ behaviour and thus, in log approximation does not depend on M (up to M -dependent pre-factor). The $\log T$ behaviour of G_T is associated with the fluctuations of $M-1$ gapless modes (spin and flavours) and is generic for all $M \geq 2$ channel “charge” Kondo models characterized by the NFL behaviour. The gate voltage oscillating dependence is described by $\sin[2\pi N(V_g)]$ factor accounting for the charge quantization. The TP scaling is given by $S^{MCK} \propto T^{1-2/M} \log T$. The TP of all MCK models vanishes at low temperature being however strongly enhanced by $\log T$ factor compared to $S^{1CK} \propto T$ FL scaling.
- [65] E. Sela, A. K. Mitchell, L. Fritz, *Phys. Rev. Lett.* **106**, 147202 (2011).
- [66] A.K. Mitchell, L.A. Landau, L. Fritz, and E. Sela, *Phys. Rev. Lett.* **116**, 157202 (2016).
- [67] L.A. Landau, E. Cornfeld, and E. Sela, *Phys. Rev. Lett.* **120**, 186801 (2018).
- [68] T.K.K. Nguyen and M.N. Kiselev (to be published).
- [69] T.K.T. Nguyen, and M.N. Kiselev, *Phys. Rev. B* **92**, 045125 (2015).
- [70] P.L.S. Lopes, I. Affleck, and E. Sela, arXiv:1911.06336, Y. Komijani, arXiv:1911.08329.
- [71] K. Snizhko, F. Buccheri, R. Egger, and Yuval Gefen, *Phys. Rev. B* **97**, 235139 (2018).
- [72] P. Lecheminant, Alexander O. Gogolin, Alexander A. Nersesyan. *Nucl. Phys. B* **639**, 502 (2002).

**S1. SUPPLEMENTAL MATERIALS TO THE MANUSCRIPT
THERMOELECTRIC TRANSPORT IN A THREE CHANNEL ‘‘CHARGE’’ KONDO CIRCUIT**

Perturbative calculations of thermoelectric coefficient G_T

We present here the details of perturbative calculations accounting for the backscattering processes. We use notations and definitions of the *Letter*. All calculations are performed for 3CK model in the spirits of Matveev-Andreev theory [A. V. Andreev and K. A. Matveev, Phys. Rev. Lett. **86**, 280 (2001); K. A. Matveev and A. V. Andreev, Phys. Rev. B **66**, 045301 (2002)], which concern the saddle-point method. We first evaluate Gaussian integral $Z(\tau) = \int \exp[-S_0 - S_C(\tau) - S'] \prod_\alpha \mathcal{D}\phi_\alpha(x, t)$ under the assumption $S' = 0$. At zero order, the saddle point, based on the principle of the action minimum, is found as

$$\phi_{c,\tau}(x, t) = \frac{\pi N}{\sqrt{3}} - \sqrt{3}TE_C \sum_{\omega_n} \frac{\exp[-|\omega_n x|/v_F]}{|\omega_n| + 3E_C/\pi} n_\tau(\omega_n) e^{-i\omega_n t}, \quad (\text{S1.1})$$

with $\omega_n = 2\pi nT$ are bosonic Matsubara frequencies and the Fourier transform of $n_\tau(t)$ is $n_\tau(\omega_n) = (e^{i\omega_n \tau} - 1)/i\omega_n$.

In the calculation $K_0(\tau) = Z(\tau)/Z(0)$, the integrals over the fluctuations of the field $\phi_c(x, t)$ about the saddle points in the numerator and the denominator cancel each other. Therefore, the value of $K_0(\tau)$ is evaluated by the integrals at the saddle point values. In the condition $\tau \gg E_C^{-1}$ and $T \ll E_C$, we find

$$[S_0 + S_C(\tau)]_{\phi=\phi_{c,\tau}(x,t)} = \frac{3E_C}{2\pi^2 T} \sum_{n=1}^{\infty} \frac{[1 - \cos(2\pi T n \tau)]}{n [n + 3E_C/2\pi^2 T]}, \quad (\text{S1.2})$$

and $[S_0 + S_C(\tau)]_{\phi=\phi_{c,0}(x,t)} = 0$. The correlator $K_0(\tau)$

$$K_0(\tau) = \left[\frac{\pi^2 T}{3\gamma E_C} \frac{1}{|\sin(\pi T \tau)|} \right]^{2/3}. \quad (\text{S1.3})$$

Plugging in the formula (S1.3) into equation (6) we find the electric conductance as a function of the temperature:

$$G = \frac{G_L \pi^{11/6} \Gamma(4/3)}{2(3\gamma)^{2/3} \Gamma(11/6)} \left[\frac{T}{E_C} \right]^{2/3}. \quad (\text{S1.4})$$

Substituting formula (S1.3) into equation (7) we find that the thermoelectric coefficient G_T vanishes. At zero order of perturbation theory, the PH symmetry is conserved (as explained in the main text). One needs to proceed the perturbation calculation at the first non-vanishing order. To proceed we express S' in terms of the products of charge mode, spin mode, and flavour mode. However, only the fluctuations of the charge mode is suppressed at low frequencies by the charging energy term. The fluctuations of the spin mode and flavour mode are not suppressed by the charging energy. Therefore, the average $\langle S' \rangle = 0$.

At second order of perturbation theory,

$$K(\tau) = K_C(\tau) \left[1 + \frac{1}{2} \left(\langle S'^2 \rangle_\tau - \langle S'^2 \rangle_0 \right) \right] \quad (\text{S1.5})$$

and therefore it is necessary to calculate $\langle S'^2 \rangle_\tau$.

$$\begin{aligned} \langle S'^2 \rangle_\tau = & \frac{D^2}{\pi^2} |r|^2 \int_0^\beta dt \int_0^\beta dt' \left\langle \cos \left[\frac{2}{\sqrt{3}} \phi_c(t) - \frac{2\sqrt{2}}{\sqrt{3}} \phi_f(t) \right] \cos \left[\frac{2}{\sqrt{3}} \phi_c(t') - \frac{2\sqrt{2}}{\sqrt{3}} \phi_f(t') \right] \right\rangle \\ & + 4 \left\langle \cos \left[\frac{2}{\sqrt{3}} \phi_c(t) + \frac{\sqrt{2}}{\sqrt{3}} \phi_f(t) \right] \cos \left[\frac{2}{\sqrt{3}} \phi_c(t') + \frac{\sqrt{2}}{\sqrt{3}} \phi_f(t') \right] \right\rangle \left\langle \cos [\sqrt{2} \phi_s(t)] \cos [\sqrt{2} \phi_s(t')] \right\rangle \end{aligned} \quad (\text{S1.6})$$

where we use shorthand notations $\phi_\alpha(t) \equiv \phi_\alpha(0, t)$. Since charge, spin, flavour modes are independent, we decouple them as

$$\begin{aligned} \langle S'^2 \rangle_\tau = & \frac{D^2}{2\pi^2} |r|^2 \int_0^\beta dt \int_0^\beta dt' \left\{ \Re \left[\kappa_c^+(t, t', \tau) \tilde{\kappa}_f^+(t, t', \tau) + \kappa_c^-(t, t', \tau) \tilde{\kappa}_f^-(t, t', \tau) \right] \right. \\ & \left. + 2\Re \left[\kappa_c^+(t, t', \tau) \kappa_f^+(t, t', \tau) + \kappa_c^-(t, t', \tau) \kappa_f^-(t, t', \tau) \right] \Re \left[\kappa_s^+(t, t', \tau) + \kappa_s^-(t, t', \tau) \right] \right\}, \end{aligned} \quad (\text{S1.7})$$

with $\kappa_j^\pm(t, t', \tau) \equiv \langle \exp [ia_j [\phi_j(t) \pm \phi_j(t')]] \rangle$. We apply the saddle-point method one more time in order to perform the integrals in the formula (S1.7)

$$\kappa_j^\pm(t, t', \tau) = \exp [ia_j (\phi_{j\tau}(t) \pm \phi_{j\tau}(t'))] \exp [-a_j^2 (\langle \varphi_j^2(t) \rangle \pm \langle \varphi_j(t) \varphi_j(t') \rangle)], \quad (\text{S1.8})$$

with $\phi_j(t) = \phi_{j\tau}(t) + \varphi_j(t)$. Calculation of the correlator $\langle \varphi_j(t) \varphi_j(t') \rangle$ is achieved by using generating functional method as follows.

To evaluate the correlator (S1.8) we introduce the generating functional

$$W[\{J_j(\omega_n)\}] = \left\langle \exp \left[-T \sum_{\omega_n} J_j(\omega_n) \varphi_j(-\omega_n) \right] \right\rangle. \quad (\text{S1.9})$$

The value of the Gaussian integral is completely defined by the saddle point

$$W[\{J_j(\omega_n)\}] = \exp \left[-\frac{T}{2} \sum_{\omega_n} J_j(\omega_n) \varphi_j^J(-\omega_n) \right], \quad (\text{S1.10})$$

in which $\varphi_j^J(t)$ is the saddle-point value of the field φ_j .

Now, we consider the charge mode: the fluctuations of $\varphi_j(t)$ coincide with those of $\phi_c(0, t)$ at $N = 0$, $n_\tau = 0$ and $n_\tau(t)$ plays the role of a source term similar to $J_c(t)$ as $J_c(t) = 2\sqrt{3}E_C n_\tau(t) / \pi$, or $n_\tau(\omega_n) = \pi J_c(\omega_n) / 2\sqrt{3}E_C$. We re-write the generating functional as follows:

$$W[\{J_c(\omega_n)\}] = \exp \left[\frac{\pi T}{4} \sum_{\omega_n} \frac{J_c(\omega_n) J_c(-\omega_n)}{|\omega_n| + \frac{3E_C}{\pi}} \right]. \quad (\text{S1.11})$$

The correlator $\langle \varphi_c(-\omega_n) \varphi_c(\omega_m) \rangle$ can be obtained by differentiating the functional W in formula (S1.9) with respect to $J_c(\omega_n)$ and $J_c(-\omega_m)$. In time representation, this correlator is written as

$$\langle \varphi_c(t) \varphi_c(t') \rangle = \frac{\pi T}{2} \sum_{\omega_n} \frac{e^{i\omega_n(t-t')} e^{-|\omega_n|/D}}{|\omega_n| + \frac{3E_C}{\pi}}.$$

At the limits we are interested in, this correlator behaves as

$$\langle \varphi_c(t) \varphi_c(t') \rangle = \begin{cases} \frac{1}{2} \ln \frac{\pi D}{3\gamma E_C \sqrt{1+[D(t-t')]^2}}, & |t-t'| \ll E_C^{-1} \\ \frac{\pi^4 T^2}{2(3E_C)^2 \sin^2[\pi T(t-t')]}, & |t-t'| \gg E_C^{-1}. \end{cases} \quad (\text{S1.12})$$

Therefore, at $T \ll E_C$, we obtain

$$\kappa_c^+(t, t', \tau) \approx \left[\frac{3\gamma E_C}{\pi D} \right]^{\frac{2}{3}} e^{i\frac{4\pi N}{3}} e^{-i\frac{2}{3}\chi_\tau(t)} e^{-i\frac{2}{3}\chi_\tau(t')}, \quad (\text{S1.13})$$

and

$$\kappa_c^-(t, t', \tau) \approx \left[\frac{3\gamma E_C}{\pi D} \right]^{\frac{2}{3}} e^{-i\frac{2}{3}\chi_\tau(t)} e^{+i\frac{2}{3}\chi_\tau(t')}, \quad (\text{S1.14})$$

with

$$\sqrt{3}\chi_\tau(t) = \frac{3E_C}{4\pi^2 T} \sum_{n=-\infty}^{\infty} \frac{e^{-i2\pi T n(t-\tau)} - e^{-i2\pi T n t}}{in \left(|n| + \frac{3E_C}{2\pi^2 T} \right)} = [\pi n_\tau(t) + \delta\chi_\tau(t)], \quad (\text{S1.15})$$

$$\begin{aligned} \delta\chi_\tau(t) &= \sum_{n=1}^{\infty} \frac{\sin[2\pi T n(t-\tau)] - \sin[2\pi T n t]}{n + \frac{3E_C}{2\pi^2 T}} \\ &\approx \frac{\pi^2 T}{3E_C} \{ \cot[\pi T(t-\tau)] - \cot[\pi T t] \}. \end{aligned} \quad (\text{S1.16})$$

Similarly, we consider flavour mode, we find that if there exists a ‘‘flavouring energy term’’ E_f and a ‘‘spinning energy term’’ E_s (in the same meaning as charging energy term E_C), the relation between $n_\tau(t)$ and $J_{f/s}(t)$ should be $J_f(t) = 2\sqrt{6}E_f n_\tau(t)/\pi$ and $J_s(t) = 2\sqrt{2}E_s n_\tau(t)/\pi$. However, after obtaining the correlator $\langle \varphi_{s/f}(t) \varphi_{s/f}(t') \rangle$, we need to take the limits $E_{s/f} = 0$. At the end, we have

$$\begin{aligned} \langle \varphi_s(t) \varphi_s(t') \rangle &= \langle \varphi_f(t) \varphi_f(t') \rangle = \frac{\pi T}{2} \sum_{\omega_n} \frac{e^{i\omega_n(t-t')} e^{-|\omega_n|/D}}{|\omega_n|} \\ &= -\frac{1}{4} \ln \left[1 + e^{-\frac{4\pi T}{D}} - 2e^{-\frac{2\pi T}{D}} \cos[2\pi T(t-t')] \right]. \end{aligned} \quad (\text{S1.17})$$

We obtain the result as shown in the second line in the limit $T \ll D$.

$$\kappa_s^-(t, t', \tau) \approx \left[\frac{\pi T}{D} \right] \frac{1}{|\sin(\pi T(t-t'))|}, \quad (\text{S1.18})$$

$$\kappa_f^-(t, t', \tau) \approx \left[\frac{\pi T}{D} \right]^{\frac{1}{3}} \frac{1}{|\sin(\pi T(t-t'))|^{\frac{1}{3}}}, \quad (\text{S1.19})$$

$$\tilde{\kappa}_f^-(t, t', \tau) \approx \left[\frac{\pi T}{D} \right]^{\frac{4}{3}} \frac{1}{|\sin(\pi T(t-t'))|^{\frac{4}{3}}}. \quad (\text{S1.20})$$

If an electron comes from QPC2, we only take into account the first term in Eq. (S1.7) and $\phi_{s\tau}(x, t) = 0$, $\phi_{f\tau}(x, t) = -2\pi N/\sqrt{6}$ then

$$\tilde{\kappa}_f^+(t, t', \tau) \approx e^{i\frac{8\pi N}{3}} \left[\frac{4\pi T}{D} \right]^{\frac{4}{3}} |\sin(\pi T(t-t'))|^{\frac{4}{3}}. \quad (\text{S1.21})$$

As we have discussed in the main text, G_T must be an odd function of the gate voltage N as well as odd function of τ . The odd N -dependent term is

$$\langle S'^2 \rangle_{\tau, \text{odd } N} \approx 4 \left[\frac{6\gamma E_C}{\pi^2} \right]^{\frac{2}{3}} T^{\frac{4}{3}} |r|^2 \sin[4\pi N] \int_0^\beta dt \sin \left[\frac{2}{3} \chi_\tau(t) \right] \int_0^\beta dt' \cos \left[\frac{2}{3} \chi_\tau(t') \right] |\sin(\pi T(t-t'))|^{\frac{4}{3}}. \quad (\text{S1.22})$$

We find that

$$\cos \left[\frac{2}{3} \chi_\tau(t') \right] = \cos \left[\frac{2}{3} \pi n_\tau(t') + \frac{2}{3} \delta \chi_\tau(t') \right] \approx \begin{cases} -\frac{1}{2} - \frac{\sqrt{3}}{3} \delta \chi_\tau(t'), & \text{if } 0 \leq t' \leq \tau, \\ 1, & \text{if } \tau < t' \leq \beta, \end{cases} \quad (\text{S1.23})$$

$$\sin \left[\frac{2}{3} \chi_\tau(t) \right] = \sin \left[\frac{2}{3} \pi n_\tau(t) + \frac{2}{3} \delta \chi_\tau(t) \right] \approx \begin{cases} \frac{\sqrt{3}}{2} - \frac{1}{3} \delta \chi_\tau(t), & \text{if } 0 \leq t \leq \tau, \\ \frac{2}{3} \delta \chi_\tau(t), & \text{if } \tau < t \leq \beta. \end{cases} \quad (\text{S1.24})$$

The function $\delta \chi_\tau(t)$ as shown in formula (S1.16) makes the integrals to diverge logarithmically. At $|t-t'| \gg E_C^{-1}$ and $T \ll E_C$, we take into account only the terms which contain the first order in $\delta \chi_\tau(t)$, we obtain

$$\langle S'_2 \rangle_{\tau, \text{odd } N} \approx -\frac{8(2-\sqrt{3})}{3} \left[\frac{6\gamma}{\pi^2} \right]^{\frac{2}{3}} |r|^2 \left[\frac{T}{E_C} \right]^{\frac{1}{3}} \ln \left[\frac{E_C}{T} \right] \sin[4\pi N] F_2(\tau), \quad (\text{S1.25})$$

with

$$F_2(t) = \cos(\pi T t) \left[F_2^1 \left[\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, \cos^2(\pi T t) \right] + 3 |\sin(\pi T t)|^{\frac{1}{3}} \right]. \quad (\text{S1.26})$$

The second order contribution to the correlator $K(\tau)$ corresponding to this process is

$$K_{\text{odd } N}(\tau) \approx -\frac{2^{\frac{2}{3}} 4(2-\sqrt{3})}{3} |r|^2 \frac{T}{E_C} \ln \left[\frac{E_C}{T} \right] \sin[4\pi N] \frac{F_2(\tau)}{|\sin(\pi T \tau)|^{\frac{2}{3}}}. \quad (\text{S1.27})$$

The equation for the thermoelectric coefficient G_T accounting for an electron coming from QPC2 is given by

$$G_T = C_2 \frac{G_L}{e} |r|^2 \frac{T}{E_C} \ln \left[\frac{E_C}{T} \right] \sin [4\pi N], \quad (\text{S1.28})$$

with

$$C_2 = \frac{2^{\frac{2}{3}} 2 (2 - \sqrt{3})}{3} \pi \int_{-\infty}^{\infty} \frac{\sinh^2(x) \left[F_2^1 \left[\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, -\sinh^2(x) \right] + 3 \cosh^{\frac{1}{3}}(x) \right]}{\cosh^{\frac{11}{3}}(x)} dx = 0.936794. \quad (\text{S1.29})$$

If an electron comes from QPC1 or QPC3, we only take into account the second term in Eq. (S1.7) and $\phi_{s\tau}(x, t) = \pm\pi N/\sqrt{2}$, $\phi_{f\tau}(x, t) = \pi N/\sqrt{6}$ then

$$\kappa_s^+(t, t', \tau) \approx e^{i2\pi N} \left[\frac{4\pi T}{D} \right] |\sin(\pi T(t - t'))|, \quad (\text{S1.30})$$

$$\kappa_f^+(t, t', \tau) \approx e^{i\frac{2\pi N}{3}} \left[\frac{4\pi T}{D} \right]^{\frac{1}{3}} |\sin(\pi T(t - t'))|^{\frac{1}{3}}. \quad (\text{S1.31})$$

Plugging all $\kappa_j^\pm(t, t', \tau)$ into formula (S1.7), we obtain the temperature scaling of $\langle S'^2 \rangle_{\tau, odd N}$. We find that only two terms $2\Re \left[\kappa_c^+(t, t', \tau) \kappa_f^+(t, t', \tau) \right] \Re [\kappa_s^+(t, t', \tau)]$ and $2\Re \left[\kappa_c^+(t, t', \tau) \kappa_f^+(t, t', \tau) \right] \Re [\kappa_s^-(t, t', \tau)]$ contribute to G_T . The first one contributes to G_T the same result as shown in Eq. (S1.28). Let us illustrate the calculations for the latter term:

$$\langle S'^2 \rangle_{\tau, odd N} \approx 2 \left[\frac{6\gamma E_C}{\pi^2} \right]^{\frac{2}{3}} T^{\frac{4}{3}} |r|^2 \sin [2\pi N] \int_0^\beta dt \sin \left[\frac{2}{3} \chi_\tau(t) \right] \int_0^\beta dt' \cos \left[\frac{2}{3} \chi_\tau(t') \right] |\sin(\pi T(t - t'))|^{-\frac{2}{3}}. \quad (\text{S1.32})$$

The function $|\sin(\pi T(t - t'))|^{-\frac{2}{3}}$ exhibits *integrable* power-law divergence at $t' \rightarrow t$. We calculate the integrals in Eq. (S1.32) in the same way as we did for Eq. (S1.22). At the end, we obtain

$$\langle S'^2 \rangle_{\tau, odd N} \approx -\frac{10}{3} \left[\frac{6\gamma}{\pi^2} \right]^{\frac{2}{3}} \left[\frac{T}{E_C} \right]^{\frac{1}{3}} |r|^2 \sin [2\pi N] \ln \left[\frac{E_C}{T} \right] F_1(\tau), \quad (\text{S1.33})$$

with

$$F_1(t) = \cos(\pi T t) F_2^1 \left[\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, \cos^2(\pi T t) \right]. \quad (\text{S1.34})$$

We plug formula (S1.33) into formula (S1.5) and obtain the correlator $K(\tau)$ at second order as

$$K_{odd N}(\tau) \approx -\frac{2^{\frac{2}{3}} 5 |r|^2 T}{3 E_C} \ln \left[\frac{E_C}{T} \right] \sin [2\pi N] \frac{F_1(\tau)}{|\sin(\pi T \tau)|^{\frac{2}{3}}}. \quad (\text{S1.35})$$

From (7) and (S1.35), we obtain the thermoelectric coefficient G_T which is contributed by $2\Re \left[\kappa_c^+(t, t', \tau) \kappa_f^+(t, t', \tau) \right] \Re [\kappa_s^-(t, t', \tau)]$. At the end, thermoelectric coefficient G_T accounting for an electron coming from QPC1 or QPC3 is given by

$$G_T = C_1 (1 + a \cos[2\pi N]) \frac{G_L}{e} |r|^2 \frac{T}{E_C} \ln \left[\frac{E_C}{T} \right] \sin [2\pi N], \quad (\text{S1.36})$$

with $a = C_2/C_1$,

$$C_1 = \frac{5}{6} 2^{\frac{2}{3}} \pi \int_{-\infty}^{\infty} \frac{\sinh^2(x) F_2^1 \left[\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, -\sinh^2(x) \right]}{\cosh^{\frac{11}{3}}(x)} dx = 0.979528. \quad (\text{S1.37})$$

We demonstrated that the temperature scaling of G_T and TP is given by $T \log T$ and $T^{1/3} \log T$ correspondingly.