

Double Circulant Self-Dual Codes From Generalized Cyclotomic Classes of Order Two

Wenpeng Gao¹ and Tongjiang Yan²

(Corresponding author: Tongjiang Yan)

College of Sciences, China University of Petroleum¹
Qingdao, 266555, China

College of Sciences, China University of Petroleum; Shandong Provincial Key Laboratory of Computer Networks²
Key Laboratory of Applied Mathematics(Putian University), Fujian Province University²
Qingdao, 266555, China; Jinan, 250014, China; Fujian Putian, 351100, China

(Email: yantoji@163.com)

(Received 2019; revised and accepted 2019)

Abstract

In this paper, constructions of some double circulant self-dual codes by generalized cyclotomic classes of order two are presented. This technique is applied to [72, 36, 12] binary highest know self-dual codes to obtain self-dual codes over GF(2) and [32, 16, 8] almost optimal self-dual codes over GF(4). Based on the properties of generalized cyclotomy, some of these codes can be proved to possess good minimum weights.

Keywords: Double circulant code; Generalized cyclotomic classes; Self-Dual code

1 Introduction

Self-dual codes have important applications in transmission [12]. Constructing self-dual codes which have good minimum weight has been an important research problem [1]. And a number of papers devoted to constructing self-dual codes [2] [3]. Most of self-dual codes which are double circulant codes have high minimum distance [11]. The generalized cyclotomic classes have wide applications in constructing sequences [9,13,20], cyclic codes [7] [8] and difference sets [10]. There are a number of results about generalized cyclotomic classes of order two [5] [19]. Motivated by the constructions of double circulant codes by cyclotomic classes of order four [18], we give constructions of double circulant codes by generalized cyclotomic classes of order two. All computations have been done by MAGMA V2.20-4 [4] on a 3.40 GHz CPU.

This paper is organized as follows. In Section II, we present definitions and some preliminaries about self-dual codes and generalized cyclotomic classes. In Section III, three infinite families of self-dual codes are given. Section IV concludes the paper.

2 Preliminaries

2.1 Self-Dual codes

A linear code C of length n and dimension k over finite field $\text{GF}(l)$ is a k -dimensional subspace of $\text{GF}(l)^n$, where l is a prime power. A generator matrix G of the code C is a $k \times n$ matrix over $\text{GF}(l)$. The elements of C are called codewords. The Euclidean inner product is defined by

$$(x, y) = \sum_{i=1}^n x_i y_i,$$

where $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$. The Euclidean dual code, denoted by C^\top , of linear code C is defined by

$$C^\top = \{x \in \text{GF}(l)^n \mid (x, c) = 0, \text{ for all } c \in C\}.$$

And we have $\dim(C) + \dim(C^\top) = n$. C is said to be Euclidean self-orthogonal if $C \subset C^\top$ and Euclidean self-dual if $C = C^\top$. If C is dual-self code, the dimension of C is $n/2$. From now on, what we mean by self-dual is Euclidean self-dual.

2.2 Hamming Distance

The Hamming distance of codewords $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$, denoted by $d(x, y)$, is defined to be the number of places at which x and y differ. The Hamming weight $wt(x)$ of a codeword $x = (x_1, \dots, x_n)$ is $wt(x) = d(x, 0)$ and the minimum Hamming distance $d(C)$ of a code C is equal to the minimum nonzero Hamming weight of all codewords in C . Then for the self-dual codes, we have the following result.

Theorem 1. [15] [16] *Let C be a self-dual code over $\text{GF}(l)$ of length n and minimum Hamming distance $d(C)$. Then we have*

- (i) If $l = 2$, then

$$d(C) \leq \begin{cases} 4 \lfloor \frac{n}{24} \rfloor + 4, & \text{if } n \not\equiv 22 \pmod{24}; \\ 4 \lfloor \frac{n}{24} \rfloor + 6, & \text{if } n \equiv 22 \pmod{24}. \end{cases}$$

(ii) If $l = 3$, then $d(C) \leq 3 \lfloor \frac{n}{12} \rfloor + 3$.

(iii) If $l = 4$, then $d(C) \leq 4 \lfloor \frac{n}{12} \rfloor + 4$.

(iv) $d(C) \leq \lfloor \frac{n}{2} \rfloor + 1$ for $l \neq 2, 3, 4$.

The code C is called extremal if the above equality holds. A self-dual code is called optimal if it has the highest possible minimum distance for its length.

2.3 Generalized Cyclotomy and Generalized Cyclotomic Number

Let $n = pq$, where p and q are distinct odd primes with $\gcd(p-1, q-1) = d$. The Chinese Remainder Theorem [6] guarantees that there exists common primitive roots of both p and q . Let g be a fixed common primitive root of both p and q . Let $e := \text{ord}_n(g)$ denote the multiplicative order of g modulo n , then

$$\begin{aligned} \text{ord}_n(g) &= \text{lcm}(\text{ord}_p(g), \text{ord}_q(g)) \\ &= \text{lcm}(p-1, q-1) = \frac{(p-1)(q-1)}{d}. \end{aligned}$$

Let x be an integer satisfying

$$x \equiv g \pmod{p}, \quad x \equiv 1 \pmod{q}.$$

Whiteman [17] defined the generalized cyclotomic classes

$$C_i = \{g^s x^i : s = 0, 1, \dots, e-1\}, i = 0, 1, \dots, d-1.$$

In this paper, we shall always assume that p, q are distinct primes with $\gcd(p-1, q-1) = 2$.

$$\begin{aligned} P &= \{p, 2p, \dots, (q-1)p\}, \\ Q &= \{q, 2q, \dots, (p-1)q\}, \\ R &= \{0\}. \end{aligned}$$

The generalized cyclotomic numbers corresponding of order two are defined by

$$(i, j) = |(C_i + 1) \cap C_j|, \text{ for all } i, j = 0, 1.$$

Lemma 1. [5] $-1 \in C_0$ if $(p-1)(q-1)/4$ is even; $-1 \in C_1$ if $(p-1)(q-1)/4$ is odd.

Lemma 2. [17] If $(p-1)(q-1)/4$ is even,

$$\begin{aligned} (0, 0) &= (1, 0) = (1, 1) = \frac{(p-2)(q-2) + 1}{4}, \\ (0, 1) &= \frac{(p-2)(q-2) - 3}{4}. \end{aligned}$$

If $(p-1)(q-1)/4$ is odd,

$$\begin{aligned} (0, 0) &= \frac{(p-2)(q-2) + 3}{4}, \\ (0, 1) &= (1, 0) = (1, 1) = \frac{(p-2)(q-2) - 1}{4}. \end{aligned}$$

2.4 General Results

Let m_0, m_1, m_2, m_3 and m_4 be elements of $\text{GF}(l)$. The matrix $C_n(m_0, m_1, m_2, m_3, m_4)$ is the $n \times n$ matrix on $\text{GF}(l)$ with components c_{ij} , $1 \leq i, j \leq n$,

$$c_{ij} = \begin{cases} m_0 & \text{if } j = i, \\ m_1 & \text{if } j - i \in P, \\ m_2 & \text{if } j - i \in Q, \\ m_3 & \text{if } j - i \in C_0, \\ m_4 & \text{if } j - i \in C_1, \end{cases}$$

Define by I_n and J_n the identity and the all-one square $n \times n$ matrices. Then $C_n(1, 0, 0, 0, 0) = I_n$ and $C_n(1, 1, 1, 1, 1) = J_n$. Denote $P_n := C_n(0, 1, 0, 0, 0)$, $Q_n := C_n(0, 0, 1, 0, 0)$, $A_1 := C_n(0, 0, 0, 1, 0)$ and $A_2 := C_n(0, 0, 0, 0, 1)$.

Lemma 3. If p is a prime of the form $4\omega + 1$ and q is a prime of the form $4\omega' + 3$. Then

$$\begin{aligned} A_1 &= A_2^\top, A_2 = A_1^\top, P_n = P_n^\top, Q_n = Q_n^\top, \\ P_n A_1 &= A_1 P_n = Q_n + A_2, P_n A_2 = A_2 P_n = Q_n + A_1, \\ Q_n A_1 &= A_1 Q_n = A_1, Q_n A_2 = A_2 Q_n = A_2, \\ P_n Q_n &= Q_n P_n = A_1 + A_2, P_n^2 = P_n, Q_n^2 = Q_n, \\ A_1^2 &= (1, 0)A_1 + (0, 1)A_2, \\ A_2^2 &= (0, 1)A_1 + (1, 0)A_2, \\ A_1 A_2 &= Q_n + (0, 0)A_1 + (1, 1)A_2, \\ A_2 A_1 &= Q_n + (1, 1)A_1 + (0, 0)A_2. \end{aligned}$$

Proof. The proof is straightforward from the definitions of these sets. \square

Lemma 4. If p is a prime of the form $4\omega + 3$ and q is a prime of the form $4\omega' + 1$. Then

$$\begin{aligned} A_1 &= A_2^\top, A_2 = A_1^\top, P_n = P_n^\top, Q_n = Q_n^\top, \\ P_n A_1 &= A_1 P_n = A_1, P_n A_2 = A_2 P_n = A_2, \\ Q_n A_1 &= A_1 Q_n = P_n + A_2, Q_n A_2 = A_2 Q_n = P_n + A_1, \\ P_n Q_n &= Q_n P_n = A_1 + A_2, P_n^2 = P_n, Q_n^2 = Q_n, \\ A_1^2 &= (1, 0)A_1 + (0, 1)A_2, \\ A_2^2 &= (0, 1)A_1 + (1, 0)A_2, \\ A_1 A_2 &= P_n + (0, 0)A_1 + (1, 1)A_2, \\ A_2 A_1 &= P_n + (1, 1)A_1 + (0, 0)A_2. \end{aligned}$$

Proof. The proof is straightforward from the definitions of these sets. \square

The following result will be needed in the sequel.

$$\begin{aligned} &(m_0 I_n + m_1 P_n + m_2 Q_n + m_3 A_1 + m_4 A_2) \\ &(m_0 I_n + m_1 P_n + m_2 Q_n + m_3 A_1 + m_4 A_2)^\top \\ &= a_0 I_n + a_1 P_n + a_2 Q_n + a_3 A_1 + a_4 A_2. \end{aligned}$$

Lemma 5. *If $p = 4\omega + 1$ and $q = 4\omega' + 3$. Then*

$$\begin{aligned} a_0 &= m_0^2, \\ a_1 &= m_1^2 + 2m_0m_1, \\ a_2 &= m_2^2 + m_3^2 + m_4^2 + 2m_0m_2 + 2m_1(m_3 + m_4), \\ a_3 &= a_4 \\ &= (m_0 + m_1 + m_2)(m_3 + m_4) + 2m_1m_2 \\ &\quad + (0, 0)(m_3^2 + m_4^2) + ((1, 0) + (0, 1))m_3m_4. \end{aligned}$$

If $p = 4\omega + 3$ and $q = 4\omega' + 1$. Then

$$\begin{aligned} a_0 &= m_0^2, \\ a_1 &= m_1^2 + m_3^2 + m_4^2 + 2m_0m_1 + 2m_2(m_3 + m_4), \\ a_2 &= m_2^2 + 2m_0m_2, \\ a_3 &= a_4 \\ &= (m_0 + m_1 + m_2)(m_3 + m_4) + 2m_1m_2 \\ &\quad + ((1, 0) + (0, 1))m_3m_4 + (0, 0)(m_3^2 + m_4^2). \end{aligned}$$

For convenience, we denote

$$\begin{aligned} \vec{m} &:= (m_0, m_1, m_2, m_3, m_4) \in GF(l)^5, \\ D_0(\vec{m}) &:= a_0, D_1(\vec{m}) := a_1, \\ D_2(\vec{m}) &:= a_2, D_3(\vec{m}) := a_3. \end{aligned}$$

Definition 1. [14] *Let $GP_n(R)$ and $GB_n(\alpha, R)$ be codes with generator matrices of the form*

$$(I_n, R) \begin{pmatrix} \alpha & 1 & \cdots & 1 \\ -1 & & & \\ I_n & \vdots & & R \\ -1 & & & \end{pmatrix},$$

where $\alpha \in GF(l)$ and R is an $n \times n$ circulant matrix. The codes $GP_n(R)$ and $GB_n(\alpha, R)$ are called **pure double circulant codes** and **bordered double circulant codes**, respectively.

For convenience, Let $GP_n(\vec{m})$ be generator matrix of $GP_n(R)$ and $GB_n(\alpha, \vec{m})$ be generator matrix of $GB_n(\alpha, R)$.

$$\begin{aligned} GP_n(\vec{m}) &:= GP_n(m_0I_n + m_1P_n + m_2Q_n + m_3A_1 + m_4A_2), \\ GB_n(\alpha, \vec{m}) &:= GB_n(\alpha, m_0I_n + m_1P_n + m_2Q_n + m_3A_1 + m_4A_2). \end{aligned}$$

The main theorem of this section is:

Theorem 2. *Let $\alpha \in GF(l)$ and $\vec{m} \in GF(l)^5$. Then*

- 1) the code with generator matrix $GP_n(\vec{m})$ is self-dual over $GF(l)$ if and only if the following holds:
 - a) $D_0(\vec{m}) = -1$,
 - b) $D_1(\vec{m}) = 0$,

- c) $D_2(\vec{m}) = 0$,
- d) $D_3(\vec{m}) = 0$,
- e) $D_4(\vec{m}) = 0$.

- 2) the code with generator matrix $GB_n(\alpha, \vec{m})$ is self-dual over $GF(l)$ if and only if the following holds:

- a) $\alpha + n = -1$,
- b) $-\alpha + m_0 + (p-1)m_1 + (q-1)m_2 + \frac{(p-1)(q-1)}{2}(m_3 + m_4) = 0$,
- c) $D_0(\vec{m}) = -2$,
- d) $D_1(\vec{m}) = -1$,
- e) $D_2(\vec{m}) = -1$,
- f) $D_3(\vec{m}) = -1$.

Proof. The result follows from

$$\begin{aligned} GP_n(\vec{m})GP_n(\vec{m})^\top &= I_n + D_0I_n + D_1P_n + D_2Q_n + D_3A_1 + D_4A_2, \end{aligned}$$

and

$$\begin{aligned} GB_n(\alpha, \vec{m})GB_n(\alpha, \vec{m})^\top &= I_{n+1} + \begin{pmatrix} \alpha + n & S \cdots S \\ S & \\ \vdots & X \\ S & \end{pmatrix}, \end{aligned}$$

where $X = J_n + D_0(\vec{m})I_n + D_1(\vec{m})P_n + D_2(\vec{m})Q_n + D_3(\vec{m})A_1 + D_4(\vec{m})A_2$ and $S = -\alpha + m_0 + (p-1)m_1 + (q-1)m_2 + \frac{(p-1)(q-1)}{2}(m_3 + m_4)$. \square

3 Double circulant self-dual codes over fields

In this section, we construct two infinite families of self-dual codes over $GF(2)$ and $GF(4)$. As a preparation, we have the following two lemmas.

Lemma 6. *Let $p = 4\omega + 3$ and $q = 4\omega' + 1$ or $p = 4\omega + 1$ and $q = 4\omega' + 3$. Then*

- 1) *If $\omega + \omega'$ is even, which means $\frac{p+q}{4}$ is odd,*
 $(0, 0) \equiv (1, 0) \equiv (1, 1) \equiv 0 \pmod{2}, (0, 1) \equiv 1 \pmod{2}.$
- 2) *If $\omega + \omega'$ is odd, which means $\frac{p+q}{4}$ is even,*
 $(0, 0) \equiv (1, 0) \equiv (1, 1) \equiv 1 \pmod{2}, (0, 1) \equiv 0 \pmod{2}.$

Proof. Let $p = 4\omega + 3$ and $q = 4\omega' + 1$. Then

$$\begin{aligned} (0, 0) &= \frac{(p-2)(q-2)+1}{4} \\ &= \frac{(4\omega+1)(4\omega'-1)+1}{4} \\ &= \frac{16\omega\omega' + 4\omega + 4\omega'}{4} \\ &= 4\omega\omega' + \omega + \omega' \\ &\equiv \omega' + \omega \pmod{2} \end{aligned}$$

$$\begin{aligned}
(0, 1) &= \frac{(p-2)(q-2)-3}{4} \\
&= \frac{(4\omega+1)(4\omega'-1)-3}{4} \\
&= \frac{16\omega\omega' - 4\omega + 4\omega' - 4}{4} \\
&= 4\omega\omega' - \omega + \omega' - 1 \\
&\equiv \omega' + \omega + 1 \pmod{2}
\end{aligned}$$

The proof for the condition of $p = 4\omega + 1$ and $q = 4\omega' + 3$ is similar. \square

3.1 Self-Dual Codes Over GF(2)

Theorem 3. Let $\frac{(p-1)(q-1)}{4}$ be even and $\frac{p+q}{4}$ be odd. Then the following holds:

1) $p = 4\omega + 1$ and $q = 4\omega' + 3$

The codes with generator matrices $GP_n(1, 0, 1, 0, 1)$ and $GP_n(1, 0, 1, 1, 0)$ over GF(2) are self-dual of length $2n$.

The codes with generator matrices $GB_n(0, 0, 1, 0, 1, 0)$ and $GB_n(0, 0, 1, 0, 0, 1)$ over GF(2) are self-dual of length $2n + 2$.

2) $p = 4\omega + 3$ and $q = 4\omega' + 1$

The codes with generator matrices $GP_n(1, 1, 0, 0, 1)$ and $GP_n(1, 1, 0, 1, 0)$ over GF(2) are self-dual of length $2n$.

The codes with generator matrices $GB_n(0, 0, 0, 1, 1, 0)$ and $GB_n(0, 0, 0, 1, 0, 1)$ over GF(2) are self-dual of length $2n + 2$.

Table 1: SOME CODES OVER GF(2)

Code	p	q	Construction	Comments
[70, 35, 10]	5	7	$GP_{35}(1, 0, 1, 0, 1)$	almost optimal
[72, 36, 12]	5	7	$GB_{36}(0, 0, 1, 0, 1, 0)$	highest know

Proof. If $p = 4\omega + 1$ and $q = 4\omega' + 3$. Then

$$\begin{aligned}
\alpha + n &= 4(4\omega\omega' + \omega + 3\omega') + 3 \equiv 1 \pmod{2}, \\
-\alpha + m_0 + (p-1)m_1 + (q-1)m_2 \\
&+ \frac{(p-1)(q-1)}{2}(m_3 + m_4) \equiv 0 \pmod{2},
\end{aligned}$$

$$D_0(0, 1, 0, 1, 0) \equiv 0 \pmod{2},$$

$$D_1(0, 1, 0, 1, 0)$$

$$= D_2(0, 1, 0, 1, 0) \equiv 1 \pmod{2},$$

$$D_3(0, 1, 0, 1, 0)$$

$$= (0, 0)(m_3^2 + m_4^2) + ((1, 0) + (0, 1))m_3m_4$$

$$= 0 \times (0 + 1) + (1 + 0) \equiv 1 \pmod{2}.$$

By Theorem 2, the code with generator matrix $GB_n(0, 0, 1, 0, 1, 0)$ over GF(2) is self-dual of length $2n + 2$. The proof that the code with generator matrix $GB_n(0, 0, 0, 1, 0, 1)$ over GF(2) is self-dual is similar.

Under the same conditions, we have

$$D_0(1, 0, 1, 0, 1) \equiv 1 \pmod{2},$$

$$D_1(1, 0, 1, 0, 1) = D_2(1, 0, 1, 0, 1) \equiv 0 \pmod{2},$$

$$D_3(1, 0, 1, 0, 1) = (0, 0)(m_3^2 + m_4^2) + ((1, 0) + (0, 1))m_3m_4 = 0 \times (0 + 1) + 0 \times (1 + 0) \equiv 0 \pmod{2}.$$

Thus the code with generator matrix $GP_n(1, 0, 1, 0, 1)$ over GF(2) is self-dual of length $2n$. The rest of this theorem can be proved similarly. \square

[72, 36, 12] codes also can be obtain by $GB_{36}(0, 0, 0, 1, 0, 1)$ with $p = 7$ and $q = 5$ over GF(2). The structures of those codes are similar. The **highest know** means that the code meets the highest know minimum distance for its parameters.

3.2 Self-Dual Codes Over GF(4)

In this subsection, we give four constructions of infinite families of self-dual codes over GF(4). Let u be the fixed primitive element of GF(4) satisfying $u^2 + u + 1 = 0$, then we have the following results.

Theorem 4. Let $\frac{(p-1)(q-1)}{4}$ be even and $\frac{p+q}{4}$ be even. Then the following holds:

1) $p = 4\omega + 1$ and $q = 4\omega' + 3$

The codes with generator matrices $GP_n(1, 0, 1, u+1, u)$ and $GP_n(1, 0, 1, u, u+1)$ over GF(4) are self-dual of length $2n$.

The codes with generator matrices $GB_n(0, 0, 1, 0, u, u+1)$ and $GB_n(0, 0, 1, 0, u+1, u)$ over GF(4) are self-dual of length $2n + 2$.

2) $p = 4\omega + 3$ and $q = 4\omega' + 1$

The codes with generator matrices $GP_n(1, 1, 0, u+1, u)$ and $GP_n(1, 1, 0, u, u+1)$ over GF(4) are self-dual of length $2n$.

The codes with generator matrices $GB_n(0, 0, 0, 1, u+1, u)$ and $GB_n(0, 0, 0, 1, u, u+1)$ over GF(4) are self-dual of length $2n + 2$.

Table 2: SOME CODES OVER GF(4)

Code	p	q	Construction	Comments
[30, 15, 6]	3	5	$GP_{15}(1, 1, 0, u+1, u)$	
[32, 16, 8]	3	5	$GB_{16}(0, 0, 0, 1, u+1, u)$	almost optimal

Proof. If $p = 4\omega + 1$ and $q = 4\omega' + 3$,

$$\begin{aligned} \alpha + N &= 4(4\omega\omega' + \omega + 3\omega') + 3 \equiv 1 \pmod{2}, \\ -\alpha + m_0 + (p-1)m_1 + (q-1)m_2 \\ &\quad + \frac{(p-1)(q-1)}{2}(m_3 + m_4) \equiv 0 \pmod{2}, \\ D_0(0, 1, 0, u, u+1) &\equiv 0 \pmod{2}, \\ D_1(0, 1, 0, u, u+1) \\ = D_2(0, 1, 0, u, u+1) &\equiv 1 \pmod{2}, \\ D_3(0, 1, 0, u, u+1) \\ = (0, 0)(m_3^2 + m_4^2) + ((1, 0) + (0, 1))m_3m_4 \\ = (u+1+u) + (1+0) \times (u+1+u) &\equiv 1 \pmod{2} \end{aligned}$$

By Theorem 2, the code with generator matrix $GB_n(0, 0, 1, 0, u, u+1)$ over $\text{GF}(4)$ is self-dual of length $2n+2$.

The rest of this theorem can be proved similarly. \square

4 Conclusions

In this paper, we construct several infinite families of self-dual codes based on the generalized cyclotomic sets of order two. Because of several good randomness properties, some of them have good parameters. These can enrich the choices of methods to construct good self-dual codes. We believe that these constructions will lead to good self-dual codes. It seems that the construction method by generalized cyclotomic classes of higher order is a rich source to obtain codes with good parameters.

Acknowledgments

The work is supported by Fundamental Research Funds for the central Universities(No.17CX02030A), Shandong Provincial Natural science Foundation of China (No.ZR2016FL01, No.ZR2017MA001, No.ZR2019MF070), Qingdao application research on special independent innovation plan project (o.16-5-1-5-jch), Key Laboratory of Applied Mathematics of Fujian Province University(Putian University)(No.SX201702, No.SX201806), Open Research Fund from Shandong provincial Key Laboratory of Computer Network, (No. SDKLCN-2017-03) and The National Natural Science Foundation of China(No.2017010754). The authors are grateful to the anonymous reviewers for valuable comments.

References

- [1] A. Alahmadi, C. Gneri, B. Ozkaya, H. Shoaib, and P. Sol. "On self-dual double negacirculant codes," *Discrete Applied Mathematics*, 222(C):205–212, 2017.
- [2] A. Alahmadi, F. Ozdemir, and P. Sol. "On self-dual double circulant codes," *Designs Codes Cryptography*, 86(6):1257–1265, 2018.
- [3] K. T. Arasu and T. A. Gulliver. "Self-dual codes over F_p and weighing matrices," *IEEE Transactions on Information Theory*, 47(5):2051–2055, 2001.
- [4] W. Bosma, J. Cannon, and C. Playoust. "The magma algebra system i: The user language," *J Symbolic Comput*, 24(3-4):235–265, 1997.
- [5] C. Ding. "Autocorrelation values of generalized cyclotomic sequences of order two," *IEEE Transactions on Information Theory*, 44(4):1699–1702, 1998.
- [6] C. Ding, D. Pei, and A. Salomaa. *Chinese remainder theorem: applications in computing, coding, cryptography*. 1996.
- [7] C. Ding. "Cyclic codes from the two-prime sequences," *IEEE Transactions on Information Theory*, 58(6):3881–3891, 2012.
- [8] C. Ding. "Cyclic codes from cyclotomic sequences of order four," *Finite Fields Their Applications*, 23(96):8–34, 2013.
- [9] X. Du and Z. Chen. "Trace representation of binary generalized cyclotomic sequences with length," *Teice Transactions on Fundamentals of Electronics Communications Computer Sciences*, 94-A(2):761–765, 2011.
- [10] T. Feng and Q. Xiang. "Cyclotomic constructions of skew hadamard difference sets," *Journal of Combinatorial Theory*, 119(1):245–256, 2012.
- [11] P. Gaborit. "Quadratic double circulant codes over fields," *Journal of Combinatorial Theory*, 97(1):85–107, 2002.
- [12] M. Garcia-Rodriguez, Y. Yanez, M. J. Garcia-Hernandez, J. Salazar, A. Turo, and J. A. Chavez. "Application of golay codes to improve the dynamic range in ultrasonic lamb waves air-coupled systems," *NDT E International*, 43(8):677–686, 2010.
- [13] L. Hu, Q. Yue, and X. Zhu. "Autocorrelation value of generalized cyclotomic sequences with period pq," *Journal of Nanjing University of Science Technology*, 2015.
- [14] F. J. Macwilliams and N. J. A. Sloane. *The theory of error-correcting codes*. 1977.
- [15] G. Nebe, E. M. Rains, and N. J. A Sloane. "Self-dual codes and invariant theory (algorithms and computation in mathematics)," In *Math Nachrichten*, 2006.
- [16] E. M. Rains. "Shadow bounds for self-dual codes," *IEEE Transactions on Information Theory*, 44(1):134–139, 1998.
- [17] R. G Stanton and D. A Sprott. "A family of difference sets," *Canadian Journal of Mathematics*, 10(1):73–77, 1958.
- [18] T. Zhang and G. Ge. "Fourth power residue double circulant self-dual codes," *IEEE Transactions on Information Theory*, 61(8):4243–4252, 2015.
- [19] Q. Wang and D. Lin. "Generalized cyclotomic numbers of order two and their applications," *Cryptography Communications*, 8(4):605–616, 2016.
- [20] T. Yan, X. Du, G. Xiao, and X. Huang. "Linear complexity of binary whitman generalized cyclotomic sequences of order $2k$," *Information Sciences*, 179(7):1019–1023, 2009.

Biography

Wenpeng Gao was born in 1994 in Shandong Province of China. He was graduated from the Department of Mathematics, China University of Petroleum, China, in 2017. He is a graduate student of China University of Petroleum.

Tongjiang Yan was born in 1973 in Shandong Province of China. He was graduated from the Department of Mathematics, Huaibei Coal-industry Teachers College, China, in 1996. He received the M.S. degree in mathematics from the Northeast Normal University, Lanzhou, China, in 1999, and the Ph.D degree in cryptography from Xidian University, Xian, China, in 2007. Now he is a associate professor of China University of Petroleum. His research interests include cryptography and algebra.