

$\{-1, 0, 1\}$ -APSP and (min,max)-Product Problems*

Hodaya Barr , Tsvi Kopelowitz , Ely Porat , Liam Roditty

Bar-Ilan University, Ramat Gan, Israel
odayaben@gmail.com, kopelot@gmail.com
porately@cs.biu.ac.il, liamr@macs.biu.ac.il

Abstract

In the $\{-1, 0, 1\}$ -APSP problem the goal is to compute all-pairs shortest paths (APSP) on a directed graph whose edge weights are all from $\{-1, 0, 1\}$. In the (min,max)-product problem the input is two $n \times n$ matrices A and B , and the goal is to output the (min,max)-product of A and B .

This paper provides a new algorithm for the $\{-1, 0, 1\}$ -APSP problem via a simple reduction to the target-(min,max)-product problem where the input is three $n \times n$ matrices A, B , and T , and the goal is to output a Boolean $n \times n$ matrix C such that the (i, j) entry of C is 1 if and only if the (i, j) entry of the (min,max)-product of A and B is exactly the (i, j) entry of the target matrix T . If (min,max)-product can be solved in $T_{MM}(n) = \Omega(n^2)$ time then it is straightforward to solve target-(min,max)-product in $O(T_{MM}(n))$ time. Thus, given the recent result of Bringmann, Künnemann, and Wegrzycki [STOC 2019], the $\{-1, 0, 1\}$ -APSP problem can be solved in the same time needed for solving approximate APSP on graphs with positive weights.

Moreover, we design a simple algorithm for target-(min,max)-product when the inputs are restricted to the family of inputs generated by our reduction. Using fast rectangular matrix multiplication, the new algorithm is faster than the current best known algorithm for (min,max)-product.

*This work is partially supported by ISF grants (no. 1278/16 and 1926/19), by a grant from the United States - Israel Binational Science Foundation (BSF) (no. 2018364), and by an ERC grant MPM under the European Union's Horizon 2020 research and innovation programme (no. 683064).

1 Introduction

The *all-pairs shortest paths* (APSP) problem is one of the most fundamental algorithmic problems in computer science, and is defined as follows. Let $G = (V, E)$ be a weighted (directed) graph with weight function $w : E \rightarrow \mathbb{R}$, where $V = \{v_1, v_2, \dots, v_n\}$. Let A be the weighted adjacency matrix of G . For $v_i, v_j \in V$, the weight of the shortest path between v_i and v_j is denoted by a_{ij}^* , and the matrix containing the weights of the shortest paths between all pairs of vertices is denoted by A^* . In the APSP problem the goal is to compute A^* .

The Floyd-Warshall APSP algorithm [8, 16, 22], is a dynamic programming algorithm, whose runtime is $O(n^3)$. After several improvements of poly-logarithmic factors in the time cost [9, 18, 12, 19, 20, 25, 4, 13, 5, 3, 14], in a recent breakthrough, Williams [23] and Chan and Williams [6] designed the current best algorithm whose time cost is $O(\frac{n^3}{2^{\Omega(\sqrt{\log n})}})$. The lack of success in designing a truly sub-cubic algorithm for APSP has led to a popular conjecture that any algorithm for APSP requires $\Omega(n^{3-o(1)})$ time [15, 21]. However, in some special cases, faster algorithms are known. We describe two examples which are strongly related to the results in this paper. Throughout the paper, let ω denote the exponent in the fastest fast matrix multiplication (FMM) algorithm; the current best upper bound on ω is roughly 2.3728639 [10].

Approximate APSP with positive real weights and (min,max)-product. The first example is the *approximate positive APSP* problem where the goal is to compute a $1 \pm \epsilon$ approximation of the distances for a graph with positive weights. A recent result by Bringmann, Künnemann, and Wegrzycki [2] shows that approximate positive APSP can be solved in $\tilde{O}\left(\frac{n^{\frac{3+\omega}{2}}}{\text{poly}(\epsilon)}\right) = \tilde{O}\left(\frac{n^{2.686}}{\text{poly}(\epsilon)}\right)$ time¹. The algorithm in [2] is obtained by showing an equivalence between approximate positive APSP and the (min,max)-product problem which is defined as follows. Throughout the paper we follow the notion that matrices are denoted with capital letters, while the entries of matrices are denoted using the same letter, just lowercase, with the appropriate indices indicated as subscripts. So, for example the (i, j) entry of matrix A is denoted by a_{ij} .

Definition 1.1 ((min,max)-product). *In the (min,max)-product problem the input is two matrices $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times n}$, and the output is a matrix $C \in \mathbb{R}^{n \times n}$ such that the (i, j) entry of C is*

$$c_{ij} = \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}.$$

We denote the (min,max)-product between A and B with the \otimes operator; that is, $C = A \otimes B$.

Duan and Pettie [7] showed that the (min,max)-product problem can be solved in $O(n^{(3+\omega)/2})$ time.

$\{-1, 0, 1\}$ -APSP. The second example is the $\{-1, 0, 1\}$ -APSP problem where the weights are from the set $\{-1, 0, 1\}$. Negative edge weights introduce a new depth to the challenges involved in solving APSP, and the $\{-1, 0, 1\}$ -APSP problem is perhaps the purest version of APSP that allows for negative weights.

Alon, Galil and Marglit [1] designed an algorithm for the $\{-1, 0, 1\}$ -APSP problem whose runtime is $O(n^{\frac{3+\omega}{2}}) = O(n^{2.686})$. Zwick [24] improved the runtime to $O(n^{2+\mu}) = O(n^{2.5286})$, where

¹We use the standard \tilde{O} notation to suppress poly-logarithmic factors.

μ satisfies $\omega(1, \mu, 1) = 1 + 2\mu$, and $\omega(1, \mu, 1)$ is the exponent of n in the fastest algorithm for multiplying two rectangular matrices of sizes $n \times n^\mu$ and $n^\mu \times n$. The current best upper bound on μ is roughly 0.5286 [10].

1.1 Our Results and Algorithmic Overview

In this paper we introduce a reduction from $\{-1, 0, 1\}$ -APSP to (\min, \max) -product, which combined with [2] implies that $\{-1, 0, 1\}$ -APSP is reducible to approximate positive APSP. Specifically, we use the framework of Alon et al. [1] who follow the paradigm of Seidel’s APSP algorithm for unweighted undirected graphs [17]. Seidel’s algorithm first recursively solves the undirected unweighted APSP problem on a specially constructed graph G' so that if the distance between vertices v_i and v_j in the original graph is a_{ij}^* , then the distance between v_i and v_j in G' is t_{ij}^* where $t_{ij}^* = \left\lceil \frac{a_{ij}^*}{2} \right\rceil$. Then, the problem of computing a_{ij}^* reduces to the problem of establishing whether a_{ij}^* is odd or even. The algorithm of [1] follows the same structure as Seidel’s algorithm, but instead of directly computing the parity of a_{ij}^* , the algorithm uses a brute-force like method.

Instead, we use a more direct approach for computing the parity of a_{ij}^* by solving two instances of the *target-(min,max)-product problem* where the input is the same as the input for the (\min, \max) -product problem together with a third *target matrix* $T \in \mathbb{R}^{n \times n}$. The output is a Boolean matrix whose (i, j) entry is an indicator of whether the (i, j) entry of the (\min, \max) -product is equal to the (i, j) entry in the target matrix. Formally:

Definition 1.2 (Target-(min,max)-product). *In the target-(min,max)-product (T -(min,max)-product) problem the input is three matrices $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times n}$ and $T \in \mathbb{R}^{n \times n}$, and the output is a Boolean matrix $C \in \{0, 1\}^{n \times n}$ such that $c_{ij} = 1$ if and only if $t_{ij} = \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. The T -(min,max)-product operation is denoted by $C = A \otimes_T B$.*

Given three matrices A, B and T it is straightforward to compute the T -(min,max)-product of A, B and T by first computing $A \otimes B$ and then spending another $O(n^2)$ time to compare each entry $A \otimes B$ with the corresponding entry in T .

Our main result is summarized by the following theorem (the proof is given in Section 3).

Theorem 1.3. *Suppose that there exists an algorithm for solving the T -(min,max)-product problem in $T_{TMM}(n) \geq n^2$ time. Then there exists an algorithm for solving $\{-1, 0, 1\}$ -APSP in $\tilde{O}(T_{TMM}(n) + n^\omega)$ time.*

A simple algorithm for restricted T-(min,max)-product. In addition to showing that $\{-1, 0, 1\}$ -APSP is reducible to T -(min,max)-product (and therefore also to (\min, \max) -product), we also design a simple algorithm for solving T -(min,max)-product for the restricted family of inputs generated by our reduction. Specifically, the entries in the second matrix B are $\pm\infty$, and the target matrix T has the property that for any² $i, j \in [n]$, $t_{ij} \leq \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. Formally, the T -(min,max)-product problem on this family of inputs is defined as follows.

Definition 1.4 (restricted T -(min,max)-product). *In the restricted T -(min,max)-product problem the input is composed of a matrix $A \in \mathbb{R}^{n \times n}$, a matrix $B \in \{-\infty, \infty\}^{n \times n}$, and a target matrix $T \in \mathbb{R}^{n \times n}$, where for every $i, j \in [n]$, $t_{ij} \leq \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. The output is a matrix $C \in \{0, 1\}^{n \times n}$, such that for every $i, j \in [n]$, $c_{ij} = 1$ if and only if $t_{ij} = \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$.*

²We use the standard notation that $[n] = \{1, 2, 3, \dots, n\}$.

In Section 4 we prove the following theorem.

Theorem 1.5. *For any $0 \leq t \leq 1$, there exists an algorithm for the restricted T -(min,max)-product problem whose time cost is $\tilde{O}(n^{2+t} + n^{\omega(2-t,1,1)})$ time.*

Using the algorithm of [11] for fast rectangular matrix multiplication, we are able to upper bound $\tilde{O}(n^{2+t} + n^{\omega(2-t,1,1)})$ by $\tilde{O}(n^{2.658044})$; see Appendix B for a detailed explanation. Thus, using our reduction from Theorem 1.3 together with the algorithm of Theorem 1.5, we obtain a new algorithm for $\{-1, 0, 1\}$ -APSP whose cost is $O(n^{2.658044})$ time. Notice that using fast squared matrix multiplication instead of fast rectangular matrix multiplication, the time bound becomes $\tilde{O}(n^{\frac{3+\omega}{2}}) = \tilde{O}(n^{2.686})$, which matches the runtime of [7] for solving (min,max)-product.

We remark that Zwick's algorithm [24] is faster and has a time cost of $\tilde{O}(n^{2.5286})$. Nevertheless, if $\omega = 2$ then the time costs of our algorithm, Zwick's algorithm [24] and the algorithm of Alon et al. [1] all become $\tilde{O}(n^{2.5})$.

2 Preliminaries

Hops and δ -regularity. If a path P in G has ℓ edges then P is said to have ℓ hops. Let $A^{\leq \ell}$ be a matrix where $a_{ij}^{\leq \ell}$ is the weight of the shortest path from v_i to v_j that has at most ℓ hops.

Definition 2.1 (δ -regularity [1]). *For any positive integer δ , a weighted adjacency matrix A of a graph $G = (V, E, w)$ with $V = \{v_1, v_2, \dots, v_n\}$ is δ -regular if for every pair of vertices $v_i, v_j \in V$: (i) if $a_{ij}^* \neq -\infty$ then $a_{ij}^* = a_{ij}^{\leq \delta}$, and (ii) if $a_{ij}^* = -\infty$ then $a_{ij}^{\leq \delta} < 0$.*

The following Lemma was proven in [1].

Lemma 2.2 (Lemma 2 in [1]). *Any weighted adjacency matrix $A \in \{-1, 0, 1, \infty\}^{n \times n}$ is n^2 -regular.*

We emphasize that for a given δ -regular matrix A , the entries of $A^{\leq \ell}$ are not necessarily the same as A^* . Specifically, since the shortest path from v_i to v_j in G could have more than ℓ hops, it is possible that $a_{ij}^{\leq \ell} \gg a_{ij}^*$. However, if A is a δ -regular matrix, then if there exists a shortest path in G between a pair of vertices u and v that does not contain a negative cycle, then there exists a shortest path in G between u and v with at most δ hops.

3 Reducing $\{-1, 0, 1\}$ -APSP to T -(min,max)-product

3.1 Canonical Graphs.

We start with the following definition of a canonical graph which is implicit in [1]:

Definition 3.1 (Canonical graph). *Let $G = (V, E, w)$ be a weighted directed graph with $w : E \rightarrow \{-1, 0, 1\}$. A weighted directed graph $G' = (V, E', w')$ is a canonical graph of G if $w' : E' \rightarrow \{-1, 0, 1\}$ and for any shortest path P from v_i to v_j in G there exists a shortest path P' from v_i to v_j in G' that satisfies the following conditions:*

1. $w(P) = w'(P')$.
2. If P has exactly ℓ hops then P' has at most ℓ hops.

3. If P' is not a single edge then P' does not contain zero weight edges.

Notice that due to the first condition in the definition of canonical graphs, the distance matrices of a graph G and its canonical graph G_c are the same. The following lemma, which is proven in [1], states that canonical graphs can be efficiently constructed.

Lemma 3.2 (Lemma 5 in [1]). *There exists an algorithm which constructs a canonical graph of a weighted directed graph $G = (V, E, w)$ where $w : E \rightarrow \{-1, 0, 1\}$ in $O(n^\omega)$.*

3.2 The Reduction Algorithm

We prove Theorem 1.3 by presenting algorithm $\text{APSP-To-TMinMax}(A, \delta)$ which solves the $\{-1, 0, 1\}$ -APSP problem on a graph whose weighted adjacency matrix A is guaranteed to be δ -regular, for an integer $\delta > 0$. Notice that, by Lemma 2.2, if A is the weighted adjacency matrix of G then A is n^2 -regular, and therefore $\text{APSP-To-TMinMax}(A, n^2)$ solves $\{-1, 0, 1\}$ -APSP on G .

We now turn to describe $\text{APSP-To-TMinMax}(A, \delta)$. (See Algorithm 1 for a pseudocode). The input is an integer value $\delta > 0$ and a δ -regular weighted adjacency matrix A . Let $G = (V, E, w)$ be the graph represented by weighted adjacency matrix A . If $\delta = 1$, and so A is 1-regular, the following lemma allows to compute A^* in $O(n^\omega)$ time.

Lemma 3.3 (Corollary of Lemmas 3 and 4 from [1]). *There exists an algorithm which solves the $\{-1, 0, 1\}$ -APSP problem on graphs whose weighted adjacency matrix is 1-regular in $O(n^\omega)$ time.*

Thus, suppose $\delta > 1$. The algorithm begins by constructing a canonical graph $G_c = (V, E_c, w_c)$ for G . Let C be the weighted adjacency matrix of G_c and let $T = \lceil C^{\leq 2}/2 \rceil$. Notice that $T \in \{-1, 0, 1, \infty\}^{n \times n}$ and computing T costs $O(n^\omega)$ time.

It was shown in [1] that T is a $\lceil \delta/2 \rceil$ -regular weighted adjacency matrix and that the distance matrix T^* of the graph represented by T is $\lceil C^*/2 \rceil$. Next, the algorithm performs a recursive call $\text{APSP-To-TMinMax}(T, \lceil \delta/2 \rceil)$, to compute T^* . From the definition of canonical graphs it follows that $C^* = A^*$. Hence, $T^* = \lceil C^*/2 \rceil = \lceil A^*/2 \rceil$.

Consider a pair of vertices v_i and v_j . Since $t_{ij}^* = \lceil a_{ij}^*/2 \rceil$ it follows that if a_{ij}^* is even then $2t_{ij}^* = a_{ij}^*$ and if a_{ij}^* is odd then $2t_{ij}^* = a_{ij}^* + 1$. Therefore, given the value of t_{ij}^* and the parity of a_{ij}^* , the algorithm is able to compute the value of a_{ij}^* . Since T^* is computed by the recursive call, the only remaining task in order to compute A^* is to compute the parity of a_{ij}^* for every $i, j \in [n]$.

Establishing parity. We now describe how to determine the parity of the entries of A^* without direct access to A^* , but with access to T^* and A . Notice that this is where our algorithm differs from the algorithm of [1].

Let

$$x_{ij}^+ = \begin{cases} -\infty, & \text{if } c_{ij} = +1 \\ \infty, & \text{otherwise} \end{cases}$$

$$x_{ij}^- = \begin{cases} -\infty, & \text{if } c_{ij} = -1 \\ \infty, & \text{otherwise} \end{cases}$$

and define a matrix M where $m_{ij} = t_{ij}^* - 1$.

Algorithm 1 APSP-To-TMinMax(A, δ)

1. **if** $\delta = 1$ **then**
 2. compute the distance matrix A^* using Lemma 3.3
 3. **return** A^*
 4. **end if**
 5. Let C be the weighted adjacency matrix of the canonical graph for the graph that A represents.
 6. $T^* \leftarrow \text{APSP-To-TMinMax}(\lceil C^{\leq 2}/2 \rceil, \lceil \delta/2 \rceil)$.
 7. $x_{ij}^- \leftarrow \begin{cases} -\infty, & \text{if } c_{ij} = -1 \\ \infty, & \text{otherwise} \end{cases}$
 $x_{ij}^+ \leftarrow \begin{cases} -\infty, & \text{if } c_{ij} = +1 \\ \infty, & \text{otherwise} \end{cases}$
 8. Let M be a matrix where $m_{ij} = t_{ij}^* - 1$
 $Z^- \leftarrow T^* \otimes_{T^*} X^-$
 $Z^+ \leftarrow T^* \otimes_M X^+$
 9. $a_{ij}^* \leftarrow \begin{cases} t_{ij}^*, & \text{if } t_{ij}^* = \pm\infty \\ 2t_{ij}^* - 1, & \text{if } z_{ij}^+ = 1 \text{ or } z_{ij}^- = 1 \\ 2t_{ij}^*, & \text{otherwise} \end{cases}$
 10. **return** A^*
-

The algorithm computes

$$\begin{aligned} Z^+ &= T^* \otimes_M X^+ \\ Z^- &= T^* \otimes_{T^*} X^-. \end{aligned}$$

The following three lemmas demonstrate the connection between both Z^+ and Z^- , and the parity of entries in A^* . For the proofs of these lemmas, notice that $t_{ij}^* \neq \infty$ if and only if there exists some path from v_i to v_j in G_c (and so, by the definition of a canonical graph, there must exist a path from v_i to v_j in G). The following observation is due to $m_{ij} = t_{ij}^* - 1$.

Observation 3.4. $t_{ij}^* = \pm\infty$ if and only if $c_{ij}^* = a_{ij}^* = m_{ij} = t_{ij}^*$.

Lemma 3.5. Let $v_i, v_j \in V$ and assume that $m_{ij} \neq \infty$, then $z_{ij}^+ = 1$ if and only if there exists a shortest path P in G_c from v_i to v_j such that the weight of the last edge of P is 1 and a_{ij}^* is odd.

Proof. The proof is a case analysis showing that the only case in which $z_{ij}^+ = 1$ is when there exists a shortest path from v_i to v_j whose last edge has weight 1 and a_{ij}^* is odd, and that in such a case, it must be that $z_{ij}^+ = 1$.

Notice that, since $m_{ij} \neq \infty$ and by the definition of X^+ , if there exists a path from v_i to v_j in G_c whose last edge has weight 1, then there exists a value $\hat{k} \in [n]$ such that $t_{i\hat{k}}^* = \min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\} \neq \infty$. Moreover, since $t_{i\hat{k}}^* \neq \infty$, there exists a path from v_i to v_j in G_c whose last edge has weight 1 and the vertex preceding v_j on this path is $v_{\hat{k}}$. Therefore, since $C^* = A^*$ and by the triangle inequality, $2t_{ij}^* - 1 \leq c_{ij}^* \leq c_{i\hat{k}}^* + 1 \leq 2t_{i\hat{k}}^* + 1$, and so $m_{ij} = t_{ij}^* - 1 \leq t_{i\hat{k}}^* = \min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\}$. Moreover, notice that, by Definition 1.2, $z_{ij}^+ = 1$ if and only if $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\} = m_{ij}$.




Figure 1: Caption goes here.

Suppose that there is no path from v_i to v_j whose last edge has weight 1. Then for every $1 \leq k \leq n$ we have that either $t_{ik}^* = \infty$ or $x_{kj}^+ = \infty$ (since otherwise there is a path from v_i to v_k in G_c and $c_{kj} = 1$). Thus, $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\} = \infty$. Since we assumed that $m_{ij} \neq \infty$ then it is guaranteed that in this case $z_{ij}^+ = 0$.

Thus, for the rest of the proof we assume that there exists a path P from v_i to v_j in G_c whose last edge is (v_k, v_j) , $c_{kj} = 1$, and without loss of generality the prefix of P from v_i to v_k is a shortest path. The following three categories cover all the possibilities for P (See Figure 1): (i) P is a shortest path and a_{ij}^* is odd. (ii) P is a shortest path and a_{ij}^* is even. (iii) P is not a shortest path.

First category. Consider the case where the weight of the last edge (v_k, v_j) of P is 1 and a_{ij}^* is odd. Since a_{ij}^* is odd we have $a_{ij}^* = 2t_{ij}^* - 1$. Since $a_{ik}^* = a_{ij}^* - 1$ we have that a_{ik}^* is even, implying that $a_{ik}^* = 2t_{ik}^*$. Therefore, $2t_{ik}^* = a_{ik}^* = a_{ij}^* - 1 = 2t_{ij}^* - 2$, and $t_{ik}^* = t_{ij}^* - 1 = m_{ij}$.

Second category. We now turn to the case where the weight of the last edge (v_k, v_j) of P is 1 and a_{ij}^* is even. Since a_{ij}^* is even, we have $a_{ij}^* = 2t_{ij}^*$. Since $a_{ik}^* = a_{ij}^* - 1$ then a_{ik}^* is odd, implying that $a_{ik}^* = 2t_{ik}^* - 1$. Therefore, $2t_{ik}^* - 1 = a_{ik}^* = a_{ij}^* - 1 = 2t_{ij}^* - 1$, and $t_{ik}^* = t_{ij}^* > m_{ij}$.

Third category. Next, consider the case where P is not a shortest path. Thus, $a_{ik}^* + 1 > a_{ij}^*$. Recall that $2t_{ij}^* - 1 \leq a_{ij}^*$ and $a_{ik}^* \leq 2t_{ik}^*$. Thus, $2t_{ij}^* - 1 \leq a_{ij}^* < a_{ik}^* + 1 \leq 2t_{ik}^* + 1$ and so $t_{ik}^* > t_{ij}^* - 1 = m_{ij}$.

Conclusion. Notice that when evaluating $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\}$, the term $\max(t_{ik}^*, x_{kj}^+) \neq \infty$ only for values of k for which there exists a path from v_i to v_k in G_c and $c_{kj} = 1$. Thus, since we assume that $m_{ij} \neq \infty$, the only values of k which need to be considered are values of k such that there exists a path from v_i to v_j whose last edge is (v_k, v_j) and $c_{kj} = 1$; all other values of k have $\max(t_{ik}^*, x_{kj}^+) = \infty$ and are therefore not relevant.

If all such paths fall into either the second or third category, then for every such path P whose last edge is (v_k, v_j) we have $\max(t_{ik}^*, x_{kj}^+) = t_{ik}^* > m_{ij}$, and so $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\} > m_{ij}$ implying

$z_{ij}^+ = 0$. On the other hand, if there exists at least one such path that falls into the first category, then for any such path P whose last edge is (v_k, v_j) , we have $\max(t_{ik}^*, x_{kj}^+) = t_{ik}^* = t_{ij}^* - 1 = m_{ij}$. Since we always have $m_{ij} \leq \min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\}$, it follows that $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^+)\} = m_{ij}$, and so $z_{kj}^+ = 1$. \square

Lemma 3.6. *Let $v_i, v_j \in V$ and assume that $m_{ij} \neq \infty$, then $z_{ij}^- = 1$ if and only if there exists a shortest path P from v_i to v_j in G_c such that the weight of the last edge of P is -1 and a_{ij}^* is odd.*

The proof of Lemma 3.6 is similar to the proof of Lemma 3.5, and so the proof is deferred to Appendix A.

Lemma 3.7. *Let $v_i, v_j \in V$. If a_{ij}^* is odd then there exists a shortest path from v_i to v_j in G_c that has a last edge with weight either 1 or -1 .*

Proof. Assume by contradiction that a_{ij}^* is odd, but the last edge in every shortest path from v_i to v_j has weight 0. Since G_c is a canonical graph, for every pair of vertices v_i and v_j , there must exist a shortest path $P \in G_c$ from v_i to v_j that either has only one edge or does not contain any edges with weight 0. Thus, if all of the shortest paths from v_i to v_j have a last edge of weight 0, then $a_{ij}^* = 0$, and so a_{ij}^* is even, which is a contradiction. \square

By Lemma 3.7, for every pair of vertices v_i and v_j , a_{ij}^* is odd if and only if there exists a shortest path from v_i to v_j in G_c such that the weight of the last edge of the path is 1 or -1 . Moreover, by Lemmas 3.5 and 3.6, if the weight of the last edge of a shortest path from v_i to v_j in G_c is 1 then $z_{ij}^+ = 1$, and if the weight of the last edge of a shortest path from v_i to v_j in G_c is -1 then $z_{ij}^- = 1$.

Corollary 3.8. *Let $v_i, v_j \in V$. Then a_{ij}^* is odd if and only if either $z_{ij}^- = 1$ or $z_{ij}^+ = 1$.*

By Lemmas 3.5 and 3.6, and by Corollary 3.8, for each pair of vertices v_i and v_j we have that

$$a_{ij}^* = \begin{cases} t_{ij}^*, & \text{if } t_{ij}^* = \pm\infty \\ 2t_{ij}^* - 1, & \text{if } z_{ij}^+ = 1 \vee z_{ij}^- = 1 \\ 2t_{ij}^*, & \text{otherwise} \end{cases}$$

Thus, constructing A^* from T^* , Z^+ , and Z^- costs $O(n^2)$ time.

Time cost. We now analyze the run time of $\text{APSP-To-TMinMax}(A, n^2)$ where A is the weighted adjacency matrix of G . In the following, the line numbers refer to the lines in Algorithm 1. Let $f(n, \delta)$ denote the run time of the algorithm on the weighted adjacency matrix of a δ -regular graph with n vertices. By Lemma 3.3, the computation in Lines 1–4 costs $O(n^\omega)$ time. By Lemma 3.2, the construction of the canonical graph in Line 5, costs $O(n^\omega)$ time. In Line 6 the algorithm first computes the matrix $\lceil C^{\leq 2}/2 \rceil$ in $O(n^\omega)$ time (using FMM), and then the algorithm makes a recursive call that costs $f(n, \delta/2)$ time. Lines 7 and 9 cost $O(n^2)$ time. Line 8 has two T-(min,max)-product computations which cost $O(T_{TMM}(n))$ time. Thus,

$$f(n, \delta) = f(n, \lceil \delta/2 \rceil) + O(n^\omega + T_{TMM}(n)).$$

Finally, since in each recursive δ is halved, the number of recursive calls is $\log \delta$. Thus, $f(n, n^2) = \tilde{O}(n^\omega + T_{TMM}(n))$.

4 A Simple Algorithm for Restricted T-(min,max)-product

Recall that Algorithm 1 (of Theorem 1.3) makes use of an algorithm for T-(min,max)-product. However, these calls are applied to a *restricted* family of inputs: the entries in the second matrix B are $\pm\infty$, and the target matrix T has the property that for any $i, j \in [n]$, $t_{ij} \leq \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. In this section we prove Theorem 1.5 by describing a simple algorithm for T-(min,max)-product when the inputs are *restricted* to the family of inputs that are seen when calling T-(min,max)-product in Algorithm 1; see Definition 1.4. Our algorithm combines *fast rectangular Boolean matrix multiplication* (BMM) with a *heavy-light* type of decomposition³.

For every $i \in [n]$, let A_i be the i^{th} row of A . The algorithm sorts the pairs (a_{ij}, j) , where $j \in [n]$, in a lexicographically increasing order and stores the result in array L_i . For A_i , we say that a value that occurs more than n^t times in A_i is *heavy* for A_i , and a value in A_i that occurs at most n^t times in A_i is *light* for A_i . It is straightforward to partition the values in A_i to heavy and light values in $O(n)$ time. Notice that there are at most $n/n^t = n^{1-t}$ heavy elements for A_i .

The algorithm constructs a rectangular matrix H as follows. For every $i \in [n]$ and every heavy value x in A_i the algorithm adds a row to H that corresponds to x . Let $\rho(i, x)$ be the index of the row in H that is added for heavy value x . Since there are at most n^{1-t} heavy values for each A_i , the number of rows in H is at most n^{2-t} . For $j \in [n]$, the j^{th} entry in the $\rho(i, x)$ 'th row is set to 1 if and only if the j^{th} entry of A_i contains the value x ; otherwise the j^{th} entry is set to 0. Thus, H is a Boolean matrix of size $O(n^{2-t}) \times n$.

The algorithm converts the matrix B into a Boolean matrix B' as follows. For every $i, j \in [n]$ let:

$$b'_{ij} = \begin{cases} 1 & \text{if } b_{ij} = -\infty \\ 0 & \text{otherwise} \end{cases}$$

Next, the algorithm computes the rectangular BMM $F = HB'$. Finally, the algorithm constructs matrix C as follows. Consider the value t_{ij} from the target matrix T , where $i, j \in [n]$. If t_{ij} is heavy in row A_i then the algorithm sets c_{ij} to f_{qj} for $q = \rho(i, t_{ij})$. If t_{ij} is light, then the algorithm uses L_i to access all the occurrences of t_{ij} in A_i . Let $I = \{k_1, \dots, k_\ell\}$ be the set of all indices of columns in row i that contain the value t_{ij} . Thus, for every $k \in I$ we have $a_{ik} = t_{ij}$. If there is $k \in I$ such that $b_{kj} = -\infty$ the algorithm sets c_{ij} to 1, otherwise the algorithm sets c_{ij} to 0.

Correctness. We now show that the matrix C computed above equals $A \otimes_T B$. By Definition 1.4 c_{ij} should be 1 if $t_{ij} = \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$, and 0 otherwise. By Definition 1.4 we also have $t_{ij} \leq \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. Thus, if there exists a k such that $t_{ij} = \max(a_{ik}, b_{kj})$ then $t_{ij} = \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. Notice that if there exists k' such that $t_{ij} > \max(a_{ik'}, b_{k'j})$, then $t_{ij} > \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$, which is a contradiction. Moreover, if for every $k \in [n]$, $t_{ij} \neq \max(a_{ik}, b_{kj})$, then $t_{ij} \neq \min_{k=1}^n \{\max(a_{ik}, b_{kj})\}$. Since $B \in \{-\infty, \infty\}^{n \times n}$, then we conclude that there exists $k \in [n]$ such that $t_{ij} = \max(a_{ik}, b_{kj})$ if and only if $a_{ik} = t_{ij}$ and $b_{kj} = -\infty$.

If t_{ij} is light in row A_i then for every $k \in [n]$ where $a_{ik} = t_{ij}$ the algorithm checks whether $b_{kj} = -\infty$ or not. Thus if $t_{ij} = \max(a_{ik}, b_{kj})$ and t_{ij} is light then the algorithm will detect this case. If t_{ij} is heavy in row A_i then row $\rho(i, x)$ in H has 1 in the same columns that contain t_{ij} in row A_i and 0 in all other columns. In matrix B' there are 1 values in all entries that correspond to entries in B with value $-\infty$ and 0 in all other locations. Hence, in F , $f_{qj} = 1$ for $q = \rho(i, t_{ij})$

³According to Definition 1.4 A and T , are both from $\mathbb{R}^{n \times n}$, while in APSP-To-TMinMax(.) the matrices can have entries from $\{-\infty, \infty\}$. However, if $t_{ij} \in \{-\infty, \infty\}$ then a_{ij}^* is set to be t_{ij} regardless the product.

if and only if there exists a $k \in [n]$ such that $a_{ik} = t_{ij}$ and $b_{kj} = -\infty$ which in turn implies that $t_{ij} = \max(a_{ik}, b_{kj})$.

Time cost. The cost of handling a light value is $O(n^t + \log n)$. Since T contains at most $O(n^2)$ light values, the total cost for handling all light values is at most $O(n^{2+t})$. Computing F using fast rectangular matrix multiplication takes $O(n^{\omega(2-t,1,1)})$ time. Thus, the total time cost of the algorithm is $O(n^{2+t} + n^{\omega(2-t,1,1)})$.

References

- [1] N. Alon, Z. Galil, and O. Margalit. On the exponent of the all pairs shortest path problem. *Journal of Computer and System Sciences*, 54(2):255–262, 1997.
- [2] K. Bringmann, M. Künnemann, and K. Wegrzycki. Approximating APSP without scaling: equivalence of approximate min-plus and exact min-max. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing, STOC*, pages 943–954, 2019.
- [3] T. C. Chan. More algorithms for all-pairs shortest paths in weighted graphs. *SIAM J. Comput.*, 39(5):2075–2089, 2010.
- [4] T. M. Chan. All-pairs shortest paths with real weights in $O(n^3/\log n)$ time. In *Algorithms and Data Structures, 9th International Workshop, WADS*, pages 318–324, 2005.
- [5] T. M. Chan. All-pairs shortest paths for unweighted undirected graphs in $o(mn)$ time. In *Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm (SODA)*, pages 514–523. SIAM, 2006.
- [6] T. M. Chan and R. R. Williams. Deterministic apsp, orthogonal vectors, and more: Quickly derandomizing razborov-smolensky. In *Proceedings of the twenty-seventh annual ACM-SIAM symposium on Discrete algorithms (SODA)*, pages 1246–1255. SIAM, 2016.
- [7] R. Duan and S. Pettie. Fast algorithms for (max, min)-matrix multiplication and bottleneck shortest paths. In *Proceedings of the twentieth annual ACM-SIAM symposium on Discrete algorithms*, pages 384–391. Society for Industrial and Applied Mathematics, 2009.
- [8] R. W. Floyd. Algorithm 97: shortest path. *Communications of the ACM*, 5(6):345, 1962.
- [9] M. L. Fredman. New bounds on the complexity of the shortest path problem. *SIAM Journal on Computing*, 5(1):83–89, 1976.
- [10] F. L. Gall. Powers of tensors and fast matrix multiplication. In *International Symposium on Symbolic and Algebraic Computation, ISSAC*, pages 296–303, 2014.
- [11] F. L. Gall and F. Urrutia. Improved rectangular matrix multiplication using powers of the coppersmith-winograd tensor. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1029–1046. SIAM, 2018.
- [12] Y. Han. Improved algorithm for all pairs shortest paths. *Inf. Process. Lett.*, 91(5):245–250, 2004.

- [13] Y. Han. An $O(n^3 (\log \log n / \log n)^{5/4})$ time algorithm for all pairs shortest paths. In *14th Annual European Symposium on Algorithms, ESA*, pages 411–417, 2006.
- [14] Y. Han and T. Takaoka. An $O(n^3 \log \log n / \log^2 n)$ time algorithm for all pairs shortest paths. In *Algorithm Theory - SWAT 2012 - 13th Scandinavian Symposium and Workshops*, pages 131–141, 2012.
- [15] L. Roditty and U. Zwick. On dynamic shortest paths problems. *Algorithmica*, 61(2):389–401, 2011.
- [16] B. Roy. Transitivité et connexité. *Comptes Rendus Hebdomadaires Des Seances De L Academie Des Sciences*, 249(2):216–218, 1959.
- [17] R. Seidel. On the all-pairs-shortest-path problem in unweighted undirected graphs. *J. Comput. Syst. Sci.*, 51(3):400–403, 1995.
- [18] T. Takaoka. A new upper bound on the complexity of the all pairs shortest path problem. In *17th International Workshop, WG '91, Fischbachau, Germany, June 17-19, 1991, Proceedings*, pages 209–213, 1991.
- [19] T. Takaoka. A faster algorithm for the all-pairs shortest path problem and its application. In *Computing and Combinatorics, 10th Annual International Conference, COCOON*, pages 278–289, 2004.
- [20] T. Takaoka. An $O(n^3 \log \log n / \log n)$ time algorithm for the all-pairs shortest path problem. *Inf. Process. Lett.*, 96(5):155–161, 2005.
- [21] V. Vassilevska Williams and R. R. Williams. Subcubic equivalences between path, matrix and triangle problems. In *51th Annual IEEE Symposium on Foundations of Computer Science, FOCS*, pages 645–654, 2010.
- [22] S. Warshall. A theorem on boolean matrices. *Journal of the ACM (JACM)*, 9(1):11–12, 1962.
- [23] R. R. Williams. Faster all-pairs shortest paths via circuit complexity. *SIAM J. Comput.*, 47(5):1965–1985, 2018.
- [24] U. Zwick. All pairs shortest paths using bridging sets and rectangular matrix multiplication. *Journal of the ACM (JACM)*, 49(3):289–317, 2002.
- [25] U. Zwick. A slightly improved sub-cubic algorithm for the all pairs shortest paths problem with real edge lengths. In *Algorithms and Computation, 15th International Symposium, ISAAC*, pages 921–932, 2004.

A Proof of Lemma 3.6

Proof of Lemma 3.6. The proof is a case analysis showing that the only case in which $z_{ij}^- = 1$ is when there exists a shortest path from v_i to v_j whose last edge has weight -1 and a_{ij}^* is odd, and that in such a case, it must be that $z_{ij}^- = 1$.

Notice that, since $t_{ij}^* \neq \infty$ and by the definition of X^- , if there exists a path from v_i to v_j in G_c whose last edge has weight -1 , then there exists a value $\hat{k} \in [n]$ such that $t_{i\hat{k}}^* = \min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\} \neq \infty$. Moreover, since $t_{i\hat{k}}^* \neq \infty$, there exists a path from v_i to v_j in G_c whose last edge has weight -1 and the vertex preceding v_j on this path is $v_{\hat{k}}$. Therefore, since $C^* = A^*$ and by the triangle inequality, $2t_{ij}^* - 1 \leq c_{ij}^* \leq c_{i\hat{k}}^* - 1 \leq 2t_{i\hat{k}}^* - 1$, and so $t_{ij}^* \leq t_{i\hat{k}}^* = \min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\}$. Moreover, notice that, by Definition 1.2, $z_{ij}^- = 1$ if and only if $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\} = t_{ij}^*$.

Suppose that there is no path from v_i to v_j whose last edge has weight -1 . Then for every $1 \leq k \leq n$ we have that either $t_{ik}^* = \infty$ or $x_{kj}^- = \infty$ (since otherwise there is a path from v_i to v_k and $c_{kj} = -1$). Thus, $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\} = \infty$. Since we assumed that $t_{ij}^* \neq \infty$ then it is guaranteed that in this case $z_{ij}^- = 0$.

Thus, for the rest of the proof we assume that there exists a path P from v_i to v_j in G_c whose last edge is (v_k, v_j) , $c_{kj} = -1$, and without loss of generality the prefix of P from v_i to v_k is a shortest path. The following three categories cover all the possibilities for P : (i) P is a shortest path and a_{ij}^* is odd. (ii) P is a shortest path and a_{ij}^* is even. (iii) P is not a shortest path.

First category. Consider the case where the weight of the last edge (v_k, v_j) of P is -1 and a_{ij}^* is odd. Since a_{ij}^* is odd we have $a_{ij}^* = 2t_{ij}^* - 1$. Since $a_{ik}^* = a_{ij}^* + 1$ we have that a_{ik}^* is even, implying that $a_{ik}^* = 2t_{ik}^*$. Therefore, $2t_{ik}^* = a_{ik}^* = a_{ij}^* + 1 = 2t_{ij}^*$, and $t_{ik}^* = t_{ij}^*$.

Second category. We now turn to the case where the weight of the last edge (v_k, v_j) of P is -1 and a_{ij}^* is even. Since a_{ij}^* is even, we have $a_{ij}^* = 2t_{ij}^*$. Since $a_{ik}^* = a_{ij}^* + 1$ then a_{ik}^* is odd, implying that $a_{ik}^* = 2t_{ik}^* - 1$. Therefore, $2t_{ik}^* - 1 = a_{ik}^* = a_{ij}^* + 1 = 2t_{ij}^* + 1$, and $t_{ik}^* = t_{ij}^* + 1 > t_{ij}^*$.

Third category. Next, consider the case where P is not a shortest path. Thus, $a_{ik}^* - 1 > a_{ij}^*$. Recall that $2t_{ij}^* - 1 \leq a_{ij}^*$ and $a_{ik}^* \leq 2t_{ik}^*$. Thus, $2t_{ij}^* - 1 \leq a_{ij}^* < a_{ik}^* - 1 \leq 2t_{ik}^* - 1$ and so $t_{ik}^* > t_{ij}^*$.

Conclusion. Notice that when evaluating $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\}$, the term $\max(t_{ik}^*, x_{kj}^-) \neq \infty$ only for values of k for which there exists a path from v_i to v_k in G_c and $c_{kj} = -1$. Thus, since we assume that $t_{ij}^* \neq \infty$, the only values of k which need to be considered are values of k such that there exists a path from v_i to v_j whose last edge is (v_k, v_j) and $c_{kj} = -1$; all other values of k have $\max(t_{ik}^*, x_{kj}^-) = \infty$ and are therefore not relevant.

If all such paths fall into either the second or third category, then for every such path P whose last edge is (v_k, v_j) we have $\max(t_{ik}^*, x_{kj}^-) = t_{ik}^* > t_{ij}^*$, and so $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\} > t_{ij}^*$ implying $z_{ij}^- = 0$. On the other hand, if there exists at least one such path that falls into the first category, then for any such path P whose last edge is (v_k, v_j) , we have $\max(t_{ik}^*, x_{kj}^-) = t_{ij}^*$. Since we always have $t_{ij}^* \leq \min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\}$, it follows that $\min_{k=1}^n \{\max(t_{ik}^*, x_{kj}^-)\} = t_{ij}^*$, and so $z_{ij}^- = 1$. \square

B Time Cost of Theorem 1.5

The goal is to find the value $0 \leq t \leq 1$ which minimizes $O(n^{2+t} + n^{\omega(2-t,1,1)})$. This goal is equivalent to finding the value $1 \leq k \leq 2$ such that $\omega(k, 1, 1) = 4 - k$. In the following we use the fact that the function $\omega(k, 1, 1)$ is convex, so we apply linear interpolation between the values of $\omega(k, 1, 1)$ for $k = 1.3$ and $k = 1.4$, which are given in [11]. Specifically, $\omega(1.3, 1, 1) = 2.621644$ and $\omega(1.4, 1, 1) = 2.708400$. Therefore, the line connecting $(1.3, 2.621644)$ and $(1.4, 2.708400)$ is above the point $(k, \omega(k, 1, 1))$ that we are searching for. This line is given by the equation

$$y = 0.86756x + 1.493816.$$

Solving $y(\hat{k}) = 4 - \hat{k}$, we have $\hat{k} = 1.3419156349$. Therefore,

$$\begin{aligned} \min_{1 \leq k \leq 2} \{\omega(k, 1, 1) + 4 - k\} &\leq \omega(\hat{k}, 1, 1) + 4 - \hat{k} \\ &\leq y(\hat{k}) + 4 - \hat{k} \\ &= y(1.3419156349) + 4 - 1.3419156349 \\ &= 2.658043651, \end{aligned}$$

and so $O(n^{2+t} + n^{\omega(2-t,1,1)}) < O(n^{2.658043651})$.