

Trans-Planckian censorship of multi-stage inflation and dark energy

Hao-Hao Li^{1*}, Gen Ye^{1†}, Yong Cai^{1‡}, and Yun-Song Piao^{1,2§}

¹ *School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China and*

² *Institute of Theoretical Physics, Chinese Academy of Sciences,*

P.O. Box 2735, Beijing 100190, China

Abstract

We explore the bound of Trans-Planckian censorship conjecture on the inflation model with multiple stages. We show that if the first inflationary stage is responsible for the primordial perturbations at CMB window, the e-folding number of each subsequent stage will be bounded by the energy scale of the first stage. This seems to imply that the lifetime of current accelerated expanding era (regarded as one of the multiple inflationary stages) might be a probe for distinguishing inflation from its alternatives. We also present a multi-stage inflation model in a landscape consisting of Anti-de Sitter vacua separated by the potential barriers.

PACS numbers:

* lihaohao18@mails.ucas.ac.cn

† yegen14@mails.ucas.ac.cn

‡ caiyong13@mails.ucas.ac.cn

§ yspiao@ucas.ac.cn

Contents

I. Introduction	2
II. Multi-stage Inflation	3
A. Multi-stage inflation with TCC	3
B. Multi-stage inflation in AdS Landscape	6
III. Implications for Dark Energy	8
IV. Discussion	10
Acknowledgments	10
References	10

I. INTRODUCTION

Inflation [1–4], which sets the initial conditions of the hot Big Bang (BB) cosmology, is a popular paradigm for the early universe. The evolution of the universe must be implemented in a UV-complete effective field theory (EFT). It was argued in [5, 6] that such EFTs should satisfy the swampland conjectures. Also importantly, it was pointed out in [7, 8] that the length scales of fluctuations we observe today might be smaller than the Planck length in the inflationary phase if inflation lasts long enough, which is the so-called “trans-Planckian” problem.

Recently, a new swampland condition, i.e., the Trans-Planckian Censorship Conjecture (TCC), has been proposed in [9], which states that the sub-Planckian fluctuations should never cross their Hubble scale to become classical, otherwise the corresponding EFT will belong to the swampland. It actually suggests that the “trans-Planckian” problem never happened in a UV-complete EFT. In other words, a cosmological model which suffers such a problem is in swampland. Requiring that the sub-Planckian perturbation will never have its length scale larger than the Hubble scale is equivalent to

$$\frac{a(t)}{a^{\text{ini}}} l_{\text{P}} < \frac{1}{H(t)} \tag{1}$$

in an expanding universe, where l_P is the Planck length. The immediate implications of TCC for inflation, the early universe and other aspects in cosmology have been studied in Refs. [10–25].

According to TCC (1), inflation can only lasts for a limited e-folding number

$$\int_{t^{\text{ini}}}^{t^{\text{end}}} H dt < \ln \frac{M_P}{H^{\text{end}}}, \quad (2)$$

where the superscript “ini” and “end” represent the beginning and ending of this stage (i.e., inflation), respectively. Thus if TCC is correct, it is not sufficiently reasonable to set the initial state of the perturbation modes as the Bunch-Davis state, which is required for explaining the observations at the CMB window. However, as showed in Ref. [11], a past-complete pre-inflation era can automatically prepare such initial states.

Furthermore, the TCC indicates that the energy scale of slow-roll inflation will reduce to $H_{\text{inf}} \sim 0.1$ GeV, which results in a tensor-to-scalar ratio $r < 10^{-30}$ at the CMB window, provided the radiation prevailed rapidly after inflation [10]. However, a non-standard post-inflationary history, in particular that of a multi-stage inflation scenario, will alleviate this TCC constraint [14, 16, 17] and might push r up to $r \lesssim 10^{-8}$, see also the multi-stage warm inflation [21]. Therefore, it is interesting to investigate the cosmological scenarios with multi-stage inflation, see also earlier studies in, e.g., Refs. [26–28].¹ The corresponding EFTs should satisfy not only the TCC, but also other swampland conjectures [5, 6].

In this paper, we explore the bound of TCC on the multi-stage inflation scenario and build a multi-stage inflation model in a landscape consisting of Anti-de Sitter (AdS) vacua. Also, if TCC is correct, it seems that the lifetime of current accelerated expansion (as one stage of multi-stage inflation scenario) might be shorter than expected in Ref. [9].

II. MULTI-STAGE INFLATION

A. Multi-stage inflation with TCC

Considering a multi-stage inflation scenario preceding the hot BB expansion, we set the Hubble parameter during the i th-stage of inflation as a constant H_i . We will assume that after the i th-stage inflation, the decelerated expanding phase has the equation of state

¹ The multi-stream inflation [29–32] might also be interesting.




FIG. 1: The sketch of multi-stage inflation confronted with the TCC. Here, $L \propto a$ denotes the physical length, H_0 is Hubble parameter at present. The black line with slope equals to 1 represents the evolution of the perturbation mode with the largest wavelength that can be observed at present. The red line represents the evolution of the perturbation mode with its physical wavelength equals to the Planck length at the beginning of the first inflationary stage. The blue polyline represents the evolution of Hubble radius, where we have assumed H_i 's are almost constant during each stage of inflation and $w_i > -1/3$.

$w_i = p_i/\rho_i$ (see Fig. 1 for a sketch), hence we have $a \sim t^{\frac{2}{3(1+w_i)}}$. Thus, the slope of $\ln(H^{-1}/H_0^{-1})$ (depicted by blue polyline) with respect to $\ln(a/a_0)$ at this stage is $3(1+w_i)/2$, where H_0 is the current Hubble constant. During the decelerated expansion with $w_i > -1/3$, the Hubble radius will rise faster than the physical length scale.

The efolding number of a stage is defined as

$$\tilde{N} = \ln \left(\frac{a^{\text{end}} H^{\text{end}}}{a^{\text{ini}} H^{\text{ini}}} \right). \quad (3)$$

We also set $N = \ln \left(\frac{a^{\text{end}}}{a^{\text{ini}}} \right)$, and hence have $\tilde{N} = N$ for $H^{\text{end}} = H^{\text{ini}}$. We will use \tilde{N}_i (or N_i) to represent the efolding number of the i th-stage inflation and \tilde{N}_{w_i} (or N_{w_i}) to represent that of the i th-stage decelerated expansion with $w_i > -1/3$. Note that $\tilde{N}_i > 0$ while $\tilde{N}_{w_i} < 0$.

From Fig. 1, we have the constraints on $\ln(H^{-1}/H_0^{-1})$ as follows.

Constraint 1: The perturbation mode with its wavelength equal to the Planck length at the beginning of the 1st-stage inflation will never cross the Hubble horizon. As a result, the red line (i.e., $\ln\left(\frac{al_{\text{P}}}{a_1^{\text{ini}}H_0^{-1}}\right)$) in Fig. 1 puts a lower bound on $\ln(H^{-1}/H_0^{-1})$, i.e.,

$$\frac{1}{aH} > \frac{l_{\text{P}}}{a^{\text{min}}}, \quad (4)$$

where ‘min’ represents the minimum throughout the history of the universe.

Constraint 2: Assuming that the 1st-stage inflation is responsible for the primordial perturbations at the CMB window, which should not be polluted by the follow-on stages, we will have an upper bound (the black line in Fig. 1) on $\ln(H^{-1}/H_0^{-1})$, i.e.,

$$\frac{1}{aH} < \frac{e^{-\tilde{N}_{\text{CMB}}}}{a_1^{\text{ini}}H_1^{\text{ini}}} \quad \text{for} \quad a_1^{\text{end}} < a < a_{\text{BB}}^{\text{ini}}, \quad (5)$$

where \tilde{N}_{CMB} is the efolds of the CMB window, $a_{\text{inf}}^{\text{end}}$ and $a_{\text{BB}}^{\text{ini}}$ are evaluated at the end of the 1st-stage inflation and the beginning of the hot BB expansion, respectively.

Considering a multi-stage inflation with $n > 1$ stages, for the j th stage, we can write down the bound (4) put by the TCC as

$$\tilde{N}_j + \sum_{i=1}^{j-1} (\tilde{N}_i + \tilde{N}_{w_i}) = \ln\left(\frac{a_j^{\text{end}}H_j^{\text{end}}}{a_1^{\text{ini}}H_1^{\text{ini}}}\right) < \ln\left(\frac{M_{\text{P}}}{H_1^{\text{ini}}}\right), \quad (6)$$

where we have used $\ln(a_j/a_0) - \ln(a_1^{\text{ini}}/a_0) < \ln(H_j^{-1}/H_0^{-1}) - \ln(l_{\text{P}}/H_0^{-1})$ (note the slope of the red line in Fig. 1 is 1) and $H_1^{\text{ini}} \simeq H_1^{\text{end}}$. We have set the Planck mass $M_{\text{P}} = l_{\text{P}}^{-1}$. Thus for $n = 1$, i.e., the single-stage inflation, we have $\tilde{N}_1 < \ln\left(\frac{M_{\text{P}}}{H_1^{\text{ini}}}\right)$ or equivalently $N_1 < \ln\left(\frac{M_{\text{P}}}{H_1^{\text{end}}}\right)$, which is consistent with the result in Refs. [9, 10].

Based on (5) and (6), for the multi-stage inflation, the efolding number of the j th-stage inflation ($j \neq 1$) can be bounded as

$$\tilde{N}_j < \ln\left(\frac{M_{\text{P}}}{H_1^{\text{ini}}}\right) - \tilde{N}_{\text{CMB}} \quad (7)$$

for $a_1^{\text{end}} < a < a_0$. Therefore, the lifetime of the j th-stage inflation is strictly bounded by the energy scale of the 1st-stage inflation and the width of the CMB window. Similarly, for a decelerated expanding stage, e.g., the stage marked by w_{j-1} , we have

$$-\tilde{N}_{w_{j-1}} = \frac{1}{2}(1 + 3w_{j-1})N_{w_{j-1}} < \ln\left(\frac{M_{\text{P}}}{H_1^{\text{ini}}}\right) - \tilde{N}_{\text{CMB}} \quad (8)$$

for $a_1^{\text{end}} < a < a_0$. Note that $\tilde{N}_{w_{j-1}} < 0$.

Therefore, if the TCC is correct and the 1st-stage inflation is responsible for the primordial perturbations at the CMB window, the e-folding number that the j th stage ($j \neq 1$, no matter accelerated or decelerated) can last is bounded by the energy scale (or H_1) of the 1st-stage inflation.

B. Multi-stage inflation in AdS Landscape

The landscape consists of all EFTs with a consistent UV-completion, which might be from various compactifications of the string theory. Otherwise, the set of the corresponding EFTs is called the swampland. As the swampland criteria for EFTs, the distance conjecture $|\Delta\phi|/M_{\text{P}} < \Delta \sim \mathcal{O}(1)$ [5] and the dS conjecture $M_{\text{P}}|\nabla_\phi V|/V > c \sim \mathcal{O}(1)$ [6] (or the refined dS conjecture $M_{\text{P}}^2 \frac{\min(\nabla_\phi \nabla_\phi V)}{V} \leq -c' \sim \mathcal{O}(1)$) [33, 34] have been proposed. Recently, many efforts on confronting the inflation scenario with the swampland conjectures have been made, see, e.g., [35–43].

The multi-stage inflation model (in which a high-scale inflation is followed by many low-scale inflations) might be interesting, since it helps to alleviate the bound of TCC on the energy scale of inflation at CMB window. Provided that the string landscape mostly consists of AdS vacua, we might have such a multi-stage inflation model, see Fig. 2(a). Each stage of slow-roll inflation only happened around the maxima of the potential barriers separated by AdS vacua ($\Delta\phi < M_{\text{P}}$ for each $V > 0$ region).² Between the j th and $(j + 1)$ th stages, the field will rapidly roll over an AdS vacuum.

As an illustrative example, we model the effective potential $V_{\text{eff}}(\phi)$ as

$$V_{\text{eff}}(\phi) = V_0 \left(1 - \cos \left(\frac{M}{\sqrt{V_0}} \phi \right) \right) e^{-\phi/\beta} - \Lambda, \quad (9)$$

which is similar to Fig. 2(a), see also [47], where $\Lambda = \text{const.} > 0$ makes the minima of the potential to be AdS, and V_0 , M and β are the positive constants as well, which set the amplitude, period and envelope curve of $V_{\text{eff}}(\phi)$, respectively. The AdS potential (9) must satisfy the swampland distance conjecture as well as the de Sitter (or refined dS) conjecture. Considering $1 - \cos \left(\frac{M}{\sqrt{V_0}} \phi \right) = 0$ and the AdS well has a width, we have $\Delta\phi < \frac{2\pi\sqrt{V_0}}{M}$ for

² See also the hilltop inflation [44–46].




FIG. 2: (a) is the sketch of AdS landscape. In (b), we plot the evolution of Hubble parameter with the potential (9), in which $\Lambda = 10^{-12}M_{\text{P}}^4$, $V_0 = 8 * 10^{-10}M_{\text{P}}^4$, $\beta = 0.642M_{\text{P}}$ and $M = 12\sqrt{V_0}M_{\text{P}}^{-1}$. These parameters satisfy the restrictions (10) and (11) of the swampland conjectures.

$V(\phi) > 0$. Hence, the swampland distance conjecture $|\Delta\phi|/M_{\text{P}} < \mathcal{O}(1)$ requires

$$\frac{\sqrt{V_0}}{M} < \mathcal{O}(1)\frac{M_{\text{P}}}{2\pi}. \quad (10)$$

Confronting the maxima of potential (9) (at which $\cos\left(\frac{M}{\sqrt{V_0}}\phi\right) \approx -1$) with the refined de Sitter conjecture $M_{\text{P}}^2 \frac{\min(\nabla_\phi \nabla_\phi V)}{V} \leq -c' \sim \mathcal{O}(1)$, we have

$$2V_0/\beta^2 - M^2 < -\mathcal{O}(1)\left(\frac{2V_0 - \Lambda}{M_{\text{P}}^2}\right). \quad (11)$$

We will show the possibility of the multi-stage inflation in such an AdS landscape (9). We assume, for simplicity, that each-stage slow-roll inflation happened around one of the maxima of the potential (which is much larger than the depth of AdS well, i.e., $V_{\text{inf}}^j \gg |\Lambda|$) and ended when inflaton rolls down the hill rapidly. When the j th-stage inflation ends, the potential energy V_{inf}^j will be thoroughly converted to $\dot{\phi}^2$, so that the field can rapidly roll over an AdS well. During this era, which corresponds to $w_j \simeq 1$, we have $\dot{\phi} \sim a^{-3}$ and $\rho_\phi \approx \dot{\phi}^2/2 \sim a^{-6}$. According to $3M_{\text{P}}^2 H^2 = \rho_\phi$, we get $\dot{\phi} \approx \sqrt{2/3}M_{\text{P}}/t$. Note that this result is valid only if $\dot{\phi}^2(t) \gg |\Lambda|$. After it rapidly rolls over the AdS well, the field must be able to slowly climb over the next maximum of the potential ($\dot{\phi}^2 \ll V_{\text{inf}}^{j+1}$), so that the $(j+1)$ th-stage inflation can happen. During the period $\dot{\phi}^2 \gg |V_{\text{eff}}|$, the shift of ϕ is approximately

$$\Delta\phi = \int_{t_{\text{ini}}}^{t_{\text{end}}} \dot{\phi} dt \approx \sqrt{\frac{2}{3}}M_{\text{P}} \ln\left(\frac{t_{\text{end}}}{t_{\text{ini}}}\right). \quad (12)$$

Base on the above analysis, we have the ratio between the potential energy of the $(j + 1)$ th and j th-stages inflation as $V_{\text{inf}}^{j+1}/V_{\text{inf}}^j \approx \left(\dot{\phi}^{\text{end}}/\dot{\phi}^{\text{ini}}\right)^2 \approx (t^{\text{ini}}/t^{\text{end}})^2$. Using Eq. (12), we find

$$\frac{V_{\text{inf}}^{j+1}}{V_{\text{inf}}^j} \approx \exp\left(-\sqrt{6}\frac{\Delta\phi}{M_{\text{P}}}\right). \quad (13)$$

Therefore, the AdS landscape (9) is viable for the multi-stage inflation. Actually, from (9) and (13), we find $\beta \approx \mathcal{O}(M_{\text{P}}/\sqrt{6}) \approx \mathcal{O}(0.4M_{\text{P}})$.

We plot the evolution of the background (the Hubble parameter) with the AdS potential (9) in Fig. 2(b), which clearly is a multi-stage inflation. The parameters V_0 , M , β and Λ used in Fig. 2(b) satisfy the bounds (10) and (11). Here, we have assumed $V_{\text{inf}}^i \gg |\Lambda|$, thus the ratio $H_{\text{inf}}^{j+1}/H_{\text{inf}}^j$ is small. Generally, the AdS minima have different values. Consequently, we may not have $V_{\text{inf}}^i \gg |\Lambda|$ for each V_{inf}^i , which will results in a larger $H_{\text{inf}}^{j+1}/H_{\text{inf}}^j$.

III. IMPLICATIONS FOR DARK ENERGY

The current universe is in an accelerated expanding phase. Nearly 70% components of the universe is dark energy (DE) [48] and its equation of state parameter is approximately $w \simeq -1$. Thus this accelerated expanding stage has similar behavior as the primordial inflation, which hence may be thought of as one stage of the multi-stage inflation scenario, see Fig. 3. According to Sec. II A, the TCC will put a bound on the lifetime of the DE era.

For simplicity, we assume that after the 1st-stage inflation ($H_1 = H_{\text{inf}}$), the universe will experience the hot BB evolution up to present DE era (the 2nd-stage inflation, $H_2 = H_{\text{DE}}$). Thus setting $j = 2$ in (6), we have the bound of TCC on the efolding number of DE era as

$$\tilde{N}_{\text{DE}} < \ln\left(\frac{M_{\text{P}}}{H_{\text{inf}}^{\text{ini}}}\right), \quad (14)$$

since $\tilde{N}_{w_1} = -\tilde{N}_1$. Here, we do not need to make allowance for the CMB window, which actually corresponds to set $a_{\text{DE}}^{\text{ini}}H_{\text{DE}}^{\text{ini}} \approx a_{\text{inf}}^{\text{ini}}H_{\text{inf}}^{\text{ini}}$. Therefore, if $H_{\text{DE}} = H_0$ is approximately constant, the lifetime of current DE era will be

$$\Delta t < \frac{1}{H_0} \ln\left(\frac{M_{\text{P}}}{H_{\text{inf}}^{\text{end}}}\right), \quad (15)$$

which is far smaller than the expected value $\frac{1}{H_0} \ln(\frac{M_{\text{P}}}{H_0})$, since $H_{\text{inf}}^{\text{end}} \gg H_0$ (even if $H_{\text{inf}}^{\text{end}}$ has reduced to $\sim 0.1\text{GeV}$ [10]). Intriguingly, if the TCC is correct, it seems that the fate of our




FIG. 3: The sketch of dark energy era confronted with the TCC.

current universe has been fixed by the property of primordial inflation, rather than that of DE itself.

It is also possible that the origin of the primordial perturbation consistent with the observations is attributed to alternatives to inflation (e.g., [49–53]). It is interesting to see what will happen to \tilde{N}_{DE} with these alternatives. Considering a nonsingular bouncing scenario, we have $a^{\text{min}} = a_{\text{b}}$, where a_{b} is evaluated at the bounce point. According to (4), assuming that the universe will start to hot-BB-like expand immediately after the bounce (i.e., $a_{\text{b}} \approx a_{\text{BB}}^{\text{ini}}$), we have

$$\tilde{N}_{\text{DE}} < \ln \left(\frac{a_{\text{b}}/l_{\text{P}}}{a_{\text{DE}}^{\text{ini}} H_{\text{DE}}^{\text{ini}}} \right) \approx \left| \tilde{N}_{\text{BB}} \right| + \ln \left(\frac{M_{\text{P}}}{H_{\text{BB}}^{\text{ini}}} \right), \quad (16)$$

where $\tilde{N}_{\text{BB}} = \ln \left(\frac{a_{\text{DE}}^{\text{ini}} H_{\text{DE}}^{\text{ini}}}{a_{\text{BB}}^{\text{ini}} H_{\text{BB}}^{\text{ini}}} \right) < 0$ is the e-folding number of the hot BB evolution before the DE era. Here, (16) is applicable no matter the scenario is ekpyrosis [49] or matter bounce [50]. Also, for the slow expansion scenario (in which the expansion is ultra-slow with $\epsilon = -\dot{H}/H^2 \ll -1$ before the hot BB evolution) [52, 53], the result is similar.

Therefore, if the universe reheated rapidly after the end of inflation or its alternatives, such that $H_{\text{inf}}^{\text{end}} = H_{\text{BB}}^{\text{ini}}$, and also experienced the same hot BB expansion, we will see that the DE era in inflation scenario will last much shorter than that in alternatives to inflation. This is because the bound of TCC on inflation is much tighter than that on its alternatives,

as pointed out in Ref. [10]. Interestingly, if the TCC is correct, the lifetime of DE era might be a probe for distinguishing different possibilities of the origins of the universe.

IV. DISCUSSION

As one possible criteria of consistent EFTs, the TCC has set an upper bound on the energy scale of inflation, which renders r at CMB window negligibly small. Recently, the implications of TCC for inflation has inspired the study for the multi-stage inflation.

We show that if the TCC is correct and the first-stage inflation is responsible for the primordial perturbations at the CMB window, the efolding number that the j th stage ($j \neq 1$) is able to last will be bounded by the energy scale of the first-stage inflation. Regarding the DE era as one stage of the multi-stage inflation scenario, we point out that in the inflation scenario the lifetime of current DE era is far smaller than expected, which might be a probe for distinguishing inflation from its alternatives. Our result might also has potential constraints on EFTs of DE. Relevant issues will be interesting for future study.

Provided that the landscape of EFTs mostly consists of AdS vacua, we present a multi-stage inflation model in AdS landscape (with AdS vacua separated by the barriers $V > 0$). Each inflationary stage only happened around the maxima of the potential barriers. The field will rapidly roll over an AdS vacuum between two stages. Such a model naturally satisfies all swampland conjectures. Note that the AdS potential is also present in past-complete pre-inflation stage [11]. Thus, if the TCC is correct, the issues related to the early universe and DE (especially in AdS landscape) will be worthy of further exploration.

Acknowledgments

YC is funded by China Postdoctoral Science Foundation, No. 2019M650810, and NSFC, No. 11905224. YSP is supported by NSFC, Nos. 11575188, 11690021.

-
- [1] A. H. Guth, Phys. Rev. D **23**, 347 (1981).
 - [2] A. A. Starobinsky, Phys. Lett. **91B**, 99 (1980).
 - [3] A. D. Linde, Phys. Lett. **108B**, 389 (1982).

- [4] A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
- [5] H. Ooguri and C. Vafa, Nucl. Phys. B **766**, 21 (2007) [hep-th/0605264].
- [6] G. Obied, H. Ooguri, L. Spodyneiko and C. Vafa, arXiv:1806.08362 [hep-th].
- [7] R. H. Brandenberger and J. Martin, Mod. Phys. Lett. A **16**, 999 (2001) [astro-ph/0005432].
- [8] J. Martin and R. H. Brandenberger, Phys. Rev. D **63**, 123501 (2001) [hep-th/0005209].
- [9] A. Bedroya and C. Vafa, arXiv:1909.11063 [hep-th].
- [10] A. Bedroya, R. Brandenberger, M. Loverde and C. Vafa, arXiv:1909.11106 [hep-th].
- [11] Y. Cai and Y. S. Piao, arXiv:1909.12719 [gr-qc].
- [12] T. Tenkanen, arXiv:1910.00521 [astro-ph.CO].
- [13] S. Das, arXiv:1910.02147 [hep-th].
- [14] S. Mizuno, S. Mukohyama, S. Pi and Y. L. Zhang, arXiv:1910.02979 [astro-ph.CO].
- [15] S. Brahma, arXiv:1910.04741 [hep-th].
- [16] M. Dhuria and G. Goswami, arXiv:1910.06233 [astro-ph.CO].
- [17] M. Torabian, arXiv:1910.06867 [hep-th].
- [18] R. G. Cai and S. J. Wang, arXiv:1910.07981 [astro-ph.CO].
- [19] K. Schmitz, arXiv:1910.08837 [hep-ph].
- [20] K. Kadota, C. S. Shin, T. Terada and G. Tumurtushaa, arXiv:1910.09460 [hep-th].
- [21] A. Berera and J. R. Caldern, arXiv:1910.10516 [hep-ph].
- [22] S. Brahma, arXiv:1910.12352 [hep-th].
- [23] N. Okada, D. Raut and Q. Shafi, arXiv:1910.14586 [hep-ph].
- [24] S. Laliberte and R. Brandenberger, arXiv:1911.00199 [hep-th].
- [25] W. C. Lin and W. H. Kinney, arXiv:1911.03736 [gr-qc].
- [26] G. Dvali and S. Kachru, In *Shifman, M. (ed.) et al.: From fields to strings, vol. 2* 1131-1155 [hep-th/0309095].
- [27] C. P. Burgess, R. Easther, A. Mazumdar, D. F. Mota and T. Multamaki, JHEP **0505**, 067 (2005) [hep-th/0501125].
- [28] Y. Liu, Y. S. Piao and Z. G. Si, JCAP **0905**, 008 (2009) [arXiv:0901.2058 [hep-th]].
- [29] M. Li and Y. Wang, JCAP **0907**, 033 (2009) [arXiv:0903.2123 [hep-th]].
- [30] S. Li, Y. Liu and Y. S. Piao, Phys. Rev. D **80**, 123535 (2009) [arXiv:0906.3608 [hep-th]].
- [31] Y. Wang, arXiv:1001.0008 [hep-th].
- [32] N. Afshordi, A. Slosar and Y. Wang, JCAP **1101**, 019 (2011) [arXiv:1006.5021 [astro-ph.CO]].

- [33] S. K. Garg and C. Krishnan, arXiv:1807.05193 [hep-th].
- [34] H. Ooguri, E. Palti, G. Shiu and C. Vafa, Phys. Lett. B **788**, 180 (2019) [arXiv:1810.05506 [hep-th]].
- [35] A. Achucarro and G. A. Palma, JCAP **1902**, 041 (2019) [arXiv:1807.04390 [hep-th]].
- [36] A. Kehagias and A. Riotto, Fortsch. Phys. **66**, no. 10, 1800052 (2018) [arXiv:1807.05445 [hep-th]].
- [37] H. Matsui and F. Takahashi, Phys. Rev. D **99**, no. 2, 023533 (2019) [arXiv:1807.11938 [hep-th]].
- [38] W. H. Kinney, S. Vagnozzi and L. Visinelli, Class. Quant. Grav. **36**, no. 11, 117001 (2019) [arXiv:1808.06424 [astro-ph.CO]].
- [39] S. Brahma and M. Wali Hossain, JHEP **1903**, 006 (2019) [arXiv:1809.01277 [hep-th]].
- [40] M. Motaharfar, V. Kamali and R. O. Ramos, Phys. Rev. D **99**, no. 6, 063513 (2019) [arXiv:1810.02816 [astro-ph.CO]].
- [41] A. Ashoorioon, Phys. Lett. B **790**, 568 (2019) [arXiv:1810.04001 [hep-th]].
- [42] S. Das, Phys. Rev. D **99**, no. 6, 063514 (2019) [arXiv:1810.05038 [hep-th]].
- [43] W. H. Kinney, Phys. Rev. Lett. **122**, no. 8, 081302 (2019) [arXiv:1811.11698 [astro-ph.CO]].
- [44] V. N. Senoguz and Q. Shafi, Phys. Lett. B **596**, 8 (2004) [hep-ph/0403294].
- [45] L. Boubekur and D. H. Lyth, JCAP **0507**, 010 (2005) [hep-ph/0502047].
- [46] K. Kohri, C. M. Lin and D. H. Lyth, JCAP **0712**, 004 (2007) [arXiv:0707.3826 [hep-ph]].
- [47] Y. S. Piao, Phys. Rev. D **70**, 101302 (2004) [hep-th/0407258].
- [48] N. Aghanim *et al.* [Planck Collaboration], arXiv:1807.06209 [astro-ph.CO].
- [49] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, Phys. Rev. D **64**, 123522 (2001) [hep-th/0103239].
- [50] F. Finelli and R. Brandenberger, Phys. Rev. D **65**, 103522 (2002) [hep-th/0112249]; D. Wands, Phys. Rev. D **60**, 023507 (1999) [gr-qc/9809062].
- [51] R. H. Brandenberger and C. Vafa, Nucl. Phys. B **316**, 391 (1989); A. Nayeri, R. H. Brandenberger and C. Vafa, Phys. Rev. Lett. **97**, 021302 (2006) [hep-th/0511140].
- [52] Y. S. Piao and E. Zhou, Phys. Rev. D **68**, 083515 (2003) [hep-th/0308080]; Z. G. Liu, J. Zhang and Y. S. Piao, Phys. Rev. D **84**, 063508 (2011) [arXiv:1105.5713].
- [53] P. Creminelli, A. Nicolis and E. Trincherini, JCAP **1011**, 021 (2010) [arXiv:1007.0027 [hep-th]].