

Flavour Anomalies

Antonio Pich*

IFIC, Universitat de València – CSIC

Parc Científic, Catedrático José Beltrán 2, E-46980 Paterna, Spain

E-mail: Antonio.Pich@ific.uv.es

The experimental data on $b \rightarrow c\tau\bar{\nu}_\tau$ and $b \rightarrow s\ell^+\ell^-$ transitions exhibit sizeable discrepancies with the Standard Model expectations. We present an overview of the present status and discuss possible interpretations within a model-independent effective Lagrangian approach. We also briefly elaborate on some other claimed flavour anomalies such as the recently observed CP asymmetry in D^0 decays or the $K^0 \rightarrow \pi\pi$ ratio $\varepsilon'_K/\varepsilon_K$. The Standard Model prediction for the direct CP-violating ratio $\varepsilon'_K/\varepsilon_K$ agrees with its measured value, once all theoretical ingredients are correctly taken into account.

*7th Annual Conference on Large Hadron Collider Physics - LHCP2019
20-25 May, 2019
Puebla, Mexico*

*Speaker.

1. Introduction

The Standard Model (SM) provides a very successful description of flavour-changing transitions in terms of the nine charged-fermion masses and the four CKM quark-mixing parameters (plus the analogous neutrino masses and mixings, when they are incorporated). However, a more fundamental understanding of flavour is still lacking. Obviously, any observed deviations from the SM predictions trigger a lot of interest as they could provide the missing hints to uncover the underlying flavour dynamics.

Many flavour-related ‘anomalies’ have been reported along the years: $W \rightarrow \tau\nu$, $\tau \rightarrow \pi K_S \nu$, $b \rightarrow c\tau\nu$, $b \rightarrow s\mu^+\mu^-$, $(g-2)_{\mu,e}$, $\varepsilon'_K/\varepsilon_K$ ($K^0 \rightarrow \pi\pi$), ε_K , ΔA_{CP} ($D^0 \rightarrow K^+K^-/\pi^+\pi^-$), V_{cd} , V_{ub} , V_{ud} , $B \rightarrow \tau\nu$, ... While some of them could indeed be true signals of new phenomena, others may just originate from statistical fluctuations, underestimated systematics or even a deficient SM prediction or measurement. For instance, the long-term discrepancy between exclusive and inclusive determinations of V_{cd} seems nowadays close to get solved through a more careful treatment of form factor parameters and data extrapolations [1, 2, 3, 4, 5, 6, 7]. The resulting (slightly larger) value of V_{cd} would also eliminate the suggested tensions in ε_K , while the claimed anomaly in $\varepsilon'_K/\varepsilon_K$ results from a poor theoretical treatment of the final-state $\pi\pi$ dynamics [8].

In fact, an easy common explanation for all anomalies does not exist, within appealing models of new physics (NP). Model builders just choose two or three anomalies, according to their preferences, in order to fit them together within the same theoretical framework. In this situation, separate analyses of the different observables are perhaps more enlightening. In the following, I will focus on those topics where there has been more activity during the last two years.

2. The kaon CP-violating ratio $\varepsilon'_K/\varepsilon_K$

A tiny difference between the CP-violating ratios $\eta_{nm} \equiv \mathcal{M}[K_L^0 \rightarrow \pi^n \pi^m]/\mathcal{M}[K_S^0 \rightarrow \pi^n \pi^m] \approx \varepsilon_K \approx 2.2 \times 10^{-3} e^{i\pi/4}$, where $nm = +-, 00$ denote the final pion charges, was first measured by the CERN NA31 experiment [9] and later confirmed at the 7.2σ level with the full data samples of NA31, NA48 and the Fermilab experiments E731 and KTeV [10]:

$$\text{Re}(\varepsilon'_K/\varepsilon_K) = \frac{1}{3} \left(1 - \left| \frac{\eta_{00}}{\eta_{+-}} \right|^2 \right) = (16.6 \pm 2.3) \times 10^{-4}. \quad (2.1)$$

This important measurement established the presence of direct CP violation in the decay amplitudes, confirming that CP violation is associated with a $\Delta S = 1$ transition as predicted by the CKM mechanism.

The first next-to-leading-order theoretical predictions gave values of $\varepsilon'_K/\varepsilon_K$ one order of magnitude smaller than (2.1), but it was soon realised that they were missing the important role of the final pion dynamics [11, 12, 13]. Once long-distance contributions are properly taken into account, the theoretical SM prediction turns out to be in good agreement with the experimental value, although the uncertainties are unfortunately large [8, 14]:

$$\text{Re}(\varepsilon'_K/\varepsilon_K)_{\text{SM}} = (14 \pm 5) \times 10^{-4}. \quad (2.2)$$

The underlying physics can be easily understood from the kaon data themselves. Owing to Bose symmetry, the two pions in the final state must be in a $I = 0$ or $I = 2$ configuration. In the absence of QCD corrections, the corresponding $K \rightarrow \pi\pi$ decay amplitudes $\mathcal{A}_I \equiv A_I e^{i\delta_I}$ are predicted to differ only by a $\sqrt{2}$ factor. However, their measured ratio is 16 times larger than that (a truly spectacular enhancement generated by the strong forces):

$$\omega \equiv \text{Re}(A_2)/\text{Re}(A_0) \approx 1/22, \quad \delta_0 - \delta_2 \approx 45^\circ. \quad (2.3)$$

Moreover, they exhibit a huge phase-shift difference that manifests the relevance of final-state interactions and, therefore, the presence of large absorptive contributions to the $K \rightarrow \pi\pi$ amplitudes, specially to the isoscalar one. Writing $\mathcal{A}_I = \text{Dis}(\mathcal{A}_I) + i\text{Abs}(\mathcal{A}_I)$ and neglecting the small CP -odd components, the measured $\pi\pi$ scattering phase shifts at $\sqrt{s} = m_K$ imply that

$$\text{Abs}(A_0)/\text{Dis}(A_0) \approx 0.82, \quad \text{Abs}(A_2)/\text{Dis}(A_2) \approx -0.15. \quad (2.4)$$

The direct CP -violating effect involves the interference between the two isospin amplitudes,

$$\text{Re}(\varepsilon'_K/\varepsilon_K) = -\frac{\omega}{\sqrt{2}|\varepsilon_K|} \left[\frac{\text{Im}A_0}{\text{Re}A_0} - \frac{\text{Im}A_2}{\text{Re}A_2} \right] = -\frac{\omega_+}{\sqrt{2}|\varepsilon_K|} \left[\frac{\text{Im}A_0^{(0)}}{\text{Re}A_0^{(0)}} (1 - \Omega_{\text{eff}}) - \frac{\text{Im}A_2^{\text{emp}}}{\text{Re}A_2^{(0)}} \right]. \quad (2.5)$$

It is suppressed by the small ratio ω and, moreover, it is very sensitive to isospin-breaking (IB) corrections [15, 16, 17], parametrized by $\Omega_{\text{eff}} = 0.11 \pm 0.09$ [14], because small IB corrections to A_0 feed into the small amplitude A_2 enhanced by the large factor $1/\omega$. In the right-hand side of Eq. (2.5), the (0) superscript indicates the isospin limit, $\omega_+ = \text{Re}(A_2^+)/\text{Re}(A_0)$ is directly extracted from $K^+ \rightarrow \pi^+\pi^0$ and A_2^{emp} contains the electromagnetic-penguin contribution to A_2 (the remaining contributions are included in Ω_{eff}).

Claims for small SM values of $\varepsilon'_K/\varepsilon_K$ usually originate from perturbative calculations that are unable to generate the physical phase shifts, *i.e.*, they predict $\delta_I = 0$ and, therefore, $\text{Abs}(\mathcal{A}_I) = 0$, failing completely to understand the empirical ratios (2.4). This unitarity pitfall implies also incorrect predictions for the dispersive components, since they are related by analyticity with the absorptive parts: a large absorptive contribution generates a large dispersive correction that is obviously missed in those calculations. This perturbative problem is more severe in $\varepsilon'_K/\varepsilon_K$ because Eq. (2.5) involves a delicate numerical balance among the three contributing terms, and naive predictions sit precisely on a nearly-exact cancellation (a 40% positive correction to the first term enhances the whole result by one order of magnitude).

The $\varepsilon'_K/\varepsilon_K$ anomaly was recently resurrected by the lattice RBC-UKQCD collaboration that reported $\text{Re}(\varepsilon'_K/\varepsilon_K) = (1.38 \pm 5.15 \pm 4.59) \times 10^{-4}$ [18, 19]. The uncertainties are still large, but the quite low central value implies a 2.1σ deviation from the experimental measurement. This has triggered a revival of the old naive estimates [20, 21], some of them making also use of the lattice data [22, 23], and a large amount of NP explanations (a list of references is given in Refs. [8, 14]). However, it is premature to derive physics implications from the current lattice simulations, since they are still unable to reproduce the known phase shifts. While the lattice determination of δ_2 is only 1σ away from its physical value, δ_0 disagrees with the experimental result by 2.9σ , a much larger discrepancy than the one quoted for $\varepsilon'_K/\varepsilon_K$. Obviously, nobody suggests a NP contribution to the $\pi\pi$ elastic scattering phase shifts. The RBC-UKQCD collaboration is actively working in order to improve the present situation.

3. Direct CP violation in charm

The first observation of CP violation in decays of charm hadrons has been recently reported by the LHCb collaboration, which has measured the difference between the CP asymmetries in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ [24]:

$$\Delta A_{CP} \equiv A_{CP}(K^+K^-) - A_{CP}(\pi^+\pi^-) = (-15.4 \pm 2.9) \times 10^{-4}, \quad (3.1)$$

where $A_{CP}(f) \equiv [\Gamma(D^0 \rightarrow f) - \Gamma(\bar{D}^0 \rightarrow f)] / [\Gamma(D^0 \rightarrow f) + \Gamma(\bar{D}^0 \rightarrow f)]$. The measured time-integrated asymmetries can be decomposed in a direct contribution $a_{CP}^{\text{dir}}(f)$ from CP violation in the decay amplitude and another component associated to CP violation either in D^0 - \bar{D}^0 mixing or in the interference between mixing and decay. These two contributions can be disentangled through measurements of the time-decay distribution of the data sample. The LHCb analysis concludes that, as expected, the CP signal comes primarily from direct CP violation: $\Delta a_{CP}^{\text{dir}} = (-15.7 \pm 2.9) \times 10^{-4}$ [24]. The HFLAV combination with previous measurements, shown in Figure 1, gives [25]

$$\Delta a_{CP}^{\text{dir}} = (-16.4 \pm 2.8) \times 10^{-4}. \quad (3.2)$$




Figure 1: World data on the direct and indirect CP-violating contributions $\Delta a_{CP}^{\text{dir}}$ and a_{CP}^{ind} [25]. The cross indicates the best fit values with their 1σ uncertainties, and the ellipses the two-dimensional 1σ , 3σ and 5σ regions. The filled circle shows the point $(0,0)$, where the two CP-violating amplitudes vanish.

The size of this asymmetry is larger than the naive SM expectation $|\Delta a_{CP}^{\text{dir}}| \leq 3 \times 10^{-4}$ [26, 27], based on perturbative QCD or light-cone sum rules estimates. However, the measured value could be explained within the SM with non-perturbative re-scattering effects of moderate size that enhance the U-spin invariant decay amplitude [28]. A direct CP asymmetry necessarily involves a non-zero strong phase-shift difference between (at least) two interfering amplitudes. Eq. (3.2) indicates the presence of a large phase-shift, which can obviously not be reproduced through perturbative calculations. A reliable SM prediction of $\Delta a_{CP}^{\text{dir}}$, including re-scattering corrections [29, 30, 31, 32], remains an interesting theoretical challenge.

4. $b \rightarrow c\tau\bar{\nu}_\tau$ transitions

Sizeable deviations from their predicted SM values have been found in the ratios

$$R(D^{(*)}) \equiv \frac{\text{Br}(\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau)}{\text{Br}(\bar{B} \rightarrow D^{(*)} \ell^- \bar{\nu}_\ell)}, \quad (4.1)$$

with $\ell = e, \mu$. The experimental averages shown in Figure 2 exhibit a 3.1σ discrepancy with the SM predictions quoted by the HFLAV group [25], which increases to 3.7σ with the more updated theoretical values given in Ref. [33]:

$$R(D)^{\text{SM}} = 0.300^{+0.005}_{-0.004}, \quad R(D^*)_{\text{SM}} = 0.251^{+0.004}_{-0.003}. \quad (4.2)$$

A large part of the hadronic form factor uncertainties cancels in these ratios, which are also independent of V_{cb} . Therefore, the measurements suggest a rather large violation of lepton-flavour universality that is quite unexpected in a tree-level SM transition. A similar 1.7σ deviation has been also found for the analogous $R(J/\psi)$ ratio of semileptonic $B_c \rightarrow J/\psi$ transitions [34]. Moreover, the measured longitudinal polarization of the D^{*-} meson in $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$ [35] also differs from its SM value by 1.6σ .




Figure 2: $R(D)$ and $R(D^*)$ measurements, their world average (red ellipse) and SM predictions (cross) [25].

From the SM inclusive prediction $\text{Br}(B \rightarrow X_c \tau \nu) / \text{Br}(B \rightarrow X_c e \nu) = (0.222 \pm 0.007)$ [36, 37], which does not involve any form factors, and $\text{Br}(B \rightarrow X_c \ell \nu) = (10.65 \pm 0.16)\%$ [10], one finds that $\text{Br}(B \rightarrow X_c \tau \nu) = (2.36 \pm 0.08)\%$, in agreement with the LEP result $\text{Br}(b \rightarrow X_c \tau \nu) = (2.41 \pm 0.23)\%$ [10]. Since the measured $R(D^{(*)})$ ratios imply $\text{Br}(B \rightarrow D \tau \nu) + \text{Br}(B \rightarrow D^* \tau \nu) = (2.33 \pm 0.11)\%$, these two final states saturate the inclusive width and there is no space left for other decay modes, such as the D^{**} , that are expected to contribute more than 0.5% [36]. Thus, there is also tension with the SM at the inclusive level.

The anomaly is dominated by the 2012–2013 BaBar results [38, 39], while the most recent Belle [40, 41, 42] and LHCb [43, 44, 45] measurements are closer to the SM values. Moreover, the normalized q^2 distributions measured by BaBar [39] and Belle [40] do not favour large deviations from the SM [37]. One must also take into account that the needed enhancement of the $b \rightarrow c\tau\nu$ transition is constrained by the cross-channel $b\bar{c} \rightarrow \tau\nu$. A conservative (more stringent) upper bound $\text{Br}(B_c \rightarrow \tau\nu) < 30\%$ (10%) can be extracted from the B_c lifetime [37, 46] (LEP data [47]).

Taking the available experimental information at face value, one can investigate the possible types of underlying NP interactions with a generic low-energy effective Hamiltonian,

$$\mathcal{H}_{\text{eff}}^{b \rightarrow c \ell \nu} = \frac{4G_F}{\sqrt{2}} V_{cb} \{ (1 + C_{V_L}) \mathcal{O}_{V_L} + C_{V_R} \mathcal{O}_{V_R} + C_{S_R} \mathcal{O}_{S_R} + C_{S_L} \mathcal{O}_{S_L} + C_T \mathcal{O}_T \} + \text{h.c.} \quad (4.3)$$

where

$$\mathcal{O}_{V_{L,R}} = (\bar{c} \gamma^\mu b_{L,R}) (\bar{\ell}_L \gamma_\mu \nu_{\ell L}), \quad \mathcal{O}_{S_{L,R}} = (\bar{c} b_{L,R}) (\bar{\ell}_R \nu_{\ell L}), \quad \mathcal{O}_T = (\bar{c} \sigma^{\mu\nu} b_L) (\bar{\ell}_R \sigma_{\mu\nu} \nu_{\ell L}). \quad (4.4)$$

The SM corresponds to $C_i = 0$. Since potential NP contributions to the light-lepton couplings are highly constrained by $b \rightarrow c(e, \mu) \nu$ data [48], one can safely consider that NP effects are only present for the τ . A global fit to all available experimental information has been recently done in Ref. [33],¹ neglecting CP-violating contributions (*i.e.*, with real C_i) and taking $C_{V_R} = 0$. The second condition follows from the assumption that the electroweak symmetry breaking is linearly realized at the electroweak scale, which implies that C_{V_R} is flavour independent [49]. The fitted results clearly indicate that NP contributions are needed (much lower χ^2 than in the SM), but they do not show any strong preference for a particular Wilson coefficient ($\chi_{\text{min}}^2/\text{d.o.f.} = 37.4/54$) [33]:

$$C_{V_L} = 0.09_{-0.12}^{+0.13}, \quad C_{S_R} = 0.09_{-0.61}^{+0.12}, \quad C_{S_L} = -0.14_{-0.07}^{+0.52}, \quad C_T = 0.008_{-0.044}^{+0.046}. \quad (4.5)$$

While none of the fitted coefficients are required to be non-zero, the simplest interpretation of this solution is a global modification of the SM. In fact, setting all coefficients but C_{V_L} to zero one also gets a good fit. In addition to this SM-like global minimum, a second local minimum ($\chi_{\text{min}}^2/\text{d.o.f.} = 40.4/54$) is found with larger non-SM contributions [33]:

$$C_{V_L} = 0.34_{-0.07}^{+0.05}, \quad C_{S_R} = -1.10_{-0.07}^{+0.48}, \quad C_{S_L} = -0.30_{-0.50}^{+0.11}, \quad C_T = 0.093_{-0.030}^{+0.029}. \quad (4.6)$$

The measured D^* longitudinal polarization fraction $F_L^{D^*}$ has a strong impact on the analysis because, with the four fitted operators, its predicted value remains always below the 1σ experimental region. Including C_{V_R} in the fit helps to remove the tension with the $B \rightarrow D^*$ data and opens new (not satisfactory) fine-tuned solutions where the SM coefficient becomes very small, its effect being substituted by several sizeable NP contributions, especially C_{V_R} . More precise experimental data is needed to clarify the current situation. If the $b \rightarrow c$ anomaly remains, an improved measurement of $F_L^{D^*}$ could have major implications in its theoretical interpretation.

5. $b \rightarrow s \ell \ell$ transitions

Several $b \rightarrow s \mu^+ \mu^-$ rates have been found at LHCb to be consistently lower than their SM predictions: $B^+ \rightarrow K^+ \mu^+ \mu^-$ [50, 51], $B^+ \rightarrow K^{*+} \mu^+ \mu^-$ [50], $B_d^0 \rightarrow K^0 \mu^+ \mu^-$ [50], $B_d^0 \rightarrow K^{*0} \mu^+ \mu^-$ [52, 53, 54, 55], $B_s^0 \rightarrow \phi \mu^+ \mu^-$ [56, 57] and $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ [58, 59]. The angular and invariant-mass distributions of the final decay products in $B \rightarrow K^* \mu^+ \mu^-$ have been also studied by ATLAS [60], BaBar [61], Belle [62, 63, 64], CDF [65], CMS [66, 67] and LHCb [52, 53, 54, 55]. The

¹An extensive list of references to previous analyses, most of them performed with a single mediator or operator and with partial data information, can be found in Ref. [33].

four-body $K\pi\mu^+\mu^-$ final state provides a rich variety of angular dependences, making possible to disentangle the different dynamical contributions. Particular attention has been devoted to specific combinations of angular observables that are free from form-factor uncertainties in the heavy-quark mass limit, the so called optimized observables $P'_i(q^2)$ [68], where q^2 is the dilepton invariant-mass squared. A sizeable discrepancy with the SM prediction [69, 70, 71], shown in Figure 3, has been identified in two adjacent bins of the P'_5 distribution, just below the J/ψ peak. Belle has also analyzed $K^*e^+e^-$ final states [62, 63, 64], finding them compatible with the SM expectations.




Figure 3: Comparison between the predicted SM values of P'_5 and the experimental measurements [72].

The SM predictions for the previous observables suffer from hadronic uncertainties that are not easy to quantify. However, LHCb has also reported sizeable violations of lepton universality, at the 2.1-2.5 σ level, through the ratios [73]

$$R_{K^{*0}} \equiv \frac{\Gamma(B_d^0 \rightarrow K^{*0}\mu^+\mu^-)}{\Gamma(B_d^0 \rightarrow K^{*0}e^+e^-)} = \begin{cases} 0.66_{-0.07}^{+0.11} \pm 0.03, & q^2 \in [0.045, 1.1] \text{ GeV}^2, \\ 0.69_{-0.07}^{+0.11} \pm 0.05, & q^2 \in [1.1, 6.0] \text{ GeV}^2, \end{cases} \quad (5.1)$$

and [74]

$$R_K \equiv \frac{\Gamma(B^+ \rightarrow K^+\mu^+\mu^-)}{\Gamma(B^+ \rightarrow K^+e^+e^-)} \Big|_{q^2 \in [1.1, 6.0] \text{ GeV}^2} = 0.846_{-0.054}^{+0.060} {}_{-0.014}^{+0.016}, \quad (5.2)$$

which constitute very clean probes of NP contributions. Owing to their larger uncertainties, the recent Belle measurements of R_{K^*} [75] and R_K [76] are compatible with the SM as well as with LHCb.

Global fits to the $b \rightarrow s\ell^+\ell^-$ data with an effective low-energy Lagrangian

$$\mathcal{L}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{td} V_{ts}^* \frac{\alpha}{\pi} \sum_{i,\ell} C_{i,\ell} O_i^\ell \quad (5.3)$$

show a clear preference for NP contributions to the operators $O_9^\ell = (\bar{s}_L \gamma_\mu b_L)(\bar{\ell} \gamma^\mu \ell)$ and $O_{10}^\ell = (\bar{s}_L \gamma_\mu b_L)(\bar{\ell} \gamma^\mu \gamma_5 \ell)$, with $\ell = \mu$ [77, 78, 79, 80, 81, 82, 83]. Although the different analyses tend to favour slightly different solutions, two main common scenarios stand out: either $\delta C_{9,\mu}^{\text{NP}} \approx -0.98$ or $\delta C_{9,\mu}^{\text{NP}} = -\delta C_{10,\mu}^{\text{NP}} \approx -0.46$. Both constitute large shifts (-24% and -11% , respectively) from the SM values: $C_{9,\mu}^{\text{SM}}(\mu_b) \approx 4.1$ and $C_{10,\mu}^{\text{SM}}(\mu_b) \approx -4.3$, at $\mu_b = 4.8$ GeV. The first possibility is slightly

preferred by the global analysis of all data, while the left-handed NP solution accommodates better the lepton-flavour-universality-violating observables [78].

The left-handed scenario is theoretically appealing because it can be easily generated through $SU(2)_L \otimes U(1)_Y$ -invariant effective operators at the electroweak scale that, moreover, could also provide an explanation to the $b \rightarrow c\tau\nu$ anomaly. This possibility emerges naturally from the so-called U_1 vector leptoquark model [84], and can be tested experimentally, since it implies a $b \rightarrow s\tau^+\tau^-$ rate three orders of magnitude larger than the SM expectation [85]. For a recent review of theoretical models with a quite complete list of references, see Ref. [86].

6. Summary

Uncovering the fundamental dynamics behind flavour-changing transitions and CP-violating phenomena is one of the main pending questions in particle physics. In the SM, flavour emerges from the Yukawa interactions with the scalar Higgs doublet, the less understood part of the electroweak Lagrangian that is more open to theoretical speculations.

Sizeable deviations from the SM expectations have been identified in $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell\ell$ data. Whether they represent the first signals of new phenomena or just result from statistical fluctuations and/or underestimated systematics remains to be understood. New experimental input from LHC and Belle-II should soon clarify the situation. A confirmation of the current flavour anomalies would constitute clear evidence of NP interactions and, moreover, would allow us to infer their low-energy structure, providing precious hints on the underlying dynamics.

Acknowledgements

I want to thank the organizers of LHCP2019 for the invitation to present this overview. I also thank V. Cirigliano, H. Gisbert, M. Jung, C. Murgui, A. Peñuelas and A. Rodríguez-Sánchez for a very productive and enjoyable collaboration. This work has been supported in part by the Spanish Government and ERDF funds from the EU Commission [grant FPA2017-84445-P], the Generalitat Valenciana [grant Prometeo/2017/053] and the Spanish Centro de Excelencia Severo Ochoa Programme [grant SEV-2014-0398].

References

- [1] D. Bigi and P. Gambino, *Revisiting $B \rightarrow D\ell\nu$* , *Phys. Rev.* **D94** (2016) 094008 [1606.08030].
- [2] D. Bigi, P. Gambino and S. Schacht, *A fresh look at the determination of $|V_{cb}|$ from $B \rightarrow D^*\ell\nu$* , *Phys. Lett.* **B769** (2017) 441 [1703.06124].
- [3] D. Bigi, P. Gambino and S. Schacht, *$R(D^*)$, $|V_{cb}|$, and the Heavy Quark Symmetry relations between form factors*, *JHEP* **11** (2017) 061 [1707.09509].
- [4] B. Grinstein and A. Kobach, *Model-Independent Extraction of $|V_{cb}|$ from $\bar{B} \rightarrow D^*\ell\nu$* , *Phys. Lett.* **B771** (2017) 359 [1703.08170].
- [5] F. U. Bernlochner, Z. Ligeti, M. Papucci and D. J. Robinson, *Combined analysis of semileptonic B decays to D and D^* : $R(D^{(*)})$, $|V_{cb}|$, and new physics*, *Phys. Rev.* **D95** (2017) 115008 [1703.05330].

- [6] F. U. Bernlochner, Z. Ligeti, M. Papucci and D. J. Robinson, *Tensions and correlations in $|V_{cb}|$ determinations*, *Phys. Rev.* **D96** (2017) 091503 [1708.07134].
- [7] P. Gambino, M. Jung and S. Schacht, *The V_{cb} puzzle: An update*, *Phys. Lett.* **B795** (2019) 386 [1905.08209].
- [8] H. Gisbert and A. Pich, *Direct CP violation in $K^0 \rightarrow \pi\pi$: Standard Model Status*, *Rept. Prog. Phys.* **81** (2018) 076201 [1712.06147].
- [9] NA31 collaboration, *First Evidence for Direct CP Violation*, *Phys. Lett.* **B206** (1988) 169.
- [10] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *Phys. Rev.* **D98** (2018) 030001.
- [11] E. Pallante and A. Pich, *Strong enhancement of ϵ'/ϵ through final state interactions*, *Phys. Rev. Lett.* **84** (2000) 2568 [hep-ph/9911233].
- [12] E. Pallante and A. Pich, *Final state interactions in kaon decays*, *Nucl. Phys.* **B592** (2001) 294 [hep-ph/0007208].
- [13] E. Pallante, A. Pich and I. Scimemi, *The Standard model prediction for ϵ'/ϵ* , *Nucl. Phys.* **B617** (2001) 441 [hep-ph/0105011].
- [14] V. Cirigliano, H. Gisbert, A. Pich and A. Rodríguez-Sánchez, *Isospin-Violating Contributions to ϵ'/ϵ* , 1911.01359.
- [15] G. Ecker, G. Muller, H. Neufeld and A. Pich, *π^0 - η mixing and CP violation*, *Phys. Lett.* **B477** (2000) 88 [hep-ph/9912264].
- [16] V. Cirigliano, A. Pich, G. Ecker and H. Neufeld, *Isospin violation in ϵ'* , *Phys. Rev. Lett.* **91** (2003) 162001 [hep-ph/0307030].
- [17] V. Cirigliano, G. Ecker, H. Neufeld and A. Pich, *Isospin breaking in $K \rightarrow \pi\pi$ decays*, *Eur. Phys. J.* **C33** (2004) 369 [hep-ph/0310351].
- [18] RBC, UKQCD collaboration, *Standard Model Prediction for Direct CP Violation in $K \rightarrow \pi\pi$ Decay*, *Phys. Rev. Lett.* **115** (2015) 212001 [1505.07863].
- [19] T. Blum et al., *$K \rightarrow \pi\pi$ $\Delta I = 3/2$ decay amplitude in the continuum limit*, *Phys. Rev.* **D91** (2015) 074502 [1502.00263].
- [20] A. J. Buras and J.-M. Gérard, *Upper bounds on ϵ'/ϵ parameters $B_6^{(1/2)}$ and $B_8^{(3/2)}$ from large N QCD and other news*, *JHEP* **12** (2015) 008 [1507.06326].
- [21] A. J. Buras and J.-M. Gerard, *Final state interactions in $K \rightarrow \pi\pi$ decays: $\Delta I = 1/2$ rule vs. ϵ'/ϵ* , *Eur. Phys. J.* **C77** (2017) 10 [1603.05686].
- [22] A. J. Buras, M. Gorbahn, S. Jäger and M. Jamin, *Improved anatomy of ϵ'/ϵ in the Standard Model*, *JHEP* **11** (2015) 202 [1507.06345].
- [23] T. Kitahara, U. Nierste and P. Tremper, *Singularity-free next-to-leading order $\Delta S = 1$ renormalization group evolution and ϵ'_K/ϵ_K in the Standard Model and beyond*, *JHEP* **12** (2016) 078 [1607.06727].
- [24] LHCb collaboration, *Observation of CP Violation in Charm Decays*, *Phys. Rev. Lett.* **122** (2019) 211803 [1903.08726].
- [25] HFLAV collaboration, *Averages of b -hadron, c -hadron, and τ -lepton properties as of 2018*, 1909.12524.
- [26] M. Chala, A. Lenz, A. V. Rusov and J. Scholtz, *ΔA_{CP} within the Standard Model and beyond*, *JHEP* **07** (2019) 161 [1903.10490].

- [27] A. Khodjamirian and A. A. Petrov, *Direct CP asymmetry in $D \rightarrow \pi^- \pi^+$ and $D \rightarrow K^- K^+$ in QCD-based approach*, *Phys. Lett.* **B774** (2017) 235 [1706.07780].
- [28] Y. Grossman and S. Schacht, *The emergence of the $\Delta U = 0$ rule in charm physics*, *JHEP* **07** (2019) 020 [1903.10952].
- [29] F. Buccella, A. Paul and P. Santorelli, *$SU(3)_F$ breaking through final state interactions and CP asymmetries in $D \rightarrow PP$ decays*, *Phys. Rev.* **D99** (2019) 113001 [1902.05564].
- [30] H.-N. Li, C.-D. Lü and F.-S. Yu, *Implications on the first observation of charm CPV at LHCb*, 1903.10638.
- [31] A. Soni, *Resonance enhancement of Charm CP*, 1905.00907.
- [32] H.-Y. Cheng and C.-W. Chiang, *Revisiting CP violation in $D \rightarrow PP$ and VP decays*, *Phys. Rev.* **D100** (2019) 093002 [1909.03063].
- [33] C. Murgui, A. Peñuelas, M. Jung and A. Pich, *Global fit to $b \rightarrow c\tau\nu$ transitions*, *JHEP* **09** (2019) 103 [1904.09311].
- [34] LHCb collaboration, *Measurement of the ratio of branching fractions $\mathcal{B}(B_c^+ \rightarrow J/\psi\tau^+\nu_\tau)/\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu)$* , *Phys. Rev. Lett.* **120** (2018) 121801 [1711.05623].
- [35] BELLE collaboration, *Measurement of the D^{*-} polarization in the decay $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$* , in *10th International Workshop on the CKM Unitarity Triangle (CKM 2018) Heidelberg, Germany, September 17-21, 2018, 2019*, 1903.03102.
- [36] M. Freytsis, Z. Ligeti and J. T. Ruderman, *Flavor models for $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev.* **D92** (2015) 054018 [1506.08896].
- [37] A. Celis, M. Jung, X.-Q. Li and A. Pich, *Scalar contributions to $b \rightarrow c(u)\tau\nu$ transitions*, *Phys. Lett.* **B771** (2017) 168 [1612.07757].
- [38] BABAR collaboration, *Evidence for an excess of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ decays*, *Phys. Rev. Lett.* **109** (2012) 101802 [1205.5442].
- [39] BABAR collaboration, *Measurement of an Excess of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ Decays and Implications for Charged Higgs Bosons*, *Phys. Rev.* **D88** (2013) 072012 [1303.0571].
- [40] BELLE collaboration, *Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}_\ell$ decays with hadronic tagging at Belle*, *Phys. Rev.* **D92** (2015) 072014 [1507.03233].
- [41] BELLE collaboration, *Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^*\tau^-\bar{\nu}_\tau$* , *Phys. Rev. Lett.* **118** (2017) 211801 [1612.00529].
- [42] BELLE collaboration, *Measurement of $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$ with a semileptonic tagging method*, 1904.08794.
- [43] LHCb collaboration, *Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau)/\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\mu^-\bar{\nu}_\mu)$* , *Phys. Rev. Lett.* **115** (2015) 111803 [1506.08614].
- [44] LHCb collaboration, *Measurement of the ratio of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ and $B^0 \rightarrow D^{*-}\mu^+\nu_\mu$ branching fractions using three-prong τ -lepton decays*, *Phys. Rev. Lett.* **120** (2018) 171802 [1708.08856].
- [45] LHCb collaboration, *Test of Lepton Flavor Universality by the measurement of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ branching fraction using three-prong τ decays*, *Phys. Rev.* **D97** (2018) 072013 [1711.02505].
- [46] R. Alonso, B. Grinstein and J. Martin Camalich, *Lifetime of B_c^- Constrains Explanations for Anomalies in $B \rightarrow D^{(*)}\tau\nu$* , *Phys. Rev. Lett.* **118** (2017) 081802 [1611.06676].

- [47] A. G. Akeroyd and C.-H. Chen, *Constraint on the branching ratio of $B_c \rightarrow \tau \bar{\nu}$ from LEP1 and consequences for $R(D^{(*)})$ anomaly*, *Phys. Rev.* **D96** (2017) 075011 [1708.04072].
- [48] M. Jung and D. M. Straub, *Constraining new physics in $b \rightarrow c \ell \nu$ transitions*, *JHEP* **01** (2019) 009 [1801.01112].
- [49] O. Catà and M. Jung, *Signatures of a nonstandard Higgs boson from flavor physics*, *Phys. Rev.* **D92** (2015) 055018 [1505.05804].
- [50] LHCb collaboration, *Differential branching fractions and isospin asymmetries of $B \rightarrow K^{(*)} \mu^+ \mu^-$ decays*, *JHEP* **06** (2014) 133 [1403.8044].
- [51] LHCb collaboration, *Differential branching fraction and angular analysis of the $B^+ \rightarrow K^+ \mu^+ \mu^-$ decay*, *JHEP* **02** (2013) 105 [1209.4284].
- [52] LHCb collaboration, *Measurements of the S-wave fraction in $B^0 \rightarrow K^+ \pi^- \mu^+ \mu^-$ decays and the $B^0 \rightarrow K^*(892)^0 \mu^+ \mu^-$ differential branching fraction*, *JHEP* **11** (2016) 047 [1606.04731].
- [53] LHCb collaboration, *Angular analysis of the $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ decay using 3 fb^{-1} of integrated luminosity*, *JHEP* **02** (2016) 104 [1512.04442].
- [54] LHCb collaboration, *Differential branching fraction and angular analysis of the decay $B^0 \rightarrow K^{*0} \mu^+ \mu^-$* , *JHEP* **08** (2013) 131 [1304.6325].
- [55] LHCb collaboration, *Measurement of Form-Factor-Independent Observables in the Decay $B^0 \rightarrow K^{*0} \mu^+ \mu^-$* , *Phys. Rev. Lett.* **111** (2013) 191801 [1308.1707].
- [56] LHCb collaboration, *Angular analysis and differential branching fraction of the decay $B_s^0 \rightarrow \phi \mu^+ \mu^-$* , *JHEP* **09** (2015) 179 [1506.08777].
- [57] LHCb collaboration, *Differential branching fraction and angular analysis of the decay $B_s^0 \rightarrow \phi \mu^+ \mu^-$* , *JHEP* **07** (2013) 084 [1305.2168].
- [58] LHCb collaboration, *Differential branching fraction and angular analysis of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ decays*, *JHEP* **06** (2015) 115 [1503.07138].
- [59] LHCb collaboration, *Measurement of the differential branching fraction of the decay $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$* , *Phys. Lett.* **B725** (2013) 25 [1306.2577].
- [60] ATLAS collaboration, *Angular analysis of $B_d^0 \rightarrow K^* \mu^+ \mu^-$ decays in pp collisions at $\sqrt{s} = 8 \text{ TeV}$ with the ATLAS detector*, *JHEP* **10** (2018) 047 [1805.04000].
- [61] BABAR collaboration, *Measurement of angular asymmetries in the decays $B \rightarrow K^* \ell^+ \ell^-$* , *Phys. Rev.* **D93** (2016) 052015 [1508.07960].
- [62] BELLE collaboration, *Lepton-Flavor-Dependent Angular Analysis of $B \rightarrow K^* \ell^+ \ell^-$* , *Phys. Rev. Lett.* **118** (2017) 111801 [1612.05014].
- [63] BELLE collaboration, *Measurement of the Differential Branching Fraction and Forward-Backward Asymmetry for $B \rightarrow K^{(*)} \ell^+ \ell^-$* , *Phys. Rev. Lett.* **103** (2009) 171801 [0904.0770].
- [64] BELLE collaboration, *Angular analysis of $B^0 \rightarrow K^*(892)^0 \ell^+ \ell^-$* , in *Proceedings, LHCSki 2016 - A First Discussion of 13 TeV Results: Obergurgl, Austria, April 10-15, 2016*, 2016, 1604.04042.
- [65] CDF collaboration, *Measurements of the Angular Distributions in the Decays $B \rightarrow K^{(*)} \mu^+ \mu^-$ at CDF*, *Phys. Rev. Lett.* **108** (2012) 081807 [1108.0695].
- [66] CMS collaboration, *Measurement of angular parameters from the decay $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ in proton-proton collisions at $\sqrt{s} = 8 \text{ TeV}$* , *Phys. Lett.* **B781** (2018) 517 [1710.02846].

- [67] CMS collaboration, *Angular analysis of the decay $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ from pp collisions at $\sqrt{s} = 8$ TeV*, *Phys. Lett.* **B753** (2016) 424 [1507.08126].
- [68] S. Descotes-Genon, J. Matias, M. Ramon and J. Virto, *Implications from clean observables for the binned analysis of $B \rightarrow K^* \mu^+ \mu^-$ at large recoil*, *JHEP* **01** (2013) 048 [1207.2753].
- [69] S. Descotes-Genon, L. Hofer, J. Matias and J. Virto, *On the impact of power corrections in the prediction of $B \rightarrow K^* \mu^+ \mu^-$ observables*, *JHEP* **12** (2014) 125 [1407.8526].
- [70] W. Altmannshofer and D. M. Straub, *New physics in $b \rightarrow s$ transitions after LHC run 1*, *Eur. Phys. J.* **C75** (2015) 382 [1411.3161].
- [71] A. Bharucha, D. M. Straub and R. Zwicky, *$B \rightarrow V \ell^+ \ell^-$ in the Standard Model from light-cone sum rules*, *JHEP* **08** (2016) 098 [1503.05534].
- [72] LHCb collaboration, *Search for new physics in $b \rightarrow q \ell \ell$ decays*, in *53rd Rencontres de Moriond on Electroweak Interactions and Unified Theories (Moriond EW 2018) La Thuile, Italy, March 10-17, 2018*, 2018, 1805.05073.
- [73] LHCb collaboration, *Test of lepton universality with $B^0 \rightarrow K^{*0} \ell^+ \ell^-$ decays*, *JHEP* **08** (2017) 055 [1705.05802].
- [74] LHCb collaboration, *Search for lepton-universality violation in $B^+ \rightarrow K^+ \ell^+ \ell^-$ decays*, *Phys. Rev. Lett.* **122** (2019) 191801 [1903.09252].
- [75] BELLE collaboration, *Test of lepton flavor universality in $B \rightarrow K^* \ell^+ \ell^-$ decays at Belle*, 1904.02440.
- [76] BELLE collaboration, *Test of lepton flavor universality in $B \rightarrow K \ell^+ \ell^-$ decays*, 1908.01848.
- [77] J. Aebischer, W. Altmannshofer, D. Guadagnoli, M. Reboud, P. Stangl and D. M. Straub, *B-decay discrepancies after Moriond 2019*, 1903.10434.
- [78] M. Algueró, B. Capdevila, A. Crivellin, S. Descotes-Genon, P. Masjuan, J. Matias et al., *Emerging patterns of New Physics with and without Lepton Flavour Universal contributions*, *Eur. Phys. J.* **C79** (2019) 714 [1903.09578].
- [79] M. Ciuchini, A. M. Coutinho, M. Fedele, E. Franco, A. Paul, L. Silvestrini et al., *New Physics in $b \rightarrow s \ell^+ \ell^-$ confronts new data on Lepton Universality*, *Eur. Phys. J.* **C79** (2019) 719 [1903.09632].
- [80] A. Datta, J. Kumar and D. London, *The B Anomalies and New Physics in $b \rightarrow s e^+ e^-$* , *Phys. Lett.* **B797** (2019) 134858 [1903.10086].
- [81] K. Kowalska, D. Kumar and E. M. Sessolo, *Implications for new physics in $b \rightarrow s \mu \mu$ transitions after recent measurements by Belle and LHCb*, *Eur. Phys. J.* **C79** (2019) 840 [1903.10932].
- [82] A. Arbey, T. Hurth, F. Mahmoudi, D. M. Santos and S. Neshatpour, *Update on the $b \rightarrow s \mu \mu$ anomalies*, *Phys. Rev.* **D100** (2019) 015045 [1904.08399].
- [83] A. K. Alok, A. Dighe, S. Gangal and D. Kumar, *Continuing search for new physics in $b \rightarrow s \mu \mu$ decays: two operators at a time*, *JHEP* **06** (2019) 089 [1903.09617].
- [84] A. Angelescu, D. Bečirević, D. A. Faroughy and O. Sumensari, *Closing the window on single leptoquark solutions to the B-physics anomalies*, *JHEP* **10** (2018) 183 [1808.08179].
- [85] B. Capdevila, A. Crivellin, S. Descotes-Genon, L. Hofer and J. Matias, *Searching for New Physics with $b \rightarrow s \tau^+ \tau^-$ processes*, *Phys. Rev. Lett.* **120** (2018) 181802 [1712.01919].
- [86] S. Bifani, S. Descotes-Genon, A. Romero Vidal and M.-H. Schune, *Review of Lepton Universality tests in B decays*, *J. Phys.* **G46** (2019) 023001 [1809.06229].