

Thouless time analysis of Anderson and many-body localization transitions

Piotr Sierant,¹ Dominique Delande,² and Jakub Zakrzewski^{1,3}

¹*Instytut Fizyki im. Mariana Smoluchowskiego, Uniwersytet Jagielloński, Łojasiewicza 11, 30-348 Kraków, Poland **

²*Laboratoire Kastler Brossel, Sorbonne Université, CNRS,*

ENS-PSL Research University, Collège de France, 4 Place Jussieu, 75005 Paris, France

³*Mark Kac Complex Systems Research Center, Uniwersytet Jagielloński, Kraków, Poland.*

(Dated: January 23, 2022)

Spectral statistics of disordered systems encode Thouless and Heisenberg time scales whose ratio determines whether the system is chaotic or localized. We show that the scaling of the Thouless time with system size and disorder strength is very similar in one-body Anderson models and in disordered quantum many-body systems. We argue that the two-parameter scaling breaks down in the vicinity of the transition to the localized phase signaling slow down of dynamics.

Introduction. The phenomenon of many-body localization (MBL) [1, 2], the robust mechanism of ergodicity breaking in quantum world [3–5] has received a lot of attention over the last decade. Investigations of MBL in lattice models, pioneered in spin systems [6–8], were extended to bosonic models [9, 10] and to systems of spinful fermions [11–14]. Remarkably, MBL, usually thought of as Anderson localization [15] in presence of interactions, was shown to occur in systems with completely delocalized single particle states either due to random interactions [16–18] or in a quasiperiodic Fibonacci chain [19]. MBL was also found in disorder-free systems as a result of gauge invariance [20, 21] or due to Wannier-Stark localization [22, 23], in systems with power-law interactions [24–27], or even with an infinite range [28] as well as driven Floquet MBL systems [29]. Local integrals of motion [30–36] provide a common framework to understand features of MBL such as area-law entanglement entropy of eigenstates [37, 38], logarithmic growth of bipartite entanglement entropy after quench from a separable state [39, 40] or Poisson statistics (PS) of energy levels.

The crossover between level statistics of an ergodic system with time reversal symmetry which follow predictions of the Gaussian Orthogonal Ensemble (GOE) of random matrices [41, 42] and PS of MBL phase seems to be well understood [43–48]. However, a recent analysis [49] of the Spectral Form Factor (SFF), $K(\tau)$, in the wide regime of slow thermalization on the ergodic side of the crossover [50–53] questions the very existence of the MBL phase in the thermodynamic limit predicting a two-parameter scaling of Thouless time

$$t_{Th} = t_0 e^{W/\Omega} L^2, \quad (1)$$

where L is system size, W is disorder strength, t_0 and Ω are constants. The Thouless time t_{Th} is defined as the time scale beyond which the SFF follows the universal GOE form. Another important time scale, the Heisenberg time $t_H = 2\pi/\Delta$ is defined by the average level spacing Δ which scales exponentially with a many-body system size L , $\Delta \propto e^{cL}$. The Heisenberg time t_H is a limit beyond which the discrete nature of the energy spectrum manifests itself and where system dependent

quantum effects are unavoidable. In the thermodynamic limit, (1) implies $t_{Th}/t_H \rightarrow 0$. Hence, [49] arrives at the surprising conclusion that disordered quantum spin chains have spectral properties following the GOE predictions regardless of the disorder strength W and that MBL is merely a finite-size effect.

In this letter we analyse the SFF in the delocalized phase and its modifications when approaching the transition to the localized phase. We show that the Thouless time scales like L^2 , in agreement with (1), in the deep delocalized phase in Anderson models as well as in disordered many-body systems. The scaling with L evolves to a larger power at the critical point of Anderson model, a phenomenon that we correlate with the diffusive and subdiffusive transport properties respectively in the delocalized phase and at the metal-insulator transition. Results obtained for 3D and 5D Anderson models with known localization properties put the conclusions of [49] about the scaling of Thouless time t_{Th} in a considerable doubt, suggesting the presence of a MBL phase at sufficiently strong disorder when finite-size effects are properly taken into account.

Thouless time. In a non-interacting system, the Thouless time was introduced as the time to diffuse through the system and reach its boundary [54]. It determines the energy scale below which the level statistics are well described by GOE [55], whereas its ratio with the Heisenberg time fixes the dimensionless conductance of the system [56] and enters the scaling theory of Anderson localization transition [57]. The Thouless time t_{Th} in disordered many-body systems can be probed by examining the behavior of the SFF [58–61] defined as

$$K(\tau) = \frac{1}{Z} \left\langle \left| \sum_{j=1}^{\mathcal{N}} g(\epsilon_j) e^{-i\epsilon_j \tau} \right|^2 \right\rangle, \quad (2)$$

where ϵ_j are eigenvalues of the system after the unfolding [62] (which sets their density to unity), $g(\epsilon)$ is a Gaussian function reducing influence of the spectrum's edges, the average is taken over disorder realizations and \mathcal{N} is the dimension of the Hilbert space. For a GOE matrix, the

SFF is known analytically: $K_{GOE}(\tau) = 2\tau - \tau \log(1+2\tau)$ for $\tau \leq 1$ and $K_{GOE}(\tau) = 2\tau - \tau \log(1+2\tau)$ for $\tau > 1$. The linear ramp $K_{GOE}(\tau) \approx 2\tau$ of SFF starting at $\tau = 0$ reflects correlations between all pairs of eigenvalues in a GOE matrix. In contrast, SFF $K(\tau)$ calculated for a physical system follows the GOE predictions $K(\tau) = K_{GOE}(\tau)$ only for $\tau > \tau_{Th}$ defining τ_{Th} , which, in turn, is proportional to the Thouless time $t_{Th} = \tau_{Th} t_H$. The proportionality factor t_H comes from the fact that unfolded eigenvalues ϵ_i enter the definition of $K(\tau)$; it is equal to the Heisenberg time t_H , determined by the inverse level spacing.

For a diffusive transport, the mean square displacement $\langle r^2(t) \rangle$ is proportional to time t . Hence, the above definition of t_{Th} coincides with the original definition of Thouless time in diffusive system provided that the $t_{Th} \sim L^2$ where L is the system size. For subdiffusion, the mean square displacement behaves as $\langle r^2(t) \rangle \sim t^\alpha$ with $0 < \alpha < 1$, thus we expect $t_{Th} \sim L^{2/\alpha}$. In the deeply localized regime where the localization length is much smaller than the system size, a particle never explores the full system size, so that the original Thouless time eventually diverges and becomes larger than the Heisenberg time. In contrast, Poisson statistics are characteristic for the localized regime where the SFF is independent of time; the Thouless time deduced from the SFF is thus equal to the Heisenberg time. This implies that the latter definition is applicable only in the delocalized regime. Before we consider interacting models, we examine first the Thouless time as defined by the SFF in Anderson models.

Thouless time in 3D and 5D Anderson models. The Hamiltonian of the Anderson model describes hopping of a particle on a D -dimensional lattice with disorder and reads

$$\hat{H} = -t \sum_{\langle i,j \rangle} (\hat{c}_i^\dagger \hat{c}_j + h.c.) + \sum_i \epsilon_i \hat{c}_i^\dagger \hat{c}_i, \quad (3)$$

where \hat{c}_i^\dagger is creation operator for particle at site i , $\langle ., . \rangle$ denotes sum over neighboring lattice sites, $t \equiv 1$ is the tunneling amplitude and $\epsilon_i \in [-W, W]$ are uniformly distributed uncorrelated random variables forming on-site potential. Numerical studies of transport properties of the 3D Anderson model [63–66] indicate that transport is diffusive for disorder strengths $W < W_C^{3D} \approx 16.54$ [67] and that the system remains insulating for $W > W_C$. Exactly at the transition, the 3D Anderson model is characterized by subdiffusion [68] and multifractal wave functions [69, 70]. Studies of transport in 5D Anderson model [71] find a localization transition, consistently with studies of level statistics [72] giving the critical disorder $W_C^{5D} = 57.3$, confirmed in [73, 74].

Level spacing distribution in the 3D Anderson model were studied in [55, 75–78]. Thouless times presented in Fig. 1 unveil a long-range correlation aspect of level




FIG. 1. Thouless time t_{Th} vs. disorder strength W extracted from the SFF for 3D (upper plot) and 5D (lower plot) Anderson models, for various system sizes L . The black solid lines denote the scaling of Eq. (1), grey vertical lines denote the critical disorder strength $W_C^{3D} = 16.54$ ($W_C^{5D} = 57.3$) in 3D (5D) model. Dashed lines denote the Heisenberg time t_H . The insets show t_{Th}/L^3 (t_{Th}/L^5) in the 3D (5D) case.

statistics in Anderson models. Examples of SFF and details on Thouless time estimation are given in [79].

At small disorder strength W , the Thouless times depend quadratically on system size L (Fig. 1), following precisely the scaling (1) which simply means that the dynamics is diffusive. For the 3D model, the $t_{Th}/L^2 \propto e^{W/O}$ behavior persists up to $W \approx 12$. For bigger disorder strength, the quadratic scaling with the system size is no longer valid. Directly at the transition, $W = W_C^{3D}$, the Thouless time should scale as the Heisenberg time i.e. $t_{Th} \propto L^3$. This is indeed the case as the inset in the upper plot in Fig. 1 demonstrates. Further increase of the disorder strength leads to a slow increase of the Thouless time t_{Th} with eventual saturation to the Heisenberg time t_H .

In the deep delocalized phase where t_{Th} scales with L^2 , the ratio t_{Th}/L^2 is nothing but - up to a constant multiplicative factor - the inverse of the diffusion coefficient $D(W)$, in accordance with the original definition of the Thouless time. The dependence of $D(W)$ with W is not known analytically, but it is known that it decreases quickly with W , vanishing at the critical point and scaling like $(W_c - W)^s$ below it, with the critical exponent $s \approx 1.574$. In any case, it is definitely not $e^{-W/O}$ as in (1). It may be that, in a limited range of W values, $D(W)$ can be approximately fitted by an exponential decrease, but other forms could do the job as well.

The 5D case is essentially identical, except that the Thouless time scales like L^5 instead of L^3 at the critical point. The growth of the Hilbert space size as L^5




FIG. 2. Time dependent $\alpha(t)$ function for 3D (left) and 5D (right) Anderson models for various disorder strengths W . In the 3D case, results for the system size $L = 80, 120, 160, 240$ are denoted by progressively thicker lines, whereas in the 5D case thin (thick) lines correspond to $L = 20$ ($L = 30$).

prevents reaching system sizes $L \geq 10$. Nevertheless, the obtained Thouless times t_{Th} , when rescaled by L^5 as suggested by the relation $t_{Th} \sim t_H$ valid at the transition, lead to a clear crossing of the t_{Th}/L^5 curves at W_C^{5D} . The W dependence of t_{Th}/L^2 in the deeply delocalized regime is again approximately reproduced by an exponential, although it certainly fails near the critical point.

Diffusion and subdiffusion in Anderson models. To demonstrate that the obtained behaviors of the Thouless time t_{Th} are related to time dynamics in Anderson systems, we consider the initial state $|\psi_0\rangle$ with a particle located at a given lattice site with periodic boundary conditions. The time evolved state $|\psi_0(t)\rangle = e^{-i\hat{H}t}|\psi_0\rangle$ is obtained employing the Chebyshev technique [80] that allows us to get results for system sizes up to $L = 240$ and $L = 30$ for the 3D and 5D cases, respectively. The mean square displacement

$$\langle r^2(t) \rangle = \langle \psi_0(t) | \sum_{i=1}^D (\hat{r}_i - \bar{r}_i)^2 | \psi_0(t) \rangle, \quad (4)$$

where r_i is i -th component of the position operator $\hat{\mathbf{r}}$ and $\bar{r}_i = \langle \psi_0(t) | \hat{r}_i | \psi_0(t) \rangle$, allows us to distinguish (considering first the $L \rightarrow \infty$ limit and then looking at times $t \gg 1$): diffusive $\langle r^2(t) \rangle \propto Dt$, subdiffusive $\langle r^2(t) \rangle \propto t^\alpha$ and localized behaviors. The latter occurs when $\langle r^2(t) \rangle$ saturates after the initial expansion of the wave packet. Time dependence of the mean square displacement is reflected by the function $\alpha(t) \equiv d \log \langle r^2(t) \rangle / d \log t$. In the case of diffusion $\alpha(t) = 1$, for subdiffusion $0 < \alpha(t) = \alpha < 1$ and in the localized case $\alpha(t) \rightarrow 0$.

On the delocalized side of the transition in 3D and 5D models, respectively for $W < W_C^{3D}$ and $W < W_C^{5D}$, we observe (Fig. 2) that $\alpha(t)$ initially increases over time reaching larger maximal values for increasing system sizes. Assuming that this trend persists with increasing system size, taking the thermodynamic limit $L \rightarrow \infty$ we end up with diffusive behavior $\alpha(t) = 1$ for $t \gg 1$. The

decrease of $\alpha(t)$ observed at the delocalized side of the transition for a given system size L occurs when the wave packet ceases to spread as its size approaches the system size. The situation is different at the transition, where, regardless of the system size, $\alpha(t)$ approaches a constant value $\alpha_{3D} = 2/3$ in the 3D case [68, 81] or $\alpha_{5D} = 2/5$ in the 5D case. Subsequently, $\alpha(t)$ decreases when the size of wavepacket approaches the system size L . This indicates that in the thermodynamic limit $L \rightarrow \infty$, for $t \gg 1$, there is a subdiffusion $\alpha(t) \rightarrow \alpha_{3D}(\alpha_{5D})$ at the transition in 3D (5D) Anderson model. Finally, for $W > W_C^{3D}$ (W_C^{5D}), $\alpha(t)$ decreases with time being nearly independent of the system size – a sign of localization.

The observed diffusion and subdiffusion for both 3D and 5D models agree with results obtained for the Thouless time t_{Th} . In diffusive system $\langle r(t)^2 \rangle \propto Dt$ which means that the time for reaching the boundary of the system $t_{Th}^B \propto L^2$. For subdiffusion, $\langle r(t)^2 \rangle \propto t^\alpha$ implies that $t_{Th}^B \propto L^{2/\alpha}$. Given the values for α_{3D} and α_{5D} we see that the obtained scalings of t_{Th}^B on the delocalized side of the transition and at the transition agree with the scalings $t_{Th} \propto L^2$ and $t_{Th} \propto L^3$ (or $t_{Th} \propto L^5$ in the 5D case) obtained from the SFF.

The results shown in Fig. 2 highlight the importance of finite size and finite time effects. The limit $L \rightarrow \infty$ followed by $t \rightarrow \infty$ has to be carefully examined to reveal the trend towards diffusion/subdiffusion in the system. For instance, if data for 3D model at $W = 15$ were available only up to time $t = 10^2$ one could incorrectly assume a subdiffusion with $\alpha \approx 0.75$. It seems plausible that the case of interacting systems is analogous suggesting that the claims about subdiffusion on the ergodic side of MBL transition [50, 51, 82, 83] might be invalid in the asymptotic limit $L \rightarrow \infty$, $t \gg 1$ [84, 85].

Thouless time in disordered many-body systems. Consider 1D disordered spin-1/2 chains with Hamiltonian:

$$H = J_1 \sum_{i=1}^L (S_i^x S_{i+1}^x + S_i^y S_{i+1}^y + \Delta S_i^z S_{i+1}^z) + \sum_{i=1}^L h_i S_i^z + J_2 \sum_{i=1}^L (S_i^x S_{i+2}^x + S_i^y S_{i+2}^y + \Delta S_i^z S_{i+2}^z), \quad (5)$$

where \vec{S}_i are spin-1/2 matrices, $J_1 = 1$ is the energy unit, periodic boundary conditions are assumed and $h_i \in [-W, W]$ are independent, uniformly distributed random variables. Setting $J_2 = 0$ and $\Delta = 1$ we arrive at a disordered XXZ model, widely studied in the MBL context [86–92], in particular, an analysis of mean gap ratio \bar{r} [93] predicts the critical value of disorder strength $W_C = 3.72(6)$ [94] for transition to a MBL phase. Similar reasoning leads to $W_C \approx 9$ for $J_1 - J_2$ model. For details of our calculations of Thouless times see [79].

In the case of $J_1 - J_2$ model, Thouless times obtained for available system sizes seem to follow the scaling (1) as for increasing system size L , the point $\tilde{W}(L)$ where




FIG. 3. Thouless time t_{Th} for $J_1 - J_2$ model (left) and XXZ model (right) extracted from the SFF. t_{Th} is divided by L^2 to emphasize the scaling with system size L . Dashed lines show the Heisenberg time t_H .

$t_{Th}/L^2(W)$ deviates from the $e^{W/\Omega}$ behavior shifts to larger disorder strength, as shown in the left panel of Fig. 3. An interpretation of this behavior along the lines of [49] is that one assumes that system size dependence of $\tilde{W}(L)$ continues indefinitely, so that the scaling (1) holds in the thermodynamic limit. This would imply that there is no transition to a MBL phase. However, the Thouless time scaling obtained for available system sizes in the 5D Anderson model, exhibited in the lower panel of Fig. 1, is very similar with larger system sizes deviating from (1) at larger disorder strength. As such a behavior occurs in the 5D Anderson model despite the localization transition taking place at W_C^{5D} , we may give a second possible interpretation of the result: the scaling (1) is not broken at available system sizes because of strong finite size effects. While it is still possible to devise the location of the critical point W_C^{5D} provided one knows the correct value of the exponent α governing the subdiffusion at the Anderson transition, it is not clear how to rescale the Thouless times t_{Th} in the many-body case since the transport properties on the delocalized side are not fully understood, with a suggestion of subdiffusive behavior with exponent α vanishing close to the transition [95]. Presumably, a sensible criterion for the transition in the many-body case would be $t_{Th} \propto t_H \propto e^{cL}$. In any case, the main observation in [49] is that t_{Th}/L^2 is approximately equal to $e^{W/\Omega}$ in the deeply delocalized regime of the $J_1 - J_2$ model. This implies, in turn, that the diffusion coefficient $D(W)$ decreases like $e^{-W/\Omega}$, exactly like in the 3D and 5D Anderson models. Concluding that $D(W)$ never vanishes is a dangerous extrapolation, which leads to incorrect results for the Anderson models. The similarity of Thouless time scaling for 5D Anderson and $J_1 - J_2$ models suggests that the conclusion of [49] about $D(W) \propto e^{-W/\Omega}$ in the $J_1 - J_2$ model for any disorder strength in the thermodynamic limit is misleading. Our results show that the apparent scaling (1) is probably a finite size effect.

The finite size effects in the $J_1 - J_2$ model are neces-

sarily enhanced by the next-to-nearest neighbor coupling term, thus we may expect weaker finite size effects for the XXZ model. The scaling of Thouless time for this model is presented in the right panel of Fig. 3 and it follows (1) only for disorder strengths $W \in [1, 2]$. We observe two important differences with the results for $J_1 - J_2$ model. Firstly, at weak disorder W , the exponential dependence of the Thouless time t_{Th} on W is weaker than in the interval $W \in [1, 2]$. This is due to the proximity of the integrable point $W = 0$ [96, 97] with Poisson level statistics and $t_{Th} = t_H$. Secondly and more importantly, we see a breakdown of (1) for the XXZ model at $W \gtrsim 2$ where the data for $L = 22$ and $L = 24$ exceed the $t_0 e^{W/\Omega}$ line even though the Thouless time is still an order of magnitude smaller than the Heisenberg time t_H . This indicates that the exponential scaling with W is a numerical observation explicitly broken in the XXZ model and likely valid only in a limited range in other systems. The data for $L = 22$ and $L = 24$ are available only for $W \geq 2$ and $W > 2.2$ [79]. Nevertheless, the breakdown of the scaling (1) for $L = 22, 24$ at $W \approx 2.2$ is apparent, indicating that the L^2 scaling of Thouless time breaks down. This reflects the slow-down of transport and approaching the MBL transition when $t_{Th} \propto t_H \propto e^{cL}$.

Conclusions. Our results show that the Thouless time, defined by the behavior of the SFF, reflects the transport properties in disordered non-interacting models as we have shown on the examples of 3D and 5D Anderson models. In particular, the scaling of the Thouless time t_{Th} at the transition encodes the subdiffusive behavior of the mean square displacement $\langle r^2(t) \rangle \sim t^\alpha$ with the exponent $\alpha_{3D} = 2/3$ and $\alpha_{5D} = 2/5$ leading to scaling $t_{Th} \sim L^{2/\alpha}$ with system size at the transition.

The scaling of Thouless time for $J_1 - J_2$ model seems to follow $t_{Th} \sim t_0 L^2 e^{W/\Omega}$, however, the behavior of t_{Th} is directly analogous to the case of 5D Anderson model. The latter undergoes a transition to a localized phase and the Thouless time does not exceed the $t_0 L^2 e^{W/\Omega}$ curve only because of strong finite size effects at available system sizes. It is plausible that the situation is the same in the $J_1 - J_2$ model, raising doubts about the claims of [49]. Our results for XXZ model demonstrate that the L^2 scaling of the Thouless time t_{Th} , valid deep in delocalized phase is evidently broken at $W \approx 2.2$, signaling a transition to a MBL phase at a strong disorder.

Finally, let us mention alternative definitions of Thouless time [98–104]. Comparison of these different approaches is in progress. While finalizing this manuscript, we became aware of the related works [105, 106].

Acknowledgments. We are most grateful to Fabien Alet for kindly sharing with us the eigenvalues for $L = 22, 24$ XXZ model as well as discussions on subjects related to this work. The computations have been performed within PL-Grid Infrastructure, its support is acknowledged. We acknowledge the support of National Science Centre (Poland)

under projects 2015/19/B/ST2/01028 (P.S. and J.Z.), 2018/28/T/ST2/00401 (doctoral scholarship – P.S.) as well as Polish-French bilateral grant Polonium 40490ZE.

* piotr.sierant@uj.edu.pl

- [1] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. Lett.* **95**, 206603 (2005).
- [2] D. Basko, I. Aleiner, and B. Altshuler, *Ann. Phys. (NY)* **321**, 1126 (2006).
- [3] R. Nandkishore and D. A. Huse, *Ann. Rev. Cond. Mat. Phys.* **6**, 15 (2015).
- [4] F. Alet and N. Laflorencie, *Comptes Rendus Physique* **19**, 498 (2018).
- [5] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, *Rev. Mod. Phys.* **91**, 021001 (2019).
- [6] L. F. Santos, G. Rigolin, and C. O. Escobar, *Phys. Rev. A* **69**, 042304 (2004).
- [7] V. Oganesyan and D. A. Huse, *Phys. Rev. B* **75**, 155111 (2007).
- [8] A. Pal and D. A. Huse, *Phys. Rev. B* **82**, 174411 (2010).
- [9] P. Sierant and J. Zakrzewski, *New Journal of Physics* **20**, 043032 (2018).
- [10] T. Orell, A. A. Michailidis, M. Serbyn, and M. Silveri, *Phys. Rev. B* **100**, 134504 (2019).
- [11] R. Mondaini and M. Rigol, *Phys. Rev. A* **92**, 041601 (2015).
- [12] P. Prelovšek, O. S. Barišić, and M. Žnidarič, *Phys. Rev. B* **94**, 241104 (2016).
- [13] J. Zakrzewski and D. Delande, *Phys. Rev. B* **98**, 014203 (2018).
- [14] M. Kozarzewski, P. Prelovšek, and M. Mierzejewski, *Phys. Rev. Lett.* **120**, 246602 (2018).
- [15] P. W. Anderson, *Phys. Rev.* **109**, 1492 (1958).
- [16] P. Sierant, D. Delande, and J. Zakrzewski, *Phys. Rev. A* **95**, 021601 (2017).
- [17] Y. Bar Lev, D. R. Reichman, and Y. Sagi, *Phys. Rev. B* **94**, 201116 (2016).
- [18] X. Li, D.-L. Deng, Y.-L. Wu, and S. Das Sarma, *Phys. Rev. B* **95**, 020201 (2017).
- [19] N. Macé, N. Laflorencie, and F. Alet, *SciPost Phys.* **6**, 50 (2019).
- [20] A. Smith, J. Knolle, R. Moessner, and D. L. Kovrizhin, *Phys. Rev. Lett.* **119**, 176601 (2017).
- [21] M. Brenes, M. Dalmonte, M. Heyl, and A. Scardicchio, *Phys. Rev. Lett.* **120**, 030601 (2018).
- [22] M. Schulz, C. A. Hooley, R. Moessner, and F. Pollmann, *Phys. Rev. Lett.* **122**, 040606 (2019).
- [23] E. van Nieuwenburg, Y. Baum, and G. Refael, *Proceedings of the National Academy of Sciences* **116**, 9269 (2019).
- [24] G. De Tomasi, *Phys. Rev. B* **99**, 054204 (2019).
- [25] A. Safavi-Naini, M. L. Wall, O. L. Acevedo, A. M. Rey, and R. M. Nandkishore, *Phys. Rev. A* **99**, 033610 (2019).
- [26] T. Botzung, D. Vodola, P. Naldesi, M. Müller, E. Ercolessi, and G. Pupillo, *Phys. Rev. B* **100**, 155136 (2019).
- [27] A. O. Maksymov and A. L. Burin, *Phys. Rev. B* **101**, 024201 (2020).
- [28] P. Sierant, K. Biedroń, G. Morigi, and J. Zakrzewski, *SciPost Phys.* **7**, 8 (2019).
- [29] P. Bordia, H. Lüschen, U. Schneider, M. Knap, and I. Bloch, *Nature Physics* **13**, 460 EP (2017), article.
- [30] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **111**, 127201 (2013).
- [31] D. A. Huse, R. Nandkishore, and V. Oganesyan, *Phys. Rev. B* **90**, 174202 (2014).
- [32] V. Ros, M. Mueller, and A. Scardicchio, *Nuclear Physics B* **891**, 420 (2015).
- [33] J. Z. Imbrie, *Phys. Rev. Lett.* **117**, 027201 (2016).
- [34] T. B. Wahl, A. Pal, and S. H. Simon, *Phys. Rev. X* **7**, 021018 (2017).
- [35] M. Mierzejewski, M. Kozarzewski, and P. Prelovšek, *Phys. Rev. B* **97**, 064204 (2018).
- [36] S. J. Thomson and M. Schiró, *Phys. Rev. B* **97**, 060201 (2018).
- [37] B. Bauer and C. Nayak, *Journal of Statistical Mechanics: Theory and Experiment* **2013**, P09005 (2013).
- [38] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **110**, 260601 (2013).
- [39] M. Žnidarič, T. Prosen, and P. Prelovšek, *Phys. Rev. B* **77**, 064426 (2008).
- [40] J. H. Bardarson, F. Pollmann, and J. E. Moore, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [41] M. L. Mehta, *Random Matrices* (Elsevier, 1990).
- [42] F. Haake, *Quantum Signatures of Chaos* (Springer, Berlin, 2010).
- [43] M. Serbyn and J. E. Moore, *Phys. Rev. B* **93**, 041424 (2016).
- [44] C. L. Bertrand and A. M. García-García, *Phys. Rev. B* **94**, 144201 (2016).
- [45] J. A. Kjäll, *Phys. Rev. B* **97**, 035163 (2018).
- [46] W. Buijsman, V. Cheianov, and V. Gritsev, *Phys. Rev. Lett.* **122**, 180601 (2019).
- [47] P. Sierant and J. Zakrzewski, *Phys. Rev. B* **99**, 104205 (2019).
- [48] P. Sierant and J. Zakrzewski, *Phys. Rev. B* **101**, 104201 (2020).
- [49] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, (2019), [arXiv:1905.06345](https://arxiv.org/abs/1905.06345).
- [50] Y. Bar Lev, G. Cohen, and D. R. Reichman, *Phys. Rev. Lett.* **114**, 100601 (2015).
- [51] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **93**, 060201 (2016).
- [52] D. J. Luitz and Y. Bar Lev, *Phys. Rev. Lett.* **117**, 170404 (2016).
- [53] M. Mierzejewski, J. Herbrych, and P. Prelovšek, *Phys. Rev. B* **94**, 224207 (2016).
- [54] D. Thouless, *Physics Reports* **13**, 93 (1974).
- [55] B. I. Shklovskii, B. Shapiro, B. R. Sears, P. Lambrianides, and H. B. Shore, *Phys. Rev. B* **47**, 11487 (1993).
- [56] J. T. Edwards and D. J. Thouless, *Journal of Physics C: Solid State Physics* **5**, 807 (1972).
- [57] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, *Phys. Rev. Lett.* **42**, 673 (1979).
- [58] J. Cotler, N. Hunter-Jones, J. Liu, and B. Yoshida, *Journal of High Energy Physics* **2017**, 48 (2017).
- [59] X. Chen and A. W. W. Ludwig, *Phys. Rev. B* **98**, 064309 (2018).
- [60] H. Gharibyan, M. Hanada, S. H. Shenker, and M. Tezuka, *Journal of High Energy Physics* **2018**, 124 (2018).
- [61] A. Chan, A. De Luca, and J. T. Chalker, *Phys. Rev. Lett.* **121**, 060601 (2018).
- [62] J. M. G. Gómez, R. A. Molina, A. Relaño, and J. Re-

- tamosa, *Phys. Rev. E* **66**, 036209 (2002).
- [63] A. MacKinnon and B. Kramer, *Phys. Rev. Lett.* **47**, 1546 (1981).
- [64] A. MacKinnon and B. Kramer, *Zeitschrift für Physik B Condensed Matter* **53**, 1 (1983).
- [65] B. Kramer, K. Broderix, A. Mackinnon, and M. Schreiber, *Physica A: Statistical Mechanics and its Applications* **167**, 163 (1990).
- [66] B. Kramer and A. MacKinnon, *Reports on Progress in Physics* **56**, 1469 (1993).
- [67] K. Slevin and T. Ohtsuki, *Journal of the Physical Society of Japan* **87**, 094703 (2018).
- [68] T. Ohtsuki and T. Kawarabayashi, *Journal of the Physical Society of Japan* **66**, 314 (1997).
- [69] A. Rodriguez, L. J. Vasquez, and R. A. Römer, *Phys. Rev. Lett.* **102**, 106406 (2009).
- [70] A. Rodriguez, L. J. Vasquez, K. Slevin, and R. A. Römer, *Phys. Rev. Lett.* **105**, 046403 (2010).
- [71] Y. Ueoka and K. Slevin, *Journal of the Physical Society of Japan* **83**, 084711 (2014).
- [72] A. M. García-García and E. Cuevas, *Phys. Rev. B* **75**, 174203 (2007).
- [73] E. Tarquini, G. Biroli, and M. Tarzia, *Phys. Rev. B* **95**, 094204 (2017).
- [74] F. Pietracaprina, V. Ros, and A. Scardicchio, *Phys. Rev. B* **93**, 054201 (2016).
- [75] E. Hofstetter and M. Schreiber, *Phys. Rev. B* **48**, 16979 (1993).
- [76] I. K. Zharekeshev and B. Kramer, *Phys. Rev. B* **51**, 17239 (1995).
- [77] I. Varga, E. Hofstetter, M. Schreiber, and J. Pipek, *Phys. Rev. B* **52**, 7783 (1995).
- [78] I. K. Zharekeshev and B. Kramer, *Phys. Rev. Lett.* **79**, 717 (1997).
- [79] See Supplemental Material at [URL will be inserted by publisher] details on extracting of the Thouless time from SFF.
- [80] H. Fehske and R. Schneider, *Computational many-particle physics* (Springer, Germany, 2008).
- [81] G. Lemarié, H. Lignier, D. Delande, P. Szriftgiser, and J. C. Garreau, *Phys. Rev. Lett.* **105**, 090601 (2010).
- [82] K. Agarwal, S. Gopalakrishnan, M. Knap, M. Müller, and E. Demler, *Phys. Rev. Lett.* **114**, 160401 (2015).
- [83] V. K. Varma, A. Lerose, F. Pietracaprina, J. Goold, and A. Scardicchio, *Journal of Statistical Mechanics: Theory and Experiment* **2017**, 053101 (2017).
- [84] S. Bera, G. De Tomasi, F. Weiner, and F. Evers, *Phys. Rev. Lett.* **118**, 196801 (2017).
- [85] F. Weiner, F. Evers, and S. Bera, *Phys. Rev. B* **100**, 104204 (2019).
- [86] T. C. Berkelbach and D. R. Reichman, *Phys. Rev. B* **81**, 224429 (2010).
- [87] S. Bera, H. Schomerus, F. Heidrich-Meisner, and J. H. Bardarson, *Phys. Rev. Lett.* **115**, 046603 (2015).
- [88] T. Enss, F. Andraschko, and J. Sirker, *Phys. Rev. B* **95**, 045121 (2017).
- [89] L. Herviou, S. Bera, and J. H. Bardarson, *Phys. Rev. B* **99**, 134205 (2019).
- [90] L. Colmenarez, P. A. McClarty, M. Haque, and D. J. Luitz, (2019), [arXiv:1906.10701](https://arxiv.org/abs/1906.10701).
- [91] T. Chanda, P. Sierant, and J. Zakrzewski, *Phys. Rev. B* **101**, 035148 (2020).
- [92] F. Pietracaprina, N. Macé, D. J. Luitz, and F. Alet, *SciPost Phys.* **5**, 45 (2018).
- [93] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, *Phys. Rev. Lett.* **110**, 084101 (2013).
- [94] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **91**, 081103 (2015).
- [95] D. J. Luitz and Y. B. Lev, *Annalen der Physik* **529**, 1600350 (2017).
- [96] H. Bethe, *Zeitschrift für Physik* **71**, 205 (1931).
- [97] F. C. Alcaraz, M. N. Barber, M. T. Batchelor, R. J. Baxter, and G. R. W. Quispel, *Journal of Physics A: Mathematical and General* **20**, 6397 (1987).
- [98] W. Beugeling, R. Moessner, and M. Haque, *Phys. Rev. E* **91**, 012144 (2015).
- [99] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. B* **96**, 104201 (2017).
- [100] E. J. Torres-Herrera and L. F. Santos, *Phys. Rev. B* **92**, 014208 (2015).
- [101] E. J. Torres-Herrera and L. F. Santos, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **375**, 20160434 (2017).
- [102] E. J. Torres-Herrera, A. M. García-García, and L. F. Santos, *Phys. Rev. B* **97**, 060303 (2018).
- [103] M. Schiulaz, E. J. Torres-Herrera, and L. F. Santos, *Phys. Rev. B* **99**, 174313 (2019).
- [104] M. Schiulaz, E. J. Torres-Herrera, F. Pérez-Bernal, and L. F. Santos, (2019), [arXiv:1906.11856](https://arxiv.org/abs/1906.11856).
- [105] D. A. Abanin, J. H. Bardarson, G. D. Tomasi, S. Gopalakrishnan, V. Khemani, S. A. Parameswaran, F. Pollmann, A. C. Potter, M. Serbyn, and R. Vasseur, (2019), [arXiv:1911.04501](https://arxiv.org/abs/1911.04501).
- [106] R. K. Panda, A. Scardicchio, M. Schulz, S. R. Taylor, and M. Žnidarič, *EPL (Europhysics Letters)* **128**, 67003 (2020).

SUPPLEMENTARY: EXTRACTING THOULESS TIME FROM SPECTRAL FORM FACTOR

The spectral form factor (SFF) is the tool employed in analysis of level statistics in Anderson models and of disordered quantum spin chains presented in the main text. In this supplementary material we recall definition of SFF, provide details of performed numerical calculations and describe the employed method of extraction of the Thouless time from SFF.

The spectral form factor (SFF) is defined as

$$K(\tau) = \frac{1}{Z} \left\langle \left| \sum_{j=1}^{\mathcal{N}} g(\epsilon_j) e^{-i\epsilon_j \tau} \right|^2 \right\rangle, \quad (\text{S.1})$$

where \mathcal{N} is the dimension of Hilbert space. The eigenvalues ϵ_j are obtained in the so called unfolding procedure. During the unfolding, a level staircase function $\sigma(E) = \sum_i \Theta(E - E_i)$ (obtained from the set of eigenvalues of the system $\{E_i\}$ ordered in an ascending manner) is separated into smooth and fluctuating parts $\sigma(E) = \bar{\sigma}(E) + \delta\sigma(E)$ and the eigenvalues are mapped via

$$E_j \rightarrow \epsilon_j = \bar{\sigma}(E_j). \quad (\text{S.2})$$

As the smooth part $\bar{\sigma}(E)$ we take a polynomial of certain small degree n_p fitted to the level staircase function $\sigma(E)$. To calculate SFF we use $g(\epsilon) \propto \exp(-(\epsilon - \bar{\epsilon})^2/2\eta\sigma_\epsilon^2)$, where $\bar{\epsilon}$ denotes the average of the unfolded eigenvalues for given disorder realization ϵ_i , σ_ϵ is the standard deviation of $\{\epsilon_i\}$ and $\eta = 0.3$. This choice of parameters follows precisely [49].

The summation in (S.1) extends over the whole spectrum of the system. To get all of the eigenvalues we perform exact diagonalization (ED) for Anderson models in 3D and 5D. We calculate SFF according to formula (S.1), using unfolding with polynomial of order $n_p = 10$ and averaging results over more than 400 disorder realizations. Similarly, we perform ED of disordered Heisenberg spin chain and $J_1 - J_2$ model of size $L \leq 16$ ($L = 18$) and average results over 1000 (500) disorder realizations.

An exemplary result for SFF of 3D Anderson model is shown in Fig. 4. One clearly observes a value $\tau_{Th} = t_{Th}/t_H$ beyond which SFF of 3D Anderson model follows the GOE prediction. To quantitatively extract Thouless time t_{Th} from the SFF, we follow [49] and consider a

function

$$\Delta K(t/t_H) = \left| \log \left(\frac{K(t/t_H)}{K_{GOE}(\tau = t/t_H)} \right) \right|. \quad (\text{S.3})$$

The Thouless time t_{Th} is the smallest positive time for which $\Delta K(t/t_H) < \epsilon$. We choose the value of cut-off $\epsilon = 0.05$. The choice of cut-off ϵ affects the obtained values of Thouless time t_{Th} , however, reasonable changes




FIG. 4. Spectral form factor $K(\tau)$ for 3D Anderson model of size $L = 40$, dots denote obtained values of Thouless time. The black dashed lines denote the SFF of GOE ensemble which is known analytically.

in value of ϵ do not affect the obtained scaling of Thouless time with system size and disorder strength.

The eigenvalues for larger sizes $L = 20, 22, 24$ of considered spin chains can be obtained with the shift-and-invert method. However, the shift-and-invert method provides only a certain number n_e eigenvalues around a specific target energy. Using $n_e = 2000, 100, 50$ eigenvalues from the middle of system spectrum for system size $L = 20, 22, 24$ and averaging results over more than 200 disordered realizations we have verified that the SFF can still be calculated if the available n_e eigenvalues is taken into account in the sum (S.1). In those cases we have used unfolding with polynomial of degree $n_p = 3$ to avoid over-fitting of the staircase function [62]. If a small fraction n_e/\mathcal{N} of eigenvalues is considered in the sum (S.1), the SFF is correctly reproduced only for $t > t_M$ with t_M depending on n_e as well as on the system size L . This can be intuitively understood: since t_{Th} is inversely proportional to E_{Th} – the energy scale at which the correlations of eigenvalues are well described by GOE, the upper bound on energy scales probed by SFF is proportional to the number of available eigenvalues n_e which translates to the lower bound t_M on accessible times. This shows that the Thouless times can be extracted from the fraction of eigenvalues from shift-and-invert method if the disorder is sufficiently strong.