

Location estimation for symmetric log-concave densities

Nilanjana Laha*

*Department of Biostatistics, Harvard University, 677 Huntington Ave, Boston, MA 02115, U.S.A**

e-mail: nlaha@hsph.harvard.edu

Abstract: We revisit the problem of estimating the center of symmetry θ of an unknown symmetric density f . Although [Stone \(1975\)](#), [Van Eeden \(1970\)](#), and [Sacks \(1975\)](#) constructed adaptive estimators of θ in this model, their estimators depend on tuning parameters. In an effort to circumvent the dependence on tuning parameters, we impose an additional assumption of log-concavity on f . We show that in this shape-restricted model, the maximum likelihood estimator (MLE) of θ exists. We also study some truncated one-step estimators and show that they are \sqrt{n} -consistent, and nearly achieve the asymptotic efficiency bound. We also show that the rate of convergence for the MLE is $O_p(n^{-2/5})$. Furthermore, we show that our estimators are robust with respect to the violation of the log-concavity assumption. In fact, we show that the one step estimators are still \sqrt{n} -consistent under some mild conditions. These analytical conclusions are supported by simulation studies.

Keywords and phrases: log-concave, shape constraint, symmetric location model, one step estimator.

1. Introduction

In this paper, we revisit the symmetric location model with an additional shape-restriction of log-concavity. We let \mathcal{P} denote the class of all densities on the real line \mathbb{R} . For any $\theta \in \mathbb{R}$, denote by \mathcal{S}_θ the class of all densities that are symmetric about θ . We take

$$\mathcal{C} := \left\{ \phi : \mathbb{R} \mapsto [-\infty, \infty) \mid \phi \text{ is concave, closed, and proper} \right\}, \quad (1.1)$$

and define $\mathcal{SC}_\theta = \mathcal{S}_\theta \cap \mathcal{C}$. Here a proper and closed concave function is as defined in [Rockafellar \(1970\)](#), page 24 and 50. Letting \mathcal{LC} denote the class of log-concave densities

$$\mathcal{LC} := \left\{ f \in \mathcal{P} \mid \phi = \log f \in \mathcal{C} \right\},$$

we set $\mathcal{SLC}_\theta = \mathcal{LC} \cap \mathcal{S}_\theta$. We aim to estimate θ in the log-concave symmetric location model

$$\mathcal{P}_0 = \left\{ f \in \mathcal{P} \mid f(x; \theta) = g(x - \theta), \theta \in \mathbb{R}, g \in \mathcal{SLC}_0, \mathcal{I}_f < \infty \right\}, \quad (1.2)$$

where \mathcal{I}_f is the Fisher information for location. It is well-established that (Huber, 1964, Theorem 3) \mathcal{I}_f is finite if and only if f is an absolutely continuous density satisfying

$$\int_{-\infty}^{\infty} \left(\frac{f'(x)}{f(x)} \right)^2 f(x) dx < \infty,$$

where f' is an L_1 -derivative of f . Also, in this case, \mathcal{I}_f takes the form

$$\mathcal{I}_f = \int_{-\infty}^{\infty} \left(\frac{f'(x)}{f(x)} \right)^2 f(x) dx.$$

Estimation of θ in the full symmetric location model

$$\mathcal{P}_s = \left\{ f \in \mathcal{P} \mid f(x; \theta) = g(x - \theta), \theta \in \mathbb{R}, g \in \mathcal{S}_0, \mathcal{I}_f < \infty \right\} \quad (1.3)$$

is an old semi-parametric problem, dating back to Stein (1956). From then on, this problem has been considered by many early authors including, but not limited to, Stone (1975), Beran (1974), Sacks (1975), and Van Eeden (1970). There are two main reasons behind the assumption of symmetry in this model. First, as Stone (1975) has noted, if f is totally unrestricted, θ is not identifiable. Second, the definition of location becomes less clear in the absence of symmetry (Takeuchi, 1975).

The appeal of the above model lies in the fact that adaptive estimation of θ is possible in this model (Stone, 1975). In other words, there exist consistent estimators of θ in \mathcal{P}_s , whose variances attain the parametric lower bound, which is \mathcal{I}_f^{-1} in this case. See Sections 3.2, 3.3, and 6.3 of Bickel *et al.* (1998) for more discussion on adaptive estimation in \mathcal{P}_s .

Now we will briefly discuss some existing methods of estimating θ in \mathcal{P}_s . Stone (1975) considers a one-step estimator. One-step estimators generally depend on the estimation of the score $-f'/f$. This paper estimates the scores using symmetrized Gaussian kernels based on truncated data, thereby incorporating two tuning parameters, one for the truncation, and another for the scaling of the Gaussian Kernel. Although Stone (1975) requires the tuning parameters to satisfy some conditions (Stone, 1975, Theorem 5.3), he does not provide any prescription for choosing them. Sacks (1975) considers a subclass of \mathcal{P}_s with some restrictions on the score function that are weaker than the assumption of log-concavity, and includes heavy-tailed distributions like Cauchy. His estimator is motivated by a linear function of the order statistics, whose coefficients depend on the score functions. The estimation procedure involves three tuning parameters. Sacks (1975) gives some examples of these tuning parameters, but does not provide any data-dependent method for choosing them. Beran (1974) uses linearized rank estimate for adaptive estimation of θ in \mathcal{P}_s . But he makes the additional assumption that the score is twice continuously differentiable, which excludes densities like Laplace. His method is based on the Fourier series expansion of the score function and requires a tuning parameter for choosing the degree of the associated complex trigonometric polynomial. Beran (1974) does not provide any data-dependent procedure to choose this tuning parameter.

Now we provide some justification for imposing the shape restriction of log-concavity. The class of log-concave densities, \mathcal{LC} , belongs to the larger class of unimodal densities. Unimodality is a reasonable assumption in our context, because, as [Takeuchi \(1975\)](#) points out, in practice, multimodal densities generally result from unimodal mixtures. Separate procedures are available for the latter class. The problem with the class of unimodal densities, however, is that it does not admit an MLE ([Birgé, 1997](#)). Many common symmetric unimodal continuous densities, notwithstanding, are in \mathcal{LC} , which, being a smaller and richer class, admit MLE (cf. [Pal *et al.*, 2007](#); [Dümbgen and Rufibach, 2009](#); [Doss and Wellner, 2019](#); [Xu and Samworth, 2019](#)). Since the MLE does not depend on tuning parameters, it is possible to estimate f without relying on any tuning parameter, which can lead to tuning parameter free estimation of θ . This is the main reason to opt for the shape-restriction of log-concavity.

Although shape-constrained estimation has a rich history, so far there has been very little to no use of shape-constraints in connection with semiparametric models. In fact, to the best of our knowledge, [Van Eeden \(1970\)](#) is the only one to incorporate shape-constraints in treating the one-sample symmetric location problem considered here. Actually [Van Eeden \(1970\)](#) uses log-concavity and considers estimation in \mathcal{P}_0 although her paper does not mention log-concavity. She restricts \mathcal{P}_s by assuming that $f'(F^{-1}(u))/f(F^{-1}(u))$ is non-increasing in u , which is equivalent to f being log-concave. [Van Eeden \(1970\)](#) considers data-partitioning to estimate scores from a small fraction of the data and constructs the Hodges-Lehmann rank estimate of location ([Hodges and Lehmann, 1963](#)) from the remaining data. The partitioning of the data and the construction of the score function involve some tuning parameters. [Van Eeden \(1970\)](#) does not explicitly describe how to choose these tuning parameters. We also want to remark about [Bhattacharyya and Bickel \(2013\)](#), who consider both location and scale estimation in an elliptical symmetry model, which albeit bearing some resemblance, is different from \mathcal{P}_0 . Also, [Bhattacharyya and Bickel \(2013\)](#)'s estimation procedure is completely different from ours.

To summarize, our goal is to bridge the gap between the symmetric location model and log-concavity. In doing so, we also try to exploit the structure in the class \mathcal{LC} to get rid of the dependence on external tuning parameters. Here is a brief account of our main contributions.

The MLE

We show that the MLE of θ and g exist in \mathcal{P}_0 . It also follows that $\hat{\theta}_n$, i.e. the MLE of θ , has the rate of convergence $O_p(n^{-2/5})$. In addition, we show that the Hellinger distance between $f \in \mathcal{P}_0$ and its MLE \hat{f}_n is of order $O_p(n^{-2/5})$, where the Hellinger distance $H(f, g)$ between two densities f and g is defined by

$$H^2(f, g) = 2^{-1} \int (\sqrt{f(x)} - \sqrt{g(x)})^2 dx.$$

The above-mentioned rate is analogous to the rate obtained by [Doss and Wellner \(2019\)](#) for the MLE of f in \mathcal{SLC}_0 , or by [Doss and Wellner \(2016\)](#) for the MLE

of f in \mathcal{LC} .

The One step estimator

We also propose four one step estimators for θ , which generally use on the mean, median or a trimmed mean as the preliminary estimator. We estimate the score using either the MLE of f in \mathcal{LC} , or in $\mathcal{SLLC}_{\tilde{\theta}_n}$, which can be efficiently computed using the R packages "logcondens" and "logcondens.mode", respectively. We establish that a truncated version of the one-step estimators is \sqrt{n} -consistent, where its variance also nearly achieves the asymptotic efficiency bound \mathcal{I}_f^{-1} . The truncated estimator $\tilde{\theta}_n$, however, use a tuning parameter.

Model misspecification

We show that our estimators are robust to the violation of the log-concavity assumption. In particular, we show that, even if $f \notin \mathcal{P}_0$, as long as f is symmetric about θ , the MLE and the truncated one step estimators are still strongly consistent. Additionally, for the specific case of the truncated one-step estimators, \sqrt{n} -consistency is also preserved. The MLE, on the other hand, may still possess a limit even when f is not symmetric, or if the data generating distribution does not even possess a density. For the analysis in the misspecified case, we rely on the log-concave projection theory developed by Dümbgen *et al.* (2011), Cule and Samworth (2010), and Xu and Samworth (2019). The early authors, Stone (1975), Beran (1974), Sacks (1975), and Van Eeden (1970) did not discuss the behavior of their estimators under model miss-specification although Stone (1975) did discuss the behavior of his estimator when $\mathcal{I}_{f_0} = \infty$.

The article is organized as follows. In Section 2.1, we introduce the one step estimators, and in 2.2, we discuss the existence of the MLE and its characterizations. We study the asymptotic properties of our estimators in Section 3. We provide a simulation study in Section 4. The subtleties of our estimators are deferred to the Discussion Section. We collect the proofs in Section 6.

1.1. Preliminaries/ Notation and terminology

We assume that X_1, \dots, X_n are independent and identically distributed (i.i.d.) random variables with density $f_0 \in \mathcal{P}_0$. Therefore, (1.2) indicates that $f_0 = g_0(\cdot - \theta_0)$, where $g_0 \in \mathcal{SLLC}_0$, the class of all log-concave densities symmetric about 0. We set $\phi_0 = \log f_0$, and $\psi_0 = \log g_0$. We let F_0 and G_0 denote the respective distribution functions of f_0 and g_0 , and denote by P_0 the measure corresponding to F_0 . We denote the empirical distribution function of the X_i 's by \mathbb{F}_n , and write \mathbb{P}_n for the corresponding empirical measure. The (classical) empirical process of the X_i 's will be denoted by $\mathbb{Z}_n = \sqrt{n}(\mathbb{F}_n - F_0)$. For any integrable function $h : \mathbb{R} \mapsto \mathbb{R}$, we write

$$\mathbb{P}_n h := n^{-1} \sum_{i=1}^n h(X_i) \quad \text{and} \quad P_0 h = \int_{-\infty}^{\infty} h(x) f_0(x) dx.$$

As usual, we denote the order statistics of a sample (Y_1, \dots, Y_n) by

$$Y_{()} = (Y_{(1)}, \dots, Y_{(n)}).$$

In context of a concave function $\psi : \mathbb{R} \mapsto \mathbb{R}$, the domain $\text{dom}(\psi)$ will be defined as in [Rockafellar \(1970\)](#) (page 40), that is, $\text{dom}(\psi) = \{x \in \mathbb{R} | \psi(x) > -\infty\}$. For any concave function ψ , we say $x \in \mathbb{R}$ is a knot of ψ , if either $\psi'(x+) \neq \psi'(x-)$, or x is at the boundary of $\text{dom}(\psi)$. We denote by $\mathcal{K}(\psi)$ the set of the knots of ψ . Unless otherwise mentioned, for a real valued function h , provided they exist, h' and $h'(\cdot-)$ will refer to the right and left derivatives of h respectively. For a distribution function F , we let $J(F)$ denote the set $\{0 < F < 1\}$. For any non-negative real-valued function h , we denote the support by $\text{supp}(h) = \{x \in \mathbb{R} | h(x) > 0\}$. As usual, we denote the set of natural numbers by \mathbb{N} .

Next, we develop some notation for maximum likelihood estimation. For $\psi \in \mathcal{SC}_0$ and $\theta \in \mathbb{R}$, following [Dümbgen et al. \(2011\)](#) and [Xu and Samworth \(2019\)](#), we define the criterion function for maximum likelihood estimation by

$$\Psi(\theta, \psi, F) = \int_{-\infty}^{\infty} \psi(x - \theta) dF(x) - \int_{-\infty}^{\infty} e^{\psi(x - \theta)} dx. \quad (1.4)$$

Following [Silverman \(1982\)](#), we included a Lagrange term to get rid of the normalizing constant involved in density estimation. This is a common device in log-concave density estimation literature (cf. [Dümbgen and Rufibach, 2009](#); [Doss and Wellner, 2019](#)). We use the notation $\Psi_n(\theta, \psi)$ to denote the sample version $\Psi(\theta, \psi, \mathbb{F}_n)$ of $\Psi(\theta, \psi, F)$. Thus,

$$\Psi_n(\theta, \psi) = \int_{-\infty}^{\infty} \psi(x - \theta) d\mathbb{F}_n(x) - \int_{-\infty}^{\infty} e^{\psi(x - \theta)} dx. \quad (1.5)$$

We denote the maximized criterion function by

$$L(F) = \sup_{\theta \in \mathbb{R}, \psi \in \mathcal{SC}_0} \Psi(\theta, \psi, F). \quad (1.6)$$

2. Estimators

2.1. One-step estimators

This section focuses on constructing the one-step estimators of θ . To this end, our first task is to choose suitable preliminary estimator $\bar{\theta}_n$ of θ . We will impose two requirements on $\bar{\theta}_n$: \sqrt{n} -consistency and strong consistency. The sample mean and the sample median also satisfy our requirements on $\bar{\theta}_n$ when $f_0 \in \mathcal{P}_0$. For a general f_0 , the Z -estimator of the shift in the logistic location shift model satisfies these requirements under minimal regularity conditions (cf. Example 5.40 and Theorem 5.23, [Van der Vaart, 1998](#)).

Suppose \tilde{g}_n is an estimator of the centered density $g_0 = f_0(\cdot + \theta_0)$. We let $\tilde{\psi}_n$ denote the respective log-density $\log \tilde{g}_n$, and write G_n for the distribution

function corresponding to \tilde{g}_n . Ideally, to construct a one-step estimator in this setting, one would first estimate the Fisher information by

$$\hat{\mathcal{I}}_n = \int_{-\infty}^{\infty} \tilde{\psi}'_n(x - \bar{\theta}_n)^2 d\mathbb{F}_n(x), \quad (2.1)$$

and set

$$\check{\theta}_n = \bar{\theta}_n - \int_{-\infty}^{\infty} \frac{\tilde{\psi}'_n(x - \bar{\theta}_n)}{\hat{\mathcal{I}}_n} d\mathbb{F}_n(x). \quad (2.2)$$

It is not hard to see that the choice of \tilde{g}_n will heavily influence the performance of $\check{\theta}_n$. We want \tilde{g}_n to be symmetric about 0, and if possible, log-concave.

A simple way of estimating $f_0 = g_0(\cdot - \theta_0)$ is to compute its MLE in \mathcal{LC} , the class of all log-concave densities. However, this log-concave MLE, which we will denote by \hat{h}_n from now on, may violate the requirement of symmetry. In what follows, we show that, this naive estimator \hat{h}_n can be improved to yield better estimators of g_0 .

Symmetrized estimator \tilde{g}_n^{sym}

The simplest way of obtaining a symmetrized estimator of g_0 from \hat{h}_n is by symmetrizing the latter about $\bar{\theta}_n$, which yields

$$\tilde{g}_n^{sym}(z) = \frac{1}{2} \left(\hat{h}_n(\bar{\theta}_n + z) + \hat{h}_n(\bar{\theta}_n - z) \right), \quad z \in \mathbb{R}. \quad (2.3)$$

Although \tilde{g}_n^{sym} is symmetric, it is not, in general, log-concave. Additionally, the performance of \tilde{g}_n^{sym} can suffer from the lack of smoothness of \hat{h}_n . Our simulations show that the one-step estimator based on \tilde{g}_n^{sym} has considerably less efficiency, especially for smaller sample sizes. Therefore, we seek to improve the performance of \tilde{g}_n^{sym} by incorporating smoothing, which leads to our next estimator of g_0 .

Smoothed symmetrized estimator $(\tilde{g}_n^{sym})^{sm}$

To perform smoothing, our first task is to choose a data-dependent smoothing parameter \hat{b}_n . Towards this end, following [Chen and Samworth \(2013\)](#), we take

$$\hat{b}_n^2 := \hat{\sigma}_n^2 - \tilde{\sigma}_n^2, \quad (2.4)$$

where $\hat{\sigma}_n^2$ is the sample variance and $\tilde{\sigma}_n^2$ is the variance corresponding to the density \hat{h}_n , i.e.

$$\hat{\sigma}_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2,$$

and

$$\tilde{\sigma}_n^2 = \int_{-\infty}^{\infty} z^2 \hat{h}_n(z) dz - \left(\int_{-\infty}^{\infty} z \hat{h}_n(z) dz \right)^2.$$

that the right hand side of (2.4) is positive follows from (2.1) of [Chen and Samworth \(2013\)](#). We define the smoothed symmetrized estimator by

$$(\tilde{g}_n^{sym})^{sm}(z) := \frac{1}{\hat{b}_n} \int_{-\infty}^{\infty} \tilde{g}_n^{sym}(z-t)\varphi(t/\hat{b}_n)dt, \quad z \in \mathbb{R}, \quad (2.5)$$

where φ is the standard normal density. The estimator $(\tilde{g}_n^{sym})^{sm}$ can also be represented in terms of the smoothed version of \hat{h}_n , which we define by

$$\hat{h}_n^{sm}(z) = \frac{1}{\hat{b}_n} \int_{-\infty}^{\infty} \hat{h}_n(z-t)\varphi(t/\hat{b}_n)dt, \quad z \in \mathbb{R}. \quad (2.6)$$

To see this, note that

$$(\tilde{g}_n^{sym})^{sm}(z) = \frac{\hat{h}_n^{sm}(\bar{\theta}_n + z) + \hat{h}_n^{sm}(\bar{\theta}_n - z)}{2}. \quad (2.7)$$

Though $(\tilde{g}_n^{sym})^{sm}$ leads to a more efficient one-step estimator of θ_0 in small samples, it is not necessarily log-concave, which leaves room for further improvement.

Partial MLE estimator $\hat{g}_{\bar{\theta}_n}$

The following estimator of g_0 is obtained by maximizing the criterion function $\Psi_n(\bar{\theta}_n, \psi)$ in (1.5) over $\psi \in \mathcal{SC}_0$. From Theorem 2.1(C) of [Doss and Wellner \(2019\)](#), it follows that the maximizer $\hat{\psi}_{\bar{\theta}_n}$ satisfies

$$\int_{-\infty}^{\infty} e^{\hat{\psi}_{\bar{\theta}_n}(x)} dx = 1,$$

which leads to the density estimator

$$\hat{g}_{\bar{\theta}_n} = e^{\hat{\psi}_{\bar{\theta}_n}}. \quad (2.8)$$

We call this estimator a *Partial MLE estimator* to distinguish it from the traditional MLE of g_0 , which we will discuss in Section 2.2. Notice that $\hat{g}_{\bar{\theta}_n}$ satisfies both requirements of symmetry and log-concavity.

From [Dümbgen and Rufibach \(2009\)](#) and [Doss and Wellner \(2019\)](#), it follows that the three symmetric density estimators \tilde{g}_n we have discussed so far share the same support $[-d, d]$, where

$$d = \max(X_{(n)} - \bar{\theta}_n, \bar{\theta}_n - X_{(1)}).$$

Geometric mean type symmetrized estimator $\tilde{g}_n^{geo,sym}$

Similar to \tilde{g}_n^{sym} , this estimator is also a symmetrized version of \hat{h}_n . However, this time the symmetrization is based on the geometric mean of $\hat{h}_n(\bar{\theta}_n + \cdot)$ and $\hat{h}_n(\bar{\theta}_n - \cdot)$. For $z \in \mathbb{R}$, we define this estimator by

$$\tilde{g}_n^{geo,sym}(z) := C_n^{geo} \left(\hat{h}_n(\bar{\theta}_n + z) \hat{h}_n(\bar{\theta}_n - z) \right)^{1/2}, \quad (2.9)$$

where C_n^{geo} is a random normalizing constant. Since the sum of two concave functions is again concave, it is not hard to see that $\tilde{g}_n^{geo,sym}$ is log-concave. However, its support

$$\text{supp}(\tilde{g}_n^{geo,sym}) = [-\min(X_{(n)} - \bar{\theta}_n, \bar{\theta}_n - X_{(1)}), \min(X_{(n)} - \bar{\theta}_n, \bar{\theta}_n - X_{(1)})]$$

is smaller than that of the previous estimators of g_0 .

Constructing the truncated one-step estimator

Ideally, one's first choice for a one step estimator will be the estimator $\check{\theta}_n$ suggested by (2.2), which we will refer to as the untruncated one-step estimator from now on. However, when ϕ'_0 is unbounded near the boundary of its domain, $\tilde{\psi}'_n$ is also unbounded on its domain (see Corollary 3 in Section 3.1). In this case, $\tilde{\psi}'_n$'s behavior can be hard to control in the tails. To avoid these difficulties, we consider a truncated one-step estimator. Groeneboom and Hendrickx (2018) used a similar idea in a current status linear regression model where a truncated log-likelihood was considered to avoid problems in the tails of the unknown error distribution.

We take the truncation parameter η to be a small fixed positive number in the interval $(0, 1/2)$, and denote by ξ_n the $(1-\eta)$ -th quantile of \tilde{G}_n , the distribution function of \tilde{g}_n . The symmetry of \tilde{g}_n about 0 implies that $\tilde{G}_n^{-1}(\eta) = -\xi_n$.

Two potential estimators of the Fisher information are

$$\hat{\mathcal{I}}_n^*(\eta) = \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n)^2 d\tilde{G}_n(x - \bar{\theta}_n),$$

and

$$\hat{\mathcal{I}}_n(\eta) = \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n)^2 d\mathbb{F}_n(x). \quad (2.10)$$

Though it can be shown that the above estimators differ by a $o_p(1)$ term, our simulations indicate that the estimator $\hat{\mathcal{I}}_n(\eta)$ yields a more efficient one-step estimator. Therefore, we take $\hat{\mathcal{I}}_n(\eta)$ to be our estimator of the Fisher information.

In the same spirit as the untruncated one-step estimator $\check{\theta}_n$ in (2.2), we now construct a truncated one-step estimator of θ_0 as follows:

$$\tilde{\theta}_n = \bar{\theta}_n - \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \frac{\tilde{\psi}'_n(x - \bar{\theta}_n)}{\hat{\mathcal{I}}_n(\eta)} d\mathbb{F}_n(x). \quad (2.11)$$

Note that we use all the data to estimate $\bar{\theta}_n$, $\tilde{\psi}'_n$, and \tilde{g}_n . The truncation comes only in the step where we set up the one step estimator. Next, observe that $\hat{\mathcal{I}}_n(\eta)$ is always smaller than $\hat{\mathcal{I}}_n$, the estimator of the untruncated Fisher information (see (2.1)). In Section 3.1, we show that the asymptotic variance of $\tilde{\theta}_n$ has an inverse relation with the Fisher information, indicating that the untruncated

estimator in (2.2) will be asymptotically more efficient. However, our simulations in Section 4 show that for sufficiently small η , this loss in efficiency is negligible for most densities. It is also evident that $\widehat{\mathcal{L}}_n(\eta)$ increases in η , so smaller values of η will lead to higher asymptotic efficiency. We will discuss about the relation between η and this asymptotic efficiency in more detail in Section 3.1.

2.2. Maximum likelihood estimator (MLE)

This section aims to establish the existence and the basic properties of the MLE of (θ_0, g_0) . We already mentioned in Section 1 that the MLE of f_0 in $\mathcal{S}\mathcal{L}\mathcal{C}_\theta$ exists for a fixed $\theta \in \mathbb{R}$. As we will see, this fact plays a major role in the derivation of the MLE $(\widehat{\theta}_n, \widehat{g}_n)$ of (θ_0, g_0) .

Let $\widehat{\psi}_n = \log \widehat{g}_n$. Recalling the definition of $\Psi_n(\theta, \psi)$ from (1.5), we observe that $(\widehat{\theta}_n, \widehat{\psi}_n)$ satisfies

$$(\widehat{\theta}_n, \widehat{\psi}_n) = \arg \max_{\theta \in \mathbb{R}, \psi \in \mathcal{S}\mathcal{C}_0} \Psi_n(\theta, \psi).$$

Let us define

$$\widehat{\psi}_\theta := \arg \max_{\psi \in \mathcal{S}\mathcal{C}_0} \Psi_n(\theta, \psi),$$

which exists for a fixed $\theta \in \mathbb{R}$ by Theorem 2.1(c) of Doss and Wellner (2019). It is not hard to see that

$$\widehat{\theta}_n = \arg \max_{\theta \in \mathbb{R}} \Psi_n(\theta, \widehat{\psi}_\theta) \quad \text{and} \quad \widehat{\psi}_n = \widehat{\psi}_{\widehat{\theta}_n}.$$

Note that $\widehat{g}_n = e^{\widehat{\psi}_n}$ is the MLE of g_0 , and $\widehat{f}_n = \widehat{g}_n(\cdot - \widehat{\theta}_n)$ is the MLE of f_0 . We let \widehat{G}_n and \widehat{F}_n denote the distribution functions corresponding to \widehat{g}_n and \widehat{f}_n , respectively. Also, we let $\widehat{\phi}_n$ denote the log-density $\log \widehat{f}_n$.

Before getting into the analysis of the existence of $(\widehat{\theta}_n, \widehat{\psi}_n)$ for a general case, let us consider a special case first. Suppose \mathbb{F}_n is degenerate, i.e. $\mathbb{F}_n\{x_0\} = 1$ for some $x_0 \in \mathbb{R}$. Intuition leads us to make the guess that in this case, $\widehat{\theta}_n = x_0$. Now for $k \geq 1$, we denote A_k to be the set $[-1/(2k), 1/(2k)]$, and consider the sequence of functions $\{\psi_k\}_{k \geq 1} \in \mathcal{S}\mathcal{C}_0$ defined by

$$\psi_k(x) = \begin{cases} \log k, & x \in A_k \\ -\infty, & \text{o.w.} \end{cases}$$

Observe that

$$\Psi_n(x_0, \psi_k) = \log k - 1 \rightarrow \infty, \quad \text{as } k \rightarrow \infty.$$

Therefore, x_0 indeed is a candidate for the MLE of θ_0 . However, the MLE of ψ_0 , i.e. $\widehat{\psi}_n$, does not exist in this case. To verify, observe that if $\widehat{\psi}_n$ does exist for some $\widehat{\theta}_n \in \mathbb{R}$, we also have

$$\widehat{\psi}_n(x_0 - \widehat{\theta}_n) - \int_{-\infty}^{\infty} e^{\widehat{\psi}_n(x)} dx = \Psi_n(x_0, \widehat{\psi}_n) \geq \lim_{k \rightarrow \infty} \Psi_n(x_0, \psi_k) = \infty,$$

leading to

$$\widehat{\psi}_n(x_0 - \widehat{\theta}_n) = \infty,$$

which contradicts the fact that $\widehat{\psi}_n$ is a proper concave function. Hence, we conclude that the MLE of (θ_0, ψ_0) does not exist when \mathbb{F}_n is degenerate. So we only need to consider the case when \mathbb{F}_n is non-degenerate. In this case, we will show that, the map $\theta \mapsto \Psi_n(\theta, \widehat{\psi}_\theta)$ is continuous, and the MLE $(\widehat{\theta}_n, \widehat{\psi}_n)$ exists.

Lemma 1. *If \mathbb{F}_n is non-degenerate, $\theta \mapsto \Psi_n(\theta, \widehat{\psi}_\theta)$ is a continuous map.*

Figure 1 illustrates the continuity of $\Psi_n(\theta, \widehat{\psi}_\theta)$ in θ , based on for two standard Gaussian samples of size 5 and 100.




Fig 1: Plot of $\Psi_n(\theta, \widehat{\psi}_\theta)$ vs θ generated from the standard Gaussian distribution. Here the blue ticks, the green line segments (dotted), and the orange dashed line represent the data points, the knots of $\widehat{\psi}_n$, and $\widehat{\theta}_n$ respectively.

Our next theorem establishes the existence of a maximizer of $\Psi_n(\theta, \widehat{\psi}_\theta)$ for non-degenerate \mathbb{F}_n .

Theorem 1. *When \mathbb{F}_n is non-degenerate, the MLE $(\widehat{\theta}_n, \widehat{\psi}_n)$ of (θ_0, ψ_0) exists. Moreover, $\widehat{\theta}_n \in [X_{(1)}, X_{(n)}]$.*

Observe that Theorem 1 does not ensure the uniqueness of $\widehat{\theta}_n$. Since $\Psi_n(\theta, \psi)$ may not be jointly concave in θ and ψ , existence of a maximizer does not lead to its uniqueness. For a particular choice of $\widehat{\theta}_n$ though, the estimator $\widehat{\psi}_n = \widehat{\psi}_{\widehat{\theta}_n}$ is unique by Theorem 2.1(c) of Doss and Wellner (2019). Therefore, if (θ, ψ_1) and (θ, ψ_2) both are MLEs of (θ_0, ψ_0) , we must have $\psi_1 = \psi_2$.

In all our simulations, $\widehat{\theta}_n$ turned out to be unique, even when the underlying density f_0 was skewed or non-log-concave. Considering this fact, in what follows, we refer to $\widehat{\theta}_n$ as “the MLE” instead of “an MLE”. We must remark that even if $\widehat{\theta}_n$ is not unique, all our theorems still hold for each version of $\widehat{\theta}_n$.

Now we discuss some finite-sample properties of $\widehat{\psi}_n$. The following theorem sheds some light on the structure of $\widehat{\psi}_n$. We find that $\widehat{\psi}_n$ is piecewise linear, and $\mathcal{K}(\widehat{\psi}_n)$, i.e. the set of the knots of $\widehat{\psi}_n$, is a subset of the dataset. This theorem is a direct consequence of Theorem 2.1(c) of Doss and Wellner (2019).

Theorem 2. For \mathbb{F}_n non-degenerate, the MLE $\widehat{\psi}_n$ of ψ_0 is piecewise linear with knots belonging to a subset of the set $\{0, \pm|X_1 - \widehat{\theta}_n|, \dots, \pm|X_n - \widehat{\theta}_n|\}$. Also, for $x \notin [-|X - \widehat{\theta}_n|_{(n)}, |X - \widehat{\theta}_n|_{(n)}]$, we have $\widehat{\psi}_n(x) = -\infty$. Moreover if $0 \notin \{\pm|X_1 - \widehat{\theta}_n|, \dots, \pm|X_n - \widehat{\theta}_n|\}$, then $\widehat{\psi}'_n(0\pm) = 0$.

There is no closed form for $\widehat{\psi}_n$, though $\widehat{\psi}_n$ can be characterized by a family of inequalities. The following two theorems provide characterizations of $\widehat{\psi}_n$.

Theorem 3. Suppose $\widehat{\theta}_n$ is the MLE of θ_0 . Consider $g = e^\psi \in \mathcal{S}\mathcal{L}\mathcal{C}_0$. Then $(\widehat{\theta}_n, g)$ is the MLE of (θ_0, g_0) if and only if

$$\int_{-\infty}^{\infty} \Delta(x - \widehat{\theta}_n) d\mathbb{F}_n(x) \leq \int_{-\infty}^{\infty} \Delta(x) g(x) dx$$

for all $\Delta : \mathbb{R} \mapsto \mathbb{R}$ such that $\psi + t\Delta \in \mathcal{S}\mathcal{C}_0$ for some $t > 0$.

Note that the MLE in $\mathcal{L}\mathcal{C}$ (Dümbgen and Rufibach, 2009) or $\mathcal{S}\mathcal{L}\mathcal{C}_0$ (Doss and Wellner, 2019) also satisfy similar characterizations. The proof of Theorem 3 follows from Theorem 2.2(c) of Doss and Wellner (2019).

Our next theorem exhibits another characterization of $\widehat{\psi}_n$. This theorem also follows as a direct consequence of Theorem 2.4(C) of Doss and Wellner (2019).

Theorem 4. Suppose $\widehat{\theta}_n$ is the MLE of θ_0 . For $g \in \mathcal{S}\mathcal{L}\mathcal{C}_0$, define

$$\widehat{G}_n^+(x) = 2 \int_x^{|X - \widehat{\theta}_n|_{(n)}} g(y) dy,$$

and

$$\mathbb{F}_n^+(x) = n^{-1} \sum_{i=1}^n 1\{|X - \widehat{\theta}_n|_{(i)} \geq x\}.$$

Then $(\widehat{\theta}_n, g)$ is the MLE of (θ_0, g_0) if and only if

$$\int_t^{|X - \widehat{\theta}_n|_{(n)}} \widehat{G}_n^+(x) dx \begin{cases} \leq \int_t^{|X - \widehat{\theta}_n|_{(n)}} \mathbb{F}_n^+(x) dx, & \text{if } t \in [0, |X - \widehat{\theta}_n|_{(n)}], \\ = \int_t^{|X - \widehat{\theta}_n|_{(n)}} \mathbb{F}_n^+(x) dx, & \text{if } t \in \mathcal{K}(\psi) \cap [0, |X - \widehat{\theta}_n|_{(n)}], \end{cases} \quad (2.12)$$

where $\psi = \log g$.

The implication of Theorem 4 is that the function

$$h_g(t) = \int_t^{|X - \widehat{\theta}_n|_{(n)}} [\mathbb{F}_n^+(x) - \widehat{G}_n^+(x)] dx, \quad t \geq 0 \quad (2.13)$$

is non-negative for $g = \widehat{g}_n$. Figure 2 displays the plot of $h_{\widehat{g}_n}$ constructed from a standard Gaussian sample of size 50.

The characterization theorems stated above lead to some interesting facts about the distribution function \widehat{G}_n of \widehat{g}_n . In fact, we show that \widehat{G}_n has a close connection with the empirical distribution function of the $|X_i - \widehat{\theta}_n|$'s.




Fig 2: Plot of the function $h_{\hat{g}_n}$ defined in (2.13) versus t for a standard Gaussian sample of size 50. The green vertical lines represents the non-negative knots of $\hat{\psi}_n$, and the blue ticks represent the $|X_i - \hat{\theta}_n|$'s.

Corollary 1. Suppose $(\hat{\theta}_n, \hat{\psi}_n)$ is the MLE of (θ_0, ψ_0) . Let us denote the empirical distribution function of the $|X_i - \hat{\theta}_n|$'s by $\mathbb{F}_{n,|X-\hat{\theta}_n|}$. Then the following holds almost surely on $\mathcal{K}(\hat{\psi}_n) \cap (0, \infty)$:

$$\mathbb{F}_{n,|X-\hat{\theta}_n|} - n^{-1} \leq 2\hat{G}_n - 1 \leq \mathbb{F}_{n,|X-\hat{\theta}_n|}.$$

The function $2\hat{G}_n - 1$ can be interpreted as the distribution function of the random variable $|Y|$, where $Y \sim \hat{G}_n$. Corollary 1 implies that at the knots, $\mathbb{F}_{n,|X-\hat{\theta}_n|}$ and $2\hat{G}_n - 1$ differ at most by $1/n$. The proof of Corollary 1 follows from The proof of Corollary 2.7(C) of Doss and Wellner (2019).

Our next corollary provides an upper bound for the second central moment of \hat{G}_n . This corollary is analogous to Corollary 2.8(C) of Doss and Wellner (2019).

Corollary 2. Suppose $(\hat{\theta}_n, \hat{\psi}_n)$ is the MLE of (θ_0, ψ_0) . Then

$$\text{Var}(\hat{G}_n) = \int_{-\infty}^{\infty} x^2 d\hat{G}_n(x) \leq \int_{-\infty}^{\infty} (x - \hat{\theta}_n)^2 d\mathbb{F}_n(x),$$

where \hat{G}_n is the distribution function corresponding to the density $\hat{g}_n = e^{\hat{\psi}_n}$.

Since $\int x d\hat{G}_n(x) = 0$, the proof follows by taking $\Delta = -x^2$ in Theorem 3.

3. Asymptotic properties

We now move on to a discussion of the asymptotic properties of the estimators proposed in sections 2.1 and 2.2. Before proceeding any further, we introduce some new notations. For any real valued function $h : \mathbb{R} \mapsto \mathbb{R}$, we let $\|h\|_p$ denote its L_p norm, i.e.

$$\|h\|_p = \left(\int_{-\infty}^{\infty} |h(x)|^p dx \right)^{1/p}, \quad p \geq 1.$$

We define the Wasserstein distance between two measures μ and ν on \mathbb{R} by

$$d_W(\mu, \nu) = \int_{-\infty}^{\infty} |F(x) - G(x)| dx, \quad (3.1)$$

where F and G are the distribution functions corresponding to μ and ν respectively. This representation of $d_W(\mu, \nu)$ follows from Villani (2003), page 75. By an abuse of notation, sometime we will denote the above distance by $d_W(F, G)$ as well. For a sequence of distribution functions $\{F_n\}_{n \geq 1}$, we say F_n converges weakly to F , and write $F_n \rightarrow_d F$, if for all bounded continuous functions $h : \mathbb{R} \mapsto \mathbb{R}$, we have $\lim_{n \rightarrow \infty} \int h dF_n = \int h dF$.

3.1. Asymptotic properties of the one step estimators

In this section, we will explore the asymptotic properties of the truncated one-step estimator $\tilde{\theta}_n$. First, we will establish \sqrt{n} -consistency of this estimator when $f_0 \in \mathcal{P}_0$. Then we will study the case when f_0 is not log-concave. In this section, we keep following the notations developed in Section 2.1. Thus, \tilde{g}_n denotes any of the density estimators of g_0 developed in Section 2.1.

3.1.1. Correctly specified model

We have already mentioned in Section 2.1 that, since $\tilde{\theta}_n$ depends on the nuisance score, its performance relies on how well \tilde{g}_n estimates g_0 . Therefore, before going into the analysis of the asymptotic efficiency of $\tilde{\theta}_n$, first we have to show that \tilde{g}_n is a reasonably good estimator of g_0 . Towards that end, first we establish that \tilde{g}_n is L_1 consistent for g_0 .

Lemma 2. *Suppose $f_0 \in \mathcal{P}_0$. Let \tilde{g}_n be one of the estimators of g_0 defined in Section 2.1. Suppose the preliminary estimator $\tilde{\theta}_n$ is a consistent estimator of θ_0 . Then it follows that*

$$\|\tilde{g}_n - g_0\|_1 \rightarrow_p 0. \quad (3.2)$$

If $\tilde{\theta}_n$ is a strongly consistent estimator of θ_0 , then “ \rightarrow_p ” can be replaced by “ $\rightarrow_{a.s.}$ ” in the above display.

Proving Lemma 2 is a key step for us because it leads to further consistency results involving $\tilde{\psi}_n, \tilde{\psi}'_n$, etc. In particular, we find that $\tilde{\psi}_n$ converges uniformly to ψ_0 over all compact subsets of $\text{int}(\text{dom}(\psi_0))$ with probability one. We can derive a very similar result concerning the convergence of $\tilde{\psi}'_n(x)$ to $\psi'_0(x)$ except that at the derivative level, the convergence can only be proved at the continuity points of ψ'_0 . The above discussion can be formulated in the following consistency theorem.

Theorem 5. *Assume $f_0 \in \mathcal{P}_0$. Suppose \tilde{g}_n is one of the estimators of g_0 defined in Section 2.1, and $\bar{\theta}_n$ is a consistent estimator of θ_0 . Let $\{y_n\}_{n \geq 1}$ be any sequence of random variables converging to 0 in probability. Then on any compact set $K \subset \text{int}(\text{dom}(\psi_0))$ we have,*

$$(A) \quad \sup_{x \in \mathbb{R}} |\tilde{g}_n(x + y_n) - g_0(x)| \rightarrow_p 0.$$

$$(B) \quad \sup_{x \in K} |\tilde{\psi}_n(x + y_n) - \psi_0(x)| \rightarrow_p 0.$$

(C) *Suppose $x \in \text{int}(\text{dom}(\psi_0))$ is a continuity point of ψ'_0 . Then*

$$\tilde{\psi}'_n(x + y_n) \rightarrow_p \psi'_0(x).$$

If $\bar{\theta}_n$ is a strongly consistent estimator of θ_0 , then “ \rightarrow_p ” can be replaced by “ $\rightarrow_{a.s.}$ ” in the above displays.

Now we present a corollary to Theorem 5. We show that if ψ_0 is unbounded, $\tilde{\psi}'_n$ imitates ψ'_0 . However, we do not know whether the sequence of concave functions $\tilde{\psi}'_n$ stays uniformly bounded inside the domain when ψ'_0 is bounded.

Corollary 3. *Suppose $f_0 \in \mathcal{P}_0$. Then under the assumptions of Theorem 5, if ψ_0 satisfies*

$$\sup_{x \in \text{int}(\text{dom}(\psi_0))} |\psi'_0(x)| = \infty,$$

we have

$$\limsup_n \sup_{x \in \text{int}(\text{dom}(\tilde{\psi}_n))} |\tilde{\psi}'_n(x)| = \infty \quad a.s.$$

Our next step is analyzing the asymptotic limit of the truncated Fisher information estimator $\hat{\mathcal{I}}_n(\eta)$. It turns out that $\hat{\mathcal{I}}_n(\eta)$ consistently estimates

$$\mathcal{I}_{f_0}(\eta) = \int_{F_0^{-1}(\eta)}^{F_0^{-1}(1-\eta)} \phi'_0(x)^2 dF_0(x), \quad (3.3)$$

which can be seen as a truncated version of the original Fisher information \mathcal{I}_{f_0} . The convergence of $\hat{\mathcal{I}}_n(\eta)$, combined with the consistency results in Lemma 2 and Theorem 5, leads to the desired consistency of $\bar{\theta}_n$.

Lemma 3. *Under the conditions of Theorem 5, we have $\widehat{\theta}_n \rightarrow_p \theta_0$ and $\widehat{\mathcal{I}}_n(\eta) \rightarrow_p \mathcal{I}_{f_0}(\eta)$, where $\mathcal{I}_{f_0}(\eta)$ was defined in (3.3). If $\bar{\theta}_n$ is a strongly consistent estimator of θ_0 , then $\widehat{\theta}_n \rightarrow_{a.s.} \theta_0$ and $\widehat{\mathcal{I}}_n(\eta) \rightarrow_{a.s.} \mathcal{I}_{f_0}(\eta)$.*

Now we focus on the rate of convergence of $\bar{\theta}_n - \theta_0$. In this case, we make the assumption that $\bar{\theta}_n$ is \sqrt{n} -consistent. A similar assumption was made by Stone (1975).

Theorem 6. *Suppose $f_0 \in \mathcal{P}_0$ and $\bar{\theta}_n$ is a \sqrt{n} -consistent estimator of θ_0 . Then the following assertion holds for the estimator $\tilde{\theta}_n$ of θ_0 defined in (2.11):*

$$\sqrt{n}(\tilde{\theta}_n - \theta_0) \rightarrow_d N(0, \mathcal{I}_{f_0}(\eta)^{-1}), \quad (3.4)$$

where $\mathcal{I}_{f_0}(\eta)$ is as defined in (3.3).

Remark 1. Although we do not know the rate of convergence of $\tilde{\theta}_n$ for a general $f_0 \in \mathcal{P}_0$, in the special case when f_0 is bounded away from 0 on its support, it can be shown that $\sqrt{n}(\tilde{\theta}_n - \theta_0)$ is asymptotically normal with variance $\mathcal{I}_{f_0}^{-1}$. We conjecture that the same holds for any $f_0 \in \mathcal{P}_0$. Our simulation study in Section 4 does not refute this conjecture.

Theorem 7 underscores the importance of $\mathcal{I}_{f_0}(\eta)$ for us in that it is inversely proportional to the asymptotic efficiency of $\tilde{\theta}_n$. Now we describe in greater detail how η affects the asymptotic efficiency of $\tilde{\theta}_n$.

It is natural to expect $\mathcal{I}_{f_0}(\eta)$ to be close to \mathcal{I}_{f_0} if $\eta \in (0, 1/2)$ is small. However, the magnitude of this closeness is likely to be controlled by the underlying density f_0 . Keeping that in mind, we seek to investigate how the ratio $\mathcal{I}_{f_0}(\eta)/\mathcal{I}_{f_0}$ behaves as a function of η for different choices of f_0 . Our choices of f_0 include the standard normal, Laplace and logistic densities. Note that all these densities are members of \mathcal{P}_0 with the center of symmetry $\theta_0 = 0$. We have plotted them in Figure 3(a) for convenience.

Another choice for f_0 is the standard symmetrized beta density with parameter $r > 2$ (see Figure 3(b)), which we define by

$$f_0(x) \equiv f_{0,r}(x) = \frac{\Gamma((3+r)/2)}{\sqrt{\pi r} \Gamma(1+r/2)} \left(1 - \frac{x^2}{r}\right)^{r/2} 1_{[-\sqrt{r}, \sqrt{r}]}(x), \quad r > 0, \quad (3.5)$$

where Γ is the usual Gamma function. It is straightforward to verify that in this case the score equals

$$\phi'_0(x) = \frac{-x}{1 - x^2/r} 1_{[-\sqrt{r}, \sqrt{r}]}(x).$$

Some computation shows that $r \leq 2$ leads to $\mathcal{I}_{f_{0,r}} = \infty$. However for $r > 2$,

$$\mathcal{I}_{f_{0,r}} = \frac{r \Gamma\left(\frac{r}{2} - 1\right) \Gamma\left(\frac{3+r}{2}\right)}{2 \Gamma\left(\frac{r}{2} + 1\right) \Gamma\left(\frac{1+r}{2}\right)} < \infty,$$

and $f_0 \in \mathcal{P}_0$. Figure 3(c) displays the plot of $\mathcal{I}_{f_{0,r}}$ versus r for the symmetrized beta density, which depicts that $\mathcal{I}_{f_{0,r}}$ decreases steeply for $r > 2$. This finding is consistent with \mathcal{I}_{f_0} being ∞ when f_0 is the uniform density on $[-1, 1]$.

Now we examine the ratio $\mathcal{I}_{f_0}(\eta)/\mathcal{I}_{f_0}$, which gives the asymptotic efficiency of the truncated one step estimators in the correctly specified model. Figure 4 illustrates the plot of $\mathcal{I}_{f_0}(\eta)/\mathcal{I}_{f_0}$ versus η for the above-mentioned densities. It may not be observable from the plot but we verified that for logistic, normal, and the Laplace distribution, the asymptotic efficiency is greater than 0.98 for $\eta \in (0, 0.001)$. However, Figure 4 illustrates that for the symmetrized beta distribution, the situation is somewhat different. We considered symmetrized beta distributions with parameter $r = 2.1, 2.5, 3.5,$ and 4.5 . Figure 4 indicates that in this case, the asymptotic efficiency is quite small even for smaller values of η . This trend becomes more extreme as r decreases to 2.1. We calculated that for $r = 2.1$, as η goes down from 0.01 to 10^{-6} , the ratio $\mathcal{I}_{f_0}(\eta)/\mathcal{I}_{f_0}$ only increases from 0.05 to 0.233.

The aberration for the symmetrized beta case can be explained inspecting Figure 3(c), which elucidates that as r decreases, the slope of the density near the boundary becomes sharper. As a result, the score, which equals f'_0/f_0 , also becomes large. In particular,

$$|\phi'_0(x)| \sim \frac{\sqrt{r}/2}{1 - |x|/\sqrt{r}} \quad \text{as } |x| \rightarrow \sqrt{r}.$$

Therefore, $\mathcal{I}_{f_{0,r}}$ grows very fast as r decreases. Since $\mathcal{I}_{f_{0,r}}$ does not depend on the value of the score near the boundary, it does not grow as fast as $r \downarrow 0$. Therefore, truncation leads to greater loss of asymptotic efficiency for symmetrized beta densities, especially for smaller values of r , which is reflected in Figure 4.




Fig 3: (a) Plot of the standard Laplace, standard normal and standard logistic densities. (b) Plot of the symmetrized beta (SB) densities $f_{0,r}$, defined in (3.5) for different values of r . (c) Plot of Fisher information $\mathcal{I}_{f_{0,r}}$ versus r where $f_{0,r}$ is the symmetrized beta density.




Fig 4: Plot of $\mathcal{I}_{f_0}(\eta)/\mathcal{I}_{f_0}$ versus η for different densities

3.1.2. Misspecified model: when f_0 may not be log-concave

The consistency properties of $\tilde{\theta}_n$ we have discussed so far rely on the log-concavity of f_0 . However, even if f_0 is not log-concave, the parameter θ_0 is still well-defined as the location parameter as long as $f_0 \in \mathcal{S}_{\theta_0}$. It is natural to ask then, in such a case, can $\tilde{\theta}_n$ be still consistent? It turns out that the answer is affirmative under some additional conditions. In what follows, we address this question in detail.

Let us assume that $f_0 \in \mathcal{S}_{\theta_0}$. First we will state the additional conditions which are required for the consistency of $\tilde{\theta}_n$ when $f_0 \notin \mathcal{P}_0$. First of all, we impose a restriction on F_0 . We require F_0 to be non-degenerate and to possess a finite first moment.

Condition A. F_0 is a non-degenerate distribution function satisfying

$$\int_{-\infty}^{\infty} |x| dF_0(x) < \infty.$$

We will see that this restriction, which also features in [Dümbgen *et al.* \(2011\)](#), [Chen and Samworth \(2013\)](#), and [Xu and Samworth \(2019\)](#), is needed for the existence of an L_1 -limit of \tilde{g}_n . The latter is an important step in proving the consistency of $\tilde{\theta}_n$. Note that, in the special case when $f_0 \in \mathcal{P}_0$, condition A is automatically satisfied ([Cule and Samworth, 2010](#), Lemma 1). We also require

f_0 to be bounded above, which is needed to ensure the pointwise convergence of \tilde{g}_n to a limit. Note that any member of \mathcal{P}_0 is bounded by Lemma 1 of [Cule and Samworth \(2010\)](#). Thus, we restrict our attention to the model

$$\mathcal{P}_1 = \left\{ f \in \mathcal{P} \mid f \in \mathcal{S}_\theta \text{ for some } \theta \in \mathbb{R}, \sup_x f(x) < \infty, \int_{-\infty}^{\infty} |x|f(x)dx < \infty \right\}. \quad (3.6)$$

Note that $\mathcal{P}_0 \subsetneq \mathcal{P}_1$ but $\mathcal{P}_1 \not\subseteq \mathcal{P}_s$, where the latter was considered by [Stone \(1975\)](#) and [Beran \(1974\)](#) (see (1.3)) because \mathcal{P}_1 allows densities with infinite Fisher information.

The log-concave projection theorem developed in [Dümbgen et al. \(2011\)](#), [Xu and Samworth \(2019\)](#), and [Chen and Samworth \(2013\)](#) plays a key role in our analysis. Therefore, we will provide some background on it. For the time being, we assume that F is any distribution function. we denote by $\tilde{\phi}$ the maximizer of the criterion function

$$\omega(\phi, F) = \int_{-\infty}^{\infty} \phi(x)dF(x) - \int_{-\infty}^{\infty} e^{\phi(x)}dx \quad (3.7)$$

subject to $\phi \in \mathcal{C}$, the class of concave functions defined in (1.1). Theorem 2.7 of [Dümbgen et al. \(2011\)](#) implies that when F satisfies Condition A, a unique maximizer $\tilde{\phi}$ exists, and $\tilde{f} = e^{\tilde{\phi}}$ integrates to one. Also, in the case when F has a density f , \tilde{f} can be interpreted as the minimizer of the Kullback-Leiber divergence

$$d_{KL}(f, h) = \int_{-\infty}^{\infty} \log \frac{f(x)}{h(x)} f(x)dx$$

over all $h \in \mathcal{LC}$. The distribution function \tilde{F} of \tilde{f} is regarded as the log-concave projection of F onto the space of all distributions with densities in \mathcal{LC} .

Now for $f_0 \in \mathcal{P}_1$ and ω as defined in (3.7), we denote

$$\tilde{\phi}_0 = \arg \max_{\phi \in \mathcal{C}} \omega(\phi, F_0), \quad \text{and} \quad \tilde{\psi}_0 = \tilde{\phi}_0(\cdot + \theta_0). \quad (3.8)$$

By our earlier arguments, $e^{\tilde{\phi}_0}$ yields a density; which we denote by \tilde{f}_0 , and write \tilde{F}_0 for the corresponding distribution function. We also set $\tilde{g}_0 = e^{\tilde{\psi}_0}$, and $\tilde{G}_0 = \tilde{F}_0(\cdot + \theta_0)$.

We are interested in \tilde{f}_0 because under Condition A, the unconstrained log-concave MLE \hat{h}_n converges almost surely to \tilde{f}_0 in L_1 . See Theorem 2.15 of [Dümbgen et al. \(2011\)](#) for a proof. Therefore, from (2.3) and (2.9), it is not hard to guess that the limit of \tilde{g}_n^{sym} and $\tilde{g}_n^{geo,sym}$ will be dependent on \tilde{f}_0 . We will show that for $f_0 \in \mathcal{P}_1$, both these density estimators are strongly L_1 consistent for \tilde{g}_0 when f_0 satisfies Condition A.

Next we consider the maximization of the criterion function $\omega(\phi, F)$ in ϕ over the constrained class \mathcal{SC}_θ for some fixed $\theta \in \mathbb{R}$. We denote

$$\bar{\phi}_\theta = \arg \max_{\phi \in \mathcal{SC}_\theta} \omega(\phi, F), \quad \text{and} \quad \bar{\psi}_\theta = \bar{\phi}_\theta(\cdot + \theta).$$

Observe that (1.4) provides an alternative representation of $\bar{\psi}_\theta$:

$$\bar{\psi}_\theta = \arg \max_{\psi \in \mathcal{SC}_0} \Psi(\theta, \psi, F).$$

Proposition 2(iii) of Xu and Samworth (2019) entails that under Condition A, $\bar{\phi}_\theta$ exists, it is unique, and its exponential yields a density. Notice that when $F = \mathbb{F}_n$, the density $e^{\bar{\psi}_{\hat{\theta}_n}}$ equals our partial MLE estimator $\hat{g}_{\hat{\theta}_n}$, which is why we study the constrained projection of F as well.

For a general F satisfying condition A, $\bar{\phi}_\theta$ need not equal $\tilde{\phi}$ since $\bar{\phi}_\theta$ is symmetric while $\tilde{\phi}$ is not necessarily symmetric. However, there is some connection between these two. Our next lemma underscores this connection.

Lemma 4. *Suppose F satisfies condition A. Define*

$$F_\theta^{sym}(x) = 2^{-1} \left(F(x) + 1 - F(2\theta - x) \right). \quad (3.9)$$

Then it follows that

$$\arg \max_{\phi \in \mathcal{SC}_\theta} \omega(\phi, F) = \arg \max_{\phi \in \mathcal{C}} \omega(\phi, F_\theta^{sym}),$$

where ω is the criterion function defined in (3.7).

Lemma 4 indicates that if F has a density in \mathcal{S}_θ , we do have $\bar{\phi}_\theta = \tilde{\phi}$. From the above, it follows that the projection of $f_0 \in \mathcal{P}_1$ onto $\mathcal{SLLC}_{\theta_0}$ yields \tilde{f}_0 .

Now let us consider the case of the partial MLE estimator $\hat{g}_{\hat{\theta}_n}$. It turns out that if additionally $\hat{\theta}_n$ is consistent, $\hat{g}_{\hat{\theta}_n}$ is an L_1 consistent estimator of \tilde{g}_0 , where \tilde{g}_0 denotes the centered density $f_0(\cdot + \theta_0)$.

Indeed for $f_0 \in \mathcal{P}_1$ and $\hat{\theta}_n \rightarrow_p \theta_0$, all the estimators \tilde{g}_n developed in Section 3.1 are L_1 consistent estimators of \tilde{g}_0 , except the smoothed symmetrized estimator $(\tilde{g}_n^{sym})^{sm}$. Before describing the limit for this case, we will elaborate a little bit on why this difference occurs.

The different behavior of $(\tilde{g}_n^{sym})^{sm}$ stems from the fact that the smoothing parameter \hat{b}_n defined in (2.4) does not necessarily decrease to 0 as n increases unless f_0 is log-concave. Also, Condition A is not sufficient for $\lim_{n \rightarrow \infty} \hat{b}_n$ to exist, for which, the finiteness of the second central moment of f_0 is also required (Chen and Samworth, 2013). Provided

$$\int_{-\infty}^{\infty} z^2 f_0(z) dz < \infty, \quad (3.10)$$

It can be shown that

$$\hat{b}_n^2 \rightarrow_{a.s.} \tilde{b}^2 = \int_{-\infty}^{\infty} z^2 (f_0(z) - \tilde{f}_0(z)) dz, \quad (3.11)$$

which is non-negative for $f_0 \in \mathcal{P}_1$ (cf. Remark 2.3 of Dümbgen et al., 2011), and 0 for $f_0 \in \mathcal{P}_0$. Denote by \tilde{g}_0^{sm} the convolution of \tilde{g}_0 and a centered Gaussian

density with variance \tilde{b}^2 . It turns out that \hat{h}_n^{sm} and $(\tilde{g}_n^{sym})^{sm}$ are L_1 consistent for $\tilde{g}_0^{sm} = \tilde{f}_0^{sm}(\cdot - \theta_0)$.

Lemma 5, which is a generalization of Lemma 2, summarizes the above discussion.

Lemma 5. *Suppose $f_0 \in \mathcal{P}_1$, which was defined in (3.6). Let \tilde{g}_n be one of the estimators of g_0 defined in Section 2.1, and suppose that the preliminary estimator $\tilde{\theta}_n$ is a consistent estimator of θ_0 . Let $\tilde{g}_0 = e^{\tilde{\psi}_0}$, where $\tilde{\psi}_0$ was defined in (3.8). Then for $\tilde{g}_n = \tilde{g}_n^{sym}$, $\hat{g}_{\tilde{\theta}_n}$, and $\hat{h}_n^{geo, sym}$, it follows that*

$$\|\tilde{g}_n - \tilde{g}_0\|_1 \rightarrow_p 0. \quad (3.12)$$

Suppose f_0 additionally satisfies (3.10). Then we also have

$$\|(\tilde{g}_n^{sym})^{sm} - \tilde{g}_0^{sm}\|_1 \rightarrow_p 0,$$

where \tilde{g}_0^{sm} is the convolution of \tilde{g}_0 and a centered Gaussian density with variance \tilde{b}^2 defined in (2.4). If $\tilde{\theta}_n$ is a strongly consistent estimator of θ_0 , then “ \rightarrow_p ” can be replaced by “ $\rightarrow_{a.s.}$ ” in the above displays.

We also have an analog of Theorem 5 for $f_0 \in \mathcal{P}_1$. (See Theorem 12 in Section 6.1.)

Now we are in a position to address the convergence of $\hat{\mathcal{I}}_n(\eta)$ defined in (2.10). It turns out that, unlike the case for $f_0 \in \mathcal{P}_0$, $\hat{\mathcal{I}}_n(\eta)$ is no longer a consistent estimator of $\mathcal{I}_{f_0}(\eta)$ because the asymptotic limit of the former now depends on $\tilde{\phi}'_0$. Let us denote $\xi_0 = \tilde{G}_0^{-1}(\eta)$. We find that, for $\tilde{g}_n \neq (\tilde{g}_n^{sym})^{sm}$, $\hat{\mathcal{I}}_n(\eta)$ consistently estimates $\tilde{\mathcal{I}}_{f_0}(\eta)$, where

$$\tilde{\mathcal{I}}_{f_0}(\eta) = \int_{\tilde{F}_0^{-1}(\eta)}^{\tilde{F}_0^{-1}(1-\eta)} \tilde{\phi}'_0(x)^2 f_0(x) dx = \int_{-\xi_0}^{\xi_0} \tilde{\psi}'_0(x)^2 g_0(z) dz. \quad (3.13)$$

$\tilde{\mathcal{I}}_{f_0}(\eta)$ can be thought of as a truncated Fisher information under model misspecification. When $\tilde{\psi}_0 = \psi_0$, $\tilde{\mathcal{I}}_{f_0}(\eta)$ equals $\mathcal{I}_{f_0}(\eta)$.

Now suppose $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$. In this case, Lemma 5 says that \tilde{g}_n consistently estimates \tilde{g}_0^{sm} in L_1 metric. Letting $\tilde{f}_0^{sm} = \tilde{g}_0^{sm}(\cdot - \tilde{\theta}_n)$, we denote the distribution function corresponding to \tilde{g}_0^{sm} and \tilde{f}_0^{sm} by \tilde{G}_0^{sm} and \tilde{F}_0^{sm} , respectively. We also denote the corresponding log-densities $\log \tilde{g}_0^{sm}$ and $\log \tilde{f}_0^{sm}$ by $\tilde{\psi}_0^{sm}$ and $\tilde{\phi}_0^{sm}$. It follows that $\hat{\mathcal{I}}_n(\eta)$ converges in probability to

$$\begin{aligned} \tilde{\mathcal{I}}_{f_0}^{sm}(\eta) &= \int_{(\tilde{F}_0^{sm})^{-1}(\eta)}^{(\tilde{F}_0^{sm})^{-1}(1-\eta)} (\tilde{\phi}_0^{sm})'(x)^2 f_0(x) dx \\ &= \int_{-\xi_0^{sm}}^{\xi_0^{sm}} (\tilde{\psi}_0^{sm})'(z)^2 g_0(z) dz, \end{aligned} \quad (3.14)$$

where ξ_0^{sm} is the $(1 - \eta)$ -th quantile of \tilde{G}_0^{sm} .

Even though the limits of \tilde{g}_n and $\hat{\mathcal{I}}_n(\eta)$ deviate from g_0 and $\mathcal{I}_{f_0}(\eta)$ in the misspecified case, it turns out that $\tilde{\theta}_n$ is still consistent for θ_0 , provided $f_0 \in \mathcal{P}_1$.

This establishes that $\tilde{\theta}_n$ is robust to the violation of log-concavity of f_0 as long as $f_0 \in \mathcal{P}_1$. Our next lemma formalizes the above discussion.

Lemma 6. *Under the conditions of Lemma 5, for $\tilde{g}_n = \tilde{g}_n^{sym}, \hat{g}_{\tilde{\theta}_n}$, and $\tilde{g}_n^{geo,sym}$, we have $\hat{\mathcal{I}}_n(\eta) \rightarrow_p \tilde{\mathcal{I}}_{f_0}(\eta)$, which was defined in (3.13). Also, $\hat{\theta}_n \rightarrow_p \theta_0$ in these cases.*

When $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$, if additionally (3.10) holds, we have $\hat{\mathcal{I}}_n(\eta) \rightarrow_p \tilde{\mathcal{I}}_{f_0}^{sm}(\eta)$ (defined in (3.14)), and $\hat{\theta}_n \rightarrow_p \theta_0$. When $\bar{\theta}_n$ is a strongly consistent estimator of θ_0 , all the above convergences hold almost surely.

Theorem 6 established the \sqrt{n} -consistency of $\tilde{\theta}_n$ under the \sqrt{n} -consistency of $\bar{\theta}_n$ when $f_0 \in \mathcal{P}_0$. Likewise, Theorem 7 provides an asymptotic representation of $\sqrt{n}(\tilde{\theta}_n - \theta_0)$ for $f_0 \in \mathcal{P}_1$.

Theorem 7. *Suppose $f_0 \in \mathcal{P}_1$ is absolutely continuous and $\bar{\theta}_n$ is a \sqrt{n} -consistent estimator of θ_0 . Let $\tilde{\theta}_n$ be the truncated one-step estimator defined in (2.11) with truncation parameter $\eta \in (0, 1/2)$. Suppose $\tilde{g}_n = \tilde{g}_n^{sym}, \hat{g}_{\tilde{\theta}_n}$ or $\tilde{g}_n^{geo,sym}$. Then under the conditions of Theorem 12, the following assertion holds:*

$$\sqrt{n}(\tilde{\theta}_n - \theta_0) = \int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{\tilde{\psi}'_0(x - \theta_0)}{\tilde{\mathcal{I}}_{f_0}(\eta)} d\mathbb{Z}_n(x) + \sqrt{n}(1 - \gamma_\eta)(\bar{\theta}_n - \theta_0) + o_p(1), \quad (3.15)$$

where $\mathbb{Z}_n = \sqrt{n}(\mathbb{F}_n - F_0)$,

$$\gamma_\eta = 1 - \frac{\int_{-\xi_0}^{\xi_0} \tilde{\psi}'_0(z) \left(\tilde{\psi}'_0(z) - \psi'_0(z) \right) g_0(z) dz}{\tilde{\mathcal{I}}_{f_0}(\eta)}, \quad (3.16)$$

$\xi_0 = \tilde{G}_0^{-1}(1 - \eta)$, and $\tilde{\mathcal{I}}_{f_0}(\eta)$ is as defined in (3.13). When $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$, we have

$$\sqrt{n}(\tilde{\theta}_n - \theta_0) = \int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{(\tilde{\psi}_0^{sm})'(x - \theta_0)}{\tilde{\mathcal{I}}_{f_0}^{sm}(\eta)} d\mathbb{Z}_n(x) + \sqrt{n}(1 - \gamma_\eta^{sm})(\bar{\theta}_n - \theta_0) + o_p(1), \quad (3.17)$$

where

$$\gamma_\eta^{sm} = 1 - \frac{\int_{-\xi_0^{sm}}^{\xi_0^{sm}} (\tilde{\psi}_0^{sm})'(z) \left((\tilde{\psi}_0^{sm})'(z) - \psi'_0(z) \right) g_0(z) dz}{\tilde{\mathcal{I}}_{f_0}^{sm}(\eta)}, \quad (3.18)$$

ξ_0^{sm} is the $(1 - \eta)$ -th quantile of \tilde{G}_0^{sm} , and $\tilde{\mathcal{I}}_{f_0}^{sm}(\eta)$ is as defined in (3.14).

Theorem 7 shows that if $\bar{\theta}_n$ is an asymptotically linear estimator of θ_0 , then so is $\tilde{\theta}_n$. Moreover, the central limit theorem, (3.13), and (3.14) entail that the first terms in the expansion of $\sqrt{n}(\tilde{\theta}_n - \theta_0)$ in (3.15) and (3.17) converge weakly to

mean zero Gaussian random variables with variances $\tilde{\mathcal{I}}_{f_0}^{-1}$ and $(\tilde{\mathcal{I}}_{f_0}^{sm})^{-1}$, respectively. From (3.15) and (3.17), it is also clear that for $f_0 \notin \mathcal{P}_0$, the asymptotic distribution of $\sqrt{n}(\hat{\theta}_n - \theta_0)$ may depend on $\bar{\theta}_n$. The coefficient γ_η or γ_η^{sm} quantifies this dependence.

For $\tilde{g}_n \neq (\tilde{g}_n^{sym})^{sm}$, using Cauchy-Schwartz inequality, we obtain that

$$\begin{aligned} |1 - \gamma_\eta| &= \frac{\int_{-\xi_0}^{\xi_0} \tilde{\psi}'_0(z)(\tilde{\psi}'_0(z) - \psi'_0(z))g_0(z)dz}{\tilde{\mathcal{I}}_{f_0}(\eta)} \\ &\leq \frac{\left(\int_{-\xi_0}^{\xi_0} \tilde{\psi}'_0(z)^2 g_0(z)dz\right)^{1/2}}{\tilde{\mathcal{I}}_{f_0}(\eta)} \left(\int_{-\xi_0}^{\xi_0} (\tilde{\psi}'_0(z) - \psi'_0(z))^2 g_0(z)dz\right)^{1/2} \\ &\leq \left(\tilde{\mathcal{I}}_{f_0}(\eta)^{-1} \int_{-\infty}^{\infty} (\tilde{\phi}'_0(z) - \phi'_0(z))^2 p_0(z)dz\right)^{1/2}, \end{aligned}$$

where the last inequality follows from (3.13). The above bound on $|1 - \gamma_\eta|$ hints that the asymptotic dependence of $\sqrt{n}(\hat{\theta}_n - \theta_0)$ on $\bar{\theta}_n$ may be controlled by the $L_2(P_0)$ distance between $\tilde{\phi}'_0$ and ϕ'_0 . We can draw a similar conclusion for γ_η^{sm} replacing $\tilde{\psi}_0$ by $\tilde{\psi}_0^{sm}$.

3.2. Asymptotic properties of the MLE

In this section, we explore the asymptotic properties of the MLE $\hat{\theta}_n$ and \hat{g}_n . First, we will consider the case when $f_0 \in \mathcal{P}_0$, and then we will move on to the discussion of the misspecified case.

3.2.1. Correctly specified model

For $f_0 \in \mathcal{P}_0$, Lemma 3 showed that the one-step estimators are consistent, whereas Lemma 2 established the L_1 -consistency of the density estimators \tilde{g}_n 's established in Section 2.1. Theorem 8 below shows that the MLEs $\hat{\theta}_n$ and \hat{g}_n satisfy similar consistency properties.

Theorem 8. *Suppose $f_0 = g_0(\cdot - \theta_0) \in \mathcal{P}_0$ for some $\theta_0 \in \mathbb{R}$. Then as $n \rightarrow \infty$, $L(\mathbb{F}_n) \rightarrow_{a.s.} L(F_0)$, and the following assertions hold:*

- (a) $\hat{\theta}_n \rightarrow_{a.s.} \theta_0$.
- (b) $\hat{\psi}_n(x) \rightarrow_{a.s.} \psi_0(x)$, for all $x \in \text{int}(\text{dom}(\psi_0))$.
- (c) $\limsup_n \hat{\psi}_n(x) \leq \psi_0(x)$, a.s. for all $x \in \mathbb{R}$.
- (d) $\hat{g}_n(x) \rightarrow_{a.s.} g_0(x)$, for all $x \in \mathbb{R}$.
- (e) There exists $\alpha > 0$ such that for all $\alpha_1 < \alpha$,

$$\int_{-\infty}^{\infty} e^{\alpha_1|x|} |\hat{g}_n(x) - g_0(x)| dx \rightarrow_{a.s.} 0.$$

Moreover,

$$\sup_{x \in \mathbb{R}} e^{\alpha_1 |x|} |\widehat{g}_n(x) - g_0(x)| \rightarrow_{a.s.} 0.$$

Since g_0 is a log-concave density, there exist α and β such that $\psi_0(x) \leq -\alpha|x| + \beta$ for all $x \in \mathbb{R}$ (Cule and Samworth, 2010, Lemma 1). This is the α that appears in Theorem 8.

Our next lemma is an analog of Theorem 12 for one step estimators, and addresses the uniform convergence of $\widehat{\phi}_n$ over compact subsets of $\text{int}(\text{dom}(\psi_0))$ and the pointwise convergence of $\widehat{\phi}'_n$.

Lemma 7. *Suppose $f_0 = g_0(\cdot - \theta_0) \in \mathcal{P}_0$ for some $\theta_0 \in \mathbb{R}$. Denoting $\phi_0 = \log f_0$, on any compact set $K \subset \text{dom}(\phi_0)$, we have,*

$$\sup_{x \in K} |\widehat{\phi}_n(x) - \phi_0(x)| \rightarrow_{a.s.} 0.$$

Suppose ϕ_0 is differentiable at $x \in K$. Then it also follows that

$$\widehat{\phi}'_n(x) \rightarrow_{a.s.} \phi'_0(x),$$

and

$$\widehat{f}'_n(x) \rightarrow_{a.s.} f'_0(x).$$

Similarly for $\psi_0 = \log g_0$, on any compact set $K \subset \text{dom}(\psi_0)$, we have,

$$\sup_{x \in K} |\widehat{\psi}_n(x) - \psi_0(x)| \rightarrow_{a.s.} 0.$$

If ψ_0 is differentiable at $x \in K$, then we also have

$$\widehat{\psi}'_n(x) \rightarrow_{a.s.} \psi'_0(x),$$

and

$$\widehat{g}'_n(x) \rightarrow_{a.s.} g'_0(x).$$

Now we turn our attention to the global convergence of \widehat{g}_n . Our Theorem 9 establishes that $H(\widehat{f}_n, f_0)^2$ and $H(\widehat{g}_n, g_0)^2$ are $O_p(n^{-4/5})$. This rate is standard for a density estimator in log-concave class \mathcal{LC} . For instance, the unrestricted log-concave MLE \widehat{h}_n satisfies $H(\widehat{h}_n, f_0)^2 = O_p(n^{-4/5})$ (Doss and Wellner, 2016, Theorem 3.2). Moreover, Theorem 1 of Kim and Samworth (2016) establishes that $n^{-4/5}$ is the minimax rate of convergence for density estimation in \mathcal{LC} . Theorem 4.1(c) of Doss and Wellner (2019) obtains the same rate in connection to the MLE in the subclass \mathcal{SLLC}_0 of \mathcal{LC} . Doss and Wellner (2019) conjectured in their Remark 4.2 that $n^{-4/5}$ is the minimax rate of convergence for estimation in \mathcal{SLLC}_0 as well. In light of the above, we conjecture that the minimax rate for estimating f_0 in \mathcal{P}_0 is also $n^{-4/5}$. As a corollary to Theorem 9, it follows that $\|\widehat{g}_n - g_0\|_1 = O_p(n^{-2/5})$. Also, for the MLE of θ_0 , we can show that $|\widehat{\theta}_n - \theta_0|$ is $O_p(n^{-2/5})$.

Theorem 9. *Suppose $\theta_0 \in \mathbb{R}$ and $f_0 = g_0(\cdot - \theta_0)$ for some $g_0 \in \mathcal{SLLC}_0$. Then the following assertions hold:*

(A) As $n \rightarrow \infty$,

$$H(\widehat{f}_n, f_0) \rightarrow_{a.s.} 0 \quad \text{and} \quad H(\widehat{g}_n, g_0) \rightarrow_{a.s.} 0.$$

Moreover,

$$H^2(\widehat{f}_n, f) = O_p(n^{-4/5}).$$

(B) We also have $|\widehat{\theta}_n - \theta_0| = O_p(n^{-2/5})$ and

$$H^2(\widehat{g}_n, g_0) = O_p(n^{-4/5}).$$

In the special case when f_0 is bounded away from 0 on its support, it can be shown that $\widehat{\theta}_n$ is \sqrt{n} -consistent. We conjecture that for any $f_0 \in \mathcal{P}_0$, the MLE $\widehat{\theta}_n$ is \sqrt{n} -consistent and asymptotically efficient.

3.2.2. Misspecified model: when f_0 may not be log-concave

In Subsection 3.2.1, we saw that even if $f_0 \notin \mathcal{P}_0$, the one step estimators are still consistent as long as $f_0 \in \mathcal{P}_1$. It turns out that, for $f_0 \in \mathcal{P}_1$, not only the one step estimators, but also the MLEs $\widehat{\theta}_n$ and \widehat{g}_n are consistent. However, in case of the MLE, it is possible to say more. In particular, if $f_0 \notin \mathcal{P}_1$, it turns out that $\widehat{\theta}_n$ converges almost surely to some limit under some minimal conditions. Much as in Section 3.1, we again appeal to the log-concave projection theory to analyze the asymptotic properties. However, in this case we need to consider projection onto the class of distributions with densities in \mathcal{P}_0 , which is not covered by the existing literature. Therefore, we construct a new framework for projection onto the above-mentioned class. The consistency results for $f_0 \in \mathcal{P}_0$ follow as a special case of the general theory developed in this subsection.

In order to study the log-concave projection in context of the class \mathcal{P}_0 , first we need to introduce some new notation. For any distribution function F , let us denote

$$(\theta^*(F), \psi^*(F)) = \arg \max_{\theta \in \mathbb{R}, \psi \in \mathcal{SC}_0} \Psi(\theta, \psi, F), \quad (3.19)$$

where Ψ was defined in (1.4). This maximization problem is different from those considered in Section 3.1 because in those cases, the maximization was only with respect to ψ . We are interested in $\theta^*(F)$ and $\psi^*(F)$ because, as we will see, under some mild conditions, $\widehat{\theta}_n \rightarrow_{a.s.} \theta^*(F)$ and $\|\widehat{g}_n - e^{\psi^*(F)}\|_1 \rightarrow_{a.s.} 0$ as $n \rightarrow \infty$ provided that $\theta^*(F)$ is unique.

Proposition 1 below entails that Condition A is sufficient for the existence of the maximizers. Following the same argument as in Theorem 2.1(c) of [Doss and Wellner \(2019\)](#), one can show that when $\psi^*(F)$ exists, $g^*(F) = e^{\psi^*(F)}$ is a density. We denote the corresponding distribution function by $G^*(F)$. We also denote $\phi^*(F) = \psi^*(F)(\cdot - \theta^*(F))$, and write $f^*(F)$ for $e^{\phi^*(F)}$. The corresponding distribution function will be denoted by $F^*(F)$, which can be interpreted as the projection of F onto the space of all distribution functions with densities in $\mathcal{SLL}_{\theta^*(F)}$.

Proposition 1. *Suppose F satisfies Condition A. Then $L(F) < \infty$, and $\theta^*(F)$ and $\psi^*(F)$ exist for F , where $\theta^*(F)$ and $\psi^*(F)$ were defined in (3.19).*

Our next lemma provides a representation of $\theta^*(F)$ and $\psi^*(F)$ in terms of the unconstrained log-concave projection in (3.8). Lemma 8 is a direct consequence of Lemma 4.

Lemma 8. *Suppose a distribution function F satisfies condition A. Then*

$$\theta^*(F) = \arg \max_{\theta \in \mathbb{R}} \left(\max_{\phi \in \mathcal{C}} \omega(\phi, F_{\theta}^{sym}) \right),$$

and

$$\psi^*(F) = \arg \max_{\phi \in \mathcal{C}} \left(\phi, F_{\theta^*(F)}^{sym} \right),$$

where F_{θ}^{sym} is defined in (3.9) for any $\theta \in \mathbb{R}$, and ω is the criterion function defined in (3.7).

The benefit of the representation of $\theta^*(F)$ and $\psi^*(F)$ in terms of the unrestricted log-concave projection lies in the fact that characterizations and tools for computing the latter are available (cf. Dümbgen *et al.*, 2011, Remark 2.3, Theorem 2.7, Remark 2.10, Remark 2.11).

Note that neither Proposition 1 nor Lemma 8 provides any information about the uniqueness of $\theta^*(F)$ and $\psi^*(F)$. However, the uniqueness of $\theta^*(F)$ is critical for the consistency of $\hat{\theta}_n$. Therefore, we only consider those distribution functions which satisfy the following condition.

Condition B. *The maximizer $\theta^*(F)$ defined in (3.19) is unique for distribution function F .*

Though Condition B does not say anything about the uniqueness of $\psi^*(F)$, from Theorem 2.1(c) of Doss and Wellner (2019) or Proposition 2 of Xu and Samworth (2019), it follows that if $\theta^*(F)$ is unique, so is $\psi^*(F)$. We have not yet discovered whether there exist some characteristics of F which enforce the uniqueness condition B. Interestingly, we have not yet found an example of F satisfying condition A but violating Condition B. However, it is easy to find distributions satisfying both Conditions A and B, a trivial example being distribution functions satisfying

$$F(x) = 1 - F(2\theta - x) \tag{3.20}$$

for some $\theta \in \mathbb{R}$. If F has a density in \mathcal{S}_{θ} , (3.20) holds.

Lemma 9. *Suppose a distribution function F satisfies (3.20) for some $\theta \in \mathbb{R}$, and set $G = F(\cdot + \theta)$. Then condition B is satisfied with*

$$\theta^*(F) = \theta,$$

and

$$\psi^*(F) = \arg \max_{\psi \in \mathcal{SC}_0} \Psi(0, \psi, G) = \arg \max_{\phi \in \mathcal{C}} \omega(\phi, G),$$

where ω is the criterion function defined in (3.7).

As a corollary to Lemma 9, it follows that the density class \mathcal{P}_1 satisfies Condition B. The following example, which is taken from Example 2.9 of Dümbgen *et al.* (2011), helps in understanding the implications of Lemma 9.

Example 1. Consider a rescaled version of Student's t_2 distribution, whose density takes the form $f(x) = g(x - \theta)$, with g satisfying

$$g(x) = 2^{-1}(1 + x^2)^{-3/2}, \quad x \in \mathbb{R}.$$

Lemma 9 implies that $\theta^*(F) = \theta$. Lemma 9 also indicates that $\psi^*(F)$ can be found using the unrestricted log-concave projection theorem developed in Dümbgen *et al.* (2011). In fact, using Example 2.9 of Dümbgen *et al.* (2011), we derive that $\psi^*(F)(x) = -|x| - \log 2$, which corresponds to the Laplace density.

Example 2. This example focuses on some bimodal densities. First, we consider a Gaussian mixture density

$$f(x) = 2^{-1}(\varphi(x - 2) + \varphi(x + 2)). \tag{3.21}$$

Here φ denotes the standard Gaussian density. Since $f \in \mathcal{S}_0$, Lemma 9 implies that $\theta^*(F) = 0$. As in Example 1, $\psi^*(F)$ can be obtained using the log-concave projection illustrated in Dümbgen *et al.* (2011). Remark 2.11 of Dümbgen *et al.* (2011) implies that there exists $z > 2$ such that the projection $f^*(F)$ equals f on $\mathbb{R} \setminus [-z, z]$, and is constant on $[-z, z]$ (see Figure 5). In addition, since $f^*(F) \in \mathcal{S}\mathcal{L}\mathcal{C}_0$, we have

$$\int_{-\infty}^{\infty} f^*(F)(x)dx = 2zf(z) + 2\left(1 - F(z)\right), \tag{3.22}$$

which implies

$$1 = z\left(\varphi(z - 2) + \varphi(z + 2)\right) + 2 - \Phi(z - 2) - \Phi(z + 2),$$

where Φ is the standard Gaussian distribution function. Solving the above equation, we obtain that $z \approx 2.83$.

Similarly, consider the following mixture of Laplace densities:

$$f(x) = 4^{-1}\left(e^{-|x-2|} + e^{-|x+2|}\right). \tag{3.23}$$

Its projection $f^*(F)$ can also be found by Remark 2.11 of Dümbgen *et al.* (2011) which implies that $f^*(F)$ equals f on $\mathbb{R} \setminus [-z, z]$, and is constant on $[-z, z]$, where $z > 2$. Using (3.22), which holds in this case as well, we find that $z = 2.61$. Figure 5 displays the above bimodal densities and their symmetric log-concave projections.

Our next lemma gives some insight into the domain of $\psi^*(F)$.




Fig 5: The plot in the left panel displays the density f (in blue) in (3.21) and its projection $f^*(F)$ (in green); the plot in the right panel displays the density f in (3.23) (drawn in blue) and the corresponding $f^*(F)$ (in green).

Lemma 10. *Suppose Conditions A and B hold for a distribution function F . Then $\theta^*(F) \in \overline{J(F)}$, where $J(F) = \{0 < F < 1\}$. Moreover, letting a and b denote the left and right end-points of the set $J(F)$ respectively, we have,*

$$\text{int}(\text{dom}(\psi^*(F))) = (-d, d),$$

where $d = (b - \theta^*(F)) \wedge (\theta^*(F) - a)$.

Lemma 10 implies that for a general F_0 , the domain of ψ_0 is contained the domain of $\psi^*(F_0)$. In the special case when $\psi_0 \in \mathcal{S}_0$, which is the case for $f_0 \in \mathcal{P}_1$, we have $b - \theta^*(F) = \theta^*(F) - a$, leading to

$$\text{int}(\text{dom}(\psi^*(F_0))) = \text{int}(\text{dom}(\psi_0)).$$

On the other hand, substituting $F = \mathbb{F}_n$ in Lemma 10, we obtain that $\hat{\theta}_n \in [X_{(1)}, X_{(n)}]$, which was already stated in Theorem 1.

Now we move on to the consistency properties of the MLE, which is connected with the continuity of the maximized criterion function $F \mapsto L(F)$ defined in (1.6). As mentioned earlier in Section 3.1, Dümbgen *et al.* (2011) considered projection of F onto the class of distributions with density in \mathcal{LC} . Dümbgen *et al.* (2011) showed that under some regularity conditions, the maximized criterion function corresponding to their projection operator is continuous with respect to the Wasserstein distance d_W . Xu and Samworth (2019) obtained similar continuity result in context of \mathcal{SLLC}_0 , which is a subclass of \mathcal{LC} . We are working with

a further subclass \mathcal{P}_0 . In light of the above, it is not unnatural to expect the same continuity result to hold in our set-up. Indeed, our Theorem 10 establishes the continuity of the map $F \mapsto L(F)$ at an F satisfying Conditions A and B. Moreover, the map $F \mapsto \theta^*(F)$ is also continuous in this case.

Theorem 10. *Let $\{F_n\}_{n \geq 1}$ be a sequence of distribution functions such that $d_W(F_n, F) \rightarrow 0$ for some distribution function F . Further suppose, F and F_n satisfy Conditions A and B for each $n \geq 1$. Then as $n \rightarrow \infty$,*

$$L(F_n) \rightarrow L(F), \quad (3.24)$$

and the following assertions hold:

- (a) $\lim_n \theta^*(F_n) = \theta^*(F)$.
- (b) $\lim_n \psi^*(F_n)(x) = \psi^*(F)(x)$ for all $x \in \text{int}(\text{dom}(\psi^*(F)))$.
- (c) $\limsup_n \psi^*(F_n)(x) \leq \psi^*(F)(x)$ for all $x \in \mathbb{R}$.
- (d) $\lim_n g^*(F_n)(x) = g^*(F)(x)$ for all $x \in \mathbb{R}$.
- (e) There exists $\alpha > 0$ such that for all $\alpha_1 < \alpha$,

$$\lim_n \int_{-\infty}^{\infty} e^{\alpha_1|x|} |g^*(F_n)(x) - g^*(F)(x)| dx = 0.$$

Moreover, if $g^*(F)$ is continuous on \mathbb{R} , then as $n \rightarrow \infty$,

$$\sup_{x \in \mathbb{R}} e^{\alpha_1|x|} |g^*(F_n)(x) - g^*(F)(x)| \rightarrow 0.$$

Now we are in a situation to address the convergence of $\hat{\theta}_n$ and \hat{g}_n . Suppose F_0 satisfies Conditions A and B. Then Theorem 11 entails that $\hat{\theta}_n \rightarrow_{a.s.} \theta^*(F_0)$ and \hat{g}_n converges to $e^{\psi^*(F_0)}$ almost surely, both pointwise and in L_1 . Theorem 11 follows as a corollary of Theorem 10. This can be verified easily observing that under Condition A, $d_W(\mathbb{F}_n, F_0) \rightarrow_{a.s.} 0$ (Villani, 2009, Theorem 6.9), and \mathbb{F}_n is non-degenerate with probability 1 for large n .

Theorem 11. *Suppose F_0 satisfies Conditions A and B. Then as $n \rightarrow \infty$, $L(\mathbb{F}_n) \rightarrow_{a.s.} L(F_0)$, and the following assertions hold:*

- (a) $\hat{\theta}_n \rightarrow_{a.s.} \theta^*(F_0)$.
- (b) $\hat{\psi}_n(x) \rightarrow_{a.s.} \psi^*(F_0)(x)$, for all $x \in \text{int}(\text{dom}(\psi^*(F_0)))$.
- (c) $\limsup_n \hat{\psi}_n(x) \leq \psi^*(F_0)(x)$, for all $x \in \mathbb{R}$ a.s.
- (d) $\hat{g}_n(x) \rightarrow_{a.s.} g^*(F_0)(x)$, for all $x \in \mathbb{R}$.
- (e) There exists $\alpha > 0$ such that for all $\alpha_1 < \alpha$,

$$\int_{-\infty}^{\infty} e^{\alpha_1|x|} |\hat{g}_n(x) - g^*(F_0)(x)| dx \rightarrow_{a.s.} 0.$$

Moreover, if $g^*(F_0)$ is continuous on \mathbb{R} ,

$$\sup_{x \in \mathbb{R}} e^{\alpha_1|x|} |\hat{g}_n(x) - g^*(F_0)(x)| \rightarrow_{a.s.} 0.$$

Theorem 11 has some important consequences for F_0 satisfying (3.20) with $\theta_0 \in \mathbb{R}$. Note that $f_0 \in \mathcal{P}_1$ corresponds to such a distribution. Lemma 9 implies that Condition B holds for F_0 satisfying (3.20) with $\theta_0 \in \mathbb{R}$, which leads to the following corollary.

Corollary 4. *If F_0 has density $f_0 \in \mathcal{P}_1$, all conclusions of Theorem 11 hold with $\theta^*(F_0) = \theta_0$ and $f^*(F_0) = f_0$.*

For $f_0 \in \mathcal{P}_1$, not only $\hat{\theta}_n$ is a consistent estimator of θ_0 , but also \hat{g}_n converges to $g_0 = f_0(\cdot + \theta_0)$ almost surely, both pointwise and in L_1 . Also, note that the results for the correctly specified model \mathcal{P}_0 in Theorem 8 follow directly from Theorem 11 and Corollary 4.

4. Simulation study

In this section, we study the performance of the one step estimators $\tilde{\theta}_n$ and the MLE $\hat{\theta}_n$ for several log-concave densities symmetric about 0. These densities include the standard normal, standard logistic, and standard Laplace density, which are plotted in Figure 3(a). We also consider the symmetrized beta distribution discussed in Section 3.1 (see Figure 3(b)), whose density is given by (3.5). Taking g_0 to be any of the above densities, we let $\theta_0 = 0$.

The general set-up of the simulations is as follows. We generate 3000 samples of size $n = 30, 100, 200,$ and 500 from each of the above-mentioned densities. Then for each sample, we construct the MLE $\hat{\theta}_n$, and the one step estimators, both truncated and untruncated, defined in (2.11) and (2.2) respectively. The truncated one step estimator requires specification of the truncation parameter η . We set η to be 0.002 because Figure 3 indicates that the value of $\mathcal{I}_{f_0}(\eta)/\mathcal{I}_{f_0}$ does not vary significantly for $\eta \in (0.0001, 0.002)$, at least for normal, logistic, and the Laplace distribution.

Different choices of the preliminary estimator $\bar{\theta}_n$ and the density estimator \tilde{g}_n in (2.2) and (2.11) lead to different one-step estimators. The preliminary estimator $\bar{\theta}_n$ is generally taken to be the mean, median, or the trimmed mean (12.5% trimming from each tail). However, for the logistic distribution, we also consider the parametric MLE as an initial estimator. The density estimator \tilde{g}_n is taken to be either $\hat{h}_n(\cdot - \bar{\theta}_n)$, or the symmetrized estimators in (2.3), (2.5), (2.8), and (2.9). The performance of the resulting estimators are compared with the preliminary estimator $\bar{\theta}_n$.

To compare the performance of different estimators, we need a measure of efficiency, which we formulate as follows:

$$\text{Efficiency}(\theta_n) = \frac{1/(n\mathcal{I}_{f_0})}{\text{Var}(\theta_n)}, \quad (4.1)$$

where θ_n is some estimator of θ_0 . The reason behind this choice is that $\mathcal{I}_{f_0}^{-1}$ is the lower bound on the asymptotic variance in this problem. Since we can not directly compute $\text{Var}(\theta_n)$, we will replace it by its Monte Carlo estimate.

Our simulations reveal that all our estimators are consistent for θ_0 . However, our simulations also suggest that among the one step estimators, the smoothed symmetrized one step estimator in (2.5) and the partial MLE estimator in (2.8) have the highest efficiency uniformly over all sample sizes. Therefore, in this section, among the one step estimators, we present the asymptotic efficiency of only these two one step estimators.

4.1. Normal density

In this case, f_0 is the standard Gaussian density, which implies $\mathcal{I}_{f_0} = 1$, and $\theta_0 = 0$. In case of the Gaussian location model, the sample mean is the parametric MLE, which is also (asymptotically) efficient. Figure 6 compares the efficiency of the MLE, the partial MLE one-step estimator in (2.8), and the smoothed symmetrized one step estimator in (2.5). Some patterns become clear from Figure 6.

1. As expected, the parametric MLE, i.e. the mean, has the highest efficiency. Our estimators, however, outperform all other preliminary estimators.
2. The untruncated one step estimators have higher efficiency than their truncated counterparts.
3. Among our estimators, the smoothed symmetrized estimator has the best efficiency for small samples. It turns out that it also slightly outperforms the other estimators in large sample ($n = 500$) as well.

4.2. Logistic density

The standard logistic density has $\mathcal{I}_{f_0} = 1/3$. In this case along with the mean, the median, and the trimmed mean, we also consider the parametric MLE as a preliminary estimator. Figure 7 compares the efficiency of the one step estimators and the MLE. We list the main observations below.

1. As usual, the parametric MLE, i.e. the mean, has the highest efficiency.
2. Surprisingly, in this case, the truncated estimator outperforms the untruncated estimator when $n = 200$, and 500, regardless of the choice of the preliminary estimator. For all the other cases, the efficiency of the untruncated estimator dominates the efficiency of the truncated estimator.
3. The smoothed symmetrized one step estimator has the highest efficiency in smaller samples (sample size $n = 40, 100$). Also, we observe that the MLE has lower efficiency than both the one step estimators, the partial MLE one-step estimator and the smoothed symmetrized one step estimator.

4.3. Laplace density

In this subsection, we simulate observations from the standard Laplace density, which has $\mathcal{I}_{f_0} = 1$. In this case, the sample median is the parametric MLE

of θ_0 . Unlike the other densities we considered, this density is not smooth. In fact, it has a kink at 0 (see Figure 3). Figure 8 indicates that the smoothed symmetrized one step estimator in (2.5) suffers from the lack of smoothness, and loses its edge to the MLE and the partial MLE one step estimator, both in small and large samples. When $n = 40$ and 500, the partial MLE one step estimator has the highest efficiency. For all other sample sizes, Figure 8 suggests that the MLE has the highest efficiency. In all cases, the untruncated one step estimator outperforms the truncated version.

4.4. Symmetrized beta density

In this subsection, we consider the symmetrized beta density in (3.5), for $r = 2.1$, 2.5, and 4.5. Figure 9 compares the efficiency of the MLE and the one step estimators with the mean as the preliminary estimator. The plots for the other preliminary estimators depict the same story, and hence we do not present them. Below we list the main observations from Figure 9.

1. The efficiency of the estimators increases with r . Also, for $r = 2.1$ and 2.5, all the estimators exhibit poor efficiency even for sample size 500.
2. The untruncated one step estimators display higher efficiency than the truncated one step estimator. We note that for $r = 2.5$ and 4.5, this gap in the efficiency is higher compared to the other densities.
3. It turns out that the smoothed untruncated one step estimator has the highest asymptotic efficiency among all the estimators.

To summarize, except for the case of standard Laplace, the smoothed symmetrized one step estimator has the highest efficiency among our estimators. We also note that in most cases the untruncated one step estimator exhibits higher efficiency than its truncated counterpart.




Fig 6: Plot of efficiency vs sample size (n) for standard Gaussian density. We plotted the efficiency of the smoothed symmetrized one step estimator in (2.5) (Smoothed OE; right column), the partial MLE one-step estimator in (2.8) (Partial MLE OE; left column), and the MLE (in purple; present in both columns). The preliminary estimators $\hat{\theta}_n$ corresponding to the one-step estimators are the mean (top), median (middle), and trimmed mean (bottom). The preliminary estimator is drawn in red, the truncated and untruncated estimators are drawn in blue and green respectively.




Fig 7: Plot of efficiency vs sample size (n) when the underlying density is standard Logistic. We calculate the efficiency of the smoothed symmetrized one step estimator in (2.5) (Smoothed OE; right column), the partial MLE one-step estimator in (2.8) (Partial MLE OE; left column), and the MLE (in purple; present in both columns). The preliminary estimators $\hat{\theta}_n$ corresponding to the one-step estimators (drawn in red) are the mean (first row), median (second row), trimmed mean (third row), and the parametric MLE (fourth MLE). The truncated and untruncated estimators are drawn in blue and green respectively.




Fig 8: Plot of efficiency vs sample size (n) for the standard Laplace density. We plotted the efficiency of the smoothed symmetrized one step estimator in (2.5) (Smoothed OE; right column), the partial MLE one-step estimator in (2.8) (Partial MLE OE; left column), and the MLE (in purple; present in both columns). The preliminary estimators $\hat{\theta}_n$ corresponding to the one-step estimators are the mean (top), median (middle), and trimmed mean (bottom). The preliminary estimator is drawn in red, and the truncated and untruncated estimators are drawn in blue and green respectively.




Fig 9: Plot of efficiency vs sample size (n) when the underlying density is symmetrized beta, which is defined in (3.5). We consider $r = 2.1$ (left), 2.5 (middle), and 4.5 (right). The MLE is drawn in purple. Here we calculated the efficiency of the smoothed symmetrized one step estimator in (2.5) (Smoothed OE; right column) and the partial MLE one-step estimator in (2.8) (Partial MLE OE; left column) using the sample mean as the preliminary estimator. The efficiency of the mean is drawn in red. The truncated and untruncated versions are drawn in blue and green respectively.

5. Discussion

In this article, we revisit the problem of estimating the location parameter θ_0 in the symmetric location model. We show that the additional assumption of log-concavity can ameliorate the dependence on tuning parameters. In particular, our one-step estimators use only one additional parameter, and the MLE does not rely on any tuning parameter.

We show that the rate of convergence of the MLE $\hat{\theta}_n$ is $O_p(n^{-2/5})$ but we conjecture that it is actually \sqrt{n} -consistent, and asymptotically efficient. In our case, the parameter of interest is bundled with the infinite dimensional nuisance parameter. Obtaining the precise rate of convergence for the MLE in such semiparametric models is typically difficult. For example, in context of the monotone current status regression model, this was noted by [Murphy et al. \(1999\)](#), who obtained $n^{-1/3}$ rate for the parametric part in this model. Whether the MLE is \sqrt{n} -consistent in this model is still unknown ([Groeneboom and Hendrickx, 2018](#)). Bundled parameters also appear in single index models ([Kuchibhotla et al., 2017](#)), where not much is known about the MLE. [Chen and Samworth \(2016\)](#) proved the consistency of the joint MLE in shape-constrained additive index model, which is more general than the single index model, but nothing is known about MLE's rate of convergence.

Although our truncated one step estimator is \sqrt{n} -consistent, it is not fully asymptotically efficient. The loss of efficiency can be attributed to the use of a fixed truncation parameter. A similarity can be drawn with [Groeneboom and Hendrickx \(2018\)](#), who study a current status regression model. They also use a fixed truncation parameter to construct a \sqrt{n} -consistent z -estimator, which, similar to our case, is not fully asymptotic efficient. However, our simulations do not exhibit a big difference between the truncated and untruncated one step estimator, where the latter is conjectured to be \sqrt{n} -consistent.

Despite having the above limitations, our estimators are easily computable because fast algorithms are available for computing log-concave density estimators (using the R packages “logcondens” and “logcondens.mode”). Moreover, we show that even if the log-concavity assumption fails, all our estimators of θ_0 are still consistent under very mild conditions. In fact, the one step estimators are still \sqrt{n} -consistent. This is a consequence of the impressive stability of log-concave density estimators ([Dümbgen et al., 2011](#); [Xu and Samworth, 2019](#)). This demonstrates the usefulness of log-concavity assumption in semiparametric models in facilitating a simplified and robust estimation procedure.

6. Proofs

In what follows, we will use the following terminologies. For two sets A and B , $A \times B$ will represent the Cartesian product. For any set $A \subset \mathbb{R}$, and $x \in \mathbb{R}$, we use the usual notation $A + x$ to denote the translated set $\{y + x : y \in A\}$. The notation \bar{A} will refer to the closure of the set A .

6.1. Proofs for the one-step estimators

Proof of Lemma 2

The proof follows from Lemma 5 because when $f_0 \in \mathcal{P}_0$, the log-concave approximation \tilde{f}_0 equals f_0 , which, combined with (3.11), also implies $\tilde{b} = 0$. Hence, $\tilde{g}_0 = \tilde{g}_0^{sm} = g_0$ follows and (3.10) is automatically satisfied (Cule and Samworth, 2010, Lemma 1).

Proof of Theorem 5

Part (A) of Theorem 5 follows from part (A) of Theorem 12 and Proposition 2(c) of Cule and Samworth (2010). Part (B) and (C) of Theorem 5 follow from part (B) and (C) of Theorem 12.

Proof of Corollary 3

Suppose that

$$\sup_{x \in \text{int}(\text{dom}(\psi_0))} |\psi'_0(x)| = \infty,$$

but there exists $M > 0$ such that $\sup_{x \in \text{int}(\text{dom}(\tilde{\psi}_n))} |\tilde{\psi}'_n(x)| < M$ for all sufficiently large n . We claim that the probability of such an event is 0. It is straightforward to see that there exists $y \in \text{int}(\text{dom}(\psi_0))$ such that $|\psi'_0(y)| > 2M$. One can also choose y such that ψ'_0 is continuous at y . Therefore, pointwise convergence of $\tilde{\psi}'_n$ to ψ'_0 fails at y . Hence, the proof follows by Theorem 12. \square

Proof of Lemma 3

The proof follows from Lemma 6 because for $f_0 \in \mathcal{P}_0$,

$$\tilde{\mathcal{I}}_{f_0}(\eta) = \tilde{\mathcal{I}}_{f_0}^{sm}(\eta) = \mathcal{I}_{f_0}(\eta). \quad (6.1)$$

Proof of Theorem 6

The proof follows from Theorem 7. In this case, $\tilde{\psi}_0 = \tilde{\psi}_0^{sm} = \psi_0$ and (6.1) holds. As a result, the first terms on the right hand sides of (3.15) and (3.17) converge weakly to a centered Gaussian random variable with variance $\mathcal{I}_{f_0}(\eta)^{-1}$ by the central limit theorem. Also, since $\gamma_\eta = \gamma_\eta^{sm} = 1$, the second terms on the right hand sides of (3.15) and (3.17) vanish, which completes the proof.

Proof of Lemma 4

Note that each $\phi \in \mathcal{SC}_\theta$ can be written as $\psi(\cdot - \theta)$ for some $\psi \in \mathcal{SC}_0$, and by (3.7) and (1.4), we also have

$$\omega(\phi, F) = \Psi(\theta, \psi, F). \quad (6.2)$$

For $\psi \in \mathcal{SC}_0$ we calculate,

$$\begin{aligned} \Psi(0, \psi, F) &= \int_{-\infty}^0 \psi(x) dF(x) + \int_0^{\infty} \psi(x) dF(x) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= - \int_0^{\infty} \psi(-x) dF(-x) + \int_0^{\infty} \psi(x) dF(x) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= \int_0^{\infty} \psi(x) d(F(x) - F(-x)) - \int_{-\infty}^{\infty} e^{\psi(x)} dx. \end{aligned}$$

It also follows that

$$\int_0^{\infty} \psi(x) d(F(x) - F(-x)) = \int_{-\infty}^0 \psi(x) d(F(x) - F(-x)).$$

Therefore,

$$\Psi(0, \psi, F) = 2^{-1} \int_{-\infty}^{\infty} \psi(x) d(F(x) - F(-x)) - \int_{-\infty}^{\infty} e^{\psi(x)} dx,$$

implying

$$\begin{aligned} \Psi(\theta, \psi, F) &= 2^{-1} \int_{-\infty}^{\infty} \psi(x) d(F(\theta + x) - F(\theta - x)) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= 2^{-1} \int_{-\infty}^{\infty} \psi(z - \theta) d(F(z) - F(2\theta - z)) - \int_{-\infty}^{\infty} e^{\psi(z)} dz \\ &= \int_{-\infty}^{\infty} \psi(z - \theta) dF_{\theta}^{sym}(z) - \int_{-\infty}^{\infty} e^{\psi(z)} dz \\ &= \Psi(\theta, \psi, F_{\theta}^{sym}). \end{aligned}$$

Thus (6.2) shows that for any $\phi \in \mathcal{SC}_\theta$,

$$\omega(\phi, F) = \omega(\phi, F_{\theta}^{sym})$$

Therefore,

$$\arg \max_{\phi \in \mathcal{SC}_\theta} \omega(\phi, F) = \arg \max_{\phi \in \mathcal{SC}_\theta} \omega(\phi, F_{\theta}^{sym}),$$

but the distribution function

$$F_{\theta}^{sym}(x) = 2^{-1} \left(F(x) + 1 - F(2\theta - x) \right)$$

satisfies Condition A and (3.20). Hence, applying Lemma 14, we obtain that

$$\arg \max_{\phi \in \mathcal{SC}_\theta} \omega(\phi, F_{\theta}^{sym}) = \arg \max_{\phi \in \mathcal{C}} \omega(\phi, F_{\theta}^{sym}),$$

which exists by Condition A, thus completing the proof. \square

Proof of Lemma 5

First note that it suffices to prove the current lemma when $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$. Suppose the strong consistency does not hold but $\bar{\theta}_n \rightarrow_p \theta_0$. Then given any subsequence of $\{\bar{\theta}_n\}_{n \geq 1}$, there exists a further subsequence $\{\bar{\theta}_{n_k}\}_{k \geq 1}$ such that $\bar{\theta}_{n_k} \rightarrow_{a.s.} \theta_0$ as $k \rightarrow \infty$. Therefore, along this subsequence $\{n_k\}_{k \geq 1}$, the L_1 distance between \tilde{g}_n and \tilde{f}_0 or \tilde{f}_0^{sym} (in case of $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$) approaches 0 almost surely. Hence, if $\|\tilde{g}_n - \tilde{g}_0\|_1$ or $\|\tilde{g}_n - \tilde{g}_0^{sm}\|_1$ has any convergent subsequence at all, the subsequence converges to 0 almost surely. In that case [Shorack \(2000\)](#) (Theorem 5.7, page 57) implies that the whole sequence converges in probability to 0. Therefore, in what follows, we assume that $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$.

For our first estimator \tilde{g}_n^{sym} , the proof is motivated by Theorem 4 of [Cule and Samworth \(2010\)](#), which states that if $\int \log f_0(x) dF_0(x) < \infty$, and Condition A holds,

$$\int_{-\infty}^{\infty} |\hat{h}_n(x) - \tilde{f}_0(x)| dx \rightarrow_{a.s.} 0. \quad (6.3)$$

Note that $f_0 \in \mathcal{P}_1$ is bounded, so $\int \log_+ f_0(x) dF_0(x) < \infty$ trivially holds, where $\log_+(y) = \max\{\log y, 0\}$. To prove the L_1 consistency of \tilde{g}_n^{sym} , we now exploit the symmetry of \tilde{f}_0 and \hat{h}_n about θ_0 and $\bar{\theta}_n$ respectively. Recalling the definition of \tilde{g}_n^{sym} from [\(2.3\)](#), we calculate that

$$\begin{aligned} & 2 \int_{-\infty}^{\infty} |\tilde{g}_n^{sym}(x) - \tilde{g}_0(x)| dx \\ &= 2 \int_{-\infty}^{\infty} |\tilde{g}_n^{sym}(x) - \tilde{f}_0(\theta_0 + x)| dx \\ &\leq \int_{-\infty}^{\infty} |\hat{h}_n(\bar{\theta}_n + x) - \tilde{f}_0(\bar{\theta}_n + x)| dx + \int_{-\infty}^{\infty} |\hat{h}_n(\bar{\theta}_n - x) - \tilde{f}_0(\bar{\theta}_n - x)| dx \\ &\quad + \int_{-\infty}^{\infty} |\tilde{f}_0(\bar{\theta}_n + x) - \tilde{f}_0(\theta_0 + x)| dx + \int_{-\infty}^{\infty} |\tilde{f}_0(\bar{\theta}_n - x) - \tilde{f}_0(\theta_0 - x)| dx, \end{aligned} \quad (6.4)$$

whose first two terms approach 0 almost surely by [\(6.3\)](#). Therefore, we only need to take care of the last two terms of [\(6.4\)](#). Since \tilde{f}_0 is log-concave, it is continuous almost everywhere on \mathbb{R} with respect to the Lebesgue measure. Therefore, $|\tilde{f}_0(\bar{\theta}_n + x) - \tilde{f}_0(\theta_0 + x)| \rightarrow_{a.s.} 0$ almost everywhere with respect to the Lebesgue measure. Application of Glick's Theorem (Theorem 2.6 [Devroye, 1987](#)) then ensures that the last two terms of [\(6.4\)](#) converge to 0 almost surely as well.

To establish the convergence of $(\tilde{g}_n^{sym})^{sm}$ to \tilde{g}_0^{sm} in L_1 , we will exploit the representation of $(\tilde{g}_n^{sym})^{sm}$ in terms of $(\hat{h}_n)^{sm}$ established in [\(2.7\)](#). Since \tilde{g}_0^{sm} and $(\tilde{g}_n^{sym})^{sm}$ are symmetric about θ_0 and $\bar{\theta}_n$ respectively, similar algebra as in

the case of $\tilde{g}_n = \tilde{g}_n^{sym}$ leads to

$$\begin{aligned}
 & 2 \int_{-\infty}^{\infty} |(\tilde{g}_n^{sym})^{sm}(x) - \tilde{g}_0^{sm}(x)| dx \\
 & \leq \int_{-\infty}^{\infty} |\widehat{h}_n^{sm}(\bar{\theta}_n + x) - \tilde{f}_0^{sm}(\bar{\theta}_n + x)| dx + \int_{-\infty}^{\infty} |\widehat{h}_n^{sm}(\bar{\theta}_n - x) - \tilde{f}_0^{sm}(\bar{\theta}_n - x)| dx \\
 & \quad + \int_{-\infty}^{\infty} |\tilde{f}_0^{sm}(\bar{\theta}_n + x) - \tilde{f}_0^{sm}(\theta_0 + x)| dx \\
 & \quad + \int_{-\infty}^{\infty} |\tilde{f}_0^{sm}(\bar{\theta}_n - x) - \tilde{f}_0^{sm}(\theta_0 - x)| dx.
 \end{aligned} \tag{6.5}$$

The first two terms can be handled by using Theorem 1 of [Chen and Samworth \(2013\)](#), which implies that when f_0 satisfies (3.10), we have

$$\int_{-\infty}^{\infty} |\widehat{h}_n^{sm}(x) - \tilde{f}_0^{sm}(x)| dx \rightarrow_{a.s.} 0. \tag{6.6}$$

Since \tilde{f}_0^{sm} is the convolution of a log-concave density and a Gaussian density, it is continuous almost everywhere on \mathbb{R} with respect to the Lebesgue measure. Therefore, imitating the proof for the case of $\tilde{g}_n = \tilde{g}_n^{sym}$, we can show that the last two terms of (6.5) are negligible.

To establish the L_1 consistency of the partial MLE estimator $\widehat{g}_{\bar{\theta}_n}$, we appeal to the projection theory developed in [Xu and Samworth \(2019\)](#). In this case $\widehat{g}_{\bar{\theta}_n}$ is the density of the projection of $\mathbb{F}_{n, X - \bar{\theta}_n}$, the empirical distribution function of the $X_i - \bar{\theta}_n$'s, onto the space of the distribution functions with density in $\mathcal{S}\mathcal{L}\mathcal{C}_0$. By Proposition 6 of [Xu and Samworth \(2019\)](#), if

$$d_W(\mathbb{F}_{n, X - \bar{\theta}_n}, G_0) \rightarrow_{a.s.} 0, \tag{6.7}$$

we have $\|\widehat{g}_{\bar{\theta}_n} - \tilde{g}_0\|_1 \rightarrow_{a.s.} 0$ as long as Condition A holds. Hence, we only need to show that $d_W(\mathbb{F}_{n, X - \bar{\theta}_n}, G_0) \rightarrow_{a.s.} 0$, for which, by Theorem 6.9 of [Villani \(2009\)](#), it suffices to show the following. First, we need to show that

$$\int_{-\infty}^{\infty} |x| d\mathbb{F}_{n, X - \bar{\theta}_n}(x) \rightarrow_{a.s.} \int_{-\infty}^{\infty} |x| dG_0(x), \tag{6.8}$$

and that $\mathbb{F}_{n, X - \bar{\theta}_n}$ converges to G_0 weakly with probability one. Since $\bar{\theta}_n$ is strongly consistent for θ_0 , and

$$\int_{-\infty}^{\infty} |x| d\mathbb{F}_{n, X - \bar{\theta}_n}(x) = \int_{-\infty}^{\infty} |x - \bar{\theta}_n| d\mathbb{F}_n(x), \tag{6.9}$$

for any $d > 0$, an application of Glivenko-Cantelli Theorem (for example, see Theorem 2.4.1 of [Van der Vaart and Wellner \(1996\)](#)) yields

$$\sup_{\bar{\theta}_n \in [\theta_0 - d, \theta_0 + d]} \left| \int_{-\infty}^{\infty} |x - \bar{\theta}_n| d(\mathbb{F}_n - F_0)(x) \right| \rightarrow_{a.s.} 0.$$

On the other hand, strong consistency of $\bar{\theta}_n$ implies $|x - \bar{\theta}_n| \leq |x - \theta_0| + 1$ with probability one for all sufficiently large n , where the latter is integrable. Therefore, the dominated convergence theorem leads to

$$\int_{-\infty}^{\infty} |x - \bar{\theta}_n| dF_0(x) \rightarrow_{a.s.} \int_{-\infty}^{\infty} |x - \theta_0| dF_0(x) = \int_{-\infty}^{\infty} |x| dG_0(x),$$

which proves (6.8).

Our next step is to prove the weak convergence of $\mathbb{F}_{n, X - \bar{\theta}_n}$ to G_0 . To this end, we note that

$$\mathbb{F}_{n, X - \bar{\theta}_n}(x) = \mathbb{F}_n(x + \bar{\theta}_n) = \int_{-\infty}^{\infty} 1_{(-\infty, x + \bar{\theta}_n]}(z) d\mathbb{F}_n(z), \quad (6.10)$$

which converges almost surely to

$$\int_{-\infty}^{\infty} 1_{(-\infty, x + \theta_0]}(z) dF_0(z) = G_0(x)$$

by an application of basic Glivenko-Cantelli Theorem (see Theorem 2.4.1 of [Van der Vaart and Wellner \(1996\)](#)), and the fact that $F_0(x + \bar{\theta}_n) \rightarrow_{a.s.} F_0(x + \theta_0)$ for all $x \in \mathbb{R}$. This completes the proof of strong L_1 consistency of $\hat{g}_{\bar{\theta}_n}$ for \tilde{g}_0 .

Finally, we consider the geometric mean estimator $\hat{f}_n^{geo, sym}$. Recall that in the course of showing the consistency of \tilde{g}_n^{sym} , we have shown that the right hand side of (6.4) converges to 0 almost surely, from which one can derive that

$$\int_{-\infty}^{\infty} |\hat{h}_n(\bar{\theta}_n \pm x) - \tilde{g}_0(x)| dx \rightarrow_{a.s.} 0. \quad (6.11)$$

Proposition 2(b) of [Cule and Samworth \(2010\)](#) shows that (6.11) leads to almost sure convergence of $\hat{h}_n(\bar{\theta}_n \pm x)$ to $\tilde{g}_0(x)$ almost everywhere on \mathbb{R} with respect to the Lebesgue measure. As a consequence, it follows that

$$\hat{f}_n^{geo, sym}(x) / C_n^{geo} \rightarrow_{a.s.} \left(\tilde{g}_0(x) \tilde{g}_0(-x) \right)^{1/2} = \tilde{g}_0(x) \quad a.e. \quad x.$$

Recall from (2.9) that

$$C_n^{geo} = \int_{-\infty}^{\infty} \sqrt{\hat{h}_n(\bar{\theta}_n + x) \hat{h}_n(\bar{\theta}_n - x)} dx.$$

From Scheffé's Lemma it follows that $C_n^{geo} \rightarrow_{a.s.} \int d\tilde{G}_0 = 1$. We have thus established that $\hat{f}_n^{geo, sym}$ converges pointwise to $\tilde{g}_0(x)$ almost surely. The desired L_1 consistency then follows from Proposition 2(c) of [Cule and Samworth \(2010\)](#). \square

Theorem 12

Theorem 12. Assume $f_0 \in \mathcal{P}_1$. Suppose \tilde{g}_n is one of the estimators of g_0 defined in Section 2.1, where $\bar{\theta}_n$ is a consistent estimator of θ_0 . Let $\{y_n\}_{n \geq 1}$ be any sequence of random variables converging to 0 in probability. Then for $\tilde{g}_n = \tilde{g}_n^{sym}, \hat{g}_{\bar{\theta}_n}, \tilde{g}_n^{geo,sym}$, or $\hat{h}_n(\bar{\theta}_n \pm \cdot)$, on any compact set $K \subset \text{int}(\text{supp}(g_0))$ we have,

$$(A) \quad \sup_{x \in K} |\tilde{g}_n(x + y_n) - \tilde{g}_0(x)| \rightarrow_p 0.$$

$$(B) \quad \sup_{x \in K} |\tilde{\psi}_n(x + y_n) - \tilde{\psi}_0(x)| \rightarrow_p 0.$$

(C) Suppose $x \in \text{int}(\text{dom}(\tilde{\psi}_0))$ is a continuity point of $\tilde{\psi}'_0$. Then

$$\tilde{\psi}'_n(x + y_n) \rightarrow_p \tilde{\psi}'_0(x).$$

Suppose f_0 also satisfies (3.10). Then for $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$ or $\hat{h}_n^{sm}(\bar{\theta}_n \pm \cdot)$, (A)-(C) hold for any compact set $K \subset \mathbb{R}$, with \tilde{g}_0 and $\tilde{\psi}_0$ replaced by \tilde{g}_0^{sm} and $\tilde{\psi}_0^{sm}$ respectively. If $\bar{\theta}_n$ is a strongly consistent estimator of θ_0 , and $y_n \rightarrow_{a.s.} 0$, then the “ \rightarrow_p ” in the above displays can be replaced by “ $\rightarrow_{a.s.}$ ”.

Proof. As in the proof of Lemma 5, one can show that it suffices to prove Theorem 12 when $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$, and $y_n \rightarrow_{a.s.} 0$. Hence, in what follows, we assume that $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$, and $y_n \rightarrow_{a.s.} 0$.

First note that by theorem 2.2 of Dümbgen *et al.* (2011),

$$\text{int}(\text{dom}(\tilde{\psi}_0)) = \text{int}(\text{supp}(g_0)).$$

Also, since $\tilde{g}_0^{sm} = e^{\tilde{\psi}_0^{sm}} > 0$ on \mathbb{R} , we find that $\text{int}(\text{dom}(\tilde{\psi}_0^{sm})) = \mathbb{R}$. Hence,

$$K \subset \text{int}(\text{dom}(\tilde{\psi}_0)) \subset \text{int}(\text{dom}(\tilde{\psi}_0^{sm})) = \mathbb{R}.$$

When \tilde{g}_n equals $\hat{g}_{\bar{\theta}_n}, \tilde{g}_n^{geo,sym}, \hat{h}_n(\bar{\theta}_n \pm \cdot)$, or $\hat{h}_n^{sm}(\bar{\theta}_n \pm \cdot)$, \tilde{g}_n is log-concave, which simplifies the proof of part A and B. We first consider the cases when $\tilde{g}_n = \hat{g}_{\bar{\theta}_n}, \tilde{g}_n^{geo,sym}$ or $\hat{h}_n(\bar{\theta}_n \pm \cdot)$. Now to prove part A, note that

$$\begin{aligned} \int_{-\infty}^{\infty} |\tilde{g}_n(x + y_n) - \tilde{g}_0(x)| dx &\leq \int_{-\infty}^{\infty} |\tilde{g}_n(x + y_n) - \tilde{g}_0(x + y_n)| dx \\ &\quad + \int_{-\infty}^{\infty} |\tilde{g}_0(x + y_n) - \tilde{g}_0(x)| dx, \end{aligned}$$

whose first term

$$\int_{-\infty}^{\infty} |\tilde{g}_n(x + y_n) - \tilde{g}_0(x + y_n)| dx = \int_{-\infty}^{\infty} |\tilde{g}_n(x) - \tilde{g}_0(x)| dx \rightarrow_{a.s.} 0,$$

which follows from either Lemma 5 or (6.11). Also, noting

$$\tilde{g}_0(x + y_n) \rightarrow_{a.s.} \tilde{g}_0(x) \quad a.e. \ x \text{ Lebesgue,}$$

and applying Glick's Theorem (Theorem 2.6 Devroye, 1987), we derive that

$$\int_{-\infty}^{\infty} |\tilde{g}_0(x + y_n) - \tilde{g}_0(x)| dx \rightarrow_{a.s.} 0.$$

Thus we obtain that

$$\int_{-\infty}^{\infty} |\tilde{g}_n(x + y_n) - \tilde{g}_0(x)| dx \rightarrow_{a.s.} 0,$$

which combined with Proposition 2(c) of Cule and Samworth (2010) yields that

$$\tilde{g}_n(x + y_n) \rightarrow_{a.s.} \tilde{g}_0(x) \quad a.e. \ x$$

with respect to Lebesgue measure. As a consequence,

$$\tilde{\psi}_n(x + y_n) = \log(\tilde{g}_n(x + y_n)) \rightarrow_{a.s.} \tilde{\psi}_0(x) \quad a.e. \ x \text{ Lebesgue.}$$

Since $\tilde{\psi}_n$ is concave for our choices of \tilde{g}_n , Theorem 10.8 of Rockafellar (1970) indicates that pointwise convergence on \mathbb{R} translates to uniform convergence on all compact sets inside $\text{int}(\text{dom}(\tilde{\psi}_0))$, leading to

$$\sup_{x \in K} |\tilde{\psi}_n(x + y_n) - \tilde{\psi}_0(x)| \rightarrow_{a.s.} 0.$$

As a corollary,

$$\sup_{x \in K} |\tilde{g}_n(x + y_n) - \tilde{g}_0(x)| \rightarrow_{a.s.} 0.$$

Thus we have established part A and part B of Theorem 12 for $\tilde{g}_n = \hat{g}_{\bar{\theta}_n}$, $\tilde{g}_n^{geo, sym}$ and $\hat{h}_n(\bar{\theta}_n \pm \cdot)$. Now consider the case when $\tilde{g}_n = \hat{h}_n^{sm}(\bar{\theta}_n \pm x)$. Note that (6.6) and continuity of \tilde{g}_0^{sm} imply that

$$\int_{-\infty}^{\infty} |\hat{h}_n^{sm}(\bar{\theta}_n \pm x) - \tilde{g}_0^{sm}(x)| dx \rightarrow_{a.s.} 0.$$

Therefore, following the same arguments as in the case of $\tilde{g}_n = \hat{g}_{\bar{\theta}_n}$, $\tilde{g}_n^{geo, sym}$ and $\hat{h}_n(\bar{\theta}_n \pm \cdot)$, we can prove part A and part B for $\tilde{g}_n = \hat{h}_n^{sm}(\bar{\theta}_n \pm \cdot)$.

Now we focus on proving part A when \tilde{g}_n is the mixture density \tilde{g}_n^{sym} . From the definition of \tilde{g}_n^{sym} in (2.5) we obtain that

$$\begin{aligned} & 2 \sup_{x \in K} |\tilde{g}_n^{sym}(x + y_n) - \tilde{g}_0(x)| \\ & \leq \sup_{x \in K} |\hat{h}_n(\bar{\theta}_n + x + y_n) - \tilde{g}_0(x)| + \sup_{x \in K} |\hat{h}_n(\bar{\theta}_n - x + y_n) - \tilde{g}_0(x)|. \end{aligned}$$

Since we have already proved part A for $\hat{h}_n(\bar{\theta}_n \pm \cdot)$, it is not hard to see that both terms on the right hand side of the above display converge to 0 almost

surely, which proves part A for \tilde{g}_n^{sym} . Part A follows for $(\tilde{g}_n^{sym})^{sm}$ in a similar way noting that (2.7) implies

$$\begin{aligned} & 2 \sup_{x \in K} |(\tilde{g}_n^{sym})^{sm}(x + y_n) - \tilde{g}_0^{sm}(x)| \\ & \leq \sup_{x \in K} |\hat{h}_n^{sm}(\bar{\theta}_n + x + y_n) - \tilde{g}_0^{sm}(x)| + \sup_{x \in K} |\hat{h}_n^{sm}(\bar{\theta}_n - x + y_n) - \tilde{g}_0^{sm}(x)|. \end{aligned}$$

Now we prove part B of Theorem 12 when $\tilde{g}_n = \tilde{g}_n^{sym}$. To this end, note that we can write

$$\begin{aligned} \sup_{x \in K} (\tilde{\psi}_n^{sym}(x + y_n) - \tilde{\psi}_0(x)) &= \sup_{x \in K} \log \left(\frac{\tilde{g}_n^{sym}(x + y_n) - \tilde{g}_0(x)}{\tilde{g}_0(x)} + 1 \right) \\ &\leq \sup_{x \in K} \frac{\tilde{g}_n^{sym}(x + y_n) - \tilde{g}_0(x)}{\tilde{g}_0(x)}, \end{aligned}$$

because $\log(1 + x) \leq x$ for any $x > -1$. Similarly we can show that

$$\sup_{x \in K} (\tilde{\psi}_0(x) - \tilde{\psi}_n^{sym}(x + y_n)) \leq \sup_{x \in K} \frac{\tilde{g}_0(x) - \tilde{g}_n^{sym}(x + y_n)}{\tilde{g}_n^{sym}(x + y_n)},$$

leading to

$$\sup_{x \in K} \left| \tilde{\psi}_n^{sym}(x + y_n) - \tilde{\psi}_0(x) \right| \leq \frac{\sup_{x \in K} |\tilde{g}_n^{sym}(x + y_n) - \tilde{g}_0(x)|}{\min \left(\inf_{x \in K} \tilde{g}_n^{sym}(x + y_n), \inf_{x \in K} \tilde{g}_0(x) \right)},$$

whose numerator converges to 0 almost surely by part A of Theorem 12. Thus, to prove part B for $\tilde{\psi}_n^{sym}$, we only need to show that the denominator of the term on the right hand side of last display is bounded away from 0. To this end, notice that since \tilde{g}_0 is unimodal, and \tilde{g}_n^{sym} is a convex combination of two unimodal densities, they attain their infimum over K at either of the endpoints. By part A, it also follows that as $n \rightarrow \infty$, the respective minimum values approach $\inf_K \tilde{g}_0$, which is bounded away from 0 as K is a closed subset of $\text{int}(\text{dom}(\tilde{\psi}_0))$. Thus part B is proved for $\tilde{\psi}_n^{sym}$.

One can prove part B for $(\tilde{g}_n^{sym})^{sm}$ in a similar fashion because \tilde{g}_0^{sm} is unimodal, and $(\tilde{g}_n^{sym})^{sm}$ is a convex combination of two unimodal densities.

Now we can proceed to prove part C of the Theorem 12. First, we consider the log-concave density estimators $\hat{g}_{\bar{\theta}_n}$, $\tilde{g}_n^{geo, sym}$, and $\hat{h}_n(\bar{\theta}_n \pm \cdot)$, all of which are strongly L_1 consistent for \tilde{g}_0 . Note that if $\tilde{\psi}'_0$ is continuous at some $x \in \text{int}(\text{dom}(\tilde{\psi}_0))$, then there exists a neighborhood $[x - \xi, x + \xi]$ where $\tilde{\psi}_0$ is continuous. It follows that for sufficiently large n , $x + y_n \in [x - \xi, x + \xi]$ with probability 1. Since $\tilde{\psi}_n$ is concave, and converges to $\tilde{\psi}_0$ uniformly over K by part B, Lemma 3.10 of Seijo and Sen (2011) entails that

$$\sup_{t \in [x - \xi, x + \xi]} |\tilde{\psi}'_n(t \pm) - \tilde{\psi}'_0(t)| \rightarrow_{a.s.} 0, \quad (6.12)$$

from which we obtain that

$$|\tilde{\psi}'_n(x + y_n) - \tilde{\psi}'_0(x + y_n)| \rightarrow_{a.s.} 0,$$

which results in

$$|\tilde{\psi}'_n(x + y_n) - \tilde{\psi}'_0(x)| \rightarrow_{a.s.} 0 \quad (6.13)$$

because $\tilde{\psi}'_0$ is continuous at x , thereby proving part C for $\tilde{g}_n = \widehat{g}_{\bar{\theta}_n}$, $\tilde{g}_n^{geo, sym}$, and $\widehat{h}_n(\bar{\theta}_n \pm \cdot)$. For $\tilde{g}_n = \widehat{h}_n^{sm}(\bar{\theta}_n \pm x)$, part C follows similarly. Therefore, we skip the proof here. To prove part C for the non-logconcave estimators, \tilde{g}_n^{sym} and $(\tilde{g}_n^{sym})^{sm}$, we have to exploit their connection to \widehat{h}_n and \widehat{h}_n^{sm} , respectively. Beginning with \tilde{g}_n^{sym} , we observe that

$$(\tilde{\psi}_n^{sym})'(x) = \varrho_n(x) \left((\log \widehat{h}_n)'(\bar{\theta}_n + x) \right) - (1 - \varrho_n(x)) \left((\log \widehat{h}_n)'(\bar{\theta}_n - x) \right) \quad (6.14)$$

where $\varrho_n(x) = \widehat{h}_n(\bar{\theta}_n + x) / 2\tilde{g}_n^{sym}(x) \leq 1$. Thus

$$\begin{aligned} |(\tilde{\psi}_n^{sym})'(x + y_n) - \tilde{\psi}'_0(x)| &\leq \varrho_n(x + y_n) |(\log \widehat{h}_n)'(\bar{\theta}_n + x + y_n) - \tilde{\psi}'_0(x)| \\ &\quad + (1 - \varrho_n(x + y_n)) |(\log \widehat{h}_n)'(\bar{\theta}_n - x - y_n) - \tilde{\psi}'_0(-x)|. \end{aligned}$$

Since ϱ_n is uniformly bounded above by 1, (6.13) (applied on $\log \widehat{h}_n(\bar{\theta}_n \pm \cdot)$) completes the proof of part C for $\tilde{\psi}_n^{sym}$. Analogously using (6.14), for $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$, one can show that

$$((\tilde{\psi}_n^{sym})^{sm})'(x) = \varrho_n^{sm}(x) \left((\log \widehat{h}_n^{sm})'(\bar{\theta}_n + x) \right) - (1 - \varrho_n^{sm}(x)) \left((\log \widehat{h}_n^{sm})'(\bar{\theta}_n - x) \right) \quad (6.15)$$

for some $\varrho_n^{sm}(x) < 1$. Therefore, following the same steps as in the case of \tilde{g}_n^{sym} , we can establish part C for $(\tilde{g}_n^{sym})^{sm}$. \square

Proof of Lemma 6

As in the proof of Lemma 5, one can show that it suffices to prove Lemma 6 when $\bar{\theta}_n$ is a strongly consistent estimator of θ_0 .

We will show that $\widehat{\mathcal{I}}_n(\eta) \rightarrow_{a.s.} \tilde{\mathcal{I}}_{f_0}(\eta)$ and $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$ when \tilde{g}_n equals \tilde{g}_n^{sym} , $\widehat{g}_{\bar{\theta}_n}$ or $\tilde{g}_n^{geo, sym}$. The proof in the case of $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$ follows in a similar way.

We will prove the consistency of $\widehat{\mathcal{I}}_n(\eta)$ first. Since Lemma 5 indicates that $\|\tilde{g}_n - \tilde{g}_0\|_1 \rightarrow_{a.s.} 0$, the corresponding distribution functions satisfy $\|\tilde{G}_n - \tilde{G}_0(\eta)\|_\infty \rightarrow_{a.s.} 0$, which combined with the continuity of \tilde{G}_0 entails that

$$-\xi_n = \tilde{G}_n^{-1}(\eta) \rightarrow_{a.s.} \tilde{G}_0^{-1}(\eta) = -\xi_0; \quad (6.16)$$

and

$$\tilde{G}_n^{-1}(1 - \eta) \rightarrow_{a.s.} \tilde{G}_0^{-1}(1 - \eta) = \xi_0. \quad (6.17)$$

Combined with strong consistency of $\bar{\theta}_n$, the above leads to

$$\bar{\theta}_n + \xi_n = \bar{\theta}_n + \tilde{G}_n^{-1}(1 - \eta) \rightarrow_{a.s.} \theta_0 + \xi_0 = \tilde{F}_0^{-1}(1 - \eta), \quad (6.18)$$

and

$$\bar{\theta}_n - \xi_n = \bar{\theta}_n + \tilde{G}_n^{-1}(\eta) \rightarrow_{a.s.} \theta_0 - \xi_0 = \tilde{F}_0^{-1}(\eta). \quad (6.19)$$

Observe that

$$\begin{aligned} |\hat{\mathcal{I}}_n(\eta) - \tilde{\mathcal{I}}_{f_0}(\eta)| &\leq \left| \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n)^2 d(\mathbb{F}_n - F_0)(x) \right| \\ &\quad + \left| \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n)^2 dF_0(x) - \tilde{\mathcal{I}}_{f_0}(\eta) \right|. \end{aligned} \quad (6.20)$$

Let us consider the first term on the right hand side of (6.20) first. To this end, note that Lemma 11 combined with (6.16) and (6.17) imply that

$$\limsup_n \sup_{z \in [-\xi_n, \xi_n]} |\tilde{\psi}'_n(z)| \leq C_{\xi_0} \quad a.s., \quad (6.21)$$

where $C(-\xi_0, \xi_0) > 0$ is a finite constant depending on the truncation parameter η via $\xi_0 = \tilde{G}_0^{-1}(1 - \eta)$. For the sake of simplicity, we denote $C(-\xi_0, \xi_0)$ by M_η . Suppose \mathcal{U}_η is the class of non-increasing functions bounded by M_η , i.e.,

$$\mathcal{U}_\eta = \left\{ h : \mathbb{R} \mapsto [-M_\eta, M_\eta] \mid h \text{ is non-increasing} \right\},$$

and denote by \mathcal{F}_η the class

$$\begin{aligned} \mathcal{F}_\eta = \left\{ h^2 : \mathbb{R} \mapsto [0, M_\eta^2] \mid h(x) = u(x)1_{[r_1, r_2]}(x), \right. \\ \left. u \in \mathcal{U}_\eta, [r_1, r_2] \subset \left(\tilde{F}_0^{-1}(\eta/2), \tilde{F}_0^{-1}(1 - \eta/2) \right) \right\}. \end{aligned}$$

Theorem 2.7.5 of Van der Vaart and Wellner (1996) (pp. 159) indicates that for any $\epsilon > 0$, the bracketing entropy

$$\log N_{[\cdot]}(\epsilon, \mathcal{U}_\eta, L_2(P_0)) \lesssim \epsilon^{-1}. \quad (6.22)$$

Also, by Example 2.5.4 of Van der Vaart and Wellner (1996), the class \mathcal{F}_I of all indicator functions of the form $1_{[z_1, z_2]}$, where $z_1 \leq z_2$ with $z_1, z_2 \in \mathbb{R}$, satisfies

$$\log N_{[\cdot]}(\epsilon, \mathcal{F}_I, L_2(P_0)) \lesssim 2/\epsilon^2, \quad (6.23)$$

implying that

$$\log N_{[\cdot]}(\epsilon, \mathcal{F}_\eta, L_2(P_0)) \lesssim \epsilon^{-1}.$$

Since $\tilde{\psi}'_n$ is non-increasing, (6.21) implies that $(\tilde{\psi}'_n)^2(\cdot - \bar{\theta}_n)$ restricted to $[\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n]$ belongs to \mathcal{F}_η . Therefore, by Glivenko-Cantelli Theorem (see Theorem 2.4.1 of Van der Vaart and Wellner (1996)),

$$\int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n)^2 d(\mathbb{F}_n - F_0)(x) \rightarrow_{a.s.} 0. \quad (6.24)$$

Now we claim that the second term in (6.20) also approaches 0 almost surely. This follows by Theorem 12(C) and the dominated convergence theorem since $\tilde{\psi}'_n$ is bounded on $[-\xi_n, \xi_n]$, completing the proof of $\widehat{\mathcal{I}}_n(\eta) \rightarrow_{a.s.} \widetilde{\mathcal{I}}_{f_0}(\eta)$.

Our next aim is to prove the consistency of $\tilde{\theta}_n$. Observe that from (2.11) it follows that

$$\begin{aligned} |\tilde{\theta}_n - \bar{\theta}_n| &\leq \widehat{\mathcal{I}}_n(\eta)^{-1} \left| \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n) d(\mathbb{F}_n - F_0)(x) \right| \\ &\quad + \widehat{\mathcal{I}}_n(\eta)^{-1} \left| \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n) dF_0(x) \right|. \end{aligned}$$

Note that $\widehat{\mathcal{I}}_n(\eta) \rightarrow_{a.s.} \widetilde{\mathcal{I}}_{f_0}(\eta)$ by the first part of the current lemma. The proof of

$$\widehat{\mathcal{I}}_n(\eta)^{-1} \left| \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n) d(\mathbb{F}_n - F_0)(x) \right| \rightarrow_{a.s.} 0$$

is very similar to the proof of (6.24), and follows by Glivenko-Cantelli Theorem (see Theorem 2.4.1 of Van der Vaart and Wellner (1996)) noting that $\tilde{\psi}'_n(\cdot - \bar{\theta}_n)$ restricted to $[-\xi_n, \xi_n]$ is a member of \mathcal{U}_η , which has finite bracketing entropy by (6.23).

Since $\tilde{\psi}'_n$ is bounded on $[-\xi_n, \xi_n]$, (6.18), (6.19), and Theorem 12(C) combined with the dominated convergence theorem further entail that

$$\int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \tilde{\psi}'_n(x - \bar{\theta}_n) dF_0(x) \rightarrow_{a.s.} \int_{\tilde{G}_0^{-1}(\eta)}^{\tilde{G}_0^{-1}(1-\eta)} \tilde{\psi}'_0(x) g_0(x) dx,$$

which is 0 because $\tilde{\psi}'_0$ is an odd function while g_0 is an even function. Therefore, strong consistency of $\tilde{\theta}_n$ follows from that of $\bar{\theta}_n$. \square

Proof of Theorem 7

As in the proof of Lemma 5, one can show that it suffices to prove Theorem 7 when $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$. Therefore, in what follows, we assume that $\bar{\theta}_n \rightarrow_{a.s.} \theta_0$.

First we consider the case when \tilde{g}_n equals \tilde{g}_n^{sym} , $\widehat{g}_{\bar{\theta}_n}$ or $\tilde{g}_n^{geo,sym}$, so that by Lemma 5, the density estimator \tilde{g}_n converges to \tilde{g}_0 in L_1 almost surely. From (2.11) we deduce that

$$-(\tilde{\theta}_n - \bar{\theta}_n) = \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \frac{\tilde{\psi}'_n(x - \bar{\theta}_n)}{\widehat{\mathcal{I}}_n(\eta)} d\mathbb{F}_n(x) = \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_n(z)}{\widehat{\mathcal{I}}_n(\eta)} d\mathbb{F}_n(z + \bar{\theta}_n).$$

Denoting $\tilde{\delta}_n = \theta_0 - \bar{\theta}_n$, we observe that the expression in the above display can also be written as

$$\begin{aligned} & \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_n(z) - \tilde{\psi}'_0(z - \tilde{\delta}_n)}{\widehat{\mathcal{I}}_n(\eta)} d(\mathbb{F}_n(z + \bar{\theta}_n) - F_0(z + \bar{\theta}_n)) \\ & + \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_n(z)}{\widehat{\mathcal{I}}_n(\eta)} \left(f_0(z + \bar{\theta}_n) - g_0(z) \right) dz + \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_n(z) - \tilde{\psi}'_0(z)}{\widehat{\mathcal{I}}_n(\eta)} g_0(z) dz \\ & + \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_0(z)}{\widehat{\mathcal{I}}_n(\eta)} g_0(z) dz + \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_0(z - \tilde{\delta}_n)}{\widehat{\mathcal{I}}_n(\eta)} d(\mathbb{F}_n(z + \bar{\theta}_n) - F_0(z + \bar{\theta}_n)) \\ & = T_{1n} + T_{2n} + T_{3n} + T_{4n} + T_{5n}. \end{aligned}$$

Observe that T_{3n} and T_{4n} vanish since $\tilde{\psi}'_n$ and $\tilde{\psi}'_0$ are odd functions while \tilde{f}_0 is an even function. Now note that to prove (3.15), it suffices to show that $\sqrt{n}T_{1n} = o_p(1)$,

$$\frac{T_{2n}}{\tilde{\delta}_n} \rightarrow_{a.s.} \int_{-\xi_0}^{\xi_0} \frac{\tilde{\psi}'_0(z)\psi'_0(z)}{\widehat{\mathcal{I}}_{f_0}(\eta)} g_0(z) dz, \quad (6.25)$$

and that

$$\sqrt{n}T_{5n} = \int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{\tilde{\psi}'_0(x - \theta_0)}{\widehat{\mathcal{I}}_{f_0}(\eta)} d\mathbb{Z}_n(x) + o_p(1).$$

When $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$, (3.17) can also be proved in a similar way, and hence we skip the proof.

We will first show that $\sqrt{n}T_{1n} = o_p(1)$. Recall that in Section 1.1 we denoted the empirical process $\sqrt{n}(\mathbb{F}_n - F_0)$ by \mathbb{Z}_n . Denote by h_n the function

$$h_n(x) = (\tilde{\psi}'_n(x - \bar{\theta}_n) - \tilde{\phi}'_0(x))1_{[\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n]}(x),$$

where $\tilde{\phi}'_0(x)$ was previously defined as $\psi'_0(x - \theta_0)$. It follows that

$$\begin{aligned} \sqrt{n}T_{1n} &= \sqrt{n} \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \frac{\tilde{\psi}'_n(x - \bar{\theta}_n) - \tilde{\phi}'_0(x)}{\widehat{\mathcal{I}}_n(\eta)} d(\mathbb{F}_n - F_0)(x) \\ &= \int_{-\infty}^{\infty} \frac{h_n(x)}{\widehat{\mathcal{I}}_n(\eta)} d\mathbb{Z}_n(x). \end{aligned} \quad (6.26)$$

Lemma 11 combined with (6.16) and (6.17) imply that

$$\limsup_n \sup_{z \in [-\xi_n, \xi_n]} |\tilde{\psi}'_n(z)| \leq C_{\xi_0} \quad a.s.,$$

where C_{ξ_0} is a positive constant depending on ξ_0 . It is not hard to see that since $\tilde{\phi}'_0$ is monotone, $|\tilde{\phi}'_0|$ attains its maxima over $[\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n]$ at either of the endpoints. Though $[\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n]$ is random, (6.17) and (6.18) indicate that with probability one, this interval is a subset of $[F_0^{-1}(\eta/2), F_0^{-1}(1 - \eta/2)]$ for all sufficiently large n . Hence, it follows that

$$\limsup_n \sup_{x \in [\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n]} |\tilde{\psi}'_0(x)| < M_\eta \quad a.s. \quad (6.27)$$

for some $M_\eta > 0$, which can be chosen in such a way so that $C_{\xi_0} \leq M_\eta$. Therefore, we obtain that

$$\limsup_n \sup_{x \in [\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n]} |h_n(x)| < M_\eta \quad a.s. \quad (6.28)$$

Now define the class \mathcal{H}_η by

$$\mathcal{H}_\eta = \left\{ h : \mathbb{R} \mapsto [-M_\eta, M_\eta] \mid \begin{aligned} &h(x) = (u(x) - \tilde{\phi}'_0(x))1_{[r_1, r_2]}(x), \\ &u \in \mathcal{U}_\eta, [r_1, r_2] \subset \left(\tilde{F}_0^{-1}(\eta/2), \tilde{F}_0^{-1}(1 - \eta/2) \right) \end{aligned} \right\},$$

where \mathcal{U}_η is the class of non-increasing functions bounded by $M_\eta > 0$, i.e.

$$\mathcal{U}_\eta = \left\{ h : \mathbb{R} \mapsto [-M_\eta, M_\eta] \mid h \text{ is non-increasing} \right\}.$$

Observe that (6.18), (6.19), and (6.28) imply $h_n \in \mathcal{H}_\eta$ almost surely for all sufficiently large n .

From (6.22) and (6.23) we derive that

$$\log N_{[\]}(\epsilon, \mathcal{H}_\eta, L_2(P_0)) \lesssim \frac{1}{\epsilon}.$$

Therefore, we conclude that the bracketing integral

$$\mathcal{J}_{[\]}(1, \mathcal{H}_\eta, L_2(P_0)) = \int_0^1 \sqrt{1 + \log N_{[\]}(M_\eta \epsilon, \mathcal{H}_\eta, L_2(P_0))} d\epsilon \lesssim \int_0^1 \epsilon^{-1/2} d\epsilon,$$

which is finite, thereby establishing that \mathcal{H}_η is P_0 -Donsker. Also, notice that $|h_n|$ is bounded above by M_η , and Theorem 12(C) indicates that h_n converges to 0 at the continuity points of $\tilde{\phi}'_0$. Since $\tilde{\phi}_0$ is concave, $\tilde{\phi}'_0$ is continuous almost everywhere with respect to Lebesgue measure on $\text{dom}(\phi_0)$. Hence, the dominated convergence theorem yields

$$\int_{-\infty}^{\infty} h_n(x)^2 dF_0(x) \rightarrow_{a.s.} 0.$$

Since \mathcal{H}_η is Donsker, Theorem 2.1 of van der Vaart and Wellner (2007) then implies that

$$\int_{-\infty}^{\infty} h_n(x) d\mathbb{Z}_n(x) = o_p(1).$$

Finally, an application of Lemma 6 leads to $\hat{\mathcal{I}}_n(\eta) \rightarrow_{a.s.} \mathcal{I}_{f_0}(\eta)$, and thus from (6.26) we conclude that $\sqrt{n}T_{1n} = o_p(1)$.

Now we turn our attention to T_{2n} . Here we aim to prove (6.25). Observe that $T_{2n}/\tilde{\delta}_n$ can be written as

$$\begin{aligned} \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_n(z)}{\tilde{\mathcal{I}}_n(\eta)} \frac{\left(g_0(z - \tilde{\delta}_n) - g_0(z)\right)}{\tilde{\delta}_n} dz &= \int_{-\xi_n}^{\xi_n} \frac{\tilde{\psi}'_n(z)}{\tilde{\mathcal{I}}_n(\eta)} \frac{\int_z^{z-\tilde{\delta}_n} g'_0(t) dt}{\tilde{\delta}_n} dz \\ &= \int_{-\xi_n - \tilde{\delta}_n}^{\xi_n} g'_0(t) \frac{\int_t^{t+\tilde{\delta}_n} \tilde{\psi}'_n(z) dz}{\tilde{\delta}_n \tilde{\mathcal{I}}_n(\eta)} dt, \end{aligned} \quad (6.29)$$

where the equality follows by Fubini's theorem since g_0 is absolutely continuous. The concavity of $\tilde{\psi}_n$ implies that $\tilde{\psi}'_n$ is non-increasing. Hence, for any $t \in \text{int}(\text{dom}(\tilde{\psi}_0))$, we have,

$$\min \left\{ \tilde{\psi}'_n(t), \tilde{\psi}'_n(t + \tilde{\delta}_n) \right\} \leq \frac{\int_t^{t+\tilde{\delta}_n} \tilde{\psi}'_n(z) dz}{\tilde{\delta}_n} \leq \max \left\{ \tilde{\psi}'_n(t), \tilde{\psi}'_n(t + \tilde{\delta}_n) \right\}. \quad (6.30)$$

Therefore, if $\tilde{\psi}'_0$ is continuous at $t \in \text{int}(\text{dom}(\tilde{\psi}_0))$, Theorem 12(C) leads to

$$\frac{\int_t^{t+\tilde{\delta}_n} \tilde{\psi}'_n(z) dz}{\tilde{\delta}_n} \rightarrow_{a.s.} \tilde{\psi}'_0(t) \quad \text{as } n \rightarrow \infty.$$

Since $\tilde{\psi}_0$ is concave, $\tilde{\psi}'_0$ is continuous almost everywhere with respect to Lebesgue measure on $\text{dom}(\tilde{\psi}_0)$. Since $\tilde{\psi}'_n$ is non-increasing, (6.30) further entails that

$$\sup_{t \in [-\xi_n - \tilde{\delta}_n, \xi_n]} \frac{\left| \int_t^{t+\tilde{\delta}_n} \tilde{\psi}'_n(z) dz \right|}{|\tilde{\delta}_n|} \leq \tilde{\psi}'_n(-\xi_n - |\tilde{\delta}_n|),$$

which can be bounded above by $\tilde{\psi}'_0(\tilde{G}_0^{-1}(\eta/2))$ using (6.16) and the fact that $\tilde{\delta}_n \rightarrow_{a.s.} 0$. Hence, we deduce that

$$|g'_0(t)| \frac{\left| \int_t^{t+\tilde{\delta}_n} \tilde{\psi}'_n(z) dz \right|}{|\tilde{\delta}_n|} 1_{[-\xi_n - \tilde{\delta}_n, \xi_n]}(z) \leq |g'_0(t)| \left(\tilde{\psi}'_0(\tilde{G}_0^{-1}(\eta/2)) \right).$$

Note that the symmetry of g_0 about 0 indicates that

$$\int_{-\infty}^{\infty} |g'_0(t)| dt = 2 \int_{-\infty}^0 g'_0(t) dt = 2g(0),$$

which is finite because of (3.6). Therefore, (6.29), Lemma 6, and the dominated convergence theorem yield

$$\frac{T_{2n}}{\bar{\delta}_n} \rightarrow_{a.s.} \int_{-\xi_0}^{\xi_0} \frac{\tilde{\psi}'_0(z)g'_0(z)}{\tilde{\mathcal{I}}_{f_0}(\eta)} dz,$$

which completes the proof of (6.25) since $g'_0(z) = \psi'_0(z)g_0(z)$.

Now it only remains to show that the last term T_{5n} converges weakly to a Gaussian random variable with variance $\tilde{\mathcal{I}}_{f_0}^{-1}$. Observe that a change of variable leads to

$$\begin{aligned} \sqrt{n}T_{5n} &= \sqrt{n} \int_{\bar{\theta}_n - \xi_n}^{\bar{\theta}_n + \xi_n} \frac{\tilde{\psi}'_0(x - \theta_0)}{\hat{\mathcal{I}}_n(\eta)} d(\mathbb{F}_n - F_0)(x) \\ &= \int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{\tilde{\psi}'_0(x - \theta_0)}{\hat{\mathcal{I}}_n(\eta)} dZ_n(x) \\ &\quad + \int_{-\infty}^{\infty} 1_{C_n}(x) \frac{\tilde{\psi}'_0(x - \theta_0)}{\hat{\mathcal{I}}_n(\eta)} dZ_n(x), \end{aligned} \quad (6.31)$$

where

$$C_n = [\bar{\theta}_n - \xi_n, \bar{\theta}_n + \xi_n] \setminus [\theta_0 - \xi_0, \theta_0 + \xi_0].$$

From Lemma 6 and Slutsky's theorem it follows that

$$\int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{\tilde{\psi}'_0(x - \theta_0)}{\hat{\mathcal{I}}_n(\eta)} dZ_n(x) = \int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{\tilde{\psi}'_0(x - \theta_0)}{\tilde{\mathcal{I}}_{f_0}(\eta)} dZ_n(x) (1 + o_p(1)),$$

which equals

$$\int_{\theta_0 - \xi_0}^{\theta_0 + \xi_0} \frac{\tilde{\psi}'_0(x - \theta_0)}{\tilde{\mathcal{I}}_{f_0}(\eta)} dZ_n(x) + o_p(1),$$

because the central limit theorem and (3.3) imply that the first term on the above display is $O_p(1)$.

To deal with the second term, observe that the indicator function 1_{C_n} belongs to \mathcal{F}_1 , the class of all indicator functions of the form $1_{[z_1, z_2] \setminus [\theta_0 - \xi_0, \theta_0 + \xi_0]}$, where $z_1 \leq z_2$ with $z_1, z_2 \in \mathbb{R}$. Since the latter class is Donsker by (6.23), Theorem 2.1 of van der Vaart and Wellner (2007) entails that the second term on the right hand side of (6.31) is of order $o_p(1)$ provided the following holds:

$$\int_{-\infty}^{\infty} 1_{C_n}(x) \frac{\tilde{\psi}'_0(x - \theta_0)^2}{\hat{\mathcal{I}}_n(\eta)^2} dF_0(x) \rightarrow_{a.s.} 0.$$

To this end note that since $\tilde{\psi}'_0$ is non-increasing, (6.18), (6.19), (6.27), and Lemma 6 entail that for all sufficiently large n ,

$$\int_{-\infty}^{\infty} 1_{C_n}(x) \frac{\tilde{\psi}'_0(x - \theta_0)^2}{\hat{\mathcal{I}}_n(\eta)^2} dF_0(x) \leq \frac{\tilde{\psi}'_0(\tilde{G}_0^{-1}(\eta/2))^2}{\tilde{\mathcal{I}}_{f_0}^2(\eta)} |\bar{\theta}_n - \theta_0| \left(\sup_x f_0(x) \right) \quad a.s.$$

Since $f_0 \in \mathcal{P}_1$, from the definition of \mathcal{P}_1 in (3.6) and the strong consistency of $\bar{\theta}_n$ we obtain that the term on the right hand side of the last display converges to 0 almost surely. Thus we conclude that $\sqrt{n}T_{5n}$ is asymptotically distributed as a centered Gaussian random variable with variance $\tilde{\mathcal{I}}_{f_0}(\eta)^{-1}$, which completes the proof. \square

6.2. Proofs for the MLE

We first state and prove a useful theorem which plays a crucial role in proving Proposition 1 and Lemma 1. This theorem concerns the continuity of the function $L(\cdot; F) : \mathbb{R} \mapsto \mathbb{R}$ given by

$$L(\theta; F) = \sup_{\psi \in \mathcal{SC}_0} \Psi(\theta, \psi, F) \quad (6.32)$$

where F is a fixed distribution function, and Ψ is as defined in (1.4).

Theorem 13. *Suppose the distribution function F satisfies Condition A. Then the map $\theta \mapsto L(\theta; F)$ is continuous on \mathbb{R} , where the function $L(\theta; F)$ was defined in (6.32).*

Proof. Suppose $\theta_k \rightarrow \theta$ as $k \rightarrow \infty$. Observe that $\Psi(\theta, \psi, F)$ can also be written as

$$\Psi(\theta, \psi, F) = \Psi(0, \psi, F(\cdot + \theta)).$$

Hence, to prove Theorem 13, it suffices to show that as $k \rightarrow \infty$,

$$L(\theta_k; F) = \sup_{\psi \in \mathcal{SC}_0} \Psi(0, \psi, F(\cdot + \theta_k)) \rightarrow \sup_{\psi \in \mathcal{SC}_0} \Psi(0, \psi, F(\cdot + \theta)) = L(\theta, F).$$

Proposition 6 of Xu and Samworth (2019) implies that under Condition A, the convergence in the above display holds if the Wasserstein distance

$$d_W(F(\cdot + \theta_k), F(\cdot + \theta)) \rightarrow 0, \quad \text{as } k \rightarrow \infty. \quad (6.33)$$

Now by Theorem 6.9 of Villani (2009) (see also Theorem 7.12 of Villani, 2003), (6.33) follows if the followings hold as $k \rightarrow \infty$:

$$F(\cdot + \theta_k) \rightarrow_d F(\cdot + \theta), \quad (6.34)$$

and

$$\int_{-\infty}^{\infty} |x| dF(x + \theta_k) \rightarrow \int_{-\infty}^{\infty} |x| dF(x + \theta). \quad (6.35)$$

To prove (6.34), note that for any bounded continuous function h ,

$$\int_{-\infty}^{\infty} h(x - \theta_k) dF(x) \rightarrow \int_{-\infty}^{\infty} h(x - \theta) dF(x)$$

by the dominated convergence theorem since $\theta_k \rightarrow \theta$ as $k \rightarrow \infty$. For proving (6.35), first notice that F has finite first moment by Condition A. Therefore, another application of the dominated convergence yields that as $\theta_k \rightarrow \theta$,

$$\int_{-\infty}^{\infty} |x| dF(x + \theta_k) = \int_{-\infty}^{\infty} |x - \theta_k| dF(x) \rightarrow \int_{-\infty}^{\infty} |x| dF(x + \theta),$$

which, combined with (6.34), leads to (6.33), and completes the proof. \square

Proof of Proposition 1

Our first step is to show that $L(F)$ is finite. From the definition of Ψ in (1.4), it is not hard to see that

$$L(F) \leq \sup_{\psi \in \mathcal{C}} \left(\int_{-\infty}^{\infty} \psi(x) dF(x) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \right),$$

where \mathcal{C} denotes the set of all real-valued concave functions. Theorem 2.2 of Dümbgen *et al.* (2011) entails that under condition A, the term on the right hand side of the above display is finite. Therefore, $L(F) < \infty$ follows. To show that $L(F) > -\infty$, we show that the map $x \mapsto -|x| \in \mathcal{SC}_0$. Therefore, (1.4) and (1.6) lead to

$$L(F) > - \int_{-\infty}^{\infty} |x| dF(x) - \int_{-\infty}^{\infty} e^{-|x|} dx > -\infty,$$

which follows from Condition A. Therefore, we conclude that $L(F) \in \mathbb{R}$.

Now we have to show that there exist $\theta^*(F) \in \mathbb{R}$ and $\psi^*(F) \in \mathcal{SLC}_0$ such that

$$\Psi(\theta^*(F), \psi^*(F), F) = \sup_{\theta \in \mathbb{R}, \psi \in \mathcal{SLC}_0} \Psi(\theta, \psi, F) = \sup_{\theta \in \mathbb{R}} L(\theta; F) = L(F).$$

Now there exists a sequence $\{\theta_k\}_{k \geq 1}$ such that $L(\theta_k; F) \uparrow L(F)$ as $k \rightarrow \infty$. Suppose the sequence $\{\theta_k\}_{k \geq 1}$ is bounded. Then we can find a subsequence $\{\theta_{k_r}\}_{r \geq 1}$ converging to some $\theta' \in \mathbb{R}$. Since the map $L(\theta; F)$ is continuous in θ by Theorem 13, we also have

$$L(\theta'; F) = \lim_{r \rightarrow \infty} L(\theta_{k_r}; F) = L(F),$$

which implies that θ' is a maximizer of $L(\theta; F)$. The rest of the proof then follows from Proposition 4(iii) of Xu and Samworth (2019), which states that for each $\theta \in \mathbb{R}$, there exists a unique log-density ψ_θ , which maximizes $\Psi(\theta, \psi, F)$ in $\psi \in \mathcal{SC}_0$ when F satisfies condition A. Hence, $(\theta', \psi_{\theta'})$ is a candidate for $(\theta^*(F), \psi^*(F))$. Thus, to complete the proof, it remains to show that $\{\theta_k\}_{k \geq 1}$ is bounded. We will show that $\theta_k \rightarrow \pm\infty$ leads to $L(\theta_k; F) \rightarrow -\infty$, which contradicts the fact that $L(\theta_k; F) \rightarrow L(F) \in \mathbb{R}$, thus completing the proof.

Consider $\theta_k \rightarrow \pm\infty$. We have already mentioned that by Proposition 4(iii) of Xu and Samworth (2019), for each θ_k , there exists a log-density $\psi_{\theta_k} \in \mathcal{SC}_0$ such that

$$L(\theta_k; F) = \Psi(\theta, \psi_{\theta_k}, F),$$

which equals

$$\int_{-\infty}^{\infty} \psi_{\theta_k}(x - \theta) dF(x) - 1 \leq - \int_{-\infty}^{\infty} \log(2|x - \theta_k|) dF(x) - 1,$$

where the last inequality follows from Lemma 12. Now if $\theta_k \rightarrow \pm\infty$, using Fatou's Lemma, we derive that

$$\limsup_{k \rightarrow \infty} L(\theta_k; F) \leq - \int_{-\infty}^{\infty} \liminf_{k \rightarrow \infty} \left(\log|x - \theta_k| \right) dF(x) - \log 2 = -\infty,$$

which leads to the desired contradiction. \square

Proof of Lemma 1

The proof follows from (1.5) and Theorem 13 noting that \mathbb{F}_n satisfies Condition A. \square

Proof of Lemma 9

First we will show that if F satisfies (3.20) and $\psi \in \mathcal{SC}_0$, there exists $\psi_1 \in \mathcal{SC}_0$ such that

$$\Psi(\theta', \psi, F) \leq \Psi(\theta, \psi_1, F), \quad (6.36)$$

leading to

$$L(\theta'; F) \leq L(\theta; F),$$

which implies $\theta^*(F) = \theta$. Note that $G = F(\cdot + \theta)$ satisfies

$$G(x) + G(-x) = 1. \quad (6.37)$$

Using the symmetry of ψ about 0 and (6.37) in the fourth step we deduce that

$$\begin{aligned} & \Psi(\theta', \psi, F) \\ &= \int_{-\infty}^{\infty} \psi(x - \theta') dG(x - \theta) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= \int_{-\infty}^0 \psi(y + \theta - \theta') dG(y) + \int_0^{\infty} \psi(y + \theta - \theta') dG(y) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= \int_{-\infty}^0 \psi(y + \theta - \theta') dG(y) - \int_{-\infty}^0 \psi(-y + \theta - \theta') dG(-y) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= \int_{-\infty}^0 \psi(y + \theta - \theta') dG(y) + \int_{-\infty}^0 \psi(y - \theta + \theta') dG(y) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= 2 \int_{-\infty}^0 \psi_1(y) dG(y) - \int_{-\infty}^{\infty} e^{\psi(x)} dx, \end{aligned}$$

where

$$\psi_1(x) = 2^{-1} \left(\psi(x + \theta - \theta') + \psi(x - \theta + \theta') \right), \quad x \in \mathbb{R}.$$

Since $\psi_1 \in \mathcal{SC}_0$, we obtain that

$$\begin{aligned} \Psi(\theta', \psi, F) &= \int_{-\infty}^{\infty} \psi_1(x) dG(x) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= \int_{-\infty}^{\infty} \psi_1(x - \theta) dF(x) - \int_{-\infty}^{\infty} e^{\psi(x)} dx \\ &= \Psi(\theta, \psi_1, F) + \int_{-\infty}^{\infty} e^{\psi_1(x)} dx - \int_{-\infty}^{\infty} e^{\psi(x)} dx. \end{aligned}$$

Also, by the arithmetic mean-geometric mean inequality,

$$\int_{-\infty}^{\infty} e^{\psi_1(x)} dx \leq \int_{-\infty}^{\infty} 2^{-1} \left(e^{\psi(x+\theta-\theta')} + e^{\psi(x-\theta+\theta')} \right) dx = \int_{-\infty}^{\infty} e^{\psi(x)} dx,$$

where equality holds if and only if $\psi(x+\theta-\theta') = \psi(x-\theta+\theta')$ almost everywhere with respect to the Lebesgue measure, which only takes place when $\theta' = \theta$. Therefore, we have established the inequality in (6.36) with equality holding if and only if $\theta' = \theta$, which proves $\theta^*(F) = \theta$. Observe that

$$\Psi(\theta, \psi, F) = \int_{-\infty}^{\infty} \psi(x) dG(x) - \int_{-\infty}^{\infty} e^{\psi(x)} dx = \Psi(0, \psi, G).$$

Finally, Lemma 14 combined with (6.37) and (6.2) entail that

$$\arg \max_{\psi \in \mathcal{SC}_0} \Psi(0, \psi, G) = \arg \max_{\phi \in \mathcal{SC}_0} \omega(\phi, G) = \arg \max_{\phi \in \mathcal{C}} \omega(\phi, G),$$

which completes the proof of Lemma 9. \square

Proof of Lemma 8

Lemma 8 follows directly from Lemma 4. \square

Proof of Lemma 10

Suppose $\overline{J(F)} = \mathbb{R}$. Then $\theta^*(F) \in \overline{J(F)}$ by Proposition 1. Otherwise, since F is non-decreasing, $\overline{J(F)}$ takes one of the following three forms: $[a, b]$, $[a, \infty)$, or $(-\infty, b]$, where $a, b \in \mathbb{R}$. Suppose $\overline{J(F)} = [a, b]$. We will show that in this case $L(\theta; F)$ defined in (6.32) is non-decreasing in θ on $(-\infty, a]$, and non-increasing in θ on $[b, \infty)$. Since $\theta^*(F)$ equals $\arg \max_{\theta \in \mathbb{R}} L(\theta; F)$, the above implies $\theta^*(F) \in [a, b]$. To show that $L(\theta; F)$ is non-decreasing in θ on $(-\infty, a]$, we first note that for $\theta < \theta' \leq a$, and $\psi \in \mathcal{SC}_0$,

$$\int_a^b \psi(x - \theta) dF(x) \leq \int_a^b \psi(x - \theta') dF(x),$$

since ψ is non-increasing on $[0, \infty)$, and

$$0 \leq x - \theta' < x - \theta$$

for $x \geq a$. Therefore, from (6.32), it is not hard to see that $L(\theta; F) \leq L(\theta'; F)$. Similarly we can show that for $\theta > \theta' \geq b$,

$$\int_a^b \psi(x - \theta) dF(x) \leq \int_a^b \psi(x - \theta') dF(x),$$

since ψ is non-decreasing on $(-\infty, 0]$, and $x - \theta < x - \theta' \leq 0$ for $x \leq b$. Therefore, $L(\theta; F) \leq L(\theta'; F)$, which completes the proof of $\theta^*(F) \in [a, b]$. When $\overline{J(F)}$ is of the form $[a, \infty)$ or $(-\infty, b]$, we can prove $\theta^*(F) \in \overline{J(F)}$ in a similar way. Hence, the first part of Lemma 10 is proved. The second part of Lemma 10 follows from Lemma 13. \square

Proof of Theorem 1

First part follows from (1.5) and Proposition 1 noting that \mathbb{F}_n satisfies Condition A. The second part follows from Lemma 10. \square

Proof of Theorem 10

Since $d_W(F_n, F) \rightarrow 0$, from Theorem 6.9 of Villani (2009) or Theorem 7.12 of Villani (2003) it follows that F_n converges to F weakly, and

$$\int_{-\infty}^{\infty} |x| dF_n(x) \rightarrow \int_{-\infty}^{\infty} |x| dF(x) < \infty. \quad (6.38)$$

Let us denote the measure corresponding to F_n by P_n .

For the sake of simplicity, we use the abbreviated notations ψ_n^* , ϕ_n^* , and θ_n^* in place of $\psi^*(F_n)$, $\phi^*(F_n)$, and $\theta^*(F_n)$ respectively.

Our proof is motivated by the proof of Proposition 6 of Xu and Samworth (2019). Since F_n satisfies condition A, $L(F_n) < \infty$ by Proposition 1. Therefore, starting with any subsequence of $\{F_n\}_{n \geq 1}$, we can extract a further subsequence $\{F_{n_k}\}_{k \geq 1}$, such that $L(F_{n_k}) \rightarrow \vartheta \in [-\infty, \infty]$, and $\theta_{n_k}^* \rightarrow \theta' \in [-\infty, \infty]$.

The proof of Theorem 10 involves the following steps:

- (A) Show that $\vartheta > -\infty$.
- (B) Show that $\vartheta < \infty$, and there exists $M > \vartheta + 1$ such that

$$\limsup_{k \uparrow \infty} \sup_{x \in \mathbb{R}} \psi_{n_k}^*(x) < M.$$

- (C) Show that $\theta' \in \mathbb{R}$.
- (D) Taking a and b to be the left and right endpoints of $J(F)$ respectively, from Lemma 10, we observe that

$$\text{int}(\text{dom}(\psi^*(F))) = (-d, d), \quad (6.39)$$

where $d = \max(b - \theta^*(F), \theta^*(F) - a)$. We now define

$$D_{\theta'} = \begin{cases} \mathbb{R}, & \text{if } a = -\infty, b = \infty, \\ \text{dom}(\psi^*(F)) + \theta^*(F) - \theta', & \text{if } b - \theta^*(F) > \theta^*(F) - a, \\ \text{dom}(\psi^*(F)) - (\theta^*(F) - \theta'), & \text{if } b - \theta^*(F) \leq \theta^*(F) - a. \end{cases} \quad (6.40)$$

We Show that $\liminf_k \psi_{n_k}^*(x) > -\infty$ for $x \in \text{int}(D_{\theta'})$, leading to $\liminf_k \psi_{n_k}^*(0) > -\infty$.

(E) Show that there exist $\alpha', \beta' > 0$ such that for all sufficiently large k ,

$$\psi_{n_k}^*(x) \leq -\alpha'|x| + \beta', \quad \text{for all } x \in \mathbb{R}.$$

(F) Show that $\vartheta = L(F)$, $\theta' = \theta$, and $\psi_{n_k}^*$ converges pointwise to $\psi^*(F)$ on $\text{int}(\text{dom}(\psi^*(F)))$.

The proof of (A) is quite straightforward. To see this, observe that the function $\psi(x) = -|x|$ is a member of \mathcal{SC}_0 . Therefore,

$$\vartheta = \lim_{k \rightarrow \infty} L(F_{n_k}) \geq \lim_{k \rightarrow \infty} \Psi(0, \psi, F_{n_k}) \geq - \lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} |x| dF_{n_k}(x) - \int_{-\infty}^{\infty} e^{-|x|} dx,$$

which equals $-\int |x| dF(x) - 2$ by (6.38). Since $\int |x| dF(x) < \infty$ by Condition A, step (A) follows.

The proof of (B) hinges on the proof of Theorem 2.2 of [Dümbgen et al. \(2011\)](#). Suppose $\limsup_k \sup_{x \in \mathbb{R}} \psi_{n_k}^*(x) = \infty$. In that case we can replace $\{n_k\}$ by a further subsequence such that $\lim_k \sup_{x \in \mathbb{R}} \psi_{n_k}^*(x) = \infty$. Hence, without loss of generality, we assume that $\lim_k \sup_{x \in \mathbb{R}} \psi_{n_k}^*(x) = \infty$. Also, note that $\sup_x \psi_{n_k}^*(x) = \psi_{n_k}^*(0) = M_k$, say.

Letting $D_{k,t}$ be the level set $\{\psi_{n_k}^* \geq t\}$, for any $c > 0$, we can bound $L(F_{n_k})$ in the following way:

$$\begin{aligned} L(F_{n_k}) &= \int_{-\infty}^{\infty} \psi_{n_k}^*(x - \theta_{n_k}^*) dF_{n_k}(x) - 1 \\ &= \int_{-\infty}^{\infty} \psi_{n_k}^*(x) dF_{n_k}(x + \theta_{n_k}^*) - 1 \\ &\leq -cM_k P_{n_k}(\mathbb{R} \setminus D_{k, -cM_k} + \theta_{n_k}^*) + M_k P_{n_k}(D_{k, -cM_k} + \theta_{n_k}^*) - 1 \\ &= -cM_k \left(1 - P_{n_k}(D_{k, -cM_k} + \theta_{n_k}^*)\right) + M_k P_{n_k}(D_{k, -cM_k} + \theta_{n_k}^*) - 1 \\ &= -(c+1)M_k \left(\frac{c}{c+1} - P_{n_k}(D_{k, -cM_k} + \theta_{n_k}^*)\right) - 1. \end{aligned} \quad (6.41)$$

Let us denote the Lebesue measure on \mathbb{R} by Leb . We intend to show that $\text{Leb}(D_{k, -cM_k}) \rightarrow 0$ as $k \rightarrow \infty$. Applying Lemma 4.1 of [Dümbgen et al. \(2011\)](#),

we obtain that as $M_k \rightarrow \infty$,

$$\begin{aligned} \text{Leb}(D_{k,-cM_k}) &\leq (1+c)M_k e^{-M_k} / \int_0^{(1+c)M_k} t e^{-t} dt \\ &= (1+c)M_k e^{-M_k} / (1+o(1)) \rightarrow 0. \end{aligned} \quad (6.42)$$

Now we will show that $P_{n_k}(D_{k,-cM_k} + \theta_{n_k}^*) \rightarrow 0$. We consider two cases, $|\theta'| < \infty$ and $\theta' = \pm\infty$. Let us focus on the case $|\theta'| < \infty$ first. Now the concavity of $\psi_{n_k}^*$ indicates that sets of the form $D_{k,-cM_k}$ are intervals including 0. Since $\text{Leb}(D_{k,-cM_k}) \rightarrow 0$ and $\theta_{n_k}^* \rightarrow \theta'$ as $k \rightarrow \infty$, for sufficiently large k , the set $D_{k,-cM_k} + \theta_{n_k}^*$ is an interval around θ' whose length shrinks to 0 as k increases. Now since F is non-degenerate, one can certainly find a $x_0 \neq \theta'$ such that $x_0 \in \text{int}(J(F))$. As a result, $x_0 \notin D_{k,-cM_k} + \theta_{n_k}^*$ for all sufficiently large k . Thus, for all sufficiently large k ,

$$D_{k,-cM_k} + \theta_{n_k}^* \in \{C \subset \mathbb{R} \text{ closed and convex, } x_0 \notin \text{int}(C)\} = D_{x_0},$$

say. However, since F_{n_k} weakly converges to F , Lemma 2.13 of [Dümbgen et al. \(2011\)](#) implies that

$$\limsup_k \sup_{C \in D_{x_0}} P_{n_k}(C) \leq \sup_{C \in D_{x_0}} P_0(C). \quad (6.43)$$

By Lemma 2.13 of [Dümbgen et al. \(2011\)](#), $\sup_{C \in D_{x_0}} P(C)$ is bounded away from 1 if

$$x_0 \in \bigcap_{C \in \mathcal{D}} C,$$

where

$$\mathcal{D} = \left\{ C \mid C \text{ is closed and convex, } P(C) = 1 \right\}.$$

Since $x_0 \in \text{int}(J(F))$, the above holds. Therefore, we conclude that the term in the left hand side of (6.43) is less than $c/(c+1)$ for sufficiently large c . Since $D_{k,-cM_k} + \theta_{n_k}^* \in D_{x_0}$, the above implies that $P_{n_k}(D_{k,-cM_k} + \theta_{n_k}^*) < c/(c+1)$ for sufficiently large k . Combined with (6.41), this result leads to

$$L(F_{n_k}) \rightarrow -\infty \quad \text{as } k \rightarrow \infty,$$

which contradicts the fact that $\vartheta > -\infty$. Therefore, for this case, we see that $\lim_k M_k = \infty$ leads to a contradiction.

At this point it is clear that for the other case also, i.e. when $\theta' = \pm\infty$, it is enough to show that $\limsup_k P_{n_k}(D_{k,-cM_k} + \theta_{n_k}^*) < c/(c+1)$ for sufficiently large c . We will consider the case $\theta' = \infty$. The proof for $\theta' = -\infty$ will follow similarly. Consider y such that $1 - F(y) < c/(c+1)$. Since $\text{Leb}(D_{k,-cM_k}) \rightarrow 0$, we observe that as $\theta_{n_k}^* \rightarrow \infty$, for all sufficiently large k ,

$$D_{k,-cM_k} + \theta_{n_k}^* \subset (y, \infty).$$

Noting $F_{n_k} \rightarrow_w F$, we obtain that

$$\limsup_k P_{n_k}(D_{k,-cM_k} + \theta_{n_k}^*) \leq \limsup_k (1 - F_{n_k}(y)) = 1 - F(y) < c/(c+1),$$

which completes the proof of (B). Therefore, we assume that $\{M_k\}_{k \geq 1}$ is bounded above by a number, say M , which can be chosen to be greater than $\vartheta + 1$.

Our next step is step (C), which follows from Lemma 12. To see this, observe that Lemma 12 implies

$$\phi_{n_k}^*(x) \leq -\log |2(x - \theta_{n_k}^*)|.$$

Suppose $\limsup_k |\theta_{n_k}^*| = \infty$. Then Fatou's Lemma implies that

$$\begin{aligned} \liminf_k \int_{-\infty}^{\infty} e^{\phi_{n_k}^*(x)} dx &\leq \liminf_k \int_{-\infty}^{\infty} e^{-\log |2(x - \theta_{n_k}^*)|} dx \\ &\leq \int_{-\infty}^{\infty} \liminf_k e^{-\log |2(x - \theta_{n_k}^*)|} dx = 0. \end{aligned}$$

However, the left hand side of the above display equals 1 since $e^{\phi_{n_k}^*}$ is a density, which leads to a contradiction. Therefore, step (C) follows.

Next we will prove (D). For $x \in D_{\theta'}$, denote $A_{n_k} = (-x + \theta_{n_k}^*, x + \theta_{n_k}^*)$. We calculate

$$\begin{aligned} L(F_{n_k}) &= \int_{-\infty}^{\infty} \psi_{n_k}^*(z) dF_{n_k}(z + \theta_{n_k}^*) - 1 \\ &= \int_{-x}^x \psi_{n_k}^*(z) dF_{n_k}(z + \theta_{n_k}^*) + \int_{-\infty}^{-x} \psi_{n_k}^*(z) dF_{n_k}(z + \theta_{n_k}^*) \\ &\quad + \int_x^{\infty} \psi_{n_k}^*(z) dF_{n_k}(z + \theta_{n_k}^*) - 1 \\ &\leq \psi_{n_k}^*(0)P_{n_k}(A_{n_k}) + \psi_{n_k}^*(x)P_{n_k}(\mathbb{R} \setminus A_{n_k}) - 1, \end{aligned}$$

which follows since $\psi_{n_k}^* \in \mathcal{SC}_0$. Now observe that the above implies

$$\begin{aligned} \liminf_{k \rightarrow \infty} \psi_{n_k}^*(x) &\geq \liminf_{k \rightarrow \infty} \frac{L(F_{n_k}) + 1 - \psi_{n_k}^*(0)P_{n_k}(A_{n_k})}{P_{n_k}(\mathbb{R} \setminus A_{n_k})} \\ &\geq -\frac{M - 1 - \vartheta}{\liminf_{k \rightarrow \infty} P_{n_k}(\mathbb{R} \setminus A_{n_k})}. \end{aligned} \tag{6.44}$$

Note that for every $\epsilon > 0$,

$$\theta' - \epsilon < \theta_{n_k}^* < \theta' + \epsilon, \quad \text{for all sufficiently large } k.$$

Therefore, we can write

$$\begin{aligned}
 & \liminf_{k \rightarrow \infty} P_{n_k}(\mathbb{R} \setminus A_{n_k}) \\
 &= \liminf_{k \rightarrow \infty} F_{n_k}(-x + \theta_{n_k}^*) + 1 - \limsup_{k \rightarrow \infty} F_{n_k}\left((x + \theta_{n_k}^*) -\right) \\
 &\geq \liminf_{k \rightarrow \infty} F_{n_k}(-x + \theta' - \epsilon) + 1 - \limsup_{k \rightarrow \infty} F_{n_k}\left((x + \theta' + \epsilon) -\right) \\
 &\geq F\left((-x + \theta' - \epsilon) -\right) + 1 - F\left((x + \theta' + \epsilon) -\right).
 \end{aligned}$$

If $J(F) = \mathbb{R}$, it is clear that the right hand side of the last display is positive. Suppose $J(F) \neq \mathbb{R}$. Then at least one of a and b , the endpoints of $J(F)$, is finite. Now suppose $b - \theta^*(F) > \theta^*(F) - a$. Noting $x \in \text{int}(D_{\theta'})$, we argue that for small enough $\epsilon > 0$, $x + \epsilon \in \text{int}(D_{\theta'})$. Then from (6.39) and (6.40) it follows that

$$x + \epsilon < b - \theta^*(F) + \theta^*(F) - \theta',$$

implying that $x + \epsilon + \theta' < b$, leading to $F((x + \theta' + \epsilon) -) < 1$ since b is the right endpoint of $J(F)$. Now consider the case when $b - \theta^*(F) \leq \theta^*(F) - a$. Analogous to the previous case, (6.39) and (6.40) yield

$$x + \epsilon < \theta^*(F) - a - \theta^*(F) + \theta',$$

or equivalently,

$$a < -x + \theta' - \epsilon,$$

implying $F((-x + \theta' - \epsilon) -) > 0$, which leads to

$$\limsup_{k \rightarrow \infty} \mathbb{P}_n(\mathbb{R} \setminus A_{n_k}) > 0.$$

Since in step (B), M was chosen such that $M > \vartheta + 1$, (6.44) entails that

$$\liminf_{k \rightarrow \infty} \widehat{\psi}_{\theta_{r_k}}(x) > -\infty$$

which proves (D).

To prove (E), we first observe that since $\psi_{n_k}^* \in \mathcal{SC}_0$, Lemma 3 of Pal *et al.* (2007) implies that

$$\psi_{n_k}^*(x) \leq \psi_{n_k}^*(0) + 1 - e^{\psi_{n_k}^*(0)|x|}, \quad x \in \mathbb{R}.$$

Observe that (B) and (D), combined with the above, imply that there exist $\alpha' > 0$ and $\beta' \in \mathbb{R}$ such that for all sufficiently large k ,

$$\psi_{n_k}^*(x) \leq \beta' - \alpha'|x| \quad \text{for all } x \in \mathbb{R}. \quad (6.45)$$

Now we move on to step (F). Since we have established (6.45) and showed that $\liminf_{k \rightarrow \infty} \widehat{\psi}_{\theta_{r_k}}(x) > -\infty$ for $x \in \text{int}(D_{\theta'})$, by Lemma 4.2 of Dümbgen *et al.*

(2011), we can replace $\{\psi_{n_k}^*\}_{k \geq 1}$, if necessary, by a subsequence such that for a concave function $\bar{\psi}$ the following conditions are met:

$$\begin{aligned} \text{int}(D_{\theta'}) &\subset \text{dom}(\bar{\psi}), \\ \bar{\psi}(x) &\leq \beta' - \alpha'|x| && \text{for all } x \in \mathbb{R}, k \in \mathbb{N}, \\ \lim_{k \rightarrow \infty, y \rightarrow x} \psi_{n_k}^*(y) &= \bar{\psi}(x) && \text{for all } x \in \text{int}(\text{dom}(\bar{\psi})), \\ \limsup_{k \rightarrow \infty, y \rightarrow x} \psi_{n_k}^*(y) &\leq \bar{\psi}(x) && \text{for all } x \in \mathbb{R}. \end{aligned} \quad (6.46)$$

We will now establish that $\bar{\psi} \in \mathcal{SC}_0$. To see this, first notice that since $\psi_{n_k}^* \in \mathcal{SC}_0$, for $x \notin \text{dom}(\bar{\psi})$,

$$\lim_{k \rightarrow \infty} \psi_{n_k}^*(-x) = \lim_{k \rightarrow \infty} \psi_{n_k}^*(x) \leq \bar{\psi}(x) = -\infty.$$

Hence, $-x \notin \text{int}(\text{dom}(\bar{\psi}))$, suggesting that either $-x \notin \text{dom}(\bar{\psi})$, or $-x \in \text{dom}(\bar{\psi})$ is a boundary point of $\text{dom}(\bar{\psi})$. Therefore, $\text{dom}(\bar{\psi})$ is of the form $(-c, c)$, $[-c, c]$, $(-c, c]$, or $[-c, c)$ for some $c > 0$. For the last two cases, if we extend $\bar{\psi}$ to $[-c, c]$ so that $\bar{\psi}$ is continuous on $[-c, c]$ and $-\infty$ everywhere else, $\bar{\psi}$ still satisfies (6.46). Therefore, without loss of generality, we assume that $\text{dom}(\bar{\psi})$ is of the form $(-c, c)$ or $[-c, c]$. For $y \in (-c, c)$ we obtain that

$$\bar{\psi}(y) = \lim_{k \rightarrow \infty} \psi_{n_k}^*(y) = \lim_{k \rightarrow \infty} \psi_{n_k}^*(-y) = \bar{\psi}(-y).$$

It remains to show that when $\text{dom}(\bar{\psi})$ is the closed interval $[-c, c]$, $\bar{\psi}(-c)$ and $\bar{\psi}(c)$ agree. Since ψ^* is concave, it is right continuous at $-c$, and left continuous at c . Therefore, the symmetry of ψ^* on $(-c, c)$ implies symmetry on $[-c, c]$, which establishes that $\bar{\psi} \in \mathcal{SC}_0$. Now we claim that $e^{\bar{\psi}} \in \mathcal{SLLC}_0$. To see this observe that since $\psi_{n_k}^*(x) \leq \beta' - \alpha'|x|$, application of the dominated convergence theorem yields

$$\int_{\text{int}(\text{dom}(\bar{\psi}))} e^{\bar{\psi}(x)} dx = \int_{\text{int}(\text{dom}(\bar{\psi}))} e^{\lim_k \psi_{n_k}^*(x)} dx = \lim_k \int_{\text{dom}(\bar{\psi})} e^{\psi_{n_k}^*(x)} dx = 1.$$

Since Condition B holds, note that if we can show

$$\vartheta = \Psi(\theta', \bar{\psi}, F) = L(F),$$

we have $\theta' = \theta^*(F)$, $\bar{\psi} = \psi^*(F)$, and (3.24) also follows. Therefore, we have $\theta_n^* \rightarrow \theta^*(F)$, and (6.46) leads to the pointwise convergence of ψ_n^* to $\psi^*(F)$. The pointwise convergence of ψ_n^* 's implies pointwise convergence of $e^{\psi_n^*}$, which, in its turn leads to the weak convergence of the corresponding distribution functions. Then the rest of the proof of Theorem 13 follows from Proposition 2 of Cule and Samworth (2010). We prove the equations in the last display by first establishing that

$$\vartheta \leq \Psi(\theta', \bar{\psi}, F) \leq L(F), \quad (6.47)$$

and then showing $\vartheta \geq L(F)$.

Observe that since F_{n_k} converges to F weakly, by Skorohod's theorem, there exists a probability space $(\Omega, \mathcal{A}, P^*)$ with random variable $X_k \sim F_{n_k}$, and $X \sim F$ such that $X_k \rightarrow_{a.s.} X$ as $k \rightarrow \infty$. Denote by E^* the expectation operator with respect to the probability space $(\Omega, \mathcal{A}, P^*)$. Denoting the positive random variable $\beta' - \alpha'|X_k| - \phi_{n_k}^*(X_k)$ by H_k , we calculate

$$\begin{aligned} \vartheta &= \limsup_{k \rightarrow \infty} \int_{-\infty}^{\infty} \phi_{n_k}^*(x) dF_{n_k}(x) - 1 \\ &= \lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} (\beta' - \alpha'|x|) dF_{n_k}(x) - \liminf_{k \rightarrow \infty} E^*[H_k] - 1 \\ &\leq \beta' - \alpha' \int_{-\infty}^{\infty} |x| dF(x) - E^* \left[\liminf_{k \rightarrow \infty} H_k \right] - 1, \end{aligned}$$

which follows using (6.38) and Fatou's Lemma. As $\theta_{n_k}^* \rightarrow \theta' \in \mathbb{R}$, and $X_k \rightarrow_{a.s.} X$, (6.46) indicates that the following holds with probability 1:

$$\liminf_{k \rightarrow \infty} H_k = \alpha' - \beta'|X| - \limsup_k \psi_{n_k}^*(X_k - \theta_{n_k}^*) \geq \beta' - \alpha'|X| - \bar{\psi}(X - \theta').$$

Since $X \sim F$, the above leads to

$$\begin{aligned} \vartheta &\leq \beta' - \alpha' \int_{-\infty}^{\infty} |x| dF(x) - E^* \left[\beta' - \alpha'|X| - \bar{\psi}(X - \theta') \right] - 1 \\ &= \int_{-\infty}^{\infty} \bar{\psi}(x - \theta') dF(x) - 1 \\ &= \Psi(\theta', \bar{\psi}, F) \leq L(F), \end{aligned}$$

which completes the proof of (6.47). Hence, it only remains to prove that $\vartheta \geq L(F)$.

Let us denote $c_0 = \psi^*(F)(0)$. From Lemma 15 it follows that for each $\epsilon > 0$, there exist $\psi^{(\epsilon)} \in \mathcal{SC}_0$ such that

$$-|x|/\epsilon + c_0 \leq \psi^{(\epsilon)}(x) \quad \text{for all } x \in \mathbb{R}, \quad (6.48)$$

and

$$\psi^*(F) \leq \psi^{(\epsilon)} \leq c_0. \quad (6.49)$$

Further, $\psi^{(\epsilon)}$ decreases to $\psi^*(F)$ pointwise as $\epsilon \downarrow 0$. Using the random variables $X_k \sim F_{n_k}$ and $X \sim F$ constructed in the proof of (6.47), we write

$$\begin{aligned} \vartheta &= \lim_{k \rightarrow \infty} L(F_{n_k}) \geq \lim_{k \rightarrow \infty} \Psi(\theta^*(F), \psi^{(\epsilon)}, F_{n_k}) \\ &= \lim_{k \rightarrow \infty} E^* \left[\psi^{(\epsilon)}(X_k - \theta^*(F)) \right] - \int_{-\infty}^{\infty} e^{\psi^{(\epsilon)}(x)} dx. \quad (6.50) \end{aligned}$$

Note that (6.48) and (6.49) indicate that $\psi^{(\epsilon)}(x)$ is bounded above and below by c_0 and $-|x|/\epsilon + c_0$ respectively. From (6.38) it follows that

$$E^*|X_k| = \int_{-\infty}^{\infty} |x| dF_{n_k}(x) \rightarrow \int_{-\infty}^{\infty} |x| dF(x), \quad \text{as } k \rightarrow \infty.$$

Moreover, as $k \rightarrow \infty$, we also have $X_k \rightarrow_{a.s.} X$. Observe that $\psi^{(\epsilon)}$ is concave, and by (6.48), $\text{dom}(\psi^{(\epsilon)}) = \mathbb{R}$, which implies that $\psi^{(\epsilon)}$ is continuous. Therefore,

$$\psi^{(\epsilon)}(X_k - \theta^*(F)) \rightarrow_{a.s.} \psi^{(\epsilon)}(X - \theta^*(F)).$$

Therefore, using Lemma 16, from (6.50) we obtain that

$$\begin{aligned} \vartheta &\geq E \left[\lim_{k \rightarrow \infty} \psi^{(\epsilon)}(X_k - \theta^*(F)) \right] - \int_{-\infty}^{\infty} e^{\psi^{(\epsilon)}(x)} dx \\ &= \int_{-\infty}^{\infty} \psi^{(\epsilon)}(x - \theta^*(F)) dF(x) - \int_{-\infty}^{\infty} e^{\psi^{(\epsilon)}(x)} dx. \end{aligned}$$

Since $\psi^{(\epsilon)}$ decreases to $\psi^*(F)$ pointwise as $\epsilon \downarrow 0$, the monotone convergence theorem and (6.49) lead to

$$\lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} \left(c_0 - \psi^{(\epsilon)}(x - \theta^*(F)) \right) dF(x) = \int_{-\infty}^{\infty} \left(c_0 - \psi^*(F)(x - \theta^*(F)) \right) dF(x),$$

which can be rewritten as

$$\lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} \psi^{(\epsilon)}(x - \theta^*(F)) dF(x) = \int_{-\infty}^{\infty} \psi^*(F)(x - \theta^*(F)) dF(x). \quad (6.51)$$

On the other hand, note that (6.49) implies that $g^*(F) \leq e^{\psi^{(\epsilon)}} \leq g^*(F)(0)$. Since $e^{\psi^{(\epsilon)}}$ decreases to $g^*(F)$ pointwise as ϵ decreases to 0, the monotone convergence theorem implies that

$$\lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} \left(g^*(F)(0) - e^{\psi^{(\epsilon)}(x)} \right) dx = \int_{-\infty}^{\infty} \left(g^*(F)(0) - g^*(F)(x) \right) dx,$$

or equivalently,

$$\lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} e^{\psi^{(\epsilon)}(x)} dx = \int_{-\infty}^{\infty} g^*(F)(x) dx = 1. \quad (6.52)$$

Therefore, from (6.51) and (6.52), we conclude that

$$\begin{aligned} \vartheta &\geq \lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} \psi^{(\epsilon)}(x - \theta^*(F)) dF(x) - \lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} e^{\psi^{(\epsilon)}(x)} dx \\ &= \int_{-\infty}^{\infty} \psi^*(F)(x - \theta^*(F)) dF(x) - 1 = L(F), \end{aligned}$$

which completes the proof. \square

Proof of Theorem 9

In their proof of Theorem 3.1, Pal *et al.* (2007) show that if a sequence of log-concave functions $\{f_n\}_{n \geq 1}$ (which can be stochastic as well) satisfies

$$\sum_{i=1}^n \log f_n(X_i) \geq \sum_{i=1}^n \log f_0(X_i) \quad a.s., \quad (6.53)$$

we have $H(f_n, f_0) \rightarrow_{a.s.} 0$, provided

$$P\left(\sup_x \log f_n(x) = o\left(\frac{\sqrt{n}}{\log n}\right)\right) = 1.$$

If we take $f_n = \hat{f}_n$, we have

$$\sup_x \log \hat{f}_n(x) = \hat{\phi}_n(\hat{\theta}_n) = \hat{\psi}_n(0).$$

Theorem 11(c) and Corollary 4 entail that

$$P\left(\limsup_n \hat{\psi}_n(0) < \infty\right) = 1.$$

Also, note that being the MLE of f_0 , \hat{f}_n automatically satisfies (6.53), which implies

$$H(\hat{f}_n, f_0) \rightarrow_{a.s.} 0. \quad (6.54)$$

Since $f_0 \in \mathcal{P}_0$ is log-concave and continuous, Proposition 2(c) of Cule and Samworth (2010) and (6.54) entail that

$$\sup_{z \in \mathbb{R}} |\hat{f}_n(z) - f_0(z)| \rightarrow_{a.s.} 0. \quad (6.55)$$

Note that

$$\begin{aligned} 2H^2(\hat{g}_n, g_0) &= \int_{-\infty}^{\infty} \left(\sqrt{\hat{g}_n(z - \hat{\theta}_n)} - \sqrt{g_0(z - \hat{\theta}_n)} \right)^2 dz \\ &\leq 2 \int_{-\infty}^{\infty} \left(\sqrt{\hat{g}_n(z - \hat{\theta}_n)} - \sqrt{g_0(z - \theta_0)} \right)^2 dz \\ &\quad + 2 \int_{-\infty}^{\infty} \left(\sqrt{g_0(z - \hat{\theta}_n)} - \sqrt{g_0(z - \theta_0)} \right)^2 dz, \end{aligned}$$

where the first term on the right hand side of the last display approaches 0 almost surely by (6.54). The second term is also bounded above by a constant multiple of $g_0(z - \hat{\theta}_n) + g_0(z - \theta_0)$, which is integrable. Therefore, using Lemma 16, Theorem 11(a), and Corollary 4, we deduce that this term also converge to 0 almost surely. Hence $H^2(\hat{g}_n, g_0) \rightarrow_{a.s.} 0$ follows.

Our next step is to find the rate of convergence of $H(\hat{f}_n, f_0)$. To do so, we first introduce the class of functions

$$\mathcal{P}_{M,0} = \left\{ f \in \mathcal{L}\mathcal{C} \mid \begin{array}{l} \sup_{x \in \mathbb{R}} f(x) < M, \quad f(x) > 1/M \text{ for all } |x| < 1, \\ \text{Supp}(f) \subset \text{Supp}(f_0) \end{array} \right\}.$$

We will show that without loss of generality, one can assume that $f_0 \in \mathcal{P}_{M,0}$ for some $M > 0$. To this end, we translate and rescale the data letting $\tilde{X}_i =$

$\alpha X_i + \beta$, where $\alpha > 0$ and $\beta \in \mathbb{R}$. Observe that the rescaled data has density $\tilde{f}_0(x) = \alpha^{-1} f_0((x - \beta)/\alpha)$. Denote by $\tilde{f}_{0,n}$ the MLE of f_0 based on the rescaled data. Note that the MLE is affine-equivalent, which entails that $\tilde{f}_{0,n}(x) = \alpha^{-1} \hat{f}_n((x - \beta)/\alpha)$. Noting Hellinger distance is invariant under affine transformations, we observe that $H(\hat{f}_n, f_0) = H(\tilde{f}_{0,n}, \tilde{f}_0)$. Therefore, it suffices to show that $H(\tilde{f}_{0,n}, \tilde{f}_0) \rightarrow_{a.s.} 0$. Note that since f_0 is log-concave, $\text{int}(\text{dom}(f_0))$ contains an interval. We can choose α and β in a way such that $(x - \beta)/\alpha$ lie inside that interval for $x = \pm 1$. Then it is possible to find $M > 0$ large enough such that

$$f_0((x - \beta)/\alpha) > \alpha/M, \quad x = \pm 1,$$

or

$$\min(\tilde{f}_0(-1), \tilde{f}_0(1)) > 1/M,$$

leading to

$$\inf_{x \in [-1, 1]} \tilde{f}_0(x) > 1/M,$$

since f_0 , or equivalently \tilde{f}_0 is unimodal. Hence, without loss of generality, we can assume that there exists $M > 0$ such that $f_0(x) > 1/M$ for $x \in [-1, 1]$. We can choose M large enough such that additionally, $\sup_{x \in \mathbb{R}} f_0(x) < M$. On the other

hand, (6.55) implies that the following inequalities hold with probability one:

$$\limsup_n \sup_{x \in \mathbb{R}} \hat{f}_n(x) < M,$$

and

$$\lim_n \hat{f}_n(\pm 1) > 1/M.$$

Therefore, $f_0 \in \mathcal{P}_{M,0}$, and with probability one, $\hat{f}_n \in \mathcal{P}_{M,0}$ as well for all sufficiently large n . Doss and Wellner (2016) obtained the bracketing entropy of the class $\mathcal{P}_{M,0}$. They showed that for any $\epsilon > 0$,

$$\log N_{[\]}(\epsilon, \mathcal{P}_{M,0}, H) \lesssim \epsilon^{-1/2}.$$

The rest of the proof now follows from an application of Theorem 3.4.1 and 3.4.4 of Van der Vaart and Wellner (1996). To this end, fixing $\epsilon > 0$, for $f \in \mathcal{P}_{M,0}$ we define the function

$$m_f(x) = \log \left(\frac{f(x) + f_0(x)}{2f_0(x)} \right),$$

and we let \mathcal{M}_ϵ denote the class

$$\mathcal{M}_\epsilon = \{m_f - m_{f_0} : H(f, f_0) \leq \epsilon\}.$$

Also, we denote

$$\|\mathbb{Z}_n\|_{\mathcal{M}_\epsilon} = \sup_{h \in \mathcal{M}_\epsilon} \int h d\mathbb{Z}_n.$$

Then Theorem 3.4.4 of [Van der Vaart and Wellner \(1996\)](#) yields

$$P(m_f - m_{f_0}) \lesssim -H^2(f, f_0),$$

and

$$E_P \|Z_n\|_{\mathcal{M}_\epsilon} \lesssim \tilde{J}_{[\cdot]}(\epsilon, \mathcal{P}_{M,0}, H) \left(1 + \frac{\tilde{J}_{[\cdot]}(\epsilon, \mathcal{P}_{M,0}, H)}{\epsilon^2 \sqrt{n}} \right),$$

where

$$\tilde{J}_{[\cdot]}(\epsilon, \mathcal{P}_{M,0}, H) = \int_0^\epsilon \sqrt{\log N_{[\cdot]}(t, \mathcal{P}_{M,0}, H)} dt \lesssim \epsilon^{3/4}.$$

Therefore, it is easy to see that

$$E_P \|Z_n\|_{\mathcal{M}_\epsilon} \lesssim \epsilon^{3/4} + \epsilon^{-1/2} n^{-1/2} = g(\epsilon),$$

where

$$g(t) = t^{3/4} + t^{-1/2} n^{-1/2}, \quad t > 0.$$

Notice that $g(t)/t^{3/4}$ is decreasing in t . Also, for any constant $c > 0$, we have

$$n^{4/5} g(cn^{-2/5}) = (c^{3/4} + c^{-1/2}) n^{1/2}.$$

Therefore, an application of Theorem 3.4.1 of [Van der Vaart and Wellner \(1996\)](#) yields

$$H(\hat{f}_n, f_0) = O_p(n^{-2/5}),$$

which completes the proof of part A of Theorem 9.

Now we turn to the proof of part B. If $x - \theta_0$ is a continuity point of g'_0 , noting $\hat{\theta}_n \rightarrow_{a.s.} \theta_0$ by Theorem 11 and Corollary 4, we obtain that

$$\frac{\sqrt{g_0(x - \hat{\theta}_n)} - \sqrt{g_0(x - \theta_0)}}{(\hat{\theta}_n - \theta_0)} \rightarrow_{a.s.} \frac{g'_0(x - \theta_0)}{2\sqrt{g_0(x - \theta_0)}}.$$

Noting g'_0 is continuous almost everywhere with respect to Lebesgue measure, and using Fatou's lemma and part A of the current theorem, we obtain that

$$\begin{aligned} & \liminf_n \frac{\int_{-\infty}^{\infty} \left(\sqrt{g_0(x - \hat{\theta}_n)} - \sqrt{g_0(x - \theta_0)} \right)^2 dx}{(\hat{\theta}_n - \theta_0)^2} \\ & \geq \int_{-\infty}^{\infty} \left(\frac{g'_0(x - \theta_0)}{2\sqrt{g_0(x - \theta_0)}} \right)^2 dx = \frac{\mathcal{I}_{f_0}}{4} \quad a.s. \end{aligned} \tag{6.56}$$

Now observe that

$$\begin{aligned} & 2H(\widehat{f}_n, f_0)^2 \\ &= \int_{-\infty}^{\infty} \left(\sqrt{\widehat{g}_n(x - \widehat{\theta}_n)} - \sqrt{g_0(x - \theta_0)} \right)^2 dx \\ &= \int_{-\infty}^{\infty} \left(\sqrt{\widehat{g}_n(x - \widehat{\theta}_n)} - \sqrt{g_0(x - \widehat{\theta}_n)} \right)^2 dx \\ &\quad + \int_{-\infty}^{\infty} \left(\sqrt{g_0(x - \widehat{\theta}_n)} - \sqrt{g_0(x - \theta_0)} \right)^2 dx + T_c, \end{aligned}$$

where

$$T_c = 2 \int_{-\infty}^{\infty} \left(\sqrt{\widehat{g}_n(x - \widehat{\theta}_n)} - \sqrt{g_0(x - \widehat{\theta}_n)} \right) \left(\sqrt{g_0(x - \widehat{\theta}_n)} - \sqrt{g_0(x - \theta_0)} \right) dx.$$

The inequality in (6.56) entails that for all sufficiently large n ,

$$2H(\widehat{f}_n, f_0)^2 \geq 2H(\widehat{g}_n, g_0)^2 + \frac{(\widehat{\theta}_n - \theta_0)^2 \mathcal{I}_{f_0}}{4} - |T_c| \quad a.s.$$

We aim to show that the cross-term $|T_c|$ is small. In fact, we show that

$$\frac{|T_c|}{|\widehat{\theta}_n - \theta_0|^2 + H(\widehat{g}_n, g_0)^2} = o_p(1). \quad (6.57)$$

Suppose (6.57) holds. Then it follows that

$$\begin{aligned} 2H(\widehat{f}_n, f_0)^2 &\geq 2H(\widehat{g}_n, g_0)^2 \\ &\quad + \frac{(\widehat{\theta}_n - \theta_0)^2 \mathcal{I}_{f_0}}{4} - o_p(1)H(\widehat{g}_n, g_0)^2 - o_p(1)(\widehat{\theta}_n - \theta_0)^2, \end{aligned}$$

which completes the proof because $\mathcal{I}_{f_0} > 0$.

Hence, it remains to prove (6.57). To this end, notice that T_c can be written as

$$T_c = 2 \int_{-\infty}^{\infty} \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right) \left(\sqrt{g_0(x)} - \sqrt{g_0(x + \widehat{\theta}_n - \theta_0)} \right) dx.$$

For the sake of simplicity, we will denote $\theta_0 - \widehat{\theta}_n$ by δ_n from now on. Since g_0 is absolutely continuous, we can write

$$|T_c| = \left| 2 \int_{-\infty}^{\infty} \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right) \left(\int_{-\delta_n}^0 \frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} dt \right) dx \right|.$$

Since $g_0 \in \mathcal{S}_0$, we have

$$\begin{aligned} |T_c| &= 2 \left| \int_0^\infty \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right) \left(\int_{-\delta_n}^0 \frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} dt \right) dx \right. \\ &\quad \left. - \int_{-\infty}^0 \left(\sqrt{\widehat{g}_n(-x)} - \sqrt{g_0(-x)} \right) \left(\int_{-\delta_n}^0 \frac{g'_0(-x-t)}{2\sqrt{g_0(-x-t)}} dt \right) dx \right| \\ &= 2 \left| \int_0^\infty \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right) \left(\int_{-\delta_n}^0 \frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} dt \right) dx \right. \\ &\quad \left. - \int_0^\infty \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right) \left(\int_{-\delta_n}^0 \frac{g'_0(x-t)}{2\sqrt{g_0(x-t)}} dt \right) dx \right|, \end{aligned}$$

yielding

$$|T_c| = 2 \left| \int_0^\infty \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right) \left(\int_{-\delta_n}^0 \left(\frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} - \frac{g'_0(x-t)}{2\sqrt{g_0(x-t)}} \right) dt \right) dx \right|. \quad (6.58)$$

Using the Cauchy-Schwarz inequality, we obtain that

$$\begin{aligned} \frac{|T_c|}{2|\delta_n|} &\leq \left(\int_0^\infty \left(\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)} \right)^2 dx \right)^{1/2} \\ &\quad \left(\int_0^\infty \left(\int_{-\delta_n}^0 \frac{1}{|\delta_n|} \left(\frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} - \frac{g'_0(x-t)}{2\sqrt{g_0(x-t)}} \right) dt \right)^2 dx \right)^{1/2}. \end{aligned}$$

Since $\sqrt{\widehat{g}_n(x)} - \sqrt{g_0(x)}$ is an even function, the first term on the right hand side of the last inequality is $\sqrt{2}H(\widehat{g}_n, g_0)$. Hence,

$$\frac{T_c^2}{8H(\widehat{g}_n, g_0)^2\delta_n^2} \leq \int_0^\infty \left(\int_{-\delta_n}^0 \frac{1}{|\delta_n|} \left(\frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} - \frac{g'_0(x-t)}{2\sqrt{g_0(x-t)}} \right) dt \right)^2 dx,$$

which, noting that

$$t \mapsto \frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} - \frac{g'_0(x-t)}{2\sqrt{g_0(x-t)}}$$

is an even function for each $x > 0$, can be bounded above by

$$\int_0^\infty |\delta_n| \left(\int_0^{|\delta_n|} \frac{1}{(\delta_n)^2} \left(\frac{g'_0(x+t)}{2\sqrt{g_0(x+t)}} - \frac{g'_0(x-t)}{2\sqrt{g_0(x-t)}} \right)^2 dt \right) dx$$

using the Cauchy-Schwarz inequality. Therefore, we obtain

$$\begin{aligned} \frac{T_c^2}{2H(\widehat{g}_n, g_0)^2\delta_n^2} &\leq \frac{1}{|\delta_n|} \int_0^{|\delta_n|} \left[\int_0^\infty \left(\frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} \right)^2 dx + \int_0^\infty \left(\frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \right)^2 dx \right. \\ &\quad \left. - 2 \int_0^\infty \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx \right] dt. \quad (6.59) \end{aligned}$$

For $t \geq 0$,

$$\int_0^\infty \left(\frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} \right)^2 dx = \int_t^\infty \left(\frac{g'_0(x)}{\sqrt{g_0(x)}} \right)^2 dx \leq \frac{\mathcal{I}_{f_0}}{2}.$$

Now observe that for $z \in (-|\delta_n|, 0)$,

$$|g'_0(z)/\sqrt{g_0(z)}| = |\psi'_0(z)|\sqrt{g_0(z)} \leq |\psi'_0(\delta_n)|\sqrt{g_0(0)} = O_p(1), \quad (6.60)$$

since $\psi_0 \in \mathcal{SC}_0$, and $\delta_n \rightarrow_{a.s.} 0$. Hence, for $t \in (0, |\delta_n|)$,

$$\begin{aligned} \int_0^\infty \left(\frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \right)^2 dx &= \int_{-t}^\infty \left(\frac{g'_0(z)}{\sqrt{g_0(z)}} \right)^2 dz \\ &= \int_{-t}^0 \left(\frac{g'_0(x)}{\sqrt{g_0(x)}} \right)^2 dx + \int_0^\infty \left(\frac{g'_0(z)}{\sqrt{g_0(z)}} \right)^2 dz \\ &\leq |\delta_n| |\psi'_0(\delta_n)|^2 \sqrt{g_0(0)} + \mathcal{I}_{f_0}/2 \\ &= |\delta_n| O_p(1) + \mathcal{I}_{f_0}/2, \end{aligned}$$

where the last step follows from (6.60). Hence, for any $t \in (0, |\delta_n|)$,

$$\int_0^\infty \left(\frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} \right)^2 dx + \int_0^\infty \left(\frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \right)^2 dx = |\delta_n| O_p(1) + \mathcal{I}_{f_0}. \quad (6.61)$$

Our objective is to apply Fatou's lemma on the third term on the right hand side of (6.59). Therefore, we want to ensure that the integrand is non-negative. Note that when $x \geq |\delta_n|$ and $t \in (0, |\delta_n|)$, we have $x > t$, which leads to

$$g'_0(x-t)g'_0(x+t) \geq 0. \quad (6.62)$$

Keeping that in mind, we partition the term

$$\begin{aligned} & - \int_0^\infty \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx \\ &= - \int_{|\delta_n|}^\infty \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx - \int_0^{|\delta_n|} \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx \\ &\leq - \int_{|\delta_n|}^\infty \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx + |\delta_n| O_p(1), \end{aligned}$$

where the last step follows from (6.60). The above combined with (6.59) and

(6.61) leads to

$$\begin{aligned}
& \limsup_n \frac{T_c^2}{2H(\widehat{g}_n, g_0)^2 \delta_n^2} \\
& \leq \limsup_n \frac{1}{|\delta_n|} \int_0^{|\delta_n|} \left[|\delta_n| O_p(1) + \mathcal{I}_{f_0} - 2 \int_{|\delta_n|}^{\infty} \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx \right] dt \\
& = O_p(1) \limsup_n |\delta_n| + \mathcal{I}_{f_0} \\
& \quad - 2 \liminf_n \frac{1}{|\delta_n|} \int_0^{|\delta_n|} \int_{|\delta_n|}^{\infty} \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} dx dt \\
& = 0 + \mathcal{I}_{f_0} - 2 \liminf_n \int_{|\delta_n|}^{\infty} \frac{\int_0^{|\delta_n|} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} dt}{|\delta_n|} dx. \tag{6.63}
\end{aligned}$$

Therefore, an application of Fatou's Lemma and (6.62) yield

$$\liminf_n \int_{|\delta_n|}^{\infty} \frac{\int_0^{|\delta_n|} \frac{g'_0(x+t)}{\sqrt{g_0(x+t)}} \frac{g'_0(x-t)}{\sqrt{g_0(x-t)}} dt}{|\delta_n|} dx \geq \int_0^{\infty} \frac{g'_0(x)^2}{g_0(x)} dx = \frac{\mathcal{I}_{f_0}}{2}.$$

Thus (6.63) leads to

$$\frac{2T_c^2}{4H(\widehat{g}_n, g_0)^2 \delta_n^2} = o_p(1).$$

from which it is obvious that

$$\frac{\sqrt{2}|T_c|}{|\delta_n|^2 + H(\widehat{g}_n, g_0)^2} \leq \frac{\sqrt{2}|T_c|}{2H(\widehat{g}_n, g_0)|\delta_n|} = o_p(1),$$

which proves (6.57) and thus completes the proof of part B of Theorem 9. \square

6.2.1. Proof of Lemma 7

From part A of Theorem 9, we obtain that \widehat{f}_n is strongly Hellinger consistent for f_0 . As a consequence, $\|\widehat{f}_n - f_0\|_1 \rightarrow_p 0$. Therefore, from proposition 2(c) of [Cule and Samworth \(2010\)](#), it follows that,

$$\sup_{z \in \mathbb{R}} |\widehat{f}_n(x) - f_0(x)| \rightarrow_{a.s.} 0. \tag{6.64}$$

Now consider any compact set $K = [b_1, b_2] \subset \text{dom}(\phi_0)$. Observe that

$$\begin{aligned} & \sup_{x \in K} |\widehat{\phi}_n(x) - \phi_0(x)| \\ &= \sup_{x \in K} \log \frac{\max(\widehat{f}_n(x), f_0(x))}{\min(\widehat{f}_n(x), f_0(x))} \\ &= \sup_{x \in K} \log \left(\frac{|\widehat{f}_n(x) - f_0(x)|}{\min(\widehat{f}_n(x), f_0(x))} + 1 \right) \\ &\leq \log \left(\frac{\sup_{x \in K} |\widehat{f}_n(x) - f_0(x)|}{\min(\widehat{f}_n(b_1), \widehat{f}_n(b_2), f_0(b_1), f_0(b_2))} + 1 \right), \end{aligned}$$

since \widehat{f}_n and f_0 are unimodal. From (6.64) it follows that $\sup_{x \in K} |\widehat{f}_n(x) - f_0(x)| \rightarrow_{a.s.} 0$. Also, since $b_1 \in \text{dom}(\phi_0)$, $\widehat{f}_n(b_1) \rightarrow_{a.s.} f_0(b_1) > 0$. The same holds for $\widehat{f}_n(b_2)$, which establishes that

$$\sup_{x \in K} |\widehat{\phi}_n(x) - \phi_0(x)| \rightarrow_{a.s.} 0.$$

Now from Lemma F.10 of Kuchibhotla *et al.* (2017), it follows that if ϕ_0 is differentiable at $x \in K$, leading to

$$\widehat{\phi}'_n(x) \rightarrow_{a.s.} \phi'_0(x),$$

which combined with (6.64) yields

$$\widehat{f}'_n(x) \rightarrow_{a.s.} f'_0(x).$$

Part A of Theorem 9 also indicates that $H(\widehat{g}_n, g_0) \rightarrow_{a.s.} 0$, from which, proceeding like above, one can show that similar results hold for \widehat{g}_n and $\widehat{\psi}_n$ as well, which completes the proof. \square

6.3. Additional lemmas

6.4. Additional lemmas for the one-step estimators

Lemma 11. Consider $\zeta, \zeta' \in \text{int}(\text{dom}(\widetilde{\psi}_0))$ such that $\zeta \leq \zeta'$. Suppose $\zeta_n \rightarrow_{a.s.} \zeta$, and $\zeta'_n \rightarrow_{a.s.} \zeta'$. Letting $\widetilde{\psi}'_n$ denote $\log \widetilde{g}_n$, for $\widetilde{g}_n = \widetilde{g}_n^{sym}, \widehat{g}_{\bar{\theta}_n}, \widetilde{g}_n^{geo, sym}$, or $\widehat{h}_n(\bar{\theta}_n \pm \cdot)$, under the conditions of Theorem 12, we have

$$\limsup_n \sup_{z \in [\zeta_n, \zeta'_n]} |\widetilde{\psi}'_n(z)| \leq C_{\zeta, \zeta'} \quad a.s.,$$

where $C_{\zeta, \zeta'}$ is a constant depending on ζ and ζ' . When $\widetilde{g}_n = (\widetilde{g}_n^{sym})^{sm}$ or $\widehat{h}_n^{sm}(\bar{\theta}_n \pm \cdot)$, the above conclusion holds for any $\zeta, \zeta' \in \mathbb{R}$.

Proof. When \tilde{g}_n equals the densities $\widehat{g}_{\bar{\theta}_n}$, $\widehat{f}_n^{geo,sym}$, $\widehat{h}_n(\bar{\theta}_n \pm \cdot)$, or $\widehat{h}_n^{sm}(\cdot + \bar{\theta}_n)$, it is clear that $\tilde{\psi}_n$ is concave. We will consider these choices of \tilde{g}_n first. For the first three cases, the pointwise limit of \tilde{g}_n is \tilde{g}_0 by Theorem 12. Note that we can choose $\zeta_1, \zeta'_1 \in \text{dom}(\tilde{\psi}_0)$ such that $\tilde{\psi}'_0$ is continuous at ζ_1 and ζ'_1 , and $[\zeta - \epsilon, \zeta' + \epsilon] \subset [\zeta_1, \zeta'_1]$ for some sufficiently small $\epsilon > 0$. Now the conditions on ζ_n and ζ'_n underscore that for sufficiently large n ,

$$[\zeta_n, \zeta'_n] \subset [\zeta_1, \zeta'_1] \quad a.s.$$

Now note that a concave $\tilde{\psi}_n$ leads to a non-increasing $\tilde{\psi}'_n$. Therefore, on any interval, $|\tilde{\psi}'_n|$ attains its maxima on either of the endpoints. As a consequence, the supremum of $|\tilde{\psi}'_n|$ on the interval $[\zeta_1, \zeta'_1]$ does not exceed $|\tilde{\psi}'_n(\zeta_1)| + |\tilde{\psi}'_n(\zeta'_1)|$, which converges almost surely to $|\tilde{\psi}'_0(\zeta_1)| + |\tilde{\psi}'_0(\zeta'_1)|$ by Theorem 12. Therefore, we conclude that

$$\limsup_n \sup_{z \in [\zeta_n, \zeta'_n]} |\tilde{\psi}'_n(z)| < |\tilde{\psi}'_0(\zeta_1)| + |\tilde{\psi}'_0(\zeta'_1)| \quad a.s., \quad (6.65)$$

when $\tilde{g}_n = \widehat{g}_{\bar{\theta}_n}$, $\widehat{f}_n^{geo,sym}$, or $\widehat{h}_n(\cdot + \bar{\theta}_n)$. Now consider $\tilde{g}_n = \widehat{h}_n^{sm}(\bar{\theta}_n \pm \cdot)$. Note that the L_1 limit of this \tilde{g}_n is \tilde{g}_0^{sm} , and $\text{dom}(\tilde{\psi}_0^{sm}) = \mathbb{R}$. Therefore, for any $\zeta, \zeta' \in \mathbb{R}$, the same arguments as the previous case will lead to

$$\limsup_n \sup_{z \in [\zeta_n, \zeta'_n]} |\tilde{\psi}'_n(z)| < |(\tilde{\psi}_0^{sm})'(\zeta_1)| + |(\tilde{\psi}_0^{sm})'(\zeta'_1)| \quad a.s. \quad (6.66)$$

Now let us consider the non-log-concave cases. Assume $\tilde{g}_n = \tilde{g}_n^{sym}$. Then from (6.14) we obtain that

$$\tilde{\psi}'_n(x) = \varrho_n(x) \left((\log \widehat{h}_n)'(\bar{\theta}_n + x) \right) - (1 - \varrho_n(x)) \left((\log \widehat{h}_n)'(\bar{\theta}_n - x) \right),$$

where \widehat{h}_n is the log-concave MLE and $\varrho_n(x) < 1$, leading to

$$\limsup_n \sup_{x \in [\zeta_n, \zeta'_n]} |\tilde{\psi}'_n(x)| \leq \sup_{x \in [\zeta_n, \zeta'_n]} |(\log \widehat{h}_n)'(\bar{\theta}_n + x)| + \sup_{x \in [\zeta_n, \zeta'_n]} |(\log \widehat{h}_n)'(\bar{\theta}_n - x)|,$$

which is bounded above by (6.65). Finally, when $\tilde{g}_n = (\tilde{g}_n^{sym})^{sm}$, from (6.15) we observe that $\tilde{\psi}'_n$ can be expressed in terms of \widehat{h}_n^{sm} as

$$\tilde{\psi}'_n(x) = \varrho_n^{sm}(x) \left((\log \widehat{h}_n^{sm})'(\bar{\theta}_n + x) \right) - (1 - \varrho_n^{sm}(x)) \left((\log \widehat{h}_n^{sm})'(\bar{\theta}_n - x) \right),$$

for some $\varrho_n^{sm}(x) < 1$. Boundedness of $\tilde{\psi}'_n$ then follows from (6.66). \square

6.5. Additional lemmas for the MLE

Lemma 12. *Suppose $e^\psi \in \mathcal{SLC}_0$. Then $\psi(x) \leq -\log |2x|$ for all $x \in \mathbb{R}$.*

Proof. Note that since $e^\psi \in \mathcal{S}\mathcal{L}\mathcal{C}_0$, for $x > 0$, we have,

$$1 = \int_{-\infty}^{\infty} e^{\psi(z)} dz \geq \int_{-x}^x e^{\psi(z)} dz,$$

which is bounded below by $2e^{\psi(x)}x$ because $\psi \in \mathcal{S}\mathcal{C}_0$. Hence the result follows. \square

Lemma 13. *Suppose F is a distribution function satisfying Condition A. Denote*

$$\text{int}(J(F)) = (a, b),$$

for some $a, b \in [-\infty, \infty]$. Suppose ψ_θ is the maximizer of $\Psi(\theta, \psi, F)$ over $\psi \in \mathcal{S}\mathcal{C}_0$ for some $\theta \in \mathbb{R}$. Then

$$\text{int}(\text{dom}(\psi_\theta)) = (-d, d),$$

where $d = (b - \theta) \vee (\theta - a)$.

Note that a or b can be ∞ , in which case $d = \infty$, and $\text{int}(\text{dom}(\psi_\theta)) = \mathbb{R}$.

Proof. Note that $\text{int}(\text{dom}(\psi_\theta)) = (-c, c)$ for some $c > 0$. We first show that $c \geq d$. Consider the set $A = \overline{(-d, d)} \setminus \text{dom}(\psi_\theta)$. Note that unless $d = \infty$, $\overline{(-d, d)} = [-d, d]$. Letting Δ denote the indicator function 1_A , we observe that $\psi_\theta + \Delta = \psi_\theta$. Also, note that

$$\psi_\theta = \arg \max_{\psi \in \mathcal{S}\mathcal{C}_0} \Psi(0, \psi, F(\cdot + \theta)) = \arg \max_{\psi \in \mathcal{S}\mathcal{C}_0} \omega(\psi, F(\cdot + \theta)),$$

which follows from (1.4) and (3.7). Hence, Proposition 5(ii) of Xu and Samworth (2019) leads to

$$\int_{-\infty}^{\infty} \Delta(x) dF(x + \theta) \leq \int_{-\infty}^{\infty} \Delta(x) e^{\psi_\theta(x)} dx.$$

Since $\Delta = 0$ on $\text{dom}(\psi_\theta)$, it follows that

$$\int_A dF(x + \theta) = 0,$$

implying $(a, b) \cap (A + \theta) = \emptyset$.

It is easy to see that if either a or b equals ∞ , $d = \infty$, and $A + \theta = \mathbb{R} \setminus \text{dom}(\psi_\theta)$. Therefore,

$$(a, b) \cap (A + \theta) = (a, b) \setminus \left(\text{dom}(\psi_\theta) + \theta \right) = \emptyset,$$

which indicates that

$$(a, b) \subset (-c, c) + \theta, \tag{6.67}$$

implying $(-c, c) = \mathbb{R}$. Hence $(-c, c) = (-d, d)$, which completes the proof for this case. Therefore, it only remains to prove Lemma 13 for the case when both a and b are finite.

When $a, b < \infty$, there can be two sub-cases. In the first case, $b - \theta \geq \theta - a$, which indicates $d = b - \theta$. In this case, $[-d, d] + \theta = [2\theta - b, b]$, and

$$[a, b] \subset [2\theta - b, b] = [-d, d] + \theta,$$

leading to

$$(a, b) \setminus \left(\text{dom}(\psi_\theta) + \theta \right) = (a, b) \cap (A + \theta) = \emptyset,$$

from which (6.67) follows. Therefore, $b \leq c + \theta$, or $d \leq c$. In the second case $b - \theta \leq \theta - a$, $d = \theta - a$ and $[-d, d] + \theta = [a, 2\theta - a]$. Proceeding similarly as in the first case, we can again show that $c \geq d$. Hence, we have established that $c \geq d$ in general.

Now suppose $c > d$, which implies that $a, b < \infty$. Observe that in this case, there exists a $\psi \in \mathcal{SC}_0$, which equals ψ_θ on $(-d, d)$ and $-\infty$ everywhere else. It is easy to verify that when $c > d$,

$$\begin{aligned} & \Psi(\theta, \psi_\theta, F) - \Psi(\theta, \psi, F) \\ &= \int_{-d}^d e^{\psi_\theta(x)} dx - \int_{-c}^c e^{\psi_\theta(x)} dx < 0. \end{aligned}$$

However, ψ_θ satisfies $\Psi(\theta, \psi_\theta, F) \geq \Psi(\theta, \psi, F)$ for all $\psi \in \mathcal{SC}_0$, which leads to a contradiction. Hence, we conclude that $c = d$, which completes the proof. \square

Lemma 14. *Suppose F satisfies Condition A and (3.20). Then*

$$\arg \max_{\phi \in \mathcal{C}} \omega(\phi, F) = \arg \max_{\phi \in \mathcal{SC}_\theta} \omega(\phi, F),$$

where the criterion function ω is defined in (3.7).

Proof. First we consider the special case $\theta = 0$. For $\phi \in \mathcal{C}$ and F symmetric about 0, using (3.20) with $\theta = 0$, the criterion function $\omega(\phi, F)$ can be written as

$$\begin{aligned} & \int_{-\infty}^0 \phi(x) dF(x) + \int_0^{\infty} \phi(x) dF(x) - \int_{-\infty}^{\infty} e^{\phi(x)} dx \\ &= - \int_0^{\infty} \phi(-x) dF(-x) + \int_0^{\infty} \phi(x) dF(x) - \int_{-\infty}^{\infty} e^{\phi(x)} dx \\ &= \int_0^{\infty} \phi(-x) dF(x) + \int_0^{\infty} \phi(x) dF(x) - \int_{-\infty}^{\infty} e^{\phi(x)} dx \\ &= \int_{-\infty}^{\infty} \frac{\phi(-x) + \phi(x)}{2} dF(x) - \int_{-\infty}^{\infty} e^{\phi(x)} dx, \end{aligned}$$

which is not larger than

$$\int_{-\infty}^{\infty} \frac{\phi(x) + \phi(-x)}{2} dF(x) - \int_{-\infty}^{\infty} e^{(\phi(x) + \phi(-x))/2} dx,$$

by the convexity of the exponential function. This proves that a $\phi \in \mathcal{SC}_0$ maximizes $\omega(\phi, F)$ over \mathcal{C} when F is symmetric about 0. The proof for a general θ follows in a similar way. \square

Lemma 15. *For any $\psi \in \mathcal{SC}_0$ with nonempty domain, and each $\epsilon > 0$, there exists a function $\psi^{(\epsilon)} \in \mathcal{SC}_0$ for each $\epsilon > 0$ such that the following conditions are met:*

$$(A) \quad \psi(x) \leq \psi^{(\epsilon)}(x) \leq \psi(0) \text{ for all } x \in \mathbb{R}.$$

$$(B)$$

$$\psi^{(\epsilon)}(x) \geq -|x|/\epsilon + \psi(0),$$

for all $x \in \mathbb{R}$.

$$(B) \quad \psi^{(\epsilon)}(x) \downarrow \psi(x) \text{ for each } x \in \mathbb{R} \text{ as } \epsilon \downarrow 0.$$

Proof. Following the construction in Lemma 4.3 of [Dümbgen et al. \(2011\)](#) we set

$$\psi^{(\epsilon)}(x) = \inf_{(t,c) \in A} (tx + c), \quad x \in \mathbb{R}, \quad (6.68)$$

where

$$A = \left\{ (t, c) \in \mathbb{R} \times \mathbb{R} \mid |t| \leq 1/\epsilon, tx + c \geq \psi(x) \text{ for all } x \in \mathbb{R} \right\}.$$

Note that $\psi^{(\epsilon)}(x)$ is concave by Lemma 4.3 of [Dümbgen et al. \(2011\)](#). To show that $\psi^{(\epsilon)}$ is symmetric about 0, observe that if $tx + c \geq \psi(x)$ for all $x \in \mathbb{R}$, the symmetry of ψ about 0 leads to

$$-t(-x) + c \geq \psi(-x) \text{ for all } x \in \mathbb{R},$$

which can be rewritten as

$$-tx + c \geq \psi(x) \text{ for all } x \in \mathbb{R}.$$

Hence, $(t, c) \in A$ implies that $(-t, c) \in A$. As a result,

$$\inf_{(t,c) \in A} (tx + c) = \inf_{(-t,c) \in A} (tx + c) = \inf_{(t,c) \in A} (-tx + c),$$

which implies that $\psi^{(\epsilon)} \in \mathcal{SC}_0$. That $\psi \leq \psi^{(\epsilon)}$ follows from Lemma 4.3 of [Dümbgen et al. \(2011\)](#). Hence, to prove part A, we need to show that $\psi^{(\epsilon)}$ is bounded above by $\psi(0)$.

Observe that since $\psi^{(\epsilon)} \in \mathcal{SC}_0$, it is bounded above by $\psi^{(\epsilon)}(0)$, which, by [\(6.68\)](#), equals the infimum of the set

$$A' = \{c : (t, c) \in A \text{ for some } t \in \mathbb{R}\}.$$

However, since $c \in A'$ satisfies $tx + c \geq \psi(x)$ for all $x \in \mathbb{R}$, and some t , we conclude that Thus we obtain

$$\psi^{(\epsilon)}(0) = \inf A' \geq \psi(0). \quad (6.69)$$

In light of (6.69), to prove part A of the current lemma, it suffices to show that $\psi(0) \in A'$. Since $\psi \in \mathcal{SC}_0$, for $x > 0$, we have $\psi(x) \leq \psi'(0+)x + \psi(0)$. When $x \leq 0$, using the fact that $\psi'(0+) \leq 0$, we calculate

$$\psi(x) = \psi(-x) \leq -\psi'(0+)x + \psi(0) \leq \psi'(0+)x + \psi(0).$$

Therefore, for all $x \in \mathbb{R}$, the following holds:

$$\psi(x) \leq \psi'(0+)x + \psi(0).$$

When $-\psi'(0+) \leq 1/\epsilon$, (6.68) thus indicates that $(-\psi'(0+), \psi(0)) \in A$. In case $-\psi'(0+) > 1/\epsilon$, we still have

$$\psi(x) \leq \psi'(0+)x + \psi(0) < -x/\epsilon + \psi(0), \quad \text{for all } x > 0.$$

For $x \leq 0$,

$$\psi(x) = \psi(-x) \leq x/\epsilon + \psi(0) \leq -x/\epsilon + \psi(0).$$

Therefore, it turns out that if $-\psi(0+) > 1/\epsilon$,

$$\psi(x) \leq -x/\epsilon + \psi(0) \text{ for all } x \in \mathbb{R},$$

which implies that $(1/\epsilon, \psi(0)) \in A$. The above calculations indicate that there always exists a $t \in \mathbb{R}$ such that $(t, \psi(0)) \in A$, entailing that $\psi(0) \in A'$. This fact, combined with (6.69), yields that $\psi^{(\epsilon)}(0) = \psi(0)$. Part A follows from (6.69).

Note that (6.68) and (6.69) imply that

$$\psi^{(\epsilon)}(x) \geq -|x|/\epsilon + \psi(0), \quad x \in \mathbb{R},$$

which settles the proof of part B.

Lemma 15(C) follows directly from Lemma 4.3 of Dümbgen *et al.* (2011). \square

The next lemma is Pratt's lemma (Pratt, 1960, Theorem 1). We state it here for convenience.

Lemma 16. *Suppose $(\Omega, \mathcal{F}, \mu)$ is a measure space and a_n, b_n, c_n are sequences of functions on Ω converging almost everywhere to functions a, b, c respectively. Also, all functions are integrable and $\int a_n d\mu \rightarrow \int a d\mu$ and $\int c_n d\mu \rightarrow \int c d\mu$. Moreover, $a_n \leq b_n \leq c_n$. Then*

$$\int b_n d\mu \rightarrow \int b d\mu.$$

References

- Beran, R. (1974). Asymptotically efficient adaptive rank estimates in location models. *Ann. Statist.*, **2**, 63–74.
- Bhattacharyya, S. and Bickel, P. J. (2013). Adaptive estimation in elliptical distributions with extensions to high dimensions by.

- Bickel, P. J., Klaassen, C. A. J., Ritov, Y., and Wellner, J. A. (1998). *Efficient and Adaptive Estimation for Semiparametric Models*. Springer-Verlag, New York.
- Birgé, L. (1997). Estimation of unimodal densities without smoothness assumptions. *Ann. Statist.*, **25**, 970–981.
- Chen, Y. and Samworth, R. J. (2013). Smoothed log-concave maximum likelihood estimation with applications. *Statist. Sinica*, **23**, 1373–1398.
- Chen, Y. and Samworth, R. J. (2016). Generalized additive and index models with shape constraints. *J R Stat Soc Series B Stat Methodol*, **78**, 729–754.
- Cule, M. and Samworth, R. (2010). Theoretical properties of the log-concave maximum likelihood estimator of a multidimensional density. *Electron. J. Statist.*, **4**, 254–270.
- Devroye, L. (1987). *A course in density estimation*. Progress in probability and statistics. Birkhäuser.
- Doss, C. R. and Wellner, J. A. (2016). Global rates of convergence of the MLEs of log-concave and s -concave densities. *Ann. Statist.*, **44**, 954–981.
- Doss, C. R. and Wellner, J. A. (2019). Univariate log-concave density estimation with symmetry or modal constraints. *Electron. J. Stat.*, **25**, 2391–2461.
- Dümbgen, L. and Rufibach, K. (2009). Maximum likelihood estimation of a log-concave density and its distribution function: Basic properties and uniform consistency. *Bernoulli*, **15**, 40–68.
- Dümbgen, L., Samworth, R., and Schuhmacher, D. (2011). Approximation by log-concave distributions, with applications to regression. *Ann. Statist.*, **39**, 702–730.
- Groeneboom, P. and Hendrickx, K. (2018). Current status linear regression. *Ann. Statist.*, **46**, 1415–1444.
- Hodges, J. L. and Lehmann, E. L. (1963). Estimates of location based on rank tests. *Ann. Math. Statist.*, **34**, 598–611.
- Huber, P. J. (1964). Robust estimation of a location parameter. *Ann. Math. Statist.*, **35**, 73–101.
- Kim, A. K. H. and Samworth, R. J. (2016). Global rates of convergence in log-concave density estimation. *Ann. Statist.*, **44**, 2756–2779.
- Kuchibhotla, A. K., Patra, R. K., and Sen, B. (2017). Efficient estimation in convex single index models. *arXiv:1708.00145v2*.
- Murphy, S. A., Van der Vaart, A., and Wellner, J. (1999). Current status regression. *Math. Methods Statist.*, **8**, 407–425.
- Pal, J. K., Woodroffe, M., and Meyer, M. (2007). Estimating a pólya frequency function₂. *Lecture Notes-Monograph Series*, **54**, 239–249.
- Pratt, J. W. (1960). On interchanging limits and integrals. *Ann. Math. Statist.*, **31**, 74–77.
- Rockafellar, R. T. (1970). *Convex Analysis*. Princeton University Press.
- Sacks, J. (1975). An asymptotically efficient sequence of estimators of a location parameter. *Ann. Statist.*, **3**, 285–298.
- Seijo, E. and Sen, B. (2011). Nonparametric least squares estimation of a multivariate convex regression function. *Ann. Statist.*, **39**, 1633–1657.
- Shorack, G. R. (2000). *Probability for Statisticians*. Springer.

- Silverman, B. W. (1982). On the estimation of a probability density function by the maximum penalized likelihood method. *Ann. Statist.*, pages 795–810.
- Stein, C. (1956). Efficient nonparametric testing and estimation. *Proc. Third Berkley Symp. Math. Statist. Prob.*, **1**, 187–196.
- Stone, C. J. (1975). Adaptive maximum likelihood estimators of a location parameter. *Ann. Statist.*, **3**, 267–284.
- Takeuchi, K. (1975). A survey of robust estimation of location: models and procedures, especially in case of measurement of a physical quantity. *Bull. Inst. Internat. Statist.*, **46**, 336–348. With discussion.
- Van der Vaart, A. (1998). *Asymptotic Statistics*. Asymptotic Statistics. Cambridge University Press.
- Van der Vaart, A. W. and Wellner, J. A. (1996). *Weak Convergence and Empirical Processes*. Springer, New York.
- van der Vaart, A. W. and Wellner, J. A. W. (2007). Empirical processes indexed by estimated functions. *Asymptotics: Particles, Processes and Inverse Problems*, **55**, 234–252.
- Van Eeden, C. (1970). Efficiency-robust estimation of location. *Ann. Math. Statist.*, **41**, 172–181.
- Villani, C. (2003). *Topics in optimal transportation*, volume 58 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI.
- Villani, C. (2009). *Optimal transport; Old and New*, volume 338 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin. Old and new.
- Xu, M. and Samworth, R. J. (2019). High-dimensional nonparametric density estimation via symmetry and shape constraints. <https://arxiv.org/abs/1903.06092v1>.