

Colourings of star systems

Iren Darijani and David A. Pike
 Department of Mathematics and Statistics
 Memorial University of Newfoundland
 St. John's, NL, Canada. A1C 5S7

July 27, 2021

Abstract

An e -star is a complete bipartite graph $K_{1,e}$. An e -star system of order $n > 1$, $S_e(n)$, is a partition of the edges of the complete graph K_n into e -stars. An e -star system is said to be k -colourable if its vertex set can be partitioned into k sets (called colour classes) such that no e -star is monochromatic. The system $S_e(n)$ is k -chromatic if $S_e(n)$ is k -colourable but is not $(k - 1)$ -colourable. If every k -colouring of an e -star system can be obtained from some k -colouring ϕ by a permutation of the colours, we say that the system is uniquely k -colourable. In this paper, we first show that for any integer $k \geq 2$, there exists a k -chromatic 3-star system of order n for all sufficiently large admissible n . Next, we generalize this result for e -star systems for any $e \geq 3$. We show that for all $k \geq 2$ and $e \geq 3$, there exists a k -chromatic e -star system of order n for all sufficiently large n such that $n \equiv 0, 1 \pmod{2e}$. Finally, we prove that for all $k \geq 2$ and $e \geq 3$, there exists a uniquely k -chromatic e -star system of order n for all sufficiently large n such that $n \equiv 0, 1 \pmod{2e}$.

Key words: star system; decomposition; colouring; chromatic number; unique colouring

AMS subject classifications: 05B30, 05C51, 05C15

1 Introduction

A G -decomposition of a graph H is a pair (V, \mathcal{B}) where V is the set of vertices of K and \mathcal{B} is a set of subgraphs of K , each isomorphic to G , whose edge sets partition the edge set of K . A G -design of order n is a G -decomposition of the complete graph K_n on n vertices. A *complete graph* is a simple graph in which every pair of distinct vertices is connected by a unique edge. A G -design (V, \mathcal{B}) is said to be *weakly k -colourable* if its vertex set can be partitioned into k sets (called colour classes) such that no subgraph belonging to \mathcal{B} is monochromatic. The G -design is *k -chromatic* if it is k -colourable but is not $(k - 1)$ -colourable. If a G -design is k -chromatic, we say that its chromatic number is k . A colouring of a G -design is said to be *equitable* if the cardinalities of the colour classes differ by at most one. It is *strongly equitable* if the colour classes are of the same size. A G -design will be called *(strongly) equitably k -chromatic* if it is

k -chromatic and admits a (strongly) equitable k -colouring. If every k -colouring of a G -design (V, \mathcal{B}) can be obtained from some k -colouring ϕ by a permutation of the colours, we say that (V, \mathcal{B}) has a *unique* k -colouring, or that (V, \mathcal{B}) is *uniquely* k -colourable.

Weak colourings of G -designs were first studied for K_3 -designs, i.e., Steiner triple systems. In particular, de Brandes, Phelps and Rödl [2] proved that for any $k \geq 3$, there exists some integer v_k such that for all admissible $v \geq v_k$, there is a k -chromatic Steiner triple system of order v . Moreover, in [3] Burgess and Pike showed that for all $k \geq 2$ and even $m \geq 4$ there exists a k -chromatic m -cycle system. An m -cycle system of order $n > 1$ is a partition of the edges of the complete graph K_n into m -cycles. Also, Horsley and Pike showed that for all $k \geq 2$ and $m \geq 3$ with $(k, m) \neq (2, 3)$, there exist k -chromatic m -cycle systems of all admissible orders greater than or equal to some integer $n_{k,m}$ [5].

A bipartite graph $G_{m,n}$ is a graph whose vertex set can be partitioned into two subsets V_1 and V_2 with m and n vertices each, such that every edge of $G_{m,n}$ has one vertex in V_1 and one vertex in V_2 . If $G_{m,n}$ contains every edge joining V_1 and V_2 , then it is called a *complete bipartite graph* and denoted by $K_{m,n}$. For an integer $e \geq 1$, an e -star is a complete bipartite graph $K_{1,e}$. An e -star with star centre x and edges $\{x, a_1\}, \{x, a_2\}, \dots, \{x, a_e\}$ is denoted by either $\{x; a_1, \dots, a_e\}$ or $\{x; A\}$ where $A = \{a_1, \dots, a_e\}$. When G is an e -star, a G -design of order n is called an e -star system of order n and denoted by $S_e(n)$. The necessary and sufficient conditions for the existence of an e -star system of order n are that e divides $\frac{n(n-1)}{2}$ and $n \geq 2e$ [6]. A positive integer n is said to be *admissible* if there exists an e -star system of order n .

A G -factor of a graph H is a spanning subgraph of H , each component of which is isomorphic to G . If the edges of H can be partitioned into G -factors, then we say H has a G -factorization. The existence of a $K_{1,e}$ -factorization of K_n was studied and completely settled in 1993 [7]. If e is even, a $K_{1,e}$ -factorization of K_n exists if and only if $n \equiv 1 \pmod{2e}$ and $n \equiv 0 \pmod{e+1}$. If e is odd, there does not exist a $K_{1,e}$ -factorization of K_n for any n .

In this paper, we investigate k -colourings of e -star systems for $e \geq 3$. Note that a 3-star system of order n exists if and only if $n \equiv 0, 1 \pmod{3}$ and $n \geq 6$. In Section 2, we first construct equitably 2-chromatic 3-star systems of all admissible orders. We then show that if there exists a k -chromatic 3-star system of order n_0 , then there exists a k -chromatic 3-star system of order n for all admissible $n > n_0$ and $k \geq 2$. Next, from a $(k-1)$ -chromatic 3-star system, we construct a k -chromatic 3-star system for all $k \geq 3$. Finally, we finish this section by showing that for any integer $k \geq 2$, there exists some integer n_k such that for all admissible $n \geq n_k$, there exists a k -chromatic 3-star system of order n .

In Section 3, we generalize the results in Section 2 for e -star systems for all $e \geq 3$. We first construct equitably 2-chromatic e -star systems of order $2e$ for all $e \geq 3$. We then show that if there exists a k -chromatic e -star system of order n_0 such that $n_0 \equiv 0, 1 \pmod{2e}$, then there exists a k -chromatic e -star system for all $n > n_0$ such that $n \equiv 0, 1 \pmod{2e}$ for all $k \geq 2$ and $e \geq 4$. Next, from a $(k-1)$ -chromatic e -star system of order $n_{k-1} \equiv 0 \pmod{2e}$, we construct a k -chromatic e -star system of order $n_k \equiv 0 \pmod{2e}$ for all $k \geq 3$. Finally, we finish this section by showing that for any integer $k \geq 2$, there exists some integer n_k where $n_k \equiv 0 \pmod{2e}$ such that for all $n \geq n_k$ where $n \equiv 0, 1 \pmod{2e}$, there exists a k -chromatic e -star system of order n .

We also study unique colourings for e -star systems. The idea of uniquely colourable designs has already arisen in the content of Steiner triple systems. In 2003, Forbes showed that for every admissible $v \geq 25$, there exists a 3-balanced Steiner triple system with a unique 3-colouring and also a Steiner triple system which has a unique, nonequitable 3-colouring [4], where a Steiner triple system is said to be *3-balanced* if every 3-colouring of it is equitable. In Section 4, we first construct a strongly equitable uniquely 2-chromatic e -star system. Next, we construct a strongly equitable k -chromatic e -star system from a strongly equitable uniquely $(k-1)$ -chromatic e -star system. We then show how to construct a strongly equitable uniquely k -chromatic e -star system from a strongly equitable k -chromatic e -star system. Finally, we finish this section by showing that for any integer $k \geq 2$, there exists some integer n_k where $n_k \equiv 0 \pmod{2e}$ such that for all $n \geq n_k$ where $n \equiv 0, 1 \pmod{2e}$, there exists a uniquely k -chromatic e -star system of order n .

2 k -colourings of 3-star systems

We initially concentrate on 3-star systems and we show that for any integer $k \geq 2$, there exists some integer n_k such that for all admissible $n \geq n_k$, there exists a k -chromatic 3-star system of order n . We first construct an equitably 2-chromatic 3-star system of order n for all admissible n .

Theorem 2.1 *For each admissible order n , there exists an equitably 2-chromatic 3-star system of order n .*

Proof. We first construct an equitably 2-chromatic 3-star system $S_3(6)$. Let $V = \{1, 2, 3, 4, 5, 6\}$, $\mathcal{B} = \{\{1; 3, 5, 6\}, \{2; 1, 3, 6\}, \{4; 1, 2, 3\}, \{5; 2, 3, 4\}, \{6; 3, 4, 5\}\}$, $R = \{1, 3, 5\}$, and $Y = \{2, 4, 6\}$. Then (V, \mathcal{B}) is an equitably 2-chromatic 3-star system of order six with colour classes R and Y .

Now, suppose that there exists an equitably 2-chromatic 3-star system $S_3(3t)$, (V, \mathcal{B}) , where $t \geq 2$, $V = \{1, \dots, 3t\}$ is the set of points which is partitioned into two subsets R and Y and

\mathcal{B} is the set of blocks. Without loss of generality, if $3t$ is odd, we can let $R = \{1, 3, 5, \dots, 3t\}$, $Y = \{2, 4, 6, \dots, 3t-1\}$ and if $3t$ is even, $R = \{1, 3, 5, \dots, 3t-1\}$, $Y = \{2, 4, 6, \dots, 3t\}$.

Next, we construct an equitably 2-chromatic 3-star system $S_3(3t+1)$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{3t+1\}$ and $\hat{\mathcal{B}} = \mathcal{B} \cup \{\{3t+1; 1, 2, 3\}, \{3t+1; 4, 5, 6\}, \dots, \{3t+1; 3t-2, 3t-1, 3t\}\}$. Let $R = \{1, 3, 5, \dots, 3t\}$, $Y = \{2, 4, 6, \dots, 3t+1\}$ if $3t$ is odd and $R = \{1, 3, 5, \dots, 3t+1\}$, $Y = \{2, 4, 6, \dots, 3t\}$ if $3t$ is even. Then $(\hat{V}, \hat{\mathcal{B}})$ is an equitably 2-chromatic 3-star system of order $3t+1$ with colour classes R and Y .

Finally, we construct an equitably 2-chromatic 3-star system $S_3(3t+3)$, $(\tilde{V}, \tilde{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\tilde{V} = V \cup \{3t+1, 3t+2, 3t+3\}$ and $\tilde{\mathcal{B}} = \mathcal{B} \cup \{\{3t+1; 1, 2, 3\}, \dots, \{3t+1; 3t-5, 3t-4, 3t-3\}, \{3t+2; 1, 2, 3\}, \dots, \{3t+2; 3t-5, 3t-4, 3t-3\}, \{3t+3; 1, 2, 3\}, \dots, \{3t+3; 3t-5, 3t-4, 3t-3\}, \{3t+1; 3t-2, 3t-1, 3t+2\}, \{3t+2; 3t-2, 3t-1, 3t+3\}, \{3t+3; 3t-2, 3t-1, 3t+1\}, \{3t; 3t+1, 3t+2, 3t+3\}\}$. Let R be the set of odd elements of \tilde{V} and Y be the set of even elements of \tilde{V} . Then $(\tilde{V}, \tilde{\mathcal{B}})$ is an equitably 2-chromatic 3-star system of order $3t+3$ with colour classes R and Y . \square

We now show how to construct a k -chromatic 3-star system from a smaller k -chromatic 3-star system.

Theorem 2.2 *Let $k \geq 2$. If there exists a k -chromatic 3-star system of order n_0 , then there exists a k -chromatic 3-star system of order n for all admissible $n > n_0$.*

Proof. Suppose that there exists a k -chromatic 3-star system of order n_0 , (V, \mathcal{B}) , where $V = \{1, \dots, n_0\}$ is the set of points and \mathcal{B} is the set of blocks. Given a k -colouring of (V, \mathcal{B}) with colours $1, 2, \dots, k$ and colour classes C_1, C_2, \dots, C_k , let $R = \bigcup_{i=1}^{\ell} C_i$ and $Y = \bigcup_{i=\ell+1}^k C_i$ for some integer ℓ such that $1 \leq \ell < k$. Let $r_1, r_2, \dots, r_{|R|}$ be the elements of R and $y_1, y_2, \dots, y_{|Y|}$ be the elements of Y . Observe that R and Y form a partition of V . Without loss of generality assume that $|R| \geq |Y|$.

Case 1. Suppose that $n_0 \equiv 0 \pmod{3}$. Then $n_0 = 3t$, $t \geq 2$.

First, we construct a k -chromatic $S_3(3t+1)$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{3t+1\}$.

If $|R| = |Y|$, let $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}$ where \mathcal{T} is the set $\{\{3t+1; r_1, r_2, y_1\}, \{3t+1; y_2, y_3, r_3\}, \dots, \{3t+1; r_{|R|-2}, r_{|R|-1}, y_{|Y|-2}\}, \{3t+1; y_{|Y|-1}, y_{|Y|}, r_{|R|}\}\}$.

Otherwise, $|R| > |Y|$. If $|Y|$ is even then let $m = \frac{|Y|}{2}$ and $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}$ where \mathcal{T} is the set $\{\{3t+1; y_1, y_2, r_1\}, \dots, \{3t+1; y_{|Y|-1}, y_{|Y|}, r_m\}, \{3t+1; r_{m+1}, r_{m+2}, r_{m+3}\}, \dots, \{3t+1; r_{|R|-2}, r_{|R|-1}, r_{|R|}\}\}$. If $|Y|$ is odd then let $m = \frac{|Y|+3}{2}$ and $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}$ where \mathcal{T} is the set $\{\{3t+1; y_1, y_2, r_1\}, \dots, \{3t+1; y_{|Y|-2}, y_{|Y|-1}, r_{m-2}\}, \{3t+1; y_{|Y|}, r_{m-1}, r_m\}, \{3t+1; r_{m+1}, r_{m+2}, r_{m+3}\}, \dots, \{3t+1; r_{|R|-2}, r_{|R|-1}, r_{|R|}\}\}$.

Note that $(\hat{V}, \hat{\mathcal{B}})$ is not $(k-1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1, \dots, C_{k-1}, C_k \cup \{3t+1\}$ are the colour classes of a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic 3-star system of order $3t+1$.

Next, we construct a k -chromatic $S_3(3t+3)$, $(\tilde{V}, \tilde{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\tilde{V} = V \cup \{3t+1, 3t+2, 3t+3\}$ and $R' = R \setminus \{r_{|R|-2}, r_{|R|-1}, r_{|R|}\}$. Note that $|R| \geq 3$ since $n_0 \geq 6$ and $|R| \geq |Y|$.

Let \mathcal{T} be the set $\{\{3t+1; r_{|R|-2}, r_{|R|-1}, 3t+2\}, \{3t+2; r_{|R|-2}, r_{|R|-1}, 3t+3\}, \{3t+3, r_{|R|-2}, r_{|R|-1}, 3t+1\}, \{r_{|R|}; 3t+1, 3t+2, 3t+3\}\}$.

If $|R'| = |Y|$, let $\tilde{\mathcal{B}} = \mathcal{B} \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=3} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; r_1, r_2, y_1\}, \{3t+i; y_2, y_3, r_3\}, \dots, \{3t+i; r_{|R|-5}, r_{|R|-4}, y_{|Y|-2}\}, \{3t+i; y_{|Y|-1}, y_{|Y|}, r_{|R|-3}\}\}$ for $i \in \{1, 2, 3\}$.

If $|R'| > |Y|$ and $|Y|$ is even, then let $m = \frac{|Y|}{2}$ and $\tilde{\mathcal{B}} = \mathcal{B} \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=3} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; y_1, y_2, r_1\}, \dots, \{3t+i; y_{|Y|-1}, y_{|Y|}, r_m\}, \{3t+i; r_{m+1}, r_{m+2}, r_{m+3}\}, \dots, \{3t+i; r_{|R|-5}, r_{|R|-4}, r_{|R|-3}\}\}$ for $i \in \{1, 2, 3\}$. The edges can be decomposed into 3-stars in a similar manner when $|R'| > |Y|$ and $|Y|$ is odd.

Otherwise, $|R'| < |Y|$. Then either $|R'| = |Y| - 1$ or $|R'| = |Y| - 2$ or $|R'| = |Y| - 3$. If $|R'| = 0$, then $|R| = |Y| = 3$. Without loss of generality, we may assume that $Y = \{1, 2, 3\}$ and $R = \{4, 5, 6\}$. Now let $\tilde{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}_0$ where \mathcal{T}_0 is the set $\{\{7; 1, 2, 4\}, \{8; 1, 2, 4\}, \{9; 1, 2, 4\}, \{7; 8, 3, 5\}, \{8; 9, 3, 5\}, \{9; 7, 3, 5\}, \{6; 7, 8, 9\}\}$. If $|R'| \geq 1$ and $|R'| = |Y| - 1$, let $\tilde{\mathcal{B}} = \mathcal{B} \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=3} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; r_1, y_1, y_2\}, \{3t+i; r_2, r_3, y_3\}, \{3t+i; y_4, y_5, r_4\}, \dots, \{3t+i; r_{|R|-2}, r_{|R|-1}, y_{|Y|-2}\}, \{3t+i; y_{|Y|-1}, y_{|Y|}, r_{|R|}\}\}$, for $i \in \{1, 2, 3\}$. The edges can be decomposed into 3-stars in a similar manner when $1 \leq |R'| < |Y|$ and $|R'| = |Y| - j$ for $j \in \{2, 3\}$.

Note that $(\tilde{V}, \tilde{\mathcal{B}})$ is not $(k-1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1, \dots, C_{k-1}, C_k \cup \{3t+1, 3t+2, 3t+3\}$ are the colour classes of a k -colouring of $(\tilde{V}, \tilde{\mathcal{B}})$. Therefore, $(\tilde{V}, \tilde{\mathcal{B}})$ is a k -chromatic 3-star system of order $3t+3$.

Case 2. Suppose that $n_0 \equiv 1 \pmod{3}$. Then $n_0 = 3t+1$, $t \geq 2$. We construct a k -chromatic $S_3(3t+3)$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{3t+2, 3t+3\}$. Note that $|R| \geq 4$ since $n_0 \geq 7$ and $|R| \geq |Y|$. So without loss of generality we may assume that $\{3t, 3t+1\} \subset R$. Moreover, since the edge $\{3t, 3t+1\}$ must be in some 3-star, we may also assume that there exists a 3-star $S = \{3t+1; 3t, 3t-1, 3t-2\} \in \mathcal{B}$. We will dismantle the 3-star S in each of the following subcases:

Case 2.1. All the vertices of the 3-star S belong to the set R . Hence R is the union of $\ell \geq 2$ distinct colour classes. Let $R' = R \setminus \{3t+1, 3t, 3t-1, 3t-2\}$.

Let \mathcal{T} be the set $\{\{3t+2; 3t+1, 3t, 3t-1\}, \{3t+3; 3t+2, 3t, 3t-1\}, \{3t-2; 3t+2, 3t+1, 3t+3\},$

$\{3t+1; 3t+3, 3t, 3t-1\}\}$.

If $|R'| = |Y|$, let $\hat{\mathcal{B}} = (\mathcal{B} \setminus \{S\}) \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=2} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; r_1, r_2, y_1\}, \{3t+i; y_2, y_3, r_3\}, \dots, \{3t+i; r_{|R|-6}, r_{|R|-5}, y_{|Y|-2}\}, \{3t+i; y_{|Y|-1}, y_{|Y|}, r_{|R|-4}\}\}$ for $i \in \{1, 2\}$.

If $|R'| > |Y|$ and $|Y|$ is even, then let $m = \frac{|Y|}{2}$ and $\hat{\mathcal{B}} = (\mathcal{B} \setminus \{S\}) \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=2} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; y_1, y_2, r_1\}, \dots, \{3t+i; y_{|Y|-1}, y_{|Y|}, r_m\}, \{3t+i; r_{m+1}, r_{m+2}, r_{m+3}\}, \dots, \{3t+i; r_{|R|-6}, r_{|R|-5}, r_{|R|-4}\}\}$ for $i \in \{1, 2\}$. The edges can be decomposed into 3-stars in a similar manner when $|R'| > |Y|$ and $|Y|$ is odd.

Otherwise $|R'| < |Y|$. Then either $|R'| = |Y| - 1$ or $|R'| = |Y| - 2$ or $|R'| = |Y| - 3$ or $|R'| = |Y| - 4$. If $|R'| = 0$, then $Y = \{1, 2, 3\}$ and $R = \{4, 5, 6, 7\}$. Let $\hat{\mathcal{B}} = (\mathcal{B} \setminus \{S\}) \cup \mathcal{T}_0$ where \mathcal{T}_0 is the set $\{\{8; 1, 2, 3\}, \{9; 1, 2, 3\}, \{8; 7, 6, 5\}, \{9; 8, 6, 5\}, \{4; 8, 7, 9\}, \{7; 9, 6, 5\}\}$. If $|R'| \geq 1$ and $|R'| = |Y| - 4$, let $\hat{\mathcal{B}} = (\mathcal{B} \setminus \{S\}) \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=2} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; r_1, y_1, y_2\}, \{3t+i; r_2, y_3, y_4\}, \{3t+i; r_3, y_5, y_6\}, \{3t+i; r_4, y_7, y_8\}, \{3t+i; r_5, r_6, y_9\}, \{3t+i; y_{10}, y_{11}, r_7\}, \dots, \{3t+i; r_{|R|-6}, r_{|R|-5}, y_{|Y|-2}\}, \{3t+i; y_{|Y|-1}, y_{|Y|}, r_{|R|-4}\}\}$ for $i \in \{1, 2\}$. The edges can be decomposed into 3-stars in a similar manner when $1 \leq |R'| < |Y|$ and $|R'| = |Y| - j$, $1 \leq j \leq 3$.

Note that $(\hat{V}, \hat{\mathcal{B}})$ is not $(k-1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1, \dots, C_{k-1}, C_k \cup \{3t+2, 3t+3\}$ are the colour classes of a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic 3-star system of order $3t+3$.

Case 2.2. Exactly three vertices of the 3-star S belong to the set R . The edges can be decomposed into 3-stars in a manner similar to the last case.

Case 2.3. Exactly two vertices of the 3-star S belong to the set R . Let $R' = R \setminus \{3t, 3t+1\}$ and $Y' = Y \setminus \{3t-1, 3t-2\}$.

If $|R'| > |Y'|$ and $|Y'|$ is even, then let $m = \frac{|Y'|}{2}$ and $\hat{\mathcal{B}} = (\mathcal{B} \setminus \{S\}) \cup \mathcal{T} \cup (\bigcup_{i=1}^{i=2} \mathcal{T}_i)$ where \mathcal{T}_i is the set $\{\{3t+i; y_1, y_2, r_1\}, \dots, \{3t+i; y_{|Y'|-3}, y_{|Y'|-2}, r_m\}, \{3t+i; r_{m+1}, r_{m+2}, r_{m+3}\}, \dots, \{3t+i; r_{|R|-4}, r_{|R|-3}, r_{|R|-2}\}\}$ for $i \in \{1, 2\}$ and \mathcal{T} is the set $\{\{3t+2; 3t+1, 3t, 3t-1\}, \{3t+3; 3t+2, 3t, 3t-1\}, \{3t-2; 3t+2, 3t+1, 3t+3\}, \{3t+1; 3t+3, 3t, 3t-1\}\}$. The edges can be decomposed into 3-stars in a similar manner when $|R'| > |Y'|$ and $|Y'|$ is odd; likewise when $|R'| = |Y'|$.

Note that $(\hat{V}, \hat{\mathcal{B}})$ is not $(k-1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1, \dots, C_{k-1}, C_k \cup \{3t+2, 3t+3\}$ are the colour classes of a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic 3-star system of order $3t+3$. \square

We now show how to iteratively construct a k -chromatic 3-star system from a $(k-1)$ -chromatic 3-star system.

Theorem 2.3 *Let $k \geq 3$. If there exists a $(k-1)$ -chromatic 3-star system then there exists a k -chromatic 3-star system.*

Proof. Given that there exists a $(k-1)$ -chromatic 3-star system, then by Theorem 2.2, there exists a $(k-1)$ -chromatic 3-star system of order n_{k-1} for some $n_{k-1} \in \mathbb{Z}$ such that $n_{k-1} \equiv 0 \pmod{3}$. Let (U_0, \mathcal{A}_0) be a $(k-1)$ -chromatic 3-star system of order n_{k-1} with colour classes C_1, \dots, C_{k-1} such that $|C_1| \leq \dots \leq |C_{k-1}|$ and each vertex of C_s has colour s , $1 \leq s \leq k-1$. For each $s \in \{1, \dots, k-1\}$, let $C_s = \{c_1^s, \dots, c_{|C_s|}^s\}$. For a positive integer ℓ that will be fixed later and for each $i \in \{1, \dots, \ell\}$, let $U_i = U_0 \times \{i\}$ and $\mathcal{A}_i = \mathcal{A}_0 \times \{i\}$ where $\mathcal{A}_0 \times \{i\}$ denotes the set $\{S \times \{i\} \mid S \in \mathcal{A}_0\}$ and when $S = \{x; a, b, c\}$, $S \times \{i\}$ denotes $\{x \times \{i\}; a \times \{i\}, b \times \{i\}, c \times \{i\}\}$. So (U_i, \mathcal{A}_i) has colour classes $C_1 \times \{i\}, \dots, C_{k-1} \times \{i\}$. If $2k \equiv 0 \pmod{3}$ then let $V = \{1, 2, 3, \dots, 2k\}$; otherwise let $V = \{1, 2, 3, \dots, 2k-1\}$. Also let $U = \bigcup_{i=1}^{\ell} U_i$ such that $V \cap U = \emptyset$. We will embed $(U_1, \mathcal{A}_1), \dots, (U_\ell, \mathcal{A}_\ell)$ into a k -chromatic 3-star system $(\hat{V}, \hat{\mathcal{B}})$ where $\hat{V} = V \cup U$. Since the edges of the complete graph on the set V admit a decomposition into 3-stars, we let (V, \mathcal{B}) be an arbitrary 3-star system of order $2k-1$. We now need to decompose the edges between V and U and the edges between U_i and U_j , for $1 \leq i < j \leq \ell$ into 3-stars such that the resulting 3-star system $(\hat{V}, \hat{\mathcal{B}})$ is k -chromatic. To do so, we will decompose the edges between V and U into 3-stars in a way such that no 3-subset in V is monochromatic in any putative $(k-1)$ -colouring. Three cases arise.

Case 1: $2k-1 \equiv 0 \pmod{3}$. Then $2k-1 = 3t$ for some $t \in \mathbb{Z}^+$. The number of 3-subsets of the set V is $\binom{2k-1}{3} = t(k-1)(2k-3)$. We now fix $\ell = (k-1)(2k-3)$. Partition the set of all 3-subsets of V into ℓ sets $\mathbb{T}_1, \dots, \mathbb{T}_\ell$ each consisting of t mutually disjoint 3-subsets. This partition is known to exist by [1]. Let $\mathbb{T}_i = \{\{x_1^i, y_1^i, z_1^i\}, \dots, \{x_t^i, y_t^i, z_t^i\}\}$, $1 \leq i \leq \ell$. Decompose the edges between V and U_i into the 3-stars of the set $\mathcal{T}_i = \bigcup_{u \in U_i} \{\{u; x_1^i, y_1^i, z_1^i\}, \dots, \{u; x_t^i, y_t^i, z_t^i\}\}$, where $1 \leq i \leq \ell$.

We now begin to decompose the edges between U_i and U_j for $1 \leq i < j \leq \ell$ into 3-stars. If $|C_1|$ is even, then let $r = \frac{|C_1|}{2}$. Observe that $2r = |C_1| \leq |C_2|$ and so for each $p \in \{1, 2, \dots, |C_1|\}$ we decompose the edges between $c_p^1 \times \{i\}$ and U_j into the 3-stars of the set: $\mathcal{S}_p^1 = \{\{c_p^1 \times \{i\}; c_1^1 \times \{j\}, c_2^1 \times \{j\}\}, \{c_p^1 \times \{i\}; c_3^1 \times \{j\}, c_4^1 \times \{j\}, c_2^2 \times \{j\}\}, \dots, \{c_p^1 \times \{i\}; c_{|C_1|-1}^1 \times \{j\}, c_{|C_1|}^1 \times \{j\}, c_r^2 \times \{j\}\} \cup \{\{c_p^1 \times \{i\}; x_t, y_t, z_t\} \mid 1 \leq t \leq \frac{n_{k-1}-3r}{3}\}$, where the sets $\{x_t, y_t, z_t\}$, $1 \leq t \leq \frac{n_{k-1}-3r}{3}$, form a partition of $U_j \setminus ((C_1 \times \{j\}) \cup \{c_1^2 \times \{j\}, c_2^2 \times \{j\}, \dots, c_r^2 \times \{j\}\})$. If $|C_1|$ is odd, then let $r = \frac{|C_1|+1}{2}$ and decompose the edges between $c_p^1 \times \{i\}$ and U_j into the 3-stars of the set: $\mathcal{S}_p^1 = \{\{c_p^1 \times \{i\}; c_1^1 \times \{j\}, c_2^1 \times \{j\}, c_1^2 \times \{j\}\}, \{c_p^1 \times \{i\}; c_3^1 \times \{j\}, c_4^1 \times \{j\}, c_2^2 \times \{j\}\}, \dots, \{c_p^1 \times \{i\}; c_{|C_1|-2}^1 \times \{j\}, c_{|C_1|-1}^1 \times \{j\}, c_{r-1}^2 \times \{j\}\}$,

$\{c_p^1 \times \{i\}; c_{|C_1|}^1 \times \{j\}, c_r^2 \times \{j\}, c_{r+1}^2 \times \{j\}\} \cup \{c_p^1 \times \{i\}; x_t, y_t, z_t\} \mid 1 \leq t \leq \frac{n_{k-1}-3r}{3}\}$, where the sets $\{x_t, y_t, z_t\}$, $1 \leq t \leq \frac{n_{k-1}-3r}{3}$, form a partition of $U_j \setminus ((C_1 \times \{j\}) \cup \{c_1^2 \times \{j\}, c_2^2 \times \{j\}, \dots, c_{r+1}^2 \times \{j\}\})$. The edges between $(C_2 \times \{i\}) \cup \dots \cup (C_{k-2} \times \{i\})$ and U_j are decomposed into sets of 3-stars $\mathcal{S}_1^2, \dots, \mathcal{S}_{|C_2|}^2, \dots, \mathcal{S}_1^{k-2}, \dots, \mathcal{S}_{|C_{k-2}|}^{k-2}$ in a similar manner.

Next, we decompose the edges between $C_{k-1} \times \{i\}$ and U_j into 3-stars. If $|C_{k-1}| = 2n$ for some $n \in \mathbb{Z}^+$, then we let $C_k = \{c_{n+1}^{k-1}, \dots, c_{|C_{k-1}|}^{k-1}\}$. If $2n$ is a multiple of 3, then for each $p \in \{1, 2, \dots, |C_{k-1}|\}$ we decompose the edges between c_p^{k-1} and U_j into the 3-stars of the set: $\mathcal{S}_p^{k-1} = \{\{c_p^{k-1} \times \{i\}; c_1^{k-1} \times \{j\}, c_2^{k-1} \times \{j\}, c_{n+1}^{k-1} \times \{j\}\}, \{c_p^{k-1} \times \{i\}; c_{n+2}^{k-1} \times \{j\}, c_{n+3}^{k-1} \times \{j\}, c_3^{k-1} \times \{j\}\}, \dots, \{c_p^{k-1} \times \{i\}; c_{n-2}^{k-1} \times \{j\}, c_{n-1}^{k-1} \times \{j\}, c_{|C_{k-1}|-2}^{k-1} \times \{j\}\}, \{c_p^{k-1} \times \{i\}; c_{|C_{k-1}|-1}^{k-1} \times \{j\}, c_{|C_{k-1}|}^{k-1} \times \{j\}, c_n^{k-1} \times \{j\}\} \cup \{c_p^{k-1} \times \{i\}; x_t, y_t, z_t\} \mid 1 \leq t \leq \frac{n_{k-1}-|C_{k-1}|}{3}\}$, where the sets $\{x_t, y_t, z_t\}$, $1 \leq t \leq \frac{n_{k-1}-|C_{k-1}|}{3}$, form a partition of $\bigcup_{s=1}^{k-2} (C_s \times \{j\})$. If $2n$ is congruent to 1 modulo 3, then the edges are decomposed into the 3-stars of the set: $\mathcal{S}_p^{k-1} = \{\{c_p^{k-1} \times \{i\}; c_1^{k-1} \times \{j\}, c_2^{k-1} \times \{j\}, c_{n+1}^{k-1} \times \{j\}\}, \{c_p^{k-1} \times \{i\}; c_{n+2}^{k-1} \times \{j\}, c_{n+3}^{k-1} \times \{j\}, c_3^{k-1} \times \{j\}\}, \dots, \{c_p^{k-1} \times \{i\}; c_{n-2}^{k-1} \times \{j\}, c_{n-1}^{k-1} \times \{j\}, c_{|C_{k-1}|-3}^{k-1} \times \{j\}\}, \{c_p^{k-1} \times \{i\}; c_{|C_{k-1}|-2}^{k-1} \times \{j\}, c_{|C_{k-1}|-1}^{k-1} \times \{j\}, c_n^{k-1} \times \{j\}\}, \{c_p^{k-1} \times \{i\}; c_{|C_{k-1}|}^{k-1} \times \{j\}, a, b\} \cup \{c_p^{k-1} \times \{i\}; x_t, y_t, z_t\} \mid 1 \leq t \leq \frac{n_{k-1}-(|C_{k-1}|+2)}{3}\}$, where a and b are distinct vertices of $\bigcup_{s=1}^{k-2} (C_s \times \{j\})$ and the sets $\{x_t, y_t, z_t\}$, $1 \leq t \leq \frac{n_{k-1}-(|C_{k-1}|+2)}{3}$, form a partition of $\bigcup_{s=1}^{k-2} (C_s \times \{j\}) \setminus \{a, b\}$. The edges are decomposed into 3-stars in a similar manner when $2n$ is congruent to 2 modulo 3. If $|C_{k-1}| = 2n + 1$ for some $n \in \mathbb{Z}^+$, then we let $C_k = \{c_{n+1}^{k-1}, \dots, c_{|C_{k-1}|}^{k-1}\}$ and decompose the edges into 3-stars in a similar manner. Now $\mathcal{U}_j^i = (\bigcup_{a=1}^{|C_1|} \mathcal{S}_a^1) \cup \dots \cup (\bigcup_{a=1}^{|C_{k-1}|} \mathcal{S}_a^{k-1})$ forms a 3-star decomposition of the edges between U_i and U_j , $1 \leq i < j \leq \ell$.

Let $\hat{\mathcal{B}} = \mathcal{B} \cup (\bigcup_{i=1}^{\ell} \mathcal{A}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{T}_i) \cup (\bigcup_{j=2}^{\ell} \mathcal{U}_j^1) \cup (\bigcup_{j=3}^{\ell} \mathcal{U}_j^2) \cup \dots \cup \mathcal{U}_{\ell}^{\ell-1}$. The k colour classes $\{1, 2\} \cup \bigcup_{i=1}^{\ell} (C_1 \times \{i\}), \{3, 4\} \cup \bigcup_{i=1}^{\ell} (C_2 \times \{i\}), \dots, \{2k-1, 2k-2\} \cup \bigcup_{i=1}^{\ell} (C_{k-1} \setminus C_k) \times \{i\}, \{2k-1\} \cup \bigcup_{i=1}^{\ell} (C_k \times \{i\})$ exhibit a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Since (U_i, \mathcal{A}_i) , $1 \leq i \leq \ell$ is a $(k-1)$ -chromatic 3-star system, there are vertices u_1^i, \dots, u_{k-1}^i in U_i such that $u_s^i \in C_s \times \{i\}$ for each $1 \leq s \leq k-1$. Since $|V| = 2k-1 > 2(k-1)$ then when attempting to colour V with $k-1$ colours, some colour must occur at least thrice within V . Thus, for some $j \in \{1, \dots, \ell\}$ and some $i \in \{1, \dots, \ell\}$ and for some $s \in \{1, \dots, k-1\}$, the three vertices x_j^i, y_j^i and z_j^i are each coloured with the same colour s . Then the 3-star $\{u_s^i, x_j^i, y_j^i, z_j^i\}$ would be monochromatic and hence $(\hat{V}, \hat{\mathcal{B}})$ cannot be coloured with $k-1$ colours.

Therefore, $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic 3-star system of order $n_k = (2k-1) + n_{k-1}(k-1)(2k-3)$.

Case 2: $2k-1 \equiv 1 \pmod{3}$. Then $2k-1 = 3t+1$ for some $t \in \mathbb{Z}^+$. The number of 3-subsets

of the set V is $\binom{2k-1}{3} = \frac{t}{2}(2k-1)(2k-3)$. Note that since $\binom{2k-1}{3} = \frac{t}{2}(2k-1)(2k-3)$ is an integer and $2k-1$ and $2k-3$ are not divisible by 2, t must be even. Let $t = 2t'$. Partition the set of all 3-subsets of V into $\ell' = (2k-1)(k-2)$ sets of disjoint 3-subsets $\mathbb{T}_1, \dots, \mathbb{T}_{\ell'}$ of size t and $\ell'' = 2k-1$ sets of disjoint 3-subsets $\mathbb{T}_{\ell'+1}, \dots, \mathbb{T}_{\ell'+\ell''}$ of size t' . Such a partition is known to exist by Theorem 1 in [1]. We now fix $\ell = \ell' + \ell''$. Let $\mathbb{T}_i = \{\{x_1^i, y_1^i, z_1^i\}, \dots, \{x_{t'}^i, y_{t'}^i, z_{t'}^i\}\}$, $1 \leq i \leq \ell'$ and $\mathbb{T}_i = \{\{x_{t'}^i, y_{t'}^i, z_{t'}^i\}, \dots, \{x_{t'}^i, y_{t'}^i, z_{t'}^i\}\}$, $\ell' + 1 \leq i \leq \ell' + \ell''$.

Now, we decompose the edges between V and U_i , $1 \leq i \leq \ell'$, into 3-stars. Let v be the single vertex of the set $V \setminus \bigcup_{j=1}^t \{x_j^i, y_j^i, z_j^i\}$. The edges between v and U_i , $1 \leq i \leq \ell'$, are decomposed into 3-stars in a manner similar to the decomposition of the edges between a vertex $u \in U_i$ and U_j , $1 \leq i < j \leq \ell$, in Case 1. Let \mathcal{T}_i^1 be the decomposition of the edges between v and U_i into 3-stars, $\mathcal{T}_i^2 = \bigcup_{u \in U_i} \{\{u, x_1^i, y_1^i, z_1^i\}, \dots, \{u, x_{t'}^i, y_{t'}^i, z_{t'}^i\}\}$ and $\mathcal{T}_i = \mathcal{T}_i^1 \cup \mathcal{T}_i^2$, $1 \leq i \leq \ell'$. Then \mathcal{T}_i is a decomposition of the edges between V and U_i , $1 \leq i \leq \ell'$, into 3-stars. The edges between V and U_i are decomposed into 3-stars \mathcal{T}_i in a similar manner for $\ell' + 1 \leq i \leq \ell' + \ell''$.

The edges between U_i and U_j , $1 \leq i < j \leq \ell$, are decomposed into sets of 3-stars \mathcal{U}_j^i in a manner similar to Case 1.

Let $\hat{\mathcal{B}} = \mathcal{B} \cup (\bigcup_{i=1}^{\ell} \mathcal{A}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{T}_i) \cup (\bigcup_{j=2}^{\ell} \mathcal{U}_j^1) \cup (\bigcup_{j=3}^{\ell} \mathcal{U}_j^2) \cup \dots \cup \mathcal{U}_{\ell}^{\ell-1}$. The k colour classes $\{1, 2\} \cup \bigcup_{i=1}^{\ell} (C_1 \times \{i\})$, $\{3, 4\} \cup \bigcup_{i=1}^{\ell} (C_2 \times \{i\})$, \dots , $\{2k-1, 2k-2\} \cup \bigcup_{i=1}^{\ell} (C_{k-1} \setminus C_k) \times \{i\}$, $\{2k-1\} \cup \bigcup_{i=1}^{\ell} (C_k \times \{i\})$ exhibit a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. It is seen that $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic 3-star system of order $n_k = (2k-1) + n_{k-1}(2k-1)(k-1)$ in a manner similar to Case 1.

Case 3: $2k-1 \equiv 2 \pmod{3}$. Then $2k = 3t$ for some $t \in \mathbb{Z}^+$. The number of 3-subsets of the set V is $\binom{2k}{3} = t(2k-1)(k-1)$. We now fix $\ell = (2k-1)(k-1)$. Partition the set of all 3-subsets of V into ℓ sets of disjoint 3-subsets $\mathbb{T}_1, \dots, \mathbb{T}_{\ell}$ of size t . Let $\mathbb{T}_i = \{\{x_1^i, y_1^i, z_1^i\}, \dots, \{x_{t'}^i, y_{t'}^i, z_{t'}^i\}\}$, $1 \leq i \leq \ell$. We decompose the edges between V and U_i into the 3-stars of the set $\mathcal{T}_i = \bigcup_{u \in U_i} \{\{u, x_1^i, y_1^i, z_1^i\}, \dots, \{u, x_{t'}^i, y_{t'}^i, z_{t'}^i\}\}$, where $1 \leq i \leq \ell$. The edges between U_i and U_j , $1 \leq i < j \leq \ell$, are decomposed into 3-stars \mathcal{U}_j^i in a manner similar to Case 1. Let $\hat{\mathcal{B}} = \mathcal{B} \cup (\bigcup_{i=1}^{\ell} \mathcal{A}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{T}_i) \cup (\bigcup_{j=2}^{\ell} \mathcal{U}_j^1) \cup (\bigcup_{j=3}^{\ell} \mathcal{U}_j^2) \cup \dots \cup \mathcal{U}_{\ell}^{\ell-1}$. The k colour classes $\{1, 2\} \cup \bigcup_{i=1}^{\ell} (C_1 \times \{i\})$, $\{3, 4\} \cup \bigcup_{i=1}^{\ell} (C_2 \times \{i\})$, \dots , $\{2k-1, 2k-2\} \cup \bigcup_{i=1}^{\ell} (C_{k-1} \setminus C_k) \times \{i\}$, $\{2k-1, 2k\} \cup \bigcup_{i=1}^{\ell} (C_k \times \{i\})$ exhibit a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. It is seen that $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic 3-star system of order $n_k = 2k + n_{k-1}(2k-1)(k-1)$ in a manner similar to Case 1. \square

We finish this section with the following corollary.

Corollary 2.1 *For any integer $k \geq 2$, there exists some integer n_k such that for all admissible $n \geq n_k$, there exists a k -chromatic 3-star system of order n .*

Proof. For $k = 2$, apply Theorem 2.1. For $k \geq 3$, apply Theorem 2.2 and Theorem 2.3. \square

Observe that the cardinalities of the colour classes of the k -chromatic 3-star systems from Theorem 2.2 and Theorem 2.3 are not excessively imbalanced. The cardinalities of the colour classes of the k -chromatic 3-star systems either differ by at most a small number or the cardinality of the largest colour class of the k -chromatic 3-star systems is almost two times that of the cardinality of the smallest colour class.

3 k -colourings of e -star systems

In this section, we generalize the results of the last section for e -star systems for all $e \geq 3$. For any arbitrary integers $e \geq 3$ and $k \geq 2$, we show that there exists some integer n_k where $n_k \equiv 0 \pmod{2e}$ such that for all $n \geq n_k$ where $n \equiv 0, 1 \pmod{2e}$, there exists a k -chromatic e -star system of order n . We first construct a strongly equitable 2-chromatic e -star system of order $2e$ for all $e \geq 3$.

Theorem 3.1 *There exists a strongly equitable 2-chromatic e -star system of order $2e$ for all $e \geq 3$.*

Proof. By Theorem 2.1, there exists an equitably 2-chromatic 3-star system of order six, (V, \mathcal{B}) , where $V = \{1, 2, 3, 4, 5, 6\}$ and $\mathcal{B} = \{\{1; 3, 5, 6\}, \{2; 1, 3, 6\}, \{4; 1, 2, 3\}, \{5; 2, 3, 4\}, \{6; 3, 4, 5\}\}$ with colour classes $R = \{1, 3, 5\}$ and $Y = \{2, 4, 6\}$. We construct a 2-chromatic 4-star system of order eight, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{7, 8\}$ and $\hat{\mathcal{B}} = \{\{1; 3, 5, 6, 8\}, \{2; 1, 3, 6, 8\}, \{4; 1, 2, 3, 8\}, \{5; 2, 3, 4, 7\}, \{6; 3, 4, 5, 7\}, \{7; 1, 2, 3, 4\}, \{8; 3, 5, 6, 7\}\}$, $\hat{R} = \{1, 3, 5, 7\}$ and $\hat{Y} = \{2, 4, 6, 8\}$. Then $(\hat{V}, \hat{\mathcal{B}})$ is an equitably 2-chromatic 4-star system of order 8 with colour classes \hat{R} and \hat{Y} .

We now generalize this construction, which can be used in an iterative manner. Suppose that there exists an equitably 2-chromatic e -star system of order $2e$, (V, \mathcal{B}) , where $V = \{1, \dots, 2e\}$ is the set of points which is partitioned into two subsets R and Y and $\mathcal{B} = \{\{1; a_1^1, \dots, a_e^1\}, \{2; a_1^2, \dots, a_e^2\}, \{4; a_1^4, \dots, a_e^4\}, \{5; a_1^5, \dots, a_e^5\}, \dots, \{2e; a_1^{2e}, \dots, a_e^{2e}\}\}$ is the set of blocks. We construct an equitably 2-chromatic $(e+1)$ -star system of order $2e+2$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{2e+1, 2e+2\}$ and $\hat{\mathcal{B}} = \{\{1; a_1^1, \dots, a_e^1, 2e+2\}, \{2; a_1^2, \dots, a_e^2, 2e+2\}, \{4; a_1^4, \dots, a_e^4, 2e+2\}, \{5; a_1^5, \dots, a_e^5, 2e+2\}, \dots, \{e+1; a_1^{e+1}, \dots, a_e^{e+1}, 2e+2\}, \{e+2; a_1^{e+2}, \dots, a_e^{e+2}, 2e+1\}, \{e+3; a_1^{e+3}, \dots, a_e^{e+3}, 2e+1\}, \dots, \{2e; a_1^{2e}, \dots, a_e^{2e}, 2e+1\}, \{2e+1; 1, 2, \dots, e+1\}, \{2e+2; 3, e+2, e+3, \dots, 2e, 2e+1\}\}$. Then $(\hat{V}, \hat{\mathcal{B}})$ is a strongly equitable 2-chromatic $(e+1)$ -star system of order $2e+2$ with colour classes $\hat{R} = \{1, 3, 5, \dots, 2e+1\}$ and $\hat{Y} = \{2, 4, 6, \dots, 2e+2\}$. \square

We now show how to construct a k -chromatic e -star system from a smaller k -chromatic e -star system.

Theorem 3.2 *Let $k \geq 2$ and $e \geq 3$. If there exists a k -chromatic e -star system of order n_0 such that $n_0 \equiv 0, 1 \pmod{2e}$, then there exists a k -chromatic e -star system for all $n > n_0$ such that $n \equiv 0, 1 \pmod{2e}$.*

Proof. Suppose that there exists a k -chromatic e -star system, (V, \mathcal{B}) , where $V = \{1, \dots, n_0\}$ is the set of points and \mathcal{B} is the set of blocks and $n_0 \equiv 0$ or $1 \pmod{2e}$. Given a k -colouring of (V, \mathcal{B}) , with colours $1, 2, \dots, k$ and colour classes C_1, C_2, \dots, C_k , let $R = \bigcup_{i=1}^{\ell} C_i$ and $Y = \bigcup_{i=\ell+1}^k C_i$ for some integer ℓ such that $1 \leq \ell < k$. Let $r_1, r_2, \dots, r_{|R|}$ be the elements of R and $y_1, y_2, \dots, y_{|Y|}$ be the elements of Y . Observe that R and Y form a partition of V . Without loss of generality assume that $|R| \geq |Y|$.

Case 1. Suppose that $n_0 \equiv 0 \pmod{2e}$. Then $n_0 = 2et$, $t \geq 1$.

First, we construct a k -chromatic $S_e(2et + 1)$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{2et + 1\}$.

If $|R| = |Y|$, let $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}$ where \mathcal{T} is the set $\{\{2et + 1; r_1, \dots, r_{e-1}, y_1\}, \{2et + 1; y_2, \dots, y_e, r_e\}, \dots, \{2et + 1; r_{|R|-(e-1)}, \dots, r_{|R|-1}, y_{|Y|-(e-1)}\}, \{2et + 1; y_{|Y|-e+2}, \dots, y_{|Y|}, r_{|R|}\}\}$.

Otherwise, $|R| > |Y|$. Then let $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}$ where \mathcal{T} is the set $\{\{2et + 1; y_1, \dots, y_{e-1}, r_1\}, \dots, \{2et + 1; y_{t'e-(t'-1)}, \dots, y_{(t'+1)e-t'-1}, r_{t'+1}\}, \{2et + 1; y_{(t'+1)e-t'}, \dots, y_{|Y|}, r_{t'+2}, \dots, r_{t'+i-1}\}, \{2et + 1; r_{t'+i}, \dots, r_{t'+i+e-1}\}, \dots, \{2et + 1; r_{|R|-(e-1)}, \dots, r_{|R|}\}\}$, where $t' = \lfloor \frac{|Y|}{e-1} \rfloor - 1$ and $i = (t' + 2)e - |Y| - t' + 1$.

Note that $(\hat{V}, \hat{\mathcal{B}})$ is not $(k-1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1, \dots, C_{k-1}, C_k \cup \{2et + 1\}$ are the colour classes of a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic e -star system of order $2et + 1$.

Next, we construct a k -chromatic $S_e(2et + 2e)$, $(\tilde{V}, \tilde{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\tilde{V} = V \cup V'$ where $V' = \{2et + 1, \dots, 2et + 2e\}$ and let (V', \mathcal{B}') be a 2-chromatic e -star system of order $2e$ constructed from Theorem 3.1 with colour classes $R' = \{2et + 1, 2et + 3, \dots, 2et + 2e - 1\}$ and $Y' = \{2et + 2, 2et + 4, \dots, 2et + 2e\}$. Let $\tilde{\mathcal{B}} = \mathcal{B} \cup \mathcal{B}' \cup \mathcal{T}$ where $\mathcal{T} = \bigcup_{v \in V} \{\{v; 2et + 1, \dots, 2et + e\}, \{v; 2et + e + 1, \dots, 2et + 2e\}\}$. Note that $(\tilde{V}, \tilde{\mathcal{B}})$ is not $(k-1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1 \cup R', C_2 \cup Y', C_3, \dots, C_k$ are the colour classes of a k -colouring of $(\tilde{V}, \tilde{\mathcal{B}})$. Therefore, $(\tilde{V}, \tilde{\mathcal{B}})$ is a k -chromatic e -star system of order $2et + 2e$.

Case 2. Suppose that $n_0 \equiv 1 \pmod{2e}$. Then $n_0 = 2et + 1$, $t \geq 1$. We construct a k -chromatic $S_e(2et + 2e)$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup V_1 \cup V_2$ where $V_1 = \{2et + 2, \dots, 2et + e + 1\}$ and $V_2 = \{2et + e + 2, \dots, 2et + 2e\}$. Let $v_0 = r_{|R|}$ and $R' = R \setminus \{v_0\}$.

If $|R'| = |Y|$, let $\mathcal{T}_1 = \bigcup_{v \in V_2} \{\{v; r_1, \dots, r_{e-1}, y_1\}, \{v; y_2, \dots, y_e, r_e\}, \dots, \{v; r_{|R|-e}, \dots, r_{|R|-2}, y_{|Y|-e+1}\}, \{v; y_{|Y|-e+2}, \dots, y_{|Y|}, r_{|R|-1}\}\}$.

If $|R'| = |Y| - 1$ or $|R'| > |Y|$, decompose the edges between V_2 and $V \setminus \{v_0\}$ into a set of e -stars \mathcal{T}_1 in a manner similar to the case $|R'| = |Y|$.

Let $V' = V_1 \cup V_2 \cup \{v_0\}$ and (V', \mathcal{B}') be a 2-chromatic e -star system of order $2e$ constructed from Theorem 3.1 with colour classes $R' = \{2et + 3, 2et + 4, \dots, 2et + e + 1\}$ and $Y' = \{2et + 2, 2et + e + 2, 2et + e + 3, \dots, 2et + 2e\}$. Let $\mathcal{T}_2 = \bigcup_{v \in V \setminus \{v_0\}} \{\{v; 2et + 2, \dots, 2et + e + 1\}\}$. Let $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{B}' \cup \mathcal{T}_1 \cup \mathcal{T}_2$. Note that $(\hat{V}, \hat{\mathcal{B}})$ is not $(k - 1)$ -colourable because it contains a copy of (V, \mathcal{B}) . Observe that $C_1 \cup R', C_2, \dots, C_{k-1}, C_k \cup Y'$ are the colour classes of a k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ is a k -chromatic e -star system of order $2et + 2e$. \square

We now show how to iteratively construct a k -chromatic e -star system from a $(k - 1)$ -chromatic e -star system.

Theorem 3.3 *Let $k \geq 3$ and $e \geq 3$. If there exists a $(k - 1)$ -chromatic e -star system of order $n_{k-1} \equiv 0 \pmod{2e}$ then there exists a k -chromatic e -star system of order $n_k \equiv 0 \pmod{2e}$.*

Proof. Let (U_0, \mathcal{A}_0) be a $(k - 1)$ -chromatic e -star system of order n_{k-1} with colour classes C_1, \dots, C_{k-1} such that $n_{k-1} \equiv 0 \pmod{2e}$, $|C_1| \leq \dots \leq |C_{k-1}|$ and each vertex of C_s has colour s , for $1 \leq s \leq k - 1$. For each $s \in \{1, \dots, k - 1\}$, let $C_s = \{c_1^s, \dots, c_{|C_s|}^s\}$. For a positive integer ℓ that will be fixed later and for each $i \in \{1, \dots, \ell\}$, let $U_i = U_0 \times \{i\}$ and $\mathcal{A}_i = \mathcal{A}_0 \times \{i\}$ where $\mathcal{A}_0 \times \{i\}$ denotes the set $\{S \times \{i\} \mid S \in \mathcal{A}_0\}$. So (U_i, \mathcal{A}_i) has colour classes $C_1 \times \{i\}, \dots, C_{k-1} \times \{i\}$. Let $W = \{w_1, w_2, \dots, w_{(e-1)(k-1)+1}\}$, $B = \{b_1, b_2, \dots, b_{k-2}\}$ and $D = \{d_1, d_2, \dots, d_e\}$ be pairwise disjoint sets. If $k - 1$ is even then let $V = W \cup B$; otherwise let $V = W \cup B \cup D$. Also let $U = \bigcup_{i=1}^{\ell} U_i$ such that $V \cap U = \emptyset$. We will embed $(U_1, \mathcal{A}_1), \dots, (U_{\ell}, \mathcal{A}_{\ell})$ into a k -chromatic e -star system $(\hat{V}, \hat{\mathcal{B}})$ where $\hat{V} = V \cup U$. Let $n_k = |\hat{V}|$ and observe that $n_k \equiv 0 \pmod{2e}$. By Theorem 3.1 and Theorem 3.2, the edges of the complete graph on the set V admit a decomposition into e -stars which is 2-chromatic. Let (V, \mathcal{B}) be a 2-chromatic e -star system on the set V (so (V, \mathcal{B}) will not obstruct the k -colouring that we will construct). We now need to decompose the edges between V and U and the edges between U_i and U_j , for $1 \leq i < j \leq \ell$ into e -stars such that the resulting e -star system $(\hat{V}, \hat{\mathcal{B}})$ is k -chromatic. To do so, we will decompose the edges between V and U into e -stars in a way such that no e -subset in V is monochromatic in any putative $(k - 1)$ -colouring.

The number of e -subsets of the set W is $N = \binom{(e-1)(k-1)+1}{e}$. Let $a = (k - 2) + \lfloor \frac{-k+2}{e} \rfloor$ and $N = aq + r$ where q is a positive integer and $0 \leq r < a$ is an integer. Partition the set of all e -

subsets of W into q sets of disjoint e -subsets $\mathbb{T}_1, \dots, \mathbb{T}_q$ of size a and one set of disjoint e -subsets \mathbb{T}_{q+1} of size r . Such a partition is known to exist by Theorem 1 in [1]. We now fix $\ell = q + 1$ if $r > 0$ and $\ell = q$ if $r = 0$. Let $\mathbb{T}_i = \{\{(x_1)_1^i, \dots, (x_e)_1^i\}, \dots, \{(x_1)_a^i, \dots, (x_e)_a^i\}\}$, $1 \leq i \leq q$ and $\mathbb{T}_{q+1} = \{\{(x_1)_1^{q+1}, \dots, (x_e)_1^{q+1}\}, \dots, \{(x_1)_r^{q+1}, \dots, (x_e)_r^{q+1}\}\}$ if $r > 0$ and $\mathbb{T}_{q+1} = \emptyset$ if $r = 0$. Let $W_i = W \setminus \bigcup_{j=1}^a (\{(x_1)_j^i, \dots, (x_e)_j^i\})$ for $1 \leq i \leq q$ and $W_{q+1} = W \setminus \bigcup_{j=1}^r (\{(x_1)_j^{q+1}, \dots, (x_e)_j^{q+1}\})$ if $r > 0$ and $W_{q+1} = \emptyset$ if $r = 0$.

For each $i \in \{1, 2, \dots, \ell\}$ and for each $w \in W_i$, we decompose the edges between w and U_i . First, we decompose all the edges between w and $C_1 \times \{i\}$ into e -stars. Let $\mathcal{R}_w^{1,2} = \{\{w; c_1^1 \times \{i\}, \dots, c_{e-1}^1 \times \{i\}, c_2^2 \times \{i\}\}, \dots, \{w; c_{(t-1)(e-1)+1}^1 \times \{i\}, \dots, c_{t(e-1)}^1 \times \{i\}, c_{t+1}^2 \times \{i\}\}, \{w; c_{t(e-1)+1}^1 \times \{i\}, \dots, c_{|C_1|}^1 \times \{i\}, c_{t+1}^2 \times \{i\}, \dots, c_{t+t'}^2 \times \{i\}\}\}$ where t and t' are integers such that $t = \lfloor \frac{|C_1|}{e-1} \rfloor$ and $t' = (t+1)e - |C_1|$. Then $\mathcal{R}_w^{1,2}$ is a decomposition of all the edges between w and $C_1 \times \{i\}$ along with the edges between w and $C'_2 \times \{i\}$ into e -stars where $C'_2 = \{c_1^2, \dots, c_{t+t'}^2\} \subseteq C_2$. For each $j = 2, 3, \dots, k-2$, iteratively proceed in a similar manner to decompose all the edges between w and $(C_j \setminus C'_j) \times \{i\}$ into a set of e -stars $\mathcal{R}_w^{j,j+1}$. So $\bigcup_{j=1}^{k-2} \mathcal{R}_w^{j,j+1}$ gives a decomposition of the edges between w and $(C_1 \cup \dots \cup C_{k-2} \cup C'_{k-1}) \times \{i\}$ where $C'_{k-1} \subseteq C_{k-1}$. Let $m = \frac{|C_{k-1} \setminus C'_{k-1}|}{e}$ and let $C''_{k-1} = C_{k-1} \setminus C'_{k-1}$ and without loss of generality we can let $C''_{k-1} = \{c_1^{k-1}, \dots, c_{me}^{k-1}\}$. Without loss of generality let $C_k = \{c_1^{k-1}, c_{e+1}^{k-1}, c_{2e+1}^{k-1}, \dots, c_{(m-1)e+1}^{k-1}\}$ and decompose the edges between w and $C''_{k-1} \times \{i\}$ into e -stars of the set: $\mathcal{R}_w^{k-1} = \{\{w; c_1^{k-1} \times \{i\}, c_2^{k-1} \times \{i\}, \dots, c_e^{k-1} \times \{i\}\}, \{w; c_{e+1}^{k-1} \times \{i\}, c_{e+2}^{k-1} \times \{i\}, \dots, c_{2e}^{k-1} \times \{i\}\}, \dots, \{w; c_{(m-1)e+1}^{k-1} \times \{i\}, c_{(m-1)e+2}^{k-1} \times \{i\}, \dots, c_{me}^{k-1} \times \{i\}\}\}$.

For each $i \in \{1, 2, \dots, q\}$, let $\mathcal{J}_i = \bigcup_{u \in U_i} \{\{u; (x_1)_1^i, \dots, (x_e)_1^i\}, \dots, \{u; (x_1)_a^i, \dots, (x_e)_a^i\}\}$ and $\mathcal{J}_{q+1} = \{\{u; (x_1)_1^{q+1}, \dots, (x_e)_1^{q+1}\}, \dots, \{u; (x_1)_r^{q+1}, \dots, (x_e)_r^{q+1}\}\}$ if $r > 0$ and $\mathcal{J}_{q+1} = \emptyset$ if $r = 0$. Then $\mathcal{P}_i = \mathcal{J}_i \cup \bigcup_{w \in W_i} (\mathcal{R}_w^{k-1} \cup (\bigcup_{j=1}^{k-2} \mathcal{R}_w^{j,j+1}))$ is a decomposition of the edges between W and U_i into e -stars. The edges between U_i and U_j , $1 \leq i < j \leq \ell$ are decomposed into sets of e -stars \mathcal{U}_j^i in a manner similar to the decomposition of the edges between W_i and U_i , $1 \leq i \leq \ell$.

We have two cases.

Case 1. $k-1$ is even. Then $V = W \cup B$. The edges between B and U_i , $1 \leq i \leq \ell$ are decomposed into sets of e -stars \mathcal{F}_i in a manner similar to the decomposition of the edges between W_i and U_i , $1 \leq i \leq \ell$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ where $\hat{\mathcal{B}} = \mathcal{B} \cup (\bigcup_{i=1}^{\ell} \mathcal{A}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{P}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{F}_i) \cup (\bigcup_{j=2}^{\ell} \mathcal{U}_j^1) \cup (\bigcup_{j=3}^{\ell} \mathcal{U}_j^2) \cup \dots \cup \mathcal{U}_{\ell}^{\ell-1}$ is an e -star system of order $n_k = (k-1)e + n_{k-1}(q+1)$. A k -colouring is given by $(\bigcup_{i=1}^{\ell} C_1 \times \{i\}) \cup \{w_1, \dots, w_{e-1}, b_1\}$, $(\bigcup_{i=1}^{\ell} C_2 \times \{i\}) \cup \{w_e, \dots, w_{2e-2}, b_2\}$, \dots , $(\bigcup_{i=1}^{\ell} C_{k-2} \times \{i\}) \cup \{w_{(e-1)(k-1)-2e-1}, \dots, w_{(e-1)(k-1)-e-1}, b_{k-2}\}$, $(\bigcup_{i=1}^{\ell} (C_{k-1} \setminus C_k) \times \{i\}) \cup$

$$\{w_{(e-1)(k-1)-e}, \dots, w_{(e-1)(k-1)}\}, \left(\bigcup_{i=1}^{\ell} C_k \times \{i\}\right) \cup \{w_{(e-1)(k-1)+1}\}.$$

Case 2. $k-1$ is odd. then $V = W \cup B \cup D$. The edges between D and U_i , $1 \leq i \leq \ell$ are decomposed into sets of e -stars \mathcal{H}_i in a manner similar to the decomposition of the edges between W_i and U_i , $1 \leq i \leq \ell$. Therefore, $(\hat{V}, \hat{\mathcal{B}})$ where $\hat{\mathcal{B}} = \mathcal{B} \cup \left(\bigcup_{i=1}^{\ell} \mathcal{A}_i\right) \cup \left(\bigcup_{i=1}^{\ell} \mathcal{P}_i\right) \cup \left(\bigcup_{i=1}^{\ell} \mathcal{F}_i\right) \cup \left(\bigcup_{i=1}^{\ell} \mathcal{H}_i\right) \cup \left(\bigcup_{j=2}^{\ell} \mathcal{U}_j^1\right) \cup \left(\bigcup_{j=3}^{\ell} \mathcal{U}_j^2\right) \cup \dots \cup \mathcal{U}_{\ell}^{\ell-1}$ is an e -star system of order $n_k = ek + n_{k-1}(q+1)$. A k -colouring is given by $\left(\bigcup_{i=1}^{\ell} C_1 \times \{i\}\right) \cup \{w_1, \dots, w_{e-1}, b_1\}, \left(\bigcup_{i=1}^{\ell} C_2 \times \{i\}\right) \cup \{w_e, \dots, w_{2e-2}, b_2\} \dots, \left(\bigcup_{i=1}^{\ell} C_{k-2} \times \{i\}\right) \cup \{w_{(e-1)(k-1)-2e-1}, \dots, w_{(e-1)(k-1)-e-1}, b_{k-2}\}, \left(\bigcup_{i=1}^{\ell} (C_{k-1} \setminus C_k) \times \{i\}\right) \cup \{w_{(e-1)(k-1)-e}, \dots, w_{(e-1)(k-1)}, d_1\}, \left(\bigcup_{i=1}^{\ell} C_k \times \{i\}\right) \cup \{w_{(e-1)(k-1)+1}, d_2, \dots, d_e\}.$

Since (U_i, \mathcal{A}_i) , $1 \leq i \leq \ell$ is a $(k-1)$ -chromatic e -star system, there are vertices u_1^i, \dots, u_{k-1}^i in U_i such that $u_s^i \in C_s \times \{i\}$ for each $1 \leq s \leq k-1$. Since $|W| = (e-1)(k-1)+1 > (e-1)(k-1)$ then when attempting to colour W with $k-1$ colours, some colour must occur at least e times within W . Thus, for some $i \in \{1, \dots, q\}$ and some $j \in \{1, \dots, a\}$ and for some $s \in \{1, \dots, k-1\}$, the vertices $(x_1)_j^i, (x_2)_j^i, \dots, (x_e)_j^i$ are each coloured with the same colour s . Also, for $i = q+1$ and some $j \in \{1, \dots, r\}$ and for some $s \in \{1, \dots, k-1\}$, the vertices $(x_1)_j^i, (x_2)_j^i, \dots, (x_e)_j^i$ are each coloured with the same colour s . Then the e -star $\{u_s^i; (x_1)_j^i, (x_2)_j^i, \dots, (x_e)_j^i\}$ would be monochromatic and hence in either case $(\hat{V}, \hat{\mathcal{B}})$ cannot be coloured with $k-1$ colours. \square

We summarize this section with the following corollary.

Corollary 3.1 *For any integers $k \geq 2$ and $e \geq 3$, there exists some integer n_k where $n_k \equiv 0 \pmod{2e}$ such that for all admissible $n \geq n_k$ where $n \equiv 0, 1 \pmod{2e}$, there exists a k -chromatic e -star system of order n .*

Proof. Apply Theorem 3.1, Theorem 3.2 and Theorem 3.3. \square

4 Unique colourings of e -star systems

We now investigate uniquely k -chromatic e -star systems for any $e \geq 3$. We commence by showing that such designs do indeed exist for $k = 2$.

Theorem 4.1 *For any $e \geq 3$, there exists a strongly equitable uniquely 2-chromatic e -star system of order n for some $n \equiv 0 \pmod{2e}$.*

Proof. let (U_0, \mathcal{A}_0) be a strongly equitable 2-chromatic e -star system of order $2e$ constructed from Theorem 3.1 with colour classes C_1 and C_2 such that each vertex of C_s has colour s ,

for $s = 1, 2$. For a positive integer ℓ that will be fixed later and for each $i \in \{1, \dots, \ell\}$, let $U_i = U_0 \times \{i\}$ and $\mathcal{A}_i = \mathcal{A}_0 \times \{i\}$ where $\mathcal{A}_0 \times \{i\}$ denotes the set $\{S \times \{i\} \mid S \in \mathcal{A}_0\}$. So (U_i, \mathcal{A}_i) has colour classes $C_1 \times \{i\}$ and $C_2 \times \{i\}$. Let $U = \bigcup_{i=1}^{\ell} U_i$. Let $A^1 = \{a_1^1, \dots, a_e^1\}$, $A^2 = \{a_1^2, \dots, a_e^2\}$ and $A = A^1 \cup A^2$. Similarly let $F^1 = \{f_1^1, \dots, f_e^1\}$, $F^2 = \{f_1^2, \dots, f_e^2\}$, $F = F^1 \cup F^2$, $G^1 = \{g_1^1, \dots, g_e^1\}$, $G^2 = \{g_1^2, \dots, g_e^2\}$, $G = G^1 \cup G^2$, $H^1 = \{h_1^1, \dots, h_e^1\}$, $H^2 = \{h_1^2, \dots, h_e^2\}$, $H = H^1 \cup H^2$, $K^1 = \{k_1^1, \dots, k_e^1\}$, $K^2 = \{k_1^2, \dots, k_e^2\}$ and $K = K^1 \cup K^2$. We construct a uniquely 2-chromatic e -star system $(\hat{V}, \hat{\mathcal{B}})$ where $\hat{V} = A \cup F \cup G \cup H \cup U \cup K$.

We need to decompose the edges between the subsets of \hat{V} into e -stars such that the resulting e -star system $(\hat{V}, \hat{\mathcal{B}})$ is uniquely 2-chromatic.

First, we decompose the edges between A and U into e -stars in a way such that no e -subset in A other than the e -subsets A^1 and A^2 are monochromatic in any putative 2-colouring of $(\hat{V}, \hat{\mathcal{B}})$. Let \mathcal{D} be the set of all e -subsets of A except A^1 and A^2 . The number of e -subsets of the set A is $N = \binom{2e}{e}$ and $|\mathcal{D}| = N - 2$. Let T_i , $1 \leq i \leq |\mathcal{D}|$, be the elements of \mathcal{D} . Let $\tilde{A}_i = A \setminus T_i$, $1 \leq i \leq |\mathcal{D}|$. We now fix $\ell = |\mathcal{D}|$.

Partition U_0 into e -subsets E_1 and E_2 such that $E_j \cap C_1 \neq \emptyset$ and $E_j \cap C_2 \neq \emptyset$, $j = 1, 2$. For each $i \in \{1, 2, \dots, \ell\}$ and for each $w \in \tilde{A}_i$, we decompose the edges between w and U_i into the set of e -stars $\mathcal{R}_w = \bigcup_{j=1}^2 \{\{w; E_j \times \{i\}\}\}$. For each $i \in \{1, \dots, \ell\}$, let $\mathcal{T}_i = \bigcup_{u \in U_i} \{\{u; T_i\}\}$. Then $\mathcal{P}_i = \mathcal{T}_i \cup \left(\bigcup_{w \in \tilde{A}_i} \mathcal{R}_w \right)$ is a decomposition of the edges between A and U_i and $\bigcup_{i=1}^{\ell} \mathcal{P}_i$ is a decomposition of the edges between A and U into e -stars.

Recall that (U_i, \mathcal{A}_i) is 2-chromatic. So for any 2-colouring of (U_i, \mathcal{A}_i) there must be a vertex u_s^i of each colour s in U_i , $1 \leq s \leq 2$. Note that for each $i \in \{1, \dots, |\mathcal{D}|\}$, $\{u_s^i; T_i\} \in \mathcal{T}_i$ and so to ultimately obtain a valid 2-colouring of $(\hat{V}, \hat{\mathcal{B}})$, the e vertices of T_i must not all have colour s , $1 \leq s \leq 2$. If a colour s occurs on more than e vertices of A then there exists a T_i with only colour s and then $\{u_s^i; T_i\}$ would be monochromatic. So in any valid 2-colouring of $(\hat{V}, \hat{\mathcal{B}})$, each colour must occur on exactly e points of A . If the points of colour s are not all in A^1 or A^2 then they are the leaves of a monochromatic e -star with centre u_s^i . Without loss of generality we therefore assign colour s to all e points of A^s , $1 \leq s \leq 2$.

We now begin to decompose the edges between A and F into e -stars. For each $i \in \{1, \dots, e\}$, let $\mathcal{K}_i^1 = \{\{f_i^1; A^2\}\}$ and $\mathcal{K}_i^1 = \{\{f_i^1; A^1\}\}$. Since each vertex of A^2 has colour 2 then each vertex of F^1 must have colour 1 or else a monochromatic e -star would now exist. Likewise, each vertex of F^2 must have colour 2. Let $\mathcal{K}^1 = \bigcup_{i=1}^e \mathcal{K}_i^1$ and $\mathcal{K}^2 = \bigcup_{i=1}^e \mathcal{K}_i^2$. We decompose the edges between A^i and F^i , $1 \leq i \leq 2$, later. Decompose the edges between A^1 and G^2 and between A^2 and G^1 into a set of e -stars \mathcal{L} in a similar manner, forcing the vertices of the subsets G^1 and G^2

to have colours 1 and 2 respectively. We decompose the edges between A^i and G^i , $1 \leq i \leq 2$, later. Also, decompose the edges between K^1 and G^2 and between K^2 and G^1 into a set of e -stars \mathcal{C} in a similar manner, forcing the vertices of the subsets K^1 and K^2 to have colours 1 and 2 respectively. We decompose the edges between K^i and G^i , $1 \leq i \leq 2$, later

Next, we decompose the edges between H and $F \cup G$. For each $i \in \{1, \dots, e\}$, let $\mathcal{E}_i^1 = \{\{h_i^1; f_1^2, \dots, f_e^2\}, \{h_i^1; f_1^1, \dots, f_{e-1}^1, g_1^2\}, \{h_i^1; f_e^1, g_1^1, \dots, g_{e-2}^1, g_2^2\}, \{h_i^1; g_{e-1}^1, g_e^1, g_3^2, \dots, g_e^2\}\}$, and $\mathcal{E}^1 = \bigcup_{i=1}^e \mathcal{E}_i^1$. Then \mathcal{E}^1 is a decomposition of the edges between H^1 and $F \cup G$ into e -stars such that all the vertices in H^1 are forced to have colour 1. Decompose the edges between H^2 and $F \cup G$ into a set of e -stars \mathcal{E}^2 in a similar manner, forcing every vertex in H^2 to have unique colour 2. Let $\mathcal{E} = \bigcup_{j=1}^2 \mathcal{E}^j$.

Partition the $6e - 6$ vertices of $F^1 \cup G^1 \cup H \cup K \setminus \{h_1^2, h_2^2, h_3^2, k_1^2, k_2^2, k_3^2\}$ into six sets S_1, \dots, S_6 of size $e - 1$. For each $1 \leq i \leq e$, let $M_i^1 = \{\{a_i^1; S_1 \cup \{h_1^2\}\}, \{a_i^1; S_2 \cup \{h_2^2\}\}, \{a_i^1; S_3 \cup \{h_3^2\}\}, \{a_i^1; S_4 \cup \{k_1^2\}\}, \{a_i^1; S_5 \cup \{k_2^2\}\}, \{a_i^1; S_6 \cup \{k_3^2\}\}\}$. Then $M^1 = \bigcup_{i=1}^e M_i^1$ is a decomposition of the edges between A^1 and $F^1 \cup G^1 \cup H \cup K$ into e -stars, none of which is monochromatic. Decompose the edges between A^2 and $F^2 \cup G^2 \cup H \cup K$ into a set of e -stars M^2 in a similar manner. Let $\mathcal{M} = \bigcup_{j=1}^2 M^j$.

Partition the $5e - 5$ vertices of $G^1 \cup F \cup H \setminus \{f_1^2, f_2^2, f_3^2, h_1^2, h_2^2\}$ into five sets R_1, \dots, R_5 of size $e - 1$. For each $1 \leq i \leq e$, let $\mathcal{F}_i^1 = \{\{k_i^1; R_1 \cup \{f_1^2\}\}, \{k_i^1; R_2 \cup \{f_2^2\}\}, \{k_i^1; R_3 \cup \{f_3^2\}\}, \{k_i^1; R_4 \cup \{h_1^2\}\}, \{k_i^1; R_5 \cup \{h_2^2\}\}\}$. Then $\mathcal{F}^1 = \bigcup_{i=1}^e \mathcal{F}_i^1$ is a decomposition of the edges between K^1 and $G^1 \cup F \cup H$ into e -stars, none of which is monochromatic. Decompose the edges between K^2 and $G^2 \cup F \cup H$ into a set of e -stars \mathcal{F}^2 in a similar manner.

Next, for each $1 \leq i \leq \ell$, we decompose the edges between $F \cup G$ and U_i into e -stars. Let u_1^s, \dots, u_e^s be the vertices of colour s , $1 \leq s \leq 2$, in an equitable 2-colouring of (U_0, \mathcal{A}_0) . For each $j \in \{1, \dots, e\}$, let $\mathcal{N}_j^1 = \{\{u_j^1 \times \{i\}; f_1^2, \dots, f_e^2\}, \{u_j^1 \times \{i\}; f_1^1, \dots, f_{e-1}^1, g_1^2\}, \{u_j^1 \times \{i\}; f_e^1, g_1^1, \dots, g_{e-2}^1, g_2^2\}, \{u_j^1 \times \{i\}; g_{e-1}^1, g_e^1, g_3^2, \dots, g_e^2\}\}$ and observe that $u_j^1 \times \{i\}$ is now forced to have colour 1.

For each $j \in \{1, \dots, e\}$, decompose the edges between $u_j^2 \times \{i\}$ and $F \cup G$ into a set of e -stars \mathcal{N}_j^2 in a similar manner so that $u_j^2 \times \{i\}$ is forced to have colour 2. Let $\mathcal{N}^s = \bigcup_{j=1}^e \mathcal{N}_j^s$ and $\mathcal{O}_i = \bigcup_{s=1}^2 \mathcal{N}^s$. Then \mathcal{O}_i is a decomposition of the edges between $F \cup G$ and U_i into e -stars which forces every vertex of U_i to have a unique colour. Let $\mathcal{O} = \bigcup_{i=1}^{\ell} \mathcal{O}_i$.

At this point every vertex of V is coloured, and the colouring is unique (up to a permutation of the colours). Moreover, the two colour classes are equal in size and so the colouring is strongly equitable. All that remains is to complete the decomposition without introducing

monochromatic e -stars.

For each $1 \leq i \leq e$, we begin to decompose the edges between F and G into e -stars. Let $Q_i^1 = \{\{f_i^1; g_1^1, \dots, g_{e-1}^1, g_1^2\}, \{f_i^1; g_e^1, g_2^2, \dots, g_e^2\}\}$. Then $Q^1 = \bigcup_{i=1}^e Q_i^1$ is a decomposition of the edges between F^1 and G into e -stars, none of which is monochromatic. Decompose the edges between F^2 and G into a set of e -stars Q^2 in a similar manner. Let $Q = \bigcup_{j=1}^2 Q^j$.

Next, for each $1 \leq i \leq \ell$, we decompose the edges between H and U_i into e -stars. Recall that u_1^s, \dots, u_e^s are vertices of colour s , $1 \leq s \leq 2$. Let $\mathcal{J}_j^1 = \{\{u_j^1 \times \{i\}; h_1^1, \dots, h_{e-1}^1, h_1^2\}, \{u_j^1 \times \{i\}; h_e^1, h_2^2, \dots, h_e^2\}\}$, $1 \leq j \leq e$. For each $j \in \{1, \dots, e\}$, decompose the edges between $u_j^2 \times \{i\}$ and H into a set of e -stars \mathcal{J}_j^2 in a similar manner. Let $\mathcal{J}^s = \bigcup_{j=1}^e \mathcal{J}_j^s$ and $\mathcal{S}_i = \bigcup_{s=1}^2 \mathcal{J}^s$.

Then \mathcal{S}_i is a decomposition of the edges between H and U_i into e -stars. Let $\mathcal{S} = \bigcup_{i=1}^\ell \mathcal{S}_i$. For each $1 \leq i \leq \ell$, decompose the edges between K and U_i into a set of e -stars \mathcal{H}_i in a similar manner so that $\bigcup_{i=1}^\ell \mathcal{H}_i$ decomposes all of the edges between K and U . The edges between U_i and U_j , $1 \leq i < j \leq \ell$, are decomposed into set of e -stars \mathcal{U}_j^i in a similar manner.

Note that $|A| = |F| = |G| = |H| = |K| = 2e$ and so we let each of $(A, \mathcal{A}), (F, \mathcal{F}), (G, \mathcal{G}), (H, \mathcal{H})$ and (K, \mathcal{K}) be an equitable 2-chromatic e -star system on the sets A, F, G, H and K respectively such that the colour classes are in agreement with the colouring that we have forced upon V .

Finally, let $\hat{B} = \mathcal{A} \cup \mathcal{F} \cup \mathcal{G} \cup \mathcal{H} \cup \mathcal{K} \cup \left(\bigcup_{i=1}^\ell \mathcal{A}_i \right) \cup \left(\bigcup_{i=1}^\ell \mathcal{P}_i \right) \cup \left(\bigcup_{j=2}^\ell \mathcal{U}_j^1 \right) \cup \left(\bigcup_{j=3}^\ell \mathcal{U}_j^2 \right) \cup \dots \cup (\mathcal{U}_\ell^{\ell-1}) \cup \left(\bigcup_{j=1}^2 \mathcal{K}^j \right) \cup \mathcal{L} \cup \mathcal{E} \cup \mathcal{M} \cup \mathcal{O} \cup \mathcal{Q} \cup \mathcal{S} \cup \mathcal{C} \cup \left(\bigcup_{j=1}^2 \mathcal{F}^j \right) \cup \left(\bigcup_{i=1}^\ell \mathcal{H}_i \right)$. Then (\hat{V}, \hat{B}) is an e -star system of order $n = 10e + \ell n_0$ which is strongly equitable uniquely 2-chromatic. \square

Observe that each of the cardinalities of the colour classes of the uniquely 2-chromatic e -star systems constructed from Theorem 4.1 is greater than e . We tacitly use this property to construct a uniquely 2-chromatic e -star system from a smaller uniquely 2-chromatic e -star system.

Theorem 4.2 *Let (V, \mathcal{B}) be a strongly equitable uniquely 2-chromatic e -star system of order n_0 constructed from Theorem 4.1 with colour classes C_1 and C_2 . Then there exists a uniquely 2-chromatic e -star system for all $n > n_0$ such that $n \equiv 0, 1 \pmod{2e}$.*

Proof. For $s \in \{1, 2\}$, let $C_s = \{c_1^s, \dots, c_{|C_s|}^s\}$. Since $n_0 \equiv 0 \pmod{2e}$, then let $n_0 = 2et$, $t \geq 1$.

First, we construct a uniquely 2-chromatic $S_e(2et + 1)$, (\hat{V}, \hat{B}) , from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{2et + 1\}$. Let $\mathcal{J}_1^{2et+1} = \{\{2et + 1; c_1^2, \dots, c_e^2\}\}$ and $\mathcal{J}_2^{2et+1} = \{\{2et + 1; c_1^1, \dots, c_{e-1}^1, c_{e+1}^2\}, \{2et + 1; c_e^1, \dots, c_{2e-2}^1, c_{e+2}^2\}, \dots, \{2et + 1; c_{(t-1)(e-1)+1}^1, \dots, c_{t(e-1)}^1, c_{e+t}^2\}, \{2et + 1; c_{t(e-1)+1}^1, \dots, c_{|C_1|}^1, c_{e+t+1}^2, \dots, c_{e+t+r}^2\}\}$ where $t = \lfloor \frac{|C_1|}{e-1} \rfloor$ and $r = e - (|C_1| - t(e-1))$.

Partition the set $C_2 \setminus \{c_1^2, \dots, c_e^2, c_{e+1}^2, c_{e+2}^2, \dots, c_{e+t+r}^2\}$ into a set \mathbb{E} of m disjoint e -subsets X_1, \dots, X_m where $m = \frac{|C_2| - (e+t+r)}{e}$. Let $\mathcal{T}_3^{2et+1} = \bigcup_{i=1}^m \{\{2et+1; X_i\}\}$.

Let $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}_1^{2et+1} \cup \mathcal{T}_2^{2et+1} \cup \mathcal{T}_3^{2et+1}$. Then $(\hat{V}, \hat{\mathcal{B}})$ is a uniquely 2-chromatic e -star system of order $2et+1$ with colour classes $C_1 \cup \{2et+1\}$ and C_2 .

Next, we construct a uniquely 2-chromatic $S_e(2et+2e)$, $(\mathring{V}, \mathring{\mathcal{B}})$, from $(\hat{V}, \hat{\mathcal{B}})$. Let $\mathring{V} = \hat{V} \cup V'$ where $V' = \{2et+2, 2et+3, \dots, 2et+2e\}$ and let $v_0 \in C_1 \setminus \{c_1^1, \dots, c_e^1\}$. Let $(V' \cup v_0, \mathcal{B}')$ be a 2-chromatic e -star system of order $2e$ constructed from Theorem 3.1. Let $\mathcal{T}_1^i = \bigcup_i \{\{i; c_1^2, \dots, c_e^2\}\}$, $i \in \{2et+3, 2et+5, \dots, 2et+2e-1\}$ and $\mathcal{T}_1^j = \bigcup_j \{\{j; c_1^1, \dots, c_e^1\}\}$, $j \in \{2et+2, 2et+4, \dots, 2et+2e\}$. For any $k \in \{2et+2, 2et+3, \dots, 2et+2e\}$, decompose the remaining edges between vertex k and set $V \setminus \{v_0\}$ into sets of e -stars \mathcal{T}_2^k and \mathcal{T}_3^k in a manner similar to the construction of $(\hat{V}, \hat{\mathcal{B}})$ from (V, \mathcal{B}) .

Let $\mathring{\mathcal{B}} = \hat{\mathcal{B}} \cup \mathcal{B}' \cup \left(\bigcup_{k=2et+2}^{2et+2e} \mathcal{T}_1^k \right) \cup \left(\bigcup_{k=2et+2}^{2et+2e} \mathcal{T}_2^k \right) \cup \left(\bigcup_{k=2et+2}^{2et+2e} \mathcal{T}_3^k \right)$. Then $(\mathring{V}, \mathring{\mathcal{B}})$ is a uniquely 2-chromatic e -star system of order $2et+2e$ with colour classes $C_1 \cup \{2et+3, 2et+5, \dots, 2et+2e-1\}$ and $C_2 \cup \{2et+2, 2et+4, \dots, 2et+2e\}$. \square

We then obtain the following corollary.

Corollary 4.1 *There exists some integer n_0 where $n_0 \equiv 0 \pmod{2e}$ such that for all admissible $n \geq n_0$ where $n \equiv 0, 1 \pmod{2e}$, there exists a uniquely 2-chromatic e -star system of order n for any $e \geq 3$.*

Proof. Apply Theorem 4.1 and Theorem 4.2. \square

We now show how to construct a strongly equitable k -chromatic e -star system from a strongly equitable uniquely $(k-1)$ -chromatic e -star system.

Theorem 4.3 *Let $k \geq 3$ and $e \geq 3$. If there exists a strongly equitable uniquely $(k-1)$ -chromatic e -star system of order $n_{k-1} \equiv 0 \pmod{2e}$ with colour classes C_1, \dots, C_{k-1} such that $|C_i| > e$, $1 \leq i \leq k-1$, then there exists a strongly equitable k -chromatic e -star system of order $n_k \equiv 0 \pmod{2e}$.*

Proof. Let (U_0, \mathcal{A}_0) be a strongly equitable uniquely $(k-1)$ -chromatic e -star system of order n_{k-1} with colour classes C_1, \dots, C_{k-1} such that $n_{k-1} \equiv 0 \pmod{2e}$, $|C_1| = \dots = |C_{k-1}| = r > e$ where r is a positive integer and each vertex of C_s has colour s , for $1 \leq s \leq k-1$. For each $s \in \{1, \dots, k-1\}$, let $C_s = \{c_1^s, \dots, c_r^s\}$. For each $i \in \{1, \dots, k\}$, let $U_i = U_0 \times \{i\}$ and $\mathcal{A}_i = \mathcal{A}_0 \times \{i\}$ where $\mathcal{A}_0 \times \{i\}$ denotes the set $\{S \times \{i\} \mid S \in \mathcal{A}_0\}$. So (U_i, \mathcal{A}_i) has colour classes $C_1 \times \{i\}, \dots, C_{k-1} \times \{i\}$. We construct a strongly equitable k -chromatic e -star system (V, \mathcal{B})

where $V = \bigcup_{i=1}^k U_i$. We need to decompose the edges between the subsets of V into e -stars such that the resulting e -star system (V, \mathcal{B}) is strongly equitably k -chromatic.

First, we decompose the edges between U_1 and U_2 . For each $s \in \{1, \dots, k-1\}$, let $D_s = \{c_1^s \times \{1\}, \dots, c_e^s \times \{1\}\}$. Let $\mathcal{F}_1^1 = \bigcup_{j=1}^r \{\{c_j^1 \times \{2\}; D_1\}, \{c_j^1 \times \{2\}; D_2\}, \dots, \{c_j^1 \times \{2\}; D_{k-1}\}\}$. Partition $U_1 \setminus \bigcup_{s=1}^{k-1} D_s$ into $m = \frac{n_{k-1}-e(k-1)}{e}$ sets G_1, \dots, G_m such that for each $\ell \in \{1, \dots, m\}$, the vertices of G_ℓ do not all have the same colour. Let $\mathcal{F}_1^2 = \bigcup_{j=1}^r \{\{c_j^1 \times \{2\}; G_1\}, \{c_j^1 \times \{2\}; G_2\}, \dots, \{c_j^1 \times \{2\}; G_m\}\}$ and $\mathcal{F}_1 = \mathcal{F}_1^1 \cup \mathcal{F}_1^2$. Then \mathcal{F}_1 is a decomposition of the edges between $C_1 \times \{2\}$ and U_1 into e -stars. In any putative $(k-1)$ -colouring of the e -star system that we are building, the colouring of U_1 is unique since (U_1, \mathcal{A}_1) is uniquely $(k-1)$ -chromatic. The uniqueness of the $(k-1)$ -colouring of (U_1, \mathcal{A}_1) is such that all vertices of D_s ($1 \leq s \leq k-1$) have colour s . But now some star of \mathcal{F}_1^1 is monochromatic if any vertex of $C_1 \times \{2\}$ has colour s for some $s \in \{1, \dots, k-1\}$. Therefore the system that we are building is not $(k-1)$ -colourable. By demonstrating a k -colouring, we establish that it is k -chromatic. Without loss of generality, actually colour each vertex of D_s with colour s and each vertex of $C_1 \times \{2\}$ with colour k .

Let $\mathcal{F}_2^1 = \bigcup_{j=1}^r \{\{c_j^2 \times \{2\}; D_1\}, \{c_j^2 \times \{2\}; D_3\}, \{c_j^2 \times \{2\}; D_4\}, \dots, \{c_j^2 \times \{2\}; D_{k-1}\}\}$. Partition $U_1 \setminus (D_1 \cup (\bigcup_{s=3}^{k-1} D_s))$ into $m = \frac{n_{k-1}-e(k-2)}{e}$ sets G_1, \dots, G_m such that for each $\ell \in \{1, \dots, m\}$, the vertices of G_ℓ do not all have the same colour. Let $\mathcal{F}_2^2 = \bigcup_{j=1}^r \{\{c_j^2 \times \{2\}; G_1\}, \dots, \{c_j^2 \times \{2\}; G_m\}\}$ and $\mathcal{F}_2 = \mathcal{F}_2^1 \cup \mathcal{F}_2^2$. Then \mathcal{F}_2 is a decomposition of the edges between $C_2 \times \{2\}$ and U_1 into e -stars such that all the vertices in $C_2 \times \{2\}$ are forced to have colour 2. For each $s \in \{3, \dots, k-1\}$, decompose the edges between $C_s \times \{2\}$ and U_1 into a set of e -stars \mathcal{F}_s in a similar manner such that all the vertices in $C_s \times \{2\}$ are forced to have colour s .

Let $\mathcal{U}_2^1 = \bigcup_{s=1}^{k-1} \mathcal{F}_s$. Then \mathcal{U}_2^1 is a decomposition of the edges between U_1 and U_2 into e -stars forcing (U_2, \mathcal{A}_2) to have colour classes $C_2^2, C_2^3, \dots, C_2^k$ of equal size with no vertex of colour 1, where $C_2^2 = C_2 \times \{2\}$, $C_2^3 = C_3 \times \{2\}$, \dots , $C_2^k = C_1 \times \{2\}$. For each $i \in \{3, \dots, k\}$, decompose the edges between U_1 and U_i into a set of e -stars \mathcal{U}_i^1 in a similar manner forcing (U_i, \mathcal{A}_i) to have colour classes $C_i^1, C_i^2, \dots, C_i^{i-2}, C_i^i, \dots, C_i^k$ of equal size with no vertex of colour $i-1$, where $C_i^1 = C_1 \times \{i\}$, $C_i^2 = C_2 \times \{i\}$, \dots , $C_i^{i-2} = C_{i-2} \times \{i\}$, $C_i^i = C_i \times \{i\}$, \dots , $C_i^k = C_{i-1} \times \{i\}$.

Next, for each $j \in \{3, \dots, k\}$, we decompose the edges between U_2 and U_j . Partition U_2 into $m = \frac{n_{k-1}}{e}$ sets G_1, \dots, G_m such that for each $\ell \in \{1, \dots, m\}$, the vertices of G_ℓ do not all have the same colour. For each $j \in \{3, \dots, k\}$, let $\mathcal{U}_j^2 = \bigcup_{u \in U_j} \{\{u; G_1\}, \{u; G_2\}, \dots, \{u; G_m\}\}$. Then \mathcal{U}_j^2 is a decomposition of the edges between U_2 and U_j into e -stars such that no e -star is monochromatic. For each $3 \leq i < j \leq k$, decompose the edges between U_i and U_j into a set of

e -stars \mathcal{U}_j^i in a similar manner.

Let $\mathcal{B} = (\bigcup_{i=1}^k \mathcal{A}_i) \cup (\bigcup_{j=2}^k \mathcal{U}_j^1) \cup (\bigcup_{j=3}^k \mathcal{U}_j^2) \cup \dots \cup \mathcal{U}_k^{k-1}$ and $C_1^1 = C_1 \times \{1\}$, $C_1^2 = C_2 \times \{1\}$, \dots , $C_1^{k-1} = C_{k-1} \times \{1\}$. Then (V, \mathcal{B}) is a strongly equitable k -chromatic e -star system of order $n_k = kn_{k-1}$ with colour classes $C_1^1 \cup C_3^1 \cup C_4^1 \cup \dots \cup C_k^1$, $C_1^2 \cup C_2^2 \cup C_4^2 \cup \dots \cup C_k^2$, \dots , $C_1^{k-1} \cup C_2^{k-1} \cup \dots \cup C_{k-1}^{k-1}$, $C_2^k \cup C_3^k \cup \dots \cup C_k^k$ where each colour class is of size $(k-1)r$. \square

We now show how to construct a strongly equitable uniquely k -chromatic e -star system from a strongly equitable k -chromatic e -star system.

Theorem 4.4 *Let $k \geq 3$ and $e \geq 3$. If there exists a strongly equitable k -chromatic e -star system of order $n_0 \equiv 0 \pmod{2e}$, then there exists a strongly equitable uniquely k -chromatic e -star system of order n for some $n \equiv 0 \pmod{2e}$.*

Proof. Let (U_0, \mathcal{A}_0) be a strongly equitable k -chromatic e -star system of order $n_0 \equiv 0 \pmod{2e}$ with colour classes C_1, \dots, C_k such that each vertex of C_s has colour s , for $s = 1, \dots, k$. For a positive integer ℓ that will be fixed later and for each $i \in \{1, \dots, \ell\}$, let $U_i = U_0 \times \{i\}$ and $\mathcal{A}_i = \mathcal{A}_0 \times \{i\}$ where $\mathcal{A}_0 \times \{i\}$ denotes the set $\{S \times \{i\} \mid S \in \mathcal{A}_0\}$. So (U_i, \mathcal{A}_i) has colour classes $C_1 \times \{i\}, \dots, C_k \times \{i\}$. Let $U = \bigcup_{i=1}^{\ell} U_i$. Let $A^1 = \{a_1^1, \dots, a_e^1\}, \dots, A^k = \{a_1^k, \dots, a_e^k\}$, $A_0^1 = \{(a_1)_0^1, \dots, (a_e)_0^1\}$, $A = \bigcup_{i=1}^k A^i$ and $A' = A \cup A_0^1$. Similarly let $F^1 = \{f_1^1, \dots, f_e^1\}, \dots, F^k = \{f_1^k, \dots, f_e^k\}$, $F_0^1 = \{(f_1)_0^1, \dots, (f_e)_0^1\}$, $F = \bigcup_{i=1}^k F^i$ and $F' = F \cup F_0^1$, $G^1 = \{g_1^1, \dots, g_e^1\}, \dots, G^k = \{g_1^k, \dots, g_e^k\}$, $G_0^1 = \{(g_1)_0^1, \dots, (g_e)_0^1\}$, $G = \bigcup_{i=1}^k G^i$ and $G' = G \cup G_0^1$, $H^1 = \{h_1^1, \dots, h_e^1\}, \dots, H^k = \{h_1^k, \dots, h_e^k\}$, $H_0^1 = \{(h_1)_0^1, \dots, (h_e)_0^1\}$, $H = \bigcup_{i=1}^k H^i$ and $H' = H \cup H_0^1$. We construct a uniquely k -chromatic e -star system $(\hat{V}, \hat{\mathcal{B}})$ where $\hat{V} = A \cup F \cup G \cup H \cup U$ if k is even and $\hat{V} = A' \cup F' \cup G' \cup H' \cup U$ if k is odd.

We need to decompose the edges between the subsets of \hat{V} into e -stars such that the resulting e -star system $(\hat{V}, \hat{\mathcal{B}})$ is uniquely k -chromatic.

First, we decompose the edges between A and U into e -stars in a way such that no e -subset in A other than the e -subsets A^1, \dots, A^k are monochromatic in any putative k -colouring of $(\hat{V}, \hat{\mathcal{B}})$. Let \mathcal{D} be the set of all e -subsets of A except A^1, \dots, A^k . The number of e -subsets of the set A is $N = \binom{k}{e}$ and $|\mathcal{D}| = N - k$. Let $a = k - 1$ and $|\mathcal{D}| = aq + r$ where q is a nonnegative integer and $0 \leq r < a$ is an integer. Partition set \mathcal{D} into q sets of disjoint e -subsets $\mathbb{T}_1, \dots, \mathbb{T}_q$ of size a and one set of disjoint e -subsets \mathbb{T}_{q+1} of size r ; such a partition is known to exist by Theorem 1 in [1]. Let $\mathbb{T}_i = \{T_1^i, \dots, T_a^i\}$, $1 \leq i \leq q$ and $\mathbb{T}_{q+1} = \{T_1^{q+1}, \dots, T_r^{q+1}\}$. Let $\tilde{A}_i = A \setminus \bigcup_{j=1}^a T_j^i$, $1 \leq i \leq q$ and $\tilde{A}_{q+1} = A \setminus \bigcup_{j=1}^r T_j^{q+1}$ if $r > 0$ and $\tilde{A}_{q+1} = \emptyset$ if $r = 0$. We now fix $\ell = q + 1$ if $r > 0$ and $\ell = q$ if $r = 0$.

Partition U_0 into e -subsets E_1, \dots, E_m where $m = \frac{n_0}{e}$ such that each E_j has nonempty intersection with at least two colour classes, $1 \leq j \leq m$. For each $i \in \{1, 2, \dots, \ell\}$ and for each $w \in \tilde{A}_i$, we decompose the edges between w and U_i into the set of e -stars $\mathcal{R}_w = \bigcup_{j=1}^m \{\{w; E_j \times \{i\}\}\}$. For each $i \in \{1, \dots, q\}$, let $\mathcal{T}_i = \bigcup_{u \in U_i} \{\{u; T_1^i\}, \dots, \{u; T_a^i\}\}$ and $\mathcal{T}_{q+1} = \bigcup_{u \in U_i} \{\{u; T_1^{q+1}\}, \dots, \{u; T_r^{q+1}\}\}$. Then $\mathcal{P}_i = \mathcal{T}_i \cup \left(\bigcup_{w \in \tilde{A}_i} \mathcal{R}_w \right)$, $1 \leq i \leq \ell$, is a decomposition of the edges between A and U_i and $\bigcup_{i=1}^{\ell} \mathcal{P}_i$ is a decomposition of the edges between A and U into e -stars.

Recall that (U_i, \mathcal{A}_i) is k -chromatic. So for any k -colouring of (U_i, \mathcal{A}_i) there must be a vertex u_s^i of each colour s in U_i , $1 \leq s \leq k$. Note that for each $i \in \{1, \dots, q\}$, $\{u_s^i; T_j^i\} \in \mathcal{T}_i$, $1 \leq j \leq a$, and $\{u_s^{q+1}; T_j^{q+1}\} \in \mathcal{T}_{q+1}$, $1 \leq j \leq r$, and so to ultimately obtain a valid k -colouring of $(\hat{V}, \hat{\mathcal{B}})$, the e vertices of T_j^i and T_j^{q+1} must not all have colour s , $1 \leq s \leq k$. If a colour s occurs on more than e vertices of A then there exists a T_j^i or T_j^{q+1} with only colour s and then $\{u_s^i; T_j^i\}$ or $\{u_s^{q+1}; T_j^{q+1}\}$ would be monochromatic. So in any valid k -colouring of $(\hat{V}, \hat{\mathcal{B}})$, each colour must occur on exactly e points of A . If the points of colour s are not all in A^1 or \dots or A^k then they are the leaves of a monochromatic e -star with centre u_s^i . Without loss of generality we therefore assign colour s to all e points of A^s , $1 \leq s \leq k$.

We now begin to decompose the edges between A and F into e -stars. For each $i \in \{1, \dots, e\}$, let $\mathcal{K}_i^1 = \{\{f_i^1; A^2\}, \dots, \{f_i^1; A^k\}\}$, $\mathcal{K}_i^2 = \{\{f_i^2; A^1\}, \{f_i^2; A^3\}, \dots, \{f_i^2; A^k\}\}$, \dots , $\mathcal{K}_i^k = \{\{f_i^k; A^1\}, \dots, \{f_i^k; A^{k-1}\}\}$. Since each vertex of A^s has colour s , $2 \leq s \leq k$, then each vertex of F^1 must have colour 1 or else a monochromatic e -star would now exist. Likewise, each vertex of F^s must have colour s , $2 \leq s \leq k$. Let $\mathcal{K}^1 = \bigcup_{i=1}^e \mathcal{K}_i^1, \dots, \mathcal{K}^k = \bigcup_{i=1}^e \mathcal{K}_i^k$. We decompose the edges between A^i and F^i , $1 \leq i \leq k$, later. For each $j \in \{1, \dots, k\}$, decompose the edges between A^j and $G^1 \cup \dots \cup G^{j-1} \cup G^{j+1} \cup \dots \cup G^k$ into a set of e -stars \mathcal{L}_j in a similar manner, forcing the vertices of the subsets G^1, \dots, G^k to have colours $1, \dots, k$ respectively. Let $\mathcal{L} = \bigcup_{j=1}^k \mathcal{L}_j$. We decompose the edges between A^i and G^i , $1 \leq i \leq k$, later.

Next, we decompose the edges between H and $F \cup G$. For each $i \in \{1, \dots, e\}$, let $\mathcal{E}_i^1 = \{\{h_i^1; f_1^2, \dots, f_e^2\}, \dots, \{h_i^1; f_1^k, \dots, f_e^k\}, \{h_i^1; f_1^1, \dots, f_{e-1}^1, g_1^2\}, \{h_i^1; f_e^1, g_1^1, \dots, g_{e-2}^1, g_2^2\}, \{h_i^1; g_{e-1}^1, g_1^2, g_2^2, \dots, g_e^2\}, \{h_i^1; g_1^3, \dots, g_e^3\}, \dots, \{h_i^1; g_1^k, \dots, g_e^k\}\}$ and $\mathcal{E}^1 = \bigcup_{i=1}^e \mathcal{E}_i^1$. Then \mathcal{E}^1 is a decomposition of the edges between H^1 and $F \cup G$ into e -stars such that all the vertices in H^1 are forced to have colour 1. For each $s \in \{2, \dots, k\}$, decompose the edges between H^s and $F \cup G$ into a set of e -stars \mathcal{E}^s in a similar manner, forcing every vertex in H^s to have unique colour s . Let $\mathcal{E} = \bigcup_{s=1}^k \mathcal{E}^s$.

Next, we decompose the edges between A^1 and $F^1 \cup G^1 \cup H$. For each $i \in \{1, \dots, e\}$, let $\mathcal{M}_i^1 =$

$\{\{a_i^1; f_1^1, \dots, f_{e-1}^1, h_1^2\}, \{a_i^1; f_e^1, g_1^1, \dots, g_{e-2}^1, h_2^2\}, \{a_i^1; g_{e-1}^1, g_e^1, h_3^2, \dots, h_e^2\}, \{a_i^1; h_1^1, \dots, h_{e-1}^1, h_1^3\}, \{a_i^1; h_e^1, h_2^3, \dots, h_e^3\}, \{a_i^1; h_1^4, \dots, h_e^4\}, \dots, \{a_i^1; h_1^k, \dots, h_e^k\}\}$. Then $\mathcal{M}^1 = \bigcup_{i=1}^e \mathcal{M}_i^1$ is a decomposition of the edges between A^1 and $F^1 \cup G^1 \cup H$ into e -stars, none of which is monochromatic. For each $j \in \{2, \dots, k\}$, decompose the edges between A^j and $F^j \cup G^j \cup H$ into a set of e -stars \mathcal{M}^j in a similar manner. Let $\mathcal{M} = \bigcup_{j=1}^k \mathcal{M}^j$.

Next, for each $1 \leq i \leq \ell$, we decompose the edges between $F \cup G$ and U_i into e -stars. Let u_1^s, \dots, u_e^s be the vertices of colour s , $1 \leq s \leq k$, in an equitable k -colouring of (U_0, \mathcal{A}_0) . For each $j \in \{1, \dots, e\}$, let $\mathcal{N}_j^1 = \{\{u_j^1 \times \{i\}; f_1^2, \dots, f_e^2\}, \dots, \{u_j^1 \times \{i\}; f_1^k, \dots, f_e^k\}, \{u_j^1 \times \{i\}; f_1^1, \dots, f_{e-1}^1, g_1^2\}, \{u_j^1 \times \{i\}; f_e^1, g_1^1, \dots, g_{e-2}^1, g_2^2\}, \{u_j^1 \times \{i\}; g_{e-1}^1, g_e^1, g_3^2, \dots, g_e^2\}, \{u_j^1 \times \{i\}; g_1^3, \dots, g_e^3\}, \dots, \{u_j^1 \times \{i\}; g_1^k, \dots, g_e^k\}\}$ and observe that $u_j^1 \times \{i\}$ is now forced to have colour 1. For each $j \in \{1, \dots, e\}$ and each $s \in \{2, \dots, k\}$, decompose the edges between $u_j^s \times \{i\}$ and $F \cup G$ into a set of e -stars \mathcal{N}_j^s in a similar manner so that $u_j^s \times \{i\}$ is forced to have colour s . Let $\mathcal{N}^s = \bigcup_{j=1}^e \mathcal{N}_j^s$ and $\mathcal{O}_i = \bigcup_{s=1}^k \mathcal{N}^s$. Then \mathcal{O}_i is a decomposition of the edges between $F \cup G$ and U_i into e -stars which forces every vertex of U_i to have a unique colour. Let $\mathcal{O} = \bigcup_{i=1}^{\ell} \mathcal{O}_i$.

At this point every vertex of \hat{V} is coloured, and the colouring is unique (up to a permutation of the colours). Moreover, the colour classes are equal in size and so the colouring is strongly equitable. All that remains is to complete the decomposition without introducing monochromatic e -stars.

For each $1 \leq i \leq e$, we begin to decompose the edges between F and G into e -stars. Let $\mathcal{Q}_i^1 = \{\{f_i^1; g_1^1, \dots, g_{e-1}^1, g_1^2\}, \{f_i^1; g_e^1, g_2^2, \dots, g_e^2\}\}, \{f_i^1; g_1^3, \dots, g_e^3\}, \dots, \{f_i^1; g_1^k, \dots, g_e^k\}\}$. Then $\mathcal{Q}^1 = \bigcup_{i=1}^e \mathcal{Q}_i^1$ is a decomposition of the edges between F^1 and G into e -stars, none of which is monochromatic. For each $j \in \{2, \dots, k\}$, decompose the edges between F^j and G into a set of e -stars \mathcal{Q}^j in a similar manner. Let $\mathcal{Q} = \bigcup_{j=1}^k \mathcal{Q}^j$.

Next, for each $1 \leq i \leq \ell$, we decompose the edges between H and U_i into e -stars. Recall that u_1^s, \dots, u_e^s are the vertices of colour s , $1 \leq s \leq k$. Let $\mathcal{J}_j^1 = \{\{u_j^1 \times \{i\}; h_1^1, \dots, h_{e-1}^1, h_1^2\}, \{u_j^1 \times \{i\}; h_e^1, h_2^2, \dots, h_e^2\}, \{u_j^1 \times \{i\}; h_1^3, \dots, h_e^3\}, \dots, \{u_j^1 \times \{i\}; h_1^k, \dots, h_e^k\}\}$, $1 \leq j \leq e$. For each $j \in \{1, \dots, e\}$ and each $s \in \{2, \dots, k\}$, decompose the edges between $u_j^s \times \{i\}$ and H into a set of e -stars \mathcal{J}_j^s in a similar manner. Let $\mathcal{J}^s = \bigcup_{j=1}^e \mathcal{J}_j^s$ and $\mathcal{S}_i = \bigcup_{s=1}^k \mathcal{J}^s$. Then \mathcal{S}_i is a decomposition of the edges between H and U_i into e -stars. Let $\mathcal{S} = \bigcup_{i=1}^{\ell} \mathcal{S}_i$. The edges between U_i and U_j , $1 \leq i < j \leq \ell$, are decomposed into set of e -stars \mathcal{U}_j^i in a similar manner.

Note that $|A| = |F| = |G| = |H| = ke$.

Case1. k is even. We let each of $(A, \mathcal{A}), (F, \mathcal{F}), (G, \mathcal{G})$, and (H, \mathcal{H}) be strongly equitable k -chromatic e -star systems on the sets A, F, G , and H respectively such that the colour classes

are in agreement with the colouring that we have forced upon \hat{V} . Let $\hat{\mathcal{B}} = \mathcal{A} \cup \mathcal{F} \cup \mathcal{G} \cup \mathcal{H} \cup (\bigcup_{i=1}^{\ell} \mathcal{A}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{P}_i) \cup (\bigcup_{j=2}^{\ell} \mathcal{U}_j^1) \cup (\bigcup_{j=3}^{\ell} \mathcal{U}_j^2) \cup \dots \cup (\mathcal{U}_{\ell}^{\ell-1}) \cup (\bigcup_{j=1}^k \mathcal{K}^j) \cup \mathcal{L} \cup \mathcal{E} \cup \mathcal{M} \cup \mathcal{O} \cup \mathcal{Q} \cup \mathcal{S}$. Then $(\hat{V}, \hat{\mathcal{B}})$ is an e -star system of order $n = 4ke + \ell n_0$ which is uniquely k -chromatic and strongly equitable.

Case 2. k is odd. For $j \in \{1, \dots, e\}$, let $\mathcal{V}_j = \{ \{(a_j)_0^1; f_1^2, \dots, f_e^2\}, \{(a_j)_0^1; f_1^3, \dots, f_e^3\}, \dots, \{(a_j)_0^1; f_1^k, \dots, f_e^k\}, \{(a_j)_0^1; f_1^1, \dots, f_{e-1}^1, g_2^2\}, \{(a_j)_0^1; f_e^1, g_1^1, \dots, g_{e-2}^1, g_2^2\}, \{(a_j)_0^1; g_{e-1}^1, g_e^1, g_3^2, \dots, g_e^2\}, \{(a_j)_0^1; g_1^3, \dots, g_e^3\}, \dots, \{(a_j)_0^1; g_1^k, \dots, g_e^k\} \cup \{ \{(a_j)_0^1; h_1^1, \dots, h_{e-1}^1, h_1^2\}, \{(a_j)_0^1; h_e^1, h_2^2, \dots, h_{e-1}^2\}, \{(a_j)_0^1; h_1^3, \dots, h_e^3\}, \dots, \{(a_j)_0^1; h_1^k, \dots, h_e^k\} \} \cup \bigcup_{i=1}^{\ell} (\{ \{(a_j)_0^1; u_1^1 \times \{i\}, u_2^1 \times \{i\}, \dots, u_{e-1}^1 \times \{i\}, u_1^2 \times \{i\} \}, \dots, \{(a_j)_0^1; u_{(t-1)(e-1)+1}^1 \times \{i\}, \dots, u_{t(e-1)}^1 \times \{i\}, u_t^2 \times \{i\} \}, \{(a_j)_0^1; u_{t(e-1)+1}^1 \times \{i\}, \dots, u_{|C_1|}^1 \times \{i\}, u_{t+1}^2 \times \{i\}, \dots, u_{t+m}^2 \times \{i\} \} \}) \cup \bigcup_{i=1}^{\ell} (\{ \{(a_j)_0^1; x_1^d \times \{i\}, \dots, x_e^d \times \{i\} \} \mid 1 \leq d \leq \frac{n_0 - (|C_1| + t + m)}{e} \})$, where t and m are positive integers and the sets $\{x_1^d, \dots, x_e^d\}$, $1 \leq d \leq \frac{n_0 - (|C_1| + t + m)}{e}$, form a partition of $U_i \setminus (C_1 \times \{i\}) \cup \{u_1^2 \times \{i\}, u_2^2 \times \{i\}, \dots, u_{t+m}^2 \times \{i\}\}$. Let $\mathcal{V} = \bigcup_{j=1}^e \mathcal{V}_j$, then \mathcal{V} is a decomposition of the edges between A_0^1 and $F \cup G \cup H \cup U$ into e -stars forcing the vertices of A_0^1 to have unique colour 1.

Decompose the edges between F_0^1 and $A' \cup G \cup H \cup U$ into set of e -stars \mathcal{W} in a similar manner so that the vertices of F_0^1 are forced to have colour 1. Also, decompose the edges between G_0^1 and $A' \cup F' \cup H \cup U$ into set of e -stars \mathcal{X} so that the vertices of G_0^1 are forced to have colour 1 and the edges between H_0^1 and $A' \cup F' \cup G' \cup U$ into set of e -stars \mathcal{Y} so that the vertices of H_0^1 are forced to have colour 1.

Now, we decompose the edges of the complete graph on set A' into e -stars. We first take 2-chromatic e -star systems $Z_1 = (A^1 \cup A^2, \mathcal{A}_2^1)$, $Z_2 = (A^3 \cup A^4, \mathcal{A}_4^3)$, \dots , $Z_{\frac{k+1}{2}} = (A^k \cup A_0^1, \mathcal{A}_4^k)$ of size $2e$. It is easy then to decompose the edges between Z_i and Z_j into set of e -stars \mathcal{Z}_{ij}^i , $1 \leq i < j \leq \frac{k+1}{2}$ such that no e -star is monochromatic. Then (A', \mathcal{A}') where $\mathcal{A}' = \mathcal{A}_2^1 \cup \mathcal{A}_4^3 \cup \dots \cup \mathcal{A}_1^k \cup (\bigcup_{j=2}^{\frac{k+1}{2}} \mathcal{Z}_{ij}^1) \cup (\bigcup_{j=3}^{\frac{k+1}{2}} \mathcal{Z}_{ij}^2) \cup \dots \cup \mathcal{Z}_{\frac{k+1}{2}}^{\frac{k+1}{2}-1}$ is an e -star system which is uniquely k -chromatic. Decompose the edges of the complete graph on the sets F' , G' and H' into set of e -stars \mathcal{F}' , \mathcal{G}' and \mathcal{H}' in a similar manner.

Let $\hat{\mathcal{B}} = \mathcal{A}' \cup \mathcal{F}' \cup \mathcal{G}' \cup \mathcal{H}' \cup (\bigcup_{i=1}^{\ell} \mathcal{A}_i) \cup (\bigcup_{i=1}^{\ell} \mathcal{P}_i) \cup (\bigcup_{j=2}^{\ell} \mathcal{U}_j^1) \cup (\bigcup_{j=3}^{\ell} \mathcal{U}_j^2) \cup \dots \cup (\mathcal{U}_{\ell}^{\ell-1}) \cup (\bigcup_{j=1}^k \mathcal{K}^j) \cup \mathcal{L} \cup \mathcal{E} \cup \mathcal{M} \cup \mathcal{O} \cup \mathcal{Q} \cup \mathcal{S} \cup \mathcal{V} \cup \mathcal{W} \cup \mathcal{X} \cup \mathcal{Y}$. Then $(\hat{V}, \hat{\mathcal{B}})$ is an e -star system of order $n = 4e(k+1) + \ell n_0$ which is uniquely k -chromatic and strongly equitable. \square

We now show how to construct a uniquely k -chromatic e -star system from a smaller uniquely k -chromatic e -star system.

Theorem 4.5 *For any $k \geq 3$, let (V, \mathcal{B}) be a strongly equitable uniquely k -chromatic e -star*

system of order n_0 constructed from Theorem 4.4 with colour classes C_1, \dots, C_k . Then there exists a uniquely k -chromatic e -star system for all $n > n_0$ such that $n \equiv 0, 1 \pmod{2e}$.

Proof. For $s \in \{1, \dots, k\}$, let $C_s = \{c_1^s, \dots, c_{|C_s|}^s\}$. Since $n_0 \equiv 0 \pmod{2e}$, then let $n_0 = 2et$, $t \geq 1$.

First, we construct a uniquely k -chromatic $S_e(2et + 1)$, $(\hat{V}, \hat{\mathcal{B}})$, from (V, \mathcal{B}) . Let $\hat{V} = V \cup \{2et + 1\}$. Let $\mathcal{T}_1^{2et+1} = \{\{2et+1; c_1^2, \dots, c_e^2\}, \{2et+1; c_1^3, \dots, c_e^3\}, \dots, \{2et+1; c_1^k, \dots, c_e^k\}\}$ and $\mathcal{T}_2^{2et+1} = \{\{2et+1; c_1^1, \dots, c_{e-1}^1, c_{e+1}^2\}, \{2et+1; c_e^1, \dots, c_{2e-2}^1, c_{e+2}^2\}, \dots, \{2et+1; c_{(t-1)(e-1)+1}^1, \dots, c_{t(e-1)}^1, c_{e+t}^2\}, \{2et+1; c_{t(e-1)+1}^1, \dots, c_{|C_1|}^1, c_{e+t+1}^2, \dots, c_{e+t+r}^2\}\}$ where $t = \lfloor \frac{|C_1|}{e-1} \rfloor$ and $r = e - (|C_1| - t(e-1))$.

Partition the set $(C_2 \setminus \{c_1^2, \dots, c_e^2, c_{e+1}^2, c_{e+2}^2, \dots, c_{e+t+r}^2\}) \cup (C_3 \setminus \{c_1^3, \dots, c_e^3\}) \cup \dots \cup (C_k \setminus \{c_1^k, \dots, c_e^k\})$ into a set \mathbb{E} of m disjoint e -subsets X_1, \dots, X_m where $m = \frac{(|C_2| - (e+t+r)) + (|C_3| - e) + \dots + (|C_k| - e)}{e}$. Let $\mathcal{T}_3^{2et+1} = \bigcup_{i=1}^m \{\{2et+1; X_i\}\}$.

Let $\hat{\mathcal{B}} = \mathcal{B} \cup \mathcal{T}_1^{2et+1} \cup \mathcal{T}_2^{2et+1} \cup \mathcal{T}_3^{2et+1}$. Then $(\hat{V}, \hat{\mathcal{B}})$ is a uniquely k -chromatic e -star system of order $2et + 1$ with colour classes $C_1 \cup \{2et + 1\}, C_2, \dots, C_k$.

Next, we construct a uniquely k -chromatic $S_e(2et + 2e)$, $(\hat{V}, \hat{\mathcal{B}})$, from $(\hat{V}, \hat{\mathcal{B}})$. Let $\hat{V} = \hat{V} \cup V'$ where $V' = \{2et + 2, 2et + 3, \dots, 2et + 2e\}$ and let $v_0 \in C_1 \setminus \{c_1^1, \dots, c_e^1\}$. Let $(V' \cup \{v_0\}, \mathcal{B}')$ be a strongly equitable 2-chromatic e -star system of order $2e$ constructed from Theorem 3.1. Let $\mathcal{T}_1^i = \bigcup_i \{\{i; c_1^2, \dots, c_e^2\}, \{i; c_1^3, \dots, c_e^3\}, \dots, \{i; c_1^k, \dots, c_e^k\}\}$, $i \in \{2et + 3, 2et + 5, \dots, 2et + 2e - 1\}$ and $\mathcal{T}_1^j = \bigcup_j \{\{j; c_1^1, \dots, c_e^1\}, \{j; c_1^3, \dots, c_e^3\}, \dots, \{j; c_1^k, \dots, c_e^k\}\}$, $j \in \{2et + 2, 2et + 4, \dots, 2et + 2e\}$. For any $k \in \{2et + 2, 2et + 3, \dots, 2et + 2e\}$, decompose the remaining edges between vertex k and set $V \setminus \{v_0\}$ into sets of e -stars \mathcal{T}_2^k and \mathcal{T}_3^k in a manner similar to the construction of \mathcal{T}_2^{2et+1} and \mathcal{T}_3^{2et+1} in $(\hat{V}, \hat{\mathcal{B}})$.

Let $\hat{\mathcal{B}} = \hat{\mathcal{B}} \cup \mathcal{B}' \cup \left(\bigcup_{k=2et+2}^{2et+2e} \mathcal{T}_1^k \right) \cup \left(\bigcup_{k=2et+2}^{2et+2e} \mathcal{T}_2^k \right) \cup \left(\bigcup_{k=2et+2}^{2et+2e} \mathcal{T}_3^k \right)$. Then $(\hat{V}, \hat{\mathcal{B}})$ is a uniquely k -chromatic e -star system of order $2et + 2e$ with colour classes $C_1 \cup \{2et + 3, 2et + 5, \dots, 2et + 2e - 1\}, C_2 \cup \{2et + 2, 2et + 4, \dots, 2et + 2e\}, C_3, \dots, C_k$. \square

We then conclude with the following corollary.

Corollary 4.2 *There exists some integer n_0 where $n_0 \equiv 0 \pmod{2e}$ such that for all admissible $n \geq n_0$ where $n \equiv 0, 1 \pmod{2e}$, there exists a uniquely k -chromatic e -star system of order n for any $e \geq 3$ and $k \geq 3$.*

Proof. Apply Theorem 4.4 and Theorem 4.5. \square

5 Acknowledgements

D.A. Pike acknowledges research support from NSERC (grant number RGPIN-04456-2016).

References

- [1] Zs. Baranyai, (1975), On the factorization of the complete uniform hypergraph, in Hajnal, A.; Rado, R.; Sós, V. T. (eds.), *Infinite and Finite Sets, Proc. Coll. Keszthely*, 1973, Colloquia Math. Soc. János Bolyai, 10, North-Holland, pp. 91–107.
- [2] M. de Brandes, K. T. Phelps and V. Rödl, Colouring Steiner triple systems, *SIAM J. Algebraic Discrete Methods*. 3 (1982), 241–249.
- [3] A. C. Burgess and D. A. Pike, Colouring even cycle systems, *Discrete Math.* 308 (2008), 962–973.
- [4] A. D. Forbes, Uniquely 3-colourable Steiner triple systems, *J. Combin. Theory Ser. A*. 101 (2003) 49–68.
- [5] D. Horsley and D. A. Pike, On cycle systems with specified weak chromatic number, *J. Combin. Theory Ser. A*. 117 (2010) 1195–1206.
- [6] S. Yamamoto, H. Ikeda, S. Shige-eda, K. Ushio, and N. Hamada, On claw-decompositions of complete graphs and complete bigraphs, *Hiroshima Math. J.* 5 (1975), 33–42.
- [7] M. L. Yu, On tree factorizations of K_n , *J. Graph Theory* 17 (1993), 713–725.