

ON EHRHART POSITIVITY OF TESLER POLYTOPES AND THEIR DEFORMATIONS

YONGGYU LEE AND FU LIU

ABSTRACT. For $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$, the Tesler polytope $\text{Tes}_n(\mathbf{a})$ is the set of upper triangular matrices with non-negative entries whose hook sum vector is \mathbf{a} . Recently, Morales conjectured that $\text{Tes}_n(1, \dots, 1)$ and $\text{Tes}_n(1, 0, \dots, 0)$ are Ehrhart positive for any positive integer n . In this paper, we consider a certain unimodular copy of $\text{Tes}_n(\mathbf{a})$ and show that the majority of the faces of this unimodular copy have positive values under a function constructed by Berline-Vergne. As a consequence, we prove that the 3rd and 4th coefficients of the Ehrhart polynomial of $\text{Tes}_n(1, \dots, 1)$ are positive for any n . Using the Reduction Theorem by Castillo and the second author, this result generalizes to any deformations of $\text{Tes}_n(1, \dots, 1)$ which includes $\text{Tes}_n(\mathbf{a})$ for all $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$. Furthermore, we give a characterization of which flow polytopes are deformations of $\text{Tes}_n(1, \dots, 1)$.

1. INTRODUCTION

A subset P of \mathbb{R}^n is a *polyhedron* if P is the intersection of finitely many half spaces, usually defined by linear inequalities. A *polytope* is a bounded polyhedron. Equivalently, a polytope $P \subset \mathbb{R}^n$ may be defined as the convex hull of finitely many points in \mathbb{R}^n . We assume that readers are familiar with the basic concepts related to polytopes, such as *face* and *dimension*, presented in [2, 27].

A polytope $P \subset \mathbb{R}^n$ is called *integral* if all of its vertices are integer points, i.e., points in \mathbb{Z}^n . In 1962, Ehrhart discovered that for any integral polytope P of dimension d , the function E_P which maps any non-negative integer $t \in \mathbb{Z}_{\geq 0}$ to the number of integer points in tP (the t -th dilate of P) is a polynomial in t of degree d . We call E_P the *Ehrhart polynomial* of P . For each $1 \leq i \leq d$, let $e_i(P)$ be the coefficient of t^i in $E_P(t)$, so

$$E_P(t) = e_d(P)t^d + e_{d-1}(P)t^{d-1} + \dots + e_0(P).$$

In [12], Ehrhart showed that for any d -dimensional polytope P , the leading coefficient $e_d(P)$ of the Ehrhart polynomial $E_P(t)$ is the normalized volume of P , the second coefficient $e_{d-1}(P)$ is one half of the sum of the normalized volumes of the facets of P , and the constant term $e_0(P)$ is 1. Thus, these three coefficients are always positive. However, the remaining coefficients of $E_P(t)$ are not always positive. We say an integral polytope P is *Ehrhart positive*, if all the coefficients of $E_P(t)$ are positive. (See [17] for a survey on the problem of Ehrhart positivity.)

An affine transformation from an affine space $A \subseteq \mathbb{R}^n$ to an affine space $B \subseteq \mathbb{R}^m$ is a *unimodular transformation* if it induces a bijection from integer points in A to integer points in B . Two polytopes $P \subset \mathbb{R}^n$ and $Q \subset \mathbb{R}^m$ are said to be *unimodularly equivalent* if there exists a unimodular transformation ϕ from the affine hull of P to the affine hull of Q such that $\phi(P) = Q$. Clearly, if two polytopes are unimodularly equivalent, they have the same Ehrhart polynomial.

Date: March 13 , 2019.

1.1. Tesler polytopes and Morales’s conjecture. Our main object of study in this paper is the family of the Tesler polytopes recently introduced and studied by Mészáros, Morales and Rhoads [19]. Let $\mathbb{U}(n)$ be the set of $n \times n$ upper triangular matrices. If we ignore the zeros below the diagonal, then $\mathbb{U}(n) \cong \mathbb{R}^{\binom{n+1}{2}}$ as a vector space. For any $M = (m_{i,j}) \in \mathbb{U}(n)$, the *hook-sum vector* of M is defined to be

$$hs(M) := (hs_1(M), hs_2(M), \dots, hs_n(M)),$$

where the k -th entry $hs_k(M)$ (for each $1 \leq k \leq n$) is called the *k -th hook-sum* of M and is computed as the sum of the entries on the k -th row minus the sum of the entries on the k -th column above the diagonal element, that is,

$$hs_k(M) := (m_{k,k} + m_{k,k+1} + \dots + m_{k,n}) - (m_{1,k} + m_{2,k} + \dots + m_{k-1,k}).$$

It is easy to see that hs_k and hs are linear functions on $\mathbb{U}(n)$. For any $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$, the *Tesler polytope of hook sum \mathbf{a}* , denoted by $\text{Tes}_n(\mathbf{a})$, is the set of all matrices $M \in \mathbb{U}(n)$ with non-negative entries satisfying the hook sum condition $hs(M) = \mathbf{a}$:

$$\text{Tes}_n(\mathbf{a}) = \{M = (m_{i,j}) \in \mathbb{U}(n) \mid hs(M) = \mathbf{a}, m_{i,j} \geq 0\}.$$

When $\mathbf{a} = \mathbf{1} := (1, \dots, 1)$, the integer points of $\text{Tes}_n(\mathbf{a})$ are called Tesler matrices, which were initially introduced by Tesler, and then rediscovered by Haglund in his work of expressing the diagonal Hilbert series as a weighted sum over these matrices [15]. Consequently, Tesler matrices play an important role in the field of diagonal harmonics [1, 13, 14, 16, 25]. Intrigued by these work, Mészáros, Morales and Rhoads [19] defined and studied the Tesler polytopes of hook sum \mathbf{a} as defined above. We remark that their definition for Tesler polytopes only allows to choose \mathbf{a} from $\mathbb{Z}_{>0}^n$. However, since important features remain the same, we extend the domain of \mathbf{a} to $\mathbb{R}_{\geq 0}^n$.

The work in the paper was motivated by a conjecture of Morales’:

Conjecture 1. ([20]) Tesler polytopes $\text{Tes}_n(\mathbf{1})$ and $\text{Tes}_n(1, 0, \dots, 0)$ are both Ehrhart positive for any positive integer n .

As we mentioned above, the integer points of $\text{Tes}_n(\mathbf{1})$ (known as the Tesler matrices) are important objects in the study of diagonal harmonics. Moreover, it was known [9] that $\text{Tes}_n(1, 0, \dots, 0)$ is unimodularly equivalent to the Chan-Robbins-Yuen (CRY) polytope, which is a face of the Birkhoff polytope and whose volume is the product of the first $n - 2$ Catalan numbers. (The volume formula was conjectured by Chan, Robbins and Yuen was proved by Zeilberger [26]). Note that the Birkhoff polytope is also a well-studied subject, and in particular computing volumes of Birkhoff polytopes is an extremely hard problem and has attracted a lot of recent research [5, 11, 21]. Therefore, both $\text{Tes}(\mathbf{1})$ and $\text{Tes}_n(1, 0, \dots, 0)$ are fascinating families of polytopes for their interesting combinatorial properties and connection to other fields of mathematics.

1.2. McMullen’s formula and α -positivity. In this paper, we will use a technique developed by Castillo and the 2nd author [6] to attack Morales’ positivity conjecture and its generalizations. The technique is based on the existence of “McMullen’s formula”. In 1975, Danilov questioned, in the context of toric varieties, whether it is possible to construct a

function α such that for any integral polytope $P \subset \mathbb{R}^n$, the following equation holds

$$|P \cap \mathbb{Z}^n| = \sum_{F : \text{face of } P} \alpha(F, P) \text{nVol}(F),$$

where $\text{nVol}(F)$ is the normalized volume of F and $\alpha(F, P)$ only depends on the normal cone of P at F [10]. McMullen was the first to confirm that it is possible to construct such a function α (in a non-constructive way). Hence, we refer to the above formula as *McMullen's formula* [18]. Since α only depends on normal cones and the normal cones are invariant under dilations, we obtain that

$$(1.1) \quad e_i(P) = \sum_{F : i\text{-dimensional face of } P} \alpha(F, P) \text{nVol}(F), \text{ for any } 1 \leq i \leq \dim(P).$$

One sees that, as a consequence of the above formula, if $\alpha(F, P) > 0$ for every i -dimensional face of P , then $e_i(P)$ is positive by the above formula. Moreover, if $\alpha(F, P) > 0$ for every face of P , then P is Ehrhart positive. We say a polytope P is α -positive if $\alpha(F, P) > 0$ for every face F of P . Currently, we know three different constructions for α given by Pommersheim and Thomas [22], by Berline and Vergne [4], by Ring and Schürmann [24]. Hence, the choice of the function α for McMullen's formula is not unique, and thus a specific construction of α has to be chosen before one discusses α -positivity. Among known constructions of the function α , we will use Berline-Vergne's construction which we will refer to as the BV - α function denoted by α^{BV} . Based on discussion above, one sees that Ehrhart positivity can be studied through α -positivity or more specifically through BV - α -positivity, which is the approach we will take. To simplify the calculation of BV - α values, we work on the projected Tesler polytopes denoted by $\text{PTes}_n(\mathbf{a})$ which is unimodularly equivalent to $\text{Tes}_n(\mathbf{a})$, and thus has the same Ehrhart polynomial (see §3.1 for the definition of $\text{PTes}_n(\mathbf{a})$). Therefore, if $\text{PTes}_n(\mathbf{a})$ is BV - α positive then $\text{Tes}_n(\mathbf{a})$ is Ehrhart positive. We have the following theorem:

Theorem 1.1. *For all positive integer n , all the codimension 2 and 3 faces of $\text{PTes}_n(\mathbf{1})$ have positive BV - α values. Therefore, $e_{d-2}(\text{Tes}_n(\mathbf{1}))$ and $e_{d-3}(\text{Tes}_n(\mathbf{1}))$ are positive, where $d = \binom{n}{2}$ is the dimension of $\text{PTes}_n(\mathbf{1})$.*

1.3. Deformations. We say a polytope Q is a *deformation* of a polytope P if the normal fan of P is a refinement of the normal fan of Q . It follows from the Reduction Theorem (Theorem 2.14) by Castillo and the second author that if all the k -dimensional faces of polytope P are BV - α positive, then the same is true for any deformations of P . Therefore, we can generalize our BV - α positivity results for codimension 2 and 3 faces of $\text{PTes}_n(\mathbf{1})$ to any deformations of $\text{PTes}_n(\mathbf{1})$:

Corollary 1.2. *Let P be a d -dimensional integral polytope that is a deformation of $\text{PTes}_n(\mathbf{1})$ and let $E_P(t) = e_d(P)t^d + e_{d-1}t^{d-1} + \dots + 1$ be its Ehrhart polynomial. Then, the followings are true:*

- (1) *If $d = \dim(\text{Tes}_n(\mathbf{1}))$, then all the codimension 2 and 3 faces of P have positive BV - α values. Therefore, $e_{d-2}(P) > 0$ and $e_{d-3}(P) > 0$.*
- (2) *If $d = \dim(\text{Tes}_n(\mathbf{1})) - 1$, then all the codimension 2 faces of P have positive BV - α values. Therefore, $e_{d-2}(P) > 0$.*

Since any invertible affine transformation preserves deformation (Lemma 3.12), we directly obtain:

Corollary 1.3. *Let P be any polytope that is a deformation of $\text{Tes}_n(\mathbf{1})$*

- (1) *If $\dim(P) = \dim(\text{Tes}_n(\mathbf{1}))$, then $e_{d-2}(P) > 0$ and $e_{d-3}(P) > 0$.*
- (2) *If $\dim(P) = \dim(\text{Tes}_n(\mathbf{1})) - 1$, then $e_{d-2}(P) > 0$.*

It is known that $\text{Tes}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{1})$ for all $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$. Therefore, the positivity result for $\text{Tes}_n(\mathbf{1})$ generalizes to $\text{Tes}_n(\mathbf{a})$ for all $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$ including $\text{Tes}_n(1, 0, \dots, 0)$, which is the other polytope in Conjecture 1.

Given the results in Corollary 1.3, it is natural to ask which polytopes are deformations of $\text{Tes}_n(\mathbf{1})$. In [19], Mészáros et al showed that for any $\mathbf{a} \in \mathbb{Z}_{>0}^n$, the Tesler polytope $\text{Tes}_n(\mathbf{a})$ is unimodularly equivalent to a certain flow polytope (See §4.2 for more details). We remark that flow polytopes have connection to several areas of mathematics such as toric geometry, representation theory, special functions and algebraic combinatorics (see the introduction of [3] and references therein). We gave the following characterization of which flow polytopes on a complete graph are deformations of $\text{Tes}_n(\mathbf{1})$:

Theorem 1.4. *For any fixed n , let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$ such that $\text{Flow}_n(\mathbf{a})$ is non-empty. Suppose a_l is the first positive entry. Then, $\text{Flow}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{1})$ if and only if $a_{l+2}, a_{l+3}, \dots, a_n \geq 0$.*

ORGANIZATION OF THE PAPER

In section 2, we provide background on polyhedra theory and Tesler polytopes, and a brief description for the construction of BV- α functions. In section 3, we define projected Tesler polytopes and develop a simple method (see section 3.3) that helps to calculate the BV- α values arising from totally unimodular polytopes. Applying this method to $\text{PTes}_n(\mathbf{1})$, we obtain a proof for Theorem 1.1 by explicitly calculating the α^{BV} -values (Lemmas 3.9 and 3.10). Then, we prove Corollaries 1.2 and 1.3 using the Reduction Theorem (Theorem 2.14). In section 4, we first give a different proof of the known fact that $\text{Tes}_n(\mathbf{a})$ are deformations of $\text{Tes}_n(\mathbf{1})$, and then show that the family of Tesler polytopes already contains all of its deformations up to translations. We finish with a proof for Theorem 1.4.

2. PRELIMINARIES

In this section, we review terminologies related to polytopes/polyhedra that are important to this article.

2.1. Basic definitions in polyhedra theory. In this section, we assume that V is a subspace of \mathbb{R}^n , and $\langle \cdot, \cdot \rangle$ is the dot product on \mathbb{R}^n . Let P be a polytope in $V \subset \mathbb{R}^n$. We denote by $\text{Vert}(P)$ the set of all vertices of P . We say P is a *simple* polytope, if each of the vertices of P are contained in exactly $\dim(P)$ many facets or equivalently contained in $\dim(P)$ many edges. If a facet F of P contains a face G , we call F a *supporting facet* of G . Two vertices v and w of P are said to be *adjacent* if they are connected by an edge. For an edge e of P connecting two vertices \mathbf{v} and \mathbf{w} , we define the *edge direction* from \mathbf{v} to \mathbf{w} to be $\mathbf{d}_{\mathbf{v},\mathbf{w}} := \mathbf{w} - \mathbf{v}$. When P is an integral polytope, we define the *primitive edge direction* from \mathbf{v} to \mathbf{w} to be $\mathbf{p}_{\mathbf{v},\mathbf{w}} := a(\mathbf{w} - \mathbf{v})$ where $a \in \mathbb{Q}_{>0}$ is chosen so that \mathbf{p} is a primitive vector. (Recall that an integer vector \mathbf{m} is *primitive* if the greatest common divisor of its components is 1.) A polyhedron defined by homogeneous linear inequalities is called a (*polyhedral*) *cone*. An equivalent definition of a cone is the set of all conic combinations of finitely many points in V . A *pointed cone* is a cone that does not contain a line.

Definition 2.1. Let K be a cone in \mathbb{R}^n that is generated by the primitive vectors $\mathbf{d}_1, \dots, \mathbf{d}_l$. We say K is *unimodular* if $\{\mathbf{d}_1, \dots, \mathbf{d}_l\}$ forms a basis of the lattice $\mathbb{Z}^n \cap \text{span}(\mathbf{d}_1, \dots, \mathbf{d}_l)$.

Definition 2.2. We say a polytope P in \mathbb{R}^n is *totally unimodular* if $\text{fcone}(P, \mathbf{v})$ is unimodular for all the vertices \mathbf{v} of P .

For any subset S of a vector space V , let $\text{lin}(S)$ denote the translation of the affine span of S to the origin, and let $V/\text{lin}(S)$ denote the orthogonal complement of $\text{lin}(S)$. For any subset or an element B of V , we use $B/\text{lin}(S)$ to denote the canonical projection of B into $V/\text{lin}(S)$.

Definition 2.3. Suppose $P \subset V$ is a non-empty polyhedron, and F is a face of P .

- (1) The *feasible cone of P at F* is:

$$\text{fcone}(F, P) := \{\mathbf{y} \in V : \mathbf{x} + \epsilon\mathbf{y} \in P \text{ for some } \epsilon > 0\},$$

where \mathbf{x} is any interior point of F . (It can be shown that the definition does not depend on the choice of \mathbf{x} .) The *pointed feasible cone* of P at F is

$$\text{fcone}^p(F, P) := \text{fcone}(F, P)/\text{lin}(F).$$

- (2) Given any face F of P , the *normal cone* of P at F with respect to V is:

$$\text{ncone}_V(F, P) := \{\mathbf{u} \in V : \langle \mathbf{u}, \mathbf{p}_1 \rangle \geq \langle \mathbf{u}, \mathbf{p}_2 \rangle, \forall \mathbf{p}_1 \in F, \forall \mathbf{p}_2 \in P\}.$$

The *normal fan* $\Sigma_V P$ of P with respect to V is the collection of all normal cones of P . (We omit the subscript V and just write $\text{ncone}(F, P)$ and ΣP , if $V = \mathbb{R}^n$ and P is full dimensional.)

- (3) Let $K \subset V$ be a cone, and let W be the subspace of V spanned by K . The *polar cone* of K is the cone

$$K^\circ = \{\mathbf{y} \in W : \langle \mathbf{x}, \mathbf{y} \rangle \leq 0, \forall \mathbf{x} \in K\}.$$

Lemma 2.4. [27] *The normal fans and the normal cones have the following properties:*

- (1) For any $C \in \Sigma_V P$, any face of C is also in $\Sigma_V P$.
- (2) There is an inclusion reversing bijection between the face lattice of P (the set of all the faces of P ordered by inclusion) and $\Sigma_V P$ given by $F \mapsto \text{ncone}_V(F, P)$.
- (3) If P is a full dimensional polytope, $\dim(\text{ncone}(F, P)) = \dim(V) - \dim(F)$ and $\text{ncone}(F, P)$ is pointed for any face F of P .
- (4) $\text{ncone}_V(F, P)^\circ = \text{fcone}^p(F, P)$.

2.2. Prior work on Tesler polytopes by Mészáros, Morales and Rhoads. In this part, we review definitions and results related to Tesler polytope given in [19] that are relevant to this paper.

For any positive integer n and $\mathbf{a} \in \mathbb{Z}_{\geq 0}^n$, Mészáros, Morales and Rhoads [19] gave the characterization for the face poset of $\text{Tes}_n(\mathbf{a})$ using the concept of support. Their characterization can be easily generalized to any $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$ by the same proof.

Definition 2.5. For any set \mathcal{A} of matrices in $\mathbb{U}(n)$, define the *support* of \mathcal{A} , denoted by $\text{supp}(\mathcal{A})$, to be the matrix $(s_{i,j}) \in \mathbb{U}(n)$,

$$\text{where } s_{i,j} = \begin{cases} 0, & \text{if the } (i,j)\text{-th entry of every matrix in } \mathcal{A} \text{ is zero,} \\ 1, & \text{otherwise.} \end{cases}$$

For $\mathcal{A} = \{\mathbf{m}\}$, containing one matrix \mathbf{m} , write $\text{supp}(\mathbf{m})$ instead of $\text{supp}(\mathcal{A})$ whenever it is convenient to do so. Let \mathcal{A} and \mathcal{B} be two subsets of $\mathbb{U}(n)$. We write $\text{supp}(\mathcal{A}) \leq \text{supp}(\mathcal{B})$, if $\text{supp}(\mathcal{A}) = (a_{i,j})$ and $\text{supp}(\mathcal{B}) = (b_{i,j})$ satisfying $a_{i,j} \leq b_{i,j}$ for any $1 \leq i \leq j \leq n$.

For $1 \leq i \leq j \leq n$, let

$$(2.1) \quad H_{i,j}^n := \{M = (m_{l,k}) \in \mathbb{U}(n) \mid m_{i,k} = 0\}.$$

We say the intersection of a collection of coordinate planes *does not make any zero rows* if the intersection is not contained in $H_{i,i}^n \cap H_{i,i+1}^n \cap \cdots \cap H_{i,n}^n$ for any $1 \leq i \leq n$.

Theorem 2.6 (Lemma 2.4 and Theorem 2.5 in [19]). *For any positive integer n , the followings are true:*

- (1) Let $\mathbf{a} \in \mathbb{R}_{>0}^n$. If F is a codimension k face of $\text{Tes}_n(\mathbf{a})$, then F is of the form

$$\text{Tes}_n(\mathbf{a}) \cap H_{i_1, j_1}^n \cap \cdots \cap H_{i_k, j_k}^n,$$

where $H_{i_1, j_1}^n \cap \cdots \cap H_{i_k, j_k}^n$ does not make any zero rows. Conversely, any such form is a codimension k face of $\text{Tes}_n(\mathbf{a})$.

- (2) Let $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$. $\mathbf{v} \in \text{Tes}_n(\mathbf{a})$ is a vertex if and only if $\text{supp}(\mathbf{v})$ has at most one 1 on each row. In particular, when $\mathbf{a} \in \mathbb{R}_{>0}^n$, $\mathbf{v} \in \text{Tes}_n(\mathbf{a})$ is a vertex if and only if each row of \mathbf{v} has exactly one 1.

Using the above theorem, one can deduce the following when $\mathbf{a} \in \mathbb{R}_{>0}^n$. In fact, it is shown to be true for any $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$.

Lemma 2.7 (Theorem 2.5 in [19]). *Let $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$, and let F, F' be two faces of $\text{Tes}_n(\mathbf{a})$. Then the followings are true.*

- (1) $F = F'$ if and only if $\text{supp}(F) = \text{supp}(F')$.
(2) $F \subseteq F'$ if and only if $\text{supp}(F) \leq \text{supp}(F')$.

We also need the following characterization for adjacent vertices of $\text{Tes}_n(\mathbf{a})$ and a specific type of vertex presented in the next two lemmas.

Lemma 2.8 (Theorem 2.7 in [19]). *Let $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$. Two vertices \mathbf{v} and \mathbf{w} of $\text{Tes}_n(\mathbf{a})$ are adjacent if and only if for every $1 \leq k \leq n$, the k -th row of $\text{supp}(\mathbf{w})$ can be obtained from the k -th of $\text{supp}(\mathbf{v})$ by one of the following operations:*

- (1) Leaving the k -th row of $\text{supp}(\mathbf{v})$ unchanged.
(2) Changing the unique 1 in k -th row of $\text{supp}(\mathbf{v})$ to 0.
(3) Changing a single 0 in the k -th row to a 1 (if k -th row of $\text{supp}(\mathbf{v})$ is a zero row).
(4) Moving the unique 1 in the k -th row of $\text{supp}(\mathbf{v})$ to a different position in the k -th row (this operation must take place in precisely one row of $\text{supp}(\mathbf{v})$).

In particular, when $\mathbf{a} \in \mathbb{R}_{>0}^n$, two vertices \mathbf{v}, \mathbf{w} are adjacent to each other if and only if $\text{supp}(\mathbf{w})$ can be obtained from $\text{supp}(\mathbf{v})$ by moving 1 on a row to a different place on the same row.

Lemma 2.9 (Lemma 2.4 of [19]). *Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$. The k -th row of a vertex \mathbf{v} of $\text{Tes}_n(\mathbf{a})$ is a zero row if and only if $a_k = 0$ and the entries of k -th column of \mathbf{v} above the diagonal (excluding the diagonal entry) are all zero.*

Example 2.10. Let $\mathbf{a} = (2, 2, 3, 4)$, and

$$\mathbf{v} = \begin{pmatrix} 0 & 2 & 0 & 0 \\ & 0 & 0 & 4 \\ & & 3 & 0 \\ & & & 8 \end{pmatrix} \text{ and } \mathbf{w} = \begin{pmatrix} 0 & 0 & 2 & 0 \\ & 0 & 0 & 2 \\ & & 5 & 0 \\ & & & 6 \end{pmatrix}$$

Then

$$\text{supp}(\mathbf{v}) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ & 0 & 0 & 1 \\ & & 1 & 0 \\ & & & 1 \end{pmatrix} \text{ and } \text{supp}(\mathbf{w}) = \begin{pmatrix} 0 & 0 & 1 & 0 \\ & 0 & 0 & 1 \\ & & 1 & 0 \\ & & & 1 \end{pmatrix}$$

Using Theorem 2.6 (1), one can easily check that \mathbf{v} and \mathbf{w} are vertices of $\text{Tes}_4(2, 2, 3, 4)$. Also, they are adjacent vertices, because $\text{supp}(\mathbf{w})$ can be obtained by moving 1 on the second row of $\text{supp}(\mathbf{v})$ to the right.

2.3. Berline-Vergne's construction. In this subsection, we provide more details about BV- α function mentioned in the introduction. In [4], Berline and Vergne associated to every rational affine cone c an analytic function $\phi(c)$ on \mathbb{R}^n which is recursively defined with respect to the dimension of cones. They showed that ϕ is a valuation and $\phi(c)$ is analytic near the origin. Then they set $\alpha^{\text{BV}}(F, P)$ to be the residue of $\phi(\text{fcone}^p(F, P))$ around 0.

One sees that this procedure is complicated. However, when $\text{fcone}^p(F, P)$ is unimodular, we can obtain a simple formula for $\alpha^{\text{BV}}(F, P)$ when $\dim(\text{fcone}^p(F, P))$ is up to 3. Note that it follows immediately from Berline-Vergne's construction that $\alpha^{\text{BV}}(F, P) = 1$ when $\dim(\text{fcone}^p(F, P)) = 0$, and $\alpha^{\text{BV}}(F, P) = \frac{1}{2}$ when $\dim(\text{fcone}^p(F, P)) = 1$ ([2, Example 19.2]). We include formulas for faces of codimensions 2 and 3 below.

Lemma 2.11 (Example 19.3 of [2]). *Let F be a codimension 2 face of a full dimensional polytope P and $C = \text{fcone}^p(F, P) = \text{cone}(u_1, u_2)$ where u_1 and u_2 form a basis for the orthogonal projection of \mathbb{Z}^n to $\mathbb{R}^n / \text{lin}(F)$. Then,*

$$\alpha^{\text{BV}}(F, P) = \frac{1}{4} + \frac{1}{12} \left(\frac{\langle u_1, u_2 \rangle}{\langle u_1, u_1 \rangle} + \frac{\langle u_1, u_2 \rangle}{\langle u_2, u_2 \rangle} \right).$$

Lemma 2.12 (Lemma 3.10 of [6]). *Let F be a codimension 3 face of a full dimensional polytope P and $C = \text{fcone}^p(F, P) = \text{cone}(u_1, u_2, u_3)$. Where u_1 and u_2 form a basis for the orthogonal projection of \mathbb{Z}^n to $\mathbb{R}^n / \text{lin}(F)$. Then,*

$$\alpha^{\text{BV}}(F, P) = \frac{1}{8} + \frac{1}{24} \left(\frac{\langle u_1, u_2 \rangle}{\langle u_1, u_1 \rangle} + \frac{\langle u_1, u_2 \rangle}{\langle u_2, u_2 \rangle} + \frac{\langle u_1, u_3 \rangle}{\langle u_1, u_1 \rangle} + \frac{\langle u_1, u_3 \rangle}{\langle u_3, u_3 \rangle} + \frac{\langle u_2, u_3 \rangle}{\langle u_2, u_2 \rangle} + \frac{\langle u_2, u_3 \rangle}{\langle u_3, u_3 \rangle} \right).$$

Notice that the formulas are in terms of the dot products between the primitive generators of $\text{fcone}^p(F, P)$.

Remark 2.13. If $\dim(\text{fcone}^p(F, P)) \geq 4$, the formula for $\alpha^{\text{BV}}(F, P)$ becomes way more complicated. However, they are still in terms of $\langle \mathbf{u}_i, \mathbf{u}_j \rangle$. Therefore, it is important to know the values of $\langle \mathbf{u}_i, \mathbf{u}_j \rangle$. In section 3.3, we develop a simple procedure of calculating the dot product values without having to calculate each \mathbf{u}_i 's (calculation of \mathbf{u}_i is not easy because as we project $\text{fcone}(F, P)$ to a subspace, the lattice structure changes).

It is also possible to obtain the formula even when $\text{fcone}^p(F, P)$ is not unimodular by decomposing it in to unimodular cones and using the valuation property of Ψ . One great

benefit of using BV- α is the Reduction Theorem which follows from the valuation property of Ψ :

Theorem 2.14 (Reduction Theorem). *Suppose P and Q are two polytopes in \mathbb{R}^n . Assume further that Q is a deformation of P . Then for any fixed k , if $\alpha^{\text{BV}}(F, P) > 0$ for every k -dimensional face F of P , then $\alpha^{\text{BV}}(G, Q) > 0$ for every k -dimensional face G of Q . Therefore, BV- α -positivity of P implies BV- α -positivity of Q .*

By the above theorem, if we show that all the k -th dimensional faces of a polytope P is BV- α positive (which together with (1.1) implies that the corresponding Ehrhar coefficient is positive), then the same is true for any deformation Q of P .

3. POSITIVITY OF THE (PROJECTED) TESLER POLYTOPE OF HOOK SUMS $(1, \dots, 1)$

We prefer working with full dimensional polytopes because facet normal vectors are uniquely determined. One can easily see that Tesler polytopes are not full dimensional in the ambient space, but it is full dimensional in the subspace defined by the hook sum conditions. In this section, we will consider suitable projections that induces unimodular transformations from the subspace defined by the hook sum conditions to $\mathbb{U}(n)$. We call the image of the Tesler polytopes $\text{Tes}_n(\mathbf{a})$ under the projection the *projected Tesler polytopes* denoted by $\text{PTes}_n(\mathbf{a})$. In order to compute the BV- α values for the faces of $\text{PTes}_n(\mathbf{a})$ or any totally unimodular polytope, we need to calculate the dot products between the generators of $\text{fcone}^p(F, P)$ (Lemmas 2.11 and 2.12). We develop a method for obtaining these dot product values without calculating the generators of $\text{fcone}^p(F, P)$. We then apply the method to $\text{PTes}_n(\mathbf{1})$ and show that the BV- α values of all the codimension 2 and codimension 3 faces of $\text{PTes}_n(\mathbf{1})$ are positive by calculating the values. As a consequence, we complete the proof for Theorem 1.1, Corollaries 1.2 and 1.3.

For any $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}$, let \mathbf{a} -hook sum space be the subspace of $\mathbb{U}(n)$ defined by hook sum conditions:

$$H_n(\mathbf{a}) := \{M \in \mathbb{U}(n) \mid \text{hs}_k(M) = a_k \text{ for } 1 \leq k \leq n\}.$$

3.1. Projection on the Tesler polytopes. Recall $[n] = \{1, 2, \dots, n\}$ and $[k, n] = \{k, k+1, \dots, n\}$.

Theorem 3.1. *For any $\mathbf{a} \in \mathbb{Z}_{\geq 0}^n$ and $\mathbf{j} = (j_1, j_2, \dots, j_{n-1}, j_n) \in [n] \times [n-1] \times [n-1] \times \dots \times [1]$, define $\psi_{\mathbf{j}} : \mathbb{U}(n) \rightarrow \mathbb{U}(n-1)$ as*

$$\psi_{\mathbf{j}}((x_{i,j})) = (y_{i,j}) \text{ where } y_{i,j} = \begin{cases} x_{i,j} & \text{if } j < j_i, \\ x_{i,j+1} & \text{if } j \geq j_i, \end{cases}$$

($\psi_{\mathbf{j}}$ deletes an entry in a row for each rows where \mathbf{j} determines which entries to be deleted). Then $\psi|_{H_n(\mathbf{a})}$ is a unimodular transformation from $H_n(\mathbf{a})$ to $\mathbb{U}(n-1)$.

Proof. One can easily see that $\psi_{\mathbf{j}}$ is a projection. Thus, it is enough to show that $\psi|_{H_n(\mathbf{a})}$ induces a bijection from the integer points of $H_n(\mathbf{a})$ to that of $\mathbb{U}(n-1)$. Let $\mathbf{z} = (z_{i,j}) \in \mathbb{U}(n-1)$. If $\mathbf{z}' = (z'_{i,j}) \in \mathbb{U}(n)$ such that $\psi_{\mathbf{j}}(\mathbf{z}') = \mathbf{z}$, then

$$z'_{i,j} = \begin{cases} z_{i,j} & \text{if } j < j_i \\ z_{i,j-1} & \text{if } j > j_i \text{ for some } x_{i,j} \in \mathbb{R}. \\ x_{i,j} & \text{if } j = j_i \end{cases}$$

Since $\mathbf{z}' \in H_n(\mathbf{a})$, one sees that \mathbf{z}' has to satisfy the hook sum conditions $hs_i(\mathbf{z}') = a_i$ for any $1 \leq i \leq n$. The first hook sum condition uniquely determines x_{1,j_1} in terms of $z_{i,j}$'s and a_1 . Since the entries of the first row of \mathbf{z}' are now all in terms of $z_{i,j}$'s and a_1 , the second hook sum condition uniquely determines x_{2,j_2} in terms of $z_{i,j}$'s, a_1 and a_2 . As we continue this process inductively (row by row), all the x_{i,j_i} 's are expressed uniquely in terms of $z_{i,j}$'s and a_i 's. The injectivity and the surjectivity follows from the above construction of \mathbf{z}' . Additionally, one sees that if \mathbf{z} was chosen to be an integer point, \mathbf{z}' is also an integer point by the construction of \mathbf{z}' . Thus, ψ is a bijection from $H_n(\mathbf{a})$ to $\mathbb{U}(n-1)$ that induces a bijection from integer points of $H_n(\mathbf{a})$ to integer points of $\mathbb{U}(n-1)$. \square

Example 3.2. The most natural way to obtain such map would be letting $j = (1, 2, \dots, n)$.

$$\begin{bmatrix} d_1 & x_{1,1} & x_{1,2} & \cdots & x_{1,n-1} \\ & d_2 & x_{2,2} & \cdots & x_{2,n-1} \\ & & \ddots & \cdots & \vdots \\ & & & d_{n-1} & x_{n-1,n-1} \\ & & & & d_n \end{bmatrix} \longrightarrow \begin{bmatrix} x_{1,1} & x_{1,2} & \cdots & x_{1,n-1} \\ & x_{2,2} & \cdots & x_{2,n-1} \\ & & \ddots & \vdots \\ & & & x_{n-1,n-1} \end{bmatrix}$$

Let ψ_{diag} be the map in the above example. We call $\psi_{\text{diag}}(\text{Tes}_n(\mathbf{a}))$ the *projected Tesler polytope of hook sum \mathbf{a}* and denote it by $\text{PTes}_n(\mathbf{a})$.

3.2. Faces and the facet normal vectors of the projected Tesler polytopes. To calculate the BV- α values of the faces of $\text{PTes}_n(\mathbf{1})$, we first find the facet normal vectors of $\text{PTes}_n(\mathbf{1})$.

Definition 3.3. For any n ,

- (1) Let $e_{i,j}^n$ be the matrix in $U(n)$ that the entries are all zero except (i, j) -th entry is 1.
- (2) For $1 \leq k \leq n$, define the *shifted k -th hook sum matrix* to be

$$sm_k^n := \sum_{j=k}^n e_{k,j}^n - \sum_{i=1}^{k-1} e_{i,k-1}^n \in \mathbb{U}(n).$$

- (3) Let $M = (m_{i,j}) \in \mathbb{U}(n)$ For $1 \leq k \leq n$ define *shifted k -th hook sum* of M to be

$$sh_k^n(M) := (m_{k,k} + m_{k,k+1} \cdots + m_{k,n}) - (m_{1,k-1} + m_{2,k-1} + \cdots + m_{k-1,k-1}).$$

We fix a notation for the facets of $\text{Tes}_n(\mathbf{a})$ and $\text{PTes}_n(\mathbf{a})$. From now on, for any $\mathbf{a} \in \mathbb{Z}_{>0}^n$, $1 \leq i < j \leq n$, we let

$$(3.1) \quad F'_{i,j}(\mathbf{a}) := \text{Tes}_n(\mathbf{a}) \cap H_{i,j}^n \text{ and } F_{i,j}(\mathbf{a}) := \phi_d(\text{Tes}_n(\mathbf{a}) \cap H_{i,j}^n).$$

where $H_{i,j}^n$ is the coordinate plane defined in (2.1) (by Theorem 2.6 (1), any facet of $\text{Tes}_n(\mathbf{a})$ can be expressed as $\text{Tes}_n(\mathbf{a}) \cap H_{i,j}^n$ and since ϕ is a unimodular transformation, it bijects the facets of $\text{Tes}_n(\mathbf{a})$ to that of $\text{PTes}_n(\mathbf{a})$). We would like to give an example of obtaining the half-space description of $\text{PTes}_n(\mathbf{a})$.

Example 3.4. Let $\mathbf{a} = (4, 5, 2, 2)$. Then

$$\text{Tes}_4(\mathbf{a}) = \left\{ \mathbf{x} = (x_{i,j}) \in \mathbb{U}(4) \mid \begin{array}{l} hs_1(\mathbf{x}) = 4, \quad hs_2(\mathbf{x}) = 5, \quad hs_3(\mathbf{x}) = 2, \quad hs_4(\mathbf{x}) = 2, \\ x_{i,j} \geq 0 \text{ for all } 1 \leq i \leq j \leq 4 \end{array} \right\}.$$

We use the following notation to describe the image of ψ_{diag} as in Theorem 3.1:

$$\psi_{\text{diag}}(\mathbf{x}) = \mathbf{y} \in \mathbb{U}(3) \text{ where } y_{i,j} = x_{i,j+1}.$$

After going through ψ_{diag} , the **diagonal components** gets erased. Since all the components are non-negative, the hook-sum conditions become inequalities.

$$\begin{aligned} hs_1(\mathbf{x}) = \mathbf{x}_{1,1} + x_{1,2} + x_{1,3} + x_{1,4} = 4 &\implies sh_1(\mathbf{y}) = y_{1,1} + y_{1,2} + y_{1,3} \leq 4. \\ hs_2(\mathbf{x}) = \mathbf{x}_{2,2} + x_{2,3} + x_{2,4} - x_{1,2} = 5 &\implies sh_2(\mathbf{y}) = y_{2,2} + y_{2,3} - y_{1,1} \leq 5. \\ hs_3(\mathbf{x}) = \mathbf{x}_{3,3} + x_{3,4} - x_{1,3} - x_{2,3} = 2 &\implies sh_3(\mathbf{y}) = y_{3,3} - y_{1,2} - y_{2,2} \leq 2. \\ hs_4(\mathbf{x}) = \mathbf{x}_{4,4} - x_{1,4} - x_{2,4} - x_{3,4} = 2 &\implies sh_4(\mathbf{y}) = -y_{1,3} - y_{2,3} - y_{3,3} \leq 2. \end{aligned}$$

The last inequality above is redundant, because $-y_{1,3} - y_{2,3} - y_{3,3} \leq 0$. As a result, we get the description for $\text{PTes}_4(\mathbf{a})$:

$$\text{PTes}_4(\mathbf{a}) = \left\{ \mathbf{y} = (y_{i,j}) \in \mathbb{U}(3) \mid \begin{array}{l} sh_1(\mathbf{y}) \leq 4, \ sh_2(\mathbf{y}) \leq 5, \ sh_3(\mathbf{y}) \leq 2, \\ y_{i,j} \geq 0 \text{ for all } 1 \leq i \leq j \leq 3 \end{array} \right\}.$$

One can easily check that all the defining inequalities above are necessary. Thus, the outer normal vectors for the facets are sm_1^4 , sm_2^4 , sm_3^4 and $-\mathbf{e}_{i,j}$ for all $1 \leq i \leq j \leq 3$.

It is easy to see that the above method of obtaining the following description of $\text{PTes}_n(\mathbf{a})$ works for any $\mathbf{a} \in \mathbb{Z}_{>0}$. So, we state the following lemma without the proof.

Lemma 3.5. *For any $n \in \mathbb{Z}_{>0}$ and $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}_{>0}^n$. Let $\mathbf{n}_{i,j} = \begin{cases} -\mathbf{e}_{i,j-1}^{n-1} & \text{if } i < j, \\ sm_k^{n-1} & \text{if } i = j = k. \end{cases}$*

Then, $\mathbf{n}_{i,j}$ is an outer facet normal vector of $F_{i,j}$.

3.3. Computing BV- α values of totally unimodular polytopes. If an integral polytope P in \mathbb{R}^n is totally unimodular (Definition 2.2), then for any face F of P , the primitive generators of $\text{fcone}^P(F, P)$ forms a basis for $\mathbb{Z}^n / \text{lin}(F)$. Thus, we can apply any known BV- α formula to calculate the BV- α value of the faces of P . In this section we will apply Lemma 2.11 and Lemma 2.12 to calculate the BV- α values of all the codimension 2 and 3 faces of $\text{PTes}_n(\mathbf{1})$. We will present a simple way to obtain the inner product values $\langle \mathbf{u}_i, \mathbf{u}_j \rangle$ in Lemma 2.11 and Lemma 2.12 which is important for computing BV- α values (recall Remark 2.13). We start with the following lemma:

Lemma 3.6. *Let $\{\mathbf{n}_1, \dots, \mathbf{n}_{k+1}\}$ be the set of linearly independent primitive vectors in \mathbb{R}^n where the matrix A formed by taking $\{\mathbf{n}_1, \dots, \mathbf{n}_{k+1}\}$ as rows is totally unimodular. Then, for any $\mathbf{c} = (c_1, \dots, c_n) \in \mathbb{Z}^n$ there exists an integer solution for the equation $AX = \mathbf{c}^t$ where $X = (x_1, \dots, x_n)$.*

Lemma 3.7. *Let P be an integral polytope in \mathbb{R}^n that is totally unimodular. Let S be a subspace of \mathbb{R}^n that is the translation of the affine subspace that P is full dimensional in. Let F be a codimension k face of P that is the intersection of the facets F_1, \dots, F_k and let \mathbf{n}_i be the primitive outer normal vector of F_i that is in S . Fix a vertex \mathbf{v} of F , let \mathbf{v}_i be a vertex of P that is adjacent to \mathbf{v} and not in F_i and \mathbf{d}_i be the primitive edge direction from \mathbf{v} to \mathbf{v}_i . If we let $\mathbf{u}_i := \mathbf{d}_i / \text{lin}(F)$, then the followings are true:*

- (1) $\langle \mathbf{d}_i, \mathbf{n}_i \rangle = -1$.
- (2) $\langle \mathbf{u}_i, \mathbf{n}_i \rangle = -1$.
- (3) $\langle \mathbf{u}_i, \mathbf{n}_j \rangle = 0$ for all $j \neq i$.
- (4) $\{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ generates $\text{fcone}^P(F, P)$.

Proof. For (1), let $H(C) := \{\mathbf{x} \in S \mid \langle \mathbf{n}_i, \mathbf{x} \rangle = C\}$ be the hyperplane in S that F_i is in. Since P is simple, all the vertices \mathbf{v}_l where $l \neq i$ are in F_i which means that the end points of the directions \mathbf{d}_l where $l \neq i$ are in $H(0)$. Since P is totally unimodular, the integer span of $\{\mathbf{d}_l \mid l \neq i\}$ is precisely all the integer points in $H(0)$. Also, by Lemma 3.6, there exists an integer point in $H(-1)$. Therefore, the end point of \mathbf{d}_i should be in $H(-1)$ (otherwise, $\{\mathbf{d}_1, \dots, \mathbf{d}_k\}$ doesn't span $\mathbb{Z}^n \cap S$). For (2), since \mathbf{u}_i is a projection of \mathbf{d}_i in to $\text{lin}(F)^\perp$, $\mathbf{u}_i - \mathbf{d}_i \in \text{lin}(F)$. Thus, $\langle \mathbf{u}_i - \mathbf{d}_i, \mathbf{n}_i \rangle = 0$ or equivalently, $\langle \mathbf{u}_i, \mathbf{n}_i \rangle = -1$ by (1). For (3), since P is simple, \mathbf{v}_i is in F_j for all $j \neq i$. Thus, $\langle \mathbf{d}_i, \mathbf{n}_j \rangle = 0$ for all $j \neq i$ which together with the fact that $\langle \mathbf{u}_i - \mathbf{d}_i, \mathbf{n}_j \rangle = 0$ for all $1 \leq j \leq k$ implies (3). The part (4) follows from Lemma 2.4 (4) and (3). \square

Since \mathbf{u}_i is in $\text{lin}(F)^\perp$, $\mathbf{u}_i = a_{i,1}\mathbf{n}_1 + \dots + a_{i,k}\mathbf{n}_k$ for some real numbers $a_{i,1}, \dots, a_{i,k}$. Let $c_{i,j} = \langle \mathbf{n}_i, \mathbf{n}_j \rangle$. We let C to be the $n \times n$ matrix with (i, j) -th entry as $c_{i,j}$ and D to be the $n \times n$ matrix with (i, j) -th entry as $a_{i,j}$.

Then we can express the the parts (2) and (3) from the above lemma as,

$$(3.2) \quad CD = -I.$$

Also, if we let \mathbf{a}_i to be the i -th column vector of D , we have,

$$(3.3) \quad \langle \mathbf{u}_i, \mathbf{u}_j \rangle = \mathbf{a}_i^t C \mathbf{a}_j = -a_{j,i}.$$

Let $M = (m_{i,j})$ be the inverse matrix of C . Then, (5.1) and (5.2) implies that

$$(3.4) \quad \langle \mathbf{u}_i, \mathbf{u}_j \rangle = m_{i,j}.$$

3.4. BV- α on $\text{PTes}_n(\mathbf{a})$. We have seen in Lemma 3.5 (2) that when $\mathbf{a} \in \mathbb{Z}_{>0}^n$, any face of $\text{PTes}_n(\mathbf{a})$ that is the intersection of facets $F_{i,j}$'s (refer to the equation (4.2)) with $i < j$ has the facet normal vector as a member of the standard basis up to sign. In this case, we get the BV- α value for free by Example 3.15 of [7].

Lemma 3.8. *Let $\mathbf{a} \in \mathbb{Z}_{>0}^n$ and F be a codimension k face of $\text{PTes}_n(\mathbf{a})$ that is an intersection of the facets $F_{i_1, j_1}, \dots, F_{i_k, j_k}$ where $i_1 < j_1, \dots, i_k < j_k$. Then, for every $1 \leq k \leq n$, $\alpha^{\text{BV}}(F) = \frac{1}{2^k}$.*

By Theorem 2.7 in [19], $\text{Tes}_n(\mathbf{a})$ is simple when $\mathbf{a} \in \mathbb{Z}_{>0}^n$. Also, one can check that the defining matrix for $\text{Tes}_n(\mathbf{a})$ is totally unimodular. Thus, $\text{Tes}_n(\mathbf{a})$ is totally unimodular for any $\mathbf{a} \in \mathbb{Z}_{>0}^n$. Since $\text{Tes}_n(\mathbf{a})$ and $\text{PTes}_n(\mathbf{a})$ are unimodularly equivalent, $\text{PTes}_n(\mathbf{a})$ is also totally unimodular. Therefore, we can apply the procedure described in the previous subsection and use Lemmas 2.11 and 2.12 to calculate the BV- α values of codimension 2 and codimension 3 faces of $\text{PTes}_n(\mathbf{1})$.

Since we already have Lemma 3.8, we only consider the case where at least one of the supporting facet for the face of consideration has form $F_{l,l}$. We calculate the BV- α values of the faces of $\text{PTes}_n(\mathbf{1})$ case by case. We will use the following terminologies to describe the cases.

We say that a position (i, j) of a matrix is *on the l -th hook* if either $i = l$ or $j = l$. More specifically, we say (i, j) is *on the row of the l -th hook* if $i = l$ and we say (i, j) is *on the column of the l -th hook* if $j = l$.

Lemma 3.9. *Let F be a codimension 2 face of $\text{PTes}_n(\mathbf{1})$ which is the intersection of facets $F_{l,l}$ and $F_{i,j}$. We consider all possible cases listed below.*

(1) *Suppose only one of the facet normal vectors is a shifted hook sum matrix (i.e, $i \neq j$). Then we have the following:*

(i) *If (i, j) is on the row of the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{4} + \frac{1}{12} \left(\frac{n}{n-1} \right)$.*

(ii) *If (i, j) is on the column of the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{6} - \frac{1}{12(n-1)}$.*

(iii) *If (i, j) is not on the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{4}$.*

(2) *Suppose both of the facet normal vectors are shifted hook sum matrices (i.e, $i = j$).*

Then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{4} - \frac{1}{36(n-1)}$.

Hence, all the values are positive for all $n \geq 3$.

Proof. We only provide a proof for case (1)/(i), but one can obtain the BV- α values for the other cases by following the same procedure. For case (1)/(i), by Lemma 3.5, the facet normal vectors are sm_l^{n-1} and $-e_{l,j}^{n-1}$. Thus, the matrix C in (3.2) is $\begin{pmatrix} n-1 & -1 \\ -1 & 1 \end{pmatrix}$ and the

inverse matrix M of C is $\begin{pmatrix} \frac{1}{n-2} & \frac{1}{n-2} \\ \frac{1}{n-2} & \frac{1}{n-2} \end{pmatrix}$. Therefore, by (3.4),

$$\langle \mathbf{u}_1, \mathbf{u}_1 \rangle = \frac{1}{n-2}, \langle \mathbf{u}_1, \mathbf{u}_2 \rangle = \frac{1}{n-2}, \text{ and } \langle \mathbf{u}_2, \mathbf{u}_2 \rangle = \frac{n-1}{n-2}.$$

By plugging in the dot product values to the formula in Lemma 2.11, we obtain

$$\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{4} + \frac{1}{12} \left(\frac{n}{n-1} \right). \quad \square$$

Lemma 3.10. *Let F be a codimension 3 face of $\text{PTes}_n(\mathbf{1})$ which is the intersection of facets $F_{l,l}, F_{i_1,j_1}$ and F_{i_2,j_2} . We consider all possible cases listed below.*

(1) *Suppose only one of the facet normal vectors is shifted hook sum matrix (i.e, $i_1 \neq j_1$ and $i_2 \neq j_2$).*

(i) *If both (i_1, j_1) and (i_2, j_2) are not on the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8}$.*

(ii) *If one of (i_2, j_2) and (i_3, j_3) is on the row of the l -th hook and the other is not on the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8} + \frac{n}{24(n-1)}$.*

(iii) *If one of (i_2, j_2) and (i_3, j_3) is on the column of the l -th hook and the other is not on the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{12} - \frac{1}{24(n-1)}$.*

(iv) *If both of (i_2, j_2) and (i_3, j_3) are on the row of the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8} + \frac{n}{12(n-2)}$.*

(v) *If one of (i_2, j_2) and (i_3, j_3) is on the column of the l -th hook and the other is on the row of the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8} - \frac{1}{12(n-2)}$.*

- (vi) If both of (i_2, j_2) and (i_3, j_3) are on the column of the l -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{24}$.
- (2) Suppose two of the facet normal vectors are shifted hook sum matrices (without loss of generality, assume $i_2 = j_2$ and let $m := i_2$).
- (i) If (i_3, j_3) is out of the l -th and m -th hook, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8} + \frac{1}{12(n-1)}$.
- (ii) If (i_3, j_3) is on the row of either the l -th hook or the m -th hook and is not on the hook of the other, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8} + \frac{n^2 + n - 3}{24(n^2 - 3n + 2)}$.
- (iii) If (i_3, j_3) is on the column of either the l -th hook or the m -th hook and is not on the hook of the other, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{12} - \frac{1}{24(n^2 - 3n + 2)}$.
- (iv) If (i_3, j_3) is on the column of either the l -th hook or the m -th hook and is on the row of the other, then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8}$.
- (3) Suppose all three of the facet normal vectors are shifted hook sum matrices (i.e., $i_2 = j_2$ and $i_3 = j_3$). Then $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1})) = \frac{1}{8} + \frac{1}{4(n-2)}$.

Hence, all the values are positive for all $n \geq 3$.

Proof. We follow the same procedure as in the proof of Lemma 3.9 (1)/(i) to obtain the formulas for $\alpha^{\text{BV}}(F, \text{PTes}_n(\mathbf{1}))$. From the formulas for each cases, it is very clear that the values are all positive for $n \geq 3$. \square

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1. By Corollary 2.6 in [19], the dimension of $\text{PTes}_n(\mathbf{1})$ is $d := \binom{n}{2}$. Thus, when $n \leq 2$, they are Ehrhart positive. Also, by Lemmas 3.9 and 3.10, and the equation (1.1), we obtain $e_{d-2}(\text{PTes}_n(\mathbf{1}))$ and $e_{d-3}(\text{PTes}_n(\mathbf{1}))$ are positive for all $n \geq 3$. \square

3.5. Proof of Corollaries 1.2 and 1.3. Using the Reduction Theorem (Theorem 2.14), we complete the proof of Corollary 1.2 which generalizes the positivity results given in Lemmas 3.9 and 3.10.

Proof of Corollary 1.2. Let P be any deformation of $\text{PTes}_n(\mathbf{1})$ with $\dim(P) = d$. If $d = \dim(\text{PTes}_n(\mathbf{1}))$, then by Lemmas 3.9, 3.10 and Theorem 2.14, $\alpha^{\text{BV}}(F, P) > 0$ for all codimension 2 and 3 faces F of P . Therefore, $e_{d-2}(P), e_{d-3}(P) > 0$ by (1.1). Similarly, if $d = \dim(\text{PTes}_n(\mathbf{1})) - 1$, Theorem 2.12 and 2.14 implies that $\alpha^{\text{BV}}(F, P) > 0$ for all codimension 3 faces F of P . Therefore, $e_{d-3}(P) > 0$ by (1.1). \square

Recall the definition for deformation in terms of normal fan. However, in the literature, there are multiple ways to define deformations of polytopes. Here, we provide one of them that we use.

Lemma 3.11. [23] *Let P_0 and Q be two polytopes in V . Then Q is a deformation of P_0 if and only if there exists a surjective map ϕ from the set of the vertices of P_0 to that of Q and $r_{i,j} \in \mathbb{R}_{\geq 0}$ such that $\phi(\mathbf{v}_i) - \phi(\mathbf{v}_j) = r_{i,j}(\mathbf{v}_i - \mathbf{v}_j)$ whenever \mathbf{v}_i and \mathbf{v}_j are adjacent vertices*

of P_0 . In particular, we say that Q is a weak deformation of P_0 if all $r_{i,j}$'s are positive and Q is a strong deformation of P_0 if some of $r_{i,j}$'s are zero.

Now, Corollary 1.3 follows directly from Corollary 1.2 and the following lemma:

Lemma 3.12. *Let Q, P be two polytopes and ψ is an invertible affine transformation from the affine hull of Q to the affine hull of P . Then Q is a deformation of P if and only if $\psi(Q)$ is a deformation of $\psi(P)$.*

Proof. For the forward direction, let \mathbf{v}, \mathbf{w} be any pair of adjacent vertices of P and \mathbf{v}', \mathbf{w}' be the corresponding vertices of Q . Then by Lemma 3.11, there exists $r \in \mathbb{R}$ such that $\mathbf{v} - \mathbf{w} = r(\mathbf{v}' - \mathbf{w}')$. Since ψ is an affine transformation, $\psi(\mathbf{v}) - \psi(\mathbf{w}) = r(\psi(\mathbf{v}') - \psi(\mathbf{w}'))$. Therefore, $\psi(Q)$ is a deformation of $\psi(P)$. The backward direction can be proven by the same argument. \square

4. TESLER POLYTOPES AND DEFORMATIONS

In this section, we explore deformations of $\text{Tes}_n(\mathbf{1})$. More specifically, we give a different proof of the known fact that for all fixed n , every tesler polytope $\text{Tes}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{1})$. Furthermore, we give a characterization for which flow polytope on the complete graph is a deformation of $\text{Tes}_n(\mathbf{1})$. Reduction theorem (Theorem 2.14) was proved using normal fan definition of a deformation but we use the equivalent definition of deformation since it works better for our case (Lemma 3.11).

4.1. For all $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$, $\text{Tes}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{1})$. To prove our result of this subsection, we will utilize a machinery that was used in the proof of Lemma 2.4 in [19].

Definition 4.1. For any $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$ and a vertex $\mathbf{v} = (v_{i,j}) \in \text{Vert}(\text{Tes}_n(\mathbf{a}))$,

(1) Define $j_{\mathbf{v}} : \{0, 1, 2, \dots, n\} \rightarrow \{0, 1, 2, \dots, n\}$ by

$$j_{\mathbf{v}}(k) = \begin{cases} l & \text{if the } k\text{-th row is not a zero row and } v_{k,l} \neq 0, \\ 0 & \text{if the } k\text{-th row is a zero row or } k = 0. \end{cases}$$

(this is well-defined by Theorem 2.6 (2)).

(2) Since \mathbf{v} is upper triangular, if we keep applying $j_{\mathbf{v}}$, the sequence $\{k, j_{\mathbf{v}}(k), j_{\mathbf{v}}^2(k), \dots\}$ will stabilize in a finite step. Assume that q is the smallest integer such that $j_{\mathbf{v}}^{q-1}(k) = j_{\mathbf{v}}^q(k)$. Define

$$\text{Dep}_{\mathbf{v}}(k) = \{(k, j_{\mathbf{v}}(k)), (j_{\mathbf{v}}(k), j_{\mathbf{v}}^2(k)) \dots, (j_{\mathbf{v}}^{q-1}(k), j_{\mathbf{v}}^q(k))\}.$$

(3) Let $D_{\mathbf{v}}(k) \in \mathbb{U}(n)$ be the matrix recording the positions appearing in $\text{Dep}_{\mathbf{v}}(k)$. More precisely,

$$D_{\mathbf{v}}(k) := (m_{i,j}) \text{ where } m_{i,j} = \begin{cases} 1, & \text{if } (i, j) \in \text{Dep}_{\mathbf{v}}(k), \\ 0, & \text{if } (i, j) \notin \text{Dep}_{\mathbf{v}}(k) \end{cases}.$$

The purpose of this definition is to describe the edge direction from a vertex of $\text{Tes}_n(\mathbf{a})$ to its adjacent vertex.

Example 4.2. Let \mathbf{v} and \mathbf{w} be as in Example 2.10. We obtain \mathbf{v} from \mathbf{w} by moving 2 on the first row to the second column and adjusting the lower rows to maintain the same hook sum. One sees that $\text{Dep}_{\mathbf{w}}(1) = \{(1, 3), (3, 3)\}$ is the set of all the positions that are affected by changing $w_{2,3}$ from 2 to 0 and $\text{Dep}_{\mathbf{v}}(1) = \{(1, 2), (2, 4), (4, 4)\}$ is the set of all the positions that are affected by the change $w_{1,2} = 0 \rightarrow 2$ (we have to add 2 to the positions to maintain the same hook sums). Thus, the edge direction from \mathbf{w} to \mathbf{v} can be described in terms of $D_{\mathbf{w}}(1)$ and $D_{\mathbf{v}}(1)$. Notice that changing the first coordinate of the hook sum vector \mathbf{a} to any other positive number would only change the length of the edge direction.

$$\mathbf{v} - \mathbf{w} = \begin{pmatrix} 0 & 2 & 0 & 0 \\ & 0 & 0 & 4 \\ & & 3 & 0 \\ & & & 8 \end{pmatrix} - \begin{pmatrix} 0 & 0 & 2 & 0 \\ & 0 & 0 & 2 \\ & & 5 & 0 \\ & & & 6 \end{pmatrix} = 2 \begin{pmatrix} 0 & 1 & -1 & 0 \\ & 0 & 0 & 1 \\ & & -1 & 0 \\ & & & 1 \end{pmatrix} = 2(D_{\mathbf{v}}(1) - D_{\mathbf{w}}(1))$$

We will show that for any $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$, the edge directions of $\text{Tes}_n(\mathbf{a})$ are always of the form as in the above example using the properties of $D_{\mathbf{v}}(k)$ stated in the Proposition below.

Proposition 4.3. *Let $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$ and $\mathbf{v} = (v_{i,j}) \in \text{Vert}(\text{Tes}_n(\mathbf{a}))$ such that the k -th row is a non-zero row. Suppose the non-zero entry in the k -th row is c and,*

$$\text{Dep}_{\mathbf{v}}(k) = \{(k, j_v(k)), (j_v(k), j_v^2(k)) \dots, (j_v^{q-1}(k), j_v^q(k))\}.$$

Then $D_{\mathbf{v}}(k)$ has the following properties:

- (1) The k -th row has one 1.
- (2) The m -th row and column are zero if $m \neq j_v^l(k)$ for $1 \leq l \leq q$.
- (3) The m -th row has exactly one 1 and the m -th column has exactly one 1 above the diagonal if $m = j_v^l(k)$ for $1 \leq l < q$.
- (4) $hs(D_{\mathbf{v}}(k)) = \mathbf{e}_k$ where \mathbf{e}_k is the k -th standard basis of \mathbb{R}^n
- (5) For any $1 \leq l \leq q - 1$, $v_{j_v^{l-1}(k), j_v^l(k)} \leq v_{j_v^l(k), j_v^{l+1}(k)}$. Thus, the entries of $\mathbf{v} - cD_{\mathbf{v}}(k)$ are non-negative.

Proof. (1),(2) and (3) are clear from the definition. For (4), let $D = D_{\mathbf{v}}(k)$. Then, by (2), $hs_m(D) = 0$ if $m \neq j_v^l(k)$ for some $1 \leq l \leq q$. If $m = j_v^l(k)$ for some $1 < l \leq q$, $hs_m(D) = 0$ by (3). Finally, when $m = k$, since k -th row has one 1 and the rows above k -th row are all zero rows, $hs_k(D) = 1$. For (5), consider the $j_v^l(k)$ -th hook sum condition $hs_{j_v^l(k)}(\mathbf{v})$. Assume $a_{j_v^l(k)} = 0$ and $v_{j_v^{l-1}(k), j_v^l(k)} > v_{j_v^l(k), j_v^{l+1}(k)}$. Then, since $v_{j_v^l(k), j_v^{l+1}(k)}$ is the only non-zero element in the row, we get $hs_{j_v^l(k)}(\mathbf{v}) = a_{j_v^l(k)} < 0$. Similarly, assume $a_{j_v^l(k)} > 0$ and $v_{j_v^{l-1}(k), j_v^l(k)} \geq v_{j_v^l(k), j_v^{l+1}(k)}$. Then the $j_v^l(k)$ -th hook sum condition would imply $a_{j_v^l(k)} \leq 0$. \square

Lemma 4.4. *For $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$, let $\mathbf{v} = (v_{i,j}), \mathbf{w} = (w_{i,j}) \in \text{Vert}(\text{Tes}_n(\mathbf{a}))$ are adjacent vertices of $\text{Tes}_n(\mathbf{a})$ where the rows of $\text{supp}(\mathbf{v})$ and $\text{supp}(\mathbf{w})$ are the same above the k -th row and differ in the k -th row. Then*

$$\mathbf{w} - \mathbf{v} = c(D_{\mathbf{w}}(k) - D_{\mathbf{v}}(k))$$

where $c \in \mathbb{R}_{> 0}$ is the unique non-zero entry in the k -th row of \mathbf{v} (or that of \mathbf{w}).

Proof. Let k be as in the above statement. Then, one of the 4 operations in Lemma 2.8 must have occurred in the k -th row of $\text{supp}(\mathbf{v})$ while getting $\text{supp}(\mathbf{w})$ from $\text{supp}(\mathbf{v})$. Obviously, (a) is impossible. Also, (b) and (c) are impossible, because if we assume that the k -th row of $\text{supp}(\mathbf{v})$ is a zero row but that of $\text{supp}(\mathbf{w})$ is not (or vice versa), since the rows above the k -th row are same, it would imply $a_k = 0$ and $a_k \neq 0$ by Lemma 2.9. Thus, the operation

(d) occurred and \mathbf{v} , \mathbf{w} each have exactly one non-zero entry in the k -th row, but in different places. Since $\text{supp}(\mathbf{v})$ and $\text{supp}(\mathbf{w})$ have the same rows above the k -th row, by the k -th hook sum condition, we obtain $v_{k,j_{\mathbf{v}}(k)} = w_{k,j_{\mathbf{w}}(k)}$. Let

$$c := v_{k,j_{\mathbf{v}}(k)} = w_{k,j_{\mathbf{w}}(k)}.$$

Then, by Proposition 4.3 (4), $c \text{hs}(D_{\mathbf{v}}(k)) = c \text{hs}(D_{\mathbf{w}}(k)) = c\mathbf{e}_k$. Thus, if we let $\mathbf{u} := \mathbf{w} + cD_{\mathbf{v}}(k) - cD_{\mathbf{w}}(k)$, then

$$\text{hs}(\mathbf{u}) = \mathbf{a} - c\mathbf{e}_k + c\mathbf{e}_k = \mathbf{a}.$$

Also, by Proposition 4.3 (5), the entries of $\mathbf{w} - cD_{\mathbf{w}}(k)$ are non-negative which implies that $\mathbf{u} = \mathbf{w} + cD_{\mathbf{v}}(k) - cD_{\mathbf{w}}(k) \in \text{Tes}_n(\mathbf{a})$. We will prove that $\mathbf{u} = \mathbf{v}$ in two steps. First, we show that \mathbf{u} is, in fact, a vertex of $\text{Tes}_n(\mathbf{a})$ by showing that each row of \mathbf{u} has at most one 1. Second, we show that $\text{supp}(\mathbf{u}) \leq \text{supp}(\mathbf{v})$.

We now show \mathbf{u} is a vertex of $\text{Tes}_n(\mathbf{a})$. Notice that since $D_{\mathbf{v}}(k)$ and $\mathbf{w} - cD_{\mathbf{w}}(k)$ have non-negative entries, we get $\text{supp}(\mathbf{u}) = \text{supp}(D_{\mathbf{v}}(k) + \mathbf{w} - cD_{\mathbf{w}}(k)) = \text{supp}(D_{\mathbf{v}}(k), \mathbf{w} - cD_{\mathbf{w}}(k))$. Also, observe that directly from the definition of $D_{\mathbf{v}}(k)$, $\text{supp}(cD_{\mathbf{v}}(k)) = \text{supp}(D_{\mathbf{v}}(k)) \leq \text{supp}(\mathbf{v})$. Similarly, $\text{supp}(cD_{\mathbf{w}}(k)) \leq \text{supp}(\mathbf{w})$ which implies $\text{supp}(\mathbf{w} - cD_{\mathbf{w}}(k)) \leq \text{supp}(\mathbf{w})$. Therefore,

$$(4.1) \quad \text{supp}(\mathbf{u}) = \text{supp}(\{D_{\mathbf{v}}(k), \mathbf{w} - cD_{\mathbf{w}}(k)\}) \leq \text{supp}(\{\mathbf{v}, \mathbf{w}\}).$$

By Lemma 2.8, the k -th row of $\text{supp}(\mathbf{v}, \mathbf{w})$ has two 1's and at most one 1 in every other rows which implies $\text{supp}(\mathbf{u})$ has at most one 1 except for the k -th row. Finally, for the k -th row of \mathbf{u} , since $v_{k,j_{\mathbf{v}}(k)} = w_{k,j_{\mathbf{w}}(k)} = c$, adding $cD_{\mathbf{v}}(k)$ and subtracting $cD_{\mathbf{w}}(k)$ from \mathbf{w} moves the unique 1 in the k -th row of $\text{supp}(\mathbf{w})$ to the position $(k, j_{\mathbf{v}}(k))$. Therefore $\text{supp}(\mathbf{u})$ has at most one 1 in each row.

For the second step, it is clear that the rows above the k -th row of \mathbf{u} are same as those of $\text{supp}(\mathbf{v})$. Also, we can easily see that the k -th row of \mathbf{v} and \mathbf{u} are the same. Thus, $\text{supp}(\mathbf{u})$ is same as $\text{supp}(\mathbf{v})$ up to the k -th row. For the lower rows, we have to show the following two statements:

- (a) There is no $l > k$ such that both l -th row of $\text{supp}(\mathbf{v})$ and $\text{supp}(\mathbf{u})$ has one 1 but in a different places.
- (b) There is no $l > k$ such that the l -th row of $\text{supp}(\mathbf{u})$ is a non-zero row but that of $\text{supp}(\mathbf{v})$ is a zero row.

The part (a) directly follow from the inequality (3.2). For (b), assume for the contradiction that we have such l 's and l_0 is the smallest of them. Since l_0 -th row of $\text{supp}(\mathbf{v})$ is a zero row, by Lemma 2.9, we have $a_{l_0} = 0$ and the entries of l_0 -th column of $\text{supp}(\mathbf{v})$ above the diagonal are all zero but at least one of the entries of l_0 -th column of $\text{supp}(\mathbf{u})$ has to be non-zero. Let u_{l_1, l_0} be one of the non-zero entry of \mathbf{u} in the l_0 -th column above the diagonal ($l_1 < l_0$). Since $\text{supp}(\mathbf{v})$ and $\text{supp}(\mathbf{u})$ are same up to k -th row, we have $k < l_1$. Thus, if the l_1 -th row of $\text{supp}(\mathbf{v})$ is a zero row, it contradicts the minimality of l_0 . Hence, the l_1 -th row of $\text{supp}(\mathbf{v})$ have exactly one 1 and the position of this 1 cannot be (l_1, l_0) -the position of the 1 in l_1 -th row of $\text{supp}(\mathbf{u})$ which is impossible by (a). Therefore, $\text{supp}(\mathbf{u}) \leq \text{supp}(\mathbf{v})$, and by Lemma 2.7, $\{\mathbf{u}\} \subset \{\mathbf{v}\}$ which implies $\mathbf{u} = \mathbf{w} + cD_{\mathbf{v}}(k) - cD_{\mathbf{w}}(k) = \mathbf{v}$. \square

We need one more lemma before proving the result of this subsection.

Lemma 4.5. *Let $\mathbf{a}_0 \in \mathbb{R}_{>0}^n$ and $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$. Then, for any $\mathbf{v} = (v_{i,j}) \in \text{Vert}(\text{Tes}_n(\mathbf{a}_0))$, there exists a unique $\mathbf{v}' \in \text{Vert}(\text{Tes}_n(\mathbf{a}))$ such that $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v})$.*

Proof. Let \mathbf{v}' be any point in $\text{Tes}_n(\mathbf{a})$ such that $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v})$. Then \mathbf{v}' has to satisfy

$$\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v}) \text{ and } hs(\mathbf{v}') = \mathbf{a}.$$

One can easily see that there is a unique solution to the above two conditions, because $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v})$ implies that there can be only one possible non-zero entry for each row of \mathbf{v}' (this also implies that \mathbf{v}' is a vertex of $\text{Tes}_n(\mathbf{a})$ by Lemma 2.6 (2)) and the values of the entries are uniquely determined by the hook sum conditions $hs(\mathbf{v}') = \mathbf{a}$. \square

Now, we are ready to prove our main result of this section.

Theorem 4.6. *Let n be any fixed positive integer and $\mathbf{a}_0 \in \mathbb{R}_{>0}^n$. Then for any $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$, $\text{Tes}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{a}_0)$. In particular, $\text{Tes}_n(\mathbf{a})$ is a weak deformation of $\text{Tes}_n(\mathbf{a}_0)$ when $\mathbf{a} \in \mathbb{R}_{>0}^n$ and is a strong deformation of $\text{Tes}_n(\mathbf{a}_0)$ when some of the coordinates of \mathbf{a} are zero.*

Proof. Since $\text{Tes}_n(0, \dots, 0)$ is a point, it is clearly a deformation of any non-empty polytope. Assume that $\mathbf{a} \in \mathbb{R}_{\geq 0}^n$ and $\mathbf{a} \neq (0, \dots, 0)$. We prove the theorem using Lemma 3.11. To use the lemma, we define a map between the set of the vertices. Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$. Define

$$\phi : \text{Vert}(\text{Tes}_n(\mathbf{a}_0)) \longrightarrow \text{Vert}(\text{Tes}_n(\mathbf{a})),$$

by letting $\phi(\mathbf{v})$ to be the unique \mathbf{v}' satisfying $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v})$ asserted by Lemma 4.5. One sees that ϕ is well-defined by Lemma 4.5.

ϕ is well-defined by Lemma 4.5. The surjectivity of ϕ comes from Lemma 2.6 (2), because for any matrix M with only 0,1 entries where each row has exactly one 1, there exist $\mathbf{v} \in \text{Tes}_n(\mathbf{a}_0)$ such that $\text{supp}(\mathbf{v}) = M$.

Now, we show that $\text{Tes}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{a}_0)$. Assume that \mathbf{v} and \mathbf{w} are adjacent vertices of $\text{Tes}_n(\mathbf{a}_0)$ where their support only differ in the k -th row. Let $\mathbf{v}' = \phi(\mathbf{v})$ and $\mathbf{w}' = \phi(\mathbf{w})$. We need to show that there exists $r_{\mathbf{w},\mathbf{v}} \in \mathbb{R}_{\geq 0}$ such that

$$\mathbf{w}' - \mathbf{v}' = r_{\mathbf{w},\mathbf{v}}(\mathbf{w} - \mathbf{v}).$$

Case1 : If the k -th row of $\text{supp}(\mathbf{v}')$ or $\text{supp}(\mathbf{w}')$ is a zero row (without loss of generality, assume that the k -th row of $\text{supp}(\mathbf{v}')$ is a zero row), then since $\text{supp}(\mathbf{v})$ and $\text{supp}(\mathbf{w})$ only differs in the k -th row, $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{w})$. Thus, $\mathbf{v}' = \mathbf{w}'$ by Lemma 4.5. Therefore, we can take $r_{\mathbf{v},\mathbf{w}} = 0$.

Case2 : If neither the k -th row of $\text{supp}(\mathbf{v}')$ nor that of $\text{supp}(\mathbf{w}')$ is a zero, since $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v})$, we have $j_{\mathbf{v}'}(k) = j_{\mathbf{v}}(k)$. Let $\mathbf{v}' = (v'_{i,j})$. Since $v'_{k,j_{\mathbf{v}'}(k)} \neq 0$, $j_{\mathbf{v}'}(k)$ -th row of \mathbf{v}' is a non-zero row by Lemma 2.9. Since $\text{supp}(\mathbf{v}') \leq \text{supp}(\mathbf{v})$, we get $j_{\mathbf{v}'}^2(k) = j_{\mathbf{v}}^2(k)$. By iterating this process, we obtain $j_{\mathbf{v}'}^l(k) = j_{\mathbf{v}}^l(k)$ for any integer $0 \leq l$. Therefore, $\text{Dep}_{\mathbf{v}'}(k) = \text{Dep}_{\mathbf{v}}(k)$ or equivalently, $D_{\mathbf{v}'}(k) = D_{\mathbf{v}}(k)$. Similarly, we obtain $D_{\mathbf{w}'}(k) = D_{\mathbf{w}}(k)$. By Lemma 4.4,

$$\mathbf{w} - \mathbf{v} = c(D_{\mathbf{w}}(k) - D_{\mathbf{v}}(k)) \text{ and } \mathbf{w}' - \mathbf{v}' = d(D_{\mathbf{w}'}(k) - D_{\mathbf{v}'}(k))$$

where c and d are the unique non-zero (positive) entries of \mathbf{v} and \mathbf{v}' respectively. Therefore, we can take $r_{\mathbf{w},\mathbf{v}} = \frac{c}{d}$.

In particular, when $\mathbf{a} \in \mathbb{R}_{>0}^n$, $r_{v,w}$ is always positive because **Case1** does not happen. However, when some of the a'_i 's are zero, one can easily see that **Case1** does happen which means the corresponding $r_{v,w}$'s are zero. □

4.2. Flow polytopes on the complete graph that are deformations of the Tesler polytopes. We first restrict ourselves to the complete graphs with $n+1$ vertices K_{n+1} . Flow polytope is defined over (complete) graphs, but in this section, we use another definition that gives a polytope that is unimodularly equivalent to the flow polytope.

Definition 4.7. For $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$, the *flow polytope* on K_{n+1} with net flow a_i on vertex i for $1 \leq i \leq n$ is defined by,

$$\text{Flow}_n(\mathbf{a}) := \{M = (m_{i,j}) \in \mathbb{U}(n) \mid hs(M) = \mathbf{a}, m_{i,j} \geq 0\}.$$

An element of $\text{Flow}_n(\mathbf{a})$ is called as *flow*.

One can show that,

$$\text{Flow}_n(\mathbf{a}) \text{ is non-empty if and only if } \sum_{i \leq l} a_i \geq 0 \text{ for all } 1 \leq l \leq n.$$

In this subsection, we will let A_n to be the set of all $\mathbf{a} \in \mathbb{R}^n$ that satisfies the above condition. To give a characterization of $\mathbf{a} \in \mathbb{R}^n$ where $\text{Flow}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{1})$, we first prove that any deformation of $\text{Tes}_n(\mathbf{1})$ is a translation of some Tesler polytope (Theorem 4.10).

Let $P(n)$ be the following set of polytopes

$$\left\{ P \subset \mathbb{R}^{\binom{n}{2}} \mid P \text{ is a deformation of } \text{Tes}_n(\mathbf{1}) \right\}.$$

By the definition of a deformation, For any $P \in P(n)$, we can write

$$P = \{M = (m_{i,j}) \in \mathbb{U}(n) : hs(M) = \boldsymbol{\beta} \text{ and } m_{i,j} \geq -b_{i,j} \text{ for every } i \leq j\}$$

where $\boldsymbol{\beta} \in \mathbb{R}^n$ and $b_{i,j} \in \mathbb{R}$ for any $i \leq j$. The only difference from Tesler polytopes are that, in this case, we use $x_{i,j} \geq -b_{i,j}$ for defining inequalities instead of $x_{i,j} \geq 0$ which corresponds to translations of facet defining planes. We know that any vertex v of $\text{Tes}_n(\mathbf{a})$ is a matrix that has at most one non-zero element in each rows. So, to find a vertex of P corresponding to v , we first replace the zero in (i,j) -th position in the matrix v by $-b_{i,j}$'s. Then hook sum conditions determines the values of the undetermined entries. One can deduce from [8], Corollary 2.14 and Proposition 2.16 that the above method of corresponding vertices yields a map from $\text{Vert}(\text{Tes}_n(\mathbf{1}))$ to $\text{Vert}(P)$ satisfying the conditions of Lemma 3.11.

Example 4.8. With P as the above remark with $n = 4$ and $\boldsymbol{\beta} = (4, 3, 8, 1)$, let us find the vertex v of P corresponding to a vertex v' of $\text{Tes}_4(1, 1, 1, 1)$. As explained above, we first change zeros with $-b_{i,j}$'s and the hook sum conditions for P determines w, x, y, z .

$$v' := \begin{bmatrix} 0 & 1 & 0 & 0 \\ & 0 & 2 & 0 \\ & & 3 & 0 \\ & & & 1 \end{bmatrix} \rightarrow v := \begin{bmatrix} -b_{1,1} & w & -b_{1,3} & -b_{1,4} \\ & -b_{2,2} & x & -b_{2,4} \\ & & y & -b_{3,4} \\ & & & z \end{bmatrix} \rightarrow \begin{cases} hs_1(v) = -b_{1,1} + w - b_{1,3} - b_{1,4} = 4 \\ hs_2(v) = -b_{2,2} + x - b_{2,4} - w = 3 \\ hs_3(v) = y - b_{3,4} - (x - b_{1,3}) = 8 \\ hs_4(v) = z - (-b_{1,4} - b_{2,4} - b_{3,4}) = 1 \end{cases}$$

By solving the system of equations, we get

$$\begin{cases} w = 4 + b_{1,1} + b_{1,3} + b_{1,4} \\ x = 3 + b_{2,2} + b_{2,4} + w \\ y = 8 + b_{3,4} + x - b_{1,3} \\ z = 1 - b_{1,4} - b_{2,4} - b_{3,4} \end{cases}$$

To state the next lemma, we introduce the following notations. Let $\widetilde{\mathbb{U}}(n)$ be the set of $(n-1) \times n$ matrices \widetilde{M} obtained by deleting the n -th row of the matrices $M \in \mathbb{U}(n)$. We define $hs_k(\widetilde{M}) = hs_k(M)$ for any $1 \leq k \leq n-1$.

Lemma 4.9. *Let*

$$P = \{M = (m_{i,j}) \in \mathbb{U}(n) \mid hs(M) = (\tilde{b}_{1,n}, \dots, \tilde{b}_{n,n}) \text{ and } m_{i,j} \geq -\tilde{b}_{i,j} \text{ for every } i \leq j\}$$

where $\tilde{B} := (\tilde{b}_{i,j}) \in \widetilde{\mathbb{U}}(n+1)$. Then $P \in P(n)$ if and only if $hs_i(\tilde{B}) \geq 0$ for all $1 \leq i \leq n$.

Proof. For the forward direction, assume $P \in P(n)$. For $1 \leq k \leq n-1$ let $d_k \in \mathbb{U}(n)$

$$\text{be } d_k = (d_{i,j}) \text{ where } d_{i,j} = \begin{cases} 1 & \text{if } k \neq i \text{ and } i = j \text{ or } k+1 \neq i \text{ and } i = j \\ 1 & \text{if } k = i = j - 1 \\ 2 & \text{if } k+1 = i = j \\ 0 & \text{otherwise} \end{cases} \quad \text{Then we see that}$$

$\mathbf{d}_k \in \text{Vert}(\text{Tes}_n(\mathbf{1}))$ by Lemma 2.6 (2). Let $\mathbf{w}_k = (w_{i,j}^k) \in \mathbb{U}(n)$ be a vertex of P corresponding to \mathbf{d}_k . To calculate the entries of \mathbf{w}_k , replace the zero in (i,j) -th position in \mathbf{d}_k with $-\tilde{b}_{i,j}$ for all $1 \leq i \leq j \leq n$ which determines all the entries of \mathbf{w}_k except one entry in each rows (Let us denote the undetermined entry in i -th row by x_i). And then determine x_i 's by using the hook sum conditions in an iterative manner from the first hook sum condition to the last hook sum condition. After solving the system of equations, (Since $hs_i(\mathbf{w}_k) = 0$ only depends on first i rows, a simple induction argument shows that the system of equations is always solvable and the solution is unique), we obtain

$$w_{k,k+1}^k = hs_k(\tilde{B}) - \tilde{b}_{k,k+1}.$$

Let I be the n by n identity matrix (I is clearly a vertex of $\text{Tes}_n(\mathbf{1})$) and $\mathbf{w} = (w_{i,j}) \in \mathbb{U}(n)$ be the corresponding vertex of P . Then,

$$(d_k - I)_{k,k+1} = 1 \text{ and } (\mathbf{w}^k - \mathbf{w})_{k,k+1} = hs_k(\tilde{B}),$$

because $w_{k,k+1} = -\tilde{b}_{k,k+1}$. By Lemma 3.11, $hs_k(\tilde{B})$ must be a non-negative. Also, since $\mathbf{w} \in P$, we have $w_{n,n} = hs_n(\tilde{B}) - \tilde{b}_{n,n} \geq -\tilde{b}_{n,n}$. Therefore, we obtain the desired result. Conversely, if we assume $hs_i(\tilde{B}) \geq 0$ for all $1 \leq i \leq n$ and let $B \in \mathbb{U}(n)$ be the matrix we obtain by removing the $n+1$ -th column of \tilde{B} , one can easily check that the polytope $P+B$ obtained by shifting P by the matrix B is:

$$P+B = \{M+B \mid M \in P\} = \text{Tes}_n(hs_1(\tilde{B}), \dots, hs_n(\tilde{B})),$$

$$\text{Case 1 : } \mathbf{f}_1 = \begin{bmatrix} a_1 - b_1 & b_1 & 0 \\ a_2 - b_2 & b_1 + b_2 & b_1 + b_2 \\ a_3 - b_3 & a_3 - b_3 & a_3 - b_3 \end{bmatrix}, \mathbf{f}_2 = \begin{bmatrix} a_1 - b_1 & 0 & b_1 \\ a_2 - b_2 & b_2 & b_2 \\ a_3 - b_3 & a_3 - b_3 & a_3 - b_3 \end{bmatrix}$$

$$\text{Case 2 : } \mathbf{f}_1 = \begin{bmatrix} 0 & b_1 & 0 & a_1 - b_1 & 0 \\ 0 & 0 & b_1 + b_2 & a_2 - b_2 & 0 \\ & & * & * & * \\ & & & * & * \\ & & & & * \end{bmatrix}, \mathbf{f}_2 = \begin{bmatrix} 0 & 0 & b_1 & a_1 - b_1 & 0 \\ 0 & 0 & b_2 & a_2 - b_2 & 0 \\ & & * & * & * \\ & & & * & * \\ & & & & * \end{bmatrix}$$

FIGURE 1. **Case 1** : is when $k = n = 3$ and **Case 2** : is when $n = 5$ and $k = 3$.

which is a deformation of $\text{Tes}_n(\mathbf{1})$ by Theorem 4.6 □

Now, we are ready to prove the following theorem.

Theorem 4.10. *Any polytope $P \in P(n)$ is a translation of some Tesler polytope.*

Proof. Let $P = \{M = (m_{i,j}) \in \mathbb{U}(n) : hs(M) = \boldsymbol{\beta} \text{ and } m_{i,j} \geq c_{i,j} \text{ for every } i \leq j\}$ be a polytope in $P(n)$ and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_n)$. Let $D = (d_{i,j})$ be a matrix such that

$$d_{i,j} = \begin{cases} \beta_i & \text{if } i=j \\ 0 & \text{if } i \neq j \end{cases} \text{ then}$$

$$P - D = \{M = (m_{i,j}) \in \mathbb{U}(n) : hs(M) = 0 \text{ and } x_{i,j} \geq b_{i,j} \text{ for every } i \leq j\}$$

$$\text{where } b_{i,j} = \begin{cases} c_{i,i} - \beta_i & \text{if } i = j \\ c_{i,j} & \text{if } i \neq j \end{cases}$$

because $hs(D) = \boldsymbol{\beta}$ and hs is a linear transformation. Since $P - D$ is just a translation of P , $P - D \in P(n)$. Thus, if we let $B := (b_{i,j}) \in \mathbb{U}(n)$ then $-hs_k(B) > 0$ for $1 \leq k \leq n - 1$ and $-hs_n(B) \geq 0$ by Lemma 4.9. Thus,

$$P - D - B = \{M = (m_{i,j}) \in \mathbb{U}(n) : hs(M) = (hs_1(-B), \dots, hs_n(-B)) \text{ and } x_{i,j} \geq 0 \text{ for every } i \leq j\}$$

is a Tesler polytope of hook sum $(hs_1(-B), \dots, hs_n(-B))$ which is a deformation of $\text{Tes}_n(\mathbf{1})$ by Theorem 4.6. □

Using the above theorem with the following lemma, we can characterize which flow polytope on a complete graph is a deformation of $\text{Tes}_n(\mathbf{1})$.

Lemma 4.11. *Let $n \geq 3$ and $\mathbf{a} = (a_1, \dots, a_n) \in A$. Assume that $a_1 > 0, a_2 \geq 0$ and a_k is the first negative component of \mathbf{a} where $3 \leq k \leq n$. Then there exists two flows $\mathbf{f}_1 = (f_{i,j}^1)$ and $\mathbf{f}_2 = (f_{i,j}^2)$ in $\text{Flow}_n(\mathbf{a})$ such that $f_{i,k}^1 = 0$ for any $1 \leq i \leq k - 2$ and $f_{k-1,k}^2 < |a_k|$.*

Proof. We will provide the result in two cases where in the first case, we will directly construct \mathbf{f}_1 and \mathbf{f}_2 and for the second case, we will construct two points $\mathbf{p}_1, \mathbf{p}_2$ and find a point \mathbf{p} where $\mathbf{p} + \mathbf{p}_1$ and $\mathbf{p} + \mathbf{p}_2$ gives our desired flows. For both of the cases, we use the following fact:

There exist b_1, \dots, b_{k-1} such that,

$$(4.2) \quad 0 < b_1 \leq a_1, \quad 0 \leq b_l \leq a_l \text{ for all } 2 \leq l \leq k - 1 \text{ and } \sum_{1 \leq l \leq k-1} b_l + a_k = 0.$$

The existence of such b_i 's is guaranteed since $\sum_{1 \leq l \leq k} a_l \geq 0$, $a_1 > 0$ and $a_2, \dots, a_{k-1} \geq 0$.

Case 1 : Suppose $k = n$,

Let $\mathbf{f}_1 = (f_{i,j}^1)$ and $\mathbf{f}_2 = (f_{i,j}^2)$ where

$$f_{i,j}^1 = \begin{cases} \sum_{1 \leq l \leq j-1} b_l & \text{if } j = i + 1 \leq k, \\ a_i - b_i & \text{if } i = j, \\ 0 & \text{otherwise,} \end{cases} \quad \text{and } f_{i,j}^2 = \begin{cases} b_i & \text{if } j = k, \\ a_i - b_i & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

It is clear that \mathbf{f}_1 and \mathbf{f}_2 are in $\text{Flow}_n(\mathbf{a})$ and have desired properties.

Case 2 : Suppose $k < n$,

We first construct \mathbf{p}_1 and \mathbf{p}_2 . Let $\mathbf{p}_1 = (p_{i,j}^1)$ and $\mathbf{p}_2 = (p_{i,j}^2)$ where

$$p_{i,j}^1 = \begin{cases} \sum_{1 \leq l \leq j-1} b_l & \text{if } j = i + 1 \leq k, \\ a_i - b_i & \text{if } j = k + 1, \\ 0 & \text{otherwise,} \end{cases} \quad \text{and } p_{i,j}^2 = \begin{cases} b_i & \text{if } j = k, \\ a_i - b_i & \text{if } j = k + 1, \\ 0 & \text{otherwise.} \end{cases}$$

Next, we will find \mathbf{p} . Since $\sum_{1 \leq l \leq m} a_l$ for all $1 \leq m \leq n$ and $\sum_{1 \leq l \leq k-1} b_l + a_k = 0$, we have,

$$\sum_{1 \leq l \leq k-1} a_l - b_l + \sum_{k+1 \leq l \leq m} a_l = \sum_{1 \leq l \leq m} a_l - \left(\sum_{1 \leq l \leq k-1} b_l + a_k \right) \geq 0 \text{ for all } k+1 \leq m \leq n.$$

By the above equation and the fact that $a_l - b_l \geq 0$ for all $1 \leq l \leq k-1$, we have that

$$\mathbf{a}' := \underbrace{(0, \dots, 0)}_k, \sum_{1 \leq l \leq k-1} a_l - b_l, a_{k+2}, \dots, a_n \in A.$$

Therefore, $\text{Flow}_n(\mathbf{a}')$ is non-empty. Let $\mathbf{p} = (p_{i,j})$ be any point in $\text{Flow}_n(\mathbf{a}')$. Then, since first k components of \mathbf{a}' are zero, we have $p_{i,j} = 0$ for any $i \leq k$. Clearly, $\mathbf{f}_1, \mathbf{f}_2 \in \text{Flow}_n(\mathbf{a})$ and $f_{i,k}^1 = 0$ for all $1 \leq i \leq k-2$. Moreover, one can check that $f_{k-1,k}^2 < |a_k|$ follows from the equation (4.2). Therefore, \mathbf{f}_1 and \mathbf{f}_2 have the desired properties. \square

we are now ready to prove Theorem 1.4.

Proof of Theorem 1.4. We first show the above statement for the case when $l = 1$ and then show that the other cases can be reduced to this case.

Assume that $l = 1$. For the backward direction, assuming that $a_3, a_4, \dots, a_n \geq 0$, we will show $\text{Flow}_n(\mathbf{a})$ is a Tesler polytope (up to a shift, which by Theorem 4.10 is a deformation of $\text{Tes}_n(\mathbf{1})$). If $a_2 \geq 0$, then $\text{Flow}_n(\mathbf{a}) = \text{Tes}_n(\mathbf{a})$. If $a_2 < 0$, then $a_1 \geq |a_2|$ and $m_{1,2} \geq -a_2$ (otherwise the condition $hs_2(M) = a_2$ cannot be satisfied). Thus, if we let $T = (t_{i,j}) \in \mathbb{U}(n)$

where $t_{i,j} = \begin{cases} a_{1,2} & (i,j) = (1,2) \\ 0 & \text{otherwise} \end{cases}$ and $\mathbf{a}' = (a_1 - |a_2|, 0, a_3, a_4, \dots, a_n)$, then

$$\text{Flow}_n(\mathbf{a}) - T = \{M = (m_{i,j}) \in \mathbb{U}(n) \mid hs(M) = \mathbf{a}' \text{ and } m_{i,j} \geq 0 \text{ for all } 1 \leq i \leq j \leq n\},$$

which is exactly $\text{Tes}_n(a_1 - |a_2|, 0, a_3, a_4, \dots, a_n)$ (notice that the above equation is still true without the assumption that $a_3, \dots, a_n \geq 0$).

For the forward direction, we assume that $\text{Flow}_n(\mathbf{a})$ is a deformation of $\text{Tes}_n(\mathbf{1})$ and therefore, a translation of $\text{Tes}_n(\mathbf{a}_0)$ for some $\mathbf{a}_0 \in \mathbb{R}_{\geq 0}^n$ by Theorem 4.10. Also, for contradiction we assume that at least one of a_3, a_4, \dots, a_n is negative. It is enough to show for the case when $a_2 \geq 0$ because in the proof of the backward direction, we showed that if $a_2 < 0$, $\text{Flow}_n(\mathbf{a})$

is a translation of $\text{Flow}_n(a_1 - |a_2|, 0, a_3, a_4, \dots, a_n)$. Let $T = (t_{i,j}) \in \mathbb{U}(n)$ be the point such that $\text{Flow}_n(\mathbf{a}) + T = \text{Tes}_n(\mathbf{a}_0)$. First of all, $t_{i,j} \leq 0$ for every (i, j) because if $t_{i,j} > 0$ for some (i, j) , then the (i, j) -th position of every point in $\text{Flow}_n(\mathbf{a}) + T = \text{Tes}_n(\mathbf{a}_0)$ is positive which is impossible by Lemma 2.6 (2) (there exists a vertex of $\text{Tes}_n(\mathbf{a}_0)$ where (i, j) -th position is 0). Let a_k be the first negative component of \mathbf{a} ($3 \leq k \leq n$). By Lemma 4.11, there exists two flows $\mathbf{f}_1 = (f_{i,j}^1)$ and $\mathbf{f}_2 = (f_{i,j}^2)$ in $\text{Flow}_n(\mathbf{a})$ such that $f_{i,k}^1 = 0$ for all $1 \leq i \leq k-2$ and $f_{k-1,k}^2 < |a_k|$. Since $\mathbf{f}_1 + T \in \text{Tes}_n(\mathbf{a}_0)$, $t_{i,k} = 0$ for all $1 \leq i \leq k-2$ because otherwise, $\mathbf{f}_1 + T$ would have a negative component which is impossible. Thus, $hs_k(T) = t_{k,k} + \dots + t_{k,n} - t_{k-1,k}$. Also, since $hs_k(\text{Flow}_n(\mathbf{a}) + T) = a_k + t_{k,k} + \dots + t_{k,n} - t_{k-1,k} = hs_k(\mathbf{a}_0) \geq 0$ and $t_{k,k}, \dots, t_{k,n} \leq 0$, we have $a_k \geq t_{k-1,k}$. Therefore, the $(k-1, k)$ -th position of $\mathbf{f}_2 + T$ is negative which is a contradiction.

Now assume $l > 1$, the proof for the backward direction follows from the proof above for $l = 1$ case because $\text{Flow}_n(\underbrace{0, \dots, 0}_{l-1}, a_l, \dots, a_n)$ and $\text{Tes}_n(\underbrace{0, \dots, 0}_{l-1}, 1, \dots, 1)$ are unimodular

equivalent to $\text{Flow}_{n-l+1}(a_l, \dots, a_n)$ and $\text{Tes}_{n-l+1}(1, \dots, 1)$ respectively which the latter is a deformation of $\text{Tes}_n(\mathbf{1})$ by Theorem 4.6. The forward direction can also be reduced to the $l = 1$ case. Similarly as before, $\text{Flow}_n(\mathbf{a}) + T = \text{Tes}_n(\mathbf{a}_0)$ for some $\mathbf{a}_0 \in \mathbb{R}_{\geq 0}^n$, $T \in \mathbb{U}(n)$, where all the entries of T are either zero or negative. Clearly, the first $(l-1)$ rows of any point in $\text{Flow}_n(\mathbf{a})$ is zero, which implies that the first $(l-1)$ rows of T is zero (otherwise, $\text{Tes}_n(\mathbf{a}_0)$ have a negative hook sum). Hence, the first $(l-1)$ rows of any points in $\text{Tes}_n(\mathbf{a}_0)$ is zero and so the first $l-1$ entries of \mathbf{a}_0 are zero. Therefore, we can instead work on $\text{Flow}_{n-l}(a_{l+1}, \dots, a_n) + T' = \text{Tes}_{n-l}(\mathbf{a}'_0)$ where $T' \in \mathbb{U}(n-l+1)$ is obtained from T by removing the first $l-1$ rows and columns and $\mathbf{a}'_0 \in \mathbb{R}_{\geq 0}^{n-l+1}$ is obtained from \mathbf{a}_0 by removing the first $l-1$ entries of \mathbf{a}_0 and obtain the same result as in $l = 1$ case. \square

REFERENCES

- [1] D. Armstrong, A. Garsia, J. Haglund, B. Rhoades, and B. Sagan. Combinatorics of tesler matrices in the theory of parking functions and diagonal harmonics. *Journal of Combinatorics*, 3(3):451–494, 2012.
- [2] A. Barvinok. *Integer points in polyhedra*, volume 452. European Mathematical Society, 2008.
- [3] C. Benedetti, R. González D’León, C. Hanusa, P. Harris, A. Khare, A. Morales, and M. Yip. A combinatorial model for computing volumes of flow polytopes. *Transactions of the American Mathematical Society*, 2019.
- [4] N. Berline and M. Vergne. Local euler–maclaurin formula for polytopes. *Moscow Mathematical Journal*, 7(3):355–386, 2007.
- [5] E. R. Canfield and B. D. McKay. The asymptotic volume of the birkhoff polytope. *arXiv preprint arXiv:0705.2422*, 2007.
- [6] F. Castillo and F. Liu. Berline-Vergne valuation and generalized permutohedra. pages 1–21, 2015.
- [7] F. Castillo and F. Liu. Berline-vergne valuation and generalized permutohedra. *arXiv preprint arXiv:1509.07884v1*, 2015.
- [8] F. Castillo and F. Liu. Deformation cones of nested braid fans. *arXiv preprint arXiv:1710.01899*, 2017.
- [9] S. Corteel, J. S. Kim, and K. Mészáros. Volumes of generalized chan–robbins–yuen polytopes. *Discrete & Computational Geometry*, pages 1–21, 2017.
- [10] V. I. Danilov. The geometry of toric varieties. *Russian Mathematical Surveys*, 33(2):97–154, 1978.
- [11] J. A. De Loera, F. Liu, and R. Yoshida. A generating function for all semi-magic squares and the volume of the birkhoff polytope. *Journal of Algebraic Combinatorics*, 30(1):113–139, 2009.
- [12] E. Ehrhart. Sur un probleme de géométrie diophantienne linéaire ii. *J. reine angew. Math*, 227(25):C49, 1967.

- [13] A. Garsia, J. Haglund, and G. Xin. Constant term methods in the theory of tesler matrices and macdonald polynomial operators. *Annals of Combinatorics*, 18(1):83–109, 2014.
- [14] E. Gorsky and A. Neguț. Refined knot invariants and hilbert schemes. *Journal de mathématiques pures et appliquées*, 104(3):403–435, 2015.
- [15] J. Haglund. A polynomial expression for the hilbert series of the quotient ring of diagonal coinvariants. *Advances in Mathematics*, 227(5):2092–2106, 2011.
- [16] J. Haglund, J. Remmel, and A. Wilson. The delta conjecture. *Transactions of the American Mathematical Society*, 370(6):4029–4057, 2018.
- [17] F. Liu. On positivity of ehrhart polynomials. In *Recent Trends in Algebraic Combinatorics*, pages 189–237. Springer, 2019.
- [18] P. McMullen. Valuations and dissections. handbook of convex geometry, vol. a, b, 933–988, 1993.
- [19] K. Mészáros, A. H. Morales, and B. Rhoades. The polytope of Tesler matrices. *Selecta Mathematica, New Series*, 23(1):425–454, 2017.
- [20] A. H. Morales. Ehrhart polynomials of examples of flow polytopes. <https://sites.google.com/site/flowpolytopes/ehrhart>.
- [21] I. Pak. Four questions on birkhoff polytope. *Annals of Combinatorics*, 4(1):83–90, 2000.
- [22] J. Pommersheim and H. Thomas. Cycles representing the todd class of a toric variety. *Journal of the American Mathematical Society*, 17(4):983–994, 2004.
- [23] A. Postnikov, V. Reiner, and L. Williams. Faces of generalized permutohedra. *Doc. Math*, 13(207-273):51, 2008.
- [24] M. H. Ring and A. Schürmann. Local formulas for ehrhart coefficients from lattice tiles. *Beiträge zur Algebra und Geometrie/Contributions to Algebra and Geometry*, pages 1–29, 2017.
- [25] A. T. Wilson. A weighted sum over generalized tesler matrices. *Journal of Algebraic Combinatorics*, 45(3):825–855, 2017.
- [26] D. Zeilberger. Proof of a conjecture of chan, robbins, and yuen. *Electron. Trans. Numer. Anal*, 9(147-148):1–2, 1999.
- [27] G. M. Ziegler. *Lectures on polytopes*, volume 152. Springer Science & Business Media, 2012.