

# New Bounds on $k$ -Planar Crossing Numbers

Alireza Shavali\*      Hamid Zarrabi-Zadeh†

November 18, 2019

## Abstract

The crossing number  $cr(G)$  of a graph  $G$  is the minimum number of crossings over all possible drawings of  $G$  in the plane. Analogously, the  $k$ -planar crossing number of  $G$ , denoted by  $cr_k(G)$ , is the minimum number of crossings over all possible drawings of the edges of  $G$  in  $k$  disjoint planes. We present new bounds on the  $k$ -planar crossing number of complete graphs and complete bipartite graphs. In particular, for the case of  $k = 2$ , we improve the current best lower bounds on biplanar crossing numbers by a factor of 1.37 for complete graphs, and by a factor of 1.34 for complete bipartite graphs. We extend our results to the  $k$ -planar crossing number of complete (bipartite) graphs, for any positive integer  $k \geq 2$ . To better understand the relation between crossing numbers and biplanar crossing numbers, we pose a new problem of finding the largest crossing number that implies biplanarity. In particular, we prove that for every graph  $G$ ,  $cr(G) \leq 10$  implies  $cr_2(G) = 0$ .

## 1 Introduction

An embedding (or drawing) of a graph  $G$  in the Euclidean plane is a mapping of the vertices of  $G$  to distinct points in the plane and a mapping of edges to smooth curves between their corresponding vertices. A planar embedding is a drawing of the graph such that no two edges cross each other, except for possibly in their endpoints. A graph that admits such a drawing is called planar. A *biplanar embedding* of a graph  $G = (V, E)$  is a decomposition of the graph into two graphs  $G_1 = (V, E_1)$  and  $G_2 = (V, E_2)$  such that  $E = E_1 \cup E_2$  and  $E_1 \cap E_2 = \emptyset$ , together with planar embeddings of  $G_1$  and  $G_2$ . In this case, we call  $G$  biplanar. Biplanar embeddings are central to the computation of thickness of graphs [13], with applications to VLSI design [14]. It is well-known that planarity can be recognized in linear time, while biplanarity testing is NP-complete [12].

Let  $cr(G)$  be the minimum number of edge crossings over all drawings of  $G$  in the plane, and let  $cr_k(G)$  be the minimum of  $cr(G_1) + cr(G_2) + \dots + cr(G_k)$  over all possible decompositions of  $G$  into  $k$  subgraphs  $G_1, G_2, \dots, G_k$ . We call  $cr(G)$  the *crossing number* of  $G$ , and  $cr_k(G)$  the  *$k$ -planar crossing number* of  $G$ . In this paper, we only consider *simple* drawings for each subgraph  $G_i$ , in which no two edges cross more than once, and no three edges cross

---

\*Department of Computer Engineering, Sharif University of Technology, Tehran 14588-89694, Iran. Email: [ashavali@ce.sharif.edu](mailto:ashavali@ce.sharif.edu).

†Department of Computer Engineering, Sharif University of Technology, Tehran 14588-89694, Iran. Email: [zarrabi@sharif.edu](mailto:zarrabi@sharif.edu).

at a point (such drawings are sometimes called nice drawings). Moreover, we denote by  $n$  the number of vertices, and by  $m$  the number of edges of a graph.

Determining the crossing number of complete graphs and complete bipartite graphs has been the subject of extensive research in graph drawing over the past few decades. In 1955, Zarankiewicz [20] conjectured that for all  $p$  and  $q$ ,  $cr(K_{p,q}) = \lfloor \frac{p}{2} \rfloor \lfloor \frac{p-1}{2} \rfloor \lfloor \frac{q}{2} \rfloor \lfloor \frac{q-1}{2} \rfloor$ . He also established a drawing with that many crossings. In 1960, Guy [9] conjectured that  $cr(K_n) = \frac{1}{4} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor$ . The problem is still open in both cases. In the biplanar case, even formulating such conjectures seems to be hard. As mentioned in [5], techniques like embedding method and bisection width method which are useful for bounding crossing numbers do not seem applicable to the biplanar case. The current best lower bounds for the biplanar crossing number of complete and complete bipartite graphs are due to Czaparka *et al.* [5]. These results are generalized to the  $k$ -planar crossing number in [18].

From Euler's formula for planar graphs, it is easy to see that for every simple graph  $G$ ,  $cr(G) \geq m - 3(n - 2)$ . Ajtai *et al.* [2] used this inequality and a probabilistic method to prove that for every simple graph  $G$  with  $m \geq 4n$ ,  $cr(G) \geq \frac{1}{64} \cdot \frac{m^3}{n^2}$ . This inequality is known as the *crossing lemma*. Pach *et al.* [15] improved the crossing lemma using the inequality  $cr(G) \geq 4m - \frac{103}{6}(n - 2)$ . The current best version of the lemma is due to Ackerman [1], which is based on the inequality  $cr(G) \geq 5m - \frac{139}{6}(n - 2)$ . These types of inequalities can be generalized to the biplanar and  $k$ -planar cases [5, 18].

One of the aims for studying biplanar crossing numbers is to get better understanding of crossing numbers. As such, the relation between crossing numbers and biplanar crossing numbers is important. In [6], Czaparka *et al.* proved that for every graph  $G$ ,

$$cr_2(G) \leq \frac{3}{8} cr(G). \quad (1)$$

They posed a question of finding the smallest constant  $c^*$  such that for every graph  $G$ , we have  $cr_2(G) \leq c^* \cdot cr(G)$ . They proved that  $\frac{3}{8} \geq c^* \geq \frac{64}{952}$ . Pach *et al.* [16] extended this inequality to the  $k$ -planar case.

In this paper, we present several new results for the  $k$ -planar crossing number of simple graphs. In particular, we provide improved lower bounds on the biplanar crossing number of complete graphs and complete bipartite graphs, using an iterative counting method. We extend our results to the  $k$ -planar crossing number of complete (bipartite) graphs, for any positive integer  $k$ . We also investigate the relation between  $cr(G)$  and  $cr_2(G)$  in more depth, and pose a new problem of finding the maximum crossing number that implies biplanarity. We prove that if  $cr(G) \leq 10$ , then  $G$  is biplanar, i.e.,  $cr_2(G) = 0$ . As a side product, we provide a biplanar embedding of  $K_{12}$  with 12 crossings, improving over the current bound of  $cr_2(K_{12}) \leq 14$ .

## 2 Lower Bounds for Complete Bipartite Graphs

In this section, we provide new lower bounds on the biplanar crossing number of complete bipartite graphs. In particular, we improve the following bound due to Czaparka *et al.* [5] which states that for all  $p, q \geq 10$ ,

$$cr_2(K_{p,q}) \geq \frac{p(p-1)q(q-1)}{290}.$$

We start by the following lemma, which is a main ingredient of our counting method.

**Lemma 1.** *Let  $\mathcal{G}$  be a hereditary class of graphs which is closed under removing edges. Let  $f$  be a linear function  $f(x) = cx$ , for some constant  $c$ , and let  $g$  be an arbitrary function. If for every graph  $G$  in  $\mathcal{G}$ ,  $cr(G) \geq f(m) + g(n)$ , then  $cr_k(G) \geq f(m) + k \cdot g(n)$  for all  $G \in \mathcal{G}$  and all  $k \geq 1$ .*

*Proof.* Fix a graph  $G$  in  $\mathcal{G}$ . Let  $G = G_1 \cup \dots \cup G_k$  be a decomposition of  $G$  into  $k$  subgraphs  $G_i = (V, E_i)$  such that  $\sum_{i=1}^k cr(G_i)$  is minimum. By the hereditary property of  $\mathcal{G}$ , each  $G_i$  is in  $\mathcal{G}$ , and hence  $cr(G_i) \geq f(m_i) + g(n)$ , where  $m_i = |E_i|$ . Thus,  $cr_k(G) = \sum_{i=1}^k cr(G_i) \geq \sum_{i=1}^k (f(m_i) + g(n)) = c \sum_{i=1}^k m_i + \sum_{i=1}^k g(n) = f(m) + k \cdot g(n)$ .  $\square$

From Euler's formula, we have  $cr(G) \geq m - 3(n - 2)$  for simple graphs, and  $cr(G) \geq m - 2(n - 2)$  for bipartite graphs. Using Lemma 1, we immediately get a lower bound of  $cr_2(G) \geq m - 6(n - 2)$  for simple graphs, and a lower bound of  $cr_2(G) \geq m - 4(n - 2)$  for bipartite graphs.

To establish stronger lower bounds, we need to incorporate more powerful ingredients. A graph is called  $k$ -planar, if it can be drawn in the plane in such a way that each edge has at most  $k$  crossings. It is known that every 1-planar drawing of any 1-planar graph has at most  $n - 2$  crossings [7]. Removing one edge per crossing yields a planar graph. Therefore, every 1-planar bipartite graph has at most  $3n - 6$  edges. Karpov [10] proved that for every 1-planar bipartite graph with at least 4 vertices, the inequality  $m \leq 3n - 8$  holds. In a recent work, Angelini *et al.* [3] proved that for every 2-planar bipartite graph we have  $m \leq 3.5n - 7$ . We use these results to obtain the following stronger lower bound.

**Lemma 2.** *For every bipartite graph  $G$  with  $n \geq 4$ ,*

$$cr_2(G) \geq 3m - 17n + 38.$$

*Proof.* Let  $G$  be a bipartite graph with  $n$  vertices and  $m$  edges. Fix a drawing of  $G$  with a minimum number of crossings. If  $m > 3.5n - 7$ , then by [3], there must be an edge in the drawing with at least three crossings. We repeatedly remove such an edge until we reach a drawing with  $\lfloor 3.5n - 7 \rfloor$  edges. Then by Karpov's result there must be an edge in the drawing with at least two crossings. Similarly we repeatedly remove such an edge until we reach a drawing with  $3n - 8$  edges. Let  $G'$  be the bipartite graph corresponding to the remaining drawing. Now,

$$\begin{aligned} cr(G) &\geq 3(m - \lfloor 3.5n - 7 \rfloor) + 2(\lfloor 3.5n - 7 \rfloor - (3n - 8)) + cr(G') \\ &\geq 3(m - \lfloor 3.5n - 7 \rfloor) + 2(\lfloor 3.5n - 7 \rfloor - (3n - 8)) + (3n - 8) - 2(n - 2) \\ &\geq 3m - \lfloor 3.5n - 7 \rfloor - (3n - 8) - 2(n - 2) \geq 3m - 8.5n + 19. \end{aligned}$$

Applying Lemma 1 yields  $cr_2(G) \geq 3m - 17n + 38$ .  $\square$

The following is a direct corollary of Lemma 2.

**Corollary 3.** *For all  $p, q \geq 2$ ,*

$$cr_2(K_{p,q}) \geq 3pq - 17(p + q) + 38.$$

We can further improve the lower bound obtained by Corollary 3 using an iterated version of a *counting method* [5, 17] described below. Let  $D$  be a biplanar drawing of  $G$  realizing  $cr_2(G)$ . If  $G$  contains  $\alpha$  copies of a graph  $H$ , and each crossing in  $D$  belongs to at most  $\beta$  copies of  $H$ , then

$$cr_2(G) \geq \left\lceil \frac{\alpha}{\beta} cr_2(H) \right\rceil.$$

Note that we have put a ceiling in the above inequality, because  $cr_2(G)$  is an integer.

**Theorem 4.** *For all  $p, q \geq 21$ ,*

$$cr_2(K_{p,q}) \geq \frac{p(p-1)q(q-1)}{216}.$$

*Proof.* Using the counting method for  $K_{n,n}$  and  $K_{n+1,n}$  we have

$$cr_2(K_{n+1,n}) \geq \left\lceil \frac{n+1}{n-1} cr_2(K_{n,n}) \right\rceil.$$

This is because  $K_{n+1,n}$  contains  $n+1$  copies of  $K_{n,n}$ , and each crossing, realized by two edges, belongs to at most  $\binom{n-1}{n-2} = n-1$  of these copies. Using a similar argument for  $K_{n+1,n}$  and  $K_{n+1,n+1}$ , we get

$$cr_2(K_{n+1,n+1}) \geq \left\lceil \frac{n+1}{n-1} \left\lceil \frac{n+1}{n-1} cr_2(K_{n,n}) \right\rceil \right\rceil. \quad (2)$$

By Corollary 3,  $cr_2(K_{15,15}) \geq 203$ . Plugging into (2), yields  $cr_2(K_{16,16}) \geq 266$ . (Note that Corollary 3 alone could only imply  $cr_2(K_{16,16}) \geq 262$ .) Now, we use the recurrence relation (2) iteratively from  $n = 15$  to 21 to get

$$cr_2(K_{21,21}) \geq 817. \quad (3)$$

We can now apply the counting method on  $K_{21,21}$  and  $K_{p,q}$  to obtain

$$cr_2(K_{p,q}) \geq \frac{\binom{p}{21} \binom{q}{21} cr_2(K_{21,21})}{\binom{p-2}{19} \binom{q-2}{19}} = \frac{p(p-1)q(q-1)cr_2(K_{21,21})}{21 \times 20 \times 21 \times 20}.$$

Replacing (3) in the above inequality implies the theorem.  $\square$

**Remark.** The exact value of the denominator obtained in our proof is around 215.911. One may continue applying the recurrence relation (2) to obtain better bounds for  $K_{n,n}$  when  $n > 21$ . This leads to a slightly improved constant in the denominator, but it does not seem to reduce the constant below 215. Indeed, the denominator seems to converge to a value around 215.131.

### 3 Improved Bounds for Complete Graphs

We now consider the biplanar crossing number of complete graphs. We start with an improved upper bound on  $cr_2(K_{12})$ , and then provide an improved lower bound for  $cr_2(K_n)$  in general.

### 3.1 A Biplanar Embedding of $K_{12}$

In 1971, Owens [14] gave a construction for a biplanar embedding of  $K_n$ . In particular, he showed that  $cr_2(K_{12}) \leq 18$ . Recently, Durocher *et al.* [8] presented a biplanar drawing of  $K_{11}$  with 6 crossings, and used this drawing to show that  $cr_2(K_{12}) \leq 14$ . We further improve this upper bound by showing that  $cr_2(K_{12}) \leq 12$ . The improved biplanar embedding is illustrated in Figure 1.




Figure 1: A biplanar embedding of  $K_{12}$  with 12 crossings.

The improved embedding is obtained by a careful investigation of possible partitionings of the edges of  $K_{12}$  into two planes, based on an optimal biplanar embeddings of  $K_{10}$  with two crossings. The OGDF library [4] is used to produce the final drawing. Combined with a lower bound presented in [6], we conclude that  $6 \leq cr_2(K_{12}) \leq 12$ .

### 3.2 Improved Lower Bound for $cr_2(K_n)$

Czabarka *et al.* [5] used a probabilistic method to prove that for large values of  $n$ ,

$$cr_2(K_n) \geq \frac{n^4}{952}.$$

We improve this lower bound using our counting method.

**Theorem 5.** *For all  $n \geq 24$*

$$cr_2(K_n) \geq \frac{n(n-1)(n-2)(n-3)}{698}.$$

*Proof.* We know from [1] that for every  $G$  with  $n \geq 3$ ,  $cr(G) \geq 5m - \frac{139}{6}(n-2)$ . Applying Lemma 1, we get

$$cr_2(G) \geq 5m - \frac{139}{3}(n-2).$$

This in particular implies that  $cr_2(K_{25}) \geq 435$ . Now we use a counting method for  $K_{25}$  and  $K_n$  to get

$$cr_2(K_n) \geq \frac{\binom{n}{25} cr_2(K_{25})}{\binom{n-4}{21}} \geq \frac{n(n-1)(n-2)(n-3)}{\frac{25 \times 24 \times 23 \times 22}{435}},$$

which implies the theorem statement.  $\square$

We can slightly improve this result, using an iterative counting method similar to what we used in the previous section.

**Theorem 6.** *For large values of  $n$ ,*

$$cr_2(K_n) \geq \frac{n^4}{694}.$$

*Proof.* Using the counting method for  $K_n$  and  $K_{n+1}$  we have,

$$cr_2(K_{n+1}) \geq \left\lceil \frac{(n+1)cr_2(K_n)}{n-3} \right\rceil. \quad (4)$$

Starting from  $cr_2(K_{25}) \geq 435$ , we use the recurrence relation (4) iteratively from  $n = 25$  to 57 to obtain  $cr_2(K_{57}) \geq 13667$ . Now, we use the counting method for  $K_{57}$  and  $K_n$  to get

$$cr_2(K_n) \geq \frac{\binom{n}{57} cr_2(K_{57})}{\binom{n-4}{53}} \geq \frac{n(n-1)(n-2)(n-3)}{\frac{57 \times 56 \times 55 \times 54}{13667}} \geq \frac{n(n-1)(n-2)(n-3)}{693.9},$$

which implies  $cr_2(K_n) \geq \frac{n^4}{694}$  for  $n$  sufficiently large.  $\square$

## 4 The Maximum Crossing Number that Implies Biplanarity

Czabarka *et al.* [6] defined  $c^*$  to be the smallest constant such that for every graph  $G$ ,  $cr_2(G) \leq c^* \cdot cr(G)$ . They proved that  $0.067 \leq c^* \leq \frac{3}{8} = 0.375$ . It is known that  $cr(K_n) \leq \frac{n^4}{64}$  [19]. By Theorem 6, for  $n$  sufficiently large,  $cr_2(K_n) \geq \frac{n^4}{694}$ . Therefore, our results from Section 3 imply an improved bound of  $c^* \geq \frac{64}{694} \approx 0.092$ . In a more general sense, we are interested in the following problem.

**Problem.** Given a positive integer  $r$ , find the largest integer  $\xi(r)$  such that for every graph  $G$ ,  $cr(G) \leq \xi(r)$  implies  $cr_2(G) \leq r$ .

For the special case of  $r = 0$ , the problem is to find the largest integer  $\xi$  such that drawing a graph with  $\xi$  crossings in the plane guarantees that the graph is biplanar. As noted in [5],  $cr_2(K_9) = 1$ , and hence,  $K_9$  is not biplanar. Moreover, we know that  $cr(K_9) = 36$  [11]. Therefore,  $\xi(0) < 36$ .

Inequality (1) implies that if  $cr(G) \leq 2$ , then  $G$  is biplanar. Therefore,  $\xi(0) \geq 2$ . We can strengthen this bound as follows. Recall that by Kuratowski's theorem, every nonplanar graph contains a subdivision of  $K_{3,3}$  or  $K_5$ . Therefore, there is no nonplanar graph with less than 9 edges. This leads to the following observation.

**Observation 1.** *Every graph with at most 8 edges is planar. The only nonplanar graph with 9 edges is  $K_{3,3}$ , and the only nonplanar graphs with 10 edges are  $K_5$ ,  $K_{3,3}$  with an extra edge, and  $K_{3,3}$  with a subdivided edge.*

From this simple observation, we can infer that  $\xi(0) \geq 4$  as follows. Suppose a graph  $G$  is drawn in the plane with at most 4 crossings. The number of edges involved in these four crossings is at most 8. If we remove these 8 edges from the drawing, the remaining drawing has no crossing. Moreover, the subgraph of  $G$  that contains only these 8 (or fewer) edges is planar by Observation 1. Therefore,  $G$  is the union of two planar graphs, and hence is biplanar. We will significantly improve this lower bound in the following theorem.

**Theorem 7.** *Every graph  $G$  with  $cr(G) \leq 10$  is biplanar. In other words,  $\xi(0) \geq 10$ .*

*Proof.* Let  $G$  be a graph with  $cr(G) \leq 10$ . Fix a drawing of  $G$  with a minimum number of crossings. We repeatedly remove an edge from the drawing that involves in a maximum number of crossings until there remains no more crossings. Let  $G_1$  be the graph corresponding to the remaining drawing, and  $G_2$  be the graph formed by the removed edges. Clearly,  $G_2$  has at most 10 edges. Moreover,  $G_1$  is planar by construction. If  $G_2$  has 8 or less edges, then it is planar by Observation 1, and we are done. Otherwise,  $G_2$  has 9 or 10 edges. Note that removing any of these edges from  $G$  has removed at least one crossing. Therefore, removing any of these edges, except possibly the first one, has removed exactly one crossing from  $G$ . By Observation 1, if  $G_2$  is not planar, then it is either  $K_5$ ,  $K_{3,3}$ ,  $K_{3,3}$  with a subdivided edge, or  $K_{3,3}$  with an extra edge. In the former two cases, let  $e$  be the last edge removed from  $G$ . Clearly,  $e$  was crossing exactly one edge  $f$  in  $G_1$  just before removal. Therefore, switching  $e$  and  $f$  between  $G_1$  and  $G_2$  keeps  $G_1$  planar. Moreover, the new  $G_2$  is planar, because it contains no subdivision of  $K_5$  and  $K_{3,3}$ . Hence,  $G$  is biplanar in the first two cases. In the latter two cases, i.e., when  $G_2$  is a  $K_{3,3}$  with a subdivided edge or a  $K_{3,3}$  with an extra edge,  $G_2$  has exactly 10 edges. Therefore, removing any of these edges from  $G$  has removed exactly one crossing, which means that any edge in  $G$  is crossing at most one edge. If  $G_2$  is a  $K_{3,3}$  with a subdivided edge, let  $e$  be any edge of  $G_2$  except the two edges forming the subdivided edge, and if  $G_2$  is a  $K_{3,3}$  with an extra edge, let  $e$  be any edge of  $G_2$  except this extra edge. We know that  $e$  was crossing exactly one edge  $f$  in  $G$ . Moreover,  $f$  was only crossing  $e$  in  $G$ , and hence, it remains in  $G_1$  after removing  $e$ . Similar to the previous case, switching  $e$  and  $f$  between  $G_1$  and  $G_2$  completes the proof.  $\square$

## 5 The $k$ -Planar Crossing Number of $K_n$ and $K_{p,q}$

In this section, we provide improved lower bounds on the  $k$ -planar crossing number of complete graphs and complete bipartite graphs. Shahrokhi *et al.* [18] proved that for any positive integer  $k$ , and sufficiently large integers  $p$ ,  $q$ , and  $n$ :

$$cr_k(K_{p,q}) \geq \frac{p(p-1)q(q-1)}{108k^2},$$

and

$$cr_k(K_n) \geq \frac{n(n-1)(n-2)(n-3)}{432k^2}.$$

We improve these results using the ideas developed in Sections 2 and 3.

**Theorem 8.** *For all  $p, q \geq 8k + 2$ ,*

$$cr_k(K_{p,q}) \geq \frac{p(p-1)q(q-1)}{\frac{512}{7}k^2}.$$

*Proof.* We use the counting method for  $K_{8k+2,8k+2}$  and  $K_{p,q}$ . As noted in the proof of Lemma 2, for every bipartite graph  $G$ ,  $cr(G) \geq 3m - 8.5n + 19$ . Therefore, by Lemma 1,  $cr_k(G) \geq 3m - (8.5n - 19)k$ . This yields

$$cr_k(K_{8k+2,8k+2}) \geq 56k^2 + 43k + 12.$$

Hence,

$$\begin{aligned} cr_k(K_{p,q}) &\geq \frac{\binom{p}{8k+2} \binom{q}{8k+2} cr_k(K_{8k+2,8k+2})}{\binom{p-2}{8k} \binom{q-2}{8k}} = \frac{p(p-1)q(q-1)cr_k(K_{8k+2,8k+2})}{(8k+2)(8k+1)(8k+2)(8k+1)} \\ &\geq \frac{p(p-1)q(q-1)}{\frac{(8k+2)^2(8k+1)^2}{56k^2+43k+12}} \geq \frac{p(p-1)q(q-1)}{\frac{512}{7}k^2}, \end{aligned}$$

which completes the proof.  $\square$

**Theorem 9.** For all  $n \geq 14k - 3$ ,

$$cr_k(K_n) \geq \frac{n(n-1)(n-2)(n-3)}{232k^2}.$$

*Proof.* We use the counting method for  $K_{14k-3}$  and  $K_n$ . Recall that for every  $G$  with  $n \geq 3$ ,  $cr(G) \geq 5m - \frac{139}{6}(n-2)$  [1]. Therefore,  $cr_k(G) \geq 5m - \frac{139}{6}(n-2)k$  by Lemma 1. Thus,

$$cr_k(K_{14k-3}) \geq \frac{497}{3}k^2 - \frac{775}{6}k + 30.$$

Therefore,

$$cr_k(K_n) \geq \frac{\binom{n}{14k-3} cr_k(K_{14k-3})}{\binom{n-4}{14k-7}} = \frac{n(n-1)(n-2)(n-3)cr_k(K_{14k-3})}{(14k-3)(14k-4)(14k-5)(14k-6)},$$

which implies the theorem.  $\square$

## 6 Conclusion

In this paper, we presented improved lower bounds on the  $k$ -planar crossing number of complete graphs and complete bipartite graphs. For the case of  $k = 2$ , we used an iterative counting method to obtain further improved bounds. Similar improvements seem possible for other values of  $k > 2$ , using the same technique. While the exact value of  $cr_2(K_{10})$  is known to be 2, the only known bounds for  $K_{11}$  and  $K_{12}$  are  $4 \leq cr_2(K_{11}) \leq 6$  and  $6 \leq cr_2(K_{12}) \leq 12$  due to [8] and this work. Determining the exact values of  $cr_2(K_{11})$  and  $cr_2(K_{12})$  would be interesting. We also posed an open problem of finding the largest positive integer  $\xi(r)$  such that  $cr(G) \leq \xi(r)$  implies  $cr_2(G) \leq r$ . In particular, we proved that  $10 \leq \xi(0) \leq 35$ . This definition can be easily generalized to the  $k$ -planar case: given positive integers  $k$  and  $r$ , find the largest integer  $\xi_k(r)$  such that  $cr(G) \leq \xi_k(r)$  implies  $cr_k(G) \leq r$ . Determining the value of  $\xi_k(r)$  is an intriguing problem, even for the special case of  $r = 0$ .

## References

- [1] E. Ackerman. On topological graphs with at most four crossings per edge. *Computational Geometry*, 85:1–37, 2019.
- [2] M. Ajtai, V. Chvátal, M. M. Newborn, and E. Szemerédi. Crossing-free subgraphs. *North-Holland Mathematics Studies*, 60(C):9–12, 1982.



- [3] P. Angelini, M. A. Bekos, M. Kaufmann, M. Pfister, and T. Ueckerdt. Beyond-planarity: Turán-type results for non-planar bipartite graphs. In *Proceedings of the 29th International Symposium on Algorithms and Computation*, 2018.
- [4] M. Chimani, C. Gutwenger, M. Jünger, G. W. Klau, K. Klein, and P. Mutzel. The open graph drawing framework (OGDF). *Handbook of Graph Drawing and Visualization*, 2011:543–569, 2013.
- [5] É. Czabarka, O. Sýkora, L. A. Székely, and I. Vrřto. Biplanar crossing numbers I: A survey of results and problems. In *More sets, graphs and numbers*, pages 57–77. 2006.
- [6] É. Czabarka, O. Sýkora, L. A. Székely, and I. Vrřto. Biplanar crossing numbers II: Comparing crossing numbers and biplanar crossing numbers using the probabilistic method. *Random Structures & Algorithms*, 33(4):480–496, 2008.
- [7] J. Czap and D. Hudák. On drawings and decompositions of 1-planar graphs. *The electronic journal of combinatorics*, 20(2):54, 2013.
- [8] S. Durocher, E. Gethner, and D. Mondal. On the biplanar crossing number of  $K_n$ . In *Proceedings of the 28th Canadian Conference on Computational Geometry*, pages 93–100, 2016.
- [9] R. K. Guy. A combinatorial problem. *Nabla (Bulletin of the Malayan Mathematical Society)*, 7:68–72, 1960.
- [10] D. Karpov. An upper bound on the number of edges in an almost planar bipartite graph. *Journal of Mathematical Sciences*, 196(6):737–746, 2014.
- [11] A. Liebers. *Methods for Planarizing Graphs: A Survey and Annotated Bibliography*. PhD thesis, 1996.
- [12] A. Mansfield. Determining the thickness of graphs is NP-hard. *Mathematical Proceedings of the Cambridge Philosophical Society*, 93(1):9–23, 1983.
- [13] P. Mutzel, T. Odenthal, and M. Scharbrodt. The thickness of graphs: a survey. *Graphs and Combinatorics*, 14(1):59–73, 1998.
- [14] A. Owens. On the biplanar crossing number. *IEEE Transactions on Circuit Theory*, 18(2):277–280, 1971.
- [15] J. Pach, R. Radoicic, G. Tardos, and G. Tóth. Improving the crossing lemma by finding more crossings in sparse graphs. *Discrete and Computational Geometry*, 36(4):527–552, 2006.
- [16] J. Pach, L. A. Székely, C. D. Tóth, and G. Tóth. Note on  $k$ -planar crossing numbers. *Computational Geometry*, 68:2–6, 2018.
- [17] F. Shahrokhi, O. Sýkora, L. A. Székely, and I. Vrřto. Crossing numbers: bounds and applications. *Intuitive geometry*, 6:179–206, 1995.

- [18] F. Shahrokhi, O. Sýkora, L. A. Székely, and I. Vrto. On  $k$ -planar crossing numbers. *Discrete Applied Mathematics*, 155(9):1106–1115, 2007.
- [19] A. T. White and L. W. Beineke. Topological graph theory. *Selected Topics in Graph Theory*, 1:15–49, 1978.
- [20] C. Zarankiewicz. On a problem of P. Turán concerning graphs. *Fundamenta Mathematicae*, 41(1):137–145, 1955.