PARTIAL SEPARABILITY/ENTANGLEMENT VIOLATES DISTRIBUTIVE RULES

KYUNG HOON HAN, SEUNG-HYEOK KYE AND SZILÁRD SZALAY

ABSTRACT. We found three qubit Greenberger-Horne-Zeilinger diagonal states which tell us that the partial separability of three qubit states violates the distributive rules with respect to the two operations of convex sum and intersection. The gaps between the convex sets involving the distributive rules are of nonzero volume.

1. INTRODUCTION

Pure states in classical probability theory are uncorrelated, which is not the case in quantum probability theory, where this nonclassical form of correlation is called entanglement [1]. Beyond the conceptual questions it raises [2, 3], entanglement plays a key role in the physics of strongly correlated many-body systems [4], and also finds direct applications in quantum information theory [5]. Mixed states in classical probability theory arise as statistical mixtures (convex combinations) of pure, hence uncorrelated states. Again, this is not the case in quantum probability theory, and states which are mixtures of uncorrelated states are called separable, while the others are entangled [6].

In the case of multipartite systems, the partitions of the total system into subsystems give rise to various notions of partial separability. In the tripartite case with the elementary subsystems A, B and C, we have three nontrivial partitions A-BC, B-CA and C-AB, and the corresponding partial separability properties are called A-BC-separability, B-CA-separability and C-AB-separability, respectively. We call these basic biseparabilities.

It is natural to consider the intersections (also called partial separability classes) and convex hulls of the three convex sets consisting of the above three kinds of basic biseparable states. For example, the intersection of them [7], the intersections of two of them [9, 8] and the convex hull of them [11, 10] have been considered. More recently, the convex hulls of two of them have also been considered together with intersections and complements of convex sets arising in the way [12, 13, 14], leading to the description of the hierarchy of the intersections [15]. See also [16, 17, 18] for further developments. Recall that the intersection and convex hull of the convex sets of three basic biseparable states give rise to *fully biseparable* and *biseparable* states, respectively. Tripartite states which are not biseparable are called genuinely multipartite entangled.

¹⁹⁹¹ Mathematics Subject Classification. 81P15, 15A30, 46L05, 46L07.

Key words and phrases. partially separable, partially entangled, lattice, distributive rule.

In this context, we consider the lattice generated by three convex sets of three qubit basic biseparable states with respect to the above mentioned two operations, intersection and convex hull. In this way, we deal with convex hulls of intersections, as well as intersections of convex hulls of convex sets arising from basic biseparability. One may go further to investigate the whole structures of partial separability and partial entanglement. Due to technical reasons, we will consider the three convex cones α , β and γ of all *un-normalized A-BC*, *B-CA* and *C-AB* biseparable three qubit states, respectively. We note that the convex hull $\sigma \lor \tau$ of two convex cones σ and τ will be denoted by $\sigma \land \tau$ following the lattice notations. In general, the two operations \land and \lor among convex sets obey associative rule and commutative rule. Furthermore, they also satisfy the following relations

$$\sigma \lor \sigma = \sigma, \qquad \sigma \land \sigma = \sigma,$$

$$\sigma \lor \tau) \land \sigma = \sigma, \qquad (\sigma \land \tau) \lor \sigma = \sigma,$$

and so they give rise to a lattice.

A lattice $(\mathcal{L}, \lor, \land)$ is distributive if the identities

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z), \qquad x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$$

hold for all $x, y, z \in \mathcal{L}$. In general, the inequalities

$$x \land (y \lor z) \ge (x \land y) \lor (x \land z), \qquad x \lor (y \land z) \le (x \lor y) \land (x \lor z)$$

always hold trivially. We denote by \mathcal{L} the lattice generated by three convex cones α , β and γ . Therefore, \mathcal{L} is the smallest lattice containing the convex cones α , β and γ in the 64-dimensional real vector space of all self-adjoint three qubit matrices. The purpose of this note is to show that the lattice \mathcal{L} does not satisfy the distributive rules. More precisely, we show that both inequalities

(1)
$$(\alpha \land \beta) \lor (\alpha \land \gamma) \le \alpha \land (\beta \lor \gamma),$$

(2)
$$\beta \lor (\gamma \land \alpha) \le (\beta \lor \gamma) \land (\beta \lor \alpha)$$

are strict. Furthermore, the gaps between the two sets are of nonzero volume, in both cases. We also show that the lattice \mathcal{L} does not satisfy the modularity which is weaker than distributivity.

We note that a state ρ in the gap $\alpha \wedge (\beta \vee \gamma) \setminus (\alpha \wedge \beta) \vee (\alpha \wedge \gamma)$ by the strict inequality in (1) has the following properties:

- ρ is A-BC biseparable, and it is a mixture of B-CA and C-AB biseparable states,
- but, it is not a mixture of a simultaneously A-BC and B-CA biseparable state and a simultaneously A-BC and C-AB biseparable state.

We will find such states among GHZ diagonal states. On the other hand, a state ρ arising by the strict inequality in (2) has the following properties:

- ρ is a mixture of *B*-*CA* and *C*-*AB* biseparable states, and it is also a mixture of a *B*-*CA* and *A*-*BC* biseparable states,
- but it is not a mixture of B-CA biseparable state and a simultaneously C-AB and A-BC biseparable state.

Examples will be found among GHZ diagonal states. After we provide backgrounds for this in the next section, we find analytic examples in Section 3. We also consider in Section 4 the lattices arising from partial separability in general multi-partite systems to see that both distributivity and modularity are violated in them, too. We close the paper to ask several questions.

The authors are grateful to the referee for bringing our attention to the general multi-partite cases.

2. X-SHAPED STATES

We will find required examples among so called X-shaped states whose entries are zeros except for diagonal and anti-diagonal entries by definition. A self-adjoint Xshaped three qubit matrix is of the form

$$\mathsf{X}(a,b,z) = \begin{pmatrix} a_1 & & & & z_1 \\ & a_2 & & & z_2 & \\ & & a_3 & & z_3 & & \\ & & & a_4 & z_4 & & \\ & & & \bar{z}_4 & b_4 & & \\ & & & \bar{z}_3 & & b_3 & & \\ & & & \bar{z}_2 & & & b_2 & \\ & & & & & & b_1 \end{pmatrix}$$

with $a, b \in \mathbb{R}^4$ and $z \in \mathbb{C}^4$. Here, $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ is identified with the vector space \mathbb{C}^8 with respect to the lexicographic order of indices. Note that X(a, b, z) is a state if and only if $a_i, b_i \geq 0$ and $\sqrt{a_i b_i} \geq |z_i|$ for each i = 1, 2, 3, 4.

We recall that every GHZ diagonal states [19] are in this form, and an X-state X(a, b, z) is GHZ diagonal if and only if a = b and $z \in \mathbb{R}^4$. In this case, we use the notation

$$\mathsf{X}\begin{pmatrix}a\\z\end{pmatrix} = \mathsf{X}(a,a,z).$$

By a pair $\{i, j\}$, we will mean an unordered set with two distinct elements for simplicity throughout this paper. For a given three qubit X-shaped state $\rho = X(a, b, z)$, we consider the inequalities

$S_1[i,j]:$	$\min\{\sqrt{a_i b_i}, \sqrt{a_j b_j}\} \ge \max\{ z_i , z_j \},\$
$S_2[i,j]$:	$\min\left\{\sqrt{a_i b_i} + \sqrt{a_j b_j}, \sqrt{a_k b_k} + \sqrt{a_\ell b_\ell}\right\} \ge \max\left\{ z_i + z_j , z_k + z_\ell \right\},\$
S_3 :	$\sum_{j \neq i} \sqrt{a_j b_j} \ge z_i , i = 1, 2, 3, 4$

for a pair $\{i, j\}$, where $\{k, \ell\}$ is chosen so that $\{i, j, k, l\} = \{1, 2, 3, 4\}$. By [22, Proposition 5.2], we have the following:

- $\rho \in \alpha$ if and only if $S_1[1, 4]$ and $S_1[2, 3]$ hold,
- $\rho \in \beta$ if and only if $S_1[1,3]$ and $S_1[2,4]$ hold,
- $\rho \in \gamma$ if and only if $S_1[1, 2]$ and $S_1[3, 4]$ hold.

We also have the following

- $\rho \in \beta \vee \gamma$ if and only if $S_2[1, 4]$ (equivalently $S_2[2, 3]$) holds,
- $\rho \in \gamma \lor \alpha$ if and only if $S_2[1,3]$ (equivalently $S_2[2,4]$) holds,
- $\rho \in \alpha \lor \beta$ if and only if $S_2[1,2]$ (equivalently $S_2[3,4]$) holds.

by [14, Theorem 5.5]. The inequality S_3 will not be used in this paper, but it is the characteristic inequality for the convex cone $\alpha \lor \beta \lor \gamma$ [20, 22, 14, 21].

We will consider the above inequality S_2 for arbitrary two pairs $\{i, j\}$ and $\{k, \ell\}$ as follows:

$$S_4[i,j|k,\ell]: \min\left\{\sqrt{a_i b_i} + \sqrt{a_j b_j}, \sqrt{a_k b_k} + \sqrt{a_\ell b_\ell}\right\} \ge \max\left\{|z_i| + |z_j|, |z_k| + |z_\ell|\right\}.$$

If $\{i, j\} = \{k, \ell\}$ then the inequality $S_4[i, j|k, \ell]$ holds automatically for any X-states $\varrho = X(a, b, z)$. If $\{i, j\} \cap \{k, \ell\} = \emptyset$ then the three inequalities $S_2[i, j]$, $S_2[k, \ell]$ and $S_4[i, j|k, \ell]$ are identical. In the other cases, the resulting inequalities are new ones.

In order to get required examples in the gaps between convex cones in the inequalities (1) and (2), we proceed to characterize the following convex cones

(3)
$$\alpha \lor (\beta \land \gamma), \qquad \beta \lor (\gamma \land \alpha), \qquad \gamma \lor (\alpha \land \beta).$$

For this purpose, we will use the duality among convex cones in a real vector space with a bi-linear pairing \langle , \rangle . For a convex cone C, the dual cone C° is defined by

 $C^{\circ} = \{ x \in V : \langle x, y \rangle \ge 0 \text{ for each } y \in C \}.$

We are now working in the real vector spaces of all three qubit self-adjoint X-shaped matrices, where the bi-linear pairing is defined by $\langle x, y \rangle = \text{Tr}(yx^{t})$, as usual. See [14]. Every closed convex cone C satisfies the relation $C = (C^{\circ})^{\circ}$, which tells us that $x \in C$ if and only if $\langle x, y \rangle \geq 0$ for every $y \in C^{\circ}$.

The dual cones of the cones in (3) have also been characterized in [14]. For a given X-shaped self-adjoint matrix W = X(s, t, u) with $s_i, t_i \ge 0$ and $u \in \mathbb{C}^4$, we have considered the inequalities W_1, W_2, W_3 given by

$$\begin{array}{ll} W_1[i,j]: & \sqrt{s_i t_i} + \sqrt{s_j t_j} \ge |u_i| + |u_j|, \\ W_2[i,j]: & \sum_{k \neq j} \sqrt{s_k t_k} \ge |u_i|, & \sum_{k \neq i} \sqrt{s_k t_k} \ge |u_j|, \\ W_3: & \sum_{i=1}^4 \sqrt{s_i t_i} \ge \sum_{i=1}^4 |u_i|, \end{array}$$

for a pair $\{i, j\}$. Then we have the following by [14, Proposition 3.3]:

- $W \in \alpha^{\circ}$ if and only if $W_1[1, 4]$ and $W_1[2, 3]$ hold,
- $W \in \beta^{\circ}$ if and only if $W_1[1,3]$ and $W_1[2,4]$ hold,
- $W \in \gamma^{\circ}$ if and only if $W_1[1, 2]$ and $W_1[3, 4]$ hold.

On the other hand, we also have the following by [14, Theorem 5.2]:

- $W \in \beta^{\circ} \vee \gamma^{\circ}$ if and only if $W_2[1, 4], W_2[2, 3]$ and W_3 hold,
- $W \in \gamma^{\circ} \lor \alpha^{\circ}$ if and only if $W_2[1,3], W_2[2,4]$ and W_3 hold,
- $W \in \alpha^{\circ} \vee \beta^{\circ}$ if and only if $W_2[1, 2], W_2[3, 4]$ and W_3 hold.

For given $a, b \in \mathbb{R}^4$ with nonzero entries, and $z \in \mathbb{C}^4$ with $\arg z_i = \theta_i$, we consider the following self-adjoint matrices:

$$W_{[i,j|k,\ell]} = \mathsf{X}\left(\sqrt{\frac{b_i}{a_i}}E_i + \sqrt{\frac{b_j}{a_j}}E_j, \sqrt{\frac{a_i}{b_i}}E_i + \sqrt{\frac{a_j}{b_j}}E_j, -e^{-\mathrm{i}\theta_k}E_k - e^{-\mathrm{i}\theta_\ell}E_l\right)$$

for pairs $\{i, j\}$ and $\{k, \ell\}$, where $\{E_i\}$ is the usual orthonormal basis of \mathbb{C}^4 . Then the inequality

$$\langle W_{[i,j|k,\ell]}, \mathsf{X}(a,b,z) \rangle \ge 0$$

gives rise to the inequality $\sqrt{a_i b_i} + \sqrt{a_j b_j} \ge |z_k| + |z_\ell|$. Now, we present the main result. By [14, Proposition 2.2], the conditions give rise to necessary criteria for general states to belong to classes in (3) in terms of diagonal and anti-diagonal entries.

Theorem 2.1. For a given three qubit X-state $\rho = X(a, b, z)$, we have the following:

- (i) $\varrho \in \alpha \lor (\beta \land \gamma)$ if and only if $S_4[i, j|k, \ell]$ holds whenever $\{i, j\}, \{k, \ell\}$ are two of $\{1, 2\}$, $\{1, 3\}$, $\{2, 4\}$, $\{3, 4\}$;
- (ii) $\varrho \in \beta \lor (\gamma \land \alpha)$ if and only if $S_4[i, j|k, \ell]$ holds whenever $\{i, j\}, \{k, \ell\}$ are two of $\{1, 2\}$, $\{1, 4\}$, $\{2, 3\}$, $\{3, 4\}$;
- (iii) $\rho \in \gamma \lor (\alpha \land \beta)$ if and only if $S_4[i, j|k, \ell]$ holds whenever $\{i, j\}, \{k, \ell\}$ are two of $\{1, 3\}$, $\{1, 4\}$, $\{2, 3\}$, $\{2, 4\}$.

Proof. We will prove (i). For the 'only if' part, we first consider the case $a_i, b_i > 0$ for i = 1, 2, 3, 4. Then we see that $W_{[i,j|k,\ell]}$ belongs to $\alpha^{\circ} \wedge (\beta^{\circ} \vee \gamma^{\circ})$ whenever $\{i, j\}$, $\{k, \ell\}$ are two of $\{1, 2\}, \{1, 3\}, \{2, 4\}, \{3, 4\}$, by checking the required inequalities $W_1[1,4], W_1[2,3], W_2[1,4], W_2[2,3]$ and W_3 . Now, the inequality $\langle W_{[i,j|k,\ell]}, \varrho \rangle \geq 0$ gives rise to $S_4[i, j|k, \ell]$ for such $\{i, j\}, \{k, \ell\}$. If a or b has a zero entry, then we consider $\rho + \varepsilon X(1, 1, 0)$ with arbitrary $\varepsilon > 0$ and take $\varepsilon \to 0$ for the conclusion, where $\mathbf{1} = (1, 1, 1, 1)$ and $\mathbf{0} = (0, 0, 0, 0)$.

For the converse, it suffices to show the following:

• if $\rho = X(a, b, z)$ satisfies $S_4[i, j|k, \ell]$ whenever $\{i, j\}, \{k, l\}$ are two of $\{1, 2\}, \{k, l\}$ $\{1,3\}, \{2,4\}, \{3,4\}, \text{ and } W = X(s,t,u) \text{ satisfies } W_1[1,4], W_1[2,3], W_2[1,4],$ $W_2[2,3]$ and W_3 , then $\langle W, \varrho \rangle \ge 0$,

by Proposition 2.2 and Corollary 2.3 of [14]. If $\rho \in \alpha$ then there is nothing to prove, and so we may assume that $\rho \notin \alpha$. Without loss of generality, we may also assume that $|z_4| > \sqrt{a_1 b_1}$. Putting $p = \max\{|z_2|, |z_3|\}$, we have

$$\min\{\sqrt{a_2b_2}, \sqrt{a_3b_3}\} \ge |z_4| + \left(p - \sqrt{a_1b_1}\right),$$

by the inequalities $S_4[i, j|k, \ell]$ for $\{i, j\} = \{1, 2\}, \{1, 3\}$ and $\{k, \ell\} = \{2, 4\}, \{3, 4\}$. Therefore, we have

$$\begin{split} \sum_{i=2}^{4} \sqrt{s_i t_i} \sqrt{a_i b_i} - |u_4| |z_4| &\geq (\sqrt{s_2 t_2} + \sqrt{s_3 t_3}) \min\{\sqrt{a_2 b_2}, \sqrt{a_3 b_3}\} + \sqrt{s_4 t_4} |z_4| - |u_4| |z_4| \\ &\geq \left(\sum_{i=2}^{4} \sqrt{s_i t_i} - |u_4|\right) |z_4| + (\sqrt{s_2 t_2} + \sqrt{s_3 t_3}) \left(p - \sqrt{a_1 b_1}\right) \\ &\geq \left(\sum_{i=2}^{4} \sqrt{s_i t_i} - |u_4|\right) \sqrt{a_1 b_1} + (\sqrt{s_2 t_2} + \sqrt{s_3 t_3}) \left(p - \sqrt{a_1 b_1}\right) \\ &= (\sqrt{s_4 t_4} - |u_4|) \sqrt{a_1 b_1} + (\sqrt{s_2 t_2} + \sqrt{s_3 t_3}) p, \end{split}$$

by the inequality $W_2[4,1]$ and the assumption $|z_4| > \sqrt{a_1 b_1}$. We also have

$$\begin{split} \sqrt{s_1 t_1} \sqrt{a_1 b_1} &- \sum_{i=1}^3 |u_i| |z_i| \ge \sqrt{s_1 t_1} \sqrt{a_1 b_1} - |u_1| \sqrt{a_1 b_1} - |u_2| |z_2| - |u_3| |z_3| \\ &\ge \sqrt{s_1 t_1} \sqrt{a_1 b_1} - |u_1| \sqrt{a_1 b_1} - (|u_2| + |u_3|) p. \end{split}$$

Summing up the above two inequalities, we have

$$\sum_{i=1}^{4} (\sqrt{s_i t_i} \sqrt{a_i b_i} - |u_i| |z_i|) \\ \ge (\sqrt{s_1 t_1} + \sqrt{s_4 t_4} - |u_1| - |u_4|) \sqrt{a_1 b_1} + (\sqrt{s_2 t_2} + \sqrt{s_3 t_3} - |u_2| - |u_3|) p,$$

which is nonnegative by the inequalities $W_1[1, 4]$ and $W_1[2, 3]$. Therefore, we have

$$\frac{1}{2} \langle \mathsf{X}(s,t,u), \mathsf{X}(a,b,z) \rangle = \frac{1}{2} \sum_{i=1}^{4} (s_i a_i + t_i b_i + 2 \operatorname{Re}(u_i z_i))$$
$$\geq \sum_{i=1}^{4} (\sqrt{s_i t_i} \sqrt{a_i b_i} - |u_i| |z_i|) \ge 0,$$

which completes the proof. \Box

3. Examples

In order to get analytic examples distinguishing the convex cones in the inequalities (1) and (2), we consider GHZ diagonal states

$$\varrho_{0,0} = \frac{1}{8} \mathsf{X} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \varrho_{1,0} = \frac{1}{8} \mathsf{X} \begin{pmatrix} 1 & 2 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \quad \varrho_{0,1} = \frac{1}{12} \mathsf{X} \begin{pmatrix} 2 & 1 & 1 & 2 \\ 2 & 1 & 1 & 0 \end{pmatrix},$$

and define

$$\begin{split} \varrho_{s,t} &= (1-s-t)\varrho_{0,0} + s\varrho_{1,0} + t\varrho_{0,1} \\ &= \frac{1}{8}\mathsf{X} \begin{pmatrix} 1 + \frac{1}{3}t & 1 + s - \frac{1}{3}t & 1 - s - \frac{1}{3}t & 1 + \frac{1}{3}t \\ s + \frac{4}{3}t & \frac{2}{3}t & \frac{2}{3}t & s \end{pmatrix}, \end{split}$$

for real numbers s and t. We consider the convex set \mathbb{D} of all three qubit states, which is a 63 affine dimensional convex body. We slice \mathbb{D} by the 2-dimensional plane



FIGURE 1. The difference between the whole quadrilateral R and the bigger hexagon H_1 shows us that the distributive rules do not hold. The states $\rho_{s,t}$ in the region labeled by β and γ belong to $\beta \setminus \gamma$ and $\gamma \setminus \beta$, respectively. The smaller hexagon represents the convex set $\alpha \wedge \beta \wedge \gamma$ which consists of PPT states.

 Π determined by $\rho_{0,0}$, $\rho_{1,0}$ and $\rho_{0,1}$ to get the pictures for various convex sets. We see that $\rho_{s,t}$ is a state if and only if

$$|s + \frac{4}{3}t| \le 1 + \frac{1}{3}t, \quad |\frac{2}{3}t| \le 1 + s - \frac{1}{3}t, \quad |\frac{2}{3}t| \le 1 - s - \frac{1}{3}t, \quad |s| \le 1 + \frac{1}{3}t$$

if and only if (s, t) belongs to the region

$$R = \{(s,t): s+t \le 1, \ -s+t \le 1, \ -s-\frac{5}{3}t \le 1, \ s-\frac{1}{3}t \le 1\}$$

which is a quadrilateral on the *st*-plane with the four vertices (1,0), $(\frac{2}{3},-1)$, (-1,0) and (0,1). Therefore, the 2-dimensional convex body $\mathbb{D} \cap \Pi$ is also a quadrilateral with the vertices

(4)
$$\varrho_{0,1}, \quad \varrho_{1,0}, \quad \varrho_{\frac{2}{3},-1} = \frac{1}{12} \mathsf{X} \begin{pmatrix} 1 & 3 & 1 & 1 \\ -1 & -1 & -1 & 1 \end{pmatrix}, \quad \varrho_{-1,0} = \frac{1}{8} \mathsf{X} \begin{pmatrix} 1 & 0 & 2 & 1 \\ -1 & 0 & 0 & -1 \end{pmatrix}.$$

See Figure 1.

It is easily checked by $S_1[1, 4]$ and $S_1[2, 3]$ that the four states in (4) belong to the convex set α , and so the convex set α on the plane Π is represented by the quadrilateral R itself. Using $S_1[1, 3]$, $S_1[2, 4]$ and $S_1[1, 2]$, $S_1[3, 4]$, it is also easy to characterize (s, t)such that $\rho_{s,t}$ belongs to to β and γ , respectively. One may check that $\rho_{s,t} \in \beta$ if and only if it is a state and satisfies both inequalities $2s + \frac{5}{3}t \leq 1$ and $-2s + \frac{1}{3}t \leq 1$. Therefore, the region for β on the plane Π is a pentagon with vertices

$$\left(-\frac{2}{5},\frac{3}{5}\right), \quad \left(-\frac{2}{11},\frac{9}{11}\right), \quad \left(\frac{6}{7},-\frac{3}{7}\right), \quad \left(\frac{2}{3},-1\right), \quad \left(-\frac{6}{11},-\frac{3}{11}\right).$$

We also see that the region for γ is determined by $\frac{5}{3}t \leq 1$, $-2s - t \leq 1$ and $2s + \frac{1}{3}t \leq 1$. This is also a pentagon with vertices

$$\left(-\frac{2}{5},\frac{3}{5}\right), \quad \left(\frac{2}{5},\frac{3}{5}\right), \quad \left(\frac{2}{3},-1\right), \quad \left(-\frac{2}{7},-\frac{3}{7}\right), \quad \left(-\frac{2}{3},\frac{1}{3}\right).$$

It is clear that the region for $\alpha \vee \beta$ or $\gamma \vee \alpha$ on the plane Π occupies all of the quadrilateral R. One may also easily check by $S_2[1,4]$ that the four states in (4) belong to $\beta \vee \gamma$, and so the region for $\beta \vee \gamma$ coincides with the quadrilateral R. More precisely, the convex sets

$$(\alpha \lor \beta) \cap \Pi = (\beta \lor \gamma) \cap \Pi = (\gamma \lor \alpha) \cap \Pi = \alpha \cap \Pi$$

are represented by the quadrilateral R. The whole quadrilateral R in Figure 1 thus represents the regions for the following convex sets

 $\alpha, \quad \alpha \lor \beta, \quad \beta \lor \gamma, \quad \gamma \lor \alpha, \quad \alpha \lor \beta \lor \gamma, \quad \alpha \land (\beta \lor \gamma), \quad \alpha \lor (\beta \land \gamma)$

on the *st*-plane. It should be noted that they are strictly bigger than the convex hull generated by $\beta \cap \Pi$ and $\gamma \cap \Pi$. For example, the state $\rho_{1,0} \in \Pi$ in Figure 1 belongs to the convex hull $\beta \vee \gamma$, but it is not a mixture of states in $\beta \cap \Pi$ and $\gamma \cap \Pi$. In fact, if $\rho_{1,0} = \rho_1 + \rho_2$ with $\rho_1 \in \beta$ and $\rho_1 \in \gamma$ then one can easily see that the X-parts of ρ_1 and ρ_2 should be of the form

$$\frac{1}{8} \mathsf{X} \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & * & 0 & 1 \end{pmatrix} \qquad \text{and} \qquad \frac{1}{8} \mathsf{X} \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & * & 0 & 0 \end{pmatrix},$$

respectively. Therefore, they never belong to the plane Π .

Now, we use Theorem 2.1 to find the region for the convex set $\beta \vee (\gamma \wedge \alpha)$. For pairs $\{i, j\} = \{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}$, the form $8\sqrt{a_ib_i} + 8\sqrt{a_jb_j}$ for the state $\rho_{s,t}$ has the values

$$2+s, \quad 2-s, \quad 2+\frac{2}{3}t, \quad 2-\frac{2}{3}t, \quad 2+s, \quad 2-s,$$

respectively. On the other hands, $8|z_i| + 8|z_j|$ becomes

m

 $|s + \frac{4}{3}t| + \frac{2}{3}|t|, \quad |s + \frac{4}{3}t| + \frac{2}{3}|t|, \quad |s + \frac{4}{3}t| + |s|, \quad \frac{4}{3}|t|, \quad |s| + \frac{2}{3}|t|, \quad |s| + \frac{2}{3}|t|.$

Therefore, we see that a state $\rho_{s,t}$ belongs to $\beta \vee (\gamma \wedge \alpha)$ if and only if it belongs to $\gamma \vee (\alpha \wedge \beta)$ if and only if the inequality

$$\begin{aligned} &\inf\{2+s, \ 2-s, \ 2+\frac{2}{3}t, \ 2-\frac{2}{3}t\} \\ &\geq \max\{|s+\frac{4}{3}t|+\frac{2}{3}|t|, \ |s+\frac{4}{3}t|+|s|, \ \frac{4}{3}|t|, \ |s|+\frac{2}{3}|t|\} \end{aligned}$$

holds. One may check a point $(s,t) \in R$ satisfies this inequality if and only if

 $\frac{1}{2}s + \frac{4}{3}t \le 1, \qquad \frac{3}{2}s + \frac{2}{3}t \le 1, \qquad -\frac{3}{2}s - \frac{2}{3}t \le 1,$

This region is represented by the bigger hexagon H_1 with the vertices

 $\left(-\frac{2}{11},\frac{9}{11}\right), \left(\frac{2}{5},\frac{3}{5}\right), \left(\frac{6}{7},-\frac{3}{7}\right), \left(\frac{2}{3},-1\right), \left(-\frac{6}{11},-\frac{3}{11}\right), \left(-\frac{10}{13},\frac{3}{13}\right)$

in Figure 1. Therefore, the difference $R \setminus H_1$ consisting of three triangles gives us examples in

(5)
$$(\beta \lor \gamma) \land (\beta \lor \alpha) \land \beta \lor (\gamma \land \alpha),$$

which shows that the strict inequality holds in (2).

In order to consider the inequality (1), we first note the inequality

(6)
$$(\alpha \wedge \beta) \vee (\alpha \wedge \gamma) \leq \alpha \wedge (\beta \vee (\gamma \wedge \alpha)) \leq \alpha \wedge (\beta \vee \gamma),$$

which holds in general. By the first inequality, the region for $(\alpha \wedge \beta) \vee (\alpha \wedge \gamma)$ is also contained in H_1 . In fact, it fills up all of H_1 . Indeed, it is clear that five vertices of H_1 belong to $(\alpha \wedge \beta) \vee (\alpha \wedge \gamma)$ except for $(-\frac{10}{13}, \frac{3}{13})$ by $S_1[i, j]$. We also see that

$$52 \,\varrho_{-\frac{10}{13},\frac{3}{13}} = \mathsf{X} \begin{pmatrix} 7 & 1 & 11 & 7 \\ -3 & 1 & 1 & -5 \end{pmatrix} = \mathsf{X} \begin{pmatrix} 3 & 1 & 3 & 3 \\ -3 & 1 & 1 & -1 \end{pmatrix} + \mathsf{X} \begin{pmatrix} 4 & 0 & 8 & 4 \\ 0 & 0 & 0 & -4 \end{pmatrix}$$

belongs to $(\alpha \wedge \beta) \vee (\alpha \wedge \gamma)$. Therefore, this coincides with $\alpha \wedge (\beta \vee (\gamma \wedge \alpha))$ for states $\rho_{s,t}$, that is, the first inequality in (6) becomes an identity on the plane Π . Therefore, the difference $R \setminus H_1$ again gives rise to examples in

(7)
$$\alpha \wedge (\beta \vee \gamma) \setminus (\alpha \wedge \beta) \vee (\alpha \wedge \gamma),$$

for the strict inequality in (1).

Now, we proceed to show that the gaps (5) and (7) arising in the distributive inequalities have nonzero volume. We first note that the convex set \mathbb{S} of all fully separable three qubit states generates the same affine manifold as the convex set \mathbb{D} of all three qubit states. See the discussion at the end of Section 7 in [23]. Therefore, all the convex sets in (5) and (7) generate the same affine manifold. We also recall that a point x_0 of a convex set C is called an *interior point* of C if it is an interior point of C with respect to the affine space generated by C. We note that the state $\varrho_{0,0}$ is a common interior point of the convex sets appearing in (5) and (7) as well as \mathbb{S} and \mathbb{D} . If we consider the line segment $x_t = (1 - t)x_0 + tx_1$ between an interior point x_0 of a convex set C and an arbitrary point $x_1 \in C$ then x_t is also an interior point of C for 0 < t < 1. See [24, Lemma 2.3]. Therefore, every interior point of $[\alpha \land (\beta \lor \gamma)] \cap \Pi$ is actually an interior point of $\alpha \land (\beta \lor \gamma)$. For example, $\varrho_{\frac{2}{3},0}$ is an interior point of $\alpha \land (\beta \lor \gamma)$ which is a boundary point of $(\alpha \land \beta) \lor (\alpha \land \gamma)$. From this, we may conclude that the difference (7) has the nonempty interior by [23, Proposition 7.4]. The exactly same argument also shows that the difference (5) also has the nonempty interior.

We also consider the smaller hexagon H_2 with vertices

 $\left(-\frac{2}{5},\frac{3}{5}\right), \quad \left(0,\frac{3}{5}\right), \quad \left(\frac{1}{2},0\right), \quad \left(\frac{2}{3},-1\right), \quad \left(-\frac{2}{7},-\frac{3}{7}\right), \quad \left(-\frac{1}{2},0\right)$

in Figure 1, which is the region for $\rho_{s,t}$ in $\beta \wedge \gamma$. This represents also $\alpha \wedge \beta \wedge \gamma$. Note that an X-shaped state belongs to $\alpha \wedge \beta \wedge \gamma$ if and only if it is of positive partial transpose by [22, Theorem 5.3]. Therefore, the region H_2 represents the region for PPT states for $\rho_{s,t}$.

Recall that a lattice is called *modular* if $x \leq z$ implies $x \vee (y \wedge z) = (x \vee y) \wedge z$. This is the case if and only if the modular identity

$$(x \land z) \lor (y \land z) = ((x \land z) \lor y) \land z$$

holds for every x, y and z. Every distributive lattice is modular. See [25, 26] for elementary properties of modular lattices. We exhibit examples of states showing that the first inequality in (6) is also strict, to conclude that the lattice \mathcal{L} is not modular. To do this, we consider $\rho_1 = \frac{1}{12} \mathsf{X} \begin{pmatrix} 2 & 1 & 1 & 2 \\ 2 & 0 & 1 & 0 \end{pmatrix}$, and put

$$\varrho_t = (1-t)\varrho_{0,0} + t\varrho_1 = \frac{1}{24} \mathsf{X} \begin{pmatrix} 3+t & 3-t & 3-t & 3+t \\ 4t & 0 & 2t & 0 \end{pmatrix}.$$

We also consider $W = \mathsf{X} \begin{pmatrix} 0 & 1 & 1 & 0 \\ -1 & 0 & -1 & 0 \end{pmatrix}$, which satisfies the inequality $W_2[i, j]$ for $\{i, j\} = \{1, 2\}, \{3, 4\}, \{1, 3\}, \{2, 4\}$ and W_3 . Therefore, we see that W belongs to the convex cone $(\alpha^{\circ} \lor \beta^{\circ}) \land (\alpha^{\circ} \lor \gamma^{\circ})$, which is the dual of the convex cone $(\alpha \land \beta) \lor (\alpha \land \gamma)$. Now, we see that $\langle W, \varrho_t \rangle = \frac{1}{24}(6-8t) \ge 0$ if and only if $t \le \frac{3}{4}$. Therefore, we conclude that ρ_t does not belong to $(\alpha \wedge \beta) \vee (\alpha \wedge \gamma)$ for $\frac{3}{4} < t \leq 1$. On the other hand, one can easily check that $\varrho_1 \in \alpha \land (\beta \lor (\gamma \land \alpha))$ by Theorem 2.1. Therefore, we see that ϱ_t also belongs to the same cone. In fact, ρ_t is an interior point of the cone $\alpha \land (\beta \lor (\gamma \land \alpha))$ for $0 \le t < 1$, because $\rho_{0,0}$ is an interior point. Hence, we also see that the gap for the first inequality in (6) has also nonzero volume.

Finally, we also characterize the full separability for the states we are considering. To do this, we summarize the results in [28, 29, 27]. See also [30, 23]. For a given GHZ diagonal state $\rho = X(a, a, c)$ with $a, c \in \mathbb{R}^4$, we consider the following:

$$\lambda_{5} = 2(+c_{1} + c_{2} + c_{3} + c_{4}), \quad \lambda_{6} = 2(-c_{1} - c_{2} + c_{3} + c_{4}),$$
$$\lambda_{7} = 2(-c_{1} + c_{2} - c_{3} + c_{4}), \quad \lambda_{8} = 2(-c_{1} + c_{2} + c_{3} - c_{4}),$$
$$t_{1} = c_{1}(-c_{1}^{2} + c_{2}^{2} + c_{3}^{2} + c_{4}^{2}) - 2c_{2}c_{3}c_{4}, \quad t_{2} = c_{2}(+c_{1}^{2} - c_{2}^{2} + c_{3}^{2} + c_{4}^{2}) - 2c_{1}c_{3}c_{4},$$
$$t_{3} = c_{3}(+c_{1}^{2} + c_{2}^{2} - c_{3}^{2} + c_{4}^{2}) - 2c_{1}c_{2}c_{4}, \quad t_{4} = c_{4}(+c_{1}^{2} + c_{2}^{2} + c_{3}^{2} - c_{4}^{2}) - 2c_{1}c_{2}c_{3}.$$

When all the following inequalities

(8)
$$\lambda_5\lambda_6\lambda_7\lambda_8 > 0, \quad t_1t_4\lambda_6\lambda_7 < 0, \quad t_2t_3\lambda_5\lambda_8 > 0$$

hold, the state $\rho = X(a, a, c)$ is fully separable if and only if the inequality

(9)
$$\min\{a_1, a_2, a_3, a_4\} \ge \frac{\sqrt{(\lambda_5\lambda_6 + \lambda_7\lambda_8)(\lambda_5\lambda_7 + \lambda_6\lambda_8)(\lambda_5\lambda_8 + \lambda_6\lambda_7)}}{8\sqrt{\lambda_5\lambda_6\lambda_7\lambda_8}}$$

is satisfied. In the other cases, ρ is fully separable if and only if it is of PPT. For the state $\rho_{s,t}$, the conditions (8) are given by

$$s(3s+4t) < 0,$$
 $(9s^2 + 18st + 4t^2)(9s^2 + 6st - 4t^2) > 0.$



FIGURE 2. Two line segments through the origin are given by the conditions (8), and two curves surrounding the region of full separability are given by (9).

We note that a point (s,t) on the line t = as satisfies this condition if and only if $-\frac{3}{4}(3+\sqrt{5}) \leq a < -\frac{3}{4}$. On the other hands, the square of right side of (9) is given by

$$\frac{t^2(9s^2+12st-4t^2)}{432s(3s+4t)}$$

respectively. See Figure 2.

4. General multi-partite cases

Now, we turn our attention to general multi-partite system $M_{A_1} \otimes M_{A_2} \otimes \cdots \otimes M_{A_n}$. For a given partition Π of local systems $\{A_1, A_2, \ldots, A_n\}$, we denote by α_{Π} the convex cone consisting of all partially separable (unnormalized) states with respect to the partition Π . When M_{A_i} is the $d_i \times d_i$ matrices, we denote by $\mathcal{L}_{d_1,d_2,\ldots,d_n}$ the lattice generated by α_{Π} through all nontrivial partitions Π with respect to two operations, intersection and convex hull. Therefore, elements of this lattice are convex cones sitting in the real vector space of all $d_1 d_2 \cdots d_n \times d_1 d_2 \cdots d_n$ Hermitian matrices. So far, we have considered the lattice $\mathcal{L}_{2,2,2}$. We close this section by showing that the lattice $\mathcal{L}_{d_1,d_2,\ldots,d_n}$ also violates distributivity and modularity.

For $d \geq 2$, we consider the canonical embedding

$$\iota_d: \varrho \in M_2 \mapsto \varrho \oplus \operatorname{tr}(\varrho) \mathbb{1}_{d-2} \in M_d$$

and the compression

$$Q_d: \varrho \in M_d \mapsto \begin{pmatrix} I_2 & 0 \end{pmatrix} \varrho \begin{pmatrix} I_2 \\ 0 \end{pmatrix} \in M_2.$$

Here, the trace is a normalized one. Then, the composition $Q_d \circ \iota_d$ is the identity map. In other words, M_2 is unital completely positive (u.c.p.) complemented in M_n . This means that every local property of $M_p \otimes M_q \otimes M_r$ is hereditary to $M_2 \otimes M_2 \otimes M_2$. We write

$$\iota = \iota_p \otimes \iota_q \otimes \iota_r : M_2 \otimes M_2 \otimes M_2 \to M_p \otimes M_q \otimes M_r$$
$$Q = Q_p \otimes Q_q \otimes Q_r : M_p \otimes M_q \otimes M_r \to M_2 \otimes M_2 \otimes M_2$$

for brevity.

We retain the notations α , β and γ for the three qubit case, that is, $\alpha = \alpha_{A-BC}$, $\beta = \alpha_{B-CA}$ and $\gamma = \alpha_{C-AB}$ generate the lattice $\mathcal{L}_{2,2,2}$. To make clear, we will use notations $\tilde{\alpha}$, $\tilde{\beta}$ and $\tilde{\gamma}$ for generators of the lattice $\mathcal{L}_{p,q,r}$. It is easily seen that $\varrho \in \alpha$ implies $\iota(\varrho) \in \tilde{\alpha}$, and $\omega \in \tilde{\alpha}$ implies $Q(\omega) \in \alpha$, and similarly for β and γ . In order to show that $\mathcal{L}_{p,q,r}$ violates the distributive rules, we first take

$$\varrho \in (\alpha \land (\beta \lor \gamma)) \land (\alpha \land \beta) \lor (\alpha \land \gamma)$$

in $M_2 \otimes M_2 \otimes M_2$. We can write $\rho = \rho_2 + \rho_3$ for $\rho_2 \in \beta$ and $\rho_3 \in \gamma$. We have

$$\iota(\varrho) \in \tilde{\alpha}, \qquad \iota(\varrho_2) \in \tilde{\beta}, \qquad \iota(\varrho_3) \in \tilde{\gamma}$$

in $M_p \otimes M_q \otimes M_r$. Thus, $\iota(\varrho)$ belongs to $\tilde{\alpha} \wedge (\tilde{\beta} \vee \tilde{\gamma})$ in $M_p \otimes M_q \otimes M_r$. Assume to the contrary that the distributive rule holds in $M_p \otimes M_q \otimes M_r$. Then, $\iota(\varrho)$ belongs to $(\tilde{\alpha} \wedge \tilde{\beta}) \vee (\tilde{\alpha} \wedge \tilde{\gamma})$, and so we can write $\iota(\varrho) = \omega_2 + \omega_3$ with $\omega_2 \in \tilde{\alpha} \wedge \tilde{\beta}$ and $\omega_3 \in \tilde{\alpha} \wedge \tilde{\gamma}$. Therefore, we have

$$\varrho = Q \circ \iota(\varrho) = Q(\omega_2) + Q(\omega_3) \in (\alpha \land \beta) \lor (\alpha \land \gamma),$$

which is a contradiction. The same argument can be applied to the other distributive rule. It should be noted that ω_2 and ω_3 themselves need not belong to the image of ι in the above argument. Non-modularity also follows from that of three qubit system in the same fashion.

For general multi-partite cases, we consider the canonical embedding

 $\iota: x \in M_{d_1} \otimes M_{d_2} \otimes M_{d_3} \mapsto x \otimes 1 \in M_{d_1} \otimes M_{d_2} \otimes M_{d_3} \otimes M_{d_4} \otimes \cdots \otimes M_{d_n}$

and the (normalized) partial trace

$$\mathrm{id}\otimes\mathrm{tr}:M_{d_1}\otimes M_{d_2}\otimes M_{d_3}\otimes M_{d_4}\otimes\cdots\otimes M_{d_n}\to M_{d_1}\otimes M_{d_2}\otimes M_{d_3}.$$

Then, the composition $(\mathrm{id} \otimes \mathrm{tr}) \circ \iota$ is the identity map. In other words, $M_{d_1} \otimes M_{d_2} \otimes M_{d_3}$ is u.c.p. complemented in $M_{d_1} \otimes M_{d_2} \otimes M_{d_3} \otimes M_{d_4} \otimes \cdots \otimes M_{d_n}$. This means that every local property of the latter is hereditary to $M_p \otimes M_q \otimes M_r$. By the exactly same argument as before, one may check that the lattice $\mathcal{L}_{d_1,\ldots,d_1}$ also violates distributivity and modularity.

5. Summary and further questions

In this paper, we have considered the lattice \mathcal{L} generated by three basic convex sets α, β and γ consisting of all A-BC biseparable, B-CA biseparable and C-AB biseparable three qubit states, respectively, with respect to the operations of convex hull \vee and intersection \wedge . In this way, we may consider convex sets of partially separable states obtaining by arbitrary convex hulls and intersections of α, β and γ , and the whole structure of partial separability may be revealed by mathematical properties of the lattice \mathcal{L} . For general theory for lattices, we refer to the monographs [25], [31] and [26].

We gave the negative answer to the first natural question asking if this lattice is distributive. The lattice \mathcal{L} is not even modular. Another interesting question is to ask if the lattice \mathcal{L} has infinitely many elements. We conjecture this is the case. This means that there are infinitely many kinds of partial separability and partial entanglement. It is known that a free lattice with three generators must have infinitely many elements. In this regard, it would be interesting to know if the lattice \mathcal{L} is free or not.

The next question is whether the lattice \mathcal{L} is complemented or not. A lattice is called *complemented* if every element x has a complement y which satisfies $x \wedge y = 0$ and $x \vee y = 1$, where 0 and 1 denote the least and greatest elements, respectively. The least and the greatest elements of \mathcal{L} are given by $\alpha \wedge \beta \wedge \gamma$ and $\alpha \vee \beta \vee \gamma$, respectively. They represent the set of all fully biseparable and biseparable states, respectively. Especially, we would like to ask if α has a complement, that is, we ask if there exist $\sigma \in \mathcal{L}$ such that $\alpha \wedge \sigma = \alpha \wedge \beta \wedge \gamma$ and $\alpha \vee \sigma = \alpha \vee \beta \vee \gamma$. Recall that the set of all closed subspaces of a Hilbert space makes a lattice, the *subspace lattice*, with respect to the closed linear hull and intersection. This plays an important role in quantum logic and theory of operator algebras. We note that the subspace lattice is non-distributive, but complemented.

Finally, we ask how the lattice $\mathcal{L}_{d_1,\ldots,d_n}$ depends on the dimensions of the local systems. In the case of tri-partite systems, we are asking if two lattices $\mathcal{L}_{2,2,2}$ and $\mathcal{L}_{p,q,r}$ are isomorphic to each other.

Acknowledgement: Both KHH and SHK were partially supported by the grant NRF-2017R1A2B4006655, Korea. SzSz was supported by the National Research, Development and Innovation Fund of Hungary within the *Researcher-initiated Research Program* (project Nr: NKFIH-K120569) and within the *Quantum Technology National Excellence Program* (project Nr: 2017-1.2.1-NKP-2017-00001), by the Ministry for Innovation and Technology within the ÚNKP-19-4 New National Excellence Program, and by the Hungarian Academy of Sciences within the János Bolyai Research Scholarship and the "Lendület" Program.

References

 R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki, *Quantum entanglement*, Rev. Mod. Phys. 81 (2009), 865–942.

- [2] E. Schrödinger, Die gegenwärtige Situation in der Quantenmechanik, Naturwissenschaften 23 (1935) 823–828.
- [3] E. Schrödinger, Discussion of Probability Relations between Separated Systems, Math. Proc. Camb. Phil. Soc. 31 (1935), 555–563.
- [4] L. Amico, R. Fazio, A. Osterloh and Vlatko Vedral, *Entanglement in many-body systems*, Rev. Mod. Phys. 80 (2008), 517–576.
- [5] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information, Cambridge University Press, 2000.
- [6] R. F. Werner, Quantum states with Einstein-Podolsky-Rosen correlations admitting a hiddenvariable model, Phys. Rev. A, 40 (1989), 4277–4281.
- [7] C. H. Bennett, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin and B. M. Terhal, Unextendible product bases and bound entanglement, Phys. Rev. Lett. 82 (1999), 5385–5388.
- [8] W. Dür, J. I. Cirac and R. Tarrach, Separability and Distillability of Multiparticle Quantum Systems, Phys. Rev. Lett. 83 (1999), 3562–3565.
- [9] W. Dür and J. I. Cirac, Classification of multi-qubit mixed states: separability and distillability properties, Phys. Rev. A 61 (2000), 042314.
- [10] M. Seevinck and J. Uffink, Partial separability and etanglement criteria for multiqubit quantum states, Phys. Rev. A 78 (2008), 032101.
- [11] A. Acin, D. Bruß, M. Lewenstein and A. Sanpera, *Classification of mixed three-qubit states*, Phys. Rev. Lett. 87 (2001), 040401.
- [12] Sz. Szalay and Z. Kökényesi, Partial separability revisited: Necessary and sufficient criteria, Phys. Rev. A 86 (2012), 032341.
- [13] K. H. Han and S.-H. Kye, Construction of three-qubit biseparable states distinguishing kinds of entanglement in a partial separability classification, Phys. Rev. A, 99 (2019), 032304.
- [14] K. H. Han and S.-H. Kye, On the convex cones arising from classifications of partial entanglement in the three qubit system, J. Phys. A: Math. Theor. 53 (2020), 015301.
- [15] Sz. Szalay, Multipartite entanglement measures, Phys. Rev. A 92 (2015), 042329.
- [16] Sz. Szalay, Separability criteria for mixed three-qubit states, Phys. Rev. A 83 (2011), 062337.
- [17] Sz. Szalay, The classification of multipartite quantum correlation, J. Phys. A: Math. Theor. 51 (2018), 485302.
- [18] Sz. Szalay, k-stretchability of entanglement, and the duality of k-separability and k-producibility, Quantum 3 (2019), 204.
- [19] D. M. Greenberger, M. A. Horne and A. Zeilinger, *Going beyond Bell's theorem*, in Kafatos M. (eds) Bells Theorem, Quantum Theory and Conceptions of the Universe. Fundamental Theories of Physics, Vol. 37, Springer, Dordrecht, 1989.
- [20] T. Gao and Y. Hong, Separability criteria for several classes of n-partite quantum states, Eur. Phys. J. D 61 (2011), 765–771.
- [21] S. M. H. Rafsanjani, M. Huber, C. J. Broadbent and J. H. Eberly, Genuinely multipartite concurrence of N-qubit X matrices, Phys. Rev. A 86 (2012), 062303.
- [22] K. H. Han and S.-H. Kye, Construction of multi-qubit optimal genuine entanglement witnesses, J. Phys. A: Math. Theor. 49 (2016), 175303.
- [23] K.-C. Ha, K. H. Han and S.-H. Kye, Separability of multi-qubit states in terms of diagonal and anti-diagonal entries, Quantum Inf. Process. 18 (2019), 34.
- [24] S.-H. Kye, Facial structures for the positive linear maps between matrix algebras, Canad. Math. Bull. 39 (1996), 74–82.
- [25] G. Birkhoff, Lattice Theory, Colloq. Publ. Vol 25, Amer. Math. Soc. 1940.
- [26] V. N. Salii, Lattices with unique complements, Trans. Math. Monog. Vol 69, Amer. Math. Soc. 1988.
- [27] A. Kay, Optimal detection of entanglement in Greenberger-Horne-Zeilinger states, Phys. Rev. A 83 (2011), 020303(R).
- [28] O. Gühne, Entanglement criteria and full separability of multi-qubit quantum states, Phys. Lett. A 375 (2011), 406–410.
- [29] K. H. Han and S.-H, Kye, Separability of three qubit Greenberger-Horne-Zeilinger diagonal states, J. Phys. A: Math. Theor. 50 (2017), 145303.
- [30] L. Chen, K. H. Han and S.-H, Kye, Separability criterion for three-qubit states with a four dimensional norm, J. Phys. A: Math. Theor. 50 (2017) 345303.

[31] R. Freese, J. Jezek and J. B. Nation, Free Lattice, Math. Surv Monog. Vol 42, Amer. Math. Soc., 1995.

Kyung Hoon Han, Department of Data Science, The University of Suwon, Gyeonggido 445-743, Korea

E-mail address: kyunghoon.han at gmail.com

SEUNG-HYEOK KYE, DEPARTMENT OF MATHEMATICS AND INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151-742, KOREA

E-mail address: kye at snu.ac.kr

Szilárd Szalay, Wigner Research Centre for Physics, 29-33, Konkoly-Thege Miklós, H-1121 Budapest, Hungary

E-mail address: szalay.szilard at wigner.mta.hu