Homogenization of oblique boundary value problems

Sunhi Choi *and Inwon C. Kim [†]

Abstract

We consider a nonlinear Neumann problem, with periodic oscillation in the elliptic operator and on the boundary condition. Our focus is on problems posed in half-spaces, but with general normal directions that may not be parallel to the directions of periodicity. As the frequency of the oscillation grows, quantitative homogenization results are derived. When the homogenized operator is rotation-invariant, we prove the Hölder continuity of the homogenized boundary data. While we follow the outline of [7], new challenges arise due to the presence of tangential derivatives on the boundary condition in our problem. In addition we improve and optimize the rate of convergence within our approach. Our result appear to be new even for the linear oblique problem.

1 Introduction

For given $\varepsilon > 0$, $\nu \in \mathcal{S}^{n-1}$ and $\tau \in \mathbb{R}^n$, let u_{ε} be a bounded solution of the following problem:

$$(P)_{\varepsilon} \qquad \left\{ \begin{array}{ll} F(D^{2}u_{\varepsilon},\frac{x}{\varepsilon}) = 0 & \text{in} \quad \Pi := \{x \in \mathbb{R}^{n} : -1 < (x-\tau) \cdot \nu < 0\} \\ u_{\varepsilon} = h(x) & \text{on} \quad H_{-1} := \{(x-\tau) \cdot \nu = -1\} \\ \partial_{\nu}u = G(Du_{\varepsilon},\frac{x}{\varepsilon}) & \text{on} \quad H_{0} := \{(x-\tau) \cdot \nu = 0\}. \end{array} \right.$$

Here F(M, y) and G(p, y) are \mathbb{Z}^n -periodic in the y variable. We also assume the boundary condition to be *oblique* and F to be uniformly elliptic: see Section 1.1 for precise assumptions on F and G.

The examples of boundary conditions we consider include the linear oblique problem

$$\vec{\gamma}(\frac{x}{\varepsilon}) \cdot Du + g(\frac{x}{\varepsilon}) = 0, \tag{1}$$

where the vector field $\vec{\gamma}$ satisfies $c(\frac{x}{\varepsilon}, x) := \vec{\gamma}(\frac{x}{\varepsilon}) \cdot \nu > 0$. In this case one can write

$$G(p, y) = (c(y))^{-1} [\vec{\gamma}(y) \cdot p_T + g(y)],$$

where p_T is the tangential component of a vector field p on H_0 . A nonlinear example is *capillarity-type* conditions, for which G is given by

$$G(p,y) = \theta(y)\sqrt{1+|p|^2},$$
 (2)

where $|\theta(x)| < 1$. This condition describes a prescribed contact angle between the graph $\Gamma := \{(x, z) : z = u(x)\}$ and the "container boundary" $H_0 \times \mathbb{R}$ with chemical inhomogeneities.

We are interested in the behavior of u_{ε} as ε tends to zero. Note that, as first pointed out by Bensoussan, Lions, and Papanicolaou [5], if ν is a multiple of a vector in \mathbb{Z}^n (i.e. if ν is rational)

^{*}Department of Mathematics, U. of Arizona, Tucson, AZ.

[†]Department of Mathematics, UCLA, LA CA. Research supported by NSF DMS-1566578

then $\tau \cdot \nu$ must be zero for u^{ε} to converge, since otherwise the Neumann boundary condition changes drastically as ε changes, and thus u_{ε} would not have a limit. When ν is irrational we expect u_{ε} to average due to the ergodic property of its Neumann data. However in this case u^{ε} is no longer periodic, and thus interesting challenges arise in dealing with the inherent lack of compactness. Compared to [7] where linear Neumann problem was considered, there is an additional challenge in our setting given by the presence of tangential derivatives on the boundary condition. We will discuss below some of the relevant literature on this issue.

Let us state a convergence result on $(P)_{\varepsilon}$ to begin the discussion. let \overline{F} be the homogenized operator of F obtained by Evans [11].

Theorem 1.1. Let ν be irratonal, or otherwise suppose $\tau = 0$. Let us assume (F1) - (F3) and (G1) - (G3) (see Section 1.1). In addition suppose that $F(\cdot, x)$ is convex when $G(\cdot, x)$ is nonlinear. Then there exists $\mu(\nu, D_T u)$, independent of τ , such that u_{ε} converges uniformly to the unique bounded solution \bar{u} of

$$(\bar{P}) \qquad \qquad \begin{cases} \bar{F}(D^2\bar{u}) = 0 & in & \Pi \\ \\ \bar{u} = h(x) & on & H_{-1} \\ \\ \partial_{\nu}\bar{u} = \mu(\nu, D_T\bar{u}) & on & H_0. \end{cases}$$

Moreover $\mu(\nu, q)$ is Lipschitz continuous with respect to q. If $\overline{F}(M)$ is rotation-invariant, then μ is also Hölder continuous over irrational directions ν with exponent $\alpha = \frac{1}{5n}$.

The proof of Theorem 1.1 will be given later in this section, based on our main result (Theorem 1.2), which establishes rates of convergence for (approximate) cell problem solutions. Our work extends the previous results in [8] on linear Neumann problems where G(p, y) = G(y). For general G(p, y) additional challenges arise due to the presence of tangential derivatives on the boundary condition, which necessitates Lipschitz regularity estimates for the solutions. As noted in [13], the continuity property of $\mu(\nu, q)$ fails when \overline{F} is not rotation-invariant, even when it is convex. When the continuity result holds for μ one can expect to proceed as in [7] to address general domains, but the analysis would require higher regularity estimates on the solutions, so we do not pursue this here.

It is unknown whether the form of the boundary condition such as (1) or (2) is preserved in the limit $\varepsilon \to 0$. With the exception of linear problems, the interaction between the operator F and the boundary condition remains to be better understood to yield further characterizations of the homogenized problems.

Literature

Before proceeding further, let us briefly describe some of relevant literature. In the classical paper of [5], the following problem was considered:

$$-\nabla \cdot (A(\frac{x}{\varepsilon})\nabla u_{\varepsilon}) = 0 \text{ in } \Omega, \qquad \nu \cdot (A(\frac{x}{\varepsilon})\nabla u_{\varepsilon})(x) = g(\frac{x}{\varepsilon}) \text{ on } \partial\Omega.$$
(3)

For this co-normal boundary value problem, explicit integral formulas have been derived for the limiting operator as well as for the limiting boundary data, under the assumption that $\partial\Omega$ does not contain any flat piece with a rational normal.

For linear elliptic systems with either Dirichlet or Neumann problem with co-normal derivatives, there has been a recent surge of development in quantitative homogenization by integral representation of solutions: we refer to [2],[15],[20] and the references therein.

For nonlinear problems, or even for linear problems with non co-normal boundary data, most available homogenization results concern half-space type domains whose boundary goes through the origin and is normal to a rational direction. In [21], Tanaka considered some model problems in half-space whose boundary is parallel to the axes of the periodicity by purely probabilistic methods. In [1] Arisawa studied specific problems in oscillatory domains near half spaces going through the origin. Generalizing the results of [1] for nonlinear boundary conditions, Barles, Da Lio, Lions and Souganidis [4] studied the problem for operators with oscillating coefficients, in half-space type domains whose boundary is parallel to the axes of periodicity. We also refer to [14] which adopts an integro-differential approach to study linear scalar problems with the specific Neumann problem G(p, y) = g(y).

For the linear Neumann problem G(p, y) = g(y) in $(P)_{\varepsilon}$, corresponding results to Theorem 1.1 -Theorem 1.2 have been recently shown in [8]. General domains has been considered in [7] based on the cell problem analysis in [8]. Corresponding results for the Dirichlet boundary data has been obtained in [12]. Lastly for general operator F, [13] discusses the generic nature of discontinuity for the homogenized boundary data, for either linear Neumann or Dirichlet problem.

Cell problem

By the formal expansion $u_{\varepsilon} = \bar{u}(x) + \varepsilon v(x, \frac{x}{\varepsilon}) + O(\varepsilon^2)$, for a rational ν , the cell problem for v was derived in [4] for a rational ν and $\tau = 0$. There they find a unique constant $\mu = \mu(\nu, q)$ for $q \in \langle \nu \rangle^{\perp}$ such that the boundary value problem

(C)
$$\begin{cases} F(D^2v, y) = 0 & \text{in} \quad \{y \cdot \nu \ge 0\}, \\ \mu = G(Dv + p, y) & \text{on} \quad H_0, \end{cases}$$

with $p = \mu \nu + q$, has a bounded periodic solution v in $\{y \cdot \nu \ge 0\}$. The existence of bounded v leads to the uniform convergence of u_{ε} to \bar{u} in the limit $\varepsilon \to 0$ with $p = D\bar{u}$ on H_0 .

For general ν and τ , an approximate cell problem needs to be derived, since v is no longer expected to be periodic and thus compactness is lost: see problem $(P)_{\varepsilon,\nu,\tau,q}$ below. In the context of (C), our result shows that for irrational ν , there exists a unique constant $\mu = \mu(\nu, q)$ for $q \in \langle \nu \rangle^{\perp}$ such that the problem

$$(\tilde{C}) \qquad \begin{cases} F(D^2v, y+\tau) = 0 & \text{in} \quad \{y \cdot \nu \ge 0\},\\ \mu = G(Dv+p, y+\tau) & \text{on} \quad H_0 \end{cases}$$

with any $\tau \in \mathbb{R}^n$ has a solution with sublinear growth at infinity. To show this, we use the ergodicity of Neumann data in a scale depending on ν , and the stability of solutions under perturbation of boundary conditions. When the homogenized operator \overline{F} is rotation-invariant, we show that v is stable as the normal direction of the domain ν varies. A quantiative version of this stability property yields the mode of continuity for μ as ν varies.

A discussion on assumptions on F and G

Our assumptions on F and G are mainly to obtain Lipschitz estimates for the solutions of (\hat{C}) . The Lipschitz estimates ensure that the solution of the cell problem has ergodic structure with respect to translations along the Neumann boundary (see Lemma 3.5), which happens when ε changes in $(P)_{\varepsilon}$ and when τ is not the origin. Already to guarantee the Lipschitz bound, available literature restricts F(M, x) to be convex with respect to M when G is a nonlinear function of Du. We refer to [3] for a detailed description of available regularity theory on nonlinear Neumann boundary problems. For the continuity properties of μ we further need $C^{1,\alpha}$ estimates for solutions of (\tilde{C}) , however this does not further restrict the class of problems we can address. To deal with domains with general geometry, the approach taken in [7] or [13] uses fundamental solutions as barriers to bound the potential singularity generated at points with rational normals. For our problem, while we suspect our result to hold in general domains, we suspect that these singular solutions may cause new challenges in dealing with perturbative arguments, due to their singularity in tangential derivatives.

1.1 Assumptions and main results

Let \mathbb{T} be the 1-periodic torus in \mathbb{R}^n , and let \mathcal{M}^n be the space of real $n \times n$ symmetric matrices. Consider the functions $F(M, y) : \mathcal{M}^n \times \mathbb{T} \to \mathbb{R}$ and $G(p, y) : \mathbb{R}^n \times \mathbb{T}$ satisfying the following properties:

(F1) (Uniform Ellipticity) There exist constants $0 < \lambda < \Lambda$ such that

$$\lambda \operatorname{Tr}(N) \le F(M, y) - F(M + N, y) \le \Lambda \operatorname{Tr}(N)$$

for all $y \in \mathbb{T}$ and $M, N \in \mathcal{M}^n$ with $N \ge 0$.

- (F2) (1-Homogeneity) F(tM, y) = tF(M, y) for all $y \in \mathbb{T}, t > 0$ and $M \in \mathcal{M}^n$.
- (F3) (Lipschitz Continuity) There exists C > 0 such that for all $y_1, y_2 \in \mathbb{T}$ and $M, N \in \mathcal{M}^n$,

$$|F(M, y_1) - F(N, y_2)| \le C(|y_1 - y_2|(1 + ||M|| + ||N||) + ||M - N||).$$

- (G1) (At most linear Growth) $|G(p, x)| \le \mu_0(1+|p|)$.
- (G2) (Lipschitz continuity) $(1+|p|)|G_p|, |G_y| \le m(1+|p|)$ for some m > 0.
- (G3) (Oblicity) $|G_p \cdot \nu| \le c < 1.$

A typical example of an operator F satisfying (F1)-(F3) is the linear elliptic operator

$$F(D^2u, x) = -\sum_{i,j} a_{ij}(x) \partial_{x_i x_j} u, \tag{4}$$

where $a_{ij} : \mathbb{R}^n \to \mathbb{R}$ is periodic and Lipschitz continuous. A nonlinear example is the Bellman-Isaacs operator arising from stochastic optimal control and differential games

$$F(D^2u, x) = \inf_{\beta \in B} \sup_{\alpha \in A} \{ \mathcal{L}^{\alpha, \beta}u \},$$
(5)

where $\mathcal{L}^{\alpha,\beta}$ is a family of uniformly elliptic operators of the form (4). In fact, all operators satisfying (F1)-(F3) can be written as (5). As for G, the ones given in (1) and (2) with Lipschitz coefficients $c^{-1}\vec{\gamma}, c^{-1}g$ and θ satisfy (G1)-(G3).

For $\tau \in \mathbb{R}^n$ and $\nu \in \mathcal{S}^{n-1}$, let us define the strip domain

$$\Pi(\tau,\nu):=\{x\in\mathbb{R}^n:-1\leq (x-\tau)\cdot\nu\leq 0\},\quad H_s:=\{(x-\tau)\cdot\nu=s\}.$$

For a given $q \in \langle \nu \rangle^{\perp}$, let u_{ε} solve the following approximate cell problem

$$(P)_{\varepsilon,\nu,\tau,q}. \qquad \begin{cases} F(D^2 u_{\varepsilon}, \frac{x}{\varepsilon}) = 0 & \text{in} \quad \Pi(\tau,\nu) \\\\ \partial_{\nu} u_{\varepsilon} = G(D u_{\varepsilon}, \frac{x}{\varepsilon}) & \text{on} \quad H_0 \\\\ u_{\varepsilon}(x) = q \cdot x & \text{on} \quad H_{-1} \end{cases}$$

Now we are ready to state the main result.

Theorem 1.2. Let u_{ε} solve $(P)_{\varepsilon,\nu,\tau,q}$. Suppose that either ν is irrational or $\tau = 0$. Then the following holds:

(a) There exists $\mu = \mu(\nu, q)$ such that u_{ε} converges uniformly to the linear profile

$$u(x) := \mu((x - \tau) \cdot \nu + 1) + q \cdot x$$

Here, $\mu(\nu,q)$ is independent of τ and Lipschitz continuous with respect to q. Moreover we have

$$|u_{\varepsilon} - u| \le C\Lambda(\varepsilon, \nu) \quad in \ \Pi(\tau, \nu), \tag{6}$$

where $\Lambda(\varepsilon,\nu)$ (as given in (22)) is an increasing function of ε such that $\lim_{\varepsilon \to 0} \Lambda(\varepsilon,\nu) = 0$.

(b) When \overline{F} is rotation-invariant, there exists a continuous extension $\overline{\mu}(\nu, q) : S^{n-1} \times \mathbb{R}^n \to \mathbb{R}$ of $\mu(\nu, q)$ over irrational directions $\nu \in S^{n-1} - \mathbb{R}\mathbb{Z}^n$. Moreover $\overline{\mu}$ is Lipschitz in q and C^{α} in ν , with $\alpha = \frac{1}{5n}$.

The proof is given in Theorem 4.1, Theorem 5.2 and Theorem 5.1.

A discussion on the rate of convergence $\Lambda(\varepsilon, \nu)$

Here we briefly describe the geometric process used in section 4 to obtain an upper bound for the rate function Λ in (6). Given $\delta > 0$, we are interested in finding $\varepsilon_0 = \varepsilon_0(\nu, \delta)$ such that $|u_{\varepsilon} - u| \leq C\delta$ for $\varepsilon \leq \varepsilon_0$.

If ν is rational and $\tau = 0$, F and G are periodic along ν -direction with period T_{ν} . Hence we expect that ε_0 needs to be smaller than $1/T_{\nu}$ for a fixed δ . In fact Theorem 4.1 (d) yields that

$$\Lambda(\varepsilon,
u) \leq \delta \text{ for } \varepsilon \leq \varepsilon_0 = \delta^2/T_{
u}$$

and thus yields a uniform bound

$$\Lambda(\varepsilon,\nu) \le C(\nu)\varepsilon^{1/2}.\tag{7}$$

If ν is irrational, for each δ we choose a reference rational direction P as follows: choose a point $P = P(\nu, \delta) \in \mathbb{Z}^n$ such that

$$|T\nu - P| \le \delta$$
 for some $T = T(\nu, \delta) > 0.$ (8)

Then F and G are periodic along P-direction with period $T + O(\delta)$. If we let $\theta = \theta(\nu, \delta)$ be the angle between ν and P, then (8) can be written as $\theta < \delta/T$. If $R < 1/\theta$, then due to the proximity of ν to P direction, $G(p, \cdot)$ takes only limited values of G on $H_0 \cap B_R(\tau)$, even though ν is irrational. In other words $G(p, \cdot)$ exhibits ergodicity on H_0 only in a neighborhood of size $R > 1/\theta$. For this reason u_{ε} homogenizes only when $\varepsilon \leq O(\theta)$. Indeed Theorem 4.1 (c) yields that

$$\Lambda(\varepsilon,\nu) \leq \delta$$
 for $\varepsilon \leq \varepsilon_0 = \delta^2 \theta$.

Since θ depends on not only ν but δ , we are not able to separate the dependence of the rate function on ε and ν , without further estimate of θ or T as δ varies. Such estimate would require better understanding of the *discrepancy* function discussed in the Appendix.

Proof of Theorem 1.1

Once Theorem 1.2 (a) is obtained, one can derive our main theorem by the *perturbed test function* arguments introduced by Evans [10].

Let u_{ε} solve $(P)_{\varepsilon}$, and define u^* and u_* as

$$u^* = \limsup^* u_{\varepsilon} := \lim_{r \to 0} \sup_{(y,\varepsilon) \in S_r^x} u_{\varepsilon}(y); \qquad u_* = \liminf_* u_{\varepsilon} := \lim_{r \to 0} \inf_{(y,\varepsilon) \in S_r^x} u_{\varepsilon}(y),$$

where $S_r^x = \{(y, \varepsilon) : y \in \Pi, |x - y| < r, 0 < \varepsilon < r\}$. First, observe that, by using a barrier of the form

$$\varphi_M(x) := M((x - \tau) \cdot \nu + 1) + f(x)$$

where f is a C^2 -approximation of h that is larger than h, one can conclude that $u_{\varepsilon} \leq \varphi_M$ in Π for any large M, and thus $u^* \leq h$ on H_{-1} . Similar arguments yield that $u_* \geq h$ on H_{-1} .

We claim that u^* and u_* are respectively a viscosity subsolution and supersolution of (P). If the claim is true, then Corollary 3.4 applies to yield that $u^* \leq u_*$. Since the opposite inequality is true from the definition, we conclude that $u^* = u_*$, which means that u_{ε} uniformly converges in $\overline{\Omega}$.

Below we will only show that u^* is a subsolution of (P), since the proof for u_* can be shown by parallel arguments. To this end, suppose that $u^* - \phi$ has a local max in $B_r(y_0) \cap \overline{\Pi}$ with a smooth test function ϕ . If y_0 is in the interior of Π , then $\overline{F}(D^2\phi)(y_0) \leq 0$ due to standard interior homogenization (see for instance [10]). Hence it remains to show that if y_0 is on the Neumann boundary then ϕ satisfies

$$\partial_{\nu}\phi \le \mu(\nu, q := D^T\phi) \text{ at } x = y_0.$$
 (9)

First suppose that ν is rational and $y_0 \cdot \nu = 0$. We may assume for simplicity that $u(y_0) = \phi(y_0) = 0$ and define $P(x) := D\phi(y_0) \cdot (x - y_0)$. Since $\Pi \subset \{x : x \cdot \nu < 0\}$, for any $\delta > 0$ we may choose r sufficiently small that $l_{\delta}(x) := P(x) - \delta(x \cdot \nu)$ is strictly larger than u^* on $B_r(0) \cap \Pi$. Then for sufficiently small choice of ε we have

$$l_{\delta} > u_{\varepsilon} \text{ on } B_r(0) \cap H_{-r\delta}, \quad \text{where } H_{-r\delta} = \{x \cdot \nu = -r\delta\}.$$
 (10)

Let $\bar{\varepsilon} := (r\delta)^{-1}\varepsilon$ and consider the re-scaled function $v_{\varepsilon}(x) := (r\delta)^{-1}u_{\varepsilon}(r\delta x) - l_{\delta}(x)$. Then v_{ε} is a subsolution of $(P)_{\bar{\varepsilon},\nu,0,q}$, in the local domain $\Pi \cap B_{\delta^{-1}}(0)$. Note that the corresponding Neumann boundary for v_{ε} remains to be H_0 since $y_0 \cdot \nu = 0$: in general it will be $\{(x - \tau) \cdot \nu = 0\}$ with

$$\tau = (\bar{\varepsilon})^{-1} y_0, \tag{11}$$

and thus the choice of τ must change as we vary $\bar{\varepsilon}$. We will compare v_{ε} with $w_{\bar{\varepsilon}}$, the unique bounded solution of $(P)_{\bar{\varepsilon},\nu,0,q}$ in Π obtained in Lemma 3.3. Due to the localization lemma (Lemma 3.2) we have

$$v_{\varepsilon} \le w_{\bar{\varepsilon}} + M\delta. \tag{12}$$

Due to Theorem 1.2 we have

$$w_{\bar{\varepsilon}} \leq \mu(\nu, q)(x \cdot \nu + 1) + q \cdot x + \Lambda(\bar{\varepsilon}, \nu)$$
 in Π .

Since $\Lambda(\varepsilon, \nu) \to 0$ as $\varepsilon \to 0$, (10) and (12) yield that

$$\limsup_{\varepsilon \to 0} (r\delta)^{-1} u_{\varepsilon}(r\delta x) = \limsup_{\varepsilon \to 0} v_{\varepsilon}(x) + l_{\delta}(-\nu) \le \mu(\nu, q)(x \cdot \nu + 1) + q \cdot x + l_{\delta}(-\nu) + M\delta \text{ in } \Pi.$$
(13)

Now suppose that (9) is false, then there exists $\delta > 0$ such that

$$\partial_{\nu}\phi(0) = \delta - l_{\delta}(-\nu) > \mu(\nu, q) + (M+1)\delta.$$

$$\tag{14}$$

This means that the right side of (13) is strictly negative at x = 0, which contradicts the assumption that $u^*(0) = 0$.

Next suppose that ν is irrational, we need to choose τ depending on $\bar{\varepsilon}$ so that (11) holds. Then we argue as above with a solution of $(P)_{\bar{\varepsilon},\nu,\tau,q}$ in Π . Here we must use the fact that ν is irrational and thus Theorem 1.2 ensures the uniform convergence of $w_{\bar{\varepsilon}}$ to the linear profile is regardless of the choice of τ .

2 Preliminaries

We adopt the following definition of viscosity solutions, which is equivalent to the one given in [9]. Let Ω be domain in \mathbb{R}^n with $\partial\Omega$ as a disjoint union of Γ_0 and Γ_1 . Let F satisfy (F1) - (F3) in the previous section, and let G satisfy (G3) with G(p, x) being uniformly continuous in p independent of the choice of x. For $f \in C(\Gamma_0)$ consider the following problem

	ſ	$F(D^2u, x) = 0$	in	Ω
(P)	J	u = f(x)	on	Γ_0
	l	$\tfrac{\partial}{\partial\nu}u=G(Du,x)$	on	Γ_1

where $\nu = \nu(x)$ is the outward unit normal at $x \in \Gamma_1$. Here we replace (G3) with

(G3)' (Oblicity) $|G_p \cdot \nu| \leq c < 1$ on $\partial \Omega$, where $\nu = \nu_x$ is the outward normal at $x \in \partial \Omega$.

Definition 2.1. (a) An upper semi-continuous function $u : \overline{\Omega} \to \mathbb{R}$ is a viscosity subsolution of (P)if u cannot cross from below any C^2 function ϕ which satisfies

$$\begin{cases} F(D^2\phi, x) > 0 \ in \ \Omega, \quad \phi > f \ on \ \Gamma_0, \\ \nu \cdot D\phi > G(D\phi, x) \ if \ \tau \in \Gamma_1. \end{cases}$$

(b) A lower semi-continuous function $u : \overline{\Omega} \to \mathbb{R}$ is a viscosity supersolution of (P) if if u cannot cross from above any C^2 function φ which satisfies

$$\left\{ \begin{array}{ll} F(D^2\phi,x) < 0 \ in \ \Omega, \quad \phi < f \ on \ \Gamma_0, \\ \nu \cdot D\phi < G(D\phi,x) \ on \ \Gamma_1. \end{array} \right.$$

(c) u is a viscosity solution of (P) if its upper semi-continuous envelope u^* is a viscosity subsolution and its lower semi-continuous envelope u_* is a viscosity supersolution of (P).

Existence and uniqueness of viscosity solutions of (P) are based on the comparison principle we state below. We refer to [9] and [16] for details on the proof of the following theorem as well as the well-posedness of the problem (P).

Theorem 2.2. Let G and F satisfy the conditions (G1) and (G3) and (F1) - (F3) in the previous section, with G being uniformly continuous in p independent of the choice of x. Let u and v be respectively bounded viscosity subsolution and supersolution of (P) in a bounded domain Ω . Then $u \leq v$ in Ω .

For a symmetric $n \times n$ matrix M, we decompose $M = M_+ - M_-$ with $M_{\pm} \ge 0$ and $M_+M_- = 0$. We define the Pucci operators as

$$\mathcal{P}^+(M) = -\Lambda tr(M_+) + \lambda tr(M_-)$$

and

$$\mathcal{P}^{-}(M) = -\lambda tr(M_{+}) + \Lambda tr(M_{-})$$

where $0 < \lambda < \Lambda$. Later in the paper we will utilize the fact that the difference of two solutions of $F(D^2u, x) = 0$ is both a subsolution of $\mathcal{P}^+ = 0$ and a supersolution of $\mathcal{P}^- = 0$. (see [6]).

Next we state some regularity results that will be used throughout this paper.

Theorem 2.3. [Chapter 8, [6], modified for our setting] Let u be a viscosity solution of $F(D^2u, x) = 0$ in a domain Ω . Then for any compact subset Ω' of Ω , we have

$$\|Du\|_{L^{\infty}(\Omega')} \le Cd^{-1}\|u\|_{L^{\infty}(\Omega)},$$

where $d = d(\Omega', \partial \Omega)$ and C > 0 depends on n, λ and Λ .

As mentioned in the introduction, regularity results for nonlinear Neumann problem is rather limited. $C^{0,\alpha}$ estimates has been obtained by Barles and Da Lio in general framework [3]. While a priori results for the gradient bounds are available for general F and G in [19], their results are based on linearization and thus require existence of classical solutions. For G(p, x) that is linear in p, regularity estimates on Du are recently obtained by Li and Zhang [18].

Theorem 2.4. [18], [19] Let u be a viscosity solution of (P) with $|u| \leq M$.

$$B_r^+ := \{ |x| < r \} \cap \{ x \cdot e_n \ge 0 \} \text{ and } \Gamma := \{ x \cdot e_n = 0 \} \cap B_1.$$

Let u be a viscosity solution of

$$\left\{ \begin{array}{ll} F(D^2u,x)=0 & in & B_1^+\\ \nu\cdot Du=G(Du,x) & on & \Gamma. \end{array} \right.$$

For F and G satisfying (F1) - (F3) and (G1) - (G3), suppose that either (A) F(M, x) is convex with respect to M, or (B) G(p, x) is linear with respect to p. Then for any $0 < \alpha < 1$ we have

$$\|u\|_{C^{0,\alpha}(B^+_{1/2})}, \|Du\|_{C^{0,\alpha}(B^+_{1/2})} \le C, \tag{15}$$

where C depends on α and M as well as the constants given in (F1) - (F3) and (G1) - (G3).

Our proof extends in general to the cases where the estimate (15) holds for some $\alpha > 0$.

Lastly we mention interior homogenization result from [7], which is a modified version of homogenization results such as in [11].

Theorem 2.5. (Theorem 2.14, [7]) Let K be a positive constant and let $f : \mathbb{R}^n \to \mathbb{R}$ be bounded and Hölder continuous. Given $\nu \in S^{n-1}$, let $u_N : \{-K \leq x \cdot \nu \leq 0\} \to \mathbb{R}$ be the unique bounded viscosity solution of

$$(P_N) \begin{cases} F(D^2 u_N, Nx) = 0 \text{ in } \{-K \le x \cdot \nu \le 0\}; \\ \nu \cdot Du_N = f(x) \text{ on } \{x \cdot \nu = 0\}, \quad u = 1 \text{ on } \{x \cdot \nu = -K\}. \end{cases}$$

Then for any $\delta > 0$, there exists N_0 depending only on K, the bound of u_N and the Hölder exponent of f, such that

$$|u_N - \bar{u}| \le \delta \text{ in } \{|x| \le K\} \quad \text{for } N \ge N_0, \tag{16}$$

where \bar{u} is the unique bounded viscosity solution of

$$\left\{ \begin{array}{l} \bar{F}(D^2\bar{u})=0 \ \ in \ \{-K \leq x \cdot \nu \leq 0\}; \\ \\ \nu \cdot D\bar{u}=f(x) \ \ on \ \{x \cdot \nu = 0\}, \qquad u=1 \ \ on \ \{x \cdot \nu = -K\}. \end{array} \right.$$

3 Localization Lemmas

In this section we prove several lemmas on perturbing and localizing the solutions, which will be used frequently throughout the paper. Below we prove a localization lemma, and as a corollary, we prove existence and uniqueness of solution u_{ε} of $(P)_{\varepsilon,\nu,\tau,q}$ with $\Pi = \Pi(\nu,\tau)$ for $\tau \in \mathbb{R}^n$ and $\nu \in S^{n-1}$. Denote $B_R(\tau) := \{|x - \tau| \leq R\}$ and recall $H_s := \{(x - \tau) \cdot \nu = s\}$.

First we state a basic lemma which will be frequently used. The proof is a direct consequence of the oblicity assumption (G3).

Lemma 3.1. There exists M = M(|q|, c) such that $q \cdot x \pm Mx \cdot \nu$ are respectively super and subsolution of $(P)_{\varepsilon,\nu,\tau,q}$.

Lemma 3.2. Let $f \in C(\mathbb{R}^n)$ be bounded. Suppose w_1 and w_2 solve, in the viscosity sense,

- (a) $F(D^2w_i, \frac{x}{\varepsilon}) = 0$ in $\Sigma_R := \Pi \cap B_R(0)$ for i = 1, 2
- (b) $\nu \cdot Dw_i = G(Dw_i, \frac{x}{\varepsilon})$ on H_0 for i = 1, 2
- (c) $w_1 = w_2$ on H_{-1}
- (d) $0 \le w_2 w_1 \le M$ on $\Pi \cap \partial B_R(0)$.

Let $L := \|G_p\|_{\infty}$ and 0 < c < 1 is the constant given in (G3). Then there exists a constant $C(\frac{\Lambda}{\lambda}, c, L) > 0$ such that

$$w_1 \le w_2 \le w_1 + \frac{CM}{(1-c)R}$$
 in $\Pi \cap B_1(0)$.

Proof. Without loss of generality, let us set $\nu = e_n$ and $\tau = 0$. The first inequality, $w_1 \leq w_2$, directly follows from Theorem 2.2. To show the second inequality, let

$$w := w_1 + M(h_1 + h_2) + C_1 h_3,$$

where

$$h_1(x) = \frac{|x|^2}{R^2}, \quad h_2(x) = \frac{C}{R^2}(1 - (x_n)^2) \text{ with } C = \frac{n\Lambda}{\lambda}, \quad h_3(x) = \frac{1 + x_n}{R}$$

and $C_1 > 0$ is a large constant depending on n, Λ , λ , L and c, which will be chosen below in the proof. Note that in Σ_R ,

$$F(D^2w, \frac{x}{\varepsilon}) = F(D^2w_1 + M(D^2h_1 + D^2h_2), \frac{x}{\varepsilon})$$

$$\geq F(D^2w_1, \frac{x}{\varepsilon}) - \mathcal{P}^+(M(D^2h_1 + D^2h_2))$$

$$= F(D^2w_1, \frac{x}{\varepsilon}) = 0.$$

Also $w_2 = w_1 \leq w$ on H_{-1} and $w_2 \leq w_1 + M \leq w$ on $\partial B_R(0) \cap \Pi$.

Hence to show that $w_2 \leq w$, it is enough to show that $\partial_{x_n} w \geq G(Dw, \frac{x}{\varepsilon})$ on H_0 . We will verify that this is true when C_1 is sufficiently large. Observe that in Σ_R

$$|D(h_1 + h_2)| \le \frac{C_0}{R}$$
 for $C_0 = C_0(n, \Lambda, \lambda).$ (17)

Hence on $H_0 \cap \Sigma_R$ we have

$$\begin{array}{lcl} \partial_{x_n}w & \geq & \partial_{x_n}w_1 + \frac{C_1}{R} - \frac{C_0}{R} \\ \\ & = & G(Dw_1, \frac{x}{\varepsilon}) + \frac{(C_1 - C_0)}{R} \\ \\ & \geq & G(Dw, \frac{x}{\varepsilon}) - \frac{cC_1}{R} + \frac{C_0L}{R} + \frac{(C_1 - C_0)}{R} \end{array}$$

where the last inequality follows from the Lipschitz property of G with (17), if $C_1 = C_1(n, \Lambda, \lambda, c)$ is chosen sufficiently large. It follows from Theorem 2.2 that $w_2 \leq w$ in Σ_R , and we obtain the lemma. \Box

As a corollary of Lemma 3.2, we prove existence and uniqueness of solutions in strip regions.

Lemma 3.3. There exists a unique solution u_{ε} of $(P)_{\varepsilon,\nu,\tau,q}$ such that

$$\|u_{\varepsilon} - q \cdot x\| \le M.$$

Proof. 1. Let Σ_R be as given in Lemma 3.2, and consider the viscosity solution $w_R(x)$ of $(P)_{\varepsilon,\nu,\tau,q}$ in Σ_R with the lateral boundary data $q \cdot x$ on $\partial B_R(\tau) \cap \Pi$. The existence and uniqueness of the viscosity solution w_R is shown, for example, in [9] and [16].

From Lemma 3.1, $q \cdot x \pm M(x - \tau + \nu) \cdot \nu$ is respectively a sub and suppressive of $(P)_{\varepsilon,\nu,\tau,q}$, and thus by comparison principle we obtain that

$$|w_R(x) - q \cdot x| \leq M$$
 for $x \in \Sigma_R$.

Due to Theorem 2.5 and the Arzela-Ascoli Theorem, w_R locally uniformly converges to a continuous function $u_{\varepsilon}(x)$. From the stability property of viscosity solutions it follows that $u_{\varepsilon}(x)$ is a viscosity solution of $(P)_{\varepsilon,\nu,\tau,q}$.

2. To show uniqueness, suppose u_1 and u_2 are both viscosity solutions of $(P)_{\varepsilon,\nu,\tau,q}$ with $|u_1 - q \cdot x|$, $|u_2 - q \cdot x| \leq M$. Then Lemma 3.2 yields that, for any point $s \in H_0$

$$|u_1 - u_2| \le O(1/R)$$
 in $B_1(s) \cap \Pi$.

Hence $u_1 = u_2$.

The following is immediate from Theorem 2.2 and the construction of u_{ε} in the above lemma.

Corollary 3.4. Suppose u, v are bounded and continuous functions in $\overline{\Pi} = \overline{\Pi(\tau, \nu)}$. In addition suppose they satisfy, for F satisfying (F1) - (F3) and G satisfying $(G_1) - (G_2)$,

(a)
$$F(D^2u, \frac{x}{\varepsilon}) \leq 0 \leq F(D^2v, \frac{x}{\varepsilon})$$
 in Π ;
(b) $u \leq v$ on H_{-1} ;
(c) $\nu \cdot Du \leq G(Du, x/\varepsilon)$; $\nu \cdot Dv \geq G(Dv, x/\varepsilon)$ on H_0 .
Then $u < v$ in Π .

Lemma 3.5. There exists C > 0 such that the following holds: let u_1 and u_2 be solutions of

$$\begin{cases} \mathcal{P}^+(D^2u_1) \le 0, \quad \mathcal{P}^-(D^2u_2) \ge 0 & in \quad \Pi \cap B_R(0) \\\\ \partial_\nu u_i = G_i(Du_i, x) & on \quad H_0 \cap B_R(0) \\\\ u_i = q \cdot x & on \quad H_{-1} \cap B_R(0) \end{cases}$$

where $\Pi = \Pi(\nu, 0)$. Furthermore suppose that G_i satisfies the assumption in Theorem 2.4 and G_1 and G_2 satisfy

$$|G_1(p,x) - G_2(p,x)| \le \delta(1+|p|) \text{ and } |u_1 - u_2| \le M.$$
(18)

Let L denote the Lipschitz bound for u_i and $G'_i s$. Then there exists $C = C(\Lambda, \lambda, n)$ such that

$$|u_1 - u_2| \le \delta(L+1) + CM/R \text{ in } \Pi \cap B_1(0)$$

Proof. By our assumption, $v := (u_1 - u_2)/M$ satisfies $|v| \le 1$ in $B_R(0)$ with

$$\begin{cases} \mathcal{P}^+(D^2v) \le 0 & \text{in} \quad \Pi \cap B_R(0) \\ v = 0 & \text{on} \quad H_{-1} \cap B_R(0) \end{cases}$$

After a change of coordinates we may assume $\nu = e_n$ so that $\Pi = \{x : -1 \le x_n \le 0\}$, and we denote $x = (x', x_n)$. Define

$$w(x) := (c_0/M + c_1/R)(x_n + 1) + 2(|x'|^2 - \frac{\Lambda n}{\lambda}(|x_n|^2 - 1))/R^2.$$

where c_0 and $c_1 > 8$ will be chosen later. Then w is a supersolution of above problem with the Neumann boundary condition

$$\partial_n w = (c_0/M + c_1/R) \ge (c_0/M + 4|x'|/R^2) = (c_0/M + |D_Tw|)$$
 on $\{x_n = 0\} \cap B_R(0)$

Now suppose v - w has positive maximum in $\Pi \cap \overline{B_R}(0)$. Then the maximum would need to be achieved at a point $\tau \in H_0 \cap B_R(0)$. At this point we should have $\partial_n(v - w) \ge 0$ and $D_T v = D_T w$. Therefore

$$\partial_n v \ge \partial_n w \ge (c_0/M + |D_T w|) = (c_0/M + |D_T v|) \text{ at } x = \tau,$$
(19)

On the other hand

$$G_1(Du_1, x) - G_2(Du_2, x) = DG_1(p^*, x) \cdot D(Mv) + G_1(Du_2, x) - G_2(Du_2, x) + G_2(Du_2, x) - G_2(Du_2, x$$

and since $|DG_1(p^*, x) \cdot e_n| \leq c$ we have, from (18) and the Lipschitz bound for u_i given in Theorem 2.4,

$$(1-c)\partial_n v \le L|D_T v| + \frac{1}{M}|G_1(Du_2, x) - G_2(Du_2, x)| \le L|D_T v| + \frac{\delta}{M}(L+1)$$
 at $x = \tau$.

Then using the fact that $|D_T w| = 4|x'|/R^2 \le 4/R$ in $B_R(0)$ it follows that

$$(1-c)|\partial_n v| \le \frac{4L}{R} + \frac{\delta(L+1)}{M}.$$
(20)

Hence from (19) we get a contradiction if $c_0/M + c_1/R$ is larger than the right handside of (30). This happens if we choose $c_1 > 4L$ and $c_0 = \delta(L+1)$. Therefore it follows that $v \leq w$ in $\Pi \cap \overline{B_R}$. We can now conclude that

$$u_1 - u_2 = Mv \le c_0 + c_1 M/R + 2M(1 + \frac{\Lambda n}{\lambda})/R^2$$
 in $\Pi \cap B_1(0)$.

The lower bound can be obtained with above argument applied to $u_2 - u_1$.

4 Homogenization in a Strip Domain

Let u_{ε} solve $(P)_{\varepsilon,\nu,\tau,q}$ given in Section 1.1. Then there is a unique linear function v_{ε} such that $v_{\varepsilon} = u_{\varepsilon}$ on $H_{-1} \cup \{z_0\}$, where $z_0 := \tau - \nu/2$ is a fixed reference point in Π . We define the average slope $\mu(u_{\varepsilon})$ of u_{ε} as follows

$$\mu(u_{\varepsilon}) := \partial_{\nu} v_{\varepsilon}. \tag{21}$$

Theorem 4.1. The followings hold for u_{ε} solving $(P)_{\varepsilon,\nu,\tau,q}$:

(a) For irrational directions ν , there exists a unique constant $\mu = \mu(\nu, q)$ such that u_{ε} converges uniformly to the linear profile

$$u(x) := \mu((x - \tau) \cdot \nu + 1) + l(x).$$

The same holds for rational directions ν with $\tau = 0$.

(b) [Error estimate] There exists a constant C > 0 depending on λ , Λ , n, and the slope of l(x) such that the following holds: if ν is an irrational direction or ν is a rational direction with $\tau = 0$, then

$$|\mu(u_{\varepsilon}) - \mu| \le C\Lambda(\varepsilon, \nu) \quad in \quad \Pi,$$

where

$$\Lambda(\varepsilon,\nu) = \begin{cases} \inf_{\substack{0 < k < 1}} \{\varepsilon^k T_\nu + \varepsilon^{1-k}\} & \text{if } \nu \text{ is a rational direction} \\ \inf_{\substack{0 < k < 1, N \in \mathbb{N}}} \{\varepsilon^k N + \omega_\nu(N) + \varepsilon^{1-k}\} & \text{if } \nu \text{ is an irrational direction.} \end{cases}$$
(22)

In (22), T_{ν} is as given in (a) of Lemma A.3 (which is the period of G(P, y) on the Neumann boundary H_0) and $\omega_{\nu}(N)$ is as given in (78) with $\omega_{\nu}(N) \to 0$ as $N \to \infty$.

(c) Let ν be an irrational direction. For any $\delta > 0$, there exist T > 0 and $P \in \mathbb{Z}^n$ such that

$$|T\nu - P| \le \delta.$$

Let $\theta = \theta(\delta, \nu)$ be the angle between ν and P, then

$$\Lambda(\varepsilon,\nu) \leq 3\delta \quad for \ \varepsilon < \delta^2\theta.$$

(d) Let ν be a rational direction, and let $\delta > 0$. Then

$$\Lambda(\varepsilon,\nu) \leq 2\delta \quad for \ \ \varepsilon < \frac{\delta^2}{T_\nu}.$$

To prove Theorem 4.1 we begin with a preliminary lemma. The following lemma states that u_{ε} looks like a linear profile (almost flat) on each hyperplane normal to ν .

Lemma 4.2. Away from the Neumann boundary H_0 , $u_{\varepsilon} - l(x)$ is almost a constant on hyperplanes parallel to H_0 . More precisely, for $x_0 \in \Pi$ we denote

$$d := \operatorname{dist}(x_0, H_0) > 0$$

and $H_d := \{(x - \tau) \cdot \nu = d\} = \{(x - x_0) \cdot \nu = 0\}$. Then the followings hold:

(a) If ν is a rational direction, there exists a constant C > 0 depending on α , λ , Λ , n, and the slope of l such that for any $x \in H_d$

$$|(u_{\varepsilon}(x) - l(x)) - (u_{\varepsilon}(x_0) - l(x_0))| \le C(d^{-1} + 1)(T_{\nu}\varepsilon).$$
(23)

where T_{ν} is a constant depending on ν , given as in (a) of Lemma A.3.

(b) If ν is an irrational direction, there exists a constant C > 0 depending on α , λ , Λ , n and the slope of l such that for any $x \in H_d$

$$|(u_{\varepsilon}(x) - l(x)) - (u_{\varepsilon}(x_0) - l(x_0))| \le C(d^{-1}\varepsilon(MN + \omega_{\nu}(N)) + \omega_{\nu}(N))$$
(24)

for any $N \in \mathbb{N}$ and $\varepsilon > 0$ with $\varepsilon(MN + \omega_{\nu}(N)) < 1$, where M is a dimensional constant given as in (b) of Lemma A.3, and $\omega_{\nu}(N)$ is given as in (78).

Proof. First, we consider a rational direction ν . By (a) of Lemma A.3, for any $x \in H_d$, there is $y \in H_d$ such that $|x - y| \leq T_{\nu}\varepsilon$ and $y = x_0 \mod \varepsilon \mathbb{Z}^n$. Then by comparison

$$u_{\varepsilon}(x) = u_{\varepsilon}(x + (y - x_0)) - l(y) + l(x_0).$$
(25)

Hence $u_{\varepsilon}(x_0) = u_{\varepsilon}(y) - l(y) + l(x_0)$ and we get

$$\begin{aligned} |(u_{\varepsilon}(x) - l(x)) - (u_{\varepsilon}(x_{0}) - l(x_{0}))| &\leq |u_{\varepsilon}(x) - u_{\varepsilon}(y)| + |l(y) - l(x)| \\ &\leq |u_{\varepsilon}(x) - u_{\varepsilon}(y)| + CT_{\nu}\varepsilon \\ &\leq Cd^{-1}T_{\nu}\varepsilon + CT_{\nu}\varepsilon \end{aligned}$$

where the third inequality follows from Theorem 2.3.

Next, we consider an irrational direction ν and let $x \in H_d$. By (b) of Lemma A.3, for any $N \in \mathbb{N}$, there exists $y \in \mathbb{R}^n$ such that $|x - y| \leq \varepsilon (MN + \omega_{\nu}(N)), y = x_0 \mod \varepsilon \mathbb{Z}^n$ and

$$\operatorname{dist}(y, H_d) < \varepsilon \omega_{\nu}(N). \tag{26}$$

Observe that

$$|(u_{\varepsilon}(x) - l(x)) - (u_{\varepsilon}(x_0) - l(x_0))| \le |u_{\varepsilon}(x) - u_{\varepsilon}(y)| + |(u_{\varepsilon}(y) - l(y)) - (u_{\varepsilon}(x_0) - l(x_0))| + |l(y) - l(x)|$$

where, from Theorem 2.3,

$$|u_{\varepsilon}(x) - u_{\varepsilon}(y)| \le Cd^{-1}\varepsilon(MN + \omega_{\nu}(N)).$$

Next we project y to $x_1 \in H_d$ and use Lemma 3.5 for $G_1 = G$ and $G_2(p, x) = G_2(p, x + (x_0 - x_1)) = G_2(p, x + (y - x_1))$ with $\delta = \omega_{\nu}(N)$ to conclude that

$$|(u_{\varepsilon}(x_0) - l(x_0)) - (u_{\varepsilon}(x_1) - l(x_1))| \le C\omega_{\nu}(N).$$

and then once again use Theorem 2.3 with (26) to compare u(y) with $u(x_1)$ to conclude that

$$|(u_{\varepsilon}(y) - l(y)) - (u_{\varepsilon}(x_0) - l(x_0))| \le C(\omega_{\nu}(N) + \varepsilon)$$

and lastly

$$|l(y) - l(x)| \le C|y - x| \le C\varepsilon(MN + \omega_{\nu}(N)) \le Cd^{-1}\varepsilon(MN + \omega_{\nu}(N))$$

where the last inequality follows if $\varepsilon(MN + \omega_{\nu}(N)) < 1$.

Since u_{ε} is flat on each hyperplanes located a constant *d*-away from the Neumann boundary, u_{ε} can be approximated well by a linear solution as in the following corollary. The proof of Corollary 4.3 follows from the comparison principle (Theorem 2.2) and Lemma 4.2 with $d = \varepsilon^{1-k}$.

Corollary 4.3. For a solution u_{ε} of $(P)_{\varepsilon,\nu,\tau,q}$, let v_{ε} be the unique linear function given as in (21). Then there exists a constant C depending on λ , Λ , n and the slope of l such that for any $N \in \mathbb{N}$ and 0 < k < 1,

$$|u_{\varepsilon}(x) - v_{\varepsilon}(x)| \leq \begin{cases} C(\varepsilon^{k}T_{\nu} + \varepsilon^{1-k}) & \text{if } \nu \text{ is a rational direction} \\ C(\varepsilon^{k}N + \omega_{\nu}(N) + \varepsilon^{1-k}) & \text{if } \nu \text{ is an irrational direction} \end{cases}$$

and hence

 $|u_{\varepsilon}(x) - v_{\varepsilon}(x)| \le C\Lambda(\varepsilon, \nu).$

Due to the uniform interior regularity of $\{u_{\varepsilon}\}$ (Theorem 2.3), along a subsequence they locally uniformly converges to u in Π . Let us choose one of the convergent subsequence u_{ε_j} and denote it by u_j , i.e., $u_j = u_{\varepsilon_j}$. Let $v_j = v_{\varepsilon_j}$ and $\mu_j = \mu(u_{\varepsilon_j})$, both as given in (21).

Corollary 4.3 implies that for any $\nu \in S^{n-1}$, $\lim u_j$ is linear. More precisely, the slope μ_j converges as $j \to \infty$ (see Lemma 4.1 of [8]), and hence by Corollary 4.3

$$\lim u_{j} = \lim v_{j} = \mu((x - \tau) \cdot \nu + 1) + l(x) = u$$

for $\mu := \lim \mu_j$.

Next, we prove that the subsequential limit is unique, i.e., μ does not depend on the subsequence $\{\varepsilon_j\}$ when ν is irrational or ν is rational with $\tau = 0$. We will also obtain a mode of convergence of μ_{ε} .

Proof of Theorem 4.1 (a) and (b) for irrational directions:

Let ν be an irrational direction and let u be a subsequential limit of u_{ε} . We claim that

$$\partial u/\partial \nu = \mu(\nu,q)$$

for a constant $\mu(\nu, q)$ which depends on ν and q, not on τ or the subsequence $\{\varepsilon_i\}$. More precisely,

$$|\mu(u_{\eta}) - \mu(u_{\varepsilon})| \le C(\Lambda(\varepsilon, \nu) + \eta).$$
(27)

For the proof of (27), let $0 < \eta < \varepsilon$ be sufficiently small. Let

$$w_{\varepsilon}(x) = \frac{u_{\varepsilon}(\varepsilon x)}{\varepsilon}, \quad w_{\eta}(x) = \frac{u_{\eta}(\eta x)}{\eta}$$

and denote by H^1 and H^2 , the corresponding Neumann boundaries of w_{ε} and w_{η} , respectively. By (c) of Lemma A.3, for $\tau \in \mathbb{R}^n$, there exist $s_1 \in H^1$ and $s_2 \in H^2$ such that

$$|\tau - s_1| \le \eta \mod \mathbb{Z}^n$$
, and $|\tau - s_2| \le \eta \mod \mathbb{Z}^n$.

Hence after translations by $\tau - s_1$ and $\tau - s_2$, we may suppose that $w_{\varepsilon}(x)$ and $w_{\eta}(x)$ are defined on the extended strips

$$\Omega_{\varepsilon} := \{ x : -\frac{1}{\varepsilon} \le (x - \tau) \cdot \nu \le 0 \} \quad \text{and} \quad \Omega_{\eta} := \{ x : -\frac{1}{\eta} \le (x - \tau) \cdot \nu \le 0 \}$$

respectively, with

$$w_{\varepsilon} = l_{\varepsilon}(x)$$
 on $\{(x - \tau) \cdot \nu = -\frac{1}{\varepsilon}\}$

and

$$w_{\eta} = l_{\eta}(x)$$
 on $\{(x - \tau) \cdot \nu = -\frac{1}{\eta}\}$

where l_{ε} and l_{η} are linear functions with the same slope as l(x). Moreover on H_0 we have

$$\partial w_{\varepsilon}/\partial \nu = G(Dw_{\varepsilon}, x - z_1)$$
 and $\partial w_{\eta}/\partial \nu = G(Dw_{\eta}, x - z_2)$

for some $|z_1|, |z_2| \leq \eta$. Observe that by Hölder continuity of G, i.e., by (G2)

$$|G(p, x - z_1) - G(p, x - z_2)| < m(1 + |p|)\eta$$
(28)

Let v_{ε} be given in (21). Then by Corollarly 4.3 (after a translation),

$$|w_{\varepsilon}(x) - \frac{v_{\varepsilon}(\varepsilon x)}{\varepsilon}| \le \frac{C\Lambda(\varepsilon, \nu)}{\varepsilon}.$$
(29)

Note that

$$\frac{v_{\varepsilon}(\varepsilon x)}{\varepsilon} = \mu_{\varepsilon}((x-\tau) \cdot \nu + \frac{1}{\varepsilon})) + l_{\varepsilon}(x).$$

From (29) and the comparison principle, it follows that

$$(\mu_{\varepsilon} - C\Lambda(\varepsilon, \nu))((x - \tau) \cdot \nu + \frac{1}{\varepsilon}) \le w_{\varepsilon}(x) - l_{\varepsilon}(x) \le (\mu_{\varepsilon} + C\Lambda(\varepsilon, \nu))((x - \tau) \cdot \nu + \frac{1}{\varepsilon}).$$
(30)

Here we denote by l_1 and l_2 , the following linear profiles

$$l_1(x) = a_1(x - \tau) \cdot \nu + b_1$$
 and $l_2(x) = a_2(x - \tau) \cdot \nu + b_2$,

whose respective slopes are $a_1 = \mu_{\varepsilon} + C\Lambda(\varepsilon, \nu)$ and $a_2 = \mu_{\varepsilon} - C\Lambda(\varepsilon, \nu)$. b_1 and b_2 are chosen so that

$$l_1(x) = l_2(x) = \omega_\eta(x) - l_\eta(x) = 0 \text{ on } \{x : (x - \tau) \cdot \nu = -\frac{1}{\eta}\}.$$
(31)

Now we define

$$\overline{w}(x) := l_{\eta}(x) + \begin{cases} l_1(x) & \text{in} \quad \{-1/\eta \le (x-\tau) \cdot \nu \le -1/\varepsilon\} \\ w_{\varepsilon}(x) - l_{\varepsilon}(x) + c_1 & \text{in} \quad \{-1/\varepsilon \le (x-\tau) \cdot \nu \le 0\} \end{cases}$$

and

$$\underline{w}(x) := l_{\eta}(x) + \begin{cases} l_2(x) & \text{in} \quad \{-1/\eta \le (x-\tau) \cdot \nu \le -1/\varepsilon\} \\ w_{\varepsilon}(x) - l_{\varepsilon}(x) + c_2 & \text{in} \quad \{-1/\varepsilon \le (x-\tau) \cdot \nu \le 0\} \end{cases}$$

where c_1 and c_2 are constants satisfying

$$l_1 = w_{\varepsilon} - l_{\varepsilon} + c_1 = c_1$$
 and $l_2 = w_{\varepsilon} - l_{\varepsilon} + c_2 = c_2$

on $\{(x - \tau) \cdot \nu = -1/\varepsilon\}$. (See Figure 2.) Note that by (31),

$$\underline{w} = \overline{w} = w_{\eta}$$
 on $\{x : (x - \tau) \cdot \nu = -\frac{1}{\eta}\}$

and also due to (30),

$$\overline{w}(x) = l_{\eta}(x) + \min(l_1(x), w_{\varepsilon}(x) - l_{\varepsilon}(x) + c_1)$$

and

$$\underline{w}(x) = l_{\eta}(x) + \max(l_2(x), w_{\varepsilon}(x) - l_{\varepsilon}(x) + c_2)$$

in $\{-\frac{1}{\varepsilon} \leq (x-\tau) \cdot \nu \leq 0\}$. Thus it follows that \overline{w} and \underline{w} are respectively viscosity super- and subsolution of (P). Hence we obtain

$$\underline{w} \le \tilde{w}_{\eta} \le \overline{w} \tag{32}$$

where \tilde{w}_{η} is a solution of (P) in Ω_{η} with $\tilde{w}_{\eta} = w_{\eta} = l_{\eta}(x)$ on $\{(x - \tau) \cdot \nu = -1/\eta\}$, and $\partial \tilde{w}_{\eta}/\partial \nu = G(D\tilde{w}_{\eta}, x - z_1)$ on H_0 . Then by (32) and Lemma 3.5 with (28),

$$|\mu_{\eta} - \mu_{\varepsilon}| \le |\mu_{\eta} - \mu(\tilde{w}_{\eta})| + |\mu(\tilde{w}_{\eta}) - \mu_{\varepsilon}| \le C(\Lambda(\varepsilon, \nu) + \eta)$$

where $\mu(\tilde{w}_{\eta})$ is the slope of the linear approximation of \tilde{w}_{ε} . The above inequality implies that the slope μ of a subsequential limit of u_{ε} depends on neither the subsequence $\{\varepsilon_j\}$ nor τ . Also sending $\eta \to 0$, we get an error estimate (d) when ν is irrational.

Proof of Theorem 4.1 (a) and (b) for rational directions: Let ν be a rational direction with $\tau = 0$. We claim that $\partial u/\partial \nu = \mu(\nu, q)$ for a constant $\mu(\nu, q)$ which depends on ν and q, not on the subsequence $\{\varepsilon_j\}$. More precisely, if $\eta \leq \varepsilon$, then

$$|\mu(u_{\eta}) - \mu(u_{\varepsilon})| \le C\Lambda(\varepsilon, \nu). \tag{33}$$

The proof of (33) is parallel to that of (27). Let w_{ε} and w_{η} be as given in the proof of (27). Note that since Ω_{ε} and Ω_{η} have their Neumann boundaries passing through the origin, $\partial w_{\varepsilon}/\partial \nu = G(x) = \partial w_{\eta}/\partial \nu$ without translation of the x variable, and thus we do not need to use the properties of hyperplanes with an irrational normal (Lemma A.3 (b)) to estimate the error between the shifted Neumann boundary datas. In other words, there exist $q_1 \in H^1$ and $q_2 \in H^2$ such that $p = q_1 = q_2 \mod \mathbb{Z}^n$, hence $G(\cdot, x - z_1) = G(\cdot, x - z_2)$ in the proof of (27). Following the proof of (27), we get an upper bound $\Lambda(\varepsilon, k)$ of $|\mu_{\eta} - \mu_{\varepsilon}|$. Note that we do not have the term η in (33) since $G(\cdot, x - z_1) = G(\cdot, x - z_2)$. Sending $\eta \to 0$ in (33), we obtain the error estimate (b) for rational directions with $\tau = 0$.

Proof of Theorem 4.1 (c) and (d): Let $\delta > 0$ and let ν be an irrational direction. Lemma A.4 implies that there is a positive number $T_{\nu}(\delta) \leq \delta^{-(n-1)}$ such that $|T_{\nu}(\delta)\nu| \leq \delta \mod \mathbb{Z}^n$. Then for some $P \in \mathbb{Z}^n$ and $T = T_{\nu}(\delta) + O(\delta)$

$$|T\nu - P| \le \delta$$

and $T\nu \in P + \langle \vec{P} \rangle^{\perp}$. Let $\theta = \theta(\delta, \nu) > 0$ be the angle between ν and \vec{P} , then

$$|T\nu - P| = T\theta \le \delta \tag{34}$$

and we can observe the set $\{mT\nu \mid 0 \leq m \leq [\frac{1}{T\theta}]\}$ is evenly distributed mod \mathbb{Z}^n . Hence we get

$$\omega_{\nu}(N) \le T\theta \text{ when } N = \left[\frac{1}{\theta}\right].$$
 (35)

Let $\varepsilon(\delta, \nu)$ be a constant depending on δ and the direction ν such that

$$\varepsilon(\delta,\nu) = \delta^2 \theta = \delta^2 \theta(\delta,\nu). \tag{36}$$

Then for $0 < \varepsilon < \varepsilon(\delta, \nu)$,

$$\Lambda(\varepsilon,\nu) = \inf_{0 < k < 1, N \in \mathbb{N}} \{\varepsilon^k N + \omega_\nu(N) + \varepsilon^{1-k}\} \le \inf_{0 < k < 1} \{\varepsilon^k / \theta + T\theta + \varepsilon^{1-k}\} \le \inf_{0 < k < 1} \{\varepsilon^k / \theta + \varepsilon^{1-k}\} + \delta$$

where the first and last inequalities follow from (35) and (34) respectively. Then by (36)

$$\inf_{0 < k < 1} \{ \varepsilon^k / \theta + \varepsilon^{1-k} \} \le \inf_{0 < k < 1} \{ (\delta^2 \theta)^k / \theta + (\delta^2 \theta)^{1-k} \}.$$

The infimum is taken when $0 < k = \ln(\theta \delta) / \ln(\theta \delta^2) < 1$ and

$$\inf_{0 < k < 1} \{ (\delta^2 \theta)^k / \theta + (\delta^2 \theta)^{1-k} \} = 2\delta.$$

Hence we can conclude $\Lambda(\varepsilon, \nu) \leq 3\delta$ for $\varepsilon < \varepsilon(\delta, \nu) = \delta^2 \theta$.

Next, we consider a rational direction ν . For $\delta > 0$, let $\varepsilon < \delta^2/T_{\nu}$. Then we can check

$$\Lambda(\varepsilon,\nu) = \inf_{0 < k < 1} \{ \varepsilon^k T_\nu + \varepsilon^{1-k} \} \le \inf_{0 < k < 1} \{ \delta^{2k} T_\nu^{1-k} + \delta^{2(1-k)} T_\nu^{k-1} \} = 2\delta.$$

The following lemma will be used in the next section.

Lemma 4.4. Let $\nu = e_n$, $\tau = 0$, and let w solve

$$\begin{cases} F(D^2w, x/\varepsilon) = 0 & in \quad \{-N\varepsilon \le x_n \le 0\};\\ \partial w/\partial x_n = G(Dw, x/\varepsilon) & on \quad H_0;\\ w = A & on \quad H_{-N\varepsilon}. \end{cases}$$

where N and A are constants. Then there is a constant $C = C(\lambda, \Lambda, n)$ such that

$$|w(x) - w(x_0)| \le C\varepsilon$$
 for $x, x_0 \in H_{-N\varepsilon}$.

Proof. For $x_0, x \in H_{-N\varepsilon}$, choose $y \in H_{-N\varepsilon}$ such that $|x - y| \leq \varepsilon$ and $y = x_0 \mod \varepsilon \mathbb{Z}^n$. Observe that then $w(y) = w(x_0)$, since G is 1-periodic on H_0 . Therefore

$$|w(x) - w(x_0))| = |w(x) - w(y)| \le C ||w - A||_{L^{\infty}} |\frac{x - y}{N\varepsilon}| \le C\varepsilon,$$

where the second inequality is from the interior Lipschitz regularity (Theorem 2.3) applied to $w(N\varepsilon x)$.

5 Continuity over normal directions

In the previous section we have shown that for an irrational direction $\nu \in S^{n-1} - \mathbb{R}\mathbb{Z}^n$, there is a unique homogenized slope $\mu(\nu, q)$ for any solution u_{ε}^{ν} of $(P)_{\varepsilon,\nu,\tau,q}$ in $\Pi(\nu,\tau)$. In this section we investigate the continuity properties of μ with respect to ν and q, as well as the mode of convergence for u_{ε}^{ν} as the normal direction ν of the domain varies.

We first show that μ is Lipschitz with respect to q, which directly follows from the 1- homogeneity of G.

Theorem 5.1. For $\nu \in S^{n-1} - \mathbb{R}\mathbb{Z}^n$, $\mu(\nu, q)$ is uniformly Lipschitz in $q \in \langle \nu \rangle^{\perp}$, independent of ν .

Proof. For $q_1, q_2 \in \langle \nu \rangle^{\perp}$, let u_{ε}^i be the unique bounded solution of $(P)_{\varepsilon,\nu,\tau,q_i}$ for i = 1, 2. Let m be the Lipschitz constant for G given in (G1) and c be as given in (G3). Then it follows that

$$w_{\pm}(x) := u_{\varepsilon}^{1}(x) + (q_{2} - q_{1}) \cdot x \pm \frac{m}{1 - c} |q_{1} - q_{2}| (x \cdot \nu)$$

is respectively a super and subsolution of $(P)_{\varepsilon,\nu,\tau,q_2}$. Hence by Corollary 3.4 we have

$$w_{-} \le u_{\varepsilon}^2 \le w_{+}.$$

From here and Theorem 4.1 it follows that

$$|\mu(\nu, q_1) - \mu(\nu, q_2)| \le \frac{m}{1-c}|q_1 - q_2|.$$

г		
L		

The dependence of μ on ν is a much more subtle matter due to the change of the domain and the resulting changes in boundary conditions on the Neumann boundary. From now on we work with a fixed choice of q and denote $\mu = \mu(\nu)$.

For $s \ge 0$, let $T_{\nu}(s)$ be the smallest positive number ≥ 1 such that

$$|T_{\nu}(s)\nu| \leq s \mod \mathbb{Z}^n.$$

Note that with this definition T_{ν} given in the Appendix corresponds to $T_{\nu}(0)$ which is larger than all $T_{\nu}(s)$. In general Lemma A.4 yields

$$T_{\nu}(s) \le \sqrt{n} \cdot s^{-(n-1)}. \tag{37}$$

Theorem 5.2. With fixed q, let us denote $\mu = \mu(\cdot, q) : (S^{n-1} - \mathbb{R}\mathbb{Z}^n) \to \mathbb{R}$ be as given in Theorem 4.1. Then μ has a continuous extension $\overline{\mu}(\nu) : S^{n-1} \to \mathbb{R}$. More precisely, let us fix a direction $\nu \in S^{n-1}$ and a constant $\delta > 0$. If ν_1 and ν_2 are irrational directions such that

$$0 < \theta_i := |\nu_i - \nu| < \frac{\delta^{5/2}}{T_\nu(\delta^{5/2})} \quad for \ i = 1, 2$$
(38)

then we have

- (a) $|\mu(\nu_1) \mu(\nu_2)| < C\delta^{1/2}$.
- (b) $\bar{\mu}(\nu)$ is Hölder continuous on S^{n-1} with a Hölder exponent of $\frac{1}{5n}$.

Remarks 5.3. In the proof we indeed show that, for any directions ν_1 and ν_2 satisfying (38), the range of $\{\mu(u_{\varepsilon}^{\nu_i})\}_{\varepsilon,i}$ fluctuates only by δ , if ε is sufficiently small. The fact that ν_i 's are irrational is only used to guarantee that there is only one subsequential limit for $\mu(u_{\varepsilon}^{\nu_i})$.

For the rest of the paper we prove (a) of Theorem 5.2. Theorem 5.2 (b) follows from (37), (38) and Theorem 5.2 (a).

5.1 Basic settings and Sketch of the proof

For notational simplicity and clarity in the proof, we assume that n = 2 and $\nu = e_2$. We will explain in the paragraph below how to modify the notations and the proof for $\nu \neq e_2$. For general dimension n, we refer to Remark 5.11. We denote

$$\Pi := \Pi(e_2, 0)$$
 and $\Pi^{\nu_i} := \Pi(\nu_i, 0)$, for $i = 1, 2$.

We also denote

$$H_0 = H_0(e_n), \quad H_0^{\nu_i} := H_0(\nu^i) \quad \text{for } i = 1, 2$$

For given

$$m \in \mathbb{N}$$
 and $\delta := 1/m > 0$,

we divide the unit strip $\mathbb{R} \times [0,1]$ by *m* number of small horizontal strips of width δ and define a family of functions $\{G_k\}_k$ so that the value of G_k at (x_1, x_2) is same as the value of G at (x_1, \tilde{x}_2) , where (x_1, \tilde{x}_2) is the projection of (x_1, x_2) onto the bottom of the k-th strip. More precisely we define

$$G_k(x_1, x_2) := G(x_1, \delta(k-1)) \text{ for } k = 1, ..., m.$$
(39)

Then G_k is a 1-periodic function with respect to x_1 .

Next we introduce the parameters

$$\theta_1 := |\nu_1 - e_2|, \ \theta_2 := |\nu_2 - e_2| \tag{40}$$

and

$$N := \left[\frac{\delta}{\theta_1}\right], \quad M := \left[\frac{\delta}{\theta_2}\right]. \tag{41}$$

Without loss of generality, assume $\theta_2 \leq \theta_1$ and thus $N \leq M$.

If θ_i 's are sufficiently small, then we will be able to approximate G on both of the Neumann boundary $H_0^{\nu_1}$ and $H_0^{\nu_2}$ using the universal boundary data G_k 's which depends only on δ , but not on the direction ν_1 nor ν_2 . In particular, in meso-scopic scale G can be approximated by many repeating pieces (N for ν_1 and M for ν_2) of G_k , for a suitable $1 \leq k \leq m$ on each pieces of $H_0^{\nu_i}$. Thus the problem already experiences averaging phenomena: we call this as the first or near-boundary homogenization. Note that in this step the only difference in the averaging phenomena between the two directions ν_1 and ν_2 , besides the errors in terms of G and G_k on $H_0^{\nu_i}$, is the number of repeating data G_k for each k. This explains the proximity of $\mu(\nu_1)$ and $\mu(\nu_2)$.

On the other hand, since $\nu'_i s$ are irrational directions, the distribution of G_k approximates the given G on $H_0^{\nu_i}$ in large scale. Since ν_1 and ν_2 are close to the rational direction e_2 , the averaging behavior of a solution $u_{\varepsilon}^{\nu_i}$ in Π^{ν_i} would appear in a very large scale, in other words only after ε gets very small. We call this as the *secondary* homogenization.

The two-scale homogenization procedure has been introduced in [7], [8]. It allows studying continuity properties of the homogenized boundary data as we approach the rational direction, which might be singular points as described in the introduction. This point of view was also employed in [12] and [13] to study homogenization for general operators, by studying the singularity of homogenized operator at rational directions. Let us also point out near the boundary the small-scale oscillation of the operator interacts with that of boundary data to create a meso-scale averaging phenomena. Due to this interaction, characterizing the homogenized boundary condition remains a challenging and interesting open problem. After the first homogenization, the boundary data changes to periodic data in a meso-scale (which will be $N\varepsilon$ below), and hence the operator is well approximated by the homogenized operator \overline{F} in the second homogenization in large scale.

Below we begin the analysis of the two-step homogenization as described above. We will work with small $\varepsilon > 0$ satisfying

$$\varepsilon \le \frac{\delta \theta_i}{T_{\nu}(\delta^{5/2})} \quad \text{for } i = 1, 2$$

$$\tag{42}$$

which can be stated as

$$0 < \varepsilon \le \delta \theta_i \quad \text{for } i = 1, 2 \tag{43}$$

since $T_{\nu}(s) \equiv 1$ when $\nu = e_2$. It follows that

$$mN\varepsilon \le mM\varepsilon \le \delta. \tag{44}$$

After the near-boundary homogenization, $u_{\varepsilon}^{\nu_1}$ will be approximated by a solution which has periodic boundary data with period $mN\varepsilon$. With (44) it follows that $u_{\varepsilon}^{\nu_1}$ fluctuates in order of δ in the interior of the strip domain.

On the other hand, (38) of Theorem 5.2 can be stated as

$$0 < \theta_1 \le \delta^{5/2}.\tag{45}$$

It follows then that

$$1/N < \delta^{3/2} \tag{46}$$

which ensures $u_{\varepsilon}^{\nu_i}$ to homogenize $N\varepsilon$ -close to the Neumann boundary.

Next define

$$I_k := [(k-1)N\varepsilon, kN\varepsilon] \times \mathbb{R} \quad \text{for} \quad k \in \mathbb{Z}.$$

$$\tag{47}$$

Then we can observe that in each I_k , the Neumann boundary $H_0^{\nu_1}$ is located within $\delta\varepsilon$ -distance from $H_0 + \delta\varepsilon(k-1)e_2$, mod $\varepsilon\mathbb{Z}^n$. Thus on each $H_0^{\nu_1} \cap I_k$, G is approximated well by G_k for $1 \le k \le m$. If we extend the definition of G_k over $k \in \mathbb{Z}$ by letting $G_k = G_{\bar{k}}$ for $k = \bar{k} \pmod{m}$ then we have

$$|G(p,\frac{x}{\varepsilon}) - G_k(p,\frac{x}{\varepsilon})| < C(1+|p|)\delta \text{ on } H_0^{\nu_1} \cap I_k \text{ for } k \in \mathbb{Z}.$$
(48)

Similarly for ν_2 , if we define $J_k := [(k-1)M\varepsilon, kM\varepsilon] \times \mathbb{R}$ for $k \in \mathbb{Z}$.

Remarks 5.4. For $\nu \neq e_2$ in \mathbb{R}^2 , there exists a rational direction $\tilde{\nu}$ such that for $T = T_{\nu}(\delta^{5/2})$,

$$T\tilde{\nu} = 0 \pmod{\mathbb{Z}^2}; \quad |\nu - \tilde{\nu}| \le \delta^{5/2}/T.$$

Observe that if Theorem 5.2 holds for the rational direction $\tilde{\nu}$, it also holds for ν . For the proof of the theorem for $\tilde{\nu}$, let $x' = x - (x \cdot \tilde{\nu})\tilde{\nu}$ and define

$$G_k = G_k(x', x - x') = G(x', \delta(k - 1)\tilde{\nu}) \text{ for } 1 \le k \le m.$$

Then G_k is a periodic function on $\{x \cdot \tilde{\nu} = 0\}$ with a period of T. The only difference between the case of $\tilde{\nu}$ and e_2 is in the periodicity of the function G_k , and it does not make any essential difference in the proof. we point out that instead of the conditions (45), (46) and (44), we will need

$$\frac{1}{TN} \le \delta^{3/2}; \quad T\theta_1 \le \delta^{5/2}; \quad mTM\varepsilon \le \delta$$

since G_k has a period of T. These conditions will be ensured if θ_i and ε satisfy the assumptions as in Theorem 5.2.

5.2 Proof of Theorem 5.2

In the first three steps we follow the heuristics above and replace the Neumann condition with the locally projected boundary data G_k . Then we go through the two-step homogenization procedures to obtain the first slope $\mu^N(G_k)$ on each I_k near the boundary, and then the global slope $\mu(\nu_1)$. While the actual first homogenization takes place in Π^{ν_1} , it turns out that its value has a small difference from $\mu^N(G_k)$ taken in Π (see Lemma 5.6). This fact is important in establishing a universal domain for both directions ν_1 and ν_2 . In fact, we rotate the middle and inner regions to compare the slopes in Π^{ν_1} and Π^{ν_2} . For this, we use the rotational invariance of the homogenized operator \overline{F} . (See Lemma 5.7 and Lemma 5.8.) The rest of steps are to verify that indeed $\mu(\nu_1)$ is the correct averaged slope for the problem $(P)_{\varepsilon,\nu_1,\tau,q}$.

Step 1. First homogenization near Boundary (N ε - away from $H_0^{\nu_1}$)

We proceed to discuss the first homogenization. Denote $x = (x_1, x_2)$ throughout this section. For a given linear function $l(x) = l(x_1)$ and $k \in \mathbb{Z}$, let $u = u^{N,\varepsilon}$ and $v_k = v_k^{N,\varepsilon}$ solve the following problem with u = l on $H_{-N\varepsilon}^{\nu_1}$ and $v_k = l(x)$ on $H_{-N\varepsilon}$:

$$F(D^{2}u, x/\varepsilon) = 0 \quad \text{in} \quad \{-N\varepsilon \le x \cdot \nu_{1} \le 0\};$$

$$\frac{\partial u}{\partial \nu_{1}}(x) = G(Du, x/\varepsilon) \quad \text{on} \quad H_{0}^{\nu_{1}}$$
(49)

and

$$\begin{cases} F(D^2 v_k, x/\varepsilon) = 0 & \text{in} \quad \{-N\varepsilon \le x_2 \le 0\};\\ \frac{\partial v_k}{\partial x_2}(x) = G_k(Dv_k, x/\varepsilon) & \text{on} \quad H_0. \end{cases}$$
(50)

Definition 5.5. For a given function $u : \{-N\varepsilon \leq x \cdot \nu \leq 0\} \to \mathbb{R}$ and I_k given as in (47), let a_k and b_k be the middle points of $I_k \cap H_{-N\varepsilon/2}$ and $I_k \cap H_{-N\varepsilon}$ respectively, and consider the unique linear function h given by h = u at $x = a_k, b_k$ and $D_T h(b_k) = D_T u(b_k)$. (Here $D_T h$ denotes the tangential derivative of h along the direction ν^{\perp} .) Then $\mu_k(u)$ is defined by

$$\mu_k(u) := \partial h / \partial \nu.$$

Note that the Neumann boundary data of v_k is G_k on each boundary pieces $H_0 \cap I_i$ $(i \in \mathbb{Z})$, and hence $\mu_i(v_k) = \mu(v_k)$. For N as given in (41), we denote

$$\mu^N(G_k) := \mu(v_k). \tag{51}$$

Lemma 5.6. For $k \in \mathbb{Z}$ and $\mu_k(u)$ as given in Definition 5.5,

$$|\mu_k(u) - \mu^N(G_k)| < C\delta^{1/2}.$$
(52)

Proof. We will prove the lemma for k = 1, i.e., we will compare $\mu_1(u)$ with $\mu(v_1)$. Let \tilde{u} and \tilde{v}_1 solve the following problem with $\tilde{u} = l$ on $H^{\nu_1}_{-\varepsilon/\delta}$ and $\tilde{v}_1 = l$ on $H_{-\varepsilon/\delta}$:

$$\begin{cases} F(D^2 \tilde{u}, x/\varepsilon) = 0 & \text{in} \quad \{-\varepsilon/\delta \le x \cdot \nu_1 \le 0\};\\ \frac{\partial \tilde{u}}{\partial \nu_1}(x) = G(D \tilde{u}, x/\varepsilon) & \text{on} \quad H_0^{\nu_1} \end{cases}$$

and

$$\begin{cases} F(D^2 \tilde{v}_1, x/\varepsilon) = 0 & \text{in} \quad \{-\varepsilon/\delta \le x_2 \le 0\} \\\\ \frac{\partial \tilde{v}_1}{\partial x_2}(x) = G_1(D\tilde{v}_1, x/\varepsilon) & \text{on} \quad H_0. \end{cases}$$

We will compare both of $\tilde{u}(x)$ and $\tilde{v}_1(x)$ to $w_1(x)$ in the ball $|x| \leq \delta^{-1-\alpha_0}\varepsilon$, where $\alpha_0 = 1/2$. For computational convenience we will call this number as α_0 . Let $w_1(x)$ solve $w_1 = l$ on $H^{\nu_1}_{-\varepsilon/\delta}$ with

$$\begin{cases}
F(D^2w_1, x/\varepsilon) = 0 & \text{in} \quad \{-\varepsilon/\delta \le x \cdot \nu_1 \le 0\}; \\
\frac{\partial w_1}{\partial \nu_1}(x) = G_1(Dw_1, x/\varepsilon) & \text{on} \quad H_0^{\nu_1}.
\end{cases}$$
(53)

Here observe that in the ball $|x| \leq \delta^{-1-\alpha_0}\varepsilon$, the hyperplanes $H_0^{\nu_1}$ and H_0 only differ by $\theta_1 \delta^{-1-\alpha_0}\varepsilon$.

Below we derive some properties of w_1 . Consider

$$\bar{w}(x) := \varepsilon^{-1} w_1(\varepsilon x)$$

Then by Theorem 2.4, \bar{w} is $C^{1,1}$ regular up to the Neumann boundary in a unit ball, if \bar{w} has a bounded oscillation in the ball $|x| \leq 1/\delta$. Observe that $(\varepsilon/\delta)^{-1}w_1(\varepsilon x/\delta)$ is defined in the strip $\{-1 \leq x \cdot \nu_1 \leq 0\}$ and it has a periodic Neumann data $G_1(\cdot, \cdot, x/\delta)$ with period δ . Since it has a periodic boundary data, it corresponds to the case of rational direction with Neumann boundary passing through the origin. Hence we can use the error estimate Theorem 4.1 (b) for the rational direction passing through the origin, with $T_{\nu} = 1$. Then we obtain

$$\left|\left(\frac{\varepsilon}{\delta}\right)^{-1}w_1\left(\frac{\varepsilon x}{\delta}\right) - h(x)\right| \le \inf_{0 \le k \le 1} C(\delta^k + \delta^{1-k}) = C\delta^{1/2}$$
(54)

where h is a linear solution approximating $(\varepsilon/\delta)^{-1}w_1(\varepsilon x/\delta)$. Then by (54)

$$w_1(\frac{\varepsilon x}{\delta}) - \frac{\varepsilon}{\delta}h(x)| \le C\delta^{-1/2}\varepsilon$$
 (55)

and hence the oscillation of \bar{w} becomes less than $C\delta^{-1/2}$ in the ball $|x| \leq 1/\delta$. Later in the proof we will use $C^{1,1}$ regularity of \bar{w} as well as the linear approximation (55) of w_1 .

First, we compare \tilde{u} to w_1 in $B_{\delta^{-1-\alpha_0}\varepsilon}(0)$. For this, we compare the boundary data of \tilde{u} , that is G, to G_1 . Observe that if $x \in H_0^{\nu_1} \cap B_{\delta^{-1-\alpha_0}\varepsilon}(0)$, then $x \in I_k$ for some $|k| \leq \delta^{-1-\alpha_0}/N = \delta^{-2-\alpha_0}\theta_1$. Hence for $x \in H_0^{\nu_1} \cap B_{\delta^{-1-\alpha_0}\varepsilon}(0)$ (i.e., for $x \in H_0^{\nu_1} \cap I_k$ with $|k| \leq \delta^{-2-\alpha_0}\theta_1$),

$$|G(p, x/\varepsilon) - G_1(p, x/\varepsilon)| \leq |G(p, x/\varepsilon) - G_k(p, x/\varepsilon)| + |G_k(p, x/\varepsilon) - G_1(p, x/\varepsilon)|$$

$$\leq C[(1+|p|)\delta + (1+|p|)|k-1|\delta]$$

$$\leq C(1+|p|)(\delta + (\theta_1\delta^{(-1-\alpha_0)}))$$

$$\leq C(1+|p|)(\delta + \delta^{(3/2-\alpha_0)})$$

$$\leq C(1+|p|)\delta,$$
(56)

where the second inequality follows from (48) and the construction of G_k , third inequality follows from $|k| \leq \delta^{-2-\alpha_0}\theta_1$, the fourth inequality follows from (45), and the last inequality follows since $\alpha_0 \leq 1/2$. This implies, by Lemma 3.5,

$$|\tilde{u}(x) - w_1(x)| \le C(\delta + \delta^{\alpha_0})(x \cdot \nu_1 + \frac{\varepsilon}{\delta}) \le C\delta^{\alpha_0}(x \cdot \nu_1 + \frac{\varepsilon}{\delta}) \text{ in } |x| \le \delta^{-1 - \alpha_0}\varepsilon.$$
(57)

Observe that (55) and (57) yield

$$|\tilde{u}(x) - L_1(x)| \le C(\delta^{\alpha_0} + \delta^{1/2})(x \cdot \nu_1 + \frac{\varepsilon}{\delta}) \le C\delta^{\alpha_0}(x \cdot \nu_1 + \frac{\varepsilon}{\delta}) \text{ in } |x| \le \delta^{-1 - \alpha_0}\varepsilon.$$

where $L_1(x) = l(x) + \mu(w_1)(x \cdot \nu_1 + \frac{\varepsilon}{\delta})$, and $\mu(w_1)$ is the average slope of w_1 . In other words, we obtain

$$|\mu_1(\tilde{u}) - \mu(w_1)| \le C\delta^{\alpha_0}.$$
(58)

Next, we compare \tilde{v}_1 and w_1 and prove

$$|\mu(\tilde{v}_1) - \mu(w_1)| \le C\delta^{\alpha_0}$$

Recall that the oscillation of \bar{w} is less than $C\delta^{-1/2}$ in the ball $|x| \leq 1/\delta$ (see (55)). If we consider $\tilde{w} = \delta^{1/2} \bar{w}$, then this function solves the boundary condition

$$\partial \tilde{w} / \partial \nu = \tilde{G}(D\tilde{w}, x) = \delta^{1/2} G(\delta^{-1/2} D\tilde{w}, x)$$

which satisfies the assumptions for the $C^{1,1}$ regularity theory, Theorem 2.4. Thus we have

$$\|\bar{w}\|_{C^{1,1}(B_1)} \le O(\delta^{-1/2})$$

For x in the $\sigma\varepsilon$ -neighborhood of $H_0^{\nu_1}$, choose \tilde{x} to be the closest point to x on H_0 . Then by (G1) and (G2) with the $C^{1,1}$ regularity of \bar{w} given above, w_1 satisfies on H_0 ,

$$|G(Dw_1(x), \frac{x}{\varepsilon}) - G(Dw_1(\tilde{x}), \frac{\tilde{x}}{\varepsilon})| \le O(\delta^{-1/2}\sigma)(1 + |Dw_1(x)|)$$

Recall that the Neumann boundaries of w_1 and v_1 $(H_0^{\nu_1} \text{ and } H_0)$ only differ in the ball $|x| \leq \delta^{-1-\alpha_0} \varepsilon$, by $\theta_1 \delta^{-1-\alpha_0} \varepsilon \leq \delta^{3/2-\alpha_0} \varepsilon$ (see (45)). So putting $\sigma = \delta^{3/2-\alpha_0}$,

$$|G(Dw_1(x), \frac{x}{\varepsilon}) - G(Dw_1(\tilde{x}), \frac{\tilde{x}}{\varepsilon})| \le O(\delta^{1-\alpha_0})(1+|Dw_1(x)|) \text{ on } H_0$$

and Lemma 3.5 yields that in $|x| \leq \delta^{-1-\alpha_0} \varepsilon$,

$$|(\tilde{v}_1 - w_1)(x)| \le C(\delta^{1-\alpha_0} + \delta^{\alpha_0})(x_n + \frac{\varepsilon}{\delta}) \le C\delta^{\alpha_0}(x_n + \frac{\varepsilon}{\delta}).$$

This and (55) yield that in $|x| \leq \delta^{-1-\alpha_0} \varepsilon$,

$$|\tilde{v}_1(x) - L(x)| \le C(\delta^{\alpha_0} + \delta^{1/2})(x_n + \frac{\varepsilon}{\delta}) \le C\delta^{\alpha_0}(x_n + \frac{\varepsilon}{\delta})$$

where $L(x) = l(x) + \mu(w_1)(x_2 + \frac{\varepsilon}{\delta})$. In other words, we obtain

$$|\mu(w_1) - \mu(\tilde{v}_1)| \le C\delta^{\alpha_0}.$$
(59)

Recalling $\alpha_0 = 1/2$ we conclude from (58) and (59) that

$$|\mu_1(\tilde{u}) - \mu(\tilde{v}_1)| \le C\delta^{1/2}.$$
(60)

In the rest of proof, we will show

$$|\mu(v_1) - \mu(\tilde{v}_1)|, \ |\mu_1(u) - \mu_1(\tilde{u})| \le C\delta^{1/2}$$

Then the above inequalities and (60) would imply

$$|\mu_1(u) - \mu(v_1)| \le |\mu_1(u) - \mu_1(\tilde{u})| + |\mu_1(\tilde{u}) - \mu(\tilde{v}_1)| + |\mu(\tilde{v}_1) - \mu(v_1)| \le C\delta^{1/2}.$$

First, observe that v_1 and \tilde{v}_1 have periodic Neumann data G_1 on H_0 . Hence by similar arguments as in the proof of (27),

$$|\mu(v_1) - \mu(\tilde{v}_1)| \le C(\Lambda(\delta, e_2) + N^{-1}) \le C(\delta^{1/2} + N^{-1}) \le C\delta^{1/2}$$
(61)

where the last inequality follows from (46).

Next, recall that

$$|\mu_1(\tilde{u}) - \mu(w_1)| \le C\delta^{1/2}$$

for a solution w_1 of (53). (See (58).) Similarly, one can prove

$$|\mu_1(u) - \mu(\tilde{w}_1)| \le CN^{-1/2} \le C\delta^{1/2}$$

where \tilde{w}_1 solves similar equations as in (53) in the domain $\{-N\varepsilon \leq x \cdot \nu_1 \leq 0\}$, and the last inequality follows from (46). Then since w_1 and \tilde{w}_1 have periodic Neumann data G_1 on $H_0^{\nu_1}$, it corresponds to the case of $\nu = e_2$. Hence by similar arguments as in (61),

$$|\mu(w_1) - \mu(\tilde{w}_1)| \le C(\Lambda(\delta, e_2) + N^{-1}) \le C(\delta^{1/2} + N^{-1}) \le C\delta^{1/2}$$

and we can conclude

$$|\mu_1(u) - \mu_1(\tilde{u})| \le |\mu_1(u) - \mu(\tilde{w}_1)| + |\mu(\tilde{w}_1) - \mu(w_1)| + |\mu(w_1) - \mu_1(\tilde{u})| \le C\delta^{1/2}.$$

Step 2. Constructing middle region barrier ω_{ε} (between $H_{-N\varepsilon/2}$ and $H_{-KmN\varepsilon}$)

In step 1 we showed that $N\varepsilon$ away from the boundary $H_0^{\nu_1}$, $u_{\varepsilon}^{\nu_1}$ is homogenized with average slope approximated by $\mu^N(G_k)$ in each vertical strip I_k . Now more than $N\varepsilon$ away from $H_0^{\nu_1}$, we obtain the second homogenization of $u_{\varepsilon}^{\nu_1}$, whose slope is determined by $\mu^N(G_k)$, k = 1, ..., m. Since the width of $I_k = N\varepsilon$, the homogenized slopes $\mu^N(G_1), ..., \mu^N(G_m)$ are repeated K times in a vertical strip of width $KmN\varepsilon$, $N\varepsilon$ -away from $H_0^{\nu_1}$. We will specify

$$K := 1/\delta,$$

but for computational clarity we will keep the symbol K.

We will construct middle region barrier ω_{ε} in the region $\{-KmN\varepsilon \leq x_2 \leq -N\varepsilon/2\}$. To ensure that ω_{ε} is regular near its Neumann boundary, we introduce a regularization of the original Neumann boundary data $\mu^N(G_k)$ as follows.

Consider a ball $B_{\delta^{-\alpha_0/2}N\varepsilon}(0)$. If $I_k \cap H_0$, $I_j \cap H_0 \subset B_{\delta^{-\alpha_0/2}N\varepsilon}(0)$, then $|k-j| \leq \delta^{-\alpha_0/2}$ and

$$|G_k(p, x/\varepsilon) - G_j(p, x/\varepsilon)| \le C(1+|p|)(|k-j|\delta) \le C(1+|p|)\delta^{(1-\alpha_0/2)}.$$
(62)

Using this fact with Lemma 3.5, we can construct a C^1 function $\Lambda(x)$ on $H_{-N\varepsilon/2}$ such that

- (a) $\Lambda \in C^1(H_{-N\varepsilon/2})$ with $\|\Lambda\|_{C^1} \leq \delta(N\varepsilon)^{-1}$;
- (b) $\mu^N(G_k) + \delta^{\alpha_0} \leq \Lambda(x) \leq \mu^N(G_k) + \delta^{\alpha_0} + \delta$ on each I_k ;

(c) $\Lambda(x)$ is periodic with period $mN\varepsilon$.

Note that when we patch the middle region barrier ω_{ε} with the near-boundary barrier f_{ε} in step 6, we will need that the average slope of ω_{ε} is "sufficiently" larger than that of f_{ε} . For this, we will make the average slope of ω_{ε} to be $\mu^{N}(G_{k}) + O(\delta^{\alpha_{0}})$, i.e., (b) is to ensure that $\mu_{k}(\omega_{\varepsilon})$ is sufficiently larger than $\mu_{k}(f_{\varepsilon})$. Also when we show the flatness of barriers in steps 4 and 5, we will localize them in a "large" ball of size $\delta^{-\alpha_{0}/2}N\varepsilon$.

Let $\Sigma := \{-KmN\varepsilon \le x_2 \le -N\varepsilon/2\}$ and ω_{ε} solve the following Neumann boundary problem

$$F(D^{2}\omega_{\varepsilon}, x/\varepsilon) = 0 \quad \text{in} \quad \Sigma$$

$$\frac{\partial\omega_{\varepsilon}}{\partial x_{2}} = \Lambda(x) \qquad \text{on} \quad H_{-N\varepsilon/2} \qquad (63)$$

$$\omega_{\varepsilon} = l(x) \qquad \text{on} \quad H_{-KmN\varepsilon}.$$

Step 3. Homogenization of the operator in the middle region

Next we show, similar to Lemma 5.6, that the second homogenization does not change too much if the domain Π is replaced by Π^{ν_1} . More precisely, we will show that ω_{ε} is close to $\tilde{\omega}_{\varepsilon}$ solving

$$\left\{ \begin{array}{ll} \bar{F}(D^2 \tilde{\omega}_{\varepsilon}) = 0 & \mbox{in} \quad \{-KmN\varepsilon \leq x \cdot \nu_1 \leq -N\varepsilon/2\} \\ \\ \frac{\partial \tilde{\omega}_{\varepsilon}}{\partial \nu_1} = \Lambda(x) & \mbox{on} \quad H^{\nu_1}_{-N\varepsilon/2} \\ \\ \\ \tilde{\omega}_{\varepsilon} = l(x) & \mbox{on} \quad H^{\nu_1}_{-KmN\varepsilon}. \end{array} \right.$$

To this end we will first compare ω_{ε} with $\bar{\omega}_{\varepsilon}$, with the same Dirichlet data l on $H_{-kmN\varepsilon}$ and solving

$$\begin{cases} \bar{F}(D^2\bar{\omega}_{\varepsilon}) = 0 & \text{in} \quad \Sigma\\ \frac{\partial\bar{\omega}_{\varepsilon}}{\partial x_2} = \Lambda(x) & \text{on} \quad H_{-N\varepsilon/2}. \end{cases}$$
(64)

Lemma 5.7. For any $\sigma > 0$, there exists N_0 such that for $N_0 > N$ we have

$$|\omega_{\varepsilon}(x) - \bar{\omega}_{\varepsilon}(x)| \le \sigma \delta N \varepsilon \quad in \quad \Sigma$$

Proof. The proof follows from Theorem 2.5 applied to $(\delta N \varepsilon)^{-1} \omega_{\varepsilon} (N \varepsilon x)$.

Next we compare $\bar{\omega}_{\varepsilon}$ to $\tilde{\omega}_{\varepsilon}$ to conclude. Here we will use the rotational invariance of \bar{F} .

Lemma 5.8. Let \mathcal{O} be the rotation matrix that maps e_2 to ν_1 . Then

$$|\tilde{\omega}_{\varepsilon}(\mathcal{O}x) - \bar{\omega}_{\varepsilon}(x)| \le \delta^{1/2}(KmN\varepsilon)$$

 $in \ \Sigma \cap \{ |x| \le \delta^{-1/2} (N\varepsilon) \}.$

Proof. Observe that $v(x) := \tilde{\omega}(\mathcal{O}x)$ solves $\bar{F}(v) = 0$ in Σ with Neumann boundary data $\Lambda(\mathcal{O}x)$ on $H_{-N\varepsilon/2}$ and Dirichlet data $l(\mathcal{O}x)$ on $H_{-KmN\varepsilon}$. Note that due to (45) and the C^1 bound of Λ we have

$$|\Lambda(KmN\varepsilon\mathcal{O}x) - \Lambda(KmN\varepsilon x)| \le \theta_1 |KmN\varepsilon x| \sup |D\Lambda| \le \delta |x|.$$

and $|l(KmN\varepsilon \mathcal{O}x) - l(KmN\varepsilon x)| \leq KmN\varepsilon \theta_1 |x| \leq \delta |x|.$

Hence one can apply Lemma 2.9 of [7] to $\tau^{-1}v(\tau x)$ and $\tau^{-1}\bar{w}(\tau x)$ in $\tau^{-1}\Sigma$, where $\tau = KmN\varepsilon$ and choose $R := \delta^{-1/2}$ and $\varepsilon = 2$ to conclude.

Step 4. Flatness of ω_{ε} on $H_{-N\varepsilon}$, and the construction of near-boundary barrier f_{ε}

Lemma 5.9. [Flatness of ω_{ε}] Let x_0 be any point on $H_{-N\varepsilon}$. Then for $x \in H_{-N\varepsilon} \cap B_{\delta^{-\alpha_0/2}N\varepsilon}(x_0)$

$$|\omega_{\varepsilon}(x) - \omega_{\varepsilon}(x_0) - \partial_1 \omega_{\varepsilon}(x_0)(x - x_0)| \le C\delta^{1 - \alpha_0} N\varepsilon.$$

Proof. Due to Lemma 5.7, it is enough to show above lemma for $\bar{\omega}_{\varepsilon}$. Let $\omega_1(x) := (KmN\varepsilon)^{-1}\bar{\omega}_{\varepsilon}((KmN\varepsilon)x)$, then it solves

$$\begin{cases} \bar{F}(D^2\omega_1) = 0 & \text{in} \quad \{-1 \le x_2 \le -\frac{1}{2Km}\} \\ \frac{\partial\omega_1}{\partial x_2} = \Lambda(KmN\varepsilon x) & \text{on} \quad H_{-\frac{1}{2Km}} \\ \omega_1(x) = l(x) + C & \text{on} \quad H_{-1} \end{cases}$$

we know that $\|\Lambda\|_{C^1} \leq \delta(N\varepsilon)^{-1}$, so the above Neumann boundary data has C^1 norm of δKm . From Theorem 2.4, we have that

$$\|\omega_1\|_{C^{1,1}} \le C\delta Km.$$

Hence

$$|\omega_1(x) - \omega_1(x_0) - \partial_1 \omega_1(x_0) \cdot (x - x_0)| \le C\delta K m |x - x_0|^2$$
(65)

which can be written in terms of $\bar{\omega}_{\varepsilon}$,

$$\begin{aligned} |\bar{\omega}_{\varepsilon}(x) - \bar{\omega}_{\varepsilon}(x_0) - \partial_1 \bar{\omega}_{\varepsilon}(x_0) \cdot (x - x_0)| &\leq C\delta(Km)^2 (N\varepsilon) |(KmN\varepsilon)^{-1} (x - x_0)|^2 \\ &\leq C\delta(\delta^{-\alpha_0/2})^2 (N\varepsilon) = C\delta^{1-\alpha_0} N\varepsilon \end{aligned}$$

in $\delta^{-\alpha_0/2} N \varepsilon$ -neighborhood of x_0 .

Now we construct the near-boundary barrier f_{ε} using ρ_{ε} . Let f_{ε} solve

$$\begin{cases} F(D^2 f_{\varepsilon}, x/\varepsilon) = 0 & \text{in} \quad \{-N\varepsilon \le x_2 \le 0\};\\ f_{\varepsilon} = \omega_{\varepsilon} + \delta^{1-\alpha_0} N\varepsilon & \text{on} \quad H_{-N\varepsilon};\\ \frac{\partial f_{\varepsilon}}{\partial x_2} = G(D f_{\varepsilon}, \frac{x}{\varepsilon}) & \text{on} \quad H_0. \end{cases}$$

Step 5. Flatness of f_{ε}

In this step we compare $\mu^N(G_k)$ given in (51) with $\mu_k(f_{\epsilon})$ given in Definition 5.5. For simplicity we put k = 1. Note that Lemma 3.2, Lemma 5.9, and Lemma 3.5 with (62) imply that

$$|\mu^N(G_1) - \mu_1(f_{\varepsilon})| \le C(\delta^{1-\alpha_0/2} + \delta + \delta^{1-\alpha_0}) \le C\delta^{1-\alpha_0}$$
(66)

Also from Lemma 5.9 and the definition of f_{ε} it follows that f_{ε} is close to a linear function

L		
L		

$$|f_{\varepsilon}(x) - L_0(x)| \le C\delta^{1-\alpha_0} N\varepsilon \quad \text{on} \quad H_{-N\varepsilon} \cap B_{\delta^{-\alpha_0/2} N\varepsilon}(0), \tag{67}$$

where $L_0(x) := f_{\varepsilon}(-N\varepsilon e_2) + \mu^N(G_1)(x_2 + N\varepsilon) + \partial_1 f_{\varepsilon}(-N\varepsilon e_2)x_1$. Then Lemma 4.4, (67) and Lemma 3.2 applied to the rescaled function $(N\varepsilon)^{-1}f_{\varepsilon}(N\varepsilon x)$ in the region $\{-1 \le x_2 \le -1/2\} \cap B_{\delta^{-\alpha_0/2}}$ yield that

$$|f_{\varepsilon} - L_0| \le C(\delta^{1-\alpha_0} + \delta^{(1-\alpha_0/2)})N\varepsilon + C\varepsilon \le C\delta^{1-\alpha_0}N\varepsilon$$
(68)

in $\{-N\varepsilon \leq x_2 \leq -N\varepsilon/2\} \cap B_{\delta^{-\alpha_0/2}N\varepsilon}(0)$, where the last inequality follows from (46).

Before we proceed to the next step, observe that the C^1 regularity of Λ , Theorem 2.4, as well as Lemma 5.7 yield that

$$|\omega_{\varepsilon}(x_1, x_2) - \omega_{\varepsilon}(x_1, -N\varepsilon) - \Lambda(x)(x_2 + N\varepsilon)| \le C\delta^{1-\alpha_0}N\varepsilon \quad \text{on } \{-N\varepsilon \le x_2 \le -\frac{N\varepsilon}{2}\}.$$
(69)

Step 6. Patching up

Let $h(x) := l(x) + (\mu(\omega_{\varepsilon}) - C\delta^{1/2})(x_2 + KmN\varepsilon)$ where C > 0 is a constant given as in (b) of Theorem 4.1, and $l(x) = l(x_1)$ is a linear function chosen so that $h(x) = q \cdot x$ on H_{-1} . We define

$$\rho_{\varepsilon} := \begin{cases} h & \text{in} \quad \{-1 \le x_2 \le -KmN\varepsilon\},\\\\ \omega_{\varepsilon} & \text{in} \quad \{-KmN\varepsilon \le x_2 \le -N\varepsilon/2\} \end{cases}$$

Since Λ is $mN\varepsilon$ -periodic, (b) of Theorem 4.1 implies that on $\{x_2 = -KmN\varepsilon\}$,

$$\partial_{x_2}\omega_{\varepsilon} \ge \mu(\omega_{\varepsilon}) - C\Lambda(1/K, e_2) = \mu(\omega_{\varepsilon}) - CK^{-1/2} = \mu(\omega_{\varepsilon}) - C\delta^{1/2} = \partial_{x_2}h.$$

Thus it follows that $F(D^2\rho_{\varepsilon}, \frac{x}{\varepsilon}) \leq 0$ in $\{-1 \leq x_2 \leq -N\varepsilon/2\}$.

Due to the flatness estimates (68) and (69), we can approximate f_{ε} and ρ_{ε} by linear functions, respectively with normal derivatives of $\mu^{N}(G_{k})$ and $\Lambda(x)$, with the error of $O(\delta^{1-\alpha_{0}}N\varepsilon)$. Here recall $\Lambda(x)$ was constructed so that $\Lambda(x) \geq \mu^{N}(G_{k}) + \delta^{\alpha_{0}}$, and α_{0} is a constant satisfying $\alpha_{0} \leq 1/2$. Then since $f_{\varepsilon} = \rho_{\varepsilon} + \delta^{1-\alpha_{0}}N\varepsilon$ on $\{x_{2} = -N\varepsilon\}$,

$$\rho_{\varepsilon} > f_{\varepsilon} \text{ on } \{x_2 = -N\varepsilon/2\} \text{ and } f_{\varepsilon} > \rho_{\varepsilon} \text{ on } \{x_2 = -N\varepsilon\}.$$
(70)

Define $\underline{\rho}_{\epsilon}$ as follows:

$$\underline{\rho}_{\varepsilon} := \begin{cases} \rho_{\varepsilon} & \text{in} \quad \{-1 \leq x_2 \leq -N\varepsilon\},\\ \min(\rho_{\varepsilon}, f_{\varepsilon}) & \text{in} \quad \{-N\varepsilon \leq x_2 \leq -N\varepsilon/2\},\\ f_{\varepsilon} & \text{in} \quad \{-N\varepsilon/2 \leq x_2 \leq 0\}, \end{cases}$$

Then by (70), $\underline{\rho}_{\varepsilon}$ is a viscosity supersolution of $(P)_{\varepsilon,e_2,0,q}$ in $\{-1 \leq x_2 \leq 0\}$. Let us mention that, due to Lemma 5.6, Lemma 5.7 and Lemma 5.8, a small perturbation of these barriers also yield a supersolution in $\{-1 \leq x \cdot \nu_1 \leq 0\}$. Similarly, one can construct a subsolution $\overline{\rho}_{\varepsilon}$ of $(P)_{\varepsilon,e_2,0,q}$ by replacing $\Lambda(x)$ given in the construction of ρ_{ε} by $\tilde{\Lambda}(x) \leq \mu^N(G_k) - \delta^{\alpha_0}$. Then by Lemma 5.6 and Lemma 5.8

$$|\mu(u_{\varepsilon}^{\nu_{1}}) - \mu(\underline{\rho}_{\varepsilon})| \le |\mu(\bar{\rho}_{\varepsilon}) - \mu(\underline{\rho}_{\varepsilon})| + C\delta^{1/2} \le C(\delta^{1/2} + \delta^{\alpha_{0}}) \le C\delta^{\alpha_{0}} = C\delta^{1/2}$$
(71)

where the last inequality follows by choosing $\alpha_0 = 1/2$.

We denote $\bar{\rho}_{\varepsilon} = \bar{\rho}_{\varepsilon}^{\nu_1}$ and $\underline{\rho}_{\varepsilon} = \underline{\rho}_{\varepsilon}^{\nu_1}$ indicating that they are obtained from the direction ν_1 , i.e., with the scale $N\varepsilon$.

Step 7. Comparing the solutions $u_{\varepsilon}^{\nu_1}$ and $u_{\varepsilon}^{\nu_2}$: Proof of Theorem 5.2 (a)

Parallel arguments as in the previous steps apply to the other direction ν_2 . Recall that

$$\theta_2 = |\nu_2 - e_2| < \theta_1, \quad M = [\frac{\delta}{\theta_2}] > N.$$

Then similarly as in the direction ν_1 , we can construct barriers $\bar{\rho}_{\varepsilon}^{\nu_2}$ and $\underline{\rho}_{\varepsilon}^{\nu_2}$ such that

$$|\mu(u_{\varepsilon}^{\nu_2}) - \mu(\underline{\rho}_{\varepsilon}^{\nu_2})| \le |\mu(\bar{\rho}_{\varepsilon}^{\nu_2}) - \mu(\underline{\rho}_{\varepsilon}^{\nu_2})| + C\delta^{1/2} \le C\delta^{1/2}.$$
(72)

Here their corresponding Neumann boundary conditions satisfy

$$\mu^{M}(G_{k}) - \delta^{\alpha_{0}} - \delta \leq \frac{\partial}{\partial x_{2}} \bar{\rho}_{\varepsilon}^{\nu_{2}} ; \frac{\partial}{\partial x_{2}} \underline{\rho}_{\varepsilon}^{\nu_{2}} \leq \mu^{M}(G_{k}) + \delta^{\alpha_{0}} + \delta \quad \text{on } H_{-M\varepsilon} \cap I_{k},$$

where $\alpha_0 = 1/2$, and the respective derivatives of $\bar{\rho}_{\varepsilon}^{\nu_2}$ and $\underline{\rho}_{\varepsilon}^{\nu_2}$ are taken as a limit from the region $\{-1 \leq x_2 < -M\varepsilon\}$.

Thus to compare $\mu(u_{\varepsilon}^{\nu_1})$ and $\mu(u_{\varepsilon}^{\nu_2})$, we compare $\mu^N(G_k)$ and $\mu^M(G_k)$. Recall that we define $\mu^M(G_k)$ similarly as $\mu^N(G_k)$. More precisely, $\mu^M(G_k)$ is the slope of the linear approximation of $v_k^{M,\varepsilon}$, where $v_k^{M,\varepsilon}$ is defined similarly as in (50) in the region $\{-M\varepsilon \leq x_2 \leq 0\}$ with the boundary condition

$$\partial_{x_2} v_k^{M,\varepsilon}(x) = G_k(Dv_k^{M,\varepsilon}, x/\varepsilon) \quad \text{ on } \quad H_0$$

and $v_k^{M,\varepsilon} = l(x)$ on $H_{-M\varepsilon}$. Since G_k is periodic on the Neumann boundary, it corresponds to the case of Neumann boundary with rational normal, passing through the origin. Hence by applying arguments as in the proof of (33),

$$|\mu^{N}(G_{k}) - \mu^{M}(G_{k})| \le C\Lambda(1/N, e_{2}) = C \inf_{0 \le k \le 1} \{1/N^{k} + 1/N^{1-k}\} = C/N^{1/2}.$$
(73)

Now we prove the following lemma using the estimate (73).

Lemma 5.10. For any ε satisfying (43),

$$|\mu(u_{\varepsilon}^{\nu_1}) - \mu(u_{\varepsilon}^{\nu_2})| \le C\delta^{1/2}$$

Proof. By the construction of the viscosity supersolution $\underline{\rho}_{\varepsilon}^{\nu_{1}}$ and Lemma 5.7,

$$|\mu(\underline{\rho}_{\varepsilon}^{\nu_{1}}) - \mu(\bar{\omega}_{\varepsilon})| \le C\delta^{1/2} \tag{74}$$

where $\bar{\omega}_{\varepsilon}$ is given as in (64). Similarly, we get

$$|\mu(\underline{\rho}_{\varepsilon}^{\nu_2}) - \mu(\bar{\omega}_{\varepsilon}^{\nu_2})| \le C\delta^{1/2} \tag{75}$$

where $\bar{\omega}_{\varepsilon}^{\nu_2}$ solves

$$\begin{split} & \bar{F}(D^2 \bar{\omega}_{\varepsilon}^{\nu_2}) = 0 \quad \text{in} \quad \{-KmM\varepsilon \le x_2 \le -M\varepsilon/2\}; \\ & \frac{\partial \bar{\omega}_{\varepsilon}^{\nu_2}}{\partial \nu} = \Lambda^{\nu_2}(x) \quad \text{on} \quad H_{-M\varepsilon/2}; \\ & \bar{\omega}_{\varepsilon}^{\nu_2} = l(x) \quad \text{on} \quad H_{-KmM\varepsilon}. \end{split}$$

Here $\Lambda^{\nu_2}(x)$ is constructed similarly as $\Lambda(x)$ with N replaced by M, i.e., with $\mu^N(G_k)$ replaced by $\mu^M(G_k)$. Then by (71), (72), (74) and (75), it suffices to prove

$$|\mu(\bar{\omega}_{\varepsilon}) - \mu(\bar{\omega}_{\varepsilon}^{\nu_2})| \le C\delta^{1/2}.$$

Recall that $|\Lambda(x) - \mu^N(G_k)| \leq \delta^{\alpha_0} + \delta$ on I_k , and similarly, $|\Lambda^{\nu_2}(x) - \mu^M(G_k)| \leq \delta^{\alpha_0} + \delta$ on I_k , with $\alpha_0 = 1/2$. Hence

$$|\mu(\bar{\omega}_{\varepsilon}) - \mu(h_1)|, \ |\mu(\bar{\omega}_{\varepsilon}^{\nu_2}) - \mu(h_2)| \le C\delta^{1/2}$$
(76)

for solutions h_1 and h_2 of

$$\begin{cases} \bar{F}(D^2h_1) = 0 & \text{in} \quad \{-KmN\varepsilon \le x_2 \le -N\varepsilon/2\} \\\\ \frac{\partial h_1}{\partial \nu} = \mu^N(G_k) & \text{on} \quad H_{-N\varepsilon/2} \cap I_k \\\\ h_1 = l(x) & \text{on} \quad H_{-KmN\varepsilon} \end{cases}$$

and

$$\begin{cases} \bar{F}(D^2h_2) = 0 & \text{in} \quad \{-KmM\varepsilon \le x_2 \le -M\varepsilon/2\} \\\\ \frac{\partial h_2}{\partial \nu} = \mu^M(G_k) & \text{on} \quad H_{-M\varepsilon/2} \cap I_k \\\\ h_2 = l(x) & \text{on} \quad H_{-KmM\varepsilon}. \end{cases}$$

Note that h_1 has a periodic Neumann condition on $H_{-N\varepsilon/2}$ with period $mN\varepsilon$, and also h_2 has a periodic Neumann condition on $H_{-M\varepsilon/2}$ with period $mM\varepsilon$. Hence they correspond to the case of periodic Neumann boundary data, i.e., the case of Neumann boundary with a normal direction e_2 , and passing through the origin. Hence by Theorem 4.1 with (73) and $K = 1/\delta$, we get

$$|\mu(h_1) - \mu(h_2)| \le \Lambda(\delta, e_2) + C/N^{1/2} \le C(\delta^{1/2} + (1/N)^{1/2}) \le C\delta^{1/2}$$
(77)

where the last inequality follows from (46). Then we can conclude from (76) and (77).

Remarks 5.11. For the dimension n > 2 and $\nu = e_n$, for a fixed $m \in \mathbb{N}$ and $\delta = \frac{1}{m}$ let us define

$$G_i(x_1, ..., x_{n-1}, x_n) := G(x_1, ..., x_{n-1}, \delta(i-1))$$
 for $i = 0, ..., m$

and

$$I_{k_1,k_2,\dots,k_{n-1}} := [(k_1 - 1)N\varepsilon, k_1N\varepsilon] \times \dots \times [(k_{n-1} - 1)N\varepsilon, k_{n-1}N\varepsilon] \times \mathbb{R}$$

Then parallel arguments as in steps 1 to 9 would apply to yield the results in \mathbb{R}^n .

A Appendix

In this section we state quantitative results on distribution of $\varepsilon \mathbb{Z}^n$ near a hyperplane. We present an improved version from those introduced in [8]. Recall that $\nu \in S^{n-1}$ is a rational direction if $\nu \in \mathbb{RZ}^n$, otherwise ν is an irrational direction. For properties of irrational directions, let us discuss the averaging property of the sequence $(nx)_n \mod 1$, for an irrational number x. We are particularly interested in the estimates on the rate of convergence of the sequence $(nx)_n$ to the uniform distribution (Definition A.2).

Note that the estimates in Lemma A.3 below are improved from the estimates in Lemma 2.7 of [8]. We begin with recalling the notion of equi-distribution.

• A bounded sequence $(x_1, x_2, x_3...)$ of real numbers is said to be *equi-distributed* on an interval [a, b] if for any $[c, d] \subset [a, b]$ we have

$$\lim_{n \to \infty} \frac{|\{x_1, ..., x_n\} \cap [c, d]|}{n} = \frac{d - c}{b - a}.$$

Here $|\{x_1, ..., x_n\} \cap [c, d]|$ denotes the number of elements of $\{x_1, ..., x_n\} \cap [c, d]$.

• The sequence $(x_1, x_2, x_3, ...)$ is said to be equi-distributed modulo 1 if $(x_1 - [x_1], x_2 - [x_2], ...)$ is equi-distributed in the interval [0, 1].

Lemma A.1 ([22], Weyl's equidistribution theorem). If a is an irrational number, the sequence (a, 2a, 3a, ...) is equi-distributed modulo 1.

To discuss quantitative versions of Lemma A.1, we introduce the notion of discrepancy. The following definition is from the book [17].

Definition A.2. Let $(x_k)_k$ be a sequence in \mathbb{R} . For a subset $E \subset [0,1]$, let

$$A(E; N) = |\{x_n : 1 \le n \le N\} \cap E|$$

i.e., A(E; N) denotes the number of points $\{x_n\}$, $1 \le n \le N$, that lie in E.

(a) The sequence $(x_n)_n$ is said to be uniformly distributed modulo 1 in \mathbb{R} if

$$\lim_{N \to \infty} \frac{A(E;N)}{N} = \mu(E)$$

for all $E = [a, b) \subset [0, 1]$. Here μ denotes the Lebesgue measure.

(b) For $x \in [0,1]$ and $N \in \mathbb{N}$, the discrepancy $D_x(N)$ is defined as follows:

$$D_x(N) = \sup_{E=[a,b)} \left| \frac{A(E;N)}{N} - \mu(E) \right|,$$

where A(E; N) is defined with the sequence $(kx)_k$ modulo 1.

It easily follows from Lemma A.1 that the sequence $(x_k)_k = (kx)_k$ is uniformly distributed modulo 1 for any irrational number $x \in \mathbb{R}$. In particular $D_x(N)$ converges to zero as $N \to \infty$.

Next, we apply the discrepancy function to multi-dimensions. For a direction $\nu = (\nu_1, ..., \nu_n) \in S^{n-1}$, let ν_i be the component with the biggest size, i.e.,

$$|\nu_i| = \max\{|\nu_j| : 1 \le j \le n\}$$

(if there are multiple components then we choose ν_i with the largest index *i*). Let *H* be the hyperplane in \mathbb{R}^n , which passes through 0 and is normal to ν , i.e.,

$$H = \{ x \in \mathbb{R}^n : x \cdot \nu = 0 \}.$$

Since $\nu_i \neq 0$, there exists $m = m(\nu)$ such that $(1, ..., 1, m, 1, ..., 1) \in H$, i.e.,

$$(1, ..., 1, m, 1, ..., 1) \cdot \nu = 0$$

where m is the i-th component of (1, ..., 1, m, 1, ..., 1). Note that m is irrational iff ν is an irrational direction. Define

$$\omega_{\nu}(N) := D_m(N). \tag{78}$$

Note that if ν is an irrational direction, then

$$\omega_{\nu}(N) \to 0 \text{ as } N \to \infty.$$

Now we are ready to state our quantitative estimate on the averaging properties of the vector sequence $(n\nu)_n$ with an irrational direction ν . Recall that for $\nu \in S^{n-1}$ and $\tau \in \mathbb{R}^n$,

$$\Pi(\nu, \tau) = \{x : -1 \le (x - \tau) \cdot \nu \le 0\}$$

and

$$H_0 = \{ x : (x - \tau) \cdot \nu = 0 \}.$$

Also for $x_0 \in \Pi(\nu, \tau)$ and $d := \operatorname{dist}(\mathbf{x}_0, \mathbf{H}_0)$, we denote

$$H_d = \{x : (x - \tau) \cdot \nu = d\} = \{x : (x - x_0) \cdot \nu = 0\}.$$

Lemma A.3. for $\nu \in S^{n-1}$ and $\tau \in \mathbb{R}^n$, let $x_0 \in \Pi(\nu, \tau)$ and let $0 < \varepsilon < d := \text{dist}(x_0, H_0)$.

(a) Suppose that ν is a rational direction. Then for any $x \in H_d$, there is $y \in H_d$ such that

$$|x-y| \leq T_{\nu}\varepsilon; \ y=x_0 \mod \varepsilon \mathbb{Z}^n$$

where T_{ν} is the smallest positive number such that $T_{\nu}\nu \in \mathbb{Z}^{n}$.

(b) Suppose that ν is an irrational direction, and let $\omega_{\nu} : \mathbb{N} \to \mathbb{R}^+$ be defined as in (78). Then there exists a dimensional constant M > 0 such that the following is true: for any $x \in H_d$ and $N \in \mathbb{N}$, there is $y \in \mathbb{R}^n$ such that

$$|x-y| \le MN\varepsilon + \varepsilon\omega_{\nu}(N); \ y = x_0 \ mod \ \varepsilon\mathbb{Z}^n$$

and

$$\operatorname{dist}(y, H_d) < \varepsilon \omega_{\nu}(N).$$

Here note that $\omega_{\nu}(N)$ converges to 0 as $N \to \infty$.

(c) If ν is an irrational direction, then for any $z \in \mathbb{R}^n$ and $\delta > 0$, there is $w \in H_d$ such that

$$|z-w| \leq \delta \mod \varepsilon \mathbb{Z}^n$$
.

Proof. Proof of (a) is immediate from the fact that $T_{\nu}\nu \in \mathbb{Z}^n$. Next, we let ν be an irrational direction in S^{n-1} . Without loss of generality, we may assume

$$|\nu_n| = \max\{|\nu_j| : 1 \le j \le n\}.$$

Let x be any point on H_d : after a translation and rotation around the x_n -axis, we may assume that $x = 0 \in H_d$ and $H_d \cap [0,1]^n \neq \emptyset$. Choose m such that

$$(1, 1, ..., 1, m) \in H_d.$$

Note that $|(1, 1, .., 1, m)| \leq M$ for a dimensional constant M > 0, since $|\nu_n|$ is the largest. Also note that

 $k\varepsilon(1,1,..,1,m) \in H_d$ for any integer k

since H_d contains the origin.

Consider the sequence $(km)_k$, then from the definition of $\omega_{\nu}(N)$ and the discrepancy function $D_m(N)$, it follows that any interval $[a, b] \subset [0, 1]$ of length $\omega_{\nu}(N)$ contains at least one point $km \pmod{1}$, for some $k \leq N$. Hence for any

$$z = (0, 0, ..., 0, z_n)$$
 with $0 \le z_n \le \varepsilon$

there exists

$$w = k\varepsilon(1, 1, .., 1, m) \in H_d$$

for some $0 \le k \le N$ such that

$$|z-w| \le \varepsilon \omega_{\nu}(N) \mod \varepsilon \mathbb{Z}^n.$$

Similarly, for any $z \in [0, \varepsilon]^n$, there exists $\tau \in H_d \cap [0, \varepsilon]^n$ such that $\tau = z + \alpha e_n$ with $|\alpha| \le \varepsilon$. Then by the above argument, we can find

$$w = k\varepsilon(1, 1, ..., 1, m) + \tau \in H_d$$

for some $0 \le k \le N$ such that

$$|z - w| \le \varepsilon \omega_{\nu}(N) \mod \varepsilon \mathbb{Z}^n.$$
⁽⁷⁹⁾

Now let \tilde{x}_0 be a point in $[0, \varepsilon]^n$ with $x_0 = \tilde{x}_0 \mod \varepsilon \mathbb{Z}^n$, and we apply the above argument for $z = \tilde{x}_0$. Then we can find $\tau \in H_d \cap [0, \varepsilon]^n$ and

$$w = k\varepsilon(1, 1, \dots, 1, m) + \tau \in H_d$$

with some $0 \le k \le N$ such that

$$x_0 - w| = |\tilde{x}_0 - w| \le \varepsilon \omega_{\nu}(N) \mod \varepsilon \mathbb{Z}^n.$$
(80)

On the other hand, recall that the coordinates are shifted so that x = 0. Thus it suffices to find $y \in \mathbb{R}^n$ such that

 $|x-y| = |y| \le MN\varepsilon; \ y = x_0 \mod \varepsilon \mathbb{Z}^n$

and

 $\operatorname{dist}(y, H_d) < \varepsilon \omega_{\nu}(N).$

By (80), there exists $w \in H_d$ such that

$$|x_0 - w| \le \varepsilon \omega_{\nu}(N) \mod \varepsilon \mathbb{Z}^n \tag{81}$$

and

$$|x - w| = |w| \le MN\varepsilon \tag{82}$$

where the last inequality follows from $|(1, 1, ..., 1, m)| \leq M$ and $0 \leq k \leq N$. Given w satisfying (81), we can take $y \in \mathbb{R}^n$ such that

$$|x_0 - y| = 0 \mod \varepsilon \mathbb{Z}^n$$
, and $|y - w| \le \varepsilon \omega_{\nu}(N)$. (83)

Then by (82) and (83)

$$|x - y| = |y| \le |y - w| + |w| \le \varepsilon \omega_{\nu}(N) + MN\varepsilon$$

Also since w is contained in H_d ,

$$\operatorname{dist}(y, H_d) \le |y - w| \le \varepsilon \omega_{\nu}(N)$$

(c) is a direct consequence of (79) since $\omega_{\nu}(N) \to 0$ as $N \to \infty$.

Next we state a version of Dirichlet's approximation theorem, whose proof is based on pigeon-hole principle.

Lemma A.4 (Lemma 2.11 in [13]). For $\alpha_1, ..., \alpha_n \in \mathbb{R}$ and $N \in \mathbb{N}$, there are integers $p_1, ..., p_n, q \in \mathbb{Z}$ with $1 \leq q \leq N$ such that

$$|q\alpha_i - p_i| \le N^{-1/n}.$$

References

- M. Arisawa, Long-time averaged reflection force and homogenization of oscillating Neumann boundary conditions, Ann. Inst. H. Poincare Anal. Lineaire 20 (2003), pp. 293-332.
- [2] S. Armstrong, T. Kuusi, C. Prange, Quantitative analysis of boundary layers in periodic homogenization, ARMA 226 (2017), no.2, pp. 695-741.
- [3] G. Barles, F. Da Lio, Local C^{0,α} estimates for viscosity solutions of Neumann-type boundary value problems, J. Diff. Eqns, 225, No. 1(2006), pp 202–241.
- [4] G. Barles, F. Da Lio, P. L. Lions and P.E. Souganidis, Ergodic problems and periodic homogenization for fully nonlinear equations in half-type domains with Neumann boundary conditions, Indiana University Mathematics Journal 57, No. 5 (2008), pp 2355-2376.
- [5] A. Bensoussan, J. L. Lions, and G. Papanicolaou. Asymptotic analysis for periodic structures, volume 5 of Studies in Mathematics and its Applications. North- Holland Publishing Co., Amsterdam, 1978.
- [6] L. A. Caffarelli and X. Cabre, *Fully Nonlinear Elliptic Equations*, AMS Colloquium Publications, Vol. 43. Providence, RI.
- [7] S. Choi and I. Kim. Homogenization for nonlinear PDEs in general domains with oscillatory Neumann boundary data, Journal de Mathématiques Pures et Appliquées 102, no. 2 (2014) pp. 419–448.
- [8] S. Choi, I. Kim and Lee, Homogenization of Neumann boundary data with fully nonlinear PDEs, Analysis and PDE 6, no. 4 (2013), pp. 951–972.
- M. Crandall, H. Ishiii and P. L. Lions, Users' Guide to viscosity solutions of second order partial differential equations. Bull. Amer. Math. Soc. 27, No.1 (1992) pp. 1–67.
- [10] L. C. Evans, The perturbed test function method for viscosity solutions of nonlinear PDE, Proc. Roy. Soc. of Edinburgh: Section A. 111, No. 3-4 (1989) pp. 359–375.
- [11] L. C. Evans Periodic homogenisation of certain fully nonlinear partial differential equations, Proc. Roy. Soc. of Edinburgh: Section A. 120, No. 3-4 (1992) pp. 245–265.
- [12] W. M. Feldman, Homogenization of the Oscillating Dirichlet boundary condition in general domains, Journal de Mathématiques Pures et Appliquées, 101 no. 5 (2014), pp. 599–622.
- [13] W. M. Feldman and I. C. Kim, Continuity and discontinuity of the boundary layer tail, Ann. Sci. Éc. Norm. Supér. (2017) 50, no.4, pp. 1017–1064.
- [14] N. Guillen and R. W. Schwab, Neumann homogenization via integro-differential operators, DCDS-A 36, no. 7 (2016), pp. 3677—3703.

- [15] D. Gerard-Varet, N. Masmoudi, Homogenization and boundary layers, Acta Math. 209, no.1 (2012) pp. 133-178.
- [16] H. Ishii, P.L. Lions, Viscosity solutions of fully nonlinear second-order elliptic partial differential equations, J. Differential Equations 83 (1) (1990) 26–78.
- [17] L. Kuipers and H. Niederreiter, Uniform Distribution of Sequences, Pure and Applied Mathematics - Wiley, 1974. New York.
- [18] D. Li and K. Zhang, Regularity for fully nonlinear elliptic equations with oblique boundary conditions, Archive for Rational Mechanics and Analysis 228 (3) (2018), 923–967.
- [19] G. M. Lieberman and Neil S. Trudinger, Nonlinear Oblique boundary value problems for Nonlinear Elliptic Equations, Trans. AMS 295 (2)(1985), pp. 509–546.
- [20] Z. Shen and J. Zhuge, Boundary Layers in Periodic Homogenization of Neumann Problems, Comm. Pure Appl. Math 71(2018), pp.2163–2219.
- [21] H. Tanaka, Homogenization of diffusion processes with boundary conditions, Stochastic Analysis and Applications, Adv. Probab. Related Topics, vol. 7, Dekker, New York, 1984, pp. 411-437.
- [22] H. Weyl, Über ein in der Theorie der säkutaren Störungen vorkommendes Problem, Rendiconti del Circolo Matematico di Palemo 330 (1910), pp. 377-407.