

MAGNITUDES REBORN: QUANTITY SPACES AS SCALABLE MONOIDS

DAN JONSSON

ABSTRACT. In ancient Greek mathematics, magnitudes such as lengths were strictly distinguished from numbers. In modern quantity calculus, a distinction is made between quantities and scalars that serve as measures of quantities. The author believes, for reasons apparent from this article, that quantities should play a major rather than a minor role in modern mathematics.

The extended Introduction includes a survey of the historical development and theoretical structure of the pre-modern theory of magnitudes and numbers. In Part 1, work, insights and controversies related to quantity calculus from Euler onward are reviewed. In Parts 2 and 3, we define scalable monoids and, as a special case, quantity spaces; both can be regarded as universal algebras. Scalable monoids are related to rings and modules, and quantity spaces are to scalable monoids as vector spaces are to modules. Subalgebras and homomorphic images of scalable monoids can be formed, and tensor products of scalable monoids can be constructed as well. We also define and investigate congruence relations on scalable monoids, unit elements of scalable monoids, basis-like substructures of scalable monoids and quantity spaces, and scalar representations of elements of quantity spaces. The mathematical theory of quantity spaces is presented with a view to metrological applications.

This article supersedes arXiv:1503.00564 and complements arXiv:1408.5024.

INTRODUCTION

Formulas such as $E = \frac{mv^2}{2}$ or $\frac{\partial T}{\partial t} = \kappa \frac{\partial^2 T}{\partial x^2}$, used to express physical laws, describe relationships between scalars, commonly real numbers. An alternative interpretation of such equations is possible, however. Since the scalars assigned to the variables in these equations are numerical measures of certain quantities, the equations express relationships between these quantities as well. For example, $E = \frac{mv^2}{2}$ can also be interpreted as describing a relation between an energy E , a mass m and a velocity v – three underlying physical quantities, whose existence and properties do not depend on the scalars that may be used to represent them. With this interpretation, though, $\frac{mv^2}{2}$ and similar expressions will have meaning only if operations on quantities, corresponding to operations on numbers, are defined. In other words, an appropriate way of calculating with quantities, a *quantity calculus*, needs to be available.

In a useful survey [3], de Boer describes the development of quantity calculus until the late 20th century, starting with Maxwell's [27] concept of a physical quantity q comprised of a unit quantity $[q]$ of the same kind as q and a scalar $\{q\}$ which is the measure of q relative to $[q]$, so that we can write $q = \{q\}[q]$. De Boer argues, however, that the notion of a physical quantity should be a primitive one, not dependent on other notions, and highlights the contributions by Wallot [35], who defined quantities independently of notions of units and measures in 1926.

The roots of quantity calculus go far deeper in the history of mathematics than to Wallot, however, or even to Maxwell or other scientists of the modern era, such as Fourier [14]; the origins of quantity calculus can be traced back to ancient Greek geometry and arithmetic, as codified in Euclid's *Elements* [11].

Of fundamental importance in the *Elements* is the distinction between *numbers* (*multitudes*) and *magnitudes*. The notion of a number (*arithmos*) is based on that of a "unit" or "monad" (*monas*); a number is "a multitude composed of units". Thus, a number is essentially a positive integer. (A collection of units containing just one unit was not, in principle, considered to be a multitude of units in Greek arithmetic, so 1 was not, strictly speaking, a number.) Numbers can be compared, added and multiplied, and a smaller number can be subtracted from a larger one, but the ratio of two numbers m, n is not itself a number but just a pair $m : n$ expressing relative size. Ratios can, however, be compared; $m : n = m' : n'$ means that $mn' = nm'$. A bigger number m is said to be measured by a smaller number n if $m = kn$ for some number k ; a prime number is a number that is not measured by any other number (or measured only by 1), and m, m' are relatively prime when there is no number (except 1) measuring both.

Magnitudes (*megethos*), on the other hand, are phenomena such as lengths, areas, volumes or times. Unlike numbers, magnitudes are of *different kinds*, and while the magnitudes of a particular kind correspond loosely to numbers, making measurement of magnitudes possible, the magnitudes form a continuum, and there is no distinguished "unit magnitude". In Greek mathematics, magnitudes of the same kind can be compared and added, and a smaller magnitude can be subtracted from a larger one of the same kind, but magnitudes cannot, in general, be multiplied or divided. One can form the ratio of two magnitudes of the same kind, p and q , but this is not a magnitude but just a pair $p : q$ expressing relative size. A greater magnitude q is said to be measured by a smaller magnitude u if there is a number n such that q is equal to u taken n times; we may write this as $q = n \times u$ here.

It is worth noticing that the first three propositions about magnitudes proved by Euclid in the *Elements* are, in the notation used here,

$$\begin{aligned} n \times (u_1 \dot{+} \cdots \dot{+} u_k) &= n \times u_1 \dot{+} \cdots \dot{+} n \times u_k, \\ (n_1 + \cdots + n_k) \times u &= n_1 \times u \dot{+} \cdots \dot{+} n_k \times u, \quad n \times (m \times u) = (nm) \times u, \end{aligned}$$

where n, m, n_1, \dots, n_k are numbers (*arithmoi*), u is a magnitude, u_1, \dots, u_k are magnitudes of the same kind, and $q_1 \dot{+} \cdots \dot{+} q_k$ is the sum of magnitudes. We will return to these identities in connection with Proposition 9.3 in Section 9.

If p and q are magnitudes of the same kind, and there is some magnitude u of this kind and some numbers m, n such that $p = m \times u$ and $q = n \times u$, then p and q are said to be "commensurable"; the ratio of magnitudes $p : q$ can then be represented by the ratio of numbers $m : n$.¹ However, magnitudes may also be "relatively prime"; it may happen that $p : q$ cannot be expressed as $m : n$ for any numbers m, n because there are no m, n, u such that $p = m \times u$ and $q = n \times u$. In view of the Pythagorean philosophical conviction of the primacy of numbers, the discovery of examples of such "incommensurable" magnitudes created a deep crisis in early Greek mathematics [17], a crisis that also affected the foundations of geometry. If ratios of *arithmoi* do not always suffice to represent ratios of magnitudes, it seems

¹It is natural to assume that if $p = m \times u = m' \times u'$ and $q = n \times u = n' \times u'$ then $m : n = m' : n'$, so that the representation of $p : q$ is unique.

that it would not always be possible to express in terms of *arithmoi* the fact that two ratios of magnitudes are equal, as are the ratios of the lengths of corresponding sides of similar triangles. This difficulty was resolved by Eudoxos, who realized that a "proportion", that is, a relation among magnitudes of the form " p is to q as p' is to q' ", conveniently denoted $p : q :: p' : q'$, can be defined numerically even if there is no pair of ratios of *arithmoi* $m : n$ and $m' : n'$ corresponding to $p : q$ and $p' : q'$, respectively, so that $p : q :: p' : q'$ cannot be inferred from $m : n = m' : n'$. Specifically, Eudoxos invented an ingenious indirect way of determining if $p : q :: p' : q'$ in terms of nothing but *arithmoi* by means of a construction similar to the Dedekind cut, as described in Book V of the *Elements*. Using modern terminology, one can say that Eudoxos defined an equivalence relation $::$ between pairs of magnitudes of the same kind numerically, and as a consequence it became possible to conceptualize in terms of positive integers not only ratios of magnitudes corresponding to rational numbers but also ratios of magnitudes corresponding to irrational numbers. Eudoxos thus reconciled the continuum of magnitudes with the discrete *arithmoi*, but in retrospect this feat reduced the incentive to rethink the Greek notion of number, to generalize the *arithmoi*.

To summarize, Greek mathematicians used two notions of muchness, and built a theoretical system around each notion. These systems were connected by relationships of the form $q = n \times u$, where q is a magnitude, n a number and u a magnitude of the same kind as q , foreboding from the distant past Maxwell's conceptualization of a physical quantity, although Euclid did not define magnitudes in terms of units and numbers.

The modern theory of numbers dramatically extends the theory of numbers in the *Elements*. Many types of numbers other than positive integers have been added, and the notion of a number as an element of an algebraic system has come to the forefront. The modern notion of number was not developed by a straight-forward extension of the concept of *arithmos*, however; the initial development of the new notion of number during the Renaissance was strongly inspired by the ancient theory of magnitudes.

The beginning of the Renaissance saw renewed interest in the classical Greek theories of magnitudes and numbers as codified by Euclid, but later these two notions gradually fused into that of a real number. Malet [26] remarks:

As far as we know, not only was the neat and consistent separation between the Euclidean notions of numbers and magnitudes preserved in Latin medieval translations [...], but these notions were still regularly taught in the major schools of Western Europe in the second half of the 15th century. By the second half of the 17th century, however, the distinction between the classical notions of (natural) numbers and continuous geometrical magnitudes was largely gone, as were the notions themselves.

The force driving this transformation was the need for a continuum of numbers as a basis for computation; the discrete *arithmoi* were not sufficient. As magnitudes of the same kind form a continuum, the idea emerged that numbers should be regarded as an aspect of magnitudes. "Number is to magnitude as wetness is to water" said Stevin in *L'Arithmétique* [33], published 1585, and defined a number as "cela, par lequel s'explique la quantité de chascune chose" (that by which one can tell the quantity of anything). Thus, numbers were seen to form a continuum by virtue of their intimate association with magnitudes.

Stevin’s definition of number is rather vague, and it is difficult to see how a magnitude can be associated with a definite number, considering that the numerical measure of a magnitude depends on a choice of a unit magnitude. The notion of number was, however, refined during the 17th century. In *La Geometrie* [8], where Descartes laid the groundwork for analytic geometry, he implicitly identified numbers with *ratios* of two magnitudes, namely lengths of line segments, one of which was considered to have unit length, and in *Universal Arithmetick* [28],² Newton, who had studied Descartes, defined a number with great precision as follows:

By *Number* we mean, not so much a Multitude of Unities, as the abstracted *Ratio* of any Quantity, to another Quantity of the same Kind, which we take for Unity.

In modern terminology, a ratio of quantities of the same kind is a “dimensionless” quantity. Systems of such quantities contain a canonical unit quantity **1**, and addition, subtraction, multiplication and division of dimensionless quantities yield dimensionless quantities.³ This means that there is no essential difference between a number and the corresponding dimensionless quantity (see Proposition 15.6). Quantities, especially so-called dimensionful quantities, that is, classical magnitudes, were thus needed only as a scaffolding for the new notion of numbers, and when this notion had been established its origins fell into oblivion and magnitudes fell out of fashion. As a consequence, quantity calculus had to be developed more or less *ex novo* in the modern era.

While the Greek theory of magnitudes derived from geometry, the new theory of quantities served the needs of modern mathematical physics, which flourished from the second half of the 18th century. In Section IX, Chapter II of *Théorie analytique de la Chaleur* [14], Fourier explained how physical quantities related to the numbers in his equations:

Pour mesurer [des quantités qui entrent dans notre analyse] et les exprimer en nombre on les compare a diverses sortes d’unités, au nombre de cinq, savoir : l’unités de longueur, l’unités de temps, celle de la temperature, celle du poids, et enfin l’unité qui sert a mesurer les quantités de chaleur.

We recognize here the ideas that there are quantities of different kinds and that the number associated with a quantity depends on the choice of a unit quantity of the same kind. Fourier also introduced the powerful notion of “*dimensions*” of quantities.

Using the modern notion of number, we can generalize relationships of the form $q = n \times u$, where n is an *arithmos* and u is a magnitude that measures (divides) q , to relationships of the form $q = \mu \cdot u$, where u is a freely chosen unit quantity of the same kind as q and μ is a number specifying the size of q relative to u , the measure of q relative to u .⁴ We may write $\mu = f(q, u)$, noting that with any two of μ , q and u given, the third is uniquely determined.

²This was a translation of the Latin original *Arithmetica Universalis*, first printed in Cambridge in 1707 and based on lecture notes by Newton for the period 1673 to 1683.

³In Greek mathematics, the product of a length and a length was an area, but Descartes argued in *La Geometrie* that it could also be another length. Descartes did not really multiply lengths, however; he multiplied the ratios of two lengths ℓ_1, ℓ_2 to a fixed length ℓ_0 to obtain the ratio of a third length ℓ to ℓ_0 , using a geometrical construction with similar triangles such that the number representing the ratio of ℓ to ℓ_0 became equal to the product of the number representing the ratio of ℓ_1 to ℓ_0 and the number representing the ratio of ℓ_2 to ℓ_0 . (See the first figure in *La Geometrie*.)

⁴Thus, originally a measure was a magnitude, but this term is now technically used to refer to a number, although a measure can still mean “a standard or unit of measurement”.

Fourier realized that the measure of a quantity may be defined in terms of measures of other quantities, in turn dependent on the units for these quantities. For example, the measure of a velocity depends on a unit of length and a unit of time since a velocity is defined in terms of a length and a time, and the measure of an area indirectly depends on a unit of length.

Specifically, let the measure μ_v of a velocity v be given by $\mu_v = F(\mu_\ell, \mu_t) = F(f(\ell, u_\ell), f(t, u_t))$, where $F(x, y) = xy^{-1}$. Generalizing the magnitude identity $m \times (n \times u) = mn \times u$, we have $\mu \cdot (\nu \cdot u) = \mu\nu \cdot u$. Thus, if $\lambda \neq 0$ then $\lambda\mu \cdot (\lambda^{-1} \cdot u) = \mu \cdot u$, so $f(q, \lambda^{-1} \cdot u) = \lambda\mu = \lambda f(q, u)$, so it follows from the definition of F that

$$LT^{-1}\mu_v = F(f(\ell, L^{-1} \cdot u_\ell), f(t, T^{-1} \cdot u_t))$$

for any non-zero numbers L, T . Also, let the measure μ_a of the area a of a rectangle be given by $\mu_a = G(\mu_\ell, \mu_w) = G(f(\ell, u_\ell), f(w, u_\ell))$, where $G(x, y) = xy$. Then,

$$L^2\mu_a = G(f(\ell, L^{-1} \cdot u_\ell), f(w, L^{-1} \cdot u_\ell)).$$

Fourier pointed out that quantity terms can be equal or combined by addition or subtraction only if they agree with respect to each *exposant de dimension*, having identical arrays of exponents in expressions such as LT^{-1} or L^2 , since otherwise the validity of numerical equations corresponding to quantity equations would depend on an arbitrary choice of units. He thus introduced the principle of dimensional homogeneity for equations that contain quantities.

Note that if $q = \mu \cdot u$ then $\lambda \cdot q = \lambda \cdot (\mu \cdot u) = \lambda\mu \cdot u$, so $f(\lambda \cdot q, u) = \lambda\mu = \lambda f(q, u)$, where u is a fixed unit. Thus, in a sense turning Fourier's argument around, we obtain the equations

$$LT^{-1}\mu_v = F(f(L \cdot \ell, u_\ell), f(T \cdot t, u_t)),$$

$$L^2\mu_a = G(f(L \cdot \ell, u_\ell), f(L \cdot w, u_\ell)).$$

Eliminating the fixed units u_ℓ and u_t , these equations can be written as

$$\Phi(L \cdot \ell, T \cdot t) = LT^{-1}\Phi(\ell, t),$$

$$\Gamma(L \cdot \ell, L \cdot w) = L^2\Gamma(\ell, w),$$

and we can regard Φ and Γ as quantity-valued functions because the units for velocity and area are fixed since they depend on the fixed units for length and time. The bilinearity properties of Φ and Γ suggest that we write $\Phi(\ell, t)$ as ℓt^{-1} and $\Gamma(\ell, w)$ as ℓw . Generalizing this, we may introduce the idea that quantities of the same or different kinds can be multiplied and divided, suggesting that we can form arbitrary expressions of the form $\prod_{i=1}^n q_i^{k_i}$, where k_i are integers. It should be emphasized, however, that Fourier did not formally define multiplication of quantities, so we are talking about a dual line of thought that he could have pursued but apparently did not.

It is clear that the Greek mathematicians' distinction between numbers and magnitudes is closely related to the modern distinction between scalars and quantities. In view of Fourier's contribution, it may be said that the foundation of a modern quantity calculus incorporating this distinction and treating quantities as mathematical objects as real as numbers was laid early in the 19th century. Subsequent progress in this area of mathematics has not been fast and straight-forward, however. In his survey from 1994, de Boer noted that the modern theory of quantities had not yet met its Euclid; he concluded that "a satisfactory axiomatic foundation for the quantity calculus" had not yet been formulated [3].

Gowers [15] points out that many mathematical constructs are not defined directly by describing their essential properties, but indirectly by *construction-definitions*, specifying constructions that can be shown to have these properties.⁵ For example, an ordered pair (x, y) may be defined by a construction-definition as a set $\{x, \{y\}\}$; it can be shown that this construction has the required properties, namely that $(x, y) = (x', y')$ if and only if $x = x'$ and $y = y'$. Many contemporary formalizations of the notion of a quantity use definitions relying on constructions, often defining quantities in terms of (something like) scalar-unit pairs, in the tradition from Maxwell. (See Sections 4 and 11 for some specifics.) However, this is rather like defining a vector as a coordinates-basis pair rather than as an element of a vector space, the modern definition.

Although magnitudes are illustrated by line segments in the *Elements*, the notion of a magnitude is abstract and general. Remarkably, Euclid dealt with this notion in a very modern way. While he carefully defined other important objects such as points, lines and numbers in terms of inherent properties, there is no statement about what a magnitude "is". Instead, magnitudes are characterized by how they relate to other magnitudes through their roles in a system of magnitudes, to paraphrase Gowers [16].

In the same spirit, that of modern algebra, quantities are defined in this article simply as elements of a *quantity space*. Thus, the focus is moved from individual quantities and operations on them to the systems to which the quantities belong, meaning that the notion of quantity calculus will give way to that of a quantity space. This article elaborates on the notion of a quantity space introduced in [19, 20].

In the conceptual framework of universal algebra, a quantity space is just a certain *scalable monoid* $(X, *, (\omega_\lambda)_{\lambda \in R}, 1_Q)$, where X is the underlying set of the algebra, $(X, *, 1_Q)$ is a monoid where we write $*(x, y)$ as xy , and $\omega_\lambda(x)$ is a scalar product $\lambda \cdot x$ such that λ belongs to a fixed ring R , $x \in X$, $\omega_1(x) = x$ for all $x \in X$, $\omega_\lambda(\omega_\kappa(x)) = \omega_{\lambda\kappa}(x)$ for all $\lambda, \kappa \in R$, $x \in X$, and $\omega_\lambda(xy) = \omega_\lambda(x)y = x\omega_\lambda(y)$ for all $\lambda \in R$, $x, y \in X$.

A scalable monoid X is partitioned into *orbit classes*, which are equivalence classes with respect to the relation \sim defined by $x \sim y$ if and only if $\omega_\alpha(x) = \omega_\beta(y)$ for some $\alpha, \beta \in R$. There is no global operation $(x, y) \mapsto x + y$ defined on X , but within each orbit class that contains a unit element addition of its elements is induced by the addition in R , and multiplication of orbit classes is induced by the multiplication of elements of X .

Quantity spaces are to scalable monoids as vector spaces are to modules. A quantity space is a *free commutative* scalable monoid over a *field* K . As mentioned, quantities are just elements of quantity spaces, and dimensions are their orbit classes.

The rest of this article is divided into three parts. Part 1 provides additional background, Part 2 deals with scalable monoids, and in Part 3 scalable monoids are specialized to quantity spaces.

⁵There are mathematical objects for which only construction-definitions are available, so that the mathematician's task becomes to find the properties of these constructions. A major example is the natural numbers, which were created by God, as Kronecker put it, leaving it to humans to discover their properties.

Part 1. From pure numbers to abstract quantities

Sections 1 and 2 below aim to clarify relations between pure and concrete numbers, concrete numbers and quantities, and concrete and abstract quantities. The chain of notions from concrete entities to measures via concrete and abstract quantities is also considered. Sections 3 and 4 contain a review of previous research on systems of abstract quantities, leading up to a list of twelve properties of such systems and three axioms from which these properties can be derived.

1. CONCRETE NUMBERS AND QUANTITIES

The numbers defined by Newton were *pure* or *abstract* numbers, without any designation, but there has always been a practical need for a notion that addresses questions about how many or how much *of what*. Reasoning in terms of magnitudes or quantities makes it possible to answer such questions, but there is an alternative notion which rose to prominence simultaneously with the demise of magnitudes [32], namely *concrete* numbers, that is, numbers associated with the things being counted or measured. For example, the Second Part of Euler's *Einleitung zur Rechen-Kunst* [12] deals with such numbers, also known as *benannten Zahlen*, in some detail.

To understand the relationship between concrete numbers and quantities, let us consider how one can use them to distinguish, for example, 1 centimeter (1 cm) from 1 meter (1 m) or 1 gram (1 g). For this purpose, we can take advantage of the fact that quantities are of different kinds. If we interpret 1 cm, 1 m and 1 g as the products $1 \cdot cm$, $1 \cdot m$ and $1 \cdot g$, respectively, where cm and m , on the one hand, and g , on the other hand, are quantities of different kinds, then $1 \cdot cm$ and $1 \cdot g$ are also quantities of different kinds, as are $1 \cdot m$ and $1 \cdot g$, so they are distinct although the associated pure number 1 is the same in both cases. Note that we have, for example, $1 \cdot cm + 2 \cdot cm = 3 \cdot cm$, $1 \cdot m + 50 \cdot cm = 1 \cdot (100 \cdot cm) + 50 \cdot cm = 100 \cdot cm + 50 \cdot cm = 150 \cdot cm$ and $(1 \cdot cm)(2 \cdot g) = 2 \cdot cm g$.

However, we can introduce a notion of different *kinds of numbers* instead of relying on the fact that quantities are of different kinds. For example, we may say that 1 cm, 1 m and 1 g are different kinds of numbers and thus different concrete numbers. To set this interpretation of 1 cm, 1 m and 1 g apart from the previous one, we may write these expressions as 1_{cm} , 1_m and 1_g , respectively, where the subscript indicates what kind of number the concrete number is. This interpretation works smoothly for addition and subtraction of numbers of the same kind; for example, $1_{cm} + 2_{cm} = 3_{cm}$. Multiplication and division of concrete numbers by pure numbers is also straight-forward; for example, $3(2_m) = 6_m$ and $2_m/3 = (2/3)_m$. There are also identities such as $m = 100_{cm}$ and $100_{cm} = 1_{(100_{cm})}$, so $1_m + 50_{cm} = 1_{(100_{cm})} + 50_{cm} = 100_{cm} + 50_{cm} = 150_{cm}$. Multiplication and division of concrete numbers by concrete numbers are more complicated operations, however; for example, $(3_{cm})(2_g) = 6_{(cm g)}$, where $cm g$ is a new kind of numbers which can be regarded as the product of the kinds of numbers cm and g . Such multiplication was not considered in Euler's textbook from 1740, and it was apparently not until the 19th century that products such as $cm g$ started to be used [32].

If kinds of numbers can be multiplied by pure numbers and any two kinds of numbers can be multiplied, one gets the the impression, however, that kinds of numbers are just quantities in disguise. In fact, extending the theory of concrete numbers to accommodate multiplication and division of any concrete numbers would require

developing a calculus of kinds of numbers essentially equivalent to a calculus of quantities, and it is conceptually cleaner to separate quantities from numbers and distinguish different kinds of quantities than to treat quantities as numbers that can be of different kinds.

2. ON CONCRETE AND ABSTRACT QUANTITIES

Recall that while Stevin defined a number as something which is a measure of a magnitude, Maxwell described a quantity as something which is measured by a number relative to a unit. Leaving this historical irony aside, Maxwell's characterization of a quantity as something which can be specified by a number relative to another quantity of the same kind is open to two somewhat different interpretations. On the one hand, the relation $q = \{q\}[u]$, where $\{q\}$ is a number and q and $[u]$ are quantities of the same kind, can be seen as analogous to $v = \lambda u$, where λ is a scalar and v and u are vectors. That is, we can see a quantity as a mathematical object which admits multiplication by a number. Equipping a system of such mathematical objects with other operations as needed and assuming that certain identities hold, we obtain a mathematical system of quantities. We may call quantities in this sense *mathematical* quantities, or alternatively *abstract* quantities.

On the other hand, Maxwell [27] talks about the quantity $[u]$ as a "standard of reference" and a "standard quantity [which] is technically called the Unit", and he discusses concrete standards of length, time and mass such as the standard *mètre* in Paris. This points to another possible interpretation of Maxwell's notion of quantity: a *concrete* quantity, or alternatively a *physical* quantity, is an *attribute* of a *concrete* or *physical entity*, such as a concrete object, process or phenomenon, which can be specified by a number relative to (a corresponding attribute of) a concrete reference entity. For example, the length $\ell(r)$ of a rigid rod r can be expressed as a multiple μ of the length $\ell(\epsilon)$ of a measuring rod ϵ ; we may write $\ell(r) = \mu \cdot \ell(\epsilon)$. More generally, if we let $a(\epsilon)$ denote the attribute of a concrete standard ϵ with respect to some aspect a then the attribute $a(x)$ of a concrete entity x with respect to a can be expressed as a multiple μ of $a(\epsilon)$, that is, $a(x) = \mu \cdot a(\epsilon)$.

It is a wide-spread practice to use the notation of concrete numbers to refer to quantities, especially concrete quantities. Specifically, we may denote $a(\epsilon)$ by $1 a(\epsilon)$ or $1 u$, and $\mu \cdot a(\epsilon)$ by $\mu a(\epsilon)$ or μu ; with this notation, we have $\mu \cdot (1 u) = \mu u$. By contrast we shall use italics for mathematical quantities; thus, the mathematical quantity corresponding to μu can be denoted by u . We conclude that a mathematical quantity *can* be specified by a number μ relative to a unit u by an expression of the form $\mu \cdot u$, whereas a concrete quantity μu is an attribute of a concrete entity that *must* be specified by a number μ relative to a concrete standard or unit $1 u$.⁶

Given a system of physical quantities, we require a corresponding system of mathematical quantities to be an abstract model of the former, meaning that there

⁶That there are two roads to quantities has led to some confusion and debate among physicists and metrologists, and to attempts at clarification. In a seminal contribution 1950, König [22] pointed out the ambiguity of the notion of quantity and called the two camps each favoring one of these interpretations *Synthetiker* and *Realists*. Distinctions similar to that between abstract and concrete quantities were subsequently suggested by several authors. Some pairs of notions corresponding to abstract and concrete quantities, respectively, are "symbolic" and "physical" quantities [31], "(abstract) physical" and "concrete" quantities [3], "abstract" and "measurable" quantities [10], and "magnitudes" and "quantities" [7].

has to exist some kind of structural similarity between the two systems. In particular, as mathematical quantities of the same kind can be added, it should be possible to add certain concrete quantities as well. As an example, two rigid rods can be concatenated, “glued” together to form one longer, straight rod, and we may define the sum of the lengths of the two original rods to be the length of the new rod. We may similarly “concatenate” two metal weights (by putting them in the same pan of a balance scale) or two processes (in a temporal sense), and define the sum of masses and times in terms of such concatenation. Specifically, the sum of the lengths, masses or durations are equal to the length, mass or duration of the concatenated rods, weights or processes. Denoting two rods, weights or processes by x and y , and their lengths, masses or durations by $a(x)$ and $a(y)$, we may write this as $a(x) \dot{+} a(y) = a(x \smile y)$, where $x \smile y$ is the physical entity obtained by concatenating x and y . It is clear that addition of concrete quantities inherits some properties from concatenation of the underlying concrete entities. In particular, we have $(x \smile y) \smile z = x \smile (y \smile z)$ and $x \smile y = y \smile x$, so

$$(a(x) \dot{+} a(y)) \dot{+} a(z) = a(x) \dot{+} (a(y) \dot{+} a(z)), \quad a(x) \dot{+} a(y) = a(y) \dot{+} a(x),$$

so addition of mathematical quantities should be defined to be associative and commutative as well.

It is less straight-forward to construct concrete quantities that can be interpreted as products or inverses of given concrete quantities, however, and this is something that critics of modern quantity calculus have focused on, since it is assumed there that mathematical quantities can be multiplied and have inverses. Hence, the structural similarity of abstract and concrete systems of quantities is called into question.

In 1922, Bridgman wrote in his well-known *Dimensional Analysis* [5] that “[i]t is meaningless to talk of dividing a length by a time: what we actually do is to operate on numbers that are measures of these quantities”. Much later, Emerson [10] argued that “[i]t is impossible to conceive of the distance between the ends of a meter standard, a unit, being divided by the half period of a seconds pendulum, another unit”. Velocity may be a badly chosen example, however. While it is not possible to “divide” a measuring rod by a pendulum, so that the technique used above to define sums of certain quantities does not work in this case, we may, given a length unit $1 u_\ell$ and a time unit $1 u_t$, define a velocity unit $1 (u_\ell \div u_t)$ as the mean velocity of a point that travels the distance $1 u_\ell$ during the time $1 u_t$. We can also define an area unit $1 (u_\ell * u_\ell)$ as the area of a square with sides of length $1 u_\ell$. Thus, $1 (u_\ell \div u_t)$ is an attribute of a movement, and $1 (u_\ell * u_\ell)$ is an attribute of a square. While the operations $*$ and \div in a system of concrete quantities are not identical to multiplication and division, respectively, in an abstract system of quantities, there is a correspondence ϕ between the two systems such that $\phi(1 (u_\ell * u_\ell)) = \phi(1 u_\ell)\phi(1 u_\ell) = u_\ell u_\ell$ and $\phi(1 (u_\ell \div u_t)) = \phi(1 u_\ell)\phi(1 u_t)^{-1} = u_\ell u_t^{-1}$, so a structural similarity exists.

The critics may be on to something nevertheless. It is not clear, for example, what kind of physical entity a physical time unit multiplied by itself, or a concrete mass unit multiplied by itself, would be an attribute of. More generally, it is not clear that any two concrete quantities can be combined in a way that corresponds to multiplication of abstract quantities, and the notion of the inverse of a concrete quantity is also not clear. Thus, the correspondence ϕ may be a partial function only. On the other hand, a partial function may establish a sufficient structural similarity between systems of physical and mathematical quantities.

We will have to leave this difficulty unresolved and unexplored here, though; a full discussion would require another article (and, I believe, another author). Instead, we will try to gain a deeper understanding of the situation by considering an analogy with a more familiar case.

Recall that abstract vectors may be represented by coordinates relative to some basis. On the other hand, abstract vectors may represent different kinds of less abstract objects, such as geometrical vectors, characterized by their direction and extension. (As we know, “vectors” originally meant geometrical vectors.) Furthermore, geometrical vectors may themselves be seen as abstract representations of more concrete geometrical phenomena such as translations or oriented line segments. These facts can be visualized as shown below:

Concrete geometrical entity \rightarrow *Geometrical vector* \rightarrow *Abstract vector* \rightarrow *Coordinates*

For example, the set T of all translations in a Euclidean space is an abelian group under composition of transformations. Thus, the corresponding set G of geometrical vectors is an abelian group under the operation inherited from the set of translations, so G is isomorphic to some abstract vector space V as an abelian group. A scalar product $(\lambda, x) \mapsto \lambda \cdot x$ can also be defined on T , making G with this scalar product inherited from T a vector space isomorphic to V .

On the other hand, if we let the concrete geometrical entities be oriented line segments then the “sum” of \overrightarrow{AB} and \overrightarrow{CD} is defined and equal to \overrightarrow{AD} only if $B = C$, so while each oriented line segment can be interpreted as a geometrical vector, the set of all oriented line segments does not form an abelian group of geometrical vectors. Every oriented line segment \overrightarrow{AB} can be associated with the unique translation that moves A to B , however, so there is still a certain structural similarity between the system of oriented line segments and the abelian group G of geometrical vectors corresponding to the set of translations T . Relations that connect concrete geometrical entities to geometrical vectors need not be strict isomorphisms, though, and this applies to relations between physical entities and physical quantities, too.

Returning to the quantities, we have seen that we can distinguish four levels of notions, corresponding to those in our vector space analogy, as shown below:

Concrete entity \rightarrow *Concrete quantity* \rightarrow *Abstract quantity* \rightarrow *Measure*

Although the arrows between the three left-most notions possibly hide certain complications, this simple scheme helps to clarify some central conceptual issues concerning quantities. It would seem that leaving out either concrete quantities or abstract quantities, using instead one of the conceptual schemes

Concrete entity \rightarrow *Concrete quantity* \rightarrow *Measure*,
Concrete entity \rightarrow *Abstract quantity* \rightarrow *Measure*,

or, even worse, mixing the two schemes, leads to a conceptually less satisfying theory, if not conceptual confusion [31].

Example 2.1. Many-to-one relations between concrete and abstract quantities.

The definition of planar angle quantities has been subject to much debate in the metrological community [4]: should a planar angle be treated as a derived quantity with unit quantity 1 m/m , or should it be treated as a base quantity with unit quantity 1 rad (radian), where the radian is the angle subtended at the center of a circle by an arc that is equal in length to the radius of the circle? That depends on whether we treat a planar angle as a physical or a mathematical quantity. As a

physical quantity, a planar angle θ rad is one expressed by a number and a physical unit quantity such as “the angle subtended at the center of a circle by an arc that is equal in length to the radius of the circle”. However, $\theta^{\text{m/m}}$ is not an attribute of any particular physical entity; in this context, $\theta^{\text{m/m}}$ is the physical quantity θ rad treated as a mathematical quantity, and as such it should be written as $\theta \cdot \text{m/m}$ or $\theta \cdot \mathbf{1}$, meaning that it is a so-called dimensionless quantity.

Similar considerations apply to many other quantities. For example, there has been a parallel debate about whether a solid angle should be treated as a derived quantity with unit quantity $1 \text{ m}^2/\text{m}^2$, or as a base quantity with unit quantity 1 sr (steradian). Again, as a physical quantity, a solid angle $\theta \text{ sr}$ is one expressed by a number and a certain physical solid angle unit, while as a mathematical quantity a solid angle is again a dimensionless quantity $\theta \cdot \text{m}^2/\text{m}^2 = \theta \cdot \mathbf{1}$.

Thus, regarded as physical quantities, planar and solid angles are quantities of different kinds; but their corresponding mathematical quantities are of the same kind. This shows that a correspondence between concrete and abstract quantities may be a many-to-one relation.

Example 2.2. Quantities and quantity vectors.

Many “quantities” in physics are vector quantities or, more correctly, *quantity vectors*; they are geometrical vectors that combine a direction and an extent given by a quantity rather than a number.

For example, a displacement D combines a direction with a length $\ell \text{ m}$ that gives the extension of the displacement, while a force F combines a direction with a quantity $f \text{ N}$ that gives the extent (strength) of the force. The *work* generated by F and D is defined as the *quantity* $FD = f\ell \cos \theta \text{ N m}$, where θ is the oriented angle between F and D .

On the other hand, we can define a *quantity vector* $F \times D$ with the same direction as that of the cross product of F and D and extent $f\ell \sin \theta \text{ N m}$; this is the *torque* created by F and D .⁷

Note that 1 N m is a unit for the physical quantity $FD = f\ell \cos \theta \text{ N m}$ as well as the physical quantity $f\ell \sin \theta \text{ N m}$ associated with the quantity vector $F \times D$, so these physical quantities are of the same kind. Thus, although work and torque are often said to illustrate a situation where “quantities having the same quantity dimension are not necessarily of the same kind” [18], they do not exemplify physical quantities of different kinds corresponding to mathematical quantities of the same kind, at least if we accept that the torque is a quantity vector rather than a vector quantity.

Example 2.3. Fractional powers of quantities.

In quantity equations we sometimes encounter fractional powers of quantities. For example, the Gaussian unit of electrical charge 1 statC is usually defined by

$$(2.1) \quad 1 \text{ statC} = 1 \frac{\text{cm}^{3/2} \text{g}^{1/2}}{\text{s}}.$$

Do we thus have to admit fractional powers of concrete and abstract quantities into the theory of quantities? Several authors do so, but this complicates the interpretation of concrete quantities and the definition of abstract quantities. Fortunately, integral powers of quantities suffice.

⁷The torque can also be defined as the geometrical bivector $F \wedge D$ whose extent is $f\ell \sin \theta \text{ N m}$.

Let c , g , ℓ and t be the abstract quantities corresponding to the concrete quantities 1 statC, 1 g, 1 cm and 1 s, respectively. Suppose that we have

$$(2.2) \quad c^2 = \frac{\ell^3 g}{t^2},$$

where fractional powers of quantities do not occur. If the measures of these quantities are positive, we can write this as

$$(2.3) \quad \kappa = \frac{\lambda^{3/2} \gamma^{1/2}}{\tau},$$

where κ , λ , γ and τ are the measures of c , ℓ , g and t , respectively, relative to some coherent system of units, so we have effectively recovered (2.1). The crucial notion here is that the relation between the concrete quantities 1 statC, 1 cm, 1 g and 1 s is described by the identity (2.2) relating the corresponding abstract quantities; (2.1) does not describe this relation directly. This example suggests that there is no need for fractional powers of quantities in a theory where a distinction is made between concrete and abstract quantities.

3. ALGEBRAIC PROPERTIES OF SYSTEMS OF ABSTRACT QUANTITIES

Some years after Maxwell had called attention to the notion of quantities, Lodge published an article called “The Multiplication and Division of Concrete Quantities” [24]. In retrospect, Lodge’s article hints at some ideas that reappear in modern quantity calculus. Using motivating examples such as

$$\frac{4\frac{1}{2} \text{ miles}}{1 \text{ hour}} \times 40 \text{ minutes} = \frac{4\frac{1}{2} \text{ miles}}{3} \times 2 = 3 \text{ miles}$$

and

$$4 \text{ feet} \times 2 \text{ yards} = 8 \times 1 \text{ foot} \times 1 \text{ yard} = 24 \times 1 \text{ foot} \times 1 \text{ foot} = 24 \text{ square feet},$$

Lodge assumes that concrete quantities can be multiplied and divided by numbers and by other quantities, and sets out to investigate the formal properties of these operations. Although Lodge deals with concrete quantities of the form αu , he takes a step towards a more abstract notion of quantities in this investigation by denoting such concrete quantities by single symbols such as u . A product of a concrete quantity by a number can then be written as $\lambda \cdot u$, and products or ratios of concrete quantities can be written as uv or u/v .

Implicit in Lodge’s examples is the assumption that multiplication of quantities by numbers is associative in the sense that $\alpha \cdot (\beta \cdot (1 u)) = \alpha\beta \cdot (1 u)$ for any numbers α , β and any quantity $1 u$. Hence, $\alpha \cdot \beta u = \alpha\beta u$ since $\gamma \cdot (1 u) = \gamma u$, so

$$(3.1) \quad 1 \cdot u = 1 \cdot \alpha u = \alpha u = u,$$

$$(3.2) \quad \alpha \cdot (\beta \cdot u) = \alpha \cdot (\beta \cdot \gamma u) = \alpha \cdot (\beta\gamma u) = \alpha\beta\gamma u = \alpha\beta \cdot (\gamma u) = \alpha\beta \cdot u,$$

although these identities are not stated explicitly.

Regarding multiplication and division of two quantities $\alpha \cdot a$ and $\beta \cdot b$, Lodge is more explicit and argues that these operations have to obey the laws

$$(3.3) \quad (\alpha \cdot a)(\beta \cdot b) = \alpha\beta \cdot ab,$$

$$(3.4) \quad (\alpha \cdot a)/(\beta \cdot b) = (\alpha/\beta) \cdot (a/b),$$

presuming that a/b and α/β exist. (3.3) is in fact equivalent to

$$(3.5) \quad \lambda \cdot ab = (\lambda \cdot a)b = a(\lambda \cdot b);$$

if (3.1) and (3.3) then trivially (3.5), and if (3.2) and (3.5) then (3.3) (see the proof of Lemma 6.1).

Lodge suggested that "[quantities] might be treated exactly as numbers"; this may be taken to mean that concrete quantities should be treated as mathematical quantities – for example, one should write $\frac{3 \text{ feet}}{1 \text{ second}}$ rather than 3 feet per second. Note, though, that this is a heuristic principle rather than a rigorous theory: physical quantities are not mathematical quantities, and mathematical quantities are not numbers.

After Lodge, no significant progress was made in the development of a rigorous theory of mathematical quantities for quite some time. In 1945, Landolt [23], supposedly encouraged by Wallot's promotion of quantity calculus in *Handbuch der Physik* [35] and inspired by the abstract "modern algebra" expounded in van der Waerden's classical textbook [34], called attention to group operations on systems of quantities. Specifically, he pointed out that the invertible quantities form a group under "*qualitative Verknüpfung*", that is, multiplication of quantities as discussed by Lodge, and that quantities of the same kind form a group under "*intensive Verknüpfung*", that is, addition of quantities.

After Landolt, but apparently independent of him, Fleischmann published an important article focusing on multiplication of *kinds of quantities* (*Größenarten*). In modern terminology, Fleischmann's idea is that a congruence relation \sim can be defined on a given set of quantities, so that the kinds of quantities correspond to the equivalence classes $[q]$, and multiplication of kinds of quantities can be defined by $[p][q] = [pq]$.

Fleischmann states seven *Grundaussagen* about kinds of quantities. First, there are four assertions to the effect that they constitute a group. Then it is asserted that this group is abelian, torsion-free and finitely generated. These assertions imply that the kinds of quantities form a finitely generated free abelian group, so there exists a finite basis for this group. This means that if $\{B_1, \dots, B_n\}$ is such a basis then any kind of quantity Q has a unique representation of the form $Q = \prod_{i=1}^n B_i^{k_i}$, where k_i are integers.

In an Appendix to his survey from 1995, de Boer [3] presented a list of properties of systems of abstract quantities, based on previous work on quantity calculus and meant to be exhaustive. The list of properties below is, in turn, based on de Boers list, but there are also elements from other sources, such as the works by Lodge, Landolt and Fleischmann surveyed in this section.

- (P1) A quantity q can be multiplied by a number λ , and we have the identities $1 \cdot q = q$ and $\alpha \cdot (\beta \cdot q) = (\alpha\beta) \cdot q$.
- (P2) A set of quantities that forms a system of abstract quantities can be partitioned into subsets of quantities of the same kind.
- (P3) $\lambda \cdot q$ is a quantity of the same kind as q .
- (P4) Quantities of the same kind can be added and form an abelian group under addition.
- (P5) If p and q are of the same kind then $\lambda \cdot (p + q) = \lambda \cdot p + \lambda \cdot q$ and $(\alpha + \beta) \cdot q = \alpha \cdot q + \beta \cdot q$.
- (P6) A quantity can be multiplied by a quantity, and the non-zero quantities form an abelian group under multiplication.
- (P7) Multiplication of quantities by numbers and by quantities are related by the identities $\lambda \cdot (pq) = (\lambda \cdot p)q$ and $\lambda \cdot (pq) = p(\lambda \cdot q)$.

- (P8) Kinds of quantities can be multiplied, forming a finitely generated free abelian group under multiplication.
- (P9) If q is a quantity of kind K and q' a quantity of kind K' then qq' is a quantity of kind KK' .
- (P10) If p and q are quantities of the same kind then $r(p+q) = rp + rq$ (and hence $(p+q)r = pr + qr$).
- (P11) For every kind of quantities there is a quantity u such that for every quantity q of this kind we have $q = \mu \cdot u$, where μ is a uniquely determined number. Such a quantity u is called a *unit*.
- (P12) It is possible to select exactly one unit from each kind of quantities in such a way that that if u is the unit of kind K and u' is the unit of kind K' then uu' is the unit of kind KK' . A set of units satisfying this condition is said to be *coherent*. A coherent set of units is a finitely generated free abelian group, isomorphic to the corresponding group of kinds of quantities.

These properties have many implications. For example, as a coherent set U of units is a finitely generated free abelian group under multiplication, there is a basis $\{u_1, \dots, u_n\}$ for U such that each unit u in U has a unique representation of the form $u = \prod_{i=1}^n u_i^{k_i}$, where k_i are integers. The units u_1, \dots, u_n are called “base units”; other units are called “derived units”. In addition, there exists a quantity $\mathbf{1}$ such that $\mathbf{1}q = q\mathbf{1} = q$ for any quantity q , there exists a kind of quantities I such that $IK = KI = K$ for any kind of quantity K , $\mathbf{1}$ is of kind I , and there is a canonical bijection $\lambda \mapsto \lambda \cdot \mathbf{1}$ between the set of numbers and the quantities of kind I .

Note that although (P10) is not mentioned by Lodge, Landolt, Fleischmann or de Boer, it is postulated by Quade [29] and Raposo [30] and derived from other properties by Whitney [36]. Conversely, though, (P1) – (P12) include or imply all items in de Boer’s list.

4. SYSTEMS OF ABSTRACT QUANTITIES FROM FIRST PRINCIPLES

Given an exhaustive list of properties of systems of mathematical quantities, it remains to design some kind of mathematical foundation from which these properties can be derived, a process analogous to Newton’s (ostentative) derivation of Kepler’s laws from first principles or Euclid’s axiomatization of Greek geometry. In reality, description and derivation were intertwined in the development of quantity calculus, but for purposes of exposition I have somewhat artificially separated these processes, focusing on the former in the preceding section and the latter here. We can let the (P1) – (P12) be the properties of quantity systems that we want to derive.

Abstract quantities will be referred to simply as quantities below since only abstract quantities will be considered henceforth.

Carlson [6] defines a quantity as a pair (r, ξ) , where r is a real number and ξ a *pre-unit*, an element of a multiplicatively written vector space over the rational numbers. Multiplication of quantities by numbers (i) is defined by setting $a \cdot (r, \xi) = (ar, \xi)$, and multiplication of quantities (ii) is defined by setting $(r, \xi)(s, \eta) = (rs, \xi\eta)$. Carlson also defines fractional powers of quantities (iii) by setting $(r, \xi)^t = (r^t, \xi^t)$ when t is a rational number; this corresponds to scalar multiplication in an additively written vector space.

Recall that Fleischmann’s *Grundaussagen* imply that there exists a finite basis $\{B_1, \dots, B_n\}$ for the group of kinds of quantities, meaning that every kind of

quantity Q has a unique representation of the form $Q = \prod_{i=1}^n B_i^{k_i}$, where k_i are integers. It similarly follows from axioms given by Carlson that for every system of quantities there exists a set of quantities $\{u_1, \dots, u_n\}$, a *system of fundamental (base) units*, such that every quantity q has a unique expansion $q = \mu \cdot \prod_{i=1}^n u_i^{k_i}$, where k_i are integers, called dimensions by Carlson. A system of fundamental units is analogous to a basis for a vector space, and is not unique. Carlson shows, however, that if the dimensions of two quantities agree relative to one fundamental system of units, then they agree relative to all fundamental systems of units, and he defines *quantities of the same kind* as quantities whose dimensions agree relative to some fundamental system of units, although he does not consider addition and subtraction of quantities of the same kind.

The definitions proposed by Drobot [9] and Whitney [36] are both based on the ideas that the set of quantities – rather than a set of pre-units – is a multiplicatively written vector space, and that this vector space contains a set \mathcal{R} of numbers, so that multiplication of type (i) is reduced to multiplication of type (ii). The exponentiation operation (iii) is present in both authors' definitions; Drobot lets \mathcal{R} be the set of positive real numbers so that r^t is always defined, but Whitney lets \mathcal{R} be all rational or real numbers and r^t be undefined for non-positive numbers r , so in his theory the set of quantities is not quite a multiplicatively written vector space but something more complicated.

Both Drobot and Whitney define, in slightly different ways, sets of quantities of the same kind, called “dimensions” by Drobot and “birays” by Whitney,⁸ and both define addition of quantities $x = \alpha u$ and $y = \beta u$ of the same kind by the identity $x + y = (\alpha + \beta)u$. Letting $[q]$ denote the dimension/biray containing q , both authors define multiplication of dimensions/birays by the identity $[x][y] = [xy]$, and exponentiation by $[x]^t = [x^t]$. Whitney also proves that $x(y + z) = xy + xz$ for any quantities.

Drobot's article [9] appeared in 1953, Whitney's [36] in 1968, and Whitney acknowledged that “[t]he essential features will be found in Drobot”. While Whitney's version of quantity theory is more complete and rigorous than Drobot's, both have peculiar features related to the overloading of numbers and quantities. In Whitney's theory, the number 0 belongs to all birays; note that $0x = 0$ for any quantity x , so this means that there exists a unique zero quantity which can be regarded as a quantity of all kinds. In Drobot's theory, there are only “positive” quantities, so a bigger quantity cannot be subtracted from a smaller one, and there is no zero quantity.

Quade [29] defines systems of quantities by means of a rather cumbersome construction with *one-dimensional vector spaces* as building blocks. As a first step, he defines a quantity system as the union of all vector spaces in a countably infinite set of pairwise disjoint vector spaces over a fixed field of real or complex numbers; the elements of these vector spaces, called *V-elements*, are in effect quantities. Quantities are of the same kind if and only if they belong to the same vector space; naturally, only quantities in the same vector space can be added. Quade also assumes that the set of quantities can be multiplied and form an abelian semigroup,

⁸We note that the term “dimension” is used to refer to a pattern of exponents of numbers by Fourier [14], an exponent of a unit by Carlson [6], a set of quantities by Drobot [9], and an object associated with a set of quantities by Raposo [30]. In this article, a dimension will be defined as a set of quantities.

that for any two one-dimensional vector spaces U and V , the set of products of quantities uv , where $u \in U$ and $v \in V$, is itself a one-dimensional vector space, denoted UV , so that the set of vector spaces is an abelian semigroup under this multiplication. Furthermore, we have $\lambda(uv) = (\lambda u)v$, where u, v are vectors and λ a scalar, and if u and v are non-zero quantities then uv is a non-zero quantity.

This scheme does not accommodate inverses of quantities and general division of quantities, however, so Quade embeds the multiplicative semigroup of quantities into a group of V -element fractions, redefining V -elements as such fractions. The set of all generalized V -elements thus obtained can be partitioned into vector spaces, corresponding to kinds of generalized quantities. The set of these vector spaces is again an abelian group, now a finitely generated free abelian group.

It is worth noting that Quade introduces a relation R_1 such that xR_1y if and only if $\alpha x = \beta y$ for some numbers α, β , not both zero. R_1 is similar to the relation \sim defined in Section 7.1 below, but the definitions are not identical and the relations are used in different ways.

Recently, Raposo [30] has proposed a definition of a system or "space" of quantities which is similar to Quade's but more concise and elegant. By this definition, a space of quantities Q is an algebraic fiber bundle, with fibers of quantities attached to dimensions (kinds of quantities) in a base space assumed to be a finitely generated free abelian group. Each fiber is again a one-dimensional vector space, with scalar product $\lambda \cdot q$. Multiplication of quantities and multiplication of dimensions are defined independently, but are assumed to be compatible in the same sense as for Quade. The quantities constitute an abelian monoid, and it is assumed that $q(r_1 + r_2) = qr_1 + qr_2$ for any quantities q and r_1, r_2 of the same kind, and $\lambda(qr) = (\lambda q)r$ for any scalar λ and quantity q . Although this is technically not part of the definition of a space of quantities, Raposo also assumes that if q and r are non-zero quantities then qr is a non-zero quantity.

Looking back, we can discern three main approaches to the definition of quantity systems. Carlson uses a two-component approach: a quantity is a pair (r, ξ) , where r is a real number and ξ a pre-unit, and operations on quantities are defined in terms of operations on their components. Quade and Raposo use a fiber-bundle approach. In Quade's conceptualization, described in fiber-bundle terms, each element of the base space is identified with the corresponding fiber and is a one-dimensional vector space whose elements are (generalized) quantities; in Raposo's version, the elements of the base space are dimensions with attached fibers which are again one-dimensional vector spaces whose elements are quantities. In Drobot's and Whitney's approach, elements of a subspace \mathcal{R} of a multiplicatively written vector space Q are identified with numbers, and elements of $Q \setminus \mathcal{R}$ are interpreted as quantities in a narrow sense.

In the three main approaches sketched above, properties of systems of quantities are derived from somewhat contrived constructions, supplemented by postulates. Most of the properties of systems of quantities listed in Section 3 can be derived from these definitions, but the tedious details will not be given here – let us instead look at the definition of quantity spaces proposed in this article. It turns out, surprisingly, that the identities (A1), (A2) and (A3) below, corresponding to items (P1), (P7) and a modified version of (P8) that concerns bases as defined by Carlson rather than by Fleischmann, suffice to derive (P1) – (P12). That is, a quantity space may be informally defined as a set of elements, called quantities, such that:

- (A1) A quantity can be multiplied by a number, and we have the identities $1 \cdot q = q$ and $\alpha \cdot (\beta \cdot q) = (\alpha\beta) \cdot q$.
- (A2) A quantity can be multiplied by a quantity, and the quantities form an abelian monoid with a finite *scalable monoid basis*, that is, a finite set of quantities $B = \{b_1, \dots, b_n\}$ such that each quantity has a unique expansion $q = \mu \cdot \prod_{i=1}^n b_i^{k_i}$, where k_i are integers.
- (A3) Multiplication of quantities by numbers and by quantities are related by the identities $\lambda \cdot (pq) = (\lambda \cdot p)q = p(\lambda \cdot q)$.

It is shown in Parts 2 and 3 how these axioms imply (P1) – (P12) and other algebraic properties of (finitely generated) quantity spaces.

Part 2. Scalable monoids

Simply stated, a scalable monoid is a monoid whose elements can be multiplied by numbers and where the identities in (A1) and (A3) hold. Scalable monoids are formally defined and compared to rings and modules in Section 5, and some basic facts about them are presented in Section 6.

Section 7 is concerned with congruences on scalable monoids and related notions such as orbit classes and other congruence classes, homomorphisms and quotient algebras. In Section 8, direct and tensor products of scalable monoids are defined.

Scalable monoids with unit elements are investigated in Section 9; addition of elements in the same orbit class is defined, and additive scalable monoids, ordered scalable monoids and coherent sets of unit elements are discussed.

A construction-definition of a scalable monoid with a coherent set of units, similar to that proposed by Carlson [6], is presented in Section 11.

5. MATHEMATICAL BACKGROUND AND MAIN DEFINITION

A unital associative algebra X over a (unital, associative) ring R is equipped with three kinds of operations on X :

- (1) *addition* of elements of X , a binary operation $+$: $(x, y) \mapsto x + y$ on X such that X equipped with $+$ is an abelian group;
- (2) *multiplication* of elements of X , a binary operation $*$: $(x, y) \mapsto xy$ on X such that X equipped with $*$ is a monoid;
- (3) *scalar multiplication* of elements of X by elements of R , a monoid action $(\alpha, x) \mapsto \alpha \cdot x$ where the multiplicative monoid of R acts on X so that $1 \cdot x = x$ and $\alpha \cdot (\beta \cdot x) = \alpha\beta \cdot x$ for all $\alpha, \beta \in R$ and $x \in X$.

These structures are linked pairwise:

- (a) addition and multiplication are linked by the distributive laws $x(y + z) = xy + xz$ and $(x + y)z = xz + yz$;
- (b) addition and scalar multiplication are linked by the distributive laws $\alpha \cdot (x + y) = \alpha \cdot x + \alpha \cdot y$ and $(\alpha + \beta) \cdot x = \alpha \cdot x + \beta \cdot x$;
- (c) multiplication and scalar multiplication are linked by the bilinearity laws $\alpha \cdot xy = (\alpha \cdot x)y$ and $\alpha \cdot xy = x(\alpha \cdot y)$ [25].

Related algebraic structures can be obtained from unital associative algebras by removing one of the three operations and hence the links between the removed operation and the two others. Two cases are very familiar. A *ring* has only addition and multiplication of elements of X , linked as described in (a). A *module* has only addition of elements of X and scalar multiplication of elements of X by elements of

R , linked as described in (b). The question arises whether it would be meaningful and useful to define an “algebra without an additive group”, with only multiplication of elements of X and scalar multiplication of elements of X by elements of R , linked as described in (c).

It would indeed. It turns out that this notion, a “sibling” of rings and modules, referred to as scalable monoids in this article, makes sense mathematically and is remarkably well suited for modeling systems of quantities. While (non-extended) numbers are elements of rings, specifically fields, quantities are elements of scalable monoids, specifically quantity spaces.

Definition 5.1. Let R be a (unital, associative) ring. A *scalable monoid over R* , or *R -scaloid*, is a monoid X equipped with a *scaling action*

$$\cdot : R \times X \rightarrow X, \quad (\alpha, x) \mapsto \alpha \cdot x,$$

such that for any $\alpha, \beta \in R$ and $x, y \in X$ we have

- (1) $1 \cdot x = x$,
- (2) $\alpha \cdot (\beta \cdot x) = \alpha\beta \cdot x$,
- (3) $\alpha \cdot xy = (\alpha \cdot x)y = x(\alpha \cdot y)$.

We denote the identity element of X by 1_X or $\mathbf{1}$, and set $x^0 = \mathbf{1}$ for any $x \in X$. An *invertible* element of a scalable monoid X is an element $x \in X$ that has a (necessarily unique) *inverse* $x^{-1} \in X$ such that $xx^{-1} = x^{-1}x = \mathbf{1}$. The product xy of invertible elements x, y is invertible with inverse $y^{-1}x^{-1}$.

6. SOME BASIC FACTS ABOUT SCALABLE MONOIDS

The following lemma will be used repeatedly.

Lemma 6.1. *Let X be a scalable monoid over R . For any $x, y \in X$ and $\alpha, \beta \in R$ we have $(\alpha \cdot x)(\beta \cdot y) = \alpha\beta \cdot xy$ and $\alpha\beta \cdot x = \alpha \cdot (\beta \cdot x) = \beta \cdot (\alpha \cdot x) = \beta\alpha \cdot x$.*

Proof. By Definition 5.1, $(\alpha \cdot x)(\beta \cdot y) = \alpha \cdot x(\beta \cdot y) = \alpha \cdot (\beta \cdot xy) = \alpha\beta \cdot xy$. Also,

$$\begin{aligned} \alpha\beta \cdot x &= \alpha \cdot (\beta \cdot x) = \alpha \cdot (\beta \cdot \mathbf{1}x) = \alpha \cdot (\beta \cdot \mathbf{1})x = \\ &(\beta \cdot \mathbf{1})(\alpha \cdot x) = \beta\alpha \cdot \mathbf{1}x = \beta\alpha \cdot x = \beta \cdot (\alpha \cdot x) \end{aligned}$$

since $(\beta \cdot \mathbf{1})(\alpha \cdot x) = \beta\alpha \cdot \mathbf{1}x$ by the first part of the lemma. \square

Let R be ring and X a monoid. It is easy to verify that the *trivial scaling action* of R on X defined by $\lambda \cdot x = x$ for all $\lambda \in R$ and $x \in X$ satisfies conditions (1) – (3) in Definition 5.1, so a monoid equipped with this function is scalable monoid, namely a *trivially scalable* monoid, though effectively just a monoid since any isomorphic monoids are isomorphic as trivially scalable monoids over a fixed ring.

Since every monoid has a unique identity element, the class of all monoids forms a variety of algebras with a binary operation $*$: $(x, y) \mapsto xy$, a nullary operation $\mathbf{1}$: $() \mapsto \mathbf{1}$ and identities

$$x(yz) = (xy)z, \quad \mathbf{1}x = x = x\mathbf{1}.$$

The class of all scalable monoids over a fixed ring R is a variety of algebras in addition equipped with a set of unary operations $\{\omega_\lambda \mid \lambda \in R\}$, derived from the external binary operation \cdot in Definition 5.1 by setting $\omega_\lambda(x) = \lambda \cdot x$ for all $\lambda \in R$ and $x \in X$, and with the additional identities

$$\omega_1(x) = x, \quad \omega_\lambda(\omega_\kappa(x)) = \omega_{\lambda\kappa}(x), \quad \omega_\lambda(xy) = \omega_\lambda(x)y = x\omega_\lambda(y) \quad (\lambda, \kappa \in R),$$

corresponding to identities (1) – (3) in Definition 5.1.

A scalable monoid is thus a universal algebra

$$(X, *, (\omega_\lambda)_{\lambda \in R}, 1_X)$$

with X as underlying set, here called a *unital magma over R* or *unital R -magma*. The general definitions of direct products, subalgebras and homomorphisms in the theory of universal algebras apply. In particular, a subalgebra of a unital R -magma X is a subset Y of X such that $1_X \in Y$ and if $x, y \in Y$ and $\lambda \in R$ then $xy, \lambda \cdot x \in Y$. Also, for given unital R -magmas X and Y , a unital R -magma homomorphism $\phi : X \rightarrow Y$ is a function such that $\phi(xy) = \phi(x)\phi(y)$, $\phi(\lambda \cdot x) = \lambda \cdot \phi(x)$ and $\phi(1_X) = 1_Y$ for any $x, y \in X$ and $\lambda \in R$.

By Birkhoff's theorem [1], varieties are preserved by the operations of forming subalgebras and homomorphic images. Thus, if a unital R -magma X is a scalable monoid over R then a subalgebra of X is also a scalable monoid over R , and a homomorphic image of X is also a scalable monoid over R .

7. CONGRUENCES AND QUOTIENTS

7.1. On the congruence \sim . In ancient Greek mathematics, the notion of a ratio between magnitudes only applied to magnitudes of the same kind, so only these could be commensurable. In this section, we introduce a more radical idea: quantities are of the same kind *if and only if* they are commensurable.

Let $R \cdot x$ denote the *orbit* of $x \in X$ with regard to the action $(\lambda, x) \mapsto \lambda \cdot x$, that is, the set $\{\lambda \cdot x \mid \lambda \in R\}$, and let \approx denote the relation on X such that $x \approx y$ if and only if there is some $t \in X$ such that $x, y \in R \cdot t$. Note that \approx is not an equivalence relation; it is reflexive since $x \in R \cdot x$ for all $x \in X$ and symmetric by construction but not transitive, meaning that the orbits of a monoid action may overlap.

Definition 7.1. Given a scalable monoid X over R , let \sim be the relation on X such that $x \sim y$ if and only if $\alpha \cdot x = \beta \cdot y$ for some $\alpha, \beta \in R$.

Note that $x \sim y$ if and only if $(R \cdot x) \cap (R \cdot y) \neq \emptyset$. We say that x and y are *commensurable* if and only if $x \sim y$; otherwise x and y are *incommensurable*.

Proposition 7.1. *The relation \sim on a scalable monoid X over R is an equivalence relation.*

Proof. The relation \sim is reflexive since $1 \cdot x = 1 \cdot x$ for all $x \in X$, symmetric by construction, and transitive because if $\alpha \cdot x = \beta \cdot y$ and $\gamma \cdot y = \delta \cdot z$ for some $x, y, z \in X$ and $\alpha, \beta, \gamma, \delta \in R$ then

$$\gamma\alpha \cdot x = \gamma \cdot (\alpha \cdot x) = \gamma \cdot (\beta \cdot y) = \beta \cdot (\gamma \cdot y) = \beta \cdot (\delta \cdot z) = \beta\delta \cdot z,$$

where $\gamma\alpha, \beta\delta \in R$. □

An *orbit class* C is an equivalence class for \sim . The orbit class that contains x is denoted $[x]$, and X/\sim denotes the set $\{[x] \mid x \in X\}$.

Proposition 7.2. *If $x \sim y$ then $\lambda \cdot x \sim y$, $x \sim \lambda \cdot y$ and $\lambda \cdot x \sim \lambda \cdot y$ for all $\lambda \in R$.*

Proof. If $x \sim y$ then $\alpha \cdot x = \beta \cdot y$ for some $\alpha, \beta \in R$, so

$$\alpha\lambda \cdot x = \alpha \cdot (\lambda \cdot x) = \lambda \cdot (\alpha \cdot x) = \lambda \cdot (\beta \cdot y) = \beta \cdot (\lambda \cdot y) = \beta\lambda \cdot y,$$

where $\alpha\lambda, \beta\lambda \in R$. □

Corollary 7.1. $\lambda \cdot x \sim x$ and $x \sim \lambda \cdot x$ for all $x \in X$ and $\lambda \in R$.

Proposition 7.3. We have $0 \cdot x = 0 \cdot y$ if and only if $x \sim y$.

Proof. If $\alpha \cdot x = \beta \cdot y$ then $0 \cdot x = 0\alpha \cdot x = 0 \cdot (\alpha \cdot x) = 0 \cdot (\beta \cdot y) = (0\beta) \cdot y = 0 \cdot y$. \square

Thus, for every orbit class C there is a unique $0_C \in C$ such that $0_C = 0 \cdot x$ for all $x \in C$, and if $0_A = 0_B$ then $A = B$; 0_C is the *zero element* of C . It is clear that $\lambda \cdot 0_C = 0_C$ for all $\lambda \in R$, and that $0_{[x]y} = 0_{[xy]}$ and $y0_{[x]} = 0_{[yx]}$ for all $x, y \in X$.

If $x = \alpha \cdot t$ and $y = \beta \cdot t$ then $\beta \cdot x = \beta \cdot (\alpha \cdot t) = \alpha \cdot (\beta \cdot t) = \alpha \cdot y$, so if $x \approx y$ then $x \sim y$. If $t \in R \cdot x$ then $t, x \in R \cdot x$, so $t \approx x$, so $t \sim x$, so $t \in [x]$; hence, $R \cdot x \subseteq [x]$ for all $x \in X$. As a consequence, $\cup_{t \in [x]} R \cdot t = [x]$.

It is instructive to relate the present notion of commensurability to the classical one. We say that x and y are *strongly commensurable* if and only if $x \approx y$; otherwise, x and y are *weakly incommensurable*. Incommensurability of magnitudes in the Pythagorean sense obviously corresponds to weak incommensurability.

We have thus weakened the classical notion of commensurability here, and this makes it possible to reasonably stipulate that two magnitudes (elements of a scalable monoid) are of the same kind if and only if they are commensurable. The deeper significance of the redefinition of commensurability may be said to be that we have shown how to replace the intuitive notion of magnitudes of the same kind by the formally defined notion of commensurable magnitudes.

Proposition 7.4. Let X be a scalable monoid over R . The relation \sim is a congruence on X with regard to the operations $(x, y) \mapsto xy$ and $(\lambda, x) \mapsto \lambda \cdot x$.

Proof. For any $x, x', y, y' \in X$ and $\alpha, \alpha', \beta, \beta' \in R$, we have that if $\alpha \cdot x = \alpha' \cdot x'$ and $\beta \cdot y = \beta' \cdot y'$ then $(\alpha \cdot x)(\beta \cdot y) = (\alpha' \cdot x')(\beta' \cdot y')$, so $\alpha\beta \cdot xy = \alpha'\beta' \cdot x'y'$ by Lemma 6.1. This means that if $x \sim x'$ and $y \sim y'$ then $xy \sim x'y'$. Also, recall that if $x \sim x'$ then $\lambda \cdot x \sim \lambda \cdot x'$ for any $\lambda \in R$. \square

We can thus define a binary operation on X/\sim by setting $[x][y] = [xy]$ (so that if $A, B \in X/\sim$, $a \in A$ and $b \in B$ then $ab \in AB \in X/\sim$). We can also set $\lambda \cdot [x] = [\lambda \cdot x]$ and $1_{X/\sim} = [1_X]$. Given these definitions, the surjective function $\phi : X \rightarrow X/\sim$ given by $\phi(x) = [x]$ satisfies the conditions

$$\phi(xy) = \phi(x)\phi(y), \quad \phi(\lambda \cdot x) = \lambda \cdot \phi(x), \quad \phi(1_X) = 1_{X/\sim}.$$

These identities induce a unital R -magma structure on X/\sim , and by Birkhoff's theorem X/\sim is an R -scaloid, so ϕ is a scalable monoid homomorphism. Thus, Proposition 7.4, which is expressed in terms of congruences, leads to Proposition 7.5, expressed in terms of homomorphisms.

Proposition 7.5. If X is a scalable monoid over R then the quotient space X/\sim is a scalable monoid over R , and the function

$$\phi : X \rightarrow X/\sim, \quad x \mapsto [x],$$

is a surjective scalable monoid homomorphism.

In this case, $\lambda \cdot [x] = [\lambda \cdot x] = [x]$ by Corollary 7.1, so we have the following fact.

Proposition 7.6. If X is a scalable monoid then X/\sim is a (trivially scalable) monoid.

7.2. On congruences of the forms $\sim_{\mathcal{M}}$ and $\sim_{\mathcal{M}}$. In a monoid we have $x(yz) = (xy)z$ and $\mathbf{1}x = x = x\mathbf{1}$, so a submonoid \mathcal{M} of a scalable monoid X can act as a monoid on X by left or right multiplication. In particular, we can define a monoid action $(m, x) \mapsto m \star x$ on a scalable monoid X by setting $m \star x = mx$ for any $m \in \mathcal{M}$ and $x \in X$. For any $x \in X$, the orbit of x with regard to this action is the right coset $\mathcal{M}x = \{mx \mid m \in \mathcal{M}\}$. Definition 7.2 below is analogous to Definition 7.1, interpreting left multiplication as a monoid action.

Definition 7.2. Let X be a scalable monoid and \mathcal{M} a submonoid of X . Then $\sim_{\mathcal{M}}$ is the relation on X such that $x \sim_{\mathcal{M}} y$ if and only if $mx = ny$ for some $m, n \in \mathcal{M}$.

Proposition 7.7. *If X is a scalable monoid and \mathcal{M} a commutative submonoid of X then $\sim_{\mathcal{M}}$ is an equivalence relation on X .*

Proof. The relation $\sim_{\mathcal{M}}$ is reflexive since $1_X x \sim_{\mathcal{M}} 1_X x$ for all $x \in X$, symmetric by construction, and transitive because if $mx = ny$ and $m'y = n'z$ for $m, n, m', n' \in \mathcal{M}$ then $m'mx = m'ny = nm'y = nn'z$, where $m'm, nn' \in \mathcal{M}$. \square

We denote the equivalence class $\{t \mid t \sim_{\mathcal{M}} x\}$ for $\sim_{\mathcal{M}}$ by $[x]_{\mathcal{M}}$, and the set of equivalence classes $\{[x]_{\mathcal{M}} \mid x \in X\}$ by X/\mathcal{M} .

The *center* of a scalable monoid X , denoted $Z(X)$, is the set of elements of X each of which commutes with all elements of X ; clearly, $1_X \in Z(X)$. A *central submonoid* of a scalable monoid X is a submonoid \mathcal{M} of X such that $\mathcal{M} \subseteq Z(X)$. We have the following corollary of Proposition 7.7.

Corollary 7.2. *If X is a scalable monoid and \mathcal{M} a central submonoid of X then $\sim_{\mathcal{M}}$ is an equivalence relation on X .*

Results analogous to Propositions 7.4 and 7.5 hold for central submonoids.

Proposition 7.8. *Let X be a scalable monoid and \mathcal{M} a central submonoid of X . Then the relation $\sim_{\mathcal{M}}$ is a congruence on X with regard to the operations $(x, y) \mapsto xy$ and $(\lambda, x) \mapsto \lambda \cdot x$.*

Proof. If $x, x', y, y' \in X$ and $m, m', n, n' \in \mathcal{M}$ then $mx = m'x'$ and $ny = n'y'$ implies $(mx)(ny) = (m'x')(n'y')$ so that $(mn)(xy) = (m'n')(x'y')$. Hence, if $x \sim_{\mathcal{M}} x'$ and $y \sim_{\mathcal{M}} y'$ then $xy \sim_{\mathcal{M}} x'y'$ since $mn, m'n' \in \mathcal{M}$.

Also, if $nx = n'x'$ for some $n, n' \in \mathcal{M}$ then $\lambda \cdot nx = \lambda \cdot n'x'$ for all $\lambda \in R$, so $n(\lambda \cdot x) = n'(\lambda \cdot x')$. Hence, if $x \sim_{\mathcal{M}} x'$ then $\lambda \cdot x \sim_{\mathcal{M}} \lambda \cdot x'$. \square

Corollary 7.3. *Let X be a commutative scalable monoid and \mathcal{M} a submonoid of X . Then the relation $\sim_{\mathcal{M}}$ is a congruence on X with regard to the operations $(x, y) \mapsto xy$ and $(\lambda, x) \mapsto \lambda \cdot x$.*

We can thus define two operations on X/\mathcal{M} by setting $[x]_{\mathcal{M}}[y]_{\mathcal{M}} = [xy]_{\mathcal{M}}$ and $\lambda \cdot [x]_{\mathcal{M}} = [\lambda \cdot x]_{\mathcal{M}}$. We also set $1_{X/\mathcal{M}} = [1_X]_{\mathcal{M}}$. Given these definitions, the surjective function $\phi_{\mathcal{M}} : X \rightarrow X/\mathcal{M}$ defined by $\phi_{\mathcal{M}}(x) = [x]_{\mathcal{M}}$ satisfies the conditions

$$\phi_{\mathcal{M}}(xy) = \phi_{\mathcal{M}}(x)\phi_{\mathcal{M}}(y), \quad \phi_{\mathcal{M}}(\lambda \cdot x) = \lambda \cdot \phi_{\mathcal{M}}(x), \quad \phi_{\mathcal{M}}(1_X) = 1_{X/\mathcal{M}}.$$

These identities induce a unital R -magma structure on X/\mathcal{M} , and by Birkhoff's theorem X/\mathcal{M} is an R -scaloid, so $\phi_{\mathcal{M}}$ is a scalable monoid homomorphism. Proposition 7.8 thus corresponds to the following result about homomorphisms.

Proposition 7.9. *If X is a scalable monoid over R and \mathcal{M} a central submonoid of X then the quotient space X/\mathcal{M} is a scalable monoid over R and the function*

$$\phi_{\mathcal{M}} : X \rightarrow X/\mathcal{M}, \quad x \mapsto [x]_{\mathcal{M}}$$

is a surjective scalable monoid homomorphism.

Recall that a subalgebra of a scalable monoid X is itself a scalable monoid, namely, a submonoid \mathcal{M} of X such that $\lambda \cdot x \in \mathcal{M}$ for every $\lambda \in R$ and $x \in \mathcal{M}$; we call \mathcal{M} a *scalable submonoid* of X . A *central* scalable submonoid of X is defined in the same way as a central submonoid of X .

For any central scalable submonoid \mathcal{M} of X we can define a congruence $\sim_{\mathcal{M}}$ on X in the same way as we defined $\sim_{\mathcal{M}}$. Hence, we can define $[x]_{\mathcal{M}}$, X/\mathcal{M} , $[x]_{\mathcal{M}}[y]_{\mathcal{M}}$, $\lambda \cdot [x]_{\mathcal{M}}$ and $1_{X/\mathcal{M}}$ by just substituting \mathcal{M} for \mathcal{M} .

For any central scalable submonoid \mathcal{M} , we have $[\lambda \cdot x]_{\mathcal{M}} = [x]_{\mathcal{M}}$ since $\mathbf{1}(\lambda \cdot x) = \lambda \cdot x = \lambda \cdot \mathbf{1}x = (\lambda \cdot \mathbf{1})x$, where $\mathbf{1}, \lambda \cdot \mathbf{1} \in \mathcal{M}$, so that $(\lambda \cdot x) \sim_{\mathcal{M}} x$ for any $x \in X$. Hence, $\lambda \cdot [x]_{\mathcal{M}} = [\lambda \cdot x]_{\mathcal{M}} = [x]_{\mathcal{M}}$ for any $\lambda \in R$ and $[x]_{\mathcal{M}} \in X/\mathcal{M}$, so while X/\mathcal{M} is a scalable monoid, X/\mathcal{M} is a (trivially scalable) monoid.

Furthermore, if $\alpha \cdot x = \beta \cdot y$ for some $\alpha, \beta \in R$ then $(\alpha \cdot \mathbf{1})x = \alpha \cdot \mathbf{1}x = \beta \cdot \mathbf{1}y = (\beta \cdot \mathbf{1})y$. Thus $x \sim y$ implies $x \sim_{\mathcal{M}} y$ for any central scalable monoid \mathcal{M} of X since $\lambda \cdot \mathbf{1} \in \mathcal{M}$ for any $\lambda \in R$ and any \mathcal{M} . Conversely, note that $R \cdot \mathbf{1}$ is a central scalable submonoid of X and if $x \sim_{R \cdot \mathbf{1}} y$ then $\alpha \cdot \mathbf{1}x = (\alpha \cdot \mathbf{1})x = (\beta \cdot \mathbf{1})y = \beta \cdot \mathbf{1}y$ for some $\alpha, \beta \in R$, so $x \sim_{R \cdot \mathbf{1}} y$ implies $x \sim y$. Thus, $x \sim_{R \cdot \mathbf{1}} y$ if and only if $x \sim y$, so $x \sim_{\mathcal{M}} y$ generalizes $x \sim y$.

8. DIRECT AND TENSOR PRODUCTS OF SCALABLE MONOIDS

Let X and Y be scalable monoids. The *direct product* of X and Y , denoted $X \boxtimes Y$, is the set $X \times Y$ equipped with the binary operation

$$\begin{aligned} * : (X \times Y) \times (X \times Y) &\rightarrow X \times Y, \\ (\langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle) &\mapsto \langle x_1, y_1 \rangle \langle x_2, y_2 \rangle := \langle x_1 x_2, y_1 y_2 \rangle \end{aligned}$$

and the external binary operation

$$\cdot : R \times (X \times Y) \rightarrow X \times Y, \quad (\lambda, \langle x, y \rangle) \mapsto \lambda \cdot \langle x, y \rangle := \langle \lambda \cdot x, \lambda \cdot y \rangle.$$

Straight-forward calculations show that $X \boxtimes Y$ is a scalable monoid with $*$ as monoid multiplication, \cdot as scaling action and $\langle 1_X, 1_Y \rangle$ as identity element.

The direct product of scalable monoids is a generic product, applicable to any universal algebra. Another kind of product, which exploits the fact that $(\lambda \cdot x)y = \lambda \cdot xy = x(\lambda \cdot y)$ in scalable monoids, namely the *tensor product*, turns out to be more useful in many cases.

Definition 8.1. Given scalable monoids X and Y over R , let \sim_{\otimes} be the binary relation on $X \times Y$ such that $(x_1, y_1) \sim_{\otimes} (x_2, y_2)$ if and only if $(\alpha \cdot x_1, \beta \cdot y_1) = (\beta \cdot x_2, \alpha \cdot y_2)$ for some $\alpha, \beta \in R$.

Proposition 8.1. *Let X and Y be scalable monoids over R . Then \sim_{\otimes} is an equivalence relation on $X \times Y$.*

Proof. \sim_{\otimes} is reflexive since $(1 \cdot x, 1 \cdot y) = (1 \cdot x, 1 \cdot y)$, and symmetric by construction. If $(\alpha \cdot x_1, \beta \cdot y_1) = (\beta \cdot x_2, \alpha \cdot y_2)$ and $(\gamma \cdot x_2, \delta \cdot y_2) = (\delta \cdot x_3, \gamma \cdot y_3)$ then

$$\begin{aligned} (\gamma \cdot (\alpha \cdot x_1), \delta \cdot (\beta \cdot y_1)) &= (\gamma \cdot (\beta \cdot x_2), \delta \cdot (\alpha \cdot y_2)), \\ (\beta \cdot (\gamma \cdot x_2), \alpha \cdot (\delta \cdot y_2)) &= (\beta \cdot (\delta \cdot x_3), \alpha \cdot (\gamma \cdot y_3)). \end{aligned}$$

Thus, we have

$$\begin{aligned} (\gamma\alpha \cdot x_1, \delta\beta \cdot y_1) &= (\gamma\beta \cdot x_2, \delta\alpha \cdot y_2) = (\beta\gamma \cdot x_2, \alpha\delta \cdot y_2) = \\ &(\beta\delta \cdot x_3, \alpha\gamma \cdot y_3) = (\delta\beta \cdot x_3, \gamma\alpha \cdot y_3), \end{aligned}$$

where $\gamma\alpha, \delta\beta \in R$, so \sim_{\otimes} is transitive as well. \square

For any $x \in X$ and $y \in Y$, let $x \otimes y$ denote the equivalence class

$$\{(s, t) \mid (s, t) \in X \times Y, (s, t) \sim_{\otimes} (x, y)\},$$

and let $X \otimes Y$ denote the set $\{x \otimes y \mid x \in X, y \in Y\}$.

Proposition 8.2. *Let X, Y be scalable monoids over R , $x \in X$ and $y \in X$. Then $(\lambda \cdot x) \otimes y = x \otimes (\lambda \cdot y)$ for every $\lambda \in R$.*

Proof. We have $(1 \cdot (\lambda \cdot x), \lambda \cdot y) = (\lambda \cdot x, 1 \cdot (\lambda \cdot y))$, so $(\lambda \cdot x, y) \sim_{\otimes} (x, \lambda \cdot y)$, meaning that $(\lambda \cdot x) \otimes y = x \otimes (\lambda \cdot y)$. \square

Proposition 8.3. *Let X, Y be scalable monoids over R , and set $(x_1 \otimes y_1)(x_2 \otimes y_2) = x_1x_2 \otimes y_1y_2$ and $\lambda \cdot x \otimes y = (\lambda \cdot x) \otimes y$. Also set $1_{X \otimes Y} = 1_X \otimes 1_Y$. With these definitions, $X \otimes Y$ is a scalable monoid over R .*

Proof. $X \otimes Y$ is a monoid since

$$\begin{aligned} (1_X \otimes 1_Y)(x \otimes y) &= 1_Xx \otimes 1_Yy = x \otimes y = x1_X \otimes y1_Y = (x \otimes y)(1_X \otimes 1_Y), \\ ((x_1 \otimes y_1)(x_2 \otimes y_2))(x_3 \otimes y_3) &= (x_1x_2 \otimes y_1y_2)(x_3 \otimes y_3) = (x_1x_2)x_3 \otimes (y_1y_2)y_3 = \\ x_1(x_2x_3) \otimes y_1(y_2y_3) &= (x_1 \otimes y_1)(x_2x_3 \otimes y_2y_3) = (x_1 \otimes y_1)((x_2 \otimes y_2)(x_3 \otimes y_3)). \end{aligned}$$

Furthermore,

$$\begin{aligned} 1 \cdot x \otimes y &= (1 \cdot x) \otimes y = x \otimes y, \\ \alpha \cdot (\beta \cdot x \otimes y) &= \alpha \cdot (\beta \cdot x) \otimes y = (\alpha \cdot (\beta \cdot x)) \otimes y = (\alpha\beta \cdot x) \otimes y = \alpha\beta \cdot x \otimes y, \\ \lambda \cdot (x_1 \otimes y_1)(x_2 \otimes y_2) &= \lambda \cdot x_1x_2 \otimes y_1y_2 = (\lambda \cdot x_1x_2) \otimes y_1y_2 = \\ (\lambda \cdot x_1)x_2 \otimes y_1y_2 &= ((\lambda \cdot x_1) \otimes y_1)(x_2 \otimes y_2) = (\lambda \cdot x_1 \otimes y_1)(x_2 \otimes y_2), \\ \lambda \cdot (x_1 \otimes y_1)(x_2 \otimes y_2) &= \lambda \cdot x_1x_2 \otimes y_1y_2 = x_1x_2 \otimes (\lambda \cdot y_1y_2) = \\ x_1x_2 \otimes y_1(\lambda \cdot y_2) &= (x_1 \otimes y_1)(x_2 \otimes (\lambda \cdot y_2)) = (x_1 \otimes y_1)(\lambda \cdot x_2 \otimes y_2), \end{aligned}$$

so $X \otimes Y$ is a scalable monoid. \square

Corollary 8.1. *If X, Y, Z are scalable monoids over R then $(X \otimes Y) \otimes Z$ and $X \otimes (Y \otimes Z)$ are scalable monoids over R .*

Proposition 8.4. *$\phi : (x \otimes y) \otimes z \mapsto x \otimes (y \otimes z)$ is a scalable monoid isomorphism $(X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$.*

Proof. We have

$$\begin{aligned} \phi(((x_1 \otimes y_1) \otimes z_1)((x_2 \otimes y_2) \otimes z_2)) &= \phi(((x_1 \otimes y_1)(x_2 \otimes y_2)) \otimes z_1z_2) = \\ \phi((x_1x_2 \otimes y_1y_2) \otimes z_1z_2) &= x_1x_2 \otimes (y_1y_2 \otimes z_1z_2) = x_1x_2 \otimes ((y_1 \otimes z_1)(y_2 \otimes z_2)) = \\ (x_1 \otimes (y_1 \otimes z_1))(x_2 \otimes (y_2 \otimes z_2)) &= \phi((x_1 \otimes y_1) \otimes z_1)\phi((x_2 \otimes y_2) \otimes z_2) \end{aligned}$$

and

$$\begin{aligned} \phi(\lambda \cdot (x \otimes y) \otimes z) &= \phi((x \otimes y) \otimes (\lambda \cdot z)) = x \otimes (y \otimes (\lambda \cdot z)) = \\ x \otimes (\lambda \cdot y \otimes z) &= \lambda \cdot x \otimes (y \otimes z) = \lambda \cdot \phi((x \otimes y) \otimes z). \end{aligned}$$

Also,

$$\phi(1_{(X \otimes Y) \otimes Z}) = \phi((1_X \otimes 1_Y) \otimes 1_Z) = 1_X \otimes (1_Y \otimes 1_Z) = 1_{X \otimes (Y \otimes Z)}.$$

Thus, ϕ is a scalable monoid homomorphism $(X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$, and similarly $\phi' : x \otimes (y \otimes z) \mapsto (x \otimes y) \otimes z$ is a scalable monoid homomorphism $X \otimes (Y \otimes Z) \rightarrow (X \otimes Y) \otimes Z$ such that $\phi' \circ \phi = \text{Id}_{(X \otimes Y) \otimes Z}$ and $\phi \circ \phi' = \text{Id}_{X \otimes (Y \otimes Z)}$, so ϕ is a scalable monoid isomorphism. \square

9. ORBIT CLASSES AND UNIT ELEMENTS

Recall the principle that magnitudes of the same kind can be added and subtracted, whereas magnitudes of different kinds cannot be combined by these operations. Also recall the idea that a quantity q can be represented by a "unit" $[q]$ and a number $\{q\}$ specifying "[how many] times the [unit] is to be taken in order to make up" q [27]. As shown below, there is a connection between these two notions.

Specifically, it may happen that $R \cdot u \supseteq [u]$ for some $u \in [u]$, and if in addition a natural uniqueness condition is satisfied we may regard u as a unit of measurement for $[u]$. If such a unit exists then a sum of magnitudes in $[u]$ can be defined by the construction described below.

Definition 9.1. Let C be an orbit class in a scalable monoid over R . A *generating element* for C is some $u \in C$ such that for every $x \in C$ there is some $\rho \in R$ such that $x = \rho \cdot u$. A *unit element* for C is a generating element u for C such that if $\rho \cdot u = \rho' \cdot u$ then $\rho = \rho'$.

By this definition, if u is a generating element for C then $R \cdot u \supseteq C$, but recall that $R \cdot u \subseteq [u]$, so actually $R \cdot u = C = [u]$, since $u \in [u]$ and $u \in C$ so that $[u] \cap C \neq \emptyset$. For any scalable monoid X and $x, u \in X$, $[x] \subseteq R \cdot u$ implies $x \sim u$ since $1 \cdot x = \rho \cdot u$ for some $\rho \in R$, and $x \sim u$ implies $[x] = [u] \supseteq R \cdot u$.

It follows from the definitions of zero elements and unit elements that a zero element 0_C cannot be a unit element for C in a scalable monoid over a non-trivial ring. Also note that Lemma 6.1, specifically the identity $\alpha\beta \cdot x = \beta\alpha \cdot x$, implies the following fact.

Proposition 9.1. *If there exists a unit element for some orbit class in a scalable monoid over R then R is commutative.*

We now turn to an argument that leads to Proposition 9.3.

Proposition 9.2. *Let C be an orbit class in a scalable monoid over R . If u and u' are unit elements for C , $\rho \cdot u = \rho' \cdot u'$ and $\sigma \cdot u = \sigma' \cdot u'$ then $(\rho + \sigma) \cdot u = (\rho' + \sigma') \cdot u'$.*

Proof. As $u' \in C$, there is a unique $\tau \in R$ such that $u' = \tau \cdot u$. Thus,

$$(\rho' + \sigma') \cdot u' = (\rho' + \sigma') \cdot (\tau \cdot u) = (\rho' + \sigma')\tau \cdot u = (\rho'\tau + \sigma'\tau) \cdot u = (\rho + \sigma) \cdot u$$

since $\rho \cdot u = \rho' \cdot u' = \rho' \cdot (\tau \cdot u) = \rho'\tau \cdot u$ and $\sigma \cdot u = \sigma' \cdot u' = \sigma' \cdot (\tau \cdot u) = \sigma'\tau \cdot u$, so that $\rho = \rho'\tau$ and $\sigma = \sigma'\tau$. \square

Hence, the sum of two elements of a scalable monoid can be defined as follows.

Definition 9.2. Let X be a scalable monoid over R , and let u be a unit element for an orbit class C . If $x = \rho \cdot u$ and $y = \sigma \cdot u$, we set

$$x + y = (\rho + \sigma) \cdot u.$$

Thus, if $x, y \in \mathbf{C}$ then $x + y = (\rho + \sigma) \cdot u \in R \cdot u = \mathbf{C}$, and if $x \in \mathbf{C}$ then $\lambda \cdot x = \lambda \cdot (\rho \cdot u) = \lambda\rho \cdot u \in R \cdot u = \mathbf{C}$. We note that the sum $x + y$ is given by Definition 9.2 if and only if x and y are commensurable and their orbit class has a unit element. This fact motivates that the notion of magnitudes of the same kind is replaced by that of commensurable magnitudes (see Section 7.1).

It follows immediately from Definition 9.2 that

$$(x + y) + z = x + (y + z), \quad x + y = y + x$$

for all $x, y, z \in \mathbf{C}$, and that

$$x + 0_{\mathbf{C}} = 0_{\mathbf{C}} + x = x$$

for any $x \in \mathbf{C}$ since $0_{\mathbf{C}} = 0 \cdot u$.

If $x = \rho \cdot u$ so that $\lambda \cdot x = \lambda\rho \cdot u$ and $\kappa \cdot x = \kappa\rho \cdot u$ then

$$(\lambda + \kappa) \cdot x = (\lambda + \kappa) \cdot (\rho \cdot u) = (\lambda + \kappa)\rho \cdot u = (\lambda\rho + \kappa\rho) \cdot u = \lambda \cdot x + \kappa \cdot x,$$

and if $x = \rho \cdot u$ and $y = \sigma \cdot u$ so that $\lambda \cdot x = \lambda\rho \cdot u$ and $\lambda \cdot y = \lambda\sigma \cdot u$ then

$$\lambda \cdot (x + y) = \lambda \cdot ((\rho + \sigma) \cdot u) = \lambda(\rho + \sigma) \cdot u = (\lambda\rho + \lambda\sigma) \cdot u = \lambda \cdot x + \lambda \cdot y.$$

A unital ring R has a unique additive inverse -1 of $1 \in R$, and we set

$$-x = (-1) \cdot x$$

for all $x \in X$. If \mathbf{C} has a unit element u and $x = \rho \cdot u$ for some $\rho \in R$ then

$$x + (-x) = -x + x = 0_{\mathbf{C}}$$

since $x + (-x) = \rho \cdot u + (-\rho) \cdot u = (\rho + (-\rho)) \cdot u = 0 \cdot u$ and $-x + x = (-\rho) \cdot u + \rho \cdot u = (-\rho + \rho) \cdot u = 0 \cdot u$, using the fact that $-x = (-1) \cdot (\rho \cdot u) = (-\rho) \cdot u$.

As usual, we may write $x + (-y)$ as $x - y$, and thus $x + (-x)$ as $x - x$.

We have thus shown the following fact.

Proposition 9.3. *If X is a scalable monoid over R and $\mathbf{C} \in X/\sim$ contains a unit element u for \mathbf{C} then \mathbf{C} is a free module over R with $\{u\}$ as a basis.*

Addition in \mathbf{C} is given by Definition 9.2, and scalar multiplication in \mathbf{C} is inherited from the scalar multiplication in X .

Thus, if every orbit class $\mathbf{C} \in X/\sim$ contains a unit element for \mathbf{C} then X is the union of disjoint isomorphic free modules over R , namely the orbit classes. This fact may be compared to Quade's and Raposo's definitions of quantity spaces [29, 30].

Recall that identities corresponding to $(\lambda + \kappa) \cdot x = \lambda \cdot x + \kappa \cdot x$, $\lambda \cdot (x + y) = \lambda \cdot x + \lambda \cdot y$ and $\lambda \cdot (\kappa \cdot x) = \lambda\kappa \cdot x$ were proved in Propositions 1 – 3 in Book V of the *Elements*, so rudiments of Proposition 9.3 were present already in the Greek theory of magnitudes.

10. SCALABLE MONOIDS AND SETS OF UNIT ELEMENTS

Definition 10.1. A *dense* set of elements of a scalable monoid X is a set U of elements of X such that for every $x \in X$ there is some $u \in U$ such that $u \sim x$. A *sparse* set of elements of X is a set U of elements of X such that $u \sim v$ implies $u = v$ for any $u, v \in U$. A *closed* set of elements of X is a set U of elements of X such that if $u, v \in U$ then $uv \in U$. A *set of unit elements* of a scalable monoid X is a set of elements of X each of which is a unit element for some $\mathbf{C} \in X/\sim$.

We call a dense sparse set of unit elements of X a *system of unit elements* for X , and a sparse set of unit elements of X a *partial system of unit elements* for X .

10.1. Additive scalable monoids.

Definition 10.2. An *additive* scalable monoid is a scalable monoid X such that for every $C \in X/\sim$ there is a binary operation

$$+ : C \times C \rightarrow C, \quad (x, y) \mapsto x + y$$

such that C equipped with $+$ is an abelian group and $x(y + z) = xy + xz$ and $(y + z)x = yx + yz$ for all $x \in A$ and $y, z \in B$ for all $A, B \in X/\sim$.

Proposition 10.1. *If a scalable monoid X is equipped with a dense closed set of unit elements U then X is an additive scalable monoid.*

Proof. By Proposition 9.3, each $C \in X/\sim$ is a module since U is dense in X .

For all $x \in A$ and $y, z \in B$ there are $u, v \in U$ such that $[x] = [u]$ and $[y] = [z] = [v]$ since U is dense in X , so $x = \rho \cdot u$, $y = \sigma \cdot v$ and $z = \tau \cdot v$ for some $\rho, \sigma, \tau \in R$, so $yx = \rho\sigma \cdot uv$, $xz = \rho\tau \cdot uv$, $yx = \sigma\rho \cdot vu$ and $zx = \tau\rho \cdot vu$, so

$$\begin{aligned} x(y + z) &= (\rho \cdot u)((\sigma + \tau) \cdot v) = \rho(\sigma + \tau) \cdot uv = (\rho\sigma + \rho\tau) \cdot uv = xy + xz, \\ (y + z)x &= ((\sigma + \tau) \cdot v)(\rho \cdot u) = (\sigma + \tau)\rho \cdot vu = (\sigma\rho + \tau\rho) \cdot vu = yx + zx, \end{aligned}$$

using the fact that uv and vu are unit elements since U is closed. \square

10.2. Ordered scalable monoids. Recall that a *total order* on a set S is a binary relation \leq such that for all $x, y, z \in S$ we have that

- (1) $x \leq y$ or $y \leq x$;
- (2) if $x \leq y$ and $y \leq x$ then $x = y$;
- (3) if $x \leq y$ and $y \leq z$ then $x \leq z$.

Also recall that a (totally) *ordered ring* is a (unital) ring R with a total order \leq such that for all $x, y, z \in R$ we also have that

- (1) if $x \leq y$ then $x + z \leq y + z$;
- (2) if $0 \leq x$ and $0 \leq y$ then $0 \leq xy$.

Well-known facts about inequalities such as $0 \leq 1$ and if $x \leq y$ and $x' \leq y'$ then $x + x' \leq y + y'$ can be derived from the definition of an ordered ring.

Let U be a dense set of unit elements of a scalable monoid X over an ordered ring R . For any $C \in X/\sim$, let \sim_{U_C} be the relation on $U \cap C$ defined by $u \sim_{U_C} v$ if and only if $u = \rho \cdot v$ for some $\rho \in R$ such that $0 \leq \rho$. Then \sim_{U_C} is reflexive since $u = 1 \cdot u$, and transitive since if $u = \rho \cdot v$ and $v = \sigma \cdot w$, where $0 \leq \rho, \sigma$, then $u = \rho\sigma \cdot w$, where $0 \leq \rho\sigma$. Also note that if $u \sim_{U_C} v$, so that $u = \rho \cdot v$, $v = \tau \cdot u$ and $0 \leq \rho$, then $1 \cdot u = \rho \cdot v = \rho\tau \cdot u$ and $1 \cdot v = \tau \cdot u = \tau\rho \cdot v$, so $\rho\tau = \tau\rho = 1$, and $0 \leq \tau$ since $0 \leq \rho$. Thus, $\tau \cdot u = \tau\rho \cdot v = v$, where $0 \leq \tau$, so \sim_{U_C} is symmetric as well. Hence, \sim_{U_C} is an equivalence relation on $U \cap C$.

Definition 10.3. A *consistent* dense set of unit elements is a dense set U of unit elements such for any $C \in X/\sim$ and any $u, v \in U \cap C$ we have $u \sim_{U_C} v$.

Proposition 10.2. *Let X be a scalable monoid over an ordered ring R , and let u, v be unit elements for $C \in X/\sim$ such that $u \sim_{U_C} v$. For any $x \in C$, if $x = \rho \cdot u = \sigma \cdot v$ for some $\rho, \sigma \in R$ and $0 \leq \rho$ then $0 \leq \sigma$.*

Proof. There is some $\tau \in R$ such that $u = \tau \cdot v$ and $0 \leq \tau$. Thus, $\sigma \cdot v = x = \rho \cdot u = \rho \cdot (\tau \cdot v) = \rho\tau \cdot v$, so $\sigma = \rho\tau$, so $0 \leq \sigma$ since $0 \leq \rho, \tau$. \square

Definition 10.4. An *ordered scalable monoid* is a scalable monoid X over an ordered ring R such that for each $C \in X/\sim$ there is a total order \leq_C on C such that for any $a \in A, b \in B, x, y \in C$ and $\lambda \in R$ we have that

- (1) if $0_A \leq_A a$ and $0_B \leq_B b$ then $0_{AB} \leq_{AB} ab$;
- (2) if $x \leq_C y$ and $0 \leq \lambda$ then $\lambda \cdot x \leq_C \lambda \cdot y$.

An ordered *additive* scalable monoid is an additive scalable monoid such that, in addition to (1) and (2), for any $x, y, z \in C$ we have that

- (3) if $x \leq_C y$ then $x + z \leq_C y + z$.

Proposition 10.3. *Let X be a scalable monoid over an ordered ring R , equipped with a consistent dense closed set U of unit elements of X . For every $C \in X/\sim$, there is a unique binary relation \leq_C on C defined by $x \leq_C y$ if and only if $y - x = \rho \cdot u$ for some $\rho \in R$ such that $0 \leq \rho$ and some $u \in U \cap C$, and X with each C ordered by \leq_C is an ordered additive scalable monoid over R .*

Proof. By Proposition 10.1, X is an additive scalable monoid, and by Proposition 10.2 the relation \leq_C does not depend on a choice of unit element in C .

We first show that \leq_C is a total order on C . If $x, y \in C$ then $y - x = \rho \cdot u$ for some $\rho \in R, u \in U \cap C$. Thus, if $0 \leq \rho$ then $x \leq_C y$; if $\rho \leq 0$ then $0 \leq (-\rho)$ and $x - y = -(y - x) = -(\rho \cdot u) = (-1) \cdot (\rho \cdot u) = (-\rho) \cdot u$, so $y \leq_C x$. If $x \leq_C y$ and $y \leq_C x$ then $0 \leq \rho$ and $\rho \leq 0$, so $\rho = 0$, so $y - x = 0_C$, so $x = y$. Also, if $x, y, z \in C$, $y - x = \rho \cdot u$ and $z - y = \sigma \cdot u$ then $z - x = (y - x) + (z - y) = (\rho + \sigma) \cdot u$, so if $x \leq_C y$ and $y \leq_C z$ so that $0 \leq \rho, \sigma$ then $x \leq_C z$ since $0 \leq \rho + \sigma$.

Furthermore, if $0_A \leq_A x$ and $0_B \leq_B y$, meaning that $x = \rho \cdot u$ and $y = \sigma \cdot v$ for some $\rho, \sigma \in R$, where $0 \leq \rho, \sigma$ and $u \in A, v \in B$, then $xy = (\rho \cdot u)(\sigma \cdot v) = \rho\sigma \cdot uv$, where $0 \leq \rho\sigma$ and $uv \in AB$, so $0_{AB} \leq_{AB} xy$. Similarly, if $x \leq_C y$, meaning that $x = \rho \cdot u$ and $y = \sigma \cdot u$ for some $\rho, \sigma \in R$ such that $0 \leq \sigma - \rho$, then $0 \leq \lambda$ implies $0 \leq \lambda(\sigma - \rho) = \lambda\sigma - \lambda\rho$, so $\lambda \cdot x \leq_C \lambda \cdot y$ since $\lambda \cdot y - \lambda \cdot x = \lambda \cdot (\sigma \cdot u) - \lambda \cdot (\rho \cdot u) = (\lambda\sigma - \lambda\rho) \cdot u$. Also, if $x \leq_C y$ then $x + z \leq_C y + z$ since $(y + z) - (x + z) = y - x = (\sigma - \rho) \cdot u$. \square

Let U and V be systems of unit elements of a scalable monoid X over an ordered ring. U and V are said to *define the same orientation* of X if and only if $U \cup V$ is a consistent set of unit elements, meaning that for all $C \in X/\sim$ we have $u \sim_{(U \cup V)_C} v$, where $u \in U \cap C$ and $v \in V \cap C$.

If U and V define the same orientation on X then U, V and $U \cup V$ are consistent, so by Proposition 10.2 each set uniquely defines, on each $C \in X/\sim$, a relation \leq_C such that $x \leq_C y$ if and only if $y - x = \rho \cdot u$ for some $\rho \in R$ such that $0 \leq \rho$ and some u in $U \cap C, V \cap C$ or $(U \cup V) \cap C$, respectively. It is clear that U and V define the same relations \leq_C , namely the same relations as $U \cup V$ defines.

10.3. Scalable monoids and coherent systems of unit elements.

Definition 10.5. A *coherent* system of unit elements for X is a submonoid of X which is a system of unit elements for X .

Proposition 10.4. *Let X be a scalable monoid, \mathcal{U} a coherent system of unit elements for X , and $\mathcal{V} \subseteq \mathcal{U}$ a central submonoid of X . Then X/\mathcal{V} is a scalable monoid, $[v]_{\mathcal{V}} = [\mathbf{1}]_{\mathcal{V}}$ for any $v \in \mathcal{V}$, and $\bar{\mathcal{U}} = \{[u]_{\mathcal{V}} \mid u \in \mathcal{U}\}$ is a coherent system of unit elements for X/\mathcal{V} .*

Proof. As \mathcal{V} is a central submonoid of X , X/\mathcal{V} is a scalable monoid, and if $v \in \mathcal{V}$ then $v \sim_{\mathcal{V}} \mathbf{1}$ since $\mathbf{1}v = v\mathbf{1}$ and $\mathbf{1} \in \mathcal{V}$, so $[v]_{\mathcal{V}} = [\mathbf{1}]_{\mathcal{V}}$.

If $\bar{u}, \bar{u}' \in \bar{\mathcal{U}}$ then $\bar{u} = [u]_{\mathcal{V}}$ and $\bar{u}' = [u']_{\mathcal{V}}$ for some $u, u' \in \mathcal{U}$, and $uu' \in \mathcal{U}$ since \mathcal{U} is a monoid, so $\bar{u}\bar{u}' = [uu']_{\mathcal{V}} \in \bar{\mathcal{U}}$. We also have

$$[\mathbf{1}]_{\mathcal{V}}[u]_{\mathcal{V}} = [\mathbf{1}u]_{\mathcal{V}} = [u]_{\mathcal{V}} = [u\mathbf{1}]_{\mathcal{V}} = [u]_{\mathcal{V}}[\mathbf{1}]_{\mathcal{V}}$$

and

$$([u][u'])[u''] = [(uu')u''] = [u(u'u'')] = [u]([u'][u'']),$$

so $\bar{\mathcal{U}}$ is a monoid. Thus, $\bar{\mathcal{U}}$ is a submonoid of X/\mathcal{V} .

For any $[x]_{\mathcal{V}} \in X/\mathcal{V}$ there is some $u \in \mathcal{U}$ and $\rho \in R$ such that $[x]_{\mathcal{V}} = [\rho \cdot u]_{\mathcal{V}} = \rho \cdot [u]_{\mathcal{V}}$; thus also $[x]_{\mathcal{V}} \sim [u]_{\mathcal{V}}$ since $\mathbf{1} \cdot [x]_{\mathcal{V}} = [\mathbf{1} \cdot x]_{\mathcal{V}} = [x]_{\mathcal{V}}$. If $[x]_{\mathcal{V}} = \rho \cdot [u]_{\mathcal{V}} = \sigma \cdot [u]_{\mathcal{V}}$ for some $\rho, \sigma \in R$ then $[\rho \cdot u]_{\mathcal{V}} = [\sigma \cdot u]_{\mathcal{V}}$, so $\rho \cdot u \sim_{\mathcal{V}} \sigma \cdot u$, so $v(\rho \cdot u) = v'(\sigma \cdot u)$ where $v, v' \in \mathcal{V}$, so $\rho \cdot vu = \sigma \cdot v'u$ where $vu, v'u \in \mathcal{U}$, so $vu \sim v'u$, so $vu = v'u$ since \mathcal{U} is sparse, so $\rho = \sigma$. Thus, $\bar{\mathcal{U}}$ is a dense set of unit elements of X/\mathcal{V} .

Also, if $u, u' \in \mathcal{U}$ and $[u]_{\mathcal{V}} \sim [u']_{\mathcal{V}}$, then $\rho \cdot [u]_{\mathcal{V}} = \sigma \cdot [u']_{\mathcal{V}}$, so $[\rho \cdot u]_{\mathcal{V}} = [\sigma \cdot u']_{\mathcal{V}}$, so $\rho \cdot u \sim_{\mathcal{V}} \sigma \cdot u'$, so $v(\rho \cdot u) = v'(\sigma \cdot u')$ for some $v, v' \in \mathcal{V}$, so $\rho \cdot vu = \sigma \cdot v'u'$, so $vu \sim v'u'$ where $vu, v'u' \in \mathcal{U}$, so $vu = v'u'$, so $[vu]_{\mathcal{V}} = [v'u']_{\mathcal{V}}$, so $[v]_{\mathcal{V}}[u]_{\mathcal{V}} = [v']_{\mathcal{V}}[u']_{\mathcal{V}}$, so $[u]_{\mathcal{V}} = [u']_{\mathcal{V}}$. Thus, $\bar{\mathcal{U}}$ is a sparse set of unit elements of X/\mathcal{V} . \square

Note that \mathcal{V} is a *partial* coherent system of unit elements of X , that is, a submonoid of X that is a partial system of unit elements of X . For any $v, v' \in \mathcal{V}$ and any $\lambda \in R$, $\lambda \cdot v$ and $\lambda \cdot v'$ in X correspond to the same element $[\lambda \cdot \mathbf{1}]_{\mathcal{V}}$ of X/\mathcal{V} ; more generally, for any $v, v' \in \mathcal{V}$, any $u, u' \in \mathcal{U}$ and any $\lambda \in R$, $\lambda \cdot uvu'$ and $\lambda \cdot uv'u'$ in X correspond to the element $[\lambda \cdot uu']_{\mathcal{V}}$ of X/\mathcal{V} .

The typical application of Proposition 10.4 in physics is described by Raposo [30]:

The mechanism of taking quotients is the algebraic tool underlying what is common practice in physics of choosing “systems of units” such that some specified universal constants become dimensionless and take on the numerical value 1. [...] But it has to be remarked that the mechanism goes beyond a change of system of units; it is indeed a change of space of quantities.

11. RING-MONOID AS SCALABLE MONOIDS

Definition 11.1. Let R be a ring and M a monoid. A *ring-monoid* $R \boxtimes M$ is a set $R \times M$ equipped with a binary operation

$$* : (R \times M) \times (R \times M) \rightarrow R \times M, \quad (\langle \alpha, x \rangle, \langle \beta, y \rangle) \mapsto \langle \alpha, x \rangle \langle \beta, y \rangle := \langle \alpha\beta, xy \rangle$$

and an external binary operation

$$\cdot : R \times (R \times M) \rightarrow R \times M, \quad (\lambda, \langle \alpha, x \rangle) \mapsto \lambda \cdot \langle \alpha, x \rangle := \langle \lambda\alpha, x \rangle.$$

Proposition 11.1. Let $R \boxtimes M$ be a ring-monoid. If R is a commutative ring, then $R \boxtimes M$ is a scalable monoid over R .

Proof. We have

$$\begin{aligned} \langle \langle \alpha, x \rangle \langle \beta, y \rangle \rangle \langle \gamma, z \rangle &= \langle (\alpha\beta)\gamma, (xy)z \rangle = \langle \alpha(\beta\gamma), x(yz) \rangle = \langle \alpha, x \rangle \langle \langle \beta, y \rangle \langle \gamma, z \rangle \rangle, \\ \langle \mathbf{1}, \mathbf{1} \rangle \langle \alpha, x \rangle &= \langle \alpha, x \rangle = \langle \alpha, x \rangle \langle \mathbf{1}, \mathbf{1} \rangle \end{aligned}$$

for any $\alpha, \beta, \gamma \in R$ and $x, y, z \in M$, so $R \boxtimes M$ is a monoid with $\langle 1, \mathbf{1} \rangle$ as identity element. Furthermore,

$$\begin{aligned} 1 \cdot \langle \alpha, x \rangle &= \langle 1\alpha, x \rangle = \langle \alpha, x \rangle, \\ \lambda \cdot (\kappa \cdot \langle \alpha, x \rangle) &= \lambda \cdot \langle \kappa\alpha, x \rangle = \langle \lambda(\kappa\alpha), x \rangle = \langle (\lambda\kappa)\alpha, x \rangle = \lambda\kappa \cdot \langle \alpha, x \rangle, \\ \lambda \cdot \langle \alpha, x \rangle \langle \beta, y \rangle &= \lambda \cdot \langle \alpha\beta, xy \rangle = \langle \lambda(\alpha\beta), xy \rangle = \\ &= \langle (\lambda\alpha)\beta, xy \rangle = \langle \lambda\alpha, x \rangle \langle \beta, y \rangle = (\lambda \cdot \langle \alpha, x \rangle) \langle \beta, y \rangle, \\ \langle (\lambda\alpha)\beta, xy \rangle &= \langle (\alpha\lambda)\beta, xy \rangle = \langle \alpha(\lambda\beta), xy \rangle = \langle \alpha, x \rangle \langle \lambda\beta, y \rangle = \langle \alpha, x \rangle (\lambda \cdot \langle \beta, y \rangle) \end{aligned}$$

for any $\alpha, \beta, \lambda, \kappa \in R$ and $x, y \in M$, so $R \boxtimes M$ is a scalable monoid with \cdot a scaling action of R on $R \times M$. \square

Proposition 11.2. *Let a ring-monoid $R \boxtimes M$ be a scalable monoid over R . Then $\mathcal{U} = \{\langle 1, x \rangle \mid x \in M\}$ is a coherent system of unit elements of $R \boxtimes M$.*

Proof. We have $\langle \alpha, x \rangle = \alpha \cdot \langle 1, x \rangle$ for any $\langle \alpha, x \rangle \in R \times M$, and if $\alpha \cdot \langle 1, x \rangle = \alpha' \cdot \langle 1, x \rangle$ then $\langle \alpha, x \rangle = \langle \alpha', x \rangle$, so $\alpha = \alpha'$. Also, $1 \cdot \langle \alpha, x \rangle = \langle \alpha, x \rangle = \alpha \cdot \langle 1, x \rangle$, so $\langle \alpha, x \rangle \sim \langle 1, x \rangle$. Furthermore, if $\langle 1, x \rangle \sim \langle 1, y \rangle$ so that $\alpha \cdot \langle 1, x \rangle = \beta \cdot \langle 1, y \rangle$ for some $\alpha, \beta \in R$ then $\langle \alpha, x \rangle = \langle \beta, y \rangle$, so $x = y$, so $\langle 1, x \rangle = \langle 1, y \rangle$. Hence, \mathcal{U} is a system of unit elements of $R \boxtimes M$.

Finally, if $\langle 1, x \rangle, \langle 1, y \rangle \in \mathcal{U}$ then $x, y, xy \in M$, so $\langle 1, x \rangle \langle 1, y \rangle = \langle 1, xy \rangle \in \mathcal{U}$, and $\langle 1, \mathbf{1} \rangle \in \mathcal{U}$ since $\mathbf{1} \in M$. Hence, \mathcal{U} is a submonoid of $R \boxtimes M$. \square

Thus, the fact that a ring-monoid $R \boxtimes M$ is a free algebraic structure in the sense that $\langle \alpha, x \rangle = \langle \beta, y \rangle$ if and only if $\alpha = \beta$ and $x = y$ implies that if x and y are distinct elements of M then $\langle 1, x \rangle$ and $\langle 1, y \rangle$ are incommensurable, and that each $\langle 1, x \rangle$, where $x \in M$, is a unit element for $R \cdot \langle 1, x \rangle = \{\langle r, x \rangle \mid r \in R\}$. In addition, \mathcal{U} is a coherent set of unit elements since M is a monoid.

Together, Propositions 11.1, 11.2, 10.1 and 9.1 imply the following fact:

Proposition 11.3. *A ring-monoid $R \boxtimes M$ is an additive scalable monoid if and only if R is commutative.*

If R is in addition an ordered ring then $R \boxtimes M$ is an ordered additive scalable monoid by Proposition 10.3.

Definition 11.1 is thus a construction-definition of a scalable monoid in the case when R is commutative. In Part 3, it will become obvious that a ring monoid $R \boxtimes M$, where R is a field and M is a free abelian group, is a quantity space. This is similar to the construction-definition of a quantity space given by Carlson [6].⁹

⁹Recall that Carlson considers ring-monoids of the form $\mathbb{R} \boxtimes G$ where G is an abelian group equipped with an external operation

$$\mathbb{Q} \times G \rightarrow G, \quad (c, x) \mapsto x^c$$

such that G is a multiplicatively written vector space over \mathbb{Q} , specifically assumed to be finite-dimensional. This is an unnecessary assumption, however; it suffices to assume that G is a free module over \mathbb{Z} , or equivalently a free abelian group. In Raposo's definition of a quantity space [30], Carlson's vector space of "pre-units" is replaced by a finitely generated free abelian group of dimensions.

Part 3. Quantity spaces

A quantity space is basically a scalable monoid equipped with property (A2) in Section 4. Section 12 below introduces the formal definition of quantity spaces; some basic facts about quantity spaces are presented in Section 13.

Systems of unit quantities for quantity spaces are discussed in Section 14. The notion of a measure of a quantity is formally defined in Section 15, and ways in which measures serve as proxies for quantities are described.

In Section 16, we show that the monoid of dimensions Q/\sim corresponding to a quantity space Q is a free abelian group and that bases in Q and Q/\sim have the same cardinality.

A construction-definition of quantity spaces in terms of Laurent monomials in n indeterminates is presented in Section 17.

12. FROM SCALABLE MONOIDS TO QUANTITY SPACES

In this section, we specialize scalable monoids in order to obtain a mathematical model suitable for calculation with quantities, a quantity space. The results in Sections 9 and 10 strongly suggest that a scalable monoid serving this purpose should be equipped with a sufficiently well-behaved set of unit elements. The simplest approach is to require that a quantity space is equipped with a coherent system of unit elements: this is a dense, sparse, closed and consistent set of unit elements.

A coherent system of unit elements of a scalable monoid corresponds to what is called a coherent system of units in metrology. There, coherent systems of units are commonly derived from sets of so-called base units, such as the three base units in the CGS system. The notion corresponding to a set of base units here is a basis in a quantity space, analogous to that of a basis in a vector space or a free abelian group.

If, for example, $B = \{u_1, u_2, u_3\}$ is a set of base units, then the set of all quantities of the form $u_1^{k_1} u_2^{k_2} u_3^{k_3}$, where k_i are integers and $u_i^0 = \mathbf{1}$, is a coherent system of units derived from B , provided that the product of $u_1^{i_1} u_2^{i_2} u_3^{i_3}$ and $u_1^{j_1} u_2^{j_2} u_3^{j_3}$ is equal to a term of the form $u_1^{k_1} u_2^{k_2} u_3^{k_3}$. It is natural to require that

$$(u_1^{i_1} u_2^{i_2} u_3^{i_3}) (u_1^{j_1} u_2^{j_2} u_3^{j_3}) = u_1^{(i_1+j_1)} u_2^{(i_2+j_2)} u_3^{(i_3+j_3)},$$

but this identity implies that the base units commute. (For example, $u_1 u_2 = u_1^1 u_2^1 u_3^0 = (u_1^0 u_2^1 u_3^0) (u_1^1 u_2^0 u_3^0) = u_2 u_1$.) Thus, if we want to include in our model the feature that a coherent system of units can be derived from a system of base units, there are good reasons to require the specialized scalable monoids to be commutative.

Recall that if u is a unit element for some orbit class and $\lambda \cdot u = \lambda' \cdot u$ then $\lambda = \lambda'$, so $\alpha\beta \cdot u = \beta\alpha \cdot u$ implies $\alpha\beta = \beta\alpha$. This suggests that only scalable monoids over commutative rings are of interest. Furthermore, if we want to deal with derived units such as $u_1^1 u_2^{-1} u_3^0$ (meter per second, etc.) then inverses of units must be admitted, and there is a close connection between inverses of quantities in a scalable monoid over R and multiplicative inverses in R . This suggests, finally, that quantity spaces should be defined as certain scalable monoids over fields.

We now come to the basic definitions motivated by the considerations above.

Definition 12.1. Let Q be a commutative scalable monoid over R . A *finite set of generators* for Q is a set $B = \{b_1, \dots, b_n\}$ of elements of Q such that every $x \in Q$ has an expansion

$$x = \mu \cdot \prod_{i=1}^n b_i^{k_i},$$

where $\mu \in R$ and k_1, \dots, k_n are integers. A *finite basis* for Q is a finite set of generators for Q such that every $x \in Q$ has a *unique* expansion of this form.

Note that the uniqueness of the expansion means that the array (μ, k_1, \dots, k_n) is unique given an indexing $B \rightarrow \{1, \dots, n\}$ of the basis elements.

Definition 12.2. A *finitely generated* commutative scalable monoid is one equipped with a finite set of generators. A finitely generated *free* commutative scalable monoid is one equipped with a finite basis. A finitely generated *quantity space* is a finitely generated free commutative scalable monoid over a *field* K .

Elements of a quantity space are called *quantities*, unit elements are called *unit quantities*, and orbit classes in a quantity space are called *dimensions*.

It is not very complicated to generalize the notion of a finite basis for a commutative scalable monoid to include possibly infinite bases, and thus to generalize finitely generated quantity spaces accordingly, but in view of the connection to metrology only the special case of finitely generated quantity spaces will be considered below.

13. SOME BASIC FACTS ABOUT QUANTITY SPACES

Proposition 13.1. *Let Q be a quantity space with basis $\{b_1, \dots, b_n\}$ and $x, y \in Q$. We have*

- (1) $\mathbf{1} = 1 \cdot \prod_{i=1}^n b_i^0$;
- (2) if $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ and $y = \nu \cdot \prod_{i=1}^n b_i^{\ell_i}$ then

$$xy = \mu\nu \cdot \prod_{i=1}^n b_i^{(k_i + \ell_i)};$$

- (3) if $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ and $\mu \neq 0$ then $x^{-1} = \frac{1}{\mu} \cdot \prod_{i=1}^n b_i^{-k_i}$.

Proof. To prove (1), note that $b_i^0 = \mathbf{1}$ for all b_i . (2) follows from Lemma 6.1 and the fact that Q is commutative. (3) follows from (1) and (2). \square

Proposition 13.2. *Every element of a basis for a (finitely generated) quantity space is non-zero.*

Proof. We have $0_C = 0 \cdot x$ and $x = 1 \cdot x$ for all $x \in C$, so if $0_C \in B$ then $0_C = 0 \cdot 0_C$ and $0_C = 1 \cdot 0_C$ are expansions of 0_C in terms of B , so 0_C does not have a unique expansion in terms of B , so B is not a basis for Q . \square

Proposition 13.3. *If Q is a quantity space with basis $\{b_1, \dots, b_n\}$ then $x \in Q$ is a non-zero quantity if and only if $\mu \neq 0$ in the expansion $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$.*

Proof. We have $0 \cdot x = 0 \cdot (\mu \cdot \prod_{i=1}^n b_i^{k_i}) = 0 \cdot \prod_{i=1}^n b_i^{k_i}$, so if $\mu = 0$ then $0 \cdot x = x$ and if $0 \cdot x = x$ then $\mu = 0$ since the expansions of $0 \cdot x$ and x are unique. \square

In particular, $\mathbf{1}$ is a non-zero quantity.

Proposition 13.4. *The product of non-zero quantities in a (finitely generated) quantity space is a non-zero quantity.*

Proof. Set $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ and $y = \nu \cdot \prod_{i=1}^n b_i^{\ell_i}$. Then $xy = \mu\nu \cdot \prod_{i=1}^n b_i^{(k_i+\ell_i)}$, and $\mu\nu \neq 0$ since there are no zero divisors in K . \square

Proposition 13.5. *An element of a (finitely generated) quantity space is invertible if and only if it is non-zero.*

Proof. Let $\{b_1, \dots, b_n\}$ be a basis for Q so that $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$. If $\mu \neq 0$ then $\frac{1}{\mu} \cdot \prod_{i=1}^n b_i^{-k_i}$ is an inverse of x ; conversely, if $\mu = 0$ then there is no ν such that $\mu\nu = 1$, so x does not have an inverse $\nu \cdot \prod_{i=1}^n b_i^{\ell_i}$. \square

Combining Propositions 13.2 and 13.5, we obtain the following result.

Proposition 13.6. *Every element of a basis for a (finitely generated) quantity space is invertible.*

Lemma 13.1. *Let Q be a quantity space over K with a basis $\{b_1, \dots, b_n\}$, and consider $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ and $y = \nu \cdot \prod_{i=1}^n b_i^{\ell_i}$. The following conditions are equivalent:*

- (1) $x \sim y$, or equivalently $\mu \cdot \prod_{i=1}^n b_i^{k_i} \sim \nu \cdot \prod_{i=1}^n b_i^{\ell_i}$;
- (2) $k^i = \ell^i$ for $i = 1, \dots, n$;
- (3) $\prod_{i=1}^n b_i^{k_i} = \prod_{i=1}^n b_i^{\ell_i}$;
- (4) $\nu \cdot x = \mu \cdot y$, or equivalently $\nu \cdot (\mu \cdot \prod_{i=1}^n b_i^{k_i}) = \mu \cdot (\nu \cdot \prod_{i=1}^n b_i^{\ell_i})$.

Proof. The implications (2) \implies (3), (3) \implies (4) and (4) \implies (1) are trivial. To prove (1) \implies (2), note that if $x \sim y$ so that $\alpha \cdot (\mu \cdot \prod_{i=1}^n b_i^{k_i}) = \beta \cdot (\nu \cdot \prod_{i=1}^n b_i^{\ell_i})$ for some $\alpha, \beta \in K$ then

$$\alpha\mu \cdot \prod_{i=1}^n b_i^{k_i} = z = \beta\nu \cdot \prod_{i=1}^n b_i^{\ell_i}.$$

As the expansion of z is unique, $k^i = \ell^i$ for $i = 1, \dots, n$. \square

Note that Lemma 13.1 implies Fourier's principle of dimensional homogeneity [14]: if not $k^i = \ell^i$ for $i = 1, \dots, n$ then $x \not\sim y$, so $x \neq y$.

14. SYSTEMS OF UNIT QUANTITIES IN QUANTITY SPACES

Proposition 14.1. *If Q is a (finitely generated) quantity space then every non-zero quantity $u \in Q$ is a unit quantity for $[u]$.*

Proof. Set $u = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ and $x = \nu \cdot \prod_{i=1}^n b_i^{\ell_i}$. Then $\mu \neq 0$ by Proposition 13.3, and if $u \sim x$ then $\nu \cdot u = \mu \cdot x$ by Lemma 13.1, so $x = \mu^{-1}\nu \cdot u = \mu^{-1} \cdot (\mu \cdot x) = \mu^{-1} \cdot (\nu \cdot u) = \mu^{-1}\nu \cdot u$.

Also, if $\lambda \cdot u = \lambda' \cdot u$ then $\lambda\mu \cdot \prod_{i=1}^n b_i^{k_i} = z = \lambda'\mu \cdot \prod_{i=1}^n b_i^{k_i}$, so $\lambda\mu = \lambda'\mu$ since the expansion of z is unique, so $\lambda = \lambda'$ since $\mu \neq 0$. \square

Corollary 14.1. *If Q is a (finitely generated) quantity space then a dense, sparse set of non-zero elements of Q is a system of unit quantities of Q , and a sparse set of non-zero elements of Q is a partial system of unit quantities of Q .*

In metrology, unit quantities are called *measurement units* [18]. A set of *base units* B is a finite set of measurement units each of which cannot be expressed as a product of powers of the other measurement units in B [18]. A finite basis B for a quantity space Q is a set of base units; if $b \in B$ is not a base unit relative to B then $1 \cdot b = b = \prod_{i=1}^n b_i^{k_i} = 1 \cdot \prod_{i=1}^n b_i^{k_i}$, where $b_i \in B$ and $b_i \neq b$ for all b_i , so b does not have a unique expansion relative to B , so B is not a basis for Q .

Proposition 14.2. *If Q is a quantity space with basis $B = \{b_1, \dots, b_n\}$ then*

$$\mathcal{U} = \left\{ 1 \cdot \prod_{i=1}^n b_i^{k_i} \mid k_i \in \mathbb{Z} \right\}$$

is a coherent system of unit quantities for Q .

Proof. All elements of B are non-zero by Proposition 13.2, so all elements of \mathcal{U} are non-zero and hence unit quantities by Proposition 14.1. Also, \mathcal{U} is dense in Q since every $x \in Q$ has an expansion $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$, so $1 \cdot x = \mu \cdot \left(1 \cdot \prod_{i=1}^n b_i^{k_i} \right)$. Lastly, if $u = 1 \cdot \prod_{i=1}^n b_i^{k_i} \sim 1 \cdot \prod_{i=1}^n b_i^{\ell_i} = v$ then $\prod_{i=1}^n b_i^{k_i} = \prod_{i=1}^n b_i^{\ell_i}$ by Lemma 13.1, so $u = v$, meaning that \mathcal{U} is sparse in Q .

It remains to prove that \mathcal{U} is a monoid. Clearly, $\mathbf{1} \in \mathcal{U}$ since $\mathbf{1} = 1 \cdot \prod_{i=1}^n b_i^0$, and we have

$$\left(1 \cdot \prod_{i=1}^n b_i^{k_i} \right) \left(1 \cdot \prod_{i=1}^n b_i^{\ell_i} \right) = 1 \cdot \prod_{i=1}^n b_i^{(k_i + \ell_i)},$$

so if $u, v \in \mathcal{U}$ then $uv \in \mathcal{U}$. Thus, \mathcal{U} is a submonoid of Q . \square

In other words, every (finite) basis B can be extended to a coherent system \mathcal{U} of unit quantities, consisting of basis quantities and other unit quantities that are expressed as products of basis quantities and their inverses.

In metrology, a *coherent system of units* U is defined essentially as a set of measurement units each of which is either a base unit $b_I \in U$ or a *coherent derived unit*, a non-base unit of the form $1 \cdot \prod_{i=1}^n b_i^{k_i}$, where each b_i is a base unit in U and k_1, \dots, k_n are integers [18]. By Proposition 14.2, a coherent system of units in this sense is a coherent system of unit quantities in the sense of Definition 10.5.

By Propositions 14.2 and 10.1, every (finitely generated) quantity space is an additive quantity space in the sense of Definition 10.2.

Also, Propositions 14.2 and 10.3 imply that if Q is a (finitely generated) quantity space over an ordered field K then Q is an ordered additive quantity space over K with \mathbb{C} ordered by $\leq_{\mathbb{C}}$ defined by $x \leq_{\mathbb{C}} y$ if and only if $y - x = \rho \cdot u$ for some $\rho \in K$ such that $0 \leq \rho$ and some $u \in \mathcal{U} \cap \mathbb{C}$, where \mathcal{U} is a coherent system of unit quantities derived from a basis for Q . Thus, every quantity space over \mathbb{Q} or \mathbb{R} can be regarded as an ordered additive quantity space since \mathbb{Q} and \mathbb{R} are ordered. We normally want quantity spaces to be ordered, and since any Dedekind-complete ordered field is isomorphic to \mathbb{R} [2], it is natural to let K be the real numbers \mathbb{R} .

15. MEASURES OF QUANTITIES

Definition 15.1. Let Q be a quantity space over K with basis $B = \{b_1, \dots, b_n\}$. The uniquely determined scalar $\mu \in K$ in the expansion

$$x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$$

is called the *measure* of x relative to B and will be denoted by $\mu_B(x)$. \square

For example, $\mathbf{1} = 1 \cdot \prod_{i=1}^n b_i^0$ for any B , so we have the following simple but useful fact.

Proposition 15.1. *Let Q be a (finitely generated) quantity space. For any basis B for Q we have $\mu_B(\mathbf{1}) = 1$.*

Relative to a fixed basis, measures of quantities can be used as proxies for the quantities themselves.

Proposition 15.2. *If Q is a (finitely generated) quantity space with basis B and $x, y \in Q$ then $\mu_B(xy) = \mu_B(x)\mu_B(y)$.*

Proof. This follows immediately from Proposition 13.1. \square

Proposition 15.3. *Let Q be a (finitely generated) quantity space with basis B . A quantity $x \in Q$ is invertible if and only if $\mu_B(x) \neq 0$, and for any invertible $x \in Q$ we have $\mu_B(x^{-1}) = \mu_B(x)^{-1}$.*

Proof. The first part of the assertion follows from Propositions 13.3 and 13.5; the second part follows Propositions 13.1, 13.3 and 13.5. \square

Proposition 15.4. *If Q is a (finitely generated) quantity space with basis B then $\mu_B(\lambda \cdot x) = \lambda \mu_B(x)$ for all $\lambda \in K$ and $x \in Q$.*

Proof. If $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ then $\lambda \cdot x = \lambda \cdot (\mu \cdot \prod_{i=1}^n b_i^{k_i}) = \lambda \mu \cdot \prod_{i=1}^n b_i^{k_i}$, so $\mu_B(\lambda \cdot x) = \lambda \mu = \lambda \mu_B(x)$. \square

Proposition 15.5. *If Q is a (finitely generated) quantity space with basis B then $\mu_B(x) + \mu_B(y) = \mu_B(x + y)$ for all $x, y \in X$ such that $x \sim y$.*

Proof. Let $x = \mu_B(x) \cdot \prod_{i=1}^n b_i^{k_i}$ and $y = \mu_B(y) \cdot \prod_{i=1}^n b_i^{k_i}$ be the expansions of x and y relative to $B = \{b_1, \dots, b_n\}$. As $\prod_{i=1}^n b_i^{k_i}$ is non-zero, and thus a unit quantity for $\left[\prod_{i=1}^n b_i^{k_i}\right]$ by Proposition 14.1, we have

$$x + y = \mu_B(x) \cdot \prod_{i=1}^n b_i^{k_i} + \mu_B(y) \cdot \prod_{i=1}^n b_i^{k_i} = (\mu_B(x) + \mu_B(y)) \cdot \prod_{i=1}^n b_i^{k_i},$$

proving the assertion. \square

In general, the measure of a quantity depends on a choice of basis, but there is an important exception to this rule.

Proposition 15.6. *Let Q be a (finitely generated) quantity space. For every $x \in [1]$, the measure $\mu_B(x)$ of x relative to a basis B for Q does not depend on B .*

Proof. $\mathbf{1}$ is a unit quantity for $[1]$ by Proposition 14.1, so there is a unique $\lambda \in K$ such that $x = \lambda \cdot \mathbf{1}$, so $\mu_B(x) = \lambda \mu_B(\mathbf{1})$ by Proposition 15.4, and $\mu_B(\mathbf{1})$ does not depend on B by Proposition 15.1. \square

It is common to refer to a quantity $x \in [\mathbf{1}]$ as a “dimensionless quantity”, although x is not really “dimensionless” – it belongs to, or “has”, the dimension $[\mathbf{1}]$. The so-called Buckingham Π theorem and hence dimensional analysis depends on the fact stated in Proposition 15.6.

16. GROUPS OF DIMENSIONS; CARDINALITY OF BASES

Recall that a trivially scalable monoid Q/\sim may also be regarded as a plain monoid. The definition of a basis for a commutative monoid differs slightly from that for a scalable commutative monoid.

Definition 16.1. Let M be a commutative monoid. A *finite basis* for M is a set $B = \{b_1, \dots, b_n\}$ of elements of M such that every $x \in M$ has a unique expansion

$$x = \prod_{i=1}^n b_i^{k_i},$$

where k_1, \dots, k_n are integers.

In this section, every quotient of the form Q/\sim , where Q is a quantity space, will be regarded as a monoid, which means that Definition 16.1 will be used instead of Definition 12.1 in these cases.

Proposition 16.1. *If Q is a (finitely generated) quantity space then Q/\sim is an abelian group.*

Proof. Q/\sim is commutative since $[x][y] = [xy] = [yx] = [y][x]$ for all $[x], [y] \in Q/\sim$. To prove that Q/\sim is a group it suffices to show that for every $[x] \in X/\sim$ there is a dimension $[x]^{-1} \in X/\sim$ such that $[x][x]^{-1} = [x]^{-1}[x] = [\mathbf{1}]$. Let $B = \{b_1, \dots, b_n\}$ be a basis for Q and let $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ be the unique expansion of x relative to B . Also set $y = 1 \cdot \prod_{i=1}^n b_i^{-k_i}$. Then $[x][y] = [xy] = [\mu \cdot \mathbf{1}] = [yx] = [y][x]$, so $[y]$ is an inverse $[x]^{-1}$ of $[x]$ since $[\mu \cdot \mathbf{1}] = [\mathbf{1}]$. \square

By a finite basis for an abelian group we mean a finite basis for the underlying commutative monoid, and a finitely generated free abelian group is an abelian group for which such a finite basis exists.

Proposition 16.2. *If Q is a quantity space over K with basis $B = \{b_1, \dots, b_n\}$, then $\mathbf{B} = \{[b_1], \dots, [b_n]\}$ is a basis for Q/\sim with the same cardinality as B .*

Proof. The unique expansions of $b_i, b_{i'} \in B$ relative to B are $b_i = 1 \cdot b_i$ and $b_{i'} = 1 \cdot b_{i'}$. Hence, $[b_i] = [b_{i'}]$ implies $b_i = b_{i'}$ since $b_i \sim b_{i'}$ implies $1 \cdot b_i = 1 \cdot b_{i'}$ by Lemma 13.1, so the surjective mapping $\phi : B \rightarrow \mathbf{B}$ given by $\phi(b_i) = [b_i]$ is injective as well and hence a bijection.

Now, let $[x]$ be an arbitrary dimension in Q/\sim . As B is a basis for Q , we have $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$ for some $\mu \in K$ and some integers k_1, \dots, k_n , so $[x] = \left[\mu \cdot \prod_{i=1}^n b_i^{k_i} \right] = \left[\prod_{i=1}^n b_i^{k_i} \right] = \prod_{i=1}^n [b_i]^{k_i}$. Also, if $[x] = \prod_{i=1}^n [b_i]^{k_i} = \prod_{i=1}^n [b_i]^{\ell_i}$, then $\left[\prod_{i=1}^n b_i^{k_i} \right] = \left[\prod_{i=1}^n b_i^{\ell_i} \right]$, so $1 \cdot \prod_{i=1}^n b_i^{k_i} \sim 1 \cdot \prod_{i=1}^n b_i^{\ell_i}$ so $k_i = \ell_i$ for $i = 1, \dots, n$ by Lemma 13.1. Hence, \mathbf{B} is a basis for Q/\sim . \square

We say that a basis $\{b_1, \dots, b_n\}$ for Q and a basis $\{\mathbf{b}_1, \dots, \mathbf{b}_m\}$ for Q/\sim are *similar* when $m = n$ and $[b_i] = \mathbf{b}_i$ for $i = 1, \dots, n$.

Corollary 16.1. *Let Q be a (finitely generated) quantity space. For every basis for Q there exists a unique similar basis for Q/\sim .*

Hence, corresponding to the fact that if X is a scalable monoid then X/\sim is a monoid, we have the following much stronger result.

Proposition 16.3. *If Q is a (finitely generated) quantity space then Q/\sim is a (finitely generated) free abelian group.*

The idea that the set of dimensions of a quantity space forms a free abelian group is present in articles by Krystek [21] and Raposo [30]. This is actually an assumption built into the definition of quantity spaces in [30]; here it is a fact derived from the definitions of quantity spaces and commensurability relations on quantity spaces.

A finitely generated abelian group may have no finite basis; in this case, a corresponding finitely generated trivially scalable commutative monoid over a field cannot have a finite basis since this would contradict Proposition 16.2. Thus, a finitely generated commutative scalable monoid over a field need not be a finitely generated quantity space. (This may be generalized to the case of infinite bases.)

Proposition 16.4. *If Q is a (finitely generated) quantity space and $B = \{b_1, \dots, b_n\}$ is a basis for Q/\sim such that for each $\mathbf{b}_i \in B$ there is a non-zero quantity $b_i \in Q$ such that $\mathbf{b}_i = [b_i]$ then $B = \{b_1, \dots, b_n\}$ is a basis for Q with the same cardinality as B .*

Proof. Consider the function $\psi : B \rightarrow \psi(B)$ given by $\psi(\mathbf{b}_i) = b_i$. ψ is surjective, and we have $\psi(B) = \{b_1, \dots, b_n\}$. Also, if $[\mathbf{b}_i] \neq [\mathbf{b}_{i'}]$ then $b_i \neq b_{i'}$ since dimensions are disjoint, meaning that ψ is injective as well and hence a bijection.

Let x be an arbitrary quantity in Q . As B is a basis for Q/\sim , we have $[x] = \prod_{i=1}^n [b_i]^{k_i} = \left[\prod_{i=1}^n b_i^{k_i} \right]$ for some integers k_1, \dots, k_n , and if $b_i \neq 0$ for each b_i then $\prod_{i=1}^n b_i^{k_i}$ is non-zero and thus a unit quantity for $[x]$ by Proposition 14.1, so there exists a unique $\mu \in K$ such that $x = \mu \cdot \prod_{i=1}^n b_i^{k_i}$. Also, if $x = \nu \cdot \prod_{i=1}^n b_i^{\ell_i} = \nu \cdot \prod_{i=1}^n b_i^{\ell_i}$ then $\left[\mu \cdot \prod_{i=1}^n b_i^{k_i} \right] = \left[\nu \cdot \prod_{i=1}^n b_i^{\ell_i} \right]$, so $\left[\prod_{i=1}^n b_i^{k_i} \right] = \left[\prod_{i=1}^n b_i^{\ell_i} \right]$, so $\prod_{i=1}^n [b_i]^{k_i} = \prod_{i=1}^n [b_i]^{\ell_i}$, so $k_i = \ell_i$ for $i = 1, \dots, n$, since B is a basis for Q/\sim , so $\nu = \mu$ by the uniqueness of μ . We have thus shown that B is a basis for Q . \square

As $\mathbf{b}_i = [\psi(\mathbf{b}_i)]$ and ψ is a bijection between B and B , Proposition 16.4 implies the following fact.

Corollary 16.2. *Let Q be a (finitely generated) quantity space. For every basis for Q/\sim there exists a similar basis for Q .*

Propositions 16.2 and 16.4, the fact that if $B = \phi(B)$ then we can set $\psi = \phi^{-1}$, so that $\psi(B) = B$, since each $b_i \in B$ is non-zero, and the fact that any two bases for a free abelian group have the same cardinality give the following result.

Proposition 16.5. *If Q is a (finitely generated) quantity space then any two bases for Q/\sim have the same cardinality, any basis for Q has the same cardinality as any basis for Q/\sim , and any two bases for Q have the same cardinality.*

Recall that a free module of rank n is a module with a basis and such that all bases have the same cardinality n . Defining the rank of a commutative scalable monoid analogously, we can say that finitely generated quantity spaces are free of finite rank, as are finite-dimensional vector spaces.

17. LAURENT MONOMIALS IN n INDETERMINATES AS QUANTITIES

Recall that if K is a field then K^n is a vector space with operations defined by

$$\begin{aligned}\lambda \cdot (x_1, \dots, x_n) &= (\lambda x_1, \dots, \lambda x_n), \\ (x_1, \dots, x_n) + (y_1, \dots, y_n) &= (x_1 + y_1, \dots, x_n + y_n),\end{aligned}$$

and every n -dimensional vector space over K is isomorphic to a vector space K^n . A quantity space has a similar representation.

We call a term of the form

$$(17.1) \quad \lambda X_1^{k_1} \dots X_n^{k_n},$$

where X_1, \dots, X_n are uninterpreted symbols, λ belongs to a field K and k_1, \dots, k_n are integers, a *Laurent monomial in n indeterminates*, and we denote the set of all terms of the form (17.1) by $K[[X_1, \dots, X_n]]$.

Proposition 17.1. *Let $s = \kappa X_1^{j_1} \dots X_n^{j_n}$ and $t = \lambda X_1^{k_1} \dots X_n^{k_n}$ be any elements of $K[[X_1, \dots, X_n]]$. For any field K , $K[[X_1, \dots, X_n]]$ is a finitely generated quantity space with operations $(\alpha, t) \mapsto \alpha \cdot t$ and $(s, t) \mapsto st$ defined by*

$$\begin{aligned}\alpha \cdot \lambda X_1^{k_1} \dots X_n^{k_n} &= \alpha \lambda X_1^{k_1} \dots X_n^{k_n}, \\ \kappa X_1^{j_1} \dots X_n^{j_n} \lambda X_1^{k_1} \dots X_n^{k_n} &= \kappa \lambda X_1^{(j_1+k_1)} \dots X_n^{(j_n+k_n)}.\end{aligned}$$

Proof. Multiplication of terms in $K[[X_1, \dots, X_n]]$ is associative and commutative because addition of integers and multiplication of elements of K is associative and commutative, and setting

$$\mathbf{1} = 1X_1^0 \dots X_n^0$$

we clearly have $\mathbf{1}t = t\mathbf{1} = t$. Furthermore,

$$\begin{aligned}1 \cdot t &= 1 \cdot \lambda X_1^{k_1} \dots X_n^{k_n} = 1\lambda X_1^{k_1} \dots X_n^{k_n} = t, \\ \alpha \cdot (\beta \cdot t) &= \alpha \cdot (\beta \cdot \lambda X_1^{k_1} \dots X_n^{k_n}) = \alpha \cdot \beta \lambda X_1^{k_1} \dots X_n^{k_n} = \\ &= \alpha \beta \lambda X_1^{k_1} \dots X_n^{k_n} = \alpha \beta \cdot \lambda X_1^{k_1} \dots X_n^{k_n} = \alpha \beta \cdot t\end{aligned}$$

and

$$\begin{aligned}\alpha \kappa X_1^{j_1} \dots X_n^{j_n} \lambda X_1^{k_1} \dots X_n^{k_n} &= \alpha \kappa \lambda X_1^{(j_1+k_1)} \dots X_n^{(j_n+k_n)} = \\ \kappa \alpha \lambda X_1^{(j_1+k_1)} \dots X_n^{(j_n+k_n)} &= \kappa X_1^{j_1} \dots X_n^{j_n} \alpha \lambda X_1^{k_1} \dots X_n^{k_n}\end{aligned}$$

since multiplication in K is associative and commutative, so $\alpha \cdot st = (\alpha \cdot s)t = s(\alpha \cdot t)$ since $\alpha \kappa X_1^{j_1} \dots X_n^{j_n} = \alpha \cdot s$ and so on.

$K[[X_1, \dots, X_n]]$ is thus a commutative scalable monoid over a field; it remains to exhibit a finite scalable monoid basis. For $i = 1, \dots, n$, set $b_i = 1X_1^{\delta_1} \dots X_n^{\delta_n}$, where $\delta_j = 1$ for $i = j$ and $\delta_j = 0$ for $i \neq j$, and $b_i^{-1} = 1X_1^{\vartheta_1} \dots X_n^{\vartheta_n}$, where $\vartheta_j = -1$ for $i = j$ and $\vartheta_j = 0$ for $i \neq j$, so that $b_i b_i^{-1} = b_i^{-1} b_i = \mathbf{1}$. It is clear that t has a unique representation of the form

$$t = \lambda \cdot \prod_{i=1}^n b_i^{k_i},$$

where $b_i^0 = \mathbf{1}$, $b_i^{k_i} = b_i^{(k_i-1)} b_i$ for $k_i > 0$ and $b_i^{k_i} = b_i^{(k_i+1)} b_i^{-1}$ for $k_i < -1$, so $\{b_1, \dots, b_n\}$ is a basis for $K[[X_1, \dots, X_n]]$. \square

Proposition 17.2. *For every finitely generated quantity space Q over K there exists an isomorphic quantity space $K[[X_1, \dots, X_n]]$.*

Proof. Let $B = \{b_1, \dots, b_n\}$ be a basis for Q and let $x \in Q$ have the unique expansion $x = \lambda \cdot \prod_{i=1}^n b_i^{k_i}$. It is easy to verify that the mapping

$$\Phi : Q \rightarrow K[[X_1, \dots, X_n]], \quad x \mapsto \lambda X_1^{k_1} \dots X_n^{k_n}$$

is an isomorphism of scalable monoids between Q and $K[[X_1, \dots, X_n]]$. \square

We have thus seen a simple construction-definition of a quantity space, in addition to the simple axiomatic definition given earlier. While a module over a field is a vector space, a commutative scalable monoid over a field is unfortunately not always a quantity space, but the theory of quantity spaces is not significantly more complicated or inaccessible than that of vector spaces, and should belong to the basic tool-set of mathematicians and physicists.

REFERENCES

- [1] Birkhoff, G. (1935). On the structure of abstract algebras. *Proceedings of the Cambridge Philosophical Society*, **31**, 433–454.
- [2] Blyth, T.S. (2005). *Lattices and Ordered Algebraic Structures*. Springer.
- [3] de Boer, J. (1994). On the history of quantity calculus and the international system, *Metrologia*, **31**, 405–429.
- [4] de Boer, J. (1979). Group properties of quantities and units. *American Journal of Physics*, **47**, 818–819.
- [5] Bridgman (1922). *Dimensional Analysis*. Yale University Press.
- [6] Carlson, D.E. (1979). A mathematical theory of physical units, dimensions and measures. *Archive for Rational Mechanics and Analysis*, **70**, 289–304.
- [7] de Courtney, N. (2015). The Double Interpretation of the Equations of Physics and the Quest for Common Meanings. In Schlaudt, O & Huber, L. (Eds.) *Standardization in Measurement*. Pickering & Chatto, London.
- [8] Descartes, R. (1637). *La Geometrie, Discours de la Méthode*. Leiden.
- [9] Drobot, S. (1953). On the foundations of dimensional analysis. *Studia Mathematica*, **14**, 84–99.
- [10] Emerson, W.H. (2008). On quantity calculus and units of measurement. *Metrologia*, **45**, 134–138.
- [11] Euclid (of Alexandria) (c 300 BC). *Stoicheia (The Elements)*.
- [12] Euler, L. (1740). *Einleitung zur Rechen-Kunst*. Bd. 2. St Petersburg.
- [13] Fleischmann, R. (1951). Die Struktur des physikalischen Begriffssystemes, *Zeitschrift für Physik*, **129**, 377–400.
- [14] Fourier, J.B.J. (1822). *Théorie analytique de la Chaleur*. Paris.
- [15] Gowers, W.T. Two definitions of 'definition'. <https://www.dpmms.cam.ac.uk/~wtg10/definition.html>, retrieved 25-09-2019.
- [16] Gowers W.T. (2002). *Mathematics. A Very Short Introduction*. Oxford University Press.
- [17] Hasse, H. & Scholz, H. (1928). Die Grundlagenkrise der griechischen Mathematik, *Kant-Studien*, **33**, 4–34.
- [18] Joint Committee for Guides in Metrology (2012). *International vocabulary of metrology – Basic and general concepts and associated terms (VIM)*. 3rd edition. Bureau International des Poids et Mesures.
- [19] Jonsson, D. (2014). *Quantities, Dimensions and Dimensional Analysis*. arXiv:1408.5024.
- [20] Jonsson, D. (2015). *On Scalable Monoids*. arXiv:1503.00564.
- [21] Krystek, M. (2015). The term 'dimension' in the international system of units. *Metrologia*, **52**, 297–300.
- [22] König, H. (1950). Über die Mehrdeutigkeit des Grössenbegriffs. *Bulletin des Schweizerischen Elektrotechnischen Vereins*, **41**, 625–629.
- [23] Landolt, M. (1945). *Grösse, Masszahl und Einheit*. Rascher Verlag, Zürich.

- [24] Lodge, A. (1888). The Multiplication and Division of Concrete Quantities. *Nature* 38, 281–283.
- [25] MacLane, S. & Birkhoff, G. (1999). *Algebra*, 3rd Edition (AMS Chelsea Publishing). American Mathematical Society.
- [26] Malet, A. (2006). Renaissance notions of number and magnitude, *Historia Mathematica*, **33**, 63–81.
- [27] Maxwell, J. (1873). *Treatise on Electricity and Magnetism*. Oxford University Press.
- [28] Newton, I. (1720). *Universal Arithmetick*. London.
- [29] Quade, W. (1961). Über die algebraische Struktur des Größenkalküls der Physik, *Abhandlungen der Braunschweigischen Wissenschaftlichen Gesellschaft*, **13**, 24–65.
- [30] Raposo, A.R. (2018). The algebraic structure of quantity calculus, *Measurement Science Review*, **18**, 147–157.
- [31] Silsbee, F.B. (1962). Systems of Electrical Units. *Journal of Research of the National Bureau of Standards-C*, 66C, 137-183.
- [32] Smith, D.E. (1923). *History of Mathematics*. Vol. 1. Reprinted 1958, Dover.
- [33] Stevin (de Bruges), S. (1585). *L'Arithmétique*. Leiden.
- [34] van der Waerden, B. L. (1930). *Moderne Algebra*. Teil I. Springer.
- [35] Wallot, J. (1926). Dimensionen, Einheiten, Masssysteme, *Handbuch der Physik II*. Springer.
- [36] Whitney, H. (1968). The mathematics of physical quantities: Part II: Quantity structures and dimensional analysis. *The American Mathematical Monthly*, 75, 227– 256.

DAN JONSSON, UNIVERSITY OF GOTHENBURG, SE-405 30 GOTHENBURG, SWEDEN
E-mail address: dan.jonsson@gu.se