

Quantifying the unextendibility of entanglement

Kun Wang,^{1,2,*} Xin Wang,^{3,4,†} and Mark M. Wilde^{5,‡}

¹*Shenzhen Institute for Quantum Science and Engineering,
Southern University of Science and Technology, Shenzhen 518055, China*

²*Center for Quantum Computing, Peng Cheng Laboratory, Shenzhen 518055, China*

³*Institute for Quantum Computing, Baidu Research, Beijing 100193, China*

⁴*Joint Center for Quantum Information and Computer Science,
University of Maryland, College Park, Maryland 20742, USA*

⁵*Hearne Institute for Theoretical Physics,
Department of Physics and Astronomy, Center for Computation and Technology,
Louisiana State University, Baton Rouge, Louisiana 70803, USA*

(Dated: December 22, 2020)

The unextendibility or monogamy of entangled states is a key property of quantum entanglement. Unlike conventional ways of expressing entanglement monogamy via entanglement measure inequalities, we develop a state-dependent resource theory to quantify the unextendibility of bipartite entangled states. First, we introduce a family of entanglement measures called *unextendible entanglement*. Given a bipartite state ρ_{AB} , the key idea behind these measures is to minimize a divergence between ρ_{AB} and any possibly reduced state $\rho_{AB'}$ of an extension $\rho_{ABB'}$ of ρ_{AB} . These measures are intuitively motivated by the fact that the more a bipartite state is entangled, the less that each of its individual systems can be entangled with a third party. Second, we show that the unextendible entanglement is an entanglement monotone under two-extendible operations, which include local operations and one-way classical communication as a special case. Unextendible entanglement has several other desirable properties, including normalization and faithfulness. As applications, we show that the unextendible entanglement provides efficiently computable benchmarks for the rate of perfect secret key distillation or entanglement distillation, as well as for the overhead of probabilistic secret key or entanglement distillation.

* nju.wangkun@gmail.com

† wangxinfelix@gmail.com

‡ mwilde@lsu.edu

CONTENTS

I.	Introduction	2
II.	Quantifying the unextendibility of entanglement	3
	A. State-dependent resource theory of unextendible entanglement	3
	B. Generalized unextendible entanglement	5
	C. α -unextendible entanglement	6
	D. Unextendible entanglement (using relative entropy)	14
III.	Efficiently computable entanglement measures	14
	A. Max-unextendible entanglement	14
	B. Min-unextendible entanglement	15
	C. Unextendible fidelity	17
IV.	Applications for secret key and entanglement distillation	19
	A. Overhead of probabilistic entanglement distillation	19
	B. Perfect entanglement distillation	21
	C. Private states and unextendible entanglement	22
	D. Overhead of probabilistic secret key distillation	25
	E. Perfect secret key distillation	25
	F. Examples	26
V.	Concluding remarks	29
	Acknowledgement	29
	References	30
A.	Proof of Lemma 8	34
B.	Proof of Lemma 9	35
C.	Dual SDP of unextendible fidelity	36

I. INTRODUCTION

Quantum entanglement is one of the most counter-intuitive phenomena of quantum theory and a key ingredient in quantum technologies [HHHH09, Hay17, Wil17]. To explore and understand the power of entanglement fully, a comprehensive characterization and quantification of entanglement is needed.

One of the remarkable features of entanglement is its unshareability or unextendibility [Wer89, DPS02, DPS04], which is the fact that the entanglement in some entangled states cannot be shared among multiple subsystems. Unextendibility underscores a fundamental difference between classical correlations and quantum correlations, given that classical correlations can be shared among many parties. This unique property is related to the “monogamy of quantum entanglement” [Ter04], which has been characterized in various ways [CKW00, Ter04, KW04, Yan06, OV06, AI07, LDH⁺16, GG18]. Unextendibility arises in the contexts of Bell inequalities [TDS03], quantum key distribution [MCL06, MRDL09, KL17], and quantum communication [NH09, KDWW19, BBFS18].

In the aforementioned work [KDWW19], a systematic and consistent resource theory of unextendibility was developed, with the goal of quantifying and characterizing unextendibility. Several quantifiers of unextendibility were defined in [KDWW19], which generalized various measures proposed in earlier work [NH09, Bra08, MCL06, HMW13].

Motivated by the central role of unextendibility in understanding quantum entanglement, this paper introduces and develops a state-dependent resource theory of unextendible entanglement. We begin by introducing a family of entanglement measures called unextendible entanglement, where the key principle is to minimize the generalized divergence between a given state ρ_{AB} and any possibly reduced state $\rho_{AB'}$ of an extension $\rho_{ABB'}$ of ρ_{AB} (Section II). This approach is motivated by the intuition that the more a bipartite state is entangled, the less can its individual systems be entangled with a third party. Second, we show that this family of measures is monotone under two-extendible operations, which include local operations and one-way classical communication (1-LOCC) as a special case (Section IIB). We furthermore prove several other desirable properties of the family of entanglement measures, including normalization and faithfulness. We also consider several key examples of unextendible entanglement based on Rényi relative entropies (Section IIC), some of which are efficiently computable (Section III). Third, we show that the unextendible entanglement can be applied to establish efficiently computable fundamental limits to the rate of perfect secret key and entanglement distillation, as well as to the overhead of probabilistic secret key and entanglement distillation (Section IV). These last results indicate that the unextendible entanglement offers a practically meaningful lookout onto the virtues of entanglement as a resource.

II. QUANTIFYING THE UNEXTENDIBILITY OF ENTANGLEMENT

Before we present our main results, let us first establish notation and review some preliminary concepts. Throughout, quantum systems are denoted by A , B , and C and have finite dimensions. Systems described by the same letter are assumed to be isomorphic: $A_1 \cong A_2$ and $A \cong A'$. Linear operators acting on system A are denoted by $L_A \in \mathcal{L}(A)$, Hermitian operators acting on system A are denoted by $H_A \in \mathcal{H}(A)$, and positive semi-definite operators by $P_A \in \mathcal{P}(A)$. Quantum states of system A are denoted by $\rho_A \in \mathcal{S}(A)$ and pure quantum states by $\Psi_A \in \mathcal{V}(A)$. A maximally entangled state Φ_{AB} of Schmidt rank d is given by $\Phi_{AB} := \frac{1}{\sqrt{d}} \sum_{i,j} |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B$, where $\{|i\rangle_A\}_i$ and $\{|i\rangle_B\}_i$ are orthonormal bases for systems A and B , respectively. Quantum channels are completely positive and trace preserving maps from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ and denoted by $\mathcal{N}_{A \rightarrow B} \in \mathcal{Q}(A \rightarrow B)$.

In Section IIB, we introduce the state-dependent resource theory of unextendible entanglement, as well as the generalized unextendible entanglement. In Section IIC, we consider special cases of the generalized unextendible entanglement, which include those based on the quantum relative entropy [Ume62], the Petz–Rényi relative entropy [Pet86], the sandwiched Rényi relative entropy [MLDS⁺13, WWY14], and the geometric Rényi relative entropy [Mat15] (see also [FF19]). We also establish several desirable properties for these measures, including the fact that they are monotone under two-extendible operations.

A. State-dependent resource theory of unextendible entanglement

Unlike the usual framework of quantum resource theories [CG19] and that established for unextendibility in [KDWW19], the free states in our resource theory are *state-dependent*. Note that state-dependent resource theories were previously considered in a different context [RNBG19]. To be specific, given a bipartite state ρ_{AB} , the free states are those bipartite states that are possibly shareable between Alice and a third party B' , where system B' is isomorphic to B . Mathematically,

the set of free states corresponding to a state ρ_{AB} is defined as follows:

$$\mathcal{F}_{\rho_{AB}} := \{\sigma_{AB'} \mid \rho_{AB} = \text{Tr}_{B'}[\rho_{ABB'}], \sigma_{AB'} = \text{Tr}_B[\rho_{ABB'}], \rho_{ABB'} \in \mathcal{S}(ABB')\}. \quad (1)$$

The free operations in this resource theory consist of two-extendible operations [KDWW19], which contain 1-LOCC operations as a special case.

First, let us define a selective 1-LOCC operation. Such an operation consists of a set of CP maps of the following form:

$$\left\{ \mathcal{L}_{AB \rightarrow A'B'}^y \equiv \sum_{x \in \mathcal{X}} \mathcal{F}_{A \rightarrow A'}^{x,y} \otimes \mathcal{G}_{B \rightarrow B'}^{x,y} \right\}_{y \in \mathcal{Y}}, \quad (2)$$

where each map $\mathcal{F}_{A \rightarrow A'}^{x,y}$ is completely positive, the sum map $\sum_{x \in \mathcal{X}, y \in \mathcal{Y}} \mathcal{F}_{A \rightarrow A'}^{x,y}$ is trace-preserving, and each map $\mathcal{G}_{B \rightarrow B'}^{x,y}$ is completely positive and trace-preserving. A 1-LOCC channel is a special case for which $|\mathcal{Y}| = 1$.

Generalizing this, we define a selective two-extendible operation to consist of a set of CP maps of the following form:

$$\{\mathcal{E}_{AB \rightarrow A'B'}^y\}_{y \in \mathcal{Y}}, \quad (3)$$

such that $\sum_{y \in \mathcal{Y}} \mathcal{E}_{AB \rightarrow A'B'}^y$ is trace-preserving, each map $\mathcal{E}_{AB \rightarrow A'B'}^y$ is completely positive and two-extendible [KDWW19], in the sense that there exists an extension map $\mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y$ satisfying

1. Permutation covariance:

$$\mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y \circ \mathcal{W}_{B_1 B_2}^\pi = \mathcal{W}_{B'_1 B'_2}^\pi \circ \mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y, \quad (4)$$

where $\mathcal{W}_{B'_1 B'_2}^\pi$ is a unitary permutation channel (either identity or swap channel).

2. Non-signaling constraint:

$$\text{Tr}_{B'_2} \circ \mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y = \text{Tr}_{B'_2} \circ \mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y \circ \mathcal{R}_{\pi_{B_2}} = \mathcal{E}_{AB \rightarrow A'B'}^y, \quad (5)$$

where $\mathcal{R}_{\pi_{B_2}}$ is a replacer channel that traces out the input on B_2 and replaces it with the maximally mixed state π_{B_2} .

The conditions above ensure that the two marginal operations

$$\mathcal{E}_{AB_1 \rightarrow A'B'_1}^y := \text{Tr}_{B'_2} \circ \mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y, \quad (6)$$

$$\mathcal{E}_{AB_2 \rightarrow A'B'_2}^y := \text{Tr}_{B'_1} \circ \mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y, \quad (7)$$

are operations that are in fact equal to the original operation:

$$\mathcal{E}_{AB \rightarrow A'B'}^y = \mathcal{E}_{AB_1 \rightarrow A'B'_1}^y = \mathcal{E}_{AB_2 \rightarrow A'B'_2}^y. \quad (8)$$

A two-extendible channel [KDWW19] is a special case of a selective two-extendible operation for which $|\mathcal{Y}| = 1$.

Selective 1-LOCC operations are a special case of selective two-extendible operations, observed by picking the extension map of $\mathcal{L}_{AB \rightarrow A'B'}^y$ in (2) as follows:

$$\mathcal{L}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y := \sum_x \mathcal{F}_{A \rightarrow A'}^{x,y} \otimes \mathcal{G}_{B_1 \rightarrow B'_1}^{x,y} \otimes \mathcal{G}_{B_2 \rightarrow B'_2}^{x,y}. \quad (9)$$

Such a choice satisfies the extension, permutation covariance, and non-signaling properties required of selective two-extendible operations. This inclusion was observed in [KDWW19] for two-extendible channels and 1-LOCC channels.

B. Generalized unextendible entanglement

A functional $\mathbf{D} : \mathcal{S}(A) \times \mathcal{S}(A) \rightarrow \mathbb{R} \cup \{+\infty\}$ is a generalized divergence [PV10, SW12] if for arbitrary Hilbert spaces \mathcal{H}_A and \mathcal{H}_B , arbitrary states $\rho_A, \sigma_A \in \mathcal{S}(A)$, and an arbitrary channel $\mathcal{N}_{A \rightarrow B} \in \mathcal{Q}(A \rightarrow B)$, the following data processing inequality holds

$$\mathbf{D}(\rho_A \|\sigma_A) \geq \mathbf{D}(\mathcal{N}_{A \rightarrow B}(\rho_A) \|\mathcal{N}_{A \rightarrow B}(\sigma_A)). \quad (10)$$

A generalized divergence \mathbf{D} is said to be faithful if the inequality $\mathbf{D}(\rho_A \|\rho_A) \leq 0$ holds for an arbitrary state ρ_A , and strongly faithful if $\mathbf{D}(\rho_A \|\sigma_A) = 0$ if and only if $\rho_A = \sigma_A$. Examples of interest are the quantum relative entropy [Ume62], the Petz-Rényi relative entropies [Pet86], the sandwiched Rényi relative entropies [MLDS⁺13, WWY14], and the geometric Rényi relative entropies [Mat15, FF19].

Definition 1 (Generalized unextendible entanglement) *The generalized unextendible entanglement of a bipartite state ρ_{AB} is defined as*

$$\mathbf{E}^u(\rho_{AB}) := \frac{1}{2} \inf_{\rho_{ABB'}} \left\{ \mathbf{D}(\rho_{AB} \|\rho_{AB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB') \right\}, \quad (11)$$

so that the infimum is with respect to all possible extensions $\rho_{ABB'}$ of ρ_{AB} .

Definition 1 is motivated by the intuition that the more a bipartite state is entangled, the less can its individual systems be entangled with a third party.

Remark 1 (Channel representation) *In [CW04], it was established that a state $\rho_{ABB'}$ is an extension of ρ_{AB} if and only if there exists a quantum channel $\mathcal{R}_{C \rightarrow B'}$ such that $\rho_{ABB'} = \mathcal{R}_{C \rightarrow B'}(\Psi_{ABC})$, where Ψ_{ABC} is a purification of ρ_{AB} . Using this correspondence, the generalized unextendible entanglement can be defined in a dynamical way as*

$$\mathbf{E}^u(\rho_{AB}) := \frac{1}{2} \inf_{\mathcal{R}_{C \rightarrow B'}} \left\{ \mathbf{D}(\rho_{AB} \|\rho_{AB'}) : \rho_{ABB'} = (\mathbb{1}_{AB} \otimes \mathcal{R}_{C \rightarrow B'}) (|\Psi\rangle\langle\Psi|_{ABC}) \right\}, \quad (12)$$

where the infimum ranges over all quantum channels $\mathcal{N}_{C \rightarrow B'}$.

Theorem 1 (Two-extendible monotonicity) *The generalized unextendible entanglement does not increase under the action of a two-extendible channel. That is, the following inequality holds*

$$\mathbf{E}^u(\rho_{AB}) \geq \mathbf{E}^u(\mathcal{E}_{AB \rightarrow A'B'}(\rho_{AB})), \quad (13)$$

for ρ_{AB} a bipartite state and $\mathcal{E}_{AB \rightarrow A'B'}$ a two-extendible channel.

Proof. Let $\rho_{AB_1 B_2}$ be an arbitrary extension of $\rho_{AB} = \rho_{AB_1}$. Since the channel $\mathcal{E}_{AB \rightarrow AB'}$ is two-extendible, there exists an extending channel $\mathcal{E}_{AB_1 B_2 \rightarrow AB'_1 B'_2}$ that satisfies the conditions stated in (4)–(5). In particular, the marginal channels $\mathcal{E}_{AB_1 \rightarrow AB'_1}$ and $\mathcal{E}_{AB_2 \rightarrow AB'_2}$ are each equal to the original channel $\mathcal{E}_{AB \rightarrow AB'}$. Furthermore, the state $\mathcal{E}_{AB_1 B_2 \rightarrow AB'_1 B'_2}(\rho_{AB_1 B_2})$ is an extension of $\mathcal{E}_{AB_1 \rightarrow AB'_1}(\rho_{AB_1})$ and $\mathcal{E}_{AB_2 \rightarrow AB'_2}(\rho_{AB_2})$. Then we conclude that

$$\mathbf{D}(\rho_{AB_1} \|\rho_{AB_2}) \geq \mathbf{D}(\mathcal{E}_{AB \rightarrow AB'}(\rho_{AB_1}) \|\mathcal{E}_{AB \rightarrow AB'}(\rho_{AB_2})) \quad (14)$$

$$= \mathbf{D}(\mathcal{E}_{AB_1 \rightarrow AB'_1}(\rho_{AB_1}) \|\mathcal{E}_{AB_2 \rightarrow AB'_2}(\rho_{AB_2})) \quad (15)$$

$$\geq 2\mathbf{E}^u(\mathcal{E}_{AB \rightarrow A'B'}(\rho_{AB})). \quad (16)$$

The first inequality follows from monotonicity of the generalized channel divergence under the action of the quantum channel $\mathcal{E}_{AB \rightarrow AB'}$. The equality follows from the observations stated above. The final inequality follows from the definition of the generalized unextendible entanglement, by applying the fact that $\mathcal{E}_{AB_1 B_2 \rightarrow AB'_1 B'_2}(\rho_{AB_1 B_2})$ is an extension of $\mathcal{E}_{AB_1 \rightarrow AB'_1}(\rho_{AB_1})$ and $\mathcal{E}_{AB_2 \rightarrow AB'_2}(\rho_{AB_2})$. Since the inequality above holds for an arbitrary extension $\rho_{AB_1 B_2}$ of ρ_{AB} , we can take an infimum over all such extensions and conclude (13). ■

Remark 2 *Due to the fact that two-extendible channels contain 1-LOCC channels, it follows that the generalized unextendible entanglement does not increase under the action of a 1-LOCC channel.*

Proposition 2 (Faithfulness) *If the underlying generalized divergence is strongly faithful, then the generalized unextendible entanglement \mathbf{E}^u is non-negative and $\mathbf{E}^u(\rho_{AB}) = 0$ if ρ_{AB} is two-extendible. If the underlying generalized divergence is also continuous, then the state ρ_{AB} is two-extendible if $\mathbf{E}^u(\rho_{AB}) = 0$.*

Proof. Non-negativity of \mathbf{E}^u follows from the assumption of strong faithfulness. Now suppose that ρ_{AB} is two-extendible. Then there exists an extension $\rho_{AB_1 B_2}$ such that $\rho_{AB} = \rho_{AB_1} = \rho_{AB_2}$. Then it follows that $\mathbf{E}^u(\rho_{AB}) = 0$ by definition and from the assumption that the underlying divergence is strongly faithful.

Now suppose that $\mathbf{E}^u(\rho_{AB}) = 0$. By the assumption of continuity, this means that there exists an extension $\rho_{AB_1 B_2}$ of ρ_{AB} such that $\rho_{AB} = \rho_{AB_1} = \rho_{AB_2}$. Then, by definition, ρ_{AB} is two-extendible. ■

C. α -unextendible entanglement

The α -unextendible entanglement of a bipartite state ρ_{AB} is defined by taking the generalized divergence in (11) to be the Petz–Rényi relative entropy [Pet86], the sandwiched Rényi relative entropy $\tilde{D}_\alpha(\omega\|\tau)$ [MLDS⁺13, WWY14], or the geometric Rényi relative entropy [Mat15]. The Petz–Rényi relative entropy is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as follows:

$$D_\alpha(\omega\|\tau) := \frac{1}{\alpha - 1} \log_2 Q_\alpha(\omega\|\tau), \quad (17)$$

$$Q_\alpha(\omega\|\tau) := \text{Tr}[\omega^\alpha \tau^{1-\alpha}]. \quad (18)$$

The sandwiched Rényi relative entropy is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as follows:

$$\tilde{D}_\alpha(\omega\|\tau) := \frac{1}{\alpha - 1} \log_2 \tilde{Q}_\alpha(\omega\|\tau), \quad (19)$$

$$\tilde{Q}_\alpha(\omega\|\tau) := \left\| \omega^{1/2} \tau^{(1-\alpha)/2\alpha} \right\|_{2\alpha}^{2\alpha}. \quad (20)$$

The geometric Rényi relative entropy is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as follows:

$$\hat{D}_\alpha(\omega\|\tau) := \frac{1}{\alpha - 1} \log_2 \hat{Q}_\alpha(\omega\|\tau), \quad (21)$$

$$\hat{Q}_\alpha(\omega\|\tau) := \text{Tr} \left[\tau \left(\tau^{-1/2} \omega \tau^{-1/2} \right)^\alpha \right]. \quad (22)$$

For all of the above quantities, if $\alpha > 1$ and $\text{supp}(\omega) \not\subseteq \text{supp}(\tau)$, then

$$D_\alpha(\omega\|\tau) = \tilde{D}_\alpha(\omega\|\tau) = \hat{D}_\alpha(\omega\|\tau) = +\infty. \quad (23)$$

The Petz–Rényi relative entropy obeys the data processing inequality for $\alpha \in [0, 2]$ [Pet86], the sandwiched Rényi relative entropy obeys the data processing inequality for $\alpha \in [1/2, \infty]$ [FL13, Bei13] (see also [Wil18]), and the geometric Rényi relative entropy obeys the data processing inequality for $\alpha \in (0, 2]$ [Mat15, HM17]. In the case that $\alpha = 1$, we take the limit $\alpha \rightarrow 1$ to find that [Pet86, MLDS⁺13, WWY14]

$$D(\omega\|\tau) := \lim_{\alpha \rightarrow 1} \tilde{D}_\alpha(\omega\|\tau) = \lim_{\alpha \rightarrow 1} D_\alpha(\omega\|\tau) = \text{Tr}[\omega [\log_2 \omega - \log_2 \tau]], \quad (24)$$

where $D(\omega\|\tau)$ is the quantum relative entropy [Ume62]. The geometric Rényi relative entropy converges to the Belavkin–Staszewski relative entropy [BS82] in the limit as $\alpha \rightarrow 1$:

$$\lim_{\alpha \rightarrow 1} \hat{D}_\alpha(\omega\|\tau) = \hat{D}(\omega\|\tau) := \text{Tr}[\rho \log_2(\rho^{1/2} \sigma^{-1} \rho^{1/2})], \quad (25)$$

which is known to obey the data processing inequality [HM17]. In the case that $\alpha = \infty$, we take the limit $\alpha \rightarrow \infty$ to find that [MLDS⁺13]

$$D_{\max}(\omega\|\tau) := \lim_{\alpha \rightarrow \infty} \tilde{D}_\alpha(\omega\|\tau) = \inf \left\{ \lambda \in \mathbb{R} : \omega \leq 2^\lambda \tau \right\}, \quad (26)$$

where $D_{\max}(\omega\|\tau)$ is the max-relative entropy [Dat09].

Lemma 3 (Faithfulness) *Let $\alpha \in (0, 1) \cup (1, \infty)$, and let ω and τ be quantum states. Then $D_\alpha(\omega\|\tau) \geq 0$ and $D_\alpha(\omega\|\tau) = 0$ if and only if $\omega = \tau$. The same is true for the sandwiched and geometric Rényi relative entropies.*

Proof. By utilizing the data processing inequality and the faithfulness of the classical Rényi relative entropy, it follows that the statements above are true for all of the quantum Rényi relative entropies for the range of α for which data processing holds (see, e.g., [WWY14] for this kind of argument). To get outside of the range for $\alpha \in (0, 1)$, we use the following inequality holding for $\alpha \in (0, 1)$ [IRS17]:

$$\alpha D_\alpha(\omega\|\tau) \leq \tilde{D}_\alpha(\omega\|\tau). \quad (27)$$

To get outside of the range for $\alpha \in (1, \infty)$, we use the following inequality holding for $\alpha \in (1, \infty)$:

$$\tilde{D}_\alpha(\omega\|\tau) \leq D_\alpha(\omega\|\tau). \quad (28)$$

This inequality was proved in [WWY14, DL14] (following from Araki–Lieb–Thirring [Ara90, LT76]). For the same range, we also make use of the following inequality:

$$\tilde{D}_\alpha(\omega\|\tau) \leq \hat{D}_\alpha(\omega\|\tau). \quad (29)$$

which also follows from Araki–Lieb–Thirring for $\alpha \in (1, \infty)$ because

$$\tilde{Q}_\alpha(\omega\|\tau) = \text{Tr}[(\tau^{(1-\alpha)/2\alpha} \omega \tau^{(1-\alpha)/2\alpha})^\alpha] \quad (30)$$

$$= \text{Tr}[(\tau^{1/2\alpha} \tau^{-1/2} \omega \tau^{-1/2} \tau^{1/2\alpha})^\alpha] \quad (31)$$

$$\leq \text{Tr}[(\tau^{1/2\alpha})^\alpha (\tau^{-1/2} \omega \tau^{-1/2})^\alpha (\tau^{1/2\alpha})^\alpha] \quad (32)$$

$$= \text{Tr}[\tau (\tau^{-1/2} \omega \tau^{-1/2})^\alpha] \quad (33)$$

$$= \hat{Q}_\alpha(\omega\|\tau). \quad (34)$$

This concludes the proof. ■

Definition 2 (α -unextendible entanglement) *The α -Petz unextendible entanglement is defined for $\alpha \in [0, \infty)$ as*

$$E_\alpha^u(\rho_{AB}) := \frac{1}{2} \inf_{\rho_{ABB'}} \left\{ D_\alpha(\rho_{AB} \parallel \rho_{AB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB') \right\}, \quad (35)$$

the α -sandwiched unextendible entanglement is defined for $\alpha \in (0, \infty]$ as

$$\tilde{E}_\alpha^u(\rho_{AB}) := \frac{1}{2} \inf_{\rho_{ABB'}} \left\{ \tilde{D}_\alpha(\rho_{AB} \parallel \rho_{AB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB') \right\}, \quad (36)$$

and the α -geometric unextendible entanglement is defined for $\alpha \in (0, \infty)$ as

$$\hat{E}_\alpha^u(\rho_{AB}) ::= \frac{1}{2} \inf_{\rho_{ABB'}} \left\{ \hat{D}_\alpha(\rho_{AB} \parallel \rho_{AB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB') \right\}. \quad (37)$$

As the following propositions state, the α -unextendible entanglement possesses several properties that are required for an entanglement measure [HHHH09], although we should clarify that we have only proven the monotonicity for two-extendible operations, which include 1-LOCC operations. We do not expect for the unextendible entanglement to be monotone under all LOCC operations.

First, it is an immediate consequence of Theorem 1 and data processing that α -unextendible entanglement is monotone under the action of a two-extendible channel for certain values of α :

Corollary 4 (Two-extendible monotonicity) *For $\alpha \in [0, 2]$, the α -Petz unextendible entanglement does not increase under the action of a two-extendible channel. For $\alpha \in [1/2, \infty]$, the α -sandwiched unextendible entanglement does not increase under the action of a two-extendible channel. For $\alpha \in (0, 2]$, the α -geometric unextendible entanglement does not increase under the action of a two-extendible channel.*

For certain values of $\alpha \geq 1$, the above monotonicity statement can be strengthened, in the sense that the α -unextendible entanglement is monotone under *selective* two-extendible operations. This property is stronger than what we previously proved in Theorem 1. As before, this implies that α -unextendible entanglement is monotone under selective 1-LOCC operations, since these are a special case of selective two-extendible operations.

Theorem 5 (Selective two-extendible monotonicity) *Let ρ_{AB} be a bipartite state, and let $\{\mathcal{E}_{AB \rightarrow A'B'}^y\}_y$ be a selective two-extendible operation. Then the following inequality holds for the α -Petz unextendible entanglement for all $\alpha \in [1, 2]$:*

$$E_\alpha^u(\rho_{AB}) \geq \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'}^y), \quad (38)$$

where

$$p(y) := \text{Tr}[\mathcal{E}_{AB \rightarrow A'B'}^y(\rho_{AB})], \quad (39)$$

$$\omega_{A'B'}^y := \frac{1}{p(y)} \mathcal{E}_{AB \rightarrow A'B'}^y(\rho_{AB}). \quad (40)$$

The following inequality holds for the α -sandwiched unextendible entanglement for all $\alpha \in [1, \infty]$:

$$\tilde{E}_\alpha^u(\rho_{AB}) \geq \sum_{y:p(y)>0} p(y) \tilde{E}_\alpha^u(\omega_{A'B'}^y). \quad (41)$$

The following inequality holds for the α -geometric unextendible entanglement for all $\alpha \in [1, 2]$:

$$\hat{E}_\alpha^u(\rho_{AB}) \geq \sum_{y:p(y)>0} p(y) \hat{E}_\alpha^u(\omega_{A'B'}^y). \quad (42)$$

Proof. We first show the proof for (41). Let $\rho_{AB_1B_2}$ be an arbitrary extension of ρ_{AB_1} . Then define

$$p(y) := \text{Tr}[\mathcal{E}_{AB_1B_2 \rightarrow A'B_1B'_2}^y(\rho_{AB_1B_2})], \quad (43)$$

$$\omega_{A'B_1B'_2}^y := \frac{1}{p(y)} \mathcal{E}_{AB_1B_2 \rightarrow A'B_1B'_2}^y(\rho_{AB_1B_2}), \quad (44)$$

where $\mathcal{E}_{AB_1B_2 \rightarrow A'B_1B'_2}^y$ is the operation extending $\mathcal{E}_{AB \rightarrow A'B'}^y$. Due to the fact that $\text{Tr}_{B'_2} \circ \mathcal{E}_{AB_1B_2 \rightarrow A'B_1B'_2}^y = \text{Tr}_{B'_2} \circ \mathcal{E}_{AB_1B_2 \rightarrow A'B_1B'_2}^y \circ \mathcal{R}_{\pi_{B_2}}$, it follows that the values of $p(y)$ in (39) and (43) are equal. Furthermore, due to the fact that the marginal operation $\mathcal{E}_{AB_1 \rightarrow A'B'_1}^y = \text{Tr}_{B'_2} \circ \mathcal{E}_{AB_1B_2 \rightarrow A'B_1B'_2}^y$ is equal to the original operation $\mathcal{E}_{AB \rightarrow A'B'}^y$, we conclude that $\omega_{A'B_1B'_2}^y$ is an extension of $\omega_{A'B'}^y$ for all y .

From the data processing inequality for the sandwiched Rényi relative entropy for $\alpha \in [1/2, 1) \cup (1, \infty)$, we conclude the following inequality:

$$\tilde{D}_\alpha(\rho_{AB_1} \|\rho_{AB_2}) \geq \tilde{D}_\alpha(\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_1}) \|\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_2})), \quad (45)$$

where the quantum channel $\mathcal{E}_{AB \rightarrow A'B'Y}$ is defined as

$$\mathcal{E}_{AB \rightarrow A'B'Y}(\cdot) := \sum_y \mathcal{E}_{AB \rightarrow A'B'}^y(\cdot) \otimes |y\rangle\langle y|_Y. \quad (46)$$

Since the marginal operations $\mathcal{E}_{AB_1 \rightarrow A'B'_1}^y$ and $\mathcal{E}_{AB_2 \rightarrow A'B'_2}^y$ are each equal to the original operation $\mathcal{E}_{AB \rightarrow A'B'}^y$, we find that

$$\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_1}) = \sum_y \mathcal{E}_{AB_1 \rightarrow A'B'_1}^y(\rho_{AB_1}) \otimes |y\rangle\langle y|_Y \quad (47)$$

$$= \sum_y p(y) \omega_{A'B'_1}^y \otimes |y\rangle\langle y|_Y \quad (48)$$

$$=: \omega_{A'B'_1Y}, \quad (49)$$

$$\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_2}) = \sum_y \mathcal{E}_{AB_2 \rightarrow A'B'_2}^y(\rho_{AB_2}) \otimes |y\rangle\langle y|_Y \quad (50)$$

$$= \sum_y p(y) \omega_{A'B'_2}^y \otimes |y\rangle\langle y|_Y \quad (51)$$

$$=: \omega_{A'B'_2Y}, \quad (52)$$

which implies that

$$\begin{aligned} & \tilde{D}_\alpha(\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_1}) \|\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_2})) \\ &= \tilde{D}_\alpha(\omega_{A'B'_1Y} \|\omega_{A'B'_2Y}) \end{aligned} \quad (53)$$

$$= \frac{1}{\alpha - 1} \log_2 \sum_{y:p(y)>0} p(y) \tilde{Q}_\alpha(\omega_{A'B'_1}^y \|\omega_{A'B'_2}^y) \quad (54)$$

$$\geq \sum_{y:p(y)>0} p(y) \left[\frac{1}{\alpha - 1} \log_2 \tilde{Q}_\alpha(\omega_{A'B'_1}^y \|\omega_{A'B'_2}^y) \right] \quad (55)$$

$$= \sum_{y:p(y)>0} p(y) \tilde{D}_\alpha(\omega_{A'B'_1}^y \|\omega_{A'B'_2}^y) \quad (56)$$

$$\geq 2 \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'_1}^y) \quad (57)$$

$$= 2 \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'}^y). \quad (58)$$

In the above, the first inequality follows from concavity of the logarithm and the fact that $\alpha > 1$. The second inequality follows because $\omega_{A'B'_1B'_2}^y$ is such that $\text{Tr}_{B'_2}[\omega_{A'B'_1B'_2}^y] = \omega_{A'B'_1}^y$, so that we can then optimize over all extensions of $\omega_{A'B'_1}^y$. Since the following inequality has been established for all extensions $\rho_{AB_1B_2}$ of ρ_{AB} :

$$\frac{1}{2} \tilde{D}_\alpha(\rho_{AB_1} \|\rho_{AB_2}) \geq \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'}^y), \quad (59)$$

we conclude the inequality in (41).

The proof of (38) follows along similar lines, but instead using data processing for the Petz–Rényi relative entropy and the quantity Q_α in (18). In the same way, the proof of (42) follows, but instead using data processing for the geometric Rényi relative entropy and the quantity \hat{Q}_α in (22).

The inequality in (41) for $\alpha \in \{1, \infty\}$ follows by taking a limit. Alternatively, we can prove them directly. To establish the inequality in (41) for $\alpha = 1$ (for quantum relative entropy), we use the same reasoning but the following property of quantum relative entropy:

$$D(\omega_{XB} \|\tau_{XB}) = \sum_x p(x) D(\omega_B^x \|\tau_B^x) + D(p \| q), \quad (60)$$

where

$$\omega_{XB} := \sum_x p(x) |x\rangle\langle x|_X \otimes \omega_B^x, \quad (61)$$

$$\tau_{XB} := \sum_x q(x) |x\rangle\langle x|_X \otimes \tau_B^x. \quad (62)$$

Similarly, the inequality in (42) for $\alpha = 1$ follows from the same reasoning and the following related property of Belavkin–Staszewski relative entropy:

$$\hat{D}(\omega_{XB} \|\tau_{XB}) = \sum_x p(x) \hat{D}(\omega_B^x \|\tau_B^x) + D(p \| q). \quad (63)$$

For $\alpha = \infty$, suppose that the optimal extendible state is $\rho_{AB_1B_2}$, which satisfies $\rho_{AB_1} = \rho_{AB}$ and $E_{\max}^u(\rho_{AB_1}) = \frac{1}{2} D_{\max}(\rho_{AB_1} \|\rho_{AB_2})$. Also, suppose that $E_{\max}^u(\rho_{AB_1}) = \frac{1}{2} \log_2 t$, which means that $\rho_{AB_1} \leq t \rho_{AB_2}$. For any y such that $p(y) > 0$, due to complete positivity and two-extendibility of the map $\mathcal{E}_{AB \rightarrow A'B'}^y$, we have that

$$t \mathcal{E}_{AB_2 \rightarrow A'B'_2}^y(\rho_{AB_2}) / p(y) \geq \mathcal{E}_{AB_1 \rightarrow A'B'_1}^y(\rho_{AB_1}) / p(y) = \omega_{A'B'}^y. \quad (64)$$

Noting that $\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y(\rho_{AB_1B_2}) / p(y)$ is an extension of the state $\omega_{A'B'}^y$, it follows that

$$E_{\max}^u(\omega_{A'B'}^y) \leq \frac{1}{2} \log_2 t. \quad (65)$$

Therefore,

$$E_{\max}^u(\rho_{AB_1}) = \frac{1}{2} \log_2 t \quad (66)$$

$$= \frac{1}{2} \log_2 \sum_{y:p(y)>0} p(y)t \quad (67)$$

$$\geq \frac{1}{2} \log_2 \sum_{y:p(y)>0} p(y) 2^{2E_{\max}^u(\omega_{A'B'}^y)} \quad (68)$$

$$\geq \sum_{y:p(y)>0} p(y) E_{\max}^u(\omega_{A'B'}^y) \quad (69)$$

In the above, (68) is due to (65), while (69) is due to the concavity of the logarithm. \blacksquare

Proposition 6 (Faithfulness) *For $\alpha \in (0, \infty)$, the α -Petz unextendible entanglement of a bipartite state ρ_{AB} satisfies $E_{\alpha}^u(\rho_{AB}) \geq 0$, and $E_{\alpha}^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible. For $\alpha \in (0, \infty]$, the α -sandwiched unextendible entanglement of a bipartite state ρ_{AB} satisfies $\tilde{E}_{\alpha}^u(\rho_{AB}) \geq 0$, and $\tilde{E}_{\alpha}^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible. For $\alpha \in (0, \infty)$, the α -geometric unextendible entanglement of a bipartite state ρ_{AB} satisfies $\hat{E}_{\alpha}^u(\rho_{AB}) \geq 0$, and $\hat{E}_{\alpha}^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible.*

Proof. This follows because the Petz, sandwiched, and geometric Rényi relative entropies satisfy all of the requirements, as given in Lemma 3, needed to invoke Proposition 2. \blacksquare

Before stating the next proposition, recall that the Petz–Rényi mutual information of a bipartite state ρ_{AB} is defined as [GW15, Eq. (6.3) and Corollary 8]

$$I_{\alpha}(A; B)_{\rho} := \min_{\sigma_B} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \sigma_B) = \frac{\alpha}{\alpha - 1} \log_2 \text{Tr} \left[\left(\text{Tr}_A[\rho_A^{1-\alpha} \rho_{AB}^{\alpha}] \right)^{1/\alpha} \right], \quad (70)$$

and the sandwiched Rényi mutual information of a bipartite state ρ_{AB} is defined as [Bei13, GW15]

$$\tilde{I}_{\alpha}(A; B)_{\rho} := \min_{\sigma_B} \tilde{D}_{\alpha}(\rho_{AB} \| \rho_A \otimes \sigma_B), \quad (71)$$

where the minimization in both cases is with respect to states σ_B . We also define the geometric Rényi mutual information of ρ_{AB} in a similar way:

$$\hat{I}_{\alpha}(A; B)_{\rho} := \min_{\sigma_B} \hat{D}_{\alpha}(\rho_{AB} \| \rho_A \otimes \sigma_B), \quad (72)$$

Also, recall that the Rényi entropy of a state ρ_A is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as

$$H_{\alpha}(A)_{\rho} := \frac{1}{1 - \alpha} \log_2 \text{Tr}[\rho_A^{\alpha}], \quad (73)$$

and it is defined for $\alpha \in \{0, 1, \infty\}$ in the limit, so that

$$H_0(A)_{\rho} = \log_2 \text{rank}(\rho_A), \quad (74)$$

$$H(A)_{\rho} := H_1(A)_{\rho} = -\text{Tr}[\rho_A \log_2 \rho_A], \quad (75)$$

$$H_{\min}(A)_{\rho} := H_{\infty}(A)_{\rho} = -\log_2 \|\rho_A\|_{\infty}. \quad (76)$$

The following lemmas relate the α -mutual information to Rényi entropy when the bipartite state is pure. The first was established in the proof of [SBW15, Proposition 13].

Lemma 7 ([SBW15]) *Let $\alpha \in (0, \infty)$. The α -Petz Rényi mutual information of a pure bipartite state ψ_{AB} reduces to twice the γ -Rényi entropy of its marginal:*

$$I_{\alpha}(A; B)_{\psi} = 2H_{\gamma}(A)_{\psi}, \quad (77)$$

where $\gamma := [2 - \alpha] / \alpha$.

Lemma 8 *Let $\alpha \in (0, \infty]$. The α -sandwiched Rényi mutual information of a pure bipartite state ψ_{AB} reduces to twice the β -Rényi entropy of its marginal:*

$$\tilde{I}_\alpha(A; B)_\psi = 2H_\beta(A)_\psi, \quad (78)$$

where $\beta := [2\alpha - 1]^{-1}$.

Proof. See Appendix A. ■

Lemma 9 *Let $\alpha \in (0, \infty)$. The α -geometric Rényi mutual information of a pure bipartite state ψ_{AB} reduces to twice the zero-Rényi entropy of its marginal:*

$$\hat{I}_\alpha(A; B)_\psi = 2H_0(A)_\psi. \quad (79)$$

Proof. See Appendix B. ■

Then the following proposition uses the above lemmas to conclude that the α -unextendible entanglement reduces to Rényi entropy of entanglement for pure states:

Proposition 10 (Reduction for pure states) *Let ψ_{AB} be a pure bipartite state. Then the α -Petz unextendible entanglement reduces to the γ -Rényi entropy of entanglement:*

$$E_\alpha(\psi_{AB}) = H_\gamma(\psi_A), \quad (80)$$

where $\gamma = [2 - \alpha]/\alpha$. The α -sandwiched unextendible entanglement reduces to the β -Rényi entropy of entanglement:

$$\tilde{E}_\alpha(\psi_{AB}) = H_\beta(\psi_A), \quad (81)$$

where $\beta = [2\alpha - 1]^{-1}$. The α -geometric unextendible entanglement reduces to the zero-Rényi entropy of entanglement:

$$\hat{E}_\alpha(\psi_{AB}) = H_0(\psi_A). \quad (82)$$

Proof. For a pure state ψ_{AB} , any extension of it has the form $\sigma_{AB_1B_2} := \psi_A \otimes \sigma_{B_2}$ for some state σ_{B_2} on system B_2 . As such, it follows that

$$E_\alpha(\psi_{AB}) = \min_{\sigma_{B_2}} \frac{1}{2} D_\alpha(\psi_{AB} \| \psi_A \otimes \sigma_{B_2}) \quad (83)$$

$$= \frac{1}{2} I_\alpha(A; B)_\psi \quad (84)$$

$$= H_\gamma(\psi_A). \quad (85)$$

The first equality follows from applying the definition of $E_\alpha(\psi_{AB})$, and the second equality follows from the definition in (70). The final equality follows from Lemma 7.

The conclusions about $\tilde{E}_\alpha(\psi_{AB})$ and $\hat{E}_\alpha(\psi_{AB})$ follow the same line of reasoning but using Lemma 8 and Lemma 9 instead, respectively. ■

Proposition 11 (Normalization) *Let Φ_{AB}^d be a maximally entangled state of Schmidt rank d . Then $E_\alpha(\Phi_{AB}^d) = \log d$ for $\alpha \in (0, \infty)$, $\tilde{E}_\alpha(\Phi_{AB}^d) = \log d$ for $\alpha \in (0, \infty]$, and $\hat{E}_\alpha(\Phi_{AB}^d) = \log d$ for $\alpha \in (0, \infty)$.*

Proof. This follows directly from Proposition 10 and the fact that the Rényi entropy of the maximally mixed state (reduced state of Φ_{AB}^d) is equal to $\log d$ for all values of β . ■

Proposition 12 (Convexity) *The α -Petz unextendible entanglement E_α^u is convex when $\alpha \in (0, 1]$ and quasi-convex when $\alpha \in (1, 2]$. The α -sandwiched unextendible entanglement \tilde{E}_α^u is convex when $\alpha \in [1/2, 1]$ and quasi-convex when $\alpha > 1$. The α -geometric unextendible entanglement \hat{E}_α^u is convex when $\alpha \in (0, 1]$ and quasi-convex when $\alpha \in (1, 2]$. That is, let $\bar{\rho}_{AB} := \sum_i p_i \rho_{AB}^i$, where $\{p_i\}_i$ is a probability distribution and $\{\rho_{AB}^i\}$ is a set of quantum states on system AB . Then*

$$E_\alpha^u(\bar{\rho}_{AB}) \leq \begin{cases} \max_i E_\alpha^u(\rho_{AB}^i) & \alpha \in (1, 2], \\ \sum_i p_i E_\alpha^u(\rho_{AB}^i) & 0 < \alpha \leq 1, \end{cases} \quad (86)$$

$$\tilde{E}_\alpha^u(\bar{\rho}_{AB}) \leq \begin{cases} \max_i \tilde{E}_\alpha^u(\rho_{AB}^i) & \alpha > 1, \\ \sum_i p_i \tilde{E}_\alpha^u(\rho_{AB}^i) & \frac{1}{2} \leq \alpha \leq 1, \end{cases} \quad (87)$$

and

$$\hat{E}_\alpha^u(\bar{\rho}_{AB}) \leq \begin{cases} \max_i \hat{E}_\alpha^u(\rho_{AB}^i) & \alpha \in (1, 2], \\ \sum_i p_i \hat{E}_\alpha^u(\rho_{AB}^i) & 0 < \alpha \leq 1. \end{cases} \quad (88)$$

Proof. This is a simple corollary of the facts that D_α is jointly quasi-convex for $\alpha \in (1, 2]$ and jointly convex for $\alpha \in (0, 1]$ [Pet86], \tilde{D}_α is jointly quasi-convex for $\alpha > 1$ and jointly convex for $\alpha \in [1/2, 1]$ [FL13], and \hat{D}_α is jointly quasi-convex for $\alpha \in (1, 2]$ and jointly convex for $\alpha \in (0, 1]$ [HM17]. ■

Aside from the above properties, the α -unextendible entanglement is also subadditive:

Proposition 13 (Subadditivity) *Let $\rho_{A_1 B_1} \in \mathcal{S}(A_1 B_1)$ and $\rho_{A_2 B_2} \in \mathcal{S}(A_2 B_2)$ be two bipartite states. Then the following subadditivity inequality holds for $\alpha \in [0, \infty)$*

$$E_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq E_\alpha^u(\rho_{A_1 B_1}) + E_\alpha^u(\rho_{A_2 B_2}), \quad (89)$$

for $\alpha \in (0, \infty]$

$$\tilde{E}_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \tilde{E}_\alpha^u(\rho_{A_1 B_1}) + \tilde{E}_\alpha^u(\rho_{A_2 B_2}), \quad (90)$$

and for $\alpha \in (0, \infty)$

$$\hat{E}_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \hat{E}_\alpha^u(\rho_{A_1 B_1}) + \hat{E}_\alpha^u(\rho_{A_2 B_2}), \quad (91)$$

where the entanglement is evaluated across the $A:B$ cut.

Proof. Let $\sigma_{A_1 B_1 B'_1}$ and $\sigma_{A_2 B_2 B'_2}$ be arbitrary extensions of $\rho_{A_1 B_1}$ and $\rho_{A_2 B_2}$, respectively. Consider the state $\sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}$. Then it follows that

$$\text{Tr}_{B'_1 B'_2}[\sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}] = \text{Tr}_{B'_1}[\sigma_{A_1 B_1 B'_1}] \otimes \text{Tr}_{B'_2}[\sigma_{A_2 B_2 B'_2}] = \rho_{A_1 B_1} \otimes \rho_{A_2 B_2}, \quad (92)$$

so that $\sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}$ is an extension of $\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}$. Thus

$$E_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \frac{1}{2} D_\alpha(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2} \| \sigma_{A_1 B'_1} \otimes \sigma_{A_2 B'_2}) \quad (93)$$

$$= \frac{1}{2} D_\alpha(\rho_{A_1 B_1} \| \sigma_{A_1 B'_1}) + \frac{1}{2} D_\alpha(\rho_{A_2 B_2} \| \sigma_{A_2 B'_2}). \quad (94)$$

Taking an infimum over all extensions $\sigma_{A_1 B_1 B'_1}$ and $\sigma_{A_2 B_2 B'_2}$ leads to the inequality in (89). The inequalities in (90) and (91) are established similarly. ■

D. Unextendible entanglement (using relative entropy)

As mentioned above, the quantum relative entropy is a particular instance of the Petz- and sandwiched Rényi relative entropy when $\alpha \rightarrow 1$. Unextendible entanglement in terms of these measures has already been defined above and several of its properties have been established, but it is notable enough that we define it explicitly here as follows:

Definition 3 (Unextendible entanglement) *For a bipartite state ρ_{AB} , the unextendible entanglement is defined as*

$$E^u(\rho_{AB}) := \frac{1}{2} \inf_{\rho_{ABB'}} \{D(\rho_{AB} \parallel \rho_{ABB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB')\}, \quad (95)$$

where D is the quantum relative entropy from (24).

As proved above, it obeys the following properties: selective two-extendible monotonicity, faithfulness, reduction to entropy of entanglement for pure bipartite states, normalization, convexity, and subadditivity.

Regarding the third property mentioned above, it is a special case of Proposition 10 for $\alpha = 1$, and it states that for pure bipartite states, the unextendible entanglement evaluates to the von Neumann entropy of the reduced state. Although the proof has already been given, we can see it with a straightforward proof consisting of a few steps: Let $|\psi\rangle_{AB}$ be a pure state. Any extension of ψ_{AB} is of the form $\sigma_{ABB'} := |\psi\rangle\langle\psi|_{AB} \otimes \sigma_{B'}$ for some state $\sigma_{B'}$ on system B' . As so, $\text{Tr}_B[\sigma_{ABB'}] = \rho_A \otimes \sigma_{B'}$, where $\rho_A = \text{Tr}_B[|\psi\rangle\langle\psi|_{AB}]$. Then it follows that

$$E^u(\psi_{AB}) = \inf_{\sigma_{B'}} \frac{1}{2} D(\psi_{AB} \parallel \rho_A \otimes \sigma_{B'}) = \frac{1}{2} I(A; B)_\psi = H(A)_\psi, \quad (96)$$

where the second equality follows from the definition of quantum mutual information.

III. EFFICIENTLY COMPUTABLE ENTANGLEMENT MEASURES

A. Max-unextendible entanglement

Another interesting instance of the sandwiched Rényi relative entropy is the max-relative entropy, as recalled in (26). The max-relative entropy was originally defined and studied in [Dat09]. Here we adopt the max-relative entropy to define the max-unextendible entanglement. It turns out that this measure is additive and can be calculated efficiently by means of a semidefinite program.

Definition 4 (Max-unextendible entanglement) *For a given bipartite state ρ_{AB} , the max-unextendible entanglement is defined as*

$$E_{\max}^u(\rho_{AB}) := \frac{1}{2} \inf_{\rho_{ABB'}} \{D_{\max}(\rho_{AB} \parallel \rho_{ABB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB')\}. \quad (97)$$

From the definition of max-unextendible entanglement, it follows that it can be computed efficiently by means of a semidefinite program (SDP). To be more specific, the following two optimization problems satisfy strong duality and both evaluate to $2^{-2E_{\max}^u(\rho_{AB})}$.

<u>primal problem</u>	<u>dual problem</u>
max λ	min $\text{Tr} [\rho_{AB} Y_{AB}]$
s.t. $\lambda \rho_{AB} \leq \sigma_{AB'}$ (98)	s.t. $\text{Tr} [\rho_{AB} X_{AB}] \geq 1$ (99)
$\rho_{AB} = \text{Tr}_{B'} \sigma_{ABB'}$	$X_{AB'} \otimes \mathbb{1}_B \leq Y_{AB} \otimes \mathbb{1}_{B'}$
$\sigma_{ABB'} \geq 0$	$X_{AB} \geq 0$

The primal SDP follows by considering that

$$D_{\max}(\omega || \tau) = \log_2 \inf\{\lambda : \omega \leq \lambda \tau\} = -\log_2 \sup\{\mu : \mu \omega \leq \tau\}. \quad (100)$$

The dual SDP follows by standard methods (Lagrange multiplier method).

By using the primal and dual expressions of $2^{-2E_{\max}^u(\rho_{AB})}$ and strong duality, it follows that $E_{\max}^u(\rho_{AB})$ is additive, which is an appealing feature that finds use in Section IV.

Proposition 14 (Additivity) *Let $\rho_{A_1 B_1} \in \mathcal{S}(A_1 B_1)$ and $\rho_{A_2 B_2} \in \mathcal{S}(A_2 B_2)$ be bipartite states. It holds that*

$$E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) = E_{\max}^u(\rho_{A_1 B_1}) + E_{\max}^u(\rho_{A_2 B_2}), \quad (101)$$

where the entanglement is evaluated across the $A|B$ cut.

Proof. Subadditivity follows from investigating the primal SDP. Suppose that $(\lambda_1, \sigma_{A_1 B_1 B'_1})$ and $(\lambda_2, \sigma_{A_2 B_2 B'_2})$ achieve $E_{\max}^u(\rho_{A_1 B_1})$ and $E_{\max}^u(\rho_{A_2 B_2})$, respectively. As

$$\lambda_1 \lambda_2 \rho_{A_1 B_1} \otimes \rho_{A_2 B_2} = \lambda_1 \rho_{A_1 B_1} \otimes \lambda_2 \rho_{A_2 B_2} \leq \sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}, \quad (102)$$

one can check that $(\lambda_1 \lambda_2, \sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2})$ forms a feasible solution. Thus

$$2^{-2E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})} \geq \lambda_1 \lambda_2 = 2^{-2E_{\max}^u(\rho_{A_1 B_1}) - 2E_{\max}^u(\rho_{A_2 B_2})},$$

which gives $E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq E_{\max}^u(\rho_{A_1 B_1}) + E_{\max}^u(\rho_{A_2 B_2})$.

The superadditivity is shown by investigating the dual SDP. Suppose that $(X_{A_1 B_1}, Y_{A_1 B_1})$ and $(X_{A_2 B_2}, Y_{A_2 B_2})$ achieve $E_{\max}^u(\rho_{A_1 B_1})$ and $E_{\max}^u(\rho_{A_2 B_2})$, respectively. As

$$\text{Tr}[(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})(X_{A_1 B_1} \otimes X_{A_2 B_2})] = \text{Tr}[\rho_{A_1 B_1} X_{A_1 B_1}] \text{Tr}[\rho_{A_2 B_2} X_{A_2 B_2}] \geq 1 \quad (103)$$

and $X_{A_1 B_1'} \otimes X_{A_2 B_2'} \otimes \mathbb{1}_{B_1 B_2} \leq Y_{A_1 B_1} \otimes Y_{A_2 B_2} \otimes \mathbb{1}_{B_1' B_2'}$, the pair $(X_{A_1 B_1} \otimes X_{A_2 B_2}, Y_{A_1 B_1} \otimes Y_{A_2 B_2})$ forms a dual feasible solution. Thus

$$2^{-2E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})} \leq \text{Tr}[\rho_{A_1 B_1} Y_{A_1 B_1}] \text{Tr}[\rho_{A_2 B_2} Y_{A_2 B_2}] = 2^{-2E_{\max}^u(\rho_{A_1 B_1}) - 2E_{\max}^u(\rho_{A_2 B_2})}, \quad (104)$$

which gives $E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \geq E_{\max}^u(\rho_{A_1 B_1}) + E_{\max}^u(\rho_{A_2 B_2})$. ■

B. Min-unextendible entanglement

In this section, we consider the limit of the Petz–Rényi relative entropy as $\alpha \rightarrow 0$, which is known as the min-relative entropy [Dat09, Definition 2]. Let us first recall the definition. Let $\rho \in \mathcal{S}(A)$ and $\sigma \in \mathcal{P}(A)$ such that the support of ρ is contained in the support of σ . Let Π_ρ denote the projection onto the support of ρ . Then the min-relative entropy of ρ to σ is defined as

$$D_{\min}(\rho || \sigma) := -\log \text{Tr}[\Pi_\rho \sigma]. \quad (105)$$

With D_{\min} , we define the min-unextendible entanglement as follows.

Definition 5 (Min-unextendible entanglement) For a given bipartite state ρ_{AB} , the min-unextendible entanglement is defined as

$$E_{\min}^u(\rho_{AB}) := \frac{1}{2} \inf \{ D_{\min}(\rho_{AB} \parallel \rho_{ABB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB') \}. \quad (106)$$

Much like the max-unextendible entanglement, the min-unextendible entanglement can also be calculated as the solution to a semidefinite program and shown to be additive. Let P_{AB} denote the projection onto the support of the state ρ_{AB} . The following two optimization problems satisfy strong duality and both evaluate to $2^{-2E_{\min}^u(\rho_{AB})}$.

$$\begin{array}{ll} \text{primal problem} & \text{dual problem} \\ \max & \text{Tr}[P_{AB}\sigma_{AB'}] \\ \text{s.t.} & \text{Tr}_{B'} \sigma_{ABB'} = \rho_{AB}, \\ & \sigma_{ABB'} \geq 0 \end{array} \quad (107)$$

$$\begin{array}{ll} \min & \text{Tr}[\rho_{AB}X_{AB}] \\ \text{s.t.} & X_{AB} \otimes \mathbb{1}_{B'} \geq P_{AB'} \otimes \mathbb{1}_B, \\ & X_{AB} = X_{AB}^\dagger \end{array} \quad (108)$$

Proposition 15 (Additivity) Let $\rho_{A_1B_1} \in \mathcal{S}(A_1B_1)$ and $\rho_{A_2B_2} \in \mathcal{S}(A_2B_2)$ be bipartite states. It holds that

$$E_{\min}^u(\rho_{A_1B_1} \otimes \rho_{A_2B_2}) = E_{\min}^u(\rho_{A_1B_1}) + E_{\min}^u(\rho_{A_2B_2}), \quad (109)$$

where the entanglement is evaluated across the $A:B$ cut.

Proof. The proof follows similarly to the proof of Proposition 14, by using the primal and dual programs, along with strong duality. \blacksquare

The min-unextendible entanglement of a pure bipartite state can be computed explicitly and possesses an operational interpretation.

Proposition 16 Let $|\psi\rangle_{AB} := \sum_{i=1}^k \sqrt{\alpha_i} |\psi_i^A\rangle |\psi_i^B\rangle$ be a pure state in \mathcal{H}_{AB} , with $\alpha_1 \geq \dots \alpha_k > 0$. Then the following equality holds

$$E_{\min}^u(\psi_{AB}) = -\log \alpha_1. \quad (110)$$

Proof. This follows from (80) by taking the limit $\alpha \rightarrow 0$. Alternatively, we briefly outline another proof as follows. For a pure state ψ_{AB} , any extension of it has the form $\sigma_{ABB'} := |\psi\rangle\langle\psi|_{AB} \otimes \sigma_{B'}$ for some state on system B' . Then $\text{Tr}_B[\sigma_{ABB'}] = \psi_A \otimes \sigma_{B'}$. Suppose that a spectral decomposition of $\sigma_{B'}$ is $\sigma_{B'} = \sum_n p_n |e_n\rangle\langle e_n|$. Then

$$\begin{aligned} & \frac{1}{2} D_{\min}(\psi_{AB} \parallel \psi_A \otimes \sigma_{B'}) \\ &= -\frac{1}{2} \log \sum_{i,j,m,n} \sqrt{\alpha_i \alpha_j} \alpha_m p_n \langle \psi_i^A \psi_i^B | (|\psi_m^A\rangle\langle\psi_m^A| \otimes |e_n\rangle\langle e_n|) | \psi_j^A \psi_j^B \rangle \end{aligned} \quad (111)$$

$$= -\frac{1}{2} \log \sum_{i,n} \alpha_i^2 p_n |\langle \psi_i^B | e_n \rangle|^2 \quad (112)$$

$$\geq -\frac{1}{2} \log \alpha_1^2 \sum_{i,n} p_n |\langle \psi_i^B | e_n \rangle|^2 \quad (113)$$

$$= -\log \alpha_1. \quad (114)$$

The inequality is achievable by choosing $\sigma_{B'} = |\psi_1^B\rangle\langle\psi_1^B|$. Thus $E_{\min}^u(\psi_{AB}) = -\log \alpha_1$. \blacksquare

The min-unextendible entanglement $E_{\min}^u(\psi_{AB})$ has an operational interpretation in terms of deterministic entanglement transformation [DFJY05], which we briefly introduce as follows. Let $|\psi_1\rangle$ and $|\psi_2\rangle$ be two pure bipartite states for systems AB . Let $m \in \mathbb{N}$ be an integer. We define $f(m)$ to be the maximum integer n such that $\psi_1^{\otimes m}$ can be transformed into $\psi_2^{\otimes n}$ by LOCC deterministically. The *deterministic entanglement transformation rate* from ψ_1 to ψ_2 , written $D(\psi_1 \rightarrow \psi_2)$, is defined as

$$D(\psi_1 \rightarrow \psi_2) := \sup_{m \geq 1} \frac{f(m)}{m}. \quad (115)$$

Intuitively, for sufficiently large m , one can transform m copies of ψ_1 exactly into $mD(\psi_1 \rightarrow \psi_2)$ copies of ψ_2 by LOCC. We have the following proposition, which is a consequence of Proposition 16 and the developments in [DFJY05]:

Proposition 17 *Let $|\psi\rangle_{AB}$ be a pure state in AB and Φ_2 be the Bell state. Then the following equality holds*

$$D(\psi \rightarrow \Phi_2) = E_{\min}^u(\psi_{AB}). \quad (116)$$

C. Unextendible fidelity

Let $\rho, \sigma \in \mathcal{S}(A)$ be two quantum states. The (root) fidelity between ρ and σ is defined as [Uhl76]

$$F(\rho, \sigma) := \|\sqrt{\rho}\sqrt{\sigma}\|_1 = \text{Tr} \left[\sqrt{\sqrt{\sigma}\rho\sqrt{\sigma}} \right]. \quad (117)$$

Here we define the unextendible fidelity of a state ρ_{AB} :

Definition 6 (Unextendible fidelity) *For a given bipartite state ρ_{AB} , the unextendible fidelity is defined as*

$$F^u(\rho_{AB}) := \max_{\rho_{ABB'}} \{F(\rho_{AB}, \rho_{AB'}) : \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}, \rho_{ABB'} \in \mathcal{S}(ABB')\}. \quad (118)$$

Suppose that Ψ_{ABC} is a purification of ρ_{AB} . By applying Remark 1, we see the unextendible fidelity can be alternatively understood as a measure of how well one can recover the state ρ_{AB} if system B is lost and a recovery operation is performed on the purification system C alone, due to the following equivalent definition

$$F^u(\rho_{AB}) = \max_{\mathcal{R}_{C \rightarrow B'}} \{F(\rho_{AB}, \rho_{AB'}) : \rho_{ABB'} = (\mathbb{1}_{AB} \otimes \mathcal{R}_{C \rightarrow B'}) (|\Psi\rangle\langle\Psi|_{ABC})\}, \quad (119)$$

where the maximum ranges over all quantum channels $\mathcal{N}_{C \rightarrow B'}$. The unextendible fidelity is thus similar in spirit to fidelity of recovery from [SW15], but one finds that it is a different measure when analyzing it in more detail.

By examining (19) and (117), one immediately finds that $\tilde{D}_{1/2}(\rho\|\sigma) = -\log F^2(\rho, \sigma)$. Thus, we establish the following equivalence between unextendible fidelity and 1/2-sandwiched unextendible entanglement:

$$F^u(\rho_{AB}) := \max_{\substack{\rho_{ABB'} \in \mathcal{S}(ABB') \\ \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}}} F(\rho_{AB}, \rho_{AB'}) \quad (120)$$

$$= \max_{\substack{\rho_{ABB'} \in \mathcal{S}(ABB') \\ \text{Tr}_{B'}[\rho_{ABB'}] = \rho_{AB}}} 2^{-\frac{1}{2} \tilde{D}_{1/2}(\rho_{AB} \| \rho_{AB'})} \quad (121)$$

$$= 2^{-\tilde{E}_{1/2}^u(\rho_{AB})}. \quad (122)$$

Since the fidelity function is SDP computable [Wat13], it follows that the unextendible fidelity can be computed efficiently by means of a semidefinite program. To be more specific, the following two optimization problems satisfy strong duality and both evaluate to $F^u(\rho_{AB})$. For completeness, we show in detail how to derive the dual problem in Appendix C.

primal problem	dual problem
$\begin{aligned} \max \quad & \frac{1}{2} \text{Tr}[X_{AB}] + \frac{1}{2} \text{Tr}[X_{AB}^\dagger] \\ \text{s.t.} \quad & \begin{pmatrix} \rho_{AB} & X_{AB} \\ X_{AB}^\dagger & \text{Tr}_B \sigma_{ABB'} \end{pmatrix} \geq 0 \\ & \rho_{AB} = \text{Tr}_{B'} \sigma_{ABB'} \\ & \sigma_{ABB'} \geq 0 \end{aligned} \quad (123)$	$\begin{aligned} \min \quad & \frac{1}{2} \text{Tr}[(W_{AB} + Z_{AB})\rho_{AB}] \\ \text{s.t.} \quad & \begin{pmatrix} W_{AB} & -\mathbf{1}_{AB} \\ -\mathbf{1}_{AB} & Y_{AB'} \end{pmatrix} \geq 0 \\ & Z_{AB} \otimes \mathbf{1}_{B'} \geq Y_{AB'} \otimes \mathbf{1}_B \end{aligned} \quad (124)$

We also establish the following equivalent dual representation of $F^u(\rho_{AB})$ in Appendix C:

$$\begin{aligned} \inf \quad & \sqrt{\text{Tr}[Y_{AB'}^{-1} \rho_{AB}] \text{Tr}[Z_{AB} \rho_{AB}]} \\ \text{s.t.} \quad & Z_{AB} \otimes \mathbf{1}_{B'} \geq Y_{AB'} \otimes \mathbf{1}_B \\ & Y_{AB'} > 0. \end{aligned} \quad (125)$$

As a direct consequence of this dual representation, we find that the extendible fidelity is multiplicative.

Proposition 18 (Multiplicativity) *Let $\rho_{A_1 B_1} \in \mathcal{S}(A_1 B_1)$ and $\rho_{A_2 B_2} \in \mathcal{S}(A_2 B_2)$ be bipartite states. The following equality holds*

$$F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) = F^u(\rho_{A_1 B_1}) F^u(\rho_{A_2 B_2}), \quad (126)$$

where the entanglement is evaluated across the $A|B$ cut.

Proof. For the “ \geq ” part, suppose that $\rho_{A_1 B_1 B'_1}$ and $\rho_{A_2 B_2 B'_2}$ achieve $F^u(\rho_{A_1 B_1})$ and $F^u(\rho_{A_2 B_2})$, respectively. Since

$$\text{Tr}_{B'_1 B'_2} [\rho_{A_1 B_1 B'_1} \otimes \rho_{A_2 B_2 B'_2}] = \rho_{A_1 B_1} \otimes \rho_{A_2 B_2}, \quad (127)$$

the state $\rho_{A_1 B_1 B'_1} \otimes \rho_{A_2 B_2 B'_2}$ is a feasible solution for $F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})$. So

$$F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \geq F(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}, \rho_{A_1 B'_1} \otimes \rho_{A_2 B'_2}) \quad (128)$$

$$= F(\rho_{A_1 B_1}, \rho_{A_1 B'_1}) F(\rho_{A_2 B_2}, \rho_{A_2 B'_2}) \quad (129)$$

$$= F^u(\rho_{A_1 B_1}) F^u(\rho_{A_2 B_2}), \quad (130)$$

where the first equality follows from the multiplicativity of the fidelity.

For the “ \leq ” part, we employ the dual representation in (125). Let $(Y_{A_1B'_1}, Z_{A_1B_1})$ and $(Y_{A_2B'_2}, Z_{A_2B_2})$ be feasible solutions for $F^u(\rho_{A_1B_1})$ and $F^u(\rho_{A_2B_2})$, respectively. Set $Y_{A_1A_2B'_1B'_2} \equiv Y_{A_1B'_1} \otimes Y_{A_2B'_2}$ and $Z_{A_1A_2B_1B_2} \equiv Z_{A_1B_1} \otimes Z_{A_2B_2}$. Since

$$Z_{A_1A_2B_1B_2} \otimes \mathbb{1}_{B'_1B'_2} = (Z_{A_1B_1} \otimes \mathbb{1}_{B'_1}) \otimes (Z_{A_2B_2} \otimes \mathbb{1}_{B'_2}) \quad (131)$$

$$\geq (Y_{A_1B'_1} \otimes \mathbb{1}_{B_1}) \otimes (Y_{A_2B'_2} \otimes \mathbb{1}_{B_2}) = Y_{A_1A_2B'_1B'_2}, \quad (132)$$

$(Y_{A_1A_2B'_1B'_2}, Z_{A_1A_2B_1B_2})$ is a feasible solution. Then

$$F^u(\rho_{A_1B_1} \otimes \rho_{A_2B_2}) \leq \sqrt{\text{Tr}[Y_{A_1A_2B'_1B'_2}^{-1}(\rho_{A_1B_1} \otimes \rho_{A_2B_2})] \text{Tr}[Z_{A_1A_2B_1B_2}(\rho_{A_1B_1} \otimes \rho_{A_2B_2})]} \quad (133)$$

$$= \sqrt{\text{Tr}[Y_{A_1B'_1}^{-1}\rho_{A_1B_1}] \text{Tr}[Z_{A_1B_1}\rho_{A_1B_1}]} \sqrt{\text{Tr}[Y_{A_2B'_2}^{-1}\rho_{A_2B_2}] \text{Tr}[Z_{A_2B_2}\rho_{A_2B_2}]}. \quad (134)$$

Since the above inequality holds for all feasible solutions, we find that

$$F^u(\rho_{A_1B_1} \otimes \rho_{A_2B_2}) \leq F^u(\rho_{A_1B_1})F^u(\rho_{A_2B_2}). \quad (135)$$

This concludes the proof. \blacksquare

Considering the relation between unextendible fidelity and the 1/2-sandwiched unextendible entanglement, Proposition 18 demonstrates that the logarithmic unextendible fidelity is an additive unextendible entanglement measure different from the min- and max-unextendible entanglements.

Corollary 19 *Let $\rho_{A_1B_1} \in \mathcal{S}(A_1B_1)$ and $\rho_{A_2B_2} \in \mathcal{S}(A_2B_2)$ be bipartite states. The following additivity relation holds*

$$\tilde{E}_{1/2}^u(\rho_{A_1B_1} \otimes \rho_{A_2B_2}) = \tilde{E}_{1/2}^u(\rho_{A_1B_1}) + \tilde{E}_{1/2}^u(\rho_{A_2B_2}), \quad (136)$$

where the entanglement is evaluated across the $A|B$ cut.

Proposition 20 *Let ψ_{AB} be a pure bipartite state. Then the following equality holds*

$$F^u(\psi_{AB}) = \lambda_{\max}(\rho_A), \quad (137)$$

where $\rho_A = \text{Tr}_B \psi_{AB}$ and $\lambda_{\max}(\rho_A)$ is the maximal eigenvalue of ρ_A .

Proof. Let ψ_{AB} be a pure bipartite state. Using the relation between unextendible fidelity and 1/2-sandwiched unextendible entanglement, we find that

$$F^u(\psi_{AB}) = 2^{-\tilde{E}_{1/2}^u(\psi_{AB})} = 2^{-H_{\min}(A)_\psi} = \lambda_{\max}(\rho_A), \quad (138)$$

where the second equality follows from Proposition 10. \blacksquare

IV. APPLICATIONS FOR SECRET KEY AND ENTANGLEMENT DISTILLATION

A. Overhead of probabilistic entanglement distillation

Entanglement distillation aims at obtaining maximally entangled states from less entangled bipartite states shared between two parties via certain free operations. As a central task in quantum information processing, various approaches [VP98, Rai99, VW02, Rai00, HHH00, CW04, WD17b,

[LDS18, WD16, FWTD19, KDWW19, RFWG19] have been developed to characterize and approximate the performance of the rates of deterministic entanglement distillation.

Here, instead of considering the rates of deterministic entanglement distillation, we consider the overhead of probabilistic entanglement distillation [BDSW96, CCGFZ99, PSBZ01, NFB14, CB07, RST+18] under selective two-extendible operations (which include 1-LOCC operations as a special case). It turns out that the relative entropy induced unextendible entanglement $E^u(\rho_{AB})$, given in Definition 3, provides a lower bound on the overhead of probabilistic entanglement distillation of a bipartite state ρ_{AB} .

We begin by defining the overhead of probabilistic entanglement distillation:

Definition 7 *The overhead of distilling m ebits from several independent copies of a bipartite state ρ_{AB} is given by the minimum number of copies of ρ_{AB} needed, on average, to produce m copies of the ebit Φ_2 using 1-LOCC operations:*

$$E_{\text{ov}}(\rho_{AB}, m) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow \Phi_2^{\otimes m} \text{ with prob. } p, \Lambda \text{ is 1-LOCC} \right\}. \quad (139)$$

We can also define the overhead when selective two-extendible operations are allowed:

$$E_{\text{ov},2}(\rho_{AB}, m) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow \Phi_2^{\otimes m} \text{ with prob. } p, \Lambda \text{ is selective two-extendible} \right\}. \quad (140)$$

The following inequality is a trivial consequence of definitions and the fact that a 1-LOCC operation is a special kind of selective two-extendible operation:

$$E_{\text{ov}}(\rho_{AB}, m) \geq E_{\text{ov},2}(\rho_{AB}, m). \quad (141)$$

Theorem 21 *For a bipartite state ρ_{AB} , the overhead of distilling m ebits from ρ_{AB} is bounded from below as*

$$E_{\text{ov},2}(\rho_{AB}, m) \geq \frac{m}{E^u(\rho_{AB})}, \quad (142)$$

where $E^u(\rho_{AB})$ is the unextendible entanglement from Definition 3.

Proof. We first show that any generalized unextendible entanglement \mathbf{E}^u satisfying selective two-extendible monotonicity, normalization, and subadditivity serves as a lower bound on the overhead. Let us suppose that the selective two-extendible operation Λ outputs $\Phi_2^{\otimes m}$ from $\rho_{AB}^{\otimes n}$ with probability p . Without loss of generality, we can assume that

$$\Lambda(\rho_{AB}^{\otimes n}) = p|0\rangle\langle 0|_{X_B} \otimes \Phi_2^{\otimes m} + (1-p)|1\rangle\langle 1|_{X_B} \otimes \tau, \quad (143)$$

where τ is some bipartite state and X_B is a flag system indicating whether the conversion is successful. We have

$$n\mathbf{E}^u(\rho_{AB}) \geq \mathbf{E}^u(\rho_{AB}^{\otimes n}) \geq p\mathbf{E}^u(\Phi_2^{\otimes m}) = pm, \quad (144)$$

where the first inequality follows from subadditivity, the second inequality follows from selective two-extendible monotonicity, and the last equality follows from normalization. Since n and p are arbitrary, we conclude that the following bound holds for all integer $n \geq 1$ and $p \in (0, 1]$:

$$\frac{n}{p} \geq \frac{m}{\mathbf{E}^u(\rho_{AB})}. \quad (145)$$

As the relative entropy induced unextendible entanglement $E^u(\rho_{AB})$ satisfies all these required properties, we conclude the lower bound in (142). \blacksquare

Remark 3 In Theorem 5, Proposition 11, and Proposition 13, we have proved that the α -Petz unextendible entanglement for $\alpha \in [1, 2]$, the α -sandwiched unextendible entanglement for $\alpha \in [1, \infty]$, and the α -geometric unextendible entanglement for $\alpha \in [1, 2]$ satisfy selective two-extendible monotonicity, normalization, and subadditivity, respectively. That is to say, all these entanglement measures can be used as lower bounds. However, due to the fact that all these divergences are monotonically increasing on α [Tom15] and $D(\omega\|\tau) \leq \widehat{D}(\omega\|\tau)$ [HP91], $E^u(\rho_{AB})$ is the smallest unextendible entanglement measure among these choices and gives the tightest lower bound. What is more, recently the authors of [FSP19] proposed a method to accurately approximate the quantum relative entropy via semidefinite programming. This enables us to estimate the lower bound using available semidefinite programming solvers.

B. Perfect entanglement distillation

We can also consider the setting in which the goal is to distill ebits perfectly from a bipartite state by using two-extendible or 1-LOCC channels. This is certainly less realistic than the above scenario, but it is still a core part of zero-error quantum information theory [GdAM16].

The one-shot 1-LOCC perfect distillable entanglement of ρ_{AB} is defined to be the maximum number of ebits achievable via a 1-LOCC channel; that is,

$$E_{1\text{-LOCC}}^{(1)}(\rho_{AB}) := \sup\{\log_2 d : \Lambda_{AB \rightarrow \hat{A}\hat{B}}(\rho_{AB}) = \Phi_{\hat{A}\hat{B}}^d, \Lambda \in 1\text{-LOCC}\}, \quad (146)$$

where $\Phi_{\hat{A}\hat{B}}^d$ is a maximally entangled state of Schmidt rank d . The 1-LOCC perfect distillable entanglement of a state ρ_{AB} is then defined as the regularization of $E_{1\text{-LOCC}}^{(1)}(\rho_{AB})$:

$$E_{1\text{-LOCC}}(\rho_{AB}) := \lim_{n \rightarrow \infty} \frac{1}{n} E_{1\text{-LOCC}}^{(1)}(\rho_{AB}^{\otimes n}). \quad (147)$$

Note that the following inequality is a direct consequence of the definitions:

$$E_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq E_{1\text{-LOCC}}(\rho_{AB}). \quad (148)$$

The one-shot two-extendible perfect distillable entanglement and two-extendible perfect distillable entanglement of ρ_{AB} are defined similarly:

$$E_{2\text{-EXT}}^{(1)}(\rho_{AB}) := \sup\{\log_2 d : \Lambda_{AB \rightarrow \hat{A}\hat{B}}(\rho_{AB}) = \Phi_{\hat{A}\hat{B}}^d, \Lambda \in 2\text{-EXT}\}, \quad (149)$$

$$E_{2\text{-EXT}}(\rho_{AB}) := \lim_{n \rightarrow \infty} \frac{1}{n} E_{2\text{-EXT}}^{(1)}(\rho_{AB}^{\otimes n}), \quad (150)$$

and the following inequality is a direct consequence of definitions:

$$E_{2\text{-EXT}}^{(1)}(\rho_{AB}) \leq E_{2\text{-EXT}}(\rho_{AB}). \quad (151)$$

Immediate consequences of definitions and the fact that 1-LOCC operations are contained in the set of two-extendible operations are the following inequalities:

$$E_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq E_{2\text{-EXT}}^{(1)}(\rho_{AB}), \quad E_{1\text{-LOCC}}(\rho_{AB}) \leq E_{2\text{-EXT}}(\rho_{AB}), \quad (152)$$

The min-unextendible entanglement $E_{\min}^u(\rho)$, as given in Definition 5, serves as an upper bound on the two-extendible perfect distillable entanglement of ρ .

Theorem 22 For bipartite state ρ_{AB} , its asymptotic perfect distillable entanglement under two-extendible operations is bounded from above by the min-unextendible entanglement:

$$E_{2\text{-EXT}}(\rho_{AB}) \leq E_{\min}^u(\rho_{AB}). \quad (153)$$

Proof. For given bipartite state ρ_{AB} , suppose that there is a two-extendible channel $\Lambda_{AB \rightarrow \hat{A}\hat{B}}$ that transforms ρ_{AB} to a maximally entangled state $\Phi_{\hat{A}\hat{B}}^d$. By the monotonicity of the min-unextendible entanglement, the following inequality holds

$$E_{\min}^u(\rho_{AB}) \geq E_{\min}^u(\Lambda_{AB \rightarrow \hat{A}\hat{B}}(\rho_{AB})) = E_{\min}^u(\Phi_{\hat{A}\hat{B}}^d) = \log_2 d. \quad (154)$$

Therefore, by optimizing over all two-extendible protocols, it follows that the one-shot perfect distillable entanglement of ρ_{AB} is bounded as

$$E_{2\text{-EXT}}^{(1)}(\rho_{AB}) \leq E_{\min}^u(\rho_{AB}). \quad (155)$$

Applying the same reasoning to the tensor-power state $\rho_{AB}^{\otimes n}$, we find that

$$E_{2\text{-EXT}}(\rho_{AB}) = \lim_{n \rightarrow \infty} \frac{1}{n} E_{2\text{-EXT}}^{(1)}(\rho_{AB}^{\otimes n}) \leq \lim_{n \rightarrow \infty} \frac{1}{n} E_{\min}^u(\rho_{AB}^{\otimes n}) = E_{\min}^u(\rho_{AB}), \quad (156)$$

where the final equality is a consequence of Proposition 15. ■

C. Private states and unextendible entanglement

In this section, we review the definition of a private state [HHHO05, HHHO09], and then we establish a bound on the number of private bits contained in a private state in terms of the state's unextendible entanglement. These results find application in the next two sections, where we investigate scenarios similar to what we considered for entanglement distillation. Namely, the next two sections consider the overhead of probabilistic secret key distillation and the distillable perfect secret key.

We first review what is meant by a private state [HHHO05, HHHO09]. Let $\rho_{ABA'B'} \in \mathcal{S}(\mathcal{H}_{ABA'B'})$ be a state shared between spatially separated parties Alice and Bob, such that

$$K \equiv \dim(\mathcal{H}_A) = \dim(\mathcal{H}_B). \quad (157)$$

Alice possesses systems A and A' and Bob possesses systems B and B' . A state $\rho_{ABA'B'}$ is called a private state [HHHO05, HHHO09] if Alice and Bob can extract a secret key from it by performing local measurements on A and B , which is product with any purifying system of $\rho_{ABA'B'}$. That is, $\rho_{ABA'B'}$ is a private state of $\log_2 K$ private bits if, for any purification $|\varphi^\rho\rangle_{ABA'B'E}$ of $\rho_{ABA'B'}$, the following holds:

$$(\mathcal{M}_A \otimes \mathcal{M}_B \otimes \text{Tr}_{A'B'}) (\varphi_{ABA'B'E}^\rho) = \frac{1}{K} \sum_{k=1}^K |k\rangle\langle k|_A \otimes |k\rangle\langle k|_B \otimes \sigma_E, \quad (158)$$

where $\mathcal{M}(\cdot) = \sum_k |k\rangle\langle k|(\cdot)|k\rangle\langle k|$ is a projective measurement channel and σ_E is some state on the purifying system. The systems A and B are known as key systems, and A' and B' are known as shield systems. Interestingly, it was shown that a private state of $\log K$ private bits can be written in the following form [HHHO05, HHHO09]

$$\gamma_{ABA'B'} := U_{ABA'B'} (\Phi_{AB} \otimes \sigma_{A'B'}) U_{ABA'B'}^\dagger, \quad (159)$$

where Φ_{AB} is a maximally entangled state of Schmidt rank K , $\sigma_{A'B'}$ is an arbitrary state, and

$$U_{ABA'B'} := \sum_{i,j=1}^K |i\rangle\langle i|_A \otimes |j\rangle\langle j|_B \otimes U_{A'B'}^{ij}. \quad (160)$$

is a controlled unitary known as a ‘‘twisting unitary’’ with each $U_{A'B'}^{ij}$ a unitary.

We now establish some bounds on the number of private bits contained in a private state, in terms of its unextendible entanglement. As it turns out, the unextendible entanglement of a private state is bounded from below by the generalized mutual information of Φ_{AB} . Recall that the generalized mutual information of a state ρ_{AB} is defined as follows [SW12]:

$$I_{\mathbf{D}}(A; B)_\rho := \inf_{\sigma_B} \mathbf{D}(\rho_{AB} \| \rho_A \otimes \sigma_B), \quad (161)$$

where \mathbf{D} is the generalized divergence discussed in Section II B.

Proposition 23 *For a γ -bipartite private state of the form in (159), the following bound holds*

$$\mathbf{E}^u(\gamma_{AA'BB'}) \geq \frac{1}{2} I_{\mathbf{D}}(A; B)_\Phi. \quad (162)$$

Proof. Since a maximally entangled state is not extendible, the only possible extension of $\gamma_{ABA'B'}$ has the following form:

$$\gamma_{ABEA'B'E'} = U_{ABA'B'} (\Phi_{AB} \otimes \sigma_{EA'B'E'}) U_{ABA'B'}^\dagger \quad (163)$$

$$= \frac{1}{d} \sum_{i,j} |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B \otimes U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{jj})^\dagger, \quad (164)$$

where $\sigma_{EA'B'E'}$ is an extension of $\sigma_{A'B'}$ such that $E \simeq B$ and $E' \simeq B'$. Observe that if we define a different twisting unitary $V_{ABA'B'}$ as

$$V_{ABA'B'} := \sum_{i,j} |i\rangle\langle i|_A \otimes |j\rangle\langle j|_B \otimes U_{A'B'}^{ii} \quad (165)$$

$$= \sum_i |i\rangle\langle i|_A \otimes I_B \otimes U_{A'B'}^{ii}, \quad (166)$$

then

$$V_{ABA'B'} (\Phi_{AB} \otimes \sigma_{EA'B'E'}) V_{ABA'B'}^\dagger = \frac{1}{d} \sum_{i,j} |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B \otimes U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{jj})^\dagger \quad (167)$$

$$= \gamma_{ABEA'B'E'} \quad (168)$$

Then we find that

$$\gamma_{AEA'E'} = \text{Tr}_{BB'} [\gamma_{ABEA'B'E'}] \quad (169)$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] \quad (170)$$

Now consider that

$$\begin{aligned} & \mathbf{D}(\gamma_{ABA'B'} \| \gamma_{AEA'E'}) \\ &= \mathbf{D}(V_{ABA'B'} (\Phi_{AB} \otimes \sigma_{A'B'}) V_{ABA'B'}^\dagger \| \gamma_{AEA'E'}) \end{aligned} \quad (171)$$

$$= \mathbf{D}(\Phi_{AB} \otimes \sigma_{A'B'} \| V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'}) \quad (172)$$

$$\geq \mathbf{D}(\Phi_{AB} \| \text{Tr}_{A'E'} [V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'}]). \quad (173)$$

where the inequality follows from monotonicity under partial trace. Observe that

$$\begin{aligned} & V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'} \\ &= \left(\sum_{i'} |i'\rangle \langle i'|_A \otimes I_E \otimes U_{A'E'}^{i'i'\dagger} \right) \left(\frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] \right) \\ & \quad \times \left(\sum_{i''} |i''\rangle \langle i''|_A \otimes I_E \otimes U_{A'E'}^{i''i''} \right) \end{aligned} \quad (174)$$

$$= \frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes U_{A'E'}^{ii\dagger} \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] U_{A'E'}^{ii}, \quad (175)$$

so that

$$\begin{aligned} & \text{Tr}_{A'E'} [V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'}] \\ &= \text{Tr}_{A'E'} \left[\frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes U_{A'E'}^{ii\dagger} \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] U_{A'E'}^{ii} \right] \end{aligned} \quad (176)$$

$$= \frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes \text{Tr}_{A'E'} \left[U_{A'E'}^{ii\dagger} \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] U_{A'E'}^{ii} \right] \quad (177)$$

$$= \frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes \text{Tr}_{A'E'} \left[\text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] \right] \quad (178)$$

$$= \frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes \text{Tr}_{A'B'E'} \left[U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger \right] \quad (179)$$

$$= \frac{1}{d} \sum_i |i\rangle \langle i|_A \otimes \sigma_E \quad (180)$$

$$= \pi_A \otimes \sigma_E. \quad (181)$$

So we find that

$$\mathbf{D}(\gamma_{ABA'B'} \| \gamma_{AEA'E'}) \geq \mathbf{D}(\Phi_{AB} \| \pi_A \otimes \sigma_E) \geq I_{\mathbf{D}}(A; B)_\Phi. \quad (182)$$

Since this bound holds for any extension $\gamma_{ABEA'B'E'}$ of $\gamma_{ABA'B'}$, we conclude (162) after normalizing by 1/2. \blacksquare

As a corollary of Proposition 23 and Lemmas 7, 8, and 9, we find that if the generalized divergence is set to be the Petz-, sandwiched, or geometric Rényi relative entropy, then the unextendible entanglement of a γ -bipartite private state is bounded from below by the amount of secret key that can be extracted from the state.

Corollary 24 *If the generalized divergence is the Petz-, sandwiched, or geometric Rényi relative entropy with α set so that data processing is satisfied, then the following bound holds*

$$\mathbf{E}_\alpha^u(\gamma_{AA'BB'}) \geq \log_2 K. \quad (183)$$

Note that the corollary above includes the relative entropy, the min-relative entropy, and the max-relative entropy.

D. Overhead of probabilistic secret key distillation

Related to the overhead of probabilistic entanglement distillation discussed previously in Section IV A, we can also consider the overhead of probabilistic secret key distillation. The basic definitions are similar to those from Section IV A.

We now define the overhead of probabilistic secret key distillation:

Definition 8 *The overhead of distilling k private bits from several independent copies of a bipartite state ρ_{AB} is given by the minimum number of copies of ρ_{AB} needed, on average, to produce some private state $\gamma_{ABA'B'}^k$ with $\log_2 K = k$ using 1-LOCC operations:*

$$K_{\text{ov}}(\rho_{AB}, k) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow \gamma_{ABA'B'}^k \text{ with prob. } p, \Lambda \text{ is 1-LOCC} \right\}. \quad (184)$$

We can also define the overhead when selective two-extendible operations are allowed:

$$K_{\text{ov},2}(\rho_{AB}, k) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow \gamma_{ABA'B'}^k \text{ with prob. } p, \Lambda \text{ is selective two-extendible} \right\}. \quad (185)$$

Note that it is not necessary to produce a particular private state $\gamma_{ABA'B'}^k$, but rather just some private state $\gamma_{ABA'B'}^k$ having k private bits.

The following inequality is a trivial consequence of definitions and the fact that a 1-LOCC operation is a special kind of selective two-extendible operation:

$$K_{\text{ov}}(\rho_{AB}, k) \geq K_{\text{ov},2}(\rho_{AB}, k). \quad (186)$$

Theorem 25 *For a bipartite state ρ_{AB} , the overhead of distilling k private bits from ρ_{AB} is bounded from below by*

$$E_{\text{ov},2}(\rho_{AB}, k) \geq \frac{k}{E^u(\rho_{AB})}, \quad (187)$$

where the unextendible entanglement is given in Definition 95.

Proof. The proof proceeds exactly the same as the proof of Theorem 21, but we instead invoke Corollary 24 in (144). \blacksquare

E. Perfect secret key distillation

Considering the importance of a γ -bipartite private state, it is natural to consider the task of secret key distillation, which aims to distill a private state $\gamma_{ABA'B'}$, defined in (159), from a given bipartite state, using available operations.

The one-shot 1-LOCC perfect distillable key of ρ_{AB} is defined to be the maximum number of private bits achievable via a 1-LOCC channel; that is,

$$K_{1\text{-LOCC}}^{(1)}(\rho_{AB}) := \sup\{k : \Lambda_{AB \rightarrow \hat{A}\hat{B}A'B'}(\rho_{AB}) = \gamma_{\hat{A}\hat{B}A'B'}^k, \Lambda \in 1\text{-LOCC}\}, \quad (188)$$

where $\gamma_{\hat{A}\hat{B}A'B'}^k$ is any private state with k private bits. The 1-LOCC perfect distillable key of a state ρ_{AB} is then defined as the regularization of $K_{1\text{-LOCC}}^{(1)}(\rho_{AB})$:

$$K_{1\text{-LOCC}}(\rho_{AB}) := \lim_{n \rightarrow \infty} \frac{1}{n} K_{1\text{-LOCC}}^{(1)}(\rho_{AB}^{\otimes n}). \quad (189)$$

Note that the following inequality holds as a direct consequence of definitions:

$$K_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq K_{1\text{-LOCC}}(\rho_{AB}). \quad (190)$$

The one-shot two-extendible perfect distillable key and two-extendible perfect distillable key of ρ_{AB} are defined similarly:

$$K_{2\text{-EXT}}^{(1)}(\rho_{AB}) := \sup\{k : \Lambda_{AB \rightarrow \hat{A}\hat{B}A'B'}(\rho_{AB}) = \gamma_{\hat{A}\hat{B}A'B'}^k, \Lambda \in 2\text{-EXT}\}, \quad (191)$$

$$K_{2\text{-EXT}}(\rho_{AB}) := \lim_{n \rightarrow \infty} \frac{1}{n} K_{2\text{-EXT}}^{(1)}(\rho_{AB}^{\otimes n}), \quad (192)$$

and the following inequality holds as a direct consequence of definitions:

$$K_{2\text{-EXT}}^{(1)}(\rho_{AB}) \leq K_{2\text{-EXT}}(\rho_{AB}). \quad (193)$$

Immediate consequences of definitions and the fact that 1-LOCC operations are contained in the set of two-extendible operations are the following inequalities:

$$K_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq K_{2\text{-EXT}}^{(1)}(\rho_{AB}), \quad K_{1\text{-LOCC}}(\rho_{AB}) \leq K_{2\text{-EXT}}(\rho_{AB}), \quad (194)$$

The min-unextendible entanglement $E_{\min}^u(\rho)$, as given in Definition 5, serves as an upper bound on the two-extendible perfect distillable key of ρ .

Theorem 26 *For a bipartite state ρ_{AB} , its asymptotic perfect distillable key under two-extendible operations is bounded from above as*

$$K_{2\text{-EXT}}(\rho_{AB}) \leq E_{\min}^u(\rho_{AB}). \quad (195)$$

Proof. The proof proceeds exactly as in the proof of Theorem 22, but we instead invoke Corollary 24 in (154). ■

F. Examples

In this section, we apply our bounds on the overhead of probabilistic entanglement or secret key distillation to two classes of states: isotropic states and erased states. We compare our lower bounds and other known estimations of the overhead to upper bounds derived from known distillation protocols. In particular, we show that our lower bound on the overhead of distillation is tight for erased states.

To begin with, recall the Rains relative entropy [Rai00, ADMVW02]:

$$R(\rho_{AB}) := \min_{\sigma_{AB} \in \text{PPT}'} D(\rho_{AB} \| \sigma_{AB}), \quad (196)$$

where $\text{PPT}'(A:B)$ is a set of subnormalized states defined as

$$\text{PPT}'(A:B) := \{\sigma_{AB} : \sigma_{AB} \geq 0, \|\sigma_{AB}^{T_B}\|_1 \leq 1\}. \quad (197)$$

Since selective LOCC preserves the set PPT' [TWW17, Eq. (23)], it follows that the Rains relative entropy $R(\rho_{AB})$ is monotone under selective LOCC by an argument similar to that given in the proof of Theorem 5. Thus, the Rains relative entropy $R(\rho_{AB})$ can be used to estimate the overhead of distillation under selective LOCC via an approach similar to that given in the proof of Theorem 21.




FIG. 1. Bounds on the overhead of distilling one entanglement bit or private bit from isotropic states. The solid line gives the lower bound for the overhead of probabilistic distillation under 1-LOCC operations while the dashed line gives the lower bound for the overhead of probabilistic distillation under LOCC operations.

Isotropic states: We investigate the class of so-called isotropic states ρ_r , defined as [HH99]

$$\rho_r = r \cdot \Phi(d) + (1 - r) \frac{I - \Phi(d)}{d^2 - 1}, \quad (198)$$

where $r \in [0, 1]$, so that ρ_r is a convex mixture of a maximally entangled state and its orthogonal complement. Let us consider the case $d = 2$ for simplicity. Figure 1 plots the unextendible entanglement and Rains relative entropy for the overhead of probabilistic entanglement or secret key distillation for this set of states.

Erased states: We also consider the class of erased states, which are the Choi states of quantum erasure channels [GBP97]:

$$\rho_{A'B}^\varepsilon = (1 - \varepsilon)\Phi_{AB} + \varepsilon|e\rangle\langle e|_{A'} \otimes \pi_B, \quad (199)$$

where $\varepsilon \in [0, 1]$, Φ_{AB} is the maximally entangled state and $|e\rangle$ is some state that is orthogonal to π_A . For simplicity, we choose $d_A = d_B = 2$ (qubit system), and $d_{A'} = d_A + 1 = 3$. This state can be obtained by Alice and Bob pre-share a two-qubit maximally entangled state Φ_{AB} , Alice transmits her local copy with probability $1 - \varepsilon$ and “erases” it (replaces it with the erasure flag state e) with probability ε .

Interestingly, there is a simple formula for the unextendible entanglement of the erased state.

Proposition 27 *The unextendible entanglement of the erased state $\rho_{A'B}^\varepsilon$ is $E^u(\rho_{A'B}^\varepsilon) = 1 - \varepsilon$.*

Proof. Note that the erased state $\rho_{A'B}^\varepsilon$ can be written in the following direct-sum form:

$$\rho_{A'B}^\varepsilon = (1 - \varepsilon)\Phi_{AB} \oplus \varepsilon\pi_B = \begin{bmatrix} (1 - \varepsilon)\Phi_{AB} & 0 \\ 0 & \varepsilon\pi_B \end{bmatrix} \quad (200)$$

From this fact, one can see that each extension $\rho_{A'BB'}$ of $\rho_{A'B}^\varepsilon$ has the form

$$\rho_{A'BB'} := \begin{bmatrix} (1 - \varepsilon)\sigma_{ABB'} & 0 \\ 0 & \varepsilon\sigma_{BB'} \end{bmatrix} \quad (201)$$

such that $\text{Tr}_{B'} \sigma_{ABB'} = \Phi_{AB}$ and $\text{Tr}_{B'} \sigma_{BB'} = \pi_B$. What is more, any extension of Φ_{AB} necessarily has the form $\sigma_{ABB'} := \Phi_{AB} \otimes \tau_{B'}$, where $\tau_{B'}$ is a state on system B' . Thus

$$E^u(\rho_{A'B}^\varepsilon) = \inf_{\rho_{A'BB'}} \text{st. Eq. (201)} \frac{1}{2} D(\rho_{A'B}^\varepsilon \| \rho_{A'B'}) \quad (202)$$

$$= \inf_{\tau_{B'}, \sigma_{B'}} \frac{1}{2} D \left(\left[\begin{array}{cc} (1-\varepsilon)\Phi_{AB} & 0 \\ 0 & \varepsilon\pi_B \end{array} \right] \left\| \left[\begin{array}{cc} (1-\varepsilon)\pi_A \otimes \tau_{B'} & 0 \\ 0 & \varepsilon\sigma_{B'} \end{array} \right] \right) \quad (203)$$

$$= (1-\varepsilon) \inf_{\tau_{B'}} \frac{1}{2} D(\Phi_{AB} \| \pi_A \otimes \tau_{B'}) + \varepsilon \inf_{\sigma_{B'}} \frac{1}{2} D(\pi_B \| \sigma_{B'}) \quad (204)$$

$$= (1-\varepsilon) \frac{1}{2} I(A; B)_\Phi \quad (205)$$

$$= (1-\varepsilon), \quad (206)$$

where $I(A; B)_\rho$ is the quantum mutual information of ρ_{AB} . ■

Theorem 28 *For the erased state $\rho_{A'B}^\varepsilon$, we have*

$$E_{\text{ov}}(\rho_{A'B}^\varepsilon, 1) = E_{\text{ov},2}(\rho_{A'B}^\varepsilon, 1) = \frac{1}{E^u(\rho_{A'B}^\varepsilon)} = \frac{1}{1-\varepsilon}. \quad (207)$$

Proof. In Theorem 21, we showed that $1/E^u(\rho_{A'B}^\varepsilon) \equiv 1/(1-\varepsilon)$ is a lower bound on the overhead of probabilistically distilling one ebit from $\rho_{A'B}^\varepsilon$.

To show the tightness of the lower bound, we propose a one-way LOCC protocol that achieves this lower bound. Given $\rho_{A'B}^\varepsilon$, Alice detects whether her local system is erased or not by performing a binary measurement $\{\mathbb{1}_A, |e\rangle\langle e|\}$. With probability $1-\varepsilon$, she finds that the system is not erased. Then she sends this information to Bob, and the protocol finishes with one ebit shared between them. ■

Therefore, our estimation on the overhead of distilling ebits from erased states under selective one-way LOCC is optimal in the sense that the upper bound on the overhead of the above protocol matches our lower bound from Theorem 21, thus characterizing the ability of probabilistic distillation for erased states. Note that this result can be generalized to multiple copies of the erased state.

In the following proposition, we further consider the overhead of distillation under LOCC operations for the erased states.

Proposition 29 *The Rains bound of the erased state $\rho_{A'B}^\varepsilon$ is $R(\rho_{A'B}^\varepsilon) = 1-\varepsilon$, which implies that the overhead of distillation under selective LOCC for $\rho_{A'B}^\varepsilon$ is given by $(1-\varepsilon)^{-1}$.*

Proof. Since $1/R(\rho)$ is a lower bound on the overhead of distillation using selective LOCC, it follows from the optimality of erased state distillation under one-way LOCC that $1/(1-\varepsilon) \geq 1/R(\rho_{A'B}^\varepsilon)$. This gives $R(\rho_{A'B}^\varepsilon) \geq 1-\varepsilon$.

It then suffices to construct a feasible state to achieve the lower bound, and here we follow an approach similar to that from [TWW17, Proposition 11]. Consider the following state

$$\sigma_{A'B} := \frac{1-\varepsilon}{2} (|00\rangle\langle 00| + |11\rangle\langle 11|) + \varepsilon |e\rangle\langle e| \otimes \pi_B. \quad (208)$$

The state $\sigma_{A'B}$ is a separable state and thus $\sigma_{A'B} \in \text{PPT}'(A':B)$. It then holds that

$$R(\rho_{A'B}^\varepsilon) \leq D(\rho_{A'B}^\varepsilon \| \sigma_{A'B}) \quad (209)$$

$$= D \left((1 - \varepsilon)\Phi_{AB} \oplus \varepsilon\pi_B \left\| \frac{1 - \varepsilon}{2} (|00\rangle\langle 00| + |11\rangle\langle 11|) \oplus \varepsilon\pi_B \right. \right) \quad (210)$$

$$= (1 - \varepsilon)D(\Phi_{AB} \| (|00\rangle\langle 00| + |11\rangle\langle 11|) / 2) \quad (211)$$

$$= 1 - \varepsilon. \quad (212)$$

This concludes the proof. \blacksquare

Proposition 29 indicates an interesting fact that the optimal protocol for distilling one ebit from erased states under one-way LOCC operations matches the lower bound $1/R(\rho_{A'B}^\varepsilon) \equiv 1/(1 - \varepsilon)$ on the overhead under LOCC operations. This reveals that one-way LOCC operations have the same power and performance as LOCC operations in probabilistically distilling ebits from erased states.

V. CONCLUDING REMARKS

We have established a state-dependent resource theory, which is useful in quantifying the unextendibility of entanglement. Our approach was inspired by the fact that the more a bipartite state is entangled, the less that each of its individual systems can be entangled with a third party. Our work has extended this fundamental feature to a quantifiable property, by considering the state-dependent free states. Our results shed light on fundamental problems in the study of manipulating entanglement as a resource. In particular, we have introduced a comprehensive set of measures called *unextendible entanglement* that directly quantify the unextendibility of entanglement, and we have shown that they are monotone under two-extendible operations, which include local operations and one-way classical communication as a special case. Unextendible entanglement has several other desirable properties—some of the unextendible entanglement quantities are simple to analyze and are efficiently computable.

Furthermore, as applications, the unextendible entanglement quantities in our work have found immediate operational applications in evaluating the overhead of probabilistic entanglement or secret key distillation, as well in bounding the rate of perfect entanglement or secret key distillation. Applying the unextendible entanglement, we in particular characterize the optimal overhead of distilling one ebit from the erased states under 1-LOCC operations. We also remark that our results on the overhead can be easily extended to the probabilistic state conversion task, which has been considered for the special pure-state case in [Vid99, AOP16].

An important problem for future work is to investigate to what extent the bounds on the asymptotic distillable secret key or entanglement can be approached. One potential approach is to explore the connections to distinguishability distillation (i.e., hypothesis testing) [Nag06, Hay07, ANSV08, MO15, WW19]. It is also of interest to consider the extension to resource theory of k -unextendibility as well as entanglement dilution [BDSW96, HHT01, VDC02, WD17a, BD11, WW18]. The techniques applied in [XFWD17] might be useful. We hope that the idea of a state-dependent resource theory could provide insights for the study of other topics in quantum resource distillation [LBT19, VCH19, RFWG19], including, for example, coherence [RFA18, FWL⁺18, LRA19, ZLY⁺19] or non-stabilizerness [BK05, BH12, CAB12, WWS18, FL19, KT19, WWS19]. Another interesting direction is to develop the continuous-variable setting, where the results of [LKAW19] could be helpful, as well as the dynamical setting of quantum channels.

ACKNOWLEDGEMENT

The authors thank Anurag Anshu, Kun Fang, Sumeet Khatri, Debbie Leung, and Dave Touchette for discussions. Part of this work was done when XW was at the Joint Center for

Quantum Information and Computer Science, University of Maryland. MMW acknowledges support from NSF under grant no. 1907615.

-
- [ADMVW02] Koenraad Audenaert, Bart De Moor, Karl Gerd H Vollbrecht, and Reinhard F Werner. Asymptotic relative entropy of entanglement for orthogonally invariant states. *Physical Review A*, 66(3):032310, 2002.
- [AI07] Gerardo Adesso and Fabrizio Illuminati. Strong monogamy of bipartite and genuine multipartite entanglement: The Gaussian case. *Physical Review Letters*, 99(15):2–5, October 2007. arXiv:quant-ph/0703277.
- [ANSV08] K. M. R. Audenaert, M. Nussbaum, A. Szkoła, and F. Verstraete. Asymptotic Error Rates in Quantum Hypothesis Testing. *Communications in Mathematical Physics*, 279(1):251–283, April 2008. arXiv:0708.4282.
- [AOP16] Álvaro M Alhambra, Jonathan Oppenheim, and Christopher Perry. Fluctuating states: What is the probability of a thermodynamical transition? *Physical Review X*, 6(4):041016, October 2016. arXiv:1504.00020.
- [Ara90] Huzihiro Araki. On an inequality of Lieb and Thirring. *Letters in Mathematical Physics*, 19(2):167–170, February 1990.
- [BBFS18] Mario Berta, Francesco Borderi, Omar Fawzi, and Volkher Scholz. Semidefinite programming hierarchies for quantum error correction. October 2018. arXiv:1810.12197.
- [BD11] Francesco Buscemi and Nilanjana Datta. Entanglement Cost in Practical Scenarios. *Physical Review Letters*, 106(13):130503, March 2011. arXiv:0906.3698.
- [BDSW96] Charles H Bennett, David P DiVincenzo, John A Smolin, and William K Wootters. Mixed-state entanglement and quantum error correction. *Physical Review A*, 54(5):3824–3851, November 1996. arXiv:quant-ph/9604024.
- [Bei13] Salman Beigi. Sandwiched Rényi divergence satisfies data processing inequality. *Journal of Mathematical Physics*, 54(12):122202, December 2013. arXiv:1306.5920.
- [BH12] Sergey Bravyi and Jeongwan Haah. Magic-state distillation with low overhead. *Physical Review A*, 86(5):052329, November 2012. arXiv:1209.2426.
- [BK05] Sergey Bravyi and Alexei Kitaev. Universal quantum computation with ideal Clifford gates and noisy ancillas. *Physical Review A*, 71(2):022316, February 2005. arXiv:quant-ph/0403025.
- [Bra08] Fernando G. S. L. Brandao. *Entanglement Theory and the Quantum Simulation of Many-Body Physics*. PhD thesis, Imperial College, August 2008. arXiv:0810.0026.
- [BS82] V. P. Belavkin and P. Staszewski. C^* -algebraic generalization of relative entropy and entropy. *Annales de l’I.H.P. Physique théorique*, 37(1):51–58, 1982.
- [BT16] Mario Berta and Marco Tomamichel. The fidelity of recovery is multiplicative. *IEEE Transactions on Information Theory*, 62(4):1758–1763, April 2016. arXiv:1502.07973.
- [CAB12] Earl T. Campbell, Hussain Anwar, and Dan E. Browne. Magic state distillation in all prime dimensions using quantum Reed-Muller codes. *Physical Review X*, 2(4):041021, May 2012. arXiv:1205.3104.
- [CB07] Earl T. Campbell and Simon C. Benjamin. Measurement based entanglement under conditions of extreme photon loss. *Physical Review Letters*, 101(13):130502, October 2007. arXiv:0710.4352.
- [CCGFZ99] C. Cabillo, J. I. Cirac, P. García-Fernández, and P. Zoller. Creation of entangled states of distant atoms by interference. *Physical Review A*, 59(2):1025–1033, February 1999. arXiv:quant-ph/9810013.
- [CG19] Eric Chitambar and Gilad Gour. Quantum resource theories. *Reviews of Modern Physics*, 91(2):025001, April 2019. arXiv:1806.06107.
- [CKW00] Valerie Coffman, Joydip Kundu, and William K. Wootters. Distributed entanglement. *Physical Review A*, 61(5):052306, April 2000. arXiv:quant-ph/9907047.
- [CW04] Matthias Christandl and Andreas Winter. “squashed entanglement”: An additive entanglement measure. *Journal of Mathematical Physics*, 45(3):829–840, March 2004. arXiv:quant-ph/0308088.
- [Dat09] Nilanjana Datta. Min- and max-relative entropies and a new entanglement monotone. *IEEE Transactions on Information Theory*, 55(6):2816–2826, June 2009. arXiv:0803.2770.

- [DFJY05] Runyao Duan, Yuan Feng, Zhengfeng Ji, and Mingsheng Ying. Efficiency of deterministic entanglement transformation. *Physical Review A*, 71(2):022305, February 2005. arXiv:quant-ph/0410046.
- [DL14] Nilanjana Datta and Felix Leditzky. A limit of the quantum Rényi divergence. *Journal of Physics A: Mathematical and Theoretical*, 47(4):045304, January 2014. arXiv:1308.5961.
- [DPS02] Andrew C. Doherty, Pablo A. Parrilo, and Federico M. Spedalieri. Distinguishing separable and entangled states. *Physical Review Letters*, 88(18):187904, April 2002. arXiv:quant-ph/0112007.
- [DPS04] Andrew C. Doherty, Pablo A. Parrilo, and Federico M. Spedalieri. Complete family of separability criteria. *Physical Review A*, 69(2):022308, February 2004. arXiv:quant-ph/0308032.
- [FF19] Kun Fang and Hamza Fawzi. Geometric Rényi divergence and its applications in quantum channel capacities. September 2019. arXiv:1909.05758.
- [FL13] Rupert L. Frank and Elliott H. Lieb. Monotonicity of a relative Rényi entropy. *Journal of Mathematical Physics*, 54(12):122201, December 2013. arXiv:1306.5358.
- [FL19] Kun Fang and Zi-Wen Liu. No-go theorems for quantum resource distillation. September 2019. arXiv:1909.02540.
- [FSP19] Hamza Fawzi, James Saunderson, and Pablo A Parrilo. Semidefinite approximations of the matrix logarithm. *Foundations of Computational Mathematics*, 19(2):259–296, April 2019. arXiv:1705.00812.
- [FWL⁺18] Kun Fang, Xin Wang, Ludovico Lami, Bartosz Regula, and Gerardo Adesso. Probabilistic distillation of quantum coherence. *Physical Review Letters*, 121(7):070404, August 2018. arXiv:1804.09500.
- [FWTD19] Kun Fang, Xin Wang, Marco Tomamichel, and Runyao Duan. Non-asymptotic entanglement distillation. *IEEE Transactions on Information Theory*, 65(10):6454–6465, June 2019. arXiv:1706.06221.
- [GBP97] Markus Grassl, Thomas Beth, and Thomas Pellizzari. Codes for the quantum erasure channel. *Physical Review A*, 56(1):33–38, July 1997. arXiv:quant-ph/9610042.
- [GdAM16] Elloa B. Guedes, Francisco Marcos de Assis, and Rex Antonio da Costa Medeiros. *Quantum Zero-Error Information Theory*. Springer, 2016.
- [GG18] Gilad Gour and Yu Guo. Monogamy of entanglement without inequalities. *Quantum*, 2:81, August 2018. arXiv:1710.03295.
- [GW15] Manish Gupta and Mark M. Wilde. Multiplicativity of completely bounded p -norms implies a strong converse for entanglement-assisted capacity. *Communications in Mathematical Physics*, 334(2):867–887, March 2015. arXiv:1310.7028.
- [Hay07] Masahito Hayashi. Error exponent in asymmetric quantum hypothesis testing and its application to classical-quantum channel coding. *Physical Review A*, 76(6):062301, December 2007. arXiv:quant-ph/0611013.
- [Hay17] Masahito Hayashi. *Quantum Information Theory*. Graduate Texts in Physics. Springer, Berlin, Heidelberg, 2017.
- [HH99] Michał Horodecki and Paweł Horodecki. Reduction criterion of separability and limits for a class of distillation protocols. *Physical Review A*, 59(6):4206–4216, June 1999. arXiv:quant-ph/9708015.
- [HHH00] Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki. Asymptotic manipulations of entanglement can exhibit genuine irreversibility. *Physical Review Letters*, 84(19):4260, May 2000.
- [HHHH09] Ryszard Horodecki, Paweł Horodecki, Michał Horodecki, and Karol Horodecki. Quantum entanglement. *Reviews of Modern Physics*, 81(2):865–942, June 2009. arXiv:quant-ph/0702225.
- [HHHO05] Karol Horodecki, Michał Horodecki, Paweł Horodecki, and Jonathan Oppenheim. Secure key from bound entanglement. *Physical Review Letters*, 94(16):160502, April 2005. arXiv:quant-ph/0309110.
- [HHHO09] Karol Horodecki, Michał Horodecki, Paweł Horodecki, and Jonathan Oppenheim. General paradigm for distilling classical key from quantum states. *IEEE Transactions on Information Theory*, 55(4):1898–1929, April 2009. arXiv:quant-ph/0506189.
- [HHT01] Patrick M. Hayden, Michał Horodecki, and Barbara M. Terhal. The asymptotic entanglement cost of preparing a quantum state. *Journal of Physics A: Mathematical and General*, 34(35):6891–6898, September 2001. arXiv:quant-ph/0008134.
- [HM17] Fumio Hiai and Milan Mosonyi. Different quantum f -divergences and the reversibility of quantum operations. *Reviews in Mathematical Physics*, 29(07):1750023, August 2017. arXiv:1604.03089.
- [HMW13] Patrick Hayden, Kevin Milner, and Mark M. Wilde. Two-message quantum interactive proofs and the quantum separability problem. In *Proceedings of the 28th IEEE Conference on Computational Complexity*, pages 156–167, Palo Alto, California, USA, June 2013. arXiv:1211.6120.

- [HP91] Fumio Hiai and Dénes Petz. The proper formula for relative entropy and its asymptotics in quantum probability. *Communications in Mathematical Physics*, 143(1):99–114, 1991.
- [IRS17] Raban Iten, Joseph M. Renes, and David Sutter. Pretty good measures in quantum information theory. *IEEE Transactions on Information Theory*, 63(2):1270–1279, February 2017. arXiv:1608.08229.
- [KDWW19] Eneet Kaur, Siddhartha Das, Mark M. Wilde, and Andreas Winter. Extendibility limits the performance of quantum processors. *Physical Review Letters*, 123(7):070502, August 2019. arXiv:1803.10710.
- [KL17] Sumeet Khatri and Norbert Lütkenhaus. Numerical evidence for bound secrecy from two-way postprocessing in quantum key distribution. *Physical Review A*, 95(4):042320, April 2017. arXiv:1612.07734.
- [KT19] Anirudh Krishna and Jean-Pierre Tillich. Towards low overhead magic state distillation. *Physical Review Letters*, 123(7):070507, August 2019. arXiv:1811.08461.
- [KW04] Masato Koashi and Andreas Winter. Monogamy of quantum entanglement and other correlations. *Physical Review A*, 69(2):22309, February 2004. arXiv:quant-ph/0310037.
- [LBT19] Zi-Wen Liu, Kaifeng Bu, and Ryuji Takagi. One-Shot Operational Quantum Resource Theory. *Physical Review Letters*, 123(2):020401, July 2019. arXiv:1904.05840.
- [LDH⁺16] Cécilia Lancien, Sara Di Martino, Marcus Huber, Marco Piani, Gerardo Adesso, and Andreas Winter. Should entanglement measures be monogamous or faithful? *Physical Review Letters*, 117(6):060501, August 2016. arXiv:1604.02189.
- [LDS18] Felix Leditzky, Nilanjana Datta, and Graeme Smith. Useful states and entanglement distillation. *IEEE Transactions on Information Theory*, 64(7):4689–4708, July 2018. arXiv:1701.03081.
- [LKAW19] Ludovico Lami, Sumeet Khatri, Gerardo Adesso, and Mark M. Wilde. Extendibility of bosonic gaussian states. *Physical Review Letters*, 123(5):050501, July 2019. arXiv:1904.02692.
- [LRA19] Ludovico Lami, Bartosz Regula, and Gerardo Adesso. Generic bound coherence under strictly incoherent operations. *Physical Review Letters*, 122(15):150402, April 2019. arXiv:1809.06880.
- [LT76] Elliott H. Lieb and Walter Thirring. *Studies in mathematical physics*, chapter Inequalities for the moments of the eigenvalues of the Schroedinger Hamiltonian and their relation to Sobolev inequalities, pages 269–297. Princeton University Press, Princeton, 1976.
- [Mat15] Keiji Matsumoto. A new quantum version of f-divergence. In *Nagoya Winter Workshop: Reality and Measurement in Algebraic Quantum Theory*, pages 229–273. Springer, 2015. arXiv:1311.4722.
- [MCL06] Tobias Moroder, Marcos Curty, and Norbert Lütkenhaus. One-way quantum key distribution: Simple upper bound on the secret key rate. *Physical Review A*, 74(5):052301, November 2006. arXiv:quant-ph/0603270.
- [MLDS⁺13] Martin Müller-Lennert, Frédéric Dupuis, Oleg Szehr, Serge Fehr, and Marco Tomamichel. On quantum rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*, 54(12):122203, December 2013. arXiv:1306.3142.
- [MO15] Milán Mosonyi and Tomohiro Ogawa. Quantum Hypothesis Testing and the Operational Interpretation of the Quantum Rényi Relative Entropies. *Communications in Mathematical Physics*, 334(3):1617–1648, March 2015. arXiv:1309.3228.
- [MRDL09] Geir Ove Myhr, Joseph M. Renes, Andrew C. Doherty, and Norbert Lütkenhaus. Symmetric extension in two-way quantum key distribution. *Physical Review A*, 79(4):042329, April 2009. arXiv:0812.3607.
- [Nag06] Hiroshi Nagaoka. The Converse Part of The Theorem for Quantum Hoeffding Bound. November 2006. arXiv:quant-ph/0611289.
- [NFB14] Naomi H. Nickerson, Joseph F. Fitzsimons, and Simon C. Benjamin. Freely scalable quantum technologies using cells of 5-to-50 qubits with very lossy and noisy photonic links. *Physical Review X*, 4(4):041041, June 2014. arXiv:1406.0880.
- [NH09] Marcin L. Nowakowski and Pawel Horodecki. A simple test for quantum channel capacity. *Journal of Physics A: Mathematical and Theoretical*, 42(13):135306, April 2009. arXiv:quant-ph/0503070.
- [OV06] Tobias J. Osborne and Frank Verstraete. General monogamy inequality for bipartite qubit entanglement. *Physical Review Letters*, 96(22):220503, June 2006. arXiv:quant-ph/0502176.
- [Pet86] Dénes Petz. Quasi-entropies for finite quantum systems. *Reports on Mathematical Physics*, 23(1):57–65, 1986.
- [PSBZ01] Jian-Wei Pan, Christoph Simon, Časlav Brukner, and Anton Zeilinger. Entanglement purification for quantum communication. *Nature*, 410(6832):1067–1070, April 2001.

- [PV10] Yury Polyanskiy and Sergio Verdú. Arimoto channel coding converse and Rényi divergence. In *2010 48th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 1327–1333. IEEE, 2010.
- [Rai99] Eric M. Rains. Bound on distillable entanglement. *Physical Review A*, 60(1):179–184, July 1999. arXiv:quant-ph/9809082.
- [Rai00] Eric M. Rains. A semidefinite program for distillable entanglement. *IEEE Transactions on Information Theory*, 47(7):2921–2933, August 2000. arXiv:quant-ph/0008047.
- [RFA18] Bartosz Regula, Kun Fang, Xin Wang, and Gerardo Adesso. One-shot coherence distillation. *Physical Review Letters*, 121(1):010401, July 2018. arXiv:1711.10512.
- [RFG19] Bartosz Regula, Kun Fang, Xin Wang, and Mile Gu. One-shot entanglement distillation beyond local operations and classical communication. *New Journal of Physics*, 21(10):103017, October 2019. arXiv:1906.01648.
- [RNB19] Bartosz Regula, Varun Narasimhachar, Francesco Buscemi, and Mile Gu. Coherence manipulation with dephasing-covariant operations. July 2019. arXiv:1907.08606.
- [RST⁺18] Filip Rozpedek, Thomas Schiet, Le Phuc Thinh, David Elkouss, Andrew C. Doherty, and Stephanie Wehner. Optimizing practical entanglement distillation. *Physical Review A*, 97(6):062333, June 2018. arXiv:1803.10111.
- [SBW15] Kaushik P. Seshadreesan, Mario Berta, and Mark M. Wilde. Rényi squashed entanglement, discord, and relative entropy differences. *Journal of Physics A: Mathematical and Theoretical*, 48(39):395303, October 2015. arXiv:1410.1443.
- [SW12] Naresh Sharma and Naqeeb Ahmad Warsi. On the strong converses for the quantum channel capacity theorems. 2012. arXiv:1205.1712.
- [SW15] Kaushik P. Seshadreesan and Mark M. Wilde. Fidelity of recovery, squashed entanglement, and measurement recoverability. *Physical Review A*, 92(4):042321, October 2015. arXiv:1410.1441.
- [TDS03] Barbara M. Terhal, Andrew C. Doherty, and David Schwab. Symmetric extensions of quantum states and local hidden variable theories. *Physical Review Letters*, 90(15):157903, April 2003. arXiv:quant-ph/0210053.
- [Ter04] Barbara M. Terhal. Is entanglement monogamous? *IBM Journal of Research and Development*, 48(1):71–78, January 2004. arXiv:quant-ph/0307120.
- [Tom15] Marco Tomamichel. *Quantum Information Processing with Finite Resources: Mathematical Foundations*, volume 5. Springer, 2015.
- [TWW17] Marco Tomamichel, Mark M Wilde, and Andreas Winter. Strong converse rates for quantum communication. *IEEE Transactions on Information Theory*, 63(1):715–727, January 2017. arXiv:1406.2946.
- [Uhl76] Armin Uhlmann. The “transition probability” in the state space of a $*$ -algebra. *Reports on Mathematical Physics*, 9(2):273–279, 1976.
- [Ume62] Hisaharu Umegaki. Conditional expectations in an operator algebra, IV (entropy and information). *Kodai Mathematical Seminar Reports*, 14(2):59–85, June 1962.
- [VCH19] Madhav Krishnan Vijayan, Eric Chitambar, and Min-Hsiu Hsieh. One-shot distillation in a general resource theory. June 2019. arXiv:1906.04959.
- [VDC02] G Vidal, W Dür, and J Ignacio Cirac. Entanglement cost of bipartite mixed states. *Physical Review Letters*, 89(2):027901, June 2002. arXiv:quant-ph/0112131.
- [Vid99] Guifré Vidal. Entanglement of pure states for a single copy. *Physical Review Letters*, 83(5):1046, August 1999. arXiv:quant-ph/9902033.
- [VP98] Vlatko Vedral and Martin B. Plenio. Entanglement measures and purification procedures. *Physical Review A*, 57(3):1619–1633, March 1998. arXiv:quant-ph/9707035.
- [VW02] Guifré Vidal and Reinhard F. Werner. Computable measure of entanglement. *Physical Review A*, 65(3):032314, February 2002. arXiv:quant-ph/0102117.
- [Wat13] John Watrous. Simpler semidefinite programs for completely bounded norms. *Chicago Journal of Theoretical Computer Science*, 8:1–19, 2013. arXiv:1207.5726.
- [WD16] Xin Wang and Runyao Duan. Improved semidefinite programming upper bound on distillable entanglement. *Physical Review A*, 94(5):050301, January 2016. arXiv:1601.07940.
- [WD17a] Xin Wang and Runyao Duan. Irreversibility of asymptotic entanglement manipulation under quantum operations completely preserving positivity of partial transpose. *Physical Review Letters*, 119(18):180506, November 2017. arXiv:1606.09421.

- [WD17b] Xin Wang and Runyao Duan. Nonadditivity of Rains' bound for distillable entanglement. *Physical Review A*, 95(6):062322, June 2017. arXiv:1605.00348.
- [Wer89] Reinhard F. Werner. An application of Bell's inequalities to a quantum state extension problem. *Letters in Mathematical Physics*, 17(4):359–363, May 1989.
- [Wil17] Mark M. Wilde. *Quantum Information Theory*. Cambridge University Press, Cambridge, 2017. arXiv:1106.1445.
- [Wil18] Mark M. Wilde. Optimized quantum f-divergences and data processing. *Journal of Physics A*, 51(37):374002, September 2018. arXiv:1710.10252.
- [WW18] Xin Wang and Mark M. Wilde. Exact entanglement cost of quantum states and channels under ppt-preserving operations. September 2018. arXiv:1809.09592.
- [WW19] Xin Wang and Mark M. Wilde. Resource theory of asymmetric distinguishability. May 2019. arXiv:1905.11629.
- [WWS18] Xin Wang, Mark M. Wilde, and Yuan Su. Efficiently computable bounds for magic state distillation. December 2018. arXiv:1812.10145.
- [WWS19] Xin Wang, Mark M. Wilde, and Yuan Su. Quantifying the magic of quantum channels. *New Journal of Physics*, 21(10):103002, October 2019. arXiv:1903.04483.
- [WWY14] Mark M. Wilde, Andreas Winter, and Dong Yang. Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy. *Communications in Mathematical Physics*, 331(2):593–622, October 2014. arXiv:1306.1586.
- [XFD17] Wei Xie, Kun Fang, Xin Wang, and Runyao Duan. Approximate broadcasting of quantum correlations. *Physical Review A*, 96(2):022302, August 2017. arXiv:1705.06071.
- [Yan06] Dong Yang. A simple proof of monogamy of entanglement. *Physics Letters A*, 360(2):249–250, December 2006. arXiv:quant-ph/0604168.
- [ZLY⁺19] Qi Zhao, Yunchao Liu, Xiao Yuan, Eric Chitambar, and Andreas Winter. One-shot coherence distillation: Towards completing the picture. *IEEE Transactions on Information Theory*, 65(10):6441–6453, October 2019. arXiv:1808.01885.

Appendix A: Proof of Lemma 8

Without loss of generality, let σ_B be an arbitrary state with support equal to the support of the reduced state ψ_B . Then we have that

$$\begin{aligned} & \tilde{D}_\alpha(\psi_{AB} \parallel \psi_A \otimes \sigma_B) \\ &= \frac{1}{\alpha - 1} \log_2 \text{Tr} \left[\left(\psi_{AB}^{1/2} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \psi_{AB}^{1/2} \right)^\alpha \right] \end{aligned} \quad (\text{A1})$$

$$= \frac{1}{\alpha - 1} \log_2 \text{Tr} \left[\left(\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \psi_{AB} \right)^\alpha \right] \quad (\text{A2})$$

$$= \frac{1}{\alpha - 1} \log_2 \text{Tr} \left[\left(|\psi\rangle\langle\psi|_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} |\psi\rangle\langle\psi|_{AB} \right)^\alpha \right] \quad (\text{A3})$$

$$= \frac{1}{\alpha - 1} \log_2 \text{Tr} \left[\left(\langle\psi|_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} |\psi\rangle\langle\psi|_{AB} \right)^\alpha \right] \quad (\text{A4})$$

$$= \frac{1}{\alpha - 1} \log_2 \text{Tr} \left[\left(\text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right] |\psi\rangle\langle\psi|_{AB} \right)^\alpha \right] \quad (\text{A5})$$

$$= \frac{1}{\alpha - 1} \log_2 \left(\text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right] \right)^\alpha \text{Tr} [|\psi\rangle\langle\psi|_{AB}^\alpha] \quad (\text{A6})$$

$$= \frac{\alpha}{\alpha - 1} \log_2 \text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right]. \quad (\text{A7})$$

For any pure bipartite state ψ_{AB} , there exists an operator X_A such that $\psi_{AB} = X_A \Gamma_{AB} X_A^\dagger$ and $\text{Tr}[X_A X_A^\dagger] = 1$, which implies that $\psi_A = X_A X_A^\dagger$. Note that we define

$$\Gamma_{AB} := |\Gamma\rangle\langle\Gamma|_{AB}, \quad |\Gamma\rangle_{AB} := \sum_{i=1}^d |i\rangle_A |i\rangle_B, \quad (\text{A8})$$

where $\{|i\rangle_A\}_i$ and $\{|i\rangle_B\}_i$ are orthonormal bases. Furthermore, by taking the polar decomposition of X_A , there exists a unitary U_A and a density operator ρ_A (having the same spectrum as ψ_A) such that $X_A = U_A \sqrt{\rho_A}$. Then consider that the last line above is equal to

$$\begin{aligned} & \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right] \\ &= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[X_A \Gamma_{AB} X_A^\dagger \left(X_A X_A^\dagger \otimes \sigma_B \right)^{\frac{1-\alpha}{\alpha}} \right] \end{aligned} \quad (\text{A9})$$

$$= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[X_A \Gamma_{AB} X_A^\dagger \left(\left(X_A X_A^\dagger \right)^{\frac{1-\alpha}{\alpha}} \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{A10})$$

$$= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\Gamma_{AB} \left(X_A^\dagger \left(X_A X_A^\dagger \right)^{\frac{1-\alpha}{\alpha}} X_A \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{A11})$$

$$= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\Gamma_{AB} \left(\sqrt{\rho_A} U_A^\dagger \left(U_A \rho_A U_A^\dagger \right)^{\frac{1-\alpha}{\alpha}} U_A \sqrt{\rho_A} \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{A12})$$

$$= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\Gamma_{AB} \left(\rho_A^{\frac{1}{\alpha}} \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{A13})$$

$$= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\Gamma_{AB} \left(\rho_A^{\frac{1}{\alpha}} T(\sigma)_A^{\frac{1-\alpha}{\alpha}} \otimes I_B \right) \right] \quad (\text{A14})$$

$$= \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\rho_A^{\frac{1}{\alpha}} T(\sigma)_A^{\frac{1-\alpha}{\alpha}} \right]. \quad (\text{A15})$$

In the above, $T(\sigma)$ denotes the transpose of the state σ . Now taking a minimum over states σ and applying [MLDS⁺13, Lemma 12], we find that

$$\min_{\sigma} \frac{\alpha}{\alpha-1} \log_2 \text{Tr} \left[\rho_A^{\frac{1}{\alpha}} T(\sigma)_A^{\frac{1-\alpha}{\alpha}} \right] = \frac{\alpha}{\alpha-1} \log_2 \left\| \rho_A^{\frac{1}{\alpha}} \right\|_{\alpha/(2\alpha-1)} \quad (\text{A16})$$

$$= \frac{\alpha}{\alpha-1} \frac{2\alpha-1}{\alpha} \log_2 \text{Tr} \left[(\rho_A^{\frac{1}{\alpha}})^{\alpha/(2\alpha-1)} \right] \quad (\text{A17})$$

$$= \frac{2\alpha-1}{\alpha-1} \log_2 \text{Tr} \left[\rho_A^{\frac{1}{2\alpha-1}} \right] \quad (\text{A18})$$

$$= \frac{2}{1 - (1/[2\alpha-1])} \log_2 \text{Tr} \left[\rho_A^{\frac{1}{2\alpha-1}} \right] \quad (\text{A19})$$

$$= 2H_{[2\alpha-1]^{-1}}(\rho_A). \quad (\text{A20})$$

The conclusion that $H_{[2\alpha-1]^{-1}}(\rho_A) = H_{[2\alpha-1]^{-1}}(\psi_A)$ follows because ρ_A and ψ_A have the same spectrum.

Appendix B: Proof of Lemma 9

Consider that

$$\psi_{AB} = \psi_A^{1/2} \Gamma_{AB} \psi_A^{1/2}, \quad (\text{B1})$$

where

$$\Gamma_{AB} := |\Gamma\rangle\langle\Gamma|_{AB}, \quad |\Gamma\rangle_{AB} := \sum_{i=1}^d |i\rangle_A |i\rangle_B, \quad (\text{B2})$$

$\{|i\rangle_A\}_i$ and $\{|i\rangle_B\}_i$ are the orthonormal bases for the Schmidt decomposition of ψ_{AB} , and d is the Schmidt rank of ψ_{AB} . Then, by noting that it suffices to optimize over states σ_B having the same support as ψ_B , we find that

$$\widehat{I}_\alpha(A; B)_\psi := \min_{\sigma_B} \widehat{D}_\alpha(\psi_{AB} \| \psi_A \otimes \sigma_B) \quad (\text{B3})$$

$$= \min_{\sigma_B} \frac{1}{\alpha - 1} \log_2 \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[(\psi_A \otimes \sigma_B)^{-1/2} \psi_{AB} (\psi_A \otimes \sigma_B)^{-1/2} \right]^\alpha \right] \quad (\text{B4})$$

Consider that

$$\begin{aligned} & \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[\left(\psi_A^{-1/2} \otimes \sigma_B^{-1/2} \right) \psi_{AB} \left(\psi_A^{-1/2} \otimes \sigma_B^{-1/2} \right) \right]^\alpha \right] \\ &= \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[\sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2} \right]^\alpha \right] \end{aligned} \quad (\text{B5})$$

$$= \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[\frac{\sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2}}{\text{Tr}[\sigma_B^{-1}]} \right]^\alpha \right] (\text{Tr}[\sigma_B^{-1}])^\alpha \quad (\text{B6})$$

$$= \text{Tr} \left[(\psi_A \otimes \sigma_B) \frac{\sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2}}{\text{Tr}[\sigma_B^{-1}]} \right] (\text{Tr}[\sigma_B^{-1}])^\alpha \quad (\text{B7})$$

$$= \text{Tr} \left[(\psi_A \otimes \sigma_B) \sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2} \right] (\text{Tr}[\sigma_B^{-1}])^{\alpha-1} \quad (\text{B8})$$

$$= \text{Tr} [\psi_A \Gamma_{AB}] (\text{Tr}[\sigma_B^{-1}])^{\alpha-1} \quad (\text{B9})$$

$$= (\text{Tr}[\sigma_B^{-1}])^{\alpha-1}. \quad (\text{B10})$$

In the fourth step, we are using the fact that $\sigma_B^{-1/2} |\Gamma\rangle_{AB}$ is a vector with norm

$$\left\| \sigma_B^{-1/2} |\Gamma\rangle_{AB} \right\|_2 = \sqrt{\langle \Gamma |_{AB} \sigma_B^{-1/2} \sigma_B^{-1/2} | \Gamma \rangle_{AB}} \quad (\text{B11})$$

$$= \sqrt{\langle \Gamma |_{AB} \sigma_B^{-1} | \Gamma \rangle_{AB}} \quad (\text{B12})$$

$$= \sqrt{\text{Tr}[\sigma_B^{-1}]}. \quad (\text{B13})$$

Then it follows that

$$\widehat{I}_\alpha(A; B)_\psi = \min_{\sigma_B} \frac{1}{\alpha - 1} \log_2 (\text{Tr}[\sigma_B^{-1}])^{\alpha-1} \quad (\text{B14})$$

$$= \min_{\sigma_B} \log_2 \text{Tr}[\sigma_B^{-1}]. \quad (\text{B15})$$

The minimum value occurs when σ_B is the maximally mixed state $\frac{1}{d} \sum_{i=1}^d |i\rangle\langle i|_B$. To conclude this final step, one can use the Lagrange multiplier method. So then the equality in the statement of Lemma 9 follows.

Appendix C: Dual SDP of unextendible fidelity

Here we derive the dual SDP of the primal SDP (123) for unextendible fidelity. We follow an argument similar to that given in [BT16]. We first bring the primal program into standard form

which expresses the primal problem as a maximization over $X \geq 0$ of the functional $\text{Tr}[XA]$ subject to $\Phi(X) = B$. Hence, we set

$$X = \begin{pmatrix} X_{11} & Z_{AB} & \cdot \\ Z_{AB}^\dagger & X_{22} & \cdot \\ \cdot & \cdot & \sigma_{ABB'} \end{pmatrix}, \quad A = \frac{1}{2} \begin{pmatrix} 0 & \mathbb{1}_{AB} & 0 \\ \mathbb{1}_{AB} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \rho_{AB} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \rho_{AB} \end{pmatrix}, \quad (\text{C1})$$

and

$$\Phi(X) := \begin{pmatrix} X_{11} & 0 & 0 \\ 0 & X_{22} - \text{Tr}_B \sigma_{ABB'} & 0 \\ 0 & 0 & \text{Tr}_{B'} \sigma_{ABB'} \end{pmatrix} \quad (\text{C2})$$

The variables with the placeholder ‘ \cdot ’ are of no interest. The dual SDP is a minimization over self-adjoint Y of the functional $\text{Tr}[YB]$ subject to $\Phi^\dagger(Y) \geq A$. The dual variables and adjoint map can be determined to be

$$Y = \begin{pmatrix} Y_{11} & \cdot & \cdot \\ \cdot & Y_{22} & \cdot \\ \cdot & \cdot & Y_{33} \end{pmatrix} \quad \text{and} \quad \Phi^\dagger(Y) := \begin{pmatrix} Y_{11} & 0 & 0 \\ 0 & Y_{22} & 0 \\ 0 & 0 & Y_{33} \otimes \mathbb{1}_{B'} - Y_{22} \otimes \mathbb{1}_B \end{pmatrix}. \quad (\text{C3})$$

This leads to the following dual problem:

$$\begin{aligned} \min \quad & \text{Tr}[(Y_{11} + Y_{33})\rho_{AB}] \\ \text{s.t.} \quad & \begin{pmatrix} Y_{11} & 0 \\ 0 & Y_{22} \end{pmatrix} \geq \frac{1}{2} \begin{pmatrix} 0 & \mathbb{1}_{AB} \\ \mathbb{1}_{AB} & 0 \end{pmatrix} \\ & Y_{33} \otimes \mathbb{1}_{B'} \geq Y_{22} \otimes \mathbb{1}_B \\ & Y_{11}, Y_{22}, Y_{33} \in \mathcal{H}(AB) \end{aligned} \quad (\text{C4})$$

The Slater condition for strong duality is satisfied, using the fact that the primal problem is feasible and the dual problem is strictly feasible. Let $\sigma_{B'}$ be a quantum state, the operator

$$X^* = \begin{pmatrix} \rho_{AB} & 0 & 0 \\ 0 & \rho_A \otimes \sigma_{B'} & 0 \\ 0 & 0 & \rho_{AB} \otimes \sigma_{B'} \end{pmatrix} \quad (\text{C5})$$

is primal feasible since $X \geq 0$ and $\Phi(X^*) = B$. For the dual problem, the operator

$$Y^* = \begin{pmatrix} \mathbb{1}_{AB} & 0 & 0 \\ 0 & \mathbb{1}_{AB} & 0 \\ 0 & 0 & 2\mathbb{1}_{AB} \end{pmatrix}, \quad (\text{C6})$$

is strictly feasible since

$$\Phi^\dagger(Y^*) = \begin{pmatrix} \mathbb{1}_{AB} & 0 & 0 \\ 0 & \mathbb{1}_{AB} & 0 \\ 0 & 0 & \mathbb{1}_{ABB'} \end{pmatrix} > \frac{1}{2} \begin{pmatrix} 0 & \mathbb{1}_{AB} & 0 \\ \mathbb{1}_{AB} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \equiv A. \quad (\text{C7})$$

The dual problem (C4) can be further simplified since the matrix inequality holds if and only if $Y_{11}, Y_{22} > 0$ and $Y_{22} \geq Y_{11}^{-1}$ [Wat13]. Without loss of generality we can choose $Y_{22} = Y_{11}^{-1}$, and the problem simplifies to

$$\begin{aligned} \inf \quad & \frac{1}{2} \text{Tr}[(Y_{AB'}^{-1} + Z_{AB})\rho_{AB}] \\ \text{s.t.} \quad & Z_{AB} \otimes \mathbb{1}_{B'} \geq Y_{AB'} \otimes \mathbb{1}_B \\ & Y_{AB'} > 0 \end{aligned} \quad (\text{C8})$$

By the arithmetic-geometric mean inequality, it holds that

$$\frac{1}{2} \operatorname{Tr} [Y_{AB'}^{-1} \rho_{AB}] + \frac{1}{2} \operatorname{Tr} [Z_{AB} \rho_{AB}] \geq \sqrt{\operatorname{Tr} [Y_{AB'}^{-1} \rho_{AB}] \operatorname{Tr} [Z_{AB} \rho_{AB}]} \quad (\text{C9})$$

for every $Y_{AB'} > 0$, with equality when the two terms are equal. For any feasible pair $(Y_{AB'}, Z_{AB})$, there exists a constant $\lambda > 0$ such that the two trace terms evaluated on $(\lambda Y_{AB'}, \lambda Z_{AB})$ are equal. Hence, we can restrict the optimization to such rescaled pairs of operators, resulting

$$\begin{aligned} \inf \quad & \sqrt{\operatorname{Tr} [Y_{AB'}^{-1} \rho_{AB}] \operatorname{Tr} [Z_{AB} \rho_{AB}]} \\ \text{s.t.} \quad & Z_{AB} \otimes \mathbf{1}_{B'} \geq Y_{AB'} \otimes \mathbf{1}_B \\ & Y_{AB'} > 0 \end{aligned} \quad (\text{C10})$$

We remark that this equivalent representation of the dual problem is essential when proving the multiplicativity of unextendible fidelity.