

Quantifying the unextendibility of entanglement^a

Kun Wang,¹ Xin Wang,^{2,1} and Mark M. Wilde^{3,4}

¹*Institute for Quantum Computing, Baidu Research, Beijing 100193, China*

²*Thrust of Artificial Intelligence, Information Hub,
The Hong Kong University of Science and Technology (Guangzhou), Guangzhou, China*

³*Hearne Institute for Theoretical Physics,
Department of Physics and Astronomy, Center for Computation and Technology,
Louisiana State University, Baton Rouge, Louisiana 70803, USA*

⁴*School of Electrical and Computer Engineering,
Cornell University, Ithaca, New York 14850, USA*

(Dated: January 8, 2024)

Entanglement is a striking feature of quantum mechanics, and it has a key property called unextendibility. In this paper, we present a framework for quantifying and investigating the unextendibility of general bipartite quantum states. First, we define the unextendible entanglement, a family of entanglement measures based on the concept of a state-dependent set of free states. The intuition behind these measures is that the more entangled a bipartite state is, the less entangled each of its individual systems is with a third party. Second, we demonstrate that the unextendible entanglement is an entanglement monotone under two-extendible quantum operations, including local operations and one-way classical communication as a special case. Normalization and faithfulness are two other desirable properties of unextendible entanglement, which we establish here. We further show that the unextendible entanglement provides efficiently computable benchmarks for the rate of exact entanglement or secret key distillation, as well as the overhead of probabilistic entanglement or secret key distillation.

CONTENTS

I. Introduction	2
II. Preliminaries	4
A. Notations	4
B. Quantum divergences	4
III. Quantifying the unextendibility of entanglement	6
A. State-dependent set of free states	6
B. Generalized unextendible entanglement	8
C. α -unextendible entanglement	9
D. Relative-entropy-induced unextendible entanglement	13
IV. Efficiently computable entanglement measures	14
A. Max-unextendible entanglement	14
B. Min-unextendible entanglement	15
C. Unextendible fidelity	16
V. Applications for secret key and entanglement distillation	18
A. Private states and unextendible entanglement	18

^a A preliminary version of this paper has been published in the Proceedings of the 2020 IEEE International Symposium on Information Theory (ISIT 2020) [WWW20].

B. Overhead of probabilistic secret key distillation	19
C. Exact secret key distillation	20
D. Overhead of probabilistic entanglement distillation	21
E. Exact entanglement distillation	21
F. Examples	22
VI. Concluding remarks	25
Acknowledgement	25
References	25
A. Proof of Lemma 1	29
B. Proof of Theorem 5	30
C. Proof of Lemma 8	32
D. Proof of Lemma 9	33
E. Proof of Proposition 13	34
F. Derivations of dual SDPs and strong duality for max- and min-unextendible entanglement	35
G. Proof of Proposition 14	36
H. Proof of Proposition 16	37
I. Dual SDP of unextendible fidelity	37
J. Proof of Proposition 18	39
K. Proof of Proposition 21	40
L. Proof of Theorem 23	41
M. Proof of Theorem 24	42
N. Proof of Proposition 27	42
O. Proof of Proposition 29	43

I. INTRODUCTION

Quantum entanglement is a striking phenomenon of quantum mechanics and a key ingredient in many quantum information processing tasks [HHHH09, Hay16, Wil17]. Unshareability or unextendibility is one of the key features of entanglement [Wer89, DPS02, DPS04]. Unextendibility asserts that the more entangled a bipartite state is, the less entangled its individual systems can be with a third party. Given that classical correlations can be shared among many parties, it emphasizes a key distinction between classical and quantum correlations. The “monogamy of quantum entanglement” [Ter04], which has been described and studied in many

ways [CKW00, Ter04, KW04, Yan06, OV06, AI07, LDMH⁺16, GG18], is also connected to this special characteristic of unextendibility.

Unextendibility of entanglement plays a key role in frontier research on Bell inequalities [TDS03], quantum key distribution [MCL06, MRDL09, KL17], and quantum communication [NH09, KDWW19, BBFS22]. Some recent works [KDWW19, KDWW21] developed a systematic resource theory of unextendibility to investigate the unextendibility of bipartite quantum states. The quantifiers of unextendibility defined in [KDWW19, KDWW21] generalize several known measures introduced in the literature [NH09, Bra08, MCL06, HMW13]. In Remark 1 below, we comment on key differences between the approach of [KDWW19, KDWW21] and our approach taken here.

In this work, we present a resource theory of unextendible entanglement by taking into account a state-dependent collection of free states, which is motivated by the fundamental importance of unextendibility in understanding quantum entanglement. In particular, our contributions are as follows:

1. We first introduce the unextendible entanglement, which is a class of entanglement measures that minimizes the generalized divergence between a given state ρ_{AB} and any possible reduced state $\rho_{AB'}$ of an extension $\rho_{ABB'}$ of ρ_{AB} (Section III). This method is based on the idea that a bipartite state's separate systems are less entangled with a third party, the more entangled the state is.
2. Second, we show that our measures of unextendible entanglement are monotone under two-extendible operations. This also means that they are monotone under local operations and one-way classical communication (1-LOCC) (Section IIIB). We also prove faithfulness and normalization of unextendible entanglement, two more desirable characteristics of the family of entanglement measures. We additionally take into account a variety of important Rényi relative entropy-based unextendible entanglement examples (Section IIIC), some of which are efficiently computable (Section IV).
3. Third, we show how the unextendible entanglement can be utilized to effectively establish fundamental limitations on both the overhead and rate of secret key and entanglement distillation (Section V). These most recent findings suggest that the unextendible entanglement provides a useful lens into the virtues of entanglement as a resource.

Remark 1 (Comparison of our results with [KDWW19, KDWW21]) *Before we delve into the rest of the paper, let us briefly remark on the similarities and differences between our present paper and [KDWW19, KDWW21]. The main aspect in which these works are similar is that they both establish ways of quantifying entanglement in terms of the notion of unextendibility. However, beyond this conceptual link, the works are completely different in their approaches and the resulting applications.*

The papers [KDWW19, KDWW21] establish measures of unextendibility in terms of a comparison between the state of interest and the set of k -extendible states, with the measure being called the relative entropy of unextendibility. The consequences of this choice are that the relative entropy of unextendibility does not give bounds on the asymptotic one-way distillable entanglement or the asymptotic quantum capacity, mainly due to [KDWW21, Lemma 10], which states that the relative entropy of k -unextendibility can never be larger than $\log_2 k$, regardless of the system size. Thus, this quantity is only applicable for providing upper bounds on the rates of these tasks in the non-asymptotic regime.

In contrast, as stated above, here we compare the bipartite state ρ_{AB} of interest, on systems AB , to another bipartite state $\sigma_{AB'}$ on systems AB' , such that there exists a tripartite state on ABB'

with marginals given by ρ_{AB} and $\sigma_{AB'}$. Thus, this measure, called *unextendible entanglement* here, is completely different from the relative entropy of unextendibility from [KDWW19, KDWW21]. Furthermore, as we show in our paper, the unextendible entanglement is applicable in the asymptotic regime, giving upper bounds on the rate of probabilistic distillation of entanglement and secret key.

Thus, the approach here and the approach taken in [KDWW19, KDWW21] are complementary approaches for quantifying the unextendibility of entanglement.

II. PRELIMINARIES

We begin in this section by establishing some notation and reviewing some definitions.

A. Notations

Throughout, quantum systems are denoted by A , B , and C , have finite dimensions, and are associated with their underlying Hilbert spaces. Systems described by the same letter are assumed to be isomorphic: $A_1 \cong A_2$ and $A \cong A'$. The set of linear operators acting on system A is denoted by $\mathcal{L}(A)$, the set of Hermitian operators acting on system A is denoted by $\text{Herm}(A)$, the set of positive semi-definite operators by $\mathcal{P}(A)$, and the set of quantum states (density operators) by $\mathcal{S}(A)$. For a linear operator L , we use L^T to denote its transpose and L^\dagger to denote its conjugate transpose. The α -norm of L is defined as $\|L\|_\alpha := (\text{Tr}[(\sqrt{L^\dagger L})^\alpha])^{1/\alpha}$. Especially, the trace norm is defined as $\|L\|_1 := \text{Tr} \sqrt{L^\dagger L}$. The identity operator is denoted by $\mathbb{1}_A$. An operator of the form $L_A \otimes \mathbb{1}_B$ is written simply L_A . Quantum states of system A are denoted by $\rho_A \in \mathcal{S}(A)$ and pure (rank-one) quantum states by Ψ_A . A maximally entangled state Φ_{AB}^d of Schmidt rank d is defined as $\Phi_{AB}^d := \frac{1}{d} \sum_{i,j=1}^d |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B$, where $\{|i\rangle_A\}_i$ and $\{|i\rangle_B\}_i$ are orthonormal bases for systems A and B , respectively. A maximally entangled state Φ_{AB}^2 is also known as an *ebit*. A purification of a quantum state $\rho_A \in \mathcal{S}(A)$ is a pure state $\Psi_{AR}^\rho \in \mathcal{S}(AR)$ such that $\text{Tr}_R \Psi_{AR}^\rho = \rho_A$, where R is called the purifying system. An extension of a quantum state $\rho_A \in \mathcal{S}(A)$ is a quantum state (not necessarily pure) $\rho_{AE} \in \mathcal{S}(AE)$ such that $\text{Tr}_E \rho_{AE} = \rho_A$, where E is called the extension system. Quantum channels are completely positive (CP) and trace preserving (TP) maps from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ and denoted by $\mathcal{N}_{A \rightarrow B} \in \mathcal{Q}(A \rightarrow B)$. The identity channel from $\mathcal{L}(A)$ to $\mathcal{L}(A)$ is denoted by id_A .

B. Quantum divergences

Let \mathbb{R} denote the field of real numbers. A functional $\mathbf{D} : \mathcal{S}(A) \times \mathcal{S}(A) \rightarrow \mathbb{R} \cup \{+\infty\}$ is a generalized divergence [PV10] if for arbitrary Hilbert spaces \mathcal{H}_A and \mathcal{H}_B , arbitrary states $\rho_A, \sigma_A \in \mathcal{S}(A)$, and an arbitrary channel $\mathcal{N}_{A \rightarrow B} \in \mathcal{Q}(A \rightarrow B)$, the following data-processing inequality holds

$$\mathbf{D}(\rho_A \| \sigma_A) \geq \mathbf{D}(\mathcal{N}_{A \rightarrow B}(\rho_A) \| \mathcal{N}_{A \rightarrow B}(\sigma_A)). \quad (1)$$

The data-processing inequality above implies that there is a constant $c \in \mathbb{R}$ such that $\mathbf{D}(\rho_A \| \sigma_A) \geq c$ for all states ρ and σ . Indeed, we can choose the channel $\mathcal{N}_{A \rightarrow B}$ in (1) to be the trace and replace channel $\rho \rightarrow \text{Tr}[\rho]\omega$, where ω is a state. Then applying (1), we conclude that $\mathbf{D}(\rho_A \| \sigma_A) \geq \mathbf{D}(\omega \| \omega)$. Furthermore, considering the trace and replace channel $\rho \rightarrow \text{Tr}[\rho]\tau$, where τ is a state, and applying the data-processing inequality twice in opposite directions implies that $\mathbf{D}(\omega \| \omega) = \mathbf{D}(\tau \| \tau)$ for all states ω and τ . Thus, we can set $c = \mathbf{D}(\omega \| \omega)$, justifying the claim, and we see that $\mathbf{D}(\rho_A \| \sigma_A)$ takes its minimal value when the two states are equal, i.e., $\rho = \sigma$. Without

loss of generality, we can then redefine the generalized divergence to be $\mathbf{D}(\rho_A \parallel \sigma_A) - c$, so that the redefined quantity satisfies $\mathbf{D}(\omega \parallel \omega) = 0$ for every state ω . We make this assumption in what follows.

We then call a generalized divergence \mathbf{D} faithful if $\mathbf{D}(\rho_A \parallel \sigma_A) = 0$ if and only if $\rho_A = \sigma_A$. Examples of interest are the quantum relative entropy [Ume62], the Petz-Rényi relative entropies [Pet86], the sandwiched Rényi relative entropies [MLDS⁺13, WWY14], and the geometric Rényi relative entropies [Mat15, FF21]. Let us recall the definitions of these relative entropies. The Petz-Rényi relative entropy is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as follows:

$$D_\alpha(\omega \parallel \tau) := \frac{1}{\alpha - 1} \log Q_\alpha(\omega \parallel \tau), \quad (2)$$

$$Q_\alpha(\omega \parallel \tau) := \text{Tr}[\omega^\alpha \tau^{1-\alpha}], \quad (3)$$

where logarithms are taken to base two throughout this paper and $D_1(\omega \parallel \tau)$ is defined as the limit of $D_\alpha(\omega \parallel \tau)$ for $\alpha \rightarrow 1$. The sandwiched Rényi relative entropy is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as

$$\tilde{D}_\alpha(\omega \parallel \tau) := \frac{1}{\alpha - 1} \log \tilde{Q}_\alpha(\omega \parallel \tau), \quad (4)$$

$$\tilde{Q}_\alpha(\omega \parallel \tau) := \left\| \omega^{1/2} \tau^{(1-\alpha)/2\alpha} \right\|_{2\alpha}^{2\alpha} = \text{Tr} \left[\left(\tau^{(1-\alpha)/2\alpha} \omega \tau^{(1-\alpha)/2\alpha} \right)^\alpha \right]. \quad (5)$$

Moreover, $\tilde{D}_1(\omega \parallel \tau)$ is defined as the limit of $\tilde{D}_\alpha(\omega \parallel \tau)$ for $\alpha \rightarrow 1$. The geometric Rényi relative entropy is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as follows:

$$\hat{D}_\alpha(\omega \parallel \tau) := \frac{1}{\alpha - 1} \log \hat{Q}_\alpha(\omega \parallel \tau), \quad (6)$$

$$\hat{Q}_\alpha(\omega \parallel \tau) := \text{Tr} \left[\tau \left(\tau^{-1/2} \omega \tau^{-1/2} \right)^\alpha \right], \quad (7)$$

and $\hat{D}_1(\omega \parallel \tau)$ is defined as the limit of $\hat{D}_\alpha(\omega \parallel \tau)$ for $\alpha \rightarrow 1$. For all of the above quantities, if $\alpha > 1$ and $\text{supp}(\omega) \not\subseteq \text{supp}(\tau)$, then

$$D_\alpha(\omega \parallel \tau) = \tilde{D}_\alpha(\omega \parallel \tau) = \hat{D}_\alpha(\omega \parallel \tau) = +\infty. \quad (8)$$

The Petz-Rényi relative entropy obeys the data-processing inequality for $\alpha \in [0, 2]$ [Pet86], the sandwiched Rényi relative entropy obeys the data-processing inequality for $\alpha \in [1/2, \infty]$ [FL13, Bei13] (see also [Wil18]), and the geometric Rényi relative entropy obeys the data-processing inequality for $\alpha \in (0, 2]$ [Mat15, HM17]. In the case that $\alpha = 1$, we take the limit $\alpha \rightarrow 1$ to find that [Pet86, MLDS⁺13, WWY14]

$$D(\omega \parallel \tau) := \lim_{\alpha \rightarrow 1} \tilde{D}_\alpha(\omega \parallel \tau) = \lim_{\alpha \rightarrow 1} D_\alpha(\omega \parallel \tau) = \text{Tr}[\omega [\log \omega - \log \tau]], \quad (9)$$

where $D(\omega \parallel \tau)$ is the well-known quantum relative entropy [Ume62]. The geometric Rényi relative entropy converges to the Belavkin-Staszewski relative entropy [BS82] in the limit $\alpha \rightarrow 1$:

$$\lim_{\alpha \rightarrow 1} \hat{D}_\alpha(\omega \parallel \tau) = \hat{D}(\omega \parallel \tau) := \text{Tr}[\omega \log(\omega^{1/2} \tau^{-1} \omega^{1/2})], \quad (10)$$

which is known to obey the data-processing inequality [HM17]. In the case that $\alpha = \infty$, we take the limit $\alpha \rightarrow \infty$ to find that [MLDS⁺13]

$$D_{\max}(\omega \parallel \tau) := \lim_{\alpha \rightarrow \infty} \tilde{D}_\alpha(\omega \parallel \tau) = \inf \left\{ \lambda \in \mathbb{R} : \omega \leq 2^\lambda \tau \right\}, \quad (11)$$

where $D_{\max}(\omega\|\tau)$ is the max-relative entropy [Dat09]. See [KW20] for a detailed review of all of these quantities.

The following faithfulness property ensures that the above divergences are non-negative for normalized states and vanishes only if both arguments are equal (proof in Appendix A). It is useful for later analysis.

Lemma 1 (Faithfulness) *Let $\alpha \in (0, \infty)$, and let ω and τ be quantum states. Then $D_\alpha(\omega\|\tau) \geq 0$ and $D_\alpha(\omega\|\tau) = 0$ if and only if $\omega = \tau$. The same is true for the sandwiched and geometric Rényi relative entropies.*

III. QUANTIFYING THE UNEXTENDIBILITY OF ENTANGLEMENT

This section is structured as follows. In Section III A, we define the set of free states determined by a given bipartite quantum state ρ_{AB} . In Section III B, we introduce the generalized unextendible entanglement. In Section III C, we consider special cases of the generalized unextendible entanglement, which include those based on the quantum relative entropy [Ume62], the Petz–Rényi relative entropy [Pet86], the sandwiched Rényi relative entropy [MLDS⁺13, WWY14], and the geometric Rényi relative entropy [Mat15] (see also [FF21]). We establish several desirable properties for these measures, including the fact that they are monotone under two-extendible operations.

A. State-dependent set of free states

Unlike the usual framework of quantum resource theories [CG19] and that which was established for unextendibility in [KDWW19], the free states in our resource theory are *state-dependent*, which is inspired by the definition and meaning of unextendibility of entanglement. Note that state-dependent resource theories were previously considered in a different context [RNBG20]. To be specific, given a bipartite state ρ_{AB} , the free states are those bipartite states that are possibly shareable between Alice and a third party B' , where the system B' is isomorphic to B . Mathematically, the set of free states corresponding to a state ρ_{AB} is defined as follows:

$$\mathcal{F}_{\rho_{AB}} := \{\sigma_{AB'} : \rho_{AB} = \text{Tr}_{B'}[\omega_{ABB'}], \sigma_{AB'} = \text{Tr}_B[\omega_{ABB'}], \omega_{ABB'} \in \mathcal{S}(ABB')\}. \quad (12)$$

Furthermore, if $\rho_{AB} \in \mathcal{F}_{\rho_{AB}}$, then ρ_{AB} is *two-extendible*, indicating that there exists an extension $\omega_{ABB'}$ of ρ_{AB} such that $\rho_{AB} = \text{Tr}_{B'}[\omega_{ABB'}] = \text{Tr}_B[\omega_{ABB'}]$. Separable states are always two-extendible. Interestingly, there are entangled states that are also two-extendible; a concrete example can be found in Eq. (7) of [Doh14]. As so, the resource theory under consideration differs from commonly studied entanglement quantifiers where separable states are usually regarded as constituting the full set of free states.

We introduce *selective two-extendible operations* as the free operations under consideration, which generalize the two-extendible channels previously defined and investigated in [KDWW19]. Mathematically, a selective two-extendible operation consists of a set of CP maps of the following form:

$$\{\mathcal{E}_{AB \rightarrow A'B'}^y\}_{y \in \mathcal{Y}}, \quad (13)$$

such that $\sum_{y \in \mathcal{Y}} \mathcal{E}_{AB \rightarrow A'B'}^y$ is trace-preserving, each map $\mathcal{E}_{AB \rightarrow A'B'}^y$ is completely positive and two-extendible, in the sense that there exists an extension map $\mathcal{E}_{AB_1 B_2 \rightarrow A' B'_1 B'_2}^y$ satisfying [KDWW19]

1. Channel extension:

$$\forall \rho_{AB_1B_2} \in \mathcal{S}(AB_1B_2) : \mathcal{E}_{AB \rightarrow A'B'}^y(\rho_{AB_1}) = \text{Tr}_{B_2} \left[\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y(\rho_{AB_1B_2}) \right], \quad (14)$$

where $\rho_{AB_1} = \text{Tr}_{B_2} \rho_{AB_1B_2}$ is the marginal quantum state of $\rho_{AB_1B_2}$.

2. Permutation covariance:

$$\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y \circ \mathcal{W}_{B_1B_2} = \mathcal{W}_{B'_1B'_2} \circ \mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y, \quad (15)$$

where $\mathcal{W}_{B_1B_2}$ is the unitary swap channel.

The two conditions above ensure that the two marginal operations

$$\mathcal{E}_{AB_1 \rightarrow A'B'_1}^y(\rho_{AB_1}) := \text{Tr}_{B'_2} \left[\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y(\rho_{AB_1B_2}) \right], \quad (16)$$

$$\mathcal{E}_{AB_2 \rightarrow A'B'_2}^y(\rho_{AB_2}) := \text{Tr}_{B'_1} \left[\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y(\rho_{AB_1B_2}) \right], \quad (17)$$

where $\rho_{AB_1} = \text{Tr}_{B_2} \rho_{AB_1B_2}$ is the marginal quantum state, are operations that are in fact equal to the original operation:

$$\mathcal{E}_{AB \rightarrow A'B'}^y = \mathcal{E}_{AB_1 \rightarrow A'B'_1}^y = \mathcal{E}_{AB_2 \rightarrow A'B'_2}^y. \quad (18)$$

We emphasize that the channel extension defined in (14) must have well defined channel marginals, which is a nontrivial property and commonly referred to as the quantum channel marginal problem. We refer interested readers to [HLA22] for comprehensive research on this problem.

A two-extendible channel [KDWW19] is a special case of a selective two-extendible operation for which $|\mathcal{Y}| = 1$.

Now we introduce selective 1-LOCC operations as a special case of selective two-extendible operations. Mathematically, a selective 1-LOCC operation consists of a set of CP maps of the following form:

$$\left\{ \mathcal{L}_{AB \rightarrow A'B'}^y \equiv \sum_{x \in \mathcal{X}} \mathcal{F}_{A \rightarrow A'}^{x,y} \otimes \mathcal{G}_{B \rightarrow B'}^{x,y} \right\}_{y \in \mathcal{Y}}, \quad (19)$$

where \mathcal{X} and \mathcal{Y} are the alphabets, each map $\mathcal{F}_{A \rightarrow A'}^{x,y}$ is completely positive, the sum map $\sum_{x \in \mathcal{X}, y \in \mathcal{Y}} \mathcal{F}_{A \rightarrow A'}^{x,y}$ is trace-preserving, and each map $\mathcal{G}_{B \rightarrow B'}^{x,y}$ is completely positive and trace-preserving. A 1-LOCC channel is a special case of a selective 1-LOCC operation for which $|\mathcal{Y}| = 1$. The fact that selective 1-LOCC operations are a special case of selective two-extendible operations can be verified by constructing the extension map of $\mathcal{L}_{AB \rightarrow A'B'}^y$ in (19) in the following way:

$$\mathcal{L}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y := \sum_x \mathcal{F}_{A \rightarrow A'}^{x,y} \otimes \mathcal{G}_{B_1 \rightarrow B'_1}^{x,y} \otimes \mathcal{G}_{B_2 \rightarrow B'_2}^{x,y}. \quad (20)$$

Such a choice satisfies the channel extension and permutation covariance properties required of selective two-extendible operations. We emphasize that similar inclusions were observed in [KDWW19] for two-extendible channels and 1-LOCC channels.

B. Generalized unextendible entanglement

We introduce a family of entanglement measures called *unextendible entanglement*. Given a bipartite state ρ_{AB} , the key idea behind these measures is to minimize the divergence between ρ_{AB} and any possible reduced state $\rho_{AB'}$ of an extension $\rho_{ABB'}$ of ρ_{AB} . These measures are intuitively motivated by the fact that the more that a bipartite state is entangled, the less that each of its individual systems is entangled with a third party.

Definition 1 (Generalized unextendible entanglement) *The generalized unextendible entanglement of a bipartite state ρ_{AB} is defined as*

$$\mathbf{E}^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} \mathbf{D}(\rho_{AB} \| \sigma_{AB'}), \quad (21)$$

where the infimum ranges over all free states in $\mathcal{F}_{\rho_{AB}}$, as defined in (12).

Remark 2 (Connection to joinability) *In [JV13, Definition 1], the concept of joinability of two bipartite states was considered. That is, two states σ_{AB} and $\tau_{AB'}$ are said to be joinable if there exists a tripartite state $\omega_{ABB'}$ such that $\text{Tr}_{B'}[\omega_{ABB'}] = \sigma_{AB}$ and $\text{Tr}_B[\omega_{ABB'}] = \tau_{AB'}$. With this in mind, the measure in (21) can be understood as quantifying how unjoinable a bipartite state ρ_{AB} is with other bipartite states of systems A and B' .*

Remark 3 (Channel representation) *In [CW04], it was established that a state $\rho_{ABB'}$ is an extension of ρ_{AB} if and only if there exists a quantum channel $\mathcal{R}_{C \rightarrow B'}$ such that $\rho_{ABB'} = \mathcal{R}_{C \rightarrow B'}(\Psi_{ABC})$, where Ψ_{ABC} is a purification of ρ_{AB} . Using this correspondence, the generalized unextendible entanglement can be defined in a dynamical way as*

$$\mathbf{E}^u(\rho_{AB}) := \frac{1}{2} \inf_{\mathcal{R}_{C \rightarrow B'}} \{ \mathbf{D}(\rho_{AB} \| \rho_{AB'}) : \rho_{ABB'} = (\text{id}_{AB} \otimes \mathcal{R}_{C \rightarrow B'}) (|\Psi\rangle\langle\Psi|_{ABC}) \}, \quad (22)$$

where the infimum ranges over every quantum channel $\mathcal{R}_{C \rightarrow B'}$.

In the following, we show that the generalized unextendible entanglement possesses the desirable monotonicity property required for a valid entanglement measure [HHHH09]. Notice that we have only proven the monotonicity for (selective) two-extendible operations, which include (selective) 1-LOCC operations as special cases. We do not expect the unextendible entanglement to be monotone under all LOCC operations.

Theorem 2 (Two-extendible monotonicity) *The generalized unextendible entanglement does not increase under the action of a two-extendible channel. That is, the following inequality holds for an arbitrary bipartite quantum state ρ_{AB} and an arbitrary two-extendible channel $\mathcal{E}_{AB \rightarrow A'B'}$:*

$$\mathbf{E}^u(\rho_{AB}) \geq \mathbf{E}^u(\mathcal{E}_{AB \rightarrow A'B'}(\rho_{AB})). \quad (23)$$

Proof. Let $\rho_{AB_1B_2}$ be an arbitrary extension of ρ_{AB} , with $\rho_{AB} = \rho_{AB_1}$. Since the channel $\mathcal{E}_{AB \rightarrow AB'}$ is two-extendible, there exists an extending channel $\mathcal{E}_{AB_1B_2 \rightarrow AB'_1B'_2}$ that satisfies the conditions stated in (14)–(15). In particular, the marginal channels $\mathcal{E}_{AB_1 \rightarrow AB'_1}$ and $\mathcal{E}_{AB_2 \rightarrow AB'_2}$ are each equal to the original channel $\mathcal{E}_{AB \rightarrow AB'}$. Furthermore, the state $\mathcal{E}_{AB_1B_2 \rightarrow AB'_1B'_2}(\rho_{AB_1B_2})$ is an extension of $\mathcal{E}_{AB_1 \rightarrow AB'_1}(\rho_{AB_1})$ and $\mathcal{E}_{AB_2 \rightarrow AB'_2}(\rho_{AB_2})$. Then we conclude that

$$\mathbf{D}(\rho_{AB_1} \| \rho_{AB_2}) \geq \mathbf{D}(\mathcal{E}_{AB \rightarrow AB'}(\rho_{AB_1}) \| \mathcal{E}_{AB \rightarrow AB'}(\rho_{AB_2})) \quad (24)$$

$$= \mathbf{D}(\mathcal{E}_{AB_1 \rightarrow AB'_1}(\rho_{AB_1}) \| \mathcal{E}_{AB_2 \rightarrow AB'_2}(\rho_{AB_2})) \quad (25)$$

$$\geq 2\mathbf{E}^u(\mathcal{E}_{AB \rightarrow A'B'}(\rho_{AB})). \quad (26)$$

The first inequality follows from monotonicity of the generalized channel divergence under the action of the quantum channel $\mathcal{E}_{AB \rightarrow AB'}$. The equality follows from the observations stated above. The final inequality follows from the definition of the generalized unextendible entanglement, by applying the fact that $\mathcal{E}_{AB_1 B_2 \rightarrow AB'_1 B'_2}(\rho_{AB_1 B_2})$ is an extension of $\mathcal{E}_{AB_1 \rightarrow AB'_1}(\rho_{AB_1})$ and $\mathcal{E}_{AB_2 \rightarrow AB'_2}(\rho_{AB_2})$. Since the inequality above holds for an arbitrary extension $\rho_{AB_1 B_2}$ of ρ_{AB} , we can take an infimum over all such extensions and conclude (23). ■

It turns out that if the underlying generalized divergence is faithful and continuous, then the generalized unextendible entanglement \mathbf{E}^u is faithful, in the sense that $\mathbf{E}^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible.

Proposition 3 (Faithfulness) *If the underlying generalized divergence is faithful, then the generalized unextendible entanglement \mathbf{E}^u is non-negative and $\mathbf{E}^u(\rho_{AB}) = 0$ if ρ_{AB} is two-extendible. If the underlying generalized divergence is also continuous, then the state ρ_{AB} is two-extendible if $\mathbf{E}^u(\rho_{AB}) = 0$.*

Proof. Non-negativity of \mathbf{E}^u follows from the assumption of faithfulness. Now suppose that ρ_{AB} is two-extendible. Then there exists an extension $\rho_{ABB'}$ such that $\rho_{AB} = \rho_{AB'}$. Then it follows that $\mathbf{E}^u(\rho_{AB}) = 0$ by definition and from the assumption that the underlying divergence is faithful.

Now suppose that $\mathbf{E}^u(\rho_{AB}) = 0$. By the assumption of continuity, this means that there exists an extension $\rho_{ABB'}$ of ρ_{AB} such that $\rho_{AB} = \rho_{AB'}$. Then, by definition, ρ_{AB} is two-extendible. ■

C. α -unextendible entanglement

The α -unextendible entanglement of a bipartite state ρ_{AB} is defined by taking the generalized divergence in (21) to be the Petz–Rényi relative entropy [Pet86], the sandwiched Rényi relative entropy $\tilde{D}_\alpha(\omega \| \tau)$ [MLDS⁺13, WWY14], or the geometric Rényi relative entropy [Mat15]. It is meaningful to consider the families of α -unextendible entanglement measures with different quantum divergences for several reasons. First of all, we are able to prove their general properties once and for all, without analyzing them case by case. Second, some special cases of these measures lead to novel applications, as we will show in Section V. It is highly possible that more of them find operational interpretations in other tasks, using the insights learned from the proved properties.

Definition 2 (α -unextendible entanglement) *The α -Petz unextendible entanglement is defined for $\alpha \in [0, \infty)$ as*

$$E_\alpha^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} D_\alpha(\rho_{AB} \| \sigma_{AB'}), \quad (27)$$

the α -sandwiched unextendible entanglement is defined for $\alpha \in (0, \infty]$ as

$$\tilde{E}_\alpha^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} \tilde{D}_\alpha(\rho_{AB} \| \sigma_{AB'}), \quad (28)$$

and the α -geometric unextendible entanglement is defined for $\alpha \in (0, \infty)$ as

$$\hat{E}_\alpha^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} \hat{D}_\alpha(\rho_{AB} \| \sigma_{AB'}). \quad (29)$$

In the following, we show that the α -unextendible entanglement possesses the desirable monotonicity property required for an entanglement measure [HHHH09]. More specifically, it is an immediate consequence of Theorem 2 and data processing that α -unextendible entanglement is monotone under the action of a two-extendible channel for certain values of α .

Corollary 4 (Two-extendible monotonicity) *For $\alpha \in [0, 2]$, the α -Petz unextendible entanglement does not increase under the action of a two-extendible channel. For $\alpha \in [1/2, \infty]$, the α -sandwiched unextendible entanglement does not increase under the action of a two-extendible channel. For $\alpha \in (0, 2]$, the α -geometric unextendible entanglement does not increase under the action of a two-extendible channel.*

Notice that we have only asserted the monotonicity for two-extendible operations, including 1-LOCC operations as special cases. As already mentioned, we do not expect the unextendible entanglement to be monotone under all LOCC operations.

For certain values of $\alpha \geq 1$, the above monotonicity statement can be strengthened, in the sense that the α -unextendible entanglement is monotone under *selective* two-extendible operations. This property is stronger than what we previously proved in Theorem 2. As before, this implies that α -unextendible entanglement is monotone under selective 1-LOCC operations because these are a particular case of selective two-extendible operations. The proof of Theorem 5 can be found in Appendix B.

Theorem 5 (Selective two-extendible monotonicity) *Let ρ_{AB} be a bipartite state, and let $\{\mathcal{E}_{AB \rightarrow A'B'}^y\}_y$ be a selective two-extendible operation. Then the following inequality holds for the α -Petz unextendible entanglement for all $\alpha \in [1, 2]$:*

$$E_\alpha^u(\rho_{AB}) \geq \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'}^y), \quad (30)$$

where

$$p(y) := \text{Tr}[\mathcal{E}_{AB \rightarrow A'B'}^y(\rho_{AB})], \quad (31)$$

$$\omega_{A'B'}^y := \frac{1}{p(y)} \mathcal{E}_{AB \rightarrow A'B'}^y(\rho_{AB}). \quad (32)$$

The following inequality holds for the α -sandwiched unextendible entanglement for all $\alpha \in [1, \infty]$:

$$\tilde{E}_\alpha^u(\rho_{AB}) \geq \sum_{y:p(y)>0} p(y) \tilde{E}_\alpha^u(\omega_{A'B'}^y). \quad (33)$$

The following inequality holds for the α -geometric unextendible entanglement for all $\alpha \in [1, 2]$:

$$\hat{E}_\alpha^u(\rho_{AB}) \geq \sum_{y:p(y)>0} p(y) \hat{E}_\alpha^u(\omega_{A'B'}^y). \quad (34)$$

Since the Petz-, sandwiched, and geometric Rényi relative entropies (and their limits in $\alpha \rightarrow 1$) satisfy all of the requirements in Lemma 1, we can invoke Proposition 3 and conclude the faithfulness property for all α -unextendible entanglement measures.

Proposition 6 (Faithfulness) *For $\alpha \in (0, \infty)$, the α -Petz unextendible entanglement of a bipartite state ρ_{AB} satisfies $E_\alpha^u(\rho_{AB}) \geq 0$, and $E_\alpha^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible. For $\alpha \in (0, \infty]$, the α -sandwiched unextendible entanglement of a bipartite state ρ_{AB} satisfies $\tilde{E}_\alpha^u(\rho_{AB}) \geq 0$, and $\tilde{E}_\alpha^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible. For $\alpha \in (0, \infty)$, the α -geometric unextendible entanglement of a bipartite state ρ_{AB} satisfies $\hat{E}_\alpha^u(\rho_{AB}) \geq 0$, and $\hat{E}_\alpha^u(\rho_{AB}) = 0$ if and only if ρ_{AB} is two-extendible.*

Before stating the next proposition, recall that the Petz–Rényi mutual information of a bipartite state ρ_{AB} is defined as [GW15, Eq. (6.3) and Corollary 8]

$$I_\alpha(A; B)_\rho := \min_{\sigma_B} D_\alpha(\rho_{AB} \| \rho_A \otimes \sigma_B) = \frac{\alpha}{\alpha - 1} \log \text{Tr} \left[\left(\text{Tr}_A [\rho_A^{1-\alpha} \rho_{AB}^\alpha] \right)^{1/\alpha} \right], \quad (35)$$

and the sandwiched Rényi mutual information of a bipartite state ρ_{AB} is defined as [Bei13, GW15]

$$\tilde{I}_\alpha(A; B)_\rho := \min_{\sigma_B} \tilde{D}_\alpha(\rho_{AB} \| \rho_A \otimes \sigma_B), \quad (36)$$

where the minimization in both cases is with respect to every quantum state σ_B . We also define the geometric Rényi mutual information of ρ_{AB} in a similar way:

$$\hat{I}_\alpha(A; B)_\rho := \min_{\sigma_B} \hat{D}_\alpha(\rho_{AB} \| \rho_A \otimes \sigma_B). \quad (37)$$

Also, recall that the Rényi entropy of a quantum state ρ_A is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as

$$H_\alpha(A)_\rho := \frac{1}{1 - \alpha} \log \text{Tr}[\rho_A^\alpha], \quad (38)$$

and it is defined for $\alpha \in \{0, 1, \infty\}$ in the limit, so that

$$H_0(A)_\rho = \log \text{rank}(\rho_A), \quad (39)$$

$$H(A)_\rho := H_1(A)_\rho = -\text{Tr}[\rho_A \log \rho_A], \quad (40)$$

$$H_{\min}(A)_\rho := H_\infty(A)_\rho = -\log \|\rho_A\|_\infty, \quad (41)$$

where $\text{rank}(\rho_A)$ is the rank of ρ_A .

The following lemmas relate the α -mutual information to Rényi entropy when the bipartite state is pure. The first was established in the proof of [SBW15, Proposition 13].

Lemma 7 ([SBW15]) *Let $\alpha \in (0, \infty)$. The α -Petz Rényi mutual information of a pure bipartite state ψ_{AB} reduces to twice the γ -Rényi entropy of its marginal:*

$$I_\alpha(A; B)_\psi = 2H_\gamma(A)_\psi, \quad (42)$$

where $\gamma := [2 - \alpha] / \alpha$.

Lemma 8 *Let $\alpha \in (0, \infty]$. The α -sandwiched Rényi mutual information of a pure bipartite state ψ_{AB} reduces to twice the β -Rényi entropy of its marginal:*

$$\tilde{I}_\alpha(A; B)_\psi = 2H_\beta(A)_\psi, \quad (43)$$

where $\beta := [2\alpha - 1]^{-1}$.

Proof. See Appendix C. ■

Lemma 9 *Let $\alpha \in (0, \infty)$. The α -geometric Rényi mutual information of a pure bipartite state ψ_{AB} reduces to twice the zero-Rényi entropy of its marginal:*

$$\hat{I}_\alpha(A; B)_\psi = 2H_0(A)_\psi. \quad (44)$$

Proof. See Appendix D. ■

The following proposition uses the lemmas above to conclude that the α -unextendible entanglement reduces to Rényi entropy of entanglement for pure states.

Proposition 10 (Reduction for pure states) *Let ψ_{AB} be a pure bipartite state. Then the α -Petz unextendible entanglement reduces to the γ -Rényi entropy of entanglement:*

$$E_\alpha(\psi_{AB}) = H_\gamma(\psi_A), \quad (45)$$

where $\gamma = [2 - \alpha]/\alpha$. The α -sandwiched unextendible entanglement reduces to the β -Rényi entropy of entanglement:

$$\tilde{E}_\alpha(\psi_{AB}) = H_\beta(\psi_A), \quad (46)$$

where $\beta = [2\alpha - 1]^{-1}$. The α -geometric unextendible entanglement reduces to the zero-Rényi entropy of entanglement:

$$\hat{E}_\alpha(\psi_{AB}) = H_0(\psi_A). \quad (47)$$

Proof. For a pure state ψ_{AB} , an arbitrary extension of it has the form $\sigma_{AB_1B_2} := \psi_{AB_1} \otimes \sigma_{B_2}$, where σ_{B_2} is a state of system B_2 . As such, it follows that

$$E_\alpha(\psi_{AB}) = \min_{\sigma_{B_2}} \frac{1}{2} D_\alpha(\psi_{AB} \| \psi_A \otimes \sigma_{B_2}) \quad (48)$$

$$= \frac{1}{2} I_\alpha(A; B)_\psi \quad (49)$$

$$= H_\gamma(\psi_A). \quad (50)$$

The first equality follows from applying the definition of $E_\alpha(\psi_{AB})$, the second equality follows from the definition in (35), and the final equality follows from Lemma 7.

The conclusions about $\tilde{E}_\alpha(\psi_{AB})$ and $\hat{E}_\alpha(\psi_{AB})$ follow the same line of reasoning but using Lemma 8 and Lemma 9 instead, respectively. ■

Following Proposition 10 and the fact that the Rényi entropy of the maximally mixed state (reduced state of Φ_{AB}^d) is equal to $\log d$ for all values of β , we conclude the normalization property and state it formally as follows.

Proposition 11 (Normalization) *Let Φ_{AB}^d be a maximally entangled state of Schmidt rank d . Then $E_\alpha(\Phi_{AB}^d) = \log d$ for $\alpha \in (0, \infty)$, $\tilde{E}_\alpha(\Phi_{AB}^d) = \log d$ for $\alpha \in (0, \infty]$, and $\hat{E}_\alpha(\Phi_{AB}^d) = \log d$ for $\alpha \in (0, \infty)$.*

Let $\{\rho_{AB}^i\}_i \subset \mathcal{S}(AB)$ be a set of bipartite quantum states, and let $\{p_i\}_i$ be a probability distribution such that $p_i \geq 0$ and $\sum_i p_i = 1$. A generalized unextendible entanglement \mathbf{E}^u is *convex* if

$$\mathbf{E}^u\left(\sum_i p_i \rho_{AB}^i\right) \leq \sum_i p_i \mathbf{E}^u(\rho_{AB}^i), \quad (51)$$

and it is *quasi-convex* if

$$\mathbf{E}^u\left(\sum_i p_i \rho_{AB}^i\right) \leq \max_i \mathbf{E}^u(\rho_{AB}^i). \quad (52)$$

If \mathbf{E}^u is convex, then it is also quasi-convex; however, the converse is not necessarily true.

Intuitively, the convexity (quasi-convexity) property implies that the amount of quantum entanglement cannot be increased by merely mixing the quantum states. Convexity is a desirable property of a valid entanglement measure from both the mathematical and physical perspectives. The α -unextendible entanglement satisfies this property for certain parameter ranges:

Proposition 12 (Convexity and quasi-convexity) *The α -Petz unextendible entanglement E_α^u is convex when $\alpha \in (0, 1]$ and quasi-convex when $\alpha \in (0, 2]$. The α -sandwiched unextendible entanglement \tilde{E}_α^u is convex when $\alpha \in [1/2, 1]$ and quasi-convex when $\alpha \geq 1/2$. The α -geometric unextendible entanglement \hat{E}_α^u is convex when $\alpha \in (0, 1]$ and quasi-convex when $\alpha \in (0, 2]$.*

Proof. This follows from the facts that D_α is jointly quasi-convex for $\alpha \in (1, 2]$ and jointly convex for $\alpha \in (0, 1]$ [Pet86]; \tilde{D}_α is jointly quasi-convex for $\alpha > 1$ and jointly convex for $\alpha \in [1/2, 1]$ [FL13]; and \hat{D}_α is jointly quasi-convex for $\alpha \in (1, 2]$ and jointly convex for $\alpha \in (0, 1]$ [HM17]. ■

Aside from the above properties, the α -unextendible entanglement is also subadditive (proof in Appendix E).

Proposition 13 (Subadditivity) *Let $\rho_{A_1 B_1} \in \mathcal{S}(A_1 B_1)$ and $\rho_{A_2 B_2} \in \mathcal{S}(A_2 B_2)$ be two bipartite states. Then the following subadditivity inequality holds for $\alpha \in [0, \infty)$*

$$E_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq E_\alpha^u(\rho_{A_1 B_1}) + E_\alpha^u(\rho_{A_2 B_2}), \quad (53)$$

for $\alpha \in (0, \infty]$

$$\tilde{E}_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \tilde{E}_\alpha^u(\rho_{A_1 B_1}) + \tilde{E}_\alpha^u(\rho_{A_2 B_2}), \quad (54)$$

and for $\alpha \in (0, \infty)$

$$\hat{E}_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \hat{E}_\alpha^u(\rho_{A_1 B_1}) + \hat{E}_\alpha^u(\rho_{A_2 B_2}), \quad (55)$$

where the entanglement is evaluated across the $A:B$ bipartition.

D. Relative-entropy-induced unextendible entanglement

As mentioned above, the quantum relative entropy is a particular instance of the Petz- and sandwiched Rényi relative entropy, recovered by taking the limit $\alpha \rightarrow 1$. The α -unextendible entanglement in terms of these measures has already been defined and investigated above. However, it is notable enough that we define the quantum relative entropy induced unextendible entanglement explicitly here.

Definition 3 (Relative-entropy-induced unextendible entanglement) *For a bipartite state ρ_{AB} , the quantum relative entropy induced unextendible entanglement is defined as*

$$E^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} D(\rho_{AB} \| \sigma_{AB'}), \quad (56)$$

where D is the quantum relative entropy defined in (9).

As proved above, E^u obeys the following properties: selective two-extendible monotonicity, faithfulness, reduction to the entropy of entanglement for pure bipartite states, normalization, convexity, and subadditivity. It is also efficiently computable by means of semi-definite programming, using the approach of [FSP19, FF18].

The third property mentioned above follows directly from Proposition 10 for $\alpha = 1$, and it asserts that for pure bipartite states, the relative-entropy-induced unextendible entanglement evaluates to the von Neumann entropy of the reduced state. Although the proof has already been given, we can see it with a straightforward proof consisting of a few steps. Let $\psi_{AB} \equiv |\psi\rangle\langle\psi|_{AB}$ be a pure state. An arbitrary extension of ψ_{AB} is of the form $\sigma_{ABB'} := \psi_{AB} \otimes \sigma_{B'}$ for some state $\sigma_{B'}$. As so, $\text{Tr}_B[\sigma_{ABB'}] = \psi_A \otimes \sigma_{B'}$, where $\psi_A = \text{Tr}_B[\psi_{AB}]$. Then

$$E^u(\psi_{AB}) = \inf_{\sigma_{B'}} \frac{1}{2} D(\psi_{AB} \| \psi_A \otimes \sigma_{B'}) = \frac{1}{2} I(A; B)_\psi = H(A)_\psi, \quad (57)$$

where the second equality follows from the definition of quantum mutual information.

IV. EFFICIENTLY COMPUTABLE ENTANGLEMENT MEASURES

A. Max-unextendible entanglement

Another interesting instance of the sandwiched Rényi relative entropy is the max-relative entropy, as recalled in (11). The max-relative entropy was originally defined and studied in [Dat09]. Here we adopt the max-relative entropy to define the max-unextendible entanglement. It turns out that this measure is additive and can be calculated efficiently by utilizing a semidefinite program.

Definition 4 (Max-unextendible entanglement) *For a given bipartite state ρ_{AB} , the max-unextendible entanglement is defined as*

$$E_{\max}^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} D_{\max}(\rho_{AB} \| \sigma_{AB'}), \quad (58)$$

where D_{\max} is the max-relative entropy defined in (11).

Note that the infimum in (58) can be replaced with a minimum.

From the definition of max-unextendible entanglement, it follows that it can be computed efficiently by means of a semidefinite program (SDP). To be more specific, the following two optimization programs satisfy strong duality and both evaluate to $2^{-2E_{\max}^u(\rho_{AB})}$.

<u>Primal Program</u>		<u>Dual Program</u>	
maximize λ		minimize $\text{Tr}[\rho_{AB} Y_{AB}]$	
subject to $\lambda \rho_{AB} \leq \sigma_{AB'}$	(59)	subject to $\text{Tr}[\rho_{AB} X_{AB}] \geq 1$	(60)
$\rho_{AB} = \text{Tr}_{B'}[\sigma_{ABB'}]$		$X_{AB'} \otimes \mathbb{1}_B \leq Y_{AB} \otimes \mathbb{1}_{B'}$	
$\sigma_{ABB'} \geq 0$		$X_{AB}, Y_{AB} \geq 0$	

The primal SDP follows by considering that

$$D_{\max}(\omega \| \tau) = \log \inf\{\lambda : \omega \leq \lambda \tau\} = -\log \sup\{\mu : \mu \omega \leq \tau\}. \quad (61)$$

The dual SDP can be obtained by standard methods (e.g., the Lagrange multiplier method). For completeness, we provide a proof in Appendix F.

By using the primal and dual expressions of $2^{-2E_{\max}^u(\rho_{AB})}$ and strong duality, it follows that $E_{\max}^u(\rho_{AB})$ is additive (proof in Appendix G), which is an appealing feature that finds use in Section V.

Proposition 14 (Additivity) *Let $\rho_{A_1 B_1} \in \mathcal{S}(A_1 B_1)$ and $\rho_{A_2 B_2} \in \mathcal{S}(A_2 B_2)$ be two bipartite states. It holds that*

$$E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) = E_{\max}^u(\rho_{A_1 B_1}) + E_{\max}^u(\rho_{A_2 B_2}), \quad (62)$$

where the entanglement is evaluated across the $A:B$ bipartition.

B. Min-unextendible entanglement

In this section, we consider the limit of the Petz–Rényi relative entropy as $\alpha \rightarrow 0$, which is known as the min-relative entropy [Dat09, Definition 2]. Let us first recall the definition. Let $\rho \in \mathcal{S}(A)$ and $\sigma \in \mathcal{P}(A)$. Let Π^ρ denote the projection onto the support of ρ . Then the min-relative entropy of ρ and σ is defined as

$$D_{\min}(\rho \parallel \sigma) := -\log \text{Tr}[\Pi^\rho \sigma]. \quad (63)$$

With D_{\min} , we define the min-unextendible entanglement as follows.

Definition 5 (Min-unextendible entanglement) *For a given bipartite state ρ_{AB} , the min-unextendible entanglement is defined as*

$$E_{\min}^u(\rho_{AB}) := \frac{1}{2} \inf_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} D_{\min}(\rho_{AB} \parallel \sigma_{AB'}). \quad (64)$$

Note that the infimum in (64) can be replaced with a minimum.

Much like the max-unextendible entanglement, the min-unextendible entanglement can also be calculated as the solution to a semidefinite program. The following two optimization programs satisfy strong duality, and both evaluate to $2^{-2E_{\min}^u(\rho_{AB})}$. We derive the dual SDP in Appendix F.

<u>Primal Program</u>		<u>Dual Program</u>
maximize $\text{Tr}[\Pi_{AB}^\rho \sigma_{AB'}]$	(65)	minimize $\text{Tr}[\rho_{AB} X_{AB}]$
subject to $\text{Tr}_{B'}[\sigma_{ABB'}] = \rho_{AB}$		subject to $X_{AB} \otimes \mathbb{1}_{B'} \geq \Pi_{AB'}^\rho \otimes \mathbb{1}_B$
$\sigma_{ABB'} \geq 0$		$X_{AB} \geq 0$

Following similarly to the proof of Proposition 14, we can show that the min-unextendible entanglement is additive.

Proposition 15 (Additivity) *Let $\rho_{A_1 B_1} \in \mathcal{S}(A_1 B_1)$ and $\rho_{A_2 B_2} \in \mathcal{S}(A_2 B_2)$ be two bipartite states. It holds that*

$$E_{\min}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) = E_{\min}^u(\rho_{A_1 B_1}) + E_{\min}^u(\rho_{A_2 B_2}), \quad (67)$$

where the entanglement is evaluated across the $A:B$ bipartition.

Furthermore, the min-unextendible entanglement of a pure bipartite state can be computed explicitly (proof in Appendix H).

Proposition 16 *Let $|\psi\rangle_{AB} := \sum_{i=1}^k \sqrt{\alpha_i} |\psi_i\rangle_A |\psi_i\rangle_B$ be a pure state in \mathcal{H}_{AB} , with $\alpha_1 \geq \dots \geq \alpha_k > 0$. Then the following equality holds*

$$E_{\min}^u(\psi_{AB}) = -\log \alpha_1. \quad (68)$$

Interestingly, $E_{\min}^u(\psi_{AB})$ has an operational interpretation in terms of deterministic entanglement transformation [DFJY05], which we briefly introduce as follows. Let $|\psi_1\rangle$ and $|\psi_2\rangle$ be two

pure bipartite states for systems AB . Let $m \in \mathbb{N}$ be an integer. We define $f(m)$ to be the maximum integer n such that $\psi_1^{\otimes m}$ can be transformed into $\psi_2^{\otimes n}$ by LOCC deterministically. The *deterministic entanglement transformation rate* from ψ_1 to ψ_2 , written $D(\psi_1 \rightarrow \psi_2)$, is defined as

$$D(\psi_1 \rightarrow \psi_2) := \sup_{m \geq 1} \frac{f(m)}{m}. \quad (69)$$

Intuitively, for sufficiently large m , one can transform m copies of ψ_1 exactly into $mD(\psi_1 \rightarrow \psi_2)$ copies of ψ_2 by LOCC. We have the following proposition, which is a consequence of Proposition 16 and the developments in [DFJY05]:

Proposition 17 *Let $|\psi\rangle_{AB}$ be a pure state in AB and Φ^2 be the ebit. Then the following equality holds*

$$D(\psi \rightarrow \Phi^2) = E_{\min}^u(\psi_{AB}). \quad (70)$$

C. Unextendible fidelity

Let $\rho, \sigma \in \mathcal{S}(A)$ be two quantum states. The (root) fidelity between ρ and σ is defined as [Uhl76]

$$F(\rho, \sigma) := \|\sqrt{\rho}\sqrt{\sigma}\|_1 = \text{Tr} \left[\sqrt{\sqrt{\sigma}\rho\sqrt{\sigma}} \right]. \quad (71)$$

Here we define the unextendible fidelity of a state ρ_{AB} :

Definition 6 (Unextendible fidelity) *For a given bipartite state ρ_{AB} , the unextendible fidelity is defined as*

$$F^u(\rho_{AB}) := \sup_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} F(\rho_{AB}, \sigma_{AB'}). \quad (72)$$

Note that the supremum in (72) can be replaced with a maximum.

Suppose that Ψ_{ABC} is a purification of ρ_{AB} . By applying Remark 3, we see that the unextendible fidelity can be alternatively understood as a measure of how well one can recover the state ρ_{AB} if system B is lost and a recovery channel is performed on the purification system C alone, due to the following equivalent formulation:

$$F^u(\rho_{AB}) = \max_{\mathcal{R}_{C \rightarrow B'}} \{F(\rho_{AB}, \rho_{AB'}) : \rho_{ABB'} = (\text{id}_{AB} \otimes \mathcal{R}_{C \rightarrow B'}) (|\Psi\rangle\langle\Psi|_{ABC})\}, \quad (73)$$

where the maximum ranges over every quantum channel $\mathcal{N}_{C \rightarrow B'}$. The unextendible fidelity is thus similar in spirit to the fidelity of recovery from [SW15], but one finds that it is a different measure when analyzing it in more detail.

By examining (4) and (71), one immediately finds that $\tilde{D}_{1/2}(\rho\|\sigma) = -\log[F(\rho, \sigma)]^2$. Thus, we establish the following equivalence between unextendible fidelity and 1/2-sandwiched unextendible entanglement:

$$F^u(\rho_{AB}) := \max_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} F(\rho_{AB}, \sigma_{AB'}) \quad (74)$$

$$= \max_{\sigma_{AB'} \in \mathcal{F}_{\rho_{AB}}} 2^{-\frac{1}{2}\tilde{D}_{1/2}(\rho_{AB}\|\sigma_{AB'})} \quad (75)$$

$$= 2^{-\tilde{E}_{1/2}^u(\rho_{AB})}. \quad (76)$$

Since the fidelity function is SDP computable [Wat13], it follows that the unextendible fidelity can be computed efficiently by means of a semidefinite program. To be more specific, the following two optimization programs satisfy strong duality and both evaluate to $F^u(\rho_{AB})$. For completeness, we show in detail how to derive the dual program in Appendix I.

Primal Program	Dual Program
maximize $\frac{1}{2} \text{Tr}[X_{AB}] + \frac{1}{2} \text{Tr}[X_{AB}^\dagger]$	minimize $\frac{1}{2} \text{Tr}[W_{AB}\rho_{AB}] + \frac{1}{2} \text{Tr}[Z_{AB}\rho_{AB}]$
subject to $\begin{pmatrix} \rho_{AB} & X_{AB} \\ X_{AB}^\dagger & \text{Tr}_B[\sigma_{ABB'}] \end{pmatrix} \geq 0$ (77)	subject to $\begin{pmatrix} W_{AB} & -\mathbb{1}_{AB} \\ -\mathbb{1}_{AB} & Y_{AB'} \end{pmatrix} \geq 0$
$\rho_{AB} = \text{Tr}_{B'}[\sigma_{ABB'}]$	$Z_{AB} \otimes \mathbb{1}_{B'} \geq Y_{AB'} \otimes \mathbb{1}_B$
$\sigma_{ABB'} \geq 0$	$W_{AB}, Y_{AB'}, Z_{AB} \geq 0$
$X_{AB} \in \mathcal{L}(AB)$	

(78)

We also establish the following equivalent dual representation of $F^u(\rho_{AB})$ in Appendix I:

$$\begin{aligned} & \text{infimum} \quad \sqrt{\text{Tr}[Y_{AB'}^{-1}\rho_{AB}] \text{Tr}[Z_{AB}\rho_{AB}]} \\ & \text{subject to} \quad Z_{AB} \otimes \mathbb{1}_{B'} \geq Y_{AB'} \otimes \mathbb{1}_B \\ & \quad \quad \quad Z_{AB}, Y_{AB'} > 0. \end{aligned} \quad (79)$$

As a direct consequence of this equivalent dual representation, we find that the extendible fidelity is multiplicative (proof in Appendix J).

Proposition 18 (Multiplicativity) *Let $\rho_{A_1B_1} \in \mathcal{S}(A_1B_1)$ and $\rho_{A_2B_2} \in \mathcal{S}(A_2B_2)$ be two bipartite states. The following equality holds*

$$F^u(\rho_{A_1B_1} \otimes \rho_{A_2B_2}) = F^u(\rho_{A_1B_1})F^u(\rho_{A_2B_2}), \quad (80)$$

where the entanglement is evaluated across the $A:B$ bipartition.

Recall (74)–(76), which relates the unextendible fidelity and the 1/2-sandwiched unextendible entanglement. Proposition 18 demonstrates that the logarithmic unextendible fidelity is an additive unextendible entanglement measure, different from both the min- and max-unextendible entanglement.

Corollary 19 *Let $\rho_{A_1B_1} \in \mathcal{S}(A_1B_1)$ and $\rho_{A_2B_2} \in \mathcal{S}(A_2B_2)$ be two bipartite states. The following additivity relation holds*

$$\tilde{E}_{1/2}^u(\rho_{A_1B_1} \otimes \rho_{A_2B_2}) = \tilde{E}_{1/2}^u(\rho_{A_1B_1}) + \tilde{E}_{1/2}^u(\rho_{A_2B_2}), \quad (81)$$

where the entanglement is evaluated across the $A:B$ bipartition.

Using the relation in (74)–(76) between the unextendible fidelity and the 1/2-sandwiched unextendible entanglement, we find that the unextendible fidelity of a pure bipartite state can be computed explicitly.

Proposition 20 *Let ψ_{AB} be a pure bipartite state. Then the following equality holds*

$$F^u(\psi_{AB}) = \lambda_{\max}(\psi_A), \quad (82)$$

where $\lambda_{\max}(\rho_A)$ is the maximal eigenvalue of ψ_A .

V. APPLICATIONS FOR SECRET KEY AND ENTANGLEMENT DISTILLATION

A. Private states and unextendible entanglement

In this section, we review the definition of a private state [HHHO05, HHHO09], and then we establish a bound on the number of private bits contained in a private state, in terms of the state's unextendible entanglement. These results find applications in the next two subsections (Sections VB and VC), where we investigate the overhead of probabilistic secret key distillation and the rate of exact secret key distillation.

We first review the definition of a private state [HHHO05, HHHO09]. Let $\rho_{ABA'B'}$ be a state shared between spatially separated parties Alice and Bob, where Alice possesses systems A and A' and Bob possesses systems B and B' , such that

$$K \equiv \dim(\mathcal{H}_A) = \dim(\mathcal{H}_B). \quad (83)$$

A state $\rho_{ABA'B'}$ is called a private state [HHHO05, HHHO09] if Alice and Bob can extract a secret key from it by performing local measurements on A and B , such that the key is product with an arbitrary purifying system of $\rho_{ABA'B'}$. That is, $\rho_{ABA'B'}$ is a private state of $\log K$ private bits if, for every purification $\varphi_{ABA'B'E}^\rho$ of $\rho_{ABA'B'}$, the following holds:

$$(\mathcal{M}_A \otimes \mathcal{M}_B \otimes \text{Tr}_{A'B'}) (\varphi_{ABA'B'E}^\rho) = \frac{1}{K} \sum_{k=1}^K |k\rangle\langle k|_A \otimes |k\rangle\langle k|_B \otimes \sigma_E, \quad (84)$$

where $\mathcal{M}(\cdot) = \sum_k |k\rangle\langle k|(\cdot)|k\rangle\langle k|$ is a projective measurement channel and σ_E is some state on the purifying system. The systems A and B are known as key systems, and A' and B' are known as shield systems. Interestingly, it was shown that a private state of $\log K$ private bits can be written in the following form [HHHO05, HHHO09]

$$\gamma_{ABA'B'} := U_{ABA'B'} (\Phi_{AB}^K \otimes \sigma_{A'B'}) U_{ABA'B'}^\dagger, \quad (85)$$

where Φ_{AB}^K is a maximally entangled state of Schmidt rank K , the state $\sigma_{A'B'}$ is an arbitrary state, and

$$U_{ABA'B'} := \sum_{i,j=1}^K |i\rangle\langle i|_A \otimes |j\rangle\langle j|_B \otimes U_{A'B'}^{ij} \quad (86)$$

is a controlled unitary known as a “twisting unitary,” with each $U_{A'B'}^{ij}$ a unitary.

We now establish some bounds on the number of private bits contained in a private state, in terms of its unextendible entanglement. Recall that the generalized mutual information of a state ρ_{AB} is defined as follows [SW12]:

$$I_{\mathbf{D}}(A; B)_\rho := \inf_{\sigma_B} \mathbf{D}(\rho_{AB} \| \rho_A \otimes \sigma_B), \quad (87)$$

where \mathbf{D} is the generalized divergence discussed in Section IIIB and the infimum is with respect to every density operator σ_B . We first show that the unextendible entanglement of a private state is bounded from below by the generalized mutual information of Φ_{AB} (proof in Appendix K).

Proposition 21 *For a γ -bipartite private state of the form in (85), the following bound holds*

$$\mathbf{E}^u(\gamma_{AA'BB'}) \geq \frac{1}{2} I_{\mathbf{D}}(A; B)_{\Phi_K}, \quad (88)$$

where $I_{\mathbf{D}}(A; B)_\Phi$ is evaluated with respect to the state Φ_{AB}^K in (85).

As a corollary of Proposition 21 and Lemmas 7, 8, and 9, we find that if the generalized divergence is set to be the Petz-, sandwiched, or geometric Rényi relative entropy, then the unextendible entanglement of a γ -bipartite private state is bounded from below by the amount of secret key that can be extracted from the state. Note that this corollary includes the relative entropy, the min-relative entropy, and the max-relative entropy as limiting cases.

Corollary 22 *If the generalized divergence is the Petz-, sandwiched, or geometric Rényi relative entropy with α set so that the data-processing inequality is satisfied, then the following bound holds*

$$\mathbf{E}_\alpha^u(\gamma_{AA'BB'}) \geq \log K. \quad (89)$$

B. Overhead of probabilistic secret key distillation

This section considers the overhead of probabilistic secret key distillation under selective two-extendible operations (which include 1-LOCC operations as a special case). We first formally define the overhead of probabilistic secret key distillation:

Definition 7 *The overhead of distilling k private bits from several independent copies of a bipartite state ρ_{AB} is given by the minimum number of copies of ρ_{AB} needed, on average, to produce some private state $\gamma_{ABA'B'}^k$ with $\log K = k$ using 1-LOCC operations:*

$$K_{\text{ov}}(\rho_{AB}, k) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow \gamma_{ABA'B'}^k \text{ with prob. } p, \Lambda \text{ is selective 1-LOCC} \right\}. \quad (90)$$

We can also define the overhead when selective two-extendible operations are allowed:

$$K_{\text{ov},2}(\rho_{AB}, k) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow \gamma_{ABA'B'}^k \text{ with prob. } p, \Lambda \text{ is selective two-extendible} \right\}. \quad (91)$$

Note that it is not necessary to produce a particular private state $\gamma_{ABA'B'}^k$, but rather just some private state $\gamma_{ABA'B'}^k$ having k private bits.

The following inequality is a trivial consequence of definitions and the fact that a selective 1-LOCC operation is a special kind of selective two-extendible operation:

$$K_{\text{ov}}(\rho_{AB}, k) \geq K_{\text{ov},2}(\rho_{AB}, k). \quad (92)$$

It turns out that the relative-entropy-induced unextendible entanglement $E^u(\rho_{AB})$, given in Definition 3, provides a lower bound on the overhead of probabilistic secret key distillation of a bipartite state ρ_{AB} (proof in Appendix L).

Theorem 23 *For a bipartite state ρ_{AB} , the overhead of distilling k private bits from ρ_{AB} is bounded from below by*

$$K_{\text{ov},2}(\rho_{AB}, k) \geq \frac{k}{E^u(\rho_{AB})}, \quad (93)$$

where the relative-entropy-induced unextendible entanglement is given in Definition 3.

In Theorem 5 and Proposition 13, we proved that the α -Petz unextendible entanglement for $\alpha \in [1, 2]$, the α -sandwiched unextendible entanglement for $\alpha \in [1, \infty]$, and the α -geometric unextendible entanglement for $\alpha \in [1, 2]$ satisfy selective two-extendible monotonicity and subadditivity, respectively. That is to say, all these entanglement measures can be used as lower bounds in (93). Since these divergences are monotonically increasing in α [Tom15] and $D(\omega\|\tau) \leq \hat{D}(\omega\|\tau)$ [HP91], $E^u(\rho_{AB})$ is the smallest unextendible entanglement measure among these choices and yields the tightest lower bound. Furthermore, the authors of [FSP19] proposed a method to accurately approximate the quantum relative entropy via semidefinite programming. This enables us to estimate the lower bound using available semidefinite programming solvers. See also [FF18] in this context.

C. Exact secret key distillation

We can also consider the setting in which the goal is to distill secret key exactly from a bipartite state by using two-extendible or 1-LOCC operations. Though exact distillation is less realistic than the above probabilistic scenario, it is still a core part of zero-error quantum information theory [GdAM16].

The one-shot 1-LOCC exact distillable key of ρ_{AB} is defined to be the maximum number of private bits achievable via a 1-LOCC channel; that is,

$$K_{1\text{-LOCC}}^{(1)}(\rho_{AB}) := \sup\{k : \Lambda_{AB \rightarrow \hat{A}\hat{B}A'B'}(\rho_{AB}) = \gamma_{\hat{A}\hat{B}A'B'}^k, \Lambda \in 1\text{-LOCC}\}, \quad (94)$$

where $\gamma_{\hat{A}\hat{B}A'B'}^k$ is any private state with k private bits. The 1-LOCC exact distillable key of a state ρ_{AB} is then defined as the regularization of $K_{1\text{-LOCC}}^{(1)}(\rho_{AB})$:

$$K_{1\text{-LOCC}}(\rho_{AB}) := \liminf_{n \rightarrow \infty} \frac{1}{n} K_{1\text{-LOCC}}^{(1)}(\rho_{AB}^{\otimes n}). \quad (95)$$

Note that the following inequality holds as a direct consequence of definitions:

$$K_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq K_{1\text{-LOCC}}(\rho_{AB}). \quad (96)$$

The one-shot two-extendible exact distillable key and two-extendible exact distillable key of ρ_{AB} are defined similarly:

$$K_{2\text{-EXT}}^{(1)}(\rho_{AB}) := \sup\{k : \Lambda_{AB \rightarrow \hat{A}\hat{B}A'B'}(\rho_{AB}) = \gamma_{\hat{A}\hat{B}A'B'}^k, \Lambda \in 2\text{-EXT}\}, \quad (97)$$

$$K_{2\text{-EXT}}(\rho_{AB}) := \liminf_{n \rightarrow \infty} \frac{1}{n} K_{2\text{-EXT}}^{(1)}(\rho_{AB}^{\otimes n}), \quad (98)$$

and the following inequality holds as a direct consequence of definitions:

$$K_{2\text{-EXT}}^{(1)}(\rho_{AB}) \leq K_{2\text{-EXT}}(\rho_{AB}). \quad (99)$$

Immediate consequences of definitions and the fact that 1-LOCC operations are contained in the set of two-extendible operations are the following inequalities:

$$K_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq K_{2\text{-EXT}}^{(1)}(\rho_{AB}), \quad K_{1\text{-LOCC}}(\rho_{AB}) \leq K_{2\text{-EXT}}(\rho_{AB}), \quad (100)$$

It turns out that the min-unextendible entanglement $E_{\min}^u(\rho_{AB})$ serves as an upper bound on the two-extendible exact distillable key of a bipartite state ρ_{AB} (proof in Appendix M).

Theorem 24 *For a bipartite state ρ_{AB} , its asymptotic exact distillable key under two-extendible operations is bounded from above as*

$$K_{2\text{-EXT}}(\rho_{AB}) \leq E_{\min}^u(\rho_{AB}). \quad (101)$$

Actually, it is evident from the proof that in Theorem 24, both $E_{\max}^u(\rho_{AB})$ and $\tilde{E}_{1/2}^u(\rho_{AB})$ are valid upper bounds on $K_{2\text{-EXT}}(\rho_{AB})$, since they are also monotonic and additive. However, since

$$E_{\min}(\rho_{AB}) \leq \tilde{E}_{1/2}^u(\rho_{AB}) \leq E_{\max}^u(\rho_{AB}), \quad (102)$$

the min-unextendible entanglement $E_{\min}^u(\rho_{AB})$ leads to the tightest upper bound. Note that the inequalities in (102) follow from the facts that the fidelity obeys data processing under the channel $\mathcal{M}(\omega) := \text{Tr}[\Pi^\rho \omega] |0\rangle\langle 0| + \text{Tr}[(1 - \Pi^\rho) \omega] |1\rangle\langle 1|$, so that $\text{Tr}[\Pi^\rho \sigma] = F^2(\mathcal{M}(\rho), \mathcal{M}(\sigma)) \geq F^2(\rho, \sigma)$ and \tilde{D}_α is monotonically increasing in α [MLDS⁺13, WWY14].

D. Overhead of probabilistic entanglement distillation

Entanglement distillation aims at obtaining maximally entangled states from less entangled bipartite states shared between two parties via certain free operations. As a central task in quantum information processing, various approaches [VP98, Rai99, VW02, Rai01, HHH00, CW04, WD17b, LDS17, WD16, FWTD19, KDWW19, RFWG19] have been developed to characterize and approximate the performance of the rates of deterministic entanglement distillation.

Here, we consider the overhead of probabilistic entanglement distillation [BDSW96, CCGFZ99, PSBZ01, NFB14, CB08, RST⁺18] under selective two-extendible operations, similar to what we considered for probabilistic secret key distillation. We begin by defining the overhead of probabilistic entanglement distillation.

Definition 8 *The overhead of distilling m ebits from several independent copies of a bipartite state ρ_{AB} is equal to the minimum number of copies of ρ_{AB} needed, on average, to produce m copies of the ebit Φ^2 using selective 1-LOCC operations:*

$$E_{\text{ov}}(\rho_{AB}, m) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow (\Phi^2)^{\otimes m} \text{ with prob. } p, \Lambda \text{ is selective 1-LOCC} \right\}. \quad (103)$$

We can also define the overhead when selective two-extendible operations are allowed:

$$E_{\text{ov},2}(\rho_{AB}, m) := \inf \left\{ \frac{n}{p} : \Lambda(\rho_{AB}^{\otimes n}) \rightarrow (\Phi^2)^{\otimes m} \text{ with prob. } p, \Lambda \text{ is selective two-extendible} \right\}. \quad (104)$$

The following inequality is a trivial consequence of definitions and the fact that a selective 1-LOCC operation is a special kind of selective two-extendible operation:

$$E_{\text{ov}}(\rho_{AB}, m) \geq E_{\text{ov},2}(\rho_{AB}, m). \quad (105)$$

Since secret key can be obtained from ebits [HHHO05] via local operations, a direct corollary of Theorem 23 is that E^u is a lower bound on the overhead of probabilistic entanglement distillation.

Corollary 25 *For a bipartite state ρ_{AB} , the overhead of distilling m ebits from ρ_{AB} is bounded from below as*

$$E_{\text{ov},2}(\rho_{AB}, m) \geq \frac{m}{E^u(\rho_{AB})}, \quad (106)$$

where $E^u(\rho_{AB})$ is the relative-entropy-induced unextendible entanglement from Definition 3.

E. Exact entanglement distillation

The one-shot 1-LOCC exact distillable entanglement of ρ_{AB} is defined to be the maximum number of ebits achievable via a 1-LOCC channel; that is,

$$E_{1\text{-LOCC}}^{(1)}(\rho_{AB}) := \sup \{ \log d : \Lambda_{AB \rightarrow \hat{A}\hat{B}}(\rho_{AB}) = \Phi_{\hat{A}\hat{B}}^d, \Lambda \in 1\text{-LOCC} \}, \quad (107)$$

where $\Phi_{\hat{A}\hat{B}}^d$ is a maximally entangled state of Schmidt rank d . The 1-LOCC exact distillable entanglement of a state ρ_{AB} is then defined as the regularization of $E_{1\text{-LOCC}}^{(1)}(\rho_{AB})$:

$$E_{1\text{-LOCC}}(\rho_{AB}) := \liminf_{n \rightarrow \infty} \frac{1}{n} E_{1\text{-LOCC}}^{(1)}(\rho_{AB}^{\otimes n}). \quad (108)$$

Note that the following inequality is a direct consequence of the definitions:

$$E_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq E_{1\text{-LOCC}}(\rho_{AB}). \quad (109)$$

The one-shot two-extendible exact distillable entanglement and two-extendible exact distillable entanglement of ρ_{AB} are defined similarly:

$$E_{2\text{-EXT}}^{(1)}(\rho_{AB}) := \sup\{\log d : \Lambda_{AB \rightarrow \hat{A}\hat{B}}(\rho_{AB}) = \Phi_{\hat{A}\hat{B}}^d, \Lambda \in 2\text{-EXT}\}, \quad (110)$$

$$E_{2\text{-EXT}}(\rho_{AB}) := \liminf_{n \rightarrow \infty} \frac{1}{n} E_{2\text{-EXT}}^{(1)}(\rho_{AB}^{\otimes n}), \quad (111)$$

and the following inequality is a direct consequence of definitions:

$$E_{2\text{-EXT}}^{(1)}(\rho_{AB}) \leq E_{2\text{-EXT}}(\rho_{AB}). \quad (112)$$

Immediate consequences of definitions and the fact that 1-LOCC operations are contained in the set of two-extendible operations are the following inequalities:

$$E_{1\text{-LOCC}}^{(1)}(\rho_{AB}) \leq E_{2\text{-EXT}}^{(1)}(\rho_{AB}), \quad E_{1\text{-LOCC}}(\rho_{AB}) \leq E_{2\text{-EXT}}(\rho_{AB}), \quad (113)$$

As a direct corollary of Theorem 24 and the relation between private states and ebits mentioned previously (just below (105)), the min-unextendible entanglement E_{\min}^u serves as an upper bound on the exact distillable entanglement via two-extendible operations.

Corollary 26 *For a bipartite state ρ_{AB} , its asymptotic exact distillable entanglement under two-extendible operations is bounded from above by its min-unextendible entanglement:*

$$E_{2\text{-EXT}}(\rho_{AB}) \leq E_{\min}^u(\rho_{AB}). \quad (114)$$

F. Examples

In this section, we apply our bounds on the overhead of probabilistic entanglement or secret key distillation to three classes of states: isotropic states, S states, and erased states. We compare our lower bounds and other known estimations of the overhead to upper bounds derived from known distillation protocols. In particular, we show that our lower bound on the overhead of distillation is tight for erased states.

To begin with, recall the relative entropy of entanglement [VP98]:

$$E_R(\rho_{AB}) := \min_{\sigma_{AB} \in \text{SEP}} D(\rho_{AB} \| \sigma_{AB}), \quad (115)$$

where SEP denotes the set of separable states. It is known that the relative entropy of entanglement is monotone under selective LOCC [VP98]. Thus, the relative entropy of entanglement $E_R(\rho_{AB})$ can be used to estimate the overhead of distillation under selective LOCC via an approach similar to that given in the proof of Corollary 25.

Isotropic states: Let us first investigate the class of isotropic states ρ_r , defined as [HH99]

$$\rho_r := r \cdot \Phi^d + (1 - r) \frac{\mathbb{1} - \Phi^d}{d^2 - 1}, \quad (116)$$

where $r \in [0, 1]$, so that ρ_r is a convex mixture of a maximally entangled state Φ^d of Schmidt rank d and its orthogonal complement. Numerous works have been carried out to study the




FIG. 1. Lower bounds on the overhead of distilling one ebit or one private bit from isotropic states. For the overhead of probabilistic distillation of entanglement or secret key under 1-LOCC operations, our lower bound (solid line) outperforms the previous lower bound (dashed line) based on relative entropy of entanglement.

distillation rate of isotropic states under various scenarios [FWTD19, WFD18, LDS17, RST⁺18]. Here we investigate probabilistic entanglement distillation under 1-LOCC operations and show the advantage of our method in estimating the overhead. We consider the case $d = 2$ for simplicity. Figure 1 plots the unextendible entanglement and relative entropy of entanglement for the overhead of probabilistic entanglement or secret key distillation for this set of states.¹ Note that the relative entropy of entanglement for qubit-qubit isotropic states ($d = 2$) was calculated in [VPRK97].

S states: An S state is a mixture of the Bell state and non-orthogonal product noise, whose distillation protocols were studied in [RST⁺18, ZZW⁺21]. Here, we define it as

$$S_p = p\Phi^2 + (1 - p)|00\rangle\langle 00|. \quad (117)$$

Figure 2 plots the unextendible entanglement and relative entropy of entanglement for the overhead of probabilistic entanglement or secret key distillation for this set of states. The result shows that our result establishes a better lower bound on the overhead of probabilistic distillation under 1-LOCC operations.

Erased states: We also consider the class of erased states, which are the Choi states of quantum erasure channels [GBP97]:

$$\rho_{A'B}^\varepsilon = (1 - \varepsilon)\Phi_{AB}^2 + \varepsilon|e\rangle\langle e|_{A'} \otimes \pi_B, \quad (118)$$

where $\varepsilon \in [0, 1]$, Φ_{AB}^2 is an ebit, and $|e\rangle$ is some state that is orthogonal to π_A . For simplicity, we choose $d_A = d_B = 2$ (qubit system), and $d_{A'} = d_A + 1 = 3$. This state can be obtained as follows. Alice and Bob share a two-qubit maximally entangled state Φ_{AB} and Alice transmits her local copy through an erasure channel, so that with probability $1 - \varepsilon$ it is unaffected and with probability ε it is “erased” (replaced with the erasure flag state e).

Interestingly, there is a simple formula for the relative-entropy-induced unextendible entanglement of the erased state (proof in Appendix N).

Proposition 27 *The unextendible entanglement of the erased state $\rho_{A'B}^\varepsilon$ is $E^u(\rho_{A'B}^\varepsilon) = 1 - \varepsilon$.*

¹ All Matlab codes used to generate these plots and calculate various unextendible entanglement measures are available with the arXiv posting of this paper as arXiv ancillary files.




FIG. 2. Bounds on the overhead of distilling one ebit or one private bit from S states. For the overhead of probabilistic distillation of entanglement or secret key under 1-LOCC operations, our lower bound (solid line) outperforms the previous lower bound (dashed line) based on relative entropy of entanglement.

Corollary 25 together with Proposition 27 shows that $1/E^u(\rho_{A'B}^\varepsilon) = 1/(1 - \varepsilon)$ is a lower bound on the overhead of probabilistically distilling one ebit from $\rho_{A'B}^\varepsilon$. Indeed, we can show the tightness of the lower bound by considering the following one-way LOCC protocol that achieves this bound. Given $\rho_{A'B}^\varepsilon$, Alice detects whether her local system is erased or not by performing a binary measurement $\{\mathbb{1}_A, |e\rangle\langle e|\}$. With probability $1 - \varepsilon$, she finds that the system is not erased. Then she sends this information to Bob, and the protocol finishes with one ebit shared between them. As such, we conclude the following.

Theorem 28 *For the erased state $\rho_{A'B}^\varepsilon$, we have*

$$E_{\text{ov}}(\rho_{A'B}^\varepsilon, 1) = E_{\text{ov},2}(\rho_{A'B}^\varepsilon, 1) = \frac{1}{E^u(\rho_{A'B}^\varepsilon)} = \frac{1}{1 - \varepsilon}. \quad (119)$$

Therefore, our estimation on the overhead of distilling ebits from erased states under selective one-way LOCC is optimal in the sense that the upper bound on the overhead of the above protocol matches our lower bound from Corollary 25, thus characterizing the ability of probabilistic distillation for erased states. Note that this result can be generalized to multiple copies of the erased state.

We further consider the overhead of distillation under LOCC operations for the erased states in the following proposition (proof in Appendix O), indicating an interesting fact that the optimal protocol for distilling one ebit from erased states under one-way LOCC operations matches the lower bound $1/E_R(\rho_{A'B}^\varepsilon) \equiv 1/(1 - \varepsilon)$ on the overhead under LOCC operations. It reveals that one-way LOCC operations have the same power and performance as LOCC operations in probabilistically distilling ebits from erased states.

Proposition 29 *The relative entropy of entanglement of the erased state $\rho_{A'B}^\varepsilon$ is $E_R(\rho_{A'B}^\varepsilon) = 1 - \varepsilon$, which implies that the overhead of distillation under selective LOCC for $\rho_{A'B}^\varepsilon$ is given by $(1 - \varepsilon)^{-1}$.*

VI. CONCLUDING REMARKS

Our work introduces a family of entanglement measures called unextendible entanglement to quantify and investigate the unextendibility of entanglement. The crucial technical contribution was intuitively motivated by the fact that the more entangled a bipartite state is, the less each of its constituent systems can be entangled with a third party. A key distinction to previous works is that our proposed measures restrict the free set to be state dependent. These entanglement measures have desirable properties, including monotonicity under selective two-extendible operations, faithfulness, and normalization. The unextendible entanglement in our work has also found direct operational applications in evaluating the overhead and rate of entanglement or secret key distillation. As a notable application example, we characterized the optimal overhead of distilling one ebit from the erased states under one-way LOCC operations. We also note that the probabilistic state conversion task, which has already been studied for the pure-state case [Vid99, AOP16], can be further explored using the results in this paper.

An important problem for future work is to investigate to what extent the bounds on the asymptotic distillable secret key or entanglement can be approached. One potential approach is to explore the connections to distinguishability distillation (i.e., hypothesis testing) [Nag06, Hay07, ANSV08, MO15, WW19]. It is also of interest to consider an extension to the resource theory of k -unextendibility and entanglement dilution [BDSW96, HHT01, VDC02, WD17a, BD11, WW23], where the techniques applied in [XFWD17] might be useful here. Moreover, we have only considered the extendibility on system B when defining generalized unextendible entanglement measure, yielding an asymmetric entanglement measure. It would be meaningful to further explore the extendibility on system A and promote our measure to a symmetric one. Another interesting direction is to develop the continuous-variable setting, where the results of [LKAW19] could be helpful, as well as the dynamical setting of quantum channels.

ACKNOWLEDGEMENT

The authors thank Anurag Anshu, Kun Fang, Debbie Leung, Vishal Singh, and Dave Touchette for discussions. Part of this work was done when KW was at the Southern University of Science and Technology and XW was at the University of Maryland. XW was partially supported by the Start-up Fund from The Hong Kong University of Science and Technology (Guangzhou), the Guangdong Quantum Science and Technology Strategic Fund (Grant No. GDZX2303007), and the Education Bureau of Guangzhou Municipality. MMW acknowledges support from NSF under grant no. 2315398.

-
- [AI07] Gerardo Adesso and Fabrizio Illuminati. Strong monogamy of bipartite and genuine multipartite entanglement: the Gaussian case. *Physical Review Letters*, 99(15):150501, 2007.
- [ANSV08] Koenraad M. R. Audenaert, Michael Nussbaum, Arleta Szkoła, and Frank Verstraete. Asymptotic error rates in quantum hypothesis testing. *Communications in Mathematical Physics*, 279(1):251–283, 2008.
- [AOP16] Álvaro M Alhambra, Jonathan Oppenheim, and Christopher Perry. Fluctuating states: What is the probability of a thermodynamical transition? *Physical Review X*, 6(4):041016, 2016.
- [Ara90] Huzihiro Araki. On an inequality of Lieb and Thirring. *Letters in Mathematical Physics*, 19:167–170, 1990.

- [BBFS22] Mario Berta, Francesco Borderi, Omar Fawzi, and Volkher Scholz. Semidefinite programming hierarchies for constrained bilinear optimization. *Mathematical Programming*, 194:781–829, July 2022. arXiv:1810.12197.
- [BD11] Francesco Buscemi and Nilanjana Datta. Entanglement cost in practical scenarios. *Physical Review Letters*, 106(13):130503, 2011.
- [BDSW96] Charles H Bennett, David P DiVincenzo, John A Smolin, and William K Wootters. Mixed-state entanglement and quantum error correction. *Physical Review A*, 54(5):3824, 1996.
- [Bei13] Salman Beigi. Sandwiched Rényi divergence satisfies data processing inequality. *Journal of Mathematical Physics*, 54(12):122202, 2013.
- [Bra08] Fernando G. S. L. Brandão. *Entanglement theory and the quantum simulation of many-body physics*. PhD thesis, Imperial College, 2008.
- [BS82] Viacheslav P. Belavkin and P. Staszewski. C^* -algebraic generalization of relative entropy and entropy. *Annales de l’IHP Physique théorique*, 37(1):51–58, 1982.
- [BT16] Mario Berta and Marco Tomamichel. The fidelity of recovery is multiplicative. *IEEE Transactions on Information Theory*, 62(4):1758–1763, 2016.
- [CB08] Earl T. Campbell and Simon C. Benjamin. Measurement-based entanglement under conditions of extreme photon loss. *Physical Review Letters*, 101(13):130502, 2008.
- [CCGFZ99] Carlos Cabrillo, J. Ignacio Cirac, Pablo Garcia-Fernandez, and Peter Zoller. Creation of entangled states of distant atoms by interference. *Physical Review A*, 59(2):1025, 1999.
- [CG19] Eric Chitambar and Gilad Gour. Quantum resource theories. *Reviews of Modern Physics*, 91(2):025001, 2019.
- [CKW00] Valerie Coffman, Joydip Kundu, and William K. Wootters. Distributed entanglement. *Physical Review A*, 61(5):052306, 2000.
- [CW04] Matthias Christandl and Andreas Winter. “Squashed entanglement”: an additive entanglement measure. *Journal of Mathematical Physics*, 45(3):829–840, 2004.
- [Dat09] Nilanjana Datta. Min- and max-relative entropies and a new entanglement monotone. *IEEE Transactions on Information Theory*, 55(6):2816–2826, 2009.
- [DFJY05] Runyao Duan, Yuan Feng, Zhengfeng Ji, and Mingsheng Ying. Efficiency of deterministic entanglement transformation. *Physical Review A*, 71(2):022305, 2005.
- [DL14] Nilanjana Datta and Felix Leditzky. A limit of the quantum Rényi divergence. *Journal of Physics A: Mathematical and Theoretical*, 47(4):045304, 2014.
- [Doh14] Andrew C. Doherty. Entanglement and the shareability of quantum states. *Journal of Physics A: Mathematical and Theoretical*, 47(42):424004, 2014.
- [DPS02] Andrew C. Doherty, Pablo A. Parrilo, and Federico M. Spedalieri. Distinguishing separable and entangled states. *Physical Review Letters*, 88(18):187904, 2002.
- [DPS04] Andrew C. Doherty, Pablo A. Parrilo, and Federico M. Spedalieri. Complete family of separability criteria. *Physical Review A*, 69(2):022308, 2004.
- [FF18] Hamza Fawzi and Omar Fawzi. Efficient optimization of the quantum relative entropy. *Journal of Physics A: Mathematical and Theoretical*, 51(15):154003, March 2018.
- [FF21] Kun Fang and Hamza Fawzi. Geometric Rényi divergence and its applications in quantum channel capacities. *Communications in Mathematical Physics*, 384:1615–1677, 2021. arXiv:1909.05758.
- [FL13] Rupert L. Frank and Elliott H. Lieb. Monotonicity of a relative Rényi entropy. *Journal of Mathematical Physics*, 54(12):122201, 2013.
- [FSP19] Hamza Fawzi, James Saunderson, and Pablo A. Parrilo. Semidefinite approximations of the matrix logarithm. *Foundations of Computational Mathematics*, 19(2):259–296, 2019.
- [FWTD19] Kun Fang, Xin Wang, Marco Tomamichel, and Runyao Duan. Non-asymptotic entanglement distillation. *IEEE Transactions on Information Theory*, 65(10):6454–6465, 2019.
- [GBP97] Markus Grassl, Thomas Beth, and Thomas Pellizzari. Codes for the quantum erasure channel. *Physical Review A*, 56(1):33, 1997.
- [GdAM16] Elloa B. Guedes, Francisco Marcos de Assis, and Rex Antonio da Costa Medeiros. *Quantum Zero-Error Information Theory*. Springer, 2016.
- [GG18] Gilad Gour and Yu Guo. Monogamy of entanglement without inequalities. *Quantum*, 2:81, 2018.
- [GW15] Manish K. Gupta and Mark M. Wilde. Multiplicativity of completely bounded p -norms implies a strong converse for entanglement-assisted capacity. *Communications in Mathematical Physics*, 334(2):867–887, 2015.

- [Hay07] Masahito Hayashi. Error exponent in asymmetric quantum hypothesis testing and its application to classical-quantum channel coding. *Physical Review A*, 76(6):062301, 2007.
- [Hay16] Masahito Hayashi. *Quantum Information Theory: Mathematical Foundation*. Springer, 2016.
- [HH99] Michał Horodecki and Paweł Horodecki. Reduction criterion of separability and limits for a class of distillation protocols. *Physical Review A*, 59(6):4206, 1999.
- [HHH00] Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki. Asymptotic manipulations of entanglement can exhibit genuine irreversibility. *Physical Review Letters*, 84(19):4260, 2000.
- [HHHH09] Ryszard Horodecki, Paweł Horodecki, Michał Horodecki, and Karol Horodecki. Quantum entanglement. *Reviews of Modern Physics*, 81(2):865, 2009.
- [HHHO05] Karol Horodecki, Michał Horodecki, Paweł Horodecki, and Jonathan Oppenheim. Secure key from bound entanglement. *Physical Review Letters*, 94(16):160502, 2005.
- [HHHO09] Karol Horodecki, Michał Horodecki, Paweł Horodecki, and Jonathan Oppenheim. General paradigm for distilling classical key from quantum states. *IEEE Transactions on Information Theory*, 55(4):1898–1929, 2009.
- [HHT01] Patrick M. Hayden, Michał Horodecki, and Barbara M. Terhal. The asymptotic entanglement cost of preparing a quantum state. *Journal of Physics A: Mathematical and General*, 34(35):6891, 2001.
- [HLA22] Chung-Yun Hsieh, Matteo Lostaglio, and Antonio Acín. Quantum channel marginal problem. *Physical Review Research*, 4(1):013249, 2022.
- [HM17] Fumio Hiai and Milán Mosonyi. Different quantum f -divergences and the reversibility of quantum operations. *Reviews in Mathematical Physics*, 29(07):1750023, 2017.
- [HMW13] Patrick Hayden, Kevin Milner, and Mark M. Wilde. Two-message quantum interactive proofs and the quantum separability problem. In *2013 IEEE Conference on Computational Complexity*, pages 156–167. IEEE, 2013.
- [HP91] Fumio Hiai and Dénes Petz. The proper formula for relative entropy and its asymptotics in quantum probability. *Communications in Mathematical Physics*, 143(1):99–114, 1991.
- [IRS16] Raban Iten, Joseph M. Renes, and David Sutter. Pretty good measures in quantum information theory. *IEEE Transactions on Information Theory*, 63(2):1270–1279, 2016.
- [JV13] Peter D. Johnson and Lorenza Viola. Compatible quantum correlations: Extension problems for werner and isotropic states. *Phys. Rev. A*, 88:032323, Sep 2013.
- [KDW19] Eneet Kaur, Siddhartha Das, Mark M. Wilde, and Andreas Winter. Extendibility limits the performance of quantum processors. *Physical Review Letters*, 123(7):070502, 2019.
- [KDW21] Eneet Kaur, Siddhartha Das, Mark M. Wilde, and Andreas Winter. Resource theory of unextendibility and nonasymptotic quantum capacity. *Physical Review A*, 104(2):022401, August 2021.
- [KL17] Sumeet Khatri and Norbert Lütkenhaus. Numerical evidence for bound secrecy from two-way postprocessing in quantum key distribution. *Physical Review A*, 95(4):042320, 2017.
- [KW04] Masato Koashi and Andreas Winter. Monogamy of quantum entanglement and other correlations. *Physical Review A*, 69(2):022309, 2004.
- [KW20] Sumeet Khatri and Mark M. Wilde. *Principles of Quantum Communication Theory: A Modern Approach*. November 2020. arXiv:2011.04672v1.
- [LDMH⁺16] Cécilia Lancien, Sara Di Martino, Marcus Huber, Marco Piani, Gerardo Adesso, and Andreas Winter. Should entanglement measures be monogamous or faithful? *Physical Review Letters*, 117(6):060501, 2016.
- [LDS17] Felix Leditzky, Nilanjana Datta, and Graeme Smith. Useful states and entanglement distillation. *IEEE Transactions on Information Theory*, 64(7):4689–4708, 2017.
- [LKAW19] Ludovico Lami, Sumeet Khatri, Gerardo Adesso, and Mark M. Wilde. Extendibility of bosonic Gaussian states. *Physical Review Letters*, 123(5):050501, 2019.
- [LT76] Elliott H. Lieb and Walter Thirring. *Studies in Mathematical Physics*, chapter Inequalities for the moments of the eigenvalues of the Schroedinger Hamiltonian and their relation to Sobolev inequalities, pages 269–297. Princeton University Press, Princeton, 1976.
- [Mat15] Keiji Matsumoto. A new quantum version of f -divergence. In *Nagoya Winter Workshop: Reality and Measurement in Algebraic Quantum Theory*, pages 229–273. Springer, 2015.
- [MCL06] Tobias Moroder, Marcos Curty, and Norbert Lütkenhaus. One-way quantum key distribution: Simple upper bound on the secret key rate. *Physical Review A*, 74(5):052301, 2006.
- [MLDS⁺13] Martin Müller-Lennert, Frédéric Dupuis, Oleg Szehr, Serge Fehr, and Marco Tomamichel. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*,

- 54(12):122203, 2013.
- [MO15] Milán Mosonyi and Tomohiro Ogawa. Quantum hypothesis testing and the operational interpretation of the quantum Rényi relative entropies. *Communications in Mathematical Physics*, 334(3):1617–1648, 2015.
- [MRDL09] Geir Ove Myhr, Joseph M. Renes, Andrew C. Doherty, and Norbert Lütkenhaus. Symmetric extension in two-way quantum key distribution. *Physical Review A*, 79(4):042329, 2009.
- [Nag06] Hiroshi Nagaoka. The converse part of the theorem for quantum Hoeffding bound. 2006. arXiv:quant-ph/0611289.
- [NFB14] Naomi H. Nickerson, Joseph F. Fitzsimons, and Simon C. Benjamin. Freely scalable quantum technologies using cells of 5-to-50 qubits with very lossy and noisy photonic links. *Physical Review X*, 4(4):041041, 2014.
- [NH09] Marcin L. Nowakowski and Pawel Horodecki. A simple test for quantum channel capacity. *Journal of Physics A: Mathematical and Theoretical*, 42(13):135306, 2009.
- [OV06] Tobias J. Osborne and Frank Verstraete. General monogamy inequality for bipartite qubit entanglement. *Physical Review Letters*, 96(22):220503, 2006.
- [Pet86] Dénes Petz. Quasi-entropies for finite quantum systems. *Reports on Mathematical Physics*, 23(1):57–65, 1986.
- [PSBZ01] Jian-Wei Pan, Christoph Simon, Časlav Brukner, and Anton Zeilinger. Entanglement purification for quantum communication. *Nature*, 410(6832):1067–1070, 2001.
- [PV10] Yury Polyanskiy and Sergio Verdú. Arimoto channel coding converse and Rényi divergence. In *2010 48th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 1327–1333. IEEE, 2010.
- [Rai99] Eric M. Rains. Bound on distillable entanglement. *Physical Review A*, 60(1):179, 1999.
- [Rai01] Eric M. Rains. A semidefinite program for distillable entanglement. *IEEE Transactions on Information Theory*, 47(7):2921–2933, 2001.
- [RFGW19] Bartosz Regula, Kun Fang, Xin Wang, and Mile Gu. One-shot entanglement distillation beyond local operations and classical communication. *New Journal of Physics*, 21(10):103017, 2019.
- [RNBG20] Bartosz Regula, Varun Narasimhachar, Francesco Buscemi, and Mile Gu. Coherence manipulation with dephasing-covariant operations. *Physical Review Research*, 2(1):013109, January 2020.
- [RST⁺18] Filip Rozpedek, Thomas Schiet, Le Phuc Thinh, David Elkouss, Andrew C. Doherty, and Stephanie Wehner. Optimizing practical entanglement distillation. *Physical Review A*, 97(6):062333, June 2018.
- [SBW15] Kaushik P. Seshadreesan, Mario Berta, and Mark M. Wilde. Rényi squashed entanglement, discord, and relative entropy differences. *Journal of Physics A: Mathematical and Theoretical*, 48(39):395303, 2015.
- [SW12] Naresh Sharma and Naqeeb Ahmad Warsi. On the strong converses for the quantum channel capacity theorems. 2012. arXiv:1205.1712.
- [SW15] Kaushik P. Seshadreesan and Mark M. Wilde. Fidelity of recovery, squashed entanglement, and measurement recoverability. *Physical Review A*, 92(4):042321, 2015.
- [TDS03] Barbara M. Terhal, Andrew C. Doherty, and David Schwab. Symmetric extensions of quantum states and local hidden variable theories. *Physical Review Letters*, 90(15):157903, 2003.
- [Ter04] Barbara M. Terhal. Is entanglement monogamous? *IBM Journal of Research and Development*, 48(1):71–78, 2004.
- [Tom15] Marco Tomamichel. *Quantum Information Processing with Finite Resources: Mathematical Foundations*, volume 5. Springer, 2015.
- [TWW16] Marco Tomamichel, Mark M. Wilde, and Andreas Winter. Strong converse rates for quantum communication. *IEEE Transactions on Information Theory*, 63(1):715–727, 2016.
- [Uhl76] Armin Uhlmann. The “transition probability” in the state space of a $*$ -algebra. *Reports on Mathematical Physics*, 9(2):273–279, 1976.
- [Ume62] Hisaharu Umegaki. Conditional expectation in an operator algebra, IV (entropy and information). *Kodai Mathematical Seminar Reports*, 14(2):59–85, 1962.
- [VDC02] Guifré Vidal, W. Dür, and J. Ignacio Cirac. Entanglement cost of bipartite mixed states. *Physical Review Letters*, 89(2):027901, 2002.
- [Vid99] Guifré Vidal. Entanglement of pure states for a single copy. *Physical Review Letters*, 83(5):1046, 1999.

- [VP98] Vlatko Vedral and Martin B. Plenio. Entanglement measures and purification procedures. *Physical Review A*, 57(3):1619, 1998.
- [VPRK97] Vlatko Vedral, Martin B. Plenio, M. A. Rippin, and Peter L. Knight. Quantifying entanglement. *Physical Review Letters*, 78(12):2275–2279, March 1997. arXiv:quant-ph/9702027.
- [VW02] Guifré Vidal and Reinhard F. Werner. Computable measure of entanglement. *Physical Review A*, 65(3):032314, 2002.
- [Wat13] John Watrous. Simpler semidefinite programs for completely bounded norms. *Chicago Journal Of Theoretical Computer Science*, 8:1–19, 2013.
- [WD16] Xin Wang and Runyao Duan. Improved semidefinite programming upper bound on distillable entanglement. *Physical Review A*, 94(5):050301, 2016.
- [WD17a] Xin Wang and Runyao Duan. Irreversibility of asymptotic entanglement manipulation under quantum operations completely preserving positivity of partial transpose. *Physical Review Letters*, 119(18):180506, 2017.
- [WD17b] Xin Wang and Runyao Duan. Nonadditivity of Rains’ bound for distillable entanglement. *Physical Review A*, 95(6):062322, 2017.
- [Wer89] Reinhard F. Werner. An application of Bell’s inequalities to a quantum state extension problem. *Letters in Mathematical Physics*, 17(4):359–363, 1989.
- [WFD18] Xin Wang, Kun Fang, and Runyao Duan. Semidefinite programming converse bounds for quantum communication. *IEEE Transactions on Information Theory*, 65(4):2583–2592, 2018.
- [Wil17] Mark M. Wilde. *Quantum Information Theory*. Cambridge University Press, second edition, 2017.
- [Wil18] Mark M. Wilde. Optimized quantum f -divergences and data processing. *Journal of Physics A: Mathematical and Theoretical*, 51(37):374002, 2018.
- [WW19] Xin Wang and Mark M. Wilde. Resource theory of asymmetric distinguishability. *Physical Review Research*, 1(3):033170, 2019.
- [WW23] Xin Wang and Mark M. Wilde. Exact entanglement cost of quantum states and channels under PPT-preserving operations. *Physical Review A*, 107(1):012429, January 2023.
- [WWW20] Kun Wang, Xin Wang, and Mark M. Wilde. Quantification of unextendible entanglement and its applications in entanglement distillation. In *2020 IEEE International Symposium on Information Theory (ISIT)*, pages 1939–1943. IEEE, 2020.
- [WWY14] Mark M. Wilde, Andreas Winter, and Dong Yang. Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy. *Communications in Mathematical Physics*, 331(2):593–622, 2014.
- [XFWD17] Wei Xie, Kun Fang, Xin Wang, and Runyao Duan. Approximate broadcasting of quantum correlations. *Physical Review A*, 96(2):022302, 2017.
- [Yan06] Dong Yang. A simple proof of monogamy of entanglement. *Physics Letters A*, 360(2):249–250, 2006.
- [ZZW⁺21] Xuanqiang Zhao, Benchu Zhao, Zihong Wang, Zhixin Song, and Xin Wang. Practical distributed quantum information processing with LOCCNet. *npj Quantum Information*, 7(1):159, dec 2021.

Appendix A: Proof of Lemma 1

By utilizing the data-processing inequality and the faithfulness of the classical Rényi relative entropy, it follows that the statements above are true for all of the quantum Rényi relative entropies for the range of α for which data processing holds (see, e.g., [WWY14] for this kind of argument). To get outside of the range for $\alpha \in (0, 1)$, we use the following inequality holding for $\alpha \in (0, 1)$ [IRS16]:

$$\alpha D_\alpha(\omega \parallel \tau) \leq \tilde{D}_\alpha(\omega \parallel \tau). \quad (\text{A1})$$

To get outside of the range for $\alpha \in (1, \infty)$, we use the following inequality holding for $\alpha \in (1, \infty)$:

$$\tilde{D}_\alpha(\omega \parallel \tau) \leq D_\alpha(\omega \parallel \tau). \quad (\text{A2})$$

This inequality was proved in [WWY14, DL14] (following from the Araki–Lieb–Thirring inequality [Ara90, LT76]). For the same range, we also make use of the following inequality:

$$\tilde{D}_\alpha(\omega\|\tau) \leq \hat{D}_\alpha(\omega\|\tau). \quad (\text{A3})$$

which also follows from the Araki–Lieb–Thirring inequality for $\alpha \in (1, \infty)$ because

$$\tilde{Q}_\alpha(\omega\|\tau) = \text{Tr}[(\tau^{(1-\alpha)/2} \omega \tau^{(1-\alpha)/2})^\alpha] \quad (\text{A4})$$

$$= \text{Tr}[(\tau^{1/2} \omega \tau^{-1/2})^\alpha] \quad (\text{A5})$$

$$\leq \text{Tr}[(\tau^{1/2} \omega \tau^{-1/2})^\alpha (\tau^{1/2} \omega \tau^{-1/2})^\alpha] \quad (\text{A6})$$

$$= \text{Tr}[\tau (\tau^{-1/2} \omega \tau^{-1/2})^\alpha] \quad (\text{A7})$$

$$= \hat{Q}_\alpha(\omega\|\tau). \quad (\text{A8})$$

This concludes the proof.

Appendix B: Proof of Theorem 5

We first show the proof for (33). Let $\rho_{AB_1B_2}$ be an arbitrary extension of ρ_{AB_1} . Then define

$$p(y) := \text{Tr}[\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y(\rho_{AB_1B_2})], \quad (\text{B1})$$

$$\omega_{A'B'_1B'_2}^y := \frac{1}{p(y)} \mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y(\rho_{AB_1B_2}), \quad (\text{B2})$$

where $\mathcal{E}_{AB_1B_2 \rightarrow A'B'_1B'_2}^y$ is the operation extending $\mathcal{E}_{AB \rightarrow A'B'}^y$. Due to the channel extension property in (14), the values of $p(y)$ in (31) and (B1) are equal and $\omega_{A'B'_1B'_2}^y$ defined in (B2) is an extension of $\omega_{A'B'}^y$, the latter defined in (32) for arbitrary y .

From the data-processing inequality for the sandwiched Rényi relative entropy for $\alpha \in [1/2, 1) \cup (1, \infty)$, we conclude the following inequality:

$$\tilde{D}_\alpha(\rho_{AB_1} \|\rho_{AB_2}) \geq \tilde{D}_\alpha(\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_1}) \|\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_2})), \quad (\text{B3})$$

where the quantum channel $\mathcal{E}_{AB \rightarrow A'B'Y}$ is defined as

$$\mathcal{E}_{AB \rightarrow A'B'Y}(\cdot) := \sum_y \mathcal{E}_{AB \rightarrow A'B'}^y(\cdot) \otimes |y\rangle\langle y|_Y. \quad (\text{B4})$$

Since the marginal operations $\mathcal{E}_{AB_1 \rightarrow A'B'_1}^y$ and $\mathcal{E}_{AB_2 \rightarrow A'B'_2}^y$ are each equal to the original operation $\mathcal{E}_{AB \rightarrow A'B'}^y$, we find that

$$\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_1}) = \sum_y \mathcal{E}_{AB_1 \rightarrow A'B'_1}^y(\rho_{AB_1}) \otimes |y\rangle\langle y|_Y \quad (\text{B5})$$

$$= \sum_y p(y) \omega_{A'B'_1}^y \otimes |y\rangle\langle y|_Y \quad (\text{B6})$$

$$=: \omega_{A'B'_1Y}, \quad (\text{B7})$$

$$\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_2}) = \sum_y \mathcal{E}_{AB_2 \rightarrow A'B'_2}^y(\rho_{AB_2}) \otimes |y\rangle\langle y|_Y \quad (\text{B8})$$

$$= \sum_y p(y) \omega_{A'B'_2}^y \otimes |y\rangle\langle y|_Y \quad (\text{B9})$$

$$=: \omega_{A'B'_2Y}, \quad (\text{B10})$$

which implies that

$$\tilde{D}_\alpha(\mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_1}) \| \mathcal{E}_{AB \rightarrow A'B'Y}(\rho_{AB_2})) = \tilde{D}_\alpha(\omega_{A'B'_1Y} \| \omega_{A'B'_2Y}) \quad (\text{B11})$$

$$= \frac{1}{\alpha - 1} \log \sum_{y:p(y)>0} p(y) \tilde{Q}_\alpha(\omega_{A'B'_1}^y \| \omega_{A'B'_2}^y) \quad (\text{B12})$$

$$\geq \sum_{y:p(y)>0} p(y) \left[\frac{1}{\alpha - 1} \log \tilde{Q}_\alpha(\omega_{A'B'_1}^y \| \omega_{A'B'_2}^y) \right] \quad (\text{B13})$$

$$= \sum_{y:p(y)>0} p(y) \tilde{D}_\alpha(\omega_{A'B'_1}^y \| \omega_{A'B'_2}^y) \quad (\text{B14})$$

$$\geq 2 \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'_1}^y) \quad (\text{B15})$$

$$= 2 \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'}^y). \quad (\text{B16})$$

In the above, the first inequality follows from concavity of the logarithm and the fact that $\alpha > 1$. The second inequality follows because $\omega_{A'B'_1B'_2}^y$ is such that $\text{Tr}_{B'_2}[\omega_{A'B'_1B'_2}^y] = \omega_{A'B'_1}^y$, so that we can then optimize over every extension of $\omega_{A'B'_1}^y$. Since the following inequality has been established for every extension $\rho_{AB_1B_2}$ of ρ_{AB} :

$$\frac{1}{2} \tilde{D}_\alpha(\rho_{AB_1} \| \rho_{AB_2}) \geq \sum_{y:p(y)>0} p(y) E_\alpha^u(\omega_{A'B'}^y), \quad (\text{B17})$$

we conclude the inequality in (33).

The proof of (30) follows along similar lines, but instead using data processing for the Petz–Rényi relative entropy and the quantity Q_α in (3). In the same way, the proof of (34) follows, but instead using data processing for the geometric Rényi relative entropy and the quantity \hat{Q}_α in (7).

The inequality in (33) for $\alpha \in \{1, \infty\}$ follows by taking a limit. Alternatively, we can prove them directly. To establish the inequality in (33) for $\alpha = 1$ (for quantum relative entropy), we use the same reasoning but the following property of quantum relative entropy:

$$D(\omega_{XB} \| \tau_{XB}) = \sum_x p(x) D(\omega_B^x \| \tau_B^x) + D(p \| q), \quad (\text{B18})$$

where

$$\omega_{XB} := \sum_x p(x) |x\rangle\langle x|_X \otimes \omega_B^x, \quad (\text{B19})$$

$$\tau_{XB} := \sum_x q(x) |x\rangle\langle x|_X \otimes \tau_B^x. \quad (\text{B20})$$

Similarly, the inequality in (34) for $\alpha = 1$ follows from the same reasoning and the following related property of Belavkin–Staszewski relative entropy:

$$\hat{D}(\omega_{XB} \| \tau_{XB}) = \sum_x p(x) \hat{D}(\omega_B^x \| \tau_B^x) + D(p \| q). \quad (\text{B21})$$

For $\alpha = \infty$, suppose that the optimal extension state is $\rho_{AB_1B_2}$, which satisfies $\rho_{AB_1} = \rho_{AB}$ and $E_{\max}^u(\rho_{AB_1}) = \frac{1}{2} D_{\max}(\rho_{AB_1} \| \rho_{AB_2})$. Also, suppose that $E_{\max}^u(\rho_{AB_1}) = \frac{1}{2} \log t$, which means that

$\rho_{AB_1} \leq t\rho_{AB_2}$ For every y such that $p(y) > 0$, due to complete positivity and two-extendibility of the map $\mathcal{E}_{AB \rightarrow A'B'}^y$, we have that

$$t \mathcal{E}_{AB_2 \rightarrow A'B'_2}^y(\rho_{AB_2})/p(y) \geq \mathcal{E}_{AB_1 \rightarrow A'B'_1}^y(\rho_{AB_1})/p(y) = \omega_{A'B'}^y. \quad (\text{B22})$$

Noting that $\mathcal{E}_{AB_1 B_2 \rightarrow A'B'_1 B'_2}^y(\rho_{AB_1 B_2})/p(y)$ is an extension of the state $\omega_{A'B'}^y$, it follows that

$$E_{\max}^u(\omega_{A'B'}^y) \leq \frac{1}{2} \log t. \quad (\text{B23})$$

Therefore,

$$E_{\max}^u(\rho_{AB_1}) = \frac{1}{2} \log t \quad (\text{B24})$$

$$= \frac{1}{2} \log \sum_{y:p(y)>0} p(y)t \quad (\text{B25})$$

$$\geq \frac{1}{2} \log \sum_{y:p(y)>0} p(y)2^{2E_{\max}^u(\omega_{A'B'}^y)} \quad (\text{B26})$$

$$\geq \sum_{y:p(y)>0} p(y)E_{\max}^u(\omega_{A'B'}^y) \quad (\text{B27})$$

In the above, (B26) is due to (B23), while (B27) is due to the concavity of the logarithm.

Appendix C: Proof of Lemma 8

Without loss of generality, let σ_B be an arbitrary state with support equal to the support of the reduced state ψ_B . Then we have that

$$\tilde{D}_\alpha(\psi_{AB} \| \psi_A \otimes \sigma_B) = \frac{1}{\alpha - 1} \log \text{Tr} \left[\left(\psi_{AB}^{1/2} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \psi_{AB}^{1/2} \right)^\alpha \right] \quad (\text{C1})$$

$$= \frac{1}{\alpha - 1} \log \text{Tr} \left[\left(\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \psi_{AB} \right)^\alpha \right] \quad (\text{C2})$$

$$= \frac{1}{\alpha - 1} \log \text{Tr} \left[\left(|\psi\rangle\langle\psi|_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} |\psi\rangle\langle\psi|_{AB} \right)^\alpha \right] \quad (\text{C3})$$

$$= \frac{1}{\alpha - 1} \log \text{Tr} \left[\left(\langle\psi|_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} |\psi\rangle\langle\psi|_{AB} \right)^\alpha \right] \quad (\text{C4})$$

$$= \frac{1}{\alpha - 1} \log \text{Tr} \left[\left(\text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right] |\psi\rangle\langle\psi|_{AB} \right)^\alpha \right] \quad (\text{C5})$$

$$= \frac{1}{\alpha - 1} \log \left(\text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right] \right)^\alpha \text{Tr} [|\psi\rangle\langle\psi|_{AB}^\alpha] \quad (\text{C6})$$

$$= \frac{\alpha}{\alpha - 1} \log \text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right]. \quad (\text{C7})$$

For every pure bipartite state ψ_{AB} , there exists an operator X_A such that $\psi_{AB} = X_A \Gamma_{AB} X_A^\dagger$ and $\text{Tr}[X_A^\dagger X_A] = 1$, which implies that $\psi_A = X_A X_A^\dagger$. Note that we define

$$\Gamma_{AB} := |\Gamma\rangle\langle\Gamma|_{AB}, \quad |\Gamma\rangle_{AB} := \sum_{i=1}^d |i\rangle_A |i\rangle_B, \quad (\text{C8})$$

where $\{|i\rangle_A\}_i$ and $\{|i\rangle_B\}_i$ are orthonormal bases. Furthermore, by taking the polar decomposition of X_A , there exists a unitary U_A and a density operator ρ_A (having the same spectrum as ψ_A) such that $X_A = U_A \sqrt{\rho_A}$. Then consider that the last line above is equal to

$$\begin{aligned} & \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\psi_{AB} (\psi_A \otimes \sigma_B)^{\frac{1-\alpha}{\alpha}} \right] \\ &= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[X_A \Gamma_{AB} X_A^\dagger \left(X_A X_A^\dagger \otimes \sigma_B \right)^{\frac{1-\alpha}{\alpha}} \right] \end{aligned} \quad (\text{C9})$$

$$= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[X_A \Gamma_{AB} X_A^\dagger \left(\left(X_A X_A^\dagger \right)^{\frac{1-\alpha}{\alpha}} \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{C10})$$

$$= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\Gamma_{AB} \left(X_A^\dagger \left(X_A X_A^\dagger \right)^{\frac{1-\alpha}{\alpha}} X_A \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{C11})$$

$$= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\Gamma_{AB} \left(\sqrt{\rho_A} U_A^\dagger \left(U_A \rho_A U_A^\dagger \right)^{\frac{1-\alpha}{\alpha}} U_A \sqrt{\rho_A} \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{C12})$$

$$= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\Gamma_{AB} \left(\rho_A^{\frac{1}{\alpha}} \otimes \sigma_B^{\frac{1-\alpha}{\alpha}} \right) \right] \quad (\text{C13})$$

$$= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\Gamma_{AB} \left(\rho_A^{\frac{1}{\alpha}} T(\sigma)_A^{\frac{1-\alpha}{\alpha}} \otimes \mathbb{1}_B \right) \right] \quad (\text{C14})$$

$$= \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\rho_A^{\frac{1}{\alpha}} T(\sigma)_A^{\frac{1-\alpha}{\alpha}} \right]. \quad (\text{C15})$$

In the above, $T(\sigma)$ denotes the transpose of the state σ . Now taking a minimum over every state σ and applying [MLDS⁺13, Lemma 12], we find that

$$\min_{\sigma} \frac{\alpha}{\alpha-1} \log \text{Tr} \left[\rho_A^{\frac{1}{\alpha}} T(\sigma)_A^{\frac{1-\alpha}{\alpha}} \right] = \frac{\alpha}{\alpha-1} \log \left\| \rho_A^{\frac{1}{\alpha}} \right\|_{\alpha/(2\alpha-1)} \quad (\text{C16})$$

$$= \frac{\alpha}{\alpha-1} \frac{2\alpha-1}{\alpha} \log \text{Tr} [(\rho_A^{\frac{1}{\alpha}})^{\alpha/(2\alpha-1)}] \quad (\text{C17})$$

$$= \frac{2\alpha-1}{\alpha-1} \log \text{Tr} [\rho_A^{\frac{1}{2\alpha-1}}] \quad (\text{C18})$$

$$= \frac{2}{1 - (1/[2\alpha-1])} \log \text{Tr} [\rho_A^{\frac{1}{2\alpha-1}}] \quad (\text{C19})$$

$$= 2H_{[2\alpha-1]^{-1}}(\rho_A). \quad (\text{C20})$$

The conclusion that $H_{[2\alpha-1]^{-1}}(\rho_A) = H_{[2\alpha-1]^{-1}}(\psi_A)$ follows because ρ_A and ψ_A have the same spectrum.

Appendix D: Proof of Lemma 9

Consider that, by similar reasoning employed in Appendix C,

$$\psi_{AB} = \psi_A^{1/2} U_A \Gamma_{AB} U_A^\dagger \psi_A^{1/2}, \quad (\text{D1})$$

where U_A is a unitary, ψ_A is the marginal density operator of ψ_{AB} on system A ,

$$\Gamma_{AB} := |\Gamma\rangle\langle\Gamma|_{AB}, \quad |\Gamma\rangle_{AB} := \sum_{i=1}^d |i\rangle_A |i\rangle_B, \quad (\text{D2})$$

$\{|i\rangle_A\}_i$ and $\{|i\rangle_B\}_i$ are the orthonormal bases for the Schmidt decomposition of ψ_{AB} , and d is the Schmidt rank of ψ_{AB} . Then, by noting that it suffices to optimize over every state σ_B having the same support as ψ_B , we find that

$$\hat{I}_\alpha(A; B)_\psi := \min_{\sigma_B} \hat{D}_\alpha(\psi_{AB} \| \psi_A \otimes \sigma_B) \quad (\text{D3})$$

$$= \min_{\sigma_B} \frac{1}{\alpha - 1} \log \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[(\psi_A \otimes \sigma_B)^{-1/2} \psi_{AB} (\psi_A \otimes \sigma_B)^{-1/2} \right]^\alpha \right] \quad (\text{D4})$$

Consider that

$$\begin{aligned} & \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[\left(\psi_A^{-1/2} \otimes \sigma_B^{-1/2} \right) \psi_{AB} \left(\psi_A^{-1/2} \otimes \sigma_B^{-1/2} \right) \right]^\alpha \right] \\ &= \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[\left(\psi_A^{-1/2} \otimes \sigma_B^{-1/2} \right) \psi_A^{1/2} U_A \Gamma_{AB} U_A^\dagger \psi_A^{1/2} \left(\psi_A^{-1/2} \otimes \sigma_B^{-1/2} \right) \right]^\alpha \right] \end{aligned} \quad (\text{D5})$$

$$= \text{Tr} \left[(\psi_A \otimes \sigma_B) \left[\left(U_A \otimes \sigma_B^{-1/2} \right) \Gamma_{AB} \left(U_A^\dagger \otimes \sigma_B^{-1/2} \right) \right]^\alpha \right] \quad (\text{D6})$$

$$= \text{Tr} \left[(\psi_A \otimes \sigma_B) U_A \left[\sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2} \right]^\alpha U_A^\dagger \right] \quad (\text{D7})$$

$$= \text{Tr} \left[\left(U_A^\dagger \psi_A U_A \otimes \sigma_B \right) \left[\frac{\sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2}}{\text{Tr}[\sigma_B^{-1}]} \right]^\alpha \right] (\text{Tr}[\sigma_B^{-1}])^\alpha \quad (\text{D8})$$

$$= \text{Tr} \left[\left(U_A^\dagger \psi_A U_A \otimes \sigma_B \right) \frac{\sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2}}{\text{Tr}[\sigma_B^{-1}]} \right] (\text{Tr}[\sigma_B^{-1}])^\alpha \quad (\text{D9})$$

$$= \text{Tr} \left[\left(U_A^\dagger \psi_A U_A \otimes \sigma_B \right) \sigma_B^{-1/2} \Gamma_{AB} \sigma_B^{-1/2} \right] (\text{Tr}[\sigma_B^{-1}])^{\alpha-1} \quad (\text{D10})$$

$$= \text{Tr} \left[U_A^\dagger \psi_A U_A \Gamma_{AB} \right] (\text{Tr}[\sigma_B^{-1}])^{\alpha-1} \quad (\text{D11})$$

$$= (\text{Tr}[\sigma_B^{-1}])^{\alpha-1}. \quad (\text{D12})$$

In the fourth step, we are using the fact that $\sigma_B^{-1/2} |\Gamma\rangle_{AB}$ is a vector with norm

$$\left\| \sigma_B^{-1/2} |\Gamma\rangle_{AB} \right\|_2 = \sqrt{\langle \Gamma |_{AB} \sigma_B^{-1/2} \sigma_B^{-1/2} | \Gamma \rangle_{AB}} \quad (\text{D13})$$

$$= \sqrt{\langle \Gamma |_{AB} \sigma_B^{-1} | \Gamma \rangle_{AB}} \quad (\text{D14})$$

$$= \sqrt{\text{Tr}[\sigma_B^{-1}]}. \quad (\text{D15})$$

Then it follows that

$$\hat{I}_\alpha(A; B)_\psi = \min_{\sigma_B} \frac{1}{\alpha - 1} \log (\text{Tr}[\sigma_B^{-1}])^{\alpha-1} \quad (\text{D16})$$

$$= \min_{\sigma_B} \log \text{Tr}[\sigma_B^{-1}]. \quad (\text{D17})$$

The minimum value occurs when $\sigma_B = \mathbb{1}_B/d_B$ (the maximally mixed state in system B , where we again note that B is restricted to the support of ψ_B so that d_B is the dimension of the support of ψ_B). To conclude this final step, one can use the Lagrange multiplier method. So then the equality in the statement of Lemma 9 follows.

Appendix E: Proof of Proposition 13

Let $\sigma_{A_1 B_1 B'_1}$ and $\sigma_{A_2 B_2 B'_2}$ be arbitrary extensions of $\rho_{A_1 B_1}$ and $\rho_{A_2 B_2}$, respectively. Consider the state $\sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}$. Then it follows that

$$\text{Tr}_{B'_1 B'_2} [\sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}] = \text{Tr}_{B'_1} [\sigma_{A_1 B_1 B'_1}] \otimes \text{Tr}_{B'_2} [\sigma_{A_2 B_2 B'_2}] = \rho_{A_1 B_1} \otimes \rho_{A_2 B_2}, \quad (\text{E1})$$

so that $\sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}$ is an extension of $\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}$. Thus

$$E_\alpha^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \frac{1}{2} D_\alpha(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2} \| \sigma_{A_1 B'_1} \otimes \sigma_{A_2 B'_2}) \quad (\text{E2})$$

$$= \frac{1}{2} D_\alpha(\rho_{A_1 B_1} \| \sigma_{A_1 B'_1}) + \frac{1}{2} D_\alpha(\rho_{A_2 B_2} \| \sigma_{A_2 B'_2}). \quad (\text{E3})$$

Taking an infimum over all extensions $\sigma_{A_1 B_1 B'_1}$ and $\sigma_{A_2 B_2 B'_2}$ leads to the inequality in (53). The inequalities in (54) and (55) are established similarly.

Appendix F: Derivations of dual SDPs and strong duality for max- and min-unextendible entanglement

We first derive the dual SDP in (60), by means of the Lagrange multiplier method. Consider that

$$\begin{aligned} & \sup_{\lambda, \sigma_{ABB'} \geq 0} \{ \lambda : \lambda \rho_{AB} \leq \text{Tr}_B[\sigma_{ABB'}], \rho_{AB} = \text{Tr}_{B'}[\sigma_{ABB'}] \} \\ &= \sup_{\lambda, \sigma_{ABB'} \geq 0} \inf_{\substack{X_{AB} \geq 0, \\ Y_{AB} \in \text{Herm}}} \left\{ \begin{aligned} & \lambda + \text{Tr}[X_{AB}(\text{Tr}_B[\sigma_{ABB'}] - \lambda \rho_{AB})] \\ & + \text{Tr}[Y_{AB}(\rho_{AB} - \text{Tr}_{B'}[\sigma_{ABB'}])] \end{aligned} \right\} \quad (\text{F1}) \end{aligned}$$

$$= \sup_{\lambda, \sigma_{ABB'} \geq 0} \inf_{\substack{X_{AB} \geq 0, \\ Y_{AB} \in \text{Herm}}} \left\{ \begin{aligned} & \text{Tr}[Y_{AB} \rho_{AB}] + \lambda(1 - \text{Tr}[X_{AB} \rho_{AB}]) \\ & + \text{Tr}[(X_{AB'} \otimes \mathbb{1}_B - Y_{AB} \otimes \mathbb{1}_{B'}) \sigma_{ABB'}] \end{aligned} \right\} \quad (\text{F2})$$

$$\leq \inf_{\substack{X_{AB} \geq 0, \\ Y_{AB} \in \text{Herm}}} \sup_{\lambda, \sigma_{ABB'} \geq 0} \left\{ \begin{aligned} & \text{Tr}[Y_{AB} \rho_{AB}] + \lambda(1 - \text{Tr}[X_{AB} \rho_{AB}]) \\ & + \text{Tr}[(X_{AB'} \otimes \mathbb{1}_B - Y_{AB} \otimes \mathbb{1}_{B'}) \sigma_{ABB'}] \end{aligned} \right\} \quad (\text{F3})$$

$$= \inf_{\substack{X_{AB} \geq 0, \\ Y_{AB} \in \text{Herm}}} \left\{ \begin{aligned} & \text{Tr}[Y_{AB} \rho_{AB}] : \text{Tr}[X_{AB} \rho_{AB}] \geq 1, \\ & X_{AB'} \otimes \mathbb{1}_B \leq Y_{AB} \otimes \mathbb{1}_{B'} \end{aligned} \right\} \quad (\text{F4})$$

$$= \inf_{X_{AB}, Y_{AB} \geq 0} \left\{ \begin{aligned} & \text{Tr}[Y_{AB} \rho_{AB}] : \text{Tr}[X_{AB} \rho_{AB}] \geq 1, \\ & X_{AB'} \otimes \mathbb{1}_B \leq Y_{AB} \otimes \mathbb{1}_{B'} \end{aligned} \right\}. \quad (\text{F5})$$

The first equality follows by introducing the Lagrange multipliers $X_{AB} \geq 0$ and $Y_{AB} \in \text{Herm}$. Indeed, the constraint $\lambda \rho_{AB} \leq \text{Tr}_B[\sigma_{ABB'}]$ does not hold if and only if

$$\inf_{X_{AB} \geq 0} \text{Tr}[X_{AB}(\text{Tr}_B[\sigma_{ABB'}] - \lambda \rho_{AB})] = -\infty, \quad (\text{F6})$$

and the constraint $\rho_{AB} = \text{Tr}_{B'}[\sigma_{ABB'}]$ does not hold if and only if

$$\inf_{Y_{AB} \in \text{Herm}} \text{Tr}[Y_{AB}(\rho_{AB} - \text{Tr}_{B'}[\sigma_{ABB'}])] = -\infty. \quad (\text{F7})$$

The second equality follows from basic algebra. The inequality follows from the max-min inequality. The third equality follows by interpreting $\lambda, \sigma_{ABB'} \geq 0$ as Lagrange multipliers. Indeed, the constraint $\text{Tr}[X_{AB} \rho_{AB}] \geq 1$ does not hold if and only if

$$\sup_{\lambda \geq 0} \lambda(1 - \text{Tr}[X_{AB} \rho_{AB}]) = +\infty, \quad (\text{F8})$$

and the constraint $X_{AB'} \otimes \mathbb{1}_B \leq Y_{AB} \otimes \mathbb{1}_{B'}$ does not hold if and only if

$$\sup_{\sigma_{ABB'} \geq 0} \text{Tr}[(X_{AB'} \otimes \mathbb{1}_B - Y_{AB} \otimes \mathbb{1}_{B'}) \sigma_{ABB'}] = +\infty. \quad (\text{F9})$$

The final equality follows because the constraint $X_{AB'} \otimes \mathbb{1}_B \leq Y_{AB} \otimes \mathbb{1}_{B'}$ implies that $Y_{AB} \geq 0$ since $X_{AB'} \otimes \mathbb{1}_B \geq 0$ and the partial trace is a completely positive map.

We can actually conclude that strong duality holds (i.e., equality in all steps above) because Slater's condition holds. Indeed, a feasible choice for the primal SDP is to pick $\sigma_{ABB'} = \rho_{AB} \otimes \pi_{B'}$, where $\pi_{B'}$ is the maximally mixed state, and $\lambda = \frac{1}{d_B^2}$. We are then guaranteed that $\lambda \rho_{AB} \leq \text{Tr}_B[\sigma_{ABB'}]$ for these choices because $\frac{1}{d_B^2} \rho_{AB} \leq \frac{1}{d_B^2} \sum_{i=1}^{d_B^2} U_B^i \rho_{AB} U_B^{i\dagger} = \rho_A \otimes \pi_{B'}$, where $\{U_B^i\}_{i=1}^{d_B^2}$ is the set of Heisenberg–Weyl unitaries. A strictly feasible choice for the dual SDP is $X_{AB} = 2\mathbb{1}_{AB}$ and $Y_{AB} = 3\mathbb{1}_{AB}$.

Let us now derive the dual SDP in (66), again by means of the Lagrange multiplier method. Consider that

$$\begin{aligned} & \sup_{\sigma_{ABB'} \geq 0} \left\{ \text{Tr}[\Pi_{AB'}^\rho, \text{Tr}_B[\sigma_{ABB'}]] : \text{Tr}_{B'}[\sigma_{ABB'}] = \rho_{AB} \right\} \\ &= \sup_{\sigma_{ABB'} \geq 0} \inf_{X_{AB} \in \text{Herm}} \left\{ \text{Tr}[\Pi_{AB'}^\rho, \text{Tr}_B[\sigma_{ABB'}]] + \text{Tr}[X_{AB}(\rho_{AB} - \text{Tr}_{B'}[\sigma_{ABB'}])] \right\} \end{aligned} \quad (\text{F10})$$

$$= \sup_{\sigma_{ABB'} \geq 0} \inf_{X_{AB} \in \text{Herm}} \left\{ \text{Tr}[X_{AB} \rho_{AB}] + \text{Tr}[(\Pi_{AB'}^\rho \otimes \mathbb{1}_B - X_{AB} \otimes \mathbb{1}_{B'}) \sigma_{ABB'}] \right\} \quad (\text{F11})$$

$$\leq \inf_{X_{AB} \in \text{Herm}} \sup_{\sigma_{ABB'} \geq 0} \left\{ \text{Tr}[X_{AB} \rho_{AB}] + \text{Tr}[(\Pi_{AB'}^\rho \otimes \mathbb{1}_B - X_{AB} \otimes \mathbb{1}_{B'}) \sigma_{ABB'}] \right\} \quad (\text{F12})$$

$$= \inf_{X_{AB} \in \text{Herm}} \left\{ \text{Tr}[X_{AB} \rho_{AB}] : \Pi_{AB'}^\rho \otimes \mathbb{1}_B \leq X_{AB} \otimes \mathbb{1}_{B'} \right\} \quad (\text{F13})$$

$$= \inf_{X_{AB} \geq 0} \left\{ \text{Tr}[X_{AB} \rho_{AB}] : \Pi_{AB'}^\rho \otimes \mathbb{1}_B \leq X_{AB} \otimes \mathbb{1}_{B'} \right\}. \quad (\text{F14})$$

The first equality follows because the constraint $\text{Tr}_{B'}[\sigma_{ABB'}] = \rho_{AB}$ does not hold if and only if

$$\inf_{X_{AB} \in \text{Herm}} \text{Tr}[X_{AB}(\rho_{AB} - \text{Tr}_{B'}[\sigma_{ABB'}])] = -\infty. \quad (\text{F15})$$

The second equality follows from basic algebra. The inequality follows from the max-min inequality. The third equality follows because the constraint $\Pi_{AB'}^\rho \otimes \mathbb{1}_B \leq X_{AB} \otimes \mathbb{1}_{B'}$ does not hold if and only if

$$\sup_{\sigma_{ABB'} \geq 0} \text{Tr}[(\Pi_{AB'}^\rho \otimes \mathbb{1}_B - X_{AB} \otimes \mathbb{1}_{B'}) \sigma_{ABB'}] = +\infty. \quad (\text{F16})$$

The final equality follows because the constraint $\Pi_{AB'}^\rho \otimes \mathbb{1}_B \leq X_{AB} \otimes \mathbb{1}_{B'}$ implies that $X_{AB} \geq 0$ since $\Pi_{AB'}^\rho \otimes \mathbb{1}_B \geq 0$ and the partial trace is a completely positive map.

Strong duality holds (i.e., equality in all steps above) because Slater's condition holds. Indeed, a feasible choice for the primal SDP is $\sigma_{ABB'} = \rho_{AB} \otimes \pi_{B'}$. A strictly feasible choice for the dual SDP is $X_{AB} = 2\mathbb{1}_{AB}$.

Appendix G: Proof of Proposition 14

Subadditivity follows from investigating the primal SDP. Suppose that $(\lambda_1, \sigma_{A_1 B_1 B'_1})$ and $(\lambda_2, \sigma_{A_2 B_2 B'_2})$ achieve $E_{\max}^u(\rho_{A_1 B_1})$ and $E_{\max}^u(\rho_{A_2 B_2})$, respectively. As

$$\lambda_1 \lambda_2 \rho_{A_1 B_1} \otimes \rho_{A_2 B_2} = \lambda_1 \rho_{A_1 B_1} \otimes \lambda_2 \rho_{A_2 B_2} \leq \sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2}, \quad (\text{G1})$$

one can check that $(\lambda_1 \lambda_2, \sigma_{A_1 B_1 B'_1} \otimes \sigma_{A_2 B_2 B'_2})$ forms a feasible solution. Thus

$$2^{-2E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})} \geq \lambda_1 \lambda_2 = 2^{-2E_{\max}^u(\rho_{A_1 B_1}) - 2E_{\max}^u(\rho_{A_2 B_2})},$$

which gives $E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq E_{\max}^u(\rho_{A_1 B_1}) + E_{\max}^u(\rho_{A_2 B_2})$.

The superadditivity is shown by investigating the dual SDP. Suppose that $(X_{A_1 B_1}, Y_{A_1 B_1})$ and $(X_{A_2 B_2}, Y_{A_2 B_2})$ achieve $E_{\max}^u(\rho_{A_1 B_1})$ and $E_{\max}^u(\rho_{A_2 B_2})$, respectively. As

$$\text{Tr}[(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})(X_{A_1 B_1} \otimes X_{A_2 B_2})] = \text{Tr}[\rho_{A_1 B_1} X_{A_1 B_1}] \text{Tr}[\rho_{A_2 B_2} X_{A_2 B_2}] \geq 1 \quad (\text{G2})$$

and $X_{A_1 B'_1} \otimes X_{A_2 B'_2} \otimes \mathbb{1}_{B_1 B_2} \leq Y_{A_1 B_1} \otimes Y_{A_2 B_2} \otimes \mathbb{1}_{B'_1 B'_2}$, the pair $(X_{A_1 B_1} \otimes X_{A_2 B_2}, Y_{A_1 B_1} \otimes Y_{A_2 B_2})$ forms a dual feasible solution. Thus

$$2^{-2E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})} \leq \text{Tr}[\rho_{A_1 B_1} Y_{A_1 B_1}] \text{Tr}[\rho_{A_2 B_2} Y_{A_2 B_2}] = 2^{-2E_{\max}^u(\rho_{A_1 B_1}) - 2E_{\max}^u(\rho_{A_2 B_2})}, \quad (\text{G3})$$

which gives $E_{\max}^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \geq E_{\max}^u(\rho_{A_1 B_1}) + E_{\max}^u(\rho_{A_2 B_2})$.

Appendix H: Proof of Proposition 16

This proposition follows from (45) by taking the limit $\alpha \rightarrow 0$. Alternatively, we briefly outline another proof as follows. For a pure state ψ_{AB} , an arbitrary extension of it has the form $\sigma_{ABB'} := |\psi\rangle\langle\psi|_{AB} \otimes \sigma_{B'}$, where $\sigma_{B'}$ is a state. Then $\text{Tr}_B[\sigma_{ABB'}] = \psi_A \otimes \sigma_{B'}$. Suppose that a spectral decomposition of $\sigma_{B'}$ is $\sigma_{B'} = \sum_n p_n |e_n\rangle\langle e_n|$. Then

$$\begin{aligned} & \frac{1}{2} D_{\min}(\psi_{AB} \| \psi_A \otimes \sigma_{B'}) \\ &= -\frac{1}{2} \log \sum_{i,j,m,n} \sqrt{\alpha_i \alpha_j} \alpha_m p_n \langle \psi_i |_A \langle \psi_i |_B (|\psi_m\rangle\langle\psi_m|_A \otimes |e_n\rangle\langle e_n|_B) |\psi_j\rangle_A |\psi_j\rangle_B \end{aligned} \quad (\text{H1})$$

$$= -\frac{1}{2} \log \sum_{i,n} \alpha_i^2 p_n |\langle \psi_i | e_n \rangle|^2 \quad (\text{H2})$$

$$\geq -\frac{1}{2} \log \alpha_1^2 \sum_{i,n} p_n |\langle \psi_i | e_n \rangle|^2 \quad (\text{H3})$$

$$= -\log \alpha_1. \quad (\text{H4})$$

The inequality is achieved by choosing $\sigma_{B'} = |\psi_1\rangle\langle\psi_1|_B$. Thus $E_{\min}^u(\psi_{AB}) = -\log \alpha_1$.

Appendix I: Dual SDP of unextendible fidelity

Here we derive the dual SDP of the primal SDP in (77) for unextendible fidelity. We follow an argument similar to that given in [BT16]. We first bring the primal program into standard form, which expresses the primal problem as a maximization of the functional $\text{Tr}[AX]$ over $X \geq 0$, subject to the constraint $\Phi(X) = B$. Hence, we set

$$X = \begin{pmatrix} X_{11} & Z_{AB} & \cdot \\ Z_{AB}^\dagger & X_{22} & \cdot \\ \cdot & \cdot & \sigma_{ABB'} \end{pmatrix}, \quad A = \frac{1}{2} \begin{pmatrix} 0 & \mathbb{1}_{AB} & 0 \\ \mathbb{1}_{AB} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \rho_{AB} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \rho_{AB} \end{pmatrix}, \quad (\text{I1})$$

and

$$\Phi(X) := \begin{pmatrix} X_{11} & 0 & 0 \\ 0 & X_{22} - \text{Tr}_B[\sigma_{ABB'}] & 0 \\ 0 & 0 & \text{Tr}_{B'}[\sigma_{ABB'}] \end{pmatrix}. \quad (\text{I2})$$

The variables with the placeholder ‘.’ are of no interest. The dual SDP is a minimization over self-adjoint Y of the functional $\text{Tr}[BY]$ subject to $\Phi^\dagger(Y) \geq A$. The dual variables and adjoint map can be determined to be

$$Y = \begin{pmatrix} Y_{11} & \cdot & \cdot \\ \cdot & Y_{22} & \cdot \\ \cdot & \cdot & Y_{33} \end{pmatrix} \quad \text{and} \quad \Phi^\dagger(Y) := \begin{pmatrix} Y_{11} & 0 & 0 \\ 0 & Y_{22} & 0 \\ 0 & 0 & Y_{33} \otimes \mathbb{1}_{B'} - Y_{22} \otimes \mathbb{1}_B \end{pmatrix}. \quad (\text{I3})$$

This leads to the following dual problem:

$$\begin{aligned} & \text{minimize} \quad \text{Tr}[(Y_{11} + Y_{33})\rho_{AB}] \\ & \text{subject to} \quad \begin{pmatrix} Y_{11} & 0 \\ 0 & Y_{22} \end{pmatrix} \geq \frac{1}{2} \begin{pmatrix} 0 & \mathbb{1}_{AB} \\ \mathbb{1}_{AB} & 0 \end{pmatrix} \\ & \quad Y_{33} \otimes \mathbb{1}_{B'} \geq Y_{22} \otimes \mathbb{1}_B \\ & \quad Y_{11}, Y_{22}, Y_{33} \in \text{Herm}(AB). \end{aligned} \quad (\text{I4})$$

The Slater condition for strong duality is satisfied, using the fact that the primal problem is feasible and the dual problem is strictly feasible. To see this, let $\sigma_{B'}$ be a quantum state. Then the operator

$$\tilde{X} = \begin{pmatrix} \rho_{AB} & 0 & 0 \\ 0 & \rho_A \otimes \sigma_{B'} & 0 \\ 0 & 0 & \rho_{AB} \otimes \sigma_{B'} \end{pmatrix} \quad (\text{I5})$$

is primal feasible since $\tilde{X} \geq 0$ and $\Phi(\tilde{X}) = B$. For the dual problem, the operator

$$\tilde{Y} = \begin{pmatrix} \mathbb{1}_{AB} & 0 & 0 \\ 0 & \mathbb{1}_{AB} & 0 \\ 0 & 0 & 2\mathbb{1}_{AB} \end{pmatrix}, \quad (\text{I6})$$

is strictly feasible since

$$\Phi^\dagger(\tilde{Y}) = \begin{pmatrix} \mathbb{1}_{AB} & 0 & 0 \\ 0 & \mathbb{1}_{AB} & 0 \\ 0 & 0 & \mathbb{1}_{ABB'} \end{pmatrix} > \frac{1}{2} \begin{pmatrix} 0 & \mathbb{1}_{AB} & 0 \\ \mathbb{1}_{AB} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \equiv A. \quad (\text{I7})$$

By performing the substitutions $Y_{11} \rightarrow \frac{1}{2}Y_{11}$ and $Y_{33} \rightarrow \frac{1}{2}Y_{33}$ and noting that

$$\begin{pmatrix} Y_{11} & 0 \\ 0 & Y_{22} \end{pmatrix} \geq \begin{pmatrix} 0 & \mathbb{1}_{AB} \\ \mathbb{1}_{AB} & 0 \end{pmatrix} \Leftrightarrow \begin{pmatrix} Y_{11} & -\mathbb{1}_{AB} \\ -\mathbb{1}_{AB} & Y_{22} \end{pmatrix} \geq 0, \quad (\text{I8})$$

we finally obtain the dual stated in (78):

$$\begin{aligned} & \text{minimize} \quad \frac{1}{2} \text{Tr}[(Y_{11} + Y_{33})\rho_{AB}] \\ & \text{subject to} \quad \begin{pmatrix} Y_{11} & -\mathbb{1}_{AB} \\ -\mathbb{1}_{AB} & Y_{22} \end{pmatrix} \geq 0 \\ & \quad Y_{33} \otimes \mathbb{1}_{B'} \geq Y_{22} \otimes \mathbb{1}_B \\ & \quad Y_{11}, Y_{22}, Y_{33} \geq 0. \end{aligned} \quad (\text{I9})$$

In this final step, we also used the facts that

$$\begin{pmatrix} Y_{11} & -\mathbb{1}_{AB} \\ -\mathbb{1}_{AB} & Y_{22} \end{pmatrix} \geq 0 \Rightarrow Y_{11}, Y_{22} \geq 0, \quad (\text{I10})$$

$$Y_{22} \geq 0, \quad Y_{33} \otimes \mathbb{1}_{B'} \geq Y_{22} \otimes \mathbb{1}_B \quad \Rightarrow \quad Y_{33} \geq 0. \quad (\text{I11})$$

The dual problem in (19) can be further simplified since the first matrix inequality therein holds if and only if $Y_{11}, Y_{22} > 0$ and $Y_{22} \geq Y_{11}^{-1}$ [Wat13]. Without loss of generality, we can choose $Y_{22} = Y_{11}^{-1}$, and the problem simplifies to

$$\begin{aligned} & \text{infimum} \quad \frac{1}{2} \text{Tr} [Y_{AB'}^{-1} \rho_{AB'}] + \frac{1}{2} \text{Tr} [Z_{AB} \rho_{AB}] \\ & \text{subject to} \quad Z_{AB} \otimes \mathbb{1}_{B'} \geq Y_{AB'} \otimes \mathbb{1}_B \\ & \quad \quad \quad Y_{AB'}, Z_{AB} > 0. \end{aligned} \quad (\text{I12})$$

By the arithmetic-geometric mean inequality, it holds that

$$\frac{1}{2} \text{Tr} [Y_{AB'}^{-1} \rho_{AB'}] + \frac{1}{2} \text{Tr} [Z_{AB} \rho_{AB}] \geq \sqrt{\text{Tr} [Y_{AB'}^{-1} \rho_{AB'}] \text{Tr} [Z_{AB} \rho_{AB}]}, \quad (\text{I13})$$

for every $Y_{AB'} > 0$, with equality when the two terms are equal. For every feasible pair $(Y_{AB'}, Z_{AB})$, there exists a constant $\lambda > 0$ such that the two trace terms evaluated on $(\lambda Y_{AB'}, \lambda Z_{AB})$ are equal. Hence, we can restrict the optimization to such rescaled pairs of operators, resulting

$$\begin{aligned} & \text{infimum} \quad \sqrt{\text{Tr} [Y_{AB'}^{-1} \rho_{AB'}] \text{Tr} [Z_{AB} \rho_{AB}]} \\ & \text{subject to} \quad Z_{AB} \otimes \mathbb{1}_{B'} \geq Y_{AB'} \otimes \mathbb{1}_B \\ & \quad \quad \quad Y_{AB'}, Z_{AB} > 0. \end{aligned} \quad (\text{I14})$$

We remark that this equivalent representation of the dual problem is essential when proving the multiplicativity of unextendible fidelity.

Appendix J: Proof of Proposition 18

For the “ \geq ” part, suppose that $\rho_{A_1 B_1 B'_1}$ and $\rho_{A_2 B_2 B'_2}$ achieve $F^u(\rho_{A_1 B_1})$ and $F^u(\rho_{A_2 B_2})$, respectively. Since

$$\text{Tr}_{B'_1 B'_2} [\rho_{A_1 B_1 B'_1} \otimes \rho_{A_2 B_2 B'_2}] = \rho_{A_1 B_1} \otimes \rho_{A_2 B_2}, \quad (\text{J1})$$

the state $\rho_{A_1 B_1 B'_1} \otimes \rho_{A_2 B_2 B'_2}$ is a feasible solution for $F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})$. So

$$F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \geq F(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}, \rho_{A_1 B'_1} \otimes \rho_{A_2 B'_2}) \quad (\text{J2})$$

$$= F(\rho_{A_1 B_1}, \rho_{A_1 B'_1}) F(\rho_{A_2 B_2}, \rho_{A_2 B'_2}) \quad (\text{J3})$$

$$= F^u(\rho_{A_1 B_1}) F^u(\rho_{A_2 B_2}), \quad (\text{J4})$$

where the first equality follows from multiplicativity of fidelity.

For the “ \leq ” part, we employ the dual representation in (79). Let $(Y_{A_1 B'_1}, Z_{A_1 B_1})$ and $(Y_{A_2 B'_2}, Z_{A_2 B_2})$ be feasible solutions for $F^u(\rho_{A_1 B_1})$ and $F^u(\rho_{A_2 B_2})$, respectively. Set $Y_{A_1 A_2 B'_1 B'_2} \equiv Y_{A_1 B'_1} \otimes Y_{A_2 B'_2}$ and $Z_{A_1 A_2 B_1 B_2} \equiv Z_{A_1 B_1} \otimes Z_{A_2 B_2}$. Since

$$Z_{A_1 A_2 B_1 B_2} \otimes \mathbb{1}_{B'_1 B'_2} = (Z_{A_1 B_1} \otimes \mathbb{1}_{B'_1}) \otimes (Z_{A_2 B_2} \otimes \mathbb{1}_{B'_2}) \quad (\text{J5})$$

$$\geq (Y_{A_1 B'_1} \otimes \mathbb{1}_{B_1}) \otimes (Y_{A_2 B'_2} \otimes \mathbb{1}_{B_2}) = Y_{A_1 A_2 B'_1 B'_2}, \quad (\text{J6})$$

it follows that $(Y_{A_1 A_2 B'_1 B'_2}, Z_{A_1 A_2 B_1 B_2})$ is a feasible solution. Then

$$F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq \sqrt{\text{Tr} [Y_{A_1 A_2 B'_1 B'_2}^{-1} (\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})] \text{Tr} [Z_{A_1 A_2 B_1 B_2} (\rho_{A_1 B_1} \otimes \rho_{A_2 B_2})]} \quad (\text{J7})$$

$$= \sqrt{\text{Tr}[Y_{A_1 B_1}^{-1} \rho_{A_1 B_1}] \text{Tr}[Z_{A_1 B_1} \rho_{A_1 B_1}]} \sqrt{\text{Tr}[Y_{A_2 B_2}^{-1} \rho_{A_2 B_2}] \text{Tr}[Z_{A_2 B_2} \rho_{A_2 B_2}]} \quad (\text{J8})$$

Since the above inequality holds for all feasible solutions, we find that

$$F^u(\rho_{A_1 B_1} \otimes \rho_{A_2 B_2}) \leq F^u(\rho_{A_1 B_1}) F^u(\rho_{A_2 B_2}). \quad (\text{J9})$$

This concludes the proof.

Appendix K: Proof of Proposition 21

Since a maximally entangled state $\Phi_{AB} \equiv \Phi_{AB}^K$ necessarily has an extension of the form $\Phi_{AB} \otimes \sigma_E$ for some quantum state σ_E , the only possible extensions of $\gamma_{ABA'B'}$ have the following form:

$$\gamma_{ABEA'B'E'} = U_{ABA'B'} (\Phi_{AB} \otimes \sigma_{EA'B'E'}) U_{ABA'B'}^\dagger \quad (\text{K1})$$

$$= \frac{1}{d} \sum_{i,j} |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B \otimes U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{jj})^\dagger, \quad (\text{K2})$$

where $\sigma_{EA'B'E'}$ is an extension of $\sigma_{A'B'}$ such that $E \simeq B$ and $E' \simeq B'$. Observe that if we define a different twisting unitary $V_{ABA'B'}$ as

$$V_{ABA'B'} := \sum_{i,j} |i\rangle\langle i|_A \otimes |j\rangle\langle j|_B \otimes U_{A'B'}^{ii} \quad (\text{K3})$$

$$= \sum_i |i\rangle\langle i|_A \otimes \mathbb{1}_B \otimes U_{A'B'}^{ii}, \quad (\text{K4})$$

then

$$V_{ABA'B'} (\Phi_{AB} \otimes \sigma_{EA'B'E'}) V_{ABA'B'}^\dagger = \frac{1}{d} \sum_{i,j} |i\rangle\langle j|_A \otimes |i\rangle\langle j|_B \otimes U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{jj})^\dagger \quad (\text{K5})$$

$$= \gamma_{ABEA'B'E'}. \quad (\text{K6})$$

We have

$$\gamma_{AEA'E'} = \text{Tr}_{BB'}[\gamma_{ABEA'B'E'}] \quad (\text{K7})$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \text{Tr}_{B'}[U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger]. \quad (\text{K8})$$

Now consider that

$$\mathbf{D}(\gamma_{ABA'B'} \| \gamma_{AEA'E'}) = \mathbf{D}(V_{ABA'B'} (\Phi_{AB} \otimes \sigma_{A'B'}) V_{ABA'B'}^\dagger \| \gamma_{AEA'E'}) \quad (\text{K9})$$

$$= \mathbf{D}(\Phi_{AB} \otimes \sigma_{A'B'} \| V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'}) \quad (\text{K10})$$

$$\geq \mathbf{D}(\Phi_{AB} \| \text{Tr}_{A'E'}[V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'}]). \quad (\text{K11})$$

where the inequality follows from monotonicity under partial trace. Observe that

$$\begin{aligned} & V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'} \\ &= \left(\sum_{i'} |i'\rangle\langle i'|_A \otimes \mathbb{1}_E \otimes U_{A'E'}^{i'i'} \right) \left(\frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \text{Tr}_{B'}[U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] \right) \end{aligned}$$

$$\times \left(\sum_{i''} |i''\rangle\langle i''|_A \otimes \mathbb{1}_E \otimes U_{A'E'}^{i''i''} \right) \quad (\text{K12})$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes U_{A'E'}^{ii\dagger} \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] U_{A'E'}^{ii}, \quad (\text{K13})$$

so that

$$\begin{aligned} & \text{Tr}_{A'E'} [V_{AEA'E'}^\dagger \gamma_{AEA'E'} V_{AEA'E'}] \\ &= \text{Tr}_{A'E'} \left[\frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes U_{A'E'}^{ii\dagger} \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] U_{A'E'}^{ii} \right] \end{aligned} \quad (\text{K14})$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \text{Tr}_{A'E'} \left[U_{A'E'}^{ii\dagger} \text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] U_{A'E'}^{ii} \right] \quad (\text{K15})$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \text{Tr}_{A'E'} \left[\text{Tr}_{B'} [U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger] \right] \quad (\text{K16})$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \text{Tr}_{A'B'E'} \left[U_{A'B'}^{ii} \sigma_{EA'B'E'} (U_{A'B'}^{ii})^\dagger \right] \quad (\text{K17})$$

$$= \frac{1}{d} \sum_i |i\rangle\langle i|_A \otimes \sigma_E \quad (\text{K18})$$

$$= \pi_A \otimes \sigma_E. \quad (\text{K19})$$

So we find that

$$\mathbf{D}(\gamma_{ABA'B'} \| \gamma_{AEA'E'}) \geq \mathbf{D}(\Phi_{AB} \| \pi_A \otimes \sigma_E) \geq I_{\mathbf{D}}(A; B)_\Phi. \quad (\text{K20})$$

Since this bound holds for an arbitrary extension $\gamma_{ABEA'B'E'}$ of $\gamma_{ABA'B'}$, we conclude (88) after normalizing by $1/2$.

Appendix L: Proof of Theorem 23

We first show that an arbitrary generalized unextendible entanglement \mathbf{E}^u , which satisfies selective two-extendible monotonicity, normalization, and subadditivity, serves as a lower bound on the overhead. Let us suppose that the selective two-extendible operation Λ outputs $\gamma_{ABA'B'}^k$ from $\rho_{AB}^{\otimes n}$ with probability p . Without loss of generality, we can assume that

$$\Lambda(\rho_{AB}^{\otimes n}) = p|0\rangle\langle 0|_{X_A} \otimes \gamma_{ABA'B'}^k + (1-p)|1\rangle\langle 1|_{X_A} \otimes \tau, \quad (\text{L1})$$

where τ is some bipartite state and X_A is a flag system in Alice's possession indicating whether the conversion is successful. Since 1-LOCC operations are allowed for free, it is possible for Alice to communicate X_A to Bob for free. We have

$$n\mathbf{E}^u(\rho_{AB}) \geq \mathbf{E}^u(\rho_{AB}^{\otimes n}) \geq p\mathbf{E}^u(\gamma_{ABA'B'}^k) \geq pk, \quad (\text{L2})$$

where the first inequality follows from subadditivity, the second inequality follows from selective two-extendible monotonicity, and the last inequality follows from Corollary 22. Since n and p are arbitrary, we conclude that the following bound holds for every integer $n \geq 1$ and $p \in (0, 1]$:

$$\frac{n}{p} \geq \frac{k}{\mathbf{E}^u(\rho_{AB})}. \quad (\text{L3})$$

As the relative-entropy-induced unextendible entanglement $E^u(\rho_{AB})$ satisfies all of these required properties, we conclude the lower bound in (93).

Appendix M: Proof of Theorem 24

For a given bipartite state ρ_{AB} , suppose that there is a two-extendible channel $\Lambda_{AB \rightarrow \hat{A}\hat{B}}$ that transforms ρ_{AB} to a private state $\gamma_{\hat{A}\hat{B}A'B'}^k$ with k private bits. By the monotonicity of the min-unextendible entanglement, the following inequality holds

$$E_{\min}^u(\rho_{AB}) \geq E_{\min}^u(\Lambda_{AB \rightarrow \hat{A}\hat{B}}(\rho_{AB})) = E_{\min}^u(\gamma_{\hat{A}\hat{B}A'B'}^k) \geq k, \quad (\text{M1})$$

where the last inequality follows from Corollary 22. Therefore, by optimizing over all two-extendible protocols, it follows that the one-shot exact distillable entanglement of ρ_{AB} is bounded as

$$K_{2\text{-EXT}}^{(1)}(\rho_{AB}) \leq E_{\min}^u(\rho_{AB}). \quad (\text{M2})$$

Applying the same reasoning to the tensor-power state $\rho_{AB}^{\otimes n}$, we find that

$$K_{2\text{-EXT}}(\rho_{AB}) = \liminf_{n \rightarrow \infty} \frac{1}{n} K_{2\text{-EXT}}^{(1)}(\rho_{AB}^{\otimes n}) \leq \liminf_{n \rightarrow \infty} \frac{1}{n} E_{\min}^u(\rho_{AB}^{\otimes n}) = E_{\min}^u(\rho_{AB}), \quad (\text{M3})$$

where the final equality is a consequence of Proposition 15.

Appendix N: Proof of Proposition 27

Note that the erased state $\rho_{A'B}^\varepsilon$ can be written in the following direct-sum form:

$$\rho_{A'B}^\varepsilon = (1 - \varepsilon)\Phi_{AB}^2 \oplus \varepsilon\pi_B = \begin{bmatrix} (1 - \varepsilon)\Phi_{AB}^2 & 0 \\ 0 & \varepsilon\pi_B \end{bmatrix}. \quad (\text{N1})$$

From this fact, one can see that each extension $\rho_{A'BB'}$ of $\rho_{A'B}^\varepsilon$ has the form

$$\rho_{A'BB'} := \begin{bmatrix} (1 - \varepsilon)\sigma_{ABB'} & 0 \\ 0 & \varepsilon\sigma_{BB'} \end{bmatrix} \quad (\text{N2})$$

such that $\text{Tr}_{B'} \sigma_{ABB'} = \Phi_{AB}^2$ and $\text{Tr}_{B'} \sigma_{BB'} = \pi_B$. Furthermore, any extension of Φ_{AB}^2 necessarily has the form $\sigma_{ABB'} := \Phi_{AB}^2 \otimes \tau_{B'}$, where $\tau_{B'}$ is a state on system B' . Thus

$$E^u(\rho_{A'B}^\varepsilon) = \min_{\rho_{A'BB'} \text{ st. Eq. (N2)}} \frac{1}{2} D(\rho_{A'B}^\varepsilon \| \rho_{A'B'}) \quad (\text{N3})$$

$$= \min_{\tau_{B'}, \sigma_{B'}} \frac{1}{2} D \left(\begin{bmatrix} (1 - \varepsilon)\Phi_{AB}^2 & 0 \\ 0 & \varepsilon\pi_B \end{bmatrix} \middle\| \begin{bmatrix} (1 - \varepsilon)\pi_A \otimes \tau_{B'} & 0 \\ 0 & \varepsilon\sigma_{B'} \end{bmatrix} \right) \quad (\text{N4})$$

$$= (1 - \varepsilon) \min_{\tau_{B'}} \frac{1}{2} D(\Phi_{AB}^2 \| \pi_A \otimes \tau_{B'}) + \varepsilon \inf_{\sigma_{B'}} \frac{1}{2} D(\pi_B \| \sigma_{B'}) \quad (\text{N5})$$

$$= (1 - \varepsilon) \frac{1}{2} I(A; B)_\Phi \quad (\text{N6})$$

$$= (1 - \varepsilon), \quad (\text{N7})$$

where $I(A; B)_\rho$ is the quantum mutual information of ρ_{AB} .

Appendix O: Proof of Proposition 29

Since $1/E_R(\rho)$ is a lower bound on the overhead of distillation using selective LOCC, it follows from the optimality of erased state distillation under one-way LOCC that $1/(1-\varepsilon) \geq 1/E_R(\rho_{A'B}^\varepsilon)$. This gives $E_R(\rho_{A'B}^\varepsilon) \geq 1-\varepsilon$.

It then suffices to construct a feasible state to achieve the lower bound, and here we follow an approach similar to that from [TWW16, Proposition 11]. Consider the following state

$$\sigma_{A'B} := \frac{1-\varepsilon}{2} (|00\rangle\langle 00| + |11\rangle\langle 11|) + \varepsilon|e\rangle\langle e| \otimes \pi_B. \quad (\text{O1})$$

The state $\sigma_{A'B}$ is a separable state. It then holds that

$$E_R(\rho_{A'B}^\varepsilon) \leq D(\rho_{A'B}^\varepsilon \| \sigma_{A'B}) \quad (\text{O2})$$

$$= D \left((1-\varepsilon)\Phi_{AB}^2 \oplus \varepsilon\pi_B \left\| \frac{1-\varepsilon}{2} (|00\rangle\langle 00| + |11\rangle\langle 11|) \oplus \varepsilon\pi_B \right. \right) \quad (\text{O3})$$

$$= (1-\varepsilon)D(\Phi_{AB}^2 \| (|00\rangle\langle 00| + |11\rangle\langle 11|)/2) \quad (\text{O4})$$

$$= 1-\varepsilon. \quad (\text{O5})$$

This concludes the proof.