

THE PICARD GROUPS OF INCLUSIONS OF C^* -ALGEBRAS INDUCED BY EQUIVALENCE BIMODULES

KAZUNORI KODAKA

ABSTRACT. Let A and B be σ -unital C^* -algebras and X and Y an $A - A$ -equivalence bimodule and a $B - B$ -equivalence bimodule, respectively. Also, let $A \rtimes_X \mathbf{Z}$ and $B \rtimes_Y \mathbf{Z}$ be the crossed products of A and B by X and Y , respectively. Furthermore, let $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ be the inclusions of C^* -algebras induced by X and Y , respectively. We suppose that $A' \cap M(A \rtimes_X \mathbf{Z}) = \mathbf{C}1$. In this paper we shall show that the inclusions $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent if and only if there is an $A - B$ -equivalence bimodule M such that $Y \cong \widetilde{M} \otimes_A X \otimes_A M$ or $\widetilde{Y} \cong \widetilde{M} \otimes_A X \otimes_A M$ as $B - B$ -equivalence bimodules, where \widetilde{M} and \widetilde{Y} are the dual $B - A$ -equivalence bimodule and the dual $B - B$ -equivalence bimodule of M and Y , respectively. Applying this result, we shall compute the Picard group of the inclusion $A \subset A \rtimes_X \mathbf{Z}$ under the assumption that $A' \cap M(A \rtimes_X \mathbf{Z}) = \mathbf{C}1$.

1. INTRODUCTION

In the previous paper [11], we discussed strong Morita equivalence for unital inclusions of unital C^* -algebras induced by involutive equivalence bimodules. That is, let A and B be unital C^* -algebras and X and Y an involutive $A - A$ -equivalence bimodule and an involutive $B - B$ -equivalence bimodule, respectively. Let C_X and C_Y be unital C^* -algebras induced by X and Y and $A \subset C_X$ and $B \subset C_Y$ the unital inclusions of unital C^* -algebras. We suppose that $A' \cap C_X = \mathbf{C}1$. In the paper [11], we showed that $A \subset C_X$ and $B \subset C_Y$ are strongly Morita equivalent if and only if there is an $A - B$ -equivalence bimodule M such that $Y \cong \widetilde{M} \otimes_A X \otimes_A M$ as $B - B$ -equivalence bimodules. In the present paper, we shall show the same result as above in the case of inclusions of C^* -algebras induced by σ -unital C^* -algebra equivalence bimodules.

Let A and B be σ -unital C^* -algebras and X and Y an $A - A$ -equivalence bimodule and a $B - B$ -equivalence bimodule, respectively. Let $A \rtimes_X \mathbf{Z}$ and $B \rtimes_Y \mathbf{Z}$ be the crossed products of A and B by X and Y , respectively, which are defined in Abadie, Eilers and Exel [1]. Then we have inclusions of C^* -algebras $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$. We note that $\overline{A(A \rtimes_X \mathbf{Z})} = A \rtimes_X \mathbf{Z}$ and $\overline{B(B \rtimes_Y \mathbf{Z})} = B \rtimes_Y \mathbf{Z}$. We suppose that $A' \cap M(A \rtimes_X \mathbf{Z}) = \mathbf{C}1$. We shall show that $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent if and only if there is an $A - B$ -equivalence bimodule M such that $Y \cong \widetilde{M} \otimes_A X \otimes_A M$ or $\widetilde{Y} \cong \widetilde{M} \otimes_A X \otimes_A M$ as $B - B$ -equivalence bimodules, where \widetilde{M} and \widetilde{Y} are the dual $B - A$ -equivalence bimodule and the dual $B - B$ -equivalence bimodule of M and Y , respectively. Applying the above result, we shall compute the Picard group of the inclusion of C^* -algebras $A \subset A \rtimes_X \mathbf{Z}$ under the assumption that $A' \cap M(A \rtimes_X \mathbf{Z}) = \mathbf{C}1$.

2010 *Mathematics Subject Classification.* 46L05.

Key words and phrases. crossed products, equivalence bimodules, inclusions of C^* -algebras, strong Morita equivalence, Picard groups.

2. PRELIMINARIES

Let \mathbf{K} be the C^* -algebra of all compact operators on a countably infinite dimensional Hilbert space and $\{e_{ij}\}_{i,j \in \mathbf{N}}$ its system of matrix units.

For each C^* -algebra A , we denote by $M(A)$ the multiplier C^* -algebra of A . Let π be an isomorphism of A onto a C^* -algebra B . Then there is the unique strictly continuous isomorphism of $M(A)$ onto $M(B)$ extending π by Jensen and Thomsen [7, Corollary 1.1.15]. We denote it by $\underline{\pi}$.

Let A and B be C^* -algebras and X an $A - B$ -bimodule. We denote its left A -action and right B -action on X by $a \cdot x$ and $x \cdot b$ for any $a \in A$, $b \in B$, $x \in X$, respectively. We denote by \tilde{X} the dual $B - A$ -bimodule of X and we denote by \tilde{x} the element in \tilde{X} induced by $x \in X$. Let $A \subset C$ and $B \subset D$ be inclusions of C^* -algebras.

Definition 2.1. We say that $A \subset C$ and $B \subset D$ are *isomorphic as inclusions of C^* -algebras* if there is an isomorphism π of C onto D such that $\pi|_A$ is an isomorphism of A onto B .

Also, we give the following definition:

Definition 2.2. Let α and β be actions of a discrete group G on A and B , respectively. We say that α and β are *strongly Morita equivalent with respect to (X, λ)* if there are an $A - B$ -equivalence bimodule X and a linear automorphism λ of X satisfying the following:

- (1) $\alpha_t(A \langle x, y \rangle) = A \langle \lambda_t(x), \lambda_t(y) \rangle$,
 - (2) $\beta_t(\langle x, y \rangle_B) = \langle \lambda_t(x), \lambda_t(y) \rangle_B$
- for any $x, y \in X$, $t \in G$.

Then we have the following:

$$\lambda_t(a \cdot x) = \alpha_t(a) \cdot \lambda_t(x), \quad \lambda_t(x \cdot b) = \lambda_t(x) \cdot \beta_t(b)$$

for any $a \in A$, $b \in B$, $x \in X$ and $t \in G$.

Let A and B be C^* -algebras and π an isomorphism of B onto A . We construct an $A - B$ -equivalence bimodule X_π as follows: Let $X_\pi = A$ as \mathbf{C} -vector spaces. For any $a \in A$, $b \in B$, $x, y \in X_\pi$,

$$\begin{aligned} a \cdot x &= ax, & x \cdot b &= x\pi(b), \\ A \langle x, y \rangle &= xy^*, & \langle x, y \rangle_B &= \pi^{-1}(x^*y). \end{aligned}$$

By easy computations, we can see that X_π is an $A - B$ -equivalence bimodule. We call X_π an $A - B$ -equivalence bimodule induced by π . Let α be an automorphism of A . Then in the same way as above, we construct X_α , as $A - B$ -equivalence bimodule. Let u_α be a unitary element in $M(A \rtimes_\alpha \mathbf{Z})$ implementing α . Hence $\alpha = \text{Ad}(u_\alpha)$. We regard Au_α as an $A - A$ -equivalence bimodule as follows:

$$\begin{aligned} a \cdot xu_\alpha &= axu_\alpha, & xu_\alpha \cdot a &= x\alpha(a), \\ A \langle xu_\alpha, yu_\alpha \rangle &, & \langle xu_\alpha, yu_\alpha \rangle_A &= \alpha^{-1}(x^*y) \end{aligned}$$

for any $a, x, y \in A$.

Lemma 2.1. *With the above notation, $X_\alpha \cong Au_\alpha$ as $A - A$ -equivalence bimodules.*

Proof. This is immediate by easy computations. □

Let A be a C^* -algebra and X an $A - A$ -equivalence bimodule. Let $A \rtimes_X \mathbf{Z}$ be the crossed product of A by X defined in [1]. We regard the C^* -algebra \mathbf{K} as the trivial $\mathbf{K} - \mathbf{K}$ -equivalence bimodule. Then we obtain an $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodule $X \otimes \mathbf{K}$ and we can also consider the crossed product

$$(A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$$

of $A \otimes \mathbf{K}$ by $X \otimes \mathbf{K}$. Hence we have the following inclusions of C^* -algebras:

$$A \subset A \rtimes_X \mathbf{Z}, \quad A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}.$$

Since there is an isomorphism π of $(A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ onto $(A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}$ such that $\pi|_{A \otimes \mathbf{K}} = \text{id}$ on $A \otimes \mathbf{K}$, we identify $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ with $A \otimes \mathbf{K} \subset (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}$. Thus $A \subset A \rtimes_X \mathbf{Z}$ and $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ are strongly Morita equivalent. Let H_A be the $A \otimes \mathbf{K} - A$ -equivalence bimodule defined as follows: Let $H_A = (A \otimes \mathbf{K})(1_{M(A)} \otimes e_{11})$ as \mathbf{C} -vector spaces. For any $a \in A$, $k \in \mathbf{K}$, $x, y \in A \otimes \mathbf{K}$,

$$\begin{aligned} (a \otimes k) \cdot x(1 \otimes e_{11}) &= (a \otimes k)x(1 \otimes e_{11}), \\ x(1 \otimes e_{11}) \cdot a &= x(a \otimes e_{11}), \\ A \otimes \mathbf{K} \langle x(1 \otimes e_{11}), y(1 \otimes e_{11}) \rangle &= x(1 \otimes e_{11})y^*, \\ \langle x(1 \otimes e_{11}), y(1 \otimes e_{11}) \rangle_A &= (1 \otimes e_{11})x^*y(1 \otimes e_{11}), \end{aligned}$$

where we identify A with $A \otimes e_{11}$. Let B be a C^* -algebra. Let H_B be as above.

Lemma 2.2. *With the above notation, let M be an $A - B$ -equivalence bimodule. Then*

$$(1 \otimes e_{11}) \cdot (M \otimes \mathbf{K}) \cdot (1 \otimes e_{11}) \cong M$$

as $A - B$ -equivalence bimodules.

Proof. Since the linear span of the set

$$\{x \otimes e_{ij} \mid x \in M, i, j \in \mathbf{N}\}$$

is dense in $M \otimes \mathbf{K}$, $(1 \otimes e_{11}) \cdot (M \otimes \mathbf{K}) \cdot (1 \otimes e_{11}) \cong M \otimes e_{11}$ as $A - B$ -equivalence bimodules. Hence

$$(1 \otimes e_{11}) \cdot (M \otimes \mathbf{K}) \cdot (1 \otimes e_{11}) \cong M$$

as $A - B$ -equivalence bimodules. \square

Lemma 2.3. *With the above notation, let M be an $A - B$ -equivalence bimodule. Then*

$$\widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B \cong M$$

as $A - B$ -equivalence bimodules.

Proof. Let π be the map from $\widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B$ to $(1 \otimes e_{11}) \cdot (M \otimes \mathbf{K}) \cdot (1 \otimes e_{11})$ defined by

$$\pi([a(1 \otimes e_{11})] \otimes x \otimes b(1 \otimes e_{11})) = (1 \otimes e_{11}) \cdot (a^* \cdot x \cdot b) \cdot (1 \otimes e_{11})$$

for any $a \in A \otimes \mathbf{K}$, $b \in B \otimes \mathbf{K}$, $x \in M \otimes \mathbf{K}$. Then by easy computations, π is an $A - B$ -equivalence bimodule isomorphism of $\widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B$ onto $(1 \otimes e_{11}) \cdot (M \otimes \mathbf{K}) \cdot (1 \otimes e_{11})$. Thus by Lemma 2.2

$$\widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B \cong M$$

as $A - B$ -equivalence bimodules. \square

We prepare the following lemma which is applied in the next section.

Lemma 2.4. *Let A and B be C^* -algebras and X and Y an $A - A$ -equivalence bimodule and $B - B$ -equivalence bimodule, respectively. Let $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ be inclusions of C^* -algebras induced by X and Y , respectively. We suppose that there is an $A - B$ -equivalence bimodule M such that $Y \cong \widetilde{M} \otimes_A X \otimes_A M$ or $\widetilde{Y} \cong \widetilde{M} \otimes_A X \otimes_A M$ as $B - B$ -equivalence bimodules. Then there is an $A \rtimes_X \mathbf{Z} - B \rtimes_Y \mathbf{Z}$ -equivalence bimodule N satisfying the following:*

- (1) M is included in N as a closed subspace,
- (2) $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent with respect to N and its closed subspace M .

Proof. Modifying the proof of [1, Theorem 4.2], we prove this lemma. We suppose that $Y \cong \widetilde{M} \otimes_A X \otimes_A M$ as $B - B$ -equivalence bimodules. Let L_M be the linking C^* -algebra for M defined by

$$L_M = \begin{bmatrix} A & M \\ \widetilde{M} & B \end{bmatrix}.$$

Also, let W be the $L_M - L_M$ -equivalence bimodule defined in the proof of [1, Theorem 4.2], which is defined by

$$W = \begin{bmatrix} X & X \otimes_A M \\ Y \otimes_B \widetilde{M} & Y \end{bmatrix}.$$

Let $L_M \rtimes_W \mathbf{Z}$ be the crossed product of L_M by W and let

$$p = \begin{bmatrix} 1_{M(A)} & 0 \\ 0 & 0 \end{bmatrix}, \quad q = \begin{bmatrix} 0 & 0 \\ 0 & 1_{M(B)} \end{bmatrix}.$$

Furthermore, let $N = p(L_M \rtimes_W \mathbf{Z})q$. Then since $M = pL_Mq$, M is a closed subspace of N . Hence by the proof of [1, Theorem 4.2], $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent with respect to N and its closed subspace M .

Next, we suppose that $\widetilde{Y} \cong \widetilde{M} \otimes_A X \otimes_A M$. Let

$$W_0 = \begin{bmatrix} X & X \otimes_A M \\ \widetilde{Y} \otimes_B \widetilde{M} & \widetilde{Y} \end{bmatrix}.$$

Then W_0 is an $L_M - L_M$ -equivalence bimodule. Let $N_0 = p(L_M \rtimes_{W_0} \mathbf{Z})q$. By the above discussions, $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_{\widetilde{Y}} \mathbf{Z}$ are strongly Morita equivalent with respect to N_0 and its closed subspace M . On the other hand, there is an isomorphism π of $B \rtimes_Y \mathbf{Z}$ onto $B \rtimes_{\widetilde{Y}} \mathbf{Z}$ such that $\pi|_B = \text{id}$ on B . Let X_π be the $B \rtimes_{\widetilde{Y}} \mathbf{Z} - B \rtimes_Y \mathbf{Z}$ -equivalence bimodule induced by π . Then B is a closed subspace of X_π and we regard B as the trivial $B - B$ -equivalence bimodule since $\pi|_B = \text{id}$ on B . Thus $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent with respect to $N_0 \otimes_{B \rtimes_{\widetilde{Y}} \mathbf{Z}} X_\pi$ and its closed subspace $M \otimes_B B (\cong M)$. Therefore, we obtain the conclusion. \square

Lemma 2.5. *With the above notation, we suppose that A is a σ -unital C^* -algebra. Then there is an automorphism α of $A \otimes \mathbf{K}$ such that $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}$ is isomorphic to $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ as inclusions of C^* -algebras.*

Proof. Since A is σ -unital, by Brown, Green and Rieffel [3, Corollary 3.5], there is an automorphism α of $A \otimes \mathbf{K}$ such that $X \otimes \mathbf{K} \cong X_\alpha$ as $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodules, where X_α is the $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodule induced by α . Let u_α be a unitary element in $M((A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$ implementing α . We regard $(A \otimes \mathbf{K})u_\alpha$ as an $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodule as above. Then by Lemma 2.1, $X_\alpha \cong (A \otimes \mathbf{K})u_\alpha$ as $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodules. Let $(A \otimes \mathbf{K}) \rtimes_{(A \otimes \mathbf{K})u_\alpha} \mathbf{Z}$ be the crossed product of $A \otimes \mathbf{K}$ by $(A \otimes \mathbf{K})u_\alpha$. Then by the definition of the crossed product of a C^* -algebra by an equivalence bimodule, we can see that

$$(A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z} \cong (A \otimes \mathbf{K}) \rtimes_{(A \otimes \mathbf{K})u_\alpha} \mathbf{Z}$$

as C^* -algebras. Since $X_\alpha \cong X \otimes \mathbf{K}$ as $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodules, we obtain that

$$(A \otimes \mathbf{K}) \rtimes_{(A \otimes \mathbf{K})u_\alpha} \mathbf{Z} \cong (A \otimes \mathbf{K}) \rtimes_{X_\alpha} \mathbf{Z} \cong (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$$

as C^* -algebras. Since the above isomorphisms leave any element in $A \otimes \mathbf{K}$ invariant, we can see that $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}$ is isomorphic to $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ as inclusions of C^* -algebras. \square

3. STRONG MORITA EQUIVALENCE

Let A and B be σ -unital C^* -algebras and X and Y an A – A -equivalence bimodule and a B – B -equivalence bimodule, respectively. Let $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ be the inclusions of C^* -algebras induced by X and Y , respectively. We suppose that $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent with respect to an $A \rtimes_X \mathbf{Z} - B \rtimes_Y \mathbf{Z}$ -equivalence bimodule N and its closed subspace M . We suppose that $A' \cap M(A \rtimes_X \mathbf{Z}) = \mathbf{C}1$. Then since the inclusion $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ is isomorphic to the inclusion $A \otimes \mathbf{K} \subset (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}$ as inclusions of C^* -algebras, by [12, Lemma 3.1],

$$(A \otimes \mathbf{K})' \cap ((A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}) = \mathbf{C}1.$$

Also, by the above assumptions, the inclusion $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ is strongly Morita equivalent to the inclusion $B \otimes \mathbf{K} \subset (B \otimes \mathbf{K}) \rtimes_{Y \otimes \mathbf{K}} \mathbf{Z}$ with respect to the $(A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z} - (B \otimes \mathbf{K}) \rtimes_{Y \otimes \mathbf{K}} \mathbf{Z}$ -equivalence bimodule $N \otimes \mathbf{K}$ and its closed subspace $M \otimes \mathbf{K}$. By Lemma 2.5, there are an automorphism α of $A \otimes \mathbf{K}$ and an automorphism β of $B \otimes \mathbf{K}$ such that $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ and $B \otimes \mathbf{K} \subset (B \otimes \mathbf{K}) \rtimes_{Y \otimes \mathbf{K}} \mathbf{Z}$ are isomorphic to $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{\alpha} \mathbf{Z}$ and $B \otimes \mathbf{K} \subset (B \otimes \mathbf{K}) \rtimes_{\beta} \mathbf{Z}$ as inclusions of C^* -algebras, respectively. Hence we can assume that $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{\alpha} \mathbf{Z}$ and $B \otimes \mathbf{K} \subset (B \otimes \mathbf{K}) \rtimes_{\beta} \mathbf{Z}$ are strongly Morita equivalent with respect to an $(A \otimes \mathbf{K}) \rtimes_{\alpha} \mathbf{Z} - (B \otimes \mathbf{K}) \rtimes_{\beta} \mathbf{Z}$ -equivalence bimodule $N \otimes \mathbf{K}$ and its closed subspace $M \otimes \mathbf{K}$. Since A and B are σ -unital, in the same way as in the proof of [10, Proposition 3.5] or [3, Proposition 3.1], there is an isomorphism θ of $(B \otimes \mathbf{K}) \rtimes_{\beta} \mathbf{Z}$ onto $(A \otimes \mathbf{K}) \rtimes_{\alpha} \mathbf{Z}$ satisfying the following:

- (1) $\theta|_{B \otimes \mathbf{K}}$ is an isomorphism of $B \otimes \mathbf{K}$ onto $A \otimes \mathbf{K}$,
- (2) There is an $(A \otimes \mathbf{K}) \rtimes_{\alpha} \mathbf{Z} - (B \otimes \mathbf{K}) \rtimes_{\beta} \mathbf{Z}$ -equivalence bimodule isomorphism Φ of $N \otimes \mathbf{K}$ onto Y_{θ} such that $\Phi|_{M \otimes \mathbf{K}}$ is an $A \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodule isomorphism of $M \otimes \mathbf{K}$ onto X_{θ} , where Y_{θ} is the $(A \otimes \mathbf{K}) \rtimes_{\alpha} \mathbf{Z} - (B \otimes \mathbf{K}) \rtimes_{\beta} \mathbf{Z}$ -equivalence bimodule induced by θ and X_{θ} is the $A \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodule induced by $\theta|_{B \otimes \mathbf{K}}$.

Let

$$\gamma = \theta|_{B \otimes \mathbf{K}} \circ \beta \circ \theta|_{B \otimes \mathbf{K}}^{-1}$$

and let λ be the linear automorphism of X_{θ} defined by $\lambda(x) = \gamma(x)$ for any $x \in X_{\theta}(= A \otimes \mathbf{K})$.

Lemma 3.1. *With the above notation, γ and β are strongly Morita equivalent with respect to (X_{θ}, λ) .*

Proof. For any $x, y \in X_{\theta}$,

$$\begin{aligned} A \otimes \mathbf{K} \langle \lambda(x), \lambda(y) \rangle &= \gamma(xy^*) = \gamma(A \otimes \mathbf{K} \langle x, y \rangle), \\ \langle \lambda(x), \lambda(y) \rangle_{B \otimes \mathbf{K}} &= \theta|_{B \otimes \mathbf{K}}^{-1}(\gamma(x^*y)) = \beta(\theta|_{B \otimes \mathbf{K}}^{-1}(x^*y)) = \beta(\langle x, y \rangle_{B \otimes \mathbf{K}}). \end{aligned}$$

Hence γ and β are strongly Morita equivalent with respect to (X_{θ}, λ) . \square

By the proof of [12, Theorem 5.5], there is an automorphism ϕ of \mathbf{Z} satisfying that γ^{ϕ} and α are exterior equivalent, that is, there is a unitary element $z \in M(A \otimes \mathbf{K})$ such that

$$\gamma^{\phi} = \text{Ad}(z) \circ \alpha, \quad \underline{\alpha}(z) = z,$$

where γ^{ϕ} is the automorphism of $A \otimes \mathbf{K}$ induced by γ and ϕ , that is, γ^{ϕ} is defined by $\gamma^{\phi} = \gamma^{\phi(1)}$. We note that $\gamma^{\phi} = \gamma$ or $\gamma^{\phi} = \gamma^{-1}$. We regard $A \otimes \mathbf{K}$ as the trivial $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodule. Let μ be the linear automorphism of $A \otimes \mathbf{K}$ defined by

$$\mu(x) = \alpha(x)z^*$$

for any $x \in A \otimes \mathbf{K}$.

Lemma 3.2. *With the above notation, α and γ^ϕ are strongly Morita equivalent with respect to $(A \otimes \mathbf{K}, \mu)$.*

Proof. For any $x, y \in A \otimes \mathbf{K}$,

$$\begin{aligned} A \otimes \mathbf{K} \langle \mu(x), \mu(y) \rangle &= A \otimes \mathbf{K} \langle \alpha(x)z^*, \alpha(y)z^* \rangle = \alpha(xy^*) = \alpha(A \otimes \mathbf{K} \langle x, y \rangle), \\ \langle \mu(x), \mu(y) \rangle_{A \otimes \mathbf{K}} &= z\alpha(x^*y)z^* = \gamma^\phi(x^*y) = \gamma^\phi(\langle x, y \rangle_{A \otimes \mathbf{K}}). \end{aligned}$$

Therefore, we obtain the conclusion. \square

Let ν be the linear automorphism of X_θ defined by

$$\nu(x) = \gamma^\phi(z^*x)$$

for any $x \in X_\theta (= A \otimes \mathbf{K})$.

Lemma 3.3. *With the above notation, α and β^ϕ are strongly Morita equivalent with respect to (X_θ, ν) , where β^ϕ is the automorphism of $B \otimes \mathbf{K}$ induced by β and ϕ .*

Proof. For any $x, y \in X_\theta$,

$$\begin{aligned} A \otimes \mathbf{K} \langle \nu(x), \nu(y) \rangle &= A \otimes \mathbf{K} \langle \gamma^\phi(z^*x), \gamma^\phi(z^*y) \rangle = \gamma^\phi(z^*xy^*z) = z\alpha(z^*xy^*z)z^* \\ &= \alpha(xy^*) = \alpha(A \otimes \mathbf{K} \langle x, y \rangle), \\ \langle \nu(x), \nu(y) \rangle_{B \otimes \mathbf{K}} &= \langle \gamma^\phi(z^*x), \gamma^\phi(z^*y) \rangle_{B \otimes \mathbf{K}} = \theta|_{B \otimes \mathbf{K}}^{-1}(\gamma^\phi(x^*y)) = \beta^\phi(\theta|_{B \otimes \mathbf{K}}^{-1}(x^*y)) \\ &= \beta^\phi(\langle x, y \rangle_{B \otimes \mathbf{K}}). \end{aligned}$$

Therefore, we obtain the conclusion. \square

Since $\beta^\phi = \beta$ or $\beta^\phi = \beta^{-1}$, by Lemma 3.3, α is strongly Morita equivalent to β or β^{-1} .

(I) We suppose that α is strongly Morita equivalent to β . Then by Lemma 3.3, there is the linear automorphism ν of X_θ satisfying the following:

- (1) $\nu(a \cdot x) = \alpha(a) \cdot \nu(x)$,
 - (2) $\nu(x \cdot b) = \nu(x) \cdot \beta(b)$,
 - (3) $A \otimes \mathbf{K} \langle \nu(x), \nu(y) \rangle = \alpha(A \otimes \mathbf{K} \langle x, y \rangle)$,
 - (4) $\langle \nu(x), \nu(y) \rangle_{B \otimes \mathbf{K}} = \beta(\langle x, y \rangle_{B \otimes \mathbf{K}})$,
- for any $a \in A \otimes \mathbf{K}$, $b \in B \otimes \mathbf{K}$, $x, y \in X_\theta$.

Lemma 3.4. *With the above notation and assumptions, let X_α and X_β be the $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodule and the $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodule induced by α and β , respectively. Then*

$$X_\beta \cong \widetilde{X_\theta} \otimes_{A \otimes \mathbf{K}} X_\alpha \otimes_{A \otimes \mathbf{K}} X_\theta$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules.

Proof. Let Ψ be the map from $\widetilde{X_\theta} \otimes_{A \otimes \mathbf{K}} X_\alpha \otimes_{A \otimes \mathbf{K}} X_\theta$ to X_β defined by

$$\Psi(\widetilde{x} \otimes a \otimes y) = \langle x, a \cdot \nu(y) \rangle_{B \otimes \mathbf{K}}$$

for any $x, y \in X_\theta$, $a \in X_\alpha$. Then for any $x, x_1, y, y_1 \in X_\theta$, $a, a_1 \in X_\alpha$,

$$\begin{aligned} B \otimes \mathbf{K} \langle \widetilde{x} \otimes a \otimes y, \widetilde{x}_1 \otimes a_1 \otimes y_1 \rangle &= B \otimes \mathbf{K} \langle \widetilde{x} \cdot A \otimes \mathbf{K} \langle a \otimes y, a_1 \otimes y_1 \rangle, \widetilde{x}_1 \rangle \\ &= \langle A \otimes \mathbf{K} \langle a_1 \otimes y_1, a \otimes y \rangle \cdot x, x_1 \rangle_{B \otimes \mathbf{K}} \\ &= \langle A \otimes \mathbf{K} \langle a_1 \cdot A \otimes \mathbf{K} \langle y_1, y \rangle, a \rangle \cdot x, x_1 \rangle_{B \otimes \mathbf{K}} \\ &= \langle A \otimes \mathbf{K} \langle a_1 \alpha(A \otimes \mathbf{K} \langle y_1, y \rangle), a \rangle \cdot x, x_1 \rangle_{B \otimes \mathbf{K}} \\ &= \langle a_1 \alpha(A \otimes \mathbf{K} \langle y_1, y \rangle) a^* \cdot x, x_1 \rangle_{B \otimes \mathbf{K}}. \end{aligned}$$

On the other hand,

$$\begin{aligned}
_{B \otimes \mathbf{K}} \langle \Psi(\widetilde{x} \otimes a \otimes y), \Psi(\widetilde{x}_1 \otimes a_1 \otimes y_1) \rangle &= _{B \otimes \mathbf{K}} \langle \langle x, a \cdot \nu(y) \rangle_{B \otimes \mathbf{K}}, \langle x_1, a_1 \cdot \nu(y_1) \rangle_{B \otimes \mathbf{K}} \rangle \\
&= \langle x, a \cdot \nu(y) \rangle_{B \otimes \mathbf{K}} \langle a_1 \cdot \nu(y_1), x_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle x, a \cdot \nu(y) \cdot \langle a_1 \cdot \nu(y_1), x_1 \rangle_{B \otimes \mathbf{K}} \rangle_{B \otimes \mathbf{K}} \\
&= \langle x, _{A \otimes \mathbf{K}} \langle a \cdot \nu(y), a_1 \cdot \nu(y_1) \rangle \cdot x_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle x, a \cdot _{A \otimes \mathbf{K}} \langle \nu(y), \nu(y_1) \rangle a_1^* \cdot x_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle a_1 \cdot _{A \otimes \mathbf{K}} \langle \nu(y_1), \nu(y) \rangle a^* \cdot x, x_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle a_1 \alpha(_{A \otimes \mathbf{K}} \langle y_1, y \rangle) a^* \cdot x, x_1 \rangle_{B \otimes \mathbf{K}}.
\end{aligned}$$

Hence the map Ψ preserves the left $B \otimes \mathbf{K}$ -valued inner products. Also,

$$\begin{aligned}
\langle \widetilde{x} \otimes a \otimes y, \widetilde{x}_1 \otimes a_1 \otimes y_1 \rangle_{B \otimes \mathbf{K}} &= \langle y, \langle \widetilde{x} \otimes a, \widetilde{x}_1 \otimes a_1 \rangle_{A \otimes \mathbf{K}} \cdot y_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle y \langle a, _{A \otimes \mathbf{K}} \langle x, x_1 \rangle \cdot a_1 \rangle_{A \otimes \mathbf{K}} \cdot y_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle y \langle a, _{A \otimes \mathbf{K}} \langle x, x_1 \rangle a_1 \rangle_{A \otimes \mathbf{K}} \cdot y_1 \rangle_{B \otimes \mathbf{K}} \\
&= \langle y, \alpha^{-1}(a^* \cdot _{A \otimes \mathbf{K}} \langle x, x_1 \rangle a_1) \cdot y_1 \rangle_{B \otimes \mathbf{K}}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\langle \Psi(\widetilde{x} \otimes a \otimes y), \Psi(\widetilde{x}_1 \otimes a_1 \otimes y_1) \rangle_{B \otimes \mathbf{K}} &= \langle \langle x, a \cdot \nu(y) \rangle_{B \otimes \mathbf{K}}, \langle x_1, a_1 \cdot \nu(y_1) \rangle_{B \otimes \mathbf{K}} \rangle_{B \otimes \mathbf{K}} \\
&= \beta^{-1}(\langle a \cdot \nu(y), x \rangle_{B \otimes \mathbf{K}} \langle x_1, a_1 \cdot \nu(y_1) \rangle_{B \otimes \mathbf{K}}) \\
&= \beta^{-1}(\langle a \cdot \nu(y), x \cdot \langle x_1, a_1 \cdot \nu(y_1) \rangle_{B \otimes \mathbf{K}} \rangle_{B \otimes \mathbf{K}}) \\
&= \beta^{-1}(\langle a \cdot \nu(y), _{A \otimes \mathbf{K}} \langle x, x_1 \rangle a_1 \cdot \nu(y_1) \rangle_{B \otimes \mathbf{K}}) \\
&= \beta^{-1}(\langle \nu(y), a^* \cdot _{A \otimes \mathbf{K}} \langle x, x_1 \rangle a_1 \cdot \nu(y_1) \rangle_{B \otimes \mathbf{K}}) \\
&= \langle y, \alpha^{-1}(a^* \cdot _{A \otimes \mathbf{K}} \langle x, x_1 \rangle a_1) \cdot y_1 \rangle_{B \otimes \mathbf{K}}.
\end{aligned}$$

Hence the map Ψ preserves the right $B \otimes \mathbf{K}$ -valued inner products. Therefore, we obtain the conclusion. \square

(II) We suppose that α is strongly Morita equivalent to β^{-1} . Then by Lemma 3.4,

$$X_{\beta^{-1}} \cong \widetilde{X_\theta} \otimes_{A \otimes \mathbf{K}} X_\alpha \otimes_{A \otimes \mathbf{K}} X_\theta$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules. Thus we obtain the following:

Lemma 3.5. *With the above notation and assumptions,*

$$\widetilde{X_\beta} \cong X_{\beta^{-1}} \cong \widetilde{X_\theta} \otimes_{A \otimes \mathbf{K}} X_\alpha \otimes_{A \otimes \mathbf{K}} X_\theta$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules.

We recall that there is an $(A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z} - (B \otimes \mathbf{K}) \rtimes_\beta \mathbf{Z}$ -equivalence bimodule isomorphism Φ of $N \otimes \mathbf{K}$ onto Y_θ such that $\Phi|_{M \otimes \mathbf{K}}$ is an $A \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodule isomorphism of $M \otimes \mathbf{K}$ onto X_θ . We identify $M \otimes \mathbf{K}$ with X_θ by $\Phi|_{M \otimes \mathbf{K}}$. Then

$$X_\beta \cong (\widetilde{M \otimes \mathbf{K}}) \otimes_{A \otimes \mathbf{K}} X_\alpha \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K})$$

or

$$\widetilde{X_\beta} \cong (\widetilde{M \otimes \mathbf{K}}) \otimes_{A \otimes \mathbf{K}} X_\alpha \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K})$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules. Also, we recall that $X_\alpha \cong X \otimes \mathbf{K}$ as $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodules and that $X_\beta \cong Y \otimes \mathbf{K}$ as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules. Thus

$$Y \otimes \mathbf{K} \cong (\widetilde{M \otimes \mathbf{K}}) \otimes_{A \otimes \mathbf{K}} (X \otimes \mathbf{K}) \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K})$$

or

$$\widetilde{Y \otimes \mathbf{K}} \cong (\widetilde{M \otimes \mathbf{K}}) \otimes_{A \otimes \mathbf{K}} (X \otimes \mathbf{K}) \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K})$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules. Furthermore, by Lemma 2.3

$$X \otimes \mathbf{K} \cong H_A \otimes_A X \otimes_A \widetilde{H_A}$$

as $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodules and

$$Y \otimes \mathbf{K} \cong H_B \otimes_B Y \otimes_B \widetilde{H_B}$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules. Thus

$$Y \cong \widetilde{H_B} \otimes_{B \otimes \mathbf{K}} (\widetilde{M \otimes \mathbf{K}}) \otimes_{A \otimes \mathbf{K}} H_A \otimes_A X \otimes_A \widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B$$

or

$$\widetilde{Y} \cong \widetilde{H_B} \otimes_{B \otimes \mathbf{K}} (\widetilde{M \otimes \mathbf{K}}) \otimes_{A \otimes \mathbf{K}} H_A \otimes_A X \otimes_A \widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B$$

as $B \otimes \mathbf{K} - B \otimes \mathbf{K}$ -equivalence bimodules. Also, by Lemma 2.3,

$$\widetilde{H_A} \otimes_{A \otimes \mathbf{K}} (M \otimes \mathbf{K}) \otimes_{B \otimes \mathbf{K}} H_B \cong M$$

$A - B$ -equivalence bimodules. Hence

$$Y \cong \widetilde{M} \otimes_A X \otimes_A M \quad \text{or} \quad \widetilde{Y} \cong \widetilde{M} \otimes_A X \otimes_A M$$

as $B - B$ -equivalence bimodules. Therefore, we obtain the following:

Theorem 3.6. *Let A and B be σ -unital C^* -algebras and X and Y an $A - A$ -equivalence bimodule and a $B - B$ -equivalence bimodule, respectively. Let $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ be the inclusions of C^* -algebras induced by X and Y , respectively. We suppose that $A' \cap M(A \rtimes_X \mathbf{Z}) \cong \mathbf{C}1$. Then the following conditions are equivalent:*

- (1) $A \subset A \rtimes_X \mathbf{Z}$ and $B \subset B \rtimes_Y \mathbf{Z}$ are strongly Morita equivalent with respect to an $A \rtimes_X \mathbf{Z} - B \rtimes_Y \mathbf{Z}$ -equivalence bimodule N and its closed subspace M ,
- (2) $Y \cong \widetilde{M} \otimes_A X \otimes_A M$ or $\widetilde{Y} \cong \widetilde{M} \otimes_A X \otimes_A M$ as $B - B$ -equivalence bimodules.

Proof. (1) \Rightarrow (2): This is immediate by the above discussions. (2) \Rightarrow (1): This is immediate by Lemma 2.4. \square

4. THE PICARD GROUPS

Let A be a unital C^* -algebra and X an $A - A$ -equivalence bimodule. Let $A \subset A \rtimes_X \mathbf{Z}$ be the inclusion of unital C^* -algebra induced by X . We suppose that $A' \cap (A \rtimes_X \mathbf{Z}) = \mathbf{C}1$. In this section, we shall compute $\text{Pic}(A, A \rtimes_X \mathbf{Z})$, the Picard group of the inclusion $A \subset A \rtimes_X \mathbf{Z}$ (See [10]).

Let G be the subgroup of $\text{Pic}(A)$ defined by

$$G = \{[M] \in \text{Pic}(A) \mid X \cong \widetilde{M} \otimes_A X \otimes_A M \text{ or } \widetilde{X} \cong \widetilde{M} \otimes_A X \otimes_A M \\ \text{as } A - A\text{-equivalence bimodules}\}$$

Let f_A be the homomorphism of $\text{Pic}(A, A \rtimes_X \mathbf{Z})$ to $\text{Pic}(A)$ defined by

$$f_A([M, N]) = [M]$$

for any $[M, N] \in \text{Pic}(A, A \rtimes_X \mathbf{Z})$. First, we show $\text{Im} f_A = G$, where $\text{Im} f_A$ is the image of f_A .

Lemma 4.1. *With the above notation, $\text{Im} f_A = G$.*

Proof. Let $[M, N] \in \text{Pic}(A, A \rtimes_X \mathbf{Z})$. Then by the definition of $\text{Pic}(A, A \rtimes_X \mathbf{Z})$, the inclusion $A \subset A \rtimes_X \mathbf{Z}$ is strongly Morita equivalent to itself with respect to an $A \rtimes_X \mathbf{Z} - A \rtimes_X \mathbf{Z}$ -equivalence bimodule N and its closed subspace M . Hence by Theorem 3.6, $X \cong \widetilde{M} \otimes_A X \otimes_A M$ or $\widetilde{X} \cong \widetilde{M} \otimes_A X \otimes_A M$ as $A - A$ -equivalence bimodules. Thus $\text{Im} f_A \subset G$. Next, let $[M] \in G$. Then by Lemma 2.4, there is an $A \rtimes_X \mathbf{Z} - A \rtimes_X \mathbf{Z}$ -equivalence bimodule N satisfying following:

- (1) M is included in N as a closed subspace,

(2) $[M, N] \in \text{Pic}(A, A \rtimes_X \mathbf{Z})$.

Hence $G \subset \text{Im} f_A$. Therefore, we obtain the conclusion. \square

Next, we compute $\text{Ker} f_A$, the kernel of f_A . Let $\text{Aut}(A, A \rtimes_X \mathbf{Z})$ be the group of all automorphisms α of $A \rtimes_X \mathbf{Z}$ such that $\alpha|_A$ is an automorphism of A . Let $\text{Aut}_0(A, A \rtimes_X \mathbf{Z})$ be the group of all automorphisms α of $A \rtimes_X \mathbf{Z}$ such that $\alpha|_A = \text{id}$ on A . It is clear that $\text{Aut}_0(A, A \rtimes_X \mathbf{Z})$ is a normal subgroup of $\text{Aut}(A, A \rtimes_X \mathbf{Z})$. Let π be the homomorphism of $\text{Aut}(A, A \rtimes_X \mathbf{Z})$ to $\text{Pic}(A, A \rtimes_X \mathbf{Z})$ defined by

$$\pi(\alpha) = [M_\alpha, N_\alpha]$$

for any $\alpha \in \text{Aut}(A, A \rtimes_X \mathbf{Z})$, where $[M_\alpha, N_\alpha]$ is an element in $\text{Pic}(A, A \rtimes_X \mathbf{Z})$ induced by α (See [10, Section 3]).

Lemma 4.2. *With the above notation,*

$$\text{Ker} f_A = \{[A, N_\beta] \in \text{Pic}(A, A \rtimes_X \mathbf{Z}) \mid \beta \in \text{Aut}_0(A, A \rtimes_X \mathbf{Z})\}.$$

Proof. Let $[M, N] \in \text{Ker} f_A$. Then $[M] = [A]$ in $\text{Pic}(A)$ and by [10, Lemma 7.5], there is a $\beta \in \text{Aut}_0(A, A \rtimes_X \mathbf{Z})$ such that

$$[M, N] = [A, N_\beta]$$

in $\text{Pic}(A, A \rtimes_X \mathbf{Z})$, where N_β is the $A \rtimes_X \mathbf{Z} - A \rtimes_X \mathbf{Z}$ -equivalence bimodule induced by β . Therefore, we obtain the conclusion. \square

Let $\text{Int}(A, A \rtimes_X \mathbf{Z})$ be the group of all $\text{Ad}(u)$ such that u is a unitary element in A . By [10, Lemma 3.4],

$$\text{Ker} \pi \cap \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) = \text{Int}(A, A \rtimes_X \mathbf{Z}) \cap \text{Aut}_0(A, A \rtimes_X \mathbf{Z}).$$

Hence

$$\begin{aligned} & \text{Ker} \pi \cap \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) \\ &= \{\text{Ad}(u) \in \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) \mid u \text{ is a unitary element in } A\} \\ &= \{\text{Ad}(u) \in \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) \mid u \text{ is a unitary element in } A' \cap A\}. \end{aligned}$$

Since $A' \cap (A \rtimes_X \mathbf{Z}) = \mathbf{C}1$, $A' \cap A = \mathbf{C}1$. Thus

$$\text{Ker} \pi \cap \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) = \{1\}.$$

It follows that we can obtain the following lemma:

Lemma 4.3. *With the above notation, $\text{Ker} f_A \cong \text{Aut}_0(A, A \rtimes_X \mathbf{Z})$.*

Proof. Since $\text{Ker} \pi \cap \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) = \{1\}$, by Lemma 4.2

$$\begin{aligned} \text{Ker} f_A &= \pi(\text{Aut}_0(A, A \rtimes_X \mathbf{Z})) \cong \text{Aut}_0(A, A \rtimes_X \mathbf{Z}) / (\text{Ker} \pi \cap \text{Aut}_0(A, A \rtimes_X \mathbf{Z})) \\ &= \text{Aut}_0(A, A \rtimes_X \mathbf{Z}). \end{aligned}$$

Therefore, we obtain the conclusion. \square

We recall that the inclusions of C^* -algebras $A \otimes \mathbf{K} \subset (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}$ and $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ are isomorphic as inclusions of C^* -algebras. Also, there is an automorphism α of $A \otimes \mathbf{K}$ such that $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{X \otimes \mathbf{K}} \mathbf{Z}$ and $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}$ are isomorphic as inclusions of C^* -algebras.

Lemma 4.4. *With the above notation, α is free, that is, for any $n \in \mathbf{Z} \setminus \{0\}$, α^n satisfies the following: If $x \in M(A \otimes \mathbf{K})$ satisfies that $xa = \alpha^n(a)x$ for any $a \in A \otimes \mathbf{K}$, then $x = 0$.*

Proof. Since $A' \cap (A \rtimes_X \mathbf{Z}) = \mathbf{C}1$, by [12, Lemma 3.1] $(A \otimes \mathbf{K})' \cap M((A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}) = \mathbf{C}1$. Hence since $A \otimes \mathbf{K} \subset (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}$ is isomorphic to $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}$ as inclusions of C^* -algebras,

$$(A \otimes \mathbf{K})' \cap M((A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}) = \mathbf{C}1.$$

Thus by [12, Corollary 4.2], α is free. \square

For any $n \in \mathbf{Z}$, let δ_n be the function on \mathbf{Z} defined by

$$\delta_n(m) = \begin{cases} 1 & m = n \\ 0 & m \neq n \end{cases}.$$

We regard δ_n as an element in $M((A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$. Let $E^{M(A \otimes \mathbf{K})}$ be the faithful conditional expectation from $M(A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}$ onto $M(A \otimes \mathbf{K})$ defined in Bédos and Conti [2, Section 3]. Let $E^{A \otimes \mathbf{K}} = E^{M(A \otimes \mathbf{K})}|_{(A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}}$. Let $\{u_i\}_{i \in I}$ be an approximate unit of $A \otimes \mathbf{K}$. We fix the approximate unit $\{u_i\}_{i \in I}$ of $A \otimes \mathbf{K}$. For any $x \in M((A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$, we define the Fourier coefficient of x at $n \in \mathbf{Z}$ and the Fourier series of x as in [12, Section 2]. We show that $\text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}) \cong \mathbf{T}$.

Let $\beta \in \text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$. For any $a \in A \otimes \mathbf{K}$,

$$\underline{\beta}(\delta_1)a\underline{\beta}(\delta_1^*) = \beta(\delta_1 a \delta_1^*) = \beta(\alpha(a)) = \alpha(a).$$

Hence $\underline{\beta}(\delta_1)a = \alpha(a)\underline{\beta}(\delta_1)$ for any $a \in A \otimes \mathbf{K}$.

Lemma 4.5. *Let $\sum_{n \in \mathbf{Z}} a_n \delta_1^n = \sum_{n \in \mathbf{Z}} a_n \delta_n$ be the Fourier series of $\underline{\beta}(\delta_1)$, where $a_n \in M(A \otimes \mathbf{K})$ for any $n \in \mathbf{Z}$. Then for any $n \in \mathbf{Z}$, $a \in A \otimes \mathbf{K}$,*

$$a_n \alpha^{n-1}(a) = a a_n.$$

Proof. For any $a \in A \otimes \mathbf{K}$, $\underline{\beta}(\delta_1)a = \alpha(a)\underline{\beta}(\delta_1)$. Hence the Fourier series of $\underline{\beta}(\delta_1)a$ is:

$$\sum_{n \in \mathbf{Z}} a_n \alpha^n(a) \delta_1^n.$$

Also, the Fourier series of $\alpha(a)\underline{\beta}(\delta_1)$ is:

$$\sum_{n \in \mathbf{Z}} \alpha(a) a_n \delta_1^n.$$

Thus $a_n \alpha^n(a) = \alpha(a) a_n$ for any $a \in A \otimes \mathbf{K}$, $n \in \mathbf{Z}$. Therefore, we obtain the conclusion. \square

Lemma 4.6. *With the above notation,*

$$\text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}) \cong \mathbf{T}.$$

Proof. Let $\beta \in \text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$ and let $\sum_{n \in \mathbf{Z}} a_n \delta_1^n$ be the Fourier series of $\underline{\beta}(\delta_1)$. Then by Lemma 4.5, $a_n \alpha^{n-1}(a) = a a_n$ for any $a \in A \otimes \mathbf{K}$, $n \in \mathbf{Z}$. Since α is free by Lemma 4.4, $a_n = 0$ for any $n \in \mathbf{Z} \setminus \{1\}$. Thus $\underline{\beta}(\delta_1) = a_1 \delta_1$. Since $\underline{\beta}(\delta_1)a\underline{\beta}(\delta_1^*) = \alpha(a)$ for any $a \in A \otimes \mathbf{K}$,

$$a_1 \delta_1 a \delta_1^* a_1^* = \alpha(a).$$

Since $\delta_1 a \delta_1^* = \alpha(a)$,

$$a_1 \alpha(a) a_1^* = \alpha(a)$$

for any $a \in A \otimes \mathbf{K}$. Since δ_1 and $\underline{\beta}(\delta_1)$ are unitary elements in $M((A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$, a_1 is a unitary element in $M(A \otimes \mathbf{K})$. Thus

$$a_1 \alpha(a) = \alpha(a) a_1$$

for any $a \in A \otimes \mathbf{K}$. Since $(A \otimes \mathbf{K})' \cap M(A \otimes \mathbf{K}) = \mathbf{C}1$, $a_1 \in \mathbf{C}1$. Since a_1 is a unitary element in $M(A \otimes \mathbf{K})$, there is the unique element $c_\beta \in \mathbf{T}$, such that $a_1 = c_\beta 1$. Let ϵ be the map from $\text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$ onto \mathbf{T} defined

by $\epsilon(\beta) = c_\beta$. By routine computations, we can see that ϵ is an isomorphism of $\text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z})$ onto \mathbf{T} . \square

Lemma 4.7. *With the above notation,*

$$\text{Aut}_0(A, A \rtimes_X \mathbf{Z}) \cong \text{Aut}_0(A \otimes \mathbf{K}, (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}).$$

Proof. Since $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_\alpha \mathbf{Z}$ and $A \otimes \mathbf{K} \subset (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}$ are isomorphic as inclusions of C^* -algebras, it suffices to show that

$$\text{Aut}_0(A, A \rtimes_X \mathbf{Z}) \cong \text{Aut}_0(A \otimes \mathbf{K}, (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K}).$$

Let κ be the homomorphism of $\text{Aut}_0(A, A \rtimes_X \mathbf{Z})$ to $\text{Aut}_0(A \otimes \mathbf{K}, (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K})$ defined by

$$\kappa(\beta) = \beta \otimes \text{id}_{\mathbf{K}}$$

for any $\beta \in \text{Aut}_0(A, A \rtimes_X \mathbf{Z})$. Then it is clear that κ is a monomorphism of $\text{Aut}_0(A, A \rtimes_X \mathbf{Z})$ to $\text{Aut}_0(A \otimes \mathbf{K}, (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K})$. We show that κ is surjective. Let $\gamma \in \text{Aut}_0(A \otimes \mathbf{K}, (A \rtimes_X \mathbf{Z}) \otimes \mathbf{K})$. Then

$$\gamma(a \otimes e_{ij}) = a \otimes e_{ij}$$

for any $a \in A$, $i, j \in \mathbf{N}$. Thus

$$\gamma(x \otimes e_{11}) = (1 \otimes e_{11})\gamma(x \otimes e_{11})(1 \otimes e_{11})$$

for any $x \in A \rtimes_X \mathbf{Z}$. Hence there is an automorphism α of $A \rtimes_X \mathbf{Z}$ such that

$$\gamma(x \otimes e_{11}) = \alpha(x) \otimes e_{11}$$

for any $x \in A \rtimes_X \mathbf{Z}$. For any $i, j \in \mathbf{N}$, $x \in A \rtimes_X \mathbf{Z}$,

$$\begin{aligned} \gamma(x \otimes e_{ij}) &= \gamma((1 \otimes e_{11})(x \otimes e_{11})(1 \otimes e_{1j})) = (1 \otimes e_{11})(\alpha(x) \otimes e_{11})(1 \otimes e_{1j}) \\ &= \alpha(x) \otimes e_{ij}. \end{aligned}$$

Especially, if $a \in A$, $\alpha(a) \otimes e_{ij} = \gamma(a \otimes e_{ij}) = a \otimes e_{ij}$ for any $i, j \in \mathbf{N}$. Thus $\alpha(a) = a$ for any $a \in A$. Therefore, $\gamma = \alpha \otimes \text{id}_{\mathbf{K}}$ and $\alpha \in \text{Aut}_0(A, A \rtimes_X \mathbf{K})$. Hence we can see that κ is surjective. \square

Lemma 4.8. *With the above notation, $\text{Ker} f_A \cong \mathbf{T}$.*

Proof. This is immediate by Lemmas 4.3, 4.6 and 4.7. \square

By Lemmas 4.1 and 4.8, we have the following exact sequence:

$$1 \longrightarrow \mathbf{T} \longrightarrow \text{Pic}(A, A \rtimes_X \mathbf{Z}) \longrightarrow G \longrightarrow 1,$$

where

$$G = \{[M] \in \text{Pic}(A) \mid X \cong \widetilde{M} \otimes_A X \otimes_A M \text{ or } \widetilde{X} \cong \widetilde{M} \otimes_A X \otimes_A M \text{ as } A - A\text{-equivalence bimodules}\}$$

Let g be the map from G to $\text{Pic}(A, A \rtimes_X \mathbf{Z})$ defined by

$$g([M]) = [M, N],$$

where N is the $A \rtimes_X \mathbf{Z} - A \rtimes_X \mathbf{Z}$ -equivalence bimodule defined in the proof of Lemma 2.4. Then g is a homomorphism of G to $\text{Pic}(A, A \rtimes_X \mathbf{Z})$ such that

$$f_A \circ g = \text{id}$$

on G . Thus we obtain the following:

Theorem 4.9. *Let A be a unital C^* -algebra and X an $A - A$ -equivalence bimodule. Let $A \subset A \rtimes_X \mathbf{Z}$ be the unital inclusion of unital C^* -algebras induced by X . We suppose that $A' \cap (A \rtimes_X \mathbf{Z}) = \mathbf{C}1$. Then $\text{Pic}(A, A \rtimes_X \mathbf{Z})$ is a semi-direct product of G by \mathbf{T} .*

REFERENCES

- [1] B. Abadie, S. Eilers and R. Exel, *Morita equivalence for crossed products by Hilbert C^* -bimodules*, Trans. Amer. Math. Soc., **350** (1998), 3043–3054.
- [2] E. Bédos and R. Conti, *On discrete twisted C^* -dynamical systems, Hilbert C^* -modules and regularity*, preprint, arXiv: 1104.1731v1.
- [3] L. G. Brown, P. Green and M. A. Rieffel, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math., **71** (1977), 349–363.
- [4] L. G. Brown, J. Mingo and N-T. Shen, *Quasi-multipliers and embeddings of Hilbert C^* -bimodules*, Can. J. Math., **46** (1994), 1150–1174.
- [5] F. Combes, *Crossed products and Morita equivalence*, Proc. London Math. Soc., **49** (1984), 289–306.
- [6] R. E. Curto, P. S. Muhly and D. P. Williams, *Cross products of strong Morita equivalent C^* -algebras*, Proc. Amer. Math. Soc., **90** (1984), 528–530.
- [7] K. K. Jensen and K. Thomsen, *Elements of KK -theory*, Birkhäuser, 1991.
- [8] M. Izumi, *Inclusions of simple C^* -algebras*, J. reine angew. Math., **547** (2002), 97–138.
- [9] K. Kodaka, *Picard groups of irrational rotation C^* -algebras*, J. London Math. Soc., **56** (1997), 179–188.
- [10] K. Kodaka, *The Picard groups for unital inclusions of unital C^* -algebras*, preprint, arXiv: 1712.09499v1, Acta Sci. Math., to appear.
- [11] K. Kodaka, *Equivalence bundles over a finite group and strong Morita equivalence for unital inclusions of unital C^* -algebras*, preprint, arXiv: 1905.10001.
- [12] K. Kodaka, *Strong Morita equivalence for inclusions of C^* -algebras induced by twisted actions of a countable discrete group*, preprint, arXiv: 1910.1003774.
- [13] K. Kodaka and T. Teruya, *The strong Morita equivalence for coactions of a finite dimensional C^* -Hopf algebra on unital C^* -algebras*, Studia Math., **228** (2015), 259–294.
- [14] K. Kodaka and T. Teruya, *The strong Morita equivalence for inclusions of C^* -algebras and conditional expectations for equivalence bimodules*, J. Aust. Math. Soc., **105** (2018), 103–144.
- [15] K. Kodaka and T. Teruya, *Coactions of a finite dimensional C^* -Hopf algebra on unital C^* -algebras, unital inclusions of unital C^* -algebras and the strong Morita equivalence*, preprint, arXiv:1706.09430.
- [16] G. K. Pedersen, *C^* -algebras and their automorphism groups*, Academic Press, London, New York, San Francisco, 1979.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, RYUKYU
UNIVERSITY, NISHIHARA-CHO, OKINAWA, 903-0213, JAPAN

E-mail address: kodaka@math.u-ryukyu.ac.jp