

Spectral order isomorphisms and AW^* -factors

Martin Bohata¹

*Department of Mathematics, Faculty of Electrical Engineering,
Czech Technical University in Prague, Technická 2,
166 27 Prague 6, Czech Republic*

Abstract: The paper deals with spectral order isomorphisms in the framework of AW^* -algebras. We establish that every spectral order isomorphism between sets of all self-adjoint operators (or between sets of all effects, or between sets of all positive operators) in AW^* -factors of Type I has a canonical form induced by a continuous function calculus and an isomorphism between projection lattices. In particular, this solves an open question about spectral order automorphisms of the set of all (bounded) self-adjoint operators on an infinite-dimensional Hilbert space. We also discuss spectral order isomorphisms preserving, in addition, orthogonality in both directions.

AMS Mathematics Subject Classification: 46L40, 47B49, 06A06

Keywords: AW^* -algebras, spectral order, spectral order isomorphisms

1 Introduction

Self-adjoint operators in a von Neumann algebra can be equipped with a number of partial orders. The most common is Löwner order induced by positive operators. If a von Neumann algebra under consideration is abelian, then its self-adjoint part endowed with Löwner order forms a conditionally complete lattice. However, this is no longer true when we consider noncommutative von Neumann algebras [14, 22].

There is a natural question whether one can find a partial order on the self-adjoint part of a von Neumann algebra which has lattice properties and coincides with Löwner order in the abelian case. An affirmative answer was given by Olson [18] who introduced so called spectral order by means of spectral families in von Neumann algebras. The definition of the spectral order can be directly extended to the more general framework of AW^* -algebras as follows. The *spectral order* is a binary relation \preceq on the self-adjoint part of an AW^* -algebra defined by setting $x \preceq y$ if $E_\lambda^y \leq E_\lambda^x$ for each $\lambda \in \mathbb{R}$, where $(E_\lambda^x)_{\lambda \in \mathbb{R}}$ and $(E_\lambda^y)_{\lambda \in \mathbb{R}}$ are spectral families of self-adjoint operators x and y , respectively, and \leq is the Löwner order.

The spectral order has been especially studied on von Neumann algebras (see, for example, [3, 8] and references therein). Nevertheless, it has been considered also in contexts of AW^* -algebra, unbounded operators, and further structures [7, 10, 12, 19, 23]. From the physical point of view, the spectral order is significant in a categorical formulation of quantum theory [4, 9, 24].

The main aim of this paper is the study of isomorphisms between posets of self-adjoint operators equipped with the spectral order. Let M and N be sets of self-adjoint elements in AW^* -algebras \mathcal{M} and \mathcal{N} , respectively. We say that a bijection $\varphi : M \rightarrow N$ is a *spectral order isomorphism* if, for all $x, y \in M$, $x \preceq y$ if and only if $\varphi(x) \preceq \varphi(y)$. In the special case $M = N$, a spectral order isomorphism φ is called *spectral order automorphism*. In this paper, we show that every spectral order isomorphism between sets of all effects (i.e. positive operators in the unit ball)

¹bohata@math.feld.cvut.cz

in AW^* -factors of Type I has the form $\varphi(x) = \Theta_\tau(f(x))$, where $f : [0, 1] \rightarrow [0, 1]$ is a strictly increasing bijection, τ is an isomorphism between projection lattices, and Θ_τ is defined by $E_\lambda^{\Theta_\tau(x)} = \tau(E_\lambda^x)$ for all $\lambda \in \mathbb{R}$. Let us remark that isomorphisms of the above form were studied for the first time by Turilova in [23]. As each AW^* -factor of Type I is precisely a von Neumann algebra of all bounded operators on a Hilbert space, our result is a direct extension of a complete description of spectral order automorphisms of the set of all effects on a Hilbert space found by Molnár, Nagy, and Šemrl [16, 17]. We also obtain a similar form for all spectral order isomorphisms between self-adjoint parts (as well as between positive parts) of AW^* -factors of Type I. This proves the conjecture of Molnár and Šemrl [17] about spectral order automorphisms on the set of all (bounded) self-adjoint operators on an infinite-dimensional Hilbert space. Moreover, our approach also covers finite-dimensional case with much simpler proof than that of [16] and [17].

The last part of this paper is devoted to spectral order orthoisomorphisms. By a *spectral order orthoisomorphism*, we mean a spectral order isomorphism $\varphi : M \rightarrow N$ preserving orthogonality in both directions (i.e., for all $x, y \in M$, $xy = 0$ if and only if $\varphi(x)\varphi(y) = 0$). A spectral order orthoisomorphism is called a *spectral order orthoautomorphism*, if $M = N$. It was proved by Hamhalter and Turilova in [13] that if φ is a spectral order orthoautomorphism of the set of all effects in an AW^* -factor that is not of Type III and of Type I_2 , then $\varphi(x) = \psi(f(x))$, where ψ is a Jordan $*$ -automorphism and $f : [0, 1] \rightarrow [0, 1]$ is a strictly increasing bijection. We generalize this theorem to spectral order orthoisomorphisms between sets of all effects in arbitrary AW^* -factors that are not of Type I_2 . Moreover, we show that a similar statement is also true when we consider spectral order orthoisomorphisms between self-adjoint parts (or between positive parts) of AW^* -factors.

2 Preliminaries

Let us start this section with summarizing some basic facts about AW^* -algebras. For a detailed treatment of AW^* -algebras, we refer the reader to [2]. An *AW^* -algebra* is a C^* -algebra \mathcal{M} such that for each nonempty set $S \subseteq \mathcal{M}$ there is a projection $p \in \mathcal{M}$ satisfying

$$\{x \in \mathcal{M} \mid sx = 0 \text{ for all } s \in S\} = p\mathcal{M}.$$

It is well known that every von Neumann algebra is AW^* -algebra but the converse is not true. More generally, every monotone complete C^* -algebra is an AW^* -algebra. However, it is an open question whether class of AW^* -algebras coincides with a class of monotone complete C^* -algebras [21]. Each AW^* -algebra is unital and is closed linear span of all its projections. The center

$$\mathcal{Z}(\mathcal{M}) = \{x \in \mathcal{M} \mid xy = yx \text{ for all } y \in \mathcal{M}\}$$

of an AW^* -algebra \mathcal{M} forms an AW^* -subalgebra of \mathcal{M} . An *AW^* -factor* is an AW^* -algebra with one-dimensional center. A classification of AW^* -algebras is analogous to that of von Neumann algebras. Kaplansky [15] proved that an AW^* -algebra of Type I is a von Neumann algebra if and only if its center is a von Neumann algebra.

In particular, every AW^* -factor of Type I is a von Neumann factor of Type I. In the sequel, we shall denote by $P(\mathcal{M})$ the set of all projections in an AW^* -algebra \mathcal{M} . The set $P(\mathcal{M})$ (endowed with the Löwner order) forms a complete lattice called the *projection lattice*. By \mathcal{M}_{sa} and $\mathcal{E}(\mathcal{M})$, we shall denote the self-adjoint part of \mathcal{M} and the set of all effects (i.e. positive operators in the unit ball of \mathcal{M}), respectively.

From the point of view of abelian algebras, AW^* -algebras are more natural than von Neumann algebras. The reason is that $C(X)$ is an AW^* -algebra if and only if X is a Stonean space (i.e. an extremally disconnected compact Hausdorff topological space). Thus every complete Boolean algebra is isomorphic to the projection lattice of an abelian AW^* -algebra. On the other hand, the projection lattice of each abelian AW^* -algebra is a complete Boolean algebra. Consequently, the projection lattice $(P(\mathcal{M}), \leq)$ of an abelian AW^* -algebra \mathcal{M} is meet-infinitely distributive (see, for example, [20, Theorem 5.13]). More specifically,

$$\bigwedge_{\lambda \in \Lambda} (p \vee q_\lambda) = p \vee \bigwedge_{\lambda \in \Lambda} q_\lambda$$

for each $e \in P(\mathcal{M})$ and each family $(q_\lambda)_{\lambda \in \Lambda}$ in $P(\mathcal{M})$. Note also that a nonempty set S of mutually commuting self-adjoint operators in an AW^* -algebra \mathcal{M} generates an abelian AW^* -subalgebra of \mathcal{M} .

Let us look at an auxiliary result concerning projections.

Lemma 2.1. *Let $I \subseteq \mathbb{R}$ be an interval. If $(p_\lambda)_{\lambda \in I}, (q_\lambda)_{\lambda \in I}$ are increasing nets of projections in an abelian AW^* -algebra, then*

$$\bigwedge_{\lambda \in I} p_\lambda \vee q_\lambda = \left(\bigwedge_{\lambda \in I} p_\lambda \right) \vee \left(\bigwedge_{\mu \in I} q_\mu \right).$$

Proof. It is easy to see that $\bigwedge_{\lambda \in I} p_\lambda \vee q_\lambda \geq \bigwedge_{(\lambda, \mu) \in I^2} p_\lambda \vee q_\mu$. On the other hand, $\bigwedge_{\lambda \in I} p_\lambda \vee q_\lambda \leq \bigwedge_{(\lambda, \mu) \in I^2} p_\lambda \vee q_\mu$ because $p_\lambda \vee q_\mu \geq p_\nu \vee q_\nu \geq \bigwedge_{\lambda \in I} p_\lambda \vee q_\lambda$ whenever $\lambda, \mu \in I$ and $\nu = \min\{\lambda, \mu\}$. This establishes that

$$\bigwedge_{\lambda \in I} p_\lambda \vee q_\lambda = \bigwedge_{(\lambda, \mu) \in I^2} p_\lambda \vee q_\mu.$$

As the projection lattice of each abelian AW^* -algebra is meet-infinitely distributive, we have

$$\bigwedge_{\lambda \in I} p_\lambda \vee q_\lambda = \bigwedge_{(\lambda, \mu) \in I^2} p_\lambda \vee q_\mu = \bigwedge_{\lambda \in I} \bigwedge_{\mu \in I} p_\lambda \vee q_\mu = \left(\bigwedge_{\lambda \in I} p_\lambda \right) \vee \left(\bigwedge_{\mu \in I} q_\mu \right).$$

□

Let \mathcal{M}, \mathcal{N} be AW^* -algebras. We say that $\varphi : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ is *orthoisomorphism* if it is a bijection preserving orthogonality in both directions. Dye's theorem [5] is an important extension theorem for orthoisomorphisms between projection lattices of von Neumann algebras. The following generalization of this famous theorem to the context of AW^* -algebras was proved by Hamhalter in [11].

Theorem 2.2 ([11, Theorem 4.3]). *Let \mathcal{M} and \mathcal{N} be AW^* -algebras and let \mathcal{M} have no Type I_2 direct summand. If $\varphi : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ is an orthoisomorphism, then there is a (unique) Jordan $*$ -isomorphism $\Phi : \mathcal{M} \rightarrow \mathcal{N}$ extending φ .*

The next proposition characterizes central projections in terms of lattice distributivity.

Proposition 2.3 ([13, Proposition 3.1]). *If \mathcal{M} is an AW^* -algebra and $z \in P(\mathcal{M})$, then following conditions are equivalent:*

- (i) z is central.
- (ii) $z = (z \wedge p) \vee (z \wedge (\mathbf{1} - p))$ for each $p \in P(\mathcal{M})$.
- (iii) $z \wedge (p \vee q) = (z \wedge p) \vee (z \wedge q)$ for all $p, q \in P(\mathcal{M})$.

Now we turn our attention to spectral families. Recall that a family $(E_\lambda)_{\lambda \in \mathbb{R}}$ of projections in an AW^* -algebra \mathcal{M} is called a *spectral family* if the following conditions hold:

- (i) $E_\lambda \leq E_\mu$ whenever $\lambda \leq \mu$.
- (ii) $E_\lambda = \bigwedge_{\mu > \lambda} E_\mu$ for every $\lambda \in \mathbb{R}$.
- (iii) There is a positive real number α such that $E_\lambda = 0$ when $\lambda < -\alpha$ and $E_\lambda = \mathbf{1}$ when $\lambda > \alpha$, where $\mathbf{1}$ is the unit of \mathcal{M} .

It is well known that there exists a bijection between set of all spectral families in \mathcal{M} and the self-adjoint part of \mathcal{M} . The spectral family $(E_\lambda)_{\lambda \in \mathbb{R}}$ corresponds to $x \in \mathcal{M}_{sa}$ if and only if $xE_\lambda \leq \lambda E_\lambda$ and $\lambda(\mathbf{1} - E_\lambda) \leq x(\mathbf{1} - E_\lambda)$ for each $\lambda \in \mathbb{R}$. In the sequel, we shall denote by $(E_\lambda^x)_{\lambda \in \mathbb{R}}$ the spectral family corresponding to $x \in \mathcal{M}_{sa}$. It is worthwhile to point out that all elements of $(E_\lambda^x)_{\lambda \in \mathbb{R}}$ belong to the abelian AW^* -subalgebra of \mathcal{M} generated by $\{\mathbf{1}, x\}$.

Lemma 2.4. *Let x be a self-adjoint element in an AW^* -algebra.*

- (i) *If α is a positive real number, then $E_\lambda^{\alpha x} = E_{\frac{\lambda}{\alpha}}^x$ for all $\lambda \in \mathbb{R}$.*
- (ii) *If $\alpha \in \mathbb{R}$, then $E_\lambda^{x+\alpha \mathbf{1}} = E_{\lambda-\alpha}^x$ for all $\lambda \in \mathbb{R}$*

Proof. The statements follow immediately from elementary computations. □

Lemma 2.5. *Let \mathcal{N} be an AW^* -subalgebra of an AW^* -algebra \mathcal{M} . Suppose that $\mathbf{1}_\mathcal{M}$ and $\mathbf{1}_\mathcal{N}$ are units of \mathcal{M} and \mathcal{N} , respectively. If $x \in \mathcal{N}_{sa}$ and $(E_\lambda^x)_{\lambda \in \mathbb{R}}$ is the spectral family of x in \mathcal{M} , then $(\mathbf{1}_\mathcal{N} E_\lambda^x)_{\lambda \in \mathbb{R}}$ is the spectral family of x in \mathcal{N} and*

$$(\mathbf{1}_\mathcal{M} - \mathbf{1}_\mathcal{N})E_\lambda^x = \begin{cases} 0 & \text{if } \lambda < 0; \\ \mathbf{1}_\mathcal{M} - \mathbf{1}_\mathcal{N} & \text{if } \lambda \geq 0. \end{cases}$$

Proof. As all elements of $(E_\lambda^x)_{\lambda \in \mathbb{R}}$ are in the abelian AW^* -algebra generated by $\{x, \mathbf{1}_M, \mathbf{1}_N\}$, $\mathbf{1}_N$ commutes with E_λ^x for every $\lambda \in \mathbb{R}$. Thus $\mathbf{1}_N E_\lambda^x \in P(\mathbf{1}_N \mathcal{M} \mathbf{1}_N)$ and

$$(\mathbf{1}_M - \mathbf{1}_N) E_\lambda^x \in P((\mathbf{1}_M - \mathbf{1}_N) \mathcal{M} (\mathbf{1}_M - \mathbf{1}_N)).$$

If $\lambda \in \mathbb{R}$, then

$$\begin{aligned} x \mathbf{1}_N E_\lambda^x &\leq \lambda \mathbf{1}_N E_\lambda^x, \\ x(\mathbf{1}_N - \mathbf{1}_N E_\lambda^x) &\geq \lambda(\mathbf{1}_N - \mathbf{1}_N E_\lambda^x), \\ (\mathbf{1}_M - \mathbf{1}_N)x[(\mathbf{1}_M - \mathbf{1}_N)E_\lambda^x] &\leq \lambda[(\mathbf{1}_M - \mathbf{1}_N)E_\lambda^x], \\ (\mathbf{1}_M - \mathbf{1}_N)x[(\mathbf{1}_M - \mathbf{1}_N) - (\mathbf{1}_M - \mathbf{1}_N)E_\lambda^x] &\geq \lambda[(\mathbf{1}_M - \mathbf{1}_N) - (\mathbf{1}_M - \mathbf{1}_N)E_\lambda^x]. \end{aligned}$$

Thus $(\mathbf{1}_N E_\lambda^x)_{\lambda \in \mathbb{R}}$ is the spectral family of x in \mathcal{N} (because $\mathcal{N} \subseteq \mathbf{1}_N \mathcal{M} \mathbf{1}_N$ and the units of $\mathbf{1}_N \mathcal{M} \mathbf{1}_N$ and \mathcal{N} are same) and $((\mathbf{1}_M - \mathbf{1}_N)E_\lambda^x)_{\lambda \in \mathbb{R}}$ is the spectral family of $(\mathbf{1}_M - \mathbf{1}_N)x = 0$ in $(\mathbf{1}_M - \mathbf{1}_N) \mathcal{M} (\mathbf{1}_M - \mathbf{1}_N)$. Hence

$$(\mathbf{1}_M - \mathbf{1}_N) E_\lambda^x = \begin{cases} 0 & \text{if } \lambda < 0; \\ \mathbf{1}_M - \mathbf{1}_N & \text{if } \lambda \geq 0. \end{cases}$$

□

At the end of this section, we consider some examples of spectral order isomorphisms. We shall see later that they are building blocks of a number of spectral order isomorphisms.

Example 2.6. Consider strictly increasing bijection $f : \mathbb{R} \rightarrow \mathbb{R}$. Suppose that x is a self-adjoint operator in an AW^* -algebra \mathcal{M} and $C(X)$ is an abelian AW^* -algebra *-isomorphic to the AW^* -algebra generated by $\{x, \mathbf{1}\}$. Let $g \in C(X)$ correspond to x . The functions E_λ^g and $E_\lambda^{f \circ g}$ are characteristic functions of clopen sets $X_\lambda = X \setminus \overline{g^{-1}((\lambda, \infty))}$ and

$$Y_\lambda = X \setminus \overline{(f \circ g)^{-1}((\lambda, \infty))} = X \setminus \overline{g^{-1}((f^{-1}(\lambda), \infty))} = X_{f^{-1}(\lambda)},$$

respectively. This shows that $E_\lambda^{f(x)} = E_{f^{-1}(\lambda)}^x$ for every $\lambda \in \mathbb{R}$. Using the definition of spectral order we conclude that $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{M}_{sa}$ given by $\varphi(x) = f(x)$ is a spectral order automorphism.

Note that $f(0) = 0$ if and only if f preserves orthogonality of self-adjoint elements in both directions. This can be shown by passing to the abelian AW^* -algebra $C(X)$ *-isomorphic to the AW^* -algebra generated by $\{\mathbf{1}, x, y\}$. Let g_1 and g_2 be functions in $C(X)$ corresponding to x and y , respectively. Suppose that $f(0) = 0$. Then $f^{-1}(0) = 0$ and so $g_1 g_2 = 0$ if and only if $(f \circ g_1)(f \circ g_2) = 0$. It remains to prove that $f(0) = 0$ whenever f preserves orthogonality. To see this let $f(0) \neq 0$. As 0 is orthogonal to itself, we have $f(0)f(0) = 0$ which is a contradiction. A simple consequence of just proved equivalence says that the map φ defined above is a spectral order orthoautomorphism if and only if $f(0) = 0$.

In a similar way, we can obtain spectral order automorphisms of $\mathcal{E}(\mathcal{M})$ and \mathcal{M}_+ by considering strictly increasing bijections on $[0, 1]$ and $[0, \infty)$, respectively. In these cases, the strictly increasing bijection f automatically satisfies $f(0) = 0$ and so every spectral order automorphism of given form is a spectral order orthoautomorphism.

Let \mathcal{M} and \mathcal{N} be AW^* -algebras. A bijection $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ is called a *projection isomorphism* if it preserves the Löwner order in both directions (i.e., for all $p, q \in P(\mathcal{M})$, $p \leq q$ if and only if $\tau(p) \leq \tau(q)$). The structure of all projection isomorphisms is well known in the case of $\mathcal{M} = B(\mathcal{H})$ and $\mathcal{N} = B(\mathcal{K})$, where \mathcal{H} and \mathcal{K} are Hilbert spaces of dimension at least 3. (As usual, $B(\mathcal{H})$ denotes the von Neumann algebra of all bounded operators on a Hilbert space \mathcal{H} .) Indeed, let $L(\mathcal{H})$ and $L(\mathcal{K})$ be lattices of all closed subspaces of \mathcal{H} and \mathcal{K} , respectively. There is a natural correspondence between τ and a lattice isomorphism $\sigma : L(\mathcal{H}) \rightarrow L(\mathcal{K})$. If \mathcal{H} and \mathcal{K} are finite-dimensional spaces (of dimension at least 3), then the fundamental theorem of projective geometry (see, for example, [1, p. 203]) says that $\sigma(X) = sX$ for some semi-linear bijection s from \mathcal{H} onto \mathcal{K} . In the infinite-dimensional case, $\sigma(X) = sX$, where s is a linear or conjugate-linear bijection from \mathcal{H} onto \mathcal{K} [6].

Example 2.7. Given a projection isomorphism $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ we define a map $\Theta_\tau : \mathcal{M}_{sa} \rightarrow \mathcal{N}_{sa}$ by

$$E_\lambda^{\Theta_\tau(x)} = \tau(E_\lambda^x)$$

for all $x \in \mathcal{M}_{sa}$ and $\lambda \in \mathbb{R}$. It is easy to verify that Θ_τ is a spectral order isomorphism.

By restrictions, we get spectral order isomorphisms between positive parts and between sets of all effects.

In the sequel, Θ_τ will always denote just described spectral order isomorphism determined by a projection isomorphism τ .

Motivated by the previous examples, we say that a spectral order isomorphism $\varphi : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{N})$ (respectively $\varphi : \mathcal{M}_+ \rightarrow \mathcal{N}_+$, $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{N}_{sa}$) is *canonical* if there are a strictly increasing bijection $f : [0, 1] \rightarrow [0, 1]$ (respectively $f : [0, \infty) \rightarrow [0, \infty)$, $f : \mathbb{R} \rightarrow \mathbb{R}$) and a projection isomorphism $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ such that

$$\varphi(x) = \Theta_\tau(f(x))$$

for all $x \in \mathcal{E}(\mathcal{M})$ (respectively $x \in \mathcal{M}_+$, $x \in \mathcal{M}_{sa}$).

Now we give an example of a spectral order orthoisomorphism of the form Θ_τ . Before doing this, we recall that a linear bijection $\psi : \mathcal{M} \rightarrow \mathcal{N}$ between AW^* -algebras \mathcal{M} and \mathcal{N} is called a *Jordan $*$ -isomorphism* if $\psi(x^2) = \psi(x)^2$ and $\psi(x^*) = \psi(x)^*$ for all $x \in \mathcal{M}$.

Example 2.8. Let $\psi : \mathcal{M} \rightarrow \mathcal{N}$ be a Jordan $*$ -isomorphism between AW^* -algebras of \mathcal{M} and \mathcal{N} . We observe that

$$E_\lambda^{\psi(x)} = \psi(E_\lambda^x)$$

for all $\lambda \in \mathbb{R}$ whenever $x \in \mathcal{M}_{sa}$. If a projection isomorphism τ is a restriction of ψ , then $\Theta_\tau(x) = \psi(x)$ for all $x \in \mathcal{M}_{sa}$. Moreover, ψ preserves orthogonality relation in both directions and so a restriction of ψ is a spectral order orthoisomorphism.

3 Spectral order on AW*-algebras

We begin this section with simple observations about spectral order. Statements in the following lemma were proved for von Neumann algebras in papers [8, 18].

Lemma 3.1. *Let \mathcal{M} be an AW*-algebra. Assume that $x, y \in \mathcal{M}_{sa}$, $\alpha, \beta \in \mathbb{R}$, and $\alpha > 0$.*

- (i) $x \preceq y$ if and only if $\alpha x + \beta \mathbf{1} \preceq \alpha y + \beta \mathbf{1}$.
- (ii) The spectral order \preceq coincides with the Löwner order \leq on $P(\mathcal{M})$.
- (iii) If $x \preceq y$, then $x \leq y$.
- (iv) Suppose that x commutes with y . Then $x \preceq y$ if and only if $x \leq y$.

Proof. Using Lemma 2.4, we immediately obtain the desired equivalence in (i).

By considering spectral families of $p, q \in P(\mathcal{M})$, $p \preceq q$ if and only if $p \leq q$. This shows (ii).

It remains to prove (iii) and (iv). Set $\alpha = \max\{\|x\|, \|y\|\}$. If $\alpha = 0$, then both statements are clear. Assume that $\alpha > 0$. Then

$$\frac{1}{2\alpha}x + \frac{1}{2}\mathbf{1}, \frac{1}{2\alpha}y + \frac{1}{2}\mathbf{1} \in \mathcal{E}(\mathcal{M}).$$

Furthermore, we see from (i) that

$$x \preceq y \Leftrightarrow \frac{1}{2\alpha}x + \frac{1}{2}\mathbf{1} \preceq \frac{1}{2\alpha}y + \frac{1}{2}\mathbf{1}.$$

Now it follows from [12, Corollary 1] and [12, Corollary 2] that (iii) and (iv) are true. \square

Proposition 3.2. *Let \mathcal{N} be an AW*-subalgebra of an AW*-algebra \mathcal{M} . Suppose that $x, y \in \mathcal{N}_{sa}$. Then $x \preceq y$ in \mathcal{M}_{sa} if and only if $x \preceq y$ in \mathcal{N}_{sa} .*

Proof. Let $(E_\lambda^x)_{\lambda \in \mathbb{R}}$ and $(E_\lambda^y)_{\lambda \in \mathbb{R}}$ be spectral families of x and y , respectively, in \mathcal{M} . If $x \preceq y$ in \mathcal{M} , then $E_\lambda^y \leq E_\lambda^x$ for all $\lambda \in \mathbb{R}$. Since the unit $\mathbf{1}_\mathcal{N}$ of \mathcal{N} commutes with E_λ^x and E_λ^y for all $\lambda \in \mathbb{R}$, $\mathbf{1}_\mathcal{N}E_\lambda^y \leq \mathbf{1}_\mathcal{N}E_\lambda^x$ for all $\lambda \in \mathbb{R}$. It follows from Lemma 2.5 that $x \preceq y$ in \mathcal{N} .

Conversely, let $x \preceq y$ in \mathcal{N} . Then $\mathbf{1}_\mathcal{N}E_\lambda^y \leq \mathbf{1}_\mathcal{N}E_\lambda^x$ for all $\lambda \in \mathbb{R}$ by Lemma 2.5. Moreover, Lemma 2.5 implies that

$$(\mathbf{1}_\mathcal{M} - \mathbf{1}_\mathcal{N})E_\lambda^x = (\mathbf{1}_\mathcal{M} - \mathbf{1}_\mathcal{N})E_\lambda^y$$

for all $\lambda \in \mathbb{R}$. Therefore,

$$E_\lambda^y = \mathbf{1}_\mathcal{N}E_\lambda^y + (\mathbf{1}_\mathcal{M} - \mathbf{1}_\mathcal{N})E_\lambda^y \leq \mathbf{1}_\mathcal{N}E_\lambda^x + (\mathbf{1}_\mathcal{M} - \mathbf{1}_\mathcal{N})E_\lambda^x = E_\lambda^x$$

for all $\lambda \in \mathbb{R}$. Hence $x \preceq y$ in \mathcal{M} . \square

It can be proved that the self-adjoint part of an AW^* -algebra equipped with the spectral order is a conditionally complete lattice (the proof is the same as for von Neumann algebras [18]). Suprema and infima are determined by their spectral families as follows. Let \mathcal{M} be an AW^* -algebra and let $M \subseteq \mathcal{M}_{sa}$ be nonempty. If M is bounded above, then its supremum $\bigvee_{x \in M} x$ in $(\mathcal{M}_{sa}, \preceq)$ is a self-adjoint element with the spectral family

$$\left(E_{\lambda}^{\bigvee_{x \in M} x}\right)_{\lambda \in \mathbb{R}} = \left(\bigwedge_{x \in M} E_{\lambda}^x\right)_{\lambda \in \mathbb{R}}.$$

If M is bounded below, then its infimum $\bigwedge_{x \in M} x$ in $(\mathcal{M}_{sa}, \preceq)$ is a self-adjoint element with the spectral family

$$\left(E_{\lambda}^{\bigwedge_{x \in M} x}\right)_{\lambda \in \mathbb{R}} = \left(\bigwedge_{\mu > \lambda} \bigvee_{x \in M} E_{\mu}^x\right)_{\lambda \in \mathbb{R}}.$$

The next two assertions state that suprema and infima with respect to the spectral order are stable under taking smaller posets. The latter is a generalization of [3, Proposition 2.2] to the case of AW^* -algebras.

Proposition 3.3. *Let \mathcal{N} be an AW^* -subalgebra of an AW^* -algebra \mathcal{M} . Suppose that $M \subseteq \mathcal{N}_{sa}$ is nonempty.*

- (i) *If M has an upper bound in $(\mathcal{M}_{sa}, \preceq)$, then both the supremum of M in $(\mathcal{M}_{sa}, \preceq)$ and the supremum of M in $(\mathcal{N}_{sa}, \preceq)$ exist and coincide.*
- (ii) *If M has a lower bound in $(\mathcal{M}_{sa}, \preceq)$, then both the infimum of M in $(\mathcal{M}_{sa}, \preceq)$ and the infimum of M in $(\mathcal{N}_{sa}, \preceq)$ exist and coincide.*

Proof. (i) The existence of suprema is ensured by the fact that $(\mathcal{M}_{sa}, \preceq)$ and $(\mathcal{N}_{sa}, \preceq)$ are conditionally complete lattices. If $x \in \mathcal{M}_{sa}$, then we denote by $(E_{\lambda}^x)_{\lambda \in \mathbb{R}}$ the spectral family of x in \mathcal{M} . Let a be the supremum of M in \mathcal{M} and let b be the supremum of M in \mathcal{N} . By Lemma 2.5,

$$\begin{aligned} E_{\lambda}^b &= \mathbf{1}_{\mathcal{N}} E_{\lambda}^b + (\mathbf{1}_{\mathcal{M}} - \mathbf{1}_{\mathcal{N}}) E_{\lambda}^b = \bigwedge_{x \in M} \mathbf{1}_{\mathcal{N}} E_{\lambda}^x + \bigwedge_{x \in M} (\mathbf{1}_{\mathcal{M}} - \mathbf{1}_{\mathcal{N}}) E_{\lambda}^x \\ &= \mathbf{1}_{\mathcal{N}} \bigwedge_{x \in M} E_{\lambda}^x + (\mathbf{1}_{\mathcal{M}} - \mathbf{1}_{\mathcal{N}}) \bigwedge_{x \in M} E_{\lambda}^x = \bigwedge_{x \in M} E_{\lambda}^x = E_{\lambda}^a \end{aligned}$$

for all $\lambda \in \mathbb{R}$. Thus $a = b$.

- (ii) This follows from (i) by considering the order-reversing map $t \mapsto -t$.

□

Proposition 3.4. *Let \mathcal{M} be an AW^* -algebra.*

- (i) *If $M \subseteq \mathcal{M}_+$ is nonempty and bounded above in $(\mathcal{M}_{sa}, \preceq)$, then the supremum of M in $(\mathcal{M}_{sa}, \preceq)$ is a positive element.*
- (ii) *If $M \subseteq \mathcal{M}_+$ is nonempty, then the infimum of M in $(\mathcal{M}_{sa}, \preceq)$ is a positive element.*

- (iii) If $L \in \{\mathcal{E}(\mathcal{M}), P(\mathcal{M})\}$, then the supremum and the infimum of every nonempty subset of L in $(\mathcal{M}_{sa}, \preceq)$ belong to L .

Proof. The proof is similar to that of [3, Proposition 2.2] and so it is omitted. \square

We shall denote suprema and infima of a set M in posets $(\mathcal{M}_{sa}, \preceq)$, (\mathcal{M}_+, \preceq) , $(\mathcal{E}(\mathcal{M}), \preceq)$, and $(P(\mathcal{M}), \leq) = (P(\mathcal{M}), \preceq)$ by symbols $\bigvee_{x \in M} x$ and $\bigwedge_{x \in M} x$, respectively. We have seen in the previous result that this simple notation cannot lead to any possible misunderstanding.

Lemma 3.5. *Let \mathcal{M} be an AW*-algebra. If $x, y \in \mathcal{M}_{sa}$ are mutually orthogonal, then*

$$x \vee y = x^+ + y^+,$$

where x^+ and y^+ are the positive parts of x and y , respectively.

Proof. The statement clearly holds for $\mathcal{M} = C(X)$, where X is Stonean. Since x commutes with y , we can reduce the general case to the above special case by Proposition 3.3. \square

The content of the following lemma appeared in the paper [17] without proof. Therefore, we present a proof for convenience of the reader.

Lemma 3.6. *Let \mathcal{M} be an AW*-algebra and let $p, q \in P(\mathcal{M})$. If real numbers α and β satisfy $0 \leq \alpha \leq \beta$, then $\alpha p \vee \beta q = \alpha(p \vee q - q) + \beta q$.*

Proof. The case $\alpha = 0$ is clear because $0 \vee \beta q = \beta q$ by Lemma 3.5.

If $0 < \alpha = \beta$, then $\alpha p \vee \beta q = \alpha(p \vee q - q) + \beta q$ because multiplication by a positive scalar is a spectral order isomorphism (see Lemma 3.1).

Suppose that $0 < \alpha < \beta$. Using Lemma 2.4, we can directly verify that

$$E_{\lambda}^{\alpha p \vee \beta q} = \begin{cases} 0, & \lambda < 0; \\ (\mathbf{1} - p) \wedge (\mathbf{1} - q), & \lambda \in [0, \alpha); \\ (\mathbf{1} - q), & \lambda \in [\alpha, \beta); \\ \mathbf{1}, & \lambda \geq \beta. \end{cases}$$

As $q \leq p \vee q$, $p \vee q$ commutes with $\mathbf{1} - q$ and so

$$p \vee q - q = (p \vee q)(\mathbf{1} - q) = (p \vee q) \wedge (\mathbf{1} - q).$$

Moreover, q is orthogonal to $p \vee q - q$. Consequently,

$$E_{\lambda}^{\alpha(p \vee q - q) + \beta q} = \begin{cases} 0, & \lambda < 0; \\ [((\mathbf{1} - p) \wedge (\mathbf{1} - q)) \vee q] \wedge (\mathbf{1} - q), & \lambda \in [0, \alpha); \\ (\mathbf{1} - q), & \lambda \in [\alpha, \beta); \\ \mathbf{1}, & \lambda \geq \beta. \end{cases}$$

Since q is orthogonal to $(\mathbf{1} - p) \wedge (\mathbf{1} - q)$ and $\mathbf{1} - q$ commutes with $[(\mathbf{1} - p) \wedge (\mathbf{1} - q)] + q$, we have

$$\begin{aligned} [((\mathbf{1} - p) \wedge (\mathbf{1} - q)) \vee q] \wedge (\mathbf{1} - q) &= [((\mathbf{1} - p) \wedge (\mathbf{1} - q)) + q] \wedge (\mathbf{1} - q) \\ &= [((\mathbf{1} - p) \wedge (\mathbf{1} - q)) + q](\mathbf{1} - q) \\ &= (\mathbf{1} - p) \wedge (\mathbf{1} - q). \end{aligned}$$

Hence

$$E_{\lambda}^{\alpha p \vee \beta q} = E_{\lambda}^{\alpha(p \vee q - q) + \beta q}$$

for all $\lambda \in \mathbb{R}$. □

A projection p in an AW^* -algebra \mathcal{M} is said to be *atomic* if there is no nonzero projection q in \mathcal{M} such that $q \neq p$ and $q \leq p$. We shall denote by $P_{at}(\mathcal{M})$ the set of all atomic projections in an AW^* -algebra \mathcal{M} . Now we characterize effects corresponding nonzero scalar multiples of atomic projections in terms of the spectral order.

Proposition 3.7. *Let \mathcal{M} be an AW^* -algebra and let $x \in \mathcal{E}(\mathcal{M})$ be nonzero. Then the following statements are equivalent:*

- (i) *There are $\lambda \in (0, 1]$ and an atomic projection $e \in \mathcal{M}$ such that $x = \lambda e$.*
- (ii) *If $y, z \in \mathcal{E}(\mathcal{M})$ satisfy $y, z \preceq x$, then $y \preceq z$ or $z \preceq y$.*

Proof. Assume that (i) holds and $y, z \in \mathcal{E}(\mathcal{M})$ satisfy $y, z \preceq x$. Then $y, z \leq \lambda e$. Since $e\mathcal{M}e$ is a hereditary AW^* -subalgebra of \mathcal{M} , we have $y, z \in e\mathcal{M}e$. From the fact that $e\mathcal{M}e$ contains only the scalar multiples of e , we see that $y = \mu e$ and $z = \nu e$ for some $\mu, \nu \in (0, 1]$. Consequently, $y \preceq z$ or $z \preceq y$.

In order to prove that (ii) implies (i), consider the abelian AW^* -algebra \mathcal{N} generated by x . Clearly, $\mathcal{N} \simeq C(X)$, where X is a Stonean space. Let $f \in C(X)$ correspond to x . Suppose that f attains two different nonzero values at points $t_1, t_2 \in X$. Take disjoint clopen subsets O_1 and O_2 of X such that $t_1 \in O_1$ and $t_2 \in O_2$. Consider $g_1 = \chi_{O_1}f$ and $g_2 = \chi_{O_2}f$, where χ_{O_1} and χ_{O_2} are characteristic functions of O_1 and O_2 , respectively. Then $g_1 \not\preceq g_2$ and $g_2 \not\preceq g_1$ which is a contradiction with (ii). Therefore, $f = \lambda \chi_O$ for some $\lambda \in (0, 1]$, where χ_O is a characteristic function of a clopen set O . Hence there is a nonzero projection $e \in P(\mathcal{M})$ such that $x = \lambda e$. Suppose that e is not atomic. Then one can find a nonzero projection $p \in P(\mathcal{M})$ such that $p \leq e$ and $p \neq e$. This implies that $\lambda p \preceq \lambda e$ and $\lambda(e - p) \preceq \lambda e$. However, $\lambda(e - p) \not\preceq \lambda p$ and $\lambda p \not\preceq \lambda(e - p)$ which is a contradiction with (ii). Thus e is atomic projection. □

Recall the notion of a distributive element in a lattice. Let (P, \leq) be a lattice. We say that an element $z \in P$ is *distributive* if

$$z \vee (x \wedge y) = (z \vee x) \wedge (z \vee y)$$

for all $x, y \in P$. We denote by $\mathcal{D}(P, \leq)$ the set of all distributive elements in (P, \leq) . In the next proposition, we characterize some classes of central elements in terms of distributivity. It turns out that this observation is essential to prove various results about spectral order isomorphisms.

Proposition 3.8. *Let \mathcal{M} be an AW^* -algebra. Then*

- (i) $\mathcal{Z}(\mathcal{M})_{sa} = \mathcal{D}(\mathcal{M}_{sa}, \preceq)$;
- (ii) $\mathcal{Z}(\mathcal{M})_+ = \mathcal{D}(\mathcal{M}_+, \preceq)$;
- (iii) $\mathcal{E}(\mathcal{Z}(\mathcal{M})) = \mathcal{D}(\mathcal{E}(\mathcal{M}), \preceq)$.

Proof. We shall prove only the statement (i). Proofs of remaining assertions (ii) and (iii) are similar. Let $z \in \mathcal{Z}(\mathcal{M})_{sa}$. Then $E_\lambda^z \in \mathcal{Z}(\mathcal{M})$ for all $\lambda \in \mathbb{R}$. Take $x, y \in \mathcal{M}_{sa}$. Applying Proposition 2.3,

$$\begin{aligned} E_\lambda^{z \vee (x \wedge y)} &= E_\lambda^z \wedge E_\lambda^{x \wedge y} = E_\lambda^z \wedge \bigwedge_{\mu > \lambda} (E_\mu^x \vee E_\mu^y) = \bigwedge_{\mu > \lambda} E_\mu^z \wedge (E_\mu^x \vee E_\mu^y) \\ &= \bigwedge_{\mu > \lambda} (E_\mu^z \wedge E_\mu^x) \vee (E_\mu^z \wedge E_\mu^y) = \bigwedge_{\mu > \lambda} E_\mu^{z \vee x} \vee E_\mu^{z \vee y} = E_\lambda^{(z \vee x) \wedge (z \vee y)} \end{aligned}$$

for all $\lambda \in \mathbb{R}$. Hence

$$z \vee (x \wedge y) = (z \vee x) \wedge (z \vee y)$$

and so $\mathcal{Z}(\mathcal{M})_{sa} \subseteq \mathcal{D}(\mathcal{M}_{sa}, \preceq)$.

It remains to prove the reverse inclusion. Assume that $z \in \mathcal{D}(\mathcal{M}_{sa}, \preceq)$ is nonzero and that $p \in P(\mathcal{M})$. It is easy to see that

$$E_\lambda^z = (E_\lambda^z \wedge p) \vee [E_\lambda^z \wedge (\mathbf{1} - p)]$$

whenever $\lambda < -\|z\|$ and $\lambda \geq \|z\|$. As $z \in \mathcal{D}(\mathcal{M}_{sa}, \preceq)$ and

$$\|z\| (2p - \mathbf{1}) \wedge \|z\| (\mathbf{1} - 2p) = -\|z\| \mathbf{1},$$

we have

$$z = z \vee [\|z\| (2p - \mathbf{1}) \wedge \|z\| (\mathbf{1} - 2p)] = (z \vee \|z\| (2p - \mathbf{1})) \wedge (z \vee \|z\| (\mathbf{1} - 2p)).$$

Hence

$$\begin{aligned} E_\lambda^z &= E_\lambda^{(z \vee \|z\| (2p - \mathbf{1})) \wedge (z \vee \|z\| (\mathbf{1} - 2p))} = \bigwedge_{\mu > \lambda} E_\mu^{(z \vee \|z\| (2p - \mathbf{1}))} \vee E_\mu^{(z \vee \|z\| (\mathbf{1} - 2p))} \\ &= \bigwedge_{\mu > \lambda} (E_\mu^z \wedge E_\mu^{\|z\| (2p - \mathbf{1})}) \vee (E_\mu^z \wedge E_\mu^{\|z\| (\mathbf{1} - 2p)}) \end{aligned}$$

for all $\lambda \in \mathbb{R}$. Since $(E_\mu^z \wedge E_\mu^{\|z\| (2p - \mathbf{1})})_{\mu \in \mathbb{R}}$ is an increasing net, $E_\mu^z \wedge E_\mu^{\|z\| (2p - \mathbf{1})}$ commutes with $E_\nu^z \wedge E_\nu^{\|z\| (2p - \mathbf{1})}$ for all $\mu, \nu \in \mathbb{R}$. In the same way, we can show that $E_\mu^z \wedge E_\mu^{\|z\| (\mathbf{1} - 2p)}$ commutes with $E_\nu^z \wedge E_\nu^{\|z\| (\mathbf{1} - 2p)}$ for all $\mu, \nu \in \mathbb{R}$. Moreover, if $\mu, \nu \in [-\|z\|, \|z\|]$, then $E_\mu^z \wedge E_\mu^{\|z\| (2p - \mathbf{1})}$ and $E_\nu^z \wedge E_\nu^{\|z\| (\mathbf{1} - 2p)}$ commute because they are mutually orthogonal. Therefore,

$$\{E_\mu^z \wedge E_\mu^{\|z\| (2p - \mathbf{1})} \mid \mu \in [-\|z\|, \|z\|]\} \cup \{E_\mu^z \wedge E_\mu^{\|z\| (\mathbf{1} - 2p)} \mid \mu \in [-\|z\|, \|z\|]\}$$

is contained in an abelian AW^* -subalgebra of \mathcal{M} . If $\lambda \in [-\|z\|, \|z\|]$ and $I_\lambda =$

$(\lambda, \|z\|)$, then we see from Lemma 2.1 that

$$\begin{aligned}
E_\lambda^z &= \bigwedge_{\mu > \lambda} (E_\mu^z \wedge E_\mu^{\|z\|(2p-1)}) \vee (E_\mu^z \wedge E_\mu^{\|z\|(1-2p)}) \\
&= \bigwedge_{\mu \in I_\lambda} (E_\mu^z \wedge E_\mu^{\|z\|(2p-1)}) \vee (E_\mu^z \wedge E_\mu^{\|z\|(1-2p)}) \\
&= \left(\bigwedge_{\mu \in I_\lambda} E_\mu^z \wedge E_\mu^{\|z\|(2p-1)} \right) \vee \left(\bigwedge_{\mu \in I_\lambda} E_\mu^z \wedge E_\mu^{\|z\|(1-2p)} \right) \\
&= \left[\left(\bigwedge_{\mu \in I_\lambda} E_\mu^z \right) \wedge \left(\bigwedge_{\mu \in I_\lambda} E_\mu^{\|z\|(2p-1)} \right) \right] \vee \left[\left(\bigwedge_{\mu \in I_\lambda} E_\mu^z \right) \wedge \left(\bigwedge_{\mu \in I_\lambda} E_\mu^{\|z\|(1-2p)} \right) \right] \\
&= (E_\lambda^z \wedge E_\lambda^{\|z\|(2p-1)}) \vee (E_\lambda^z \wedge E_\lambda^{\|z\|(1-2p)}) = (E_\lambda^z \wedge E_{\frac{\lambda+\|z\|}{2\|z\|}}^p) \vee (E_\lambda^z \wedge E_{\frac{\lambda+\|z\|}{2\|z\|}}^{1-p}) \\
&= [E_\lambda^z \wedge (1-p)] \vee (E_\lambda^z \wedge p).
\end{aligned}$$

According to Proposition 2.3, E_λ^z is a central projection for each $\lambda \in \mathbb{R}$. As each projection in the spectral family of z is central, $z \in \mathcal{Z}(\mathcal{M})$. \square

4 Spectral order isomorphisms

It was proved in [13, Theorem 4.3] that canonical spectral order automorphisms are characterized by preserving scalar multiples of projections. We generalize this result to spectral order isomorphisms.

Theorem 4.1. *Let \mathcal{M} and \mathcal{N} be AW^* -algebras. Suppose that $\varphi : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{N})$ is a spectral order isomorphism. Then the following statements are equivalent:*

- (i) *For each $p \in P(\mathcal{M})$, $\varphi(p) \in P(\mathcal{N})$ and there is a strictly increasing bijection $f_p : [0, 1] \rightarrow [0, 1]$ such that $\varphi(\lambda p) = f_p(\lambda)\varphi(p)$ for all $\lambda \in [0, 1]$.*
- (ii) *φ is canonical.*

Proof. (i) \Rightarrow (ii). Define the map $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ by $\tau(p) = \varphi(p)$. The map τ is a projection isomorphism. Thus $\Theta_\tau^{-1} \circ \varphi : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{M})$ is a spectral order isomorphism. By [13, Theorem 4.3], $\Theta_\tau^{-1} \circ \varphi$ is canonical. Therefore, φ is canonical.

(ii) \Rightarrow (i). By the assumption, there are a strictly increasing bijection $f : [0, 1] \rightarrow [0, 1]$ and a projection isomorphism $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ such that $\varphi(x) = \Theta_\tau(f(x))$ for all $x \in \mathcal{E}(\mathcal{M})$. A direct computation shows that

$$\varphi(\lambda p) = \Theta_\tau(f(\lambda)p) = f(\lambda)\Theta_\tau(p) = f(\lambda)\varphi(p)$$

for all $\lambda \in [0, 1]$. \square

Using the previous theorem, we obtain in Corollary 4.2 that every spectral order isomorphism between sets of all effects in AW^* -factors of Type I is canonical. This statement is a slight generalization of known results concerning spectral order automorphisms [16, 17]. In contrast with the form of automorphisms in [16, 17], it shows that the main structure of a spectral order isomorphism between sets of all

effects does not depend on a particular choice of AW^* -factors of Type I. We give the proof of Corollary 4.2 for completeness of discussion. Note that an advantage of our approach is that we need not discuss the case of AW^* -factor of Type I_2 separately.

Corollary 4.2. *If \mathcal{M} and \mathcal{N} are AW^* -factors of Type I, then every spectral order isomorphism $\varphi : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{N})$ is canonical.*

Proof. It follows from Proposition 3.7 that if $p \in P_{at}(\mathcal{M})$ and $\lambda \in (0, 1]$, then there are $q \in P_{at}(\mathcal{N})$ and $\mu \in (0, 1]$ such that $\varphi(\lambda p) = \mu q$. Take $\lambda_1, \lambda_2 \in (0, 1]$ and $p \in P_{at}(\mathcal{M})$. Then $\varphi(\lambda_1 p) = \mu_1 q_1$ and $\varphi(\lambda_2 p) = \mu_2 q_2$ for some $\mu_1, \mu_2 \in (0, 1]$ and $q_1, q_2 \in P_{at}(\mathcal{N})$. Let us show that $q_1 = q_2$. It is not difficult to verify that $\lambda_1 p \vee \lambda_2 p = \max\{\lambda_1, \lambda_2\}p$. Thus

$$\varphi(\max\{\lambda_1, \lambda_2\}p) = \varphi(\lambda_1 p \vee \lambda_2 p) = \mu_1 q_1 \vee \mu_2 q_2.$$

Suppose that $q_1 \neq q_2$. If $\mu_1 q_1 \preceq \mu_2 q_2$, then $E_\nu^{\mu_2 q_2} \leq E_\nu^{\mu_1 q_1}$ for every $\nu \in \mathbb{R}$. In particular, $E_0^{\mu_2 q_2} \leq E_0^{\mu_1 q_1}$ and so $q_1 \leq q_2$. This is a contradiction because q_1 and q_2 are two distinct atoms. Hence $\mu_1 q_1 \not\preceq \mu_2 q_2$. Interchanging the roles of $\mu_1 q_1$ and $\mu_2 q_2$, we see that $\mu_2 q_2 \not\preceq \mu_1 q_1$. According to Proposition 3.7, there are no $\mu \in (0, 1]$ and $e \in P_{at}(\mathcal{N})$ such that $\varphi(\max\{\lambda_1, \lambda_2\}p) = \mu e$ which is a contradiction. Consequently, $q_1 = q_2$. As φ^{-1} is also a spectral order isomorphism, there are a strictly increasing bijection $f_p : [0, 1] \rightarrow [0, 1]$ and $q \in P_{at}(\mathcal{N})$ such that $\varphi(\lambda p) = f_p(\lambda)q$ for all $\lambda \in [0, 1]$.

If $p, q \in P_{at}(\mathcal{M})$ are different and $\lambda \in (0, 1]$, then there is $e \in P_{at}(\mathcal{M})$ different from p and q such that $e \leq p \vee q$. Multiplication by λ is a spectral order automorphism of \mathcal{M}_+ which maps $\mathcal{E}(\mathcal{M})$ into itself. This implies that

$$\varphi(\lambda p \vee \lambda q) = \varphi(\lambda p \vee \lambda e) = \varphi(\lambda e \vee \lambda q).$$

Hence

$$f_p(\lambda)\varphi(p) \vee f_q(\lambda)\varphi(q) = f_p(\lambda)\varphi(p) \vee f_e(\lambda)\varphi(e) = f_e(\lambda)\varphi(e) \vee f_q(\lambda)\varphi(q).$$

Since the spectra of these three operators have to coincide, we see from Lemma 3.6 that $f_p(\lambda) = f_e(\lambda) = f_q(\lambda)$. Therefore, there is a strictly increasing function $f : [0, 1] \rightarrow [0, 1]$ such that $\varphi(\lambda p) = f(\lambda)\varphi(p)$ for all $p \in P_{at}(\mathcal{M})$ and $\lambda \in [0, 1]$. As every projection in \mathcal{M} is the supremum of a set of atomic projections in \mathcal{M} , $\varphi(\lambda p) = f(\lambda)\varphi(p)$ for all $p \in P(\mathcal{M})$ and $\lambda \in [0, 1]$. According to Theorem 4.1, φ is canonical. \square

Now we would like to extend the preceding corollary to positive operators and also to self-adjoint operators. We start with the case of positive operators. For this, we need the following lemmas.

Lemma 4.3. *Let \mathcal{M} and \mathcal{N} be AW^* -algebras and let $\alpha \in [1, \infty)$. If $f : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing bijection and $\varphi : \mathcal{M}_+ \rightarrow \mathcal{N}_+$ is a spectral order isomorphism such that $\varphi(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in [0, \infty)$, then*

$$\frac{1}{\alpha} \varphi(f^{-1}(\alpha \mathcal{E}(\mathcal{M}))) = \mathcal{E}(\mathcal{N}).$$

Proof. Let $x \in \mathcal{E}(\mathcal{M})$. Lemma 3.1 establishes that $0 \preceq x \preceq \mathbf{1}_{\mathcal{M}}$. Since multiplication by a positive number is a spectral order isomorphism, we have

$$0 \preceq \frac{1}{\alpha} \varphi(f^{-1}(\alpha x)) \preceq \frac{1}{\alpha} \varphi(f^{-1}(\alpha \mathbf{1}_{\mathcal{M}})) = \mathbf{1}_{\mathcal{N}}.$$

By Lemma 3.1, $\frac{1}{\alpha} \varphi(f^{-1}(\alpha x)) \in \mathcal{E}(\mathcal{N})$.

Now take $y \in \mathcal{E}(\mathcal{N})$. Consider $x = \frac{1}{\alpha} f(\varphi^{-1}(\alpha y))$. Similar arguments as above shows that $x \in \mathcal{E}(\mathcal{M})$. Moreover, $\frac{1}{\alpha} \varphi(f^{-1}(\alpha x)) = y$. \square

Lemma 4.4. *Let $\varphi : \mathcal{M}_+ \rightarrow \mathcal{N}_+$ be a spectral order isomorphism between positive parts of AW^* -algebras \mathcal{M} and \mathcal{N} such that there exists a strictly increasing bijection $f : [0, \infty) \rightarrow [0, \infty)$ satisfying $\varphi(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in [0, \infty)$. Suppose that, for each $\alpha \in [1, \infty)$, the equation*

$$\varphi_{\alpha}(x) = \frac{1}{\alpha} \varphi(f^{-1}(\alpha x))$$

defines a canonical spectral order isomorphism $\varphi_{\alpha} : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{N})$.

- (i) *There exists a projection isomorphism $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ such that $\varphi(x) = \Theta_{\tau}(f(x))$ for all $x \in \mathcal{M}_+$.*
- (ii) *If, in addition, there is a Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that $\psi(x) = \varphi(f^{-1}(x))$ for all $x \in \mathcal{E}(\mathcal{M})$, then $\varphi(x) = \psi(f(x))$ for all $x \in \mathcal{M}_+$.*

Proof. (i) Maps φ_{α} are canonical spectral order isomorphisms and so, for each $\alpha \in [1, \infty)$, there are a strictly increasing bijection $g_{\alpha} : [0, 1] \rightarrow [0, 1]$ and a projection isomorphism $\tau_{\alpha} : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ such that $\varphi_{\alpha}(x) = \Theta_{\tau_{\alpha}}(g_{\alpha}(x))$ whenever $x \in \mathcal{E}(\mathcal{M})$.

Let $\alpha \in [1, \infty)$ and $\lambda \in [0, 1]$ be arbitrary. Then

$$\alpha g_{\alpha}(\lambda) \mathbf{1}_{\mathcal{N}} = \alpha \varphi_{\alpha}(\lambda \mathbf{1}_{\mathcal{M}}) = \varphi(f^{-1}(\alpha \lambda \mathbf{1}_{\mathcal{M}})) = \alpha \lambda \mathbf{1}_{\mathcal{N}}.$$

Thus $g_{\alpha}(\lambda) = \lambda$ and so $\varphi_{\alpha}(x) = \Theta_{\tau_{\alpha}}(x)$ for all $x \in \mathcal{E}(\mathcal{M})$. Set $\tau = \tau_1$. If $0 \leq x \leq \frac{1}{\alpha} \mathbf{1}_{\mathcal{M}}$, then

$$\Theta_{\tau_{\alpha}}(\alpha x) = \alpha \Theta_{\tau_{\alpha}}(x) = \alpha \varphi_{\alpha}(x) = \varphi(f^{-1}(\alpha x)) = \Theta_{\tau}(\alpha x).$$

Consequently, $\Theta_{\tau_{\alpha}} = \Theta_{\tau}$ on \mathcal{M}_+ .

If $y \in \mathcal{M}_+$, then there are $x \in \mathcal{E}(\mathcal{M})$ and $\beta \in [1, \infty)$ such that $y = \beta x$. Hence

$$\varphi(f^{-1}(y)) = \varphi(f^{-1}(\beta x)) = \beta \Theta_{\tau_{\beta}}(x) = \beta \Theta_{\tau}(x) = \Theta_{\tau}(y).$$

- (ii) According to the proof of (i), $\Theta_{\tau}(x) = \psi(x)$ for all $x \in \mathcal{E}(\mathcal{M})$. Hence Θ_{τ} and ψ coincide on \mathcal{M}_+ . \square

Theorem 4.5. *If \mathcal{M} and \mathcal{N} are AW^* -factors of Type I, then every spectral order isomorphism $\varphi : \mathcal{M}_+ \rightarrow \mathcal{N}_+$ is canonical.*

Proof. By Proposition 3.8, φ preserves central positive operators. Thus there is a strictly increasing bijection $f : [0, \infty) \rightarrow [0, \infty)$ such that $\varphi(\lambda \mathbf{1}) = f(\lambda) \mathbf{1}$ for every $\lambda \in [0, \infty)$. Let $\alpha \in [1, \infty)$. Set $\varphi_\alpha(x) = \frac{1}{\alpha} \varphi(f^{-1}(\alpha x))$ for every $x \in \mathcal{E}(\mathcal{M})$. It follows from Lemma 4.3 and Corollary 4.2 that φ_α is a canonical spectral order isomorphism from $\mathcal{E}(\mathcal{M})$ onto $\mathcal{E}(\mathcal{N})$. Therefore, we obtain from Lemma 4.4 that φ is canonical. \square

Lemma 4.6. *Let $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{M}_{sa}$ be a spectral order automorphism of the self-adjoint part of an AW^* -algebra \mathcal{M} such that*

- (i) $\varphi(\lambda \mathbf{1}) = \lambda \mathbf{1}$ for all $\lambda \in \mathbb{R}$;
- (ii) $\varphi(x) = x$ for all $x \in \mathcal{M}_+$.

If, for each $\alpha \in (0, \infty)$, the equation

$$\varphi_\alpha(x) = \varphi(x - \alpha \mathbf{1}) + \alpha \mathbf{1}$$

defines a canonical spectral order automorphism $\varphi_\alpha : \mathcal{M}_+ \rightarrow \mathcal{M}_+$, then $\varphi(x) = x$ for every $x \in \mathcal{M}_{sa}$.

Proof. Let $x \in \mathcal{M}_{sa}$. Then there exists $\alpha \in (0, \infty)$ such that $x + \alpha \mathbf{1}$ is positive invertible element. Hence $\beta \mathbf{1} \leq x + \alpha \mathbf{1}$ for some $\beta \in (0, \infty)$ and so $\alpha \mathbf{1} \leq \frac{\alpha}{\beta}(x + \alpha \mathbf{1})$. As φ is the identity map on positive operators, we have

$$\frac{\alpha}{\beta}(x + \alpha \mathbf{1}) = \varphi\left(\frac{\alpha}{\beta}(x + \alpha \mathbf{1}) - \alpha \mathbf{1}\right) + \alpha \mathbf{1} = \varphi_\alpha\left(\frac{\alpha}{\beta}(x + \alpha \mathbf{1})\right).$$

Moreover, $\varphi_\alpha(x) = \Theta_{\tau_\alpha}(x)$ for some projection isomorphism $\tau_\alpha : P(\mathcal{M}) \rightarrow P(\mathcal{M})$ because φ_α is canonical and $\varphi_\alpha(\lambda \mathbf{1}) = \lambda \mathbf{1}$ for all $\lambda \in [0, \infty)$. Hence

$$\frac{\alpha}{\beta}(x + \alpha \mathbf{1}) = \Theta_{\tau_\alpha}\left(\frac{\alpha}{\beta}(x + \alpha \mathbf{1})\right) = \frac{\alpha}{\beta} \Theta_{\tau_\alpha}(x + \alpha \mathbf{1}) = \frac{\alpha}{\beta} \varphi_\alpha(x + \alpha \mathbf{1}).$$

Consequently,

$$x = \varphi_\alpha(x + \alpha \mathbf{1}) - \alpha \mathbf{1} = \varphi(x).$$

\square

Theorem 4.7. *If \mathcal{M} and \mathcal{N} are AW^* -factors of Type I, then every spectral order isomorphism $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{N}_{sa}$ is canonical.*

Proof. According to Proposition 3.8, $\varphi(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in \mathbb{R}$, where $f : \mathbb{R} \rightarrow \mathbb{R}$ is a strictly increasing bijection $f : \mathbb{R} \rightarrow \mathbb{R}$. We may assume without loss of generality that f is identity on \mathbb{R} . Then $f(0) = 0$ and so $\varphi(0) = 0$. Hence $\varphi(\mathcal{M}_+) = \mathcal{N}_+$. Theorem 4.5 ensures the existence of a projection isomorphism $\tau : P(\mathcal{M}) \rightarrow P(\mathcal{N})$ such that $\varphi(x) = \Theta_\tau(x)$ for all $x \in \mathcal{M}_+$.

Set $\psi(x) = \varphi(\Theta_\tau^{-1}(x))$, $x \in \mathcal{N}_{sa}$. Then $\psi : \mathcal{N}_{sa} \rightarrow \mathcal{N}_{sa}$ is a spectral order automorphism satisfying $\psi(x) = x$ for all $x \in \mathcal{N}_+$ and $\psi(\lambda \mathbf{1}) = \lambda \mathbf{1}$ for all $\lambda \in \mathbb{R}$. Take $\alpha \in (0, \infty)$ and put

$$\psi_\alpha(x) = \psi(x - \alpha \mathbf{1}) + \alpha \mathbf{1}$$

for each $x \in \mathcal{N}_+$. It is easy to see that $\psi_\alpha : \mathcal{N}_+ \rightarrow \mathcal{N}_+$ is a spectral order automorphism. By Theorem 4.5, ψ_α is canonical. From Lemma 4.6, $\psi(x) = x$ for all $x \in \mathcal{N}_{sa}$. This completes the proof. \square

5 Spectral order orthoisomorphisms

Let \mathcal{M} be an AW^* -algebra having no Type I_2 direct summand. In [12, Theorem 5], Hamhalter and Turilova proved that every spectral order orthoautomorphism φ on $\mathcal{E}(\mathcal{M})$ is given by the continuous function calculus and a Jordan $*$ -isomorphism provided that φ preserves scalar multiples of the unit. In the following theorem, we reformulate this result for spectral order orthoisomorphisms between sets of all effects. We present a proof here for the sake of completeness.

Theorem 5.1. *Let \mathcal{M} be an AW^* -algebra having no Type I_2 direct summand and let \mathcal{N} be an AW^* -algebra. If a spectral order orthoisomorphism $\varphi : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{N})$ and a strictly increasing bijection $f : [0, 1] \rightarrow [0, 1]$ satisfy $\varphi(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in [0, 1]$, then there is a unique Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that*

$$\varphi(x) = \psi(f(x))$$

for all $x \in \mathcal{E}(\mathcal{M})$.

Proof. Let p be a projection in \mathcal{M} . Since φ preserves orthogonality, $\varphi(p)$ is orthogonal to $\varphi(\mathbf{1} - p)$. Hence

$$\mathbf{1} = \varphi(\mathbf{1}) = \varphi(p \vee (\mathbf{1} - p)) = \varphi(p) \vee \varphi(\mathbf{1} - p) = \varphi(p) + \varphi(\mathbf{1} - p),$$

where we have used Lemma 3.5. Multiplying both sides by $\varphi(p)$, we see that $\varphi(p)$ is a projection. Consequently, $\varphi(P(\mathcal{M})) = P(\mathcal{N})$ and so φ restricts to an orthoisomorphism from $P(\mathcal{M})$ onto $P(\mathcal{N})$. By Dye's theorem (see Theorem 2.2), there is a Jordan $*$ -isomorphism $\varrho : \mathcal{M} \rightarrow \mathcal{N}$ such that φ and ϱ coincide on $P(\mathcal{M})$. Consider $\theta = \varrho^{-1} \circ \varphi$. The map θ is a spectral order orthoautomorphism from $\mathcal{E}(\mathcal{M})$ onto itself such that $\theta(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in [0, 1]$. According to [12, Theorem 5], we obtain the required conclusion. \square

In the case of an AW^* -factor \mathcal{M} that is not of Type I_2 and Type III, it was proved by Hamhalter and Turilova [13] that every spectral order orthoautomorphism φ on $\mathcal{E}(\mathcal{M})$ automatically preserves scalar multiples of the unit and so we have a complete description of such automorphisms. However, the question of Type III factors was left open in [13]. It turns out that the case of Type III factors does not lead to any different behavior.

Corollary 5.2. *Let \mathcal{M} and \mathcal{N} be AW^* -factors that are not of Type I_2 . Let $\varphi : \mathcal{E}(\mathcal{M}) \rightarrow \mathcal{E}(\mathcal{N})$ be a spectral order orthoisomorphism. Then there are a unique strictly increasing bijection $f : [0, 1] \rightarrow [0, 1]$ and a unique Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that*

$$\varphi(x) = \psi(f(x))$$

for all $x \in \mathcal{E}(\mathcal{M})$.

Proof. The statement is a direct consequence of Proposition 3.8 and Theorem 5.1. \square

The previous results can be generalized to spectral order isomorphisms between positive parts (or between self-adjoint parts) of AW^* -algebras as follows.

Theorem 5.3. *Let \mathcal{M} be an AW^* -algebra having no Type I_2 direct summand and let \mathcal{N} be an AW^* -algebra. If a spectral order orthoisomorphism $\varphi : \mathcal{M}_+ \rightarrow \mathcal{N}_+$ and a strictly increasing bijection $f : [0, \infty) \rightarrow [0, \infty)$ satisfy $\varphi(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in [0, \infty)$, then there is a unique Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that*

$$\varphi(x) = \psi(f(x))$$

for all $x \in \mathcal{M}_+$.

Proof. Let $\alpha \in [1, \infty)$. We see from Lemma 4.3 that

$$\varphi_\alpha(x) = \frac{1}{\alpha} \varphi(f^{-1}(\alpha x))$$

defines a spectral order orthoisomorphism from $\mathcal{E}(\mathcal{M})$ onto $\mathcal{E}(\mathcal{N})$. It follows from Theorem 5.1 that φ_α coincides with a Jordan $*$ -isomorphism on $\mathcal{E}(\mathcal{M})$. Thus Lemma 4.4 ensures the existence of a Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that $\varphi(x) = \psi(f(x))$ for all $x \in \mathcal{M}_+$. \square

Corollary 5.4. *Let \mathcal{M} and \mathcal{N} be AW^* -factors that are not of Type I_2 . Let $\varphi : \mathcal{M}_+ \rightarrow \mathcal{N}_+$ be a spectral order orthoisomorphism. Then there are a unique strictly increasing bijection $f : [0, \infty) \rightarrow [0, \infty)$ and a unique Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that*

$$\varphi(x) = \psi(f(x))$$

for all $x \in \mathcal{M}_+$.

Proof. It follows from Proposition 3.8 and Theorem 5.3. \square

Let us fix a notation. We denote by x^+ and x^- , respectively, the positive part and the negative part of a self-adjoint operator x in an AW^* -algebra.

Lemma 5.5. *Let $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{N}_{sa}$ be a spectral order isomorphism between self-adjoint parts of AW^* -algebras \mathcal{M} and \mathcal{N} with $\varphi(0) = 0$. If $x \in \mathcal{M}_{sa}$, then $\varphi(x)^+ = \varphi(x^+)$ and $\varphi(x)^- = -\varphi(-(x^-))$.*

Proof. Using Lemma 3.5 and the fact that multiplication by -1 is order-reversing map, $y \vee 0 = y^+$ and $y \wedge 0 = -y^-$ for each $y \in \mathcal{M}_{sa}$. Hence $\varphi(x)^+ = \varphi(x^+)$ and $\varphi(x)^- = -\varphi(-(x^-))$. \square

Theorem 5.6. *Let \mathcal{M} be an AW^* -algebra having no Type I_2 direct summand and let \mathcal{N} be an AW^* -algebra. If a spectral order orthoisomorphism $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{N}_{sa}$ and a strictly increasing bijection $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfy $\varphi(\lambda \mathbf{1}_{\mathcal{M}}) = f(\lambda) \mathbf{1}_{\mathcal{N}}$ for all $\lambda \in \mathbb{R}$, then there is a unique Jordan $*$ -isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that*

$$\varphi(x) = \psi(f(x))$$

for all $x \in \mathcal{M}_{sa}$.

Proof. As φ preserves orthogonality and 0 is orthogonal to $\lambda \mathbf{1}_{\mathcal{M}}$ for all $\lambda \in \mathbb{R}$, we have $\varphi(0) = f(0) \mathbf{1}_{\mathcal{N}} = 0$. We infer from this that $\varphi(\mathcal{M}_+) = \mathcal{N}_+$. By Theorem 5.3, there is a unique Jordan *-isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that $\varphi(x) = \psi(f(x))$ for all $x \in \mathcal{M}_+$.

Consider a spectral order orthoautomorphism $\varrho : \mathcal{N}_{sa} \rightarrow \mathcal{N}_{sa}$ defined by

$$\varrho(x) = -\varphi(f^{-1}(\psi^{-1}(-x))).$$

In order to complete the proof, we need to show that $\varrho(x) = x$ for all $x \in \mathcal{N}_{sa}$. It follows from the above discussion that ϱ is the identity map on $\mathcal{N}_- = -\mathcal{N}_+$, $\varrho(\mathcal{N}_+) = \mathcal{N}_+$, and $\varrho(\lambda \mathbf{1}) = \lambda \mathbf{1}$ for all $\lambda \in \mathbb{R}$. By virtue of Theorem 5.3, there exists a unique Jordan *-isomorphism $\tilde{\psi} : \mathcal{N} \rightarrow \mathcal{N}$ such that $\varrho(x) = \tilde{\psi}(x)$ for all $x \in \mathcal{N}_+$. Let $p \in P(\mathcal{N})$. Entirely similar discussion to that of the proof of Theorem 5.1 leads to the equation $\mathbf{1} = \rho(p) + \rho(\mathbf{1} - p)$. Upon multiplying both sides of the equation by $\varrho(-p)$, we get

$$\varrho(-p) = \varrho(p)\varrho(-p).$$

Moreover, ϱ is the identity map on \mathcal{N}_- and so

$$-\mathbf{1} = \varrho(-p) + \varrho(-(\mathbf{1} - p)).$$

We multiply both sides of the last equation by $\varrho(p)$ to obtain

$$-\varrho(p) = \varrho(p)\varrho(-p) = \varrho(-p).$$

Hence $p = \tilde{\psi}(p)$. As $p \in P(\mathcal{N})$ is arbitrary, $\tilde{\psi}(x) = x$ for all $x \in \mathcal{N}$. This means that $\varrho(x) = x$ for all $x \in \mathcal{N}_+ \cup \mathcal{N}_-$.

Now we assume that $x \in \mathcal{N}_{sa}$. We see from Lemma 5.5 that

$$\varrho(x) = \varrho(x)^+ - \varrho(x)^- = \varrho(x^+) + \varrho(-(x^-)) = x^+ - x^- = x.$$

□

Corollary 5.7. *Let \mathcal{M} and \mathcal{N} be AW*-factors that are not of Type I_2 . Let $\varphi : \mathcal{M}_{sa} \rightarrow \mathcal{N}_{sa}$ be a spectral order orthoisomorphism. Then there are a unique strictly increasing bijection $f : \mathbb{R} \rightarrow \mathbb{R}$ and a unique Jordan *-isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{N}$ such that*

$$\varphi(x) = \psi(f(x))$$

for all $x \in \mathcal{M}_{sa}$.

Proof. The statement is a direct consequence of Proposition 3.8 and Theorem 5.6.

□

Acknowledgement

This work was supported by the project OPVVV Center for Advanced Applied Science CZ.02.1.01/0.0/0.0/16.019/0000778 and the “Grant Agency of the Czech Republic” grant number 17-00941S, “Topological and geometrical properties of Banach spaces and operator algebras II”.

References

- [1] M. K. Bennett: *Affine and Projective Geometry*, Wiley, New York, 1995.
- [2] S. K. Berberian: *Bear \ast -rings*, Springer, Berlin, 2011.
- [3] M. Bohata: *Vigiers theorem for the spectral order and its applications*, J. Math. Anal. Appl. **476**, 801–810 (2019).
- [4] A. Döring and B. Dewitt: *Self-adjoint operators as functions I*, Commun. Math. Phys. **328**, 499–525 (2014).
- [5] H. A. Dye: *On the geometry of projections in certain operator algebras*, Ann. of Math. **61** (1955). 73–89.
- [6] P. A. Fillmore and W. E. Longstaff: *On isomorphisms of lattices of closed subspaces*, Can. J. Math. **36**, 820–829 (1984).
- [7] D. J. Foulis and S. Pulmannová: *Spectral order on a synaptic algebras*, Order **36**, 1–17 (2019).
- [8] H. F. de Groote: *On a canonical lattice structure on the effect algebra of a von Neumann algebra*, arXiv:math-ph/0410018v2.
- [9] H. Halvarson: *Deep Beauty*, Cambridge University Press, New York, 2011.
- [10] J. Hamhalter: *Spectral order of operators and range projections*, J. Math. Anal. Appl. **331**, 1122–1134 (2007).
- [11] J. Hamhalter: *Dye’s theorem and Gleason’s theorem for AW^* -algebras*, J. Math. Anal. Appl. **422**, 1103–1115 (2015).
- [12] J. Hamhalter and E. Turilova: *Spectral order on AW^* -algebras and its preservers*, Lobachevskii J. Math. **37**, 439–448 (2016).
- [13] J. Hamhalter and E. Turilova: *Quantum spectral symmetries*, Int. J. Theor. Phys. **56**, 3807–3818 (2017).
- [14] R. V. Kadison: *Order properties of bounded self-adjoint operators*, Proc. Amer. Math. Soc. **2**, 505–510 (1951).
- [15] I. Kaplansky: *Algebras of Type I*, Ann. Math. **56**, 460–472 (1952).
- [16] L. Molnár and G. Nagy: *Spectral order automorphisms of the spaces of Hilbert space effects and observables: The 2-dimensional case*, Lett. Math. Phys. **106**, 535–544 (2016).
- [17] L. Molnár and P. Šemrl: *Spectral order automorphisms of the spaces of Hilbert space effects and observables*, Lett. Math. Phys. **80**, 239–255 (2007).
- [18] M. P. Olson: *The self-adjoint operators of a von Neumann algebra form a conditionally complete lattice*, Proc. Amer. Math. Soc. **28**, 537–544 (1971).

- [19] A. Planetta and J. Stochel: *Spectral order for unbounded operators*, J. Math. Anal. Appl. **389**, 1029–1045 (2012).
- [20] S. Roman: *Lattices and Ordered Sets*, Springer, New York, 2008.
- [21] K. Saitô and J.D.M. Wright: *Monotone Complete C^* -algebras and Generic Dynamics*, Springer, London, 2015.
- [22] S. Sherman: *Order in operator algebras*, Amer. J. Math. **73**, 227–232 (1951).
- [23] E. Turilova: *Automorphisms of spectral lattices of unbounded positive operators*, Lobachevskii J. Math. **35**, 259–263 (2014).
- [24] S. Wolters: *Topos models for physics and topos theory*, J. Math. Phys. **55**, 082110 (2014).