Rényi and von Neumann entropies for various Bipartite Gaussian States

DaeKil Park[1,2*]

[1] *Department of Electronic Engineering, Kyungnam University, Changwon 631-701, Korea*

[2] *Department of Physics, Kyungnam University, Changwon 631-701, Korea*

arXiv:1911.07955v1 [quant-ph] 18 Nov 2019

Abstract

The Rényi and von Neumann entropies of various bipartite Gaussian states are derived analytically. We also discuss on the tripartite purification for the bipartite states when some particular conditions hold. The generalization to non-Gaussian states is briefly discussed.

* dkpark@kyungnam.ac.kr

I. INTRODUCTION

As IC (integrated circuit) becomes smaller and smaller, the effect of quantum mechanics becomes prominent more and more. As a result, quantum technology (technology based on quantum information theories[1]) becomes important more and more recently. The representative constructed by quantum technology is a quantum computer[2], which was realized recently by making use of superconducting qubits.

In quantum information processing entanglement[3, 4] plays an important role as a physical resource. It is used in various quantum information processing, such as quantum teleportation[5, 6], superdense coding[7], quantum cloning[8], quantum cryptography[9, 10], quantum metrology[11], and quantum computer[2, 12, 13]. Furthermore, with many researchers trying to realize such quantum information processing in the laboratory for the last few decades, quantum cryptography and quantum computer seem to approaching the commercial level[14, 15].

For last few decades many entanglement measures have been developed. For example, the entanglement of formation (EoF)[16], one of the measure for bipartite system, is defined as a von Neumann entropy of the substate in bipartite quantum state. In two qubit system the EoF of any pure and mixed states can be computed if one follows the Wootters procedure[17, 18]. However, computation of EoF in qudit or continuous-variable (CV) system is extremely difficult problem except few rare cases. In this reason recent quantum technology is developed by making use of qubits generated through various physical setup.

Although quantum information theories have been developed mainly with qubit or qudit system, entanglement of CV states was also studied for last few years. In qubit and qudit systems the necessary and sufficient criterion of separability for given quantum state is a positive partial transposition (PPT)[19–21], which is valid for only 2×2 qubit-qubit and 2×3 qubit-qudit states. However, as Ref. [22, 23] have shown, PPT also provides a necessary and sufficient criterion for the separability of Gaussian CV quantum states. Thus, the entanglement measure like negativity[24] can be used as a quantification of entanglement in the Gaussian states. Furthermore, the distillation protocols for partially entangled Gaussian state to a maximally entangled state have already been suggested in Duan et al.[25] and Giedke et al.[26]. Quantum information processing with CV was reviewed from both theoretical and experimental aspects[27, 28].

The analytical expression of von Neumann entropy was derived for a general real Gaussian density matrix in Ref. [29] and it was generalized to massless scalar field in Ref. [30]. Putting the scalar field system in the spherical box, the author in Ref. [30] has shown that the total entropy of the system is proportional to surface area. This may give some insight into the reason why the black hole entropy is proportional to the horizon area. Recently, the entanglement was computed in the coupled harmonic oscillator system using a Schmidt decomposition[31]. The von Neumann and Rényi entropies were also explicitly computed in the similar system, called two site Bose-Hubbard model[32]. More recently, the Rényi and von Neumann entropies in bipartite[33], tripartite[34], and thermal[35] harmonic oscillator systems were computed when the spring and coupling constants are arbitrarily time-dependent.

In this paper we will compute the Rényi and von Neumann entropies of various bipartite Gaussian states, which can be substates of muitipartite Gaussian states or their partial transpose states. Thus, the bipartite states we consider in this paper can be physically relevant or irrelevant. We categorize the bipartite Gaussian states into four types and compute their Rényi and von Neumann entropies. Probably, many bipartite Gaussian states belong to one of four types.

The paper is organized as follows. In section II we compute the Rényi and von Neumann entropies for the most general single-party Gaussian states. The results of this section are used in next section if the eigenvalue equation of the bipartite density matrix can be factorized into two single-party eigenvalue equations. In section III we compute the Rényi and von Neumann entropies of four types of bipartite Gaussian states analytically. For type I, II, and III the entropies are exactly derived. For the case of type IV we cannot find a suitable change of variables, which factorizes the eigenvalue equation into two single-party eigenvalue equations. We compute the entropies of type IV under some assumption (see Eq. (3.2)). In Sec. IV a brief conclusion is given.

II. THE MOST GENERAL SINGLE-PARTY GAUSSIAN STATE

Before we consider various bipartite Gaussian states in next section, we derive the entropies and spectral decomposition of the most general single-party Gaussian quantum state

given by

$$\rho_0[x', x] = Ae^{-a_1x^2 - a_2x'^2 + 2bxx'} \tag{2.1}$$

where $A = \sqrt{\frac{a_1+a_2-2b}{\pi}}$. In order to compute the Rényi and von Neumann entropies for ρ_0 we should solve the following eigenvalue equation:

$$\int \rho_0[x', x] f_n(x) dx = \lambda_n f_n(x') \tag{2.2}$$

with $n = 0, 1, 2, \cdots$. One can show directly that the eigenfunction is

$$f_n(x) = \mathcal{C}_n^{-1} H_n(\sqrt{\epsilon_0}x) e^{-\frac{\alpha_0}{2}x^2} \tag{2.3}$$

where $\epsilon_0 = \sqrt{(a_1+a_2)^2 - 4b^2}$ and $\alpha_0 = \epsilon_0 - (a_1 - a_2)$, and $H_n(z)$ is n^{th}-order Hermite polynomial. By making use of integral formula[36]

$$\begin{aligned} &\int_{-\infty}^{\infty} e^{-(x-y)^2} H_m(cx) H_n(cx) dx \\ &= \sqrt{\pi} \sum_{k=0}^{\min(m,n)} 2^k k! \begin{pmatrix} m \\ k \end{pmatrix} \begin{pmatrix} n \\ k \end{pmatrix} (1-c^2)^{\frac{m+n}{2}-k} H_{m+n-2k}\left(\frac{cy}{\sqrt{1-c^2}}\right) \end{aligned} \tag{2.4}$$

and various properties of Gamma function[37], the normalization constant $\mathcal{C}_n$ can be written in a form

$$\mathcal{C}_n^2 = \frac{1}{\sqrt{\alpha_0}} \sum_{k=0}^{n} 2^{2n-k} \left(\frac{\epsilon_0}{\alpha_0} - 1\right)^{n-k} \frac{\Gamma^2(n+1)\Gamma\left(n-k+\frac{1}{2}\right)}{\Gamma(k+1)\Gamma^2(n-k+1)}. \tag{2.5}$$

If $a_1 = a_2$, $\alpha_0 = \epsilon_0$, which makes nonzero in k-summation of Eq. (2.5) only when $k = n$. Then, $\mathcal{C}_n$ becomes usual harmonic oscillator normalization constant

$$\mathcal{C}_n^{-1} = \frac{1}{\sqrt{2^n n!}} \left(\frac{\epsilon_0}{\pi}\right)^{1/4}. \tag{2.6}$$

It is straightforward to show that the eigenvalues in Eq. (2.2) become

$$\lambda_n = A\sqrt{\frac{2\pi}{(a_1+a_2)+\epsilon_0}} \left[\frac{(a_1+a_2)-\epsilon_0}{(a_1+a_2)+\epsilon_0}\right]^{n/2} = (1-\xi_0)\xi_0^n \tag{2.7}$$

where

$$\xi_0 = \frac{2b}{a_1+a_2+\epsilon_0} = \frac{\sqrt{a_1+a_2+2b} - \sqrt{a_1+a_2-2b}}{\sqrt{a_1+a_2+2b} + \sqrt{a_1+a_2-2b}}. \tag{2.8}$$

Eq. (2.7) is consistent with $\mathrm{tr}\rho_0 = 1$ if $|\xi_0| < 1$. Thus, ρ_0 is physical quantum state if $0 < \xi_0 < 1$. If this condition does not hold, ρ_0 is unphysical Gaussian state. Anyway, the Rényi and von Neumann entropies for ρ_0 are

$$S_{0,\alpha} = \frac{1}{1-\alpha} \ln \frac{(1-\xi_0)^\alpha}{1-\xi_0^\alpha} \qquad S_{0,von} = -\ln(1-\xi_0) - \frac{\xi_0}{1-\xi_0} \ln \xi_0. \tag{2.9}$$

Finally, the spectral decomposition of ρ_0 can be written as

$$\rho_0[x', x] = \sum_n \lambda_n f_n(x') f_n^*(x). \tag{2.10}$$

III. SEVERAL BIPARTITE GAUSSIAN STATES

In this section we consider four types of bipartite Gaussian states $\rho_j[x_1', x_2' : x_1, x_2]$ with j = I, II, III, IV. They can be physically relevant or irrelevant depending on their eigenvalues. In order to compute the Rényi and von Neumann entropies of ρ_j, we should solve the following eigenvalue equation

$$\int dx_1 dx_2 \rho_j[x_1', x_2' : x_1, x_2] f_{mn}(x_1, x_2) = \lambda_{mn} f_{mn}(x_1', x_2') \tag{3.1}$$

with $m, n = 0, 1, 2, \cdots$. Since $\mathrm{tr}\rho_j = 1$, the eigenvalue λ_{mn} obeys $\sum_{m,n} \lambda_{mn} = 1$. In this reason the eigenvalues of ρ_j can be represented as

$$\lambda_{mn} = (1 - \xi_1)\xi_1^m (1 - \xi_2)\xi_2^n, \tag{3.2}$$

where $0 < |\xi_1|, |\xi_2| < 1$. If $0 < \xi_1, \xi_2 < 1$, the bipartite Gaussian states are physically relevant. If not, they are unphysical states. Then, the Rényi and von Neumann entropies of ρ_j become

$$S_\alpha \equiv \frac{1}{1-\alpha} \ln \mathrm{tr}\rho_j^\alpha = S_{1,\alpha} + S_{2,\alpha} \qquad S_{von} \equiv \lim_{\alpha \to 1} S_\alpha = S_{1,von} + S_{2,von} \tag{3.3}$$

where

$$S_{k,\alpha} = \frac{1}{1-\alpha} \ln \frac{(1-\xi_k)^\alpha}{1-\xi_k^\alpha} \qquad S_{k,von} = -\ln(1-\xi_k) - \frac{\xi_k}{1-\xi_k} \ln \xi_k \tag{3.4}$$

with $k = 1, 2$. Thus, in the following we will derive ξ_1 and ξ_2 for each ρ_j analytically.

A. Type I Bipartite State

In this subsection we examine the following bipartite Gaussian state:

$$\rho_I[x_1', x_2' : x_1, x_2] = A \exp\Big[-a_1(x_1'^2 + x_1^2) - a_2(x_2'^2 + x_2^2) + 2b(x_1'x_2' + x_1x_2) \\ +2c(x_1x_1' + x_2x_2') + 2f(x_1x_2' + x_2x_1') \Big] \tag{3.5}$$

where $A = 2\sqrt{(a_1-c)(a_2-c)-(b+f)^2}/\pi$. We will call this type of Gaussian states by type I. The purity function of ρ_I is

$$\mathrm{tr}\rho_I^2 = \sqrt{\frac{(a_1-c)(a_2-c)-(b+f)^2}{(a_1+c)(a_2+c)-(b-f)^2}}. \tag{3.6}$$

Thus, if $c = f = 0$, ρ_I becomes pure state. For other cases it is mixed state.

Now, let us solve the eigenvalue equation (3.1) (with $j = \mathrm{I}$) explicitly. First, we change the variables as

$$\begin{pmatrix} X_1 \\ X_2 \end{pmatrix} = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \tag{3.7}$$

where

$$\theta = \tan^{-1}\frac{2b}{\sqrt{(a_1-a_2)^2+4b^2}+(a_1-a_2)}. \tag{3.8}$$

In terms of new variables the eigenvalue equation (3.1) can be written as

$$\begin{aligned} Ae^{-\mu_+X_1'^2-\mu_-X_2'^2}\int dX_1dX_2e^{-\mu_+X_1^2-\mu_-X_2^2+2C_1X_1X_1'+2C_2X_2X_2'+2F(X_1X_2'+X_2X_1')}f_{mn}(X_1,X_2) \\ = \lambda_{mn}f_{mn}(X_1',X_2') \end{aligned} \tag{3.9}$$

where

$$\begin{aligned} \mu_\pm &= \frac{1}{2}\left[(a_1+a_2)\pm\sqrt{(a_1-a_2)^2+4b^2}\right] \\ C_1 = c-f\sin 2\theta \qquad C_2 &= c+f\sin 2\theta \qquad F = f\cos 2\theta. \end{aligned} \tag{3.10}$$

In order to simplify Eq. (3.9) some more we define

$$y_1 = \sqrt{\mu_+}X_1 \qquad\qquad y_2 = \sqrt{\mu_-}X_2. \tag{3.11}$$

Then, Eq. (3.9) reduces to

$$\begin{aligned} \frac{A}{\sqrt{\mu_+\mu_-}}e^{-y_1'^2-y_2'^2}\int dy_1dy_2\exp\left[-y_1^2-y_2^2+\frac{2C_1}{\mu_+}y_1y_1'+\frac{2C_2}{\mu_-}y_2y_2'+\frac{2F}{\sqrt{\mu_+\mu_-}}(y_1y_2'+y_2y_1')\right] \\ \times f_{mn}(y_1,y_2) = \lambda_{mn}f_{mn}(y_1',y_2'). \end{aligned} \tag{3.12}$$

Finally, we change the variables again as

$$\begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix} = \begin{pmatrix} \cos\varphi & -\sin\varphi \\ \sin\varphi & \cos\varphi \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \tag{3.13}$$

where

$$\varphi = \tan^{-1}\left[\frac{-2F/\sqrt{\mu_+\mu_-}}{\sqrt{\left(\frac{C_1}{\mu_+}-\frac{C_2}{\mu_-}\right)^2+\frac{4F^2}{\mu_+\mu_-}}+\left(\frac{C_1}{\mu_+}-\frac{C_2}{\mu_-}\right)}\right]. \tag{3.14}$$

In terms of the new coordinates Eq. (3.12) is simplified in a form

$$\frac{A}{\sqrt{\mu_+\mu_-}}e^{-Y_1'^2-Y_2'^2}\int dY_1 dY_2 e^{-Y_1^2-Y_2^2+2\nu_+Y_1Y_1'+2\nu_-Y_2Y_2'}f_{mn}(Y_1,Y_2)=\lambda_{mn}f_{mn}(Y_1',Y_2') \tag{3.15}$$

where

$$\nu_\pm = \frac{1}{2}\left[\left(\frac{C_1}{\mu_+}+\frac{C_2}{\mu_-}\right)\pm\sqrt{\left(\frac{C_1}{\mu_+}-\frac{C_2}{\mu_-}\right)^2+\frac{4F^2}{\mu_+\mu_-}}\right]. \tag{3.16}$$

Now, we define

$$f_{mn}(Y_1,Y_2)=G_{1,m}(Y_1)G_{2,n}(Y_2). \tag{3.17}$$

Then, the eigenvalue equation (3.15) is solved if one solves the following two single-party eigenvalue equations:

$$\begin{aligned} e^{-Y_1'^2}\int dY_1 e^{-Y_1^2+2\nu_+Y_1Y_1'}G_{1,m}(Y_1)&=p_mG_{1,m}(Y_1') \\ e^{-Y_2'^2}\int dY_2 e^{-Y_2^2+2\nu_-Y_2Y_2'}G_{2,n}(Y_2)&=q_nG_{2,n}(Y_2'). \end{aligned} \tag{3.18}$$

Using the results of the previous section it is easy to show that the eigenvalues p_m and q_n are

$$p_m=\sqrt{\frac{\pi}{1+\epsilon_+}}\left(\frac{1-\epsilon_+}{1+\epsilon_+}\right)^{m/2} \qquad q_n=\sqrt{\frac{\pi}{1+\epsilon_-}}\left(\frac{1-\epsilon_-}{1+\epsilon_-}\right)^{n/2} \tag{3.19}$$

where $\epsilon_\pm=\sqrt{1-\nu_\pm^2}$. Then, the eigenvalue $\lambda_{mn}=Ap_mq_n/\sqrt{\mu_+\mu_-}$ can be represented as Eq. (3.2), where

$$\xi_1=\frac{\nu_+}{1+\epsilon_+}=\frac{\sqrt{1+\nu_+}-\sqrt{1-\nu_+}}{\sqrt{1+\nu_+}+\sqrt{1-\nu_+}} \qquad \xi_2=\frac{\nu_-}{1+\epsilon_-}=\frac{\sqrt{1+\nu_-}-\sqrt{1-\nu_-}}{\sqrt{1+\nu_-}+\sqrt{1-\nu_-}}. \tag{3.20}$$

Thus, the Rényi and von Neumann entropies of ρ_I can be explicitly computed by inserting ξ_1 and ξ_2 in Eq. (3.20) into Eq. (3.3) and (3.4). Applying the results of the previous section one can derive the normalized eigenfunction, whose explicit expression is

$$f_{mn}(x_1,x_2)=\left[\frac{1}{\sqrt{2^m m!}}\left(\frac{2\epsilon_+}{\pi}\right)^{1/4}H_m(\sqrt{2\epsilon_+}Y_1)e^{-\epsilon_+Y_1^2}\right]\left[\frac{1}{\sqrt{2^n n!}}\left(\frac{2\epsilon_-}{\pi}\right)^{1/4}H_n(\sqrt{2\epsilon_-}Y_2)e^{-\epsilon_-Y_2^2}\right]. \tag{3.21}$$

Thus, the spectral decomposition of ρ_I is

$$\rho_I[x_1', x_2' : x_1, x_2] = \sum_{m,n} \lambda_{mn} f_{mn}(x_1', x_2') f_{mn}^*(x_1, x_2). \tag{3.22}$$

If $c = \pm f$, either ξ_1 or ξ_2 is zero depending on the signs of c, f, and $(\mu_+ + \mu_-) \pm (\mu_+ - \mu_-)\sin 2\theta$. This fact implies from Schmidt decomposition that ρ_I can be purified to tripartite pure state. If, for example, $c = f \equiv z$, the tripartite pure state can be constructed as

$$\sigma_{ABC}[x_1', x_2', x_3' : x_1, x_2, x_3] = \psi(x_1', x_2', x_3')\psi^*(x_1, x_2, x_3) \tag{3.23}$$

where

$$\psi(x_1, x_2, x_3) = \frac{1}{\pi^{3/4}} \exp\Bigg[-(a_1 + z)x_1^2 - (a_2 + z)x_2^2 - \frac{x_3^3}{8[(a_1 - z)(a_2 - z) - (b + z)^2]} + 2(b - z)x_1 x_2 - \sqrt{\frac{z}{(a_1 - z)(a_2 - z) - (b + z)^2}}(x_1 + x_2)x_3 \Bigg]. \tag{3.24}$$

Then, ρ_I is constructed by $\rho_I = \mathrm{tr}_C \sigma_{ABC}$. If $c = -f \equiv z$, $\psi(x_1, x_2, x_3)$ in Eq. (3.24) is changed into

$$\psi(x_1, x_2, x_3) = \frac{1}{\pi^{3/4}} \exp\Bigg[-(a_1 + z)x_1^2 - (a_2 + z)x_2^2 - \frac{x_3^3}{8[(a_1 - z)(a_2 - z) - (b - z)^2]} + 2(b + z)x_1 x_2 - \sqrt{\frac{z}{(a_1 - z)(a_2 - z) - (b - z)^2}}(x_1 - x_2)x_3 \Bigg]. \tag{3.25}$$

B. Type II Bipartite State

In this subsection we examine the following bipartite Gaussian state:

$$\rho_{II}[x_1', x_2' : x_1, x_2] = A \exp\Big[-a_1(x_1'^2 + x_2'^2) - a_2(x_1^2 + x_2^2) + 2b_1 x_1' x_2' + 2b_2 x_1 x_2 + 2c(x_1 x_1' + x_2 x_2') + 2f(x_1 x_2' + x_2 x_1') \Big] \tag{3.26}$$

where $A = \sqrt{(a_1 + a_2 - 2c)^2 - (b_1 + b_2 + 2f)^2}/\pi$. We will call this type of Gaussian states by type II. This type of Gaussian state does not arise in the usual harmonic oscillator systems. This means that this type of states cannot be obtained as a substate of mutipartite pure state in the usual harmonic oscillator system. However, this type appears in Ref.[35] as a thermal state in the two-coupled harmonic oscillator system. The purity function of ρ_{II} is

$$\mathrm{tr}\rho_{II}^2 = \sqrt{\frac{(a_1 + a_2 - 2c)^2 - (b_1 + b_2 + 2f)^2}{(a_1 + a_2 + 2c)^2 - (b_1 + b_2 - 2f)^2}}. \tag{3.27}$$

Thus, if $c = f = 0$, ρ_{II} is pure Gaussian state. If not, it is mixed state.

Now, let us solve the eigenvalue equation (3.1) (with $j = \mathrm{II}$) analytically. First we change the variables as

$$y_1 = \frac{1}{\sqrt{2}}(x_1 + x_2) \qquad\qquad y_2 = \frac{1}{\sqrt{2}}(x_1 - x_2). \tag{3.28}$$

Then, Eq. (3.1) is simplified as

$$\begin{aligned} Ae^{-(a_1-b_1)y_1'^2-(a_1+b_1)y_2'^2} \int dy_1 dy_2 e^{-(a_2-b_2)y_1^2-(a_2+b_2)y_2^2+2(c+f)y_1'y_1+2(c-f)y_2'y_2} f_{mn}(y_1, y_2) \\ = \lambda_{mn} f_{mn}(y_1', y_2'). \end{aligned} \tag{3.29}$$

If we define

$$f_{mn}(y_1, y_2) = g_m(y_1)h_n(y_2), \tag{3.30}$$

Eq. (3.29) can be solved if one solves the following two single-party eigenvalue equations:

$$\begin{aligned} e^{-(a_1-b_1)y_1'^2} \int dy_1 e^{-(a_2-b_2)y_1^2+2(c+f)y_1'y_1} g_m(y_1) = p_m g_m(y_1') \\ e^{-(a_1+b_1)y_2'^2} \int dy_2 e^{-(a_2+b_2)y_2^2+2(c-f)y_2'y_2} h_n(y_2) = q_n h_n(y_2'). \end{aligned} \tag{3.31}$$

The solutions of Eq. (3.31) are analytically derived by using the results of the previous section. Then, eigenvalue of Eq. (3.1) is $\lambda_{mn} = Ap_m q_n$, which can be represented as Eq. (3.2), where

$$\begin{aligned} \xi_1 &= \frac{2(c+f)}{(a_1+a_2-b_1-b_2)+\epsilon_1} \\ &= \frac{\sqrt{(a_1+a_2-b_1-b_2)+2(c+f)} - \sqrt{(a_1+a_2-b_1-b_2)-2(c+f)}}{\sqrt{(a_1+a_2-b_1-b_2)+2(c+f)} + \sqrt{(a_1+a_2-b_1-b_2)-2(c+f)}} \\ \xi_2 &= \frac{2(c-f)}{(a_1+a_2+b_1+b_2)+\epsilon_2} \\ &= \frac{\sqrt{(a_1+a_2+b_1+b_2)+2(c-f)} - \sqrt{(a_1+a_2+b_1+b_2)-2(c-f)}}{\sqrt{(a_1+a_2+b_1+b_2)+2(c-f)} + \sqrt{(a_1+a_2+b_1+b_2)-2(c-f)}} \end{aligned} \tag{3.32}$$

with

$$\epsilon_1 = \sqrt{(a_1+a_2-b_1-b_2)^2 - 4(c+f)^2} \qquad \epsilon_2 = \sqrt{(a_1+a_2+b_1+b_2)^2 - 4(c-f)^2}. \tag{3.33}$$

Thus, the Rényi and von Neumann entropies of ρ_{II} can be explicitly computed by inserting ξ_1 and ξ_2 in Eq. (3.32) into Eq. (3.3) and (3.4).

Following the analysis in section II one can also derive the normalized eigenfunction, whose explicit expression is

$$f_{mn}(x_1,x_2)=\left(\mathcal{C}_{1,m}^{-1}H_m(\sqrt{\epsilon_1}y_1)e^{-\frac{\alpha_1}{2}y_1^2}\right)\left(\mathcal{C}_{2,n}^{-1}H_n(\sqrt{\epsilon_2}y_2)e^{-\frac{\alpha_2}{2}y_2^2}\right) \tag{3.34}$$

where

$$\alpha_1=\epsilon_1+(a_1-a_2)-(b_1-b_2)\qquad \alpha_2=\epsilon_2+(a_1-a_2)+(b_1-b_2) \tag{3.35}$$

and the normalization constants $\mathcal{C}_{1,m}$ and $\mathcal{C}_{2,n}$ are

$$\begin{aligned}\mathcal{C}_{1,m}^2&=\frac{1}{\sqrt{\alpha_1}}\sum_{k=0}^{m}2^{2m-k}\left(\frac{\epsilon_1}{\alpha_1}-1\right)^{m-k}\frac{\Gamma^2(m+1)\Gamma(m-k+1/2)}{\Gamma(k+1)\Gamma^2(m-k+1)}\\ \mathcal{C}_{2,n}^2&=\frac{1}{\sqrt{\alpha_2}}\sum_{k=0}^{n}2^{2n-k}\left(\frac{\epsilon_2}{\alpha_2}-1\right)^{n-k}\frac{\Gamma^2(n+1)\Gamma(n-k+1/2)}{\Gamma(k+1)\Gamma^2(n-k+1)}.\end{aligned} \tag{3.36}$$

Thus, the spectral decomposition of ρ_{II} is

$$\rho_{II}[x_1',x_2':x_1,x_2]=\sum_{m,n}\lambda_{mn}f_{mn}(x_1',x_2')f_{mn}^*(x_1,x_2). \tag{3.37}$$

Now, let us consider a case of $a_1=a_2\equiv a$ and $b_1=b_2\equiv b$. In this case ϵ_1 and ϵ_2 become $\epsilon_1=2\sqrt{(a-b)^2-(c+f)^2}$ and $\epsilon_2=2\sqrt{(a+b)^2-(c-f)^2}$. Thus, ξ_1, ξ_2, $\mathcal{C}_{1,m}$, and $\mathcal{C}_{2,n}$ become

$$\begin{aligned}\xi_1&=\frac{c+f}{(a-b)+\frac{\epsilon_1}{2}}=\frac{\sqrt{(a-b)+(c+f)}-\sqrt{(a-b)-(c+f)}}{\sqrt{(a-b)+(c+f)}+\sqrt{(a-b)-(c+f)}}\\ \xi_2&=\frac{c-f}{(a+b)+\frac{\epsilon_2}{2}}=\frac{\sqrt{(a+b)+(c-f)}-\sqrt{(a+b)-(c-f)}}{\sqrt{(a+b)+(c-f)}+\sqrt{(a+b)-(c-f)}}\\ \mathcal{C}_{1,m}^{-1}&=\frac{1}{\sqrt{2^m m!}}\left(\frac{\epsilon_1}{\pi}\right)^{1/4}\qquad \mathcal{C}_{2,n}^{-1}=\frac{1}{\sqrt{2^n n!}}\left(\frac{\epsilon_2}{\pi}\right)^{1/4}.\end{aligned} \tag{3.38}$$

Eq. (3.38) implies that either ξ_1 or ξ_2 is zero if $c=\pm f$, which may indicate that the tripartite purification of ρ_{II} is possible. In fact, one can find a following tripartite pure state, whose substate coincides with ρ_{II}. For $c=f\equiv z$ the pure state is

$$\begin{aligned}\psi(x_1,x_2,x_3)=\frac{1}{\pi^{3/4}}\exp\Bigg[&-(a+z)(x_1^2+x_2^2)-\frac{x_3^2}{8[(a-z)^2-(b+z)^2]}\\ &+2(b-z)x_1x_2-\sqrt{\frac{z}{(a-z)^2-(b+z)^2}}(x_1+x_2)x_3\Bigg].\end{aligned} \tag{3.39}$$

If $c=-f\equiv z$, the purification is changed into

$$\begin{aligned}\psi(x_1,x_2,x_3)=\frac{1}{\pi^{3/4}}\exp\Bigg[&-(a+z)(x_1^2+x_2^2)-\frac{x_3^2}{8[(a-z)^2-(b-z)^2]}\\ &+2(b+z)x_1x_2-\sqrt{\frac{z}{(a-z)^2-(b-z)^2}}(x_1-x_2)x_3\Bigg].\end{aligned} \tag{3.40}$$

Even without the restriction $a_1 = a_2 \equiv a$ and $b_1 = b_2 \equiv b$, Eq. (3.32) implies that either ξ_1 or ξ_2 is zero at $c = \pm f$. Thus, one may conjecture that the tripartite purification of ρ_{II} might be possible at $c = \pm f$ even in this case. However, we cannot find any pure state σ_{ABC}, which gives ρ_{II} as $\rho_{II} = \mathrm{tr}_C \sigma_{ABC}$. We think this is because of the fact that ρ_{II} cannot arise in the usual zero-temperature harmonic oscillator system.

C. Type III Bipartite State

In this subsection we examine the following bipartite Gaussian state:

$$\rho_{III}\left[x_1', x_2' : x_1, x_2\right] = A \exp\Big[- a_1(x_1'^2 + x_1^2) - a_2(x_2'^2 + x_2^2) \\ +2b(x_1 x_2 + x_1' x_2') + 2c_1 x_1 x_1' + 2c_2 x_2 x_2' + 2f x_1' x_2 + 2f^* x_1 x_2' \Big] \tag{3.41}$$

where $A = 2\sqrt{(a_1 - c_1)(a_2 - c_2) - (b + f_R)^2}/\pi$, with $f_R = \mathrm{Re} f$. We will call this type of Gaussian state by type III. If $c_1 = c_2 \equiv c$ and $f = f^*$, ρ_{III} reduces to ρ_I. The type III Gaussian state arises in the reduced state of three coupled harmonic oscillator system[34]. In this reference the eigenvalue equation (3.1) (with $j = \mathrm{III}$) is analytically solved by introducing the appropriate change of variables as we did in the previous sections. In this paper, however, we will compute the entropy of ρ_{III} without solving the eigenvalue equation explicitly as follows. This method is useful in the next subsection.

First, we note that $\beta_1 = \mathrm{tr}\rho_{III}^2$ and $\beta_2 = \mathrm{tr}\rho_{III}^3$ become

$$\beta_1 = \sqrt{\frac{X_1}{X_2}} \qquad \beta_2 = \frac{4X_1}{X_1 + 3X_2 - 12x} \tag{3.42}$$

where

$$X_1 = 4\left[(a_1 - c_1)(a_2 - c_2) - (b + f_R)^2\right] \tag{3.43}$$
$$X_2 = 4\left[(a_1 + c_1)(a_2 + c_2) - (b - f_R)^2\right] \qquad x = c_1 c_2 - |f|^2.$$

It is easy to show $X_2 \pm X_1 = 8A_\pm$ and $X_1 X_2 = 16\tilde{A}$, where

$$A_+ = a_1 a_2 + c_1 c_2 - b^2 - f_R^2 \qquad A_- = a_1 c_2 + a_2 c_1 + 2b f_R \tag{3.44}$$
$$\tilde{A} = (a_1^2 - c_1^2)(a_2^2 - c_2^2) + (b^2 - f_R^2)^2 - 2(a_1 a_2 + c_1 c_2)(b^2 + f_R^2) - 4b f_R (a_1 c_2 + a_2 c_1).$$

The eigenvalue of Eq. (3.1) is shown to be represented as Eq. (3.2) in Ref. [34]. Thus, the remaining task we should do is to compute ξ_1 and ξ_2 explicitly. Putting $\alpha = 2$ or 3 in

S_α of Eq. (3.3), it is possible to derive the following equations:

$$\frac{(1-\xi_1)(1-\xi_2)}{(1+\xi_1)(1+\xi_2)} = \beta_1 \qquad \frac{(1-\xi_1)^2(1-\xi_2)^2}{(1+\xi_1+\xi_1^2)(1+\xi_2+\xi_2^2)} = \beta_2. \tag{3.45}$$

Solving Eq. (3.45), we get

$$u = \xi_1 + \xi_2 = \frac{1-\beta_1}{2(4\beta_1^2 - \beta_1^2\beta_2 - 3\beta_2)}\left[-3\beta_2(1+\beta_1) + \sqrt{3\beta_2\left[16\beta_1^2 - \beta_2(3-\beta_1)^2\right]}\right] \tag{3.46}$$
$$v = \xi_1\xi_2 = -1 + \frac{1+\beta_1}{2(4\beta_1^2 - \beta_1^2\beta_2 - 3\beta_2)}\left[-3\beta_2(1+\beta_1) + \sqrt{3\beta_2\left[16\beta_1^2 - \beta_2(3-\beta_1)^2\right]}\right].$$

Thus, ξ_1 and ξ_2 for ρ_{III} become

$$\xi_1 = \frac{u + \sqrt{u^2 - 4v}}{2} \qquad \xi_2 = \frac{u - \sqrt{u^2 - 4v}}{2}. \tag{3.47}$$

Then, the Rényi and von Neumann entropies of ρ_{III} can be explicitly computed by inserting ξ_1 and ξ_2 in Eq. (3.47) into Eq. (3.3) and (3.4).

Inserting Eq. (3.42) into Eq. (3.46), one can show straightforwardly

$$u = \frac{A_-}{x}\left[1 - \sqrt{1 - \frac{2x}{A_+ + \sqrt{\tilde{A}}}}\right] \qquad v = -1 + \frac{A_+ + \sqrt{\tilde{A}}}{x}\left[1 - \sqrt{1 - \frac{2x}{A_+ + \sqrt{\tilde{A}}}}\right]. \tag{3.48}$$

If $c_1c_2 = |f|^2$, it is easy to show $v = 0$. Thus, Eq. (3.47) indicates that either ξ_1 or ξ_2 is zero depending on sign of u when $c_1c_2 = |f|^2$. For example, if u is positive, $\xi_2 = 0$ and $\xi_1 = A_-/(A_+ + \sqrt{\tilde{A}})$. If, in this case, $c_1 = c_2 = f_R \equiv c$ with $\mathrm{Im}f = 0$, ξ_1 reduces to

$$\xi_1 = \frac{c(a_1 + a_2 + 2b)}{(a_1a_2 - b^2) + \sqrt{(a_1a_2 - b^2)^2 - c^2(a_1 + a_2 + 2b)^2}}, \tag{3.49}$$

which is exactly the same with ξ_1 of type I when $c = f$.

Above discussion implies that when $c_1c_2 = |f|^2$, there may exist a tripartite pure state $\sigma_{ABC} = \psi(x_1', x_2', x_3')\psi^*(x_1, x_2, x_3)$, which satisfies $\rho_{III} = \mathrm{tr}\sigma_{ABC}$. In fact, one can construct two parameter (θ, $\bar{x}$) family of $\psi(x_1, x_2, x_3)$, whose explicit form is

$$\begin{aligned}&\psi(x_1, x_2, x_3 : \theta, \bar{x}) \\ &= \sqrt{\mathcal{N}}\exp\left[-r_1e^{-i\phi_1}x_1^2 - r_2e^{-i\phi_2}x_2^2 - \frac{\bar{x}}{2}x_3^2 - 2r_3e^{-i\phi_3}x_1x_2 - 2\sqrt{c_1\bar{x}}e^{-i\theta}x_1x_3 - 2\sqrt{c_2\bar{x}}e^{-i(\theta+\theta_f)}x_2x_3\right],\end{aligned} \tag{3.50}$$

where $\theta_f = \arg f$ and

$$\begin{aligned}&\mathcal{N} = \frac{\sqrt{\bar{x}}\sqrt{4(a_1 - c_1)(a_2 - c_2) - (2b + f + f^*)^2}}{\pi^{3/2}} \qquad r_1^2 = (a_1 - c_1)^2 + 4a_1c_1\cos^2\theta \\ &r_2^2 = (a_2 - c_2)^2 + 4a_2c_2\cos^2(\theta + \theta_f) \qquad r_3^2 = c_1c_2 + b^2 - 2b\sqrt{c_1c_2}\cos(2\theta + \theta_f) \\ &\tan\phi_1 = \frac{c_1\sin 2\theta}{c_1\cos 2\theta + a_1} \quad \tan\phi_2 = \frac{c_2\sin(2\theta + 2\theta_f)}{c_2\cos(2\theta + 2\theta_f) + a_2} \quad \tan\phi_3 = \frac{\sqrt{c_1c_2}\sin(2\theta + \theta_f)}{\sqrt{c_1c_2}\cos(2\theta + \theta_f) - b}.\end{aligned} \tag{3.51}$$

If we choose

$$\bar{x} = \frac{1}{4(a_1 - c_1)(a_2 - c_2) - (2b + f + f^*)^2} \qquad \theta = 0, \tag{3.52}$$

the parameters in Eq. (3.51) reduce to

$$\begin{aligned} &\mathcal{N} = \frac{1}{\pi^{3/2}} \quad r_1 = a_1 + c_1 \quad r_2 = \sqrt{(a_2 - c_2)^2 + 4a_2 c_2 \cos^2\theta_f} \\ &r_3 = \sqrt{b^2 + c_1 c_2 - 2b\sqrt{c_1 c_2}\cos\theta_f} \qquad \phi_1 = 0 \\ &\phi_2 = \tan^{-1}\left[\frac{c_2 \sin 2\theta_f}{c_2 \cos 2\theta_f + a_2}\right] \quad \phi_3 = \tan^{-1}\left[\frac{\sqrt{c_1 c_2}\sin\theta_f}{\sqrt{c_2 c_2}\cos\theta_f - b}\right]. \end{aligned} \tag{3.53}$$

If $c_1 = c_2 \equiv c$ and $\theta_f = 0$, it is easy to show that the pure state in Eq. (3.50) exactly coincides with Eq. (3.24).

D. Type IV Bipartite State

In this subsection we examine the following bipartite Gaussian state:

$$\begin{aligned} \rho_{IV}[x_1', x_2', : x_1, x_2] = A \exp\Bigg[&- a_1(x_1^2 + x_2'^2) - a_2(x_1'^2 + x_2^2) \\ &+ 2b(x_1' x_2' + x_1 x_2) + 2c(x_1 x_1' + x_2 x_2') + 2f_1 x_1 x_2' + 2f_2 x_1' x_2 \Bigg] \end{aligned} \tag{3.54}$$

where $A = \sqrt{(a_1 + a_2 - 2c)^2 - (2b + f_1 + f_2)^2}/\pi$. The density matrix ρ_{IV} is obtained from ρ_{II} by taking a partial transposition ($x_1 \leftrightarrow x_1'$). In Ref. [35] it is shown that solving the eigenvalue equation (3.1) (with $j = \mathrm{IV}$) is highly difficult because we cannot find suitable change of variables to factorize Eq. (3.1) into the two single-party eigenvalue equations. In fact, we do not know whether the factorization is possible or not. Thus, we will compute the Rényi and von Neumann entropies of ρ_{IV} without solving the eigenvalue equation explicitly as we did in the previous section. Here, we assume that the eigenvalues of ρ_{IV} is also represented as Eq. (3.2).

First, we note

$$\tilde{\beta}_1 \equiv \mathrm{tr}\rho_{IV}^2 = \sqrt{\frac{Y_1}{Y_2}} \qquad \tilde{\beta}_2 \equiv \mathrm{tr}\rho_{IV}^3 = \frac{4Y_1}{Y_1 + 3Y_2 - 12y} \tag{3.55}$$

where

$$Y_1 = (a_1{+}a_2{-}2c)^2{-}(f_1{+}f_2{+}2b)^2 \quad Y_2 = (a_1{+}a_2{+}2c)^2{-}(f_1{+}f_2{-}2b)^2 \quad y = c^2{-}f_1 f_2. \tag{3.56}$$

Let $Y_2 \pm Y_1 = 2B_\pm$ and $\tilde{B} = Y_1 Y_2$, where $B_+ = (a_1 + a_2)^2 + 4c^2 - (f_1 + f_2)^2 - 4b^2$ and $B_- = 4c(a_1 + a_2) + 4b(f_1 + f_2)$. Then, if one follows the procedure of previous subsection, ξ_1 and ξ_2 of ρ_{IV} are given by

$$\xi_1 = \frac{\bar{u} + \sqrt{\bar{u}^2 - 4\bar{v}}}{2} \qquad \xi_2 = \frac{\bar{u} - \sqrt{\bar{u}^2 - 4\bar{v}}}{2} \tag{3.57}$$

with

$$\bar{u} = \frac{B_-}{4y}\left[1 - \sqrt{1 - \frac{8y}{B_+ + \sqrt{\tilde{B}}}}\right] \quad \bar{v} = -1 + \frac{B_+ + \sqrt{\tilde{B}}}{4y}\left[1 - \sqrt{1 - \frac{8y}{B_+ + \sqrt{\tilde{B}}}}\right]. \tag{3.58}$$

If $c^2 = f_1 f_2$, $\bar{u}$ and $\bar{v}$ become

$$\bar{u} = \frac{B_-}{B_+ + \sqrt{\tilde{B}}} \qquad \bar{v} = 0. \tag{3.59}$$

Thus, in this case, either ξ_1 or ξ_2 is zero depending on the sign of $\bar{u}$. In spite of this fact we cannot find any tripartite pure state σ_{ABC} whose substate coincides with ρ_{IV}. This seems to be due to the fact that ρ_{IV} cannot arise in the usual zero-temperature harmonic oscillator system like ρ_{II}.

IV. CONCLUSIONS

In this paper we derive the Rényi and von Neumann entropies for four types of the bipartite Gaussian states. For the case of type I and type II we solve the eigenvalue equation (3.1) (with $j = \mathrm{I}, \mathrm{II}$) analytically by factorizing it into two single-party eigenvalue equations. For the case of type III we use a different method. Since the derivation of Rényi and von Neumann entropies does not need the normalized eigenfunctions, we derive the eigenvalues of Eq. (3.1) (with $j = \mathrm{III}$) without deriving the eigenfunction explicitly. This method is valid because of the fact that the eigenvalue of ρ_{III} is represented as Eq. (3.2) as shown in Ref. [34]. For the case of type IV we derive the entropies under the assumption that the eigenvalue of Eq. (3.1) (with $j = \mathrm{IV}$) is also represented as Eq. (3.2).

Of course, there might be many bipartite Gaussian states, which do not belong to these four types. We do not know whether the Rényi and von Neumann entropies of all bipartite Gaussian states can be exactly derived or not. Furthermore, we do not know how to solve the eigenvalue equation (3.1) for non-Gaussian state. For example, let us consider the general

Gaussian or non-Gaussian bipartite state $\rho[x_1', x_2' : x_1, x_2]$. Then, its eigenvalues λ_{mn} satisfy

$$\det\Big[\rho[x_1', x_2' : x_1, x_2] - \lambda_{mn}\delta(x_1 - x_1')\delta(x_2 - x_2')\Big] = 0. \tag{4.1}$$

If the determinant of Eq. (4.1) can be computed, it is possible to derive the Rényi and von Neumann entropies for general state ρ without deriving the normalized eigenfunctions. Probably, the computation of the determinant may need to adopt some regularization scheme to escape the infinity arising due to δ-functions. We hope to explore this issue in the future.

---

[1] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge University Press, Cambridge, England, 2000).

[2] F. Arute et al.,Quantum supremacy using a programmable superconducting processor, Nature **574** (2019) 505. Its supplementary information is given in arXiv:1910.11333.

[3] E. Schrödinger, *Die gegenwärtige Situation in der Quantenmechanik*, Naturwissenschaften, **23** (1935) 807.

[4] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Quantum Entanglement*, Rev. Mod. Phys. **81** (2009) 865 [quant-ph/0702225] and references therein.

[5] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres and W. K. Wootters, *Teleporting an Unknown Quantum State via Dual Classical and Einstein-Podolsky-Rosen Channles*, Phys.Rev. Lett. **70** (1993) 1895.

[6] Y. H. Luo et al., *Quantum Teleportation in High Dimensions*, Phys. Rev. Lett. **123** (2019) 070505 [arXiv:1906.09697 (quant-ph)].

[7] C. H. Bennett and S. J. Wiesner, *Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states*, Phys. Rev. Lett. **69** (1992) 2881.

[8] V. Scarani, S. Lblisdir, N. Gisin and A. Acin, *Quantum cloning*, Rev. Mod. Phys. **77** (2005) 1225 [quant-ph/0511088] and references therein.

[9] A. K. Ekert , *Quantum Cryptography Based on Bells Theorem*, Phys. Rev. Lett. **67** (1991) 661.

[10] C. Kollmitzer and M. Pivk, Applied Quantum Cryptography (Springer, Heidelberg, Germany, 2010).

[11] K. Wang, X. Wang, X. Zhan, Z. Bian, J. Li, B. C. Sanders, and P. Xue, *Entanglement-enhanced*

quantum metrology in a noisy environment, Phys. Rev. **A97** (2018) 042112 [arXiv:1707.08790 (quant-ph)].

[12] T. D. Ladd, F. Jelezko, R. Laflamme, Y. Nakamura, C. Monroe, and J. L. O'Brien, *Quantum Computers*, Nature, **464** (2010) 45 [arXiv:1009.2267 (quant-ph)].

[13] G. Vidal, *Efficient classical simulation of slightly entangled quantum computations*, Phys. Rev. Lett. **91** (2003) 147902 [quant-ph/0301063].

[14] S. Ghernaouti-Helie, I. Tashi, T. Laenger, and C. Monyk, *SECOQC Business White Paper*, arXiv:0904.4073 (quant-ph).

[15] see https://www.technologyreview.com/s/609451/ibm-raises-the-bar-with-a-50-qubit-quantum-computer/.

[16] C. H. Bennett, D. P. DiVincenzo, J. A. Smokin and W. K. Wootters, *Mixed-state entanglement and quantum error correction*, Phys. Rev. **A 54** (1996) 3824 [quant-ph/9604024].

[17] S. Hill and W. K. Wootters, *Entanglement of a Pair of Quantum Bits*, Phys. Rev. Lett. **78** (1997) 5022 [quant-ph/9703041].

[18] W. K. Wootters, *Entanglement of Formation of an Arbitrary State of Two Qubits*, Phys. Rev. Lett. **80** (1998) 2245 [quant-ph/9709029].

[19] A. Peres, Phys. Rev. Lett. *Separability Criterion for Density Matrices*, **77** (1996) 1413 [quant-ph/9604005].

[20] M. Horodecki, P. Horodecki, and R. Horodecki, *Separability of mixed states: necessary and sufficient conditions*, Phys. Lett. **223** (1996) 1 [quant-ph/9605038].

[21] P. Horodecki, *Separability criterion and inseparable mixed states with positive partial transposition*, Phys. Lett. **A 232** (1997) 333 [quant-ph/9703004].

[22] L. M. Duan, G. Giedke, J. I. Cirac, and P. Zoller, *Inseparability criterion for continuous variable systems*, Phys. Rev. Lett. **84** (2000) 2722 [quant-ph/9908056].

[23] R. Simon, *Peres-Horodecki separability criterion for continuous variable systems*, Phys. Rev. Lett. **84** (2000) 2726 (quant-ph/9909044).

[24] G. Vida and R. F. Werner, A computable measure of entanglement, Phys. Rev. **A 65** (2002) 032314 [quant-ph/0102117].

[25] L. M. Duan, G. Giedke, J. I. Cirac, and P. Zoller, *Entanglement purification of Gaussian variable quantum states*, Phys. Rev. Lett. **84** (2000) 4002 (quant-ph/9912017).

[26] G. Giedke, L. M. Duan, J. I. Cirac, and P. Zoller, *All inseparable two-mode Gaussian contin-*

uous variable states are distillable, quant-ph/0007061.

[27] S. L. Braunstein and P. van Loock, *Quantum Information with continuous variables*, Rev. Mod. Phys. **77** (2005) 513 quant-ph/0410100) and references therein.

[28] G. Adesso, S. Ragy, and A. R. Lee, *Continuous variable quantum information: Gaussian states and beyond*, Open. Syst. Inf. Dyn., **21** (2014) 1440001 [arXiv:1401.4679 (quant-ph)].

[29] L. Bombelli, R. K. Koul, J. Lee, and R. D. Sorkin, *Quantum source of entropy for black holes*, Phys. Rev. **D 34** (1986) 373.

[30] M. Srednicki, *Entropy and Area*, Phys. Rev. Lett. **71** (1993) 666.

[31] D. N. Makarov, *Coupled harmonic oscillators and their quantum entanglement*, arXiv:1710.01158 (quant-ph).

[32] S. Ghosh, K. S. Gupta, S. C. L. Srivastava, *Entanglement dynamics following a sudden quench: an exact solution*, arXiv:1709.02202 [quant-ph].

[33] DaeKil Park, *Dynamics of entanglement and uncertainty relation in coupled harmonic oscillator system: exact results*, Quant. Inf. Proc. **17** (2018) 147 [arXiv:1801.07070 (quant-ph)].

[34] DaeKil Park, *Dynamics of Entanglement in Three Coupled Harmonic Oscillator System with Arbitrary Time-Dependent Frequency and Coupling Constants*, Quant. Inf. Proc. **18** (2019) 282 [arXiv:1811.06256 (quan-ph)].

[35] DaeKil Park, *Thermal Entanglement Phase Transition in Coupled Harmonic Oscillators with Arbitrary Time-Dependent Frequencies*, arXiv:1903.03297 [quant-ph].

[36] A. P. Prudnikov, Y. A. Brychkov, and O. I. Marichev, *Integrals and Series*, (Gordon and Breach Science Publishers, 1983, New York).

[37] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover Publications, 1972, New York).