

SUPERNOVA REMNANTS II
AN ODYSSEY IN SPACE AFTER STELLAR DEATH
3-8 JUNE 2019, CHANIA, CRETE, GREECE

Evolution of High-energy Particle Distribution in Supernova Remnants

Houdun Zeng¹, Yuliang Xin¹, Qi Fu² and Siming Liu¹

¹ Key Laboratory of Dark Matter and Space Astronomy, Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210008, China

² School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China

Abstract

The spectra fits to a sample of 34 supernova remnants (Zeng et al., 2019) are updated. γ -ray spectra of 20 supernova remnants (SNRs) with a soft TeV spectrum are further analyzed. We found that 17 of them can be fitted in the hadronic scenario with a single power-law ion distribution with an index of ~ 2.6 , which is significantly softer than the ion distribution inferred from γ -ray observations of star-burst galaxies. If Galactic cosmic rays are mostly produced by SNRs, this result suggests that SNRs in star-burst galaxies may never reach the phase with a soft γ -ray spectra or escape of high-energy particles from SNRs before they reach this phase with a soft γ -ray spectrum dominates the contribution of SNRs to Galactic cosmic rays.

1 Introduction

As early as 1930s, supernova remnants (SNRs) have been suggested as the dominant contributors to the galactic cosmic rays (Baade & Zwicky, 1934). However, it is still not clear how the galactic cosmic ray spectrum is related to energetic particle distribution in individual SNRs.

2 Sample and Model

The sample and model are introduced in Zeng et al. (2019). Note that we here update the gamma-ray data and fitted results of SNR W30 (Liu et al., 2019) and G205.5+0.5, and remove SNR G150.3+4.5 whose radio emission may be produced by secondary leptons in the hadronic scenario for the γ -ray emission.

3 The Results

Figure. 1 shows the main results of correlations of model parameters, and the evolution of high-energy particle distribution in SNRs. There are the detailed introductions for those in Zeng et al. (2019).




Figure 1: Upper: the electron synchrotron cooling time at the cutoff energy and the break energy of particle distribution decreases with the age of the corresponding SNR. Lower: correlation between low-energy spectral indices and the ages of SNRs, and the scenario of SNR origin of Galactic cosmic rays.

Figure. 2 shows the results of fitted SED of every hadronic original SNR with a single power-law distribution, as well as the fitted index. The fitted index Γ and E_{min} are compared with the high-end index ($\alpha + 1$) and break energy (E_{br}) of particle distribution in Zeng et al. (2019).

Figure. 3 and 4 show the results of fitted SED of SNRs and/or Starburst galaxies with different energy bands. Specially, the left panel in Figure. 3 indicates that the reduced χ^2 and index vary with the fitting energy band, implying that high-energy particles may have been escaped from SNR efficiently so that only particles with relatively low-energies are still trapped.

Acknowledgments

Support for this work was provided by National Key R&D Program of China: 2018YFA0404203, NSFC grants: U1738122, 11761131007 and by the International Partnership Program of Chinese Academy of Sciences, grant No. 114332KYSB20170008

References

- Baade, W., & Zwicky, F. 1934, Proceedings of the National Academy of Science, 20, 259
- Liu, B., Yang, R.-z., Sun, X.-n., Aharonian, F., & Chen, Y. 2019, arXiv e-prints, arXiv:1907.00180
- Zeng, H., Xin, Y., & Liu, S. 2019, apj, 874, 50




Figure 2: Red point: the results of fitted SED of hadronic original SNRs with a single power-law distribution. Here α is the index and E_{\min} is the minimum energy of fitted SED. Black point: the values from Zeng et al. (2019).




Figure 3: Left: the result of fitted SEDs with $E > 100$ GeV using the π^0 decay for 9 SNRs. Middle: the result of fitted SEDs with $E < 1$ GeV using the π^0 decay for 16 SNRs. Right: the result of fitted SEDs with the whole spectrum using the π^0 decay for 17 SNRs.




Figure 4: Left: the reduced χ^2 and index vary with the fitting energy bands. The red, green and blue points represent the best-fit values of high- and low-end SEDs for SNRs and high-end SEDs for Starburst galaxies. Right: the result of fitted SEDs with $E > 0.1$ GeV using the π^0 decay for 7 Starburst galaxies.