

Constructions of mutually unbiased entangled bases

Fei Shi,^{1,*} Xiande Zhang,^{2,†} and Lin Chen^{3,4,‡}

¹*School of Cyber Security, University of Science and Technology of China, Hefei, 230026, People's Republic of China.*

²*School of Mathematical Sciences, University of Science and Technology of China, Hefei, 230026, People's Republic of China.*

³*School of Mathematical Sciences, Beihang University, Beijing 100191, China*

⁴*International Research Institute for Multidisciplinary Science, Beihang University, Beijing 100191, China*

We construct two mutually unbiased bases by maximally entangled states (MUMEBs) in $\mathbb{C}^2 \otimes \mathbb{C}^3$. This is the first example of MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ when $d \nmid d'$, namely d' is not divisible by d . We show that they cannot be extended to four MUBs in \mathbb{C}^6 . We propose a recursive construction of mutually unbiased bases formed by special entangled states with a fixed Schmidt number k (MUSEBks). It shows that $\min\{t_1, t_2\}$ MUSEB $k_1 k_2$ s in $\mathbb{C}^{pd} \otimes \mathbb{C}^{qd'}$ can be constructed from t_1 MUSEB k_1 s in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ and t_2 MUSEB k_2 s in $\mathbb{C}^p \otimes \mathbb{C}^q$ for any d, d', p, q . Further, we show that three MUMEBs exist in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ for any d, d' with $d \mid d'$, and two MUMEBs exist in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ for infinitely many d, d' with $d \nmid d'$.

Keywords: mutually unbiased bases, maximally entangled states, special entangled states

I. INTRODUCTION

The notion of mutually unbiased bases (MUBs) became an essential feature of quantum mechanics in 1960 in the works of Schwinger [1]. In his work, Schwinger realized that no information can be retrieved when a quantum state which is prepared in a basis state is measured with respect to the basis mutually unbiased with the prepared one. This observation has a striking application in the well-known BB84 quantum key distribution (QKD) protocol [2].

Estimating an unknown physical transformation of a state, which is described by a unitary operator, was studied in [3]. It is a more challenging problem compared to state estimation because it requires optimal measurements of states both post and prior to transformation. Such estimation has an immediate application in the alignment of reference frames using quantum spins [4], and in the field of quantum cryptography but with a lower security threshold [5, 6]. To improve the security, a notion of “mutually unbiased unitary-operator bases” (MUUBs) was put forward in [7] for unitary operators acting on a d -dimensional Hilbert space as a set of pairwise *mutually unbiased* unitary operator bases. The concept MUUBs was shown to be equivalent to MUBs formed by maximally entangled states (MUMEBs) [8]. In the same paper, Shaari et al. [8] generalized the original MUUBs for unitary operators acting on *subspaces* of a Hilbert space, and considered the distinguishability of unitaries selected from a set of MUUBs and its use in a QKD setup.

In this paper, we consider another generalization of MUUBs. Motivated by the equivalence between MUUBs in \mathbb{C}^d and MUBs formed by maximally entangled states (MUMEBs) in $\mathbb{C}^d \otimes \mathbb{C}^d$, we study mutually unbiased bases formed by special entangled states with a fixed Schmidt number k (MUSEBks) in a general bipartite state space $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ ($1 \leq k \leq d, d'$). Let $M_k(d, d')$ denote the maximum size of MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$, i.e., the maximum number of mutually unbiased bases formed by special entangled states with Schmidt number k . The central problem here is to determine $M_k(d, d')$ in a given bipartite state space and provide bounds for this value.

There have been results in [7–15] in which MUMEBs were known only when $d \mid d'$, namely d' is divisible by d . We give the first example of MUMEBs when $d \nmid d'$, and propose a simple construction of MUSEBks for general systems via matrix spaces. Consequently, we improve all previous results on MUSEBks (MUMEBs). See Table I for a summary and comparison with known results.

The rest of this paper is organized as follows. In Sec. II, we introduce some related notations and terminologies, and give a one-to-one correspondence between MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ and MUSEBks in $\mathcal{M}_{d \times d'}$. Sec. III provides a construction of two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$, which is the first MUMEBs when $d \nmid d'$. In Sec. IV, we show that the two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$ constructed in Sec. III cannot be extended to 4 MUBs in \mathbb{C}^6 . In Sec. V, we establish a construction of $\min\{t_1, t_2\}$ MUSEB $k_1 k_2$ s in $\mathbb{C}^{pd} \otimes \mathbb{C}^{qd'}$ from t_1 MUSEB k_1 s in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ and t_2 MUSEB k_2 s in $\mathbb{C}^p \otimes \mathbb{C}^q$, which yields several new lower bounds in Sec. VI. We give the main conclusion and some open problems in Sec. VII.

*Electronic address: shifei@mail.ustc.edu.cn

†Electronic address: drzhangx@ustc.edu.cn

‡Electronic address: linchen@buaa.edu.cn

TABLE I: Results about MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$

(d, q have a factorization of prime power, $p_1^{a_1} \leq \dots \leq p_s^{a_s}, (p'_1)^{a'_1} \leq \dots \leq (p'_s)^{a'_s}$)		
Conditions	Bounds	References
$d \geq 2$	$M_d(d, d) \leq d^2 - 1$	[7]
$d = 2, 3, 5, 7, 11$	$M_d(d, d) = d^2 - 1$	[7]
$d = p_1^{a_1} \dots p_s^{a_s}, p_1^{a_1} \geq 3$	$M_d(d, d) \geq p_1^{a_1} - 1$	[11]
$d \geq 3$, prime power	$M_d(d, d) \geq 2(d - 1)$	[12]
$d = p_1^{a_1} \dots p_s^{a_s}$, odd	$M_d(d, d) \geq 2(p_1^{a_1} - 1)$	[14]
d , odd prime power	$M_d(d, d) \geq (d^2 - 1)/2$	[13]
$d = 2^t p_1^{a_1} \dots p_s^{a_s}, t = 0$	$M_d(d, d) \geq (p_1^{2a_1} - 1)/2$	This paper
$d = 2^t p_1^{a_1} \dots p_s^{a_s}, t = 1$	$M_d(d, d) \geq 3$	This paper
$d = 2^t p_1^{a_1} \dots p_s^{a_s}, t \geq 2$	$M_d(d, d) \geq \min\{2(2^t - 1), (p_1^{2a_1} - 1)/2\}$	This paper
$M_d(d, d') \geq 2, \ell \geq 1$	$M_d(d, 2^\ell d') \geq 2$	[10]
q and d , both prime powers	$M_d(d, qd) \geq \min\{q, M_d(d, d)\}$	[12]
$q = (p'_1)^{a'_1} \dots (p'_s)^{a'_s}, d$ odd	$M_d(d, qd) \geq \min\{(p'_1)^{a'_1} + 1, M_d(d, d)\}$	[14]
$d, q \geq 2$	$M_d(d, qd) \geq \min\{N(q), M_d(d, d)\} \geq 3$	This paper
$d, q \geq 1$	$M_{2d}(2d, 3qd) \geq 2$.	This paper
$k = 2^\ell$	$M_2(3, 4k) \geq 2$	[15]
$k \geq 1$	$M_2(3, 2k) \geq 2$	This paper
$d, p \geq 2, q \geq 1$	$M_d(pd, qd) \geq \min\{N(p), M_d(d, qd)\} \geq 3$	This paper

II. DEFINITION AND PRELIMINARY

Assume that $1 \leq k \leq \min\{d, d'\}$. A (pure) state $|\psi\rangle \in \mathbb{C}^d \otimes \mathbb{C}^{d'}$ is called a *special entangled state with Schmidt number k* if it has Schmidt decomposition:

$$\psi = \frac{1}{\sqrt{k}} \sum_{i=0}^{k-1} |i\rangle |i'\rangle,$$

where $\{|i\rangle\}$ and $\{|i'\rangle\}$ are orthonormal sets of \mathbb{C}^d and $\mathbb{C}^{d'}$, respectively [16]. In particular, a special entangled state with Schmidt number k is a product state when $k = 1$, and a maximally entangled state when $k = \min\{d, d'\}$. A basis \mathcal{B} is called a special entangled basis with Schmidt number k (SEBk) in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ if it contains dd' pairwise orthogonal special entangled states with Schmidt number k [16, 17]. In particular, \mathcal{B} is a maximally entangled state basis (MEB) when $k = \min\{d, d'\}$.

Two orthonormal bases $\mathcal{B}_1 = \{|\phi_i\rangle\}_{i=1}^{dd'}$ and $\mathcal{B}_2 = \{|\psi_i\rangle\}_{i=1}^{dd'}$ of $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ are said to be *mutually unbiased* if

$$|\langle \phi_i | \psi_j \rangle| = \frac{1}{\sqrt{dd'}}, \quad (1 \leq i, j \leq dd').$$

A set of SEBks $\mathcal{A} = \{\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_m\}$ in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ is called mutually unbiased (MUSEBks) if every pair \mathcal{B}_i and \mathcal{B}_j are mutually unbiased. MUSEBks are MUUBs when $k = d = d'$, and MUMEBs when $k = \min\{d, d'\}$. When $k = d' = 1$, MUSEBks are just MUBs [18], for which the maximum size is denoted by $N(d)$ conventionally. It is shown that $N(d) \leq d + 1$ and $N(d) = d + 1$ when d is a prime power [18]. An open problem is to determine $N(d)$ when d is not a prime power, which has an elementary lower bound ($d = p_1^{a_1} \dots p_s^{a_s}, p_1^{a_1} \leq \dots \leq p_s^{a_s}$):

$$N(d) \geq \min\{N(p_1^{a_1}), \dots, N(p_s^{a_s})\} = p_1^{a_1} + 1. \quad (1)$$

Actually, this bound can be improved in some special cases [19]. It is shown that a complete set of MUBs of a bipartite system contains a fixed amount of entanglement [20].

Now we define similar concepts in matrix spaces. Let $\mathcal{M}_{d \times d'}$ be the Hilbert space of all $d \times d'$ complex matrices equipped with inner product

$$(A, B) = \text{Tr}(A^\dagger B).$$

There is a one-to-one relation between $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ and $\mathcal{M}_{d \times d'}$ [17]:

$$|\psi_i\rangle = \sum_{k,l} a_{k,l}^{(i)} |k\rangle |l'\rangle \in \mathbb{C}^d \otimes \mathbb{C}^{d'} \iff A_i = [a_{k,l}^{(i)}] \in \mathcal{M}_{d \times d'},$$

$$Sr(|\psi_i\rangle) = \text{rank}(A_i), \quad \langle \psi_i | \psi_j \rangle = \text{Tr}(A_i^\dagger A_j),$$

where $\{k\}$ and $\{l'\}$ are the computational bases of \mathbb{C}^d and $\mathbb{C}^{d'}$, respectively, and $Sr(|\psi_i\rangle)$ denotes the Schmidt number of $|\psi_i\rangle$.

Let $\{\frac{1}{\sqrt{k}}, \frac{1}{\sqrt{k}}, \dots, \frac{1}{\sqrt{k}}\}_k$ denote k values of $\frac{1}{\sqrt{k}}$. A $d \times d'$ matrix is called a k -singular-value- $\frac{1}{\sqrt{k}}$ matrix if its nonzero singular values are $\{\frac{1}{\sqrt{k}}, \frac{1}{\sqrt{k}}, \dots, \frac{1}{\sqrt{k}}\}_k$. Then $|\psi_i\rangle$ is a special entangled state with Schmidt number k if and only if A_i is a k -singular-value- $\frac{1}{\sqrt{k}}$ matrix. Especially, when $k = d = d'$, $|\psi_i\rangle$ is a maximally entangled state if and only if $\sqrt{d}A_i$ is a unitary matrix. We give the definitions of SEBks and MUSEBks in $\mathcal{M}_{d \times d'}$.

Definition 1. A set of k -singular-value- $\frac{1}{\sqrt{k}}$ matrices $\{A_i\}_{i=1}^{dd'}$ of $\mathcal{M}_{d \times d'}$ is called an SEBk if $\text{Tr}(A_i^\dagger A_j) = \delta_{ij}$.

Definition 2. Two SEBks $\mathcal{B}_1 = \{A_i\}_{i=1}^{dd'}$ and $\mathcal{B}_2 = \{B_i\}_{i=1}^{dd'}$ in $\mathcal{M}_{d \times d'}$ are said to be mutually unbiased if

$$|\text{Tr}(A_i^\dagger B_j)| = \frac{1}{\sqrt{dd'}}, \quad (1 \leq i, j \leq dd').$$

A set of SEBks $\mathcal{A} = \{\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_m\}$ in $\mathcal{M}_{d \times d'}$ is called MUSEBks if every pair \mathcal{B}_i and \mathcal{B}_j are mutually unbiased.

Due to the one-to-one relation, an SEBk of $\mathcal{M}_{d \times d'}$ is an SEBk of $\mathbb{C}^d \otimes \mathbb{C}^{d'}$; MUSEBks in $\mathcal{M}_{d \times d'}$ are MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$. When $k = d = d'$, Definition 2 gives MUMEBs in $\mathcal{M}_{d \times d}$, which corresponds to MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^d$. In this case, multiplying each matrix by a scalar \sqrt{d} gives a set of mutually unbiased unitary bases (MUUBs) in \mathbb{C}^d defined in [7]. So Definition 2 can be viewed as a generation of MUUBs in [7], from which we know $M_d(d, d) \leq d^2 - 1$ and $M_d(d, d) = d^2 - 1$ when $d = 2, 3, 5, 7, 11$.

Remark 1. In [8], the authors gave another definition of MUUBs: Consider two distinct orthogonal bases, \mathcal{A}_0 and \mathcal{A}_1 , composed of unitary transformations for some subspace of the vector space \mathbb{C}^d . \mathcal{A}_0 and \mathcal{A}_1 are MUUBs provided that

$$|\text{Tr}(A_0^{(i)\dagger} A_1^{(j)})|^2 = c, \quad \forall A_0^{(i)} \in \mathcal{A}_0, A_1^{(j)} \in \mathcal{A}_1,$$

for $i, j = 1, \dots, n$ and some constant $c \neq 0$.

When \mathcal{A}_0 and \mathcal{A}_1 are composed of unitary transformations for the whole vector space \mathbb{C}^d , and $c = 1$, then this definition reduces to Definition 2 for the case $k = d = d'$ by multiplying each matrix by a scalar $1/\sqrt{d}$.

III. THE FIRST MUMEBs WITH $d \nmid d'$

In previous works of [9–14], all constructions about MUMEBs are for $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ with $d \mid d'$. Are there MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ when $d \nmid d'$? We next give an example in this case.

Theorem 1. There are two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$. Namely, $M_2(2, 3) \geq 2$.

Proof. Let

$$|\phi_{m,n}\rangle = \frac{1}{\sqrt{2}} \sum_{p=0}^1 (-1)^{np} |p\rangle |(p \oplus m)'\rangle,$$

where $n = 0, 1$, $m = 0, 1, 2$, and $p \oplus m := p + m \pmod{3}$. It is easy to check that $\{|\phi_{m,n}\rangle\}$ is an MEB in $\mathbb{C}^2 \otimes \mathbb{C}^3$. Let A be a 2×2 unitary matrix

$$A = \begin{pmatrix} A_{00} & A_{01} \\ A_{10} & A_{11} \end{pmatrix},$$

and let

$$\begin{pmatrix} |a_0\rangle \\ |a_1\rangle \end{pmatrix} = \begin{pmatrix} A_{00} & A_{01} \\ A_{10} & A_{11} \end{pmatrix} \begin{pmatrix} |0\rangle \\ |1\rangle \end{pmatrix}.$$

Denote

$$|\psi_{m,n}\rangle = \frac{1}{\sqrt{2}} \sum_{p=0}^1 (-1)^{np} |a_p\rangle |(p \oplus m)'\rangle,$$

where $n = 0, 1$ and $m = 0, 1, 2$. Then $\{|\psi_{m,n}\rangle\}$ is also an MEB in $\mathbb{C}^2 \otimes \mathbb{C}^3$. By definition, $\{|\phi_{m,n}\rangle\}$ and $\{|\psi_{m,n}\rangle\}$ are two MUMEBs if and only if

$$\left| \sum_{p_2=0}^1 \sum_{p_1=0}^1 (-1)^\lambda \langle (p_1 \oplus m_1)' | (p_2 \oplus m_2)' \rangle A_{p_2, p_1} \right| = \frac{2}{\sqrt{6}},$$

for all $\lambda = 0, 1$ and $m_1, m_2 = 0, 1, 2$. When $m_1 = m_2$, we have $|A_{0,0} \pm A_{1,1}| = \frac{2}{\sqrt{6}}$, which implies $A_{0,0} \perp A_{1,1}$ and $|A_{0,0}| = |A_{1,1}| = \frac{1}{\sqrt{3}}$; when $m_2 \equiv m_1 + 1 \pmod{3}$, we have $|A_{0,1}| = \frac{2}{\sqrt{6}}$; when $m_1 \equiv m_2 + 1 \pmod{3}$, we have $|A_{1,0}| = \frac{2}{\sqrt{6}}$. So we can assume that

$$A = \frac{1}{\sqrt{3}} \begin{pmatrix} e^{i\theta_1} & \sqrt{2}e^{i\theta_2} \\ \sqrt{2}e^{i\theta_3} & e^{i(\theta_1 + \frac{\pi}{2})} \end{pmatrix}, \quad (2)$$

for some θ_i , $i = 1, 2, 3$. Since A is a unitary matrix, θ_i must satisfy the following condition:

$$\theta_2 + \theta_3 - 2\theta_1 = 2k\pi + \frac{3\pi}{2}, \text{ for some } k \in Z. \quad (3)$$

Taking

$$\theta_1 = 0, \theta_2 = \frac{3\pi}{2}, \theta_3 = 0,$$

we have

$$A = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & -\sqrt{2}i \\ \sqrt{2} & i \end{pmatrix} \text{ and}$$

$$|a_0\rangle = \frac{1}{\sqrt{3}}(|0\rangle - \sqrt{2}i|1\rangle), \quad |a_1\rangle = \frac{1}{\sqrt{3}}(\sqrt{2}|0\rangle + i|1\rangle).$$

Then $\{|\phi_{m,n}\rangle\}$ and $\{|\psi_{m,n}\rangle\}$ are as follows:

$$\begin{cases} |\phi_{0,0}\rangle = \frac{1}{\sqrt{2}}(|00'\rangle + |11'\rangle) \\ |\phi_{0,1}\rangle = \frac{1}{\sqrt{2}}(|00'\rangle - |11'\rangle) \\ |\phi_{1,0}\rangle = \frac{1}{\sqrt{2}}(|01'\rangle + |12'\rangle) \\ |\phi_{1,1}\rangle = \frac{1}{\sqrt{2}}(|01'\rangle - |12'\rangle) \\ |\phi_{2,0}\rangle = \frac{1}{\sqrt{2}}(|02'\rangle + |10'\rangle) \\ |\phi_{2,1}\rangle = \frac{1}{\sqrt{2}}(|02'\rangle - |10'\rangle) \end{cases}, \quad (4)$$

$$\left\{ \begin{array}{l} |\psi_{0,0}\rangle = \frac{1}{\sqrt{6}}(|00'\rangle + \sqrt{2}|01'\rangle - \sqrt{2}i|10'\rangle + i|11'\rangle) \\ |\psi_{0,1}\rangle = \frac{1}{\sqrt{6}}(|00'\rangle - \sqrt{2}|01'\rangle - \sqrt{2}i|10'\rangle - i|11'\rangle) \\ |\psi_{1,0}\rangle = \frac{1}{\sqrt{6}}(|01'\rangle + \sqrt{2}|02'\rangle - \sqrt{2}i|11'\rangle + i|12'\rangle) \\ |\psi_{1,1}\rangle = \frac{1}{\sqrt{6}}(|01'\rangle - \sqrt{2}|02'\rangle - \sqrt{2}i|11'\rangle - i|12'\rangle) \\ |\psi_{2,0}\rangle = \frac{1}{\sqrt{6}}(|02'\rangle + \sqrt{2}|00'\rangle - \sqrt{2}i|12'\rangle + i|10'\rangle) \\ |\psi_{2,1}\rangle = \frac{1}{\sqrt{6}}(|02'\rangle - \sqrt{2}|00'\rangle - \sqrt{2}i|12'\rangle - i|10'\rangle) \end{array} \right. \quad (5)$$

Thus the above $\{|\phi_{m,n}\rangle\}$ and $\{|\psi_{m,n}\rangle\}$ are two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$. □

Note that we can also take $\theta_1 = \frac{\pi}{4}$, $\theta_2 = \frac{3\pi}{4}$, $\theta_3 = \frac{5\pi}{4}$ in Eq. (2), then

$$A = \frac{1}{\sqrt{3}} \begin{pmatrix} \frac{\sqrt{2}}{2}(1+i) & -1+i \\ -1-i & \frac{\sqrt{2}}{2}(-1+i) \end{pmatrix} \text{ and}$$

$$|a_0\rangle = \frac{1}{\sqrt{3}} \left(\frac{\sqrt{2}}{2}(1+i)|0\rangle + (-1+i)|1\rangle \right), |a_1\rangle = \frac{1}{\sqrt{3}} \left((-1-i)|0\rangle + \frac{\sqrt{2}}{2}(-1+i)|1\rangle \right). \quad (6)$$

The above $\{|\phi_{m,n}\rangle\}$ and $\{|\psi_{m,n}\rangle\}$ are also two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$. Unfortunately, if A_1 and A_2 are two matrices that satisfy the conditions of Eq. (2) and Eq. (3), then $A_1 A_2$ cannot satisfy these conditions. It means we cannot construct three MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$ by this way.

We should mention that our method of constructing two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$ in Theorem 1 is similar to that of constructing MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{kd}$ from [9], where a unitary matrix A of size $kd \times kd$ is applied on the latter subspace \mathbb{C}^{kd} . However, using their method, we find that such a 3×3 unitary matrix A does not exist. So we cannot construct MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$ by the method in [9].

Before closing this section, we convert $\{|\phi_{m,n}\rangle\}$ and $\{|\psi_{m,n}\rangle\}$ of Eq. (4) and Eq. (5) to MUMEBs \mathcal{R}_1 and \mathcal{R}_2 in $\mathcal{M}_{2 \times 3}$, respectively.

$$\begin{aligned} \mathcal{R}_1 = & \left\{ \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \right. \\ & \left. \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix} \right\}, \\ \mathcal{R}_2 = & \left\{ \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & \sqrt{2} & 0 \\ -\sqrt{2}i & i & 0 \end{pmatrix}, \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & -\sqrt{2} & 0 \\ -\sqrt{2}i & -i & 0 \end{pmatrix}, \right. \\ & \frac{1}{\sqrt{6}} \begin{pmatrix} 0 & 1 & \sqrt{2} \\ 0 & -\sqrt{2}i & i \end{pmatrix}, \frac{1}{\sqrt{6}} \begin{pmatrix} 0 & 1 & -\sqrt{2} \\ 0 & -\sqrt{2}i & -i \end{pmatrix}, \\ & \left. \frac{1}{\sqrt{6}} \begin{pmatrix} \sqrt{2} & 0 & 1 \\ i & 0 & -\sqrt{2}i \end{pmatrix}, \frac{1}{\sqrt{6}} \begin{pmatrix} -\sqrt{2} & 0 & 1 \\ -i & 0 & -\sqrt{2}i \end{pmatrix} \right\}. \end{aligned}$$

IV. APPLICATION

In this section, we propose an application of the two MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$ constructed in Eqs. (4) and (5) in Theorem 3. This is to tackle the special case of a long-standing open problem, namely whether four six-dimensional MUBs exist. The study of MUBs has extensive physical applications in quantum cryptography, tomography, and the construction of Wigner functions. It has been shown that there exist three six-dimensional MUBs, and widely believed

that four six-dimensional MUBs may not exist in spite of great efforts from mathematical and physical community [20–38].

To explain the application, we refer to the vectors in an orthonormal basis in \mathbb{C}^n as the column vectors of a unitary matrix. In particular we refer to the computational basis as the identity matrix. Hence, the existence of four six-dimensional MUBs is equivalent to the existence of the identity matrix, and three complex Hadamard matrices (CHMs) which are MU. If they exist then we refer to the three CHMs as the so-called MUB trio [36]. From Lemma 11 (ii) and matrix Y_6 of [36], we have found the following criterion of determining when a CHM belongs to an MUB trio.

Lemma 2. (i) *A CHM belongs to an MUB trio if and only if so does its transpose.*

(ii) *If a CHM has a 2×3 real submatrix then it does not belong to any MUB trio.*

Now we are in a position to present the application of the two MUMEBs in Eqs. (4) and (5).

Theorem 3. *The two MUMEBs in Eqs. (4) and (5) cannot be extended to four MUBs.*

Proof. In the following, we express the two MUMEBs as two unitary matrices U and V , respectively.

$$U = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 \end{pmatrix},$$

and

$$V = \begin{pmatrix} \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & 0 & 0 & \frac{\sqrt{2}}{\sqrt{6}} & -\frac{\sqrt{2}}{\sqrt{6}} \\ \frac{\sqrt{2}}{\sqrt{6}} & -\frac{\sqrt{2}}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & 0 & 0 \\ 0 & 0 & \frac{\sqrt{2}}{\sqrt{6}} & -\frac{\sqrt{2}}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ -\frac{\sqrt{2}i}{\sqrt{6}} & -\frac{\sqrt{2}i}{\sqrt{6}} & 0 & 0 & \frac{i}{\sqrt{6}} & -\frac{i}{\sqrt{6}} \\ \frac{i}{\sqrt{6}} & -\frac{i}{\sqrt{6}} & -\frac{\sqrt{2}i}{\sqrt{6}} & -\frac{\sqrt{2}i}{\sqrt{6}} & 0 & 0 \\ 0 & 0 & \frac{i}{\sqrt{6}} & -\frac{i}{\sqrt{6}} & -\frac{\sqrt{2}i}{\sqrt{6}} & -\frac{\sqrt{2}i}{\sqrt{6}} \end{pmatrix}.$$

One can verify that the lower left 2×3 submatrix of $U^\dagger V Q$ is $\begin{pmatrix} -\frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} \end{pmatrix}$, where the diagonal unitary $Q = \text{diag}(-i, -i, 1, 1, 1, 1)$. Hence $U^\dagger V Q$ does not belong to any MUB trio in terms of Lemma 2. We have proven the assertion. \square

Our results restrict the form of MUB trio, if it really exists. Following the proof of Theorem 3, one can similarly show that the other pair of MUMEBs constructed at Eq. (6) cannot be extended to four MUBs too. We may conjecture that, any two MUMEBs cannot be extended to four MUBs.

V. A RECURSIVE CONSTRUCTION OF MUSEBKs

In this section, we give a construction of MUSEBK₁k₂s in $\mathbb{C}^{pd} \otimes \mathbb{C}^{qd'}$ from MUSEBK₁s in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ and MUSEBK₂s in $\mathbb{C}^p \otimes \mathbb{C}^q$ for any positive integers d, d', p and q .

For any two sets $\mathcal{S} = \{A_1, A_2, \dots, A_n\}$ and $\mathcal{T} = \{B_1, B_2, \dots, B_m\}$, define $\mathcal{S}^T := \{A_1^T, A_2^T, \dots, A_n^T\}$, and

$$\mathcal{S} \otimes \mathcal{T} := \{A_i \otimes B_j \mid i = 1, \dots, n, j = 1, \dots, m\}.$$

Theorem 4. *If there are t_1 MUSEBK₁s in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ and t_2 MUSEBK₂s in $\mathbb{C}^p \otimes \mathbb{C}^q$, then there are $\min\{t_1, t_2\}$ MUSEBK₁k₂s in $\mathbb{C}^{pd} \otimes \mathbb{C}^{qd'}$. Namely,*

$$M_{k_1 k_2}(pd, qd') \geq \min\{M_{k_1}(d, d'), M_{k_2}(p, q)\}. \quad (7)$$

Proof. We prove it by using the language of MUSEBks in $\mathcal{M}_{d \times d'}$. Let $\mathcal{S}_1 = \{A_1, A_2, \dots, A_{dd'}\}$ and $\mathcal{S}_2 = \{B_1, B_2, \dots, B_{dd'}\}$ be two MUSEB k_1 s in $\mathcal{M}_{d \times d'}$, that is, A_i, B_j are k_1 -singular-value- $\frac{1}{\sqrt{k_1}}$ matrices, $\text{Tr}(A_i^\dagger A_j) = \delta_{i,j}$, $\text{Tr}(B_i^\dagger B_j) = \delta_{i,j}$ and $|\text{Tr}(A_i^\dagger B_j)| = \frac{1}{\sqrt{dd'}}$.

Similarly, let $\mathcal{T}_1 = \{C_1, C_2, \dots, C_{pq}\}$ and $\mathcal{T}_2 = \{D_1, D_2, \dots, D_{pq}\}$ be two MUSEB k_2 s in $\mathcal{M}_{p \times q}$, then C_i, D_j each have k_2 singular values $\frac{1}{\sqrt{k_2}}$, $\text{Tr}(C_i^\dagger C_j) = \delta_{i,j}$, $\text{Tr}(D_i^\dagger D_j) = \delta_{i,j}$ and $|\text{Tr}(C_i^\dagger D_j)| = \frac{1}{\sqrt{pq}}$.

Then $\mathcal{S}_1 \otimes \mathcal{T}_1$ and $\mathcal{S}_2 \otimes \mathcal{T}_2$ are two sets of $dd'pq$ matrices in $\mathcal{M}_{pd \times qd'}$. We prove that they are two MUSEB $k_1 k_2$ s in $\mathcal{M}_{pd \times qd'}$.

First, we show that $\mathcal{S}_1 \otimes \mathcal{T}_1$ and $\mathcal{S}_2 \otimes \mathcal{T}_2$ are two SEB $k_1 k_2$ s. We only prove it for $\mathcal{S}_1 \otimes \mathcal{T}_1$, the other one is similar. Since each $A_i \otimes C_j \in \mathcal{S}_1 \otimes \mathcal{T}_1$ has nonzero singular values $\{\frac{1}{\sqrt{k_1 k_2}}, \frac{1}{\sqrt{k_1 k_2}}, \dots, \frac{1}{\sqrt{k_1 k_2}}\}_{k_1 k_2}$, it is a $k_1 k_2$ -singular-value- $\frac{1}{\sqrt{k_1 k_2}}$ matrix. Next, $\text{Tr}[(A_i \otimes C_j)^\dagger (A_{i'} \otimes C_{j'})] = \text{Tr}(A_i^\dagger A_{i'}) \cdot \text{Tr}(C_j^\dagger C_{j'}) = \delta_{i,i'} \delta_{j,j'}$, so they are orthogonal if $(i, j) \neq (i', j')$.

Finally, the mutually unbiased property follows from the fact that $|\text{Tr}[(A_i \otimes C_j)^\dagger (B_{i'} \otimes D_{j'})]| = |\text{Tr}(A_i^\dagger B_{i'})| \cdot |\text{Tr}(C_j^\dagger D_{j'})| = \frac{1}{\sqrt{pdqd'}}$.

Let $t = \min\{t_1, t_2\}$. Assume that $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_t$ are t MUSEB k_1 s in $\mathcal{M}_{d \times d'}$, and $\mathcal{T}_1, \mathcal{T}_2, \dots, \mathcal{T}_t$ are t MUSEB k_2 s in $\mathcal{M}_{p \times q}$. It is easy to see that $\{\mathcal{S}_1 \otimes \mathcal{T}_1, \mathcal{S}_2 \otimes \mathcal{T}_2, \dots, \mathcal{S}_t \otimes \mathcal{T}_t\}$ is a set of t MUSEB $k_1 k_2$ s in $\mathcal{M}_{pd \times qd'}$ from the above analysis. \square

Theorem 4 gives a useful generic construction of MUSEBks (MUMEBs). We illustrate its importance in several cases.

When $k_1 = d \leq d'$ and $k_2 = p \leq q$, we obtain the case for MUMEBs,

$$M_{pd}(pd, qd') \geq \min\{M_d(d, d'), M_p(p, q)\}. \quad (8)$$

Example 1. There are five MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^4$ and three MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^6$ [9], then we can construct at least three MUMEBs in $\mathbb{C}^4 \otimes \mathbb{C}^{24}$, that is, $M_4(4, 24) \geq \min\{M_2(2, 4), M_2(2, 6)\} \geq 3$.

Observing that Theorem 4 does not require $d \leq d'$ and $p \leq q$, and the fact that $M_k(d, d') = M_k(d', d)$, we can get the following different example.

Example 2. From five MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^4$ and three MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^6$, we can also construct three MUSEB4s in $\mathbb{C}^8 \otimes \mathbb{C}^{12}$, that is, $M_4(8, 12) \geq \min\{M_2(4, 2), M_2(2, 6)\} \geq 3$.

Now we consider Theorem 4 when some parameters of d, d', p, q equal one. Observing that $\mathcal{M}_{1 \times d'}$ is exactly the same space as $\mathbb{C}^{d'}$, an SEB1 in $\mathcal{M}_{1 \times d'}$ is an orthonormal basis in $\mathbb{C}^{d'}$, and MUSEB1s in $\mathcal{M}_{1 \times d'}$ are indeed MUBs in $\mathbb{C}^{d'}$. Namely, $M_1(1, d') = N(d')$. Hence, Eq (1) is a special case of Theorem 4 when $k_1 = k_2 = d = p = 1$. When $p = 1$ (or $q = 1$), we have the following corollary.

Corollary 5. *If there are t_1 MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$, t_2 MUBs in \mathbb{C}^q , then there are at least $\min\{t_1, t_2\}$ MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{qd'}$. Namely,*

$$M_k(d, qd') \geq \min\{N(q), M_k(d, d')\}. \quad (9)$$

If $k = d \leq d'$, then

$$M_d(d, qd') \geq \min\{N(q), M_d(d, d')\}, \quad (10)$$

and

$$M_d(pd, d') \geq \min\{N(p), M_d(d, d')\}. \quad (11)$$

Remark 2. As $N(q) \geq 3$ for general $q \geq 2$ [18], by Eq. (10), if $M_d(d, d') \geq 2$, then $M_d(d, qd') \geq 2$ for any $q \geq 2$. This improves the result in [10], which only showed that if $M_d(d, d') \geq 2$, then $M_d(d, 2^l d') \geq 2$.

From Eq. (11), we give a method to construct MUSEBks from MUBs and MUMEBs. In fact, we can construct MUSEBds in $\mathbb{C}^{pd} \otimes \mathbb{C}^{d'}$ from MUBs in \mathbb{C}^p and MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ for any $p \geq 2$.

Example 3. We can construct three MUSEB3s in $\mathbb{C}^6 \otimes \mathbb{C}^6$ from three MUMEBs in $\mathbb{C}^3 \otimes \mathbb{C}^3$ and three MUBs in \mathbb{C}^2 . Namely, $M_3(6, 6) \geq 3$. For three MUMEBs in $\mathbb{C}^3 \otimes \mathbb{C}^3$, we construct the following three bases in $\mathcal{M}_{3 \times 3}$ with $w = e^{\frac{2\pi i}{3}}$ and $i = \sqrt{-1}$.

$$\begin{aligned}
\mathcal{S}_1 = & \left\{ \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & w+2 & w+2 \\ 2w^2+1 & w+2 & w^2+2w \\ w^2+2w & w+2 & 2w^2+1 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & w^2+2w & 2w^2+1 \\ 2w^2+1 & w^2+2w & w+2 \\ w^2+2w & w^2+2w & w^2+2w \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & 2w^2+1 & w^2+2w \\ 2w^2+1 & 2w^2+1 & 2w^2+1 \\ w^2+2w & 2w^2+1 & w+2 \end{pmatrix}, \right. \\
& \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & w+2 & w+2 \\ w+2 & w^2+2w & 2w^2+1 \\ w+2 & 2w^2+1 & w^2+2w \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & w^2+2w & 2w^2+1 \\ w+2 & 2w^2+1 & w^2+2w \\ w+2 & w+2 & w+2 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & 2w^2+1 & w^2+2w \\ w+2 & w+2 & w+2 \\ w+2 & w^2+2w & 2w^2+1 \end{pmatrix}, \\
& \left. \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & w+2 & w+2 \\ w^2+2w & 2w^2+1 & w+2 \\ 2w^2+1 & w^2+2w & w+2 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & w^2+2w & 2w^2+1 \\ w^2+2w & w+2 & 2w^2+1 \\ 2w^2+1 & 2w^2+1 & 2w^2+1 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} w+2 & 2w^2+1 & w^2+2w \\ w^2+2w & w^2+2w & w^2+2w \\ 2w^2+1 & w+2 & w^2+2w \end{pmatrix} \right\}, \\
\mathcal{S}_2 = & \left\{ \frac{1}{3\sqrt{3}} \begin{pmatrix} 0 & 3w^2 & 0 \\ 3w & 0 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 0 & 3 & 0 \\ 3w & 0 & 0 \\ 0 & 0 & 3w^2 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 0 & 3w & 0 \\ 3w & 0 & 0 \\ 0 & 0 & 3w \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 3w^2 & 0 & 0 \\ 0 & 0 & 3w \\ 0 & 3 & 0 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 3w^2 & 0 & 0 \\ 0 & 0 & 3 \\ 0 & 3w & 0 \end{pmatrix}, \right. \\
& \left. \frac{1}{3\sqrt{3}} \begin{pmatrix} 3w^2 & 0 & 0 \\ 0 & 0 & 3w^2 \\ 0 & 3w^2 & 0 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 0 & 0 & 3w^2 \\ 0 & 3w & 0 \\ 3 & 0 & 0 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 0 & 0 & 3w \\ 0 & 3w^2 & 0 \\ 3 & 0 & 0 \end{pmatrix}, \frac{1}{3\sqrt{3}} \begin{pmatrix} 0 & 0 & 3 \\ 0 & 3 & 0 \\ 3 & 0 & 0 \end{pmatrix} \right\}, \\
\mathcal{S}_3 = & \left\{ \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & w & 0 \\ 0 & 0 & w^2 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & w^2 & 0 \\ 0 & 0 & w \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & w^2 \\ 1 & 0 & 0 \\ 0 & w & 0 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & w \\ 1 & 0 & 0 \\ 0 & w^2 & 0 \end{pmatrix}, \right. \\
& \left. \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & w & 0 \\ 0 & 0 & w^2 \\ 1 & 0 & 0 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & w^2 & 0 \\ 0 & 0 & w \\ 1 & 0 & 0 \end{pmatrix} \right\}.
\end{aligned}$$

For three MUBs in \mathbb{C}^2 , choose

$$\mathcal{T}_1 = \{(0, 1), (1, 0)\}, \mathcal{T}_2 = \left\{ \frac{1}{\sqrt{2}}(1, 1), \frac{1}{\sqrt{2}}(1, -1) \right\}, \mathcal{T}_3 = \left\{ \frac{1}{\sqrt{2}}(1, i), \frac{1}{\sqrt{2}}(1, -i) \right\}.$$

Then $\{\mathcal{S}_1 \otimes \mathcal{T}_1, \mathcal{S}_2 \otimes \mathcal{T}_2, \mathcal{S}_3 \otimes \mathcal{T}_3\}$ is a set of three MUMEBs in $\mathbb{C}^3 \otimes \mathbb{C}^6$ by Eq. (10), and $\{\mathcal{T}_1^T \otimes \mathcal{S}_1 \otimes \mathcal{T}_1, \mathcal{T}_2^T \otimes \mathcal{S}_2 \otimes \mathcal{T}_2, \mathcal{T}_3^T \otimes \mathcal{S}_3 \otimes \mathcal{T}_3\}$ is a set of three MUSEB3s in $\mathbb{C}^6 \otimes \mathbb{C}^6$ by Eq. (11).

VI. LOWER BOUNDS FOR MUMEBs

Through out this section, we always assume that $d = 2^t p_1^{a_1} p_2^{a_2} \cdots p_s^{a_s}$, where p_1, p_2, \dots, p_s are distinct odd primes such that $3 \leq p_1^{a_1} \leq p_2^{a_2} \leq \cdots \leq p_s^{a_s}$. For convenience, we define $M_1(1, 1) := \infty$.

Theorem 6. *For any positive integer $d \geq 2$, we have $M_d(d, d) \geq 3$. In particular,*

- (i) $M_d(d, d) \geq \frac{p_1^{2a_1} - 1}{2}$ when $t = 0$;
- (ii) $M_d(d, d) \geq 3$ when $t = 1$;
- (iii) $M_d(d, d) \geq \min\{2(2^t - 1), \frac{p_1^{2a_1} - 1}{2}\}$ when $t \geq 2$.

Proof. By Eq. (8), we have

$$M_d(d, d) \geq \min\{M_{2^t}(2^t, 2^t), M_{p_1^{a_1}}(p_1^{a_1}, p_1^{a_1}), \dots, M_{p_s^{a_s}}(p_s^{a_s}, p_s^{a_s})\}.$$

Since $M_{p_i^{a_i}}(p_i^{a_i}, p_i^{a_i}) \geq \frac{p_i^{2a_i} - 1}{2}$ by [13], $M_d(d, d) \geq \min\{M_{2^t}(2^t, 2^t), \frac{p_1^{2a_1} - 1}{2}\}$. Then the three cases follow from the fact that $M_1(1, 1) = \infty$, $M_2(2, 2) = 3$ and $M_{2^t}(2^t, 2^t) \geq 2(2^t - 1)$ by [12]. \square

Theorem 6 gives a general lower bound $M_d(d, d) \geq 3$ for any $d \geq 2$, which is similar to the case of MUBs $N(q) \geq 3$ for any $q \geq 2$.

By Eq. (10) and Eq. (11), we have the following corollary.

Corollary 7. *For all integers $d, p, q \geq 2$, we have*

$$M_d(d, qd) \geq \min\{N(q), M_d(d, d)\} \geq 3, \quad (12)$$

$$M_d(pd, qd) \geq \min\{N(p), M_d(d, qd)\} \geq 3. \quad (13)$$

Remark 3. In [14], it was shown that $M_d(d, qd) \geq \min\{(p'_1)^{a'_1} + 1, M_d(d, d)\}$ when d is an *odd* number. Eq. (12) extends this result for any number $d \geq 2$.

We have the following corollary by applying Eq. (8) with Theorem 1 and Eq. (12).

Corollary 8. *There are at least two MUMEBs in $\mathbb{C}^{2d} \otimes \mathbb{C}^{3qd}$ for any positive integers q and d . Namely,*

$$M_{2d}(2d, 3qd) \geq 2. \quad (14)$$

There are at least two MUSEB2s in $\mathbb{C}^3 \otimes \mathbb{C}^{2k}$ for any $k \geq 1$. Namely,

$$M_2(3, 2k) \geq 2. \quad (15)$$

Remark 4. Corollary 8 shows that two MUMEBs exist in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ for infinitely many parameters d, d' satisfying $d \nmid d'$. The result of Eq. (15) is better than that in [15], which only showed that $M_2(3, 4k) \geq 2$ where $k = 2^l$.

Example 4. There are two MUMEBs in $\mathbb{C}^6 \otimes \mathbb{C}^9$, $\mathcal{R}_1 \otimes \mathcal{S}_1, \mathcal{R}_2 \otimes \mathcal{S}_2$. Namely, $M_6(6, 9) \geq 2$. Here, $\mathcal{R}_1, \mathcal{R}_2$ are from Sec. III, and $\mathcal{S}_1, \mathcal{S}_2$ are constructed in Example 3.

VII. CONCLUSION

In this paper we studied constructions of MUSEBks (MUMEBs) and provided several better lower bounds on the maximum size of MUSEBks. See Table I for a conclusion of all known results. By similar arguments as in [8], one can consider the issue of distinguishability of SEBks selected from a set of MUSEBks and the use of MUSEBks in a QKD setup. We can also consider the mutually unbiased measurements consisting of MUSEBks, which plays a special role in the problem of state determination [39].

For the existence of MUSEBks, there are still many open questions. For example, which d can achieve the upper bound $M_d(d, d) \leq d^2 - 1$ besides 2, 3, 5, 7, 11? Can we improve the lower bound of $M_d(d, qd) \geq 3$? Are there three MUMEBs in $\mathbb{C}^2 \otimes \mathbb{C}^3$, while there are three MUBs in \mathbb{C}^6 . Are there MUSEBks in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ when $k \nmid dd'$? Are there MUMEB in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ for any $d \nmid d'$? The minimum unsolved case is $\mathbb{C}^2 \otimes \mathbb{C}^5$.

Acknowledgments

FS and XZ were supported by NSFC under Grant No. 11771419, the Fundamental Research Funds for the Central Universities, and Anhui Initiative in Quantum Information Technologies under Grant No. AHY150200. LC was supported by the NNSF of China (Grant No. 11871089), and the Fundamental Research Funds for the Central Universities (Grant Nos. KG12080401 and ZG216S1902).

-
- [1] Julian Schwinger. Unitary operator bases. *Proceedings of the national academy of sciences of the United States Of America*, 46(4):570, 1960.
 - [2] Charles H Bennett and Gilles Brassard. Proceedings of the ieee international conference on computers, systems and signal processing.
 - [3] A Acin, E Jané, and G Vidal. Optimal estimation of quantum dynamics. *Physical Review A*, 64(5):050302, 2001.
 - [4] Giulio Chiribella, GM DAriano, P Perinotti, and MF Sacchi. Efficient use of quantum resources for the transmission of a reference frame. *Physical review letters*, 93(18):180503, 2004.

- [5] Marco Lucamarini and Stefano Mancini. Secure deterministic communication without entanglement. *Physical review letters*, 94(14):140501, 2005.
- [6] Giulio Chiribella, Giacomo Mauro D’Ariano, and Paolo Perinotti. Optimal cloning of unitary transformation. *Physical review letters*, 101(18):180504, 2008.
- [7] Andrew James Scott. Optimizing quantum process tomography with unitary 2-designs. *Journal of Physics A: Mathematical and Theoretical*, 41(5):055308, 2008.
- [8] Jesni Shamsul Shaari, Rinie NM Nasir, and Stefano Mancini. Mutually unbiased unitary bases. *Physical Review A*, 94(5):052328, 2016.
- [9] Yuan-Hong Tao, Hua Nan, Jun Zhang, and Shao-Ming Fei. Mutually unbiased maximally entangled bases in $\mathbb{C}^d \otimes \mathbb{C}^{kd}$. 2015.
- [10] Jun Zhang, Yuan-Hong Tao, Hua Nan, and Shao-Ming Fei. Construction of mutually unbiased bases in $\mathbb{C}^d \otimes \mathbb{C}^{2^l d}$. 2015.
- [11] Junying Liu, Minghui Yang, and Keqin Feng. Mutually unbiased maximally entangled bases in $\mathbb{C}^d \otimes \mathbb{C}^d$. *Quantum Information Processing*, 16(6):159, 2017.
- [12] Dengming Xu. Construction of mutually unbiased maximally entangled bases through permutations of hadamard matrices. *Quantum Information Processing*, 16:1–11, 2017.
- [13] Dengming Xu. Trace-2 excluded subsets of special linear groups over finite fields and mutually unbiased maximally entangled bases. *Quantum Information Processing*, 18(7):213, 2019.
- [14] Xiaoya Cheng and Yun Shang. New bounds of mutually unbiased maximally entangled bases in $\mathbb{C}^d \otimes \mathbb{C}^{kd}$. *Quantum Information & Computation*, 18(13-14):1152–1164, 2018.
- [15] Yi-Fan Han, Gui-Jun Zhang, Xin-Lei Yong, Ling-Shan Xu, and Yuan-Hong Tao. Mutually unbiased special entangled bases with schmidt number 2 in $\mathbb{C}^3 \otimes \mathbb{C}^{4k}$. *Quantum Information Processing*, 17(3):58, 2018.
- [16] Fei Shi, Xiande Zhang, and Yu Guo. Constructions of unextendible entangled bases. *Quantum Information Processing*, 18(10):324, 2019.
- [17] Yu Guo, Shuanping Du, Xiulan Li, and Shengjun Wu. Entangled bases with fixed schmidt number. *Journal of Physics A: Mathematical and Theoretical*, 48(24):245301, 2015.
- [18] Andreas Klappenecker and Martin Rötteler. Constructions of mutually unbiased bases. In *International Conference on Finite Fields and Applications*, pages 137–144. Springer, 2003.
- [19] Pawel Wocjan and Thomas Beth. New construction of mutually unbiased bases in square dimensions. *arXiv preprint quant-ph/0407081*, 2004.
- [20] M Wiesniak, T Paterek, and A Zeilinger. Entanglement in mutually unbiased bases. *New Journal of Physics*, 13(5):053047, 2011.
- [21] P. O. Boykin, M. Sitharam, P. H. Tiep, and P. Wocjan. Mutually Unbiased Bases and Orthogonal Decompositions of Lie Algebras, June 2005.
- [22] Stephen Brierley and Stefan Weigert. Maximal sets of mutually unbiased quantum states in dimension 6. *Phys. Rev. A*, 78:042312, Oct 2008.
- [23] Stephen Brierley and Stefan Weigert. Constructing mutually unbiased bases in dimension six. *Phys. Rev. A*, 79:052316, May 2009.
- [24] Philippe Jaming, Máté Matolcsi, Péter Móra, Ferenc Szöllösi, and Mihály Weiner. A generalized pauli problem and an infinite family of mub-triplets in dimension 6. *Journal of Physics A: Mathematical and Theoretical*, 42(24):245305, 2009.
- [25] Stephen Brierley. *Mutually Unbiased Bases in Low Dimensions*. PhD thesis, University of York, Department of Mathematics. 2009.
- [26] Stephen Brierley and Stefan Weigert. Mutually Unbiased Bases and Semi-definite Programming. *Journal of Physics: Conference Series*, 254:012008, June 2010.
- [27] T. Durt, B.-G. Englert, I. Bengtsson, and K. Życzkowski. On mutually unbiased bases. *Int. J. Quantum Information*, 8(4):535–640, 2010.
- [28] Daniel McNulty and Stefan Weigert. On the impossibility to extend triples of mutually unbiased product bases in dimension six. *International Journal of Quantum Information*, 10(05):1250056, 2012.
- [29] Daniel McNulty and Stefan Weigert. All mutually unbiased product bases in dimension 6. *Journal of Physics A: Mathematical and Theoretical*, 45(13):135307, 2012.
- [30] Philippe Raynal, Xin Lü, and Berthold-Georg Englert. Mutually unbiased bases in six dimensions: The four most distant bases. *Phys. Rev. A*, 83:062303, Jun 2011.
- [31] Ferenc Szöllösi. Complex hadamard matrices of order 6: a four-parameter family. *Journal of the London Mathematical Society*, 85(3):616–632, 2012.
- [32] D Goyeneche. Mutually unbiased triplets from non-affine families of complex hadamard matrices in dimension 6. *Journal of Physics A: Mathematical and Theoretical*, 46(10):105301, 2013.
- [33] Daniel McNulty and Stefan Weigert. The limited role of mutually unbiased product bases in dimension 6. *Journal of Physics A: Mathematical and Theoretical*, 45(10):102001, 2012.
- [34] Andrew S. Maxwell and Stephen Brierley. On properties of Karlsson Hadamards and sets of mutually unbiased bases in dimension six. *Linear Algebra and its Applications*, 466(0):296 – 306, 2015.
- [35] Daniel McNulty, Bogdan Pammer, and Stefan Weigert. Mutually unbiased product bases for multiple qudits. *Journal of Mathematical Physics*, 57(3), 2016.
- [36] Lin Chen and Li Yu. Product states and schmidt rank of mutually unbiased bases in dimension six. *Journal of Physics A: Mathematical and Theoretical*, 50(47):475304, 2017.
- [37] Lin Chen and Li Yu. Mutually unbiased bases in dimension six containing a product-vector basis. *Quantum Information*

Processing, 17(8):198, 2018.

- [38] Mengfan Liang, Mengyao Hu, Lin Chen, and Xiaoyu Chen. The h_2 -reducible matrix in four six-dimensional mutually unbiased bases. *Quantum Information Processing*, 18(11):352, 2019.
- [39] William K Wootters and Brian David Fields. Optimal state-determination by mutually unbiased measurements. *Annals of Physics*, 191(2):363–381, 1989.