

Group Theoretical Description of Mendeleev Periodic System

V. V. Varlamov*

Abstract

The group theoretical description of the periodic system of elements in the framework of the Rumer-Fet model is considered. We introduce the concept of a single quantum system, the generating core of which is an abstract C^* -algebra. It is shown that various concrete implementations of the operator algebra depend on the structure of generators of the fundamental symmetry group attached to the energy operator. In the case of generators of the complex shell of a group algebra of a conformal group, the spectrum of states of a single quantum system is given in the framework of the basic representation of the Rumer-Fet group, which leads to a group-theoretic interpretation of the Mendeleev's periodic system of elements. A mass formula is introduced that allows to give the termwise mass splitting for the main multiplet of the Rumer-Fet group. The masses of elements of the Seaborg table (eight-periodic extension of the Mendeleev table) are calculated starting from the atomic number $Z = 121$ to $Z = 220$. The continuation of Seaborg homology between lanthanides and actinides is established to the group of superactinides. A 10-periodic extension of the periodic table is introduced in the framework of the group-theoretic approach. The multiplet structure of the extended table periods is considered in detail. It is shown that the period lengths of the system of elements are determined by the structure of the basic representation of the Rumer-Fet group. The theoretical masses of the elements of 10th and 11th periods are calculated starting from $Z = 221$ to $Z = 364$. The concept of hypertwistor is introduced.

Keywords: periodic table, Bohr's model, Rumer-Fet model, conformal group, single quantum system, mass formulae, Seaborg table, homological series, twistors

1 Introduction

In 2019 marks the 150th anniversary of the discovery of the periodic law of chemical elements by Dmitry Ivanovich Mendeleev. Mendeleev's periodic table shed light on a huge number of experimental facts and allowed to predict the existence and basic properties of new, previously unknown, elements. However, the reasons (more precisely, root causes) for periodicity, in particular, the reasons for the periodic recurrence of similar electronic configurations of atoms, are still not clear to the end. Also, the limits of applicability of the periodic law have not yet been delineated – the controversy regarding the specifics of nuclear and electronic properties of atoms of heavy elements continues.

The now generally accepted structure of the periodic system, based on the Bohr model, proceeds from the fact that the arrangement of elements in the system with increasing their atomic numbers is uniquely determined by the individual features of the electronic structure of atoms

*Siberian State Industrial University, Kirova 42, Novokuznetsk 654007, Russia, e-mail: varlamov@sibsiu.ru

described in the framework of one-electronic approximation (Hartree method), and directly reflects the energy sequence of atomic orbitals of s -, p -, d -, f -shells populated by electrons with increasing their total number as the charge of the nucleus of the atom increases in accordance with the principle of minimum energy. However, this is only possible in the simplest version of Hartree approximation, but in the variant of the Hartree-Fock approximation total energy of an atom is not equal to the sum of orbital energies, and electron configuration of an atom is determined by the minimum of its full energy. As noted in the book [1], the traditional interpretation of the structure of the periodic system on the basis of the sequence of filling of electronic atomic orbitals in accordance with their relative energies ε_{nl} is very approximate, has, of course, a number of drawbacks and has narrow limits of applicability. There is no universal sequence of orbital energies ε_{nl} , moreover, such a sequence does not completely determine the order of the atomic orbitals settling by electrons, since it is necessary to take into account the configuration interactions (superposition of configurations in the multi-configuration approximation). And, of course, periodicity is not only and not completely orbital-energy effects. The reason for the repetition of similar electronic configurations of atoms in their ground states escapes us, and within one-electronic approximation can hardly be revealed at all. Moreover, it is possible that the theory of periodicity in general awaits a fate somewhat reminiscent of the fate of the theory of planetary retrogressions in the Ptolemaic system after the creation of the Copernican system. It is quite possible that what we call the periodicity principle is the result of the nonspatial symmetries of the atom (permutation and dynamical symmetries).

In 1971 academician V.A. Fock in his work [2] put the main question for the doctrine of the principle of periodicity and the theory of the periodic system: “Do the properties of atoms and their constituent parts fit into the framework of purely spatial representations, or do we need to somehow expand the concepts of space and spatial symmetry to accommodate the inherent degrees of freedom of atoms and their constituent parts?” [2, p. 108]. As is known, Bohr’s model in its original formulation uses quantum numbers relating to electrons in a field with spherical symmetry, which allowed Bohr to introduce the concept of closed electron shells and bring this concept closer to the periods of the Mendeleev’s table. Despite this success, the problem of explaining the periodic system was far from solved. Moreover, for all the depth and radicality of these new ideas, they still fit into the framework of conventional spatial representations. A further important step was associated with the discovery of the internal, not spatial, degree of freedom of the electron – spin, which is not a mechanical concept. The discovery of spin is closely related to the discovery of Pauli principle, which was formulated before quantum mechanics as requiring that each orbit, characterized by certain quantum numbers, contains no more than two electrons. At the end of the article [2] Fock himself answers his own question: “Purely spatial degrees of freedom of the electron is not enough to describe the properties of the electron shell of the atom and need to go beyond purely spatial concepts to express the laws that underlie the periodic system. The new degree of freedom of the electron – its spin – allows us to describe the properties of physical systems that are alien to classical concepts. This internal degree of electron freedom is essential for the formulation of the properties of multi-electron systems, thus for the theoretical justification of the Mendeleev’s periodic system” [2, c. 116].

In this paper, we consider the group theoretical description of the periodic system in the framework of the Rumer-Fet model. Unlike Bohr’s model, in which spatial and internal (spin) symmetries are combined on the basis of a classical composite system borrowed from celestial

mechanics¹, the Rumer-Fet group G describes non-spatial symmetries². Moreover, the Rumer-Fet model is entirely based on the mathematical apparatus of quantum mechanics and group theory without involving any classical analogies, such as the concept of a composite system. The concept of a composite system, which directly follows from the principle of separability (the basic principle of reductionism), is known to have limited application in quantum mechanics, since in the microcosm, in contrast to the composite structure of the macrocosm, the superposition structure prevails. Heisenberg argued that the concept of “consists of” does not work in particle physics. On the other hand, the problem of “critical” elements of Bohr’s model is also a consequence of visual spatial representations. Feynmann’s solution, representing the atomic nucleus as a point, leads to the Klein paradox for the element **Uts** (Untriseptium) with atomic number $Z = 137$. Another spatial image, used in the Greiner-Reinhardt solution, represents the atomic nucleus as a charged ball, resulting in a loss of electroneutrality for atoms above the value $Z = 173$ (see section 6).

The most important characteristic feature of the Rumer-Fet model is the representation of the periodic table of elements as a single quantum system. While Bohr’s model considers an atom of one element as a separate quantum system (and the atomic number is included in the theory as a parameter, so that there are as many quantum systems as there are elements), in the Rumer-Fet model atoms of various elements are considered as states of a single quantum system, connected to each other by the action of the symmetry group. A peculiar feature of the Rumer-Fet model is that it “ignores” the atomic structure underlying Bohr’s model. In contrast to Bohr’s model, which represents each atom as a composite aggregate of protons, neutrons and electrons, the Rumer-Fet model is distracted from the internal structure of each single atom, presenting the entire set of elements of the periodic table as a single quantum system of structureless states³. In this paper, the single quantum system \mathbf{U} is defined by a C^* -algebra consisting of the *energy operator* H and the generators of the *fundamental symmetry group* G_f attached to H . The states of the system \mathbf{U} are formed within the framework of the Gelfand-Naimark-Segal construction (GNS) [3, 4], that is, as cyclic representations of the operator algebra. Due to the generality of the task of the system \mathbf{U} and the flexibility of the GNS-construction for the each particular implementation of the operator algebra (the so-called “dressing” of the C^* -algebra), we obtain our (corresponding to this implementation) spectrum of states of the system \mathbf{U} ⁴. In section 2 of this article, a conformal group is considered as a fundamental symmetry group ($G_f = \text{SO}(2, 4)$). In this case, the concrete

¹It is obvious that the visual spatial image used in Bohr’s model is a vestige of classical representations. So, in the middle of 19th century, numerous attempts were made to build mechanical models of electromagnetic phenomena, even Maxwell’s treatise contains a large number of mechanical analogies. As time has shown, all mechanical models of electromagnetism turned out to be nothing more than auxiliary scaffolding, which were later discarded as unnecessary.

²The group G also contains the Lorentz group (rotation group of the Minkowski space-time) as a subgroup.

³The notion of the atom as a “structureless” state does not mean that there is no structure at all behind the concept. This only means that this structure is of a different order, not imported from the outside, from the “repertoire of classical physics”, but a structure that naturally follows from the mathematical apparatus of quantum mechanics (state vectors, symmetry group, Hilbert space, tensor products of Hilbert (\mathbb{K} -Hilbert) spaces, and so on).

⁴Thus, in the case when the generators of the fundamental symmetry group ($G_f = \text{SO}_0(1, 3)$ is the Lorentz group) attached to H are generators of the complex shell of the group algebra $\mathfrak{sl}(2, \mathbb{C})$ (see Appendix A), we obtain a linearly growing spectrum of state masses (“elementary particles”) [5]. In this case, the “dressing” of the operator algebra and the construction of the cyclic representations of the GNS-construction are carried out in the framework of spinor structure (charged, neutral, truly neutral (Majorana) states and their discrete symmetries set through morphisms of the spinor structure, see [6, 7, 9, 10, 11]). In [5] it is shown that the masses of “elementary particles” are multiples of the mass of the electron with an accuracy of 0.41%. Here there is a direct analogy with the electric charge. Any electric charge is a multiple of the charge of the electron, and a multiple of exactly. If any electric charge is absolutely a multiple of the electron charge, then in the case of masses this multiplicity takes place with an accuracy of 0.41% (on average).

implementation of the operator C^* -algebra is given by means of the generators of the complex shell of the group algebra $\mathfrak{so}(2, 4)$ attached to H and the twistor structure associated with the group $SU(2, 2)$ (the double covering of the conformal group). The complex shell of the algebra $\mathfrak{so}(2, 4)$ leads to a representation $F_{ss'}^+$ of the Rumer-Fet group, within which a group theoretical description of the periodic system of elements is given (section 3–4). At this point, atoms are considered as states (discrete stationary states) of the *matter spectrum*⁵, each atom is given by a state vector of the physical Hilbert space, in which a symmetry group acts, translating some state vectors into others (that is, a group that specifies quantum transitions between elements of the periodic system). In the section 6, Seaborg table (eight-periodic extension of the Mendeleev table) is formulated in the framework of the basic representation of the Rumer-Fet group for two different group chains, which specify the split of the main multiplet into smaller multiplets. It also calculates the average mass of the multiplets included in the Seaborg table (in addition to those multiplets that belong to the Mendeleev table with exception of elements **Uue** and **Ubn**). In subsection 6.1 the mass formula is introduced to allow a termwise mass splitting for the basic representation of the Rumer-Fet group. The masses of elements are calculated starting from the atomic number $Z = 121$ to $Z = 220$. In section 7 the 10-periodic extension of the Mendeleev table is studied. The multiplet structure of the extended table is considered in detail. It is shown that the period lengths of the system of elements are determined by the structure of the basic representation of the Rumer-Fet group. The theoretical masses of the elements of the 10th and 11th periods are calculated. In section 8 quantum transitions between state vectors of the physical Hilbert space, formed by the set of elements of periodic system, are considered.

It is possible to imagine an electron in any way: whether as a point (particle or wave), or a charged ball, or as an electron cloud on an atomic orbital, all these mental images only obscure the essence of the matter, because they remain within the framework of visual spatial representations. However, there is a mathematical structure that is far from visualization, and yet accurately describes the electron: it is a two-component spinor, the vector of the fundamental representation of the double covering $SL(2, \mathbb{C}) \simeq \mathbf{Spin}_+(1, 3)$ of the Lorentz group. Similarly, apart from any visual representations of the atom, it can be argued that the meaning is only the mathematical structure, which is directly derived from the symmetry group of the periodic system. In the section 9 it is shown that such structure is a hypertwistor acting in the \mathbb{K} -Hilbert space $\mathbf{H}_8 \otimes \mathbf{H}_\infty$.

Bohr's model does not explain the periodicity, but only approximates it within the framework of one-electronic Hartree approximation. Apparently, the explanation of the periodic law lies on the path indicated by academician Fock, that is, it is necessary to go beyond the classical (space-time) representations in the description of the periodic system of elements. It is obvious that the most suitable scheme of description in this way is the group-theoretic approach.

2 Single Quantum System

As already noted in the introduction, the starting point of the construction of the group theoretical description of the periodic system of elements is the concept of a single quantum system **U**. Following Heisenberg, we assume that at the fundamental level, the definition of the system **U** is based on two concepts: *energy* and *symmetry*. Let us define a single quantum system **U** by means of the following axiom system:

A.I (Energy and fundamental symmetry) *A single quantum system **U** at the fundamental level is characterized by a C^* -algebra \mathfrak{A} with a unit consisting of the energy operator H and the generators of the fundamental symmetry group G_f attached to H , forming a common system of eigenfunctions with H .*

⁵A term introduced by Heisenberg in the book [12] with reference to particle physics.

A.II (States and GNS construction) *The physical state of a C^* -algebra \mathfrak{A} is determined by the cyclic vector $|\Phi\rangle$ of the representation π of a C^* -algebra in a separable Hilbert space \mathbf{H}_∞ :*

$$\omega_\Phi(H) = \frac{\langle \Phi | \pi(H) \Phi \rangle}{\langle \Phi | \Phi \rangle}.$$

The set $PS(\mathfrak{A})$ of all pure states of a C^ -algebra \mathfrak{A} coincides with the set of all states $\omega_\Phi(H)$ associated with all irreducible cyclic representations π of an algebra \mathfrak{A} , $|\Phi\rangle \in \mathbf{H}_\infty$ (Gelfand-Naimark-Segal construction).*

A.III (Physical Hilbert space) *The set of all pure states $\omega_\Phi(H)$ under the condition $\omega_\Phi(H) \geq 0$ forms a physical Hilbert space \mathbf{H}_{phys} (in general, the space \mathbf{H}_{phys} is nonseparable). For each state vector $|\Psi\rangle \in \mathbf{H}_{\text{phys}}$ there is a unit ray $\Psi = e^{i\alpha} |\Psi\rangle$, where α runs through all real numbers and $\sqrt{\langle \Psi | \Psi \rangle} = 1$. The ray space is a quotient-space $\hat{H} = \mathbf{H}_{\text{phys}}/S^1$, that is, the projective space of one-dimensional subspaces of \mathbf{H}_{phys} . All states of a single quantum system \mathbf{U} are described by the unit rays.*

A.IV (Axiom of spectrality) *In \hat{H} there is a complete system of states with non-negative energy.*

A.V (Superposition principle) *The basic correspondence between physical states and elements of space \hat{H} involves the superposition principle of quantum theory, that is, there is a set of basic states such that arbitrary states can be constructed from them using linear superpositions.*

We choose a conformal group as the fundamental symmetry. The conformal group occurs in modern physics in a wide variety of situations and is essentially as universal as the Lorentz group, there are many relativistic theories and, similarly, conformal ones⁶.

3 Rumer-Fet group

In the previous section we define conformal group as the fundamental symmetry. The next logical step is to construct a concrete implementation of the operator algebra. We begin this construction by defining a complex shell of a group algebra $\mathfrak{so}(2, 4)$ that leads to a basic representation $F_{ss'}^+$ of the Rumer-Fet group⁷.

As is known [18], a system of fifteen generators of the conformal group $\text{SO}_0(2, 4)$ satisfies the following commutativity relations:

$$[\mathbf{L}_{\alpha\beta}, \mathbf{L}_{\gamma\delta}] = i(g_{\alpha\delta}\mathbf{L}_{\beta\gamma} + g_{\beta\gamma}\mathbf{L}_{\alpha\delta} - g_{\alpha\gamma}\mathbf{L}_{\beta\delta} - g_{\beta\delta}\mathbf{L}_{\alpha\gamma}),$$

$$(\alpha, \beta, \gamma, \delta = 1, \dots, 6, \alpha \neq \beta, \gamma \neq \delta).$$

⁶Moreover, as Segal showed [13], the Lie algebra of an inhomogeneous Lorentz group (that is, a Poincaré group) can be obtained by deformation from a conformal Lie algebra. In turn, the conformal Lie algebra is “rigid”, that is, cannot be obtained by deforming another Lie algebra. Because of this property, the conformal algebra (the algebra of a non-compact real pseudo-orthogonal group in a six-dimensional space with signature $(-, -, -, -, +, +)$) has a unique (complete) character and occupies a special place among other algebras.

⁷The first work in this direction is [14], where it was noted a striking similarity between the structure of the system of chemical elements and the structure of the energy spectrum of the hydrogen atom. This similarity is explained in [14] within the framework of the Fock representation F [15] for the group $\mathbf{Spin}(4)$ (the double covering of the group $\text{SO}(4)$). However, the main drawback of the description in [14] is the reducibility of the representation F , which did not allow to consider the system as “elementary” in the sense of group mechanics. In 1972, Konopelchenko [16] extended the Fock representation F to the representation F^+ of the conformal group, thus eliminating the above drawback. Further, based on the connection with the Madelung numbering, Fet defines the F_s^+ and $F_{ss'}^+$ representations (to define the $F_{ss'}^+$ representation, Madelung’s “lexicographic rule” had to be changed). After a rather long period of oblivion (totally undeserved), interest in the Rumer-Fet model will resume (see Kibler [17]).

Generators $\mathbf{L}_{\alpha\beta}$ form a basis of the group algebra $\mathfrak{so}(2, 4)$. In order to move to the complex shell of the algebra $\mathfrak{so}(2, 4)$, consider another system of generators proposed by Tsu Yao [19, 18]. Let

$$\begin{aligned}
\mathbf{J}_1 &= 1/2 (\mathbf{L}_{23} - \mathbf{L}_{14}), & \mathbf{J}_2 &= 1/2 (\mathbf{L}_{31} - \mathbf{L}_{24}), & \mathbf{J}_3 &= 1/2 (\mathbf{L}_{12} - \mathbf{L}_{34}), \\
\mathbf{K}_1 &= 1/2 (\mathbf{L}_{23} + \mathbf{L}_{14}), & \mathbf{K}_2 &= 1/2 (\mathbf{L}_{31} + \mathbf{L}_{24}), & \mathbf{K}_3 &= 1/2 (\mathbf{L}_{12} + \mathbf{L}_{34}), \\
\mathbf{P}_1 &= 1/2 (-\mathbf{L}_{35} - \mathbf{L}_{16}), & \mathbf{P}_2 &= 1/2 (\mathbf{L}_{45} - \mathbf{L}_{36}), & \mathbf{P}_0 &= 1/2 (-\mathbf{L}_{34} - \mathbf{L}_{56}), \\
\mathbf{Q}_1 &= 1/2 (\mathbf{L}_{35} - \mathbf{L}_{46}), & \mathbf{Q}_2 &= 1/2 (\mathbf{L}_{45} + \mathbf{L}_{36}), & \mathbf{Q}_0 &= 1/2 (\mathbf{L}_{34} - \mathbf{L}_{56}), \\
\mathbf{S}_1 &= 1/2 (-\mathbf{L}_{15} + \mathbf{L}_{26}), & \mathbf{S}_2 &= 1/2 (-\mathbf{L}_{25} - \mathbf{L}_{16}), & \mathbf{S}_0 &= 1/2 (\mathbf{L}_{12} - \mathbf{L}_{56}), \\
\mathbf{T}_1 &= 1/2 (-\mathbf{L}_{15} - \mathbf{L}_{26}), & \mathbf{T}_2 &= 1/2 (\mathbf{L}_{25} - \mathbf{L}_{16}), & \mathbf{T}_0 &= 1/2 (-\mathbf{L}_{12} - \mathbf{L}_{56}).
\end{aligned} \tag{1}$$

This system of eighteen generators is tied by the three relations

$$\mathbf{J}_3 - \mathbf{K}_3 = \mathbf{P}_0 - \mathbf{Q}_0, \quad \mathbf{J}_3 + \mathbf{K}_3 = \mathbf{S}_0 - \mathbf{T}_0, \quad \mathbf{P}_0 + \mathbf{Q}_0 = \mathbf{S}_0 + \mathbf{T}_0. \tag{2}$$

In virtue of independence of the generators $\mathbf{L}_{\alpha\beta}$ ($\alpha < \beta$), the system (1) defines a surplus system of generators of $\text{SO}_0(2, 4)$, from which we can obtain the basis of $\mathfrak{so}(2, 4)$ excluding three generators by means of (2).

Introducing the generators

$$\begin{aligned}
\mathbf{J}_{\pm} &= \mathbf{J}_1 \pm i\mathbf{J}_2, & \mathbf{P}_{\pm} &= \mathbf{P}_1 \pm i\mathbf{P}_2, & \mathbf{S}_{\pm} &= \mathbf{S}_1 \pm i\mathbf{S}_2, \\
\mathbf{K}_{\pm} &= \mathbf{K}_1 \pm i\mathbf{K}_2, & \mathbf{Q}_{\pm} &= \mathbf{Q}_1 \pm i\mathbf{Q}_2, & \mathbf{T}_{\pm} &= \mathbf{T}_1 \pm i\mathbf{T}_2,
\end{aligned} \tag{3}$$

we come to the complex shell of the algebra $\mathfrak{so}(2, 4)$.

Then

$$\begin{aligned}
[\mathbf{J}_3, \mathbf{J}_+] &= \mathbf{J}_+, & [\mathbf{J}_3, \mathbf{J}_-] &= -\mathbf{J}_-, & [\mathbf{J}_+, \mathbf{J}_-] &= 2\mathbf{J}_3, \\
[\mathbf{K}_3, \mathbf{K}_+] &= \mathbf{K}_+, & [\mathbf{K}_3, \mathbf{K}_-] &= -\mathbf{K}_-, & [\mathbf{K}_+, \mathbf{K}_-] &= 2\mathbf{K}_3, \\
[\mathbf{P}_0, \mathbf{P}_+] &= \mathbf{P}_+, & [\mathbf{P}_0, \mathbf{P}_-] &= -\mathbf{P}_-, & [\mathbf{P}_+, \mathbf{P}_-] &= -2\mathbf{P}_0, \\
[\mathbf{Q}_0, \mathbf{Q}_+] &= \mathbf{Q}_+, & [\mathbf{Q}_0, \mathbf{Q}_-] &= -\mathbf{Q}_-, & [\mathbf{Q}_+, \mathbf{Q}_-] &= -2\mathbf{Q}_0, \\
[\mathbf{S}_0, \mathbf{S}_+] &= \mathbf{S}_+, & [\mathbf{S}_0, \mathbf{S}_-] &= -\mathbf{S}_-, & [\mathbf{S}_+, \mathbf{S}_-] &= -2\mathbf{S}_0, \\
[\mathbf{T}_0, \mathbf{T}_+] &= \mathbf{T}_+, & [\mathbf{T}_0, \mathbf{T}_-] &= -\mathbf{T}_-, & [\mathbf{T}_+, \mathbf{T}_-] &= -2\mathbf{T}_0, \\
[\mathbf{J}_i, \mathbf{K}_j] &= 0 \quad (i, j = +, -, 3), \\
[\mathbf{J}_+, \mathbf{P}_+] &= 0, & [\mathbf{J}_+, \mathbf{P}_-] &= -\mathbf{T}_-, & [\mathbf{J}_+, \mathbf{P}_0] &= -1/2\mathbf{J}_+, \\
[\mathbf{J}_-, \mathbf{P}_+] &= \mathbf{T}_+, & [\mathbf{J}_-, \mathbf{P}_-] &= 0, & [\mathbf{J}_-, \mathbf{P}_0] &= 1/2\mathbf{J}_-, \\
[\mathbf{J}_3, \mathbf{P}_+] &= 1/2\mathbf{P}_+, & [\mathbf{J}_3, \mathbf{P}_-] &= -1/2\mathbf{P}_-, & [\mathbf{J}_3, \mathbf{P}_0] &= 0, \\
[\mathbf{J}_+, \mathbf{Q}_+] &= \mathbf{S}_+, & [\mathbf{J}_+, \mathbf{Q}_-] &= 0, & [\mathbf{J}_+, \mathbf{Q}_0] &= 1/2\mathbf{J}_+, \\
[\mathbf{J}_-, \mathbf{Q}_+] &= 0, & [\mathbf{J}_-, \mathbf{Q}_-] &= -\mathbf{S}_-, & [\mathbf{J}_-, \mathbf{Q}_0] &= -1/2\mathbf{J}_-, \\
[\mathbf{J}_3, \mathbf{Q}_+] &= -1/2\mathbf{Q}_+, & [\mathbf{J}_3, \mathbf{Q}_-] &= 1/2\mathbf{Q}_-, & [\mathbf{J}_3, \mathbf{Q}_0] &= 0, \\
[\mathbf{J}_+, \mathbf{S}_+] &= 0, & [\mathbf{J}_+, \mathbf{S}_-] &= -\mathbf{Q}_-, & [\mathbf{J}_+, \mathbf{S}_0] &= -1/2\mathbf{J}_+, \\
[\mathbf{J}_-, \mathbf{S}_+] &= \mathbf{Q}_+, & [\mathbf{J}_-, \mathbf{S}_-] &= 0, & [\mathbf{J}_-, \mathbf{S}_0] &= 1/2\mathbf{J}_-, \\
[\mathbf{J}_3, \mathbf{S}_+] &= 1/2\mathbf{S}_+, & [\mathbf{J}_3, \mathbf{Q}_-] &= -1/2\mathbf{S}_-, & [\mathbf{J}_3, \mathbf{S}_0] &= 0, \\
[\mathbf{J}_+, \mathbf{T}_+] &= \mathbf{P}_+, & [\mathbf{J}_+, \mathbf{T}_-] &= 0, & [\mathbf{J}_+, \mathbf{T}_0] &= 1/2\mathbf{J}_+,
\end{aligned}$$

$$\begin{aligned}
[\mathbf{J}_-, \mathbf{T}_+] &= 0, & [\mathbf{J}_-, \mathbf{T}_-] &= -\mathbf{P}_-, & [\mathbf{J}_-, \mathbf{T}_0] &= -1/2\mathbf{J}_-, \\
[\mathbf{J}_3, \mathbf{T}_+] &= -1/2\mathbf{T}_+, & [\mathbf{J}_3, \mathbf{T}_-] &= 1/2\mathbf{T}_-, & [\mathbf{J}_3, \mathbf{T}_0] &= 0, \\
[\mathbf{K}_+, \mathbf{P}_+] &= -\mathbf{S}_+, & [\mathbf{K}_+, \mathbf{P}_-] &= 0, & [\mathbf{K}_+, \mathbf{P}_0] &= 1/2\mathbf{K}_+, \\
[\mathbf{K}_-, \mathbf{P}_+] &= 0, & [\mathbf{K}_-, \mathbf{P}_-] &= \mathbf{S}_-, & [\mathbf{K}_-, \mathbf{P}_0] &= -1/2\mathbf{K}_-, \\
[\mathbf{K}_3, \mathbf{P}_+] &= -1/2\mathbf{P}_+, & [\mathbf{K}_3, \mathbf{P}_-] &= 1/2\mathbf{P}_-, & [\mathbf{K}_3, \mathbf{P}_0] &= 0, \\
[\mathbf{K}_+, \mathbf{Q}_+] &= 0, & [\mathbf{K}_+, \mathbf{Q}_-] &= \mathbf{T}_-, & [\mathbf{K}_+, \mathbf{Q}_0] &= -1/2\mathbf{K}_+, \\
[\mathbf{K}_-, \mathbf{Q}_+] &= -\mathbf{T}_+, & [\mathbf{K}_-, \mathbf{Q}_-] &= 0, & [\mathbf{K}_-, \mathbf{Q}_0] &= 1/2\mathbf{K}_-, \\
[\mathbf{K}_3, \mathbf{Q}_+] &= 1/2\mathbf{Q}_+, & [\mathbf{K}_3, \mathbf{Q}_-] &= -1/2\mathbf{Q}_-, & [\mathbf{K}_3, \mathbf{Q}_0] &= 0, \\
[\mathbf{K}_+, \mathbf{S}_+] &= 0, & [\mathbf{K}_+, \mathbf{S}_-] &= \mathbf{P}_-, & [\mathbf{K}_+, \mathbf{S}_0] &= -1/2\mathbf{K}_+, \\
[\mathbf{K}_-, \mathbf{S}_+] &= -\mathbf{P}_+, & [\mathbf{K}_-, \mathbf{S}_-] &= 0, & [\mathbf{K}_-, \mathbf{S}_0] &= 1/2\mathbf{K}_-, \\
[\mathbf{K}_3, \mathbf{S}_+] &= 1/2\mathbf{S}_+, & [\mathbf{K}_3, \mathbf{S}_-] &= -1/2\mathbf{S}_-, & [\mathbf{K}_3, \mathbf{S}_0] &= 0, \\
[\mathbf{K}_+, \mathbf{T}_+] &= -\mathbf{Q}_+, & [\mathbf{K}_+, \mathbf{T}_-] &= 0, & [\mathbf{K}_+, \mathbf{T}_0] &= 1/2\mathbf{K}_+, \\
[\mathbf{K}_-, \mathbf{T}_+] &= 0, & [\mathbf{K}_-, \mathbf{T}_-] &= \mathbf{Q}_-, & [\mathbf{K}_-, \mathbf{T}_0] &= -1/2\mathbf{K}_-, \\
[\mathbf{K}_3, \mathbf{T}_+] &= -1/2\mathbf{T}_+, & [\mathbf{K}_3, \mathbf{T}_-] &= 1/2\mathbf{T}_-, & [\mathbf{K}_3, \mathbf{T}_0] &= 0, \\
[\mathbf{P}_i, \mathbf{Q}_j] &= 0 \quad (i, j = +, -, 0), \\
[\mathbf{P}_+, \mathbf{S}_+] &= 0, & [\mathbf{P}_+, \mathbf{S}_-] &= \mathbf{K}_-, & [\mathbf{P}_+, \mathbf{S}_0] &= -1/2\mathbf{P}_+, \\
[\mathbf{P}_-, \mathbf{S}_+] &= -\mathbf{K}_+, & [\mathbf{P}_-, \mathbf{S}_-] &= 0, & [\mathbf{P}_-, \mathbf{S}_0] &= 1/2\mathbf{P}_-, \\
[\mathbf{P}_0, \mathbf{S}_+] &= 1/2\mathbf{S}_+, & [\mathbf{P}_0, \mathbf{S}_-] &= -1/2\mathbf{S}_-, & [\mathbf{P}_0, \mathbf{S}_0] &= 0, \\
[\mathbf{P}_+, \mathbf{T}_+] &= 0, & [\mathbf{P}_+, \mathbf{T}_-] &= -\mathbf{J}_+, & [\mathbf{P}_+, \mathbf{T}_0] &= -1/2\mathbf{P}_+, \\
[\mathbf{P}_-, \mathbf{T}_+] &= \mathbf{J}_-, & [\mathbf{P}_-, \mathbf{T}_-] &= 0, & [\mathbf{P}_-, \mathbf{T}_0] &= 1/2\mathbf{P}_-, \\
[\mathbf{P}_0, \mathbf{T}_+] &= 1/2\mathbf{T}_+, & [\mathbf{P}_0, \mathbf{T}_-] &= -1/2\mathbf{T}_-, & [\mathbf{P}_0, \mathbf{T}_0] &= 0, \\
[\mathbf{Q}_+, \mathbf{S}_+] &= 0, & [\mathbf{Q}_+, \mathbf{S}_-] &= -\mathbf{J}_-, & [\mathbf{Q}_+, \mathbf{S}_0] &= -1/2\mathbf{Q}_+, \\
[\mathbf{Q}_-, \mathbf{S}_+] &= \mathbf{J}_+, & [\mathbf{Q}_-, \mathbf{S}_-] &= 0, & [\mathbf{Q}_-, \mathbf{S}_0] &= 1/2\mathbf{Q}_-, \\
[\mathbf{Q}_0, \mathbf{S}_+] &= 1/2\mathbf{S}_+, & [\mathbf{Q}_0, \mathbf{S}_-] &= -1/2\mathbf{S}_-, & [\mathbf{Q}_0, \mathbf{S}_0] &= 0, \\
[\mathbf{Q}_+, \mathbf{T}_+] &= 0, & [\mathbf{Q}_+, \mathbf{T}_-] &= \mathbf{K}_+, & [\mathbf{Q}_+, \mathbf{T}_0] &= -1/2\mathbf{Q}_+, \\
[\mathbf{Q}_-, \mathbf{T}_+] &= -\mathbf{K}_-, & [\mathbf{Q}_-, \mathbf{T}_-] &= 0, & [\mathbf{Q}_-, \mathbf{T}_0] &= 1/2\mathbf{Q}_-, \\
[\mathbf{Q}_0, \mathbf{T}_+] &= 1/2\mathbf{T}_+, & [\mathbf{Q}_0, \mathbf{T}_-] &= -1/2\mathbf{T}_-, & [\mathbf{Q}_0, \mathbf{T}_0] &= 0, \\
[\mathbf{S}_i, \mathbf{T}_j] &= 0 \quad (i, j = +, -, 0).
\end{aligned}$$

Let us consider a special representation of the conformal group $\text{SO}_0(2, 4)$, which is analogous to the van der Waerden representation (A.5) for the Lorentz group $\text{SO}_0(1, 3)$ (see Appendix A). This *local* representation of $\text{SO}_0(2, 4)$ is related immediately with the Fock representation for the group $\text{SO}(4)$ (see Appendix B). In essence this representation is an extension of the Fock representation

for $\text{SO}(4)$ to the unitary representation of the conformal group $\text{SO}_0(2, 4)$ in the *Fock space* \mathfrak{F} by the basis (B.2). Using generators (3) of the complex shell of the algebra $\mathfrak{so}(2, 4)$, we obtain

$$\begin{aligned}
\mathbf{J}_-|j, \sigma, \tau\rangle &= \sqrt{(j+\sigma)(j-\sigma+1)}|j, \sigma-1, \tau\rangle, \\
\mathbf{J}_+|j, \sigma, \tau\rangle &= \sqrt{(j-\sigma)(j+\sigma+1)}|j, \sigma+1, \tau\rangle, \\
\mathbf{J}_3|j, \sigma, \tau\rangle &= \sigma|j, \sigma, \tau\rangle, \\
\mathbf{K}_-|j, \sigma, \tau\rangle &= \sqrt{(j+\tau)(j-\tau+1)}|j, \sigma, \tau-1\rangle, \\
\mathbf{K}_+|j, \sigma, \tau\rangle &= \sqrt{(j-\tau)(j+\tau+1)}|j, \sigma, \tau+1\rangle, \\
\mathbf{K}_3|j, \sigma, \tau\rangle &= \tau|j, \sigma, \tau\rangle, \\
\\
\mathbf{P}_-|j, \sigma, \tau\rangle &= -i\sqrt{(j+\sigma)(j-\tau)}\left|j-\frac{1}{2}, \sigma-\frac{1}{2}, \tau+\frac{1}{2}\right\rangle, \\
\mathbf{P}_+|j, \sigma, \tau\rangle &= i\sqrt{(j-\tau+1)(j+\sigma+1)}\left|j+\frac{1}{2}, \sigma+\frac{1}{2}, \tau-\frac{1}{2}\right\rangle, \\
\mathbf{P}_0|j, \sigma, \tau\rangle &= \left(j+\frac{\sigma-\tau+1}{2}\right)\sigma|j, \sigma, \tau\rangle, \\
\mathbf{Q}_-|j, \sigma, \tau\rangle &= i\sqrt{(j+\tau)(j-\sigma)}\left|j-\frac{1}{2}, \sigma+\frac{1}{2}, \tau-\frac{1}{2}\right\rangle, \\
\mathbf{Q}_+|j, \sigma, \tau\rangle &= -i\sqrt{(j-\sigma+1)(j+\tau+1)}\left|j+\frac{1}{2}, \sigma-\frac{1}{2}, \tau+\frac{1}{2}\right\rangle, \\
\mathbf{Q}_0|j, \sigma, \tau\rangle &= \left(j-\frac{\sigma-\tau-1}{2}\right)|j, \sigma, \tau\rangle, \\
\\
\mathbf{S}_-|j, \sigma, \tau\rangle &= i\sqrt{(j+\sigma)(j+\tau)}\left|j-\frac{1}{2}, \sigma-\frac{1}{2}, \tau-\frac{1}{2}\right\rangle, \\
\mathbf{S}_+|j, \sigma, \tau\rangle &= -i\sqrt{(j+\tau+1)(j+\sigma+1)}\left|j+\frac{1}{2}, \sigma+\frac{1}{2}, \tau+\frac{1}{2}\right\rangle, \\
\mathbf{S}_0|j, \sigma, \tau\rangle &= \left(j+\frac{\sigma+\tau+1}{2}\right)\sigma|j, \sigma, \tau\rangle, \\
\mathbf{T}_-|j, \sigma, \tau\rangle &= -i\sqrt{(j-\tau)(j-\sigma)}\left|j-\frac{1}{2}, \sigma+\frac{1}{2}, \tau+\frac{1}{2}\right\rangle, \\
\mathbf{T}_+|j, \sigma, \tau\rangle &= i\sqrt{(j-\sigma+1)(j-\tau+1)}\left|j+\frac{1}{2}, \sigma-\frac{1}{2}, \tau-\frac{1}{2}\right\rangle, \\
\mathbf{T}_0|j, \sigma, \tau\rangle &= \left(j-\frac{\sigma+\tau-1}{2}\right)|j, \sigma, \tau\rangle.
\end{aligned} \tag{4}$$

The formulas (4) define an unitary representation of the conformal group $\text{SO}_0(2, 4)$ in the Fock space \mathfrak{F} . The formulas (4) include the formulas for \mathbf{J}_k , \mathbf{K}_k , giving representations Φ_n in subspaces \mathfrak{F}_n (see (B.1), where $j_1 = j_2 = j$) and thus the Fock representation Φ on the subgroup $\text{SO}(4)$. Moreover, if we restrict $\text{SO}_0(2, 4)$ to a subgroup $\text{SO}_0(1, 3)$ (Lorentz group), we obtain the van der Waerden representation (A.5) given by the generators \mathbf{X}_k , \mathbf{Y}_k , which proves the similarity of the complex shells of the group algebras $\mathfrak{so}(2, 4)$ and $\mathfrak{sl}(2, \mathbb{C})$. The representation, defined by (4), is called *an extension F^+ of the Fock representation on the conformal group* [18]. The representation F^+ is insufficient yet for the description of periodic system of elements. With that end in view it

is necessary to include a fourth Madelung number s (which is analogous to a spin) that leads to a group (the first “doubling”)

$$\text{SO}(2, 4) \otimes \text{SU}(2). \quad (5)$$

A representation $F_s^+ = \varphi_2 \otimes F^+$ of the group (5), where φ_2 is an unitary representation of the group $\text{SU}(2)$ in the space $C(2)$, already satisfies to this requirement (inclusion of the Madelung number s). A basis of the space $\mathfrak{F}^2 = C(2) \otimes \mathfrak{F}$ of the representation F_s^+ has the form

$$|n, l, m, s\rangle, \quad n = 1, 2, \dots; \quad l = 0, 1, \dots, n-1; \quad m = -l, -l+1, \dots, l-1, l; \quad s = -1/2, 1/2. \quad (6)$$

Here n, l, m are quantum numbers of the conformal group.

Let τ_k be generators of the Lie algebra of $\text{SU}(2)$, then a generator τ_3 commutes with the all generators of the subgroup $\text{SO}(2, 4) \otimes \mathbf{1}$. For that reason generators $\mathbf{R}_0, \mathbf{L}^2, \mathbf{J}_3 + \mathbf{K}_3, \tau_3$ commute with the each other. Eigenvectors of the operators, representing these generators in the space \mathfrak{F}^2 , have the form

$$\left| n, l, m, \frac{1}{2} \right\rangle = \begin{bmatrix} \Psi_{nlm}^1 \\ 0 \end{bmatrix}, \quad \left| n, l, m, -\frac{1}{2} \right\rangle = \begin{bmatrix} 0 \\ \Psi_{nlm}^2 \end{bmatrix}$$

with eigenvalues $n, l(l+1), m, \frac{1}{2}$ and $n, l(l+1), m, -\frac{1}{2}$. An action of the operators, representing generators τ_+, τ_-, τ_3 in the space \mathfrak{F}^2 , is defined by the following formulas:

$$\tau_+ \left| n, l, m, -\frac{1}{2} \right\rangle = \left| n, l, m, \frac{1}{2} \right\rangle, \quad \tau_- \left| n, l, m, \frac{1}{2} \right\rangle = \left| n, l, m, -\frac{1}{2} \right\rangle,$$

$$\tau_3 |n, l, m, s\rangle = s |n, l, m, s\rangle.$$

In virtue of the Madelung numbering the basis $|n, l, m, s\rangle$ stands in one-to-one correspondence with the elements of periodic system. A relation between the arrangement of the element in the Mendeleev table and a number collection (n, l, m, s) is defined by a so-called Madelung “lexicographic rule” $Z \leftrightarrow (n, l, m, s)$ ⁸:

- 1) elements are arranged in increasing order of atomic number Z ;
- 2) collections (n, l, m, s) are arranged in increasing order of $n+l$; at the given $n+l$ in increasing order of n ; at the given $n+l, n$ in increasing order of m ; at the given $n+l, n, m$ in increasing order of s ;
- 3) Z -th element corresponds to Z -th collection.

In Madelung numbering the sum $n+l$ does not have a group sense: it is a sum of quantum number n (an eigenvalue of the operator $\mathbf{R}_0 = -\mathbf{L}_{56}$) and the number l which is not quantum number. In this case, a quantum number is $l(l+1)$ (an eigenvalue of the operator $\mathbf{L}^2 = \mathbf{L}_{12}^2 + \mathbf{L}_{23}^2 + \mathbf{L}_{31}^2$), and l only defines this quantum number. Therefore, in accordance with group theoretical viewpoint the number $n+l$ should be excluded from the formulation of “lexicographic rule”. In [18] Fet introduced a new quantum number ν , which equal to $\nu = 1/2(n+l+1)$ for the odd value of $n+l$ and $\nu = 1/2(n+l)$ for the even value of $n+l$. Introduction of the quantum number ν allows us to change Madelung numbering, that leads to the following “lexicographic rule” (Fet rule):

⁸Erwin Madelung was the first to apply “hydrogen” quantum numbers n, l, m, s to the numbering of elements of the periodic table. It should be noted that the numbers n, l, m, s are not quantum numbers in Bohr’s model, because in this model there is no single quantum-mechanical description of the system of elements, the latter is assigned an atomic number Z , distinguishing rather than combining individual quantum systems. The resulting classification of elements Madelung called “empirical”, because he could not connect it with the Bohr model. Apparently, it was because of the lack of theoretical justification at the time (20-ies of the last century), he published it as a reference material in [20]. Theoretical justification (understanding of the nature of Madelung numbers) was later given by Fet [18] from the position of group-theoretic vision. The history of Madelung numbers was also developed in a slightly different direction in the works of Kleczkowski [21], where the filling of the electronic levels of the atom was considered according to the rule of sequential filling of $(n+l)$ -groups (the so-called Madelung-Kleczkowski groups).

- 1) elements are arranged in increasing order of atomic number Z ;
- 2) collections $(\nu, \lambda, \mu, s, s')$ are arranged in increasing order of ν ; at the given ν in increasing order of s' ; at the given ν, s' in decreasing order of λ ; at the given ν, s', λ in increasing order of μ ; at the given ν, s', λ, μ in increasing order of s ;
- 3) Z -th element corresponds to Z -th collection.

The introduction of the fifth quantum number leads to another (second) “doubling” of the representation space. In this space we have the Rumer-Fet group

$$\text{SO}(2, 4) \otimes \text{SU}(2) \otimes \text{SU}(2)'. \quad (7)$$

A representation $F_{ss'}^+ = \varphi_2' \otimes F_s^+$ of the group (7), where φ_2' is an unitary representation of the group $\text{SU}(2)'$ in the space $C(2)$, satisfies to the requirement of inclusion of the fifth quantum number. A basis of the space $\mathfrak{F}^4 = C(2) \otimes \mathfrak{F}^2$ of the representation $F_{ss'}^+$ has the form

$$\begin{aligned} |\nu, \lambda, \mu, s, s'\rangle, \quad \nu = 1, 2, \dots; \lambda = 0, 1, \dots, \nu - 1; \\ \mu = -\lambda, -\lambda + 1, \dots, \lambda - 1, \lambda; s = -1/2, 1/2, s' = -1/2, 1/2. \end{aligned} \quad (8)$$

Let τ'_k be generators of the Lie algebra $\text{SU}(2)'$, then the generator τ'_3 commutes with the all generators of the subgroup $\text{SO}(2, 4) \otimes \text{SU}(2) \otimes \mathbf{1}$. Therefore, generators $\mathbf{R}_0 = -\mathbf{L}_{56} = \mathbf{P}_0 + \mathbf{Q}_0 = \mathbf{S}_0 + \mathbf{T}_0, \mathbf{L}^2, \mathbf{J}_3 + \mathbf{K}_3, \tau_3, \tau'_3$ commute with the each other. Common eigenvectors of the operators, which represent these generators in the space \mathfrak{F}^4 , have the form

$$\left| \nu, \lambda, \mu, s, +\frac{1}{2} \right\rangle = \begin{bmatrix} \Psi_{\nu\lambda,\mu}^1 \\ \Psi_{\nu\lambda,\mu}^2 \\ 0 \\ 0 \end{bmatrix}, \quad \left| \nu, \lambda, \mu, s, -\frac{1}{2} \right\rangle = \begin{bmatrix} 0 \\ 0 \\ \Psi_{\nu\lambda,\mu}^3 \\ \Psi_{\nu\lambda,\mu}^4 \end{bmatrix}$$

with eigenvalues $\nu, \lambda(\lambda + 1), \mu, s, \frac{1}{2}$ and $\nu, \lambda(\lambda + 1), \mu, s, -\frac{1}{2}$.

An action of the operators, representing the generators τ_+, τ_-, τ_3 and $\tau'_+, \tau'_-, \tau'_3$ in the space \mathfrak{F}^4 , is defined by the following formulas:

$$\begin{aligned} \tau_+ \left| \nu, \lambda, \mu, -\frac{1}{2}, s' \right\rangle &= \left| \nu, \lambda, \mu, \frac{1}{2}, s' \right\rangle, \quad \tau_- \left| \nu, \lambda, \mu, \frac{1}{2}, s' \right\rangle = \left| \nu, \lambda, \mu, -\frac{1}{2}, s' \right\rangle, \\ \tau_3 \left| \nu, \lambda, \mu, s, s' \right\rangle &= s \left| \nu, \lambda, \mu, s, s' \right\rangle. \\ \tau'_+ \left| \nu, \lambda, \mu, s, -\frac{1}{2} \right\rangle &= \left| \nu, \lambda, \mu, s, \frac{1}{2} \right\rangle, \quad \tau'_- \left| \nu, \lambda, \mu, s, \frac{1}{2} \right\rangle = \left| \nu, \lambda, \mu, s, -\frac{1}{2} \right\rangle, \\ \tau'_3 \left| \nu, \lambda, \mu, s, s' \right\rangle &= s' \left| \nu, \lambda, \mu, s, s' \right\rangle. \end{aligned}$$

“Addresses” of the elements (see Fig. 1) are defined by a collection of quantum numbers of the Rumer-Fet group (7), which numbering basis vectors $|\nu, \lambda, \mu, s, s'\rangle$ of the space \mathfrak{F}^4 . Generators of the Lie algebra of $\text{SO}(2, 4)$ act on the quantum numbers ν, λ, μ by means of the formulas (4) with a replacement of n, l, m via ν, λ, μ .

There is the following chain of groups:

$$G \supset G_1 \supset G_2 \longmapsto \text{SO}(2, 4) \otimes \text{SU}(2) \otimes \text{SU}(2) \supset \text{SO}(4) \otimes \text{SU}(2) \supset \text{SO}(3) \otimes \text{SU}(2). \quad (9)$$

	$\overbrace{s'=-1/2 \quad s'=1/2}^{\nu=1}$	$\overbrace{s'=-1/2 \quad s'=1/2}^{\nu=2}$	$\overbrace{s'=-1/2 \quad s'=1/2}^{\nu=3}$	$\overbrace{s'=-1/2 \quad s'=1/2}^{\nu=4}$				
$\lambda = 0$	<div>H He</div>	<div>Li Be</div>	<div>Na Mg</div>	<div>K Ca</div>	<div>Rb Sr</div>	<div>Cs Ba</div>	<div>Fr Ra</div>	<div>Uue Ubn</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 0$
$\lambda = 1$	{	<div>B C</div>	<div>Al Si</div>	<div>Ga Ge</div>	<div>In Sn</div>	<div>Tl Pb</div>	<div>Nh Fl</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = -1$	
		<div>N O</div>	<div>P S</div>	<div>As Se</div>	<div>Sb Te</div>	<div>Bi Po</div>	<div>Mc Lv</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 0$	
		<div>F Ne</div>	<div>Cl Ar</div>	<div>Br Kr</div>	<div>I Xe</div>	<div>At Rn</div>	<div>Ts Og</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 1$	
$\lambda = 2$	{		<div>Sc Ti</div>	<div>Y Zr</div>	<div>Lu Hf</div>	<div>Lr Rf</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = -2$		
		<div>V Cr</div>	<div>Nb Mo</div>	<div>Ta W</div>	<div>Db Sg</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = -1$			
		<div>Mn Fe</div>	<div>Tc Ru</div>	<div>Re Os</div>	<div>Bh Hs</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 0$			
		<div>Co Ni</div>	<div>Rh Pd</div>	<div>Ir Pt</div>	<div>Mt Ds</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 1$			
		<div>Cu Zn</div>	<div>Ag Cd</div>	<div>Au Hg</div>	<div>Rg Cn</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 2$			
$\lambda = 3$	{		<div>La Ce</div>	<div>Ac Th</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = -3$				
		<div>Pr Nd</div>	<div>Pa U</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = -2$					
		<div>Pm Sm</div>	<div>Np Pu</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = -1$					
		<div>Eu Gd</div>	<div>Am Cm</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 0$					
		<div>Tb Dy</div>	<div>Bk Cf</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 1$					
		<div>Ho Er</div>	<div>Es Fm</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 2$					
		<div>Tm Yb</div>	<div>Md No</div> $\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu = 3$					

Fig. 1. Mendeleev table in the form of the basic representation $F_{ss'}^+$ of the Rumer-Fet group $\text{SO}(2, 4) \otimes \text{SU}(2) \otimes \text{SU}(2)'$.

The reduction of the basic representation $F_{ss'}^+$ of the Rumer-Fet group on the subgroups is realized in accordance with the chain (9). So, multiplets of a subgroup $\text{SO}(3) \otimes \text{SU}(2)$ are represented by vertical rectangles of Fig. 1, and their elements compose well-known s -, p -, d - and f -families (in particular, lanthanides and actinides are selected as multiplets of the subgroup $\text{SO}(3) \otimes \text{SU}(2)$). The each element occupies a quite definite place, which is defined by its “address” in the table

$(\nu, \lambda, \mu, s, s')$, that is, by corresponding quantum numbers of the symmetry group. Thus, atoms of all possible elements stand in one-to-one correspondence with the vectors of the basis (8).

4 Mendeleev Table

In Fig. 1 we mark by bold script the elements that had not yet been discovered or received their official names during lifetime of Rumer and Fet. These elements belong the last column of the Fig. 1 with the quantum numbers $\nu = 4$ and $s' = 1/2$: **Db** – Dubnium, **Sg** – Seaborgium, **Bh** – Bohrium, **Hs** – Hassium (eka-osmium), **Mt** – Meitnerium, **Ds** – Darmstadtium, **Rg** – Roentgenium, **Cn** – Copernicium (eka-mercury). All these elements belong to a multiplet with a quantum number $\lambda = 2$. A multiplet with $\lambda = 1$ ($\nu = 4, s' = 1/2$) consists of recently discovered elements: **Nh** – Nihonium (eka-thallium), **Fl** – Flerovium (eka-lead), **Mc** – Moscovium (eka-bismuth), **Lv** – Livermorium (eka-polonium), **Ts** – Tennessine (eka-astatine), **Og** – Oganesson (eka-radon). Further, a multiplet with a quantum number $\lambda = 0$ ($\nu = 4, s' = 1/2$) is formed by undetected yet hypothetical elements **Uue** – Ununennium (eka-francium) with supposed atomic mass 316 a.u. and **Ubn** – Unbinillium (eka-radium). All the enumerated elements accomplish the filling of the Mendeleev table (from 1-th to 120-th number) inclusive the value of quantum number $\nu = 4$.

Tab. 1. Average masses of “heavy” multiplets.

	Multiplet	Mass (exp.)	Mass (theor.)	Approx. %
1.	$(\nu = 3, s' = -1/2, \lambda = 2)$	55, 31	53	−4, 36
2.	$(\nu = 3, s' = -1/2, \lambda = 1)$	76, 65	75	−2, 2
3.	$(\nu = 3, s' = -1/2, \lambda = 0)$	86, 54	86	−0, 63
4.	$(\nu = 3, s' = 1/2, \lambda = 2)$	99, 76	104	4, 07
5.	$(\nu = 3, s' = 1/2, \lambda = 1)$	123, 51	126	1, 98
6.	$(\nu = 3, s' = 1/2, \lambda = 0)$	135, 12	137	1, 37
7.	$(\nu = 4, s' = -1/2, \lambda = 3)$	154, 59	156	0, 90
8.	$(\nu = 4, s' = -1/2, \lambda = 2)$	187, 96	189	0, 55
9.	$(\nu = 4, s' = -1/2, \lambda = 1)$	210, 21	211	0, 37
10.	$(\nu = 4, s' = -1/2, \lambda = 0)$	224, 52	222	−1, 13
11.	$(\nu = 4, s' = 1/2, \lambda = 3)$	244, 56	241	−1, 48
12.	$(\nu = 4, s' = 1/2, \lambda = 2)$	273, 10	274	0, 33
13.	$(\nu = 4, s' = 1/2, \lambda = 1)$	290, 83	296	1, 75
14.	$(\nu = 4, s' = 1/2, \lambda = 0)$	318*	307	−3, 58*

Let us calculate masses of the hypothetical elements **Uue** and **Ubn**. Before to proceed this, we calculate average masses of the multiplets belonging to the Mendeleev table. With this aim in view we use a mass formula⁹ proposed in [18]:

$$m = m_0 + a \left[s'(2\nu - 3) - 5\nu + \frac{11}{2} + 2(\nu^2 - 1) \right] - b \cdot \lambda(\lambda + 1), \quad (10)$$

where m_0 , a , b are the coefficients which are underivable from the theory. The formula (10) is analogous to the “first perturbation” in SU(3)- and SU(6)-theories which allows to calculate an average mass of the elements of the multiplet (an analogue of the “second perturbation” for the Rumer-Fet group, which leads to the mass splitting inside the multiplet, we will give

⁹This formula is analogous to a Gell-Mann–Okubo formula for the hadrons in SU(3)-theory [22, 23], and also to a Bég-Singh formula in SU(6)-theory [24].

in section 6). The Tab.1 contains average masses of “heavy” multiplets ($\nu = 3, 4$) calculated according the formula (10) at $m_0 = 1$, $a = 17$, $b = 5, 5$. From the Tab.1 we see that an accuracy between experimental and theoretical masses rises with growth of the “weight” of the multiplet, therefore, the formula (10) is asymptotic. An exception is the last multiplet ($\nu = 4, s' = 1/2, \lambda = 0$), consisting of the hypothetical elements **Uue** and **Ubn**, which have masses unconfirmed by experiment.

5 Implementation of the Operator Algebra

As is known, in the foundation of an algebraic formulation of quantum theory we have a Gelfand-Naimark-Segal (GNS) construction, which is defined by a canonical correspondence $\omega \leftrightarrow \pi_\omega$ between states and cyclic representations of C^* -algebra [25, 26, 27].

Let us suppose that according to the axiom **A.I** (section 2) generators of the conformal group (a fundamental symmetry $G_f = \text{SO}(2, 4)$ in this context) are attached to the energy operator H . Therefore, the each eigensubspace \mathbf{H}_E of the energy operator is invariant with respect to the operators of the representation F^+ of the conformal group¹⁰. It allows us to obtain a concrete implementation (“dressing”) of the operator algebra $\pi(\mathfrak{A}) \rightarrow \pi(H)$, where $\pi \equiv F^+$. Thus, the each possible value of energy (an energy level) is a vector state of the form (axiom **A.II**):

$$\omega_\Phi(H) = \frac{\langle \Phi | \pi(H)\Phi \rangle}{\langle \Phi | \Phi \rangle} = \frac{\langle \Phi | F^+(H)\Phi \rangle}{\langle \Phi | \Phi \rangle},$$

where $|\Phi\rangle$ is a cyclic vector of the Hilbert space \mathbf{H}_∞ .

Further, in virtue of an isomorphism $\text{SU}(2, 2) \simeq \mathbf{Spin}_+(2, 4)$ (see Appendix C) we will consider an universal covering \tilde{G}_f as a *spinor group*. It allows us to associate in addition a *twistor structure* with the each cyclic vector $|\Phi\rangle \in \mathbf{H}_\infty$. Spintensor representations of the group $\tilde{G}_f = \mathbf{Spin}_+(2, 4)$ form a substrate of finite-dimensional representations $\boldsymbol{\tau}_{k/2, r/2}$, $\overline{\boldsymbol{\tau}}_{k/2, r/2}$ of the conformal group realized in the spaces $\text{Sym}_{(k, r)} \subset \mathbb{S}_{2k+r}$ and $\overline{\text{Sym}}_{(k, r)} \subset \overline{\mathbb{S}}_{2k+r}$, where \mathbb{S}_{2k+r} is a spinspace. Indeed, a twistor $\mathbf{Z}^\alpha = (s^\alpha, s_{\dot{\alpha}})$ is a vector of the fundamental representation of the group $\mathbf{Spin}_+(2, 4)$, where $\alpha, \dot{\alpha} = 0, 1$. A vector of the general spintensor representation of the group $\mathbf{Spin}_+(2, 4)$ is

$$\mathbf{Z} = \begin{bmatrix} \mathbf{S} \\ \overline{\mathbf{S}} \end{bmatrix}, \quad (11)$$

where \mathbf{S} is a spintensor of the form

$$\mathbf{S} = s_{\dot{\alpha}_1 \dot{\alpha}_2 \dots \dot{\alpha}_r}^{\alpha_1 \alpha_2 \dots \alpha_k} = \sum s^{\alpha_1} \otimes s^{\alpha_2} \otimes \dots \otimes s^{\alpha_k} \otimes s_{\dot{\alpha}_1} \otimes s_{\dot{\alpha}_2} \otimes \dots \otimes s_{\dot{\alpha}_r}, \quad \alpha_i, \dot{\alpha}_i = 0, 1; \quad (12)$$

that is, the vector of the spinspace $\mathbb{S}_{2k+r} = \mathbb{S}_{2k} \otimes \dot{\mathbb{S}}_{2r}$, where $\dot{\mathbb{S}}_{2r}$ is a dual spinspace. $\overline{\mathbf{S}}$ is a spintensor from the conjugated spinspace $\overline{\mathbb{S}}_{2k+r}$. Symmetrizing the each spintensor \mathbf{S} and $\overline{\mathbf{S}}$ in (11), we obtain symmetric *twisttensor* \mathbf{Z} . In turn, as is known [28], spinspace is a minimal left ideal of the Clifford algebra $\mathcal{C}_{p, q}$, that is, there is an isomorphism $\mathbb{S}_{2^m}(\mathbb{K}) \simeq I_{p, q} = \mathcal{C}_{p, q} f$, where f is a primitive idempotent of the algebra $\mathcal{C}_{p, q}$, $\mathbb{K} = f \mathcal{C}_{p, q} f$ is a division ring for $\mathcal{C}_{p, q}$, $m = (p + q)/2$. A complex spinspace $\mathbb{S}_{2^m}(\mathbb{C})$ is a complexification $\mathbb{C} \otimes I_{p, q}$ of the minimal left ideal $I_{p, q}$ of a real subalgebra $\mathcal{C}_{p, q}$. Hence, \mathbb{S}_{2k+r} is a minimal left ideal of the complex algebra $\mathbb{C}_{2k} \otimes \dot{\mathbb{C}}_{2r} \simeq \mathbb{C}_{2(k+r)}$ (for more details see [29, 30]).

¹⁰It follows from a similarity of the complex shells of the group algebras $\mathfrak{sl}(2, \mathbb{C})$, $\mathfrak{so}(4)$ and $\mathfrak{so}(2, 4)$.

Now we are in a position that allows us to define a system of *basic cyclic vectors* endowed with a complex twistor structure (these vectors correspond to the system of finite-dimensional representations of the conformal group). Let

$$\begin{aligned}
& | \mathbb{C}_0, \tau_{0,0}(H)\Phi \rangle; \\
& | \mathbb{C}_2, \tau_{1/2,0}(H)\Phi \rangle, \quad | \dot{\mathbb{C}}_2, \tau_{0,1/2}(H)\Phi \rangle; \\
& | \mathbb{C}_2 \otimes \mathbb{C}_2, \tau_{1,0}(H)\Phi \rangle, \quad | \mathbb{C}_2 \otimes \dot{\mathbb{C}}_2, \tau_{1/2,1/2}(H)\Phi \rangle, \quad | \dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2, \tau_{0,1}(H)\Phi \rangle; \\
& | \mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \mathbb{C}_2, \tau_{3/2,0}(H)\Phi \rangle, \quad | \mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \dot{\mathbb{C}}_2, \tau_{1,1/2}(H)\Phi \rangle, \\
& \quad | \mathbb{C}_2 \otimes \dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2, \tau_{1/2,1}(H)\Phi \rangle, \quad | \dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2, \tau_{0,3/2}(H)\Phi \rangle; \\
& \dots\dots\dots \\
& | \overline{\mathbb{C}_0, \tau_{0,0}(H)\Phi} \rangle; \\
& | \overline{\mathbb{C}_2, \tau_{1/2,0}(H)\Phi} \rangle, \quad | \overline{\dot{\mathbb{C}}_2, \tau_{0,1/2}(H)\Phi} \rangle; \\
& | \overline{\mathbb{C}_2 \otimes \mathbb{C}_2, \tau_{1,0}(H)\Phi} \rangle, \quad | \overline{\mathbb{C}_2 \otimes \dot{\mathbb{C}}_2, \tau_{1/2,1/2}(H)\Phi} \rangle, \quad | \overline{\dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2, \tau_{0,1}(H)\Phi} \rangle; \\
& | \overline{\mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \mathbb{C}_2, \tau_{3/2,0}(H)\Phi} \rangle, \quad | \overline{\mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \dot{\mathbb{C}}_2, \tau_{1,1/2}(H)\Phi} \rangle, \\
& \quad | \overline{\mathbb{C}_2 \otimes \dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2, \tau_{1/2,1}(H)\Phi} \rangle, \quad | \overline{\dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2 \otimes \dot{\mathbb{C}}_2, \tau_{0,3/2}(H)\Phi} \rangle; \\
& \dots\dots\dots
\end{aligned}$$

Therefore, in accordance with GNS-construction (axiom **A.II**) we have complex vector states of the form

$$\begin{aligned}
\omega_{\Phi}^c(H) &= \frac{\langle \Phi | \mathbb{C}_{2(k+r)}, \tau_{k/2,r/2}(H)\Phi \rangle}{\langle \Phi | \Phi \rangle}, \\
\bar{\omega}_{\Phi}^c(H) &= \frac{\langle \Phi | \overline{\mathbb{C}_{2(k+r)}, \tau_{k/2,r/2}(H)\Phi} \rangle}{\langle \Phi | \Phi \rangle}.
\end{aligned}$$

According to (11), the pairs $(\omega_{\Phi}^c(H), \bar{\omega}_{\Phi}^c(H))$ form *neutral states*. Further, at execution of the condition $\omega_{\Phi}^c(H) \geq 0$ ($\bar{\omega}_{\Phi}^c(H) \geq 0$) a set of all pure states $(\omega_{\Phi}^c(H), \bar{\omega}_{\Phi}^c(H))$ forms a physical Hilbert space \mathbf{H}_{phys} (axiom **A.III**) and, correspondingly, a space of rays $\hat{H} = \mathbf{H}_{\text{phys}}/S^1$. All the pure states of the physical quantum system \mathbf{U} are described by the unit rays and at this realization of the operator algebra these states correspond to atoms of the periodic system of elements. At this point, there is superposition principle (axiom **A.V**).

Following to Heisenberg's classification [31], the all set of symmetry groups G should be divided on the two classes: 1) *groups of fundamental (primary) symmetries* G_f , which participate in construction of state vectors of the quantum system \mathbf{U} ; 2) *groups of dynamic (secondary) symmetries* G_d , which describe approximate symmetries between state vectors of \mathbf{U} . Dynamic symmetries G_d relate different states (state vectors $|\Psi\rangle \in \mathbf{H}_{\text{phys}}$) among the quantum system \mathbf{U} . The symmetry G_d of the system \mathbf{U} can be represented as a *quantum transition* between its states (levels of state spectrum of \mathbf{U}).

We now show that Rumer-Fet group is a dynamic symmetry. Indeed, the group (7) is equivalent to $\widetilde{\text{SO}}(2, 4) \otimes \text{SU}(2) = \text{SU}(2, 2) \otimes \text{SU}(2)$ (see [32]), since one “doubling” in (7) already actually described by the two-sheeted covering $\text{SU}(2, 2)$ of the conformal group¹¹. At this point, atoms of different elements stand in one-to-one correspondence with the vectors belonging the basis

¹¹Throughout the article, the term “doubling” occurs many times. “Doubling” (or Pauli’s “doubling and decreasing symmetry” [31]) is one of the leading principles of group-theoretic description. Heisenberg notes [33] that all the real symmetries of nature arose as a consequence of such doubling. “Symmetry decreasing” should be understood as group reduction, that is, if there is a chain of nested groups $G \supset G_1 \supset G_2 \supset \dots \supset G_k$ and an irreducible

(8) of the space of the representation $F_{ss'}^+$. Here we have a direct analogue with the physics of “elementary particles”. According to [36], a quantum system, described by an irreducible unitary *representation* of the Poincaré group \mathcal{P} , is called elementary particle. On the other hand, in accordance with SU(3)- and SU(6)-theories elementary particle is described by a *vector* of an irreducible representation of the group SU(3) (or SU(6)). Therefore, we have two group theoretical interpretations of the elementary particle: as a *representation* of the group \mathcal{P} (group of fundamental symmetry) and as a *vector* of the representation of the group of dynamic symmetry SU(3) (or SU(6)). Besides, the structure of the mass formula (10) for the Rumer-Fet group is analogous to Gell-Mann-Okubo and Bég-Singh mass formulas for the groups SU(3) and SU(6). An action of the group $G_d = \text{SU}(2, 2) \otimes \text{SU}(2)$, which lifted into \mathbf{H}_{phys} via a central extension (see, for example, [11]), moves state vectors $|\Psi\rangle \in \mathbf{H}_{\text{phys}}$, corresponding to different atoms of the periodic system, into each other.

6 Seaborg Table

As is known, Mendeleev table includes 118 elements, from which at present time 118 elements have been discovered (the last detected element **Og** – Oganesson (eka-radon) with the atomic number $Z = 118$). Two as yet undiscovered hypothetical elements **Uue** – Ununennium (eka-francium) with $Z = 119$ and **Ubn** – Unbinillium (eka-radium) with $Z = 120$ begin to fill the eight period. According to Bohr model, both elements belong to s -shell. Mendeleev table contains seven periods (rows), including s -, p -, d - and f -families (shells). The next (eight) period involves the construction of the g -shell. In 1969, Glenn Seaborg [38] proposed an eight-periodic table containing g -shell. The first element of g -shell is **Ubu** (Unbiunium) with the atomic number $Z = 121$ (superactinide group starts also with this element). The full number of elements of Seaborg table is equal to 218.

No one knows how many elements can be in the periodic system. The Reserford-Bohr structural model leads to the following restriction (so-called “Bohr model breakdown”) on the number of physically possible elements. So, for elements with atomic number greater than 137 a “speed” of an electron in $1s$ orbital is given by

$$v = Z\alpha c \approx \frac{Zc}{137,036},$$

where α is the fine structure constant. Under this approximation, any element with $Z > 137$ would require $1s$ electrons to be travelling faster than c . On the other hand, Feynman pointed out that a relativistic Dirac equation leads also into problems with $Z > 137$, since a ground state energy for the electron on $1s$ -subshell is given by an expression $E = m_0 c^2 \sqrt{1 - Z^2 \alpha^2}$, where m_0 is the rest mass of the electron. In the case of $Z > 137$, an energy value becomes an imaginary number, and, therefore, the wave function of the ground state is oscillatory, that is, there is no gap between the positive and negative energy spectra, as in the Klein paradox. For that reason 137-th element **Uts** (Untriseptium) was proclaimed as the “end” of the periodic system, and also in honour of Feynman this element was called Feynmanium (symbol: **Fy**). As is known, Feynman derived this result with the assumption that the atomic nucleus is point-like.

unitary representation \mathfrak{P} of group G in space \mathbf{H}_{phys} is given, then the reduction G/G_1 of the representation \mathfrak{P} of group G by subgroup G_1 leads to the decomposition of \mathfrak{P} into orthogonal sum of irreducible representations $\mathfrak{P}_i^{(1)}$ of subgroup G_1 . In turn, the reduction G_1/G_2 of the representation of the group G_1 over the subgroup G_2 leads to the decomposition of the representations $\mathfrak{P}_i^{(1)}$ into irreducible representations $\mathfrak{P}_{ij}^{(2)}$ of the group G_2 and so on (see [34, 35]). Thus there is a reduction (“symmetry decreasing” of Pauli) of the group G with high symmetry to lower symmetries of the subgroups.

	$\nu=1$ $s'=-1/2$ $s'=1/2$		$\nu=2$ $s'=-1/2$ $s'=1/2$		$\nu=3$ $s'=-1/2$ $s'=1/2$		$\nu=4$ $s'=-1/2$ $s'=1/2$		$\nu=5$ $s'=-1/2$ $s'=1/2$		
$\lambda = 0$	<div>H He</div>	<div>Li Be</div>	<div>Na Mg</div>	<div>K Ca</div>	<div>Rb Sr</div>	<div>Cs Ba</div>	<div>Fr Ra</div>	<div>Uue Ubn</div>	<div>Uhe Usn</div>	<div>Bue Bbn</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=0$
$\lambda = 1$	{		<div>B C</div>	<div>Al Si</div>	<div>Ga Ge</div>	<div>In Sn</div>	<div>Tl Pb</div>	<div>Nh Fl</div>	<div>Uht Uhq</div>	<div>But Buq</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-1$
			<div>N O</div>	<div>P S</div>	<div>As Se</div>	<div>Sb Te</div>	<div>Bi Po</div>	<div>Mc Lv</div>	<div>Uhp Uhn</div>	<div>Bup Buh</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=0$
			<div>F Ne</div>	<div>Cl Ar</div>	<div>Br Kr</div>	<div>I Xe</div>	<div>At Rn</div>	<div>Ts Og</div>	<div>Uhs Uho</div>	<div>Bus Buo</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=1$
$\lambda = 2$	{				<div>Sc Ti</div>	<div>Y Zr</div>	<div>Lu Hf</div>	<div>Lr Rf</div>	<div>Upt Upq</div>	<div>Bnt Bnq</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-2$
					<div>V Cr</div>	<div>Nb Mo</div>	<div>Ta W</div>	<div>Db Sg</div>	<div>Upp Uph</div>	<div>Bnp Bnh</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-1$
					<div>Mn Fe</div>	<div>Tc Ru</div>	<div>Re Os</div>	<div>Bh Hs</div>	<div>Ups Upo</div>	<div>Bns Bno</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=0$
					<div>Co Ni</div>	<div>Rh Pd</div>	<div>Ir Pt</div>	<div>Mt Ds</div>	<div>Upe Uhn</div>	<div>Bne Bun</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=1$
					<div>Cu Zn</div>	<div>Ag Cd</div>	<div>Au Hg</div>	<div>Rg Cn</div>	<div>Uhu Uhb</div>	<div>Buu Bub</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=2$
$\lambda = 3$	{						<div>La Ce</div>	<div>Ac Th</div>	<div>Ute Uqn</div>	<div>Uoe Uen</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-3$
							<div>Pr Nd</div>	<div>Pa U</div>	<div>Uqu Uqb</div>	<div>Ueu Ueb</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-2$
							<div>Pm Sm</div>	<div>Np Pu</div>	<div>Uqt Uqq</div>	<div>Uet Ueq</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-1$
							<div>Eu Gd</div>	<div>Am Cm</div>	<div>Uqp Uqh</div>	<div>Uep Ueh</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=0$
							<div>Tb Dy</div>	<div>Bk Cf</div>	<div>Uqs Uqo</div>	<div>Ues Ueo</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=1$
							<div>Ho Er</div>	<div>Es Fm</div>	<div>Uqe Upn</div>	<div>Uee Bnn</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=2$
							<div>Tm Yb</div>	<div>Md No</div>	<div>Upu Upb</div>	<div>Bnu Bnb</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=3$
	M										
$\lambda = 4$	{								<div>Ubu Ubb</div>	<div>Usu Usb</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-4$
									<div>Ubt Ubq</div>	<div>Ust Usq</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-3$
									<div>Ubp Ubn</div>	<div>Usp Ush</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-2$
									<div>Ubs Ubo</div>	<div>Uss Uso</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=-1$
									<div>Ube Utn</div>	<div>Use Uon</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=0$
									<div>Utu Utb</div>	<div>Uou Uob</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=1$
									<div>Utt Utq</div>	<div>Uot Uoq</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=2$
									<div>Utp Uth</div>	<div>Uop Uoh</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=3$
									<div>Uts Uto</div>	<div>Uos Uoo</div>	$\left. \begin{matrix} s=-1/2 \\ s=1/2 \end{matrix} \right\} \mu=4$

Fig. 2. Seaborg table in the form of the basic representation $F_{ss'}^+$ of the Rumer-Fet group (basis $|\nu, \lambda, \mu, s, s'\rangle$).

Further, Greiner-Reinhardt solution [39], representing the atomic nucleus by a charged ball of the radius $R = 1, 2A^{1/3}fm$, where A is the atomic mass, moves aside the Feynman limit to the value $Z = 173$. For $Z \approx 173$ under action of the electric field of the nucleus $1s$ -subshell “dives” into the negative continuum (Dirac sea), that leads to spontaneous emission of electron-positron pairs and, as a consequence, to the absence of neutral atoms above the element **Ust** (Unsepttrium)

with $Z = 173$. Atoms with $Z > Z_{cr} \approx 173$ are called *supercritical* atoms. It is supposed that elements with $Z > Z_{cr}$ could only exist as ions.

As shown earlier, the Seaborg table is an eight-periodic extension of the Mendeleev table (from 119-th to the 218-th element). The Seaborg table contains both “critical” elements of the Bohr model: **Uts** (Untriseptium, $Z = 137$) and **Ust** (Unsepttrium, $Z = 173$). According to Bohr model, the filling of the g -shell (formation of g -family) begins with the 121-th element. In the Rumer-Fet model [18] g -shell corresponds to quantum numbers $\nu = 5$ and $\lambda = 4$ of the symmetry group $SO(2, 4) \otimes SU(2) \otimes SU(2)'$. The Seaborg table is presented on the Fig. 2 in the form of the basic representation $F_{ss'}^+$ of the Rumer-Fet group. The Mendeleev table (as part of the Seaborg table) is highlighted by a dotted border. Within eight-periodic extension (quantum numbers $\nu = 5$, $\lambda = 4$) in addition to 20 multiplets of the Mendeleev table we have 10 multiplets.

Let us calculate average masses of these multiplets. With this aim in view we use the mass formula (10). The formula (10) corresponds to the chain of groups (9), according to which we have a reduction of the basic representation $F_{ss'}^+$ on the subgroups of this chain, that is, a partition of the basic multiplet into the lesser multiplets. As noted above, the formula (10) is analogous to the “first perturbation” in $SU(3)$ - and $SU(6)$ -theories, which allows us to calculate an average mass of the elements belonging to a given multiplet¹². At $m_0 = 1$, $a = 17$, $b = 5, 5$ from (10) we obtain the average masses of multiplets (see Tab. 2).

Tab. 2. Average masses of multiplets of Seaborg table.

	Multiplet	Mass (theor.)
1.	$(\nu = 5, s' = -1/2, \lambda = 4)$	316
2.	$(\nu = 5, s' = -1/2, \lambda = 3)$	360
3.	$(\nu = 5, s' = -1/2, \lambda = 2)$	393
4.	$(\nu = 5, s' = -1/2, \lambda = 1)$	415
5.	$(\nu = 5, s' = -1/2, \lambda = 0)$	426
6.	$(\nu = 5, s' = 1/2, \lambda = 4)$	435
7.	$(\nu = 5, s' = 1/2, \lambda = 3)$	479
8.	$(\nu = 5, s' = 1/2, \lambda = 2)$	512
9.	$(\nu = 5, s' = 1/2, \lambda = 1)$	534
10.	$(\nu = 5, s' = 1/2, \lambda = 0)$	545

With the aim to obtain an analogue of the “second perturbation”, which leads to a mass splitting inside the multiplets of the group $G_2 = SO(3) \otimes SU(2)$, it needs to find a subsequent lengthening of the group chain $G \supset G_1 \supset G_2$ (9). Therefore, we need to find another subgroup G_3 . Then G_2/G_3 -reduction gives a termwise mass splitting. As is known, a representation $\{u_O\}$ of the group $SU(2)$ compares the each rotation O from $SO(3)$ with the matrix u_O from $SU(2)$ and thereby the pair (O, u_O) , that is, the element of G_2 . At the multiplication (O, u_O) give the pairs of the same form: $(O_1, u_{O_1})(O_2, u_{O_2}) = (O_1 O_2, u_{O_1} u_{O_2}) = (O_1 O_2, u_{O_1 O_2})$, reverse pairs are analogous: $(O, u_O)^{-1} = (O^{-1}, u_{O^{-1}})$. Therefore, such pairs form a subgroup G_3 in G_2 . The subgroup G_3 is

¹²So, in $SU(3)$ -theory we have a Gell-Mann–Okubo mass formula

$$m = m_0 + \alpha + \beta Y + \gamma \left[I(I+1) - \frac{1}{4} Y^2 \right] + \alpha' - \beta' Q + \gamma' \left[U(U+1) - \frac{1}{4} Q^2 \right],$$

in which, according to $SU(3)/SU(2)$ -reduction, quantum numbers (isospin I , hypercharge Y), standing in the first square bracket, define the “first perturbation” that leads to a so-called *hypercharge* mass splitting, that is, a partition of the multiplet of $SU(3)$ into the lesser multiplets of the subgroup $SU(2)$. A “second perturbation” is defined by the quantum numbers, standing in the second square bracket (charge Q and isospin U which in difference from I corresponds to other choice of the basis in the subgroup $SU(2)$), that leads to a *charge* mass splitting inside the multiplets of $SU(2)$.

locally isomorphic to $\text{SO}(3)$. Following to Fet, we will denote it via $\text{SO}(3)_c$. Further, one-parameter subgroups of the group $\text{SO}(3)$ have the form $\{e^{-i\alpha A_k}\}$ ($k = 1, 2, 3$); since the representation $\{u_O\}$ converts them into one-parameter subgroups $\{e^{-i\alpha \tau_k}\}$ of the group $\text{SU}(2)$, then corresponding one-parameter subgroups in $\text{SO}(3)_c$ have the form

$$(e^{-i\alpha A_k}, e^{-i\alpha \tau_k}) = (e^{-i\alpha A_k}, 1) (1, e^{-i\alpha \tau_k}).$$

Since the matrices A_1, A_2, A_3 correspond to rotations $\mathbf{L}_{23} = \mathbf{J}_1 + \mathbf{K}_1$, $\mathbf{L}_{31} = \mathbf{J}_2 + \mathbf{K}_2$, $\mathbf{L}_{12} = \mathbf{J}_3 + \mathbf{K}_3$ in the group G (in the basic representation of the group G , see section 3), the pair $(e^{-i\alpha A_k}, 1)$ is represented by the operator $e^{-i\alpha(\mathbf{J}_k + \mathbf{K}_k)}$; the pair $(1, e^{-i\alpha \tau_k})$ is represented by $e^{-i\alpha \tau_k}$. Thus, one-parameter subgroups of $G_3 = \text{SO}(3)_c$ correspond to subgroups of operators $e^{-i\alpha(\mathbf{J}_k + \mathbf{K}_k)} e^{-i\alpha \tau_k}$ ($k = 1, 2, 3$). Irreducible representations of the group $G_2 = \text{SO}(3) \otimes \text{SU}(2)$ are numbered by the collections of quantum numbers (ν, s', λ) . These representations form vertical rectangles in Fig. 2. Each of them is defined by a fundamental representation of $\text{SU}(2)$ and $(2\lambda + 1)$ -dimensional irreducible representation of the group $\text{SO}(3)$. At G_2/G_3 -reduction from such representation we obtain an irreducible representation of the subgroup $G_3 = \text{SO}(3)_c$, for which a Clebsch-Gordan sequence $|j_1 - j_2|, \dots, j_1 + j_2$ with the values $j_1 = 1/2$ and $j_2 = \lambda$ is reduced to two terms $\lambda - 1/2$, $\lambda + 1/2$ at $\lambda > 0$ and to one term $1/2$ at $\lambda = 0$. Therefore, at $\lambda > 0$ the representation (ν, s', λ) of the group G_2 is reduced to two irreducible representations of the subgroup G_3 with dimensionality $2(\lambda - 1/2) + 1 = 2\lambda$ and $2(\lambda + 1/2) + 1 = 2\lambda + 2$, and at $\lambda = 0$ to one two-dimensional irreducible representation. Thus, at $\lambda > 0$ multiplets of the subgroup G_2 are reduced to two multiplets of the subgroup G_3 . G_2/G_3 -reduction leads to the following (lengthened) chain of groups:

$$G \supset G_1 \supset G_2 \supset G_3 \longmapsto \text{SO}(2, 4) \otimes \text{SU}(2) \otimes \text{SU}(2) \supset \text{SO}(4) \otimes \text{SU}(2) \supset \text{SO}(3) \otimes \text{SU}(2) \supset \text{SO}(3)_c, \quad (13)$$

The lengthening of the group chain requires the introduction of a new basis whose vectors belong to the smallest multiplets of symmetry, that is, multiplets of the subgroup G_3 . The vectors $|\nu, \lambda, \mu, s, s'\rangle$ of the basis (8), corresponding to the group chain (9), do not compose already a chosen (well-defined) basis, since μ, s do not belong to quantum numbers of the symmetry group, that is, these vectors do not belong to irreducible spaces of the group G_3 . The new basis is defined as follows. Since ν, s', λ are related with the groups G, G_1, G_2 , they remain quantum numbers of the chain (9), and instead μ, s we have new quantum numbers related with G_3 . First, quantum number ι_λ relates with the Casimir operator of the subgroup G_3 , which equals to $\sum_{k=1}^3 (\tau_k + \mathbf{J}_k + \mathbf{K}_k)^2$. At this point, two multiplets of G_3 , obtained at G_2/G_3 -reduction, correspond to $\iota_\lambda = \lambda - 1/2$ and $\iota_\lambda = \lambda + 1/2$, whence $2\lambda = 2\iota_\lambda + 1$, $2\lambda + 2 = 2\iota_\lambda + 1$. Other quantum number κ is an eigenvalue of the operator $q_3 = \tau_3 + \mathbf{J}_3 + \mathbf{K}_3$, which belongs to the Lie algebra of the group $G_3 = \text{SO}(3)_c$. Thus, the new basis, corresponding to the group chain (13), has the form

$$|\nu, s', \lambda, \iota_\lambda, \kappa\rangle, \quad \nu = 1, 2, \dots; \quad s' = -1/2, 1/2 \quad \lambda = 0, 1, \dots, \nu - 1; \\ \iota_\lambda = \lambda - 1/2, \lambda + 1/2 \quad \kappa = -\iota_\lambda, -\iota_\lambda + 1, \dots, \iota_\lambda - 1, \iota_\lambda. \quad (14)$$

The Seaborg table recorded in basis (14), is shown in Fig. 3.

		$\nu=1$ $s'=-1/2$ $s'=1/2$		$\nu=2$ $s'=-1/2$ $s'=1/2$		$\nu=3$ $s'=-1/2$ $s'=1/2$		$\nu=4$ $s'=-1/2$ $s'=1/2$		$\nu=5$ $s'=-1/2$ $s'=1/2$		
$\lambda = 0$		H He	Li Be	Na Mg	K Ca	Rb Sr	Cs Ba	Fr Ra	Uue Ubn	Uhe Usn	Bue Bbn	$\left. \begin{matrix} \kappa=-1/2 \\ \kappa=1/2 \end{matrix} \right\} \iota_\lambda=1/2$
$\lambda = 1$	{			B C	Al Si	Ga Ge	In Sn	Tl Pb	Nh Fl	Uht Uhq	But Buq	$\left. \begin{matrix} \kappa=-1/2 \\ \kappa=1/2 \end{matrix} \right\} \iota_\lambda=1/2$
				N O	P S	As Se	Sb Te	Bi Po	Mc Lv	Uhp Uhn	Bup Buh	$\left. \begin{matrix} \kappa=-3/2 \\ \kappa=-1/2 \end{matrix} \right\} \iota_\lambda=3/2$
				F Ne	Cl Ar	Br Kr	I Xe	At Rn	Ts Og	Uhs Uho	Bus Buo	$\left. \begin{matrix} \kappa=1/2 \\ \kappa=3/2 \end{matrix} \right\}$
$\lambda = 2$	{					Sc Ti	Y Zr	Lu Hf	Lr Rf	Upt Upq	Bnt Bnq	$\left. \begin{matrix} \kappa=-3/2 \\ \kappa=-1/2 \end{matrix} \right\} \iota_\lambda=3/2$
						V Cr	Nb Mo	Ta W	Db Sg	Upp Uph	Bnp Bnh	$\left. \begin{matrix} \kappa=1/2 \\ \kappa=3/2 \end{matrix} \right\}$
						Mn Fe	Tc Ru	Re Os	Bh Hs	Ups Upo	Bns Bno	$\left. \begin{matrix} \kappa=-5/2 \\ \kappa=-3/2 \end{matrix} \right\}$
						Co Ni	Rh Pd	Ir Pt	Mt Ds	Upe Uhn	Bne Bun	$\left. \begin{matrix} \kappa=-1/2 \\ \kappa=1/2 \end{matrix} \right\} \iota_\lambda=5/2$
						Cu Zn	Ag Cd	Au Hg	Rg Cn	Uhu Uhb	Buu Bub	$\left. \begin{matrix} \kappa=3/2 \\ \kappa=5/2 \end{matrix} \right\}$
$\lambda = 3$	{							La Ce	Ac Th	Ute Uqn	Uoe Uen	$\left. \begin{matrix} \kappa=-5/2 \\ \kappa=-3/2 \end{matrix} \right\} \iota_\lambda=5/2$
								Pr Nd	Pa U	Uqu Uqb	Ueu Ueb	$\left. \begin{matrix} \kappa=-1/2 \\ \kappa=1/2 \end{matrix} \right\}$
								Pm Sm	Np Pu	Uqt Uqq	Uet Ueq	$\left. \begin{matrix} \kappa=3/2 \\ \kappa=5/2 \end{matrix} \right\}$
								Eu Gd	Am Cm	Uqp Uqh	Uep Ueh	$\left. \begin{matrix} \kappa=-7/2 \\ \kappa=-5/2 \end{matrix} \right\}$
								Tb Dy	Bk Cf	Uqs Uqo	Ues Ueo	$\left. \begin{matrix} \kappa=-3/2 \\ \kappa=-1/2 \end{matrix} \right\} \iota_\lambda=7/2$
								Ho Er	Es Fm	Uqe Upn	Uee Bnn	$\left. \begin{matrix} \kappa=1/2 \\ \kappa=3/2 \end{matrix} \right\}$
								Tm Yb	Md No	Upu Upb	Bnu Bnb	$\left. \begin{matrix} \kappa=5/2 \\ \kappa=7/2 \end{matrix} \right\}$
M												
$\lambda = 4$	{									Ubu Ubb	Usu Usb	$\left. \begin{matrix} \kappa=-7/2 \\ \kappa=-5/2 \end{matrix} \right\}$
										Ubt Ubq	Ust Usq	$\left. \begin{matrix} \kappa=-3/2 \\ \kappa=-1/2 \end{matrix} \right\} \iota_\lambda=7/2$
										Ubp Ubn	Usp Ush	$\left. \begin{matrix} \kappa=1/2 \\ \kappa=3/2 \end{matrix} \right\}$
										Ubs Ubo	Uss Uso	$\left. \begin{matrix} \kappa=5/2 \\ \kappa=7/2 \end{matrix} \right\}$
										Ube Utn	Use Uon	$\left. \begin{matrix} \kappa=-9/2 \\ \kappa=-7/2 \end{matrix} \right\}$
										Utu Utb	Uou Uob	$\left. \begin{matrix} \kappa=-5/2 \\ \kappa=-3/2 \end{matrix} \right\}$
										Utt Utq	Uot Uoq	$\left. \begin{matrix} \kappa=-1/2 \\ \kappa=1/2 \end{matrix} \right\} \iota_\lambda=9/2$
										Utp Uth	Uop Uoh	$\left. \begin{matrix} \kappa=3/2 \\ \kappa=5/2 \end{matrix} \right\}$
										Uts Uto	Uos Uoo	$\left. \begin{matrix} \kappa=7/2 \\ \kappa=9/2 \end{matrix} \right\}$

Fig. 3. Seaborg table in the form of the basic representation $F_{ss'}^+$ of the Rumer-Fet group (basis $|\nu, s', \lambda, \iota_\lambda, \kappa\rangle$).

6.1 Masses of Elements

The lengthened group chain $G \supset G_1 \supset G_2 \supset G_3$ (13) allows us to provide a termwise mass splitting of the basic representation $F_{ss'}^+$ of the Rumer-Fet group. With this aim in view we introduce the

following mass formula:

$$m = m_0 + a \left[s'(2\nu - 3) - 5\nu + \frac{11}{2} + 2(\nu^2 - 1) \right] - b \cdot \lambda(\lambda + 1) + a' [2\kappa - 0, 1666\kappa^3 + 0, 0083\kappa^5 - 0, 0001\kappa^7] + (b'\iota_\lambda)^p - 1, \quad (15)$$

where

$$p = \begin{cases} 0, & \text{if } \iota_\lambda = \lambda - 1/2; \\ 1, & \text{if } \iota_\lambda = \lambda + 1/2. \end{cases}$$

As the “first perturbation” we have in (15) the Fet formula (10) corresponding to the group chain (9), where the basic representation $F_{ss'}^+$ is divided into the multiplets (ν, s', λ) with average masses of (ν, s', λ) . An analogue of the “second perturbation” in the formula (15) is defined by quantum numbers ι_λ, κ , that, according to the chain (13), leads to a partition of the multiplets (ν, s', λ) into the pair of multiplets of the subgroup G_3 (G_2/G_3 -reduction), and thereby we have here a termwise mass splitting. The masses of elements of the Seaborg table, starting from atomic number $Z = 121$ to $Z = 220$ ¹³, are calculated according to the mass formula (15) at the values $m_0 = 1$, $a = 17$, $b = 5, 5$, $a' = 2, 15$, $b' = 5, 3$ (see Tab. 3). The first column of Tab. 3 contains atomic number of the element; in the second column we have a generally accepted (according to IUPAC¹⁴) designation of the element; the third column contains quantum numbers of the element defining the vector $|\nu, s', \lambda, \iota_\lambda, \kappa\rangle$ of the basis (14)¹⁵; the fourth column contains the mass of the element calculated via the formula (15).

Tab 3. Masses of elements of the Seaborg table.

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
121	Ubu	$ 5, -1/2, 4, 7/2, -7/2\rangle$	308,3181
122	Ubb	$ 5, -1/2, 4, 7/2, -5/2\rangle$	309,2352
123	Ubt	$ 5, -1/2, 4, 7/2, -3/2\rangle$	310,6271
124	Ubq	$ 5, -1/2, 4, 7/2, -1/2\rangle$	313,8942
125	Ubp	$ 5, -1/2, 4, 7/2, 1/2\rangle$	318,1057
126	Ubn	$ 5, -1/2, 4, 7/2, 3/2\rangle$	321,3729
127	Ubs	$ 5, -1/2, 4, 7/2, 5/2\rangle$	322,7647
128	Ubo	$ 5, -1/2, 4, 7/2, 7/2\rangle$	323,6818
129	Ube	$ 5, -1/2, 4, 9/2, -9/2\rangle$	327,2491
130	Utn	$ 5, -1/2, 4, 9/2, -7/2\rangle$	331,1681
131	Utu	$ 5, -1/2, 4, 9/2, -5/2\rangle$	332,0852
132	Utb	$ 5, -1/2, 4, 9/2, -3/2\rangle$	333,4771
133	Utt	$ 5, -1/2, 4, 9/2, -1/2\rangle$	336,7421
134	Utq	$ 5, -1/2, 4, 9/2, 1/2\rangle$	340,9557
135	Utp	$ 5, -1/2, 4, 9/2, 3/2\rangle$	344,2229
136	Uth	$ 5, -1/2, 4, 9/2, 5/2\rangle$	345,6147
137	Uts	$ 5, -1/2, 4, 9/2, 7/2\rangle$	346,5318
138	Uto	$ 5, -1/2, 4, 9/2, 9/2\rangle$	350,4551

¹³As noted above, the Seaborg table is an extension of the Mendeleev table, highlighted in the Fig. 3 by a dotted border. The Table 2 shows the masses of the elements outside the dotted frame. In turn, the Mendeleev table contains two not yet discovered elements **Uue** ($Z = 119$) and **Ubn** ($Z = 120$) corresponding to the basis vectors $|4, 1/2, 0, 1/2, -1/2\rangle$ and $|4, 1/2, 0, 1/2, 1/2\rangle$. The masses of **Uue** and **Ubn**, calculated according to (15), are equal respectively to 304,8942 and 309,1057.

¹⁴IUPAC – International Union of Pure and Applied Chemistry.

¹⁵Recall, that according to group-theoretical description, the each element of periodic system corresponds to the vector $|\nu, s', \lambda, \iota_\lambda, \kappa\rangle$ of the basis (14), forming thereby a single quantum system.

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
139	Ute	$ 5, -1/2, 3, 5/2, -5/2\rangle$	353,2352
140	Uqn	$ 5, -1/2, 3, 5/2, -3/2\rangle$	354,6271
141	Uqu	$ 5, -1/2, 3, 5/2, -1/2\rangle$	357,8942
142	Uqb	$ 5, -1/2, 3, 5/2, 1/2\rangle$	362,1057
143	Uqt	$ 5, -1/2, 3, 5/2, 3/2\rangle$	365,3729
144	Uqq	$ 5, -1/2, 3, 5/2, 5/2\rangle$	366,7647
145	Uqp	$ 5, -1/2, 3, 7/2, -7/2\rangle$	369,8681
146	Uqh	$ 5, -1/2, 3, 7/2, -5/2\rangle$	370,7852
147	Uqs	$ 5, -1/2, 3, 7/2, -3/2\rangle$	372,1771
148	Uqo	$ 5, -1/2, 3, 7/2, -1/2\rangle$	375,4442
149	Uqe	$ 5, -1/2, 3, 7/2, 1/2\rangle$	379,6557
150	Upn	$ 5, -1/2, 3, 7/2, 3/2\rangle$	382,9229
151	Upu	$ 5, -1/2, 3, 7/2, 5/2\rangle$	384,3147
152	Uqp	$ 5, -1/2, 3, 7/2, 7/2\rangle$	385,2318
153	Upt	$ 5, -1/2, 2, 3/2, -3/2\rangle$	387,6271
154	Upq	$ 5, -1/2, 2, 3/2, -1/2\rangle$	390,8942
155	Upp	$ 5, -1/2, 2, 3/2, 1/2\rangle$	395,1057
156	Uph	$ 5, -1/2, 2, 3/2, 3/2\rangle$	398,3729
157	Ups	$ 5, -1/2, 2, 5/2, -5/2\rangle$	398,4852
158	Upo	$ 5, -1/2, 2, 5/2, -3/2\rangle$	399,8771
159	Upe	$ 5, -1/2, 2, 5/2, -1/2\rangle$	403,1442
160	Uhn	$ 5, -1/2, 2, 5/2, 1/2\rangle$	407,3557
161	Uhu	$ 5, -1/2, 2, 5/2, 3/2\rangle$	410,6229
162	Uhb	$ 5, -1/2, 2, 5/2, 5/2\rangle$	412,0147
163	Uht	$ 5, -1/2, 1, 1/2, -1/2\rangle$	412,8942
164	Uhq	$ 5, -1/2, 1, 1/2, 1/2\rangle$	417,1057
165	Uhp	$ 5, -1/2, 1, 3/2, -3/2\rangle$	416,5771
166	Uhn	$ 5, -1/2, 1, 3/2, -1/2\rangle$	419,8442
167	Uhs	$ 5, -1/2, 1, 3/2, 1/2\rangle$	424,0557
168	Uho	$ 5, -1/2, 1, 3/2, 3/2\rangle$	427,3229
169	Uhe	$ 5, -1/2, 0, 1/2, -1/2\rangle$	425,5452
170	Usn	$ 5, -1/2, 0, 1/2, 1/2\rangle$	429,7557
171	Usu	$ 5, 1/2, 4, 7/2, -7/2\rangle$	427,3181
172	Usb	$ 5, 1/2, 4, 7/2, -5/2\rangle$	428,2352
173	Ust	$ 5, 1/2, 4, 7/2, -3/2\rangle$	429,6271
174	Usq	$ 5, 1/2, 4, 7/2, -1/2\rangle$	432,8942
175	Usp	$ 5, 1/2, 4, 7/2, 1/2\rangle$	437,1057
176	Ush	$ 5, 1/2, 4, 7/2, 3/2\rangle$	440,3729
177	Uss	$ 5, 1/2, 4, 7/2, 5/2\rangle$	441,7647
178	Uso	$ 5, 1/2, 4, 7/2, 7/2\rangle$	442,6818
179	Use	$ 5, 1/2, 4, 9/2, -9/2\rangle$	446,2449
180	Uon	$ 5, 1/2, 4, 9/2, -7/2\rangle$	450,1681
181	Uou	$ 5, 1/2, 4, 9/2, -5/2\rangle$	451,0852
182	Uob	$ 5, 1/2, 4, 9/2, -3/2\rangle$	452,4771
183	Uot	$ 5, 1/2, 4, 9/2, -1/2\rangle$	455,7442

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
184	Uoq	$ \bar{5}, 1/2, 4, 9/2, 1/2\rangle$	459,9557
185	Uop	$ \bar{5}, 1/2, 4, 9/2, 3/2\rangle$	463,2229
186	Uoh	$ \bar{5}, 1/2, 4, 9/2, 5/2\rangle$	464,6147
187	Uos	$ \bar{5}, 1/2, 4, 9/2, 7/2\rangle$	465,5318
188	Uoo	$ \bar{5}, 1/2, 4, 9/2, 9/2\rangle$	469,4551
189	Uoe	$ \bar{5}, 1/2, 3, 5/2, -5/2\rangle$	472,2352
190	Uen	$ \bar{5}, 1/2, 3, 5/2, -3/2\rangle$	473,6271
191	Ueu	$ \bar{5}, 1/2, 3, 5/2, -1/2\rangle$	476,8942
192	Ueb	$ \bar{5}, 1/2, 3, 5/2, 1/2\rangle$	481,1057
193	Uet	$ \bar{5}, 1/2, 3, 5/2, 3/2\rangle$	484,3729
194	Ueq	$ \bar{5}, 1/2, 3, 5/2, 5/2\rangle$	485,7647
195	Uep	$ \bar{5}, 1/2, 3, 7/2, -7/2\rangle$	488,8681
196	Ueh	$ \bar{5}, 1/2, 3, 7/2, -5/2\rangle$	489,7852
197	Ues	$ \bar{5}, 1/2, 3, 7/2, -3/2\rangle$	491,177
198	Ueo	$ \bar{5}, 1/2, 3, 7/2, -1/2\rangle$	494,4442
199	Uee	$ \bar{5}, 1/2, 3, 7/2, 1/2\rangle$	498,6557
200	Bnn	$ \bar{5}, 1/2, 3, 7/2, 3/2\rangle$	501,9229
201	Bnu	$ \bar{5}, 1/2, 3, 7/2, 5/2\rangle$	503,3147
202	Bnb	$ \bar{5}, 1/2, 3, 7/2, 7/2\rangle$	504,2318
203	Bnt	$ \bar{5}, 1/2, 2, 3/2, -3/2\rangle$	506,6271
204	Bnq	$ \bar{5}, 1/2, 2, 3/2, -1/2\rangle$	509,8942
205	Bnp	$ \bar{5}, 1/2, 2, 3/2, 1/2\rangle$	514,1057
206	Bnh	$ \bar{5}, 1/2, 2, 3/2, 3/2\rangle$	517,3729
207	Bns	$ \bar{5}, 1/2, 2, 5/2, -5/2\rangle$	517,4852
208	Bno	$ \bar{5}, 1/2, 2, 5/2, -3/2\rangle$	518,8771
209	Bne	$ \bar{5}, 1/2, 2, 5/2, -1/2\rangle$	522,1442
210	Bun	$ \bar{5}, 1/2, 2, 5/2, 1/2\rangle$	526,3557
211	Buu	$ \bar{5}, 1/2, 2, 5/2, 3/2\rangle$	529,6229
212	Bub	$ \bar{5}, 1/2, 2, 5/2, 5/2\rangle$	531,0147
213	But	$ \bar{5}, 1/2, 1, 1/2, -1/2\rangle$	531,8942
214	Buq	$ \bar{5}, 1/2, 1, 1/2, 1/2\rangle$	536,1057
215	Bup	$ \bar{5}, 1/2, 1, 3/2, -3/2\rangle$	535,5771
216	Buh	$ \bar{5}, 1/2, 1, 3/2, -1/2\rangle$	538,8442
217	Bus	$ \bar{5}, 1/2, 1, 3/2, 1/2\rangle$	543,0557
218	Buo	$ \bar{5}, 1/2, 1, 3/2, 3/2\rangle$	546,3229
219	Bue	$ \bar{5}, 1/2, 0, 1/2, -1/2\rangle$	544,5442
220	Bbn	$ \bar{5}, 1/2, 0, 1/2, 1/2\rangle$	548,7557

7 10-periodic Extension

Fig. 4 shows a 10-periodic extension of the Mendeleev table in the form of the basic representation $F_{ss'}^+$ of the Rumer-Fet group G for the basis (14). The Mendeleev and Seaborg tables are separated by dotted frames with the symbols **M** and **S**, respectively. The first period of the Mendeleev table, including hydrogen **H** and helium **He**, corresponds to the simplest multiplet ($\nu = 1, s' = -1/2, \lambda = 0, \iota_\lambda = 1/2$) of the group G . The second period consists of three multiplets: lithium **Li** and beryllium **Be** ($\nu = 1, s' = 1/2, \lambda = 0, \iota_\lambda = 1/2$), boron **B** and carbon **C** ($\nu = 2, s' = -1/2, \lambda = 0, \iota_\lambda = 1/2$), elements **N**, **O**, **F**, **Ne** form a quadruplet ($\nu = 2, s' = -1/2, \lambda = 1, \iota_\lambda = 3/2$).

($\nu = 3, s' = -1/2, \lambda = 2, \iota_\lambda = 3/2$), and also sextet ($\nu = 3, s' = -1/2, \lambda = 2, \iota_\lambda = 5/2$) formed by the elements **Mn**, ..., **Zn**. This sextet and quadruplet ($\nu = 3, s' = -1/2, \lambda = 2, \iota_\lambda = 3/2$) form the first insertion decade (transitional elements). The fifth period has the analogous structure: doublets **Rb**, **Sr** ($\nu = 3, s' = -1/2, \lambda = 0, \iota_\lambda = 1/2$), **In**, **Sn** ($\nu = 3, s' = 1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplet **Sb**, **Te**, **I**, **Xe** ($\nu = 3, s' = 1/2, \lambda = 1, \iota_\lambda = 3/2$), quadruplet **Y**, **Zr**, **Nb**, **Mo** ($\nu = 3, s' = 1/2, \lambda = 2, \iota_\lambda = 3/2$) and sextet **Tc**, ..., **Cd** ($\nu = 3, s' = 1/2, \lambda = 2, \iota_\lambda = 5/2$) (the second insertion decade). The sixth period consists of seven multiplets: doublets **Cs**, **Ba** ($\nu = 3, s' = 1/2, \lambda = 0, \iota_\lambda = 1/2$) and **Tl**, **Pb** ($\nu = 4, s' = -1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplets **Bi**, **Po**, **At**, **Rn** ($\nu = 4, s' = -1/2, \lambda = 1, \iota_\lambda = 3/2$) and **Lu**, **Hf**, **Ta**, **W** ($\nu = 4, s' = -1/2, \lambda = 2, \iota_\lambda = 3/2$), sextets **Re**, ..., **Hg** ($\nu = 4, s' = -1/2, \lambda = 2, \iota_\lambda = 5/2$) and **La**, ..., **Sm** ($\nu = 4, s' = -1/2, \lambda = 3, \iota_\lambda = 5/2$), octet **Eu**, ..., **Yb** ($\nu = 4, s' = -1/2, \lambda = 3, \iota_\lambda = 7/2$). The seventh period (the last period of the Mendeleev table) duplicates the structure of the sixth period: doublets **Fr**, **Ra** ($\nu = 4, s' = -1/2, \lambda = 0, \iota_\lambda = 1/2$) and **Nh**, **Fl** ($\nu = 4, s' = 1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplets **Mc**, **Lv**, **Ts**, **Og** ($\nu = 4, s' = 1/2, \lambda = 1, \iota_\lambda = 3/2$) and **Lr**, **Rf**, **Db**, **Sg** ($\nu = 4, s' = 1/2, \lambda = 2, \iota_\lambda = 3/2$), sextets **Bh**, ..., **Cn** ($\nu = 4, s' = 1/2, \lambda = 2, \iota_\lambda = 5/2$) and **Ac**, ..., **Pu** ($\nu = 4, s' = 1/2, \lambda = 3, \iota_\lambda = 5/2$), octet **Am**, ..., **No** ($\nu = 4, s' = 1/2, \lambda = 3, \iota_\lambda = 7/2$). The eighth period¹⁶, forming an 8-periodic extension of the Mendeleev table (Seaborg table), consists of nine multiplets: doublets **Uue**, **Ubn** ($\nu = 4, s' = 1/2, \lambda = 0, \iota_\lambda = 1/2$) and **Uht**, **Uhq** ($\nu = 5, s' = -1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplets **Uhp**, ..., **Uho** ($\nu = 5, s' = -1/2, \lambda = 1, \iota_\lambda = 3/2$) and **Upt**, ..., **Uph** ($\nu = 5, s' = -1/2, \lambda = 2, \iota_\lambda = 3/2$), sextets **Ups**, ..., **Uhb** ($\nu = 5, s' = -1/2, \lambda = 2, \iota_\lambda = 5/2$) and **Ute**, ..., **Uqq** ($\nu = 5, s' = -1/2, \lambda = 3, \iota_\lambda = 5/2$), octets **Uqp**, ..., **Upb** ($\nu = 5, s' = -1/2, \lambda = 3, \iota_\lambda = 7/2$) and **Ubu**¹⁷, ..., **Ubo** ($\nu = 5, s' = -1/2, \lambda = 4, \iota_\lambda = 7/2$), decuplet **Ube**, ..., **Uto** ($\nu = 5, s' = -1/2, \lambda = 4, \iota_\lambda = 9/2$). The eighth period contains 50 elements. The ninth period, finishing Seaborg table, contains also nine multiplets: doublets **Uhe**, **Usn** ($\nu = 5, s' = -1/2, \lambda = 0, \iota_\lambda = 1/2$) and **But**, **Buq** ($\nu = 5, s' = 1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplets **Bup**, ..., **Buo** ($\nu = 5, s' = 1/2, \lambda = 1, \iota_\lambda = 3/2$) and **Bnt**, ..., **Bnh** ($\nu = 5, s' = 1/2, \lambda = 2, \iota_\lambda = 3/2$), sextets **Bns**, ..., **Bub** ($\nu = 5, s' = 1/2, \lambda = 2, \iota_\lambda = 5/2$) and **Uoe**, ..., **Ueq** ($\nu = 5, s' = 1/2, \lambda = 3, \iota_\lambda = 5/2$), octets **Uep**, ..., **Bnb** ($\nu = 5, s' = 1/2, \lambda = 3, \iota_\lambda = 7/2$) and **Usu**, ..., **Uso** ($\nu = 5, s' = 1/2, \lambda = 4, \iota_\lambda = 7/2$), decuplet **Use**, ..., **Uoo** ($\nu = 5, s' = 1/2, \lambda = 4, \iota_\lambda = 9/2$). The construction of a family of multiplets with the quantum number $\lambda = 5$ of the group G is started with the tenth period (in the Bohr's model it corresponds to the formation of h -shell). The tenth period consists of 11 multiplets: doublets **Bue**, **Bbn** ($\nu = 5, s' = 1/2, \lambda = 0, \iota_\lambda = 1/2$) and **Bop**, **Boh** ($\nu = 6, s' = -1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplets **Bos**, ..., **Ben** ($\nu = 6, s' = -1/2, \lambda = 1, \iota_\lambda = 3/2$) and **Bsp**, ..., **Bso** ($\nu = 6, s' = -1/2, \lambda = 2, \iota_\lambda = 3/2$), sextets **Bse**, ..., **Boq** ($\nu = 6, s' = -1/2, \lambda = 2, \iota_\lambda = 5/2$) and **Bhu**, ..., **Bhh** ($\nu = 6, s' = -1/2, \lambda = 3, \iota_\lambda = 5/2$), octets **Bhs**, ..., **Bsq** ($\nu = 6, s' = -1/2, \lambda = 3, \iota_\lambda = 7/2$) and **Bqt**, ..., **Bpn** ($\nu = 6, s' = -1/2, \lambda = 4, \iota_\lambda = 7/2$), decuplets **Bpu**, ..., **Bhn** ($\nu = 6, s' = -1/2, \lambda = 4, \iota_\lambda = 9/2$) and **Bbu**, ..., **Bth** ($\nu = 6, s' = -1/2, \lambda = 5, \iota_\lambda = 9/2$), 12-plet **Btu**, ..., **Bqb** ($\nu = 6, s' = -1/2, \lambda = 5, \iota_\lambda = 11/2$). The eleventh period has the analogous structure: doublets **Beu**, **Beb** ($\nu = 6, s' = -1/2, \lambda = 0, \iota_\lambda = 1/2$) and **Tps**, **Tpo** ($\nu = 6, s' = 1/2, \lambda = 1, \iota_\lambda = 1/2$), quadruplets **Tpe**, ..., **Thb** ($\nu = 6, s' = 1/2, \lambda = 1, \iota_\lambda = 3/2$) and **Tqs**, ..., **Tpn** ($\nu = 6, s' = 1/2, \lambda = 2, \iota_\lambda = 3/2$), sextets **Tpu**, ..., **Tph** ($\nu = 6, s' = 1/2, \lambda = 2, \iota_\lambda = 5/2$) and **Ttt**, ..., **Tto** ($\nu = 6, s' = 1/2, \lambda = 3, \iota_\lambda = 5/2$), octets **Tte**, ..., **Tqh** ($\nu = 6, s' = 1/2, \lambda = 3, \iota_\lambda = 7/2$) and **Tup**, ..., **Tbb** ($\nu = 6, s' = 1/2, \lambda = 4, \iota_\lambda = 7/2$), decuplets **Tbt**, ..., **Ttb** ($\nu = 6, s' = 1/2, \lambda = 4, \iota_\lambda = 9/2$) and **Bet**, ..., **Tnb** ($\nu = 6, s' = 1/2, \lambda = 5, \iota_\lambda = 9/2$), 12-plet **Tnt**, ..., **Tuq** ($\nu = 6, s' = 1/2, \lambda = 5, \iota_\lambda = 11/2$). The tenth and eleventh periods each contain

¹⁶The domain of hypothetical (undiscovered) elements of the periodic system begins with the eighth period.

¹⁷According to Bohr model, filling of g -shell is started with the element **Ubu** (Unbiunium). An analogue of g -shell in the Rumer-Fet model is a family of multiplets with quantum number $\lambda = 4$ of the group G .

72 elements. The lengths of periods form the following number sequence:

$$2, 8, 8, 18, 18, 32, 32, 50, 50, 72, 72, \dots \quad (16)$$

The numbers of this sequence are defined by the famous Rydberg formula $2p^2$ (p is an integer number). Rydberg series

$$R = 2(1^2 + 1^2 + 2^2 + 2^2 + 3^2 + 3^2 + 4^2 + 4^2 + \dots)$$

contains a doubled first period, which is somewhat inconsistent with reality, that is, sequence (16).

Further, 12-th period begins with the elements **Tht** (Trihexitrium, $Z = 363$) and **Thq** (Trihexiquadium, $Z = 364$), forming a doublet ($\nu = 6, s' = 1/2, \lambda = 0, \iota_\lambda = 1/2$). This period, already beyond the table in Fig. 4, contains 13 multiplets. The length of 12-th period is equal to 98 (in exact correspondence with the sequence (16)). A new family of multiplets with quantum number $\lambda = 6$ of group G starts from the 12-th period. This family corresponds to the i -shell filling. The 13-th period has the analogous structure.

Obviously, as the quantum number ν increases, we will see new “steps” (doubled periods) and corresponding λ -families of multiplets (shells) in Fig. 4.

7.1 Masses of elements of 10-th and 11-th periods

The table in Fig. 4 corresponds to the reduction chain (13). Theoretical masses of elements of 10-th and 11-th periods, starting from $Z = 221$ to $Z = 364$, are calculated according to the mass formula (15) at the values $m_0 = 1$, $a = 17$, $b = 5, 5$, $a' = 2, 15$, $b' = 5, 3$ (see Tab. 4).

Tab. 4. Masses of elements of 10-th and 11-th periods.

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
221	Bbu	$ 6, -1/2, 5, 9/2, -9/2\rangle$	521,3949
222	Bbb	$ 6, -1/2, 5, 9/2, -7/2\rangle$	525,3181
223	Bbt	$ 6, -1/2, 5, 9/2, -5/2\rangle$	526,2352
224	Bbq	$ 6, -1/2, 5, 9/2, -3/2\rangle$	527,6270
225	Bbp	$ 6, -1/2, 5, 9/2, -1/2\rangle$	530,8342
226	Bbh	$ 6, -1/2, 5, 9/2, 1/2\rangle$	535,1057
227	Bbs	$ 6, -1/2, 5, 9/2, 3/2\rangle$	538,3729
228	Bbo	$ 6, -1/2, 5, 9/2, 5/2\rangle$	539,7647
229	Bbe	$ 6, -1/2, 5, 9/2, 7/2\rangle$	540,6818
230	Btn	$ 6, -1/2, 5, 9/2, 9/2\rangle$	544,6050
231	Btu	$ 6, -1/2, 5, 11/2, -11/2\rangle$	545,0151
232	Btb	$ 6, -1/2, 5, 11/2, -9/2\rangle$	549,5449
233	Btt	$ 6, -1/2, 5, 11/2, -7/2\rangle$	553,4681
234	Btq	$ 6, -1/2, 5, 11/2, -5/2\rangle$	554,3852
235	Btp	$ 6, -1/2, 5, 11/2, -3/2\rangle$	555,7770
236	Bth	$ 6, -1/2, 5, 11/2, -1/2\rangle$	559,0442
237	Bts	$ 6, -1/2, 5, 11/2, 1/2\rangle$	563,2557
238	Bto	$ 6, -1/2, 5, 11/2, 3/2\rangle$	566,5229
239	Bte	$ 6, -1/2, 5, 11/2, 5/2\rangle$	567,9147
240	Bqn	$ 6, -1/2, 5, 11/2, 7/2\rangle$	568,8318

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
241	Bqu	$ 6, -1/2, 5, 11/2, 9/2\rangle$	572,7556
242	Bqb	$ 6, -1/2, 5, 11/2, 11/2\rangle$	582,2848
243	Bqt	$ 6, -1/2, 4, 7/2, -7/2\rangle$	580,3181
244	Bqq	$ 6, -1/2, 4, 7/2, -5/2\rangle$	581,2352
245	Bqp	$ 6, -1/2, 4, 7/2, -3/2\rangle$	582,6270
246	Bqh	$ 6, -1/2, 4, 7/2, -1/2\rangle$	585,8942
247	Bqs	$ 6, -1/2, 4, 7/2, 1/2\rangle$	590,1057
248	Bqo	$ 6, -1/2, 4, 7/2, 3/2\rangle$	593,3729
249	Bqe	$ 6, -1/2, 4, 7/2, 5/2\rangle$	594,7647
250	Bpn	$ 6, -1/2, 4, 7/2, 7/2\rangle$	595,6818
251	Bpu	$ 6, -1/2, 4, 9/2, -9/2\rangle$	599,2449
252	Bpb	$ 6, -1/2, 4, 9/2, -7/2\rangle$	603,1681
253	Bpt	$ 6, -1/2, 4, 9/2, -5/2\rangle$	604,0852
254	Bpq	$ 6, -1/2, 4, 9/2, -3/2\rangle$	605,4770
255	Bpp	$ 6, -1/2, 4, 9/2, -1/2\rangle$	608,7442
256	Bph	$ 6, -1/2, 4, 9/2, 1/2\rangle$	612,9557
257	Bps	$ 6, -1/2, 4, 9/2, 3/2\rangle$	616,2229
258	Bpo	$ 6, -1/2, 4, 9/2, 5/2\rangle$	617,6147
259	Bpe	$ 6, -1/2, 4, 9/2, 7/2\rangle$	618,5318
260	Bhn	$ 6, -1/2, 4, 9/2, 9/2\rangle$	622,4550
261	Bhu	$ 6, -1/2, 3, 5/2, -5/2\rangle$	625,2352
262	Bhb	$ 6, -1/2, 3, 5/2, -3/2\rangle$	626,6270
263	Bht	$ 6, -1/2, 3, 5/2, -1/2\rangle$	629,8942
264	Bhq	$ 6, -1/2, 3, 5/2, 1/2\rangle$	634,1057
265	Bhp	$ 6, -1/2, 3, 5/2, 3/2\rangle$	637,3729
266	Bhh	$ 6, -1/2, 3, 5/2, 5/2\rangle$	638,7647
267	Bhs	$ 6, -1/2, 3, 7/2, -7/2\rangle$	641,8681
268	Bho	$ 6, -1/2, 3, 7/2, -5/2\rangle$	642,7852
269	Bhe	$ 6, -1/2, 3, 7/2, -3/2\rangle$	644,1770
270	Bsn	$ 6, -1/2, 3, 7/2, -1/2\rangle$	647,4442
271	Bsu	$ 6, -1/2, 3, 7/2, 1/2\rangle$	651,6557
272	Bsb	$ 6, -1/2, 3, 7/2, 3/2\rangle$	654,9229
273	Bst	$ 6, -1/2, 3, 7/2, 5/2\rangle$	656,3147
274	Bsq	$ 6, -1/2, 3, 7/2, 7/2\rangle$	657,2318
275	Bsp	$ 6, -1/2, 2, 3/2, -3/2\rangle$	659,6270
276	Bsh	$ 6, -1/2, 2, 3/2, -1/2\rangle$	662,8942
277	Bss	$ 6, -1/2, 2, 3/2, 1/2\rangle$	667,1057
278	Bso	$ 6, -1/2, 2, 3/2, 3/2\rangle$	670,3729
279	Bse	$ 6, -1/2, 2, 5/2, -5/2\rangle$	670,4852
280	Bon	$ 6, -1/2, 2, 5/2, -3/2\rangle$	671,8770
281	Bou	$ 6, -1/2, 2, 5/2, -1/2\rangle$	675,1442
282	Bob	$ 6, -1/2, 2, 5/2, 1/2\rangle$	679,3557
283	Bot	$ 6, -1/2, 2, 5/2, 3/2\rangle$	682,6229
284	Boq	$ 6, -1/2, 2, 5/2, 5/2\rangle$	684,0147
285	Bop	$ 6, -1/2, 1, 1/2, -1/2\rangle$	684,8942
286	Boh	$ 6, -1/2, 1, 1/2, 1/2\rangle$	689,1057

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
287	Bos	$ 6, -1/2, 1, 3/2, -3/2\rangle$	689,5770
288	Boo	$ 6, -1/2, 1, 3/2, -1/2\rangle$	691,8442
289	Boe	$ 6, -1/2, 1, 3/2, 1/2\rangle$	696,0557
290	Ben	$ 6, -1/2, 1, 3/2, 3/2\rangle$	699,3229
291	Beu	$ 6, -1/2, 0, 1/2, -1/2\rangle$	699,8942
292	Beb	$ 6, -1/2, 0, 1/2, 1/2\rangle$	700,1037
293	Bet	$ 6, 1/2, 5, 9/2, -9/2\rangle$	674,3949
294	Beq	$ 6, 1/2, 5, 9/2, -7/2\rangle$	678,3181
295	Bep	$ 6, 1/2, 5, 9/2, -5/2\rangle$	679,2352
296	Beh	$ 6, 1/2, 5, 9/2, -3/2\rangle$	680,6270
297	Bes	$ 6, 1/2, 5, 9/2, -1/2\rangle$	683,8942
298	Beo	$ 6, 1/2, 5, 9/2, 1/2\rangle$	688,1097
299	Bee	$ 6, 1/2, 5, 9/2, 3/2\rangle$	691,3729
300	Tnn	$ 6, 1/2, 5, 9/2, 5/2\rangle$	692,7647
301	Tnu	$ 6, 1/2, 5, 9/2, 7/2\rangle$	693,6818
302	Tnb	$ 6, 1/2, 5, 9/2, 9/2\rangle$	697,6050
303	Tnt	$ 6, 1/2, 5, 11/2, -11/2\rangle$	693,0151
304	Tnq	$ 6, 1/2, 5, 11/2, -9/2\rangle$	702,5449
305	Tnp	$ 6, 1/2, 5, 11/2, -7/2\rangle$	706,4681
306	Tnh	$ 6, 1/2, 5, 11/2, -5/2\rangle$	707,3852
307	Tns	$ 6, 1/2, 5, 11/2, -3/2\rangle$	708,7770
308	Tno	$ 6, 1/2, 5, 11/2, -1/2\rangle$	712,0442
309	Tne	$ 6, 1/2, 5, 11/2, 1/2\rangle$	716,2557
310	Tun	$ 6, 1/2, 5, 11/2, 3/2\rangle$	719,5229
311	Tuu	$ 6, 1/2, 5, 11/2, 5/2\rangle$	720,9147
312	Tub	$ 6, 1/2, 5, 11/2, 7/2\rangle$	721,8318
313	Tut	$ 6, 1/2, 5, 11/2, 9/2\rangle$	725,7550
314	Tuq	$ 6, 1/2, 5, 11/2, 11/2\rangle$	735,2848
315	Tup	$ 6, 1/2, 4, 7/2, -7/2\rangle$	733,3181
316	Tuh	$ 6, 1/2, 4, 7/2, -5/2\rangle$	734,2352
317	Tus	$ 6, 1/2, 4, 7/2, -3/2\rangle$	735,6270
318	Tuo	$ 6, 1/2, 4, 7/2, -1/2\rangle$	738,8942
319	Tue	$ 6, 1/2, 4, 7/2, 1/2\rangle$	743,1057
320	Tbn	$ 6, 1/2, 4, 7/2, 3/2\rangle$	746,3729
321	Tbu	$ 6, 1/2, 4, 7/2, 5/2\rangle$	747,7647
322	Tbb	$ 6, 1/2, 4, 7/2, 7/2\rangle$	748,6818
323	Tbt	$ 6, 1/2, 4, 9/2, -9/2\rangle$	752,2449
324	Tbq	$ 6, 1/2, 4, 9/2, -7/2\rangle$	756,1681
325	Tbp	$ 6, 1/2, 4, 9/2, -5/2\rangle$	757,0852
326	Tbh	$ 6, 1/2, 4, 9/2, -3/2\rangle$	758,4770
327	Tbs	$ 6, 1/2, 4, 9/2, -1/2\rangle$	761,7442
328	Tbo	$ 6, 1/2, 4, 9/2, 1/2\rangle$	765,9557
329	Tbe	$ 6, 1/2, 4, 9/2, 3/2\rangle$	769,2229
330	Ttn	$ 6, 1/2, 4, 9/2, 5/2\rangle$	770,6147
331	Ttu	$ 6, 1/2, 4, 9/2, 7/2\rangle$	771,5318
332	Ttb	$ 6, 1/2, 4, 9/2, 9/2\rangle$	775,4550

Z	Element	Vector $ \nu, s', \lambda, \iota_\lambda, \kappa\rangle$	Mass
333	Ttt	$ 6, 1/2, 3, 5/2, -5/2\rangle$	778,2352
334	Ttq	$ 6, 1/2, 3, 5/2, -3/2\rangle$	779,6270
335	Ttp	$ 6, 1/2, 3, 5/2, -1/2\rangle$	782,8942
336	Tth	$ 6, 1/2, 3, 5/2, 1/2\rangle$	787,1057
337	Tts	$ 6, 1/2, 3, 5/2, 3/2\rangle$	790,3729
338	Tto	$ 6, 1/2, 3, 5/2, 5/2\rangle$	791,7647
339	Tte	$ 6, 1/2, 3, 7/2, -7/2\rangle$	794,8681
340	Tqn	$ 6, 1/2, 3, 7/2, -5/2\rangle$	795,7852
341	Tqu	$ 6, 1/2, 3, 7/2, -3/2\rangle$	797,1770
342	Tqb	$ 6, 1/2, 3, 7/2, -1/2\rangle$	800,4442
343	Tqt	$ 6, 1/2, 3, 7/2, 1/2\rangle$	804,6557
344	Tqq	$ 6, 1/2, 3, 7/2, 3/2\rangle$	807,9229
345	Tqp	$ 6, 1/2, 3, 7/2, 5/2\rangle$	809,3147
346	Tqh	$ 6, 1/2, 3, 7/2, 7/2\rangle$	810,2318
347	Tqs	$ 6, 1/2, 2, 3/2, -3/2\rangle$	812,6270
348	Tqo	$ 6, 1/2, 2, 3/2, -1/2\rangle$	815,8942
349	Tqe	$ 6, 1/2, 2, 3/2, 1/2\rangle$	820,1057
350	Tpn	$ 6, 1/2, 2, 3/2, 3/2\rangle$	823,3729
351	Tpu	$ 6, 1/2, 2, 5/2, -5/2\rangle$	823,4852
352	Tpb	$ 6, 1/2, 2, 5/2, -3/2\rangle$	824,8770
353	Tpt	$ 6, 1/2, 2, 5/2, -1/2\rangle$	828,1442
354	Tpq	$ 6, 1/2, 2, 5/2, 1/2\rangle$	832,3557
355	Tpp	$ 6, 1/2, 2, 5/2, 3/2\rangle$	835,6224
356	Tph	$ 6, 1/2, 2, 5/2, 5/2\rangle$	837,0147
357	Tps	$ 6, 1/2, 1, 1/2, -1/2\rangle$	837,8942
358	Tpo	$ 6, 1/2, 1, 1/2, 1/2\rangle$	842,1057
359	Tpe	$ 6, 1/2, 1, 3/2, -3/2\rangle$	841,5770
360	Thn	$ 6, 1/2, 1, 3/2, -1/2\rangle$	844,8442
361	Thu	$ 6, 1/2, 1, 3/2, 1/2\rangle$	849,0557
362	Thb	$ 6, 1/2, 1, 3/2, 3/2\rangle$	852,3229
363	Tht	$ 6, 1/2, 0, 1/2, -1/2\rangle$	848,8942
364	Thq	$ 6, 1/2, 0, 1/2, 1/2\rangle$	853,1057

8 Homological Series

All elements of the extended table, starting with hydrogen **H** ($Z = 1$) and ending with **Thq** (Trihexiquadium, $Z = 364$), form a single quantum system. The each element of the periodic system corresponds to the basis vector $|\nu, s', \lambda, \iota_\lambda, \kappa\rangle$, where $\nu, s', \lambda, \iota_\lambda, \kappa$ are quantum numbers

of the symmetry group G (Rumer-Fet group). Thus, we have the following set of state vectors:

$$\begin{aligned}
|\mathbf{H}\rangle &= \left| 1, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle, \\
|\mathbf{He}\rangle &= \left| 1, -\frac{1}{2}, 0, \frac{1}{2}, \frac{1}{2} \right\rangle, \\
|\mathbf{Li}\rangle &= \left| 1, \frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle, \\
&\vdots \\
|\mathbf{Thq}\rangle &= \left| 6, \frac{1}{2}, 0, \frac{1}{2}, \frac{1}{2} \right\rangle.
\end{aligned} \tag{17}$$

In accordance with quantum mechanical laws, in the aggregate (17), which forms a Hilbert space, we have linear superpositions of state vectors, as well as quantum transitions between different state vectors, that is, transitions between elements of the periodic system.

Let us now consider the operators that determine quantum transitions between the state vectors of the system (17):

$$\mathbf{\Gamma}_+ = \mathbf{P}_+ + \mathbf{Q}_+, \quad \mathbf{\Gamma}_- = \mathbf{P}_- + \mathbf{Q}_-. \tag{18}$$

Operators (18) connect subspaces \mathfrak{F}_n of the unitary representation F^+ of the conformal group $\text{SO}(2, 4)$ in the Fock space \mathfrak{F} . Indeed, the action of these operators on the basis vectors $|j, \sigma, \tau\rangle$ of \mathfrak{F} has the form

$$\begin{aligned}
\mathbf{\Gamma}_+ |j, \sigma, \tau\rangle &= i\sqrt{(j + \sigma + 1)(j - \tau + 1)} \left| j + \frac{1}{2}, \sigma + \frac{1}{2}, \tau - \frac{1}{2} \right\rangle - \\
&\quad - i\sqrt{(j - \sigma + 1)(j + \tau + 1)} \left| j + \frac{1}{2}, \sigma - \frac{1}{2}, \tau + \frac{1}{2} \right\rangle, \\
\mathbf{\Gamma}_- |j, \sigma, \tau\rangle &= -i\sqrt{(j + \sigma)(j - \tau)} \left| j - \frac{1}{2}, \sigma - \frac{1}{2}, \tau + \frac{1}{2} \right\rangle + \\
&\quad + i\sqrt{(j - \sigma)(j + \tau)} \left| j - \frac{1}{2}, \sigma + \frac{1}{2}, \tau - \frac{1}{2} \right\rangle.
\end{aligned}$$

Hence it follows that $\mathbf{\Gamma}_+$ transforms vectors of the subspace \mathfrak{F}_n into vectors of \mathfrak{F}_{n+1} , since for the Fock representation $\Phi_n = D_{\frac{n-1}{2}, \frac{n-1}{2}}$ in the subspace \mathfrak{F}_n , where $j = \frac{n-1}{2}$, increasing the number j by $1/2$ means increasing the number n by 1 (see Appendix B). Analogously, the operator $\mathbf{\Gamma}_-$ transforms vectors of the subspace \mathfrak{F}_n into vectors of \mathfrak{F}_{n-1} . Operators $\mathbf{\Gamma}_+$, $\mathbf{\Gamma}_-$ commute with the subgroup $G_2 = \text{SO}(3) \otimes \text{SU}(2)$ belonging the group chain (13). Indeed, in virtue of commutation relations of the conformal group $\text{SO}(2, 4)$ (see section 3) it follows that

$$[\mathbf{P}_\pm + \mathbf{Q}_\pm, \mathbf{J}_+ + \mathbf{K}_+] = [\mathbf{P}_\pm + \mathbf{Q}_\pm, \mathbf{J}_- + \mathbf{K}_-] = [\mathbf{P}_\pm + \mathbf{Q}_\pm, \mathbf{J}_3 + \mathbf{K}_3] = 0.$$

Therefore, operators $\mathbf{\Gamma}_+$, $\mathbf{\Gamma}_-$ save quantum number μ . Further, $\mathbf{\Gamma}_+$, $\mathbf{\Gamma}_-$ commute with a Casimir operator $(\mathbf{J}_1 + \mathbf{K}_1)^2 + (\mathbf{J}_2 + \mathbf{K}_2)^2 + (\mathbf{J}_3 + \mathbf{K}_3)^2$ of the subgroup $\text{SO}(3)$ and thereby save quantum number λ . It is easy to see that $\mathbf{\Gamma}_+$, $\mathbf{\Gamma}_-$ commute with the operators $\boldsymbol{\tau}_k$ ($k = 1, 2, 3$) of the subgroup $\text{SU}(2)$ and, therefore, they save quantum number s . Since $\mathbf{\Gamma}_+$ and $\mathbf{\Gamma}_-$ commute with $\mathbf{J}_k + \mathbf{K}_k$ and $\boldsymbol{\tau}_k$ separately, then they commute with the all subgroup $G_2 = \text{SO}(3) \otimes \text{SU}(2)$. Further, the operators $\mathbf{\Gamma}_+$, $\mathbf{\Gamma}_-$ commute with the subgroup $\text{SU}(2)'$, which defines second “doubling”, and, therefore, they

save quantum number s' . Since Γ_+ transforms \mathfrak{F}_n into \mathfrak{F}_{n+1} , and Γ_- transforms \mathfrak{F}_n into \mathfrak{F}_{n-1} , then in the space $\mathfrak{F}^4 = C(2) \otimes \mathfrak{F}^2 = C(2) \otimes [C(2) \otimes \mathfrak{F}]$ of the representation $F_{ss'}^+$, the operator Γ_+ , Γ_- raises, correspondingly lowers quantum number ν by 1. Thus, for the basis (8) the operator Γ_+ saves quantum numbers s' , λ , μ , s , raising ν by the unit, therefore, $\Gamma_+ |\nu, s', \lambda, \mu, s\rangle = \eta |\nu+1, s', \lambda, \mu, s\rangle$, where $\eta \neq 0$. Analogously, $\Gamma_- |\nu, s', \lambda, \mu, s\rangle = \eta' |\nu-1, s', \lambda, \mu, s\rangle$, where $\eta' \neq 0$. Since Γ_+ (correspondingly Γ_-) defines an isomorphic mapping of the space of (ν, s', λ) onto the space of $(\nu+1, s', \lambda)$ (corresp. $(\nu-1, s', \lambda)$), then η (corresp. η') does not depend on quantum numbers μ , s . Therefore, for the vectors $|\nu, s', \lambda, \iota_\lambda, \kappa\rangle$ of the basis (14) we have

$$\Gamma_+ |\nu, s', \lambda, \iota_\lambda, \kappa\rangle = \eta |\nu+1, s', \lambda, \iota_\lambda, \kappa\rangle, \quad (19)$$

$$\Gamma_- |\nu, s', \lambda, \iota_\lambda, \kappa\rangle = \eta' |\nu-1, s', \lambda, \iota_\lambda, \kappa\rangle. \quad (20)$$

The equality (20) holds at $0 \leq \lambda \leq \nu-2$. A visual sense of the operators Γ_+ , Γ_- is that they move basic vectors, represented by cells in Fig. 4, to the right, correspondingly to the left through horizontal columns of the table. At this point, Γ_+ always transfers basic vector of the column (ν, s') into basic vector of the same parity $(\nu+1, s')$ with multiplication by some non-null factor η . In turn, the operator Γ_- transfers basis vector of the column (ν, s') into basis vector of the same parity $(\nu-1, s')$ with multiplication by non-null factor η' when the column $(\nu-1, s')$ contains a vector on the same horizontal (otherwise we have zero).

Further, operators $\tau'_+ = \tau'_1 + i\tau'_2$, $\tau'_- = \tau'_1 - i\tau'_2$ of the subgroup $SU(2)'$ also define quantum transitions between state vectors (17). Since these operators commute with the subgroup $G_1 = SO(4) \otimes SU(2)$, they save quantum numbers ν , λ , ι_λ , κ , related with G_1 , and change only quantum number s' :

$$\tau'_+ |\nu, -\frac{1}{2}, \lambda, \iota_\lambda, \kappa\rangle = |\nu, \frac{1}{2}, \lambda, \iota_\lambda, \kappa\rangle, \quad (21)$$

$$\tau'_- |\nu, \frac{1}{2}, \lambda, \iota_\lambda, \kappa\rangle = |\nu, -\frac{1}{2}, \lambda, \iota_\lambda, \kappa\rangle. \quad (22)$$

A visual sense of the operators τ'_+ , τ'_- is that τ'_+ moves basis vectors of the each odd column (see Fig. 4) on horizontal into basis vectors of neighboring right column; in turn, τ'_- moves basis vectors of the each even column on horizontal into basis vectors of neighboring left column. Thus, operators (19)–(22) define quantum transitions between state vectors of the system (17).

It is easy to see that on the horizontals of Fig. 4 we have *Mendeleev homological series*, that is, families of elements with similar properties. Therefore, operators (19)–(22) define quantum transitions between elements of homological series. For example,

$$\begin{aligned} \Gamma_+ |\mathbf{H}\rangle &= \Gamma_+ \left| 1, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle = \eta_1 |\mathbf{Na}\rangle = \eta_1 \left| 2, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle \mapsto \\ &\eta_1 \Gamma_+ |\mathbf{Na}\rangle = \eta_1 \eta_2 |\mathbf{Rb}\rangle = \eta_1 \eta_2 \left| 3, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle \mapsto \\ &\eta_1 \eta_2 \Gamma_+ |\mathbf{Rb}\rangle = \eta_1 \eta_2 \eta_3 |\mathbf{Fr}\rangle = \eta_1 \eta_2 \eta_3 \left| 4, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle \mapsto \\ &\eta_1 \eta_2 \eta_3 \Gamma_+ |\mathbf{Fr}\rangle = \eta_1 \eta_2 \eta_3 \eta_4 |\mathbf{Uhe}\rangle = \eta_1 \eta_2 \eta_3 \eta_4 \left| 5, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right\rangle. \end{aligned}$$

Further, operators τ'_+ , τ'_- establish homology between lanthanides and actinides¹⁸:

$$\tau'_+ |\mathbf{La}\rangle = \tau'_+ \left| 4, -\frac{1}{2}, 3, \frac{5}{2}, -\frac{5}{2} \right\rangle = |\mathbf{Ac}\rangle = \left| 4, \frac{1}{2}, 3, \frac{5}{2}, -\frac{5}{2} \right\rangle,$$

¹⁸This homology was first discovered by Seaborg. It is obvious that Seaborg homology is a particular case of the Mendeleev homology.

$$\begin{aligned}
\tau'_+|\mathbf{Ce}\rangle &= \tau'_+ \left| 4, -\frac{1}{2}, 3, \frac{5}{2}, -\frac{3}{2} \right\rangle = |\mathbf{Th}\rangle = \left| 4, \frac{1}{2}, 3, \frac{5}{2}, -\frac{3}{2} \right\rangle, \\
&\vdots \\
\tau'_+|\mathbf{Yb}\rangle &= \tau'_+ \left| 4, -\frac{1}{2}, 3, \frac{7}{2}, \frac{7}{2} \right\rangle = |\mathbf{No}\rangle = \left| 4, \frac{1}{2}, 3, \frac{7}{2}, \frac{7}{2} \right\rangle,
\end{aligned}$$

By means of operators Γ_+ , Γ_- we can continue the Seaborg homology to a superactinide group:

$$\begin{aligned}
\Gamma_+|\mathbf{La}\rangle &= \Gamma_+ \left| 4, -\frac{1}{2}, 3, \frac{5}{2}, -\frac{5}{2} \right\rangle = \eta|\mathbf{Ute}\rangle = \eta \left| 5, -\frac{1}{2}, 3, \frac{5}{2}, -\frac{5}{2} \right\rangle, \\
\Gamma_+|\mathbf{Ce}\rangle &= \Gamma_+ \left| 4, -\frac{1}{2}, 3, \frac{5}{2}, -\frac{3}{2} \right\rangle = \eta|\mathbf{Uqn}\rangle = \eta \left| 5, -\frac{1}{2}, 3, \frac{5}{2}, -\frac{3}{2} \right\rangle, \\
&\vdots \\
\Gamma_+|\mathbf{Yb}\rangle &= \Gamma_+ \left| 4, -\frac{1}{2}, 3, \frac{7}{2}, \frac{7}{2} \right\rangle = \eta|\mathbf{Upb}\rangle = \eta \left| 5, -\frac{1}{2}, 3, \frac{7}{2}, \frac{7}{2} \right\rangle.
\end{aligned}$$

Correspondingly,

$$\begin{aligned}
\Gamma_+|\mathbf{Ac}\rangle &= \Gamma_+ \left| 4, \frac{1}{2}, 3, \frac{5}{2}, -\frac{5}{2} \right\rangle = \eta|\mathbf{Uoe}\rangle = \eta \left| 5, \frac{1}{2}, 3, \frac{5}{2}, -\frac{5}{2} \right\rangle, \\
\Gamma_+|\mathbf{Th}\rangle &= \Gamma_+ \left| 4, \frac{1}{2}, 3, \frac{5}{2}, -\frac{3}{2} \right\rangle = \eta|\mathbf{Uen}\rangle = \eta \left| 5, \frac{1}{2}, 3, \frac{5}{2}, -\frac{3}{2} \right\rangle, \\
&\vdots \\
\Gamma_+|\mathbf{No}\rangle &= \Gamma_+ \left| 4, \frac{1}{2}, 3, \frac{7}{2}, \frac{7}{2} \right\rangle = \eta|\mathbf{Bnb}\rangle = \eta \left| 5, \frac{1}{2}, 3, \frac{7}{2}, \frac{7}{2} \right\rangle.
\end{aligned}$$

In conclusion of this paragraph we will say a few words about the *principle of superposition* in relation to the system (17). Apparently, the situation here is similar to the Wigner's *superselection principle* [40] in particle physics, according to which not every superposition of physically possible states leads again to a physically possible state. The Wigner's principle limits (*superselection rules*) the existence of superpositions of states. According to the superselection rules, superpositions of physically possible states exist only in the coherent subspaces of the physical Hilbert space. Thus, the problem of determining coherent subspaces for the system of states (17) arises.

9 Hypertwistors

The Rumer-Fet group is constructed in many respects by analogy with the groups of internal (dynamic) symmetries, such as $SU(3)$ and $SU(6)$. Using quark model and $SU(3)$ -symmetry, we continue this analogy. As is known, quark is a vector of fundamental representation of the group $SU(3)$. Let us define a vector of “fundamental” representation of the Rumer-Fet group.

The Rumer-Fet group

$$SO(2, 4) \otimes SU(2) \otimes SU(2)'$$

is equivalent to

$$\widetilde{SO}(2, 4) \otimes SU(2) \simeq SU(2, 2) \otimes SU(2),$$

where $SU(2, 2)$ is a double covering of the conformal group (the group of pseudounitary unimodular 4×4 matrices). Further, in virtue of the isomorphism (C.4) (see Appendix C) we will consider the double covering $SU(2, 2)$ as a *spinor group*¹⁹. Spintensor representations of the group $\mathbf{Spin}_+(2, 4)$ form a substratum of finite-dimensional representations $\tau_{k/2, r/2}, \bar{\tau}_{k/2, r/2}$ of the conformal group, which realized in the spaces $\text{Sym}_{(k, r)} \subset \mathbb{S}_{2k+r}$ and $\overline{\text{Sym}}_{(k, r)} \subset \overline{\mathbb{S}}_{2k+r}$, where \mathbb{S}_{2k+r} is a spinspace. Twistor $\mathbf{Z}^\alpha = (\mathbf{s}^\alpha, \mathbf{s}_{\dot{\alpha}})^\top$ is a vector of fundamental representation of the group $\mathbf{Spin}_+(2, 4)$, where $\alpha, \dot{\alpha} = 0, 1$ and $\mathbf{s}^\alpha, \mathbf{s}_{\dot{\alpha}}$ are 2-component mutually conjugated spinors. Hence it immediately follows that a doubled twistor

$$\mathbf{z} = \begin{bmatrix} \mathbf{z}_+ \\ \mathbf{z}_- \end{bmatrix}, \quad (23)$$

or *hypertwistor*, is a vector of fundamental representation of the group $SU(2, 2) \otimes SU(2)$. Further, the twistor $\mathbf{Z} = [\mathbf{S}, \bar{\mathbf{S}}]^\top$ is a vector of general spintensor representation of the group $\mathbf{Spin}_+(2, 4)$, where \mathbf{S} is a spintensor of the form (12). Therefore, a general hypertwistor is defined by the expression of the form (23), where $\mathbf{z}_+ = [\mathbf{S}, \bar{\mathbf{S}}]^\top$, $\mathbf{z}_- = [\bar{\mathbf{S}}, \mathbf{S}]^\top$.

Applying GNS-construction, we obtain vector states

$$\omega_\Phi(H) = \frac{\langle \Phi | \pi(H) \Phi \rangle}{\langle \Phi | \Phi \rangle} = \frac{\langle \Phi | F_{ss'}^+(H) \Phi \rangle}{\langle \Phi | \Phi \rangle},$$

where H is an energy operator, $|\Phi\rangle$ is a cyclic vector of the Hilbert space \mathbf{H}_∞ . A set of all pure states $\omega_\Phi(H)$ forms a *physical Hilbert space* $\mathbf{H}_{\text{phys}} = \mathbf{H}_8 \otimes \mathbf{H}_\infty$ ²⁰ and, correspondingly, a *space of rays* $\hat{H} = \mathbf{H}_{\text{phys}}/S^1$.

Further, with the aim of observance of electroneutrality and inclusion of discrete symmetries it is necessary to expand the double covering $SU(2, 2) \simeq \mathbf{Spin}_+(2, 4)$ up to an *universal covering* $\mathbf{Pin}(2, 4)$. In general form (for arbitrary orthogonal groups) such extension has been given in the works [7, 44, 9, 11]. At this point, a pseudo-automorphism $\mathcal{A} \rightarrow \bar{\mathcal{A}}$ of the complex Clifford algebra \mathbb{C}_n [45] plays a central role, where \mathcal{A} is an arbitrary element of the algebra \mathbb{C}_n . Since the real spinor structure appears as a result of reduction $\mathbb{C}_{2(k+r)} \rightarrow \mathcal{O}_{p,q}$, then, as a consequence, the *charge conjugation* C (pseudo-automorphism $\mathcal{A} \rightarrow \bar{\mathcal{A}}$) for algebras $\mathcal{O}_{p,q}$ over the real number field $\mathbb{F} = \mathbb{R}$ and the quaternion division ring $\mathbb{K} \simeq \mathbb{H}$ (types $p - q \equiv 4, 6 \pmod{8}$) is reduced to a *particle-antiparticle exchange* C' . As is known, there are two classes of neutral particles: 1) particles that have antiparticles, such as neutrons, neutrinos, and so on; 2) particles that coincide with their antiparticles (for example, photons, π^0 -mesons, and so on), that is, the so-called *truly neutral particle*. The first class is described by neutral states $\omega_\Phi^i(H)$ with algebras $\mathcal{O}_{p,q}$ over the field $\mathbb{F} = \mathbb{R}$ with the rings $\mathbb{K} \simeq \mathbb{H}$ and $\mathbb{K} \simeq \mathbb{H} \oplus \mathbb{H}$ (types $p - q \equiv 4, 6 \pmod{8}$ and $p - q \equiv 5 \pmod{8}$). To describe the second class of neutral particles we introduce *truly neutral states* $\omega_\Phi^{r_0}(H)$ with algebras $\mathcal{O}_{p,q}$ over the real number field $\mathbb{F} = \mathbb{R}$ and real division rings $\mathbb{K} \simeq \mathbb{R}$ and $\mathbb{K} \simeq \mathbb{R} \oplus \mathbb{R}$ (types $p - q \equiv 0, 2 \pmod{8}$ and $p - q \equiv 1 \pmod{8}$). In the case of states $\omega_\Phi^{r_0}(H)$ the pseudo-automorphism $\mathcal{A} \rightarrow \bar{\mathcal{A}}$ is reduced to the identical transformation (the particle coincides with its antiparticle).

Following to [46], we define $\mathbf{H}_{\text{phys}} = \mathbf{H}_8 \otimes \mathbf{H}_\infty$ as a \mathbb{K} -Hilbert space, that is, as a space endowed with a $*$ -ring structure, where $*$ -ring is isomorphic to a division ring $\mathbb{K} = \mathbb{R}, \mathbb{C}, \mathbb{H}$. Thus, the

¹⁹The elements of group $\mathbf{Spin}_+(2, 4)$ are 15 bivectors $\mathbf{e}_i \mathbf{e}_j = \mathbf{e}_{ij}$, where $i, j = 1, \dots, 6$. The explicit form of all fifteen generators leads through Cartan decomposition for group $SU(2, 2)$ to biquaternion angles, that is, to generalization of complex and quaternion angles for groups $SL(2, \mathbb{C})$ and $Sp(1, 1)$, where $Sp(1, 1)$ is a double covering of the de Sitter group [41, 42, 43].

²⁰At the restriction of the group G onto the Lorentz subgroup $SO_0(1, 3)$ and application of GNS-construction within double covering $SL(2, \mathbb{C}) \simeq \mathbf{Spin}_+(1, 3)$, we obtain a *spinor* (vector of fundamental representation of the group $\mathbf{Spin}_+(1, 3)$), acting in a doubled Hilbert space $\mathbf{H}_2 \otimes \mathbf{H}_\infty$ (Pauli space). Spinor is a particular case of hypertwistor.

hypertwistor has a tensor structure (energy, mass) and a \mathbb{K} -linear structure (charge), and the connection of these two structures leads to a dynamic change in charge and mass.

Appendix A: Lorentz group and van der Waerden representation

As is known, an universal covering of the proper orthochronous Lorentz group $SO_0(1, 3)$ (rotation group of the Minkowski space-time $\mathbb{R}^{1,3}$) is the spinor group

$$\mathbf{Spin}_+(1, 3) \simeq \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathbb{C}_2 : \det \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = 1 \right\} = \mathrm{SL}(2, \mathbb{C}).$$

Let $\mathfrak{g} \rightarrow T_{\mathfrak{g}}$ be an arbitrary linear representation of the proper orthochronous Lorentz group $SO_0(1, 3)$ and let $A_i(t) = T_{a_i(t)}$ be an infinitesimal operator corresponding to the rotation $a_i(t) \in SO_0(1, 3)$. Analogously, let $B_i(t) = T_{b_i(t)}$, where $b_i(t) \in SO_0(1, 3)$ is the hyperbolic rotation. The elements A_i and B_i form a basis of the group algebra $\mathfrak{sl}(2, \mathbb{C})$ and satisfy the relations

$$\left. \begin{aligned} [A_1, A_2] &= A_3, & [A_2, A_3] &= A_1, & [A_3, A_1] &= A_2, \\ [B_1, B_2] &= -A_3, & [B_2, B_3] &= -A_1, & [B_3, B_1] &= -A_2, \\ [A_1, B_1] &= 0, & [A_2, B_2] &= 0, & [A_3, B_3] &= 0, \\ [A_1, B_2] &= B_3, & [A_1, B_3] &= -B_2, \\ [A_2, B_3] &= B_1, & [A_2, B_1] &= -B_3, \\ [A_3, B_1] &= B_2, & [A_3, B_2] &= -B_1. \end{aligned} \right\} \quad (\text{A.1})$$

Defining the operators

$$\begin{aligned} X_l &= \frac{1}{2}i(A_l + iB_l), & Y_l &= \frac{1}{2}i(A_l - iB_l), \\ (l &= 1, 2, 3), \end{aligned} \quad (\text{A.2})$$

we come to a *complex shell* of the group algebra $\mathfrak{sl}(2, \mathbb{C})$. Using the relations (A.1), we find

$$[X_k, X_l] = i\varepsilon_{klm}X_m, \quad [Y_l, Y_m] = i\varepsilon_{lmn}Y_n, \quad [X_l, Y_m] = 0. \quad (\text{A.3})$$

From the relations (A.3) it follows that each of the sets of infinitesimal operators X and Y generates the group $SU(2)$ and these two groups commute with each other. Thus, from the relations (A.3) it follows that the group algebra $\mathfrak{sl}(2, \mathbb{C})$ (within the complex shell) is algebraically isomorphic to the direct sum $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ ²¹.

Further, introducing operators of the form ('rising' and 'lowering' operators of the group $SL(2, \mathbb{C})$)

$$\left. \begin{aligned} X_+ &= X_1 + iX_2, & X_- &= X_1 - iX_2, \\ Y_+ &= Y_1 + iY_2, & Y_- &= Y_1 - iY_2, \end{aligned} \right\} \quad (\text{A.4})$$

²¹In a sense, it allows one to represent the group $SL(2, \mathbb{C})$ by a product $SU(2) \otimes SU(2)$ as it done by Ryder in his textbook [47]. Moreover, in the works [48, 49] the Lorentz group is represented by a product $SU_R(2) \otimes SU_L(2)$, where the spinors $\psi(p^\mu) = \begin{pmatrix} \phi_R(p^\mu) \\ \phi_L(p^\mu) \end{pmatrix}$ ($\phi_R(p^\mu)$ and $\phi_L(p^\mu)$ are the right- and left-handed spinors) are transformed within $(j, 0) \oplus (0, j)$ representation space, in our case $j = l = \frac{1}{2}$. However, the isomorphism $SL(2, \mathbb{C}) \simeq SU(2) \otimes SU(2)$ is not correct from group-theoretical viewpoint. Indeed, the groups $SO_0(1, 3)$ and $SO(4)$ are real forms of the complex 6-dimensional Lie group $SO(4, \mathbb{C})$ with complex Lie algebra $D_2 = A_1 + A_1$. Real Lie algebras are compact iff the Killing form is negative definite [50]. That is the case for Lie algebra of $SO(4)$, not for $SO_0(1, 3)$.

we see that

$$\begin{aligned} [\mathbf{X}_3, \mathbf{X}_+] &= \mathbf{X}_+, & [\mathbf{X}_3, \mathbf{X}_-] &= -\mathbf{X}_-, & [\mathbf{X}_+, \mathbf{X}_-] &= 2\mathbf{X}_3, \\ [\mathbf{Y}_3, \mathbf{Y}_+] &= \mathbf{Y}_+, & [\mathbf{Y}_3, \mathbf{Y}_-] &= -\mathbf{Y}_-, & [\mathbf{Y}_+, \mathbf{Y}_-] &= 2\mathbf{Y}_3. \end{aligned}$$

In virtue of commutativity of the relations (A.3) a space of an irreducible finite-dimensional representation of the group $\text{SL}(2, \mathbb{C})$ can be spanned on the totality of $(2l+1)(2\dot{l}+1)$ basis ket-vectors $|l, m; \dot{l}, \dot{m}\rangle$ and basis bra-vectors $\langle l, m; \dot{l}, \dot{m}|$, where l, m, \dot{l}, \dot{m} are integer or half-integer numbers, $-l \leq m \leq l$, $-\dot{l} \leq \dot{m} \leq \dot{l}$. Therefore,

$$\begin{aligned} \mathbf{X}_- |l, m; \dot{l}, \dot{m}\rangle &= \sqrt{(l+m)(l-m+1)} |l, m-1; \dot{l}, \dot{m}\rangle \quad (m > -l), \\ \mathbf{X}_+ |l, m; \dot{l}, \dot{m}\rangle &= \sqrt{(l-m)(l+m+1)} |l, m+1; \dot{l}, \dot{m}\rangle \quad (m < l), \\ \mathbf{X}_3 |l, m; \dot{l}, \dot{m}\rangle &= m |l, m; \dot{l}, \dot{m}\rangle, \\ \langle l, m; \dot{l}, \dot{m}| \mathbf{Y}_- &= \langle l, m; \dot{l}, \dot{m}-1| \sqrt{(\dot{l}+\dot{m})(\dot{l}-\dot{m}+1)} \quad (\dot{m} > -\dot{l}), \\ \langle l, m; \dot{l}, \dot{m}| \mathbf{Y}_+ &= \langle l, m; \dot{l}, \dot{m}+1| \sqrt{(\dot{l}-\dot{m})(\dot{l}+\dot{m}+1)} \quad (\dot{m} < \dot{l}), \\ \langle l, m; \dot{l}, \dot{m}| \mathbf{Y}_3 &= \langle l, m; \dot{l}, \dot{m}| \dot{m}. \end{aligned} \tag{A.5}$$

In contrast to the Gelfand-Naimark representation for the Lorentz group [51, 52], which does not find a wide application in physics, a representation (A.5) is a most useful in theoretical physics (see, for example, [53, 54, 55, 47]). This representation for the Lorentz group was first given by van der Waerden in his brilliant book [56]. It should be noted here that the representation basis, defined by the formulae (A.2)–(A.5), has an evident physical meaning. For example, in the case of $(1, 0) \oplus (0, 1)$ -representation space there is an analogy with the photon spin states. Namely, the operators \mathbf{X} and \mathbf{Y} correspond to the right and left polarization states of the photon. For that reason we will call the canonical basis consisting of the vectors $|lm; \dot{l}\dot{m}\rangle$ as a *helicity basis*.

Thus, the complex shell of the group algebra $\mathfrak{sl}(2, \mathbb{C})$, generating complex momentum, leads to a *duality* which is mirrored in the appearance of the two spaces: a space of ket-vectors $|l, m; \dot{l}, \dot{m}\rangle$ and a dual space of bra-vectors $\langle l, m; \dot{l}, \dot{m}|$.

Appendix B: The group $\text{SO}(4)$ and Fock representation

As is known, the group $\text{SO}(4)$ is a maximal compact subgroup of the conformal group $\text{SO}_0(2, 4)$. $\text{SO}(4)$ corresponds to basis elements $\mathbf{J} = (\mathbf{J}_1, \mathbf{J}_2, \mathbf{J}_3)$ and $\mathbf{K} = (\mathbf{K}_1, \mathbf{K}_2, \mathbf{K}_3)$ of the algebra $\mathfrak{so}(2, 4)$:

$$[\mathbf{J}_k, \mathbf{J}_l] = i\varepsilon_{klm} \mathbf{J}_m, \quad [\mathbf{J}_k, \mathbf{K}_l] = i\varepsilon_{klm} \mathbf{K}_m, \quad [\mathbf{K}_k, \mathbf{K}_l] = i\varepsilon_{klm} \mathbf{J}_m.$$

Introducing linear combinations $\mathbf{V} = (\mathbf{J} + \mathbf{K})/2$ and $\mathbf{V}' = (\mathbf{J} - \mathbf{K})/2$, we obtain

$$[\mathbf{V}_k, \mathbf{V}_l] = i\varepsilon_{klm} \mathbf{V}_m, \quad [\mathbf{V}'_k, \mathbf{V}'_l] = i\varepsilon_{klm} \mathbf{V}'_m.$$

Generators \mathbf{V} and \mathbf{V}' form bases of the two independent algebras $\mathfrak{so}(3)$. It means that the group $\text{SO}(4)$ is isomorphic to the product $\text{SO}(3) \otimes \text{SO}(3)$ ²².

An universal covering of the rotation group $\text{SO}(4)$ of the four-dimensional Euclidean space \mathbb{R}^4 is a spinor group

$$\mathbf{Spin}(4) \simeq \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathbb{H} \oplus \mathbb{H} : \det \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = 1 \right\} = \text{SU}(2) \otimes \text{SU}(2).$$

²²Such state of affairs is explained by the following definition: the group $\text{SO}(4)$ is *locally* decomposed into a direct product of subgroups $\text{SO}(3)$. *In whole* (that is, without the supposition that all the matrices similar the unit matrix) this decomposition is ambiguous. $\text{SO}(4)$ is an unique group (among the all orthogonal groups $\text{SO}(n)$) which admits such local decomposition.

Let $\text{SO}(3)_{\mathbf{J}}$ and $\text{SO}(3)_{\mathbf{K}}$ be the subgroups of $\text{SO}(4)$ with the generators \mathbf{J}_k and \mathbf{K}_k ($k = 1, 2, 3$), respectively. Then the each irreducible representation T of the group $\text{SO}(4)$ has the following structure: a space \mathfrak{R} of the representation T is a tensor product of spaces \mathfrak{R}_1 and \mathfrak{R}_2 in which we have irreducible representations D_{j_1} and D_{j_2} of the subgroups $\text{SO}(3)_{\mathbf{J}}$ and $\text{SO}(3)_{\mathbf{K}}$ with dimensionality $2j_1 + 1$ and $2j_2 + 1$. Thus, a dimensionality of T is equal to $(2j_1 + 1)(2j_2 + 1)$, where j_1 and j_2 are integer or half-integer numbers. An action of \mathbf{J}_k , \mathbf{K}_k on the basis vectors is defined by the formulas

$$\begin{aligned} \mathbf{J}_-|\sigma, \tau\rangle &= \sqrt{(j_1 + \sigma)(j_1 - \sigma + 1)}|\sigma - 1, \tau\rangle, \\ \mathbf{J}_+|\sigma, \tau\rangle &= \sqrt{(j_1 - \sigma)(j_1 + \sigma + 1)}|\sigma + 1, \tau\rangle, \\ \mathbf{J}_3|\sigma, \tau\rangle &= \sigma|\sigma, \tau\rangle, \\ \mathbf{K}_-|\sigma, \tau\rangle &= \sqrt{(j_2 + \tau)(j_2 - \tau + 1)}|\sigma, \tau - 1\rangle, \\ \mathbf{K}_+|\sigma, \tau\rangle &= \sqrt{(j_2 - \tau)(j_2 + \tau + 1)}|\sigma, \tau + 1\rangle, \\ \mathbf{K}_3|\sigma, \tau\rangle &= \tau|\sigma, \tau\rangle. \end{aligned} \tag{B.1}$$

This representation of the group $\text{SO}(4)$, denoted via $D_{j_1 j_2}$, is irreducible and unitary.

A structure of the Fock representation Φ of $\text{SO}(4)$ is defined by the decomposition onto irreducible components Φ_n in the spaces \mathfrak{F}_n . As is known [18], at the reduction on the subgroup $\text{SO}(3)$ an irreducible representation Φ_n is decomposed into a sum of irreducible representations of $\text{SO}(3)$ with dimensionality $1, 3, \dots, 2n - 1$. Taking into account the structure of irreducible representations of the group $\text{SO}(4)$, we see that a smallest dimensionality 1 should be equal to $2|j_1 - j_2| + 1$, from which it follows that $j_1 = j_2$. Therefore, the representation Φ_n has the form $D_{j_1 j_2}$, and since for the biggest dimensionality should be $2n - 1 = 2(j_1 + j_2) + 1 = 4j + 1$, then $j = (n - 1)/2$. Thus, for the Fock representation we have the following structure:

$$\Phi = \Phi_1 \oplus \Phi_2 \oplus \dots \oplus \Phi_n \oplus \dots, \quad \Phi_n = D_{\frac{n-1}{2}, \frac{n-1}{2}}.$$

It means that in the each space \mathfrak{F}_n there is an orthonormal basis $|j, \sigma, \tau\rangle$, where $j = (n - 1)/2$. All these bases form together an orthonormal basis of the Fock space \mathfrak{F} :

$$\begin{aligned} |j, \sigma, \tau\rangle \quad (j = 0, 1/2, 1, 3/2, \dots; \\ \sigma = -j, -j + 1, \dots, j - 1, j; \tau = -j, -j + 1, \dots, j - 1, j), \end{aligned} \tag{B.2}$$

in which Lie algebra of the group $\text{SO}(4)$ acts via the formulas (B.1) with $j_1 = j_2 = j$.

Appendix C: Twistor structure and group $\text{SU}(2, 2)$

The main idea of the Penrose twistor program [57, 58] consists in the representation of classical space-time as some secondary construction obtained from more primary notions. As more primary notions we have here two-component (complex) spinors, moreover, the pairs of two-component spinors. In the Penrose program they called *twistors*.

So, a twistor \mathbf{Z}^α is defined by the pair of two-component spinors: spinor ω^s and covariant spinor $\pi_{\dot{s}}$ from a conjugated space, that is, $\mathbf{Z}^\alpha = (\omega^s, \pi_{\dot{s}})$. In twistor theory momentum ($\vec{\omega}$) and impulse ($\vec{\pi}$) of the particle are constructed from the quantities ω^s and $\pi_{\dot{s}}$. One of the most important aspects of this theory is a *transition from twistors to coordinate space-time*. Penrose described this transition by means of so-called *basic relation of twistor theory*

$$\omega^s = ix^{s\dot{r}} \pi_{\dot{r}}, \tag{C.1}$$

where x^{sr} is a mixed spintensor of second rank. In more detailed record this relation has the form

$$\begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix} = \frac{i}{\sqrt{2}} \begin{bmatrix} x^0 + x^3 & x^1 + ix^2 \\ x^1 + ix^2 & x^0 - x^3 \end{bmatrix} \begin{bmatrix} \pi_1 \\ \pi_2 \end{bmatrix}.$$

From (C.1) it immediately follows that points of space-time $\mathbb{R}^{1,3}$ are reconstructed over the twistor space \mathbb{C}^4 (these points correspond to linear subspaces of the twistor space \mathbb{C}^4). Therefore, points of $\mathbb{R}^{1,3}$ present secondary (derivative) construction with respect to twistors.

In fact, twistors can be defined as “reduced spinors” of pseudounitary group $\text{SO}_0(2, 4)$ acting in a six-dimensional space with the signature $(+, +, -, -, -, -)$. These reduced spinors are derived as follows. General spinors are elements of minimal left ideal of a *conformal algebra* $\mathcal{C}_{2,4}$:

$$I_{2,4} = \mathcal{C}_{2,4} f_{24} = \mathcal{C}_{2,4} \frac{1}{2} (1 + \mathbf{e}_{15}) \frac{1}{2} (1 + \mathbf{e}_{26}).$$

Reduced spinors (twistors) are formulated within an even subalgebra $\mathcal{C}_{2,4}^+ \simeq \mathcal{C}_{4,1}$ (*de Sitter algebra*). The minimal left ideal of the algebra $\mathcal{C}_{4,1} \simeq \mathbb{C}_4$ is defined by the following expression [29]:

$$I_{4,1} = \mathcal{C}_{4,1} f_{4,1} = \mathcal{C}_{4,1} \frac{1}{2} (1 + \mathbf{e}_0) \frac{1}{2} (1 + i\mathbf{e}_{12}).$$

Therefore, after reduction $I_{2,4} \rightarrow I_{4,1}$, generated by the isomorphism $\mathcal{C}_{2,4}^+ \simeq \mathcal{C}_{4,1}$, we see that twistors \mathbf{Z}^α are elements of the ideal $I_{4,1}$ which leads to the group $\text{SU}(2, 2) \simeq \mathbf{Spin}_+(2, 4) \in \mathcal{C}_{2,4}^+$ (see further (C.2) and (C.3)). Indeed, let us consider the algebra $\mathcal{C}_{2,4}$ associated with a six-dimensional pseudoeuclidean space $\mathbb{R}^{2,4}$. A double covering $\mathbf{Spin}_+(2, 4)$ of the rotation group $\text{SO}_0(2, 4)$ of the space $\mathbb{R}^{2,4}$ is described within the even subalgebra $\mathcal{C}_{2,4}^+$. The algebra $\mathcal{C}_{2,4}$ has the type $p - q \equiv 6 \pmod{8}$, therefore, according to $\mathcal{C}_{p,q}^+ \simeq \mathcal{C}_{q,p-1}$ we have $\mathcal{C}_{2,4}^+ \simeq \mathcal{C}_{4,1}$, where $\mathcal{C}_{4,1}$ is the de Sitter algebra associated with the space $\mathbb{R}^{4,1}$. In its turn, the algebra $\mathcal{C}_{4,1}$ has the type $p - q \equiv 3 \pmod{8}$ and, therefore, there is an isomorphism $\mathcal{C}_{4,1} \simeq \mathbb{C}_4$, where \mathbb{C}_4 is a *Dirac algebra*. The algebra \mathbb{C}_4 is a complexification of the space-time algebra: $\mathbb{C}_4 \simeq \mathbb{C} \otimes \mathcal{C}_{1,3}$. Further, $\mathcal{C}_{1,3}$ admits the following factorization: $\mathcal{C}_{1,3} \simeq \mathcal{C}_{1,1} \otimes \mathcal{C}_{0,2}$. Hence it immediately follows that $\mathbb{C}_4 \simeq \mathbb{C} \otimes \mathcal{C}_{1,1} \otimes \mathcal{C}_{0,2}$. Thus,

$$\mathbf{Spin}_+(2, 4) = \{s \in \mathbb{C} \otimes \mathcal{C}_{1,1} \otimes \mathcal{C}_{0,2} \mid N(s) = 1\}. \quad (\text{C.2})$$

On the other hand, in virtue of $\mathcal{C}_{1,3} \simeq \mathcal{C}_{1,1} \otimes \mathcal{C}_{0,2}$ a general element of the algebra $\mathcal{C}_{1,3}$ can be written in the form

$$\mathcal{A}_{\mathcal{C}_{1,3}} = \mathcal{C}_{1,1}^0 \mathbf{e}_0 + \mathcal{C}_{1,1}^1 \phi + \mathcal{C}_{1,1}^2 \psi + \mathcal{C}_{1,1}^3 \phi\psi,$$

where $\phi = \mathbf{e}_{123}$, $\psi = \mathbf{e}_{124}$ are quaternion units. Therefore,

$$\mathbf{Spin}_+(2, 4) = \left\{ s \in \begin{bmatrix} \mathbb{C} \otimes \mathcal{C}_{1,1}^0 - i\mathbb{C} \otimes \mathcal{C}_{1,1}^3 & -\mathbb{C} \otimes \mathcal{C}_{1,1}^1 + i\mathbb{C} \otimes \mathcal{C}_{1,1}^2 \\ \mathbb{C} \otimes \mathcal{C}_{1,1}^1 + i\mathbb{C} \otimes \mathcal{C}_{1,1}^2 & \mathbb{C} \otimes \mathcal{C}_{1,1}^0 + i\mathbb{C} \otimes \mathcal{C}_{1,1}^3 \end{bmatrix} \mid N(s) = 1 \right\}. \quad (\text{C.3})$$

Mappings of the space $\mathbb{R}^{1,3}$, generated by the group $\text{SO}_0(2, 4)$, induce linear transformations of the twistor space \mathbb{C}^4 with preservation of the form $\mathbf{Z}^\alpha \bar{\mathbf{Z}}_\alpha$ of the signature $(+, +, -, -)$. Hence it follows that a corresponding group in the twistor space is $\text{SU}(2, 2)$ (the group of pseudo-unitary unimodular 4×4 matrices, see (C.3)):

$$\text{SU}(2, 2) = \left\{ \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{C}_4 : \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = 1 \right\} \simeq \mathbf{Spin}_+(2, 4). \quad (\text{C.4})$$

References

- [1] Korableva, T. P., Korolkov, D. V.: Theory of the Periodic System. Saint-Petersburg University Press, Saint-Petersburg (2005) [in Russian].
- [2] Fock, V. A.: *Do the chemical properties of atoms fit into the framework of purely spatial representations?* The Periodic Law and the Structure of the Atom. Atomizdat, Moscow (1971), P. 107–117 [in Russian].
- [3] Gelfand, I., Neumark, M.: *On the Imbedding of Normed Rings into the Ring of Operators in Hilbert Space.* Rec. Math. [Mat. Sbornik] N.S. **12**(54), 197–217 (1943).
- [4] Segal, I.: *Postulates for general quantum mechanics.* Ann. Math. **48**, 930–948 (1947).
- [5] Varlamov, V. V.: *Lorentz Group and Mass Spectrum of Elementary Particles.* arXiv: 1705.02227 [physics.gen-ph] (2017).
- [6] Varlamov, V. V.: *Fundamental Automorphisms of Clifford Algebras and an Extension of Dqbrowski Pin Groups.* Hadronic J. **22**, 497–535 (1999).
- [7] Varlamov, V. V.: *Discrete Symmetries and Clifford Algebras.* Int. J. Theor. Phys. **40**, 4, 769–805 (2001); arXiv:math-ph/0009026 (2000).
- [8] Varlamov, V. V.: *Discrete symmetries on the spaces of quotient representations of the Lorentz group.* Mathematical structures and modeling. **7**, 115–128 (2001).
- [9] Varlamov, V. V.: *CPT groups for spinor field in de Sitter space.* Phys. Lett. B. **631**, 187–191 (2005); arXiv:math-ph/0508050 (2005).
- [10] Varlamov, V. V.: *CPT Groups of Higher Spin Fields.* Int. J. Theor. Phys. **51**, 1453–1481 (2012); arXiv: 1107.4156 [math-ph] (2011).
- [11] Varlamov, V. V.: *Spinor Structure and Internal Symmetries.* Int. J. Theor. Phys. **54**, 10, 3533–3576 (2015); arXiv:1409.1400 [math-ph] (2014).
- [12] Heisenberg, W.: Introduction to the Unified Field Theory of Elementary Particles. Interscience Publishers, London, New York, Sydney (1966).
- [13] Segal, I.: *A class of operator algebras which are determined by groups.* Duke Math. J. **18**, 221–265 (1951).
- [14] Rumer, Yu. B., Fet, A. I.: *Spin(4) group and the Mendeleev system.* Teor. Mat. Fiz. **9**, 203–209 (1971).
- [15] Fock, V. A.: *Hydrogen atom and non-Euclidean geometry.* Izv. AN SSSR. Ser. VII. **2**, 169–179 (1935).
- [16] Konopelchenko, B. G.: *Group $SO(2, 4)+R$ and Mendeleev system.* SO RAN. Institute of Nuclear Physics. Prepr. Novosibirsk (1972).
- [17] Kibler, M. R.: *From the Mendeleev periodic table to particle physics and back to periodic table.* Foundations of Chemistry. **9**, 221–234 (2006).
- [18] Fet, A. I.: Symmetry group of chemical elements. Novosibirsk, Nauka (2010) [in Russian].

- [19] Yao Tsu: *Unitary Irreducible Representations of $SU(2,2)$* . J. Math. Phys. **8**, 1931–1954 (1967).
- [20] Madelung, E.: *Die Mathematischen Hilfsmittel des Physikers*. Springer-Verlag, Berlin (1957).
- [21] Kleczkowski, V.M.: Atomic electron distribution and the rule of sequential filling of $(n + l)$ -groups. Atomizdat, Moscow (1968) [in Russian].
- [22] Gell-Mann, M.: *Symmetries of Baryons and Mesons*. Phys. Rev. **125**, 1067–1084 (1962).
- [23] Okubo, S., Ryan, C.: *Quadratic mass formula in $SU(3)$* . Nuovo Cimento **34**, 776–779 (1964).
- [24] Bég, M., Singh, V.: *Splitting of the 70-Plet of $SU(6)$* . Phys. Rev. Lett. **13**, 509–511 (1964).
- [25] Emch, G.G.: *Algebraic methods in statistical mechanics and quantum field theory*. Wiley-Interscience, New York (1972).
- [26] Bogoljubov, N.N., Logunov, A.A., Oksak, A.I., Todorov, I.T.: *General Principles of Quantum Field Theory*. Nauka, Moscow (1987) [in Russian].
- [27] Horuzhy, S.S.: *Introduction to Algebraic Quantum Field Theory*. Nauka, Moscow (1986) [in Russian].
- [28] Lounesto, P.: *Clifford Algebras and Spinors*. Cambridge Univ. Press, Cambridge (2001).
- [29] Varlamov, V.V.: *Generalized Weierstrass representation for surfaces in terms of Dirac-Hestenes spinor field*. J. Geometry and Physics. 2000. **32**, 241–251 (2000).
- [30] Varlamov, V.V.: *Spinor Structure and Periodicity of Clifford Algebras*. Mathematical structures and modeling. **35**, 4–20 (2015).
- [31] Heisenberg, W.: *Schritte über Grenzen*. München (1977).
- [32] Fet, A.I.: *Conformal symmetry of chemical elements*. Teor. Mat. Fiz. **22**, 323–334 (1975).
- [33] Heisenberg, W.: *Der Teil und das Ganze*. München (1969).
- [34] Varlamov, V.V.: *Spinor Structure and $SU(3)$ -Symmetry*. Mathematical structures and modeling. **39**, 5–23 (2016).
- [35] Varlamov, V.V.: *Heisenberg Matter Spectrum in Abstract Algebraic Approach*. Mathematical structures and modeling. **33**, 18–33 (2015).
- [36] Wigner, E.P.: *On unitary representations of the inhomogeneous Lorentz group*. Ann. Math. **40**, 149–204 (1939).
- [37] Varlamov, V.V.: *CPT groups of spinor fields in de Sitter and anti-de Sitter spaces*. Adv. Appl. Clifford Algebras. **25**, 487–516 (2015); arXiv: 1401.7723 [math-ph] (2014).
- [38] Seaborg, G.: *Expansion of the limits of the periodic system*. One hundred years of the periodic law of the chemical elements. Nauka, Moscow (1971), P. 21–39 [in Russian].
- [39] Greiner, W., Reinhardt, J.: *Quantum Electrodynamics*. Springer, Berlin (2009).
- [40] Wick, G.G., Wigner, E.P., Wightman, A.S.: *Intrinsic Parity of Elementary Particles*. Phys. Rev. **88**, 101 (1952).

- [41] Varlamov, V.V.: *Relativistic wavefunctions on the Poincaré group*. J. Phys. A: Math. Gen. **37**, 5467–5476 (2004); arXiv:math-ph/0308038 (2003).
- [42] Varlamov, V.V.: *Relativistic spherical functions on the Lorentz group*. J. Phys. A: Math. Gen. **39**, 805–822 (2006).
- [43] Varlamov, V.V.: *Spherical functions on the de Sitter group*. J. Phys. A: Math. Theor. **40**, 163–201 (2007).
- [44] Varlamov, V.V.: *Universal Coverings of Orthogonal Groups*. Adv. Appl. Clifford Algebras. **14**, 81–168 (2004); arXiv:math-ph/0405040 (2004).
- [45] Rashevskii, P.K.: *The Theory of Spinors*. Uspekhi Mat. Nauk. **10**, 3–110 (1955); English translation in Amer. Math. Soc. Transl. (Ser. 2). **6**, 1 (1957).
- [46] Baez, J.C.: *Division Algebras and Quantum Mechanics*. Found. Phys. **42**, 819–855 (2012).
- [47] Ryder, L.: *Quantum Field Theory*. Cambridge University Press, Cambridge (1985).
- [48] Ahluwalia, D. V., Ernst, D. J.: $(j, 0) \oplus (0, j)$ covariant spinors and causal propagators based on Weinberg formalism. Int. J. Mod. Phys. E. **2**, 397–422 (1993).
- [49] Dvoeglazov, V. V.: *Extra Dirac equations*. Nuovo Cimento B. **111**, 483–496 (1996).
- [50] Knapp, A.W. Representation Theory of Semisimple Groups. Princeton University Press, Princeton (1986).
- [51] Gelfand, I. M., Minlos, R. A., Shapiro, Z. Ya.: Representations of the Rotation and Lorentz Groups and their Applications. Pergamon Press, Oxford (1963).
- [52] Naimark, M. A.: Linear Representations of the Lorentz Group. Pergamon, London (1964).
- [53] Akhiezer, A. I., Berestetskii, V. B.: Quantum Electrodynamics. John Wiley & Sons, New York (1965).
- [54] Schweber, S. S.: An Introduction to Relativistic Quantum Field Theory. Harper & Row, New York (1961).
- [55] Rumer, Yu. B., Fet, A. I.: Group Theory and Quantized Fields. — Nauka, Moscow (1977) [in Russian].
- [56] van der Waerden, B. L.: Die Gruppentheoretische Methode in der Quantenmechanik. Springer, Berlin (1932).
- [57] Penrose, R.: *The twistor programme*. Rep. Math. Phys. **12**, 65–76 (1977).
- [58] Penrose, R., MacCallum, M. A. H.: *Twistor theory: an approach to the quantization of fields and space-time*. Physics Reports. **6**, 241–316 (1972).