

# The Batchelor spectrum of passive scalar turbulence in stochastic fluid mechanics

Jacob Bedrossian\*

Alex Blumenthal†

Sam Punshon-Smith‡

November 26, 2019

## Abstract

In 1959, Batchelor predicted that passive scalars advected in fluids at finite Reynolds number with small diffusivity  $\kappa$  should display a  $|k|^{-1}$  power spectrum over a small-scale inertial range in a statistically stationary experiment. This prediction has been experimentally and numerically tested extensively in the physics and engineering literature and is a core prediction of *passive scalar turbulence*.

In this article we provide the first mathematically rigorous proof of Batchelor’s prediction on the cumulative power spectrum in the  $\kappa \rightarrow 0$  limit. We consider fluids governed by the 2D Navier-Stokes equations and 3D hyperviscous Navier-Stokes equations in  $\mathbb{T}^d$  forced by sufficiently regular, nondegenerate stochastic forcing at fixed (arbitrary) Reynolds number. The scalar is subjected to a smooth-in-space, white-in-time stochastic source, and evolves by advection-diffusion with diffusivity  $\kappa > 0$ . Our results rely on the quantitative understanding of Lagrangian chaos and passive scalar mixing established in our recent works. In the  $\kappa \rightarrow 0$  limit, we obtain statistically stationary, weak solutions in  $H^{-\epsilon}$  to the stochastically-forced advection problem *without diffusivity*. These solutions are almost-surely not locally integrable functions, have a non-vanishing average anomalous flux, and satisfy the Batchelor spectrum at all sufficiently small scales. We also prove an Onsager-type criticality result which shows that no such dissipative, weak solutions with a little more regularity can exist.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Passive scalar turbulence . . . . .	3
1.2	Main results . . . . .	5
1.3	A simple physical mechanism for Batchelor’s Law . . . . .	12
1.4	Open problems in Batchelor-regime passive scalar turbulence . . . . .	14
<b>2</b>	<b>Preliminaries</b>	<b>16</b>
2.1	Fluid models . . . . .	16
2.2	$C_t^k C_x^\infty$ fluid models governed by finite dimensional SDE . . . . .	17
2.3	Lyapunov functions and exponential estimates . . . . .	19
2.4	Uniform mixing and enhanced dissipation . . . . .	20
2.5	Unique stationary measure for $(u_t, g_t^k)$ . . . . .	22

\*Department of Mathematics, University of Maryland, College Park, MD 20742, USA [jacob@math.umd.edu](mailto:jacob@math.umd.edu). J.B. was supported by NSF CAREER grant DMS-1552826 and NSF RNMS #1107444 (Ki-Net)

†Department of Mathematics, University of Maryland, College Park, MD 20742, USA [alex123@math.umd.edu](mailto:alex123@math.umd.edu). This material was based upon work supported by the National Science Foundation under Award No. DMS-1604805.

‡Division of Applied Mathematics, Brown University, Providence, RI 02906, USA [punshs@brown.edu](mailto:punshs@brown.edu). This material was based upon work supported by the National Science Foundation under Award No. DMS-1803481.

<b>3</b>	<b>Proof of Batchelor’s law</b>	<b>23</b>
3.1	Upper bound in the inertial range . . . . .	23
3.2	Lower bound in the inertial range . . . . .	25
3.3	Upper bound in the dissipative range . . . . .	27
<b>4</b>	<b>Vanishing diffusivity limit</b>	<b>29</b>
4.1	Existence of weak limits of $\{\mu^\kappa\}$ . . . . .	29
4.2	The scalar process $(u_t, g_t^0)$ at $\kappa = 0$ . . . . .	30
4.3	Stationarity of $\mu^0$ : restricted Feller property of $(u_t, g_t^0)$ . . . . .	31
4.4	Stationarity of $\mu^0$ : completing the proof . . . . .	33
4.5	Non-vanishing flux . . . . .	35
<b>5</b>	<b>Irregularity of the limiting statistically stationary solutions</b>	<b>35</b>
5.1	$\mu^0$ generic functions cannot be $L^1$ . . . . .	35
5.2	Onsager-type criticality of $L_t^2 B_{2,\infty}^0$ . . . . .	38

# 1 Introduction

In physics, turbulence usually refers to the observed phenomena in dynamics of weakly-damped, infinite-dimensional, nonlinear conservative systems in a “generic” setting. The most canonical example is the 3D Navier-Stokes equations in the high Reynolds number limit [68], studied by Kolmogorov in his foundational 1941 works [88–90]. Since then, analogous dynamics have been studied in a wide range of physical systems: 2D Navier-Stokes [26], passive scalars advected by fluids [123, 126], compressible fluids [97], magneto-hydrodynamics [22], nonlinear dispersive equations [106, 135], and both fully kinetic and hybrid-kinetic models of plasmas (see e.g. [74] and the references therein).

A setting which captures a large proportion of phenomena in both nature and experiments is the *statistically stationary case*<sup>1</sup>, wherein fluctuations are injected at a fixed characteristic scale (the *integral scale*) and the experiment is run long enough to measure ensemble statistics in equilibrium. Many general settings can be reasonably approximated as statistically stationary, as done originally by Kolmogorov and in many studies since then; see [68, 106, 135] for discussions.

In all of the above examples, there are a few basic, common themes that are observed:

- (a) **Unique long-time statistics:** Time averages are observed to converge to ensemble averages that do not depend on initial condition.
- (b) **Chaos:** Experiments are observed to be chaotic in the sense of having a positive Lyapunov exponent, i.e. exponential divergence of nearby states in phase space.
- (c) **Anomalous dissipation:** As the damping/dissipation vanishes, one or more ranges of length-scales form (called *inertial range(s)*) wherein conserved quantities undergo a constant scale-by-scale flux extending from the integral scale to dissipative scale(s), where the quantities are subsequently dissipated at a constant rate independent of the dissipative parameter(s).
- (d) **Universality in the inertial range:** Over the inertial ranges, certain statistics are observed to be essentially universal, in the sense that they appear to be independent of the details of the experiment.

In much of the physics and engineering literature, these properties are usually, to varying degrees, taken for granted, though we are lacking a predictive theory from first principles (see e.g. [28, 68]). Similarly, we are lacking satisfactory theories for the inertial-range statistics. Formal arguments often yield predictions

---

<sup>1</sup>A stochastic process  $u(t)$  is called statistically stationary if the law of the paths  $u(\cdot)$  and  $u(\cdot + \tau)$  are equal for any  $\tau > 0$ .

that do not agree with experiments, for example, the famous Kolmogorov  $-5/3$  spectrum of 3D Navier-Stokes (see e.g. [3, 68] and the references therein) and the Kolmogorov-Zakharov spectra of wave turbulence (see e.g. [134] and the references therein).

It has been known for some time that turbulence is deeply connected with the theory of dynamical systems; see [28] and the references therein. For systems subjected to deterministic forcing, one expects that the statistically stationary regime should be described by a unique, generically observable, invariant measure on which the dynamics are chaotic, also known as an SRB measure [57, 132]. Although there is by now a well-developed abstract theory of SRB measures (see e.g. [12, 96, 114]), even for systems posed in infinite-dimensions (see e.g. [24, 25, 116]), it is notoriously challenging to prove the existence and uniqueness of SRB measures in practice, even for very low-dimensional systems, and so this problem seems very far out of reach for now— see, e.g., the discussion in [112, 131].

Nevertheless, varied progress has been made, for instance: in bounding dynamical complexity from above, in the sense of upper bounds on attractor dimension and closely related upper bounds on determining modes (see e.g. [36, 37, 40, 69, 84, 120]); estimation of possible power spectra (see e.g. [38, 41, 111, 119]); characterization of sufficient conditions for predictions relating to turbulence (see e.g. [34, 46, 52, 55, 107]); and recent progress on wave turbulence [31, 32]. Another place where significant progress has been made is the work surrounding *Onsager's conjecture* [33, 35, 39, 47, 60, 80].

In the presence of stochastic forcing, conditions for existence, uniqueness, and geometric ergodicity of stationary measures of the associated Markov process are now well-understood in many settings: see e.g. [66, 71, 75–77, 92, 94]. However, we are mostly lacking major progress on properties (b)–(d) for real physical systems. Some progress has been made: see [20, 21, 64] for conditional theorems and [67] for a study carrying out (a), (c), and (d) for a highly idealized shell model. For the remainder of this work, we will consider only stochastically-forced, statistically stationary turbulence.

## 1.1 Passive scalar turbulence

Let  $u_t(x)$  be a velocity field evolving according to a fluid evolution equation in the  $d$ -dimensional periodic box  $\mathbb{T}^d = [0, 2\pi]^d$ , e.g. Navier-Stokes. *Passive scalar turbulence* refers to the dynamics of a passive tracer  $g_t$  undergoing advection-diffusion in the fluid described by  $(u_t)$  as the diffusivity  $\kappa > 0$  is taken to zero. See e.g. [62, 108, 123, 126] for more discussions of the physical background. As discussed above, we consider the statistically stationary setting, where the fluid is stochastically forced and the scalar has a stochastic source. In  $d = 2$  we will consider:

$$\begin{aligned} \partial_t u_t + u_t \cdot \nabla u_t + \nabla p_t - \nu \Delta u_t &= Q\dot{W}_t \\ \partial_t g_t^\kappa + u_t \cdot \nabla g_t^\kappa - \kappa \Delta g_t^\kappa &= b\dot{\xi}_t. \end{aligned} \tag{1.1}$$

Here,  $Q\dot{W}_t$  is a white-in-time, sufficiently-regular-in-space, Gaussian process (see Section 2 for details),  $b \in C^\infty(\mathbb{T}^d)$ , and  $\xi_t$  is an independent 1D Brownian motion. Our methods apply equally well when the random source is given by  $\sum_{j=1}^\infty b_j \xi_t^j$  for a sequence of smooth,  $L^2$  orthogonal  $b_j$ 's that decay sufficiently fast in  $j$  and independent Brownian motions  $\xi_t^j$ . For simplicity we will consider the case  $x \in \mathbb{T}^d$ ; see Section 1.4 for discussion. In the statistically stationary regime, Itô's formula implies an on-average constant dissipation of  $L^2$  density:

$$\kappa \mathbf{E} \|\nabla g^\kappa\|_{L^2}^2 = \frac{1}{2} \|b\|_{L^2}^2 =: \chi.$$

See Proposition 1.1 for mathematical details.

As  $\kappa \rightarrow 0$ , physicists observe a number of different regimes, depending on the rate of growth or decay of the Schmidt number  $\text{Sc} := \nu/\kappa$ . In particular, if  $\nu, \kappa \rightarrow 0$  simultaneously, then the fluid itself is becoming turbulent. Essentially nothing has been established mathematically for the (very singular) turbulent limit

$\nu \rightarrow 0$  for statistically stationary solutions  $(u_t)$  to the Navier-Stokes equations, and this appears to be out of reach of mathematical analysis for the time being. One setting where this kind of ‘turbulent advection’ in (1.1) has been studied is in the *Kraichnan model*, in which  $(u_t)$  is replaced by an idealized rough-in-space, white-in-time Gaussian velocity field, and the equation for  $(g_t)$  is interpreted as a Stratonovich stochastic transport equation. This model was introduced for this purpose in [91] and there is now a wide literature on this model in physics [43, 44, 123] and the references therein. Properties of stochastic transport with rough velocities have also been studied mathematically as well in the work on isotropic stochastic flows [14, 83, 100] and well-posedness by stochastic perturbation [63, 65, 104]. See also the recent preprint [53] for a deterministic work on turbulent advection by rough velocities.

In this paper, we are concerned with the so-called *Batchelor regime*: where  $\nu$  is considered fixed and  $\kappa \rightarrow 0$ . In his foundational paper [13], Batchelor predicted that for systems of type (1.1) the ensemble statistics for a passive tracer in the inertial range exhibits a  $|k|^{-1}$  *power spectrum*:

$$|k|^{d-1} \mathbf{E} |\hat{g}(k)|^2 \approx \chi |k|^{-1} \quad \text{for} \quad 1 \ll |k| \lesssim \kappa^{-1/2}, \quad (1.2)$$

where  $\hat{g}(k)$ ,  $k \in \mathbb{Z}^d$  denotes the Fourier transform of  $g$ . This prediction is now called *Batchelor’s law* or the *Batchelor spectrum*. Batchelor made this prediction in the general case of  $Sc \rightarrow \infty$ , which is different from fixing  $\nu = O(1)$  and sending  $\kappa \rightarrow 0$ . In the former regime, for length-scales larger than the dissipative range of the Navier-Stokes equations, one expects hydrodynamic turbulence to dominate and the passive scalar is observed to follow Obukhov-Corrsin statistics [42, 109]; see e.g. [4, 123]. See Figure 1 and the associated caption for a description of these two different regimes.




Figure 1: The dotted curve denotes the predictions for the power spectrum  $E(k) = |k|^{d-1} \mathbf{E} |\hat{u}(k)|^2$  of the fluid and the solid curve denotes the predictions for the power spectrum of the passive scalar  $\Gamma(k) = |k|^{d-1} \mathbf{E} |\hat{g}(k)|^2$ , viewed on a log-log plot. The Batchelor regime when  $\kappa \ll \nu$  is shown in the range between  $\ell_D^{NSE}$  (the dissipative scale for Navier-Stokes) and  $\kappa^{1/2}$  (the dissipative scale for Batchelor’s regime), where it is theorized that Batchelor’s  $|k|^{-1}$  law holds. Between  $\ell_I$  and  $\ell_D^{NSE}$  one expects Obukhov-Corrsin statistics, e.g., the  $|k|^{-5/3}$  law in dimension  $d = 3$ . No universality is expected at length-scales in the integral range above  $\ell_I$ . We emphasize that in the setting of the present manuscript,  $\nu$  is fixed  $O(1)$  and  $\kappa \ll 0$ . In this regime, the scale  $\ell_I$  is basically the same order as  $\ell_D^{NSE}$  so that the Obukhov-Corrsin regime is negligible and indistinguishable from the integral range.

Since Batchelor’s predictions, engineers and physicists have been making measurements of the Batchelor spectrum in nature, e.g., temperature and salinity variations in the ocean (see e.g. [49, 73, 108] and the

references therein), in laboratory experiments (see e.g. [2, 70, 85, 102, 127, 128] and the references therein), and in numerical studies (see e.g. [4, 27, 51, 78, 133] and the references therein). As discussed in e.g. [4, 51] it remains an open problem in physics to determine the settings in which the Batchelor spectrum is expected and to what degree of accuracy it holds. We will briefly discuss some theoretical studies in the physics literature in Section 1.3.

## 1.2 Main results

In all our results we will consider our domain to be the  $d$ -dimensional periodic box  $\mathbb{T}^d = [0, 2\pi]^d$  for  $d = \{2, 3\}$ . The velocity field  $(u_t)$  will take values in the space  $\mathbf{H}$  consisting mean zero divergence free velocity fields belongs to  $H^\sigma = H^\sigma(\mathbb{T}^d; \mathbb{R}^d)$ , the Sobolev space of mappings from  $\mathbb{T}^d \rightarrow \mathbb{R}^d$  with regularity  $\sigma > \frac{d}{2} + 3$ ; note that this implies velocities are always at least  $C^3$  in space.

The *velocity process*  $(u_t)$  will evolve in  $\mathbf{H}$  according to one of the following stochastic PDEs depending on whether  $d = 2$  or 3:

**System 1** (2D Navier-Stokes equations). When  $d = 2$ ,  $(u_t)$  solves

$$\begin{aligned} \partial_t u_t + u_t \cdot \nabla u_t &= -\nabla p_t + \nu \Delta u_t + Q \dot{W}_t \\ \operatorname{div} u_t &= 0, \end{aligned}$$

where  $u_0 = u \in \mathbf{H}$ . Here, the viscosity  $\nu > 0$  is a fixed constant.

**System 2** (3D hyper-viscous Navier-Stokes). When  $d = 3$ ,  $(u_t)$  solves

$$\begin{aligned} \partial_t u_t + u_t \cdot \nabla u_t &= -\nabla p_t + \nu' \Delta u_t - \nu \Delta^2 u_t + Q \dot{W}_t \\ \operatorname{div} u_t &= 0, \end{aligned}$$

where  $u_0 = u \in \mathbf{H}$ . Here, the viscosity  $\nu' \geq 0$  and hyperviscosity  $\nu > 0$  are fixed constants.

In the above systems,  $W_t$  is a cylindrical Wiener process on mean-zero, divergence free  $L^2$  vector fields with respect to an associated canonical stochastic basis  $(\Omega_W, \mathcal{F}^W, (\mathcal{F}_t^W), \mathbf{P}_W)$  and  $Q$  a positive Hilbert-Schmidt operator on mean-zero, divergence free  $L^2$  vector fields satisfying suitable non-degeneracy and regularity assumptions. See Section 2.1 for full details. We couple systems 1 or 2 to the advection-diffusion equation

$$\begin{aligned} \partial_t g_t^\kappa + u_t \cdot \nabla g_t^\kappa &= \kappa \Delta g_t^\kappa + b \dot{\xi}_t \\ g_0^\kappa &= g. \end{aligned} \tag{1.3}$$

where  $\xi_t$  is a Wiener process on another canonical probability space  $(\Omega_\xi, \mathcal{F}^\xi, (\mathcal{F}_t^\xi), \mathbf{P}_\xi)$ . We denote the product measure  $\mathbf{P} = \mathbf{P}_W \times \mathbf{P}_\xi$  our main probability measure on the associated product space  $\Omega = \Omega_W \times \Omega_\xi$  with standard product sigma-algebra  $\mathcal{F} = \mathcal{F}^W \otimes \mathcal{F}^\xi$  and filtration  $\mathcal{F}_t = \mathcal{F}_t^W \otimes \mathcal{F}_t^\xi$ . Equation (1.3) has a  $\mathbf{P}$  almost-sure, unique,  $\mathcal{F}_t$ -adapted weak solution for every initial  $(u, g) \in \mathbf{H} \times L^2$  (see Proposition 2.1 below) and defines a Markov semigroup  $P_t^\kappa$  on bounded, measurable observables  $\varphi : \mathbf{H} \times L^2 \rightarrow \mathbb{R}$  via

$$P_t^\kappa \varphi(u, g) = \mathbf{E}_{(u, g)} \varphi(u_t, g_t^\kappa) := \mathbf{E}[\varphi(u_t, g_t^\kappa) | (u_0, g_0^\kappa) = (u, g)].$$

### 1.2.1 The Batchelor spectrum

Recall, a probability measure  $\mu$  on  $\mathbf{H} \times L^2$  is called *stationary* for the Markov semigroup  $P_t^\kappa$  if for all bounded, measurable  $\varphi : \mathbf{H} \times L^2 \rightarrow \mathbb{R}$  and all  $t > 0$ ,

$$\int P_t^\kappa \varphi \, d\mu = \int \varphi \, d\mu.$$

For observables  $\varphi : \mathbf{H} \times L^2 \rightarrow \mathbb{R}$ , we frequently write  $\mathbf{E}_\mu \varphi(u, g) := \int \varphi d\mu$ . We will also say a probability measure is stationary for a given process if it is stationary for the corresponding Markov semi-group.

Due to the infinite-dimensionality of  $\mathbf{H}$ , uniqueness of stationary measures for  $(u_t)$  is in general a subtle question: on the domain  $\mathbb{T}^2$  this was first proved in [66] in the setting of completely nondegenerate noise, and has by now been established even for highly degenerate noise [75]. In comparison, the homogeneous part of the evolution equation for  $g_t^\kappa$  is linear, and so uniqueness of stationary measures for  $(u_t, g_t^\kappa)$  is relatively straightforward due to the contracting nature on  $L^2$ .

**Proposition 1.1.** *Assume  $(u_t)$  admits a unique stationary probability measure  $\mu$  on  $\mathbf{H}$ . For all  $\kappa > 0$ , there exists a unique stationary measure  $\mu^\kappa$  for  $(u_t, g_t^\kappa)$  on  $\mathbf{H} \times L^2$ . Moreover,*

$$\kappa \mathbf{E}_{\mu^\kappa} \|\nabla g\|_{L^2}^2 = \frac{1}{2} \|b\|_{L^2}^2 =: \chi. \quad (1.4)$$

Uniqueness of  $\mu^\kappa$  is proved in Section 2.5, and has the following consequence by the Birkhoff ergodic theorem: for  $\mu^\kappa$ -typical initial  $(u, g) \in \mathbf{H} \times L^2$  and any continuous observable  $\phi \in L^1(\mu^\kappa)$  on  $\mathbf{H} \times L^2$ , we have  $\mathbf{P}$ -almost-surely<sup>2</sup>

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u_t, g_t^\kappa) dt = \int \phi d\mu^\kappa. \quad (1.5)$$

Equation (1.4) follows from (1.5) and Itô's formula. The convergence of time averages to ensemble averages in (1.5) for typical initial data  $(u, g)$  confirms the statistically stationary setting described above: typical time-asymptotic behavior is captured by a unique stationary measure  $\mu^\kappa$ .

Our main result is Batchelor's law on the *cumulative* power spectrum (see Remark 1.8). Let  $\Pi_{\leq N}$  denote the  $L^2$  projection to the Fourier basis functions with frequency  $|k| \leq N$  (see Section 2.1).

**Theorem 1.2** (Batchelor's law on the cumulative power spectrum). *There exists an  $N_0$  (independent of  $\kappa$ ) such that for all  $\kappa \in (0, 1)$  sufficiently small and  $p \in [1, \infty)$ , the following holds with implicit constants<sup>3</sup> independent of  $\kappa$  and  $N$ :*

$$\left( \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \approx_p \log N \quad \text{for all } N_0 \leq N \leq \kappa^{-1/2}. \quad (1.6)$$

Moreover, for all  $s \in [0, \sigma - \frac{3d}{2} - 1)$  and  $\forall p \in [1, \infty)$ ,

$$\left( \mathbf{E}_{\mu^\kappa} \|g\|_{H^s}^{2p} \right)^{1/p} \lesssim_{p,s} \kappa^{-s} |\log \kappa|. \quad (1.7)$$

**Remark 1.3.** Batchelor's law is often stated with a constant proportional to  $\chi = \kappa \mathbf{E} \|\nabla g^\kappa\|_{L^2}^2 = \frac{1}{2} \|b\|_{L^2}^2$ , as in (1.2). A careful reading of our proof provides a simple estimate on the constants in (1.6) in terms of  $b$ , specifically,  $\forall s > 0$ , the following holds with implicit constants independent of  $b$ :

$$\log N \frac{\chi^2}{\|b\|_{H^s}^2} \lesssim_{p,s} \left( \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \lesssim_{p,s} \|b\|_{H^s}^2 \log N \quad \text{for all } N_0 \leq N \leq \kappa^{-1/2}, \quad (1.8)$$

<sup>2</sup>When  $(u_t)$  is forced with nondegenerate noise, as we do in this paper, it is possible to promote the convergence in (1.5) to *all* initial  $(u, g) \in \mathbf{H} \times L^2$  when one considers bounded, uniformly continuous observables  $\varphi : \mathbf{H} \times L^2 \rightarrow \mathbb{R}$ . This follows from (1) the strong Feller property for the Markov semigroup associated to  $u_t$  (see [66]) and (2) a small variation of the proof of Proposition 1.1 given in Section 2.5. Details are omitted for brevity.

<sup>3</sup>We denote  $f \lesssim_{p,q,\dots} h$  if there exists a constant  $C > 0$  depending on  $p, q, \dots$  but independent of the other parameters of interest such that  $f \leq Ch$  and  $f \approx h$  when  $f \lesssim h$  and  $h \lesssim f$ . For the entire paper, these implicit constants will *never* depend on  $\kappa, N$  or  $t$ .

and that it suffices to take

$$N_0 \approx_{s,\delta} \left( \frac{\|b\|_{H^s}^{2/s}}{\chi^{1/s}} \right)^{1+\delta}$$

for any  $\delta > 0$  (the implicit constants in (1.8) will also depend on  $\delta$ ). The use of regularity is because mixing estimates require the source to have *some* positive regularity in order to get quantitative,  $\kappa$ -independent decay rates in negative Sobolev spaces. It is unclear if Batchelor's law as stated in Theorem 1.2 is still true if  $b$  has no more than  $L^2$  regularity. If  $b$  takes values in any  $\Lambda_C = \{b \in H^s : C^{-1} \|b\|_{L^2} \leq \|b\|_{H^s} \leq C \|b\|_{L^2}\}$ , then both implicit constants in (1.6) are proportional to  $\chi$  and  $N_0$  depends only on  $C$ .

Theorem 1.2 implies the following uniform in  $\kappa$  estimate:

**Corollary 1.4.** *For all  $s > 0$  and  $p \geq 1$  the following holds*

$$\sup_{\kappa} \mathbf{E}_{\mu^\kappa} \|g\|_{H^{-s}}^{2p} \lesssim_{p,s} 1. \quad (1.9)$$

*Proof.* Note that by Minkowski's inequality, (denoting  $\Pi_N = \Pi_{\leq N} - \Pi_{\leq N/2}$  for  $N \geq 2$  and  $\Pi_1 = \Pi_{\leq 1}$ ),

$$\left( \mathbf{E}_{\mu^\kappa} \|g\|_{H^{-s}}^{2p} \right)^{1/p} \approx \left( \mathbf{E}_{\mu^\kappa} \left( \sum_{j \geq 0} 2^{-2sj} \|\Pi_{2^j} g\|_{L^2}^2 \right)^p \right)^{1/p} \lesssim \sum_{j \geq 0} 2^{-2sj} \left( \mathbf{E}_{\mu^\kappa} \|\Pi_{2^j} g\|_{L^2}^{2p} \right)^{1/p},$$

which is bounded  $\lesssim 1$  by (1.6). □

**Remark 1.5.** Note that Theorem 1.2 (with  $s = 0$ ) implies the logarithmic divergence in  $L^2$ :

$$\mathbf{E}_{\mu^\kappa} \|g\|_{L^2}^2 \approx |\log \kappa|. \quad (1.10)$$

**Remark 1.6.** The cumulative power spectrum estimate in (1.6) comes in two parts. the lower bound is *easier* to establish, and contains relatively little dynamical information: we show in Section 3.2 that it follows from the fact that Lipschitz velocity fields cannot mix a scalar faster than exponential (provided that one has suitable exponential moment estimates to control large deviations). The upper bound on the other hand is much more difficult. It make crucial use of the optimal, almost-sure uniform in  $\kappa$  mixing estimates obtained in our recent work [18], which in turn depends heavily on the results and methods of our earlier works on mixing and Lagrangian chaos [17, 19]. Without these results, we would not be able to obtain an upper bound. See below in Section 1.3 for more discussion on this sequence of works.

**Remark 1.7.** The proof of the upper bound in (1.6) holds directly for all  $N < \infty$ . By itself, the estimates on the cumulative power spectrum in (1.6) are not precise enough to localize the dissipative range. Indeed, it is easy to check that (1.6) implies for all  $\zeta > 0$ ,

$$\left( \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \approx_{\zeta,p} \log N \quad \text{for all } N_0 \leq N \leq \kappa^{-\zeta}.$$

However, (1.4) provides the additional information needed to localize the dissipative scale to approximately  $\kappa^{-1/2}$ , as one should expect from parabolic regularity. Indeed,

$$\mathbf{E}_{\mu^\kappa} \|(I - \Pi_{\leq N}) g\|_{L^2}^2 \leq \frac{1}{N^2} \mathbf{E}_{\mu^\kappa} \|\nabla g\|_{L^2}^2 = \frac{\chi}{\kappa N^2}.$$

The estimate (1.7) provides more precise high frequency moment control, albeit with a logarithmic deviation from (1.4).

**Remark 1.8.** The cumulative power spectrum estimate (1.6) is a little weaker than (1.2) or the dyadic-shell averaged version in (1.19) below. However, if  $\mathbf{E}_{\mu^\kappa} |\hat{g}(k)^2| \approx F(k)$  for  $F(k)$  a monotone function, then (1.6) implies that  $F(k) \approx |k|^{-d}$  (and analogously for any dyadic-shell averaged version). In a dyadic-shell averaged version, the only kind of violations of Batchelor’s law not ruled out by (1.6) are if a sparse set (i.e. zero asymptotic density) of dyadic shells have too little or too much  $L^2$  mass (but never more than  $\log N$ ). The pointwise version in (1.2) could also be violated if the  $L^2$  mass was not distributed evenly enough in angle (as in, for example, the CAT map example in Section 1.3). See also 1.4.

### 1.2.2 Yaglom’s law

Yaglom’s law was predicted in 1949 in [129] for all passive scalar turbulence regimes, which is a reformulation of the constant scale-by-scale  $L^2$  flux characteristic of anomalous dissipation. Every turbulent system satisfies at least one analogue of this; for 3D Navier-Stokes it is the Kolmogorov 4/5 law (see [21, 68, 107] and the references therein). In [17], we showed that Yaglom’s law holds over *some* inertial range with an unspecified lower bound  $\ell_\kappa$  with  $\ell_\kappa \rightarrow 0$  as  $\kappa \rightarrow 0$ . Theorem 1.2 allows us to refine this and show that this law essentially holds over the entire inertial range  $\ell \gtrsim \kappa^{1/2}$ . The proof is an easy application of the methods of [17, 21] and the a priori estimate (1.10) from Theorem 1.2 and so is omitted for brevity.

**Corollary 1.9** (Yaglom’s law over sharp inertial range). *Denote the finite increment for  $h \in \mathbb{R}^d$ ,*

$$\delta_h g(x) = g(x+h) - g(x).$$

Let  $\ell_\kappa > 0$  be such that  $\ell_\kappa^{-1} = o((\kappa |\log \kappa|)^{-1/2})$  as  $\kappa \rightarrow 0$ , then

$$\lim_{\ell_I \rightarrow 0} \limsup_{\kappa \rightarrow 0} \sup_{\ell \in (\ell_\kappa, \ell_I)} \left| \frac{1}{\ell} \mathbf{E}_{\mu^\kappa} \int_{\mathbb{T}^d} \int_{\mathbb{S}^{d-1}} |\delta_{\ell n} g|^2 \delta_{\ell n} u \cdot n \, dndx + \frac{4}{d} \chi \right| = 0. \quad (1.11)$$

Note that by 1.10, statistically stationary solutions are blowing up in  $L^2$ , and so (1.11) is only possible with the assistance of a large amount of cancellations.

**Remark 1.10.** For 3D Navier-Stokes, the Kolmogorov 4/5 law together with statistical self-similarity formally predicts the Kolmogorov  $-5/3$  power spectrum ( $|k|^2 \mathbf{E} |\hat{u}(k)|^2 \approx |k|^{-5/3}$ ). However, most experimental evidence indicates that the  $-5/3$  power law scaling is not exact [3, 68, 86, 124], and corrections are expected due to intermittency effects (see e.g. [105] for dissenting opinion). While passive scalar turbulence in the Batchelor regime is expected to display intermittency [78, 110, 113, 122], we nevertheless do not see intermittency corrections to Batchelor’s prediction on the power spectrum. Heuristically, one might guess this from Yaglom’s law (1.11): the regularity of the velocity implies that (1.11) formally scales like the second order structure function  $\mathbf{E}_{\mu^\kappa} \|\delta_\ell g\|_{L^2}^2$  which formally scales as the power spectrum by the Wiener-Khinchin theorem. Hence, Yaglom’s law suggests a certain rigidity to Batchelor-regime passive scalars that is not present in many other turbulent systems.

### 1.2.3 The vanishing diffusivity limit

A further property of general turbulent systems is that we expect to be able to pass to the vanishing diffusivity limit in the statistically stationary solution for  $(u_t, g_t^\kappa)$  and obtain statistically stationary weak solutions of the  $\kappa = 0$  problem. One expects that these limiting statistics, supported on very low regularity spaces, capture the dissipation of conserved quantities at a constant rate, i.e., the scale-by-scale flux of conserved quantities is on average constant through *all* sufficiently small scales. The limit itself is sometimes called *ideal turbulence* [61]. See [67] for an example of a simplified shell model where such a program has been carried out successfully.

Batchelor’s law in Theorem 1.2 provides a fundamental regularity estimate needed to complete this program in our setting. To our knowledge, this is the first time such a program has been carried out for a



fundamental physical model. More specifically, Corollary 1.4 implies that for each  $s > 0$  we have a uniform in  $\kappa$  bound in negative Sobolev spaces. Prokhorov's theorem then implies that  $\{\mu^\kappa\}_{\kappa \in (0,1)}$  has at least one weak limit measure  $\mu^0$  as  $\kappa \rightarrow 0$ , which lives on the intersection of these spaces

$$H^- := \bigcap_{s \in (0,1]} H^{-s};$$

see Section 4.1 for details. By their very nature, the limiting measures are necessarily pathological and in fact assigns zero measure to  $\mathbf{H} \times L^1$  and therefore the passive scalar  $g$  is almost-surely not even a locally-integrable function (see Theorem 1.14 below). With this in mind, it is natural to study solutions  $(u_t, g_t^0)$  of the limiting  $\kappa = 0$  system

$$\begin{aligned} \partial_t u_t + u_t \cdot \nabla u_t + \nabla p_t - \nu \Delta u_t &= Q\dot{W}_t, \\ \partial_t g_t^0 + u_t \cdot \nabla g_t^0 &= b\dot{\xi}_t, \end{aligned}$$

in a negative Sobolev space  $H^{-s}$ . For sufficiently regular velocity fields such as those in  $\mathbf{H}$ , weak solutions to the transport equation are well-posed on  $H^{-s}$  by duality with respect to  $H^s$  for all  $s > 0$ . After extracting the limit  $\mu^0$ , we pass  $\kappa \rightarrow 0$  in the invariance relation for  $(u_t, g_t^\kappa)$  to show that any limit point of  $\{\mu^\kappa\}_{\kappa \in (0,1)}$  is in fact a stationary measure for  $(u_t, g_t^0)$ . This is summarized in the following theorem.

**Theorem 1.11** (Vanishing dissipation limit). *There exists a subsequence  $\{\mu^{\kappa_n}\}$  and a limit measure  $\mu^0$  on  $\mathbf{H} \times H^-$  such that for each  $s > 0$ ,  $\mu^{\kappa_n}$  converges weakly to  $\mu^0$  as a measure on  $\mathbf{H} \times H^{-s}$ . Moreover, any limit point  $\mu^0$  is a stationary measure for  $(u_t, g_t^0)$  and the limit satisfies Batchelor's law over an infinite inertial range: for  $N_0$  as in Theorem 1.2 there holds*

$$\left( \mathbf{E}_{\mu^0} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \approx_p \log N \quad \text{for all } N \geq N_0, \quad (1.12)$$

and moreover, there is non-vanishing  $L^2$  flux:

$$\mathbf{E}_{\mu^0} \langle \Pi_{\leq N}(ug), \nabla \Pi_{\leq N} g \rangle = -\frac{1}{2} \|\Pi_{\leq N} b\|_{L^2}^2. \quad (1.13)$$

**Remark 1.12.** Note that the absence of the  $\kappa \Delta$  dissipation in the transport equation makes the existence of non-trivial stationary solutions far from obvious. The Markov process  $(u_t, g_t^0)$  has bad regularity properties on  $\mathbf{H} \times H^{-s}$  (it is not Feller) due to a lack of stability in the transport equation with respect to  $\mathbf{H}$  perturbations of the velocity field, and so a Krylov-Bogoliubov argument using the uniform mixing bound do not apply. See Section 4.3 for more discussion.

**Remark 1.13.** The non-vanishing flux (1.13) is analogous to Yaglom's law (1.11) for the  $\kappa = 0$  equation, in that both results say something about the constancy of the  $L^2$  flux in the  $\kappa \rightarrow 0$  limit. However, as we show below in Theorem 1.14,  $\mu^0$  generic  $g$  are not even integrable on  $\mathbb{T}^d$ , and so we are unsure how to pass the  $\kappa \rightarrow 0$  the limit in Yaglom's law directly.

#### 1.2.4 Irregularity of $\kappa = 0$ stationary statistics

As mentioned in Section 1.2.3, the limiting stationary statistics associated to  $\mu^0$  are very irregular. Indeed, the uniform bound (1.9) used to extract the limit  $\mu^0$  suggests that the measure is concentrated on a subspace of irregular distributions in  $H^-$ , while the limiting version of Batchelor's law (1.12) implies that

$$\mathbf{E}_{\mu^0} \|g\|_{L^2}^2 = \infty, \quad (1.14)$$

so that  $\mu^0$  cannot give second moments to functions in  $L^2$ . The primary reason for this irregularity is that, in the absence of a Laplacian, stationary solutions to the zero diffusivity equation

$$\partial_t g_t^0 + u_t \cdot \nabla g_t^0 = b\dot{\xi}_t \quad (1.15)$$

must “anomalously” dissipate the input from the noise through the advection term. Since the velocity field  $(u_t)$  is smooth we cannot rely on roughness of the velocity field to dissipate and must rely solely on the mixing properties of  $(u_t)$ . Indeed, since the velocity field  $(u_t)$  is at least Lipschitz, one can see that the semi-group  $S_t^0(\omega, u)$  conserves the  $L^1$  norm and therefore one should expect that  $L^1$  solutions to (1.15) cannot dissipate. This conservation  $L^1$  for Lipschitz velocity fields follows from the DiPerna/Lions theory [50] for the transport equation. In this theory an important role is played by the commutator

$$(u \cdot \nabla g)_\epsilon - u \cdot \nabla(g)_\epsilon, \quad (1.16)$$

where  $(\cdot)_\epsilon$  denotes mollification to scale  $\epsilon > 0$ . The equation (1.15) cannot dissipate if (1.16) vanishes, and since the velocity field is regular, the only way that such a commutator would not vanish is if the associated scalar  $(g_t)$  is sufficiently irregular. One can check that if the velocity field is Lipschitz and  $(g_t)$  belongs to  $L^1$ , then (1.16) vanishes in  $L^1$  as  $\epsilon \rightarrow 0$ . As a consequence of this type of analysis, we obtain the following theorem.

**Theorem 1.14** (Limiting solutions are not  $L^1$ ). *Any limit measure  $\mu^0$  from Theorem 1.11 satisfies*

$$\mu^0(\mathbf{H} \times L^1) = 0.$$

**Remark 1.15.** It is important to note that contrary to (1.14), which only states that  $\mu^0$  can’t have second moments on  $L^2$ , Theorem (1.14) is much stronger in that it implies that  $\mu^0$  generic scalars  $g$  are “true” distributions in the sense that they do not take values in any space of integrable functions, regardless of moments.

**Remark 1.16.** In general, DiPerna/Lions theory predicts that the the semi-group  $S_t^0(\omega, u)$  conserves the  $L^p$  norm if the velocity field is in the Sobolev space  $W^{1,q}(\mathbb{T}^d)$  for  $q = p/(p-1)$ . Velocity fields  $(u_t)$  which are not Lipschitz, in general do not propagate  $L^1$  and should not be considered as belonging to the “Batchelor Regime”. See [103] for an example of a continuous  $W^{1,p}$  velocity field which does not propagate  $L^1$  and also [53] for a related example.

Theorem 1.14 is based on understanding the formal conservation law  $L^1$  of the inviscid equation (1.15). However, another clear question is to determine how irregular  $g$  must be to have a non-vanishing  $L^2$  flux as in (1.13), i.e. studying violation of the  $L^2$  conservation law. This is analogous to the problem for weak solutions of the 3D Euler equations known as *Onsager’s conjecture* which has received a significant amount of mathematical attention in recent years (see e.g. [47, 48, 80] and the review [33]). In the context of the 3D Euler equations, Onsager’s conjecture states that weak solutions can dissipate energy when not in  $C_{t,x}^{0,1/3-}$  and cannot dissipate energy if in  $C_{t,x}^{0,1/3+}$ . The easier direction, that weak solutions of 3D Euler conserve energy with sufficient regularity, was studied previously in [35, 39, 60], which provided progressively sharper regularity. Specifically, in [35] it was shown that the Onsager-critical space (the space which divides dissipative from conservative) is the Besov space  $L_t^3 B_{3,\infty}^{1/3}$ .

In the spirit of [35], we show below that the space  $L_t^2 B_{2,\infty}^0$  is Onsager-type critical for the passive scalar turbulence problem, where  $B_{2,\infty}^0 \subseteq H^-$  is the Besov space of tempered distributions  $f \in H^-$  such that

$$\sup_{N \in \{2^j : j \in \mathbb{N}_*\}} \|\Pi_N f\|_{L^2} < \infty,$$

(recall  $\Pi_N = \Pi_{\leq N} - \Pi_{\leq N/2}$  is projection onto the dyadic shell of frequencies between  $N$  and  $N/2$  and  $\mathbb{N}_* = \{0\} \cup \mathbb{N}$ ). Time-integrability will be connected to moments with respect to the stationary measure  $\mu^0$  since for statistically stationary processes, expectations of time averages are exactly equal to ensemble

averages. Understanding when the  $L^2$  flux vanishes reduces to essentially the same commutator as in the  $L^1$  case above in (1.16):

$$\langle \Pi_{\leq N}(u \cdot \nabla g), \Pi_{\leq N} g \rangle_{L^2} = \langle \Pi_{\leq N}(u \cdot \nabla g) - u \cdot \nabla \Pi_{\leq N} g, \Pi_{\leq N} g \rangle_{L^2}.$$

Indeed, the projection  $\Pi_{\leq N}$  is a particular type of mollifier where here  $N$  plays the role of  $\epsilon^{-1}$ . We show that this commutator vanishes as  $N \rightarrow \infty$  in the subspace  $B_{2,c}^0 \subset B_{2,\infty}^0$  of distributions such that

$$\limsup_{j \rightarrow \infty} \|\Pi_{2^j} f\|_{L^2} = 0.$$

In order to quantify regularity in this space, we introduce generalized fractional derivative norms, akin to studying the modulus of continuity to generalize Hölder regularity.

**Definition 1.17.** We call a multiplier  $M : [0, \infty) \rightarrow [1, \infty)$   $B_{2,c}^0$ -suitable if

- (i)  $M$  is monotone increasing and  $\lim_{k \rightarrow \infty} M(k) = \infty$ ;
- (ii)  $M$  is (globally) Lipschitz continuous and  $\forall C > 0, \exists c > 0$  such that if  $C^{-1}|k| \leq |\ell| \leq C|k|$ ,

$$|M(k) - M(\ell)| \leq \frac{c}{\sqrt{1 + |k|^2}} M(\ell) |k - \ell|.$$

Given such an  $M$ , we define the generalized Besov norm

$$\|f\|_{B_{2,\infty}^M} := \sup_{N \in \{2^j : j \in \mathbb{N}_*\}} M(N) \|\Pi_N f\|_{L^2}.$$

We show below in Lemma 5.3 that  $f \in B_{2,c}^0$  if and only if  $\|f\|_{B_{2,\infty}^M} < \infty$  for some  $B_{2,c}^0$ -suitable  $M$ . We prove the following theorem by contradiction with (1.13) in Section 5.2.

**Theorem 1.18** (Onsager-type criticality of  $L_t^2 B_{2,\infty}^0$ ). *Let  $\mu^0$  be a stationary measure for the  $\kappa = 0$  limit process extended to  $\mathbf{H} \times H^{-1}$ . Then for every  $p > 2$  and all  $B_{2,c}^0$ -suitable  $M$ ,*

$$\mathbf{E}_{\mu^0} \|g\|_{B_{2,\infty}^M}^p = +\infty.$$

**Remark 1.19.** Note that we are not able to show that  $\mu^0$  assigns zero mass to  $B_{2,c}^0$  as we could for  $L^1$ , and instead can only show that moments with  $p > 2$  cannot be finite. This obstruction is related to the fact that moments in  $\mu^0$  are related to time integrability of stationary solutions and the fact that the critical space is  $L_t^2 B_{2,\infty}^0$ . It is unclear if it is possible for  $\mu^0$  to assign positive measure to  $\mathbf{H} \times B_{2,c}^0$ .

Using Theorems 1.18 and 1.14 we can deduce the following about the solutions studied in Theorem 1.2. In particular, we show that moments of solutions diverge in  $L^1$  and  $B_{2,c}^0$ , analogous to (1.10).

**Corollary 1.20.** *The unique stationary measure  $\mu^\kappa$  for the  $(u_t, g_t)$  process on  $\mathbf{H} \times L^2$  satisfies the following  $\forall \delta > 0$*

$$\lim_{\kappa \rightarrow 0} \mathbf{E}_{\mu^\kappa} \|g\|_{L^1}^\delta = \infty \quad \text{and} \quad \lim_{\kappa \rightarrow 0} \mathbf{E}_{\mu^\kappa} \|g\|_{B_{2,\infty}^M}^{2+\delta} = \infty$$

for any  $B_{2,c}^0$ -suitable multiplier  $M$ .

### 1.3 A simple physical mechanism for Batchelor’s Law

Analogous to the energy cascade in hydrodynamic turbulence, the mechanism responsible for the power law scaling in Batchelor’s law is the transfer of scalar mass from low to high Fourier modes.

Recall that in the absence of diffusivity ( $\kappa = 0$ ) and sources ( $b = 0$ ), the solution  $g_t$  to passive scalar advection is given by  $g_t(x) = g_0 \circ (\phi^t)^{-1}$ , where the Lagrangian flow  $\phi^t : \mathbb{T}^d \rightarrow \mathbb{T}^d$  solves the ODE

$$\frac{d}{dt}\phi^t(x) = u_t(\phi^t(x)), \quad \phi^0(x) = x. \quad (1.17)$$

This implies

$$\nabla g_t(x) = (D_{(\phi^t)^{-1}x}\phi^t)^{-\top} \nabla g_0((\phi^t)^{-1}(x)).$$

As physicists have understood for some time now [5, 6, 11, 28, 62], this hints that exponential growth in  $|D_x\phi^t| = |(D_x\phi^t)^{-\top}|$  is responsible for shifting the scalar from low to high modes. We discuss such physical theories further below.

#### Hyperbolicity and scalar advection: case studies

An explicit example of the role played by hyperbolicity is given in Batchelor’s original formal argument [13] for (1.2), which considered the effect of a local linear straining flow on a pure Fourier mode of passive scalar. A version of this is as follows: consider the Lagrangian flow  $\phi^t : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$\phi^t(x) = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix} x.$$

In the absence of diffusion or random sources, the initial scalar  $g_0(x) = e^{ik \cdot x}$ ,  $k \in \mathbb{Z}^2$  is advected by  $\phi^t$  to  $g_t(x) = g_0 \circ (\phi^t)^{-1}(x) = e^{i(e^{-t}k_1, e^t k_2) \cdot x}$ , resulting in an exponential-in-time increase in frequency for all  $k$  such that  $k_2 \neq 0$ .

The following example is in some ways closer to passive scalar advection on a bounded domain, and provides a simple heuristic for the exponent  $-1$  in (1.2). Consider the following discrete-time model, where Lagrangian flow is replaced by iterations of Arnold’s CAT<sup>4</sup> map  $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$  defined by

$$f(x) = Ax \pmod{1}, \quad A := \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix},$$

where  $x = (x_1, x_2) \in \mathbb{T}^2$  is parametrized by points in  $[0, 1) \times [0, 1)$ , and “mod 1” means that we take the fractional part of each coordinate individually. Note  $D_x f = A$ ; since  $A$  is symmetric with determinant 1, we see that  $f$  preserves volume (i.e., is incompressible), and that  $D_x f = A$  has two distinct real eigenvalues  $\lambda > 1 > \lambda^{-1} > 0$ . This yields, for every  $x \in \mathbb{T}^2$ , a splitting of tangent space  $T_x \mathbb{T}^2 = E_x^u \oplus E_x^s$  into a stable space  $E_x^s$  (eigenspace for  $\lambda^{-1}$ ) along which  $D_x f$  contracts exponentially, and the unstable space  $E_x^u$  (eigenspace for  $\lambda$ ) along which  $D_x f$  expands<sup>5</sup>. Viewing  $f$  as a model of the time-1 map of Lagrangian flow, we see that all trajectories of  $f$  exhibit the shear-straining behavior from Batchelor’s original argument.

Passive scalar advection (ignoring diffusivity for now) can be modeled as follows: given an initial scalar  $g_0 : \mathbb{T} \rightarrow \mathbb{R}$ , the time evolution  $g_n \mapsto g_{n+1}$  for the scalar is

$$g_{n+1}(x) = g_n \circ f^{-1}(x) + \omega_{n+1} b(x),$$

<sup>4</sup>Here CAT is an acronym for “continuous automorphism of the torus”.

<sup>5</sup>This behavior of exponential stretching and contracting infinitesimally in phase space is called *hyperbolicity*, and is the core mechanism behind the emergence of chaotic or seemingly random behavior in otherwise deterministic dynamical systems. For a review, see, e.g., [30].

where  $\omega_i \sim N(0, 1), i \geq 1$  are IID Gaussian random variables (playing the role of  $\xi_i$  in (1.1)), and  $b(x) = \sin(2\pi x_1)$  (playing the role of  $b$  in (1.1)). If  $g_0 = 0$ , the time- $n$  scalar is given by

$$g_n(x) = \sum_{i=0}^{n-1} \omega_{n-i} b \circ f^{-i} = \sum_{i=0}^{n-1} \omega_{n-i} \sin \left( 2\pi \left\langle (A^{-\top})^i \hat{e}_1, x \right\rangle \right). \quad (1.18)$$

where  $\hat{e}_1 = (1, 0) \in \mathbb{R}^2$ . The eigenvalues of  $A^{-\top}$  are again  $\lambda > 1 > \lambda^{-1}$  and  $|(A^{-\top})^n \hat{e}_1| \approx \lambda^n$ , signaling an exponential transfer of energy from large to small scales. Indeed, for all  $j \geq 1, n \gg 2^{j+1}$  we have

$$\mathbf{E} \sum_{2^j \leq |k| < 2^{j+1}} |\hat{g}_n(k)|^2 \approx 1, \quad (1.19)$$

where  $g_n(x) = \sum_k \hat{g}_n(k) \sin(2\pi k \cdot x), k \in \mathbb{Z}^2$ . This power spectrum corresponds precisely to (1.2), provided one sums over dyadic shells.

### Lagrangian chaos and $\kappa$ -uniform almost-sure exponential mixing

Exponential growth in time of  $\|D_x \phi^t\|$  for typical  $x$  is associated with hyperbolicity for the Lagrangian flow due to the incompressibility. This is sometimes referred to as *Lagrangian chaos*, specifically, the positivity of the Lyapunov exponent

$$\lambda_1(x) = \limsup_{t \rightarrow \infty} \frac{1}{t} \log |D_x \phi^t| > 0 \quad (1.20)$$

for ‘typical’  $x \in \mathbb{T}^d$ . Before our work, Lagrangian chaos had been proved for stationary, white-in-time velocity fields in [15, 16]. In [17], we proved that the Lagrangian flow associated to solutions  $(u_t)$  of the stochastically-forced 2D Navier-Stokes and 3D hyperviscous Navier-Stokes in the sense that there is a deterministic constant  $\lambda_1 > 0$  such that (1.20) holds with lim-sup replaced with lim for all  $x$  and all initial fluid configurations, almost surely (see [17] for rigorous statements). We emphasize that the results of [17] rely crucially upon the stochastic framework: it is often very difficult to provide rigorous proofs of positivity of Lyapunov exponents, even for deceptively simple models such as the Chirikov standard map family [54, 72] (a discrete-time toy model of the Lagrangian flow [43]). See [112, 131] for more discussions. In Navier-Stokes, one of the enemies is the formation of coherent vortices inside of which hyperbolicity is halted (see e.g. [10]). The arguments in our paper [17] imply that with probability 1, vortices of this kind cannot permanently trap any particles.

However, a positive Lyapunov exponent alone is not enough to obtain the kind of frequency cascade that led to Batchelor’s law in the CAT map example above. To provide a proof of Batchelor’s law here, we use a much stronger property: uniform-in- $\kappa$ , almost-sure exponential mixing<sup>6</sup>. In [18], we proved that if  $g_t^\kappa$  is mean zero and solves the initial value problem

$$\begin{aligned} \partial_t g_t^\kappa + u_t \nabla g_t^\kappa &= \kappa \Delta g_t^\kappa \\ g_0^\kappa &= g, \end{aligned}$$

then there exists a *deterministic* constant  $\gamma > 0$  independent of  $\kappa$  and random constant  $D_\kappa$  (depending on initial fluid configuration) such that the negative Sobolev norm

$$\|g_t^\kappa\|_{H^{-1}} := \sup_{\|f\|_{H^1}=1} \left| \int f g_t^\kappa dx \right| \leq D_\kappa e^{-\gamma t} \|g\|_{H^1},$$

<sup>6</sup>One can construct dynamical systems with a positive Lyapunov exponent but arbitrarily slow (e.g., polynomial or logarithmic) mixing/decay of Lagrangian correlations, for example, Pommeau-Manneville maps (see, e.g., [99, 117]).

where  $D_\kappa$  also has suitable moment bounds *independent of  $\kappa$*  (see Definition 2.9 and Theorem 2.12 for precise statements). See [125] for a discussion of using negative Sobolev norms to quantify mixing of scalar fields.

To see why uniform-in- $\kappa$  scalar mixing implies an exponential increase of frequency scale, note that

$$\|\Pi_{\leq N} g_t^\kappa\|_{L^2} \leq N \|g_t^\kappa\|_{H^{-1}} \leq DN e^{-\gamma t} \|g\|_{H^1},$$

where  $\Pi_{\leq N}$  denotes projection to Fourier modes of frequency  $\leq N$ . From this, one can see that for times  $t \gg \frac{\log N}{\gamma}$ , most of the scalar has been transferred from frequencies  $\leq N$  to higher frequencies. As in the CAT map example, this exponentially fast transfer from low to high frequencies is exactly the mechanism which gives rise to Batchelor’s law when  $u_t$  is given by Navier-Stokes – one difficulty is of course dealing with potential unboundedness of the random constant  $D$ , which captures fluctuations in the mixing time. There is a large mathematical literature on scalar mixing in the mathematics literature; see e.g. [1, 29, 53, 59, 81, 98, 118, 130] and the references therein.

Our work [18] builds on our earlier work [19] that proved the corresponding statement for  $\kappa = 0$ ; similarly, this latter work uses the Lagrangian chaos result of [17] as a lemma. Our works [18, 19] are based on analyzing two-point statistics of the Lagrangian flow and a key step is to upgrade the positivity of the Lyapunov exponent to positivity of the *moment Lyapunov exponents* for  $0 < p \ll 1$ .

$$\Lambda(p) = - \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbf{E} |D_x \phi^t|^{-p}.$$

This is deeply related to fluctuations of the finite-time Lagrangian Lyapunov exponents as  $t \rightarrow \infty$  (for the relation of moment Lyapunov exponents to large deviations in the convergence of Lyapunov exponents, see, e.g., [7–9]). It had already been realized by physicists that fluctuations of Lagrangian Lyapunov exponents should play a key role in Batchelor’s law (see discussions in [5, 6, 11] and the references therein) and so our works are in some ways a mathematically rigorous completion of some of these ideas.

#### 1.4 Open problems in Batchelor-regime passive scalar turbulence

Theorems 1.2–1.20 provide a satisfactory foundation and starting point for a mathematical understanding of Batchelor-regime passive scalar turbulence. However, there are many remaining open questions, all of which are potentially accessible in the near future using a combination of stochastic PDEs, harmonic analysis, and random dynamical systems. Let us briefly outline these here.

- **Intermittency** Certainly the most important set of open questions regard intermittency. Following discussions in e.g. [34, 68] we can begin to study intermittency by looking at the flatness parameters

$$F_p(N) = \frac{\mathbf{E} \|\Pi_N g\|_{L^{2p}}^{2p}}{\left(\mathbf{E} \|\Pi_N g\|_{L^2}^2\right)^p}.$$

A non-intermittent field would satisfy  $F_p(N) \leq C(p)$  as  $N \lesssim \kappa^{-1/2}$ ,  $N \rightarrow \infty$ ,  $\kappa \rightarrow 0$ . An example of such a field is white noise. At the opposite extreme is a maximally intermittent random field consisting of single Dirac delta function placed with uniform probability on  $\mathbb{T}^d$ , which satisfies  $F_p(N) \approx_p N^{p-1}$ . Passive scalar turbulence is expected to be intermittent [78, 110, 113, 122]. A major step in our understanding would be to provide an analytic derivation of powers  $\zeta(p)$  such that  $F_p(N) \approx N^{\zeta(p)}$  for  $N \rightarrow \infty$ ,  $\kappa \rightarrow 0$ , if such powers exist. It is instructive to see that the CAT map (1.18) example is in fact *not* intermittent, in contrast to expectations regarding passive scalar turbulence. RDS toy systems extending (1.18), such as the CAT map replaced with the Chirikov standard map with noise<sup>7</sup> [23], might be able to shed some further light on intermittency effects.

<sup>7</sup>Note that the presence of noise makes this tractable, unlike the deterministic case which is notoriously difficult [54, 72].

- **How universal is “universal”?** Another set of important problems is to study how widely applicable the Batchelor spectrum, and other tenants of the theory, such as uniqueness of stationary measures, Lagrangian chaos etc, to different and more realistic settings.
  - Problems on  $\mathbb{T}^d$  with body forcing are far removed from any real physical applications. Extending existing theories to include boundaries (exterior or interior domains) and replacing body forces with boundary driving is probably the most physically important extension. Even relatively basic questions, such as uniqueness of stationary measures of the Navier-Stokes equations, are to our knowledge, quite challenging and still open for most questions of this type. See [121] for some progress in this direction. A related direction is to study spatially homogeneous solutions on  $\mathbb{R}^d$ .
  - Even on  $\mathbb{T}^d$  with stochastic forcing, in the case of the Navier-Stokes equations, our results on Lagrangian chaos and scalar mixing [17–19] cannot handle  $C_x^\infty$  forcing yet. It is natural to seek to extend this to  $C_x^\infty$  forcing and further to include non-white-in-time forcing such as OU tower forcing (see Section 2.2 and [19]) and the class of bounded forcing studied in [82, 92, 93]. Note that the hypoellipticity theory of Hairer and Mattingly [77] applies to the one-point Lagrangian flow ( $\dot{x}_t = u_t(x_t)$ ) generated by 2D Navier-Stokes with OU tower forcing (and so the  $(u_t, Z_t, x_t)$  Markov process has a unique stationary measure), however, our Lagrangian chaos results require strong Feller in order to use the infinite-dimensional Furstenberg criterion [17].
- **Sharper regularity estimates and structure function renormalization.** Even just concerning basic questions related to Batchelor’s law (1.6), there are still remaining questions.
  - There are three basic levels of precision when discussing the power spectrum. After the cumulative spectrum, the next most difficult is the dyadic shell-by-shell estimate as in (1.19). The next most difficult after that is the pointwise estimate

$$\mathbf{E}_{\mu^\kappa} |\hat{g}(k)|^2 \approx |k|^{-d} \quad N_0 < |k| < \kappa^{-1/2}.$$

In fact, one can even try to search for an estimate of the type (see discussion in the physics literature e.g. [51]), where of course we mean that the error is uniformly controlled in  $\kappa$ ,

$$\mathbf{E}_{\mu^\kappa} |\hat{g}(k)|^2 = C_B |k|^{-d} + o_{k \rightarrow \infty}(|k|^{-d}) \quad N_0 < |k| < \kappa^{-1/2}.$$

These estimates are definitely not equivalent, as seen from (1.19). Simple RDS systems such as hyperbolic toral automorphisms drawn from random distributions might be able to shed some light on the subtle differences between these spectral characterizations.

- It is an interesting and subtle question to determine if the limiting solutions we obtain in Theorem 1.2 are *exactly* in the Onsager critical space  $L_t^2 B_{2,\infty}^0$ , or more to the point, whether the  $\kappa > 0$  approximations are uniformly bounded in this space, that is,

$$\mathbf{E}_{\mu^\kappa} \sup_{N \in 2^{\mathbb{N}}} \|\Pi_N g\|_{L^2}^2 \lesssim 1.$$

This is almost the same as the shell-by-shell Batchelor spectrum as in (1.19), but the  $\mathbf{E}$  and  $\sup_{N \in 2^{\mathbb{N}}}$  are reversed.

- Batchelor’s law should concern the second order structure function  $\mathbf{E}_{\mu^\kappa} \|\delta_\ell g\|_{L^2}^2$ . However, as the scalar does not even remain a locally integrable function as  $\kappa \rightarrow 0$ , it is hard to make sense of exactly how the second order structure function will behave in and at the limit  $\kappa \rightarrow 0$ ; instead, this may require a suitable renormalization. Similarly, Yaglom’s law on the  $L^2$  flux (1.11) is hard to make sense of rigorously at the  $\kappa = 0$  limit.

## 2 Preliminaries

### 2.1 Fluid models

Let  $d = 2$  or  $3$ . We fix a real Fourier basis of  $\mathbf{L}^2 = \{u \in L^2(\mathbb{T}^d, \mathbb{R}^d) : \int u dx = 0, \operatorname{div} u = 0\}$  as follows: for  $m = (k, i) \in \mathbb{K} := \mathbb{Z}_0^d \times \{1, \dots, d-1\}$ , we set

$$e_m(x) = \begin{cases} c_d \gamma_k^i \sin(k \cdot x) & k \in \mathbb{Z}_+^d \\ c_d \gamma_k^i \cos(k \cdot x) & k \in \mathbb{Z}_-^d. \end{cases}$$

Here,  $\mathbb{Z}_0^d := \mathbb{Z}^d \setminus \{0\}$ , and  $\mathbb{Z}_0^d = \mathbb{Z}_+^d \cup \mathbb{Z}_-^d$  is the partition defined by  $\mathbb{Z}_+^d = \{k = (k^{(1)}, \dots, k^{(d)}) \in \mathbb{Z}_0^d : k^{(d)} > 0\} \cup \{k \in \mathbb{Z}_0^d : k^{(1)} > 0, k^{(d)} = 0\}$  and  $\mathbb{Z}_-^d = -\mathbb{Z}_+^d$ . For each  $k \in \mathbb{Z}_0^d$ , we have fixed a set  $\{\gamma_k^i\}_{i=1}^{d-1}$  of orthonormal vectors spanning the complement of the line spanned by  $k \in \mathbb{R}^d$ ; these are assumed to satisfy  $\gamma_{-k}^i = -\gamma_k^i$ . The coefficients  $c_d > 0$  are normalization constants. Note that if  $d = 2$ ,  $\gamma_k = \gamma_k^1$  spans the perpendicular to  $k$ , and may therefore be taken to be  $\gamma_k = k^\perp/|k|$ ,  $k^\perp := (k^{(2)}, -k^{(1)})$ .

In terms of this Fourier basis, we consider the white-in-time, spatially Sobolev stochastic forcing term

$$QW_t := \sum_{m \in \mathbb{K}} q_m e_m(x) W_t^m, \quad (2.1)$$

where  $W_t^m, m \in \mathbb{K}$  are a family of independent standard one-dimensional Wiener processes with respect to the canonical stochastic basis  $(\Omega_W, \mathcal{F}^W, (\mathcal{F}_t^W), \mathbf{P}_W)$ . The following decay and non-degeneracy assumption is made throughout this and our previous works [17–19].

**Assumption 1.** *There exists  $\alpha > \frac{5d}{2}$  such that for all  $m = (k, i) \in \mathbb{K}$ , we have*

$$q_m \approx \frac{1}{|k|^\alpha}.$$

The state space for our fluid velocity fields is

$$\mathbf{H} := \left\{ u \in H^\sigma(\mathbb{T}^d, \mathbb{R}^d) : \int u dx = 0, \operatorname{div} u = 0 \right\},$$

where  $\sigma \in (\alpha - 2(d-1), \alpha - \frac{d}{2})$ . Note that by our choice of  $\alpha$ , we have  $\sigma > \frac{d}{2} + 3$ , so  $\mathbf{H} \hookrightarrow C^3$ . We will write the Navier-Stokes system as an abstract evolution equation on  $\mathbf{H}$  by

$$\partial_t u + B(u, u) + Au = Q\dot{W} = \sum_{m \in \mathbb{K}} q_m e_m \dot{W}^m, \quad (2.2)$$

where

$$B(u, v) = (\operatorname{Id} - \nabla(-\Delta)^{-1}\nabla \cdot) \nabla \cdot (u \otimes v)$$

$$Au = \begin{cases} -\nu \Delta u & \text{if } d = 2 \\ -\nu' \Delta u + \nu \Delta^2 u & \text{if } d = 3. \end{cases}$$

The  $(u_t)$  process with initial data  $u$  is defined as the solution to (2.2) in the mild sense [45, 94]:

$$u_t = e^{-tA}u - \int_0^t e^{-(t-s)A} B(u_s, u_s) ds + \int_0^t e^{-(t-s)A} Q dW(s), \quad (2.3)$$

where the above identity holds  $\mathbf{P}_W$  almost surely for all  $t > 0$ . We have the following well-posedness theorem:



**Proposition 2.1** ([45, 94]). *Let  $d = 2$  or  $3$ . Under Assumption 1, for all initial  $u \in \mathbf{H} \cap \mathbf{H}^{\sigma'}$  with  $\sigma' < \alpha - \frac{d}{2}$  and all  $T > 0, p \geq 1$ , there exists a  $\mathbf{P}_W$ -a.s. unique solution  $(u_t)$  to (2.3) which is  $\mathcal{F}_t^W$ -adapted, and belongs to  $L^p(\Omega_W; C([0, T]; \mathbf{H} \cap \mathbf{H}^{\sigma'})) \cap L^2(\Omega_W; L^2(0, T; \mathbf{H}^{\sigma'+(d-1)}))$ . Additionally, for all  $p \geq 1$  and  $0 \leq \sigma' < \sigma'' < \alpha - \frac{d}{2}$ ,*

$$\begin{aligned} \mathbf{E}_W \sup_{t \in [0, T]} \|u_t\|_{\mathbf{H}^{\sigma'}}^p &\lesssim_{T, p, \sigma'} 1 + \|u\|_{\mathbf{H} \cap \mathbf{H}^{\sigma'}}^p \\ \mathbf{E}_W \int_0^T \|u_s\|_{\mathbf{H}^{\sigma'+(d-1)}}^2 ds &\lesssim_{T, \delta} 1 + \|u\|_{\mathbf{H}^{\sigma'}}^2 \\ \mathbf{E}_W \sup_{t \in [0, T]} \left( t^{\frac{\sigma'' - \sigma'}{2(d-1)}} \|u_t\|_{\mathbf{H}^{\sigma''}} \right)^p &\lesssim_{p, T, \sigma', \sigma''} 1 + \|u\|_{\mathbf{H}^{\sigma'}}^p. \end{aligned}$$

**Proposition 2.2.** *Under Assumption 1, the process  $(u_t)$  solving 2.2 admits a unique stationary measure  $\mu$ .*

Proposition 2.2 was first proved for  $d = 2$  in [66] under Assumption 1; the generalization to  $d = 3$  is a straightforward extension.

**Remark 2.3.** Uniqueness of stationary measures is known for Navier-Stokes under much weaker nondegeneracy conditions than Assumption 1, e.g., the *truly hypoelliptic* setting of [75] with  $d = 2$  which only requires to force modes  $|m|_\infty \leq 1$ . However, Assumption 1 is necessary for the Lagrangian chaos and scalar mixing results in [17–19] because our methods require, for now, strong Feller regularity of the semigroup  $t \mapsto u_t$ . See Remark 2.6 in [17] and Remark 2.19 in [19] for more discussion.

Because we require Assumption 1, we are not aware of how to extend our results to fluid models solving the Navier-Stokes equations which are spatially  $C^\infty$  or differentiable in time. However, we are able to apply our results to a class of finite-dimensional fluid models for which solutions are  $C_t^k C_x^\infty$ . This is the subject of the following short section.

## 2.2 $C_t^k C_x^\infty$ fluid models governed by finite dimensional SDE

If the fluid evolves according to a finite-dimensional SDE, then the methods involved are significantly simpler at a technical level and the strength of Hörmander’s theorem [79] allows us to impose much weaker conditions on the noise models we consider. Consequently, we can produce fluid models which have better spatial and time regularity.

Below, for  $QW_t$  given by (2.1), we define  $\mathcal{K}_0 = \{m \in \mathbb{K} : q_m \neq 0\}$ . For  $m = (k, i) \in \mathbb{K} = \mathbb{Z}_0^d \times \{1, \dots, d-1\}$ , we define  $|m|_\infty = \max_j |k^{(j)}|$ .

For the following finite dimensional stochastic fluids models we will make the the following less restrictive assumption to Assumption 1.

**Assumption 2.** *Assume  $m \in \mathcal{K}_0$  if  $|m|_\infty \leq 2$ .*

For  $\mathcal{K} \subset \mathbb{K}$  we define  $\mathbf{H}_{\mathcal{K}}$  to be the linear span of the Fourier modes  $\{e_m\}_{m \in \mathcal{K}}$ . One model we consider is the Stokes system on  $\mathbb{T}^d, d = 2, 3$ , prescribing the time evolution  $(u_t)$  on the state space  $\mathbf{H}_{\mathcal{K}_0}$  for fixed initial  $u_0 \in \mathbf{H}_{\mathcal{K}_0}$  by

$$\begin{aligned} \partial_t u_t &= -\nabla p_t + \Delta u_t + Q\dot{W}_t \\ \operatorname{div} u_t &= 0. \end{aligned} \tag{2.4}$$

When  $\mathcal{K}_0$  is finite, (2.4) is a finite-dimensional SDE on  $\mathbf{H}_{\mathcal{K}_0} \cong \mathbb{R}^{|\mathcal{K}_0|}$ ; indeed, it is essentially a product of independent Ornstein-Uhlenbeck processes on  $\mathbb{R}^{|\mathcal{K}_0|}$ , and in particular an elliptic diffusion on  $\mathbb{R}^{|\mathcal{K}_0|}$ .

For  $N \geq 1$  we define  $\mathbf{H}_N \subset \mathbf{H}$  to be the linear span of the  $e_m$  with  $|m|_\infty \leq N$ . Define  $\Pi_{\leq N} : \mathbf{H} \rightarrow \mathbf{H}_N$  to be the orthogonal projection. Another model we consider for  $(u_t)$  is the Galerkin-Navier-Stokes system, defined for fixed  $u_0 \in \mathbf{H}_N$  by

$$\begin{aligned} \partial_t u_t &= -\Pi_{\leq N}(\nabla p_t + u_t \cdot \nabla u_t) + \nu \Delta u_t + Q \dot{W}_t \\ \operatorname{div} u_t &= 0, \end{aligned} \tag{2.5}$$

where implicitly we assume that  $q_m \neq 0$  only if  $|m|_\infty \leq N$ . As with the Stokes system, Galerkin-Navier-Stokes is an SDE on the finite-dimensional space  $\mathbf{H}_N$ ; under Assumption 2, equation (2.5) is known to satisfy the parabolic Hörmander condition [56, 115], and so results in a hypoelliptic diffusion.

Standard finite-dimensional stochastic analysis [95] applies to each of (2.4), (2.5), yielding velocity field processes  $(u_t)$  which are  $C^{\frac{1}{2}-}$  in time and spatially  $C^\infty$ . In addition to these, our methods also apply to a class of models including those which vary  $C^k$  in time for any fixed  $k \geq 1$ . These models are effectively driven by the projection of a coupled system of Ornstein-Uhlenbeck processes. More precisely: fix  $2 \leq N \leq M$  and let  $\mathcal{A} : \mathbf{H}_M \times \mathbf{H}_M$  be diagonalizable with strictly positive spectrum. Let

$$\Gamma W_t = \sum_{|m|_\infty \leq M} \Gamma_m e_m W_t^m$$

and let  $X : \mathbf{H}_N \times \mathbf{H}_N \rightarrow \mathbf{H}_N$  be a bilinear mapping with  $u \cdot X(u, u) = 0$  and  $X(e_m, e_m) = 0$  for all  $|m|_\infty \leq N$ . We consider the following *generalized Galerkin-Navier-Stokes system with OU tower noise*, defined by

$$\begin{aligned} \partial_t u_t &= -X(u_t, u_t) + \nu \Delta u_t + Q Z_t \\ \partial_t Z_t &= -\mathcal{A} Z_t + \Gamma W_t. \end{aligned} \tag{2.6}$$

Here, the noise term applied to the  $(u_t)$  process is  $Q Z_t := \sum_{|m|_\infty \leq N} q_m Z_t^m e_m(x)$ , where  $t \mapsto Z_t^m \in \mathbb{R}$  are the Fourier coefficients of  $Z_t$ , i.e.,  $Z_t(x) = \sum_m Z_t^m e_m(x)$ . Note that  $(u_t)$  is not a Markov process, but  $(Z_t)$  alone and  $(u_t, Z_t)$  are. The appropriate nondegeneracy assumption in this setting is as follows:

**Assumption 3.** *Assume that the coefficients  $\{q_m\}$  satisfy Assumption 2, and additionally that the parabolic Hörmander condition holds for the  $(u_t, Z_t)$  process on  $\mathbf{H}_N \times \mathbf{H}_M$ .*

**Theorem 2.4.** *All of the main results in Section 1.2 hold whenever  $(u_t)$  evolves according to (2.4) or (2.5) under Assumption 2, or (2.6) under Assumption 3.*

**Remark 2.5** (see Remark 1.10 in [19]). Consider the following example of a system in the setting of (2.6). Fix  $n \geq 1$  and consider the model

$$u_t(x) = \sum_{|m|_\infty \leq 2} u_t^m e_m(x), \tag{2.7}$$

where the coefficients  $u_t^m$  evolve according to

$$\begin{aligned} \partial_t u_t^m &= -u_t^m + Z_t^{m,0} \\ \partial_t Z_t^{m,\ell} &= -Z_t^{m,\ell} + Z_t^{m,\ell+1} \quad 1 \leq \ell \leq n-1 \\ \partial_t Z_t^{m,n} &= -Z_t^{m,n} + \dot{W}_t^m. \end{aligned}$$

Up to re-indexing, this fits into the framework of (2.6) with  $X \equiv 0$ . The parabolic Hörmander condition for  $(u_t, Z_t)$  is satisfied, and so Theorem 2.4 holds for  $(u_t)$  as above. Notably, solutions  $(u_t)$  to (2.7) are  $C^{n+1}$ -differentiable in time and smooth (indeed, analytic) in space. The authors hope this serves as an indication that although the methods in this paper rely strongly on the stochastic framework, they are not inherently restricted to the rough time regularity of white-in-time noise.

Going forward, we will assume  $(u_t)$  solves (2.2) with Assumption 1 for remainder of the paper. The application of these arguments to the finite dimensional models (2.4), (2.5) and (2.6) is straightforward and omitted for brevity.

### 2.3 Lyapunov functions and exponential estimates

The results in our previous series of papers [18, 19] require the use of the family of Lyapunov functions

$$V(u) = V_{\beta, \eta}(u) := (1 + \|u\|_{\mathbf{H}}^2)^\beta e^{\eta \|u\|_{\mathbf{W}}^2} \quad (2.8)$$

where  $\beta \geq 0, \eta > 0$ . Here, for velocity fields  $u$  we define

$$\|u\|_{\mathbf{W}} := \begin{cases} \|\operatorname{curl} u\|_{L^2} & d = 2 \\ \|u\|_{L^2} & d = 3. \end{cases}$$

Below, we formulate a drift condition for the family  $V = V_{\beta, \eta}$ , ensuring that trajectories of the  $(u_t)$  process frequently visit the sublevel sets of  $V$ . Define  $\eta_* = \nu/\mathcal{Q}$ , where

$$\mathcal{Q} = 64 \begin{cases} \sup_{m \in \mathbb{K}} |k| |q_m| & d = 2 \\ \sup_{m \in \mathbb{K}} |q_m| & d = 3. \end{cases}$$

**Lemma 2.6** (Lemma 3.7 in [19]). *Let  $(u_t)$  be a solution to the stochastic Navier Stokes equations (2.2) with initial data  $u \in \mathbf{H}$ . For all  $0 \leq \gamma < \nu/8$ ,  $r \in (0, 3)$ ,  $C_0 \geq 0$ , and  $V(u) = V_{\beta, \eta}$  where  $\beta \geq 0$  and  $0 < e^{\gamma T} \eta < \eta_*$ , there exist constants  $c = c(\gamma, r, C_0, \beta, \eta) > 0$ ,  $C = C(\gamma, r, C_0, \beta, \eta) \geq 1$  such that the following estimate holds for any  $T > 0$ :*

$$\mathbf{E}_W \exp \left( C_0 \int_0^T \|u_s\|_{\mathbf{H}^r} ds \right) \sup_{0 \leq t \leq T} V^{e^{\gamma t}}(u_t) \leq C e^{cT} V(u). \quad (2.9)$$

**Remark 2.7.** To connect (2.9) with more standard drift conditions given in, e.g., [101]: write  $P_1$  for the Markov semi-group for the Navier-Stokes process  $(u_t)$  and apply Jensen's inequality to (2.9) to deduce that  $\exists C_L > 0$  such that  $P_1 V \leq (e^{C_L} V)^{e^{-\gamma}}$ . In particular, we have the following drift condition:

$$\forall \delta > 0, \exists C_\delta > 0 \quad \text{such that} \quad P_1 V \leq \delta V + C_\delta.$$

Note that the contraction constant  $\delta > 0$  above can be made *arbitrarily small*. Iterating this bound, it is straightforward to show that  $\forall \lambda > 0, \exists K_\lambda$  such that for all  $t > 0$  there holds

$$P_t V \leq e^{-\lambda t} V + K_\lambda. \quad (2.10)$$

We will also need the following basic stability estimate for solutions to the Navier-Stokes equations.

**Lemma 2.8.** *For all  $u, u' \in \mathbf{H}$  and let  $(u_t)$  and  $(u'_t)$  be the corresponding solutions with the same noise-path  $\omega$ . Then  $\forall p \in [1, \infty)$ , there exists a deterministic  $K > 0$  such that following stability estimate holds for  $V(u)$  as in (2.8),*

$$\mathbf{E} \| |u_t - u'_t| \|_{\mathbf{H}}^p \lesssim_p e^{pKt} (V(u)^p + V(u')^p) \| |u - u'| \|_{\mathbf{H}}^p.$$

*Proof.* Define  $w_t = u_t - u'_t$ ,

$$\partial_t w_t + B(u_t, w_t) + B(w_t, u'_t) + Aw_t = 0.$$

The proof proceeds as a variation of e.g. [Lemma 3.10; [19]]. Analogous to the calculations therein we have for some  $C > 0$  and some  $r \in (\frac{d}{2} + 1, 3)$

$$\|w_t\|_{\mathbf{W}} \leq \exp\left(C \int_0^t \|u_\tau\|_{H^r} + \|u'_\tau\|_{H^r} d\tau\right) \|u - u'\|_{\mathbf{W}}$$

and for some  $q, C > 0$  (possibly a different  $C$ ),

$$\|w_t\|_{\mathbf{H}}^2 \lesssim \exp\left(C \int_0^t \|u_\tau\|_{H^r} + \|u'_\tau\|_{H^r} d\tau\right) \left(\sup_{0 < s < t} (\|u_\tau\|_{\mathbf{H}}^q + \|u'_\tau\|_{\mathbf{H}}^q)\right) \|u - u'\|_{\mathbf{H}}^2.$$

The result then follows from Lemma 2.6. □

## 2.4 Uniform mixing and enhanced dissipation

In this section we will summarize the results of [17–19] which are used to prove Batchelor’s law (Theorem 1.2) and the other results of this paper. Throughout, assume  $(u_t)$  solves (2.2) and that Assumption 1 holds. Consider the advection-diffusion equation with diffusivity  $0 \leq \kappa \ll 1$ ,

$$\partial_t \bar{g}_t + u_t \cdot \nabla \bar{g}_t = \kappa \Delta \bar{g}_t \tag{2.11}$$

for fixed initial  $\bar{g}_0 = g \in L^2$ ,  $\int g dx = 0$ . This defines the following (random) two-parameter semigroup of linear operators on  $L^2$ : for  $0 \leq s \leq t$ ,  $\omega \in \Omega_W$  and initial  $u = u_0 \in \mathbf{H}$ , define  $S_{t,s}^\kappa(\omega, u) : L^2 \rightarrow L^2$  by

$$\bar{g}_t = S_{t,s}^\kappa(\omega, u) \bar{g}_s.$$

When starting from  $\tau = 0$  we will write  $S_t^\kappa(\omega, u) = S_{t,0}^\kappa(\omega, u)$ . Note that  $S_0^\kappa(\omega, u) = I$ , the identity on  $L^2$ . Note as well the following cocycle property: for any  $s, t > 0$ , we have

$$S_{t+s}^\kappa(\omega, u) = S_s^\kappa(\theta_t \omega, u_t) S_t^\kappa(\omega, u),$$

where,  $\theta_t : \Omega_W \rightarrow \Omega_W$  is the standard time-shift on Wiener space defined for each  $\omega \in \Omega_W$  by

$$\theta_t \omega(\cdot) := \omega(\cdot + t) - \omega(t).$$

This implies that we can write the two parameter semi-group  $S_{t,s}^\kappa(\omega, u)$  as

$$S_{t,s}^\kappa(\omega, u) = S_t^\kappa(\theta_s \omega, u_s). \tag{2.12}$$

Note that for any  $p \geq 1$ ,  $S_t^\kappa$  extends in a natural way to a defined mapping on  $L^p$ , with range contained in  $L^p$ . The same is true of Sobolev spaces  $H^s$ ,  $s > 0$ .

Several of the important results in this section involve random constants of the form  $D : \Omega_W \times \mathbf{H} \rightarrow \mathbb{R}_{\geq 1}$ , the  $\mathbf{P}$ -law of which are controlled in terms of  $V(u)$ . Since random constants of this type appear many times in this paper, we introduce the following definition.

**Definition 2.9.** Let  $D : \Omega_W \times \mathbf{H} \rightarrow \mathbb{R}_{\geq 1}$  be measurable. We say that  $D$  has  $V$  bounded  $p$ -th moment if  $\exists \beta \geq 0$  such that  $\forall 0 < \eta < \eta_*$ , we have for  $V = V_{\beta, \eta}(u)$

$$\mathbf{E} D^p(\cdot, u) \lesssim_\eta V(u).$$

The following Lemma is very important since it provides us some control on the fluctuations of the mixing time in terms of the Lyapunov function  $V(u)$ .

**Lemma 2.10.** *Let  $D : \Omega_W \times \mathbf{H} \rightarrow [1, \infty)$  have  $V$ -bounded  $p$ -th moment. Then,  $\forall \lambda > 0, \exists K_\lambda > 0$  such that*

$$\mathbf{E}D^p(\theta_t\omega, u_t) \lesssim_{p,\lambda} e^{-\lambda t}V(u) + K_\lambda. \quad (2.13)$$

*Proof.* Using, in sequence, the tower property of conditional expectation, the fact that  $u_t$  is  $\mathcal{F}_t^W$  measurable and the increment  $\theta_t\omega$  is independent of  $\mathcal{F}_t^W$ , and equations (2.13) and (2.10), we have:

$$\mathbf{E}D^p(\theta_t\omega, u_t) = \mathbf{E} \left( \mathbf{E} [D^p(\theta_t\omega, u_t) | \mathcal{F}_t^W] \right) \lesssim \mathbf{E}V(u_t) \lesssim e^{-\lambda t}V(u) + K_\lambda.$$

□

The following Lemma is a useful corollary of (2.9).

**Lemma 2.11.** *For all  $p \in [1, \infty)$ , there exists a (deterministic)  $C_0 > 0$  and a random constant  $D_0 : \mathbf{H} \times \Omega_W \rightarrow \mathbb{R}_{\geq 1}$  with  $V$ -bounded  $p$ -th moment such that*

$$\exp \left( \int_0^t \|\nabla u_\tau\|_{L^\infty} d\tau \right) \leq D_0(\omega, u)e^{C_0 t}.$$

*Proof.* We provide the proof when  $p = 1$ ; other values of  $p$  require straightforward adjustments.

Set  $V = V_{0,\eta}$ , where  $\eta \in (0, \eta_*)$  is arbitrary. To start, note that by Lemma 2.6 and Chebyshev's inequality,  $\exists c = c(\eta) > 0$  such that

$$\mathbf{P} \left( \exp \left( 2 \int_0^t \|\nabla u_s\|_{L^\infty} ds \right) > V(u)e^{4ct} \right) \leq \frac{\mathbf{E} \exp \left( 2 \int_0^t \|\nabla u_s\|_{L^\infty} ds \right)}{V(u)e^{4ct}} \lesssim_\eta e^{-3ct}.$$

By Borel-Cantelli, there exists  $N(\omega, u) \geq 1$  with  $\mathbf{P}(N(\cdot, u) \geq n) \lesssim_\eta e^{-3cn}$  such that

$$\exp \left( \int_0^n \|\nabla u_s\|_{L^\infty} ds \right) \leq V(u)e^{4cn} \quad \text{for } n \geq N(\omega, u).$$

To bound when  $n < N(\omega, u)$ , we find

$$\exp \left( \int_0^n \|\nabla u_s\|_{L^\infty} ds \right) \leq D_0(\omega, u) := \exp \left( \int_0^{N(\omega, u)} \|\nabla u_s\|_{L^\infty} ds \right)$$

and note that by Cauchy-Schwarz,

$$\begin{aligned} \mathbf{E}D_0(\cdot, u) &\leq \sum_n (\mathbf{P}(N = n))^{1/2} \left[ \mathbf{E} \exp \left( 2 \int_0^n \|\nabla u_s\|_{L^\infty} ds \right) \right]^{1/2} \\ &\leq \sum_n (e^{-3cn})^{1/2} (V(u)e^{4cn})^{1/2} \leq V(u) \sum_n e^{-cn} \lesssim V(u). \end{aligned}$$

□

With these preparations in place, we now state the main results of [17–19].

**Theorem 2.12** (Theorem 1.2 in [18]). *There exists  $\kappa_0 > 0$  for which the following holds for all  $\kappa \in [0, \kappa_0]$ . Let  $(u_t)$  solve (2.2) for an arbitrary initial condition  $u_0 = u \in \mathbf{H}$ . For all  $s > 0, p \geq 1$ , there exists a deterministic  $\gamma = \gamma(s, p) > 0$  (depending only on  $s, p$  and the parameters  $Q, \nu$  etc) and a random constant  $D_\kappa(\omega, u) : \Omega \times \mathbf{H} \rightarrow [1, \infty)$  (also depending on  $p, s$ , as well as  $\kappa$ ) such that for all  $g \in H^s$ ,*

$$\|S_t^\kappa(\omega, u)g\|_{H^{-s}} \leq D_\kappa(\omega, u)e^{-\gamma t} \|g\|_{H^s}.$$

*The random constant  $D_\kappa$  has  $V$ -bounded  $p$ -th moment with implicit constant independent of  $\kappa$ .*

Theorem 2.12 is proved in [19] for  $\kappa = 0$  and [18] for  $\kappa \in [0, \kappa_0]$ . Both papers rely heavily on Lagrangian chaos as proved in [17]. From Theorem 2.12, it is relatively straightforward to prove the following enhanced dissipation result.

**Theorem 2.13** (Theorem 1.3 in [18]). *Let  $\kappa_0 > 0$  and  $\gamma = \gamma(1, p)$  be as in Theorem 2.12, where  $p \geq 2$  is arbitrary. Let  $(u_t)$  solve (2.2) for an arbitrary initial condition  $u_0 = u \in \mathbf{H}$ . Then, there is a random constant  $D'_\kappa(\omega, u)$  such that for all  $g \in L^2$*

$$\|S_t^\kappa(\omega, u)g\|_{L^2} \leq \min\{1, D'_\kappa(\omega, u)\kappa^{-1}e^{-\gamma t}\}\|g\|_{L^2}.$$

The random constant  $D'_\kappa$  has  $V$ -bounded  $p$ -th moment with implicit constant independent of  $\kappa$ .

**Theorem 2.14** (Theorem 1.5 in [18]). *In the setting of Theorem 2.12, let*

$$\tau_* = \tau_*(\omega, u, g) = \inf\{t : \|g_t\|_{L^2} < \frac{1}{2}\|g\|_{L^2}\}.$$

Then, there exists a  $\kappa_0 > 0$  a sufficiently small universal constant such that for all  $\kappa \in (0, \kappa_0]$ , one has

$$\tau_*(\omega, u, g) \geq \delta(g, u, \omega)|\log \kappa| \quad \text{with probability } 1,$$

where  $\delta(g, u, \omega) \in (0, 1)$  is a  $\kappa$ -independent random constant with the property that there exists a  $\beta \geq 1$  such that for all  $p \geq 1$  and  $\eta > 0$  with  $V(u) = V_{\beta, \eta}(u)$  we have

$$\mathbf{E}\delta^{-p}(\cdot, u) \lesssim_{p, \eta, \beta} \frac{\|g\|_{H^1}^p}{\|g\|_{L^2}^p} V(u)^p.$$

## 2.5 Unique stationary measure for $(u_t, g_t^\kappa)$

We provide here an argument proving uniqueness of the stationary measure for the passive scalar process  $(u_t, g_t^\kappa)$  on  $\mathbf{H} \times L^2$ . The following argument is quite general, and applies in a variety of cases outside the scope of the main results; see Remark 2.16 below.

**Proposition 2.15.** *For any  $\kappa > 0$ , the Markov process  $(u_t, g_t^\kappa)$  admits a unique stationary measure  $\mu^\kappa$  on  $\mathbf{H} \times L^2$ .*

*Proof.* It suffices to show that any two ergodic stationary measures  $\mu_1^\kappa, \mu_2^\kappa$  for  $(u_t, g_t^\kappa)$  coincide. By standard ergodic theory for random dynamical systems [87] and the ergodic decomposition theorem (see, e.g., [58]), uniqueness of ergodic stationary measures for  $(u_t, g_t^\kappa)$  implies uniqueness of stationary measures.

Assume now that  $\mu_i^\kappa$  are two ergodic stationary measures,  $i = 1, 2$ . To prove  $\mu_1^\kappa = \mu_2^\kappa$ , it suffices to show that for each bounded, globally Lipschitz  $\psi : \mathbf{H} \times L^2 \rightarrow \mathbb{R}$ , we have that  $\int \psi d\mu_1^\kappa = \int \psi d\mu_2^\kappa$ . Without loss of generality, we can assume  $\|\psi\|_{\text{Lip}} = 1$ .

By ergodicity of  $\mu_i^\kappa$  and the Birkhoff ergodic theorem, we have the following for  $i = 1, 2$ : there is a set  $\hat{\mathcal{G}}_i = \hat{\mathcal{G}}_i(\psi) \subset \mathbf{H} \times L^2$  of full  $\mu_i^\kappa$ -measure such that for all initial  $(u, g) \in \hat{\mathcal{G}}_i$ , we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{t=0}^{n-1} \psi(u_t, g_t^\kappa) = \int \psi d\mu_i^\kappa$$

with probability 1. Next, observe that  $\mu_i^\kappa$  projects to the unique stationary measure  $\mu$  for the  $(u_t)$  process (Proposition 2.2). It follows from Fubini's theorem that for  $i = 1, 2$ , we have  $\mu(\mathcal{G}_i) = 1$ , where  $\mathcal{G}_i$  is the projection of  $\hat{\mathcal{G}}_i$  to  $\mathbf{H}$ , i.e.,  $\mathcal{G}_i = \{u \in \mathbf{H} : \exists g \in L^2 \text{ such that } (u, g) \in \hat{\mathcal{G}}_i\}$ . Since  $\mu(\mathcal{G}_i) = 1, i = 1, 2$ , we conclude that  $\mathcal{G}_1 \cap \mathcal{G}_2$  is nonempty. Fix  $u \in \mathcal{G}_1 \cap \mathcal{G}_2$  and let  $g, h \in L^2$  be such that  $(u, g) \in \hat{\mathcal{G}}_1, (u, h) \in \hat{\mathcal{G}}_2$ . Observe that

$$g_t - h_t = S_t^\kappa(\omega, u)(g - h),$$

i.e.,  $g_t - h_t$  solves (2.11). Theorem 2.13 implies  $\|g_t - h_t\|_{L^2} \rightarrow 0$  as  $t \rightarrow \infty$ . Given  $\delta > 0$ , let  $N_0 = N_0(\delta')$  be such that  $\|g_t - h_t\|_{L^2} < \delta$  for all  $t \geq N_0$ .

With these preparations in place, fix  $u \in \mathcal{G}_1 \cap \mathcal{G}_2$  and  $g, h \in L^2$  such that  $(u, g) \in \hat{\mathcal{G}}_1, (u, h) \in \hat{\mathcal{G}}_2$ . With  $\delta > 0$  as above, let  $N_0 = N_0(\delta)$ . We have

$$\left| \int \psi d\mu_1^\kappa - \int \psi d\mu_2^\kappa \right| \leq \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{t=0}^{n-1} |\psi(u_t, g_t^\kappa) - \psi(u_t, h_t^\kappa)| = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{t=N_0}^{n-1} |\psi(u_t, g_t^\kappa) - \psi(u_t, h_t^\kappa)|$$

Each summand on the RHS is  $\leq \delta$ , and so the whole limit is  $\leq \delta$ . Since  $\delta > 0$  was arbitrary, this completes the proof.  $\square$

**Remark 2.16.** The proof given above has two main ingredients: (1) uniqueness for the stationary measure  $\mu$  on  $\mathbf{H}$ , and (2)  $L^2$  dissipation estimates for  $S_t^\kappa(\omega, u)$  for  $\kappa > 0$ . Item (1) is known to hold for 2D Navier-Stokes under very weak nondegeneracy condition on the noise [75], while item (2) holds for a wide variety of fluid models by standard estimates. Thus, the proof given above is applicable at a much higher level of generality than that of Assumption 1, including the setting of 2D Navier-Stokes with “truly hypoelliptic” forcing as in [75].

### 3 Proof of Batchelor’s law

In this section we prove the main result of this paper, Theorem 1.2. To summarize the main approach, observe that by uniqueness of the stationary measure  $\mu^\kappa$  for (1.1) (Proposition 2.15), the Birkhoff ergodic theorem implies

$$\mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^2 = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \|\Pi_{\leq N} g_t^\kappa\|_{L^2}^2 dt$$

for  $\mu^\kappa$ -generic initial  $u = u_0 \in \mathbf{H}, g = g_0^\kappa \in L^2$ . On the other hand, the mild formulation for (1.1) reads as

$$g_t^\kappa = S_t^\kappa(\omega, u)g + \int_0^t S_{t,s}^\kappa(\omega, u)b d\xi_s. \quad (3.1)$$

Using the fact that

$$\mathbf{E}_\xi \left\langle S_t^\kappa(\omega, u)g, \int_0^t \Pi_{\leq N} S_{t,s}^\kappa(\omega, u)b d\xi_s \right\rangle_{L^2} = 0$$

results in the identity

$$\mathbf{E} \|\Pi_{\leq N} g_t^\kappa\|_{L^2}^2 = \mathbf{E} \|\Pi_{\leq N} S_t^\kappa(\omega, u)g\|_{L^2}^2 + \mathbf{E} \int_0^t \|\Pi_{\leq N} S_{t,s}^\kappa(\omega, u)b\|_{L^2}^2 ds \quad (3.2)$$

after taking an  $L^2$  norm in  $\mathbb{T}^d$ , then an expectation, and finally the Itô isometry. By Theorem 2.13, the first such term vanishes exponentially fast in  $L^2$  as  $t \rightarrow \infty$ , and so it is the second term that dominates the long-time behavior of  $\mathbf{E} \|g_t^\kappa\|_{L^2}^2$ , hence the value of  $\mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^2$ . While the above is stated for  $p = 1$ , the case  $p > 1$  is handled similarly using the Burkholder-Davis-Gundy inequality.

The upper bound for (1.6) in the inertial range is carried out in Section 3.1, and the lower bound in Section 3.2. The dissipative-range upper bound is done in Section 3.3.

#### 3.1 Upper bound in the inertial range

In this section we prove the  $\lesssim$  direction of (1.6), that is,

**Lemma 3.1.** *Under the conditions of Theorem 1.2, there holds for all  $p \in [1, \infty)$  and for all  $2 \leq N < \infty$ ,*

$$\left( \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \lesssim_p \log N. \quad (3.3)$$

We begin with the following lemma.

**Lemma 3.2.** For all  $p \in [1, \infty)$  and  $2 \leq N < \infty$ ,

$$\limsup_{t \rightarrow \infty} \int_0^t \left( \mathbf{E} \left\| \Pi_{\leq N} S_{t,\tau}^\kappa(\omega, u) b \right\|_{L^2}^{2p} \right)^{1/p} d\tau \lesssim_p \log N.$$

*Proof.* Recall that  $\|\Pi_{\leq N} h\|_{L^2} \lesssim N \|\Pi_{\leq N} h\|_{H^{-1}}$  for  $h \in L^2$ . By (2.12) and Theorem 2.12 followed by Lemma 2.10,

$$\begin{aligned} \int_0^t \left( \mathbf{E} \left\| \Pi_{\leq N} S_{t,\tau}^\kappa(\omega, u) b \right\|_{L^2}^{2p} \right)^{1/p} d\tau &= \int_0^t \left( \mathbf{E} \left\| \Pi_{\leq N} S_{t-\tau}^\kappa(\theta_\tau \omega, u_\tau) b \right\|_{L^2}^{2p} \right)^{1/p} d\tau \\ &\lesssim \int_0^t \left( \mathbf{E} D^{2p}(\theta_\tau \omega, u_\tau) \right)^{1/p} \min(1, N^2 e^{-2\lambda(t-\tau)}) d\tau \\ &\lesssim \int_0^t \left( e^{-\lambda\tau} V(u) + K_\lambda \right) \min(1, N^2 e^{-2\lambda(t-\tau)}) d\tau \\ &\lesssim \left( e^{-\lambda t} V(u) + 1 \right) \log N. \end{aligned}$$

The proof then follows by sending  $t \rightarrow \infty$ . □

We are now ready to complete the proof of Lemma 3.1.

**Proof of Lemma 3.1.** First observe that, for all  $(u, g) \in \mathbf{H} \times L^2$  there holds by (3.1)

$$\|\Pi_{\leq N} g t\|_{L^2}^{2p} \lesssim_p \|\Pi_{\leq N} S_t^\kappa(\omega, u) g\|_{L^2}^{2p} + \left\| \int_0^t \Pi_{\leq N} S_{t,\tau}^\kappa(\omega, u) b d\xi_\tau \right\|_{L^2}^{2p}.$$

From the initial data term, by Theorem 2.13,  $\exists \lambda$  (depending on  $p$ ) such that for a suitable  $V$  as in (2.8),

$$\left( \mathbf{E} \left\| \Pi_{\leq N} S_t^\kappa(\omega, u) g \right\|_{L^2}^{2p} \right)^{1/p} \lesssim \kappa^{-1} V(u) e^{-\lambda t} \|g\|_{L^2}^2.$$

By Burkholder-Davis-Gundy [e.g. Theorem 5.2.4 [45]] followed by Minkowski's inequality,

$$\begin{aligned} \left( \mathbf{E} \left\| \int_0^t \Pi_{\leq N} S_{t,\tau}^\kappa(\omega, u) b d\xi_\tau \right\|_{L^2}^{2p} \right)^{1/p} &\lesssim_p \left( \mathbf{E} \left( \int_0^t \|\Pi_{\leq N} S_{t,\tau}^\kappa(\omega, u)\|_{L^2}^2 d\tau \right)^p \right)^{1/p} \\ &\lesssim \int_0^t \left( \mathbf{E} \left\| \Pi_{\leq N} S_{t,\tau}^\kappa(\omega, u) b \right\|_{L^2}^{2p} \right)^{1/p} d\tau. \end{aligned}$$

Hence, Lemma 3.2 implies

$$\limsup_{t \rightarrow \infty} \left( \mathbf{E} \left\| \Pi_{\leq N} g t \right\|_{L^2}^{2p} \right)^{1/p} \lesssim \log N.$$

Note that by standard moment estimates (see e.g. [94]), for all  $p, \kappa > 0$ ,  $\mathbf{E}_{\mu^\kappa} \|g\|_{L^2}^p < \infty$ . Hence, by the Birkhoff ergodic theorem, for  $\mu^\kappa$ -a.e.  $(u, g) \in \mathbf{H} \times L^2$  we have

$$\left( \mathbf{E}_{\mu^\kappa} \left\| \Pi_{\leq N} g \right\|_{L^2}^{2p} \right)^{1/p} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \left( \mathbf{E}_{(u,g)} \left\| \Pi_{\leq N} g t \right\|_{L^2}^{2p} \right)^{1/p} dt \lesssim \log N.$$

This completes the proof of Lemma 3.1. □



### 3.2 Lower bound in the inertial range

In this section we prove the  $\gtrsim$  direction of (1.6).

**Lemma 3.3.** *Under the conditions of Theorem 1.2, there exists an  $N_0 \geq 2$  chosen sufficiently large (depending on  $b$  and the fluid parameters only) such that for all  $\kappa \in (0, \kappa_0]$  the following holds for all  $p \in [1, \infty)$  and all  $N \in [N_0, \kappa^{-1/2}]$ ,*

$$\left( \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \gtrsim_p \log N.$$

We begin with the following, which follows from an  $H^1$  energy estimate and Lemma 2.11.

**Lemma 3.4.** *There holds (with  $C_0$  and  $D_0$  as in Lemma 2.11)*

$$\|S_t^\kappa(\omega, u)g\|_{H^1} \leq D_0(\omega, u)e^{C_0 t} \|g\|_{H^1}.$$

The next lemma provides the lower bound on the stochastic integral contribution.

**Lemma 3.5.** *There exists an  $N_0 \geq 2$  chosen sufficiently large (depending on  $b$  and the fluid parameters only) such that for all  $\kappa \in (0, \kappa_0]$  the following limit holds for all  $N \in [N_0, \kappa^{-1/2}]$ ,*

$$\liminf_{t \rightarrow \infty} \int_0^t \mathbf{E} \|\Pi_{\leq N} S_{t-\tau}^\kappa(\omega, u)b\|_{L^2}^2 d\tau \gtrsim \log N.$$

*Proof.* Fix  $u \in \mathbf{H}$  and let  $\gamma \in (0, 1)$  to be chosen later. Write for  $t \geq \gamma \log N$  and  $N \geq 2$ ,

$$\begin{aligned} \int_0^t \|\Pi_{\leq N} S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr &\geq \int_{t-\gamma \log N}^t \|S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr \\ &\quad - \int_{t-\gamma \log N}^t \|\Pi_{>N} S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr. \end{aligned} \quad (3.4)$$

The first term is controlled by the  $H^1$  norm growth bound in Lemma 3.4

$$\mathbf{E} \int_{t-\gamma \log N}^t \|\Pi_{>N} S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr \lesssim \int_{t-\gamma \log N}^t N^{-2} \mathbf{E} D_0^2(\theta_r \omega, u_r) e^{\lambda(t-r)} dr.$$

By Lemma 2.10

$$\mathbf{E} D_0^2(\theta_r \omega, u_r) \lesssim e^{-\lambda r} V(u) + K_\lambda,$$

and hence, choosing  $\gamma < \lambda^{-1}$ , we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \mathbf{E} \int_{t-\gamma \log N}^t \|\Pi_{>N} S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr &\lesssim \gamma N^{\gamma \lambda - 2} \log N \limsup_{t \rightarrow \infty} (e^{-\lambda t} V(u) + K_\lambda) \\ &\leq C_0 N^{-1} \gamma \log N, \end{aligned}$$

for some constant  $C_0$  universal independent of  $\gamma, \kappa, t, u$ . This completes the second term in (3.4). For the first term in (3.4), we use the stopping time defined in Theorem 2.14 with initial data  $b$ :

$$\mathbf{E} \|S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 \geq \frac{1}{2} \|b\|_{L^2}^2 \mathbf{P}(\tau_*(\theta_r \omega, u_r) > t - r),$$

where (analogously to Theorem 2.14)  $\tau_*(\omega, u)$  is defined by

$$\tau_*(\omega, u) = \inf \left\{ t > 0 : \|S_t^\kappa(\omega, u)b\|_{L^2}^2 \leq \frac{\|b\|_{L^2}^2}{2} \right\}.$$

Theorem 2.14 then implies that  $\exists C_1 > 0$  (independent of  $t, r, \kappa, u$ ) and  $\delta(\omega, u) : \Omega \times \mathbf{H} \rightarrow \mathbb{R}_{>0}$  with  $\mathbf{E}\delta^{-1}(\cdot, u) \leq C_1 V(u)$  such that  $\tau_* \geq \delta |\log \kappa|$ . Using Chebyshev's inequality and an argument similar to that used to prove Lemma 2.10

$$\begin{aligned} \mathbf{P}(\tau_*(\theta_r \omega, u_r) > t - r) &\geq 1 - \mathbf{P}(\delta^{-1}(\theta_r \omega, u_r) > |\log \kappa|/(t - r)) \\ &\geq 1 - \frac{t - r}{|\log \kappa|} \mathbf{E}(\mathbf{E}(\delta^{-1}(\theta_r \omega, u_r) | \mathcal{F}_r)) \\ &\geq 1 - C_1 \frac{t - r}{|\log \kappa|} (e^{-\lambda r} V(u) + K_\lambda). \end{aligned}$$

It follows that for  $2 \leq N \leq \kappa^{-1/2}$ ,

$$\begin{aligned} \liminf_{t \rightarrow \infty} \mathbf{E} \int_{t-\gamma \log N}^t \|S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr \\ \geq \liminf_{t \rightarrow \infty} \frac{1}{2} \|b\|_{L^2}^2 \gamma \log N \left( 1 - \frac{C_1 \gamma \log N}{2 |\log \kappa|} (e^{-\lambda t} V(u) + K_\lambda) \right) \\ \geq \frac{1}{2} \|b\|_{L^2}^2 \gamma \log N \left( 1 - \frac{C_1 K_\lambda \gamma}{4} \right). \end{aligned}$$

Therefore, for  $K_\lambda C_1 \gamma < 1/8$  and  $N > N_0 := 8C_0$ .

$$\begin{aligned} \liminf_{t \rightarrow \infty} \mathbf{E} \int_0^t \|\Pi_{\leq N} S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr &\gtrsim (1 - \frac{\gamma}{4} C_1 K_\lambda - 2C_0 N^{-1}) \log N \\ &\gtrsim \log N. \end{aligned}$$

□

**Proof of Lemma 3.3:** By Jensen's inequality,  $\mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \geq (\mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^2)^p$  for  $p \geq 1$ , and so in the following argument it suffices to consider the  $p = 1$  case. For arbitrary initial data  $g \in L^2$ , we have by (3.2) that

$$\mathbf{E} \|\Pi_{\leq N} g_t\|_{L^2}^2 \geq \mathbf{E} \int_0^t \|\Pi_{\leq N} S_{t-r}^\kappa(\theta_r \omega, u_r)b\|_{L^2}^2 dr,$$

hence  $\liminf_{t \rightarrow \infty} \mathbf{E} \|\Pi_{\leq N} g_t\|_{L^2}^2 \gtrsim \log N$  by Lemma 3.5 for  $N_0 < N < \kappa^{-1/2}$ . By the Birkhoff ergodic theorem (as in Lemma 3.1 above),

$$\mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g_t\|_{L^2}^2 = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \mathbf{E} \|\Pi_{\leq N} g_t\|_{L^2}^2 dt \gtrsim \log N.$$

This completes the proof of Lemma 3.3. □

### 3.3 Upper bound in the dissipative range

First, we derive the upper bound on the  $L^2$  norm stated in (1.7). If we were not also interested in higher moments we could use the following simple observation using the  $L^2$  balance (1.4) and the upper bound in Batchelor's law (3.3)

$$\mathbf{E}_{\mu^\kappa} \|g\|_{L^2}^2 \lesssim \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq \kappa^{-1/2}} g\|_{L^2}^2 + \kappa \mathbf{E}_{\mu^\kappa} \|\Pi_{> \kappa^{-1/2}} \nabla g\|_{L^2}^2 \lesssim 1 + |\log \kappa|.$$

The following argument provides also estimates on moments  $p > 2$ .

**Lemma 3.6.** *Under the conditions of Theorem 1.2, there holds for all  $p \in [1, \infty)$ , for all  $\kappa \in (0, \kappa_0]$*

$$\left( \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} \right)^{1/p} \lesssim_p |\log \kappa|.$$

*Proof.* The proof proceeds similar to Lemma 3.3 with the enhanced dissipation estimate (Theorem 2.13) in place of the uniform mixing estimate (Theorem 2.12). As in Lemma 3.3, by Burkholder-Davis-Gundy and Minkowski's inequality,

$$\left( \mathbf{E} \|g_t\|_{L^2}^{2p} \right)^{1/p} \lesssim \left( \mathbf{E} \|S_t^\kappa(\omega, u)g\|_{L^2}^{2p} \right)^{1/p} + \int_0^t \left( \mathbf{E} \|S_{t,\tau}^\kappa(\omega, u)b\|_{L^2}^{2p} \right)^{1/p} d\tau.$$

By Theorem 2.13,  $\exists \lambda > 0$  and  $D'_\kappa(\omega, u)$  with  $V$ -bounded  $2p$ -th moments we have for any  $(u, g) \in \mathbf{H} \times L^2$ ,

$$\begin{aligned} \left( \mathbf{E} \|g_t\|_{L^2}^{2p} \right)^{1/p} &\lesssim \left( \mathbf{E} \|S_t^\kappa(\omega, u)g\|_{L^2}^{2p} \right)^{1/p} + \int_0^t \left( \mathbf{E} \|S_{t,\tau}^\kappa(\omega, u)b\|_{L^2}^{2p} \right)^{1/p} d\tau \\ &\lesssim \left( \mathbf{E} (D'_\kappa(\omega, u))^{2p} \right)^{1/p} \kappa^{-1} e^{-\lambda t} \|g\|_{L^2}^2 \\ &\quad + \int_0^t \left( \mathbf{E} (D'_\kappa(\theta_\tau \omega, u_\tau))^{2p} \right)^{1/p} \min(1, \kappa^{-1} e^{-\lambda(t-\tau)}) d\tau. \end{aligned}$$

By Lemma 2.10, we therefore have

$$\begin{aligned} \left( \mathbf{E} \|g_t\|_{L^2}^{2p} \right)^{1/p} &\lesssim_p V(u) \kappa^{-1} e^{-\lambda t} \|g\|_{L^2}^2 + \int_0^t (e^{-\lambda \tau} V(u) + 1) \min(1, \kappa^{-1} e^{-\lambda(t-\tau)}) d\tau \\ &\lesssim_p V(u) \kappa^{-1} e^{-\lambda t} (1 + \|g\|_{L^2}^2) + |\log \kappa|. \end{aligned}$$

Hence for all  $(u, g) \in \mathbf{H} \times L^2$ ,

$$\limsup_{t \rightarrow \infty} \mathbf{E} \|g_t\|_{L^2}^{2p} \lesssim |\log \kappa|.$$

Finally, by the Birkhoff ergodic theorem, for  $\mu^\kappa$  a.e.  $(u, g) \in \mathbf{H} \times L^2$

$$\mathbf{E}_{\mu^\kappa} \|g\|_{L^2}^{2p} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \mathbf{E}_{(u,g)} \|g_t\|_{L^2}^{2p} dt \lesssim |\log \kappa|^p.$$

□

Finally we turn to the proof of (1.7) for  $q > 0$ , which is a relatively straightforward consequence of parabolic regularity and the  $L^2$  a priori estimate in Lemma 3.6. First we prove the following quantitative  $H^s$  regularization estimate. Below we use the Fourier multiplier notation: for  $m : \mathbb{C}^d \rightarrow \mathbb{R}$  measurable, we define the operator

$$m(\nabla)f = \left( m(ik) \hat{f}(k) \right)^\vee.$$

**Lemma 3.7.** For all initial data  $(u, g) \in \mathbf{H} \times L^2$ , for all  $\gamma \in (0, \alpha - d - 1)$ ,  $\exists \beta \geq 0$ , such that  $\forall p \in [1, \infty)$  and  $\forall \eta > 0$  there holds for  $V = V_{\beta, \eta}$

$$\mathbf{E} \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^\gamma S_1^\kappa(\omega, u)g \right\|_{L^2}^p \lesssim_{p, \gamma} V^p(u) \|g\|_{L^2}^p.$$

*Proof.* We first deduce a pathwise a priori estimate assuming  $g$  is sufficiently smooth. To this end, by Plancherel's identity we have for  $t \in [0, 1]$ ,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa \right\|_{L^2}^2 &= \sum_{k \in \mathbb{Z}_*^d} \left( \gamma \log(1 + \sqrt{\kappa} \langle k \rangle) - \kappa |k|^2 \right) (1 + \sqrt{\kappa} \langle k \rangle)^{2\gamma t} |\hat{g}_t^\kappa(k)|^2 \\ &\quad + \left\langle (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa, (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} \nabla \cdot (u_t g_t^\kappa) \right\rangle_{L^2} \\ &=: \mathcal{I}_1 + \mathcal{I}_2. \end{aligned}$$

For  $\mathcal{I}_1$ , by  $\gamma \log(1 + x) \leq x^2 + C(\gamma)$ ,

$$\mathcal{I}_1 \lesssim_\gamma \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa \right\|_{L^2}^2.$$

By incompressibility and Plancherel's identity

$$\begin{aligned} \mathcal{I}_2 &= \left\langle (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa, (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} \nabla \cdot (u_t g_t^\kappa) - \nabla \cdot (u_t (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa) \right\rangle_{L^2} \\ &= \sum_{k, \ell \in \mathbb{Z}_*^d} (1 + \sqrt{\kappa} \langle k \rangle)^{\gamma t} \overline{\hat{g}_t^\kappa(k)} \left( (1 + \sqrt{\kappa} \langle k \rangle)^{\gamma t} - (1 + \sqrt{\kappa} \langle k - \ell \rangle)^{\gamma t} \right) ik \cdot \hat{u}_t(\ell) \hat{g}_t^\kappa(k - \ell) \\ &= \sum_{k, \ell \in \mathbb{Z}_*^d} \left( \mathbf{1}_{|k - \ell| \geq \frac{1}{2}|\ell|} + \mathbf{1}_{|k - \ell| < \frac{1}{2}|\ell|} \right) (1 + \sqrt{\kappa} \langle k \rangle)^{\gamma t} \overline{\hat{g}_t^\kappa(k)} \\ &\quad \times \left( (1 + \sqrt{\kappa} \langle k \rangle)^{\gamma t} - (1 + \sqrt{\kappa} \langle \ell \rangle)^{\gamma t} \right) ik \cdot \hat{u}_t(k - \ell) \hat{g}_t^\kappa(\ell) \\ &= \mathcal{I}_{2;HL} + \mathcal{I}_{2;LH}. \end{aligned}$$

On the support of the  $\mathcal{I}_{2;HL}$  (here  $HL$  stands for ‘‘high-low’’), we use that  $|k| + |\ell| \lesssim |k - \ell|$ , together with Cauchy-Schwarz, Young's inequality,  $H^r \hookrightarrow \widehat{L}^1$  for  $r > \frac{d}{2}$ , to deduce that for any  $r \in (\frac{d}{2} + 1, 3)$ ,

$$|\mathcal{I}_{2;HL}| \lesssim \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa \right\|_{L^2} \|g_t^\kappa\|_{L^2} \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} u_t \right\|_{H^r}.$$

On the support of the  $\mathcal{I}_{2;LH}$ , we use the mean value theorem and  $|k| \lesssim |\ell|$  to deduce

$$(1 + \sqrt{\kappa} \langle k \rangle)^{\gamma t} - (1 + \sqrt{\kappa} \langle k - \ell \rangle)^{\gamma t} \lesssim \sqrt{\kappa} \gamma t (1 + \sqrt{\kappa} \langle k - \ell \rangle)^{\gamma t - 1} |\ell|.$$

Therefore, by Cauchy-Schwarz, Young's inequality, and  $H^r \hookrightarrow \widehat{L}^1$  for  $r > \frac{d}{2}$ , for any  $r \in (\frac{d}{2} + 1, 3)$  we have

$$|\mathcal{I}_{2;LH}| \lesssim \|u_t\|_{H^r} \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa \right\|_{L^2}^2.$$

By Grönwall's inequality, there is some  $C > 0$  (independent of  $g, \kappa, u, t$ ) such that the following holds,

$$\begin{aligned} \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma t} g_t^\kappa \right\|_{L^2}^2 &\leq \exp \left( Ct + \int_0^t \|u_\tau\|_{H^r} d\tau \right) \|g\|_{L^2}^2 \\ &\quad + \exp \left( Ct + \int_0^t \|u_\tau\|_{H^r} d\tau \right) \int_0^t \left\| (1 + \sqrt{\kappa} \langle \nabla \rangle)^{\gamma \tau} u_\tau \right\|_{H^r}^2 \|g_\tau\|_{L^2}^2 d\tau. \end{aligned}$$

Taking expectations and applying Lemma 2.6 completes the a priori estimate (3.7). For an arbitrary  $g \in L^2$ , the desired result follows by density.  $\square$

**Lemma 3.8.** *Under the assumptions of Theorem 1.2, (1.7) holds.*

*Proof.* By standard moment estimates (see e.g. [94]),  $\mathbf{E}_{\mu^\kappa} \|g\|_{H^s}^{2p} < \infty$  for  $s \in [0, \sigma - \frac{3d}{2} - 1)$  and  $p < \infty$ . By stationarity,

$$\begin{aligned} \mathbf{E}_{\mu^\kappa} \|g\|_{H^s}^{2p} &= \int \mathbf{E}_{(u,g)} \|g_1^\kappa\|_{H^s}^{2p} d\mu^\kappa(u, g) \\ &\lesssim_p \int \mathbf{E} \|S_1^\kappa(\omega, u)g\|_{H^s}^{2p} d\mu^\kappa(u, g) + \int \mathbf{E} \left\| \int_0^1 S_{1-\tau}^\kappa(\theta_\tau\omega, u_\tau)b d\xi_\tau \right\|_{H^s}^{2p} d\mu^\kappa(u, g). \end{aligned}$$

By Lemma 3.7,  $\exists \beta \geq 0$  such that for  $\forall \eta > 0$ ,  $V = V_{\beta, \eta}$ , and all  $(u, g) \in \mathbf{H} \times L^2$ ,

$$\mathbf{E} \|S_1^\kappa(\omega, u)g\|_{H^s}^{2p} \lesssim \kappa^{-sp} V^p(u) \|g\|_{L^2}^{2p}.$$

Because  $b \in C^\infty$ , it follows from Burkholder-Davis-Gundy, Minkowski's inequality, and classical  $H^s$  estimates for the transport equation (together with Lemma 2.10) that  $\exists \beta \geq 0$  such that  $\forall \eta > 0$ , there holds for suitable  $V = V_{\beta, \eta}$ ,

$$\mathbf{E} \left\| \int_0^1 S_{1-\tau}^\kappa(\theta_\tau\omega, u_\tau)b d\xi_\tau \right\|_{H^s}^{2p} \lesssim \int_0^1 \mathbf{E} \|S_{1-\tau}^\kappa(\theta_\tau\omega, u_\tau)b\|_{H^s}^{2p} d\tau \lesssim V^p(u).$$

Therefore by Lemmas 3.6, 3.7, and Hölder's inequality,

$$\mathbf{E}_{\mu^\kappa} \|g\|_{H^s}^{2p} \lesssim \mathbf{E}_{\mu^\kappa} \left[ V^p(u) \left( 1 + \kappa^{-sp} \|g\|_{L^2}^{2p} \right) \right] \lesssim_p \kappa^{-sp} |\log \kappa|^p.$$

This completes the proof of (1.7).  $\square$

## 4 Vanishing diffusivity limit

This section is devoted to the proof of Theorem 1.11 describing weak limits of the stationary measures  $\mu^\kappa$  for the passive scalar process  $(u_t, g_t^\kappa)$ . In Section 4.1 we show that such weak limits exist in the weak topology of measures on  $\mathbf{H} \times H^{-s}$  and satisfy Batchelor's Law (1.12) over an infinite inertial range.

Once these are established, it remains to show that  $\mu^0$  is a stationary measure for the passive scalar process  $(u_t, g_t^0)$ . As we check in Section 4.2, the latter can be extended to a Markov process defined by a random dynamical system on  $\mathbf{H} \times H^{-1}$ , the trajectories of which are weak solutions to the passive scalar advection equation at  $\kappa = 0$ . However, to check that weak limits  $\mu^0$  are stationary is a non-trivial task due to poor continuity properties of  $(u_t, g_t^0)$  in  $H^{-1}$  with respect to perturbations of the initial data  $u \in \mathbf{H}$  for the velocity field  $(u_t)$ , and so stationarity of  $\mu^0$  requires careful justification—see Section 4.3 for more discussion. The proof is completed in Section 4.4.

Finally we complete this section with a proof of the  $L^2$  non-vanishing flux relation (1.13).

### 4.1 Existence of weak limits of $\{\mu^\kappa\}$

Corollary 1.4 of Batchelor's law in Theorem 1.11 is that for each  $s \in (0, 1]$  we have the uniform in  $\kappa$  estimate

$$\sup_{\kappa > 0} \mathbf{E}_{\mu^\kappa} \|g\|_{H^{-s}}^{2p} \lesssim_{s,p} 1. \quad (4.1)$$

Since  $H^{-s/2}$  is compactly embedded in  $H^{-s}$  for all  $s > 0$ , we can use this to show tightness of  $\{\mu^\kappa\}$ , which permits us to take weak limits by Prokhorov's theorem. Precisely, we have the following.

**Lemma 4.1.** *There exists a sequence  $\{\kappa_n\}$ ,  $\kappa_n \rightarrow 0$  as  $n \rightarrow \infty$  and a measure  $\mu^0$  on  $\mathbf{H} \times L^2$  such that for each  $s \in (0, 1]$ , the following holds:  $\mu^\kappa \rightarrow \mu^0$  weakly as a measure on  $\mathbf{H} \times H^{-s}$ , i.e., for each  $\phi \in C_b(\mathbf{H} \times H^{-s})$  we have*

$$\int \phi d\mu^{\kappa_n} \rightarrow \int \phi d\mu^0.$$

*Proof.* Fix  $s, \epsilon > 0$  and define

$$K = \{h \in H^{-s} : \|h\|_{H^{-s/2}}^2 \leq \epsilon^{-2} \mathbf{E}_{\mu^\kappa} \|g\|_{H^{-s/2}}^2\}.$$

In light of the compact embedding  $H^{-s/2} \hookrightarrow H^{-s}$ ,  $K$  is a compact subset of  $H^{-s}$ ; by (4.1) and Chebyshev's inequality, we have

$$\sup_{\kappa \in (0, \kappa_0]} \mu^\kappa(\mathbf{H} \times (H^{-s} \setminus K)) \leq \epsilon.$$

It follows by Prokhorov's theorem that  $(\mu^\kappa)_{\kappa > 0}$  has a subsequence that converges weakly as a measure on  $\mathbf{H} \times H^{-s}$ . Choosing  $s = s_j = 2^{-j}$  we can extract a diagonal subsequence  $\kappa_n$  and a limit measure  $\mu^0$  such that  $\mu^{\kappa_n} \rightarrow \mu^0$  weakly as a measure on  $\mathbf{H} \times H^{-s_j}$  for every  $j > 0$ , hence on  $\mathbf{H} \times H^{-s}$  for every  $s \in (0, 1]$ .  $\square$

It is now straightforward to verify the infinite-inertial range version of Batchelor's law as in (1.12). With  $\mu^0 = \lim_n \mu^{\kappa_n}$  fixed as above, observe that  $g \mapsto \|\Pi_{\leq N} g\|_{L^2}$  varies continuously in  $H^{-s}$  and by a straight forward truncation argument along with the fact that

$$\sup_{\kappa \in [0, 1]} \mathbf{E}_{\mu^\kappa} \|\Pi_{\leq N} g\|_{L^2}^{2p} < \infty$$

to conclude that

$$\mathbf{E}_{\mu^0} \|\Pi_{\leq N} g\|_{L^2}^{2p} = \lim_{n \rightarrow \infty} \mathbf{E}_{\mu^{\kappa_n}} \|\Pi_{\leq N} g\|_{L^2}^{2p} \approx_p (\log N)^p,$$

for all  $N \geq N_0$ , using the fact that the inertial range  $N_0 \leq N \lesssim \kappa^{-1/2}$  grows to all of  $\{N \geq N_0\}$  as  $\kappa \rightarrow 0$ .

## 4.2 The scalar process $(u_t, g_t^0)$ at $\kappa = 0$

In this section we study the properties of the passive scalar process  $(u_t, g_t^0)$  which solves the advection equation

$$\partial_t g_t^0 + u_t \cdot \nabla g_t^0 = b \dot{\xi}_t, \quad (4.2)$$

in the absence of any diffusion term. Since the limiting measure  $\mu^0$  assigns full measure to  $H^{-1}$ , it is natural to consider  $t \mapsto g_t^0$  as a process in the negative Sobolev space  $H^{-1}$ , for which it will be necessary to show that weak solutions to (4.2) are well-posed for  $H^{-1}$  initial data.

To formulate this, recall that the solution operator  $S_t^0(\omega, u) : L^2 \rightarrow L^2$ , defined by

$$S_t^0(\omega, u)g = g \circ (\phi_{\omega, u}^t)^{-1}$$

solves (4.2) in the absence of the noise term  $b \dot{\xi}_t$ , where  $\phi_{\omega, u}^t : \mathbb{T}^d \rightarrow \mathbb{T}^d$  is the Lagrangian flow (1.17) associated to  $(u_t)$ . By incompressibility, for  $f \in H^1, g \in L^2$  we have

$$\int f S_t^0(\omega, u)g dx = \int (f \circ \phi_{\omega, u}^t)g dx. \quad (4.3)$$

**Lemma 4.2.** *The following holds*

- (a) *For each  $t > 0$  operator  $S_t^0(\omega, u)$  defined by (4.2) admits a unique extension to a bounded linear operator on  $H^{-1}$  satisfying (4.3) for all  $g \in L^2, f \in H^{-1}$  and for all  $g \in H^{-1}, t \mapsto S_t(\omega, u)g$  is strongly continuous in  $H^{-1}$ .*

(b) For initial data  $g \in H^{-1}$  the unique weak solution  $g_t^0$  to equation (4.2) is given by

$$g_t^0 = S_t^0(\omega, u)g + \int_0^t S_{t-s}^0(\theta_s\omega, u_s)b \, d\xi_s.$$

*Proof.* Part (b) follows from (a), which in turn follows from a density argument and the following estimate for  $g \in L^2$

$$\|S_t^0(\omega, u)g\|_{H^{-1}} \leq \bar{D}(\omega, u)e^{ct}\|g\|_{H^{-1}}, \quad (4.4)$$

where  $c > 0$  and for each  $p \in [1, \infty)$  we can take  $\bar{D}(\omega, u) : \Omega \times \mathbf{H} \rightarrow \mathbb{R}_{\geq 1}$  to have  $V$ -bounded  $p$ -th moment. To see this, fix  $f \in H^1, g \in L^2$ , and observe that by Grönwall's inequality and Lemma 2.11

$$\begin{aligned} |\nabla(f \circ \phi_t^{\omega, u})(x)| &\leq \exp\left(\int_0^t \|\nabla u_s\|_{L^\infty} ds\right) |\nabla f(\phi_{\omega, u}^t(x))| \\ &\leq D_0(\omega, u)e^{ct} |\nabla f(\phi_{\omega, u}^t(x))|, \end{aligned}$$

where  $c > 0$  and  $D_0(\omega, u)$  has  $V$ -bounded  $p$ -th moment. Using incompressibility of  $\phi_{\omega, u}^t$ , we obtain

$$\|f \circ \phi_{\omega, u}^t\|_{H^1} \leq D_0(\omega, u)e^{ct}\|f\|_{H^1},$$

hence

$$\left| \int S_t^0(\omega, u)gf \, dx \right| = \left| \int g(f \circ \phi_{\omega, u}^t) \, dx \right| \leq \|g\|_{H^{-1}} \|f \circ \phi_{\omega, u}^t\|_{H^1} \leq D_0(\omega, u)e^{ct}\|g\|_{H^{-1}}\|f\|_{H^1}.$$

We conclude  $\|S_t^0(\omega, u)g\|_{H^{-1}} \leq D_0(\omega, u)e^{ct}\|g\|_{H^{-1}}$ , as desired. Strong continuity of  $t \mapsto S_t^0$  follows by density of  $C^\infty(\mathbb{T}^d)$  in  $H^{-1}$ .  $\square$

### 4.3 Stationarity of $\mu^0$ : restricted Feller property of $(u_t, g_t^0)$

With the evolution  $t \mapsto g_t^0$  defined, let  $P_t^0$  denote its corresponding Markov semigroup:

$$P_t^0\phi(u, g) := \mathbf{E}_{(u, g)}\phi(u_t, g_t^0)$$

for each bounded measurable  $\phi : \mathbf{H} \times H^{-1} \rightarrow \mathbb{R}$ . We seek to show that if  $\mu^0$  is a weak limit of  $\mu^{\kappa_n}$ ,  $\kappa_n \rightarrow 0$ , then  $\mu^0$  is stationary for  $(u_t, g_t^0)$ ; equivalently it suffices to show that for all globally Lipschitz, bounded  $\phi : \mathbf{H} \times H^{-1} \rightarrow \mathbb{R}$ , we have

$$\int P_1^0\phi \, d\mu^0 = \int \phi \, d\mu^0.$$

By the weak convergence  $\mu^{\kappa_n} \rightarrow \mu^0$  and stationarity of  $\mu^{\kappa_n}$ , it suffices to show that

$$\lim_{n \rightarrow \infty} \int P_1^{\kappa_n}\phi \, d\mu^{\kappa_n} = \int P_1^0\phi \, d\mu^0,$$

where  $P_t^\kappa$  is the Markov semigroup on  $\mathbf{H} \times L^2$  for the process  $(u_t, g_t^\kappa)$ . To this end the strategy is to write

$$\left| \int P_1^{\kappa_n}\phi \, d\mu^{\kappa_n} - \int P_1^0\phi \, d\mu^0 \right| \leq \left| \int (P_1^{\kappa_n}\phi - P_1^0\phi) \, d\mu^{\kappa_n} \right| + \left| \int P_1^0\phi \, (d\mu^{\kappa_n} - d\mu^0) \right| \quad (4.5)$$

and show that each term vanishes as  $n \rightarrow \infty$ . The first term is relatively straightforward to bound, having to do with the nearness of  $g_t^{\kappa_n}, g_t^0$  when initiated at the same initial condition  $g \in L^2$  and subjected to the same noise sample  $b\xi_t$ . This is dealt with by Lemma 4.4 below.

The second term in (4.5) is more challenging. One would like to use weak convergence to justify passing the limit using weak convergence  $\mu^{\kappa_n} \rightarrow \mu^0$ , but this will not work: to the best of our knowledge, the semigroup  $P_t^0$  for  $(u_t, g_t^0)$  does not send continuous functions on  $\mathbf{H} \times H^{-1}$  to continuous functions  $\mathbf{H} \times H^{-1}$ , namely  $P_t^0$  is not *Feller* on  $\mathbf{H} \times H^{-1}$ . When  $\kappa > 0$  it is a straightforward consequence of parabolic regularity that the semigroup  $P_t^\kappa$  is Feller on  $\mathbf{H} \times L^2$ . However, when  $\kappa = 0$  this mechanism is not available and consequently  $P_t^0$  is not necessarily Feller on  $\mathbf{H} \times H^{-1}$  due to a lack of stability of the transport equation in  $H^{-1}$  under perturbations of the velocity field.

Instead, we will show the following *restricted Feller property*: namely  $P_t^0$  has the Feller property in  $\mathbf{H} \times H^{-1}$  when restricted to initial  $g$  in a bounded subset of  $L^2$ . We state this property below for Lipschitz observables.

**Lemma 4.3** (Restricted Feller). *Let  $\phi \in \text{Lip}(\mathbf{H} \times H^{-1})$ , then for all  $u, u' \in \mathbf{H}$  and  $g, g' \in L^2$  we have*

$$|P_1^0 \phi(u, g) - P_1^0 \phi(u', g')| \lesssim \|\phi\|_{\text{Lip}} W(u, g, u', g') (\|u - u'\|_{\mathbf{H}} + \|g - g'\|_{H^{-1}}),$$

where

$$W(u, g, u', g') = (V(u) + V(u'))(1 + \|g\|_{L^2})(1 + \|g'\|_{L^2}),$$

and  $V = V_{\beta, \eta}$  for a sufficiently large universal  $\beta > 0$  and all  $\eta > 0$ .

*Proof.* To prove this it suffices to consider two cases: (i)  $u = u'$  and  $g \neq g'$  and (ii)  $u \neq u'$ ,  $g' = g$ .

Case (i) is straightforward since the difference between any two solutions  $g_t$  and  $g'_t$  to (4.2) with different initial data and the same noise and same velocity field  $(u_t)$  immediately satisfy

$$g_t - g'_t = S_t^0(\omega, u)(g - g'),$$

hence by (4.4),

$$\begin{aligned} |P_1^0 \phi(u, g) - P_1^0 \phi(u, g')| &\leq \|\phi\|_{\text{Lip}} \mathbf{E} \bar{D}(u, \cdot) \|g - g'\|_{H^{-1}} \\ &\lesssim \|\phi\|_{\text{Lip}} V(u) \|g - g'\|_{H^{-1}}. \end{aligned}$$

For case (ii) let  $u_t$  and  $u'_t$  be two solutions to (1) with initial data  $u$  and  $u'$  respectively and  $g_t$  and  $g'_t$  the solutions to (4.2) with velocity fields  $(u_t)$  and  $(u'_t)$  respectively and the same initial data  $g \in H^1$ . Then the difference  $\tilde{g}_t = g_t - g'_t$  satisfies

$$\partial_t \tilde{g}_t + u_t \cdot \nabla \tilde{g}_t + (u_t - u'_t) \cdot \nabla g'_t = 0, \quad \tilde{g}_0 = 0,$$

and therefore can be written as

$$\tilde{g}_t = \int_0^t S_{t-s}^0(\theta_s \omega, u_s) [(u_s - u'_s) \cdot \nabla g'_s] ds.$$

It follows from (4.4) (choosing  $\bar{D}$  to have a  $V$ -bounded 4th moment) that for  $0 \leq s < t \leq 1$

$$\begin{aligned} \|S_{t-s}^0(\theta_s \omega, u_s) [(u_s - u'_s) \cdot \nabla g'_s]\|_{H^{-1}} &\leq \bar{D}(\theta_s \omega, u_s) \|(u_s - u'_s) \cdot \nabla g'_s\|_{H^{-1}} \\ &\leq \bar{D}(\theta_s \omega, u_s) \|u_s - u'_s\|_{W^{1, \infty}} \|g'_s\|_{L^2}, \end{aligned}$$

where in the last inequality we used that if  $f \in W^{1, \infty}$  and  $h \in H^{-1}$  then  $\|fh\|_{H^{-1}} \leq \|f\|_{W^{1, \infty}} \|h\|_{H^{-1}}$ , which can be seen by duality between  $H^1$  and  $H^{-1}$ . Consequently we have

$$\mathbf{E} \|\tilde{g}_1\|_{H^{-1}} \leq \int_0^1 \mathbf{E} [\bar{D}(\theta_s \omega, u_s) \|u_s - u'_s\|_{\mathbf{H}} \|g'_s\|_{L^2}] ds$$



We use Hölder's inequality under the expectation  $\mathbf{E}$ . The 4th moment  $\mathbf{E}\bar{D}(\omega, u)^4$  is bounded  $\lesssim V(u)$  by construction, while by Lemma 2.8 we have  $\mathbf{E}\|u_s - u'_s\|_{\mathbf{H}}^4 \lesssim (V(u) + V(u'))\|u - u'\|_{\mathbf{H}}$ . To bound  $\mathbf{E}\|g'_s\|_{L^2}^2$ , we estimate as follows: by Itô's formula,

$$\|g'_s\|_{L^2}^2 = \|g\|_{L^2}^2 + 2\chi s + \int_0^s \langle g'_r, b \rangle_{L^2} d\xi_r.$$

By taking  $\mathbf{E}$  of both sides, the stochastic integral vanishes, hence  $\mathbf{E}\|g'_s\|_{L^2}^2 = \|g\|_{L^2}^2 + 2\chi s$ . Putting this together, we conclude for suitable  $V(u)$

$$|P_1\phi(u, g) - P_1\phi(u', g)| \leq \|\phi\|_{\text{Lip}} \mathbf{E}\|\tilde{g}_1\|_{H^{-1}} \lesssim_p \|\phi\|_{\text{Lip}} (V(u) + V(u'))(1 + \|g\|_{L^2})\|u - u'\|_{\mathbf{H}}.$$

□

#### 4.4 Stationarity of $\mu^0$ : completing the proof

It suffices to show that the right-hand side of (4.5) vanishes. The first term  $|\int (P_1^{\kappa n}\phi - P_1^0\phi)d\mu^{\kappa n}|$  in (4.5) will be estimated using the following.

**Lemma 4.4.** *Let  $g_t^\kappa$  and  $g_t^0$  have the same initial data  $g \in L^2$ , the same velocity field  $(u_t)$ , and the same noise sample  $b\xi_t$ . Then,*

$$\mathbf{E}\|g_1^\kappa - g_1^0\|_{H^{-1}} \lesssim V(u) \int_0^1 \kappa \mathbf{E}\|g_s^\kappa\|_{H^1} ds.$$

*Proof.* Note that  $\tilde{g}_t^\kappa = g_t^\kappa - g_t^0$  satisfies

$$\partial_t \tilde{g}_t^\kappa + u_t \cdot \nabla \tilde{g}_t^\kappa - \kappa \Delta g_t^\kappa = 0.$$

with  $\tilde{g}_0^\kappa = 0$ . This implies that

$$\tilde{g}_1^\kappa = \kappa \int_0^1 S_{t-s}^0(\theta_s \omega, u_s) \Delta g_s^\kappa ds,$$

and therefore using the  $H^{-1}$  bound (4.4) on  $S_{t-s}^0(\theta_s \omega, u_s)$  we find

$$\|\tilde{g}_1^\kappa\|_{H^{-1}} \lesssim \kappa \int_0^1 \bar{D}(\theta_s \omega, u_s) \|g_s^\kappa\|_{H^1} ds,$$

where  $\bar{D}(\omega, u)$  has  $V$ -bounded 1st moment. Taking an expectation w.r.t.  $\omega$  and applying Lemma 2.6 completes the proof. □

From here, to prove convergence  $\int (P_1^{\kappa n}\phi - P_1^0\phi)d\mu^{\kappa n} \rightarrow 0$ , we have from Lemma 4.4 and the Lipschitz property of  $\phi$  that

$$|(P_1^\kappa\phi - P_1^0\phi)(u, g)| \leq \|\phi\|_{\text{Lip}} \mathbf{E}\|g_1^\kappa - g_1^0\|_{H^{-1}} \lesssim V(u) \int_0^1 \kappa \mathbf{E}\|g_s^\kappa\|_{H^1} ds,$$

hence upon integrating in  $\mu^\kappa$  and using Cauchy-Schwarz,

$$\int |P_1^\kappa\phi - P_1^0\phi| d\mu^\kappa \lesssim \kappa \int_0^1 (\mathbf{E}_{\mu^\kappa} \|g\|_{H^1}^2)^{1/2} ds \lesssim \sqrt{\kappa}$$

by the energy balance relation  $\kappa \mathbf{E}_{\mu^\kappa} \|g_t\|_{H^1}^2 = \chi$  (see (1.4)).

It remains to show that  $|\int P_1^0\phi(d\mu^{\kappa n} - d\mu^0)| \rightarrow 0$  in (4.5). As observed already, this does not follow immediately from weak convergence since  $P_1^0\phi$  need not be continuous. Instead, we will use the restricted

Feller property from Lemma 4.3, which guarantees continuity of  $P_1^0\phi$  along  $g \in L^2$ . To bridge the gap from  $L^2$  to  $H^{-1}$ , we use the following mollification argument: for  $\delta \in (0, 1)$  and  $g \in H^{-s}$ ,  $s \in (0, 1]$ , define  $T_\delta g$  to be the mollification<sup>8</sup> of  $g$ , and define the regularized semigroup

$$(P_1^0\phi)_\delta(u, g) := P_1^0\phi(u, T_\delta g).$$

Note that a straight-forward duality argument shows that

$$\|T_\delta g - g\|_{H^{-1}} \lesssim \delta^{1-s} \|g\|_{H^{-s}}.$$

This, in turn gives rise to the following approximation property for  $(P_1^0\phi)_\delta$ .

**Lemma 4.5.** *For each  $\delta \in (0, 1)$ ,  $s \in (0, 1]$ ,  $(P_1^0\phi)_\delta \in C_b(\mathbf{H} \times H^{-1})$  and for each globally Lipschitz  $\phi : \mathbf{H} \times H^{-1} \rightarrow \mathbb{R}$  and  $g \in H^{-s}$ , we have*

$$|(P_1\phi)_\delta - P_1\phi|(u, g) \lesssim V(u)\delta^{1-s} \|g\|_{H^{-s}},$$

where  $V = V_{\beta, \eta}$  for any  $\beta > 0$  and  $\eta > 0$ .

*Proof.* Note that Lemma 4.3 implies  $(P_1^0\phi)_\delta \in C_b(\mathbf{H} \times H^{-1})$ . For  $g \in H^{-s}$ , denote  $g_1^\delta$  and  $g_1$  solutions at time  $t = 1$  of the transport equation (4.2) with the same velocity path  $(u_t)$  and noise  $b\xi_t$  but with different initial data  $T_\delta g$  and  $g$  respectively. We have

$$\begin{aligned} |((P_1^0\phi)_\delta - P_1^0\phi)(u, g)| &\leq \mathbf{E}|\phi(u_1, g_1^\delta) - \phi(u_1, g_1)| \\ &\leq \|\phi\|_{\text{Lip}} \mathbf{E}\|S_1^0(u, \omega)(T_\delta g - g)\|_{H^{-1}} \\ &\lesssim V(u) \|T_\delta g - g\|_{H^{-1}} \\ &\lesssim V(u) \delta^{1-s} \|g\|_{H^{-s}} \end{aligned}$$

having used (4.4) and the  $V$ -boundedness of the 1st moment of  $\bar{D}$ . □

To complete the proof: for  $\delta > 0$ , we estimate

$$\left| \int P_1^0\phi(d\mu^{\kappa_n} - d\mu^0) \right| \leq \left| \int (P_1^0\phi)_\delta(d\mu^{\kappa_n} - d\mu^0) \right| + \int |(P_1^0\phi)_\delta - P_1^0\phi|(d\mu^{\kappa_n} + d\mu^0)$$

The first term on the right vanishes as  $n \rightarrow \infty$  since  $(P_1\phi)_\delta \in C_b(\mathbf{H} \times H^{-1})$  by Lemma 4.2 and  $\mu^{\kappa_n} \rightarrow \mu^0$  weakly. For the second term, Lemma 4.5 and Cauchy-Schwarz imply

$$\int |(P_1\phi)_\delta - P_1\phi|(d\mu^{\kappa_n} + d\mu^0) \lesssim \delta^{1-s} \left( (\mathbf{E}_{\mu^{\kappa_n}} \|g\|_{H^{-s}}^2)^{1/2} + (\mathbf{E}_{\mu^0} \|g\|_{H^{-s}}^2)^{1/2} \right)$$

which converges to 0 as  $\delta \rightarrow 0$  uniformly in  $\kappa_n$  by (4.1). This completes the proof of stationarity of  $\mu^0$  for  $P_t^0$ , the last item remaining from Theorem 1.11.

<sup>8</sup>Here mollification is defined by convolution with  $\eta_\delta = \delta^{-d}\eta(\cdot/\delta)$ , where  $\eta$  is a smooth, symmetric, compactly supported test function with  $\int \eta = 1$ .

## 4.5 Non-vanishing flux

We conclude this section by proving the non-vanishing flux law (1.13). To do this, we remark that  $\Pi_{\leq N} g_t^\kappa$  satisfies

$$\Pi_{\leq N} g_t^\kappa = \Pi_{\leq N} g_0^\kappa - \int_0^t \Pi_{\leq N} (u_s \cdot \nabla g_s^\kappa) ds + \Pi_{\leq N} b \xi_t,$$

and therefore applying Itô's formula gives

$$\begin{aligned} \|\Pi_{\leq N} g_t^0\|_{L^2}^2 &= \|\Pi_{\leq N} g_0^0\|_{L^2}^2 - 2 \int_0^t \langle \Pi_{\leq N} g_s, \Pi_{\leq N} (u_s \cdot \nabla g_s) \rangle_{L^2} ds \\ &\quad + \|\Pi_{\leq N} b\|_{L^2}^2 t + \int_0^t 2 \langle \Pi_{\leq N} g_s, \Pi_{\leq N} b \rangle_{L^2} d\xi_s. \end{aligned}$$

Taking expectation, integrating the initial data  $(u, g)$  with respect to the stationary measure  $\mu^0$  and using the stationarity of  $(u_t, g_t^0)$  with respect to  $\mu^0$  we readily obtain the flux balance

$$\mathbf{E}_{\mu^\kappa} \langle \Pi_{\leq N} g, \Pi_{\leq N} (u \cdot \nabla g) \rangle_{L^2} = \frac{1}{2} \|\Pi_{\leq N} b\|_{L^2}^2.$$

Using the divergence free property and integrating by parts in the  $L^2$  inner product gives (1.13).

## 5 Irregularity of the limiting statistically stationary solutions

In the previous section, we produced a  $\kappa \rightarrow 0$  subsequential limit of  $\mu^\kappa$  which converges to a stationary measure  $\mu^0$  of the Markov semigroup associated with  $(u_t, g_t^0)$ , where the scalar solves

$$\partial_t g_t^0 + u_t \cdot \nabla g_t^0 = b \dot{\xi}_t. \quad (5.1)$$

Since there is no dissipation in the equation, the input from the noise must be ‘‘anomalously’’ dissipated by the mixing mechanism. As discussed in Section 1.2.4, this requires a degree of roughness of statistically stationary  $g_t^0$ . Specifically, in this section we prove

- (a)  $\mu^0$ -generic  $(u, g) \in \mathbf{H} \times H^- = \mathbf{H} \times \cap_{\delta>0} H^{-\delta}$  are such that  $g$  is not locally integrable, hence are ‘strictly’ distributions (Theorem 1.14);
- (b) the Besov space  $L_t^2 B_{2,\infty}^0$  is ‘‘Onsager-type critical’’, in the sense that no statistically stationary solution  $g_t^0$  can have even a small amount of additional regularity (Theorem 1.18).

Part (a) is carried out by applying a variant of the DiPerna-Lions theory of renormalized solutions for the transport equation, and is carried out in Section 5.1, while part (b) is carried out in Section 5.2.

### 5.1 $\mu^0$ generic functions cannot be $L^1$

Our proof is by contradiction, based on invariance of  $\mathbf{H} \times L^1$  by solutions to (5.1).

**Lemma 5.1.** *The set  $\mathbf{H} \times L^1$  is almost surely invariant for  $(u_t, g_t^0)$ : if  $u_0 \in \mathbf{H}$  and  $g_0^0 \in L^1$ , then  $g_t^0 \in L^1$  for all  $t > 0$ .*

*Proof.* This follows from Lemma 4.2(b) by noting that due to smoothness of  $b$ , the stochastic convolution

$$\int_0^t S_{t-s}^0(\theta_s \omega, u_s) b d\xi_s$$

takes values in (at least)  $L^2$ , and  $S_t^0$  propagates  $L^1$  regularity. In fact, we have

$$\|S_t^0(\omega, u)g\|_{L^1} = \|g \circ (\phi_{\omega, u}^t)^{-1}\|_{L^1} = \|g\|_{L^1}$$

by the fact that  $\phi_{\omega, u}^t$  is volume preserving. □

To continue, we will find it convenient to show that any  $L^1$  valued solutions to (5.1) can be renormalized in the sense of DiPerna and Lions [50], meaning that we show that  $\beta(g_t^0)$  solves another transport-type equation for some suitably regular function  $\beta : \mathbb{R} \rightarrow \mathbb{R}$ . In our setting the velocity field  $(u_t)$  is not rough, which allows for the transport part of the equation to be easily renormalized, but the presence of noise introduces Itô corrections and imposes higher regularity requirements on  $\beta(z)$ . In what follows we define

$$\beta(z) := \sqrt{1 + |z|^2}.$$

**Lemma 5.2.** *Let  $g^0 \in L^1$ , and  $g_t^0$  be a mild solution to (5.1), then the following holds almost surely*

$$\int \beta(g_t^0) dx = \int \beta(g_0^0) dx + \int_0^t \left( \int \frac{b^2}{2\beta(g_s^0)^3} dx \right) ds + \int_0^t \left( \int \frac{g_s^0 b}{\beta(g_s^0)} dx \right) d\xi_s. \quad (5.2)$$

*Proof.* For a given function  $f \in L^1$ , we denote  $(f)_\delta$  the mollification of  $f$  (see the proof of Lemma 4.5). It is straightforward to show that  $(f)_\delta \rightarrow f$  both in  $L^1$  and pointwise a.e.. Since  $g_t^0$  is a weak solution to (5.1) we see that  $(g_t^0)_\delta$  solves the following equation almost surely:

$$(g_t^0)_\delta = (g_0^0)_\delta - \int_0^t u_s \cdot \nabla (g_s^0)_\delta ds + (b)_\delta \xi_t + \int_0^t R_\delta(u_s, g_s^0) ds,$$

where, we write

$$R_\delta(u_t, g_t^0) = u_t \cdot \nabla (g_t^0)_\delta - (u_t \cdot \nabla g_t^0)_\delta$$

for the commutator term, noting that convolution  $(\cdot)_\delta$  does not commute with  $u_t \cdot \nabla$ . Since the equation for  $(g_t^0)_\delta$  holds pointwise as an identity on smooth functions for each  $x \in \mathbb{T}^d$ , we can apply Itô's formula to  $\beta((g_t^0)_\delta(x))$  to deduce the pointwise identity

$$\begin{aligned} \beta((g_t^0)_\delta) &= \beta((g_0^0)_\delta) - \int_0^t u_s \cdot \nabla \beta((g_s^0)_\delta) ds + \int_0^t \frac{1}{2} \beta''((g_s^0)_\delta) (b)_\delta^2 ds + \int_0^t \beta'((g_s^0)_\delta) (b)_\delta d\xi_s \\ &\quad + \int_0^t \beta'((g_s^0)_\delta) R_\delta(u_s, g_s^0) ds. \end{aligned} \quad (5.3)$$

Above, we have used the fact  $(g_t^0)_\delta$  is smooth and therefore  $\beta'(g_t^0) u_t \cdot \nabla (g_t^0)_\delta = u_t \cdot \nabla \beta((g_t^0)_\delta)$ . Itô's formula for the unbounded function  $\beta(z) = \sqrt{1 + |z|^2}$  is justified in this case because  $\beta((g_t^0)_\delta) \lesssim_\delta 1 + \|g_t^0\|_{L^1}$ , hence for each  $t \in \mathbb{R}_+$ ,

$$\mathbf{P} \left( \sup_{s \in [0, t]} \|\beta((g_s^0)_\delta)\|_{L^\infty} < \infty \right) = 1.$$

Next, we note that since  $\operatorname{div} u_t = 0$ , it is standard that (see [50])

$$\int |R_\delta(u_t, g_t^0)| dx \leq \|u_t\|_{W^{1, \infty}} \left( \int_{B_\delta} |y \cdot \nabla \eta_\delta| dy \right) \sup_{|y| < \delta} \|g_t^0(\cdot + y) - g_t^0(\cdot)\|_{L^1}.$$

Since  $\int |y \cdot \nabla \eta_\delta| dy = \int |y \cdot \nabla \eta| dy$ , and  $g_t^0$  belongs to  $L^1$  almost surely we conclude by the  $L^1$ -continuity of spatial translations that for each  $t \in \mathbb{R}_+$

$$R_\delta(u_t, g_t^0) \rightarrow 0 \quad \text{in } L^1 \text{ as } \delta \rightarrow 0 \text{ almost surely.} \quad (5.4)$$

Using the fact

$$\beta'(z) = \frac{z}{\beta(z)}, \quad \beta''(z) = \frac{1}{\beta(z)^3}$$

we can integrate (5.3) and use the fact that  $u_t$  is divergence free to obtain

$$\begin{aligned} \int \beta((g_t^0)_\delta) dx &= \int \beta((g_0^0)_\delta) dx + \int_0^t \left( \int \frac{1}{2} \frac{(b)_\delta^2}{\beta((g_s^0)_\delta)^3} dx \right) ds + \int_0^t \left( \int \frac{(b)_\delta (g_t^0)_\delta}{\beta((g_s^0)_\delta)} dx \right) d\xi_t \\ &\quad + \int_0^t \left( \int \frac{(g_t^0)_\delta}{\beta((g_t^0)_\delta)} R_\delta(u_s, g_s^0) dx \right) ds. \end{aligned} \quad (5.5)$$

The proof will be complete if we can pass the  $\delta \rightarrow 0$  limit in both sides of (5.5) almost-surely for each fixed  $t$ . Since we have  $(g_t^0)_\delta \rightarrow g_t^0$  in  $L^1$  and almost everywhere on  $\mathbb{T}^d$ , as well as the convergence of the commutator (5.4) in  $L^1$  and  $(b)_\delta \rightarrow b$  in  $L^2$ , we can use bounded convergence to pass the  $\delta \rightarrow 0$  limits almost-surely in every term of (5.5) except for the stochastic integral. To deal with the stochastic integral, we note that by Itô's formula

$$\mathbf{E} \left| \int_0^t \left( \int \frac{(b)_\delta (g_t^0)_\delta}{\beta((g_s^0)_\delta)} - \frac{b g_t^0}{\beta(g_s^0)} dx \right) d\xi_t \right|^2 = \mathbf{E} \int_0^t \int \left| \frac{(b)_\delta (g_t^0)_\delta}{\beta((g_s^0)_\delta)} - \frac{b g_t^0}{\beta(g_s^0)} \right|^2 dx ds \rightarrow 0$$

as  $\delta \rightarrow 0$  by bounded convergence. □

We are now ready to prove Theorem 1.14.

**Proof of Theorem 1.14.** In pursuit of a contradiction, we assume that  $\mu^0(\mathbf{H} \times L^1) > 0$ . By Lemma 5.1,  $\mathbf{H} \times L^1$  is an invariant set for  $(u_t, g_t^0)$ , and so the conditional measure

$$\hat{\mu}^0 := \frac{\mu^0(\cdot \cap (\mathbf{H} \times L^1))}{\mu^0(\mathbf{H} \times L^1)}$$

is another stationary measure for  $(u_t, g_t^0)$  that assigns full measure to  $\mathbf{H} \times L^1$ .

We would like to conclude the proof by taking expectations of both sides of (5.2) using stationarity with respect to  $\hat{\mu}^0$ . However, we are unable to justify this, since a priori we do not know whether or not  $\beta(g_t^0)$  has finite moments. To get around this, we apply Itô's formula to  $\phi\left(\int \beta(g_t^0) dx\right)$ , where  $\phi(z)$  is a bounded  $C^2$  function, to deduce

$$\begin{aligned} \phi\left(\int \beta(g_t^0) dx\right) &= \phi\left(\int \beta(g_0^0) dx\right) + \int_0^t \left( \int \frac{b^2}{2\beta(g_s^0)^3} dx \right) \phi'\left(\int \beta(g_s^0) dx\right) ds \\ &\quad + \frac{1}{2} \int_0^t \left( \int \frac{g_s^0 b}{\beta(g_s^0)} dx \right)^2 \phi''\left(\int \beta(g_s^0) dx\right) ds \\ &\quad + \int_0^t \left( \int \frac{g_s^0 b}{\beta(g_s^0)} dx \right) \phi'\left(\int \beta(g_s^0) dx\right) d\xi_s. \end{aligned}$$

Taking expectation and integrating the initial data with respect to  $\hat{\mu}^0$  (crucially using that  $\hat{\mu}^0$  assigns full measure to  $L^1$ ), we find

$$\mathbf{E}_{\hat{\mu}^0} \left[ \left( \int \frac{b^2}{\beta(g)^3} dx \right) \phi'\left(\int \beta(g) dx\right) + \left( \int \frac{gb}{\beta(g)} dx \right)^2 \phi''\left(\int \beta(g) dx\right) \right] = 0.$$

For  $\epsilon > 0$ , define  $\phi(z) = \frac{z}{1+\epsilon z}$ , so that

$$\mathbf{E}_{\hat{\mu}^0} \frac{\int \frac{b^2}{\beta(g)^3} dx}{(1 + \epsilon \int \beta(g) dx)^2} = \mathbf{E}_{\hat{\mu}^0} \frac{2\epsilon \left( \int \frac{gb}{\beta(g)} dx \right)^2}{(1 + \epsilon \int \beta(g) dx)^3} \leq 2\epsilon \int b^2 dx.$$

Sending  $\epsilon \rightarrow 0$  and applying monotone convergence gives

$$\mathbf{E}_{\hat{\mu}^0} \int \frac{b^2}{\beta(g)^3} dx = 0,$$

which contradicts  $\hat{\mu}^0(\mathbf{H} \times L^1) = 1$ . Therefore  $\mu^0(\mathbf{H} \times L^1) = 0$ , which completes the proof.  $\square$

## 5.2 Onsager-type criticality of $L_t^2 B_{2,\infty}^0$

In this section we prove Theorem 1.18. Recall Definition 1.17 of a  $B_{2,c}^0$ -suitable multiplier  $M$ , and the associated generalized Besov norm

$$\|f\|_{B_{2,\infty}^M} = \sup_{N \in \{2^j : j \in \mathbb{N}_*\}} M(N) \|\Pi_N f\|_{L^2}.$$

We denote  $B_{2,\infty}^M$  the associated space of tempered distributions  $f$  with  $\|f\|_{B_{2,\infty}^M} < \infty$ . The following Lemma shows that we can exhaust  $B_{2,c}^0$  in terms of the spaces  $\{B_{2,\infty}^M\}$ .

**Lemma 5.3.** *The following holds*

$$B_{2,c}^0 = \bigcup_M B_{2,\infty}^M,$$

where the union is over all  $B_{2,c}^0$ -suitable multipliers  $M$ .

*Proof.* That  $\|f\|_{B_{2,\infty}^M} < \infty \Rightarrow f \in B_{2,c}^0$  follows from  $\lim_{k \rightarrow \infty} M(k) = \infty$ .

Conversely, let  $g \in B_{2,c}^0$ . Let  $C_0$  be such that  $\sup_{j \in \mathbb{N}_*} \|\Pi_{2^j} g\|_{L^2} \leq C_0$ . By  $g \in B_{2,c}^0$ , there exists a strictly increasing sequence  $\{N_k\}_{k=1}^\infty \subseteq \{2^j : j \in \mathbb{N}\}$  such that  $\forall N \geq N_k$ ,  $\|\Pi_N g\|_{L^2} < 2^{-k}$ . Define  $M(k) : [1, \infty) \rightarrow [1, \infty)$  to be the monotone increasing multiplier

$$M(k) = \begin{cases} 2 & k < N_1 \\ \frac{2^j}{N_{j+1}-N_j} (N_{j+1} - k) + \frac{2^{j+1}}{N_{j+1}-N_j} (k - N_j) & N_j \leq k \leq N_{j+1}. \end{cases}$$

We see that  $M$  is piecewise linear with slope  $\leq 1$ , and therefore satisfies conditions (i) and (ii) of Definition 1.17.  $\square$

We introduce Littlewood-Paley decomposition for future use. Let  $\zeta \in C_0^\infty(\mathbb{R}; \mathbb{R})$  be such that  $\zeta(\xi) = 1$  for  $|\xi| \leq 1$  and  $\zeta(\xi) = 0$  for  $|\xi| \geq 3/2$  and define  $\psi(\xi) = \zeta(\xi/2) - \zeta(\xi)$ , supported in the range  $\xi \in (1, 3)$ . Denote for each  $N > 0$ ,  $\zeta_N(\xi) := \zeta(N^{-1}\xi)$  and  $\psi_N(\xi) := \psi(N^{-1}\xi)$ . For  $f \in L^2(\mathbb{R})$  we define the Littlewood-Paley projections

$$\pi_N f := \psi_N(\nabla) f \quad \text{and} \quad \pi_{\leq N} f := \zeta_N(\nabla) f.$$

Note by Definition 1.17, (denoting  $\mathbb{N}_* = \{0\} \cup \mathbb{N}$  and analogously for  $\mathbb{Z}_*^d = \{0\} \cup \mathbb{Z}^d$  below).

$$\sup_{N \in \{2^j : j \in \mathbb{N}_*\}} M(N) \|\Pi_N g\|_{L^2} \approx \|M(|\nabla|) \pi_{\leq 1} g\|_{L^2} + \sup_{N \in \{2^j : j \in \mathbb{N}_*\}} \|M(|\nabla|) \pi_N g\|_{L^2}.$$

**Lemma 5.4.** *For any  $(u, g) \in \mathbf{H} \times H^{-1}$ , if  $\|g\|_{B_{2,\infty}^M} < \infty$  for some  $B_{2,c}^0$ -suitable  $M$ , then*

$$|\langle \pi_{\leq N} g, \pi_{\leq N} \nabla \cdot (ug) \rangle| \lesssim \frac{1}{M^2(N)} \|g\|_{B_{2,\infty}^M}^2 \|u\|_{\mathbf{H}}.$$

*Proof.* By incompressibility of  $u$  and Plancherel's identity

$$\begin{aligned}
& \langle \pi_{\leq N} g, \pi_{\leq N} \nabla \cdot (ug) \rangle \\
&= \langle \pi_{\leq N} g, \pi_{\leq N} \nabla \cdot (ug) - \nabla \cdot (u\pi_{\leq N} g) \rangle \\
&= \sum_{k, \ell \in \mathbb{Z}_*^d} (\mathbf{1}_{|\ell| < 2|k-\ell|} + \mathbf{1}_{|\ell| \geq 2|k-\ell|}) \zeta_N(k) \overline{\hat{g}_t(k)} (\zeta_N(k) - \zeta_N(\ell)) ik \cdot \hat{u}_t(k-\ell) \hat{g}_t(\ell) \\
&=: \mathcal{I}_{HL} + \mathcal{I}_{LH}.
\end{aligned}$$

Note that due to the presence of the cutoffs  $(\zeta_N(k) - \zeta_N(\ell))$  we see that one of  $|k|$  or  $|\ell|$  must be larger or equal to  $\frac{1}{2}N$  for the corresponding term in the summation to be non-zero. The ‘‘high-low’’ term is treated by noting that on the support of the summation,  $|k| + |\ell| \lesssim |k - \ell|$ , and therefore for any  $\delta > 0$ ,

$$\begin{aligned}
|\mathcal{I}_{HL}| &\lesssim \frac{1}{N^2} \sum_{k, \ell \in \mathbb{Z}_*^d} \mathbf{1}_{|\ell| < 2|k-\ell|} \left| \zeta_N(k) \hat{g}_t(k) \langle k - \ell \rangle^3 \hat{u}_t(k - \ell) \hat{g}_t(\ell) \right| \\
&\lesssim \frac{1}{N^2} \sum_{k, \ell \in \mathbb{Z}_*^d} \mathbf{1}_{|\ell| < 2|k-\ell|} \frac{1}{\langle k \rangle^\delta \langle \ell \rangle^\delta} \left| \zeta_N(k) \hat{g}_t(k) \langle k - \ell \rangle^{3+2\delta} \hat{u}_t(k - \ell) \hat{g}_t(\ell) \right|.
\end{aligned}$$

Then, using Cauchy-Schwarz, Young's inequality, and  $H^r \hookrightarrow \widehat{L}^1$  for  $r > d/2$ , (recall  $\sigma > 3 + \frac{d}{2}$ ),

$$|\mathcal{I}_{HL}| \lesssim_\delta \frac{1}{N^2} \|g\|_{H^{-\delta}}^2 \|u\|_{\mathbf{H}} \lesssim \frac{1}{M(N)^2} \|g\|_{B_{2,\infty}^M}^2 \|u\|_{\mathbf{H}},$$

where the last inequality followed from  $M(N) \lesssim N$  by Definition 1.17 and the embedding  $H^{-\delta} \hookrightarrow B_{2,\infty}^0$ .

We turn next to the ‘‘low-high’’ term. First, by the mean value theorem there holds,

$$|\zeta_N(k) - \zeta_N(\ell)| \lesssim \frac{1}{N} |k - \ell|. \quad (5.6)$$

Second, observe that on the support of the summation, we have  $N \approx |k| \approx |\ell|$  because of the frequency cut-off and that at least one of  $|k|$  or  $|\ell|$  must be larger or equal to  $\frac{1}{2}N$  but one of  $|k|$  and  $|\ell|$  must also be less than  $3N$ . Therefore, we deduce from part (ii) of Definition 1.17 and (5.6),

$$|\mathcal{I}_{LH}| \lesssim \frac{1}{M(N)^2} \sum_{k, \ell \in \mathbb{Z}_*^d} \mathbf{1}_{|\ell| \geq 2|k-\ell|} |\zeta_N(k) M(|k|) \hat{g}_t(k) \langle k - \ell \rangle \hat{u}_t(k - \ell) M(|\ell|) \hat{g}_t(\ell)|.$$

Again by using Cauchy-Schwarz, Young's inequality, and  $H^r \hookrightarrow \widehat{L}^1$  for  $r > d/2$ ,

$$|\mathcal{I}_{LH}| \lesssim \frac{1}{M(N)^2} \|g\|_{B_{2,\infty}^M}^2 \|u\|_{\mathbf{H}}.$$

This completes the proof.  $\square$

**Proof of Theorem 1.18.** We proceed by contradiction and assume  $\mathbf{E}_{\mu^0} \|g\|_{B_{2,\infty}^M}^p < \infty$  for some  $p > 2$ . By repeating the proof of the flux balance (1.13) as in Section 4.5 with  $\pi_{\leq N}$  replacing  $\Pi_{\leq N}$  we have

$$\mathbf{E}_{\mu^0} \langle \pi_{\leq N} g, \pi_{\leq N} \nabla \cdot (ug) \rangle = \frac{1}{2} \|\pi_{\leq N} b\|_{L^2}^2. \quad (5.7)$$

However Lemma 5.4 and the assumption that  $\mathbf{E}_{\mu^0} \|g\|_{B_{2,\infty}^M}^p < \infty$  for some  $p > 2$  implies that

$$\begin{aligned} |\mathbf{E}_{\mu^0} \langle \pi_{\leq N} g, \pi_{\leq N} \nabla \cdot (ug) \rangle| &\lesssim \frac{1}{M(N)^2} \mathbf{E}_{\mu^0} \left[ \|g\|_{B_{2,\infty}^M}^2 \|u\|_{\mathbf{H}} \right] \\ &\lesssim \frac{1}{M(N)^2} \left( \mathbf{E}_{\mu^0} \|g\|_{B_{2,\infty}^M}^p \right)^{2/p} \left( \mathbf{E} \|u\|_{\mathbf{H}}^{\frac{p}{p-2}} \right)^{\frac{p-2}{p}} \\ &\lesssim \frac{1}{M(N)^2}. \end{aligned}$$

On the other hand,  $\lim_{N \rightarrow \infty} \frac{1}{2} \|\pi_{\leq N} b\|_{L^2}^2 = \chi$  and so by choosing  $N$  sufficiently large in (5.7) gives the desired contradiction.  $\square$

## References

- [1] G. Alberti, G. Crippa, and A. Mazzucato, *Exponential self-similar mixing by incompressible flows*, Journal of the American Mathematical Society **32** (2019), no. 2, 445–490.
- [2] Y. Amarouchene and H. Kellay, *Batchelor scaling in fast-flowing soap films*, Physical review letters **93** (2004), no. 21, 214504.
- [3] F. Anselmetti, Y. Gagne, E. Hopfinger, and R. Antonia, *High-order velocity structure functions in turbulent shear flows*, J. of Fluid Mech. **140** (1984), 63–89.
- [4] R. Antonia and P. Orlandi, *Effect of Schmidt number on small-scale passive scalar turbulence*, Appl. Mech. Rev. **56** (2003), no. 6, 615–632.
- [5] T. M. Antonsen Jr, Z. Fan, E. Ott, and E. Garcia-Lopez, *The role of chaotic orbits in the determination of power spectra of passive scalars*, Physics of Fluids **8** (1996), no. 11, 3094–3104.
- [6] T. M. Antonsen Jr and E. Ott, *Multifractal power spectra of passive scalars convected by chaotic fluid flows*, Physical Review A **44** (1991), no. 2, 851.
- [7] L. Arnold, *A formula connecting sample and moment stability of linear stochastic systems*, SIAM Journal on Applied Mathematics **44** (1984), no. 4, 793–802.
- [8] L. Arnold, W. Kliemann, and E. Oeljeklaus, *Lyapunov exponents of linear stochastic systems*, Lyapunov exponents, 1986, pp. 85–125.
- [9] L. Arnold and W. Kliemann, *Large deviations of linear stochastic differential equations*, Stochastic differential systems, 1987, pp. 115–151.
- [10] A. Babiano and A. Provenzale, *Coherent vortices and tracer cascades in two-dimensional turbulence*, Journal of Fluid Mechanics **574** (2007), 429–448.
- [11] E. Balkovsky and A. Fouxon, *Universal long-time properties of Lagrangian statistics in the Batchelor regime and their application to the passive scalar problem*, Physical Review E **60** (1999), no. 4, 4164.
- [12] L. Barreira and Y. B. Pesin, *Lyapunov exponents and smooth ergodic theory*, Vol. 23, American Mathematical Soc., 2002.
- [13] G. K. Batchelor, *Small-scale variation of convected quantities like temperature in turbulent fluid part 1. general discussion and the case of small conductivity*, Journal of Fluid Mechanics **5** (1959), no. 1, 113–133.
- [14] P. Baxendale and T. E. Harris, *Isotropic stochastic flows*, Ann. Probab. **14** (1986), no. 4, 1155–1179.
- [15] P. H. Baxendale, *Lyapunov exponents and relative entropy for a stochastic flow of diffeomorphisms*, Probability Theory and Related Fields **81** (1989), no. 4, 521–554.
- [16] P. Baxendale and B. Rozovsky, *Kinematic dynamo and intermittence in a turbulent flow*, Geophysical & Astrophysical Fluid Dynamics **73** (1993), no. 1-4, 33–60.
- [17] J. Bedrossian, A. Blumenthal, and S. Punshon-Smith, *Lagrangian chaos and scalar advection in stochastic fluid mechanics*, arXiv preprint arXiv:1809.06484 (2018).
- [18] J. Bedrossian, A. Blumenthal, and S. Punshon-Smith, *Almost-sure enhanced dissipation and uniform-in-diffusivity exponential mixing for advection-diffusion by stochastic Navier-Stokes*, arXiv preprint arXiv:1911.01561 (2019).
- [19] J. Bedrossian, A. Blumenthal, and S. Punshon-Smith, *Almost-sure exponential mixing of passive scalars by the stochastic Navier-Stokes equations*, arXiv preprint arXiv:1905.03869 (2019).



- [20] J. Bedrossian, M. Coti Zelati, S. Punshon-Smith, and F. Weber, *Sufficient conditions for dual cascade flux laws in the stochastic 2D Navier-Stokes equations*, arXiv preprint arXiv:1905.03299 (2019).
- [21] J. Bedrossian, M. Coti Zelati, S. Punshon-Smith, and F. Weber, *A sufficient condition for the Kolmogorov 4/5 law for stationary martingale solutions to the 3D Navier–Stokes equations*, *Communications in Mathematical Physics* **367** (2019), no. 3, 1045–1075.
- [22] D. Biskamp, *Magnetohydrodynamic turbulence*, Cambridge University Press, 2003.
- [23] A. Blumenthal, J. Xue, and L.-S. Young, *Lyapunov exponents for random perturbations of some area-preserving maps including the standard map*, *Annals of Mathematics* (2017), 285–310.
- [24] A. Blumenthal and L.-S. Young, *Absolute continuity of stable foliations for mappings of Banach spaces*, *Communications in Mathematical Physics* **354** (2017), no. 2, 591–619.
- [25] A. Blumenthal and L.-S. Young, *Entropy, volume growth and SRB measures for Banach space mappings*, *Inventiones mathematicae* **207** (2017), no. 2, 833–893.
- [26] G. Boffetta and R. E Ecke, *Two-dimensional turbulence*, *Annual Review of Fluid Mechanics* **44** (2012), 427–451.
- [27] D. Bogucki, J. A. Domaradzki, and P. Yeung, *Direct numerical simulations of passive scalars with  $Pr > 1$  advected by turbulent flow*, *Journal of Fluid Mechanics* **343** (1997), 111–130.
- [28] T. Bohr, M. H Jensen, G. Paladin, and A. Vulpiani, *Dynamical systems approach to turbulence*, Cambridge University Press, 2005.
- [29] A. Bressan, *A lemma and a conjecture on the cost of rearrangements*, *Rend. Sem. Mat. Univ. Padova* **110** (2003), 97–102.
- [30] M. Brin and G. Stuck, *Introduction to dynamical systems*, Cambridge university press, 2002.
- [31] T. Buckmaster, P. Germain, Z. Hani, and J. Shatah, *Effective dynamics of the nonlinear Schrödinger equation on large domains*, *Communications on Pure and Applied Mathematics* **71** (2018), no. 7, 1407–1460.
- [32] T. Buckmaster, P. Germain, Z. Hani, and J. Shatah, *Onset of the wave turbulence description of the longtime behavior of the nonlinear Schrödinger equation*, arXiv preprint arXiv:1907.03667 (2019).
- [33] T. Buckmaster and V. Vicol, *Convex integration and phenomenologies in turbulence*, arXiv preprint arXiv:1901.09023 (2019).
- [34] A. Cheskidov and R. Shvydkoy, *Euler equations and turbulence: analytical approach to intermittency*, *SIAM Journal on Mathematical Analysis* **46** (2014), no. 1, 353–374.
- [35] A. Cheskidov, P. Constantin, S. Friedlander, and R. Shvydkoy, *Energy conservation and Onsager’s conjecture for the Euler equations*, *Nonlinearity* **21** (2008), no. 6, 1233.
- [36] A. Cheskidov, M. Dai, and L. Kavlie, *Determining modes for the 3D Navier–Stokes equations*, *Physica D: Nonlinear Phenomena* **374** (2018), 1–9.
- [37] P. Constantin, C. Foias, O. Manley, and R. Temam, *Determining modes and fractal dimension of turbulent flows*, *Journal of Fluid Mechanics* **150** (1985), 427–440.
- [38] P. Constantin, *The Littlewood–Paley spectrum in two-dimensional turbulence*, *Theoretical and computational fluid dynamics* **9** (1997), no. 3-4, 183–189.
- [39] P. Constantin, W. E, and E. S. Titi, *Onsager’s conjecture on the energy conservation for solutions of Euler’s equation*, *Comm. Math. Phys.* **165** (1994), no. 1, 207–209.
- [40] P. Constantin and C. Foias, *Global Lyapunov exponents, Kaplan-Yorke formulas and the dimension of the attractors for 2d Navier-Stokes equations*, *Communications on pure and applied mathematics* **38** (1985), no. 1, 1–27.
- [41] P. Constantin, Q. Nie, and S. Tanveer, *Bounds for second order structure functions and energy spectrum in turbulence*, *Physics of fluids* **11** (1999), no. 8, 2251–2256.
- [42] S. Corrsin, *On the spectrum of isotropic temperature fluctuations in an isotropic turbulence*, *Journal of Applied Physics* **22** (1951), no. 4, 469–473.
- [43] A. Crisanti, M. Falcioni, A. Vulpiani, and G. Paladin, *Lagrangian chaos: transport, mixing and diffusion in fluids*, *La Rivista del Nuovo Cimento* (1978-1999) **14** (1991), no. 12, 1–80.
- [44] A. Crisanti, G. Paladin, and A. Vulpiani, *Products of random matrices: in statistical physics*, Vol. 104, Springer Science & Business Media, 2012.
- [45] G. Da Prato and J. Zabczyk, *Ergodicity for infinite-dimensional systems*, London Mathematical Society Lecture Note Series, vol. 229, Cambridge University Press, Cambridge, 1996.

- [46] R. Dascaliuc and Z. Grujić, *Anomalous dissipation and energy cascade in 3d inviscid flows*, Communications in Mathematical Physics **309** (2012), no. 3, 757–770.
- [47] C. De Lellis and L. Székelyhidi, *Dissipative continuous Euler flows*, Inventiones mathematicae **193** (2013), no. 2, 377–407.
- [48] C. De Lellis and L. Székelyhidi Jr., *The Euler equations as a differential inclusion*, Ann. of Math. (2) **170** (2009), no. 3, 1417–1436.
- [49] T. M. Dillon and D. R. Caldwell, *The Batchelor spectrum and dissipation in the upper ocean*, Journal of Geophysical Research: Oceans **85** (1980), no. C4, 1910–1916.
- [50] R. J. DiPerna and P.-L. Lions, *Ordinary differential equations, transport theory and Sobolev spaces*, Inventiones mathematicae **98** (1989), no. 3, 511–547.
- [51] D. A. Donzis, K. Sreenivasan, and P. Yeung, *The Batchelor spectrum for mixing of passive scalars in isotropic turbulence*, Flow, turbulence and combustion **85** (2010), no. 3-4, 549–566.
- [52] T. D. Drivas, *Turbulent cascade direction and Lagrangian time-asymmetry*, J. Nonlinear Sci. **29** (2019), no. 1, 65–88.
- [53] T. D. Drivas, T. M. Elgindi, G. Iyer, and I.-J. Jeong, *Anomalous dissipation in passive scalar transport*, arXiv preprint arXiv:1911.03271 (2019).
- [54] P. Duarte, *Plenty of elliptic islands for the standard family of area preserving maps*, Annales de l’institut henri poincaré (c) non linear analysis, 1994, pp. 359–409.
- [55] J. Duchon and R. Robert, *Inertial energy dissipation for weak solutions of incompressible Euler and Navier-Stokes equations*, Nonlinearity **13** (2000), no. 1, 249–255.
- [56] W. E and J. C. Mattingly, *Ergodicity for the Navier-Stokes equation with degenerate random forcing: Finite-dimensional approximation*, Commun. Pure Appl. Math. **54** (Nov. 2001), no. 11, 1386–1402.
- [57] J.-P. Eckmann and D. Ruelle, *Ergodic theory of chaos and strange attractors*, The theory of chaotic attractors, 1985, pp. 273–312.
- [58] M. Einsiedler and T. Ward, *Ergodic theory*, Springer, 2013.
- [59] T. M. Elgindi and A. Zlatoš, *Universal mixers in all dimensions*, Advances in Mathematics **356** (2019), 106807.
- [60] G. L. Eyink, *Energy dissipation without viscosity in ideal hydrodynamics. I. Fourier analysis and local energy transfer*, Phys. D **78** (1994), no. 3-4, 222–240.
- [61] G. L. Eyink, *Review of the Onsager “ideal turbulence” theory*, arXiv preprint arXiv:1803.02223 (2018).
- [62] G. Falkovich, K. Gawdzki, and M. Vergassola, *Particles and fields in fluid turbulence*, Reviews of modern Physics **73** (2001), no. 4, 913.
- [63] E. Fedrizzi and F. Flandoli, *Noise prevents singularities in linear transport equations*, J. Funct. Anal. **264** (Mar. 2013), no. 6, 1329–1354.
- [64] F. Flandoli, M. Gubinelli, M. Hairer, and M. Romito, *Rigorous remarks about scaling laws in turbulent fluids*, Comm. Math. Phys. **278** (2008), no. 1, 1–29. MR2367196
- [65] F. Flandoli, M. Gubinelli, and E. Priola, *Well-posedness of the transport equation by stochastic perturbation*, Invent. Math. **180** (Nov. 2009), no. 1, 1–53.
- [66] F. Flandoli and B. Maslowski, *Ergodicity of the 2-D Navier-Stokes equation under random perturbations*, Comm. in Math. Phys. **172** (1995), no. 1, 119–141.
- [67] S. Friedlander, N. Glatt-Holtz, and V. Vicol, *Inviscid limits for a stochastically forced shell model of turbulent flow*, Annales de l’institut henri Poincaré, probabilités et statistiques, 2016, pp. 1217–1247.
- [68] U. Frisch, *Turbulence: the legacy of A.N. Kolmogorov*, Cambridge university press, 1995.
- [69] J. D. Gibbon and E. Titi, *Attractor dimension and small length scale estimates for the three-dimensional Navier-Stokes equations*, Nonlinearity **10** (1997), no. 1, 109.
- [70] C. Gibson and W. Schwarz, *The universal equilibrium spectra of turbulent velocity and scalar fields*, Journal of Fluid Mechanics **16** (1963), no. 3, 365–384.
- [71] B. Goldys and B. Maslowski, *Exponential ergodicity for stochastic Burgers and 2D Navier–Stokes equations*, Journal of Functional Analysis **226** (2005), no. 1, 230–255.
- [72] A. Gorodetski, *On stochastic sea of the standard map*, Communications in Mathematical Physics **309** (2012), no. 1, 155–192.
- [73] H. Grant, B. Hughes, W. Vogel, and A. Moilliet, *The spectrum of temperature fluctuations in turbulent flow*, Journal of Fluid Mechanics **34** (1968), no. 3, 423–442.

- [74] D. Grošelj, S. S Cerri, A. B. Navarro, C. Willmott, D. Told, N. F Loureiro, F. Califano, and F. Jenko, *Fully kinetic versus reduced-kinetic modeling of collisionless plasma turbulence*, The Astrophysical Journal **847** (2017), no. 1, 28.
- [75] M. Hairer and J. C. Mattingly, *Ergodicity of the 2D Navier-Stokes equations with degenerate stochastic forcing*, Ann. of Math. **164** (2006), no. 3, 993–1032.
- [76] M. Hairer and J. C. Mattingly, *Spectral gaps in Wasserstein distances and the 2D stochastic Navier-Stokes equations*, Ann. Probab. **36** (2008), no. 6, 2050–2091.
- [77] M. Hairer and J. C. Mattingly, *A theory of hypoellipticity and unique ergodicity for semilinear stochastic PDEs*, Electron. J. Probab. **16** (2011), no. 23, 658–738.
- [78] M. Holzer and E. D Siggia, *Turbulent mixing of a passive scalar*, Physics of Fluids **6** (1994), no. 5, 1820–1837.
- [79] L. Hörmander, *The analysis of linear partial differential operators I–IV*, Springer, New York, 1985.
- [80] P. Isett, *A proof of Onsager’s conjecture*, Annals of Mathematics **188** (2018), no. 3, 871–963.
- [81] G. Iyer, A. Kiselev, and X. Xu, *Lower bounds on the mix norm of passive scalars advected by incompressible enstrophy-constrained flows*, Nonlinearity **27** (2014), no. 5, 973.
- [82] V. Jaksic, V. Nersesyan, C.-A. Pillet, and A. Shirikyan, *Large deviations and entropy production in viscous fluid flows*, arXiv preprint arXiv:1902.03278 (2019).
- [83] Y. L. Jan and O. Raimond, *Integration of brownian vector fields*, Ann. Probab. **30** (Apr. 2002), no. 2, 826–873.
- [84] D. A Jones and E. S Titi, *Upper bounds on the number of determining modes, nodes, and volume elements for the navier-stokes equations*, Indiana University Mathematics Journal (1993), 875–887.
- [85] M.-C. Jullien, P. Castiglione, and P. Tabeling, *Experimental observation of Batchelor dispersion of passive tracers*, Physical review letters **85** (2000), no. 17, 3636.
- [86] Y. Kaneda, T. Ishihara, M. Yokokawa, K. Itakura, and A. Uno, *Energy dissipation rate and energy spectrum in high resolution direct numerical simulations of turbulence in a periodic box*, Physics of Fluids **15** (2003), no. 2, L21–L24.
- [87] Y. Kifer, *Ergodic theory of random transformations*, Vol. 10, Springer Science & Business Media, 2012.
- [88] A. Kolmogoroff, *The local structure of turbulence in incompressible viscous fluid for very large Reynold’s numbers*, C. R. (Doklady) Acad. Sci. URSS (N.S.) **30** (1941), 301–305. MR0004146
- [89] A. N. Kolmogoroff, *Dissipation of energy in the locally isotropic turbulence*, C. R. (Doklady) Acad. Sci. URSS (N.S.) **32** (1941), 16–18. MR0005851
- [90] A. N. Kolmogoroff, *On degeneration of isotropic turbulence in an incompressible viscous liquid*, C. R. (Doklady) Acad. Sci. URSS (N. S.) **31** (1941), 538–540. MR0004568
- [91] R. H Kraichnan, *Small-scale structure of a scalar field convected by turbulence*, The Physics of Fluids **11** (1968), no. 5, 945–953.
- [92] S. Kuksin, V. Nersesyan, and A. Shirikyan, *Exponential mixing for a class of dissipative pdes with bounded degenerate noise*, arXiv preprint arXiv:1802.03250 (2018).
- [93] S. Kuksin, V. Nersesyan, and A. Shirikyan, *Mixing via controllability for randomly forced nonlinear dissipative pdes*, arXiv preprint arXiv:1902.00494 (2019).
- [94] S. Kuksin and A. Shirikyan, *Mathematics of two-dimensional turbulence*, Vol. 194, Cambridge University Press, 2012.
- [95] H. Kunita, *Stochastic differential equations with jumps and stochastic flows of diffeomorphisms*, Itô’s stochastic calculus and probability theory, 1996, pp. 197–211.
- [96] F. Ledrappier and L-S Young, *The metric entropy of diffeomorphisms: Part i: Characterization of measures satisfying pesin’s entropy formula*, Annals of Mathematics (1985), 509–539.
- [97] S. K Lele, *Compressibility effects on turbulence*, Annual review of fluid mechanics **26** (1994), no. 1, 211–254.
- [98] Z. Lin, J.-L. Thiffeault, and C. R Doering, *Optimal stirring strategies for passive scalar mixing*, J. of Fluid Mech. **675** (2011), 465–476.
- [99] C. Liverani, B. Saussol, and S. Vaienti, *A probabilistic approach to intermittency*, Ergodic theory and dynamical systems **19** (1999), no. 3, 671–685.
- [100] D. Luo, *Isotropic stochastic flow of homeomorphisms on  $\mathbb{R}^d$  associated with the critical sobolev exponent*, Stochastic Process. Appl. **118** (Aug. 2008), no. 8, 1463–1488.
- [101] S. P Meyn and R. L Tweedie, *Markov chains and stochastic stability*, Springer Science & Business Media, 2012.
- [102] P. L Miller and P. E Dimotakis, *Measurements of scalar power spectra in high schmidt number turbulent jets*, Journal of Fluid Mechanics **308** (1996), 129–146.

- [103] S. Modena and L. Székelyhidi, *Non-renormalized solutions to the continuity equation*, Calc. Var. Partial Differential Equations **58** (2019), no. 6, Paper No. 208. MR4029736
- [104] S.-E. A Mohammed, T. K Nilssen, and F. N Proske, *Sobolev differentiable stochastic flows for SDEs with singular coefficients: Applications to the transport equation*, Ann. Probab. **43** (May 2015), no. 3, 1535–1576 (en).
- [105] L. Mydlarski and Z. Warhaft, *On the onset of high-reynolds-number grid-generated wind tunnel turbulence*, Journal of Fluid Mechanics **320** (1996), 331–368.
- [106] S. Nazarenko, *Wave turbulence*, Vol. 825, Springer Science & Business Media, 2011.
- [107] Q. Nie and S. Tanveer, *A note on third-order structure functions in turbulence*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **455** (1999), no. 1985, 1615–1635.
- [108] J. O Nye and R. S Brodkey, *The scalar spectrum in the viscous-convective subrange*, Journal of Fluid Mechanics **29** (1967), no. 1, 151–163.
- [109] A. Obukhov, *The structure of the temperature field in a turbulent flow*, Dokl. Akad. Nauk. SSSR **39** (1949).
- [110] E. Ott and T. M Antonsen Jr, *Chaotic fluid convection and the fractal nature of passive scalar gradients*, Physical review letters **61** (1988), no. 25, 2839.
- [111] F. Otto and F. Ramos, *Universal bounds for the Littlewood-Paley first-order moments of the 3D Navier-Stokes equations*, Communications in mathematical physics **300** (2010), no. 2, 301–315.
- [112] Y. Pesin and V. Climenhaga, *Open problems in the theory of non-uniform hyperbolicity*, Discrete Contin. Dyn. Syst **27** (2010), no. 2, 589–607.
- [113] R. Prasad and K. Sreenivasan, *The measurement and interpretation of fractal dimensions of the scalar interface in turbulent flows*, Physics of Fluids A: Fluid Dynamics **2** (1990), no. 5, 792–807.
- [114] C. Pugh and M. Shub, *Ergodic attractors*, Transactions of the American Mathematical Society **312** (1989), no. 1, 1–54.
- [115] M. Romito, *Ergodicity of the finite dimensional approximation of the 3D Navier–Stokes equations forced by a degenerate noise*, J. Stat. Phys. **114** (Jan. 2004), no. 1, 155–177.
- [116] D. Ruelle, *Characteristic exponents and invariant manifolds in hilbert space*, Annals of Mathematics (1982), 243–290.
- [117] O. Sarig, *Subexponential decay of correlations*, Inventiones mathematicae **150** (2002), no. 3, 629–653.
- [118] C. Seis, *Maximal mixing by incompressible fluid flows*, Nonlinearity **26** (2013), no. 12, 3279.
- [119] C. Seis, *On the Littlewood–Paley spectrum for passive scalar transport equations*, Journal of Nonlinear Science (2019), 1–12.
- [120] G. R Sell, *Global attractors for the three-dimensional navier-stokes equations*, Journal of Dynamics and Differential Equations **8** (1996), no. 1, 1–33.
- [121] A. Shirikyan, *Controllability implies mixing ii. convergence in the dual-lipschitz metric*, arXiv preprint arXiv:1803.01893 (2018).
- [122] B. I Shraiman and E. D Siggia, *Anomalous scaling for a passive scalar near the Batchelor limit*, Physical Review E **57** (1998), no. 3, 2965.
- [123] B. I Shraiman and E. D Siggia, *Scalar turbulence*, Nature **405** (2000), no. 6787, 639.
- [124] K. R. Sreenivasan and P. Kailasnath, *An update on the intermittency exponent in turbulence*, Physics of Fluids A: Fluid Dynamics **5** (1993), no. 2, 512–514.
- [125] J.-L. Thiffeault, *Using multiscale norms to quantify mixing and transport*, Nonlinearity **25** (Feb. 2012), no. 2, R1–R44.
- [126] Z. Warhaft, *Passive scalars in turbulent flows*, Annual Review of Fluid Mechanics **32** (2000), no. 1, 203–240.
- [127] B. S Williams, D Marteau, and J. P Gollub, *Mixing of a passive scalar in magnetically forced two-dimensional turbulence*, Physics of Fluids **9** (1997), no. 7, 2061–2080.
- [128] X. Wu, B Martin, H Kellay, and W. Goldburg, *Hydrodynamic convection in a two-dimensional couette cell*, Physical review letters **75** (1995), no. 2, 236.
- [129] A. Yaglom, *On the local structure of a temperature field in a turbulent flow*, Dokl. akad. nauk sssr, 1949, pp. 743–746.
- [130] Y. Yao and A. Zlatoš, *Mixing and un-mixing by incompressible flows*, J. Euro. Math. Soc. **19** (2017), no. 7, 1911–1948.
- [131] L.-S. Young, *Ergodic theory of differentiable dynamical systems*, Real and complex dynamical systems, 1995, pp. 293–336.
- [132] L.-S. Young, *What are SRB measures, and which dynamical systems have them?*, Journal of Statistical Physics **108** (2002), no. 5, 733–754.

- [133] G.-C. Yuan, K. Nam, T. M. Antonsen Jr, E. Ott, and P. N. Guzdar, *Power spectrum of passive scalars in two dimensional chaotic flows*, *Chaos: An Interdisciplinary Journal of Nonlinear Science* **10** (2000), no. 1, 39–49.
- [134] V. Zakharov, F. Dias, and A. Pushkarev, *One-dimensional wave turbulence*, *Physics Reports* **398** (2004), no. 1, 1–65.
- [135] V. E. Zakharov, V. S. L'vov, and G. Falkovich, *Kolmogorov spectra of turbulence I: Wave turbulence*, Springer Science & Business Media, 2012.