

High-precision nuclear forces from chiral EFT: State-of-the-art, challenges and outlook

E. Epelbaum¹, H. Krebs¹ and P. Reinert¹

¹ Ruhr-Universität Bochum, Fakultät für Physik und Astronomie, Institut für Theoretische Physik II, D-44780 Bochum, Germany

emails: evgeny.epelbaum@rub.de, hermann.krebs@rub.de, patrick.reinert@rub.de

ABSTRACT

We review a new generation of nuclear forces derived in chiral effective field theory using the recently proposed semilocal regularization method. We outline the conceptual foundations of nuclear chiral effective field theory, discuss all steps needed to compute nuclear observables starting from the effective chiral Lagrangian and consider selected applications in the two- and few-nucleon sectors. We highlight key challenges in developing high-precision tree-body forces, such as the need to maintain consistency between two- and many-body interactions and constraints placed by the chiral and gauge symmetries after regularization.

Keywords: Nuclear forces, effective field theory, chiral perturbation theory, regularization, few-body systems

1 INTRODUCTION

Almost thirty years ago, Weinberg put forward his groundbreaking idea to apply chiral perturbation theory (ChPT), the low-energy effective field theory (EFT) of QCD, to the derivation of nuclear interactions [1, 2]. This seminal work has revolutionized the whole field of nuclear physics by providing a solid theoretical basis and offering a systematically improvable approach to low-energy nuclear structure and reactions.

So where do we stand today in the implementation of the program initiated by Weinberg? Much has been learned about specific features of the nuclear interactions and currents and about the role of many-body forces from the point of view of the effective chiral Lagrangian, see Refs. [3, 4, 5, 6] for review articles covering different research areas, while some issues are still under debate [7, 6]. Meanwhile, the interactions derived in chiral EFT, sometimes referred to as “chiral forces”, have largely replaced phenomenological potentials developed in the nineties of the last century. They are nowadays commonly used in *ab initio* nuclear structure calculations, see Refs. [8, 9, 10, 11, 12, 13] for recent examples using a variety of continuum *ab initio* methods and [14, 15, 16] for selected highlights from nuclear lattice simulations. With the most recent chiral nucleon-nucleon (NN) potentials [17] providing a nearly perfect description of the mutually consistent neutron-proton (np) and proton-proton (pp) scattering data below pion production threshold from the Granada-2013 database [18], the two-nucleon sector is already in a very good shape. On the other hand, three-nucleon forces (3NF) are much less understood at the quantitative level [19] and constitute an important frontier in nuclear physics [20].

In this article we focus on the latest generation of chiral nuclear forces based on an improved regularization approach [21, 22, 17], which allows one to maintain the long-range part of the interaction. We review our recent work along these lines in the two-nucleon sector, describe the ongoing efforts by the Low-Energy

Nuclear Physics International Collaboration (LENPIC) towards developing consistent many-body forces and solving the structure and reactions of light nuclei, and discuss selected applications.

Our paper is organized as follows. In section 2, we outline the foundations of the employed theoretical framework. Section 3 gives an overview of various methods to derive nuclear forces and currents from the effective chiral Lagrangian. It also summarizes the available results for nuclear potentials derived using dimensional regularization (DR). In section 4 we present the improved semilocal regularization approach, which is utilized in the most accurate and precise NN potentials of Ref. [17]. We also discuss the challenges that need to be addressed to construct *consistently regularized* 3NFs and exchange current operators beyond tree level, which are not restricted to any particular type of cutoff regularization. Section 5 is devoted to uncertainty quantification in chiral EFT. Selected results for the NN system, three-nucleon scattering and light nuclei are presented in section 6. We conclude with a short summary and outlook in section 7.

2 THE FRAMEWORK IN A NUTSHELL

Throughout this work, we restrict ourselves to the two-flavor case of the light up- and down-quarks and employ the simplest version of the effective chiral Lagrangian with pions and nucleons as the only active degrees of freedom. Contributions of the $\Delta(1232)$ isobar to the nuclear potentials are discussed in Refs. [23, 24, 25, 26, 27, 28]. The effective Lagrangian involves all possible interactions between pions and nucleons compatible with the symmetries of QCD and is organized in powers of derivatives and quark (or equivalently pion) masses. Pions correspond to the (pseudo) Nambu-Goldstone bosons of the spontaneously broken axial generators and thus transform nonlinearly with respect to chiral $SU(2)_L \times SU(2)_R$ transformations. The effective Lagrangian can be constructed in a straightforward way using covariantly transforming building blocks defined in terms of the pion fields [29, 30]. All applications reviewed in this paper rely on a nonrelativistic treatment of the nucleon fields and make use of the heavy-baryon formalism to eliminate the nucleon mass m from the leading-order Lagrangian. The individual terms in the effective Lagrangian are multiplied by the corresponding coupling constants, commonly referred to as low-energy constants (LECs), which are not fixed by the symmetry and typically need to be determined from experimental data. The most accurate currently available nuclear potentials at fifth order in the chiral expansion, i.e. at $N^4\text{LO}$, require input from the following effective Lagrangians (with each line containing the contributions with a fixed number of the nucleon fields)

$$\begin{aligned}
\mathcal{L}_{\text{eff}} &= \mathcal{L}_{\pi}^{(2)}(M_{\pi}, F_{\pi}) + \mathcal{L}_{\pi}^{(4)}(l_{1,\dots,7}) \\
&+ \mathcal{L}_{\pi N}^{(1)}(g_A) + \mathcal{L}_{\pi N}^{(2)}(m, c_{1,\dots,7}) + \mathcal{L}_{\pi N}^{(3)}(d_{1,\dots,23}) + \mathcal{L}_{\pi N}^{(4)}(e_{1,\dots,118}) \\
&+ \mathcal{L}_{NN}^{(0)}(C_S, C_T) + \mathcal{L}_{NN}^{(2)}(C_{1,\dots,7}) + \mathcal{L}_{NN}^{(4)}(D_{1,\dots,12}) + \mathcal{L}_{\pi NN}^{(1)}(D) + \dots \\
&+ \mathcal{L}_{NNN}^{(0)}(E) + \mathcal{L}_{NNN}^{(2)}(E_{1,\dots,10}), \tag{1}
\end{aligned}$$

where M_{π} and F_{π} are the pion mass and decay constant¹, g_A is the nucleon axial-vector coupling while l_i , c_i , d_i , e_i , C_i , D , D_i , E and E_i are further LECs. The superscript n of $\mathcal{L}^{(n)}$ denotes the number of derivatives and/or M_{π} -insertions and is sometimes referred to as the chiral dimension. Notice that we only show new LECs that appear in the corresponding Lagrangians and suppress the dependence on the LECs appearing at lower orders. The pionic Lagrangian can be found in Ref. [31], $\mathcal{L}_{\pi N}$ is given in

¹ Strictly speaking, M_{π} is to be understood as the pion mass to leading order in the chiral expansion while F_{π} and other parameters in the effective Lagrangian refer to the corresponding LECs in the chiral limit of vanishing light quark masses.

Refs. [32, 33], $\mathcal{L}_{\text{NN}}^{(0)}$ was introduced in Refs. [1, 2], $\mathcal{L}_{\text{NN}}^{(2)}$ can be found in Refs. [34, 35], the minimal form of $\mathcal{L}_{\text{NN}}^{(4)}$ is given in Ref. [17], $\mathcal{L}_{\pi\text{NN}}^{(1)}$ and $\mathcal{L}_{\text{NNN}}^{(0)}$ are discussed in Ref. [36] while $\mathcal{L}_{\text{NNN}}^{(2)}$ was constructed in Ref. [37]. Notice further that the chiral symmetry breaking terms $\propto M_\pi^2$ are not shown explicitly in $\mathcal{L}_{\text{NN}}$ and $\mathcal{L}_{\text{NNN}}$. For calculations at the physical value of the quark masses, their contributions are absorbed into the LECs listed in Eq. (1). We have, furthermore, restricted ourselves in this equation to isospin-invariant terms for the Lagrangians involving two and three nucleons. The single-nucleon Lagrangian $\mathcal{L}_{\pi\text{N}}$ does involve isospin-breaking contributions due to the quark mass difference and can be extended to include virtual photon effects Refs. [38, 39]. The ellipses in the second-to-last line of Eq. (1) refer to higher-order Lagrangians $\mathcal{L}_{\pi\text{NN}}$, which have not been worked out yet and would be needed to finalize the derivation of the 3NF at N^4LO .

The long-range parts of the nuclear forces emerge from pion exchange diagrams and can be derived from $\mathcal{L}_\pi$ and $\mathcal{L}_{\pi\text{N}}$. Fortunately, only a very restricted set of (linear combinations of) LECs from these Lagrangians contributes to the $\pi N \rightarrow \pi N$ and $\pi N \rightarrow \pi\pi N$ scattering amplitudes, which enter as subprocesses when deriving the long-range nuclear interactions up to N^4LO , namely $c_{1,\dots,4}$ from $\mathcal{L}_{\pi\text{N}}^{(2)}$, $d_1 + d_2$, $d_{3,5,18}$ and $d_{14} - d_{15}$ from $\mathcal{L}_{\pi\text{N}}^{(3)}$ and $e_{14,\dots,18}$ from $\mathcal{L}_{\pi\text{N}}^{(4)}$. Here, we made use of the fact that the contributions from the LECs l_3 , $e_{19,\dots,22}$ and $e_{35,\dots,38}$ can be absorbed into the appropriate shifts of the LECs c_i [40]. All these πN LECs can nowadays be reliably extracted by matching the πN scattering amplitude from the recent Roy-Steiner equation analysis [41] with ChPT at the subthreshold point [42], see also Ref. [43] for an alternative strategy. Thus, *the long-range nuclear interactions are completely determined by the spontaneously broken approximate chiral symmetry of QCD and experimental/empirical information on the πN system in a parameter-free way*. The two- and three-nucleon interactions in the last two lines of Eq. (1) parametrize the short-range part of the nuclear forces, and the corresponding LECs have to be determined from NN scattering and three- or more-nucleon observables.

In the single-nucleon sector, the effective Lagrangian $\mathcal{L}_\pi + \mathcal{L}_{\pi\text{N}}$ can be used to systematically compute the scattering amplitude in perturbation theory by applying the chiral expansion, a simultaneous expansion in particles' external three-momenta $p \equiv |\vec{p}|$ and around the chiral limit $M_\pi \rightarrow 0$. The importance of every Feynman diagram is estimated by counting powers of the soft scales and applying the rules of naive dimensional analysis (NDA). The expansion parameter $Q \in \{p/\Lambda_b, M_\pi/\Lambda_b\}$ is determined by the breakdown scale Λ_b , which may (optimistically) be expected to be of the order of the ρ -meson mass.² At every order in the chiral expansion only a finite number of Feynman diagrams need to be evaluated. For more details on ChPT in the 1N sector see the review article [45].

Contrary to the 1N case, the NN S-wave scattering amplitude exhibits poles in the near-threshold region corresponding to the bound state (deuteron) and the virtual state in the $^1\text{S}_0$ channel, which signal the breakdown of perturbation theory. In this context, it was pointed out by Weinberg that the contributions of multi-nucleon ladder diagrams are enhanced compared to the estimation based on the chiral power counting due to the appearance of pinch singularities (in the $m \rightarrow \infty$ limit) [1, 2]. Weinberg also argued that the nucleon mass needs to be counted as $m \sim \Lambda_b^2/M_\pi \gg \Lambda_b$ in order to formally justify the need to perform a nonperturbative resummation of the ladder contributions. Given that the ladder diagrams are automatically resummed by solving the few-nucleon Schrödinger equation, Weinberg's chiral EFT approach to low-energy nuclear systems, perhaps not surprisingly, resembles the quantum mechanical

² An upper bound for Λ_b is set by the scale $4\pi F_\pi$ emerging from pion loops [44].




Figure 1. Representation of the on-shell scattering amplitude from the one-pion-two-pion-exchange Feynman diagram (left) in terms of iterations of the Faddeev equation (right). Gray-shaded rectangles visualize the corresponding two- and three-nucleon potentials $V_{2N}^{1\pi}$, $V_{3N}^{2\pi, 1/m}$ and $V_{3N}^{2\pi-1\pi}$ while G_0 denotes the free resolvent operator for nonrelativistic nucleons.

A-body problem

$$\left[\left(\sum_{i=1}^A \frac{-\Delta_i}{2m} + \mathcal{O}(m^{-3}) \right) + V_{2N} + V_{3N} + V_{4N} + \dots \right] |\Psi\rangle = E|\Psi\rangle, \quad (2)$$

where Δ_i is the Laplace operator acting on the nucleon i . The nuclear potentials V_{2N} , V_{3N} , $\dots$ receive contributions from diagrams that cannot be reduced to ladder iterations and are calculable in a systematically improvable way within ChPT.

Among the many attractive features, the approach outlined above allows one to maintain *consistency* between nuclear forces and exchange current operators which are scheme-dependent quantities. To illustrate the meaning and importance of consistency consider the Feynman diagram on the left-hand side (l.h.s.) of the equality shown in Fig. 1 as an example. The corresponding (on-shell) contribution to the scattering amplitude features both a reducible (i.e. of a ladder-type) and irreducible pieces as visualized in the figure. Reducible contributions to the amplitude are resummed up to an infinite order when solving the Faddeev equation corresponding to Eq. (2). In doing so, the diagrams corresponding to its zeroth and first iterations shown in the figure *must* match the result obtained from the Feynman diagram when taken on the energy shell. The iterative contribution from the first graph on the right-hand side of the depicted equality, however, involves NN and 3N potentials, whose off-shell behavior is scheme dependent. Also the 3NF corresponding to the last diagram is scheme dependent (even on the energy shell) [46], and only a *consistent* choice of the involved two- and three-nucleon potentials guarantees the validity of matching for the scattering amplitude. This can indeed be verified explicitly using the expressions for the 3NFs $V_{3N}^{2\pi, 1/m}$ from Eqs. (4.9)-(4.11) of Ref. [47] and $V_{3N}^{2\pi-1\pi}$ from Eqs. (2.16)-(2.20) of Ref. [46] and employing DR to evaluate loop integrals.³

Clearly, DR is impractical for a numerical solution of the A -body problem and is usually replaced by cutoff regularization. Renormalization of the Schrödinger equation in the context of chiral EFT is a controversial and heavily debated topic, see Refs. [48, 49, 50, 51, 52, 53, 54, 55, 7, 6] for a range of opinions. The essence of the problem is related to the nonrenormalizable nature of the Lippmann-Schwinger (LS) equation for NN potentials truncated at a finite order in the chiral expansion. Except for the leading-order (LO) equation in pionless EFT and in chiral EFT in spin-singlet channels, ultraviolet (UV) divergences emerging from the loop expansion of the scattering amplitude cannot be absorbed into

³ Notice that the contributions from diagrams shown in Fig. 1 are finite in DR.

redefinitions of parameters appearing in the truncated potentials [7, 56]. The problem can be avoided by treating the one-pion exchange (OPE) and higher-order contributions to the potential in perturbation theory using e.g. the systematic power counting scheme proposed by Kaplan, Savage and Wise [57], but the resulting approach unfortunately fails to converge (at least) in certain spin-triplet channels [58, 59], see also Refs. [60] for a recent discussion. A renormalizable framework with the one-pion exchange potential (OPEP) treated nonperturbatively was proposed in [61] based on a manifestly Lorentz invariant form of the effective Lagrangian. This approach requires a perturbative inclusion of higher-order contributions to the potential in order to maintain renormalizability (which may lead to convergence issues in some channels [62]) but has not been systematically explored beyond LO yet.

Throughout this work we employ a finite-cutoff version of nuclear chiral EFT in the formulation of Ref. [48], which is utilized in most of the applications available today. This is so far the only scheme, that has been advanced to high chiral orders and successfully applied to a broad range of few- and many-nucleon systems. Below, we briefly summarize the basic steps involved in the calculation of nuclear observables within this framework. In the following sections, all four steps outlined below will be discussed in detail.

- i. *Derivation of nuclear forces and current operators from the effective chiral Lagrangian.* This can be achieved by separating out irreducible contributions to the A -nucleon scattering amplitude that cannot be generated by iterations of the dynamical equation using various methods outlined in section 3. The derivations are carried out in perturbation theory using the standard chiral power counting. In contrast to ChPT for the scattering amplitude, special efforts are needed to arrive at renormalized nuclear potentials. This requires that all UV divergences from *irreducible* loop diagrams are cancelled by the corresponding counter terms. The renormalizability requirement imposes strong constraints on the unitary ambiguity of nuclear forces and currents [40, 63, 64, 65, 66].
- ii. *Introduction of a regulator for external (off-shell) momenta of the nucleons* in order to make the A -body Schrödinger equation well-behaved. Given the lack of counter terms needed to absorb all UV divergences from iterations of the dynamical equation with a truncated potential, the (momentum-space) cutoff Λ must not be set to arbitrarily high values but should be kept of the order of the breakdown scale, $\Lambda \sim \Lambda_b$ [48, 53, 7]. The accessible cutoff window is, in practice, further restricted by the need to avoid the appearance of spurious deeply bound states which provide a severe complication for applications beyond the NN system [67] and a preference for soft interactions in order to optimize convergence of *ab initio* many-body methods. Given the rather restricted available cutoff window, it is important to employ regulators that minimize the amount of finite-cutoff artifacts, see section 4 for discussion. While the regulator choice for V_{2N} still features a high degree of ambiguity, maintaining the relevant symmetries and consistency with *regularized* many-body forces and exchange currents beyond tree level represents a highly nontrivial task [68, 69], see section 4 for an example and discussion.
- iii. *Renormalization of the few-nucleon amplitude by fixing the short-range multi-nucleon interactions from low-energy experimental data*, see section 6 for details. This allows one to express the calculated scattering amplitude in terms of observable quantities instead of the *bare* LECs $C_{S,T}(\Lambda)$, $C_i(\Lambda)$, $D_i(\Lambda)$, $D(\Lambda)$, $E(\Lambda)$, $E_i(\Lambda)$, $\dots$, and amounts to implicit renormalization of the amplitude. Notice that in the pion and 1N sectors of ChPT, renormalization is usually carried out *explicitly* by splitting the bare LECs l_i , d_i , e_i , $\dots$, into the (finite) renormalized ones and counter terms, e.g. $d_i = d_i^r(\mu) + R_i(\mu)$. Here, μ denotes the renormalization scale while R_i are the corresponding counter terms, which diverge in the limit of a removed regulator (i.e. $\Lambda \rightarrow \infty$ in the cutoff regularization or the number of dimensions $d \rightarrow 4$ in DR). Such a splitting is not unique as reflected by the scale μ , and the appropriate choice of renormalization conditions is essential to maintain the desired power counting, i.e. to ensure the

appropriate scaling behavior of renormalized contributions to the amplitude leading to a systematic and self-consistent scheme, see e.g. Refs. [70, 71]. In the few-nucleon sector, the nonperturbative resummation of pion-exchange potentials via Eq. (2) can only be carried out numerically⁴, which leaves the *implicit* renormalization outlined above as the only available option. Notice that contrary to the renormalized LECs $l_i^r(\mu)$, $d_i^r(\mu)$, $\dots$, the bare LECs $C_{S,T}(\Lambda)$, $C_i(\Lambda)$, $\dots$, must be re-determined at every order in the expansion.

- iv. *Estimation of the truncation uncertainty and a-posteriori consistency checks of the obtained results.* These include, among others, testing the naturalness of the extracted LECs [17], making error plots for phase shifts as suggested in Refs. [48, 72], verifying a reduced residual Λ -dependence of observables (within a specified cutoff range) upon including higher-order short-range interactions, see e.g. Fig. 4 of Ref. [73], and confronting the contributions of many-body interactions and/or exchange currents with estimations based on the assumed power counting [74, 75]. Our approach to error analysis is outlined in section 5, while selected consistency checks are discussed in section 6.

Before closing this section, several remarks are in order. First, we emphasize that the approach outlined above is applicable at the physical quark masses. Quark mass dependence of nuclear observables can be studied more efficiently in the renormalizable chiral EFT framework of Refs. [61, 76], see also Refs. [77, 78, 79] for an alternative method. Secondly, the validity (in the EFT sense) of the finite-cutoff EFT formulation outlined above has been demonstrated numerically by means of the error plots [48, 80] and analytically [53] for toy-models with long-range interactions. It can also be easily verified in pionless EFT. For the case of exactly known non-singular long-range potentials, the employed approach reduces in the NN sector to the well-known modified effective range expansion [81]. The relation between the choice of renormalization conditions and power counting is discussed within pionless EFT in Ref. [82].⁵ That paper provides an explicit example of the choice of subtraction scheme (i.e. renormalization conditions), which leads to a self-consistent EFT approach for two particles with both a natural and unnaturally large scattering length, while respecting the NDA scaling of renormalized LECs. Notice that in all applications reviewed in this article, few-nucleon short-range interactions are counted according to NDA. A number of authors advocate alternative approaches, in particular by inferring the importance of short-range operators from the requirement of Λ -independence of the scattering amplitude at arbitrarily large values of Λ as articulated in detail in Ref. [6]. However, performing the loop expansion of the solution of the LS equation in spin-triplet channels for the resummed OPEP shows that the scattering amplitude is only *partially* renormalized in spite of the fact that it admits, in some cases, a finite $\Lambda \rightarrow \infty$ limit at a fixed energy [7]. The danger of choosing $\Lambda \gg \Lambda_b$ in such partially renormalized nonperturbative expressions is demonstrated using an exactly solvable model in Ref. [53].

3 CHIRAL PERTURBATION THEORY FOR NUCLEAR POTENTIALS

One method to decouple pion-nucleon and purely nucleonic subspaces of the Fock space, thereby reducing a quantum field theoretic problem to a quantum mechanical one, is the unitary transformation (UT) technique. Let η and λ be the projection operators onto the purely nucleonic subspace of the Fock space and the rest, respectively. The time-independent Schrödinger equation can be written in the form

$$\begin{pmatrix} \eta H \eta & \eta H \lambda \\ \lambda H \eta & \lambda H \lambda \end{pmatrix} \begin{pmatrix} \eta |\Psi\rangle \\ \lambda |\Psi\rangle \end{pmatrix} = E \begin{pmatrix} \eta |\Psi\rangle \\ \lambda |\Psi\rangle \end{pmatrix}, \quad (3)$$

⁴ See, however, Ref. [60] for analytical results in the chiral limit.

⁵ For pionless EFT or chiral EFT with perturbative pions, the NN amplitude can be calculated analytically, and renormalization can be carried out explicitly.

where E denotes the eigenenergy of the πN system. The idea is to apply a UT to the Hamilton operator H in order to block diagonalize the matrix on the l.h.s. of Eq. (3) leading to

$$\left[U^\dagger H U \right] U^\dagger |\Psi\rangle = E U^\dagger |\Psi\rangle. \quad (4)$$

The decoupling requirement is given by

$$\eta U^\dagger H U \lambda = \lambda U^\dagger H U \eta = 0. \quad (5)$$

To construct the UT U we first introduce a Møller operator Ω [83], which is defined by

$$|\Psi\rangle = \Omega \eta |\Psi\rangle \quad (6)$$

with the requirement

$$\Omega = \Omega \eta. \quad (7)$$

The Møller operator reproduces the original state out of projected state. By projecting Eq. (6) onto the model space η one obtains the identity

$$\eta \Omega = \eta. \quad (8)$$

Using Eq. (6), we can write the time-independent Schrödinger equation in the form

$$(E - H_0) \Omega \eta |\Psi\rangle = V |\Psi\rangle, \quad (9)$$

where H_0 denotes a free Hamiltonian. On the other hand, projecting the original Schrödinger equation Eq. (3) onto the model space and applying on the resulting equation the operator Ω , we obtain

$$(E \Omega - \Omega H_0) \eta |\Psi\rangle = \Omega \eta V |\Psi\rangle. \quad (10)$$

Subtracting Eq. (10) from Eq. (9) leads to

$$[\Omega, H_0] \eta |\Psi\rangle = (V - \Omega \eta V) |\Psi\rangle = (V - \Omega \eta V) \Omega \eta |\Psi\rangle.$$

This way we obtain a nonlinear equation for the Møller operator Ω

$$[\Omega, H_0] - V \Omega + \Omega V \Omega = 0. \quad (11)$$

Defining the operator A via $\Omega =: \eta + A$ with $A = \lambda A \eta$, as follows from Eqs. (7) and (8), we rewrite Eq. (11) in the form

$$\lambda (H + [H, A] - A V A) \eta = 0. \quad (12)$$

The UT U was parametrized by Okubo [84] in terms of the operator A via

$$U = \begin{pmatrix} \eta (1 + A^\dagger A)^{-1/2} & -A^\dagger (1 + A A^\dagger)^{-1/2} \\ A (1 + A^\dagger A)^{-1/2} & \lambda (1 + A A^\dagger)^{-1/2} \end{pmatrix}. \quad (13)$$

The resulting transformed Hamiltonian

$$\eta U^\dagger H U \eta = (\Omega^\dagger \Omega)^{1/2} \eta H \Omega (\Omega^\dagger \Omega)^{-1/2}, \quad (14)$$

leads to the effective potential defined via

$$V_{\text{eff}}^{\text{UT}} := \eta U^\dagger H U \eta - H_0. \quad (15)$$

Obviously, the Okubo transformation in Eq. (13) is not the only possibility to obtain a block-diagonalized Hamiltonian. On top of the transformation U one can always apply e.g. a UT acting nontrivially on the η -space, thus leaving the Hamiltonian block-diagonal. This freedom has been exploited in a systematic manner to construct renormalizable/factorizable 3NFs and four-nucleon forces (4NFs) in chiral EFT in Refs. [63, 46, 47, 40, 85].

To derive the potential $V_{\text{eff}}^{\text{UT}}$ from the effective chiral Lagrangian in Eq. (1) one needs to solve the nonlinear decoupling equation (12) for the operator A . This can be done perturbatively using NDA [3] to count powers of three-momenta and pion masses, denoted collectively by Q . For the sake of definiteness, we restrict ourselves in the following to nuclear potentials in the absence of external sources. The extension to the current operators is straightforward and discussed in details in Ref. [66]. The irreducible contributions of any connected Feynman diagram scale as Q^ν with $\nu = -2 + \sum_i V_i \kappa_i$, where V_i denotes the number of vertices of type i and κ_i is the inverse mass dimension of the corresponding coupling constant, $\kappa_i = d_i + \frac{3}{2}n_i + p_i - 4$. Here, d_i is the number of derivatives and/or M_π -insertions, while n_i and p_i denote the number of nucleon and pion fields, respectively.⁶ This particular form of the power counting allows one to formulate the chiral expansion in the form that is completely analogous to the expansion in powers of coupling constants. It is thus particularly well suited for algebraic approaches such as the method of UT. Once the operator A is available, one can perform the chiral expansion of Eq. (14) to construct the effective potential order-by-order.

The chiral expansion of the nuclear forces is visualized in Fig. 2. Below, we briefly discuss isospin symmetric contributions starting from the leading order (LO) Q^0 . The only contributions at this order emerge from the OPEP and two contact interactions $\propto C_{S,T}$ [1, 2]. The first corrections at order Q^2 (NLO) involve the leading two-pion exchange potential (TPEP) [23, 86, 87] and 7 short range interactions $\propto C_i$. At order Q^3 (N²LO), further corrections to the TPEP $\propto c_i$ need to be taken into account [86]. At the same order one has the first nonvanishing contributions to the 3NF. They are given by the two-pion exchange diagram involving the LECs c_i and two shorter-range tree-level diagrams involving the LECs D and E [88, 36]. At order Q^4 (N³LO), the NN potential receives the contributions from the leading three-pion exchange [89, 90, 91], further corrections to the TPEP [92, 93] and 12 new short-range interactions $\propto D_i$ [17]. At the same order, there are various one-loop corrections to the 3NF [46, 47, 94] and the first contributions to the 4NFs [63, 64], which do not involve unknown parameters. Finally, at order Q^5 (N⁴LO), the NN potential receives corrections to the three-pion exchange $\propto c_i$ [91] and further contributions to the TPEP [95]. No additional unknown parameters appear in the isospin-conserving part of the NN force at this order. The 3NF also receives corrections at N⁴LO, most of which have already been worked out using DR [40, 85, 37]. Notice that the 3NF involves at this order a number of new short-range operators. Work is still in progress to derive the remaining 3NF and 4NF at N⁴LO. We further emphasize that all calculations mentioned above are carried out using DR or equivalent schemes.

⁶ Alternatively (but equivalently), the chiral order ν of a connected, N -nucleon irreducible diagram with L loops can be expressed as $\nu = -4 + 2N + 2L + \sum_i V_i \Delta_i$ with $\Delta_i = d_i + n_i/2 - 2$.

	Two-nucleon force	Three-nucleon force	Four-nucleon force
LO (Q^0)			
NLO (Q^2)			
N ² LO (Q^3)			
N ³ LO (Q^4)			
N ⁴ LO (Q^5)			

Figure 2. Hierarchy of nuclear forces at increasing orders in chiral expansion in the Weinberg scheme. Solid and dashed lines refer to nucleons and pions, respectively. Solid dots, filled circles, filled squares, filled diamonds and open squares refer to vertices from the Lagrangian in Eq. (1) of dimension $\Delta = 0, 1, 2, 3$ and 4, respectively.

The effective potential $V_{\text{eff}}^{\text{UT}}$ leads, by construction, to the same spectrum and on-shell scattering matrix as the original untransformed potential V [96, 97]. There are, however, other possibilities to define the effective potential without changing on-shell physics. One example is an energy-independent potential defined by

$$V_{\text{eff}}^{\text{EI}} = \eta V \Omega \eta. \quad (16)$$

The proof that $V_{\text{eff}}^{\text{EI}}$ of Eq. (16) leads to the same spectrum is trivial:

$$\begin{aligned} (\eta H_0 \eta + \eta V \eta + \eta V \lambda) |\Psi\rangle &= E \eta |\Psi\rangle, \\ (\eta H_0 \eta + \eta V \eta + \eta V \lambda \Omega) \eta |\Psi\rangle &= E \eta |\Psi\rangle, \\ (\eta H_0 \eta + \eta V \Omega \eta) \eta |\Psi\rangle &= E \eta |\Psi\rangle, \end{aligned} \quad (17)$$

where we used Eqs. (6) and (8) in the first and second lines, respectively. Note that the potential $V_{\text{eff}}^{\text{EI}}$ is manifestly non-hermitian. However, due to its simplicity, it is widely used in the literature [83]. This example shows that there is a considerable freedom to define nuclear potentials. Nuclear forces and current operators constructed by the Bochum-Bonn group, see e.g. Refs. [87, 64, 46, 47, 65, 40, 85, 21, 22, 66, 17], are obtained using the method of UT. The JLab-Pisa group utilizes a different approach by starting with the on-shell transfer matrix T and “inverting” it to obtain the effective potential, see e.g. Refs. [98, 99, 100, 101].

This is carried out in perturbation theory by counting the nucleon mass via $m \sim \Lambda_b$

$$T = T^{(0)} + T^{(1)} + T^{(2)} + \dots, \quad (18)$$

where the superscripts indicate the chiral order Q^n . The same counting scheme is used to organize the contributions to effective potential:

$$v = v^{(0)} + v^{(1)} + v^{(2)} + \dots. \quad (19)$$

The inversion of the LS equation is carried out iteratively to yield

$$v^{(0)} = T^{(0)}, \quad v^{(1)} = T^{(1)} - v^{(0)} G_0 v^{(0)}, \quad \dots \quad (20)$$

Obviously, the knowledge of the on-shell transfer matrix is insufficient to perform the inversion, and one needs to specify its off-shell extension. Notice that the potentials constructed in this way are not necessarily hermitian, and thus there is no guarantee that they are unitarily equivalent to the ones derived using the UT technique. It should, however, always be possible to find a similarity transformation that relates one potential to another. This is exemplified with the potential $v_{2\pi}^{(3)}(\nu = 1)$ in Eq. (20) of Ref. [99], where ν is an arbitrary phase, which is manifestly non-hermitian. Using the similarity transformation in Eq. (28) of that paper⁷, it can be transformed to the hermitian potential $v_{2\pi}^{(3)}(\nu = 0)$, that is actually employed in the current version of the interactions developed by the JLab-Pisa group. With this choice, their potentials are unitarily equivalent to the ones of the Bochum-Bonn group.

4 REGULARIZATION

4.1 Semilocal momentum-space regularization of the NN potential

In this review article we focus on the semilocal regularization approach of the chiral nuclear potentials carried out in momentum space [17]. For the purpose of regularization we will consider the two-nucleon interaction consisting of two distinct parts: the short-range contact interaction part and the long-range pion-exchange part. In this context, the term "semilocal" refers to the application of a nonlocal regulator for the former and a local regulator for the latter. In particular, the momentum-space matrix elements of the contact potential are multiplied by a simple nonlocal Gaussian regulator

$$\langle \vec{p}' | V_{\text{cont}} | \vec{p} \rangle_{\text{reg}} = \langle \vec{p}' | V_{\text{cont}} | \vec{p} \rangle e^{-\frac{p'^2 + p^2}{\Lambda^2}}. \quad (21)$$

Here and in what follows, $p \equiv |\vec{p}|$ and $p' \equiv |\vec{p}'|$. Such kinds of nonlocal regulators (albeit with different powers of p , p' and Λ) have been and still are employed as the main method of regularization for the entire potential including the long-range interactions, see e.g. Refs. [102, 103, 104, 105, 106, 107].⁸

However, in Refs. [21, 22] it was shown that the amount of long-range cutoff artifacts can be significantly reduced by employing a local regulator for pion-exchange potentials. Notice that pion-exchange contributions, except for some relativistic corrections, give rise to local potentials. We *require* the regulator to preserve the analytic structure of the scattering amplitude and, in particular, the left-hand cuts

⁷ The claim in Ref. [99] that the transformation e^{iU} in Eq. (28) of that paper is unitary is incorrect since the operator $iU^{(1)}(\nu)$ from Eq. (28) is not antihermitian.

⁸ Notice that the aforementioned potentials (except the one of Ref. [102]) additionally apply spectral function regularization (SFR) [108, 109] of the TPEP in the form of a sharply cut-off spectral integral in order to suppress its remaining unphysical short-distance behavior. Notice, however, that the application of a nonlocal regulator $\exp(-(p^{2n} + p'^{2n})/\Lambda^{2n})$ with suitably chosen n is sufficient to arrive at UV-finite iterations of the potential.

in the complex energy plane related to the long-range part of the interaction, which are unambiguously determined in chiral EFT. Inspired by Ref. [110], this is achieved in our momentum-space approach by regularizing the static propagators of pions exchanged between different nucleons with a local Gaussian cutoff via

$$\frac{1}{l^2 + M_\pi^2} \rightarrow \frac{1}{l^2 + M_\pi^2} e^{-\frac{l^2 + M_\pi^2}{\Lambda^2}}, \quad (22)$$

with $l = |\vec{l}|$ and $\vec{l}$ denoting the three-momentum of the exchanged pion. The introduction of the Gaussian form factor in the pion propagators leads to properly regularized long-range potentials that are finite at short distances in coordinate space. In order to have a clean separation of the long-range pion-exchange potential from the short-range contact interactions, we made use of the available contact interactions to subtract out the remaining (finite) admixtures of short-range interactions [17]. The fixed coefficients of these subtractions are determined from the requirement that the corresponding coordinate-space potential and as many derivatives thereof as allowed by power counting vanish at the origin. This convention leads to a qualitatively similar regularization as the coordinate-space regulator previously employed in Refs. [21, 22].

Application of these ideas to the OPEP is straightforward and leads, in the limit of exact isospin symmetry, to

$$V_{2N,\Lambda}^{1\pi}(M_\pi) = -\frac{g_A^2}{4F_\pi^2} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \left(\frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{q^2 + M_\pi^2} + C(M_\pi) \vec{\sigma}_1 \cdot \vec{\sigma}_2 \right) e^{-\frac{q^2 + M_\pi^2}{\Lambda^2}}, \quad (23)$$

where $q \equiv |\vec{q}| \equiv |\vec{p}' - \vec{p}|$ and $\vec{\sigma}_i$ ($\boldsymbol{\tau}_i$) are the Pauli spin (isospin) matrices of the i -th nucleon. Here, the static pion propagator has been regularized according to Eq. (22) and a likewise-regularized LO contact interaction has been added to the OPEP. Its coefficient $C(M_\pi)$,

$$C(M_\pi) = -\frac{\Lambda (\Lambda^2 - 2M_\pi^2) + 2\sqrt{\pi} M_\pi^3 e^{\frac{M_\pi^2}{\Lambda^2}} \text{erfc}\left(\frac{M_\pi}{\Lambda}\right)}{3\Lambda^3}, \quad (24)$$

with $\text{erfc}(z)$ denoting the complementary error function, is fixed by the requirement that the spin-spin part of the OPEP in coordinate space vanishes at the origin. For the regularization of the TPEP, we start with a generic three-dimensional loop integral $I(\vec{q})$ arising in the derivation of the TPEP using e.g. the method of unitary transformation as detailed in the previous section or comparable approaches like time-ordered perturbation theory or S-matrix-based methods [86]. As discussed in Ref. [110], the pion energy denominators in the corresponding 1-loop expressions can always be rewritten into an integral over a mass parameter λ involving a product of two static pion propagators with mass $\sqrt{M_\pi^2 + \lambda^2}$

$$I(\vec{q}) = \int_0^\infty d\lambda \int \frac{d^3 l_1}{(2\pi)^3} \frac{d^3 l_2}{(2\pi)^3} (2\pi)^3 \delta(\vec{q} - \vec{l}_1 - \vec{l}_2) \frac{1}{(l_1^2 + M_\pi^2 + \lambda^2)(l_2^2 + M_\pi^2 + \lambda^2)} \times \dots, \quad (25)$$

where $\vec{l}_1$ and $\vec{l}_2$ denote the three-momenta of the exchanged pions and the ellipses refer to additional momentum-spin-isospin structures arising from the vertices of a particular diagram. With the pion propagators factorized in this a way, we can regularize them by applying the prescription specified in Eq. (22) to each of them. Although the introduction of the regulator obviously affects the resulting expression for the TPEP, there is no need to rederive them explicitly. Indeed, the scalar functions accompanying the spin-isospin operators in the *unregularized* TPEP can be expressed using the dispersive representation

$$V_{2N}^{2\pi}(q) = \frac{2}{\pi} \int_{2M_\pi}^\infty \mu d\mu \frac{\rho(\mu)}{q^2 + \mu^2}, \quad (26)$$

with the spectral functions $\rho(\mu) = \Im(V_{2\pi}(q))|_{q=0+i\mu}$ which are readily available up to N⁴LO. For the explicit expressions of the TPEP, additional subtractions of short-range terms have to be performed to arrive at a convergent spectral integral in Eq. (26) whose number depends on the chiral order of the contribution at hand. Introducing the pion propagator regulators in Eq. (25), the regularized generic spectral integral of Eq. (26) takes the form

$$V_{2N,\Lambda}^{2\pi}(q) = e^{-\frac{q^2}{2\Lambda^2}} \frac{2}{\pi} \int_{2M_\pi}^{\infty} \mu d\mu \frac{\rho(\mu)}{q^2 + \mu^2} e^{-\frac{\mu^2}{2\Lambda^2}}. \quad (27)$$

As the spectral representation in Eq. (26) is a dispersion relation, it makes the analytic structure of the TPEP as a function of the momentum-transfer $\vec{q}$ manifest. Indeed, one could immediately infer the form of the regulator as a function of q^2 and μ^2 by the requirement to preserve the analytic structure of the TPEP. The additional factor of 1/2 in the exponent of Eq. (27) would, however, not have been taken into account correctly if we had naively applied our prescription Eq. (22) to the propagator-like denominator of the spectral integral representation.

Expanding the exponentials in inverse powers of the cutoff in either Eq. (23) or Eq. (27), one observes that the regulator indeed does not affect the long-range part of the potential to any order, but generates an infinite series of short-range terms polynomial in q^2 . Since an increasing number of contact interactions of this form with freely adjustable LECs become available with increasing chiral order, the perturbative restoration of cutoff-independence is also obvious in this scheme.

The expressions of the regularized and subtracted TPEP can be found in Ref. [17]. Here we restrict ourselves to the example of the isospin-independent central part of the leading TPEP at NLO which is given by

$$W_{C,\Lambda}^{(2)}(q) = e^{-\frac{q^2}{2\Lambda^2}} \frac{2}{\pi} \int_{2M_\pi}^{\infty} \frac{d\mu}{\mu^3} \rho_C^{(2)}(\mu) \left(\frac{q^4}{\mu^2 + q^2} + C_{C,1}^2(\mu) + C_{C,2}^2(\mu) q^2 \right) e^{-\frac{\mu^2}{2\Lambda^2}}, \quad (28)$$

with the spectral function

$$\rho_C^{(2)}(\mu) = \frac{\sqrt{\mu^2 - 4M_\pi^2}}{768\pi F_\pi^4 \mu} \left(4M_\pi^2(5g_A^4 - 4g_A^2 - 1) - \mu^2(23g_A^4 - 10g_A^2 - 1) + \frac{48g_A^4 M_\pi^4}{4M_\pi^2 - \mu^2} \right). \quad (29)$$

Two subtractions have been performed in order to render the unregularized spectral integral in Eq. (28) convergent and according to our convention, we have additionally fixed the subtraction coefficients $C_{C,1}^2(\mu)$ and $C_{C,2}^2(\mu)$ by the requirement that $W_{C,\Lambda}^{(2)}(r)|_{r=0} = \frac{d^2}{dr^2} W_{C,\Lambda}^{(2)}(r)|_{r=0} = 0$. (The first derivative of $W_{C,\Lambda}^{(2)}(r)$ vanishes at the origin regardless of the subtraction coefficients.) Figure 3 shows the ratio of the regularized and unregularized expressions in Eq. (28) in coordinate space. As one can see, the behavior of the regularized potential is smoother when fixing the subtraction coefficients by the convention explained above. Also note that the potential with $C_{C,1}^2(\mu) = C_{C,2}^2(\mu) = 0$ does not vanish at the origin.⁹

4.2 Regularization and consistency of nuclear forces

Having defined the regularization scheme in the NN sector, we now turn to regularization of the 3NF. The expressions for the 3NFs described in section 3 have been worked out completely through N³LO using DR. They are off-shell consistent with the unregularized NN interactions reviewed in that section in the way explained in section 2. To arrive at regularized 3NFs, it is tempting to apply some kind of multiplicative

⁹ This is not visible in Fig. 3 since the unregularized potential $W_{C,\infty}(r)$ is singular at $r = 0$.




Figure 3. Ratio of the regularized and unregularized central part of the leading TPEP in coordinate space for $C_{C,1}^2(\mu)$, $C_{C,2}^2(\mu)$ fixed as discussed in the text and $C_{C,1}^2(\mu) = C_{C,2}^2(\mu) = 0$.

regulators to the expressions of the 3NF derived using DR. Such a naive approach, however, leads to a violation of the chiral symmetry at N³LO and destroys the consistency between two- and three-nucleon forces after regularization.

To illustrate the problem consider the diagrams shown in Fig. 1, which have already been discussed in section 2. The 3NF entering the first graph on the right-hand side (r.h.s.) is given by [47]

$$V_{3N}^{2\pi, 1/m} = i \frac{g_A^2}{32mF_\pi^4} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_3 \cdot \vec{q}_3}{(q_1^2 + M_\pi^2)(q_3^2 + M_\pi^2)} \boldsymbol{\tau}_1 \cdot (\boldsymbol{\tau}_2 \times \boldsymbol{\tau}_3) (2\vec{k}_1 \cdot \vec{q}_3 + 4\vec{k}_3 \cdot \vec{q}_3 + i [\vec{q}_1 \times \vec{q}_3] \cdot \vec{\sigma}_2), \quad (30)$$

with $\vec{q}_i = \vec{p}'_i - \vec{p}_i$, $\vec{k}_i = 1/2(\vec{p}'_i - \vec{p}_i)$ and $\vec{p}_i$, $(\vec{p}'_i)$ the initial (final) momenta of the i -th nucleon. The complete expression for the relativistic corrections to the 3NF at N³LO can be found in Ref. [47]. We now consider the first iteration of $V_{3N}^{2\pi, 1/m}$ with the static OPEP

$$V_{2N}^{1\pi} = - \left(\frac{g_A}{2F_\pi} \right)^2 \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{q^2 + M_\pi^2} \quad (31)$$

as shown by the first diagram on the r.h.s. of Fig. 1. By simply counting the powers of momenta in the loop integration one observes that the loop integral is linearly divergent, which leads to a finite result in DR. As already pointed out in section 2, adding the DR expression for the 3NF $V_{3N}^{2\pi-1\pi}$ from Eqs. (2.16)-(2.20) of Ref. [46] yields (on-shell) the same result as obtained from calculating the Feynman diagram on the l.h.s. of Fig. 1 as expected for consistent two- and three-nucleon forces.

We now repeat this exercise using the semilocally regularized nuclear potentials

$$V_{3N,\Lambda}^{2\pi, 1/m} = V_{3N}^{2\pi, 1/m} e^{-\frac{q_1^2 + M_\pi^2}{\Lambda^2}} e^{-\frac{q_3^2 + M_\pi^2}{\Lambda^2}}, \quad V_{2N,\Lambda}^{1\pi} = V_{2N}^{1\pi} e^{-\frac{q^2 + M_\pi^2}{\Lambda^2}}, \quad (32)$$

in the calculation of the first diagram on the r.h.s. of Fig. 1. This leads to

$$\begin{aligned} V_{3N,\Lambda}^{2\pi,1/m} G_0 V_{2N,\Lambda}^{1\pi} + V_{2N,\Lambda}^{1\pi} G_0 V_{3N,\Lambda}^{2\pi,1/m} &= \Lambda \frac{g_A^4}{128\sqrt{2}\pi^{3/2}F_\pi^6} (\boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 - \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3) \frac{\vec{q}_2 \cdot \vec{\sigma}_2 \vec{q}_3 \cdot \vec{\sigma}_3}{q_3^2 + M_\pi^2} \\ &- \Lambda \frac{g_A^4}{96\sqrt{2}\pi^{3/2}F_\pi^6} \frac{\vec{q}_3 \cdot \vec{\sigma}_3 \vec{q}_3 \cdot \vec{\sigma}_1}{q_3^2 + M_\pi^2} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3 + \dots, \end{aligned} \quad (33)$$

where the ellipses refer to all permutations of the nucleon labels and terms finite in the $\Lambda \rightarrow \infty$ -limit. The linear divergence $\propto \vec{q}_3 \cdot \vec{\sigma}_3 \vec{q}_3 \cdot \vec{\sigma}_1$ is cancelled by the D counter term in the second 3NF diagram at N²LO in Fig. 2. To cancel the linearly divergent contribution $\propto \vec{q}_2 \cdot \vec{\sigma}_2$ one would, however, need to introduce a vertex in $\mathcal{L}_{\pi NN}^{(1)}$ corresponding to a derivative-less coupling of the pion to the NN systems. Such vertices violate the chiral symmetry and, being suppressed by powers of M_π^2 , cannot appear in $\mathcal{L}_{\pi NN}^{(1)}$. As a consequence, this linear divergence can *not* be absorbed into redefinition of the LECs, and the amplitude on the r.h.s. of Fig. 1 can seemingly not be renormalized (i.e. made finite in the $\Lambda \rightarrow \infty$ limit). The r.h.s. of the shown equation, therefore, apparently cannot match the (renormalizable) on-shell scattering amplitude from the Feynman diagram on the l.h.s.. The problem can be traced back to mixing the DR when calculating the 3NF $V_{3N}^{2\pi-1\pi}$ with a cutoff regularization for the iterative contributions in Eq. (33), see Ref. [68] for another example with the NN axial vector current operator at N³LO. Indeed, recalculating the loop integral in $V_{3N}^{2\pi-1\pi}$ using the cutoff-regularized pion propagators leads to

$$V_{3N,\Lambda}^{2\pi-1\pi} = -\Lambda \frac{g_A^4}{128\sqrt{2}\pi^{3/2}F_\pi^6} (\boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 - \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3) \frac{\vec{q}_2 \cdot \vec{\sigma}_2 \vec{q}_3 \cdot \vec{\sigma}_3}{q_3^2 + M_\pi^2} - \Lambda \frac{g_A^4}{32\sqrt{2}\pi^{3/2}F_\pi^6} \frac{\vec{q}_3 \cdot \vec{\sigma}_3 \vec{q}_3 \cdot \vec{\sigma}_1}{q_3^2 + M_\pi^2} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3 + \dots, \quad (34)$$

where the ellipses refer to the finite terms in the $\Lambda \rightarrow \infty$ -limit. The problematic linear divergence cancels exactly and the agreement with the on-shell amplitude from the Feynman diagram is restored when both *consistently regularized* contributions on the r.h.s. of Fig. 1 are added together.

One may worry whether the regularization issues discussed above could also be relevant for NN interactions. Fortunately, this is not the case since the momentum structure of the NN contact interactions is not restricted by the chiral symmetry. UV divergences emerging from iterations of the LS equation can, therefore, always be absorbed into redefinition of the bare LECs $C_{S,T}(\Lambda)$, $C_i(\Lambda)$, $\dots$

In the considered example with the 3N amplitude, the consistently regularized 3NF could be obtained by simply recalculating $V_{3N}^{2\pi-1\pi}$ with all pion propagators being regularized according to Eq. (22). This would indeed solve the problem with the cancelation of linear divergencies at N³LO, but it would still lead to a violation of the chiral symmetry in diagrams involving three- and four-pion vertices, which depend on the parametrization of the pion field. For vertices involving up to four pion fields, this freedom is represented by a single real parameter α . In the effective chiral Lagrangian, all pion fields are collected in an SU(2) matrix $U(\boldsymbol{\pi})$, whose most general expression, expanded in powers of the pion fields, takes the form

$$U(\boldsymbol{\pi}) = 1 + \frac{i}{F} \boldsymbol{\pi} \cdot \boldsymbol{\tau} - \frac{1}{2F^2} (\boldsymbol{\pi} \cdot \boldsymbol{\tau})^2 - \alpha \frac{i}{F^3} (\boldsymbol{\pi} \cdot \boldsymbol{\tau})^3 + \left(\alpha - \frac{1}{8} \right) \frac{1}{F^4} (\boldsymbol{\pi} \cdot \boldsymbol{\tau})^4 + O(\boldsymbol{\pi}^5). \quad (35)$$

Clearly, the on-shell amplitude must be independent of the arbitrary parameter α . Evaluating the 3NF and 4NF with the regularized pion propagators, however, leads to α -dependent expressions (for finite values of Λ). This shows, perhaps not surprisingly, that the simplistic approach by regularizing all pion propagators as described above violates the chiral symmetry. A possible solution of this problem is provided

by the symmetry preserving higher derivative regularization method introduced by Slavnov [111], see also Refs. [112, 113] for recent applications in chiral EFT.

To summarize, we have shown that a naive regularization of the three- and more-nucleon forces by multiplying the expressions derived in DR with regulator functions leads to inconsistencies starting from $N^3\text{LO}$, see Ref. [68] for the same conclusion for two- and more-nucleon charge and current operators. This problem is by no means restricted to semilocal cutoffs. To derive many-body forces and currents regularized *consistently* with the NN potentials of Ref. [17], the expressions for the 3NF of Refs. [46, 47, 40], 4NF of Ref. [64] and exchange charge and current operators of Refs. [114, 65, 66, 115] need to be recalculated using e.g. an appropriately chosen higher derivative regulator at the level of the effective Lagrangian.

5 TRUNCATION ERROR ANALYSIS

Estimating the uncertainty associated with truncations of the EFT expansion, which typically dominates the error budget (see section 6), is an important task – in particular since chiral EFT is being developed into a precision tool. In the past, truncation errors were often estimated in few-nucleon calculations from a residual cutoff dependence. This approach, however, suffers from several drawbacks and does not allow for a reliable estimation of truncation errors [103]. In Ref. [21], we have formulated a simple algorithm to estimate the size of neglected higher-order terms based on the available information about the EFT expansion pattern for any given observable. To be specific, consider an arbitrary NN scattering observable X at the center of mass momentum p , which is calculated in chiral EFT up to the order Q^k

$$X(p) = X^{(0)} + \Delta X^{(2)} + \Delta X^{(3)} + \dots + \Delta X^{(k)} + \Delta X^{(k+1)} + \dots \equiv X^{(k)} + \delta X^{(k)}. \quad (36)$$

The corrections $\Delta X^{(i)}$, $\Delta X^{(i)} = \mathcal{O}(X^{(0)}Q^i)$, are assumed to be known explicitly up to the order $i = k$. The goal is to estimate the size of neglected higher-order terms $\delta X^{(k)} = \sum_{n>k} \Delta X^{(n)}$. We, furthermore, assume that the expansion parameter Q is given by

$$Q = \max\left(\frac{M_\pi^{\text{eff}}}{\Lambda_b}, \frac{p}{\Lambda_b}\right). \quad (37)$$

This simple ansatz is motivated by the expectation that at very low energies, the errors are dominated by the expansion around the chiral limit. The scale M_π^{eff} , which will be specified below, is related to the pion mass and controls the convergence rate of the expansion around the chiral limit. At higher energies one would, on the other hand, expect the expansion to be dominated by powers of momenta. This simple picture is in qualitative agreement with the error plots for NN phase shifts [21], which show clearly the two different regimes mentioned above, see Ref. [73] for a discussion.

The algorithm proposed by Epelbaum, Krebs and Meißner (EKM) in Ref. [21] employs $M_\pi^{\text{eff}} = M_\pi$ and $\Lambda_b = 600$ MeV based on the estimation from the error plots. It also assumes the truncation error $\delta X^{(k)}$ to be dominated by the first neglected term. The truncation errors at orders Q^i , $0 \leq i \leq k$, are then estimated via

$$\delta X^{(0)} = Q^2 |X^{(0)}|, \quad \delta X^{(i)} = \max_{2 \leq j \leq i} \left(Q^{i+1} |X^{(0)}|, Q^{i+1-j} |\Delta X^{(j)}| \right) \text{ for } i \geq 2, \quad (38)$$

subject to the additional constraint

$$\delta X^{(i)} \geq \max_{j, m=i, \dots, k} \left(|X^{(j)} - X^{(m)}| \right), \quad (39)$$

which ensures that the estimated errors cannot be smaller than the known actual higher-order contributions. Notice that this relation leads, per construction, to overlapping errors at different orders. In Ref. [74], the method was adjusted to make it applicable to incomplete calculations of few-body observables based on NN interactions only. The EKM approach was applied to a broad range of low-energy reactions in the single-baryon [116, 117, 43, 118] as well as few- and many-nucleon [22, 119, 120, 8, 121] sectors. The robustness of this method and some alternative algorithms are discussed in Ref. [122]. The obvious drawback of the EKM approach is that the estimated uncertainties do not offer a statistical interpretation.

In Refs. [123, 124, 125], a more general and statistically well-founded Bayesian approach was developed to calculate the probability distribution function (pdf) for truncation errors in chiral EFT. The EKM approach was then shown to correspond to one particular choice of prior probability distribution for the coefficients in the chiral expansion of $X(p)$. In Ref. [123], the EKM uncertainties for the np total cross section were found to be consistent with 68% degree-of-belief (DoB) intervals. The authors of that paper, furthermore, found using the semilocal coordinate-space regularized (SCS) potentials of Refs. [21, 22] the assumed value of the breakdown scale of $\Lambda_b = 600$ MeV to be statistically consistent for not too soft regulator values, see also Ref. [124] for a related discussion. Recently, a slightly modified version of the Bayesian approach developed in Refs. [123, 124] was applied by the LENPIC Collaboration to study NN and 3N scattering [75]. Below, we briefly outline the Bayesian model $\bar{C}_{0.5-10}^{650}$ proposed in that paper, which will be employed throughout section 6. For more details on the Bayesian approach the reader is referred to the original publications [123, 124].

We begin with rewriting Eq. (36) in terms of dimensionless expansion coefficients c_i by introducing a (generally dimensionfull) scale X_{ref} via

$$X = X_{\text{ref}} (c_0 + c_2 Q^2 + c_3 Q^3 + c_4 Q^4 + \dots), \quad (40)$$

where¹⁰

$$X_{\text{ref}} = \begin{cases} X_{\text{ref}} = \max(|X^{(0)}|, Q^{-2}|\Delta X^{(2)}|) & \text{for } k = 2, \\ X_{\text{ref}} = \max(|X^{(0)}|, Q^{-2}|\Delta X^{(2)}|, Q^{-3}|\Delta X^{(3)}|) & \text{for } k \geq 3. \end{cases} \quad (41)$$

This choice of the reference scale was found in Ref. [75] to be more robust for observables that depend on continuously varying parameters, as compared with the choice of $X_{\text{ref}} = |X^{(0)}|$ adopted in Ref. [124]. Alternatively, correlations between observables (and thus the coefficients c_i) evaluated at different values of continuously varying parameters can be taken into account using Gaussian processes [125]. Except for the coefficient $c_m = 1$, $m \in \{0, 2, 3\}$, used to set the scale X_{ref} , the expansion coefficients c_i are assumed to be distributed according to some common pdf $\text{pr}(c_i|\bar{c})$ with a hyperparameter $\bar{c}$. Performing marginalization over h chiral orders $k+1, \dots, k+h$, which are assumed to dominate the truncation error, the probability distribution for the dimensionless residual $\Delta_k \equiv \sum_{n=k+1}^{\infty} c_n Q^n \simeq \sum_{n=k+1}^{k+h} c_n Q^n$ to take a value $\Delta_k = \Delta$, given the knowledge of $\{c_{i \leq k}\}$, is given by [124]

$$\text{pr}_h(\Delta|\{c_{i \leq k}\}) = \frac{\int_0^\infty d\bar{c} \text{pr}_h(\Delta|\bar{c}) \text{pr}(\bar{c}) \prod_{i \in A} \text{pr}(c_i|\bar{c})}{\int_0^\infty d\bar{c} \text{pr}(\bar{c}) \prod_{i \in A} \text{pr}(c_i|\bar{c})}, \quad (42)$$

¹⁰ No meaningful uncertainty estimation can be carried out within the Bayesian approach at LO.

where the set A is defined as $A = \{n \in \mathbb{N}_0 \mid n \leq k \wedge n \neq 1 \wedge n \neq m\}$ and

$$\text{pr}_h(\Delta|\bar{c}) \equiv \left[\prod_{i=k+1}^{k+h} \int_{-\infty}^{\infty} dc_i \text{pr}(c_i|\bar{c}) \right] \delta\left(\Delta - \sum_{j=k+1}^{k+h} c_j Q^j\right). \quad (43)$$

The model $\bar{C}_{0.5-10}^{650}$ utilizes the Gaussian prior of “set C” from Ref. [124],

$$\text{pr}(c_i|\bar{c}) = \frac{1}{\sqrt{2\pi\bar{c}}} e^{-c_i^2/(2\bar{c}^2)}, \quad \text{pr}(\bar{c}) = \frac{1}{\ln(\bar{c}_>/\bar{c}_<)} \frac{1}{\bar{c}} \theta(\bar{c} - \bar{c}_<) \theta(\bar{c}_> - \bar{c}), \quad (44)$$

for which the integrals in Eq. (42) can be performed analytically [124], and uses the values of $h = 10$, $\bar{c}_< = 0.5$ and $\bar{c}_> = 10$. Following Ref. [126], the scales that control the expansion parameter are set to $M_\pi^{\text{eff}} = 200$ MeV and $\Lambda_b = 650$ MeV. The sensitivity of the estimated uncertainties to the choice of prior pdf is discussed in Refs. [123, 124, 75]. One generally finds minor dependence on the prior pdf if a sufficient amount of information on the coefficients c_i is available.

6 SELECTED RESULTS

6.1 The two-nucleon system

We now turn to the calculation of phase shifts and observables in the two-nucleon system. While the derivation and regularization of the nuclear forces have been outlined in the previous sections, we also need to specify the numerical values of all relevant physical quantities and LECs to perform actual calculations. For pion-exchange contributions to the potential, all LECs can be extracted from processes involving at most one nucleon, making it parameter-free. In the TPEP, we use the values of the πN LECs as determined recently by matching the πN scattering amplitude from chiral perturbation theory to a Roy-Steiner equations analysis of πN scattering data at the subthreshold point [42].

We account for the isospin-breaking effects due to the different pion masses in the OPEP and employ the physical masses of the charged and neutral pions $M_{\pi^\pm} = 139.57$ MeV and $M_{\pi^0} = 134.98$ MeV, while we use the isospin-averaged value of $M_\pi = 138.03$ MeV in the TPEP. We adopt an effective value of $g_A = 1.29$ for the nucleon axial coupling constant which is slightly larger than the current experimental average value of $g_A = 1.2723(23)$ [127] because it already accounts for the Goldberger-Treiman discrepancy, see Ref. [128] for a related discussion. The employed value of the pion decay constant is $F_\pi = 92.4$ MeV.

In contrast to the parameter-free long-range potential, the short-range contact interactions in the two-nucleon force have to be determined from experimental NN data. In order to achieve a proper reproduction of pp and, to a lesser extent, np scattering data, it is crucial to also include electromagnetic interactions between the nucleons. Although these interactions are accompanied by powers of a numerically small coupling constant $\alpha \sim 1/137$, they are enhanced at low energies and/or forward angles due to the infrared singularity of the photon propagator or, equivalently, due to their long-range nature. Here, we follow the treatment of the Nijmegen group [129] and include the so-called improved Coulomb potential [130], the magnetic-moment interaction [131] as well as the vacuum-polarization potential [132] in the calculation of proton-proton observables. The magnetic moment interaction is also taken into account in neutron-proton scattering. To the best of our knowledge, these effects have been included in every partial-wave analysis (PWA) of or fit of a high-quality potential model from NN data since the Nijmegen PWA of Ref. [129],

Table 1. χ^2/datum for the description of the neutron-proton and proton-proton scattering data at various orders in the chiral expansion for $\Lambda = 450$ MeV. The numbers in brackets after the order indicate the number of parameters entering the neutron-proton and proton-proton potentials.

E_{lab} bin	LO ₍₃₎	NLO ₍₁₀₎	N ² LO ₍₁₀₎	N ³ LO ₍₂₂₎	N ⁴ LO ₍₂₃₎	N ⁴ LO ⁺ ₍₂₇₎
neutron-proton scattering data						
0–100	73	2.2	1.2	1.07	1.07	1.07
0–200	62	5.4	1.7	1.09	1.08	1.06
0–300	75	14	4.2	2.01	1.16	1.06
proton-proton scattering data						
0–100	2290	10	2.2	0.90	0.88	0.86
0–200	1770	90	37	1.99	1.42	0.95
0–300	1380	90	41	3.43	1.67	1.00

so that differences in their predictions stem from modeling the strong interaction, the experimental input and/or details of the fitting procedure itself.

For scattering data we use the Granada-2013 database [18] which consists of experimental data for NN elastic scattering up to $E_{\text{lab}} = 350$ MeV from 1950 up to 2013.¹¹ The database contains the data that have been found to be mutually compatible by means of a 3σ rejection criterion in the corresponding phase shift analysis of Ref. [18]. The presence of very precisely measured proton-proton data in the database, such as those of Ref. [134], motivated us to introduce the N⁴LO⁺ version of the potential. As the proper description of these data requires a precise reproduction of F-waves, the N⁴LO⁺ potential adds the four leading F-wave contact interactions

$$\langle {}^S F_j, p' | V_{\text{cont}} | {}^S F_j, p \rangle = E_{SF_j} p^3 p'^3, \quad (45)$$

formally appearing at N⁵LO and entering the 3F_2 , 1F_3 , 3F_3 and 3F_4 partial waves, to the N⁴LO potential.

The fits have been performed for all cutoffs $\Lambda = 400, 450, 500$ and 550 MeV as well as for all orders from LO up to N⁴LO⁺.¹² When determining the values of the contact LECs, one has to decide up to which energy E_{lab} the experimental data should be taken into account. One is faced with the two competing features: on the one hand, the inclusion of as many data as possible is desirable from a data fitting point of view. On the other hand, the truncation errors for the chiral interactions become larger at high energies. We therefore chose the maximum energy E_{lab} of data to be included to be $E_{\text{max}} = 260$ MeV for N⁴LO and N⁴LO⁺, while we reduced the energy to $E_{\text{max}} = 25, 100, 125$ and 200 MeV at the orders LO, NLO, N²LO and N³LO, respectively. From N³LO on, we also adjust the deuteron binding energy B_d and the coherent neutron-proton scattering length b_{np} to reproduce their experimental values of $B_d = 2.224575(9)$ MeV [135] and $b_{\text{np}} = -3.7405(9)$ fm [136].

Table 1 shows the reproduction of neutron-proton and proton-proton scattering data in terms of χ^2/datum values at all considered orders for the cutoff $\Lambda = 450$ MeV.¹³ As expected, a clear order-by-order improvement in the description of the scattering data can be seen. Table 1 also gives the number of

¹¹ Strictly speaking, our database differs from the one of Ref. [18] by the omission of the data set from Ref. [133], see Ref. [17] for more details.

¹² In our paper [17] also the cutoff $\Lambda = 350$ MeV was considered. Given the sizable finite- Λ artifacts for this very soft cutoff choice, we do not consider this case in the following discussion.

¹³ We have corrected the last figures in the values for χ^2/datum for np data in the E_{lab} bins of 0–100 MeV and 0–200 MeV at N³LO and N⁴LO⁺ quoted in Table 3 of Ref. [17].

adjustable parameters at each order which also includes isospin-breaking LECs contributing to the 1S_0 partial wave. It should be noted that no new contact interactions are added when going from NLO to N²LO and that the observed improvement of the χ^2/datum values is entirely due to the N²LO contributions to the parameter-free TPEP. A similar situation occurs when going from N³LO to N⁴LO, although here we also allow for additional isospin-breaking of the C_{1S0} contact LEC splitting it into two independently adjustable parameters for the neutron-proton and proton-proton/neutron-neutron systems. These improvements demonstrate both the importance of the chiral TPEP in the nuclear force and the predictive power of chiral perturbation theory, which allows to use LECs extracted in one process for making parameter-free predictions for (parts of) another.

Starting from N³LO, a satisfactory description of the neutron-proton data in the energy range of $E_{\text{lab}} = 0 - 200$ MeV and the proton-proton data for $E_{\text{lab}} = 0 - 100$ MeV is achieved. Although the N⁴LO potential improves on this, especially at intermediate and higher energies, it does not achieve a $\chi^2/\text{datum} \sim 1$ description of the proton-proton data for $E_{\text{lab}} \geq 100$ MeV. In the intermediate region, this value is significantly affected by the already mentioned high-precision data which requires an accurate description of F-waves. At N⁴LO the differential cross section data set of Ref. [134] at $E_{\text{lab}} = 144.1$ MeV, although well described within the Bayesian truncation errors, yields a χ^2/datum value of 27.88.

The situation is much improved once we switch to the N⁴LO⁺ potential and short-range interactions in F-waves are added. The description of scattering data at higher energies is generally improved and also the high-precision proton-proton data at intermediate energies is accurately reproduced leading to a $\chi^2/\text{datum} \sim 1$ description of the complete scattering database. Throughout the orders LO – N⁴LO the χ^2/datum value for proton-proton scattering up to 200 or 300 MeV has been larger than the one for neutron-proton scattering. This is plausible as proton-proton data is in general more precise than neutron-proton data and because only isovector partial waves contribute to it and hence only roughly half of the total number of parameters. However, at N⁴LO⁺, the reproduction of proton-proton data becomes very accurate while the slightly larger χ^2/datum values for the neutron-proton data as compared to proton-proton data reflects the larger statistical fluctuations among different data sets. This can be seen as an indication for reaching the threshold where the model accuracy approaches the precision of the data. In fact, the description of the scattering data at N⁴LO⁺ and $\Lambda = 450$ MeV is comparable to or exceeds that of the high-quality semi-phenomenological potentials such as CD-Bonn [137], Nijm I, II [138] and Reid93 [138]. Thanks to the parameter-free effects of the TPEP this is achieved with only 27 adjustable short-range parameters instead of the $\sim 40 - 50$ parameters used in those potentials.

Indeed, due to the excellent description of the data, the obtained results at $\Lambda = 450$ MeV qualify to be considered a partial-wave analysis. In Figs. 4 and 5 we show the obtained N⁴LO⁺ neutron-proton and proton-proton phase shifts for $\Lambda = 450$ MeV, respectively. We compare them to the 2013 Granada analysis [18] and in the case of neutron-proton scattering also to the corresponding 2008 analysis by Gross and Stadler [139]. Furthermore, we also show the predictions from the N⁴LO⁺ potential of Ref. [106] at the intermediate cutoff $\Lambda = 500$ MeV.

In general, there is good agreement between the shown N⁴LO⁺ phase shifts and the results obtained by the considered phase shift analyses. This is especially true for the case of proton-proton phase shifts which are more strongly constrained by the precise experimental data. Some discrepancies among the different results remain e.g. around the maximum of the 3P_0 phase shift where the N⁴LO⁺ prediction for the proton-proton phase is slightly larger than the ones of the Nijmegen and Granada PWAs, resulting in a $\sim 3\sigma$ deviation from the former at $E_{\text{lab}} = 50$ MeV. On the other hand, our neutron-proton phase shifts fall in between the results of the two PWAs. The study of isospin-breaking effects in P-waves beyond the ones




Figure 4. Neutron-proton phase shifts with respect to Riccati-Bessel functions in comparison with the Nijmegen [129] (solid dots), the Granada [140] (blue open triangles) and Gross-Stadler [139] (green open squares) PWA. Red solid lines and peach-colored bands denote the central results and 68% DoB truncation errors at the order $N^4\text{LO}^+$ for the cutoff $\Lambda = 450$ MeV. Black dashed lines denote the result of the nonlocal $N^4\text{LO}^+$ potential of Ref. [106] for the cutoff $\Lambda = 500$ MeV. The shown uncertainties of the Nijmegen PWA correspond to systematic errors defined in Eq. (32) of Ref. [21].

included in the two PWAs and the current version of the semilocal momentum-space regularized (SMS) interaction of Ref. [17] is expected to shed some light on this issue. We can also compare our results at $N^4\text{LO}^+$ to the ones of Ref. [106]. Similar to the comparison with the PWAs, agreement with proton-proton phases is better than with neutron-proton ones. There are, however, notable differences in the 3P_0 , 3P_2 and 3D_2 waves starting at low or intermediate energies. At higher energies around $E_{\text{lab}} = 250 - 300$ MeV, a change in curvature of the phase shift as a function of energy is visible e.g. in the 1P_1 and 3P_1 waves, which is presumably caused by the regulator employed in Ref. [106]. The effects of regulator artifacts can be observed particularly well in the 1G_4 , 3H_4 and ϵ_4 phase shifts and mixing angle shown in Fig. 5 since they do not involve any adjustable short-range parameters at $N^4\text{LO}^+$ but are solely determined by the long-range pion-exchange potential. Here, the local regulator of Eq. (22) leads to an undistorted reproduction of the peripheral phase shifts.

Selected proton-proton scattering observables and their estimated truncation error at various orders are shown in Fig. 6 for E_{lab} around ~ 143 MeV. In particular, we show our predictions for the differential cross section at $E_{\text{lab}} = 144.1$ MeV and compare them with two high-precision data sets, most notably




Figure 5. Proton-proton phase shifts with respect to Coulomb wave functions in comparison with the Nijmegen [129] (solid dots) and the Granada [140] (blue open triangles) PWA. Red solid lines and peach-colored bands denote the central result and 68% DoB truncation errors at the order $N^4\text{LO}^+$ for the cutoff $\Lambda = 450$ MeV. Black dashed lines denote the result of the nonlocal $N^4\text{LO}^+$ potential of Ref. [106] for the cutoff $\Lambda = 500$ MeV. The shown uncertainties of the Nijmegen PWA correspond to systematic errors defined in Eq. (32) of Ref. [21].

the one of Ref. [134], which motivated the introduction of the $N^4\text{LO}^+$ potential as discussed above. The data are well described within the given truncation error for all considered orders, but the $N^4\text{LO}^+$ clearly allows for a proper quantitative description. Likewise, the reproduction of the spin observables in Fig. 6 is excellent already at $N^3\text{LO}$ with a good convergence pattern. Notice however, that the error bands at lower orders for D (A) at the minimum (maximum) around $\Theta_{\text{CM}} = 150^\circ$ do not overlap with the ones for $N^{\geq 3}\text{LO}$ and are indeed underestimating the uncertainty. Here we find that the value of the observable in that particular angular region is notably shifted starting at $N^3\text{LO}$ while lower-order corrections are small, such that the overall scale in Eq. (41) is still underestimated. Using a more sophisticated Bayesian approach of Ref. [125] would likely allow for a more reliable estimation of the truncation errors at $\text{LO}-N^2\text{LO}$ in these particular cases.

There are various a posteriori checks that can be performed to test the self-consistency and quality of the fit. First, the values of the LECs have to be of natural size assuming the cutoff is kept below the hard scale. The expected sizes of the spectroscopic contact LECs can be estimated to be [21]

$$|\tilde{C}_i| \sim \frac{4\pi}{F_\pi^2}, \quad |C_i| \sim \frac{4\pi}{F_\pi^2 \Lambda_b^2}, \quad |D_i| \sim \frac{4\pi}{F_\pi^2 \Lambda_b^4}, \quad |E_i| \sim \frac{4\pi}{F_\pi^2 \Lambda_b^6}, \quad (46)$$

where the LECs $\tilde{C}_i$, C_i , D_i and E_i start to contribute at order Q^0 , Q^2 , Q^4 and Q^6 , respectively. Λ_b is the breakdown scale of the chiral expansion discussed in Sec. 5. Furthermore, the factor of 4π emerges from the angular integration of the partial-wave decomposition and has been included in the definition of




Figure 6. Selected proton-proton observables around $E_{\text{lab}} = 143$ MeV: Differential cross section $d\sigma/d\Omega$ at $E_{\text{lab}} = 144.1$ MeV with experimental data taken from Ref. [134] and Ref. [141]. The data sets have been corrected for their estimated norms of 0.988 and 1.001, respectively. Analyzing power P at $E_{\text{lab}} = 142$ MeV with experimental data taken from Ref. [142]. The data have been floated and multiplied by an estimated norm of 0.942. Depolarization D , rotation parameter A , polarization transfer coefficient D_t and spin-correlation parameter C_{kp} at $E_{\text{lab}} = 143$ MeV with experimental data taken from Refs. [143] and [144]. The light- (dark-) shaded green, blue and red bands depict the 68% (95%) DoB truncation errors at $N^2\text{LO}$, $N^3\text{LO}$ and $N^4\text{LO}^+$, respectively. Open circles show the predictions of the Nijmegen partial-wave analysis [129].

the spectroscopic LECs. If we now divide the contact LECs obtained in the fit by their expected sizes in Eq. (46), we consequently should obtain values of unit magnitude. Fig. 7 shows the absolute values of the LECs at $N^4\text{LO}^+$ in these natural units for all considered values of the cutoff Λ using $\Lambda_b = 650$ MeV. As can be seen, all LECs are indeed of natural size with D_{1S0} and D_{3S1} being among the largest in magnitude. This is especially true for the softest cutoff $\Lambda = 400$ MeV, for which also most of the other- Q^4 LECs turn out to be slightly larger than at higher values of the cutoff. This indicates that at $\Lambda = 400$ MeV and below, finite-cutoff artifacts start to increase, leading to a lower effective breakdown scale compared to the other considered cutoffs. Notice further that the values for the Q^6 LECs E_i included at $N^4\text{LO}^+$ turn out to be of a perfectly natural size. Therefore, even though we have emphasized their importance in describing some high-precision proton-proton data and achieving a $\chi^2/\text{datum} \sim 1$ description of the database, their actual contributions agree with the expectations from naive dimensional analysis (i.e. Weinberg) power counting, and there is no need to promote them to a lower order.

In addition to the absolute of the central values, Fig. 7 also shows the statistical uncertainties of the contact LECs as determined from the covariance matrix of the fit (expressed in their natural units). When going from $\tilde{C}_i$, C_i , D_i to E_i the statistical relative errors tend to increase. This is in accordance with the decreasing importance of higher-order contributions as predicted by power counting. One also notices that errors are smaller for LECs entering isovector partial waves, because these parameters are mainly




Figure 7. Absolute values of the contact interaction LECs in natural units at the order $N^4\text{LO}^+$ for all considered cutoffs. Error bars represent the statistical errors of the LECs.

constrained by the more precise proton-proton data. Since we perform a combined fit of neutron-proton and proton-proton data, the isovector partial waves are not only constrained by more precise data but also by more data in general compared to the isoscalar partial waves which have to be extracted from neutron-proton data alone. The covariance matrix also gives access to the correlations among the LECs. As to be expected, correlations mostly occur among LECs entering the same partial waves with the largest ones arising in the channels with the most parameters, namely in the 1S_0 and $^3S_1 - ^3D_1$ channels. Nevertheless, all LECs are well-constrained as can already be seen by looking at the errors in Fig. 7. We can further look at the largest eigenvalue of the covariance matrix of the natural LECs as a measure of how well-determined the parameters are. Throughout the considered range of the cutoff $\Lambda = 400 - 550$ MeV, the largest eigenvalue of the covariance matrix does not exceed 0.1 and is ~ 0.08 for $\Lambda = 450$ MeV.

From the point of view of data fitting, another check concerns the statistical assumptions underlying a χ^2 fit. One usually assumes that the residuals $r_i = (O_i^{\text{exp}} - O_i^{\text{th}})/\Delta O_i$ follow a normal distribution $\mathcal{N}(0, 1)$ with zero mean and unit standard deviation. Here O_i^{exp} and ΔO_i are the experimental value and its error of an observable and O_i^{th} is its calculated "theoretical" value. If the assumptions on the normally-distributed residuals can be verified, this confirms that the data are described sufficiently well by the theoretical model. An easy and often employed check is the value of χ^2 per degree of freedom. For the $N^4\text{LO}^+$ fit with $\Lambda = 450$ MeV we get $\chi^2 = 4708.65$ in the fitting range of $E_{\text{lab}} = 0 - 260$ MeV with the number of data $N_{\text{dat}} = 4616$ and the number of parameters $N_{\text{par}} = 27$. Consequently, we obtain $\chi^2/\nu = 1.026$ with $\nu = N_{\text{dat}} - N_{\text{par}}$. If the residuals are indeed normal-distributed then χ^2/ν should follow the χ^2 -distribution and yields $\chi^2/\nu = 1 \pm \sqrt{2/\nu} = 1 \pm 0.021$ as the 68% confidence interval.

We can go one step beyond this simple check and plot the quantiles of the empirical distribution of residuals r_i that we obtain against the quantiles of the assumed normal distribution $\mathcal{N}(0, 1)$. If they are the same, they should lie on the diagonal line $x = y$. In order to statistically quantify deviations from the diagonal, confidence bands have been derived with one of the most recent and most sensitive being the ones by Aldor-Noiman et al. [145]. This graphical test for normal-distributed residuals has been first applied to the analysis of nucleon-nucleon scattering by Navarro Pérez et al. [146] and named the "tail-sensitive test" in that publication. Fig. 8 shows a rotated quantile-quantile plot for the $N^4\text{LO}^+$ residuals at $\Lambda = 450$ MeV where the theoretical quantiles have been subtracted from the empirical ones on the y -axis, turning the diagonal line into a horizontal one. As evident from the figure, the empirical distribution of residuals




Figure 8. Rotated Quantile-Quantile plot of the empirical quantiles at $N^4\text{LO}^+$ and $\Lambda = 450$ MeV versus the quantiles of the normal distribution $\mathcal{N}(0, 1)$. Dotted (solid) red bands denote the 68% (95%) confidence bands of the tail-sensitive test by Aldor-Noiman et al. [145].

lies within the 68% confidence region of the tail-sensitive test signaling that the residuals are indeed normal-distributed. The quantile-quantile plot for the other values of the cutoff turn out to be overall similar, but perform slightly worse. For $\Lambda = 500$ MeV and $\Lambda = 550$ MeV the quantiles that are already close to the edge of the 68% confidence region in Fig. 8 cross these limits but still stay well within the 95% confidence region. The increased cutoff-artifacts at $\Lambda = 400$ MeV manifest themselves in a stronger deviation from normality as the plotted quantiles also cross the 95% confidence limits at the spike at $Q_{\text{th}} = 2$ in Fig. 8.

We now turn to the extended error analysis for observable predictions. While the truncation of the chiral expansion is clearly the dominant source of uncertainty at higher energies, other sources of uncertainty can become relevant at $N^4\text{LO}^+$. In particular we account for the following sources of uncertainty:

- *Statistical uncertainties of NN LECs:* As already mentioned, Fig. 7 shows the statistical errors of the contact LECs as determined from the covariance matrix of the fit. The uncertainties of the parameters can then be propagated from the covariance matrix to the observable of interest. While it is always possible to do this via a Monte Carlo sampling of the corresponding multivariate Gaussian probability distribution, it is computationally much more convenient to do a Taylor expansion of the desired observable with respect to the LECs and evaluate the moments of the LECs analytically. While a linear expansion is commonly employed, it has been argued in Ref. [104] that some observables require a second order expansion for an accurate reproduction of their uncertainties. In the case of large second-order contributions, the error bars become asymmetric and we usually give both the upper and lower error to accommodate for this possibility. Notice that in such a case, the probability density of the observable is not Gaussian and the quoted uncertainties do not necessarily correspond to a 68% degree-of-belief.
- *Statistical uncertainties of πN LECs:* In addition to the central values, the authors of Ref. [42] also give the covariance matrix as determined from πN scattering data. Propagation of these uncertainties to NN observables is, however, less straightforward, because the values of the NN contact interactions depend




Figure 9. Neutron-proton total cross section in the range of $E_{\text{lab}} = 0 - 300$ MeV. The plot on the left shows the results divided by the predictions of the Nijmegen PWA. The red line and peach-colored band show the central values and truncation errors (the 68% DoB interval) at the order $N^4\text{LO}^+$ and for $\Lambda = 450$ MeV. The experimental data are taken from Ref. [147] and have been corrected for their estimated norm of 0.999. The plot on the right shows the relative uncertainties as discussed in the text.

on the values of πN LECs. We thus resort to some Monte Carlo sampling of the multivariate Gaussian probability distribution of the πN LECs given by their central values and their covariance matrix. For each of the sampled sets of LECs, we refit the NN contact LECs before calculating any observables. The uncertainty of a given observable can then be estimated in a standard way from the variance of the results calculated with different πN LEC sets. Due to the need to refit the contact interactions for each sampled set of πN LECs and the computational overhead related to it, we have restricted the total number of such sets to 50. Although this is a quite low statistics for a Monte Carlo approach, it should give an idea of the order of magnitude of the uncertainty. It indeed turns out that the uncertainty related to the statistical error of the πN LECs is small compared to the other sources of uncertainty. However, the aforementioned approach does not probe the systematic errors in the determination of the πN LECs emerging from the truncation of the chiral expansion and thus does not represent the full uncertainty related to these LECs.

- *Uncertainty due to the choice of the maximum fit energy:* The extracted values of the contact LECs also depend on details of the fitting protocol. In particular, we probe the impact of the choice for the maximum laboratory energy $E_{\text{max}} = 260$ MeV up to which scattering data is included in the $N^4\text{LO}^+$ fit. This is achieved by performing additional fits with $E_{\text{max}} = 220$ MeV and $E_{\text{max}} = 300$ MeV and determining the maximum deviation of the observables from the $E_{\text{max}} = 260$ MeV predictions. Unlike the aforementioned uncertainties, the error estimated via this simple procedure does not reflect any particular degree-of-belief.

As an example, Fig. 9 shows the neutron-proton total cross section and the corresponding uncertainties in the energy range $E_{\text{lab}} = 0 - 300$ MeV. The plot on the left in Fig. 9 shows the ratio of our predictions using the $N^4\text{LO}^+$ potential at $\Lambda = 450$ MeV and the result of the Nijmegen partial-wave analysis [129]. In the right panel, the different relative errors stemming from the various sources discussed above are shown. For the case of the statistical errors of the NN contact interactions, second order effects and resulting

Table 2. Deuteron binding energy B_d , asymptotic S-state normalization A_S , asymptotic D/S-state ratio η , radius r_d , quadrupole moment Q and D-state probability P_D as predicted by various high-quality potentials. The binding energy has been calculated with the nonrelativistic energy-momentum relation for the potentials of Ref. [106] and with the relativistic relation for the SMS potential of Ref. [17] and the CD Bonn potential Ref. [137].

	Granada [140]	CD Bonn [137]	EMN $N^4\text{LO}^+$ [106] $\Lambda = 500 \text{ MeV}$	SMS $N^4\text{LO}^+$ [17] $\Lambda = 450 \text{ MeV}$	Empirical
B_d (MeV)	2.2246*	2.2246*	2.2246*	2.2246*	2.224575(9) [135]
A_S ($\text{fm}^{-1/2}$)	0.8829	0.8846	0.8852	0.8847	0.8846(8) [148]
η	0.0249	0.0256	0.0258	0.0255	0.0256(4) [149]
r_d (fm)	1.965	1.966	1.973	1.966	1.97535(85) [†] [150]
Q (fm^2)	0.268	0.270	0.273	0.270	0.2859(3) [151]
P_D (%)	5.62	4.85	4.10	4.59	—

*The deuteron binding energy has been taken as input in the fit.

[†]This value corresponds to the so-called deuteron structure radius, which is defined as a square root of the difference of the deuteron, proton and neutron mean square charge radii.

asymmetries in the error bands turn out to be small for the total cross section, and the plotted uncertainty corresponds to the average of upper and lower statistical errors. As expected, the dominant contribution to the uncertainty at higher energies ($E_{\text{lab}} > 100 \text{ MeV}$) arises from the truncation of the chiral expansion. At lower energies, however, other sources of uncertainty become relevant and indeed both the statistical errors of the NN contact LECs and the uncertainty due to the maximum fitting energy are larger than the truncation error in the range of $E_{\text{lab}} = 30 - 100 \text{ MeV}$. When quantitatively comparing the different errors, one has to keep in mind that the uncertainty due to the maximum fitting energy does not correspond to a particular degree-of-belief. The uncertainty arising from the statistical errors of the πN LECs is found to be significantly smaller throughout the whole considered energy range and is negligible for the total cross section. Finally, we would like to comment on the origin of the existing kinks in the right-hand-side plot of Fig. 9. In particular, the kink in the E_{max} -error at around 200 MeV arises because of the maximum operation. Below 200 MeV, the error is dominated by the deviation of the $E_{\text{max}} = 220 \text{ MeV}$ fit while it is given by the deviation of the $E_{\text{max}} = 300 \text{ MeV}$ fit above 200 MeV. The second kink present in the truncation error, on the other hand, is caused by the transition of Q from $M_\pi^{\text{eff}}/\Lambda_b$ to p/Λ_b .

Tab. 2 shows the deuteron properties as predicted by various high-quality potentials. Clearly, the error analysis can also be applied to the bound state properties of Tab. 2. However, the obtained uncertainties are only meaningful for a complete calculation of an unconstrained observable. This excludes the deuteron binding energy B_d (as it is a fitted quantity), the quadrupole moment Q and deuteron radius r_d (as meson exchange currents and relativistic corrections are not taken into account) as well as the D-state probability P_D (which is not observable). On the other hand, we can perform the uncertainty quantification for the asymptotic S-state normalization A_S and the asymptotic D/S-state ratio η for which we obtain at $N^4\text{LO}^+$ and for $\Lambda = 450 \text{ MeV}$ the values of $A_S = 0.8847_{(-3)}^{(+3)}(5)(0)(1)$ and $\eta = 0.02553_{(-9)}^{(+11)}(4)(3)(8)$, respectively. Here the first, second, third and fourth error refer to the NN statistical, truncation, πN statistical and E_{max} uncertainty, respectively. Notice that the quoted truncation errors estimated using the Bayesian model of section 5 are fairly similar to the ones given in Ref. [17], which were obtained using the EKM method. On the other hand, the πN statistical uncertainties are much smaller than the corresponding

errors quoted in Ref. [17], where an attempt was made to also include systematic effects by using the values of these LECs determined in the physical region of πN scattering.

Finally, let us discuss the treatment of isospin-breaking effects in the two-nucleon interaction. Like all modern high-precision potentials, the SMS interactions include isospin-breaking in the OPEP due to the different physical pion masses $M_{\pi^\pm}$ and M_{π^0} and charge dependence of the short-range potential in the 1S_0 partial wave. These are the dominant and well-understood isospin-breaking effects necessary to arrive at e.g. a correct description of the charge-dependence of the 1S_0 scattering length. For the calculation of scattering observables in the two-nucleon system, the isospin-breaking due to long-range electromagnetic interactions is taken into account as discussed at the beginning of this section. This treatment of strong and electromagnetic isospin-breaking effects is identical to the Nijmegen PWA [129]. On the other hand, chiral EFT allows for a systematic inclusion of isospin-breaking effects beyond the ones previously considered. In fact, expressions for the leading isospin-breaking TPEP [152, 153], the subleading isospin-breaking TPEP [154] and irreducible $\pi\gamma$ exchange [155], which are (mostly) parameter-free in the two-nucleon system, have been available for some time. The long-standing question regarding the charge-dependence of the πNN coupling constant also re-emerges in a systematic treatment of isospin-breaking effects in the framework of chiral EFT. While the Nijmegen group did not find evidence for charge-dependence, the issue does not seem to be settled, see Ref. [156] for a recent determination. Last but not least, charge-dependence in the short-range potential entering P-waves should also be taken into account starting from $N^4\text{LO}$ [154].

6.2 Three-nucleon scattering

As discussed in the previous subsection, the $N^4\text{LO}^+$ SMS potentials of Ref. [17] lead to excellent and in fact a nearly perfect description of np and pp scattering data below pion production threshold. Moreover, an order-by-order comparison of the results for various observables along with the Bayesian error analysis indicate a generally good convergence of the chiral expansion in the NN sector. On the other hand, a description of nucleon-deuteron elastic and breakup scattering data at a comparable level of accuracy is not available yet. Extensive calculations performed in the last decades using high-precision phenomenological NN potentials and 3NF models in the framework of the Faddeev equations [157] and using other *ab initio* methods [158] have revealed the following picture (see Ref. [19] and references therein):

- Calculations based on high-precision NN potentials alone (including the $N^4\text{LO}^+$ ones of Ref. [17]) tend to underestimate the ^3H and ^3He binding energy by ~ 0.5 MeV and generally lead to similar predictions in Nd scattering observables.
- At low energies, the resulting description of Nd data appears to be rather good apart from a few exceptions such as the underprediction of the nucleon analyzing power A_y , known as the A_y puzzle [159], and the discrepancy for the cross section for the symmetric space star deuteron breakup configuration [160]. 3NF effects in this energy range are found to be small in agreement with qualitative arguments based on the chiral power counting as explained below.
- Starting from $E_N \sim 50$ MeV, discrepancies between theory and experimental data set in and become large at $E_N \sim 200$ MeV and above. Except for the cross section, the inclusion of the phenomenological 3NFs like the Tucson-Melbourne (TM99) [161] and Urbana-IX [162] models does not globally reduce the discrepancies between theory and data [19]. Relativistic effects have also been studied, see Ref. [163] and references therein, and found to be small at energies below the pion production threshold.

Assuming that the discrepancies between theory and experimental data in the 3N system are to be resolved by 3NFs, these findings demonstrate that the currently available phenomenological models do not provide an appropriate description of the 3NF. This should not come as a surprise given the enormously rich and complex spin-isospin-momentum structure of a most general 3NF [85, 164, 165, 166]. Here, chiral EFT offers a decisive advantage over more phenomenological approaches by predicting the long-range part of the 3NF in a model-independent way, establishing a clear importance hierarchy of short-range terms and providing a solid theoretical framework for maintaining consistency between two- and three-nucleon forces and ensuring scheme independence of the calculated observables.

As already mentioned in section 3, three-body contributions to the nuclear Hamiltonian first appear at $N^2\text{LO}$ in the chiral expansion and are, therefore, suppressed by Q^3 relative to the dominant pairwise NN interaction. It is instructive to estimate the expected magnitude of 3NF effects for various observables solely on the basis of the chiral power counting (i.e. using NDA). For ^3H and ^4He , one can use the typical expectation values of the NN potential energy of $\langle V_{\text{NN}} \rangle_{^3\text{H}} \sim 50 \text{ MeV}$ and $\langle V_{\text{NN}} \rangle_{^4\text{He}} \sim 100 \text{ MeV}$ [74], along with the estimation of the expansion parameter $Q \sim M_\pi^{\text{eff}}/\Lambda_b$ with $M_\pi^{\text{eff}} = 200 \text{ MeV}$ and $\Lambda_b = 650 \text{ MeV}$, in order to estimate the expected 3NF contributions to the binding energy to be $\langle V_{3\text{N}} \rangle_{^3\text{H}} \sim Q^3 \langle V_{\text{NN}} \rangle_{^3\text{H}} \sim 1.5 \text{ MeV}$ and $\langle V_{3\text{N}} \rangle_{^4\text{He}} \sim Q^3 \langle V_{\text{NN}} \rangle_{^4\text{He}} \sim 3 \text{ MeV}$. These qualitative estimations agree well with the actual underprediction of the ^3H and ^4He by the NN interactions alone which, using the AV18 [167], CD Bonn [137], $N^2\text{LO}$ [103] and Idaho $N^3\text{LO}$ [102] potentials as representative examples, amounts to $0.5 \dots 0.9 \text{ MeV}$ and $2.1 \dots 4.1 \text{ MeV}$, respectively. The shallow nature of few-nucleon bound states indicates that there are large cancellations between the kinetic and potential energies. Because of this fine tuning, 3NF contributions to the binding energies are enhanced beyond the naive estimation of $Q^3 \sim 3\%$ and actually reach $10 \dots 15\%$. On the other hand, there is generally no reason to expect a similar enhancement for Nd scattering observables at low energy except for some fine-tuned polarization observables such as A_y . It is well known that tiny changes of the NN interaction in the triplet P -waves amount to large relative changes in the Nd A_y [159]. On the other hand, starting from $E_N \sim 60 \text{ MeV}$, the expansion parameter Q in Eq. (37) is dominated by the momentum scale $p = \sqrt{2/3mE_N}$ [75]. At e.g. the energies of $E_N \sim 100 \text{ MeV}$ and $E_N \sim 200 \text{ MeV}$, the expansion parameter becomes $Q \sim 0.40$ and $Q \sim 0.55$, and the relative contributions of the 3NF to a generic scattering observable are expected to increase to $\sim 6\%$ and $\sim 16\%$, respectively. Clearly, these simplistic back-of-envelope estimations only yield qualitative insights into the role of the 3NF. Nevertheless, they agree remarkably well with the observed trend of discrepancies between theoretical predictions based solely on the NN interactions and experimental data, which tend to increase with energy. For further examples and a more quantitative analysis along this line of Nd scattering, selected properties of light and medium-mass nuclei and the equation of state of nuclear matter see Refs. [74, 122, 119, 168]. We further emphasize that it is not entirely clear how to estimate the relevant momentum scale, that determines the expansion parameter in heavy nuclei, and how to quantify truncation errors for excited states, see Ref. [122] for an extended discussion.

As discussed in section 3 and visualized in Fig. 2, the leading contributions to the 3NF at $N^2\text{LO}$ emerge from the two-pion exchange, one-pion-exchange-contact and purely contact tree-level diagrams, leading to the well-known expressions [88, 36]

$$\begin{aligned}
 V_{3\text{N}} = & \frac{g_A^2}{8F_\pi^4} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_3 \cdot \vec{q}_3}{(\vec{q}_1^2 + M_\pi^2)(\vec{q}_3^2 + M_\pi^2)} \left[\vec{\tau}_1 \cdot \vec{\tau}_3 (-4c_1 M_\pi^2 + 2c_3 \vec{q}_1 \cdot \vec{q}_3) + c_4 \vec{\tau}_1 \times \vec{\tau}_3 \cdot \vec{\tau}_2 \vec{q}_1 \times \vec{q}_3 \cdot \vec{\sigma}_2 \right] \\
 & - \frac{g_A D}{8F_\pi^2} \frac{\vec{\sigma}_3 \cdot \vec{q}_3}{\vec{q}_3^2 + M_\pi^2} \vec{\tau}_1 \cdot \vec{\tau}_3 \vec{\sigma}_1 \cdot \vec{q}_3 + \frac{1}{2} E \vec{\tau}_1 \cdot \vec{\tau}_2 + 5 \text{ permutations} , \quad (47)
 \end{aligned}$$

where $\vec{q}_i = \vec{p}_i' - \vec{p}_i$ with $\vec{p}_i'$ and $\vec{p}_i$ being the final and initial momenta of the nucleon i . The LECs D and E are usually expressed in terms of the corresponding dimensionless coefficients c_D and c_E via $D = c_D/(F_\pi^2 \Lambda_\chi)$ and $E = c_E/(F_\pi^4 \Lambda_\chi)$ [36]. In Refs. [8] and [75], semilocal coordinate- and momentum-space regularized 3NF expressions in combination with the corresponding chiral NN potentials from Refs. [21, 22] and [17], respectively, were employed by the LENPIC Collaboration to analyze Nd scattering observables at N²LO. The numerical implementation of the 3NF in the Faddeev equations is carried out in the partial wave basis. Partial wave decomposition (PWD) of a general 3NF can be carried out numerically using the machinery developed in Ref. [169] by performing five-dimensional angular integrations. Given the required number of partial waves and grid points for the four Jacobi momenta to reach converged results for Nd scattering observables, such a numerical PWD requires substantial computational resources. In Ref. [170], a more efficient approach was introduced, that exploits the local nature of the bulk of the 3NF.

To make predictions for few-nucleon observables, one first needs to determine the LECs c_D and c_E entering the 3NF. A broad range of few- and many-body observables including the binding energies and radii of ^3H , ^4He and heavier nuclei, nucleon-deuteron doublet scattering length 2a , n - α scattering, triton β -decay and the saturation properties of nuclear matter have been proposed and employed in the past to determine these two LECs [36, 171, 172, 9, 173, 105]. A reliable determination of c_D , c_E is complicated by the existence of strong correlations between some of the low-energy observables, see e.g. [174], which originate from the large S-wave scattering lengths in the NN system. Furthermore, going beyond the 3N system may require, depending on the observable and the chiral order, the inclusion of 4NF and exchange current contributions. In Ref. [8] we, therefore, restricted ourselves to 3N observables in the determination of c_D , c_E . Specifically, we employed the ^3H binding energy of $B_{^3\text{H}} = 8.482$ MeV to fix the LECs c_E for a given value of c_D . The remaining LEC c_D was determined by considering a number of observables including $^2a = 0.645 \pm 0.008$ fm [136], nd total cross section data from [175] and precisely measured pd differential cross section in the minimum region at $E_N = 70$ MeV [176], 108 MeV [177] and 135 MeV [176]. In the left panel of Fig. 10, we show the extracted values of c_D for the SCS interactions with the cutoff $R = 0.9$ fm. It is reassuring to see that the considered 3N observables lead to consistent values of c_D . In addition, these results show that the strongest constraint on c_D , given the experimental *and the estimated truncation uncertainty*, is imposed by the pd differential cross section data at $E_N = 70$ MeV from Ref. [176] as visualized in the right panel of Fig. 10. We also found no correlations between this observable and the ^3H binding energy. In particular, the resulting value of the LEC c_D is largely determined by the differential cross section and almost insensitive to a variation of the triton binding energy.

In Ref. [75], we have analyzed Nd scattering observables using the most recent SMS NN potentials from Ref. [17] in combination with the N²LO 3NF regularized in the same way. Motivated by the experience with the SCS interactions [8], the LECs c_D and c_E were determined from the ^3H binding energy and the pd cross section minimum at $E_N = 70$ MeV. In Fig. 11 we show, as a representative example, our N²LO predictions for selected Nd scattering observables at $E_N = 135$ MeV, along with the experimental data and calculations based on the CD Bonn NN potential with and without the TM99 3NF model. As an important internal consistency check of the calculations, we have verified that the predictions obtained using different cutoff values are consistent with each other (within errors), see Fig. 5 of Ref. [75].

It is reassuring to see that the experimental data are generally well described by the theory. On the other hand, while accurate, our predictions at N²LO have obviously rather low precision at this energy. In fact, the N²LO truncation errors are comparable with or even larger than the observed deviations between experimental data and calculations based on phenomenological high-precision NN and 3NF models, see the dotted and dashed-dotted lines in Fig. 11. Based on the experience in the NN sector as discussed in




Figure 10. Left: Determination of the LEC c_D at N^2 LO from selected Nd scattering observables. The smaller (blue) error bars correspond to the experimental uncertainty while the larger (orange) error bars also take into account the truncation error at N^2 LO estimated using the EKM approach of Ref. [21]. The green (violet) bands show standard error intervals of c_D resulting from a combined least squares single-parameter fit to all observables (to observables up to $E_N = 108$ MeV) using the orange error bars. Right: Nd cross section in the minimum region ($\theta = 130^\circ$) at $E_N = 70$ MeV as function of the LEC c_D . For each c_D value, the LEC c_E is adjusted to the ^3H binding energy. Dotted lines show the statistical uncertainty of the experimental data from Ref. [176], while the yellow band also takes into account the quoted systematic uncertainty of 2%. All results are obtained using the N^2 LO SCS NN potential from Ref. [21] in combination with the N^2 LO SCS 3NF for the coordinate-space cutoff $R = 0.9$ fm.

section 6.1, it is conceivable that a high-precision description of Nd scattering data will require the chiral expansion of the 3NF to be pushed to (at least) N^4 LO. At the energy of $E_N = 135$ MeV, the uncertainty bands at N^4 LO are expected to become 4-5 times more narrow as compared with the N^2 LO ones shown in Fig. 11.

So where do we stand in terms of efforts to include 3NF corrections beyond N^2 LO? As explained in section 4.2, the main obstacle for the inclusion of higher order contributions to the 3NF is the lack of their *consistently regularized* expressions. Starting from N^3 LO, it is not sufficient anymore to naively regularize the available expressions for the 3NF from Refs. [46, 47, 40, 85] derived using DR, since such an approach violates constraints imposed by the chiral symmetry. Rather, the N^3 LO and N^4 LO corrections to the 3NF need to be re-derived using the consistent finite-cutoff regularization approach. Work along these lines is in progress. Another challenge, that will have to be addressed at N^4 LO, is the determination of the LECs appearing in the 3NF at this order. While the N^3 LO contributions do not involve free parameters, the short-range part of the 3NF at N^4 LO depends on 10 unknown LECs [37], from which 9 contribute to the isospin-1/2 channel and thus can, in principle, be determined in Nd scattering. Furthermore, the yet-to-be-derived one-pion-exchange-contact contributions to the 3NF at N^4 LO will also involve unknown LECs. Given the still rather significant computational cost of solving the Faddeev equations in the 3N continuum, the complicated treatment of the Coulomb interaction [178] and the lack of partial wave analyses in the 3N sector, the determination of these LECs from 3N scattering data will certainly be a challenging task.




Figure 11. Predictions for the differential cross section, nucleon and deuteron analyzing powers A_y^n and A_y^d , deuteron tensor analyzing powers A_{yy} , A_{xz} , A_{xx} , polarization transfer coefficients K_{xx}^y , K_y^y , K_{yy}^y , K_{xz}^y , $K_{xx}^y - K_{yy}^y$ and the induced polarization P^y in elastic Nd scattering at laboratory energy of $E_N = 135$ MeV at NLO (yellow bands) and N^2 LO (green bands). The light- (dark-) shaded bands indicate 95% (68%) DoB intervals using the Bayesian model $\bar{C}_{0.5-10}^{650}$ introduced in section 5. Open circles are proton-deuteron data from Ref. [176]. The dotted (dashed-dotted) lines show the results based on the CD Bonn NN potential [137] (CD Bonn NN potential in combination with the Tucson-Melbourne 3NF [161]). All results are obtained using the N^2 LO SMS NN potential from Ref. [17] in combination with the N^2 LO SMS 3NF for the momentum-space cutoff $\Lambda = 500$ MeV.

While a complete analysis of Nd scattering is currently not available beyond N^2 LO, it is instructive to explore the role of subleading short-range 3NF interactions. In Ref. [179], it was shown within a hybrid phenomenological approach that the 3N contact operators at N^4 LO can be tuned to reproduce the ^3H binding energy, nd scattering lengths, cross section and polarization observables of pd scattering at 2 MeV center-of-mass energy. The resulting models were shown to lead to a satisfactory description of pd polarization observables below the deuteron breakup. On the other hand, 3NF effects are expected to be much more visible at intermediate and higher energies. In Ref. [75], we explored the impact of the short-range 3NF operators of the central and spin-orbit types proportional to the LECs E_1 and E_7 , respectively,

$$V_{3N} = E_1 \vec{q}_1^2 + iE_7 \vec{q}_1 \times (\vec{K}_1 - \vec{K}_2) \cdot (\vec{\sigma}_1 + \vec{\sigma}_2) + 5 \text{ permutations}, \quad (48)$$

where $\vec{K}_i = (\vec{p}_i' + \vec{p}_i)/2$. Parametrizing the dimension-full LECs E_1 , E_7 in terms of the corresponding dimensionless parameters via $E_i = c_{E_i}/(F_\pi^4 \Lambda_\chi^3)$ with $\Lambda_\chi = 700$ MeV, we studied the impact of these N^4 LO terms on selected Nd scattering observables for the fixed values of the LECs of $c_{E_i} = \pm 2$. Based on




Figure 12. Results for the differential cross section, nucleon analyzing powers A_y^n as well as deuteron tensor analyzing powers A_{xx} and A_{yy} in elastic nucleon-deuteron scattering at laboratory energy of $E_{\text{lab}}^N = 10$ MeV based on the SMS NN potentials of Ref. [17] at $N^4\text{LO}^+$ in combination with the SMS 3NF at $N^2\text{LO}$ using $\Lambda = 450$ MeV. Blue light- (dark-) shaded bands show the expected truncation uncertainty for a *complete* $N^3\text{LO}$ calculation and are obtained by multiplying the $N^2\text{LO}$ truncation error corresponding to 95% (68%) DoB intervals for the model $\bar{C}_{0.5-10}^{650}$ with the corresponding value of the expansion parameter Q . Short-dashed-dotted and long-dashed-dotted red lines show the impact of the $N^4\text{LO}$ central short-range 3NF $\propto c_{E_1}$ with $c_{E_1} = -2$ and $c_{E_1} = 2$, respectively. Similarly, short-dashed and long-dashed blue lines show the impact of the $N^4\text{LO}$ spin-orbit short-range 3NF $\propto c_{E_7}$ with $c_{E_7} = -2$ and $c_{E_7} = 2$, respectively. Open circles are neutron-deuteron data from Ref. [180] and proton-deuteron data from Ref. [181, 182, 183], corrected for the Coulomb effects, see Ref. [36] for details.

the experience in the NN sector and with the $N^2\text{LO}$ 3NF, we expect the actual values of these LECs to lie well within this range. The expectation values of various contributions to the 3NF in the triton state indicate that the employed values $c_{E_7} = \pm 2$ may already overestimate the expected natural range of this LEC.

In order to compute the contributions of the c_{E_i} -terms to 3N observables in a meaningful way, one needs to perform (implicit) renormalization as explained in section 2. This was achieved in Ref. [75] by simultaneously adjusting the values of the $N^2\text{LO}$ LECs c_D , c_E to the triton binding energy and the cross section minimum at $E_N = 70$ MeV for all considered values of the LECs c_{E_i} . The calculations have been performed using the $N^4\text{LO}^+$ SMS NN potential from Ref. [17] in combination with the SMS $N^2\text{LO}$ 3NF. In Fig. 12, we show the resulting predictions at the lowest considered energy of $E_N = 10$ MeV. The blue bands show the estimated truncation error at $N^3\text{LO}$, obtained by rescaling the $N^2\text{LO}$ Bayesian truncation uncertainty with the expansion parameter Q ¹⁴, and visualize the expected impact of $N^4\text{LO}$ terms. In agreement with the expectations, 3NF effects generally appear to be rather small at such low energies. This figure also provides a nice illustration of the fine tuned nature of the nucleon vector analyzing power A_y , which shows a strong sensitivity to small changes in the Hamiltonian. What has been traditionally referred to as the A_y -puzzle thus appears to be just a consequence of the fine-tuned nature of this observable, and the “puzzle” may be expected to be resolved by 3NF contributions beyond $N^2\text{LO}$. While the A_y is well known to be particularly sensitive to spin-orbit types of 3NFs [184] such as the one proportional to c_{E_7} , our results also show an unexpectedly strong sensitivity to the subleading central interaction of the c_{E_1} -type.

At higher energies, the effects of the considered $N^4\text{LO}$ 3NF terms become more significant as visualized in Fig. 13 for the case of selected spin-correlation parameters. More results for the cross section, vector and tensor analyzing powers and polarization transfer coefficients at $E_N = 135$ MeV can be found in Ref. [75]. It is comforting to see that the impact of the c_{E_i} -terms on Nd scattering observables is, in general,

¹⁴ We cannot estimate the $N^3\text{LO}$ truncation error using the Bayesian approach described in section 5 since no complete $N^3\text{LO}$ results are available for Nd scattering.




Figure 13. Same as Fig. 12 but for the deuteron-nucleon spin-correlation parameters $C_{z,x}$, $C_{y,y}$, $C_{z,z}$ and $C_{x,z}$ for Nd elastic scattering at $E_N = 135$ MeV (left panel) and $E_N = 200$ MeV (right panel). Open circles are proton-deuteron data from Ref. [185].

consistent with the estimated $N^3\text{LO}$ truncation errors. One should, however, keep in mind that the employed Bayesian approach may, under certain circumstances, become unreliable. This is, in particular, the case for observables that depend on a continuously varying parameter in the kinematical regions where the LO results and higher-order corrections change sign, see Ref. [75] for a detailed discussion. One such failure of the Bayesian model is shown in Fig. 13 for the spin-correlation coefficient $C_{x,z}$ at $E_N = 200$ MeV around $\theta = 120^\circ$. In such problematic cases, the approach proposed in Ref. [125] and based on Gaussian processes is expected to provide more reliable estimations of the truncation uncertainty.

6.3 Light nuclei

While no results for light nuclei using SMS chiral interactions are available yet, we briefly review here some recent highlights obtained by the LENPIC Collaboration using the SCS NN potentials of Refs. [21, 22] with and without the corresponding 3NFs at $N^2\text{LO}$. In Refs. [74, 122], we have calculated the ground state energies and selected properties of light and medium-mass nuclei up to ^{48}Ca using the SCS NN interactions at various chiral orders. Specifically, $A = 3, 4$ nuclei were analyzed in the framework of the Faddeev-Yakubovsky equations while light p-shell nuclei were calculated using the No-Core Configuration Interaction (NCCI) method [186, 187, 188] and employing Similarity Renormalization Group (SRG) transformed interactions [189, 190, 191, 192] to improve the convergence. The results for $^{16,24}\text{O}$ and $^{40,48}\text{Ca}$ were obtained within the coupled cluster and in-medium SRG group frameworks, see Refs. [193, 194, 195, 12, 196] and references therein. A qualitatively similar convergence pattern was observed in all considered cases, namely a significant overbinding at LO, results close to the experimental values at NLO and $N^2\text{LO}$ and underbinding at $N^3\text{LO}$ and $N^4\text{LO}$. Notice that the strongly repulsive nature of the $N^3\text{LO}$ contributions to the SCS NN interactions of [21, 22] was shown to be caused by the employed unnaturally large values of the redundant short-range operators [17]. The SMS interactions of Ref. [17] utilize a soft choice for these contact terms, which leads to more perturbative interactions at and beyond $N^3\text{LO}$. No large gap between the $N^2\text{LO}$ and $N^3\text{LO}$ results for the ground state energies is, therefore, expected for the new SMS NN interactions. The calculated charge radii of the considered medium-mass




Figure 14. Calculated ground state energies in MeV using chiral SCS NN interactions from Ref. [8] in combination with the SCS 3NF at $R = 1.0$ fm (open and solid dots) in comparison with experimental values (red levels). For each nucleus the NLO and N²LO results are the left and right symbols and bars, respectively. The open blue symbols correspond to incomplete calculations at N²LO using NN-only interactions. Blue and green error bars indicate the NCCI extrapolation uncertainty and, where applicable, an estimate of the SRG dependence. The shaded bars indicate the truncation error at each chiral order corresponding to 68% DoB intervals using the Bayesian model $\bar{C}_{0.5-10}^{650}$ with the expansion parameter $Q = M_{\pi}^{\text{eff}}/\Lambda_b$.

nuclei were found to show a systematic improvement with the chiral order, but remain underestimated using the NN interaction at the highest available order N⁴LO⁺.

In Ref. [8], a *complete* N²LO analysis of p-shell nuclei was presented by the LENPIC Collaboration using the SCS NN and 3N interactions. In Fig. 14, we show the NLO and N²LO results from that paper for nuclei up to $A = 16$. We emphasize that since the Hamiltonian has been completely determined in the NN and 3N system as described in sections 6.1 and 6.2, the ground-state energies shown in that figure are parameter-free predictions. In Fig. 14, we have updated the corresponding figure from Ref. [8] by replacing the truncation errors, that have been estimated in that paper using the EKM approach of Refs. [21, 122], with the Bayesian uncertainties calculated as described in section 5. The 68% DoB Bayesian truncation errors are similar to those quoted in Ref. [8] at N²LO but appear to be significantly larger at NLO. We also calculated in that paper the excitation energies for selected states of $A = 6 - 12$ nuclei. For almost all considered cases, adding the 3NF to the NN interaction was found to lead to a significant improvement in

the description of experimental data. The predicted ground state energies of p-shell nuclei show a good agreement with the data except for ^{16}O , which appears to be overbound. Notice that the deviation between the predicted and experimental values of the ^{16}O binding energy agrees with the 95% DoB Bayesian error at N^2LO . It will be very interesting to repeat the calculations for the newest SMS interactions and to extend them to higher orders.

7 SUMMARY AND OUTLOOK

In this review article we have presented a snapshot of the current state-of-the-art in low-energy nuclear theory with a focus on the latest generation of semilocal nuclear potentials from chiral EFT. We now summarize some of the key conclusions of our paper.

- We have presented a concise and self-contained introduction to the conceptual foundations of chiral effective field theory in the few-nucleon sector and described in some detail all steps needed to compute low-energy observables from the effective chiral Lagrangians (including error analysis). Special emphasis was given to clarify the notion of *consistency* of nuclear forces and current operators in terms of a perturbative matching to the unambiguously defined on-shell scattering amplitude. In particular, few-nucleon potentials from Refs. [64, 46, 47, 40, 85] and electroweak current operators from Refs. [114, 65, 66, 115] at N^3LO and beyond, derived using DR, are off-shell consistent with each other provided DR is also used to compute loop integrals arising from iterations of the dynamical equations.
- We have reviewed the semilocal momentum-space regularized potentials of Ref. [17], which are currently the most precise chiral EFT NN forces on the market. These are the only NN interactions derived in chiral EFT, which – from the statistical point of view – qualify to be regarded as PWA of NN data below pion production threshold, see section 6.1 for details. At the highest considered order N^4LO^+ , these interactions describe the np and pp data from the self-consistent Granada-2013 database with a precision that is at least comparable to the one reached by modern phenomenological potentials with a much larger number of adjustable parameters. The significantly better description of the scattering data by the SMS N^4LO^+ interactions of Ref. [17] as compared to the nonlocal potentials of Ref. [106] at the same chiral order, and their much smaller residual cutoff dependence, see Fig. 17 of Ref. [17], can presumably be traced back to the improved semilocal regulator, which maintains the long-range part of the interaction as described in section 4.1. We also addressed in detail the issue of uncertainty quantification in the NN sector. In particular, we discussed statistical uncertainties of NN and πN LECs and their propagation to selected observables as well as uncertainty introduced by fixing the maximum fit energy in the determination of the NN LECs. We also estimated truncation errors at various chiral orders using the Bayesian model specified in section 5.
- Beyond the NN sector, calculations based on the SMS interactions have so far been carried out up to N^2LO [75]. The LECs c_D and c_E , which enter the 3NF at this order, have been determined from the ^3H binding energy and the very precise pd cross section data at $E_{\text{lab}}^N = 70$ MeV from Ref. [176]. Using the employed Bayesian model to estimate truncation uncertainties, the predicted ground state energies of p-shell nuclei up to $A = 16$ are generally in a good agreement with the data. Also the predicted Nd scattering observables including the vector analyzing power A_y are consistent with experimental data within errors. We performed an additional test of the employed Bayesian model for truncation errors by exploring the impact of selected short-range 3NF terms at N^4LO on observables in Nd elastic scattering. Our results suggest that a high-precision description of Nd scattering data will likely require the chiral expansion of the 3NF to be pushed to N^4LO .

- The novel semilocal nuclear forces, derived in the finite-cutoff formulation of chiral EFT with short-range interactions counted according to NDA (i.e. the Weinberg scheme), have already been successfully confronted with few-nucleon data and passed a number of a-posteriori consistency checks as briefly summarized below:
 - Using the minimal basis of the order- Q^4 NN contact interactions as detailed in section 6, the LECs determined from the np and pp scattering data come out of a natural size, see Fig. 7. The same is true for the LECs c_D and c_E entering the leading 3NF, as can be seen e.g. from the corresponding expectation values in the ^3H state [75].
 - The residual cutoff-dependence of NN phase shifts is strongly reduced at $\text{N}^{3,4}\text{LO}$ as compared to $\text{N}^{1,2}\text{LO}$ within the considered Λ -range, see e.g. Fig. 4 of Ref. [73].
 - There is a clear systematic improvement in the description of np and pp data with increasing chiral orders, see Tab. 1. At order Q^3 (i.e. N^2LO), this improvement results solely from taking into account the parameter-free subleading TPEP contributions. Notice that certain alternative power counting schemes suggest that some of the contact interactions that appear at order Q^4 in the Weinberg scheme are enhanced and should yield contributions to observables larger than the order- Q^3 TPEP, see e.g. Table 1 of Ref. [72]. The clear evidence of the chiral TPEP at orders Q^3 and Q^5 observed in Refs. [21, 22, 17] does, however, not support such alternative scenarios.
 - The resulting convergence pattern of the EFT expansion for selected NN observables was scrutinized using Bayesian statistical methods, see section 5 for details. For not too soft cutoffs, the assumed breakdown scale of the EFT expansion of $\Lambda_b \sim 600$ MeV [21] was found to be statistically consistent [123], see also Refs. [124, 75] for a related discussion.
 - Scheme-dependence of nuclear potentials offers yet another way to perform nontrivial consistency checks of the theoretical framework by explicitly verifying (approximate) scheme-independence of observables. In the formulation we employ, scheme dependence of the nuclear forces first appears at N^3LO and manifests itself in their dependence on arbitrary real phases $\bar{\beta}_8, \bar{\beta}_9$, which parametrize the unitary ambiguity of the leading relativistic corrections [65, 21], and the appearance of three off-shell short-range operators in the $^1\text{S}_0$ and $^3\text{S}_1$ - $^3\text{D}_1$ channels proportional to the LECs $D_{1\text{S}0}^{\text{off}}$, $D_{3\text{S}1}^{\text{off}}$ and $D_{\epsilon 1}^{\text{off}}$ [197, 198, 17]. The SMS potentials of Ref. [17] make use of the standard choice for $\bar{\beta}_{8,9}$ namely $\bar{\beta}_8 = -\bar{\beta}_9 = 1/4$, which minimizes the amount of $1/m^2$ -corrections to the OPEP, and employ $D_{1\text{S}0}^{\text{off}} = D_{3\text{S}1}^{\text{off}} = D_{\epsilon 1}^{\text{off}} = 0$. Different choices of these parameters lead to different off-shell behaviors of the potential. They are related to each other by unitary transformations which, however, also induce an infinite tower of higher-order terms beyond the order one is working. The residual dependence of observables on $\bar{\beta}_{8,9}$ and D_i^{off} , therefore, probes the impact of neglected higher-order terms and should lie within the estimated truncation errors. We have redone the fits at N^4LO^+ for $\Lambda = 450$ MeV using alternative choices of D_i^{off} [17] and also developed a version of the potential with $\bar{\beta}_8 = \bar{\beta}_9 = 1/2$ [199]. The latter choice is motivated by the vanishing isoscalar exchange charge density operator at N^3LO . In all considered cases, we found negligibly small changes in observables in spite of strong changes at the interaction level.
 - Calculations of three- and more-nucleon observables based on solely NN interactions are incomplete beyond second order. They do, however, provide information about the magnitude of the missing 3NF contributions by assessing the spread in results at different orders $Q^{\geq 3}$ and via a comparison of such incomplete predictions with experimental data. In Ref. [74], such an analysis was performed for Nd scattering observables and selected properties of light nuclei using the SCS NN interactions of Refs. [21, 22]. The sizes of the 3NF contributions required to bring such incomplete results in

agreement with experimental data were found to agree well with expectations based on Weinberg's power counting. Furthermore, recent calculations by the LENPIC Collaboration which include the leading 3NF [8, 75] show that the resulting N²LO predictions for observables that have not been used in the determination of the LECs c_D , c_E are generally in a good agreement with the data, see section 6. No indications are found for enhanced contributions of the 3NF in general and of the c_D -term in particular as suggested in Ref. [200].

To summarize, major progress has been achieved in recent years towards developing chiral EFT into a precision tool for low-energy nuclear physics. In the NN sector, the latest SMS interactions at fifth chiral order have already reached the accuracy at or even below permille level for low-energy observables such as e.g. the deuteron asymptotic S-state normalization A_S , see section 6.1 for details and further examples. The only essential missing step in the NN sector concerns the inclusion of isospin-breaking interactions up to fifth chiral order. Work along this line is in progress.

Pushing the precision frontier beyond the NN system opens exciting perspectives for low-energy nuclear theory and will allow one to confront chiral EFT with currently unsolved problems, such as a quantitative description of 3N scattering observables [19]. This, however, will require to address the two core challenges:

- (i) Derivation of *consistent regularized* three- and four-nucleon forces and exchange charge and current operators at and beyond N³LO as detailed in section 4.2. This issue has not been paid attention to in the recent calculations involving the 3NFs [201, 202, 203, 204] and exchange electroweak currents [100, 205, 206] at N³LO. As explained in section 4.2, using ad hoc regularization approaches at N³LO and beyond generally leads to incorrect results for the scattering amplitude and other observables due to the appearance of uncontrolled short-range artifacts, which violate chiral symmetry and are not suppressed by inverse powers of Λ . This puts the findings of these studies into question.
- (ii) Determination of the LECs in the 3NF at N⁴LO. While the N³LO contributions to the 3NF and 4NF do not involve unknown parameters, the N⁴LO corrections to the 3NF involve 10 LEC accompanying purely short-range operators [37] and one or more LECs entering the one-pion-exchange-contact topology, which has not been worked out yet. As discussed in section 6.2, the determination of these LECs from 3N data will require a computationally challenging analysis.

As a first example of a precision calculation not restricted to NN scattering, we have recently determined the deuteron structure radius with an accuracy below the permille level, $r_{\text{str}} = 1.9731^{+0.0013}_{-0.0018}$ fm, by pushing the chiral expansion of the electromagnetic exchange charge density beyond N³LO and performing a thorough analysis of various types of uncertainty [199]. By combining the predicted value for r_{str} with the very accurate atomic data from isotope shift measurements, it was, for the first time, possible to extract the neutron charge radius from experimental data on light nuclei. This study was facilitated by the absence of loop contributions in the isoscalar exchange charge density at N³LO [114, 65], which allowed for a trivial construction of the corresponding consistently regularized expressions for the charge operator. Rederivation of the contributions to 3NFs, 4NFs and exchange currents at and beyond N³LO using a regulator, consistent with the one employed in Ref. [17], would open the way for performing similar precision calculations for a broad class of low-energy few-nucleon reactions.

FUNDING

This work was supported by BMBF (Grant No. 05P18PCFP1) and by DFG through funds provided to the Sino-German CRC 110 110 “Symmetries and the Emergence of Structure in QCD” (Grant No. TRR110).

ACKNOWLEDGMENTS

We are grateful to Ashot Gasparyan, Jambul Gegelia and Ulf-G. Meißner for a careful reading of the manuscript and useful comments and to all members of the LENPIC Collaboration for sharing their insights into the topics addressed in this review article. We also thank Pieter Maris for providing us with a source file of Fig. 8 from Ref. [8].

REFERENCES

- [1] S. Weinberg, Phys. Lett. B **251**, 288 (1990).
- [2] S. Weinberg, Nucl. Phys. B **363**, 3 (1991).
- [3] E. Epelbaum, Prog. Part. Nucl. Phys. **57**, 654 (2006) [nucl-th/0509032].
- [4] E. Epelbaum, H. W. Hammer and U.-G. Meißner, Rev. Mod. Phys. **81**, 1773 (2009) [arXiv:0811.1338 [nucl-th]].
- [5] R. Machleidt and D. R. Entem, Phys. Rept. **503**, 1 (2011) [arXiv:1105.2919 [nucl-th]].
- [6] H.-W. Hammer, S. König and U. van Kolck, arXiv:1906.12122 [nucl-th].
- [7] E. Epelbaum, A. M. Gasparyan, J. Gegelia and U.-G. Meißner, Eur. Phys. J. A **54**, no. 11, 186 (2018) [arXiv:1810.02646 [nucl-th]].
- [8] E. Epelbaum *et al.* [LENPIC Collaboration], Phys. Rev. C **99**, no. 2, 024313 (2019) [arXiv:1807.02848 [nucl-th]].
- [9] M. Piarulli *et al.*, Phys. Rev. Lett. **120**, no. 5, 052503 (2018) [arXiv:1707.02883 [nucl-th]].
- [10] D. Lonardonì, J. Carlson, S. Gandolfi, J. E. Lynn, K. E. Schmidt, A. Schwenk and X. Wang, Phys. Rev. Lett. **120**, no. 12, 122502 (2018) [arXiv:1709.09143 [nucl-th]].
- [11] G. Hagen, G. R. Jansen and T. Papenbrock, Phys. Rev. Lett. **117**, no. 17, 172501 (2016) [arXiv:1605.01477 [nucl-th]].
- [12] E. Gebrerufael, K. Vobig, H. Hergert and R. Roth, Phys. Rev. Lett. **118**, no. 15, 152503 (2017).
- [13] A. Cipollone, C. Barbieri and P. Navrátil, Phys. Rev. C **92**, no. 1, 014306 (2015) [arXiv:1412.0491 [nucl-th]].
- [14] E. Epelbaum, H. Krebs, D. Lee and U.-G. Meißner, Phys. Rev. Lett. **106**, 192501 (2011) [arXiv:1101.2547 [nucl-th]].
- [15] S. Elhatisari, D. Lee, G. Rupak, E. Epelbaum, H. Krebs, T. A. Lähde, T. Luu and U.-G. Meißner, Nature **528**, 111 (2015) [arXiv:1506.03513 [nucl-th]].
- [16] T. A. Lähde and U.-G. Meißner, Lect. Notes Phys. **957**, 1 (2019).
- [17] P. Reinert, H. Krebs and E. Epelbaum, Eur. Phys. J. A **54**, no. 5, 86 (2018) [arXiv:1711.08821 [nucl-th]].
- [18] R. Navarro Pérez, J. E. Amaro and E. Ruiz Arriola, Phys. Rev. C **88**, no. 6, 064002 (2013) Erratum: [Phys. Rev. C **91**, no. 2, 029901 (2015)] [arXiv:1310.2536 [nucl-th]].
- [19] N. Kalantar-Nayestanaki, E. Epelbaum, J. G. Messchendorp and A. Nogga, Rept. Prog. Phys. **75**, 016301 (2012) [arXiv:1108.1227 [nucl-th]].
- [20] H. W. Hammer, A. Nogga and A. Schwenk, Rev. Mod. Phys. **85**, 197 (2013) [arXiv:1210.4273 [nucl-th]].

- [21] E. Epelbaum, H. Krebs and U.-G. Meißner, *Eur. Phys. J. A* **51**, no. 5, 53 (2015) [arXiv:1412.0142 [nucl-th]].
- [22] E. Epelbaum, H. Krebs and U.-G. Meißner, *Phys. Rev. Lett.* **115**, no. 12, 122301 (2015) [arXiv:1412.4623 [nucl-th]].
- [23] C. Ordonez, L. Ray and U. van Kolck, *Phys. Rev. C* **53**, 2086 (1996) [hep-ph/9511380].
- [24] N. Kaiser, S. Gerstendorfer and W. Weise, *Nucl. Phys. A* **637**, 395 (1998) [nucl-th/9802071].
- [25] H. Krebs, E. Epelbaum and U.-G. Meißner, *Eur. Phys. J. A* **32**, 127 (2007) [nucl-th/0703087].
- [26] E. Epelbaum, H. Krebs and U.-G. Meißner, *Nucl. Phys. A* **806**, 65 (2008) [arXiv:0712.1969 [nucl-th]].
- [27] E. Epelbaum, H. Krebs and U.-G. Meißner, *Phys. Rev. C* **77**, 034006 (2008) [arXiv:0801.1299 [nucl-th]].
- [28] H. Krebs, A. M. Gasparyan and E. Epelbaum, *Phys. Rev. C* **98**, no. 1, 014003 (2018) [arXiv:1803.09613 [nucl-th]].
- [29] S. R. Coleman, J. Wess and B. Zumino, *Phys. Rev.* **177**, 2239 (1969).
- [30] C. G. Callan, Jr., S. R. Coleman, J. Wess and B. Zumino, *Phys. Rev.* **177**, 2247 (1969).
- [31] J. Gasser and H. Leutwyler, *Annals Phys.* **158**, 142 (1984).
- [32] V. Bernard, N. Kaiser and U.-G. Meißner, *Int. J. Mod. Phys. E* **4**, 193 (1995) [hep-ph/9501384].
- [33] N. Fettes, U.-G. Meißner, M. Mojzis and S. Steininger, *Annals Phys.* **283**, 273 (2000) Erratum: [*Annals Phys.* **288**, 249 (2001)] [hep-ph/0001308].
- [34] E. Epelbaum, U.-G. Meißner, W. Glöckle and C. Elster, *Phys. Rev. C* **65**, 044001 (2002) [nucl-th/0106007].
- [35] L. Girlanda, S. Pastore, R. Schiavilla and M. Viviani, *Phys. Rev. C* **81**, 034005 (2010) [arXiv:1001.3676 [nucl-th]].
- [36] E. Epelbaum, A. Nogga, W. Glöckle, H. Kamada, U.-G. Meißner and H. Witała, *Phys. Rev. C* **66**, 064001 (2002) [nucl-th/0208023].
- [37] L. Girlanda, A. Kievsky and M. Viviani, *Phys. Rev. C* **84**, 014001 (2011) [arXiv:1102.4799 [nucl-th]].
- [38] G. Müller and U.-G. Meißner, *Nucl. Phys. B* **556**, 265 (1999) [hep-ph/9903375].
- [39] J. Gasser, M. A. Ivanov, E. Lipartia, M. Mojzis and A. Rusetsky, *Eur. Phys. J. C* **26**, 13 (2002) [hep-ph/0206068].
- [40] H. Krebs, A. Gasparyan and E. Epelbaum, *Phys. Rev. C* **85**, 054006 (2012) [arXiv:1203.0067 [nucl-th]].
- [41] M. Hoferichter, J. Ruiz de Elvira, B. Kubis and U.-G. Meißner, *Phys. Rept.* **625**, 1 (2016) [arXiv:1510.06039 [hep-ph]].
- [42] M. Hoferichter, J. Ruiz de Elvira, B. Kubis and U.-G. Meißner, *Phys. Rev. Lett.* **115**, no. 19, 192301 (2015) [arXiv:1507.07552 [nucl-th]].
- [43] D. Siemens, V. Bernard, E. Epelbaum, A. M. Gasparyan, H. Krebs and U.-G. Meißner, *Phys. Rev. C* **96**, no. 5, 055205 (2017) [arXiv:1704.08988 [nucl-th]].
- [44] A. Manohar and H. Georgi, *Nucl. Phys. B* **234**, 189 (1984).
- [45] V. Bernard, *Prog. Part. Nucl. Phys.* **60**, 82 (2008) [arXiv:0706.0312 [hep-ph]].
- [46] V. Bernard, E. Epelbaum, H. Krebs and U.-G. Meißner, *Phys. Rev. C* **77**, 064004 (2008) [arXiv:0712.1967 [nucl-th]].
- [47] V. Bernard, E. Epelbaum, H. Krebs and U.-G. Meißner, *Phys. Rev. C* **84**, 054001 (2011) [arXiv:1108.3816 [nucl-th]].
- [48] G. P. Lepage, *nucl-th/9706029*.
- [49] M. Pavon Valderrama and E. Ruiz Arriola, *Phys. Rev. C* **70**, 044006 (2004) [nucl-th/0405057].
- [50] A. Nogga, R. G. E. Timmermans and U. van Kolck, *Phys. Rev. C* **72**, 054006 (2005) [nucl-th/0506005].

- [51] M. C. Birse, Phys. Rev. C **74**, 014003 (2006) [nucl-th/0507077].
- [52] E. Epelbaum and U.-G. Meißner, Few Body Syst. **54**, 2175 (2013) [nucl-th/0609037].
- [53] E. Epelbaum and J. Gegelia, Eur. Phys. J. A **41**, 341 (2009) [arXiv:0906.3822 [nucl-th]].
- [54] B. Long and C. J. Yang, Phys. Rev. C **85**, 034002 (2012) [arXiv:1111.3993 [nucl-th]].
- [55] M. P. Valderrama, Int. J. Mod. Phys. E **25**, no. 05, 1641007 (2016) [arXiv:1604.01332 [nucl-th]].
- [56] M. J. Savage, In *Pasadena 1998, Nuclear physics with effective field theory* 247-267 [nucl-th/9804034].
- [57] D. B. Kaplan, M. J. Savage and M. B. Wise, Phys. Lett. B **424**, 390 (1998) [nucl-th/9801034].
- [58] T. D. Cohen and J. M. Hansen, Phys. Rev. C **59**, 3047 (1999) [nucl-th/9901065].
- [59] S. Fleming, T. Mehen and I. W. Stewart, Nucl. Phys. A **677**, 313 (2000) [nucl-th/9911001].
- [60] D. B. Kaplan, arXiv:1905.07485 [nucl-th].
- [61] E. Epelbaum and J. Gegelia, Phys. Lett. B **716**, 338 (2012) [arXiv:1207.2420 [nucl-th]].
- [62] E. Epelbaum, A. M. Gasparyan, J. Gegelia and H. Krebs, Eur. Phys. J. A **51**, no. 6, 71 (2015) [arXiv:1501.01191 [nucl-th]].
- [63] E. Epelbaum, Phys. Lett. B **639**, 456 (2006) [nucl-th/0511025].
- [64] E. Epelbaum, Eur. Phys. J. A **34**, 197 (2007) [arXiv:0710.4250 [nucl-th]].
- [65] S. Kölling, E. Epelbaum, H. Krebs and U.-G. Meißner, Phys. Rev. C **84**, 054008 (2011) [arXiv:1107.0602 [nucl-th]].
- [66] H. Krebs, E. Epelbaum and U.-G. Meißner, Annals Phys. **378**, 317 (2017) [arXiv:1610.03569 [nucl-th]].
- [67] E. Epelbaum, A. Nogga, W. Glöckle, H. Kamada, U.-G. Meißner and H. Witała, Eur. Phys. J. A **15**, 543 (2002) [nucl-th/0201064].
- [68] H. Krebs, arXiv:1908.01538 [nucl-th].
- [69] E. Epelbaum, arXiv:1908.09349 [nucl-th].
- [70] J. Gegelia, G. Japaridze and X. Q. Wang, J. Phys. G **29**, 2303 (2003) [hep-ph/9910260].
- [71] T. Fuchs, J. Gegelia, G. Japaridze and S. Scherer, Phys. Rev. D **68**, 056005 (2003) [hep-ph/0302117].
- [72] H. W. Griebhammer, PoS CD **15**, 104 (2016) [arXiv:1511.00490 [nucl-th]].
- [73] E. Epelbaum, PoS CD **15**, 014 (2016) [arXiv:1510.07036 [nucl-th]].
- [74] S. Binder *et al.* [LENPIC Collaboration], Phys. Rev. C **93**, no. 4, 044002 (2016) [arXiv:1505.07218 [nucl-th]].
- [75] E. Epelbaum *et al.*, arXiv:1907.03608 [nucl-th].
- [76] E. Epelbaum and J. Gegelia, PoS CD **12**, 090 (2013) [arXiv:1301.6134 [nucl-th]].
- [77] V. Baru, E. Epelbaum, A. A. Filin and J. Gegelia, Phys. Rev. C **92**, no. 1, 014001 (2015) [arXiv:1504.07852 [nucl-th]].
- [78] V. Baru, E. Epelbaum and A. A. Filin, Phys. Rev. C **94**, no. 1, 014001 (2016) [arXiv:1604.02551 [nucl-th]].
- [79] T. A. Lähde, U.-G. Meißner and E. Epelbaum, arXiv:1906.00607 [nucl-th].
- [80] E. Epelbaum, arXiv:1001.3229 [nucl-th].
- [81] H. van Haeringen and L. P. Kok, Phys. Rev. A **26**, 1218 (1982).
- [82] E. Epelbaum, J. Gegelia and U.-G. Meißner, Nucl. Phys. B **925**, 161 (2017) [arXiv:1705.02524 [nucl-th]].
- [83] K. Suzuki, R. Okamoto, Prog. Theor. Phys. **70**, 439 (1983).
- [84] S. Okubo, Prog. Theor. Phys. **12**, 603 (1954).
- [85] H. Krebs, A. Gasparyan and E. Epelbaum, Phys. Rev. C **87**, no. 5, 054007 (2013) [arXiv:1302.2872 [nucl-th]].

- [86] N. Kaiser, R. Brockmann and W. Weise, Nucl. Phys. A **625**, 758 (1997) [nucl-th/9706045].
- [87] E. Epelbaum, W. Glöckle and U.-G. Meißner, Nucl. Phys. A **637**, 107 (1998) [nucl-th/9801064].
- [88] U. van Kolck, Phys. Rev. C **49**, 2932 (1994).
- [89] N. Kaiser, Phys. Rev. C **62**, 024001 (2000) [nucl-th/9912054].
- [90] N. Kaiser, Phys. Rev. C **61**, 014003 (2000) [nucl-th/9910044].
- [91] N. Kaiser, Phys. Rev. C **63**, 044010 (2001) [nucl-th/0101052].
- [92] N. Kaiser, Phys. Rev. C **64**, 057001 (2001) [nucl-th/0107064].
- [93] N. Kaiser, Phys. Rev. C **65**, 017001 (2002) [nucl-th/0109071].
- [94] S. Ishikawa and M. R. Robilotta, Phys. Rev. C **76**, 014006 (2007) [arXiv:0704.0711 [nucl-th]].
- [95] D. R. Entem, N. Kaiser, R. Machleidt and Y. Nosyk, Phys. Rev. C **91**, no. 1, 014002 (2015) [arXiv:1411.5335 [nucl-th]].
- [96] E. Epelbaum, W. Glöckle, A. Krüger and U.-G. Meißner, Nucl. Phys. A **645**, 413 (1999) [nucl-th/9809084].
- [97] W. N. Polyzou, Phys. Rev. C **82**, 014002 (2010) [arXiv:1001.4523 [math-ph]].
- [98] S. Pastore, L. Girlanda, R. Schiavilla, M. Viviani and R. B. Wiringa, Phys. Rev. C **80**, 034004 (2009) [arXiv:0906.1800 [nucl-th]].
- [99] S. Pastore, L. Girlanda, R. Schiavilla and M. Viviani, Phys. Rev. C **84**, 024001 (2011) [arXiv:1106.4539 [nucl-th]].
- [100] M. Piarulli, L. Girlanda, L. E. Marcucci, S. Pastore, R. Schiavilla and M. Viviani, Phys. Rev. C **87**, no. 1, 014006 (2013) [arXiv:1212.1105 [nucl-th]].
- [101] A. Baroni, L. Girlanda, S. Pastore, R. Schiavilla and M. Viviani, Phys. Rev. C **93**, no. 1, 015501 (2016) Erratum: [Phys. Rev. C **93**, no. 4, 049902 (2016)] Erratum: [Phys. Rev. C **95**, no. 5, 059901 (2017)] [arXiv:1509.07039 [nucl-th]].
- [102] D. R. Entem and R. Machleidt, Phys. Rev. C **68**, 041001 (2003) [nucl-th/0304018].
- [103] E. Epelbaum, W. Glöckle and U.-G. Meißner, Nucl. Phys. A **747**, 362 (2005) [nucl-th/0405048].
- [104] B. D. Carlsson *et al.*, Phys. Rev. X **6**, no. 1, 011019 (2016) [arXiv:1506.02466 [nucl-th]].
- [105] A. Ekström *et al.*, Phys. Rev. C **91**, no. 5, 051301 (2015) [arXiv:1502.04682 [nucl-th]].
- [106] D. R. Entem, R. Machleidt and Y. Nosyk, Phys. Rev. C **96**, no. 2, 024004 (2017) [arXiv:1703.05454 [nucl-th]].
- [107] A. Ekström, G. Hagen, T. D. Morris, T. Papenbrock and P. D. Schwartz, Phys. Rev. C **97**, no. 2, 024332 (2018) [arXiv:1707.09028 [nucl-th]].
- [108] E. Epelbaum, W. Glöckle and U.-G. Meißner, Eur. Phys. J. A **19**, 125 (2004) [nucl-th/0304037].
- [109] E. Epelbaum, W. Glöckle and U.-G. Meißner, Eur. Phys. J. A **19**, 401 (2004) [nucl-th/0308010].
- [110] T. A. Rijken, Annals Phys. **208**, 253 (1991).
- [111] A. A. Slavnov, Nucl. Phys. B **31**, 301 (1971).
- [112] D. Djukanovic, M. R. Schindler, J. Gegelia and S. Scherer, Phys. Rev. D **72**, 045002 (2005) [hep-ph/0407170].
- [113] B. Long and Y. Mei, Phys. Rev. C **93**, no. 4, 044003 (2016) [arXiv:1605.02153 [nucl-th]].
- [114] S. Kölling, E. Epelbaum, H. Krebs and U.-G. Meißner, Phys. Rev. C **80**, 045502 (2009) [arXiv:0907.3437 [nucl-th]].
- [115] H. Krebs, E. Epelbaum and U.-G. Meißner, Few Body Syst. **60**, no. 2, 31 (2019) [arXiv:1902.06839 [nucl-th]].
- [116] D. Siemens, V. Bernard, E. Epelbaum, A. Gasparyan, H. Krebs and U.-G. Meißner, Phys. Rev. C **94**, no. 1, 014620 (2016).

- [117] D. L. Yao, D. Siemens, V. Bernard, E. Epelbaum, A. M. Gasparyan, J. Gegelia, H. Krebs and U.-G. Meißner, *JHEP* **1605**, 038 (2016).
- [118] A. N. Hiller Blin, Z. F. Sun and M. J. Vicente Vacas, *Phys. Rev. D* **98**, no. 5, 054025 (2018).
- [119] J. Hu, Y. Zhang, E. Epelbaum, U.-G. Meißner and J. Meng, *Phys. Rev. C* **96**, no. 3, 034307 (2017).
- [120] R. Skibiński *et al.*, *Phys. Rev. C* **93**, no. 6, 064002 (2016).
- [121] N. Nevo Dinur, O. J. Hernandez, S. Bacca, N. Barnea, C. Ji, S. Pastore, M. Piarulli and R. B. Wiringa, *Phys. Rev. C* **99**, no. 3, 034004 (2019).
- [122] S. Binder *et al.* [LENPIC Collaboration], *Phys. Rev. C* **98**, no. 1, 014002 (2018) [arXiv:1802.08584 [nucl-th]].
- [123] R. J. Furnstahl, N. Klco, D. R. Phillips and S. Wesolowski, *Phys. Rev. C* **92**, no. 2, 024005 (2015).
- [124] J. A. Melendez, S. Wesolowski and R. J. Furnstahl, *Phys. Rev. C* **96**, no. 2, 024003 (2017).
- [125] J. A. Melendez, R. J. Furnstahl, D. R. Phillips, M. T. Pratola and S. Wesolowski, *Phys. Rev. C* **100**, no. 4, 044001 (2019) [arXiv:1904.10581 [nucl-th]].
- [126] E. Epelbaum, *High-precision nuclear forces: Where do we stand?*, to appear in proceedings of the 9th International workshop on Chiral Dynamics, 17-21 September 2018, Durham, NC, USA.
- [127] M. Tanabashi *et al.* [Particle Data Group], *Phys. Rev. D* **98**, no. 3, 030001 (2018).
- [128] N. Fettes, U.-G. Meißner and S. Steininger, *Nucl. Phys. A* **640**, 199 (1998) [hep-ph/9803266].
- [129] V. G. J. Stoks, R. A. M. Klomp, M. C. M. Rentmeester and J. J. de Swart, *Phys. Rev. C* **48**, 792 (1993).
- [130] G. J. M. Austen and J. J. de Swart, *Phys. Rev. Lett.* **50**, 2039 (1983).
- [131] V. G. J. Stoks and J. J. De Swart, *Phys. Rev. C* **42**, 1235 (1990).
- [132] L. Durand, *Phys. Rev.* **108**, 1597 (1957).
- [133] B. H. Daub, V. Henzl, M. A. Kovash, J. L. Matthews, Z. W. Miller, K. Shoniyozov and H. Yang, *Phys. Rev. C* **87**, no. 1, 014005 (2013) [arXiv:1211.2203 [nucl-ex]].
- [134] G. F. Cox, G. H. Eaton, C. P. Van Zyl, O. N. Jarvis and B. Rose, *Nucl. Phys. B* **4**, 353 (1968).
- [135] C. Van Der Leun and C. Alderliesten, *Nucl. Phys. A* **380**, 261 (1982).
- [136] K. Schoen *et al.*, *Phys. Rev. C* **67**, 044005 (2003).
- [137] R. Machleidt, *Phys. Rev. C* **63**, 024001 (2001) [nucl-th/0006014].
- [138] V. G. J. Stoks, R. A. M. Klomp, C. P. F. Terheggen and J. J. de Swart, *Phys. Rev. C* **49**, 2950 (1994) [nucl-th/9406039].
- [139] F. Gross and A. Stadler, *Phys. Rev. C* **78**, 014005 (2008) [arXiv:0802.1552 [nucl-th]].
- [140] R. Navarro Pérez, J. E. Amaro and E. Ruiz Arriola, *Phys. Rev. C* **88**, 024002 (2013) Erratum: [*Phys. Rev. C* **88**, no. 6, 069902 (2013)] [arXiv:1304.0895 [nucl-th]].
- [141] O. N. Jarvis, C. Whitehead and M. Shah, *Phys. Lett.* **36B**, no. 4, 409 (1971).
- [142] A. E. Taylor, E. Wood and L. Bird, *Nucl. Phys.* **16**, 320 (1960).
- [143] L. Bird, P. Christmas, A. Taylor and E. Wood, *Nucl. Phys.* **27**, 586 (1961).
- [144] O. Jarvis, B. Rose, J. Scanlon and E. Wood, *Nucl. Phys.* **42**, 294 (1963).
- [145] S. Aldor-Noiman, L. D. Brown, A. Buja, W. Rolke and R. A. Stine, *Am. Stat.* **67**, 249 (2013) Erratum: [*Am. Stat.* **68**, 318 (2014)].
- [146] R. Navarro Pérez, J. E. Amaro and E. Ruiz Arriola, *Phys. Rev. C* **89**, no. 6, 064006 (2014) [arXiv:1404.0314 [nucl-th]].
- [147] P. W. Lisowski, R. E. Shamu, G. F. Auchampaugh, N. S. P. King, M. S. Moore, G. L. Morgan and T. S. Singleton, *Phys. Rev. Lett.* **49**, 255 (1982).
- [148] T. E. O. Ericson and M. Rosa-Clot, *Nucl. Phys. A* **405**, 497 (1983).
- [149] N. L. Rodning and L. D. Knutson, *Phys. Rev. C* **41**, 898 (1990).

- [150] A. Huber, T. Udem, B. Gross, J. Reichert, M. Kourogi, K. Pachucki, M. Weitz and T. W. Hansch, *Phys. Rev. Lett.* **80**, 468 (1998).
- [151] D. M. Bishop and L. M. Cheung, *Phys. Rev. A* **20**, 381 (1979).
- [152] J. L. Friar and U. van Kolck, *Phys. Rev. C* **60**, 034006 (1999) [nucl-th/9906048].
- [153] J. L. Friar, U. van Kolck, G. L. Payne and S. A. Coon, *Phys. Rev. C* **68**, 024003 (2003) [nucl-th/0303058].
- [154] E. Epelbaum and U.-G. Meißner, *Phys. Rev. C* **72**, 044001 (2005) [nucl-th/0502052].
- [155] U. van Kolck, M. C. M. Rentmeester, J. L. Friar, J. T. Goldman and J. J. de Swart, *Phys. Rev. Lett.* **80**, 4386 (1998) [nucl-th/9710067].
- [156] R. Navarro Pérez, J. E. Amaro and E. Ruiz Arriola, *Phys. Rev. C* **95**, no. 6, 064001 (2017) [arXiv:1606.00592 [nucl-th]].
- [157] W. Glöckle, H. Witała, D. Hüber, H. Kamada and J. Golak, *Phys. Rept.* **274**, 107 (1996).
- [158] A. Kievsky, M. Viviani, S. Rosati, D. Huber, W. Glöckle, H. Kamada, H. Witała and J. Golak, *Phys. Rev. C* **58**, 3085 (1998) [nucl-th/9807061].
- [159] H. Witała and W. Glöckle, *Nucl. Phys. A* **528**, 48 (1991).
- [160] H. Witała and W. Glöckle, *J. Phys. G* **37**, 064003 (2010) [arXiv:0909.0495 [nucl-th]].
- [161] S. A. Coon and H. K. Han, *Few Body Syst.* **30**, 131 (2001) [nucl-th/0101003].
- [162] B. S. Pudliner, V. R. Pandharipande, J. Carlson and R. B. Wiringa, *Phys. Rev. Lett.* **74**, 4396 (1995) [nucl-th/9502031].
- [163] H. Witała, J. Golak, R. Skibiński, W. Glöckle, H. Kamada and W. N. Polyzou, *Phys. Rev. C* **83**, 044001 (2011) Erratum: [*Phys. Rev. C* **88**, no. 6, 069904 (2013)] [arXiv:1101.4053 [nucl-th]].
- [164] D. R. Phillips and C. Schat, *Phys. Rev. C* **88**, no. 3, 034002 (2013) [arXiv:1307.6274 [nucl-th]].
- [165] E. Epelbaum, A. M. Gasparyan, H. Krebs and C. Schat, *Eur. Phys. J. A* **51**, no. 3, 26 (2015) [arXiv:1411.3612 [nucl-th]].
- [166] K. Topolnicki, *Eur. Phys. J. A* **53**, no. 9, 181 (2017).
- [167] R. B. Wiringa, V. G. J. Stoks and R. Schiavilla, *Phys. Rev. C* **51**, 38 (1995) [nucl-th/9408016].
- [168] J. Hu, P. Wei and Y. Zhang, *Phys. Lett. B* **798**, 134982 (2019) [arXiv:1909.11826 [nucl-th]].
- [169] J. Golak *et al.*, *Eur. Phys. J. A* **43**, 241 (2010) [arXiv:0911.4173 [nucl-th]].
- [170] K. Hebeler, H. Krebs, E. Epelbaum, J. Golak and R. Skibiński, *Phys. Rev. C* **91**, no. 4, 044001 (2015) [arXiv:1502.02977 [nucl-th]].
- [171] P. Navratil, V. G. Gueorguiev, J. P. Vary, W. E. Ormand and A. Nogga, *Phys. Rev. Lett.* **99**, 042501 (2007) [nucl-th/0701038].
- [172] D. Gazit, S. Quaglioni and P. Navratil, *Phys. Rev. Lett.* **103**, 102502 (2009) Erratum: [*Phys. Rev. Lett.* **122**, no. 2, 029901 (2019)] [arXiv:0812.4444 [nucl-th]].
- [173] J. E. Lynn, I. Tews, J. Carlson, S. Gandolfi, A. Gezerlis, K. E. Schmidt and A. Schwenk, *Phys. Rev. C* **96**, no. 5, 054007 (2017) [arXiv:1706.07668 [nucl-th]].
- [174] L. Platter, H.-W. Hammer and U.-G. Meißner, *Phys. Lett. B* **607**, 254 (2005) [nucl-th/0409040].
- [175] W. P. Abfalterer, F. B. Bateman, F. S. Dietrich, R. W. Finlay, R. C. Haight and G. L. Morgan, *Phys. Rev. C* **63**, 044608 (2001).
- [176] K. Sekiguchi *et al.*, *Phys. Rev. C* **65**, 034003 (2002).
- [177] K. Ermisch *et al.*, *Phys. Rev. C* **68**, 051001 (2003).
- [178] A. Deltuva, A. C. Fonseca and P. U. Sauer, *Phys. Rev. C* **71**, 054005 (2005) [nucl-th/0503012].
- [179] L. Girlanda, A. Kievsky, M. Viviani and L. E. Marcucci, *Phys. Rev. C* **99**, no. 5, 054003 (2019) [arXiv:1811.09398 [nucl-th]].
- [180] C.R. Howell *et al.*, *Few. Body Syst.* **2** (1987) 19.

- [181] K. Sagara *et al.*, Phys. Rev. C **50** (1994) 576.
- [182] G. Rauprich *et al.*, Few. Body Syst. **5** (1988) 67.
- [183] F. Sperisen *et al.*, Nucl. Phys. **A422** (1984) 81.
- [184] A. Kievsky, Phys. Rev. C **60**, 034001 (1999) [nucl-th/9905045].
- [185] B. von Przewoski *et al.*, Phys. Rev. C **74**, 064003 (2006) [nucl-ex/0411019].
- [186] B. R. Barrett, P. Navratil and J. P. Vary, Prog. Part. Nucl. Phys. **69**, 131 (2013).
- [187] P. Maris, J. P. Vary and A. M. Shirokov, Phys. Rev. C **79**, 014308 (2009).
- [188] P. Maris and J. P. Vary, Int. J. Mod. Phys. E **22**, 1330016 (2013).
- [189] S. D. Glazek and K. G. Wilson, Phys. Rev. D **48**, 5863 (1993).
- [190] F. Wegner, Annalen der Physik 506, 77 (1994).
- [191] S. K. Bogner, R. J. Furnstahl, P. Maris, R. J. Perry, A. Schwenk and J. P. Vary, Nucl. Phys. A **801**, 21 (2008).
- [192] S. K. Bogner, R. J. Furnstahl and A. Schwenk, Prog. Part. Nucl. Phys. **65**, 94 (2010).
- [193] G. Hagen, T. Papenbrock, D. J. Dean, A. Schwenk, A. Nogga, M. Wloch and P. Piecuch, Phys. Rev. C **76**, 034302 (2007).
- [194] R. Roth, S. Binder, K. Vobig, A. Calci, J. Langhammer and P. Navratil, Phys. Rev. Lett. **109**, 052501 (2012).
- [195] S. Binder, P. Piecuch, A. Calci, J. Langhammer, P. Navratil and R. Roth, Phys. Rev. C **88**, no. 5, 054319 (2013).
- [196] S. R. Stroberg, A. Calci, H. Hergert, J. D. Holt, S. K. Bogner, R. Roth and A. Schwenk, Phys. Rev. Lett. **118**, no. 3, 032502 (2017).
- [197] S. R. Beane and M. J. Savage, Nucl. Phys. A **694**, 511 (2001) [nucl-th/0011067].
- [198] R. J. Furnstahl, H. W. Hammer and N. Tirfessa, Nucl. Phys. A **689**, 846 (2001) [nucl-th/0010078].
- [199] A. A. Filin, V. Baru, E. Epelbaum, H. Krebs, D. Möller and P. Reinert, arXiv:1911.04877 [nucl-th].
- [200] M. C. Birse, PoS CD **09**, 078 (2009) [arXiv:0909.4641 [nucl-th]].
- [201] J. Golak *et al.*, Eur. Phys. J. A **50**, 177 (2014) [arXiv:1410.0756 [nucl-th]].
- [202] C. Drischler, K. Hebeler and A. Schwenk, Phys. Rev. Lett. **122**, no. 4, 042501 (2019) [arXiv:1710.08220 [nucl-th]].
- [203] J. Hoppe, C. Drischler, K. Hebeler, A. Schwenk and J. Simonis, Phys. Rev. C **100**, no. 2, 024318 (2019) [arXiv:1904.12611 [nucl-th]].
- [204] T. H  ther, K. Vobig, K. Hebeler, R. Machleidt and R. Roth, arXiv:1911.04955 [nucl-th].
- [205] A. Baroni, L. Girlanda, A. Kievsky, L. E. Marcucci, R. Schiavilla and M. Viviani, Phys. Rev. C **94**, no. 2, 024003 (2016) Erratum: [Phys. Rev. C **95**, no. 5, 059902 (2017)] [arXiv:1605.01620 [nucl-th]].
- [206] A. Baroni *et al.*, Phys. Rev. C **98**, no. 4, 044003 (2018) [arXiv:1806.10245 [nucl-th]].